To Study Properties of a Known Procedure in Adaptive Sequential Sampling Design

Sampurna Kundu, Jayant Jha, Subir Kumar Bhandari

Interdisciplinary Statistical Research Unit, Indian Statistical Institute, Kolkata

sampurna.kundu58@gmail.com, jayantjha@gmail.com, subirkumar.bhandari@gmail.com

Abstract

We revisit the procedure proposed by Bhandari et al. (2009) in the context of twotreatment clinical trials, with the objective of minimizing the applications of a less effective drug to the least number of patients. Our focus is on an adaptive sequential procedure that is both simple and intuitive. Our findings show that the expected number of applications of the less effective drug remains finite. In contrast, Bhandari et al. (2009) observed that this number increases logarithmically with the total sample size. We attribute this discrepancy to differences in their choice of starting sample size and the method of analysis employed.

Keywords: Adaptive allocation, Average sample number, Composite hypothesis, Probability of incorrect inference, Number of applications of less effective drug.

1 Introduction

The field of adaptive sequential design has a long history of research, particularly in optimizing multiple objective functions. Berry and Fristedt (1985) in their book on Bandit Problems, explored Bayesian methods to achieve optimal designs in this context.

In conventional sequential analysis for the two-population problem, samples are drawn one-by-one at each stage from both populations, ensuring that the sample sizes remain equal. The primary objectives in this case are to make inferences about the population parameters with less error while minimizing the average sample number. In contrast, adaptive sampling allows for unequal sampling from the two populations at each stage (sometimes selecting a sample exactly from one population). The choice of which population to sample at each stage depends on the performance of previously collected data. This approach has a natural connection with the context of most studied clinical trials, where two treatments are applied sequentially to a series of patients. The goal in such trials is to balance the need for statistical accuracy with ethical considerations - specifically, applying the less effective treatment to a minimum number of patients.

In this context, Bhandari et al. (2007) considered the case with known parameters and concluded that the expected number of applications of the less effective drug is finite under an adaptive sequential design. However, where the parameters are unknown, Bhandari et al. (2009) found that the expected number of applications of the less effective drug grows logarithmically with the total sample size. This result was derived under the assumption of a large initial sample size.

In the present work, we modify the procedure proposed in Bhandari et al. (2009) and provide a more refined analysis to elucidate that the expected number of applications of the less effective drug remains finite.

2 Preliminaries

Let $\mathcal{X} = X_1, X_2, X_3, \ldots$ follow i.i.d. $N(\theta_0, \sigma_0^2)$ and $\mathcal{Y} = Y_1, Y_2, Y_3, \ldots$ follow i.i.d. $N(\theta_1, \sigma_1^2)$ and they be two independent data streams. We will use population 0 and population 1 for \mathcal{X} and \mathcal{Y} respectively. We draw samples adaptively, i.e., we draw samples sequentially, and at stage *n* after drawing a total of *n* samples, we define the following two count variables.

> $N'_{0,n} :=$ number of samples drawn from \mathcal{X} $N'_{1,n} :=$ number of samples drawn from \mathcal{Y} .

Let $\hat{\theta}_{0,n}$ and $\hat{\theta}_{1,n}$ denote the sample means at stage n for \mathcal{X} and \mathcal{Y} respectively. We adopt the following allocation rule:

- (i) If $\hat{\theta}_{0,n} \hat{\theta}_{1,n} > 0$, we increase $N'_{0,n}$ by 1
- (ii) If $\hat{\theta}_{0,n} \hat{\theta}_{1,n} < 0$, we increase $N'_{1,n}$ by 1

(iii) If $\hat{\theta}_{0,n} - \hat{\theta}_{1,n} = 0$, one unit observation is allocated to each population with probability $\frac{1}{2}$ each.

There is a total N number of patients. Finally, when n = N, we accept $H_0: \theta_0 > \theta_1$, with probability 1, if $\hat{\theta}_{0,N} - \hat{\theta}_{1,N} > 0$ and, with probability $\frac{1}{2}$, if $\hat{\theta}_{0,N} - \hat{\theta}_{1,N} = 0$.

Remark 1. The procedure described above was extensively studied by Bhandari et al. (2009), where it was referred to as Procedure III.

Remark 2. In the proof of the Main Result, we assume σ_0^2 and σ_1^2 to be unity and $\theta_0 > \theta_1$ without any loss of generality.

3 Main Result

Let us define the following:

 $N_{1,n} := \min\{N'_{1,n}, N'_{0,n}\}$ and $N_{1,N} := \min\{N'_{1,N}, N'_{0,N}\}.$

Theorem 3.1. We have the following:

- (i) $\lim_{N\to\infty} N_{1,N} < \infty$, i.e., $N_{1,N}$ is a finite random variable.
- (ii) The first four moments of $N_{1,N}$ are bounded (uniformly over N).

Remark 3. Bhandari et al. (2009) found that both $\frac{N'_{1,N}}{\log(N)}$ and $\frac{\mathbb{E}(N'_{1,N})}{\log(N)}$ tend to positive constants, contrary to our main result. This variation arises because they started with a large initial sample size and employed a different method of (mathematical) analysis.

Remark 4. Note that, $\lim_{N\to\infty} N_{1,N} \neq \lim_{N\to\infty} N'_{1,N}$ if and only if there is incorrect inference about $H_0: \theta_0 > \theta_1$. The probability of incorrect inference is negligible and it is less than or equal to

$$\Phi\left(-\left(\theta_0-\theta_1+\varepsilon\right)\cdot N_{1,N}\right)<\Phi\left(-\left(\theta_0-\theta_1+\varepsilon\right)\cdot M\right),$$

for some $\varepsilon > 0$, where M is a finite natural number. We start with an initial sample size of M from each population at the beginning of the procedure.

4 Proof of Main Result

The following two results, as stated in Chapter 1, Section 9 of Billingsley (1995), are presented here without proof.

Result 4.1. (Law of Iterated Logarithm) Let $\{X_n\}$ be i.i.d. random variables with mean zero and unit variance. Let, $S_n = X_1 + X_2 + X_3 + \ldots + X_n$. Then,

$$\limsup_{n \to \infty} \frac{|S_n|}{\sqrt{2n \log \log n}} = 1 \quad almost \ surely.$$

Result 4.2. (Chernoff's Theorem in the context of Normal Distribution) Let $\{X_n\}$ be the *i.i.d.* random variables following $N(\mu, 1)$ with $\mu < 0$, with MGF M(t). If $\rho = \min_t M(t)$, then, $\rho < 1$ and,

$$\lim_{n \to \infty} \frac{1}{n} \log \left[\mathbb{P} \left(X_1 + X_2 + X_3 + \ldots + X_n \ge 0 \right) \right] = \log(\rho),$$

which implies, $\mathbb{P}(\bar{X}_n > 0) < C \cdot \rho^n \forall n$, for some constant C (> 0),

where,
$$\bar{X}_n = \frac{1}{n} (X_1 + X_2 + X_3 + \ldots + X_n).$$

Using Result 4.1, we find that for any given $\varepsilon > 0$, there exist a natural number N_{ε}^* , and subset of sample space A, with $\mathbb{P}(A) > 1 - \varepsilon$, such that,

$$\forall \ \omega \in A, \quad |\hat{\theta}_{0,n}(\omega) - \theta_0| < \varepsilon \text{ and } |\hat{\theta}_{1,n}(\omega) - \theta_1| < \varepsilon, \text{ (for all } n > N_{\varepsilon}^*).$$

Using the above, we find that with probability $\geq (1 - \varepsilon)$,

 $\lim_{N \to \infty} N_{1,N} < N_{\varepsilon}^*$, i.e., $N_{1,N}$ is a finite random variable.

This proves part (i) of Theorem 3.1.

Let us define

$$M_{\varepsilon}^* = \inf\left\{k : \sup_{n \ge k} \bar{Z}_n \le \varepsilon\right\},\$$

where, $Z_1, Z_2, Z_3, \ldots, Z_n$ are i.i.d. N(0, 1) and $\overline{Z}_n = \frac{1}{n} \sum_{i=1}^n Z_i$. Define $X_i = Z_i - \varepsilon$ for small $\varepsilon > 0$, $i = 1, 2, 3, \ldots$ Then, Result 4.2 holds for this sequence of X_i .

The proof of part (ii) of Theorem 3.1 will be complete upon proving Lemma 4.1.

Lemma 4.1. The first four moments of the random variable M_{ε}^* are finite.

Proof. We will prove the finiteness of the first moment only, i.e., $\mathbb{E}(M_{\varepsilon}^*) < \infty$, as the proofs for the remaining three moments are analogous.

$$\mathbb{P}(M_{\varepsilon}^{*} > k) = \mathbb{P}\left[\sup_{n \ge k} \bar{Z}_{n} > \varepsilon\right]$$

$$\leq \mathbb{P}\left[\bigcup_{n \ge k} \left(\bar{Z}_{n} \ge \varepsilon\right)\right]$$

$$= \sum_{n=k}^{\infty} \mathbb{P}\left(\bar{X}_{n} \ge 0\right) \text{ [since } \bar{X}_{n} = \bar{Z}_{n} - \varepsilon\text{]}$$

$$\leq \sum_{n=k}^{\infty} C \cdot \rho^{n} \quad \text{[using Result 4.2]}$$

$$= \frac{C \cdot \rho^{k}}{1 - \rho} \quad \text{[this holds for large } k\text{]}$$

Hence, for large k, $\mathbb{E}(M_{\varepsilon}^*) \leq k + \frac{C}{(1-\rho)^2}\rho^k < \infty$.

5 Conclusion

In this study, we have shown that, with correct inference, the number of applications of the less effective drug can be kept finite. Given that the procedure is both simple and natural, we expect it to be extendable to other response distributions, which could have meaningful practical implications.

Although Chernoff's Theorem has been applied here in the context of the Normal distribution, it is applicable to a broader range of response distributions. Therefore, we anticipate that our results will hold for other distributions as well.

References

- Berry, D. A. and Fristedt, B. (1985). *Bandit Problems: Sequential Allocation of Experiments*. Chapman and Hall, London.
- Bhandari, S. K., De, S. K., Mandal, S., Pradhan, S., and Ghosh, B. (2009). Study of optimal adaptive rule in testing composite hypothesis. *Sequential Analysis*, 28(3):394– 405. https://doi.org/10.1080/07474940903041852.

Bhandari, S. K., Dutta, R., and Niyogi, R. G. (2007). Study of optimal adaptive rule in testing problem. In *Advances in Multivariate Statistical Methods*. World Scientific.

Billingsley, P. (1995). Probability and Measure. Wiley.