A mixing time bound for Gibbs sampling from log-smooth log-concave distributions

Neha S. Wadia*

Center for Computational Mathematics, Flatiron Institute, New York NY 10010, USA

December 25, 2024

Abstract

The Gibbs sampler, also known as the coordinate hit-and-run algorithm, is a Markov chain that is widely used to draw samples from probability distributions in arbitrary dimensions. At each iteration of the algorithm, a randomly selected coordinate is resampled from the distribution that results from conditioning on all the other coordinates. We study the behavior of the Gibbs sampler on the class of log-smooth and strongly log-concave target distributions supported on \mathbb{R}^n . Assuming the initial distribution is *M*-warm with respect to the target, we show that the Gibbs sampler requires at most $O^{\star}\left(\kappa^2 n^{7.5}\left(\max\left\{1,\sqrt{\frac{1}{n}\log\frac{2M}{\gamma}}\right\}\right)^2\right)$ steps to produce a sample with error no more than γ in total variation distance from a distribution with condition number κ .

1 Introduction

Sampling from probability distributions in high dimensional spaces is a fundamental computational primitive; it forms the basis of efficient numerical methods for approximating arbitrary integrals. The problem statement is the following: given a density function π , compute a point x with density proportional to $\pi(x)$.

A general approach to solving this problem is to design a reversible, ergodic Markov chain with a unique stationary distribution that is equal to the target distribution from which samples are needed. It is often possible to design relatively simple chains with low per-iteration computational complexity that are fit for purpose by implementing the Metropolis-Hastings filter [1, 2], a rule by which to either accept the next step in the dynamics or remain put and so tailor the dynamics toward a specific stationary distribution. The resulting *Metropolized* or *Markov Chain Monte Carlo* algorithms are known to converge asymptotically to their stationary distributions under mild regularity conditions. Non-asymptotic rates of convergence or *mixing times* are comparatively few in number and are both algorithm- and target-specific. They are important because downstream estimators computed using samples drawn from a dynamics that has not converged will suffer from bias.

The class of log-concave target distributions is of particular interest. These are distributions with density functions of the form $e^{-f(x)}$, $x \in \mathbb{R}^n$, where f is convex in x. The mixing times of a number of Markov chains are polynomial in n for this class of target distributions. Such chains are considered to be *rapidly mixing*. Examples are the ball walk [3, 4], the grid walk [5, 6], and the hit-and-run walk [7, 8, 9]. An expository survey of the associated results may be found in [10]. In

^{*}neha.wadia@berkeley.edu

this paper, we upper bound the mixing time of the Gibbs sampler, also known as coordinate hitand-run (CHAR), for sampling from the class of log-smooth and strongly log-concave distributions supported on \mathbb{R}^n . That is, in addition to convexity we require f to be such that the spectrum of its Hessian is uniformly bounded both above and below. We show that the Gibbs sampler also mixes rapidly on this class of distributions.

At each iteration of the dynamics of the Gibbs sampler, a single coordinate x_i of x is chosen uniformly at random and replaced with a sample from its conditional distribution, i.e., the onedimensional distribution over x_i that results from conditioning on all the other n-1 coordinates. This dynamics can be interpreted as a limiting case of a Metropolized algorithm in which every move is accepted. An advantage of the Gibbs sampler is that it reduces the *n*-dimensional sampling problem to a one-dimensional problem at each iteration. Another advantage is that it has no hyperparameters to tune; in particular, there is no notion of a step size in the dynamics.

The Gibbs sampler has been in practice long enough to have been reintroduced to the literature several times and is difficult to trace accurately. It was introduced in the theoretical computer science literature by Turchin in 1971 [11], in the image processing literature by Geman and Geman in 1984 [12], and popularized in the statistics literature by Gelfand and Smith, who demonstrated that it could be implemented efficiently on a wide variety of problems of interest in statistics at the time, in 1990 [13]. Long before this, it was being used in statistical physics to sample from models of magnetic materials.

An important proof technique for establishing mixing time bounds relies on the following two ingredients: a result that gives the dependence of the mixing time on a quantity called the *conductance*, which measures how well or poorly the dynamics explores the state space, and an isoperimetric inequality that enables a lower bound on the conductance. Isoperimetric inequalities that are useful in this context are dynamics-dependent. The fact that the Gibbs sampler is constrained to move only along the *n* coordinate directions prevents the use of a general technique (the localization lemma [14]) for proving the relevant isoperimetric inequality. Until recently, this was a barrier to establishing mixing times for target distributions supported on continuous state spaces. (The case in discrete state spaces, where the Gibbs sampler is often termed "Glauber dynamics", is different, [15].)

Recently, Laddha and Vempala [16] and Narayanan and Srivastava [17] concurrently gave bounds on the mixing time of the Gibbs sampler for sampling from a uniform distribution supported on a convex body K in \mathbb{R}^n from a warm start. Laddha and Vempala prove an isoperimetric theorem using a first-principles approach. They obtain a bound of $O^*(R^2n^8)$ on the mixing time where R^2 is the expected squared distance of a random point from the centroid of K. The notation O^* suppresses dependence on logarithmic factors and other problem parameters. Narayanan and Srivastava instead establish a mixing time bound for an auxiliary dynamics that they show dominates the mixing time of the Gibbs sampler. They give a bound of $O^*(R_1^4n^7)$, where R_1 is the smallest number such that K is sandwiched between an L_{∞} ball and an L_{∞} ball scaled by R_1 . Narayanan, Rajaraman, and Srivastava [18] have since also demonstrated that the Gibbs sampler mixes in polynomial time for the same class of target distributions from a cold start.

We give an upper bound of $O^*(\kappa^2 n^{7.5})$ on the mixing time for Gibbs sampling from log-smooth and strongly log-concave distributions supported on \mathbb{R}^n from a warm start. κ is the ratio of the upper and lower bounds on the spectrum of the Hessian of f. The following theorem is our main contribution:

Theorem 1.1. Consider a probability distribution supported on \mathbb{R}^n with density function proportional to $e^{-f(x)}$ where f is μ -strongly convex in x and has L-Lipschitz gradients. Let $\kappa = L/\mu$. Let π^k be the distribution of the k^{th} iterate produced by the Gibbs sampler. For some universal constant C and any $\gamma \in (0, 1/2)$, if the initial distribution π^0 is M-warm with respect to π , then the total variation distance between π^{τ} and π is no more than γ when τ is no more than

$$C\kappa^2 n^{7.5} \log^2 n \left(\max\left\{ 1, \sqrt{\frac{1}{n} \log \frac{2M}{\gamma}} \right\} \right)^2 \log \frac{2M}{\gamma}.$$
⁽¹⁾

The difference of 1/2 in the exponent of n between (1) and the Laddha and Vempala result is due to the difference in concentration behavior between strongly log-concave distributions and the uniform distribution, which is log-concave but not strongly log-concave.

To prove Theorem 1.1, we establish a conductance bound in a high-probability region of \mathbb{R}^n , specifically, a mode-centered Euclidean ball of sufficiently large radius. Building on the proof technique of Laddha and Vempala, we prove an isoperimetric inequality for subsets of the Euclidean ball. The crucial insight that enables us to do so is the fact that smooth functions are approximately uniform on small-enough domains. In order to state the isoperimetric inequality, we will need the following definition.

Definition 1 (Axis-disjoint sets). Two sets S_1 and S_2 are axis-disjoint if, for every pair of vectors x, y where $x \in S_1$ and $y \in S_2$, $|\{i \in [n] \ s.t. \ x_i \neq y_i\}| \ge 2$.

The Gibbs sampler cannot transition between two axis-disjoint sets in a single iteration. The following isoperimetric inequality gives a worst-case lower bound on the measure of the region of the state space that *is* accessible to the dynamics from either of two axis-disjoint regions, and is the main technical contribution of this paper.

Lemma 1.2. Consider a probability density function $\pi(x) = Z^{-1}e^{-f(x)}$, $x \in \mathbb{R}^n$, where f is μ -strongly convex in x and has L-Lipschitz gradients. Let Π be the associated probability measure. Let $x^* = \arg \max_x \pi(x)$. For $\varepsilon \in (0, 1/2)$, define the function

$$r(\varepsilon) = 2 + 2 \max\left\{ \left(\frac{1}{n}\log\frac{1}{\varepsilon}\right)^{1/4}, \left(\frac{1}{n}\log\frac{1}{\varepsilon}\right)^{1/2} \right\}.$$

Consider a Euclidean ball K centered at x^* with radius strictly greater than $r(\varepsilon)\sqrt{n/\mu}$ and no larger than twice that quantity. Let $S_1 \cup S_2 \cup S_3$ be a partition of K such that S_1 and S_2 are axis-disjoint. Then there is a positive quantity Ψ such that

$$\Pi(S_3) \ge \Psi \min\left\{\frac{\Pi(S_1)}{5} - \varepsilon, \frac{\Pi(S_2)}{5} - \varepsilon\right\},\tag{2}$$

where

$$\Psi \ge C \frac{1}{\kappa n^{2+\frac{3}{4}} \log n \max\left\{1, \sqrt{\frac{1}{n} \log \frac{1}{\varepsilon}}\right\}} \tag{3}$$

for some universal constant C.

In independent and contemporaneous work, Ascolani, Lavenant, and Zanella proved a periteration contraction of the Kullback-Leibler divergence between the law of the sampler and the target under the dynamics of the Gibbs sampler for the same class of target distributions as in Theorem 1.1 [19]. Their analysis implies linear dependence of the mixing time on both the condition number and n. For several technical reasons, it is unclear how to directly compare this with Theorem 1.1. One of these is that the proof of Theorem 1.1 fundamentally treats the sampler as being constrained to a convex subset of \mathbb{R}^n . This is not the case in [19]. We leave a careful comparison of the two types of results as an interesting open problem.

The remainder of the paper is devoted to the proofs of Theorem 1.1 and Lemma 1.2, followed by a discussion.

2 Preliminaries

Denote by $\pi(x) : \mathbb{R}^n \to \mathbb{R}_+$ the normalized density function of a distribution that can be written in the form $e^{-f(x)}/Z$ where f is μ -strongly convex and L-smooth. That is, for any $x, y \in \mathbb{R}^n$ and some $L \ge \mu > 0$, f satisfies the following inequalities:

$$f(y) \ge f(x) + \langle \nabla f(x), y - x \rangle + \frac{\mu}{2} ||x - y||^2, \tag{4}$$

$$f(y) \le f(x) + \langle \nabla f(x), y - x \rangle + \frac{L}{2} ||x - y||^2.$$
(5)

Z is the partition function $\int_{\mathbb{R}^n} e^{-f(x)} dx$. π is absolutely continuous with respect to the Lebesgue measure, so we can and will safely replace $d\pi(x)$ with $\pi(x)dx$ in all integrals.

Let $x^* = \arg \max_{x \in \mathbb{R}^n} \pi(x)$ be the mode of π . Define the condition number $\kappa = L/\mu$.

We write Π for the probability measure associated with π , and Π_{n-1} for the measure induced by Π on an n-1-dimensional set. We write vol for the Lebesgue measure of a set.

Superscripts on x indicate iteration number and subscripts index vector components. [n] denotes the set of natural numbers up to and including n. Let e_i , $i \in [n]$, denote a unit vector in the *i*th coordinate direction. We use the notation x_{-i} to represent the n-1-dimensional vector that results from leaving out the *i*th coordinate of x.

We are now ready to formally describe the algorithm.

Algorithm 1 Gibbs sampler

Given $x^0 \in \mathbb{R}^n$, $\tau \in \mathbb{N}$: for t=1... τ do Choose $i \in [n]$ uniformly at random. Draw a sample y from $\pi(x_i|x_{-i})$. $x_i^t \to y$. end for

The stationary distribution of Algorithm 1 is π .

In preparation for the proof of Theorem 1.1, we give some basic facts about discrete-time Markov chains in continuous state spaces.

A discrete-time Markov chain on \mathbb{R}^n is specified by an initial state $x^0 \in \mathbb{R}^n$ and a conditional distribution $P_x(A)$. For any measurable subset A of \mathbb{R}^n , $P_x(A)$ gives the probability of transitioning to a state $y \in A$ given that the chain is currently in the state x. A distribution π is *stationary* for this chain if $P_x(A)$ is given by $\Pi(A)$ when x is distributed according to π :

$$\int_{\mathbb{R}^n} P_x(A)\pi(x)dx = \Pi(A).$$
(6)

A Markov chain is *reversible* if the probability of a transition between two states is equally likely in both directions. That is, for any two measurable sets A, B, we must have

$$\int_{B} P_x(A)\pi(x)dx = \int_{A} P_x(B)\pi(x)dx.$$
(7)

Algorithm 1 is reversible.

The ergodic flow p(A) of a set A measures how likely the dynamics is to exit it in a single step:

$$p(A) = \int_{A} P_x \left(\mathbb{R}^n \backslash A \right) \pi(x) dx.$$
(8)

The ratio of the ergodic flow of a set to either its measure or the measure of its complement, whichever is smaller, is its *conductance* $\phi(A)$:

$$\phi(A) = \frac{p(A)}{\min\{\Pi(A), \Pi(\mathbb{R}^n \setminus A)\}}.$$
(9)

The conductance ϕ of the chain is the infimum over all measurable sets of their conductances. It is convenient to restrict to sets of measure not exceeding a half so we may drop the minimum in (9). Then the conductance is given by the following expression:

$$\phi \equiv \inf_{A: \Pi(A) \le 1/2} \frac{p(A)}{\Pi(A)}.$$
(10)

The conductance quantifies bottlenecks to the dynamics. If it is small, there is at least one region of the state space that is not easy to escape from, limiting the ability of the sampler to explore the state space. In what follows, we will work with a weakening of the definition of the conductance, called the *s*-conductance, which allows us to neglect sets of measure no more than s in the infimum in (10). For any 0 < s < 1/2, we have

$$\phi_s \equiv \inf_{A: s < \Pi(A) \le 1/2} \frac{p(A)}{\Pi(A) - s}.$$
(11)

At any specific iteration number t, the current state of the chain, x^t , is distributed according to some distribution π^t . We will refer to π^t as the law of the sampler at time t. We use the total variation distance to measure the distance between the law of the sampler and its stationary distribution, defined as follows:

$$d_{TV}(\pi^{t},\pi) = \sup_{A} |\pi^{t}(A) - \pi(A)|.$$
(12)

The supremum in (12) is taken over all measurable sets.

The *mixing time* of the chain is the smallest number of iterations $\tau(\epsilon)$ needed to drive this distance below some small ϵ :

$$\tau(\epsilon) \equiv \inf\{t \in \mathbb{N} \ s.t. \ d_{TV}(\pi^t, \pi) \le \epsilon\}.$$
(13)

We will work with a *lazy* version of Algorithm 1, where at every iteration, with probability 1/2 we do nothing. This ensures that π is the *unique* stationary distribution for Algorithm 1.

The last basic notion we shall need is one that characterizes the quality of the initial distribution π^0 . π^0 is a warm start for π if, for some M > 0,

$$\sup_{A} \frac{\pi^0(A)}{\pi(A)} \le M,\tag{14}$$

where the supremum is taken over all measurable sets. A result due to Altschuler and Chewi [20] guarantees that a warm start to any log-concave distribution can be computed in $O(\sqrt{n})$ iterations. Since this is dwarfed by the mixing time in Theorem 1.1, the warm start is a mild assumption to make for Gibbs sampling from log-concave distributions.

Our goal is to upper bound $\tau(\epsilon)$ for Algorithm 1. In particular, we are interested in identifying the *n*-dependence of $\tau(\epsilon)$. To do so, we will lower bound the s-conductance and then invert the following result due to Lovász and Simonovits [21] to upper bound the mixing time.

Theorem 2.1. Consider a lazy, reversible Markov chain. Let $s \in (0, 1/2)$. If the initial distribution is M-warm with respect to the stationary distribution, then after t iterations,

$$d_{TV}(\pi^t, \pi) \le Ms + M\left(1 - \frac{\phi_s^2}{2}\right)^t.$$
 (15)

Theorem 2.1 is itself a generalization to arbitrary state spaces of a result due to Jerrum and Sinclair [22]. Both results are related to Cheeger's inequality [23] in differential geometry (see [24] for a treatment of the relationship). The presence of the warmness parameter M in (15) is the price to be paid for working not with the conductance but with the s-conductance.

3 Upper bounding the mixing time

The proof of Theorem 1.1, presented in this section, proceeds via a lower bound on the s-conductance of the Gibbs sampler inside a convex high-probability region within which the gradient norm of f can be bounded. The argument illustrates the use of Lemma 1.2, which is proved in the following section.

See Figure 1 for an illustration of the overall construction.

We will need the following concentration result for strongly log-concave distributions.

Lemma 3.1. Let Π be a μ -strongly log-concave measure on \mathbb{R}^n and π the associated density. Let $x^* = \arg \max_{x \in \mathbb{R}^n} \pi(x)$. Let $\mathcal{B}(x, r)$ denote a Euclidean ball of radius r centered at x. For any $\varepsilon \in (0, 1/2)$ and

$$r(\varepsilon) = 2 + 2\max\left\{\left(\frac{1}{n}\log\frac{1}{\varepsilon}\right)^{1/4}, \left(\frac{1}{n}\log\frac{1}{\varepsilon}\right)^{1/2}\right\},\tag{16}$$

we have

$$\Pi\left(\mathcal{B}\left(x^{\star}, r(\varepsilon)\sqrt{\frac{n}{\mu}}\right)\right) \ge 1 - \varepsilon.$$
(17)

This lemma is due to Dwivedi, Chen, Wainwright, and Yu [25].

Let $s \in (0, 1/2)$. Take $\varepsilon = s/11$ in Lemma 3.1, and consider a Euclidean ball K centered ar x^* with radius R such that

$$r\left(\frac{s}{11}\right)\sqrt{\frac{n}{\mu}} < R \le 2r\left(\frac{s}{11}\right)\sqrt{\frac{n}{\mu}},\tag{18}$$

where r is the function defined in (16). (The reason for defining R thus via two inequalities will be made clear in the proof of Lemma 1.2.) Then by Lemma 3.1, $\Pi(K) > 1-s/11$ and $\Pi(\mathbb{R}^n \setminus K) < s/11$.

The following lemma enables us to construct a pair of axis-disjoint subsets of \mathbb{R}^n .

Lemma 3.2. Let $A_1 \cup A_2$ be a partition of \mathbb{R}^n , and let $A'_1 = \{x \in A_1 \text{ s.t. } P_x(A_2) < \frac{1}{2n}\}$ and $A'_2 = \{x \in A_2 \text{ s.t. } P_x(A_1) < \frac{1}{2n}\}$. Then A'_1 and A'_2 are axis-disjoint.

Proof. The proof is by contradiction and relies on the fact that the probability of finding x^{t+1} along each of the *n* lines that pass through x^t and run parallel to the coordinate axes is 1/n.

Assume that A'_1 and A'_2 are not axis-disjoint. Then there must be a line running parallel to one of the coordinate axes that passes through both sets. Without loss of generality, let this line, call it ℓ_j , run along the coordinate axis e_j . Consider a point $y \in \ell_j \cap A'_1$ and a point $z \in \ell_j \cap A'_2$. By construction, we have

$$P_y(\ell_j \cap A_2) \le P_y(A_2) < \frac{1}{2n}$$
, and $P_z(\ell_j \cap A_1) \le P_z(A_1) < \frac{1}{2n}$.

This implies

$$P_y(\ell_j \cap A_2) + P_z(\ell_j \cap A_1) < \frac{1}{n}.$$

Furthermore,

$$P_y(\ell_j \cap A_2) + P_z(\ell_j \cap A_1) = \Pi(\ell_j \cap A_2 | y_{-j}) + \Pi(\ell_j \cap A_1 | z_{-j}) = \Pi(\ell_j) = 1/n$$

In the second equality we have used the fact that conditioning on z_{-j} is equivalent to conditioning on y_{-j} because y and z differ only in their j^{th} coordinate. Thus we arrive at the contradiction 1 < 1.



Figure 1: Illustrated here is a partition $A_1 \cup A_2$ of \mathbb{R}^2 such that for some $s \in (0, 1/2)$, $s < \Pi(A_1) \le 1/2$. The Euclidean ball K is centered at the mode x^* of π and is large enough so that its measure differs from unity by a fraction of s. A'_1 (in light pink) and A'_2 (in light blue) are subsets of A_1 and A_2 , respectively. A'_1 and A'_2 are axis disjoint. $S_1 \cup S_2 \cup S_3$ is a partition of K such that $S_i = K \cap A'_i$ for i = 1, 2. $\Pi(S_1) \le \Pi(S_2)$.

Consider a partition $A_1 \cup A_2$ of \mathbb{R}^n such that $s < \Pi(A_1) \le 1/2$, and subsets A'_1 and A'_2 defined as in Lemma 3.2. A'_1 and A'_2 are axis-disjoint. We have

$$\Pi(A_i) = \Pi(A_i \setminus K) + \Pi(A_i \cap K) < \frac{s}{11} + \Pi(A_i \cap K),$$

$$\Rightarrow \Pi(A_i \cap K) > \Pi(A_i) - \frac{s}{11}$$
(19)

for i = 1, 2.

Let $S_i = K \cap A'_i$ for i = 1, 2 and $S_3 = K \setminus S_1 \setminus S_2$. S_1 and S_2 are axis-disjoint. We write down two lower bounds on the ergodic flow of A_1 .

$$p(A_{1}) = \int_{A_{1}} P_{x}(A_{2}) \pi(x) dx$$

= $\int_{A_{1}\setminus K} P_{x}(A_{2}) \pi(x) dx + \int_{S_{1}} P_{x}(A_{2}) \pi(x) dx + \int_{A_{1}\cap K\setminus S_{1}} P_{x}(A_{2}) \pi(x) dx$
$$\geq \int_{A_{1}\cap K\setminus S_{1}} P_{x}(A_{2}) \pi(x) dx$$

$$\geq \frac{1}{2n} \Pi(A_{1}\cap K\setminus S_{1}).$$
 (20)

In the fourth line we have applied the definition of A'_1 . Similarly,

$$p(A_1) = \int_{A_1} P_x(A_2) d\pi(x)$$

$$= \int_{A_1} \left[P_x(A_2 \setminus K) + P_x(S_2) + P_x(A_2 \cap K \setminus S_2) \right] \pi(x) dx$$

$$\geq \int_{A_1} P_x(A_2 \cap K \setminus S_2) \pi(x) dx$$

$$= \int_{A_2 \cap K \setminus S_2} P_x(A_1) \pi(x) dx$$

$$\geq \frac{1}{2n} \Pi(A_2 \cap K \setminus S_2).$$
(21)

In the penultimate line above, we have applied the reversibility property (7) of Algorithm 1. In the last line, we have applied the definition of A'_2 .

There are two cases to consider depending on the relative sizes of S_i in $A_i \cap K$, i = 1, 2.

Case 1. If $\Pi(S_1) < \Pi(A_1 \cap K)/2$, then $\Pi(A_1 \cap K \setminus S_1) \ge \Pi(A_1 \cap K)/2$. From (20) we have

$$p(A_1) \ge \frac{1}{2n} \Pi(A_1 \cap K \setminus S_1) \ge \frac{1}{4n} \Pi(A_1 \cap K) > \frac{1}{4n} \left(\Pi(A_1) - \frac{s}{11} \right) > \frac{1}{4n} \left(\Pi(A_1) - s \right).$$
(22)

Similarly, if $\Pi(S_2) < \Pi(A_2 \cap K)/2$, then $\Pi(A_2 \cap K \setminus S_2) \ge \Pi(A_2 \cap K)/2$, and (21) gives the bound

$$p(A_1) \ge \frac{1}{2n} \Pi(A_2 \cap K \setminus S_2) \ge \frac{1}{4n} \Pi(A_2 \cap K) > \frac{1}{4n} \left(\Pi(A_2) - \frac{s}{11} \right) \ge \frac{1}{4n} \left(\Pi(A_1) - \frac{s}{11} \right) \ge \frac{1}{4n} \left(\Pi(A_1) - s \right)$$
(23)

where the penultimate inequality follows by construction.

Thus if either either S_1 or S_2 is small with respect to $A_1 \cap K$ or $A_2 \cap K$, respectively, the ergodic flow of A_1 is lower bounded by $\frac{1}{4n} (\Pi(A_1) - s)$.

Case 2. When both S_1 and S_2 are large in measure with respect to $A_1 \cap K$ and $A_2 \cap K$, respectively, Lemma 1.2 is needed to lower bound the measure of $K \setminus S_1 \setminus S_2$.

In particular, if $\Pi(S_1) \ge \Pi(A_1 \cap K)/2$ and $\Pi(S_2) \ge \Pi(A_2 \cap K)/2$, then, summing (20) and (21), we have

$$p(A_1) > \frac{1}{2} \left(\frac{1}{2n} \Pi(A_1 \cap K \setminus S_1) + \frac{1}{2n} \Pi(A_2 \cap K \setminus S_2) \right)$$

= $\frac{1}{4n} \Pi(S_3)$
> $\frac{\Psi}{4n} \left(\frac{\Pi(S_1)}{5} - \frac{s}{11} \right) \ge \frac{\Psi}{4n} \left(\frac{1}{5} \frac{\Pi(A_1 \cap K)}{2} - \frac{s}{11} \right) \ge \frac{\Psi}{40n} \left(\Pi(A_1) - s \right).$ (24)

In the third line we have applied Lemma 1.2 with $\varepsilon = s/11$.

This concludes discussion of Case 2.

We note that

$$\min\left\{\frac{1}{4n}, \frac{\Psi}{40n}\right\} = \frac{\Psi}{40n} \quad \forall n \ge 2.$$
(25)

Thus for any set $A_1 \subset \mathbb{R}^n$ such that $s < \Pi(A_1) \le 1/2$ for $s \in (0, 1/2)$,

$$\frac{p(A_1)}{\Pi(A_1) - s} > \frac{\Psi}{40n}.$$
(26)

This implies the following lower bound on the the s-conductance of the Gibbs sampler:

$$\phi_s > \frac{\Psi}{40n}.\tag{27}$$

We are now ready to prove Theorem 1.1.

Proof of Theorem 1.1. Making the assignment $s = \gamma/2M$ in (15), we have

$$d_{TV}(\pi^{\tau},\pi) \le \frac{\gamma}{2} + M\left(1 - \frac{\phi_s^2}{2}\right)^{\tau} \le \frac{\gamma}{2} + Me^{-\tau\phi_s^2/2}.$$
(28)

To ensure $d_{TV}(\pi^{\tau},\pi) \leq \gamma$, it is therefore enough to choose τ such that

$$2Me^{-\tau\phi_s^2/2} = \gamma \Rightarrow \tau = \frac{2}{\phi_s^2}\log\frac{2M}{\gamma}.$$
(29)

Applying the lower bound (27) to ϕ_s , we have

$$\tau < 2^5 10^2 \frac{n^2}{\Psi^2} \log \frac{2M}{\gamma}.$$
(30)

4 Isoperimetry

In this section we prove Lemma 1.2. The proof is bipartite. In the first part, we write down an isoperimetric inequality on an *n*-dimensional cube of side δ contained in *K*. This inequality relies on a similar result proved by Laddha and Vempala for a uniform distribution on the cube, and on the fact that smooth functions are approximately uniform on small domains. δ must be chosen carefully to ensure a good approximation. In the second part, we tile S_1 (see Figure 1) with these cubes to establish the lower bound on Ψ .

The proof of Lemma 1.2 is given in Section 4.2. In Section 4.1 we develop the necessary approximation results.

4.1 Approximation theory

K is a Euclidean ball centered at x^* . For $\varepsilon \in (0, 1/2)$, the radius of K is lower and upper bounded as follows:

$$r(\varepsilon)\sqrt{\frac{n}{\mu}} < R \le 2r(\varepsilon)\sqrt{\frac{n}{\mu}},\tag{31}$$

where r is the function defined in Lemma 3.1.

We work with cubes that are *axis-aligned*:

Definition 2 (Axis-aligned cubes.). An axis-aligned cube of side δ is the set $\{x \in \mathbb{R}^n : ||x-x_0||_{\infty} \leq \delta\}$, where x_0 is any reference point in the cube.

4.1.1 Control of f in a cube

Consider an *n*-dimensional axis-aligned cube $C \subset K$ of side δ .

For any two points $x, y \in C$, Taylor's theorem guarantees the existence of a point z, also in C, such that

$$f(y) = f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2}(y - x)^{\top} \nabla^2 f(z)(y - x).$$

Rearranging and applying the triangle inequality, we have

$$|f(y) - f(x)| \le ||\nabla f(x)|| \ ||y - x|| + \frac{1}{2} ||\nabla^2 f(z)||_{\text{op}} \ ||y - x||^2.$$
(32)

The maximal Euclidean distance between any two points in C is bounded above by $\delta\sqrt{n}$. Smoothness guarantees that the operator norm of $\nabla^2 f(z)$ is bounded by L for any $z \in \mathbb{R}^n$. Moreover, for any $x \in K$,

$$||\nabla f(x)|| = ||\nabla f(x) - \nabla f(x^{\star})|| \le L||x - x^{\star}|| \le 2Lr(\varepsilon)\sqrt{\frac{n}{\mu}},\tag{33}$$

where $\varepsilon \in (0, 1/2)$. The last inequality follows from the upper bound (31) on the radius of K. Thus we arrive at the bound

$$|f(y) - f(x)| \le 2\frac{L}{\sqrt{\mu}}\delta r(\varepsilon)n + \frac{1}{2}L\delta^2 n.$$
(34)

In order to control the right-hand side of (34), we will need δ to scale inversely in n and L with the appropriate powers. We note that

$$\frac{r(\varepsilon)}{\max\left\{1, \sqrt{\frac{1}{n}\log\frac{1}{\varepsilon}}\right\}} \le 4.$$
(35)

For some positive constant c and function $\sigma > 0$, we make the assignment

$$\delta = \left(8c\sqrt{\kappa L}n^{1+\sigma}\max\left\{1,\sqrt{\frac{1}{n}\log\frac{1}{\varepsilon}}\right\}\right)^{-1},\tag{36}$$

in terms of which we have the bound

$$|f(y) - f(x)| \le \frac{1}{cn^{\sigma}} + \frac{1}{128c^{2}\kappa n^{1+2\sigma}} \left(\max\left\{1, \sqrt{\frac{1}{n}\log\frac{1}{\varepsilon}}\right\} \right)^{2} \le \frac{1}{cn^{\sigma}} + \frac{1}{128c^{2}n^{1+2\sigma}}.$$
 (37)

 σ and c must be large enough to ensure that |f(y) - f(x)| is both small $\forall n \geq 2$ and goes to zero as $n \to \infty$, but no larger than necessary because the final mixing time bound will have $c^2 n^{-2\sigma}$ dependence. We will find it useful to enforce the specific constraint

$$|f(y) - f(x)| \le \log \frac{6}{5} \quad \forall n \ge 2$$

Taking $\sigma = \frac{\log \log n}{\log n}$ and c = 8 is enough to do so. We finally arrive at the bound

$$|f(y) - f(x)| \le \frac{1}{8\log n} + \frac{1}{2^{13}n\log n} \equiv \nu(n) \le \log \frac{6}{5} \ \forall n \ge 2.$$
(38)

for the difference in f between any two points in C. We have defined the function $\nu(n)$ for the sake of notational convenience. We will omit writing its n-dependence in the sequel.

(38) enables the following bound on how much the ratio $\pi(x)/\pi(y)$ deviates from unity within C.

Fact 4.1. For any $x, y \in C \subset K$,

$$\left| e^{-(f(x) - f(y))} - 1 \right| \le e^{\nu} - 1.$$
 (39)

Proof. If $f(x) \ge f(y)$, then (38) implies $e^{f(x)-f(y)} \le e^{\nu}$ and therefore

$$\left| e^{-(f(x) - f(y))} - 1 \right| = 1 - e^{-(f(x) - f(y))} \le 1 - e^{-\nu}.$$
(40)

If f(x) < f(y), then (38) implies $e^{-(f(x)-f(y))} < e^{\nu}$ and we have

$$\left| e^{-(f(x) - f(y))} - 1 \right| = e^{-(f(x) - f(y))} - 1 \le e^{\nu} - 1.$$
(41)

 $1 - e^{-\nu} = e^{\nu} - 1$ for $\nu = 0$, and $\forall \nu > 0$,

$$\frac{d}{d\nu}(1 - e^{-\nu}) < \frac{d}{d\nu}(e^{\nu} - 1),$$

which implies $1 - e^{-\nu} < e^{\nu} - 1$.

In Sections 4.1.2 and 4.1.3 we use Fact 4.1 to write down uniform approximations to Π and Π_{n-1} for subsets of C.

A uniform approximation for Π 4.1.2

We continue discussing an axis-aligned cube $C \subset K$ of side δ . Consider any $S \subseteq C$. Let

$$w \equiv \underset{x \in C}{\arg\min} f(x) = \underset{x \in C}{\arg\max} \pi(x).$$
(42)

C is a bounded set and f is strongly convex, and so we are guaranteed that w exists. We proceed to give an error bound for the uniform approximation $\pi(w) \operatorname{vol}(S)$ to $\Pi(S)$.

Since $\pi(w) \ge \pi(x) \ \forall x \in C$, we have the simple upper bound $\Pi(S) \le \pi(w) \operatorname{vol}(S)$. For a lower bound on $\Pi(S)$, we have

$$\begin{aligned} \pi(w) \operatorname{vol}(S) - \Pi(S) &= \int_{S} dx \left(\pi(w) - \pi(x) \right) \\ &\leq \pi(w) \int_{S} dx \left(1 - e^{-(f(x) - f(w))} \right) \\ &\leq \pi(w) \left(1 - e^{-\nu} \right) \int_{S} dx \\ &= \pi(w) (1 - e^{-\nu}) \operatorname{vol}(S). \end{aligned}$$

In the penultimate step above we have applied (40).

It will be useful to summarize the discussion so far in the following Fact:



Figure 2: Pictured here is a cube of side δ . β , shaded in light yellow, is a facet normal to the e_3 coordinate axis. β_a , shaded in gray, is the set that results from translating β along the e_3 axis to $x_3 = a$. ω is the subset of β shaded in a darker yellow, and ω_a is the subset of β_a shaded in a darker gray. Outlined in black is the extension B of ω along e_3 in the cube.

Fact 4.2. Let $w \equiv \arg \min_{x \in C} f(x)$. For any $S \subseteq C \subset K$, we have

$$\frac{\Pi(S)}{\pi(w)\mathrm{vol}(S)} \in [e^{-\nu}, 1],\tag{43}$$

where ν is as defined in (38).

We are not limited to choosing w as a reference point in C. We do so for convenience, but any choice of reference point is sufficient.

4.1.3 Approximating Π by $\delta \cdot \Pi_{n-1}$

Let β be a facet of C normal to the coordinate direction e_i and let β_a be the n-1 dimensional set in C that results from translating β to the location $x_i = a$ in C. See Figure 4 for an illustration in three dimensions. Let $\omega \subseteq \beta$. Let ω_a be the translation of ω to $x_i = a$, and let B be the ndimensional subset of C that is swept out by ω as it is translated along e_i from β all the way to the opposite facet. We will call this the *extension of* ω *in* C. Without loss of generality, let $x_i = 0$ in β .

We can write $\Pi(B)$ as follows:

$$\Pi(B) = \int_0^\delta dx_i \,\Pi_{n-1}(\omega_{x_i}). \tag{44}$$

Denote by x_i^a any vector with i^{th} coordinate fixed to a. We will now give an error bound for approximating $\Pi(B)$ with $\delta \cdot \Pi_{n-1}(\omega_b)$ for any $x_i^b \in B$.

Fix any two values a and b of x_i in B. We have

$$|\Pi_{n-1}(\omega_a) - \Pi_{n-1}(\omega_b)| = \left| \int_B dx_{-i} \left[\pi(x_i^a) - \pi(x_i^b) \right] \right|$$

$$\leq \int_B dx_{-i} \pi \left(x_i^b \right) \left| e^{-(f(x_i^a) - f(x_i^b))} - 1 \right|$$

$$\leq (e^{\nu} - 1) \int_B dx_{-i} \pi(x_i^b)$$

$$= (e^{\nu} - 1) \Pi_{n-1}(\omega_b), \qquad (45)$$

where we have applied Fact 4.1 in the penultimate line. Therefore,

$$|\Pi(B) - \delta \cdot \Pi_{n-1}(\omega_b)| = \left| \int_0^\delta dx_i \left[\Pi_{n-1}(\omega_{x_i}) - \Pi_{n-1}(\omega_b) \right] \right|$$

$$\leq \int_0^\delta dx_i \left| \Pi_{n-1}(\omega_{x_i}) - \Pi_{n-1}(\omega_b) \right|$$

$$\leq (e^\nu - 1)\Pi_{n-1}(\omega_b) \int_0^\delta dx_i$$

$$= (e^\nu - 1) \cdot \delta \cdot \Pi_{n-1}(\omega_b).$$
(46)

Since this bound holds for any value of x_i^b in C, we are free to choose b to be the value of x_i on one of the facets β of the cube. We have argued the following fact.

Fact 4.3. Consider an axis-aligned cube $C \subset K$ of side δ and a log-smooth strongly log-concave probability measure Π with density π . Let β be any n-1-dimensional facet of C and ω any subset of β . Let B be the extension of ω in C along the coordinate direction normal to β . Then,

$$|\Pi(B) - \delta \cdot \Pi_{n-1}(\omega)| \le (e^{\nu} - 1) \cdot \delta \cdot \Pi_{n-1}(\omega), \tag{47}$$

where ν is as defined in (38).

4.2 L_0 isoperimetry

4.2.1 Cube isoperimetry

The following lemma is due to Laddha and Vempala [16].

Lemma 4.4. Consider an axis-aligned cube C of side length δ . Let S_1 and S_2 be two axis-disjoint subsets of C such that $\operatorname{vol}(S_1) \leq (2/3)\operatorname{vol}(C)$. Let $S_3 = C \setminus \{S_1 \cup S_2\}$. Then

$$\operatorname{vol}(S_3) \ge \frac{\psi_c}{4} \operatorname{vol}(S_1)$$
$$\psi_c \ge \frac{\log 2}{\sqrt{n}}.$$
(48)

where

Laddha and Vempala originally gave a lower bound of
$$n^{-1} \log 2$$
 for ψ_c . This was later improved
to $n^{-1/2} \log 2$ by Fernandez [26], who also showed that the inverse square root scaling of n is optimal.

We prove a similar result for π that relies on Lemma 4.4.

Lemma 4.5. Consider an axis-aligned cube C of side length δ and a log-smooth strongly log-concave measure Π supported on \mathbb{R}^n with associated density π . Let S_1 and S_2 be two axis-disjoint subsets of C such that

$$\Pi(S_1) \le \frac{2}{3}e^{-\nu}\Pi(C),$$

where ν is as defined in (38). Let $S_3 = C \setminus \{S_1 \cup S_2\}$. Then,

$$\Pi(S_3) \ge \frac{\psi_c}{4} e^{-\nu} \Pi(S_1)$$

Proof. Recall the quantity $w = \arg \min_{x \in C} f(x)$. From Fact 4.2 we have the following two inequalities:

$$\pi(w)e^{-\nu}\operatorname{vol}(S_1) \le \Pi(S_1),\tag{49}$$

$$\pi(w)e^{-\nu}\operatorname{vol}(S_3) \le \Pi(S_3). \tag{50}$$

We check the condition in Lemma 4.4:

$$\pi(w)e^{-\nu}\operatorname{vol}(S_1) \stackrel{(i)}{\leq} \Pi(S_1) \stackrel{(ii)}{\leq} \frac{2}{3}e^{-\nu}\Pi(C) \stackrel{(iii)}{\leq} \frac{2}{3}e^{-\nu}\pi(w)\operatorname{vol}(C)$$

$$\Rightarrow \operatorname{vol}(S_1) \leq \frac{2}{3}\operatorname{vol}(C).$$

In (i), we used (49). (ii) is by assumption. In (iii) we used the definition of w. Thus the condition of Lemma 4.4 is satisfied and we are guaranteed

$$\operatorname{vol}(S_3) \ge \frac{\psi_c}{4} \operatorname{vol}(S_1).$$
(51)

To translate this into the statement of the lemma, we apply (51) to (50) and then use the definition of w. We have

$$\frac{\Pi(S_3)}{\pi(w)}e^{\nu} \ge \operatorname{vol}(S_3) \ge \frac{\psi_c}{4}\operatorname{vol}(S_1) \ge \frac{\psi_c}{4}\frac{\Pi(S_1)}{\pi(w)}$$
$$\Rightarrow \Pi(S_3) \ge \frac{\psi_c}{4}e^{-\nu}\Pi(S_1).$$

4.2.2 Proof of Lemma 1.2

We begin by stating an isoperimetric theorem for strongly log-concave distributions that will be needed in the course of the argument.

Theorem 4.6. Consider a convex body $A \subset \mathbb{R}^n$ and a subset $I \subseteq A$. Let $\partial_A I$ be the internal boundary of I in A. Let Π be a strongly log-concave probability measure supported on \mathbb{R}^n and let $||\Sigma||_{op}$ be the operator norm (the largest eigenvalue) of its covariance matrix Σ . Let $\Pi_{n-1}(\partial_K I)$ represent the measure of the boundary set $\partial_K I$ induced by Π . Then there exists a universal constant c' such that

$$\Pi_{n-1}(\partial_A I) \ge \psi_{\pi} \min \left\{ \Pi(I), \Pi(A \setminus I) \right\}$$

where

$$\psi_{\pi} \ge \frac{1}{c' \left(n \, ||\Sigma||_{op}^2\right)^{1/4}}.\tag{52}$$

This theorem is due to Lee and Vempala [27]. For a μ -strongly log-concave distribution, $||\Sigma||_{op} \leq \frac{1}{\mu}$.

We are now ready to prove Lemma 1.2.

Proof of Lemma 1.2. Consider a Euclidean ball K of radius R in \mathbb{R}^n embedded in a grid of axisaligned cubes of side

$$\delta = \frac{1}{64\sqrt{\kappa L}n\log n \max\left\{1, \sqrt{\frac{1}{n}\log\frac{1}{\varepsilon}}\right\}}$$

for some $0 < \varepsilon < 1/2$. K is centered at the mode x^* of π .

 S_1 and S_2 are two axis-disjoint subsets of K, and let $S_3 = K \setminus \{S_1 \cup S_2\}$. Without loss of generality, let $\Pi(S_1) \leq \Pi(S_2)$.

Let $K' = (1 - \alpha)K$ be the α -shrinkage of K, which we must consider in order to avoid overestimating the lower bound on $\Pi(S_3)$. We require α to be such that it can fit at least two cube diagonals, i.e., we need $(\sqrt{n\delta})^{-1} \alpha \ge 2$. This choice of ratio between α and δ ensures that any cube intersecting K' as well as all its neighbors are contained in K. We take

$$\alpha = \frac{1}{4\sqrt{\kappa L}\sqrt{n}\log n \max\left\{1, \sqrt{\frac{1}{n}\log\frac{1}{\varepsilon}}\right\}}.$$
(53)

 α must be less than unity. In fact, we will enforce $\alpha \leq 1/2$ and place the following mild constraint on L to ensure it is satisfied:

$$L > \max\left\{\frac{1}{n\log^2 n}, \mu\right\}.$$
(54)

Let R' be the radius of K'. we make the assignment

$$R' = r(\varepsilon) \sqrt{\frac{n}{\mu}}.$$
(55)

Together with $\alpha \leq 1/2$, (55) implies $R' < R \leq 2R'$. Moreover, Lemma 3.1 guarantees $\Pi(K') \geq 1 - \varepsilon$, and therefore

$$\Pi(K \setminus K') < \Pi(\mathbb{R}^n \setminus K') < \varepsilon.$$
(56)

Let $S'_i = S_i \cap K'$ for i = 1, 2. For any $X \subseteq K$, we have

$$\Pi(X \cap K') = \Pi(X) - \Pi(X \cap K \setminus K') \ge \Pi(X) - \Pi(K \setminus K') > \Pi(X) - \varepsilon.$$
(57)

Let C be the set of all cubes that intersect S_1 . We partition it into two sets, the boundary set C_1 and the bulk set C_2 :

$$C_1 = \left\{ c \in C \ s.t. \ \Pi(c \cap S_1) \le \frac{2}{3} e^{-\nu} \Pi(c) \right\},\tag{58}$$

$$C_2 = \left\{ c \in C \ s.t. \ \Pi(c \cap S_1) > \frac{2}{3} e^{-\nu} \Pi(c) \right\}.$$
(59)

To preserve the bulk/boundary identities of these sets, we must have $\nu < \log \frac{4}{3}$. We have already ensured this (see the discussion surrounding (38)).

We use curly notation, C_i , to denote the *n*-dimensional body that is formed by taking the union of all the cubes in C_i for i = 1, 2.

Depending on whether most of the measure of S_1 is contained in the bulk C_2 or the boundary C_1 , there are two cases to consider.

Case 1. $\Pi(C_1 \cap S_1) \ge \frac{1}{2} \Pi(S_1).$

By definition, we have $\Pi(c \cap S_1) \leq \frac{2}{3}e^{-\nu}\Pi(c)$ for every $c \in C_1$, and so we can apply Lemma 4.5 to each cube in C_1 and sum the resulting contribution over all the cubes in C_1 to lower bound $\Pi(S_3)$. However, there being no guarantee that all cubes in C_1 are fully contained in K, we must correct for the possibility of overcounting contributions to S_3 . We do so by considering only the subset of cubes in C_1 that intersect K',

$$C_1' = \{ c \in C_1 \ s.t. \ c \cap K' \neq \emptyset \}.$$

$$(60)$$

Denote by C'_1 the solid body that is formed by taking the union of all the cubes in C'_1 . Due to our choice of α , we are guaranteed that $C'_1 \subseteq K$. By Lemma 4.5, each cube c in C'_1 makes the following minimal contribution to the measure of S_3 :

$$\Pi(c \cap S_3) \ge \frac{\psi_c}{4} e^{-\nu} \Pi(c \cap S_1).$$
(61)



Figure 3: The Euclidean ball K is embedded in a grid of cubes of side δ . Shaded in purple is the set of cubes in the bulk set C_2 covering S_1 . In orange is the boundary set C_1 .

Summing (61) over all $c \in C'_1$, we have

$$\Pi(S_3) \ge \sum_{c \in C_1'} \Pi(c \cap S_3) \ge \frac{\psi_c}{4} e^{-\nu} \sum_{c \in C_1'} \Pi(c \cap S_1) = \frac{\psi_c}{4} e^{-\nu} \Pi(\mathcal{C}_1' \cap S_1).$$

To lower bound $\Pi(\mathcal{C}'_1 \cap S_1)$, note that while \mathcal{C}'_1 may not be fully contained in K', the set $\mathcal{C}_1 \cap K'$ is, and therefore $\Pi(\mathcal{C}'_1) \geq \Pi(\mathcal{C}_1 \cap K')$. Using this fact, we have

$$\Pi(\mathcal{C}'_1 \cap S_1) \ge \Pi(\mathcal{C}_1 \cap S_1 \cap K') \ge \Pi(\mathcal{C}_1 \cap S_1) - \varepsilon.$$
(62)

The second inequality follows by taking $X = \mathcal{C}_1 \cap S_1$ in (57). Thus,

$$\Pi(S_3) \ge \frac{\psi_c}{4} e^{-\nu} \left(\Pi(\mathcal{C}_1 \cap S_1) - \varepsilon \right) \ge \frac{\psi_c}{4} e^{-\nu} \left(\frac{1}{2} \Pi(S_1) - \varepsilon \right) > \frac{5\psi_c}{24} \left(\frac{1}{2} \Pi(S_1) - \varepsilon \right), \tag{63}$$

where the second inequality follows from the condition $\Pi(\mathcal{C}_1 \cap S_1) \geq \frac{1}{2} \Pi(S_1)$ and the third from $\nu < \log 6/5.$

Case 2. $\Pi(\mathcal{C}_1 \cap S_1) < \frac{1}{2}\Pi(S_1) \Rightarrow \Pi(\mathcal{C}_2 \cap S_1) \geq \frac{1}{2}\Pi(S_1).$ Let $\partial_K \mathcal{C}_2$ denote the internal boundary of \mathcal{C}_2 in K, i.e., the intersection of K and the boundary of C_2 . Let \mathcal{B} be the set of facets (of cubes) that intersect $\partial_K C_2$. Consider a facet β in \mathcal{B} and the two cubes on either side of β . Since β is on the boundary of C_2 , one of these cubes, call it v, is in C_2 , and the other, u, is not. See Figure 4 for a picture of this construction.



Figure 4: The part of β underneath the dashed line is $\mathcal{P}_{\beta}(S_1 \cap v)$, the projection of $S_1 \cap v$ onto β . The part of u underneath the dashed line is $\mathcal{E}_u(\mathcal{P}_{\beta})$, the extension of $\mathcal{P}_{\beta}(S_1 \cap v)$ in u along the coordinate direction normal to β . The subset of this region shaded with lines is the precisely the part of $S_3 \cap u$ that the dynamics can escape to through β from $S_1 \cap v$. A worst-case lower bound on its measure is derived in (73).

In what follows, we will lower bound $\Pi(S_3 \cap u)$, thus quantifying the minimal measure of the region to which the sampler can escape S_1 through β . The bound will be in terms of $\Pi(u)$. We will rewrite it in terms of $\Pi_{n-1}(\beta)$, gather contributions from all the facets in \mathcal{B} , and then apply Theorem 4.6 to obtain a lower bound on $\Pi(S_3)$ in terms of $\Pi(S_1)$.

Let e_i be the coordinate vector perpendicular to β . Let $\mathcal{P}_{\beta}(S_1 \cap v)$ be the projection of $S_1 \cap v$ onto β , and let $\mathcal{E}_u(\mathcal{P}_{\beta}(S_1 \cap v))$ be the extension of this projection along e_i in u.

$$\mathcal{P}_{\beta}(S_1 \cap v) = \{ x \in \beta \ s.t. \ \exists y \in v \ s.t. \ y_j = x_j \ \forall j \in [n] \setminus i \}, \tag{64}$$

$$\mathcal{E}_u(\mathcal{P}_\beta(S_1 \cap v)) = \{ x \in u \ s.t. \ \exists y \in \mathcal{P}_\beta(S_1 \cap v) \ s.t. \ x_j = y_j \ \forall j \in [n] \setminus i \}.$$
(65)

 $\mathcal{E}_u(\mathcal{P}_\beta(S_1 \cap v))$ is precisely the region of u that the dynamics can move to from v in the direction e_i in one step. For notational simplicity, we will drop the argument of \mathcal{P}_β in the sequel.

 S_1 and S_2 are axis disjoint. Thus we have

$$\mathcal{E}_u(\mathcal{P}_\beta) \cap S_2 = \emptyset.$$

Consequently,

$$\Pi(\mathcal{E}_{u}(\mathcal{P}_{\beta})) = \Pi(\mathcal{E}_{u}(\mathcal{P}_{\beta}) \cap S_{1}) + \Pi(\mathcal{E}_{u}(\mathcal{P}_{\beta}) \cap S_{3}) \stackrel{(i)}{\leq} \Pi(u \cap S_{1}) + \Pi(u \cap S_{3}),$$

$$\Rightarrow \Pi(u \cap S_{3}) \geq \Pi(\mathcal{E}_{u}(\mathcal{P}_{\beta})) - \Pi(u \cap S_{1}).$$
(66)

(i) follows from the fact that $\mathcal{E}_u(\mathcal{P}_\beta) \subseteq u$.

Next, we develop a lower bound on $\Pi(\mathcal{E}_u(\mathcal{P}_\beta))$ in terms of $\Pi(u)$. By construction,

$$\operatorname{vol}(\mathcal{E}_u(\mathcal{P}_\beta)) \ge \operatorname{vol}(S_1 \cap v)$$

Let $w_v = \arg \min_{x \in v} f(x)$. Then

$$\operatorname{vol}(S_1 \cap v) \ge \frac{\Pi(S_1 \cap v)}{\pi(w_v)} > \frac{2}{3}e^{-\nu}\frac{\Pi(v)}{\pi(w_v)}$$

where the second inequality follows from the fact that v is an element of C_2 . Now, using Fact 4.2,

$$\Pi(v) \ge e^{-\nu} \pi(w_v) \operatorname{vol}(v) = e^{-\nu} \pi(w_v) \operatorname{vol}(u).$$

Thus,

$$\operatorname{vol}(S_1 \cap v) \ge \frac{2}{3}e^{-2\nu}\operatorname{vol}(u)$$

i.e.,

$$\operatorname{vol}(\mathcal{E}_u(\mathcal{P}_\beta)) \ge \frac{2}{3}e^{-2\nu}\operatorname{vol}(u).$$
(67)

Let $w_u = \arg \min_{x \in u} f(x)$. Then

$$\operatorname{vol}(u) \ge \frac{\Pi(u)}{\pi(w_u)}$$

and by Fact 4.2,

$$\frac{\Pi(\mathcal{E}_u(\mathcal{P}_\beta))}{\pi(w_u)}e^{\nu} \ge \operatorname{vol}(\mathcal{E}_u(\mathcal{P}_\beta)).$$

Applying the last two bounds to (67), we have the following lower bound on the measure of $\mathcal{E}_u(\mathcal{P}_\beta)$:

$$\Pi(\mathcal{E}_u(\mathcal{P}_\beta)) \ge \frac{2}{3} e^{-3\nu} \Pi(u).$$
(68)

Substituting (68) in (66), we have the following lower bound on $\Pi(S_3 \cap u)$ in terms of $\Pi(u)$ and $\Pi(S_1 \cap u)$:

$$\Pi(u \cap S_3) \ge \frac{2}{3}e^{-3\nu}\Pi(u) - \Pi(u \cap S_1).$$
(69)

Now, if $u \notin C_1$ (i.e., $\Pi(u \cap S_1) = 0$), the second term on the right-hand side of (69) vanishes, and we are left with a bound on $\Pi(S_3 \cap u)$ that depends only on $\Pi(u)$, as was our goal. However, if $u \in C_1$, then there is more work to be done, which we split into two cases depending on the proportion of u occupied by S_1 . Let this proportion be represented by a quantity h, which we will choose later. Note that h cannot exceed $\frac{2}{3}e^{-\nu}$.

• If $\Pi(u \cap S_1) \leq h \Pi(u)$, then from (69), we have

$$\Pi(u \cap S_3) \ge \Pi(\mathcal{E}_u(\mathcal{P}_\beta)) - h\Pi(u) \ge \left(\frac{2}{3}e^{-3\nu} - h\right)\Pi(u).$$
(70)

• If $h\Pi(u) < \Pi(u \cap S_1) \le \frac{2}{3}e^{-\nu}\Pi(u)$, we can apply Lemma 4.5 to $u \cap S_3$ to directly lower bound its measure:

$$\Pi(u \cap S_3) \ge \frac{\psi_c}{4} e^{-\nu} \Pi(S_1 \cap u) \ge \frac{\psi_c}{4} e^{-\nu} h \Pi(u).$$
(71)

The minimal bound on $\Pi(u \cap S_3)$. The contribution of u to the measure of S_3 is at least the minimum of the three bounds we have so far written for it (given by (69) when $\Pi(S_1 \cap u) = 0$ and by (70) and (71) when $\Pi(S_1 \cap u) \neq 0$):

$$\Pi(S_3 \cap u) \ge \min\left\{\frac{2}{3}e^{-3\nu}, \frac{2}{3}e^{-3\nu} - h, \frac{\psi_c}{4}e^{-\nu}h\right\}\Pi(u).$$
(72)

The first term in (72) is clearly larger than the other two. To distinguish between the latter, we must make a choice of h.

Subject to the constraint that the second term in (72) be positive for all n, it is not possible to choose h so that this term is smaller than the third term for all n.¹ It is therefore sufficient to take h to be a constant strictly smaller than $\frac{2}{3}e^{-3\nu}$. Since $e^{-\nu} \ge 5/6$ (see the discussion surrounding (38)), any $h < \frac{2}{3}\left(\frac{5}{6}\right)^3 = \frac{125}{324}$ will suffice. We will take h = 1/3. Thus we arrive at the bound

$$\Pi(S_3 \cap u) \ge \frac{\psi_c}{12} e^{-\nu} \Pi(u). \tag{73}$$

To write this bound in terms of $\Pi_{n-1}(\beta)$, we apply Fact 4.3 to $\Pi(u)$ with $\omega = \beta$. Then,

$$\Pi(S_3 \cap u) \ge \frac{\psi_c}{12} \cdot (2e^{-\nu} - 1) \cdot \delta \cdot \Pi_{n-1}(\beta) \ge \frac{\psi_c}{12} \cdot \frac{2}{3} \cdot \delta \cdot \Pi_{n-1}(\beta).$$

$$\tag{74}$$

The cube u has 2n facets, which means that there are up to 2n facets in \mathcal{B} that can contribute to $\Pi(S_3 \cap u)$. We will shortly sum (74) over all $\beta \in \mathcal{B}$. To avoid overcounting contributions to $\Pi(S_3)$ when we do so, we normalize the right-hand side of (74) by a factor of 1/2n. Every facet $\beta \in \mathcal{B}$ therefore makes at least the contribution

$$\frac{1}{2n} \cdot \frac{\psi_c}{18} \cdot \delta \cdot \Pi_{n-1}(\beta) \tag{75}$$

to $\Pi(S_3)$.

Now we must correct for the possibility that not all the cubes that contribute facets to \mathcal{B} are fully contained in K. To do so, we define the following subset of C_2 :

$$C'_{2} = \{ c \in C_{2} \ s.t. \ c \cap K' \neq \emptyset \}.$$

$$(76)$$

By construction, every cube in C'_2 and its immediate neighbors are fully contained in K. Define the set $I = K' \cap \mathcal{C}_2$. $I \subseteq \mathcal{C}'_2$. Let \mathcal{B}' be the set of facets of cubes that intersect the internal boundary of I in K', $\partial_{K'}I$. \mathcal{B}' is the set of facets out of which the dynamics can escape S_1 and that are contributed by cubes that are fully contained in K. $\mathcal{B}' \subseteq \mathcal{B}$. Summing (75) over all facets in \mathcal{B}' , we arrive at the bound

$$\Pi(S_3) \ge \frac{1}{2n} \frac{\psi_c}{18} \cdot \delta \sum_{\beta \in \mathcal{B}'} \Pi_{n-1}(\beta) = \frac{1}{2n} \frac{\psi_c}{18} \cdot \delta \cdot \Pi\left(\partial_{K'}I\right).$$
(77)

 $\Pi(\partial_{K'}I)$ is lower bounded via Theorem 4.6 as follows:

$$\Pi\left(\partial_{K'}I\right) \ge \psi_{\pi} \min\left\{\Pi(I), \Pi(K' \setminus I)\right\}.$$
(78)

We evaluate the minimum of $\Pi(I)$ and $\Pi(K' \setminus I)$. We have

$$\Pi(I) = \Pi(K' \cap \mathcal{C}_2) > \Pi(\mathcal{C}_2) - \varepsilon > \frac{1}{2}\Pi(S_1) - \varepsilon.$$
(79)

The first inequality follows from (57), and the second by construction: $\Pi(S_1) = \Pi(\mathcal{C}_2 \cap S_1) + \Pi(\mathcal{C}_1 \cap S_1) \leq 2\Pi(\mathcal{C}_2 \cap S_1) \leq 2\Pi(\mathcal{C}_2)$. In addition, we have the lower bound

$$\Pi(I) \le \Pi(\mathcal{C}'_2) < \frac{3}{2} e^{\nu} \Pi(\mathcal{C}'_2 \cap S_1) \le \frac{9}{5} \Pi(\mathcal{C}'_2 \cap S_1) \le \frac{9}{10} \Pi(K),$$
(80)

where the second inequality follows from the definition of C_2 , the third from (38), and the last from $\Pi(S_1) \leq \Pi(K)/2$. Using (80), we have the following lower bound for $\Pi(K' \setminus I)$:

$$\Pi(K' \setminus I) = \Pi(K') - \Pi(I) \ge \Pi(K') - \frac{9}{10}\Pi(K) = \frac{1}{10}\Pi(K) - \Pi(K \setminus K') \ge \frac{1}{5}\Pi(S_1) - \varepsilon.$$
(81)

¹For any fixed n', if we make the assignment $h = \iota e^{-3\nu}$ where ι is a constant strictly less than 2/3, we can choose ι to be such that the second term in (72) is smaller than the third term for all $n \leq n'$.

And thus

$$\min\left\{\Pi(I), \Pi(K' \setminus I)\right\} \ge \frac{1}{5}\Pi(S_1) - \varepsilon.$$
(82)

Putting together (77), (78), and (82), and using the definition of δ , we finally arrive at the following bound on $\Pi(S_3)$:

$$\Pi(S_3) \ge \frac{1}{2n} \frac{\psi_c}{18} \psi_\pi \frac{1}{8\sqrt{\kappa L} n \log n \max\left\{1, \sqrt{\frac{1}{n} \log \frac{1}{\varepsilon}}\right\}} \left(\frac{1}{5} \Pi(S_1) - \varepsilon\right).$$
(83)

This concludes the discussion of Case 2.

The minimum of (63) and (83) is clearly the latter. Applying the lower bounds (48) and (52) on ψ_c and ψ_{π} , respectively, we finally arrive at the bound

$$\Pi(S_3) \ge \Psi\left(\frac{1}{5}\Pi(S_1) - \varepsilon\right),\,$$

where, for a universal constant c,

$$\Psi \ge \frac{\log 2}{2^5 \cdot 3^3 \cdot c} \frac{1}{\kappa n^{2+\frac{3}{4}} \log n \max\left\{1, \sqrt{\frac{1}{n} \log \frac{1}{\varepsilon}}\right\}}.$$

5 Discussion

For very large n, the exponent of n in (1) can be improved by applying an alternative lower bound on the log-concave isoperimetric coefficient in Theorem 4.6 due to Chen [28]:

$$\psi_{\pi} \ge \frac{\mu^{\frac{1}{2}}}{n^{c'}\sqrt{\frac{\log\log n}{\log n}}}.$$
(84)

c' is a universal constant. Applying this bound on ψ_{π} in Ψ instead of (52), we have

$$\tau(\gamma) < C\kappa^2 n^{7+2c'} \sqrt{\frac{\log \log n}{\log n}} \log^2 n \left(\max\left\{ 1, \sqrt{\frac{1}{n} \log \frac{2M}{\gamma}} \right\} \right)^2 \log \frac{2M}{\gamma}$$
(85)

for universal constants C and c'. The function $\sqrt{\frac{\log \log n}{\log n}}$ goes to zero as n goes to infinity. When n is large enough that $2c'\sqrt{\frac{\log \log n}{\log n}} < 0.5$, (85) is a tighter bound on the mixing time than (1).

It is sometimes of interest to consider sampling from an *isotropic* log-concave distribution, i.e., a distribution for which $\kappa = 1$ and $x^* = 0$. Any strongly log-concave distribution can be brought into isotropic position in $O^*(n^5)$ steps [4]. After this preconditioning step, which need be done only once, samples may be drawn using a sampling algorithm of choice. If we were to consider only the subclass of isotropic log-concave distributions in the proof of Theorem 1.1, we could apply concentration results for isotropic log-concave measures that have slightly better *n*-dependence than does Lemma 3.1, such as Theorem 16 in [27], when choosing the radius of *K*. Doing so enables us to improve the *n*-dependence of δ to $\left(\sqrt{n} \log n \sqrt{\log \frac{1}{\varepsilon}} \max\left\{n^{1/4}, \sqrt{\log \frac{1}{\varepsilon}}\right\}\right)^{-1}$ for $\varepsilon \in (0, 1/2)$. This leads to a mixing time bound with a better exponent in *n* but worse dependence on M/γ :

$$\tau(\gamma) < Cn^{6.5} \log^2 n \left(\max\left\{ n^{\frac{1}{4}}, \sqrt{\frac{1}{n} \log \frac{2M}{\gamma}} \right\} \right)^2 \log^2 \frac{2M}{\gamma}, \tag{86}$$

where C is a universal constant.

We suspect that neither the bound in Theorem 1.1 nor (85) is tight. Establishing an isoperimetric inequality with better *n*-dependence than Lemma 1.2 is therefore an important problem for the future. A different proof technique than the one used to prove Lemma 1.2 will likely be needed.

Acknowledgements

NSW is grateful to Yuansi Chen for suggesting the possibility of exploiting the approximate local uniformity of smooth functions to prove Lemma 1.2, and to Andre Wibisono, Santosh Vempala, Aditi Laddha, Samuel Livingstone, and Andrew Wade for useful discussions. NSW would like to thank the Isaac Newton Institute for Mathematical Sciences for support and hospitality during the program Stochastic Systems for Anomalous Diffusion where a part of this work was completed. This work was supported by: EPSRC grant number EP/R014604/1. The Flatiron Institute is a division of the Simons Foundation.

References

- Nicholas Metropolis, Arianna W. Rosenbluth, Marshall N. Rosenbluth, Augusta H. Teller, and Edward Teller. Equation of state calculations by fast computing machines. J. Chem. Phys. 21, 1087, 21:1087–1092, 1953.
- [2] W. K. Hastings. Monte Carlo sampling methods using Markov chains and their applications. *Biometrika*, 57:97–109, 1970.
- [3] Ravi Kannan, László Miklós Lovász, and Miklós Simonovits. Random walks and an O*(n⁵) volume algorithm for convex bodies. *Random Struct. Algorithms*, 11:1–50, 1997.
- [4] Laszló Lovász and Santosh Vempala. The geometry of logconcave functions and sampling algorithms. Random Structures & Algorithms, 30:307–358, 2007.
- [5] M. Dyer, A. Frieze, and R. Kannan. A random polynomial-time algorithm for approximating the volume of convex bodies. *Journal of the Association for Computing Machinery*, 38(1):1–17, 1991.
- [6] L. Lovász and S. Vempala. Simulated annealing in convex bodies and an O^{*}(n⁴) volume algorithm. Journal of Computer and System Sciences, 72:392–417, 2006.
- [7] Arnon Boneh. Preduce a probabilistic algorithm identifying redundancy by a random feasible point generator (RFPG). In *Redundancy in Mathematical Programming*, pages 108–134, Berlin, Heidelberg, 1983. Springer Berlin Heidelberg.
- [8] Robert L. Smith. Efficient Monte Carlo procedures for generating points uniformly distributed over bounded regions. Operations Research, 32(6):1296–1308, 1984.
- [9] László Lovász and Santosh Vempala. Hit-and-Run from a corner. SIAM Journal on Computing, 35:310–314, 2004.
- [10] Santosh S. Vempala. Geometric random walks: a survey. Combinatorial and Computational Geometry, 52, 2005.
- [11] V. F. Turchin. On the computation of multidimensional integrals by the Monte-Carlo method. Theory of Probability & Its Applications, 16(4):720–724, 1971.

- [12] Stuart Geman and Donald Geman. Stochastic relaxation, Gibbs distributions, and the Bayesian restoration of images. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, PAMI-6(6):721–741, 1984.
- [13] Alan E. Gelfand and Adrian F. M. Smith. Sampling-based approaches to calculating marginal densities. *Journal of the American Statistical Association*, 85:398–409, 1990.
- [14] R. Kannan, L. Lovász, and M. Simonovits. Isoperimetric problems for convex bodies and a localization lemma. Discrete & Computational Geometry, 13:541—-559, 1995.
- [15] Fabio Martinelli. Lectures on glauber dynamics for discrete spin models. In Pierre Bernard, editor, Lectures on Probability Theory and Statistics: Ecole d'Eté de Probailités de Saint-Flour XXVII - 1997, pages 93–191. Springer Berlin Heidelberg, Berlin, Heidelberg, 1999.
- [16] Aditi Laddha and Santosh S. Vempala. Convergence of Gibbs sampling: Coordinate hit-and-run mixes fast. Discrete Comput. Geom., 70(2):406–425, apr 2023.
- [17] Hariharan Narayanan and Piyush Srivastava. On the mixing time of coordinate hit-and-run. Combinatorics, Probability and Computing, 31(2):320-332, 2022.
- [18] Hariharan Narayanan, Amit Rajaraman, and Piyush Srivastava. Sampling from convex sets with a cold start using multiscale decompositions. In *Proceedings of the 55th Annual ACM Symposium on Theory of Computing*, STOC 2023, pages 117—130, New York, NY, USA, 2023. Association for Computing Machinery.
- [19] Filippo Ascolani, Hugo Lavenant, and Giacomo Zanella. Entropy contraction of the Gibbs sampler under log-concavity. arXiv:2410.00858, 2024.
- [20] Jason M. Altschuler and Sinho Chewi. Faster high-accuracy log-concave sampling via algorithmic warm starts. J. ACM, 71:1–55, 2024.
- [21] L. Lovász and M. Simonovits. Random walks in a convex body and an improved volume algorithm. Random Structures & Algorithms, 4(4):359–412, 1993.
- [22] Mark Jerrum and Alistair Sinclair. Conductance and the rapid mixing property for Markov chains: the approximation of permanent resolved. In *Proceedings of the Twentieth Annual ACM Symposium on Theory of Computing*, STOC '88, page 235–244, New York, NY, USA, 1988. Association for Computing Machinery.
- [23] Jeff Cheeger. A lower bound for the smallest eigenvalue of the laplacian. In Problems in Analysis, pages 195–199. Princeton University Press, 1970.
- [24] László Lovász. Random walks on graphs: A survey. In Combinatorics, Paul Erdős is Eighty, volume 2, pages 1–46. János Bolyai Mathematical Society, 1993.
- [25] Raaz Dwivedi, Yuansi Chen, Martin J. Wainwright, and Bin Yu. Log-concave sampling: Metropolis-Hastings algorithms are fast. *Journal of Machine Learning Research*, 20(183):1–42, 2019.
- [26] Manuel Fernandez V. On the ℓ_0 isoperimetric coefficient of measurable sets. arXiv:2312.00015, 2024.
- [27] Yin Tat Lee and Santosh S. Vempala. Eldan's stochastic localization and the KLS conjecture: Isoperimetry, concentration and mixing. *Annals of Mathematics*, 199:1043–1092, 2024.
- [28] Yuansi Chen. An almost-constant lower bound of the isoperimetric coefficient in the KLS conjecture. *Geometric and Functional Analysis*, 31:34–61, 2021.