Multilevel Method with Low-Order Equations of Mixed Types and Two Grids in Photon Energy for Thermal Radiative Transfer

Dmitriy Y. Anistratov^a, Terry S. Haut^b

^aDepartment of Nuclear Engineering, North Carolina State University Raleigh, NC ^bLawrence Livermore National Laboratory, Livermore, CA

Abstract

Thermal radiative transfer (TRT) is an essential piece of physics in inertial confinement fusion, high-energy density physics, astrophysics etc. The physical models of this type of problem are defined by strongly coupled differential equations describing multiphysics phenomena. This paper presents a new nonlinear multilevel iterative method with two photon energy grids for solving the multigroup radiative transfer equation (RTE) coupled with the material energy balance equation (MEB). The multilevel system of equations of the method is formulated by means of a nonlinear projection approach. The RTE is projected over elements of phase space to derive the low-order equations of different types. The hierarchy of equations consists of (1) multigroup weighted flux equations which can be interpreted as the multigroup RTE averaged over subintervals of angular range and (2) the effective grey (one-group) equations. The system of RTE, low-order and MEB equations is approximated by the fully implicit Euler time-integration method in which absorption coefficient and emission term are evaluated at the current time step. Numerical results are presented to demonstrate convergence of a multilevel iteration algorithm in the Fleck-Cummings test problem with Marshak wave solved with large number of photon energy groups.

Keywords: high-energy-density physics, radiative transfer, multilevel iteration methods, multigrid methods, multiphysics problems.

1. Introduction

Radiative transfer plays an important part in high-temperature phenomena of high-energy density physics, inertial confinement fusion, and astrophysics [1]. High-energy photons affect redistribution of energy in such physical systems. The state of the physical system is significantly impacted by absorption of photons by matter and their emission, leading to change in the material energy. The Boltzmann radiative transfer equation (RTE) describes photon transport in a medium and the interaction of particles with matter. The absorption coefficient and emission source term in the RTE nonlinearly depend on the material temperature.

The fundamental thermal radiative transfer (TRT) model in 1D slab geometry is defined by the time-dependent multigroup RTE for the group intensity

$$\frac{1}{c}\partial_t I_g + \mathcal{L}I_g = \sigma_g B_g, \quad g \in \mathbb{N}(n_\nu), \tag{1}$$

Email addresses: anistratov@ncsu.edu (Dmitriy Y. Anistratov), haut3@llnl.gov (Terry S. Haut) Preprint

D. Anistratov & T. Haut, Multilevel Method for Thermal Radiative Transfer

$$\mathcal{L}I_g(x,\mu,t) \equiv \mu \partial_x I_g(x,\mu,t) + \sigma_g(x;T) I_g(x,\mu,t) , \qquad (2)$$

$$I_g|_{t=0} = I_g^0, \quad I_g|_{\substack{\mu>0\\x=0}} = I_g^+, \quad I_g|_{\substack{\mu<0\\x=X}} = I_g^-,$$
(3)

and the material energy balance (MEB) equation

$$\partial_t \varepsilon(T) = \sum_{g=1}^{n_\nu} \int_{-1}^1 \sigma_g(x;T) \Big(I_g(x,\mu,t) - B_g(T) \Big) d\mu \,, \tag{4}$$

$$T\Big|_{t=0} = T^0.$$
 (5)

Here x is the spatial position; μ is the directional cosine of particle motion; t is time; g is the index of the photon energy group; T is the material temperature; ε is the material energy density;

$$I_g = \int_{\nu_g}^{\nu_{g+1}} I_\nu d\nu \tag{6}$$

is the group intensity, ν is the photon frequency,

$$B_g(T) = \int_{\nu_g}^{\nu_{g+1}} B_\nu(T) d\nu$$
(7)

is the group Planckian function,

$$\sigma_g(T) = \frac{\int_{\nu_g}^{\nu_{g+1}} \sigma_\nu(T) B_\nu(T) d\nu}{\int_{\nu_g}^{\nu_{g+1}} B_\nu(T) d\nu}$$
(8)

is the group averaged absorption coefficient. The group structure is defined by the photon energy grid

$$\Lambda_{h\nu} = \{h\nu_g, g \in \mathbb{N}(n_\nu + 1): \ h\nu_{min} = h\nu_1 < \ldots < h\nu_g < h\nu_{g+1} < \ldots < h\nu_{n_\nu + 1} = h\nu_{max}\}.$$
 (9)

This TRT model neglects material motion, scattering, heat conduction, and external sources. It describes the phenomena of a supersonic radiation wave. The general model is formulated by the system of radiation hydrodynamics equations, which involves conservation of mass and momentum equations and accounts for the radiation effect on matter [2].

There exist a family of computational methods for TRT problems which apply linearizaton to the multigroup RTE resulting in the RTE with a pseudo-scattering term. To accelerate transport iterations due to pseudo-scattering, efficient algorithms are used [3–6]. A different group of methods follows the moment-based approach to formulate iteration schemes based high-order and low-order equations [7–9].

In this paper, we apply a nonlinear projection approach to formulate a new multilevel iteration method for solving coupled RTE and MEB equation [10-15]. The hierarchy of equations of the method is defined for different moments of the radiation intensity. The RTE is projected sequentially over elements of phase space to derive nonlinear moment equations with exact closures. One part of the hierarchy of moment equations is defined by multigroup low-order nonlinear weighted flux (LONWF) equations, which can be viewed as the RTE averaged over subintervals of angular range with certain weight functions [12, 16–22]. Another part of the hierarchy is the effective grey low-order quasidiffusion (GLOQD), aka variable Eddington factor (VEF), equations [10, 23–27]. The LONWF

2

equations are defined for partial angular moments of the intensity. They mimic the structure of the RTE and lead to a coupled set of Cauchy problems in space. The GLOQD equations are formulated for the full-range angular moments of the intensity and have a structure similar to the P_1 equations, resulting in a boundary value problem in space. Thus, the proposed multilevel method is based on low-order problems of mixed types. Parallel algorithms developed for solving the RTE on GPUs can be adopted for solving the LONWF equations. This provides an opportunity for developing fast and efficient algorithms for TRT problems on advanced high-performance computing system with GPUs.

Our approach possesses the following major potential advantages over more traditional discretization schemes: (i) the low-order system for the radiation energy is hyperbolic and this naturally avoids some pathological issues that can arrive when coupling to diffusion near highly heterogeneous material interfaces, (ii) our formulation can result in positivity for both the angular intensity and the angle-averaged solutions through positivity-preserving sweeps, (iii) our method naturally extends to using a hierarchy of levels in both angle and frequency, which opens the way for potentially much more efficient iterative methods, (iv) our approach can dynamically adapt the multi-level hierarchy of equations for different physics regimes, potentially resulting in greater efficiency.

The remainder of this paper is organized as follows. In Sec. 2, the multilevel method is formulated. The iteration algorithm is described in Sec. 3. In Sec. 4, the numerical results are presented. We conclude with a brief summary in Sec. 5.

2. Formulation of the Multilevel Method

The method is formulated by a multiscale system of differential equations consisting of

- the multigroup high-order RTE (Eq. (1)) on the grid $\Lambda_{h\nu}$ for the group intensities I_q ,
- the multigroup LONWF equations on the grid $\Lambda_{h\nu}$ for the group partial angular moments of radiation intensities

$$\phi_g^{\pm} = \pm \int_0^{\pm 1} I_g d\mu \,, \tag{10}$$

• effective GLOQD equations on the coarse photon energy grid

$$\Lambda_{h\nu}^* = \{h\nu_1 = h\nu_{min}, h\nu_2 = h\nu_{max}\}$$
(11)

for the total radiation energy density and flux

$$E = \sum_{g=1}^{n_{\nu}} E_g , \quad F = \sum_{g=1}^{n_{\nu}} F_g , \qquad (12)$$

where the group quantities are defined by

$$E_g = \frac{1}{c} \int_{-1}^{1} I_g d\mu \,, \quad F_g = \int_{-1}^{1} \mu I_g d\mu \,. \tag{13}$$

The multigroup LONWF equations for ϕ_g^{\pm} are derived by taking angular moments of the RTE with weight $w(\mu) = \gamma(1 + \sqrt{3}|\mu|)$ over half angular ranges, and are given by [19, 20]

$$\frac{1}{c}\partial_t (K_g^{\pm}\phi_g^{\pm}) \pm \partial_x (H_g^{\pm}\phi_g^{\pm}) + \sigma_g K_g^{\pm}\phi_g^{\pm} = \sigma_g B_g \,, \tag{14}$$

D. Anistratov & T. Haut, Multilevel Method for Thermal Radiative Transfer

$$\gamma = \left(1 + \frac{1}{2}\sqrt{3}\right)^{-1}.\tag{15}$$

The closures for the low-order equations (14) are formulated by the factors

$$K_g^{\pm} = \gamma \frac{\int_0^{\pm 1} \left(1 + \sqrt{3}|\mu|I_g\right) d\mu}{\int_0^{\pm 1} I_g d\mu}, \quad H_g^{\pm} = \gamma \frac{\int_0^{\pm 1} |\mu| \left(1 + \sqrt{3}|\mu|I_g\right) d\mu}{\int_0^{\pm 1} I_g d\mu}.$$
 (16)

The group radiation energy density and flux (Eq. (13)) are defined with the LONWF solution as follows:

$$E_g = \frac{1}{c}(\phi_g^- + \phi_g^+), \quad F_g = G_g^- \phi_g^- + G_g^+ \phi_g^+, \tag{17}$$

where

$$G_g^{\pm} = \frac{\int_0^{\pm 1} \mu I_g d\mu}{\int_0^{\pm 1} I_g d\mu}.$$
 (18)

The projection with the angular weight $w(\mu)$ leads to the low-order equations which meet the asymptotic diffusion limit in the interior of a diffusive domain with asymptotic boundary conditions. The low-order equations (14) and high-order RTE can be approximated by independent discretization schemes.

The first step in deriving the effective GLOQD equations is integrating the RTE over $-1 \le \mu \le 1$ with weigh 1 and μ and introducing exact closure by means of the group Eddington factor

$$f_g = \frac{\int_{-1}^{1} \mu^2 I_g d\mu}{\int_{-1}^{1} I_g d\mu}$$
(19)

to obtain multigroup moment equations for E_g and F_g . On the next step, the moment equations are summed over photon energy groups to get the effective grey moment equations on the coarse photon energy grid $\Lambda_{h\nu}^*$ with just one group $(n_{\nu} = 1)$ for the total radiation energy density and flux (Eq. (12)). The effective GLOQD equations are defined by [24, 26–28]:

$$\partial_t E + \partial_x F + c\bar{\sigma}_E E = c\bar{\sigma}_B a_R T^4 , \qquad (20a)$$

$$\frac{1}{c}\partial_t F + c\partial_x(\bar{f}E) + \bar{\sigma}_R F + \bar{\eta}E = 0, \qquad (20b)$$

where

$$\bar{\sigma}_E = \frac{\sum_{g=1}^{n_\nu} \sigma_g E_g}{\sum_{g=1}^{n_\nu} E_g}, \quad \bar{\sigma}_B = \frac{\sum_{g=1}^{n_\nu} \sigma_g B_g}{\sum_{g=1}^{n_\nu} B_g}, \quad \bar{\sigma}_R = \frac{\sum_{g=1}^{n_\nu} \sigma_g |F_g|}{\sum_{g=1}^{G} |F_g|}$$
(21)

are different grey absorption coefficients.

$$\bar{f} = \frac{\sum_{g=1}^{n_{\nu}} f_g E_g}{\sum_{g=1}^{n_{\nu}} E_g}$$
(22)

is the grey Eddington factor. It is a spectrum average of group Eddington factors. The compensation term in Eq. (20b) is defined by

$$\bar{\eta} = \frac{\sum_{g=1}^{n_{\nu}} (\sigma_g - \bar{\sigma}_R) F_g}{\sum_{g=1}^{n_{\nu}} E_g} \,.$$
(23)

4

The MEB equation in the effective grey form is given by

$$\partial_t \varepsilon(T) = c \left(\bar{\sigma}_E E - \bar{\sigma}_B a_R T^4 \right). \tag{24}$$

Finally, the multilevel method is defined by the hierarchy of equations the general form of which is the following:

• the multigroup RTE

$$\frac{1}{c}\partial_t I_g + \mathcal{L}I_g = q_g, \quad g \in \mathbb{N}(n_\nu), \qquad (25)$$
$$\mathcal{L} = \mathcal{L}(T), \quad q_g = q_g(T),$$

• the multigroup LONWF equations

$$\partial_t (\mathcal{K} \mathbf{\Phi}_g) + \mathcal{P} \mathbf{\Phi}_g = \mathbf{Q}_g, \quad \mathbf{\Phi}_g = (\phi_g^-, \phi_g^+)^\top, \quad \mathbf{Q}_g = \mathbf{Q}_g(T),$$
(26)
 $\mathcal{P} = \mathcal{P}(T),$

• the GLOQD equations

$$\partial_t \mathbf{Y} + \mathcal{M} \mathbf{Y} = \mathbf{Q}, \quad \mathbf{Y} = (E, F)^{\top}, \qquad (27)$$
$$\mathcal{M} = \mathcal{M}(\mathbf{\Phi}, T), \quad \mathbf{Q} = \mathbf{Q}(T), \quad \mathbf{\Phi} = \{\mathbf{\Phi}_g\}_{g=1}^{n_{\nu}}$$

• the MEB equation

$$\partial_t \varepsilon = \mathcal{R} \mathbf{Y} - Q_{pl},$$
(28)
 $\mathcal{R} = \mathcal{R}(\mathbf{\Phi}, T), \quad Q_{pl} = Q_{pl}(T).$

In this study, we discretize this system of equations with a fully-implicit temporal scheme based on the backward Euler time-integration method. The RTE is approximated in space by the simple corner balance scheme [29]. The lumped linear discontinuous (LLD) scheme is used to discretize the LONWF equations [19, 20, 30]. The GLOQD equations are approximated by the second-order finite volume method [28]. We note that independent spatial discretization schemes are applied for approximating the RTE, LONWF, and GLOQD equations [12].

3. Iteration Algorithm

Algorithm 1 presents the multilevel method with two photon energy grids for solving the low-order equations (26) and (27) coupled with MEB equation (28). On each time step, the outer iteration is the transport iteration. s is the index of the outer iterations. Note that there are no transport sweeps for s = 0. The inner iteration cycles consist of solving (i) the system of the group LONWF equations on the given grid $\Lambda_{h\nu}$ and (ii) the coupled GLOQD and MEB equations on the coarse grid $\Lambda^*_{h\nu}$. ℓ is the index of the inner iteration cycles. This inner iteration scheme is the multigrid V-cycle over two photon energy grids [13, 31]. Figure 1 shows the diagram of the V-cycle. The coupled GLOQD and MEB are linearized and use the Fréchet derivative of $\bar{\sigma}_E$ to account for variation in group absorption coefficients and spectrum due to change in temperature [13]. On every cycle, the linearized system of the GLOQD and MEB equations is solved with one Newton iteration. Algorithm 1: The multilevel method with two photon energy groups and V-cycle

while
$$t^{j} \leq t^{end}$$
 do
 $s = 0, T^{(0)} = T^{j-1}, K_{g}^{\pm (1/2)} = K_{g}^{\pm j-1}$
while $||T^{(s)} - T^{(s-1)}|| > \epsilon ||T^{(s)}|| \vee ||E^{(s)} - E^{(s-1)}|| > \epsilon ||E^{(s)}||$ do
if $s > 0$ then
update group absorption coefficients $\sigma_{g}(T^{(s)})$
solve $c^{-1}\partial_{t}I_{g} + \mathcal{L}I_{g} = q_{g}$ for $I_{g}^{(s+1/2)}$ on $\Lambda_{h\nu}$ for all g
compute group factors $K_{g}^{\pm (s+1/2)}, H_{g}^{\pm (s+1/2)}, G_{g}^{\pm (s+1/2)}, \text{and } f_{g}^{(s+1/2)}$
while $\ell \leq \ell_{max} \wedge \left(||T^{(\ell,s)} - T^{(\ell-1,s)}|| > \tilde{\epsilon}||T^{(\ell,s)}|| \vee ||E^{(\ell,s)} - E^{(\ell-1,s)}|| > \tilde{\epsilon}||E^{(\ell,s)}|| \right)$
do
given: $T^{(\ell,s)}$
update group absorption coefficients $\sigma_{g}(T^{(\ell,s)})$
solve $\overline{\partial_{t}(\mathcal{K}_{g}\Phi_{g}) + \mathcal{P}_{g}\Phi_{g} = \mathbf{Q}_{g}}$ for $\phi_{g}^{\pm (\ell,s)}$ on $\Lambda_{h\nu}$ for all g
compute $E_{g}^{(\ell,s)}$ and $F_{g}^{(\ell,s)}$
compute $\overline{\sigma}_{E}^{(\ell,s)}, \overline{\sigma}_{R}^{(\ell,s)}, \overline{f}^{(\ell,s+1/2)}, \overline{\eta}^{(\ell,s+1/2)},$ and Fréchet derivative $\delta_{T}\overline{\sigma}_{E}^{(\ell,s)}$
solve $\overline{\partial_{t}\mathbf{Y} + \overline{\mathcal{M}}\mathbf{Y} = \mathbf{Q} - \partial_{t}\varepsilon = \mathcal{R}\mathbf{Y} - Q_{pl}$ on $\Lambda_{h\nu}^{*}$ for $T^{(\ell+1,s)}, E^{(\ell+1,s)}, F^{(\ell+1,s)}$



Figure 1: Diagram of the V-cycle of inner iterations. Φ_g - calculation of spectrum functions on the fine grid by solving the multigroup LONWF equations, T - calculation of temperature by solving coupled GLOQD and MEB equations on the coarse grid.

4. Numerical Results

The convergence of the presented iteration algorithm is analyzed on the Fleck-Cummings (FC) test [32]. It is defined for a slab ($0 \le x \le 4$ cm). The analytic spectral absorption coefficient of the material is given by

$$\sigma_{\nu}(T) = \frac{27}{(h\nu)^3} \left(1 - e^{-\frac{h\nu}{kT}} \right).$$
(29)

The incoming radiation on the left boundary has the black-body spectrum at $kT_b=1$ keV and hence $I_g^+ = B_g(T_b)$. The right boundary is vacuum. At t = 0, the temperature in the slab equals

 $kT_0=10^{-3}$ keV and the group intensity at the initial instant is $I_g^0 = B_g(T_0)$. The material energy is a linear function and given by $\varepsilon(T) = c_v T$, where $c_v = 0.5917 a_R T_b^3$. The spatial mesh is uniform with 10 intervals. The time interval of the problem is $t \in [0, 0.3 \text{ sh}]$ (1 $sh = 10^{-8}$ sec). The photon energy grid consists of $n_{\nu} = 256$ groups over the interval $h\nu \in [0, 10^7]$ keV. $n_{\nu} - 2$ groups are evenly spaced in logarithmic scale between $h\nu_a = 10^{-4}$ keV and $h\nu_b = 10$ keV. The angular grid has 16 angular directions defined by the double S₈ Gauss-Legendre quadrature set. The parameters of convergence criteria are $\epsilon = 10^{-6}$ and $\tilde{\epsilon} = 10^{-7}$. Figure 2 shows temperature and total energy density at various instants of time obtained with $\Delta t = 2 \times 10^{-3}$ sh and illustrates evolution of heat and radiation waves.



Figure 2: Numerical solution of the FC test with 256 groups using $\Delta t = 2 \times 10^{-3}$ sh.

Table 1 presents the total numbers of transport iterations, N_{ti} , and cycles, N_c , accumulated over all time steps when the problems is solved with $\Delta t = 1 \times 10^{-3}, 2 \times 10^{-3}, 4 \times 10^{-3}$ sh.

- $\Delta t = 1 \times 10^{-3}$ sh. The algorithm with 4 cycles per transport iteration yields the minimum number of transport iterations ($N_{ti} = 653$) over the time range of the problem. The algorithm with 3 cycles per transport iteration leads to extra 5 transport iterations with smaller total number of cycles. These two algorithms have comparable computational costs.
- $\Delta t = 2 \times 10^{-3}$ sh. The algorithm with 4 cycles per transport iteration has the optimum number of transport iterations ($N_{ti} = 384$). The algorithm with 3 cycles needs extra 15 transport iterations.
- $\Delta t = 4 \times 10^{-3}$ sh. The algorithm with 5 cycles per transport iteration requires the minimum number of transport iterations ($N_{ti} = 232$).

Figures 3-5 show extra details of performance of algorithm with different values of ℓ_{max} . The plots present the total numbers of transport iterations, M_{ti} , and cycles, M_c , at each time step. There are several different stages in Marshak wave evolution in this test problem. At the initial stage, the rate of change of the solution is large. This leads to more transport iterations and cycles compared to the next stage during which we observe a well developed wave. At the last stage, the solution approaches a steady-state regime. The number of transport iterations is very small. Figures 3-5 demonstrate these effects as well as influence of the time step value.

	$\Delta t = 1 \times 10^{-2} sh$		$\Delta t = 2 \times 10^{-2} sh$		$\Delta t = 4 \times 10^{-2} sh$	
ℓ_{max}	N_{ti}	N_c	N_{ti}	N_c	N _{ti}	N_c
2	1208	2970	653	1605	432	1014
3	658	2408	399	1554	300	1109
4	653	3010	384	1789	249	1215
5	653	3297	384	2023	232	1278
6	653	3509	384	2283	232	1490
7	653	3534	384	2443	232	1700

Table 1: Performance of algorithms in the FC test with 256 groups using different time steps Δt



Figure 3: Number of transport iterations (M_{ti}) and number of cycles (M_c) at each time step in the FC test with 256 groups and $\Delta t = 1 \times 10^{-3}$ sh over $t \in [0, 0.3 \text{ sh}]$.



Figure 4: Number of transport iterations (M_{ti}) and number of cycles (M_c) at each time step in the FC test with 256 groups and $\Delta t = 2 \times 10^{-3}$ sh over $t \in [0, 0.3 \text{ sh}]$.



Figure 5: Number of transport iterations (M_{ti}) and number of cycles (M_c) at each time step in the FC test with 256 groups and $\Delta t = 4 \times 10^{-3}$ sh over $t \in [0, 0.3 \text{ sh}]$.

5. Conclusions

We developed a new multilevel iteration method for solving multigroup TRT problems in 1D slab geometry. The obtained results are promising. The algorithm is rapidly convergent. It is stable with independent discretization of the high-order RTE and low-order equations at each level. Efficiency of the algorithm can be optimized by choosing the number of V-cycles per outer (transport) iteration. Further detailed analysis of iteration algorithm is necessary. Future work will include development of methods with the mixed type low-order problems using multiple grids in photon energy and various multigrid cycles to further optimize performance of iteration schemes and computational cost. Various discretization schemes will be considered. The proposed multilevel method can be extended to multidimensional geometries, and is suitable for computations on high-performance computing systems with GPUs.

Acknowledgements

This manuscript has been authored by Lawrence Livermore National Security, LLC under Contract No. DE-AC52-07NA27344 with the US. Department of Energy. The United States Government retains, and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes. LLNL-CONF-2001051.

References

- R. P. Drake, High Energy Density Physics: Fundamentals, Inertial Fusion and Experimental Astrophysics, Springer, 2006.
- [2] Y. B. Zeldovich, Y. P. Razier, Physics of Shock Waves and High Temperature Hydrodynamic Phenomena, Academic, New York, 1966.
- [3] J. E. Morel, E. W. Larsen, M. K. Matzen, A synthetic acceleration scheme for radiative diffusion calculations, J. Quantitative Spectroscopy and Radiative Transfer 34 (1985) 243–261.

- [4] E. W. Larsen, A grey transport acceleration method for time-dependent radiative transfer problems, Journal of Computational Physics 78 (1988) 459–480.
- [5] J. E. Morel, T.-Y. B. Yang, J. S. Warsa, Linear multifrequency-grey acceleration recast for preconditioned Krylov iterations, Journal Computational Physics 227 (2007) 244–263.
- [6] A. T. Till, M. L. Adams, J. E. Morel, Application of nonlinear Krylov acceleration to radiative transfer problems, in: Proc. of Int. Conf. on Math. and Comp., M&C 2013, Sun Valley, Idaho, 2013, pp. 2690–2701.
- [7] H. Park, D. A. Knoll, R. M. Rauenzahn, A. B. Wollaber, J. D. Densmore, Moment-based, multiscale solution approach for thermal radiative transfer problems, Transport Theory and Statistical Physics 41 (2012) 284–303.
- [8] B. C. Yee, A. B. Wollaber, T. S. Haut, H. Park, A stable 1D multigroup high-order low-order method, Journal of Computational and Theoretical Transport 46 (2016) 46–76.
- [9] A. B. Wollaber, H. Park, R. B. Lowrie, R. M. Rauenzahn, M. A. Cleveland, Multigroup radiation hydrodynamics with a high-order-low-order method, Nuclear Science and Engineering 185 (2017) 117–129.
- [10] V. Y. Gol'din, A quasi-diffusion method of solving the kinetic equation, USSR Comp. Math. and Math. Phys. 4 (1964) 136–149.
- [11] V. Y. Gol'din, On mathematical modeling of problems of non-equilibrium transfer in physical systems, in: Modern Problems of Mathematical Physics and Computational Mathematics, Nauka, Moscow, 1982, pp. 113–127, in Russian.
- [12] D. Y. Anistratov, V. Ya. Gol'din, Nonlinear methods for solving particle transport problems, Transport Theory and Statistical Physics 22 (1993) 42–77.
- [13] D. Y. Anistratov, Nonlinear iterative projection methods with multigrid in photon frequency for thermal radiative transfer, Journal of Computational Physics 444 (2021) 110568.
- [14] S. Olivier, W. Pazner, T. S. Haut, B. C. Yee, A family of independent variable Eddington factor methods with efficient preconditioned iterative solvers, Journal of Computational Physics 473 (2023) 111747.
- [15] S. Olivier, T. S. Haut, High-order mixed finite element variable Eddington factor methods, Journal of Computational and Theoretical Transport 52 (2023) 79–142.
- [16] T. A. Germogenova, Convergence of some approximate methods of solving the transport equation, Soviet Mathematics - Doklady 9 (1968) 855–858.
- [17] V. Ya. Gol'din, G. V. Danilova, B. N. Chetverushkin, Approximate method of solving the time-dependent kinetic equation, Computational Methods in Transport Theory 3 (1969) 34–46, in Russian.
- [18] V. V. Gorskii, S.T. Surzhikov, Use of the semimoment method to solve the shock layer radiative heat-transfer problem, J. Eng. Physics 42 (1982) 108–111.
- [19] D. Y. Anistratov, E. W. Larsen, Nonlinear and linear α-weighted methods for particle transport problems, Journal Computational Physics 173 (2001) 664–684.
- [20] L. Roberts, D. Y. Anistratov, Nonlinear weighted flux methods for particle transport problems, Transport Theory and Statistical Physics 36 (2007) 589–608.
- [21] L. Roberts, D. Y. Anistratov, Nonlinear weighted flux methods for solving the transport equation in 2D Cartesian geometry, Nuclear Science and Engineering 165 (2010) 133–148.
- [22] E. Shands, J. Hanophy, J. E. Morel, A modified S₂ method for calculating the uncollided flux from a point source without ray effects, in: Proc. of M&C 2023, Int. Conf. on Math. and Comp. (M&C), to Nuclear Science and Engineering, Niagara Falls, Ontario, Canada, 2023, p. 9.
- [23] L. H. Auer, D. Mihalas, On the use of variable Eddington factors in non-LTE stellar atmospheres computations, Monthly Notices of the Royal Astronomical Society 149 (1970) 65–74.
- [24] V. Y. Gol'din, D. A. Gol'dina, A. V. Kolpakov, A. V. Shilkov, Mathematical modeling of hydrodynamics processes with high-energy density radiation, Problems of Atomic Sci. & Eng.: Methods and Codes for Numerical Solution of Math. Physics Problems 2 (1986) 59–88, in Russian.
- [25] K.-L. A. Winkler, M. L. Norman, D. Mihalas, Implicit adaptive-grid radiation hydrodynamics, in: Multiple Time Scales, Academic Press, 1985, pp. 145–184.
- [26] D. Y. Anistratov, E. N. Aristova, V. Y. Gol'din, A nonlinear method for solving problems of radiation transfer in a physical system, Mathematical Modeling 8 (1996) 3–28, in Russian.
- [27] E. N. Aristova, V. Ya. Gol'din, A. V. Kolpakov, Multidimensional calculations of radiation transport by nonlinear quasi-diffusion method, in: Proc. of Int. Conf. on Math. and Comp., M&C 1999, Madrid, Spain, 1999, pp. 667–676.
- [28] D. Y. Anistratov, Stability analysis of a multilevel quasidiffusion method for thermal radiative transfer problems, Journal of Computational Physics 376 (2019) 186–209.
- [29] M. L. Adams, Subcell balance methods for radiative transfer on arbitrary grids, Transport Theory and Statistical Physics 26 (1997) 385–431.

- [30] E. W. Larsen, J.E. Morel, Asymptotic solutions of numerical transport problems in optically thick, diffusive regimes II, Journal of Computational Physics 83 (1989) 212–236.
- [31] U. Trottenberg, C. Oosterlee, A. Schuller, Multigrid, Academic Press, San Diego San Francisco New York Boston London Sydney Tokyo, 2000.
- [32] J. A. Fleck, J. D. Cummings, An implicit Monte Carlo scheme for calculating time and frequency dependent nonlinear radiation transport, Journal of Computational Physics 8 (1971) 313–342.