Algorithmic Universality, Low-Degree Polynomials, and Max-Cut in Sparse Random Graphs

Houssam El Cheairi^{*}, David Gamarnik[†]

December 25, 2024

Abstract

Universality, namely distributional invariance, is a well-known property for many random structures. For example it is known to hold for a broad range of variational problems with random input. Much less is known about the universality of the performance of specific algorithms for solving such variational problems. Namely, do algorithms tuned to specific variational tasks produce the same asymptotic answer regardless of the underlying distribution?

In this paper we show that the answer is yes for a class of models, which includes spin glass models and constraint satisfaction problems on sparse graphs, provided that an algorithm can be coded as a lowdegree polynomial (LDP). We illustrate this specifically for the case of the Max-Cut problem in sparse Erdös-Rényi graph $\mathcal{G}(n, d/n)$. We use the fact that the Approximate Message Passing (AMP) algorithm, which is an effective algorithm for finding near-ground state of the Sherrington-Kirkpatrick (SK) model, is well approximated by an LDP. We then establish our main universality result: the performance of the LDP based algorithms exhibiting certain connectivity property, is the same in the mean-field (SK) and in the random graph $\mathcal{G}(n, d/n)$ setting, up to an appropriate rescaling. The main technical challenge which we address in this paper is showing that the output of the LDP algorithm on $\mathcal{G}(n, d/n)$ is truly discrete, namely it is close to the set of points in the binary cube. Our result in particular recovers an analogue of the main result in [EMS23]. There the authors construct a local near optimum algorithm for the Max-Cut problem in regular graphs with large girth. Here we establish a similar result for the $\mathcal{G}(n, d/n)$ random graph model.

Contents

1	Introduction and Problem Statement	2
	1.1 Algorithmic universality. The main result	3
	1.2 Notation	5
2	2 Universality of Low-Degree Polynomials. Main Results	6
3	Preliminary Results	8
	3.1 Some elementary estimates	8
	3.2 Probabilistic properties of LDP	13
4	Proof of Theorem 2.1	20
	4.1 Proof of (2.5)	20
	4.2 Proof of (2.6)	25
	4.3 Proof of (2.7)	
	*MIT Emoil house of the du	

*MIT. Email: houssamc@mit.edu

[†]MIT. Email : gamarnik@mit.edu

5	IAM	IP Representation and Proof of Theorem 2.2	29
	5.1	Approximation of IAMP by Polynomials	29
	5.2	Approximation of AMP with Connected Tree-based Low-Degree Polynomials	32
	5.3	Proof of Theorem 2.2.	34
	5.4	Proof of Proposition 5.5	35

1 Introduction and Problem Statement

Universality in random structures is a well known phenomena stating that many large scale observables of the system are distributionally invariant. Namely they remain the same upon changing the distribution, provided that the first order parameters such as first two moments remain invariant. The Central Limit Theorem is the textbook example of universality.

It also well known that universality holds for variational problems (optimization values) associated with random structures. Loosely speaking it says that given some objective function (Hamiltonian) $H(\boldsymbol{\sigma}, \mathbf{X})$ which is a function of decision variables σ and random input X distributed according to some distribution F, say Gaussian, the value $\max_{\sigma} H(\sigma, \mathbf{X})$ remains asymptotically the same when F changes to a different distribution F', with the same first two moments as F (and some mild condition on higher moments). An example which is the most relevant to the present paper is the ground state of the Sherrington-Kirkpatrick (SK) model. Namely, let X be an $n \times n$ random symmetric matrix with i.i.d. mean zero, variance 1/n Gaussian entries in the upper triangular part and 2/n variance on diagonal, also known as Gaussian Orthogonal Ensemble, which we denote by GOE(n). Let also $H(\boldsymbol{\sigma}, \mathbf{X}) = \boldsymbol{\sigma}^T \mathbf{X} \boldsymbol{\sigma}$, where $\boldsymbol{\sigma}$ ranges in $\{\pm 1\}^n$. The associated ground state value is then defined as $\frac{1}{2n} \max_{\sigma} \sigma^T \mathbf{X} \sigma$ which is known to converge to some limit \mathcal{P}^* called the *Parisi* constant [Par79],[Tal06],[Pan13], with high probability (whp) as $n \to \infty$. The universality property refers to the fact that the convergence to the same constant \mathcal{P}^* takes place regardless of the distribution of the entries of \mathbf{X} provided matching of the first two moments and some additional mild conditions. This fact can be established in a relatively straightforward way when the distributions F and F' governing entries of **X** do not depend on *n*. In particular, in this case one can use the Lindeberg's method applied to "softened" max function, namely log-partition function $\frac{1}{\beta} \log \left(\sum_{\boldsymbol{\sigma}} \exp(\beta \boldsymbol{\sigma}^T \mathbf{X} \boldsymbol{\sigma}) \right)$, which approximates the max function for large β . Lindeberg's method is based on interpolating between two distributions entry by entry of **X** and using 2nd order Taylor approximation, see for example [Cha06].

The universality property and its proof based on Lindeberg's interpolation also carries over to a far less trivial case when the modified distribution F' is *n*-dependent, for example when it corresponds to sparse matrices. The case in point is when F' is defined by $\mathbf{Y}_{ij} = 1$ with probability d/n and = 0 otherwise, for some *n*-independent parameter *d*. In other words when \mathbf{Y} is the adjacency matrix of a sparse Erdös-Rényi graph $\mathcal{G}(n, d/n)$. To match the mean and variance, one considers instead $\hat{\mathbf{Y}}$ with

$$\hat{\mathbf{Y}}_{ij} = -\frac{1}{\sqrt{d(1-\frac{d}{n})}} \left(\mathbf{Y}_{ij} - \frac{d}{n} \right).$$
(1.1)

It is known [AMS17] and [Sen18] that in this case

$$\underset{d \to \infty, n \to \infty}{\text{p-lim}} \max_{\boldsymbol{\sigma}} \boldsymbol{\sigma}^T \hat{\mathbf{Y}} \boldsymbol{\sigma} = \mathcal{P}^*$$
(1.2)

as well. Furthermore, this result can be used to obtain an asymptotic value of the Maximum Cut (Max-Cut) problem in the graph $\mathcal{G}(n, d/n)$. To explain this connection lets recall the Max-Cut problem.

Given a graph G = (V, E) with node set V = [n], with $[n] \triangleq 1, 2, ..., n$, and edge set E, the Max-Cut problem on G is the following constrained optimization task

$$\begin{array}{ll} \text{maximize} & \sum_{(i,j)\in E} \frac{1-\sigma_i \sigma_j}{2} \\ \text{subject to} & \boldsymbol{\sigma} \in \{+1,-1\}^n. \end{array}$$

$$(1.3)$$

We note that the objective function above is $|E|/2 - (1/4)H(\boldsymbol{\sigma}, \mathbf{Y})$. The Max-Cut problem can then be reformulated as

maximize
$$\frac{|E|}{2} - \frac{1}{4}H(\boldsymbol{\sigma}, \mathbf{Y})$$

subject to $\boldsymbol{\sigma} \in \{+1, -1\}^n$. (1.4)

Namely Max-Cut is equivalent to minimizing the Hamiltonian $H(\cdot, \mathbf{Y})$ over $\{\pm 1\}^n$. The identity (1.2) then implies the following asymptotic value of the Max-Cut problem in $\mathcal{G}(n, d/n)$: with high probability (whp)

$$\frac{\operatorname{MaxCut}(\mathcal{G}(n,d/n))}{n} = \frac{d}{4} + \mathcal{P}^* \sqrt{\frac{d}{4}} + o_d(\sqrt{d}).$$

1.1 Algorithmic universality. The main result

A totally different matter is the universality of algorithms. Suppose one runs an algorithm \mathcal{A} for some input $\mathbf{X} = \mathbf{X}(F)$ distributed as F and obtains a solution $\sigma_{ALG}(\mathbf{X})$. Suppose now \mathbf{Y} is generated according to distribution F' which is different from F, but shares the first two moments with F. Is it the case that

$$H(\sigma_{ALG}(\mathbf{Y})) \approx H(\sigma_{ALG}(\mathbf{X}))$$
? (1.5)

In particular is $\sigma_{ALG}(\mathbf{Y})$ near optimal when $\sigma_{ALG}(\mathbf{X})$ is? An evidence supporting this conjecture is found in [EMS23]. Specifically, there the authors adopt the algorithm first developed in [Mon21] for the SK model, to the same model but defined now on sparse regular locally tree-like graphs.

Both the original and the adopted algorithm is based on the so-called Approximate Message Passing (AMP) scheme (see the next section), which found to be a very effective in a broad range of models [Fen22]. In [EMS23] the authors reproduce the AMP scheme and prove that the output of their algorithm is near optimum as well in the double limit $n \to \infty, d \to \infty$. In particular (1.5) is validated in this setting. This required however reproducing the optimality proof for their algorithm in the new setting. A similar universality result was established recently in the quantum setting in [Bas22]. There the universality was also verified directly by computing the performance of an algorithm for the SK model on the one hand, and a random regular graph on the other hand, and verifying that the results match. If the algorithmic universality (1.5) was known to hold for these model though, the need for reproducing the proof would have been eliminated.

In this paper we establish the algorithmic universality (1.5) for the specific class of algorithms known as Low-Degree-Polynomials (LDP), exhibiting certain connectivity property to be defined. Our universality result is established specifically for the SK and $\mathcal{G}(n, d/n)$ settings above, but we believe that our technique can be easily extended to a broader range of Hamiltonians H. We will elaborate on this after we state our main result below informally. Next, we use the fact that AMP can be approximated by an LDP-based algorithm. While variants of this are known in the literature [Mon22],[IS23], we prove this fact in this paper for completeness. Combining these two results, we obtain an LDP-based algorithm for solving the Max-Cut problem to near optimality in $\mathcal{G}(n, d/n)$, similarly to the main result in [EMS23] which concerned sparse regular graphs. Our analysis unfortunately does not carry through in the case of regular graphs for technical reasons. Specifically, while for a regular graph, we may also naturally define rescaling variables $\hat{\mathbf{Y}}_{ij} = \frac{\mathbf{Y}_{ij} - \frac{d}{n-1}}{\sqrt{\frac{nd}{n-1}(1-\frac{d}{n-1})}}$, these terms are no longer independent which creates difficulties in bounding moments $\mathbb{E}[\prod_{(i,j)\in S} \hat{\mathbf{Y}}_{ij}]$ for subsets of edges S.

Before we state our main result informally we comment on some existing literature for algorithmic universality, which to the best of our knowledge is rather limited. It is known that AMP is a universal algorithm [BMLM15]. However this was established for a simpler setting of the AMP than the one appearing in [EMS23], and importantly for the *n*-independent case (in particular not applicable to the $\mathcal{G}(n, d/n)$). We leave it as a separate interesting question as to whether one can find a simple and direct argument proving universality of AMP in our *n*-dependent case.

As mentioned earlier, algorithmic universality was also established in the quantum setting for the SK and $\mathcal{G}(n, d/n)$ models in the same context of ground states/Max-Cut problem for the class of quantum algorithms known as Quantum Approximate Optimization Algorithms (QAOA) [Bas22]. This was achieved by a direct

computation of the performance of this algorithm in the two settings and verifying that the outputs are nearly identical. It is an interesting challenge to establish the same universality but from general principles. Unfortunately, the quantum nature of the QAOA algorithm precludes setting it up as a LDP, though a step in this direction was taken in [AM23].

Regarding the non-algorithmic/non-variational universality results, the literature is naturally much broader. Close to our setting, universality for low degree polynomials was established in [MOO10] in the context of the so-called Majority is Stablest conjecture. There the universality was also proven in a simpler *n*-independent setting, and for the case when the LDP is the value function itself rather than an algorithmic tool (that is, not in a variational setting). One can view our result as a generalization of results in [MOO10] to the more challenging *n*-dependent distributional setting. Finally, a reference loosely related to algorithmic universality is [Dei14], where the universality of the stopping time of a numerical procedure for solving a differential equation system is established.

We now describe our main result informally. We begin by describing the class of LDP-based algorithms. This is a class of algorithms viewed as mapping $\mathbb{R}^{n \times n} \to \{\pm 1\}^n$ constructed as follows. One fixes a vector $p = (p_i, i \in [n])$ of n multi-variable polynomials $p_i, i \in [n]$ in variables $\mathbf{x}_{ij}, 1 \leq i \leq j \leq n$. The degree $\Delta = \Delta(p)$ of this polynomial is the largest size (the number of terms) of its monomials. Let $\mathbf{x} = (\mathbf{x}_{ij}, 1 \leq i < j \leq n)$. $i < j \leq n$. The vector p induces a mapping $\mathbb{R}^{n \times n} \to \mathbb{R}^n$ via $\mathbf{x} \to p(\mathbf{x}) = (p_i(\mathbf{x}), 1 \leq i \leq n)$.

The projection $\mathbb{R}^n \to \{\pm 1\}^n$ is obtained first by truncating values outside [-1, 1] to ± 1 and then increasing values inside (-1, 1) to ± 1 sequentially while increasing the objective value. The resulting map is denoted by $\sigma(\operatorname{ALG}(p), \mathbf{x})$. We say that p is a low degree (LDP) connected tree-based polynomial if (a) Δ is bounded by n-independent constant, (b) the degree of every variable \mathbf{x}_{ij} in every monomial is at most 2, and (c) the graphs on the node set [n] induced by the monomials in $p_i, i \in [n]$ are connected trees. Part (c) will be made more precise in the next section. We now state informally the main result of this paper. We denote by p-lim, p-lim convergence in probability.

Theorem 1.1. [Informal] Let $\mathbf{X} = (\mathbf{X}_{ij}, 1 \le i < j \le n)$ consist of i.i.d. Gaussian entries with mean zero and variance 1/n. Let \mathbf{Y} consist of i.i.d. random variables with $\mathbb{P}(\mathbf{Y}_{ij} = 1) = d/n, \mathbb{P}(\mathbf{Y}_{ij} = 0) = 1 - d/n$, and define $\hat{\mathbf{Y}} = (\hat{\mathbf{Y}}_{ij}, 1 \le i < j \le n)$ as in (1.1). Let p be a connected tree-based LDP. Suppose

$$\operatorname{p-lim}_{n \to \infty} \frac{1}{2n} \sigma(\operatorname{ALG}(p), \mathbf{X})^\top \mathbf{X} \sigma(\operatorname{ALG}(p), \mathbf{X}) \geq \mathcal{P},$$

for some constant \mathcal{P} . Then also

$$\underset{d\to\infty,n\to\infty}{\text{p-lim}}\frac{1}{2n}\sigma(\text{ALG}(p),\hat{\mathbf{Y}})^{\top}\hat{\mathbf{Y}}\sigma(\text{ALG}(p),\hat{\mathbf{Y}}) \geq \mathcal{P}.$$

The next result states that roughly speaking connected tree-based LDP achieve near optimality value \mathcal{P}^* .

Theorem 1.2. [Informal] For every $\epsilon > 0$ there exists a connected tree-based LDP such that

$$\operatorname{p-lim}_{n \to \infty} \frac{1}{2n} \sigma(\operatorname{ALG}(p), \mathbf{X})^{\top} \mathbf{X} \sigma(\operatorname{ALG}(p), \mathbf{X}) \geq \mathcal{P}^* - \epsilon,$$

where \mathbf{X} is as in Theorem 1.1.

Combining these two results we obtain

Corollary 1.3. For every $\epsilon > 0$ there exists a connected tree-based LDP such that

$$\underset{d\to\infty,n\to\infty}{\text{p-lim}}\frac{1}{2n}\sigma(\text{ALG}(p),\hat{\mathbf{Y}})^{\top}\hat{\mathbf{Y}}\sigma(\text{ALG}(p),\hat{\mathbf{Y}}) \geq \mathcal{P}^* - \epsilon,$$

where $\hat{\mathbf{Y}}$ is as in Theorem 1.1. In particular there exists a LDP-based algorithm which solves the Max-Cut problem for $\mathcal{G}(n, d/n)$ near optimally as n and d increase. Namely, if $\hat{\mathbf{Y}}$ is the normalized adjacency matrix of the sparse graph, and $\operatorname{CUT}(\sigma(\operatorname{ALG}(p), \hat{\mathbf{Y}}))$ is the cut value obtained from $\sigma(\operatorname{ALG}(p), \hat{\mathbf{Y}})$, then

$$\underset{d \to \infty, n \to \infty}{\text{p-lim}} \frac{\text{CUT}(\sigma(\text{ALG}(p), \hat{\mathbf{Y}}))}{n} \ge \frac{d}{4} + (\mathcal{P}^* - \epsilon) \sqrt{\frac{d}{4}}.$$

We now describe the proof technique underlying our main result. The first element of the proof of our main result, Theorem 1.1 is the Lindeberg's interpolation method. The method is based on expanding the underlying function using second order Taylor expansion, taking advantage of the matching of the first two moments and then showing that the terms depending on higher moments are vanishing. This needs to be done both for the objective function H evaluated at the solution produced by LDP and importantly for the constraints which ensure that the solution remains close to $[-1, 1]^n$. While it is relatively easy to show that the objective value remains roughly the same in expectation when switching from the Gaussian to the sparse Bernoulli distribution underlying the $\mathcal{G}(n, d/n)$ model for the LDP based solutions, showing that this takes place with high probability is more challenging. Likewise, establishing that the vector of solutions remains close to $[-1,1]^n$ who is also challenging. We resolve this by introducing penalty functions, one penalty for deviating from the target value \mathcal{P} in the evaluation of the polynomial, and one penalty for deviating from the cube $[-1,1]^n$. We then conduct 3rd order Taylor expansion on these penalties to obtain the result.

While the derivation of our bounds is technically involved, it is fairly elementary and simple conceptually. Also, importantly, this analysis is conducted generically, *regardless* of the details of the algorithm giving rise to the underlying LDP. Namely, the analysis of the algorithmic performance is also universal in some sense.

We also stress that while our proof is laid out for the special case of the objective function H of the form $H(\boldsymbol{\sigma}, \mathbf{X}) = \boldsymbol{\sigma}^T \mathbf{X} \boldsymbol{\sigma}$, in fact our proof technique applies to a broader set of objectives, so long that they can be written as polynomials in σ themselves (the case above corresponding to the quadratic case). In particular, we anticipate that the same proof goes through for the case of p-spin models and their counterparts without change.

The proof of Theorem 1.2 is not particularly novel and versions of it can be found in [Mon22], [IS23]. Although it is quite lengthy we include it for convenience. The novel technical part of this result involves showing that the approximating polynomials can be taken to satisfy the required properties such as connectivity.

1.2Notation

We end the section with some notational conventions. We denote matrices and vectors by bold letters. We denote $[n] = \{1, ..., n\}$ for $n \in \mathbb{Z}_{\geq 0}$, and let $[0, n] = \{0\} \cup [n]$. For $\Delta \geq 0, i \in [n]$ we denote by $\mathcal{T}, \mathcal{T}_{\leq \Delta}, \mathcal{T}^i_{\leq \Delta}$ the set of trees on [n], the set of trees with at most Δ edges, and the set of rooted trees at i with at most Δ edges, respectively. Given $(i, j) \in [n]^2$, we denote by [ij] the elementary matrix $\mathbf{E}_{ij} \in \mathbb{R}^{n \times n}$, i.e the $n \times n$ matrix with 1 at entry (i, j) and 0 elsewhere. We write $k | \boldsymbol{\alpha}$ for $k \in \mathbb{Z}_{\geq 0}, \boldsymbol{\alpha} \in \mathbb{Z}_{\geq 0}^{n \times n}$ if and only if k is a divisor of $\boldsymbol{\alpha}_{ij} + \boldsymbol{\alpha}_{ji}$ for all $1 \leq i < j \leq n$. We denote by $\max(\boldsymbol{\alpha})$ the maximum of $\boldsymbol{\alpha}_{ij} + \boldsymbol{\alpha}_{ji}, 1 \leq i < j \leq n$. We let $\operatorname{Supp}(\boldsymbol{\alpha})$ denote the support of $\boldsymbol{\alpha}$, namely set of pairs (i, j) with $\boldsymbol{\alpha}_{i,j} > 0$. Given $\boldsymbol{\alpha} \in \mathbf{Z}_{\geq 0}^{n \times n}$ we denote by $\mathbf{X}^{\boldsymbol{\alpha}}$ the monomial $\prod_{1 \leq i < j \leq n} \mathbf{X}_{ij}^{\boldsymbol{\alpha}_{ij}}$.

We use the notation p-lim to denote convergence in probability. The asymptotic notations $o(.), O(.), \omega(.), \omega(.)$ and $\Omega(.)$ hide constants in n, d where n is the dimension of the disorder matrices we consider and d is a parameter for sparse disorders. We use $\tilde{o}(.), \hat{O}(.), \tilde{\omega}(.), \hat{\Omega}(.)$ to hide polylog asymptotic terms in d. We use \lesssim to hide polylog asymptotic terms in d in inequalities. We write $a_{n,d} \sim b_{n,d}$ if $a_{n,d} = (1 + o(1))b_{n,d}$. We denote by $|||_p, p \ge 0$ the usual ℓ_p norms, by $|||_{\text{fro}}$ the Frobenius norm, by $|||_{\text{op}}$ the operator norm, and by $d(\mathbf{z}, K)$ the ℓ_2 distance between z and the set K, whenever $K \neq \emptyset$.

Given a random variable X with distribution \mathcal{D} , we write $X \stackrel{\mathsf{d}}{=} \mathcal{D}$. We denote by $\operatorname{GOE}(n)$ the distribution on $n \times n$ symmetric matrices **X** where $\mathbf{X}_{ij} = \mathbf{X}_{ji}$ are independent and identically distributed as $\mathcal{N}(0, \frac{1}{n})$ for $1 \le i < j \le n$ and **X** has null diagonal. We denote by $\mathbf{Y} \stackrel{\mathsf{d}}{=} \mathcal{G}(n, d/n)$ a symmetric matrix with zeros on the diagonal and $\mathbb{P}(\mathbf{Y}_{ij} = 1) = d/n$, $\mathbb{P}(\mathbf{Y}_{ij} = 0) = 1 - d/n$, i.i.d. across $1 \le i < j \le n$. When $\hat{\mathbf{Y}}$ is defined via (1.1) we also write $\hat{\mathcal{G}}(n, d/n)$ for the distribution of $\hat{\mathbf{Y}}$. As we will not be working directly with the $\mathcal{G}(n, d/n)$ model, from this point on we use $\mathcal{G}(n, d/n)$ to denote $\hat{\mathcal{G}}(n, d/n)$ and **Y** to denote $\hat{\mathbf{Y}}$.

Given $m \in \mathbb{Z}_{\geq 0}$, we denote by $\mathcal{C}^m(\mathcal{D}, \mathbb{R})$ the set of *m*-times continuously differentiable functions from \mathcal{D} to \mathbb{R} . Given a differentiable function $f \in \mathcal{C}^m(\mathcal{D}, \mathbb{R})$ and $k \in [m]$ we denote by $f^{(k)}$ the k-th derivative of f. Given $\ell \in \mathbb{Z}_{\geq 0}$, we say that a function $f : \mathbb{R}^k \to \mathbb{R}$ is pseudo-Lipschitz of order ℓ (and write $f \in PL(\ell)$) if $|f(\mathbf{x}) - f(\mathbf{y})| \leq C(1 + ||\mathbf{x}||_2^{\ell-1} + ||\mathbf{y}||_2^{\ell-1}) ||\mathbf{x} - \mathbf{y}||_2$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^k$. For any matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$ and vector $\mathbf{x} \in \mathbb{R}^n$, we define $\sigma(\mathbf{x}, \mathbf{X}) \in \{\pm 1\}^n$ as follows. First set $\mathbf{x}' = (\mathbf{x}'_i, i \in [n])$ via $\mathbf{x}'_i = \min(1, \max(-1, \mathbf{x}_i))$. Then obtain $\mathbf{x}'' = \sigma(\mathbf{x}, \mathbf{X})$ by sequentially replacing $\mathbf{x}'_i \in (-1, 1)$

by $\mathbf{x}_i'' = \pm 1$ so that the value $(\mathbf{x}')^T \mathbf{X} \mathbf{x}'$ increases or stays the same.

2 Universality of Low-Degree Polynomials. Main Results

We next formally define the class of Low-Degree Polynomials (LDP) we use and then restate our main results formally. For a symmetric matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$ we denote by $\mathbb{R}[\mathbf{X}]_{\leq \Delta}$ the set of multivariate polynomials in $\{\mathbf{X}_{ij}, 1 \leq i < j \leq n\}$ with degree at most Δ and denote the degree of a polynomial p by Deg(p). When Δ is constant independent of n, we refer to these polynomials with the abbreviation LDP. We denote by $\mathcal{U}^{n \times n}$ the set of strictly upper triangluar matrices in $\mathbb{Z}_{>0}^{n \times n}$ and introduce

$$\mathcal{U}_{1}^{n \times n} \triangleq \{ \boldsymbol{\alpha} \in \mathcal{U}^{n \times n} | \boldsymbol{\alpha}_{ij} = 1, \text{ for some } 1 \le i < j \le n \},$$

$$\mathcal{U}_{*}^{n \times n} \triangleq \{ \boldsymbol{\alpha} \in \mathcal{U}^{n \times n} | \boldsymbol{\alpha}_{ij} \ne 1, \text{ for all } 1 \le i < j \le n \},$$
(2.1)

so that $\mathcal{U}^{n \times n} = \mathcal{U}_1^{n \times n} \cup \mathcal{U}_*^{n \times n}$. To each polynomial p in \mathbf{X} we associate its coefficients $c_{p, \boldsymbol{\alpha}}, \boldsymbol{\alpha} \in \mathcal{U}^{n \times n}$ so that

$$p(\mathbf{X}) = \sum_{\boldsymbol{\alpha} \in \mathcal{U}^{n \times n}} c_{p, \boldsymbol{\alpha}} \mathbf{X}^{\boldsymbol{\alpha}}$$

Note that the degree of p is given by $\max \|\boldsymbol{\alpha}\|_1$, where the maximum is over $\boldsymbol{\alpha}$ such that $c_{p,\boldsymbol{\alpha}} \neq 0$. Thus $p \in \mathbb{R}[\mathbf{X}]_{\leq \Delta}$ if $\|\boldsymbol{\alpha}\|_1 \leq \Delta$ for all $\boldsymbol{\alpha}$ such that $c_{p,\boldsymbol{\alpha}} \neq 0$. Define

$$\|p\| \triangleq \max_{\boldsymbol{\alpha} \in \mathcal{U}^{n \times n}} |c_{p,\boldsymbol{\alpha}}|$$

Each $\boldsymbol{\alpha} \in \mathcal{U}^{n \times n}$ is associated with a graph $G_{\boldsymbol{\alpha}} = (V_{\boldsymbol{\alpha}}, E_{\boldsymbol{\alpha}})$ which we call a factor graph as follows

$$V_{\boldsymbol{\alpha}} \triangleq \{k \in [n] | \exists j \in [n] \setminus \{k\}, \quad \boldsymbol{\alpha}_{kj} + \boldsymbol{\alpha}_{jk} > 0\}, E_{\boldsymbol{\alpha}} \triangleq \{(i,j) \in [n]^2 | i \neq j, \quad \boldsymbol{\alpha}_{ij} + \boldsymbol{\alpha}_{ji} > 0\}.$$
(2.2)

We say that $\mathbf{X}^{\boldsymbol{\alpha}}$ is connected if $G_{\boldsymbol{\alpha}}$ is a connected graph and write $\boldsymbol{\alpha} \in \mathcal{C}$. For a LDP $p \in \mathbb{R}[\mathbf{X}]_{\leq\Delta}$, we say that p is connected if for all $\boldsymbol{\alpha}$, $c_{p,\boldsymbol{\alpha}} = 0$ when $\mathbf{X}^{\boldsymbol{\alpha}}$ is not connected. For $k \in [n]$ we write $k \in p$ when $k \in V_{\boldsymbol{\alpha}}$ for all $\boldsymbol{\alpha}$ such that $c_{p,\boldsymbol{\alpha}} \neq 0$. The constant monomial associated with $\boldsymbol{\alpha} = \mathbf{0}_{\mathbb{Z}_{\geq 0}^{n \times n}}$ is assumed to be connected, and $k \in V_{\boldsymbol{\alpha}}$ is assumed for all $k \in [n]$.

Given $\boldsymbol{\alpha} \in \mathcal{U}^{n \times n}$, we say that $\mathbf{X}^{\boldsymbol{\alpha}}$ is a tree and write $\boldsymbol{\alpha} \in \mathcal{T}$ if $G_{\boldsymbol{\alpha}}$ is a tree. Given an LDP $p \in \mathbb{R}[\mathbf{X}]_{\leq \Delta}$ we say that p is a tree if its coefficients $c_{p,\boldsymbol{\alpha}}$ are null when $\mathbf{X}^{\boldsymbol{\alpha}}$ is not a tree. Note that every tree polynomial p is connected. The constant monomial is assumed to be a tree. Given a subset $S \subset [n]$, we use the slightly abused notation $S \subset \boldsymbol{\alpha}$ when $S \subset V_{\boldsymbol{\alpha}}$. If every monomial $\mathbf{X}^{\boldsymbol{\alpha}}$ in p with non-zero $c_{p,\boldsymbol{\alpha}}$ satisfies $S \subset V_{\boldsymbol{\alpha}}$, then we write $S \subset p$. Introduce the sets

$$\mathcal{T}_{n,2,\Delta} \triangleq \{ \boldsymbol{\alpha} \in \mathcal{U}^{n \times n} | \boldsymbol{\alpha} \in \mathcal{T}, \max(\boldsymbol{\alpha}) \le 2, \|\boldsymbol{\alpha}\|_1 \le \Delta \}, \\ \mathcal{C}_{n,2,\Delta} \triangleq \{ \boldsymbol{\alpha} \in \mathcal{U}^{n \times n} | \boldsymbol{\alpha} \in \mathcal{C}, \max(\boldsymbol{\alpha}) \le 2, \|\boldsymbol{\alpha}\|_1 \le \Delta \}.$$

We define p^{Tr} as the restriction of p to monomials $\mathbf{X}^{\boldsymbol{\alpha}}$ with $\boldsymbol{\alpha} \in \mathcal{T}_{n,2,\mathrm{Deg}(p)}$, i.e.

$$p^{\mathrm{Tr}}(\mathbf{X}) \triangleq \sum_{\boldsymbol{\alpha} \in \mathcal{T}_{n,2,\Delta}} c_{p,\alpha} \mathbf{X}^{\boldsymbol{\alpha}}.$$

We write $p \in \mathcal{T}_{n,2,\Delta}$ if $\boldsymbol{\alpha} \in \mathcal{T}_{n,2,\Delta}$ whenever $c_{p,\boldsymbol{\alpha}} \neq 0$. By definition $p^{\mathrm{Tr}} \in \mathcal{T}_{n,2,\mathrm{Deg}(p)}$ for every polynomial p. We will use the notations $\mathcal{T}_{n,2}, \mathcal{C}_{n,2}$ when the constant Δ is clear from context. We note that for any LDP $p \in \mathbb{R}[X]_{<\Delta}$, the projection p^{Tr} can be computed in $n^{O(\Delta)}$ time complexity.

We now restate our main result Theorem 1.1 formally. Recall the operator $\sigma(\mathbf{x}, \mathbf{X})$ defined in the notations section.

Theorem 2.1. Suppose $\mathbf{X} \stackrel{\mathsf{d}}{=} \operatorname{GOE}(n)$, $\mathbf{Y} \stackrel{\mathsf{d}}{=} \mathcal{G}(n, d/n)$, and $\Delta > 0$. Suppose an n-sequence of vectors of polynomials $p = (p_i, i \in [n])$ is such that $p_i \in \mathcal{T}_{n,2,\Delta}$, $||p_i|| \leq c(\Delta)$ for some constant $c(\Delta)$, and $i \in p_i$ for all $i \in [n]$. Suppose furthermore

$$\operatorname{p-lim}_{n \to \infty} \frac{1}{2n} p(\mathbf{X})^{\top} \mathbf{X} p(\mathbf{X}) \ge \mathcal{P},$$
(2.3)

$$\underset{n \to \infty}{\text{p-lim}} \frac{d(p(\mathbf{X}), [-1, 1]^n)}{\sqrt{n}} = 0.$$
 (2.4)

for some constant \mathcal{P} . Then

$$\underset{d \to \infty, n \to \infty}{\text{p-lim}} \frac{1}{2n} p(\mathbf{Y})^{\top} \mathbf{Y} p(\mathbf{Y}) \ge \mathcal{P},$$
(2.5)

$$\underset{d \to \infty, n \to \infty}{\text{p-lim}} \frac{d(p(\mathbf{Y}), [-1, 1]^n)}{\sqrt{n}} = 0.$$
(2.6)

Furthermore

$$\underset{n \to \infty}{\text{p-lim}} \, \frac{1}{2n} \sigma(p(\mathbf{Y}), \mathbf{Y})^\top \mathbf{Y} \sigma(p(\mathbf{Y}), \mathbf{Y}) \ge \mathcal{P}.$$
(2.7)

Next we move to stating formally our second result regarding approximating AMP via LDP. For this we begin by introducing AMP formally. There is a multitude of ways to describe AMP iterations, and each AMP algorithm is typically problem specific. For our purposes, we will use an appropriate variant of the general AMP iteration given in (6) in [Mon19] (namely, we consider the setting where the denoisers do not depend on \mathbf{y}), which we restate here. Consider a sequence of *denoisers* functions $f^t : \mathbb{R}^{t+1} \to \mathbb{R}$ and an initialization $\mathbf{u}^0 \in \mathbb{R}^n$. The AMP iteration is given by

$$\mathbf{u}^{t+1} = \mathbf{X} f^{t}(\mathbf{u}^{0}, ..., \mathbf{u}^{t}) - \sum_{j=1}^{t} b_{t,j} f^{j-1}(\mathbf{u}^{0}, ..., \mathbf{u}^{j-1}),$$

$$b_{t,j} = \frac{1}{n} \sum_{i=1}^{n} \frac{\partial f^{t}}{\partial \mathbf{u}_{i}^{j}} (\mathbf{u}_{i}^{0}, ..., \mathbf{u}_{i}^{t}),$$
(2.8)

where $f^j = 0$ for j < 0. Here $f^t(\mathbf{u}^1, \ldots, \mathbf{u}^t)$ is applied coordinate-wise. The AMP iterations satisfy a certain state evolution property formulated in terms of the centered Gaussian process $(U_j), j \ge 1$ with covariance specified recursively as $\mathbf{Q} = (\mathbf{Q}_{kj})_{k,j \ge 1}$ with

$$\mathbf{Q}_{k+1,j+1} = \mathbb{E}[f^k(U_0, ..., U_k)f^j(U_0, ..., U_j)], \quad \forall k, j \ge 0.$$
(2.9)

The output of the AMP is $\sigma(\mathbf{v}, \mathbf{X})$, where \mathbf{v} is defined by

$$\mathbf{v} = \sqrt{\delta} \sum_{k=1}^{\lfloor \bar{q}/\delta \rfloor} f^k(\mathbf{u}^0, ..., \mathbf{u}^k), \qquad (2.10)$$

where δ, \bar{q} are constant parameters driving the optimality gap, and the number of iterations of the AMP satisfies $T \geq \lfloor \bar{q}/\delta \rfloor$. The main result of [Mon19] asserts the near optimality of this algorithm for appropriate choice of denoisers f^t . Specifically, it asserts the existence for any $\epsilon > 0$ of the denoisers f^t and parameters \bar{q}, δ , so that whp

$$\frac{1}{2n} \mathbf{v}^{\top} \mathbf{X} \mathbf{v} \ge \mathcal{P}^* - \epsilon, \qquad (2.11)$$

$$\frac{d(\mathbf{v}, [-1, 1]^n)}{\sqrt{n}} \le \epsilon. \tag{2.12}$$

This version of AMP is called Incremental AMP which for the remainder of this paper, we call IAMP. Naturally, the IAMP algorithm depends on the parameter $\epsilon > 0$. This dependency will be clear from context whenever we refer to IAMP. We will show that the output \mathbf{v} of IAMP can be well approximated by LDP, and, as a result, the implied objective value can be reached via LDP as well. This is the essence of our intermediate result Theorem 1.2 stated now formally below as follows.

Theorem 2.2. Given $\epsilon > 0, T$, consider a sequence of denoisers $f^t, t \in [T]$ satisfying,

- 1. There exists a constant L s.t f^t are L-Lipschitz.
- 2. f^t have weak derivatives that are either Pseudo-Lipschitz or indicators.
- 3. The covariance matrix $\mathbf{Q}_{\leq t}$ defined in (2.9) satisfies $\mathbf{Q}_{\leq t} \succeq \mathbf{I}_t$ and $\|\mathbf{Q}_{\leq t}\|_{\infty} \leq 2$ for all $t \in [T]$.

Suppose there exists a constant M > 0 independent of n such that $\|\mathbf{u}^0\|_{\infty} \leq M$ whp. Then there exists constants $\Delta = \Delta(\epsilon, T), c = c(\epsilon, T)$, and a sequence $p = (p_i, i \in [n]), p_i \in \mathcal{T}_{n,2,\Delta}$, such that whp

$$\frac{\|\mathbf{v} - p(\mathbf{X})\|_2}{\sqrt{n}} \le \epsilon$$
$$\frac{|\mathbf{v}^\top \mathbf{X} \mathbf{v} - p(\mathbf{X})^\top \mathbf{X} p(\mathbf{X})|}{n} \le \epsilon,$$

and p satisfy $\forall i \in [n], i \in p_i \text{ and } ||p_i|| \leq c$.

As a result, for every $\epsilon > 0$ there exists Δ and a sequence $p = (p_i, i \in [n]), p_i \in \mathcal{T}_{n,2,\Delta}$ such that whp

$$\frac{1}{2n} p(\mathbf{X})^{\top} \mathbf{X} p(\mathbf{X}) \ge \mathcal{P}^* - \epsilon,$$
$$\frac{d(p(\mathbf{X}), [-1, 1]^n)}{\sqrt{n}} \le \epsilon.$$

We will show that IAMP in [Mon19] can be adopted to satisfy the assumptions of the theorem above. Near optimality of the LDP after rounding, namely Corollary 1.3 is now an immediate corollary of Theorems 2.1, 2.2 and is restated as follows.

Corollary 2.3. For every $\epsilon > 0$ there exists a sequence $p = (p_i, i \in [n]), p_i \in \mathcal{T}_{n,2,\Delta}$ such that

$$\operatorname{p-lim}_{d\to\infty,n\to\infty}\frac{1}{2n}\sigma(p(\mathbf{Y}),\mathbf{Y})^{\top}\mathbf{Y}\sigma(p(\mathbf{Y}),\mathbf{Y}) \geq \mathcal{P}^* - \epsilon.$$

3 Preliminary Results

In this section we establish some preliminary technical results.

3.1 Some elementary estimates

We begin by establishing some non-probabilistic bounds on functions and their derivatives appearing in our analysis.

Lemma 3.1 (Faà di Bruno [Arb]). Let $g \in C^m(\mathcal{D}_g, \mathbb{R}), f \in C^m(\mathcal{D}_f, \mathbb{R})$ be m-times differentiable scalar functions with $\operatorname{Im}(g) \subset \mathcal{D}_f$, and let w(z) = f(g(z)). The following holds for all $z \in \mathcal{D}_g$ and $k \leq m$

$$w^{(k)}(z) = \sum_{\substack{\ell_1 + 2\ell_2 + \dots + k\ell_k = k \\ \ell_1, \dots, \ell_k \ge 0}} \frac{k!}{\ell_1! \dots \ell_k!} \cdot f^{\ell_1 + \dots + \ell_k}(g(z)) \cdot \prod_{j=1}^k \left(\frac{g^{(j)}(z)}{j!}\right)^{\ell_j}.$$

From Lemma 3.1 we obtain the following corollary.

Corollary 3.2. In the setting of Lemma 3.1 with $m \ge 4$, we have the following

$$\begin{split} & w^{(1)}(z) = g^{(1)}(z) \cdot f^{(1)}((g(z))), \\ & w^{(2)}(z) = (g^{(1)}(z))^2 \cdot f^{(2)}((g(z))) + g^{(2)}(z) \cdot f^{(1)}((g(z))), \\ & w^{(3)}(z) = (g^{(1)}(z))^3 \cdot f^{(3)}((g(z))) + 3g^{(1)}(z) \cdot g^{(2)}(z) \cdot f^{(2)}((g(z))) + g^{(3)}(z) \cdot f^{(1)}(g(z)), \\ & w^{(4)}(z) = (g^{(1)}(z))^4 \cdot f^{(4)}(g(z)) + 6(g^{(1)}(z))^2 \cdot g^{(2)}(z) \cdot f^{(3)}(g(z)) \\ & \quad + (4g^{(1)}(z) \cdot g^{(3)}(z) + 3(g^{(2)}(z))^2) \cdot f^{(2)}(g(z)) + g^{(4)}(z) \cdot f^{(1)}(g(z)). \end{split}$$

In particular, if $g: \mathbb{R} \to \mathbb{R}_{>0}$ and $f: x \mapsto \sqrt{x}$, we have the following

$$\begin{split} w^{(1)} &= \frac{g^{(1)}}{2\sqrt{g}}, \\ w^{(2)} &= \frac{g^{(2)}}{2\sqrt{g}} - \frac{(g^{(1)})^2}{4g^{\frac{3}{2}}}, \\ w^{(3)} &= \frac{g^{(3)}}{2\sqrt{g}} - \frac{3}{4}\frac{g^{(1)}g^{(2)}}{g^{\frac{3}{2}}} + \frac{3}{8}\frac{(g^{(1)})^3}{g^{\frac{5}{2}}}, \\ w^{(4)} &= \frac{g^{(4)}}{2\sqrt{g}} - \frac{g^{(1)}g^{(3)}}{g^{\frac{3}{2}}} - \frac{3}{4}\frac{(g^{(2)})^2}{g^{\frac{3}{2}}} + \frac{9}{4}\frac{(g^{(1)})^2g^{(2)}}{g^{\frac{5}{2}}} - \frac{15}{16}\frac{(g^{(1)})^4}{g^{\frac{7}{2}}}. \end{split}$$

Lemma 3.3. Given $a, b \in \mathbb{R}$ let

$$\begin{split} \phi : & \mathbb{R} \to \mathbb{R}, \quad z \mapsto \frac{e^z}{1 + e^z}, \\ \psi_{a,b} : & \mathbb{R} \to \mathbb{R}, \quad z \mapsto \phi \left(az - b \right), \\ \pi_a : & \mathbb{R} \to \mathbb{R}, \quad z \mapsto \frac{\log(1 + e^{a(x-1)}) + \log(1 + e^{a(-x-1)})}{a}. \end{split}$$

Then

1.
$$\psi_{a,b}$$
 is $|a|-Lipschitz_{a,b}$

- 2. $\|\psi_{a,b}^{(j)}\|_{\infty} \le |a|^j, \forall j \in \{1, 2, 3, 4\},\$
- 3. $\forall a > 0, \sup_{z \in \mathbb{R}} |d(z, [-1, 1]) \pi_a(z)| \le \frac{2}{a},$
- $4. \ \|\pi_a^{(j)}\|_\infty \leq 2|a|^{j-1}, \forall j \in \{1,2,3,4\},$
- 5. $\forall z \ge 0, \pi_a(z) \ge \frac{1}{2ae^a},$
- 6. If $a \ge 2$ then $\forall z \ge 0$, $\pi_a(z) \le z^2 + 1$.

Proof. The first four derivatives of ϕ are

$$\begin{split} \phi^{(1)}(z) &= \frac{e^z}{(1+e^z)^2}, \\ \phi^{(2)}(z) &= \frac{e^z(1-e^z)}{(1+e^z)^3}, \\ \phi^{(3)}(z) &= \frac{e^z(-4e^z+e^{2z}+1)}{(1+e^z)^4}, \\ \phi^{(4)}(z) &= -\frac{e^z(11e^z-11e^{2z}+e^{3z}-1)}{(e^z+1)^5}. \end{split}$$

In particular, elementary calculus yields $\|\phi^{(j)}\|_{\infty} \leq 1$ for $j \leq 4$.

- 1. The first claim follows from noting that $\psi_{a,b}^{(1)}(z) = a\phi^{(1)}(az-b)$, and thus $\|\psi_{a,b}^{(1)}\|_{\infty} \leq |a|$.
- 2. The second claim follows from $\psi^{(j)}(z) = a^j \phi^{(j)}(az b)$ combined with the previous bounds on $\|\phi^{(j)}\|_{\infty}$ for $j \leq 4$.
- 3. We first note that π_a is symmetric, and thus we only need to bound $|d(z, [-1, 1]) \pi_a(z)|$ for $z \ge 0$. If $z \in [0, 1]$, then

$$|\pi_a(z) - d(z, [-1, 1])| = |\pi_a(z)| \le \frac{\log 2 + \log(1 + e^{-a})}{a} \le \frac{2}{a}.$$

If z > 1, then

$$\begin{aligned} |\pi_a(z) - d(z, [-1, 1])| &= \left| \frac{\log(1 + e^{a(-z-1)}) + \log(1 + e^{a(z-1)})}{a} - (z-1) \right| \\ &\leq \left| \frac{\log(1 + e^{a(-z-1)})}{a} \right| + \left| \frac{\log(1 + e^{a(z-1)}) - a(z-1)}{a} \right| \\ &\leq \frac{\log 2}{a} + \frac{\log(e^{-a(z-1)} + 1)}{a} \\ &\leq \frac{2\log 2}{a} \\ &\leq \frac{2}{a}. \end{aligned}$$

Therefore $\sup_{z \in \mathbb{R}} |d(z, [-1, 1]) - \pi_a(z)| \leq \frac{2}{a}$.

4. Let $\xi(z) \triangleq \log(1+e^z)$ and note that $\xi^{(j)} = \phi^{(j-1)}$ for $j \ge 1$, and $\pi_a(z) = \frac{\xi(a(-z-1)) + \xi(a(z-1))}{a}$. We then have for $j \in [4]$,

$$|\pi_a^{(j)}(z)| \le |a^{j-1}\xi^{(j)}(a(-z-1))| + |a^{j-1}\xi^{(j)}(a(z-1))| \le 2|a|^{j-1} \|\phi^{(j-1)}\|_{\infty} \le 2|a|^{j-1},$$

which yields the result of the claim.

5. Let $z \ge 0$, then

$$\pi_a(z) \ge \frac{\log(1 + e^{a(z-1)})}{a} \ge \frac{\log(1 + e^{-a})}{a} \ge \frac{e^{-a}}{2a}$$

where the last inequality follows from noting that $\log(1 + e^{-x}) \ge e^{-x}/2$, $\forall x \ge 0$. Indeed, the function $x \mapsto \log(1 + e^{-x}) - e^{-x}/2$ has derivative $\frac{e^{-x}-1}{2(1+e^x)} \le 0$, thus it is non-increasing. Since the latter function has limit 0 at $x \to +\infty$ it remains nonnegative on \mathbb{R}_+ , which yields the inequality $\log(1 + e^{-x}) \ge e^{-x}/2, \forall x \ge 0$. The case z < 0 is immediate by symmetry of π_a .

6. By item (3) π_a is a 2/a-uniform approximation of $z \mapsto d(z, [-1, 1]) = (|z| - 1)_+$. Therefore, if $|z| \le 1$ then $|\pi_a(z)| \le \frac{2}{a} \le 1$, and if |z| > 1 then

$$|\pi_a(z)| \le \frac{2}{a} + (|z| - 1)_+ = \frac{2}{a} + |z| - 1 \le |z|^2.$$

Combining both cases yields $|\pi_a(z)| \leq |z|^2 + 1$ for all $z \in \mathbb{R}$.

Lemma 3.4. Let $\mathbf{x} \in \mathbb{R}^n$ and $\psi_{\frac{\beta_1}{\alpha_n},c}, \pi_{\beta_2}$ as in Lemma 3.3 with $\beta_1, \beta_2 > 0$ and $c \in \mathbb{R}$. Then

$$\left|\psi_{\frac{\beta_1}{\sqrt{n}},c}\left(d(\mathbf{x},[-1,1]^n)\right) - \psi_{\frac{\beta_1}{\sqrt{n}},c}\left(\sqrt{\sum_{i=1}^n \pi_{\beta_2}(\mathbf{x}_i)^2}\right)\right| \le \frac{2\beta_1}{\beta_2}.$$

Proof. By Lemma 3.3 we have that $\psi_{\frac{\beta_1}{\sqrt{n}},c}$ is $\frac{\beta_1}{\sqrt{n}}$ -Lipschitz. Therefore

$$\left| \psi_{\frac{\beta_{1}}{\sqrt{n}},c} \left(d(\mathbf{x}, [-1,1]^{n}) \right) - \psi_{\frac{\beta_{1}}{\sqrt{n}},c} \left(\sqrt{\sum_{i=1}^{n} \pi_{\beta_{2}}(\mathbf{x}_{i})^{2}} \right) \right| \\ \leq \frac{\beta_{1}}{\sqrt{n}} \left| \sqrt{\sum_{i=1}^{n} d(\mathbf{x}_{i}, [-1,1])^{2}} - \sqrt{\sum_{i=1}^{n} \pi_{\beta_{2}}(\mathbf{x}_{i})^{2}} \right| \\ \leq \frac{\beta_{1}}{\sqrt{n}} \frac{\sum_{i=1}^{n} |d(\mathbf{x}_{i}, [-1,1]) - \pi_{\beta_{2}}(\mathbf{x}_{i})| (d(\mathbf{x}_{i}, [-1,1]) + \pi_{\beta_{2}}(\mathbf{x}_{i}))}{\sqrt{\sum_{i=1}^{n} d(\mathbf{x}_{i}, [-1,1])^{2}} + \sqrt{\sum_{i=1}^{n} \pi_{\beta_{2}}(\mathbf{x}_{i})^{2}} \\ \leq \frac{\beta_{1}}{\sqrt{n}} \sup_{z \in \mathbb{R}} |d(z, [-1,1] - \pi_{\beta_{2}}(z))| \frac{\sum_{i=1}^{n} d(\mathbf{x}_{i}, [-1,1]) + \pi_{\beta_{2}}(\mathbf{x}_{i})}{\sqrt{\sum_{i=1}^{n} d(\mathbf{x}_{i}, [-1,1])^{2}} + \sqrt{\sum_{i=1}^{n} \pi_{\beta_{2}}(\mathbf{x}_{i})^{2}} \\ \leq \frac{\beta_{1}}{\sqrt{n}} \sup_{z \in \mathbb{R}} |d(z, [-1,1] - \pi_{\beta_{2}}(z))| \sqrt{n} \tag{3.1}$$

$$=\frac{2\beta_1}{\beta_2},\tag{3.2}$$

where (3.1) follows from Cauchy-Schwartz inequality and (3.2) from part 3 of Lemma 3.3.

Lemma 3.5. Let $\mathbf{X} \in \mathbb{R}^{n \times n}$, $p \in \mathbb{R}[\mathbf{X}]_{\leq \Delta_1}$ and $q \in \mathbb{R}[\mathbf{X}]_{\leq \Delta_2}$, Then

$$\|pq\| \le 2^{\Delta_1 + \Delta_2} \|p\| \|q\|$$

Proof. For $\boldsymbol{\alpha} \in \mathcal{U}^{n \times n}$ with $c_{pq,\boldsymbol{\alpha}} \neq 0$, the coefficient of the monomial \mathbf{X}^{α} in pq is given by

$$c_{pq,\boldsymbol{\alpha}} = \sum_{\substack{\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2 \in \mathcal{U}^{n \times n} \\ \boldsymbol{\alpha}_1 + \boldsymbol{\alpha}_2 = \boldsymbol{\alpha}}} c_{p,\boldsymbol{\alpha}_1} c_{q,\boldsymbol{\alpha}_2}$$

Each of the summands has absolute value bounded by ||p|| ||q||. Since the degree of pq is at most $\Delta_1 + \Delta_2$ we have $||\boldsymbol{\alpha}||_1 \leq \Delta_1 + \Delta_2$, therefore the number of pairs $(\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2) \in (\mathcal{U}^{n \times n})^2$ summing to $\boldsymbol{\alpha}$ is bounded by $2^{||\boldsymbol{\alpha}||_1} \leq 2^{\Delta_1 + \Delta_2}$.

Lemma 3.6. Let $\mathbf{X} \in \mathbb{R}^{n \times n}$ and let $p_i \in \mathbb{R}[\mathbf{X}]_{\leq \Delta_i}$, i = 1, ..., m. Suppose p_i are connected and $c_{p_i, \mathbf{0}_{\mathbb{R}^{n \times n}}} = 0$ (i.e. $p(\mathbf{0}_{\mathbb{R}^{n \times n}}) = 0$) for all $i \in [m]$. Assume there exists subsets $S_i \subset [n], i = 1, ..., m$ such that

1. $\forall i \in [m], \quad S_i \subset p_i.$

2. For all $j \in [m]$ and $1 \le k_1 < \ldots < k_j \le m$, $|S_{k_1} \cup \ldots \cup S_{k_j}| \ge cj^{\gamma}$, where c, γ are positive constants.

Then

$$\left\|\sum_{i=1}^{m} p_i\right\| \le \left(\frac{1 + \max_{i \in [m]} \Delta_i}{c}\right)^{\frac{1}{\gamma}} \max_{i \in [m]} \|p_i\|.$$

Proof. For $\boldsymbol{\alpha} \in \mathcal{U}^{n \times n}$, the coefficient of the monomial $\mathbf{X}^{\boldsymbol{\alpha}}$ in $s(\mathbf{X}) \triangleq \sum_{i=1}^{m} p_i(\mathbf{X})$ is given by

$$c_{s,\boldsymbol{\alpha}} = \sum_{i \in [m]} c_{p_i,\boldsymbol{\alpha}}.$$

Each of the summands has absolute value bounded by $\max_{i \in [m]} ||p_i||$. If $c_{p_i, \alpha} \neq 0$ then $S_i \subset V_{\alpha}$. Let $I \triangleq \{i \in [m] | c_{p_i, \alpha} \neq 0\}$. Then $\forall i \in I, S_i \subset V_{\alpha}$, and thus $\bigcup_{i \in I} S_i \subset V_{\alpha}$, which implies

$$|V_{\boldsymbol{\alpha}}| \ge |\cup_{i \in I} S_i| \ge c|I|^{\gamma},$$

where the last inequality follows from the assumption of the lemma. Since the number of nodes of a connected graph is at most one plus the number of edges, we have $|V_{\alpha}| \leq 1 + \max_{i \in [m]} \Delta_i$. We obtain

$$|I| \le \left(\frac{1 + \max_{i \in [m]} \Delta_i}{c}\right)^{\frac{1}{\gamma}}.$$

Henceforth

$$|c_{s,\boldsymbol{\alpha}}| \le |I| \max_{i \in [m]} ||p_i|| \le \left(\frac{1 + \max_{i \in [m]} \Delta_i}{c}\right)^{\frac{1}{\gamma}} \max_{i \in [m]} ||p_i||,$$

which concludes the proof.

Lemma 3.7. Suppose $\beta \geq 1$, $u, a, b \in \mathbb{R}$, $v(z) = u + za + z^2b$ and

$$k(z) = \pi_{\beta}(v(z))^2,$$

where π_{β} is given in Lemma 3.3. There exists a universal constant c such that for all $j \in [4]$, and $|z^*| \leq z$ it holds

$$|k^{(j)}(z^*)| \le c\beta^3 \sum_{\substack{j_1, j_2, j_3, j_4 \le 4+j\\ j_2+j_3 \ge \lceil j/2 \rceil}} |z|^{j_1} |a_i|^{j_2} |b_i|^{j_3} |u_i|^{j_4}.$$

Proof. We drop the subscript in π_{β} and write π . Note the following elementary properties of π from Lemma 3.3

$$\forall j \in [4], \quad \|\pi^{(j)}\|_{\infty} \le 2\beta^{j-1},$$
(3.3)

$$\forall z \in \mathbb{R}, \quad |\pi(z)| \le z^2 + 1. \tag{3.4}$$

We have

$$k^{(1)}(z) = 2(a_i + 2zb_i)\pi^{(1)}(v_i(z)) \cdot \pi(v_i(z)).$$

Using (3.3), (3.4), and $|z^*| \le |z|$, we have

$$k^{(1)}(z^*) \le 8 \left(|a_i| + |z| |b_i| \right) \left(v_i(z^*)^2 + 1 \right).$$
(3.5)

Using $|z^*| \leq z$ it follows

$$(v(z^*))^2 = u^2 + (z^*)^2 a^2 + (z^*)^4 b^2 + 2z^* u a + 2(z^*)^2 b u + 2(z^*)^3 b a,$$

$$\leq 2 \left(u^2 + z^2 a^2 + z^4 b^2 + |z| |u| |a| + z^2 |b| |u| + |z|^3 |b| |a| \right).$$
(3.6)

Plugging (3.6) in (3.5) and expanding yields

$$|k^{(1)}(z^*)| \le c \sum_{\substack{j_1, j_2, j_3, j_4 \le 5\\ j_2+j_3 \ge 1}} |z|^{j_1} |a|^{j_2} |b|^{j_3} |u|^{j_4},$$

for some universal constant c. The proof for $j \in \{2,3,4\}$ is identical using (3.3), (3.4) and the following higher order derivatives expressions

$$\begin{split} k^{(2)}(z) &= 4b\pi^{(1)}(v(z))\pi(v(z)) + 2(a+2zb)^2 \left[\pi^{(2)}(v(z))\pi(v(z)) + \pi^{(1)}(v(z))^2\right],\\ k^{(3)}(z) &= 12b(a+2zb) \left[\pi^{(2)}(v(z))\pi(v(z)) + \pi^{(1)}(v(z))^2\right] \\ &\quad + 2(a+2zb)^3 \left[\pi^{(3)}(v(z))\pi(v(z)) + 3\pi^{(2)}(v(z))\pi^{(1)}(v(z))\right],\\ k^{(4)}(z) &= 24b^2 \left[\pi^{(2)}(v(z))\pi(v(z)) + \pi^{(1)}(v(z))^2\right] \\ &\quad + 12b(a+2zb)^2 \left[2\pi^{(3)}(v(z))\pi(v(z)) + 6\pi^{(1)}(v(z))\pi^{(2)}(v(z))\right] \\ &\quad + 2(a+2zb)^4 \left[\pi^{(4)}(v(z))\pi(v(z)) + 4\pi^{(3)}(v(z))\pi^{(1)}(v(z)) + 3\pi^{(2)}(v(z))^2\right]. \end{split}$$

3.2 Probabilistic properties of LDP

We now establish some preliminary properties of LDP, including some probabilistic bounds.

Lemma 3.8. Let $\mathbf{X} \in \mathbb{R}^n$, $p, q \in \mathbb{R}[\mathbf{X}]$. Then

- $(p+q)^{\mathrm{Tr}} = p^{\mathrm{Tr}} + q^{\mathrm{Tr}}$.
- If $r \in \{p,q\}$ is connected and satisfies $r^{\mathrm{Tr}} = 0$ then $(pq)^{\mathrm{Tr}} = 0$.

Proof. The first property is straightforward from the definition of projection into $\mathcal{T}_{n,2}$. For the second assume without loss of generality r = p, i.e p is connected and $p^{\mathrm{Tr}} = 0$. Assume that $c_{(pq)^{\mathrm{Tr}}, \alpha} \neq 0$ for some $\alpha \in \mathcal{T}_{n,2}$. Since $c_{(pq)^{\mathrm{Tr}}, \alpha} = \sum_{\alpha_1 + \alpha_2 = \alpha}^{\alpha_1, \alpha_2} c_{p,\alpha_1} c_{q,\alpha_2}$, there must exist α_1, α_2 such that $\alpha_1 + \alpha_2 = \alpha, c_{p,\alpha_1} \neq 0, c_{q,\alpha_2} \neq 0$. Since $p^{\mathrm{Tr}} = 0$, it follows that $\alpha_1 \notin \mathcal{T}_{n,2}$ which is equivalent to : $\max(\alpha_1) > 2$ or $\alpha_1 \notin \mathcal{T}$. In the former case we have $\max(\alpha_1) > 2 \implies \max(\alpha) > 2$ and thus $\alpha \notin \mathcal{T}_{n,2}$ which contradicts the claim $c_{(pq)^{\mathrm{Tr}}, \alpha} \neq 0$, while in the latter case it follows that G_{α_1} has a cycle (since p is connected), and therefore $G_{\alpha} = G_{\alpha_1} \cup G_{\alpha_2}$ has a cycle, which contradicts $\alpha \in \mathcal{T}_{n,2}$ and concludes the proof.

Lemma 3.9. Let $\mathbf{Y} \stackrel{\mathsf{d}}{=} \mathcal{G}(n, d/n)$. Then

- 1. $\mathbb{E}[\mathbf{Y}_{ij}] = 0, \mathbb{E}[\mathbf{Y}_{ij}^2] = \frac{1}{n}, \forall 1 \le i < j \le n$.
- 2. $\mathbb{E}[\mathbf{Y}_{ij}^k] = \frac{(-1)^k + o(1)}{nd^{\frac{k}{2}-1}}, \mathbb{E}[|\mathbf{Y}_{ij}|^k] = \frac{1 + o(1)}{nd^{\frac{k}{2}-1}}, \forall k \ge 2, \forall 1 \le i < j \le n.$
- 3. $\forall \boldsymbol{\alpha} \in \mathcal{U}^{n \times n}$ such that $\forall 1 \leq i < j \leq n, \boldsymbol{\alpha}_{ij} \neq 1$ and $\|\boldsymbol{\alpha}\|_1$ bounded by a constant independent of n, it holds

$$\mathbb{E}[\mathbf{Y}^{\boldsymbol{\alpha}}] = ((-1)^{\|\boldsymbol{\alpha}\|_1} + o(1))n^{-\|\boldsymbol{\alpha}\|_0}d^{-\frac{\|\boldsymbol{\alpha}\|_1}{2} + \|\boldsymbol{\alpha}\|_0}$$
$$\mathbb{E}[|\mathbf{Y}^{\boldsymbol{\alpha}}|] = (1 + o(1))n^{-\|\boldsymbol{\alpha}\|_0}d^{-\frac{\|\boldsymbol{\alpha}\|_1}{2} + \|\boldsymbol{\alpha}\|_0}.$$

Proof. The first property is immediate from the definition of sparse disorder $\mathcal{G}(n, d/n)$. For the second property, we have for $k \geq 2$

$$\mathbb{E}[\mathbf{Y}_{ij}^k] = \frac{(-1)^k}{(d(1-\frac{d}{n}))^{\frac{k}{2}}} \left(\left(-\frac{d}{n} \right)^k \left(1 - \frac{d}{n} \right) + \left(1 - \frac{d}{n} \right)^k \frac{d}{n} \right) \sim (-1)^k \frac{\frac{d}{n} (1 - \frac{d}{n})^k}{(d(1-\frac{d}{n}))^{\frac{k}{2}}} = \frac{(-1)^k + o(1)}{nd^{\frac{k}{2}-1}}.$$

Similar computations yield the same asymptotic for $\mathbb{E}[|\mathbf{Y}_{ij}|^k]$. Finally, the second property yields

$$\mathbb{E}[\mathbf{Y}^{\boldsymbol{\alpha}}] = \prod_{\boldsymbol{\alpha}_{ij}>0} \frac{(-1)^{\boldsymbol{\alpha}_{ij}} + o(1)}{nd^{\frac{\boldsymbol{\alpha}_{ij}}{2}} - 1} = ((-1)^{\|\boldsymbol{\alpha}\|_{1}} + o(1))n^{-\|\boldsymbol{\alpha}\|_{0}}d^{-\frac{\|\boldsymbol{\alpha}\|_{1}}{2} + \|\boldsymbol{\alpha}\|_{0}}.$$

A similar computation yields the asymptotic bound for $\mathbb{E}[|\mathbf{Y}|^{\boldsymbol{\alpha}}]$. This concludes the proof of the Lemma.

Lemma 3.10. Let $\mathbf{Y} \stackrel{\mathsf{d}}{=} \mathcal{G}(n, d/n)$. There exists a constant c > 0 such that $\|\mathbf{Y}\|_{\text{op}} \leq c$ with high probability as $n \to \infty, d \to \infty$.

Proof. The Lemma is a Corollary of Theorem 2 in [FK81]. While the statement of the previous theorem assumes that the entries of \mathbf{Y} are bounded by K and have variance σ where K, σ do not depend on n, the proof in [FK81] extends naturally in the setting of $\mathcal{G}(n, d/n)$ for sufficiently large d.

Lemma 3.11. Let $\mathbf{X} \stackrel{\mathsf{d}}{=} \operatorname{GOE}(n)$ and $\mathbf{Y} \stackrel{\mathsf{d}}{=} \mathcal{G}(n, d/n)$. Suppose $p \in \mathbb{R}[\mathbf{X}]_{\leq \Delta}$ and $S \subset [n]$ satisfy

- 1. $S \subset p$.
- 2. For each $\boldsymbol{\alpha}$ with $c_{p,\boldsymbol{\alpha}} \neq 0$ either of the following holds
 - α is connected.

• There exists $\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2 \in \mathcal{U}^{n \times n} \setminus \mathcal{T}_{n,2}$ with disjoint supports satisfying $\boldsymbol{\alpha} = \boldsymbol{\alpha}_1 + \boldsymbol{\alpha}_2$ and $\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2$ are connected.

For every Δ , there exists $c = c(\Delta)$ independent of n, d such that

$$\left|\mathbb{E}[p(\mathbf{X}) - p^{\mathrm{Tr}}(\mathbf{X})]\right| \le \frac{c||p||}{n^{|S|}},\tag{3.7}$$

$$\left|\mathbb{E}[p(\mathbf{Y}) - p^{\mathrm{Tr}}(\mathbf{Y})]\right| \le \frac{c||p||}{n^{|S|-2}d}.$$
(3.8)

Furthermore, when p is connected, the following tighter concentration for \mathbf{Y} holds

$$|\mathbb{E}[p(\mathbf{Y}) - p^{\mathrm{Tr}}(\mathbf{Y})]| \le \frac{c||p||}{n^{|S|-1}\sqrt{d}}$$

We note that property (3.7) is known in the literature (see [BMLM15] for a reference). We provide a proof here for completeness.

Proof. We first show (3.7). Introduce

$$\mathcal{K} \triangleq \{ \boldsymbol{\alpha} \in \mathcal{U}^{n \times n} \setminus \mathcal{T}_{n,2} | \boldsymbol{\alpha} \text{ is connected } \}, \\ \mathcal{W} \triangleq \{ (\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2) \in (\mathcal{U}^{n \times n} \setminus \mathcal{T}_{n,2})^2 | \boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2 \text{ are connected, } \operatorname{Supp}(\boldsymbol{\alpha}_1) \cap \operatorname{Supp}(\boldsymbol{\alpha}_2) = \emptyset \}.$$

We have

$$|\mathbb{E}[p(\mathbf{X}) - p^{\mathrm{Tr}}(\mathbf{X})]| \leq \left| \sum_{\boldsymbol{\alpha} \in \mathcal{K}} c_{p,\boldsymbol{\alpha}} \mathbb{E}[\mathbf{X}^{\boldsymbol{\alpha}}] \right| + \left| \sum_{(\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2) \in \mathcal{W}} c_{p,\boldsymbol{\alpha}_1 + \boldsymbol{\alpha}_2} \mathbb{E}[\mathbf{X}^{\boldsymbol{\alpha}_1 + \boldsymbol{\alpha}_2}] \right|.$$

Note that $\forall \boldsymbol{\alpha} \in \mathcal{U}^{n \times n}, \mathbb{E}[\mathbf{X}^{\boldsymbol{\alpha}}] = \gamma_{\boldsymbol{\alpha}} n^{-\frac{\|\boldsymbol{\alpha}\|_{1}}{2}} \mathbf{1}_{2|\boldsymbol{\alpha}}$, where $\gamma_{\boldsymbol{\alpha}} = \prod_{\substack{1 \leq i < j \leq n \\ 2|\boldsymbol{\alpha}_{ij}, \boldsymbol{\alpha}_{ij} > 0}} (\boldsymbol{\alpha}_{ij} - 1)!!$. Moreover $\gamma_{\boldsymbol{\alpha}} \leq c_{\Delta}$ for some constant c_{Δ} depending only on Δ . We then have

$$|\mathbb{E}[p(\mathbf{X}) - p^{\mathrm{Tr}}(\mathbf{X})]| \le c_{\Delta} ||p|| \left(\sum_{\substack{\boldsymbol{\alpha} \in \mathcal{K} \\ 2|\boldsymbol{\alpha}}} n^{-\frac{||\boldsymbol{\alpha}||_{1}}{2}} + \sum_{\substack{(\boldsymbol{\alpha}_{1}, \boldsymbol{\alpha}_{2}) \in \mathcal{W} \\ 2|\boldsymbol{\alpha}_{1}, 2|\boldsymbol{\alpha}_{2}}} n^{-\frac{||\boldsymbol{\alpha}_{1}||_{1} + ||\boldsymbol{\alpha}_{2}||_{1}}{2}} \right)$$

We claim that $\forall \boldsymbol{\alpha}' \in \{\boldsymbol{\alpha}, \boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2\}$ appearing in the sums above, we have $|V_{\boldsymbol{\alpha}'}| \leq \frac{\|\boldsymbol{\alpha}'\|_1}{2}$. There are two cases

• Case 1: $\alpha' \notin \mathcal{T}$.

As $G_{\boldsymbol{\alpha}'}$ is connected but is not a tree, it follows that $|V_{\boldsymbol{\alpha}'}| \leq |E_{\boldsymbol{\alpha}'}| = \|\boldsymbol{\alpha}'\|_0$. Using $2|\boldsymbol{\alpha}'$, we have $\|\boldsymbol{\alpha}'\|_0 \leq \frac{\|\boldsymbol{\alpha}'\|_1}{2}$ and thus $|V_{\boldsymbol{\alpha}'}| \leq \frac{\|\boldsymbol{\alpha}'\|_1}{2}$.

• Case 2: $\boldsymbol{\alpha}' \in \mathcal{T}$. We have $|V_{\boldsymbol{\alpha}'}| = |E_{\boldsymbol{\alpha}'}| + 1 = \|\boldsymbol{\alpha}'\|_0 + 1$, moreover $2|\boldsymbol{\alpha}' \text{ and } \boldsymbol{\alpha}' \notin \mathcal{T}_{n,2}, \boldsymbol{\alpha}' \in \mathcal{T}$ implies that $\max(\boldsymbol{\alpha}') \ge 4$, therefore $\|\boldsymbol{\alpha}'\|_0 \le \frac{\|\boldsymbol{\alpha}'\|_1 - 2}{2} = \frac{\|\boldsymbol{\alpha}'\|_1}{2} - 1$. Hence $|V_{\boldsymbol{\alpha}'}| \le \frac{\|\boldsymbol{\alpha}'\|_1}{2}$.

Thus the claim is verified. Given $a \in [0, \Delta]$ and a graph H introduce the set

 $\mathcal{F}_{a,H} \triangleq \{ \boldsymbol{\alpha} \in \mathcal{K} | \| \boldsymbol{\alpha} \|_1 = a, G_{\boldsymbol{\alpha}} \sim H, 2 | \boldsymbol{\alpha} \} \cup \{ \boldsymbol{\alpha} \in \mathcal{U}^{n \times n} | \| \boldsymbol{\alpha} \|_1 = a, G_{\boldsymbol{\alpha}} \sim H, \exists (\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2) \in \mathcal{W}, \boldsymbol{\alpha} = \boldsymbol{\alpha}_1 + \boldsymbol{\alpha}_2, 2 | \boldsymbol{\alpha}_1, 2 | \boldsymbol{\alpha}_2 \},$ where $G_{\boldsymbol{\alpha}} \sim H$ means $G_{\boldsymbol{\alpha}}$ is isomorphic to H. Recalling $S \subset G_{\boldsymbol{\alpha}}$, it follows $|\mathcal{F}_{a,H}| \leq \beta_{\Delta} n^{|V_H| - |S|}$ where β_{Δ} is a constant depending on Δ . From the previous analysis, it holds that $\forall \boldsymbol{\alpha} \in \mathcal{F}_{a,H}, |V_H| = |V_{\boldsymbol{\alpha}}| \leq \frac{\|\boldsymbol{\alpha}\|_1}{2},$ therefore $|\mathcal{F}_{a,H}| \leq \beta_{\Delta} n^{\frac{\|\boldsymbol{\alpha}\|_1}{2} - |S|}$. Hence

$$\begin{aligned} |\mathbb{E}[p(\mathbf{X}) - p^{\mathrm{Tr}}(\mathbf{X})]| &\leq c_{\Delta} \|p\| \sum \sum_{\boldsymbol{\alpha} \in \mathcal{F}_{a,H}} n^{-\frac{\|\boldsymbol{\alpha}\|\|}{2}} \\ &\leq c_{\Delta} \|p\| \sum_{\boldsymbol{\alpha} \in \mathcal{F}_{a,H}} \beta_{\Delta} n^{-|S|} \\ &\leq c \|p\| n^{-|S|}, \end{aligned}$$

where the first sum is taken over all pairs (a, H) such that $a \in [0, \Delta]$ and H is a graph in the orbit generated by G_{α} with $\alpha \in \mathcal{U}^{n \times n} \setminus \mathcal{T}_{n,2}, c_{p,\alpha} \neq 0$, and c is a constant depending on Δ . This concludes the proof of the first part of the lemma.

We now prove the second part. We have $\mathcal{U}^{n\times n} = \mathcal{U}_1^{n\times n} \cup \mathcal{U}_*^{n\times n}$ where $\mathcal{U}_1^{n\times n}, \mathcal{U}_*^{n\times n}$ are given in (2.1). Note in particular that $\forall \boldsymbol{\alpha} \in \mathcal{U}_1^{n\times n}, \mathbb{E}[\mathbf{Y}^{\boldsymbol{\alpha}}] = 0$. Furthermore, we have by Lemma 3.9 that $\forall \boldsymbol{\alpha} \in \mathcal{U}_*^{n\times n}, \mathbb{E}[\mathbf{Y}^{\boldsymbol{\alpha}}] = (1 + o(1))n^{-\|\boldsymbol{\alpha}\|_0} d^{-\frac{\|\boldsymbol{\alpha}\|_1}{2} + \|\boldsymbol{\alpha}\|_0}$. Hence

$$\begin{split} |\mathbb{E}[p(\mathbf{Y}) - p^{\mathrm{Tr}}(\mathbf{Y})]| &= \left| \sum_{\boldsymbol{\alpha} \in \mathcal{U}^{n \times n} \setminus \mathcal{T}_{n,2}} c_{p,\boldsymbol{\alpha}} \mathbb{E}[\mathbf{Y}^{\boldsymbol{\alpha}}] \right| \\ &= \left| \sum_{\boldsymbol{\alpha} \in \mathcal{U}^{n \times n}_{*} \setminus \mathcal{T}_{n,2}} c_{p,\boldsymbol{\alpha}} \mathbb{E}[\mathbf{Y}^{\boldsymbol{\alpha}}] \right| \\ &\leq \left| \sum_{\boldsymbol{\alpha} \in \mathcal{K} \cap \mathcal{U}^{n \times n}_{*} c_{p,\boldsymbol{\alpha}} \mathbb{E}[\mathbf{Y}^{\boldsymbol{\alpha}}] \right| + \left| \sum_{\substack{(\boldsymbol{\alpha}_{1}, \boldsymbol{\alpha}_{2}) \in \mathcal{W} \\ \boldsymbol{\alpha}_{1} + \boldsymbol{\alpha}_{2} \in \mathcal{U}^{n \times n}} c_{p,\boldsymbol{\alpha}_{1} + \boldsymbol{\alpha}_{2}} \mathbb{E}[\mathbf{Y}^{\boldsymbol{\alpha}_{1} + \boldsymbol{\alpha}_{2}}] \right| \\ &\leq \left\| p \| (1 + o(1)) \left(\underbrace{\sum_{\substack{\boldsymbol{\alpha} \in \mathcal{K} \cap \mathcal{U}^{n \times n}_{*} \\ \boldsymbol{\alpha} \in \mathcal{K} \cap \mathcal{U}^{n \times n}_{*}} n^{-\|\boldsymbol{\alpha}\|_{0}} d^{-\frac{\|\boldsymbol{\alpha}\|_{1}}{2} + \|\boldsymbol{\alpha}\|_{0}} + \sum_{\substack{(\boldsymbol{\alpha}_{1}, \boldsymbol{\alpha}_{2}) \in \mathcal{W} \\ \boldsymbol{\alpha}_{1} + \boldsymbol{\alpha}_{2} \in \mathcal{U}^{n \times n}_{*}} n^{-\|\boldsymbol{\alpha}\|_{0} - \frac{\|\boldsymbol{\alpha}_{2}\|_{1}}{2} + \|\boldsymbol{\alpha}\|_{0}} \right) \right\|_{\boldsymbol{\alpha}_{1} = \mathbf{1} + \mathbf{1} +$$

We will bound the above summation similarly to the Gaussian case. We first consider the summands in A indexed by $\boldsymbol{\alpha} \in \mathcal{K} \cap \mathcal{U}_*^{n \times n}$. We distinguish between two cases

• Case 1: $\boldsymbol{\alpha} \in \mathcal{T}$.

Then $|V_{\boldsymbol{\alpha}}| = |E_{\boldsymbol{\alpha}}| + 1 = \|\boldsymbol{\alpha}\|_0 + 1$. Furthermore $\boldsymbol{\alpha} \in \mathcal{U}_*^{n \times n} \implies \|\boldsymbol{\alpha}\|_0 \leq \frac{\|\boldsymbol{\alpha}\|_1}{2}$, and the equality occurs if and only if every nonzero entry in $\boldsymbol{\alpha}$ equals 2, but that would imply $\boldsymbol{\alpha} \in \mathcal{T}_{n,2}$. Therefore the inequality is strict, i.e. $\|\boldsymbol{\alpha}\|_0 \leq \frac{\|\boldsymbol{\alpha}\|_1 - 1}{2}$.

• Case 2: $\alpha \notin \mathcal{T}$.

Then $|V_{\boldsymbol{\alpha}}| \leq |E_{\boldsymbol{\alpha}}| = \|\boldsymbol{\alpha}\|_0$. Furthermore $\boldsymbol{\alpha} \in \mathcal{U}_*^{n \times n} \implies \|\boldsymbol{\alpha}\|_0 \leq \frac{\|\boldsymbol{\alpha}\|_1}{2}$.

Given $a \in [0, \Delta]$ and a graph H introduce the sets

$$\mathcal{F}_{a,H}^{1} \triangleq \{ \boldsymbol{\alpha} \in (\mathcal{K} \cap \mathcal{U}_{*}^{n \times n} \cap \mathcal{T}) \setminus \mathcal{T}_{n,2}, \|\boldsymbol{\alpha}\|_{1} = a, G_{\boldsymbol{\alpha}} \sim H \}, \mathcal{F}_{a,H}^{2} \triangleq \{ \boldsymbol{\alpha} \in \mathcal{K} \cap \mathcal{U}_{*}^{n \times n} \setminus (\mathcal{T}_{n,2} \cup \mathcal{T}), \|\boldsymbol{\alpha}\|_{1} = a, G_{\boldsymbol{\alpha}} \sim H \}.$$

$$(3.9)$$

Recalling $S \subset G_{\boldsymbol{\alpha}}$ it follows that there exists constants $\beta_{\Delta}^1, \beta_{\Delta}^2$ depending on Δ such that, $\forall \boldsymbol{\alpha} \in \mathcal{F}_{a,H}^1, |\mathcal{F}_{a,H}^1| \leq \beta_{\Delta}^1 n^{\|\boldsymbol{\alpha}\|_0 + 1 - |S|}$ and $\forall \boldsymbol{\alpha} \in \mathcal{F}_{a,H}^2, |\mathcal{F}_{a,H}^2| \leq \beta_{\Delta}^2 n^{\|\boldsymbol{\alpha}\|_0 - |S|}$. We then have

$$\begin{split} A &= \sum \sum_{\pmb{\alpha} \in \mathcal{F}_{a,H}^1} n^{-\|\pmb{\alpha}\|_0} d^{-\frac{\|\pmb{\alpha}\|_1}{2} + \|\pmb{\alpha}\|_0} + \sum \sum_{\pmb{\alpha} \in \mathcal{F}_{a,H}^2} n^{-\|\pmb{\alpha}\|_0} d^{-\frac{\|\pmb{\alpha}\|_1}{2} + \|\pmb{\alpha}\|_0} \\ &\leq \sum \sum_{\pmb{\alpha} \in \mathcal{F}_{a,H}^1} n^{-\|\pmb{\alpha}\|_0} d^{-\frac{1}{2}} + \sum \sum_{\pmb{\alpha} \in \mathcal{F}_{a,H}^2} n^{-\|\pmb{\alpha}\|_0} \\ &\leq \sum \beta_{\Delta}^1 n^{-|S|+1} d^{-\frac{1}{2}} + \sum \beta_{\Delta}^2 n^{-|S|} \\ &= O\left(n^{-|S|+1} d^{-\frac{1}{2}} \left(1 + \frac{d^{\frac{1}{2}}}{n}\right)\right) \end{split}$$

$$= O\left(n^{-|S|+1}d^{-\frac{1}{2}}\right).$$

where the first sums are taken over all pairs (a, H) such that $a \in [0, \Delta]$ and H is a graph in the orbit generated by G_{α} with $\alpha \in \mathcal{U}_*^{n \times n} \setminus \mathcal{T}_{n,2}, c_{p,\alpha} \neq 0$.

We now bound B similarly. Let $\boldsymbol{\alpha} \triangleq \boldsymbol{\alpha}_1 + \boldsymbol{\alpha}_2$ with $\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2$ as in B. We consider three cases

• Case 1: $\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2 \in \mathcal{T}$.

We have $|V_{\boldsymbol{\alpha}_1}| = |E_{\boldsymbol{\alpha}_1}| + 1 = \|\boldsymbol{\alpha}_1\|_0 + 1$ and $|V_{\boldsymbol{\alpha}_2}| = |E_{\boldsymbol{\alpha}_2}| + 1 = \|\boldsymbol{\alpha}_2\|_0 + 1$. Therefore $|V_{\boldsymbol{\alpha}}| = \|\boldsymbol{\alpha}_1\|_0 + \|\boldsymbol{\alpha}_2\|_0 + 2$. Furthermore, similarly to Case 1 in the previous analysis of A, we obtain $\|\boldsymbol{\alpha}_1\|_0 \leq \frac{\|\boldsymbol{\alpha}_1\|_{1-1}}{2}$, $\|\boldsymbol{\alpha}_2\|_0 \leq \frac{\|\boldsymbol{\alpha}_2\|_{1-1}}{2}$.

• Case 2: $\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2 \notin \mathcal{T}$.

Since $G_{\boldsymbol{\alpha}_1}$ is connected, it follows that $|V_{\boldsymbol{\alpha}_1}| \leq |E_{\boldsymbol{\alpha}_1}| = \|\boldsymbol{\alpha}_1\|_0$. Similarly $|V_{\boldsymbol{\alpha}_2}| \leq \|\boldsymbol{\alpha}_2\|_0$. Therefore $|V_{\boldsymbol{\alpha}}| = |V_{\boldsymbol{\alpha}_1}| + |V_{\boldsymbol{\alpha}_2}| \leq \|\boldsymbol{\alpha}_1\|_0 + \|\boldsymbol{\alpha}_2\|_0$. Furthermore, similarly to Case 2 in the previous analysis of A, we obtain $\|\boldsymbol{\alpha}_1\|_0 \leq \frac{\|\boldsymbol{\alpha}_1\|_1}{2}, \|\boldsymbol{\alpha}_2\|_0 \leq \frac{\|\boldsymbol{\alpha}_2\|_1}{2}$.

• Case 3: $\boldsymbol{\alpha}_1 \in \mathcal{T}, \boldsymbol{\alpha}_2 \notin \mathcal{T}.$

Combining the arguments of Cases 1, 2 above we obtain $|V_{\boldsymbol{\alpha}}| \leq \|\boldsymbol{\alpha}_1\|_0 + \|\boldsymbol{\alpha}_2\|_0 + 1$, and $\|\boldsymbol{\alpha}_1\|_0 \leq \frac{\|\boldsymbol{\alpha}_1\|_{1-1}}{2}, \|\boldsymbol{\alpha}_2\|_0 \leq \frac{\|\boldsymbol{\alpha}_2\|_1}{2}$.

Similarly to A, introduce the sets

$$\mathcal{F}_{a,H}^{1} \triangleq \{ (\boldsymbol{\alpha}_{1},\boldsymbol{\alpha}_{2}) \in \mathcal{W} | \boldsymbol{\alpha}_{1} + \boldsymbol{\alpha}_{2} \in \mathcal{U}_{*}^{n \times n}, \boldsymbol{\alpha}_{1}, \boldsymbol{\alpha}_{2} \in \mathcal{T}, \|\boldsymbol{\alpha}\|_{1} = a, G_{\boldsymbol{\alpha}} \sim H \},
\mathcal{F}_{a,H}^{2} \triangleq \{ (\boldsymbol{\alpha}_{1},\boldsymbol{\alpha}_{2}) \in \mathcal{W} | \boldsymbol{\alpha}_{1} + \boldsymbol{\alpha}_{2} \in \mathcal{U}_{*}^{n \times n}, \boldsymbol{\alpha}_{1}, \boldsymbol{\alpha}_{2} \notin \mathcal{T}, \|\boldsymbol{\alpha}\|_{1} = a, G_{\boldsymbol{\alpha}} \sim H \},
\mathcal{F}_{a,H}^{3} \triangleq \{ (\boldsymbol{\alpha}_{1},\boldsymbol{\alpha}_{2}) \in \mathcal{W} | \boldsymbol{\alpha}_{1} + \boldsymbol{\alpha}_{2} \in \mathcal{U}_{*}^{n \times n}, \boldsymbol{\alpha}_{1} \in \mathcal{T}, \boldsymbol{\alpha}_{2} \notin \mathcal{T}, \|\boldsymbol{\alpha}\|_{1} = a, G_{\boldsymbol{\alpha}} \sim H \}.$$
(3.10)

There exists constants $\beta_{\Delta}^1, \beta_{\Delta}^2, \beta_{\Delta}^3$ depending on Δ such that, $\forall (\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2) \in \mathcal{F}_{a,H}^1, |\mathcal{F}_{a,H}^1| \leq \beta_{\Delta}^1 n^{\|\boldsymbol{\alpha}_1\|_0 + \|\boldsymbol{\alpha}_2\|_0 + 2 - |S|}$, $\forall (\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2) \in \mathcal{F}_{a,H}^3, |\mathcal{F}_{a,H}^3| \leq \beta_{\Delta}^2 n^{\|\boldsymbol{\alpha}_1\|_0 + \|\boldsymbol{\alpha}_2\|_0 - |S|}$, and $\forall (\boldsymbol{\alpha}_1, \boldsymbol{\alpha}_2) \in \mathcal{F}_{a,H}^3, |\mathcal{F}_{a,H}^3| \leq \beta_{\Delta}^2 n^{\|\boldsymbol{\alpha}_1\|_0 + \|\boldsymbol{\alpha}_2\|_0 + 1 - |S|}$. We then have

$$\begin{split} B &= \sum \sum_{(\pmb{\alpha}_1,\pmb{\alpha}_2)\in \mathcal{F}^1_{a,H}} n^{-\|\pmb{\alpha}_1\|_0 + \pmb{\alpha}_2\|_0} d^{-\frac{\|\pmb{\alpha}_1\|_1 + \|\pmb{\alpha}_2\|_1}{2}} + \|\pmb{\alpha}_1\|_0 + \|\pmb{\alpha}_2\|_0} \\ &+ \sum \sum_{(\pmb{\alpha}_1,\pmb{\alpha}_2)\in \mathcal{F}^2_{a,H}} n^{-\|\pmb{\alpha}_1\|_0 + \pmb{\alpha}_2\|_0} d^{-\frac{\|\pmb{\alpha}_1\|_1 + \|\pmb{\alpha}_2\|_1}{2}} + \|\pmb{\alpha}_1\|_0 + \|\pmb{\alpha}_2\|_0} \\ &+ \sum \sum_{(\pmb{\alpha}_1,\pmb{\alpha}_2)\in \mathcal{F}^3_{a,H}} n^{-\|\pmb{\alpha}_1\|_0 + \pmb{\alpha}_2\|_0} d^{-\frac{\|\pmb{\alpha}_1\|_1 + \|\pmb{\alpha}_2\|_1}{2}} + \|\pmb{\alpha}_1\|_0 + \|\pmb{\alpha}_2\|_0} \\ &\leq \sum \beta_{\Delta}^1 n^{-|S|+2} d^{-1} + \sum \beta_{\Delta}^2 n^{-|S|} + \sum \beta_{\Delta}^3 n^{-|S|+1} d^{-\frac{1}{2}} \\ &= O\left(n^{-|S|+2} d^{-1} \left(1 + \frac{d}{n^2} + \frac{d^{\frac{1}{2}}}{n}\right)\right) \\ &= O\left(n^{-|S|+2} d^{-1}\right). \end{split}$$

Combining the bounds on A and B yields

$$|\mathbb{E}[p(\mathbf{Y}) - p^{\mathrm{Tr}}(\mathbf{Y})]| \le c ||p|| n^{-|S|+2} d^{-1},$$

where c is a constant. Finally, note that if we further assume p connected, then B is removed from our analysis. Therefore we obtain the tighter bound

$$|\mathbb{E}[p(\mathbf{Y}) - p^{\mathrm{Tr}}(\mathbf{Y})]| \le c ||p|| n^{-|S|+1} d^{-\frac{1}{2}},$$

from the analysis of A only. This concludes the proof of the second part of the lemma.

Lemma 3.12. Let $\mathbf{X} \stackrel{d}{=} \operatorname{GOE}(n)$ and $\mathbf{Y} \stackrel{d}{=} \mathcal{G}(n, d/n)$. Suppose $L_1 \cup L_2$ is a partition of the ordered pairs $1 \leq i < j \leq n$. Consider the (symmetric) matrix \mathbf{Z} given by

$$\mathbf{Z}_{ij} = \mathbf{X}_{ij} \quad if \quad (i,j) \in L_1,
\mathbf{Z}_{ij} = \mathbf{Y}_{ij} \quad if \quad (i,j) \in L_2,$$
(3.11)

and $\mathbf{Z}_{ii} = 0$. Let $S \subset [n]$, $q \in \mathbb{R}[\mathbf{X}]_{\leq \Delta}$, and e = (i, j) with $1 \leq i < j \leq n$. Suppose $\mathbf{Z}_e q(\mathbf{Z})$ is connected and contains all nodes in S. For every Δ , there exists $c = c(\Delta)$ independent of n, d such that

$$\begin{aligned} |\mathbb{E}[|\mathbf{Z}_{e}|^{\ell}q(\mathbf{Z})]| &= \frac{c(\Delta)||q||}{n^{|S| + \frac{\ell}{2} - 2}} & \text{if } e \in L_{1}, and \quad \ell \geq 2, \\ |\mathbb{E}[|\mathbf{Z}_{e}|^{\ell}q(\mathbf{Z})]| &= \frac{c(\Delta)||q||}{n^{|S| - 1}d^{\frac{\ell}{2} - 1}} & \text{if } e \in L_{2}, and \quad \ell \geq 2. \end{aligned}$$
(3.12)

Proof. For $\boldsymbol{\alpha} \in \mathcal{U}^{n \times n}$, we will write $\boldsymbol{\alpha} = \boldsymbol{\alpha}^1 + \boldsymbol{\alpha}^2$ where $\boldsymbol{\alpha}_{ij}^r = \boldsymbol{\alpha}_{ij}$ if $(i, j) \in L_r$, and $\boldsymbol{\alpha}_{ij}^r = 0$ otherwise for r = 1, 2. Consider the case $e \in L_1$ in (3.12) and denote by $w(\mathbf{Z})$ the LDP $\mathbf{Z}_e^{\ell}q(\mathbf{Z})$. We will use the same notations introduced in the proof of Lemma 3.11. For $\boldsymbol{\alpha} \in \mathcal{U}^{n \times n}$, we let $\bar{\boldsymbol{\alpha}} \in \mathcal{U}^{n \times n}$ where $\bar{\boldsymbol{\alpha}}_{ij} = \boldsymbol{\alpha}_{ij}, \forall (ij) \neq e$ and $\bar{\boldsymbol{\alpha}}_e = 0$. Note that $\mathbb{E}[\mathbf{X}^{\boldsymbol{\alpha}}] = \gamma_{\boldsymbol{\alpha}} n^{-\|\boldsymbol{\alpha}\|_1/2}$ where $\gamma_{\boldsymbol{\alpha}} = \prod_{\substack{1 \le i < j \le n \\ 2|\boldsymbol{\alpha}_{ij}, \boldsymbol{\alpha}_{ij} > 0}} (\boldsymbol{\alpha}_{ij} - 1)!!$ and $|\gamma_{\boldsymbol{\alpha}}| \le c_{\Delta}$ for some constant c_{Δ} , and similarly $\mathbb{E}[|\mathbf{X}|^{\boldsymbol{\alpha}}] \le c'_{\Delta} n^{-\|\boldsymbol{\alpha}\|_1/2}$ for some constant c'_{Δ} . Using the latter and Lemma 3.9

we have

$$\begin{split} |\mathbb{E}\left[w(\mathbf{Z})\right]| &= \left|\sum_{\boldsymbol{\alpha}} c_{q,\boldsymbol{\alpha}} \mathbb{E}\left[|\mathbf{Z}_{e}|^{\ell} \mathbf{Z}^{\boldsymbol{\alpha}}\right]\right| \\ &= \left|\sum_{\boldsymbol{\alpha}} c_{q,\boldsymbol{\alpha}} \mathbb{E}\left[|\mathbf{X}_{e}|^{\ell+\boldsymbol{\alpha}_{e}^{1}}\right] \mathbb{E}\left[\mathbf{X}^{\bar{\boldsymbol{\alpha}}^{1}}\right] \mathbb{E}\left[\mathbf{Y}^{\boldsymbol{\alpha}^{2}}\right]\right| \\ &\leq \|q\|c_{\Delta}c_{\Delta}'\left|\sum_{\boldsymbol{\alpha},c_{q,\boldsymbol{\alpha}}\neq 0} n^{-\frac{\ell+|\boldsymbol{\alpha}_{e}^{1}|+\|\bar{\boldsymbol{\alpha}}^{1}\|_{1}}{2}} \mathbf{1}_{2|\bar{\boldsymbol{\alpha}}^{1}} \mathbf{1}_{\boldsymbol{\alpha}^{2}\in\mathcal{U}_{*}^{n\times n}} ((-1)^{\|\boldsymbol{\alpha}^{2}\|_{1}} + o(1))n^{-\|\boldsymbol{\alpha}^{2}\|_{0}} d^{-\frac{\|\boldsymbol{\alpha}^{2}\|_{1}}{2}} + \|\boldsymbol{\alpha}^{2}\|_{0}}\right|, \\ &\leq 2\|q\|c_{\Delta}c_{\Delta}'\sum_{\boldsymbol{\alpha},c_{q,\boldsymbol{\alpha}}\neq 0} \mathbf{1}_{2|\bar{\boldsymbol{\alpha}}^{1}} \mathbf{1}_{\boldsymbol{\alpha}^{2}\in\mathcal{U}_{*}^{n\times n}} n^{-\frac{\ell+\|\boldsymbol{\alpha}^{1}\|_{1}}{2}} - \|\boldsymbol{\alpha}^{2}\|_{0} d^{-\frac{\|\boldsymbol{\alpha}^{2}\|_{1}}{2}} + \|\boldsymbol{\alpha}^{2}\|_{0}, \end{split}$$

where we upper bounded 1 + o(1) by 2 for large enough n, d. Since $\boldsymbol{\alpha}^2 \in \mathcal{U}_*^{n \times n}$, we have $-\frac{\|\boldsymbol{\alpha}^2\|_1}{2} + \|\boldsymbol{\alpha}^2\|_0 \leq 0$. Therefore

$$\left|\mathbb{E}\left[w(\mathbf{Z})\right]\right| \leq 2\|q\|c_{\Delta}c'_{\Delta}\sum_{\boldsymbol{\alpha}\in\mathcal{C}}\mathbf{1}_{2|\bar{\boldsymbol{\alpha}}^{1}}\mathbf{1}_{\boldsymbol{\alpha}^{2}\in\mathcal{U}_{*}^{n\times n}}n^{-\frac{\ell+\|\boldsymbol{\alpha}^{1}\|_{1}}{2}-\|\boldsymbol{\alpha}^{2}\|_{0}}$$

Let $\hat{\boldsymbol{\alpha}} = \ell[e] + \boldsymbol{\alpha}$ (so that $\mathbf{Z}^{\hat{\boldsymbol{\alpha}}} = \mathbf{Z}_{e}^{\ell} \cdot \mathbf{Z}^{\boldsymbol{\alpha}}$). Since $2|\bar{\boldsymbol{\alpha}}^{1}$, it follows that $\|\hat{\boldsymbol{\alpha}}^{1}\|_{0} \leq \frac{\|\hat{\boldsymbol{\alpha}}^{1}\|_{1} - (\ell-2)}{2}$. Since the LDP w is connected, it follows that $\hat{\boldsymbol{\alpha}}$ is connected, and furthermore $|V_{\hat{\boldsymbol{\alpha}}}| \leq |E_{\hat{\boldsymbol{\alpha}}}| + 1 = \|\hat{\boldsymbol{\alpha}}\|_{0} + 1 = \|\hat{\boldsymbol{\alpha}}^{1}\|_{0} + \|\boldsymbol{\alpha}^{2}\|_{0} + 1 \leq \frac{\|\boldsymbol{\alpha}^{1}\|_{1} - (\ell-2)}{2} + \|\boldsymbol{\alpha}^{2}\|_{0} + 1 = \frac{\|\hat{\boldsymbol{\alpha}}^{1}\|_{1}}{2} + \|\boldsymbol{\alpha}^{2}\|_{0} + 2 - \frac{\ell}{2}$. For $a \in [0, \Delta]$ and a graph H introduce the set

$$\mathcal{F}_{a,H} \triangleq \{ \boldsymbol{\alpha} | c_{q,\boldsymbol{\alpha}} \neq 0, \| \boldsymbol{\alpha} \|_{1} = a, G_{\boldsymbol{\alpha}} \sim H, 2 | \bar{\boldsymbol{\alpha}}^{1}, \boldsymbol{\alpha}^{2} \in \mathcal{U}_{*}^{n \times n} \}.$$
(3.13)

Recalling that $S \subset w$, it follows that $\forall \boldsymbol{\alpha} \in \mathcal{F}_{a,H}, |\mathcal{F}_{a,H}| \leq \beta_{\Delta} n^{|V_{\hat{\boldsymbol{\alpha}}}| - |S|} \leq \beta_{\Delta} n^{\frac{\|\hat{\boldsymbol{\alpha}}^{\perp}\|_{1}}{2} + \|\boldsymbol{\alpha}^{2}\|_{0} + 2 - \frac{\ell}{2} - |S|}$, where β_{Δ} is a constant depending on Δ . Hence

$$\begin{split} |\mathbb{E}\left[w(\mathbf{Z})\right]| &\leq 2 \|q\| c_{\Delta} c_{\Delta}' \sum \sum_{\boldsymbol{\alpha} \in \mathcal{F}_{a,H}} \mathbf{1}_{2|\bar{\boldsymbol{\alpha}}^{1}} \mathbf{1}_{\boldsymbol{\alpha}^{2} \in \mathcal{U}_{*}^{n \times n}} n^{-\frac{\ell + \|\boldsymbol{\alpha}^{1}\|_{1}}{2} - \|\boldsymbol{\alpha}^{2}\|_{0}} \\ &= 2 \|q\| c_{\Delta} c_{\Delta}' \sum \sum_{\boldsymbol{\alpha} \in \mathcal{F}_{a,H}} \mathbf{1}_{2|\bar{\boldsymbol{\alpha}}^{1}} \mathbf{1}_{\boldsymbol{\alpha}^{2} \in \mathcal{U}_{*}^{n \times n}} n^{-\frac{\|\boldsymbol{\hat{\alpha}}^{1}\|_{1}}{2} - \|\boldsymbol{\alpha}^{2}\|_{0}} \\ &\leq 2 \|q\| c_{\Delta} c_{\Delta}' \sum \beta_{\Delta} n^{2 - \frac{\ell}{2} - |S|} \\ &= \frac{O(\|q\|)}{n^{|S| + \frac{\ell}{2} - 2}}, \end{split}$$

where the first sum is taken over over all pairs (a, H) such that $a \in [0, \Delta]$ and H is a graph in the orbit generated by G_{α} with $c_{q,\alpha} \neq 0$. Denoting by $c(\Delta)$ the constant hidden in $O(\cdot)$, we obtain the proof for the case $e \in L_1$.

Consider now the case $e \in L_2$ and denote again by $w(\mathbf{Z})$ the LDP $\mathbf{Z}_e^{\ell}q(\mathbf{Z})$. We have similarly to the previous case

$$|\mathbb{E}\left[w(\mathbf{Z})\right]| \leq 2 \|q\| c_{\Delta} c_{\Delta}' \sum_{\boldsymbol{\alpha} \in \mathcal{C}} \mathbf{1}_{2|\boldsymbol{\alpha}^1} \mathbf{1}_{\bar{\boldsymbol{\alpha}}^2 \in \mathcal{U}^{n \times n}_*} n^{-\frac{\|\boldsymbol{\alpha}^1\|_1}{2} - \|\hat{\boldsymbol{\alpha}}^2\|_0} d^{-\frac{\|\hat{\boldsymbol{\alpha}}^2\|_1}{2} + \|\hat{\boldsymbol{\alpha}}^2\|_0}.$$

As $\ell \geq 2$, it follows that $\bar{\boldsymbol{\alpha}}^2 \in \mathcal{U}_*^{n \times n}$ if and only if $\hat{\boldsymbol{\alpha}}^2 \in \mathcal{U}_*^{n \times n}$. Let $\boldsymbol{\alpha}$ satisfy $2|\boldsymbol{\alpha}^1, \bar{\boldsymbol{\alpha}}^2 \in \mathcal{U}_*^{n \times n}$. Since $\hat{\boldsymbol{\alpha}}^2 \in \mathcal{U}_*^{n \times n}$, all the nonzero values in $\hat{\boldsymbol{\alpha}}^2$ are at least 2. It follows that $\|\hat{\boldsymbol{\alpha}}^2\|_0 \leq \frac{\|\hat{\boldsymbol{\alpha}}^2\|_1 - (\ell-2)}{2}$ and thus $-\frac{\|\hat{\boldsymbol{\alpha}}^2\|_1}{2} + \|\hat{\boldsymbol{\alpha}}^2\|_0 \leq -\frac{\ell}{2} + 1$. Hence

$$\left|\mathbb{E}\left[w(\mathbf{Z})\right]\right| \leq 2\|q\|c_{\Delta}c_{\Delta}'d^{-\frac{\ell}{2}+1}\sum_{\boldsymbol{\alpha},c_{q,\boldsymbol{\alpha}}\neq 0}\mathbf{1}_{2|\boldsymbol{\alpha}^{1}}\mathbf{1}_{\bar{\boldsymbol{\alpha}}^{2}\in\mathcal{U}_{*}^{n\times n}}n^{-\frac{\|\boldsymbol{\alpha}^{1}\|_{1}}{2}-\|\hat{\boldsymbol{\alpha}}^{2}\|_{0}}$$

Since w is connected, it follows that $\hat{\boldsymbol{\alpha}}$ is connected. Therefore, we have $|V_{\hat{\boldsymbol{\alpha}}}| \leq |E_{\hat{\boldsymbol{\alpha}}}| + 1 = \|\hat{\boldsymbol{\alpha}}\|_0 + 1 = \|\boldsymbol{\alpha}^1\|_0 + \|\boldsymbol{\alpha}^2\|_0 + 1 \leq \frac{\|\boldsymbol{\alpha}^1\|_1}{2} + \|\hat{\boldsymbol{\alpha}}^2\|_0 + 1$, where the last inequality follows from $2|\boldsymbol{\alpha}^1$. Recalling that $S \subset w$ and using the same notation for $\mathcal{F}_{a,\Delta}$, it follows that $\forall \boldsymbol{\alpha} \in \mathcal{F}_{a,H}, |\mathcal{F}_{a,H}| \leq \beta_{\Delta} n^{|V_{\hat{\boldsymbol{\alpha}}}| - |S|} \leq \beta_{\Delta} n^{\frac{\|\boldsymbol{\alpha}^1\|_1}{2} + \|\hat{\boldsymbol{\alpha}}^2\|_0 + 1 - |S|}$. Hence

$$\begin{split} |\mathbb{E}[w(\mathbf{Z})]| &\leq 2 \|q\| c_{\Delta} c'_{\Delta} d^{-\frac{\ell}{2}+1} \sum \sum_{\boldsymbol{\alpha} \in \mathcal{F}_{a,H}} \mathbf{1}_{2|\boldsymbol{\alpha}^{1}} \mathbf{1}_{\bar{\boldsymbol{\alpha}}^{2} \in \mathcal{U}_{*}^{n \times n}} n^{-\frac{\|\boldsymbol{\alpha}^{1}\|_{1}}{2} - \|\hat{\boldsymbol{\alpha}}^{2}\|_{0}} \\ &\leq 2 \|q\| c_{\Delta} c'_{\Delta} d^{-\frac{\ell}{2}+1} \sum \beta_{\Delta} n^{1-|S|} \\ &= \frac{O(\|q\|)}{n^{|S|-1} d^{\frac{\ell}{2}-1}}, \end{split}$$

where the first sum is taken over all pairs (a, H) such that $a \in [0, \Delta]$ and H is a graph in the orbit generated by G_{α} with $\alpha \in \mathcal{C}, c_{q,\alpha} \neq 0$. Denoting by $c(\Delta)$ the constant hidden in $O(\cdot)$, we obtain the proof for the case $e \in L_2$. This concludes the proof.

Lemma 3.13. Fix $\ell \in [\binom{n}{2}]$ Suppose $L_1 \cup L_2$ is a partition of pairs $1 \leq i < j \leq n, (i, j) \neq \ell$ with $L_1 \neq \emptyset$. Given $i \in [n]$ let $u_{i,\ell}, a_{i,\ell}, b_{i,\ell}$ be LDP in variables $\mathbf{x} = (\mathbf{x}_{ij}, 1 \leq i < j \leq n)$. Let $\beta \geq 2$ satisfy $\beta = \tilde{O}(1)$, and let for $z \in \mathbb{R}$

$$v_{i,\ell}(z) = u_{i,\ell} + za_{i,\ell} + z^2 b_{i,\ell}, \qquad i \in [n],$$
$$k(z) = \sum_{i=1}^n \pi_\beta (v_{i,\ell}(z))^2,$$

where π_{β} is defined in Lemma 3.3. Suppose that in addition

- 1. $\forall i \in [n], u_{i,\ell}(\mathbf{x}), \mathbf{x}_{\ell} a_{i,\ell}(\mathbf{x}), \mathbf{x}_{\ell} b_{i,\ell}(\mathbf{x})$ are connected and contain node *i*.
- 2. $\forall i \in [n], u_{i,\ell}(\mathbf{x}), a_{i,\ell}(\mathbf{x}), b_{i,\ell}(\mathbf{x})$ do not depend on the variable \mathbf{x}_{ℓ} .
- 3. $\max_{i \in [n]} \|u_{i,\ell}\| + \|a_{i,\ell}\| + \|b_{i,\ell}\| = O(1).$

Let $\mathbf{X} \stackrel{\mathsf{d}}{=} \operatorname{GOE}(n), \mathbf{Y} \stackrel{\mathsf{d}}{=} \mathcal{G}(n, d/n)$. Let \mathbf{Z}^{ℓ} be given by $\mathbf{Z}_{ii}^{\ell} = 0, i \in [n]$ and

$$\begin{aligned} [\mathbf{Z}^{\ell}]_{ij} &= \mathbf{X}_{ij} \qquad if \quad (i,j) \in L_1, \\ [\mathbf{Z}^{\ell}]_{ij} &= \mathbf{Y}_{ij} \qquad if \quad (i,j) \in L_2. \end{aligned}$$

$$(3.14)$$

Then for every $(s_1, m_1, s_2, m_2) \in \mathbb{Z}_{\geq 0}^4$ such that $s_1m_1 + s_2m_2 = 3$ it holds

$$\forall |x_{\ell}| \le |\mathbf{X}_{\ell}|, \quad \mathbb{E}\left[\left|\mathbf{X}_{\ell}^{3}(k^{(s_{1})})^{m_{1}}(x_{\ell})(k^{(s_{2})})^{m_{2}}(x_{\ell})\right|\right] = \tilde{O}(n^{m_{1}+m_{2}-\frac{5}{2}}), \tag{3.15}$$

$$\forall |y_{\ell}| \le |\mathbf{Y}_{\ell}|, \quad \mathbb{E}\left[\left|\mathbf{Y}_{\ell}^{3}(k^{(s_{1})})^{m_{1}}(y_{\ell})(k^{(s_{2})})^{m_{2}}(y_{\ell})\right|\right] = \tilde{O}(n^{m_{1}+m_{2}-2}d^{-\frac{1}{2}}). \tag{3.16}$$

Proof. We use the slightly abused notation $u_{i,\ell}, a_{i,\ell}, b_{i,\ell}$ to denote $u_{i,\ell}(\mathbf{Z}^{\ell}), a_{i,\ell}(\mathbf{Z}^{\ell}), b_{i,\ell}(\mathbf{Z}^{\ell})$. We first show (3.15), (3.16). We explicit the proof here for the particular case $(s_1, m_1, s_2, m_2) = (2, 1, 1, 1)$, the remaining cases will follow from similar arguments. Using Lemma 3.7

$$\begin{aligned} |\mathbf{X}_{\ell}^{3} \cdot k^{(2)}(x_{\ell}) \cdot k^{(1)}(x_{\ell})| &\lesssim \sum \sum |\mathbf{X}_{\ell}|^{3+j_{1}^{1}+j_{1}^{2}} |a_{i_{1},\ell}|^{j_{2}^{1}} |b_{i_{1},\ell}|^{j_{3}^{1}} |u_{i_{1},\ell}|^{j_{4}^{1}} |a_{i_{2},\ell}|^{j_{2}^{2}} |b_{i_{2},\ell}|^{j_{3}^{2}} |u_{i_{2},\ell}|^{j_{4}^{2}} \\ &\lesssim \sum \sum |\mathbf{X}_{\ell}|^{3+2j_{1}^{1}} (a_{i_{1},\ell}^{j_{2}^{1}} b_{i_{1},\ell}^{j_{3}^{1}} u_{i_{1},\ell}^{j_{4}^{1}})^{2} + |\mathbf{X}_{\ell}|^{3+2j_{1}^{2}} (a_{i_{2},\ell}^{j_{2}^{2}} b_{i_{2},\ell}^{j_{3}^{2}} u_{i_{2},\ell}^{j_{4}^{2}})^{2}. \end{aligned}$$
(3.17)

Where the first summations are taken over the set $\mathcal{I} \triangleq \{(i_1, i_2), 1 \leq i_1, i_2 \leq n\}$, and the second summations are taken over the set

$$\mathcal{I} \triangleq \left\{ j_s^t, t \in [2], s \in [4] | \max_{s \in [4]} (j_s^1) \le 6, \max_{s \in [4]} (j_s^2) \le 5, \min_{t \in [2]} (j_2^t + j_3^t) \ge 1 \right\}.$$

Furthermore, line (3.17) follows from the arithmetic geometric inequality. Using Lemma 3.5, it follows that $\mathbf{X}_{\ell}^{3+2j_1^t}(a_{i_t,\ell}^{j_2^t}b_{i_t,\ell}^{j_t^t}a_{i_t,\ell}^{j_t^t})^2$ for t = 1, 2 are LDPs with O(1) coefficients. Moreover, these LDPs are connected polynomials containing node i_t . Indeed, these LDPs are product of powers of the following terms: $\mathbf{X}_{\ell}a_{i_t,\ell}, \mathbf{X}_{\ell}b_{i_t,\ell}, \mathbf{X}_{\ell}a_{i_t,\ell}b_{i_t,\ell}, u_{i,\ell}$, all of which are connected and share node i_t by assumption. (Note that the connectivity of $\mathbf{X}_{\ell}a_{i_t,\ell}b_{i_t,\ell}$ follows trivially from the connectivity of $\mathbf{X}_{\ell}a_{i_t,\ell}, \mathbf{X}_{\ell}b_{i_t,\ell}$ and the fact that the latter two LDPs share node i_t).

Since the LDP $\mathbf{X}_{\ell}^{3+2j_1^t}(a_{i_t,\ell}^{j_2^t}b_{i_t,\ell}^{j_t^t}u_{i_t,\ell}^{j_t^t})^2$ contains node i_t for t = 1, 2, it follows by Lemma 3.6 (using $S_1 = \{i_1\}$ and $S_2 = \{i_2\}$ respectively), that the resulting LDP from summing the LDPs above over \mathcal{I}, \mathcal{J} has O(n) coefficients, where the O(n) is due to double counting from the sum on \mathcal{I} . In particular, we have

$$\begin{aligned} |\mathbf{X}_{\ell}^{3} \cdot k^{(2)}(x_{\ell}) \cdot k^{(1)}(x_{\ell})| &\lesssim n \mathbf{X}_{\ell}^{3} w(\mathbf{Z}^{\ell}), \\ |\mathbf{Y}_{\ell}^{3} \cdot k^{(2)}(y_{\ell}) \cdot k^{(1)}(y_{\ell})| &\lesssim n \mathbf{Y}_{\ell}^{3} w(\mathbf{Z}^{\ell+1}), \end{aligned}$$

where w is a LDP with ||w|| = O(1) and $\mathbf{X}^{3}_{\ell}w(\mathbf{Z}^{\ell}), \mathbf{Y}^{3}_{\ell}w(\mathbf{Z}^{\ell})$ are connected. Applying (3.12) in Lemma 3.12 (with S the set of endpoints of edge ℓ) yields

$$\left| \mathbb{E} \left[n \mathbf{X}_{\ell}^{3} w(\mathbf{Z}^{\ell}) \right] \right| = \tilde{O}(n^{-\frac{1}{2}})$$
$$\left| \mathbb{E} \left[n \mathbf{Y}_{\ell}^{3} w(\mathbf{Z}^{\ell+1}) \right] \right| = \tilde{O}(d^{-\frac{1}{2}}).$$

Therefore

$$\mathbb{E}\left[|\mathbf{X}_{\ell}^{3} \cdot k^{(2)}(x_{\ell}) \cdot k^{(1)}(x_{\ell})|\right] = \tilde{O}(n^{-\frac{1}{2}}) = \tilde{O}(n^{m_{1}+m_{2}-\frac{5}{2}}),$$
$$\mathbb{E}\left[|\mathbf{Y}_{\ell}^{3} \cdot k^{(2)}(y_{\ell}) \cdot k^{(1)}(y_{\ell})|\right] = \tilde{O}(d^{-\frac{1}{2}}) = \tilde{O}(n^{m_{1}+m_{2}-2}d^{-\frac{1}{2}}),$$

which ends the proof of (3.15), (3.16) with $(s_1, m_1, s_2, m_2) = (2, 1, 1, 1)$. The case given by $(s_1, m_1, s_2, m_2) = (1, 3, 0, 0)$ is identical in treatment, specifically, we can use Lemma 3.7 to obtain the following bound,

$$|\mathbf{X}_{\ell}^{3} \cdot k^{(1)}(x_{\ell})^{3}| \lesssim \sum_{\mathcal{I}} \sum_{\mathcal{J}} \sum_{t=1}^{3} |\mathbf{X}_{\ell}|^{3 + (4\mathbf{1}_{t \leq 2} + 2\mathbf{1}_{t=3})j_{1}^{t}} (a_{i_{t},\ell}^{j_{2}} b_{i_{t},\ell}^{j_{3}^{t}} u_{i_{t},\ell}^{j_{4}^{t}})^{4\mathbf{1}_{t \leq 2} + 2\mathbf{1}_{t=3}},$$

where

$$\mathcal{I} \triangleq \{i_t, t \in [3] | i_t \in [n]\}, \mathcal{J} \triangleq \{j_s^t, t \in [3], s \in [4] | \max_{s \in [4], t \in [3]} (j_t) \le 5, \min_{t \in [3]} (j_2^t + j_3^t) \ge 1\}.$$

Moreover, the same bound holds for **Y**. It suffices then to notice that each summand in the above sum is a connected LDP containing the corresponding i_t node. Applying Lemma 3.6 yields that each summation can be written as $n^{m_1+m_2-1}|\mathbf{X}_{\ell}|^3 w(\mathbf{Z}^{\ell})$ where $\mathbf{X}_{\ell}^3 w(\mathbf{Z}^{\ell})$ is a connected LDP with O(1) coefficients, and the $n^{m_1+m_2-1}$ accounts for double counting in the summation over all indices in $\{i_1, i_2, i_3\}$. The result of (3.15) and (3.16) then readily follows in the above cases by applying (3.12) in Lemma 3.12 (with S the set of endpoints of edge ℓ).

It remains to deal with the case $(s_1, m_1, s_2, m_2) = (3, 1, 0, 0)$ which we do next. We have from Lemma 3.7

$$\begin{split} |\mathbf{X}_{\ell}^{3} \cdot k^{(3)}(x_{\ell})| &\lesssim |\mathbf{X}_{\ell}|^{3} \sum_{i=1}^{n} \sum_{\substack{j_{1}, j_{2}, j_{3}, j_{4} \leq 8\\ j_{2}+j_{3} \geq 2}} |\mathbf{X}_{\ell}|^{j_{1}} |a_{i,\ell}|^{j_{2}} |b_{i,\ell}|^{j_{3}} |u_{i,\ell}|^{j_{4}} \\ &\lesssim \sum_{i=1}^{n} \sum_{\substack{j_{1}, j_{2}, j_{3}, j_{4} \leq 8\\ j_{2}+j_{3} \geq 2}} |\mathbf{X}_{\ell}|^{3+2j_{1}} \left(a_{i,\ell}^{2j_{2}} \mathbf{1}_{j_{2}, j_{3} \geq 1} + b_{i,\ell}^{2} \mathbf{1}_{j_{2}=0} + a_{i,\ell}^{2j_{2}-2} \mathbf{1}_{j_{3}=0}\right) |u_{i,\ell}|^{2j_{4}} \\ &+ |\mathbf{X}_{\ell}|^{3} \left(b_{i,\ell}^{2j_{3}} \mathbf{1}_{j_{2}, j_{3} \geq 1} + b_{i,\ell}^{2j_{3}-2} \mathbf{1}_{j_{2}=0} + a_{i,\ell}^{2} \mathbf{1}_{j_{3}=0}\right), \end{split}$$

where the last line follows from the arithmetic geometric inequality. Note in particular that each term in $a_{i,\ell}, b_{i,\ell}$ in the summand has a positive even exponent. Moreover, all the terms in the summand are connected LDPs containing node *i*. Using Lemma 3.6, it follows that the summation can then be written as $|\mathbf{X}_{\ell}|^3 w(\mathbf{Z}^{\ell})$ where $\mathbf{X}_{\ell}^3 w(\mathbf{Z}^{\ell})$ is a connected LDP with O(1) coefficients. We can then apply (3.12) in Lemma 3.12 to obtain

$$\mathbb{E}\left[\left|\mathbf{X}_{\ell}^{3} \cdot k^{(3)}(x_{\ell})\right|\right] = \tilde{O}(n^{-\frac{3}{2}}) = \tilde{O}(n^{m_{1}+m_{2}-\frac{5}{2}}),\\ \mathbb{E}\left[\left|\mathbf{Y}_{\ell}^{3} \cdot k^{(3)}(y_{\ell})\right|\right] = \tilde{O}(n^{-1}d^{-\frac{1}{2}}) = \tilde{O}(n^{m_{1}+m_{2}-2}d^{-\frac{1}{2}}).$$

This concludes the proof of (3.15) and (3.16).

4 Proof of Theorem 2.1

In this section we complete the proof of our first main result, Theorem 2.1. Thus fix a sequence of polynomials $p = (p_i, i \in [n])$ satisfying the assumptions of the theorem. We split proving (2.5) and (2.6) in two separate subsections.

4.1 Proof of (2.5)

Proof. For $1 \le i < j \le n$, introduce the LDP

$$r_{ij}(\mathbf{X}) \triangleq p_i(\mathbf{X}) \mathbf{X}_{ij} p_j(\mathbf{X}).$$

Since p_i, p_j contain nodes i, j respectively, it follows that r_{ij} is connected and contains both nodes i and j. Indeed the factor graphs of the monomials in r_{ij} are the union of: a tree rooted at i, a tree rooted at j, and the edge-tree [ij]. Furthermore, using Lemma 3.5, it follows that

$$||r_{ij}|| \le 2^{2\Delta+1} ||p_i|| ||p_j|| = O(1)$$

where the last part follows from the assumption $||p_i|| = O(1)$. Thus r_{ij} is a connected LDP with constant coefficients. In fact the following stronger property holds, as an implication of which the contribution of the non-tree parts of r_{ij} is asymptotically negligible.

Lemma 4.1. The following hold

$$\left\|\sum_{1 \le i < j \le n} r_{ij}\right\| = O(1). \tag{4.1}$$

Furthermore, for every $\epsilon > 0$

$$\frac{1}{n} \sum_{1 \le i < j \le n} r_{ij}^{\text{Tr}}(\mathbf{X}) \ge \mathcal{P} - \epsilon,$$
(4.2)

whp as $n \to \infty$.

Proof of Lemma 4.1. Let $q \triangleq \sum_{1 \leq i < j \leq n} (r_{ij} - r_{ij}^{\text{Tr}})$. Since $q^{\text{Tr}} = 0$ and q is connected, it follows that $(q^2)^{\text{Tr}} = 0$ by Lemma 3.8. Applying (3.7) from Lemma 3.11 to q^2 with $S \triangleq \emptyset$, it follows that

$$\mathbb{E}\left[q^2(\mathbf{X})\right] = \mathbb{E}\left[q^2(\mathbf{X}) - (q^2)^{\mathrm{Tr}}(\mathbf{X})\right] = O(||q^2||).$$
(4.3)

Since the degree of each p_i is at most Δ , it follows that the degree of r_{ij} is at most $2\Delta + 1$, and hence the degree of q is at most $2\Delta + 1$. Using Lemma 3.5, it follows

$$\|q^2\| \le 2^{4\Delta+2} \|q\|^2. \tag{4.4}$$

By definition of q

$$\|q\| \le \left\|\sum_{1 \le i < j \le n} r_{ij}\right\| + \left\|\sum_{1 \le i < j \le n} r_{ij}^{\mathrm{Tr}}\right\| \le 2 \left\|\sum_{1 \le i < j \le n} r_{ij}\right\|.$$

$$(4.5)$$

Using Lemma 3.6 with sets $S_{i,j} \triangleq \{i, j\}$ for $1 \le i < j \le n$ and $\gamma = \frac{1}{2}, c = \sqrt{2}$ (any set of s edges touches at least $\sqrt{2s}$ nodes), it follows that

$$\left\| \sum_{1 \le i < j \le n} r_{ij} \right\| \le \left(\frac{1 + \Delta}{\sqrt{2}} \right)^2 \max_{i \in [n]} \|p_i\|^2 = O(1),$$
(4.6)

and (4.1) is established. Combining (4.1) and (4.5) in (4.4) yields $||q^2|| = O(1)$. Plugging the latter in (4.3)

$$\mathbb{E}\left[q^2(\mathbf{X})\right] = O(1)$$

By Chebyshev's inequality it follows that

$$\forall \epsilon' > 0, \mathbb{P}\left(\left| \sum_{1 \le i < j \le n} \left[r_{ij}(\mathbf{X}) - r_{ij}^{\mathrm{Tr}}(\mathbf{X}) \right] \right| \ge n\epsilon' \right) = \mathbb{P}\left(q^2(\mathbf{X}) \ge n^2 \epsilon'^2 \right) = \frac{O(1)}{n^2 \epsilon'^2}.$$

The claim follows then from (2.3).

In light of the Lemma above, we first show that property (4.2) extends to the sparse disorder Y. **Proposition 4.2.** For every $\epsilon > 0$ whp as $n \to \infty$ and $d \to \infty$

$$\frac{1}{n} \sum_{1 \le i < j \le n} r_{ij}^{\text{Tr}}(\mathbf{Y}) \ge \mathcal{P} - \epsilon.$$
(4.7)

Proof of Proposition 4.2. Fix $c, \beta > 0$ and recall $\phi, \psi_{\beta,c}$ from Lemma 3.3. For convenience we redefine $\psi_{\beta,c}$ as

$$\psi_{\beta,c}(z) = \phi\left(\beta\left(\frac{z}{n} - c\right)\right).$$

Consider an interpolation path from **X** to **Y** given by $\mathbf{Z}^0, ..., \mathbf{Z}^N$ where $\mathbf{Z}^0 = \mathbf{X}, \mathbf{Z}^N = \mathbf{Y}$ and $N \triangleq \binom{n}{2}$. For $t \in [N]$ we will identify t with its corresponding (enumerated) edge (ij), and thus write $\mathbf{X}_t, \mathbf{Y}_t$ instead of $\mathbf{X}_{ij}, \mathbf{Y}_{ij}$. We have

$$\mathbb{E}\left[\psi_{\beta,c}\left(\sum_{1\leq i< j\leq n} r_{ij}^{\mathrm{Tr}}(\mathbf{Y})\right)\right] - \mathbb{E}\left[\psi_{\beta,c}\left(\sum_{1\leq i< j\leq n} r_{ij}^{\mathrm{Tr}}(\mathbf{X})\right)\right]$$
$$=\sum_{\ell=0}^{N-1}\left[\psi_{\beta,c}\left(\sum_{1\leq i< j\leq n} r_{ij}^{\mathrm{Tr}}(\mathbf{Z}^{\ell+1})\right) - \psi_{\beta,c}\left(\sum_{1\leq i< j\leq n} r_{ij}^{\mathrm{Tr}}(\mathbf{Z}^{\ell})\right)\right]$$

By definition of $\mathcal{T}_{n,2}$, the monomials $\mathbf{Z}^{\boldsymbol{\alpha}}$ in $\sum_{1 \leq i < j \leq n} r_{ij}^{\mathrm{Tr}}(\mathbf{Z})$ satisfy $\max(\boldsymbol{\alpha}) \leq 2$. In particular, we can write for $\ell \leq N-1$

$$\sum_{\leq i < j \leq n} r_{ij}^{\mathrm{Tr}}(\mathbf{Z}^{\ell}) = u_{\ell}(\mathbf{Z}^{\ell}) + \mathbf{X}_{\ell} a_{\ell}(\mathbf{Z}^{\ell}) + \mathbf{X}_{\ell}^{2} b_{\ell}(\mathbf{Z}^{\ell}),$$
(4.8)

where $u_{\ell}(\mathbf{x}), a_{\ell}(\mathbf{x}), b_{\ell}(\mathbf{x})$ are LDPs in the variables $\mathbf{x} = (\mathbf{x}_{ij}, 1 \leq i < j \leq n)$, and do not depend on the variable associated with edge ℓ , i.e. the LDPs do not depend on \mathbf{x}_{ℓ} . Furthermore, since we switch the distribution of $[\mathbf{Z}^{\ell}]_{\ell}$ from \mathbf{X}_{ℓ} to \mathbf{Y}_{ℓ} to obtain $\mathbf{Z}^{\ell+1}$ (and keep everything else unchanged), it follows that

$$\sum_{1 \le i < j \le n} r_{ij}^{\mathrm{Tr}}(\mathbf{Z}^{\ell+1}) = u_{\ell}(\mathbf{Z}^{\ell}) + \mathbf{Y}_{\ell}a_{\ell}(\mathbf{Z}^{\ell}) + \mathbf{Y}_{\ell}^{2}b_{\ell}(\mathbf{Z}^{\ell}).$$
(4.9)

Two important elementary properties of $u_{\ell}, a_{\ell}, b_{\ell}$ are

$$\forall \ell \in [0, N-1], \quad u_{\ell}(\mathbf{x}), \mathbf{x}_{\ell} a_{\ell}(\mathbf{x}), \mathbf{x}_{\ell} b_{\ell}(\mathbf{x}) \text{ are connected and contain node } i.$$
(4.10)

$$\max_{\ell \in [0, N-1]} \|u_\ell\| + \|a_\ell\| + \|b_\ell\| = O(1).$$
(4.11)

Property (4.11) follows immediately from (4.1). The connectedness of $\mathbf{x}_{\ell}b_{\ell}(\mathbf{x}), \mathbf{x}_{\ell}b_{\ell}(\mathbf{x})$ in (4.10) follows from first noting that $u_{\ell}(\mathbf{x}), \mathbf{x}_{\ell}^{2}b_{\ell}(\mathbf{x}), \mathbf{x}_{\ell}^{2}b_{\ell}(\mathbf{x})$ are connected (as $\sum r_{ij}^{\mathrm{Tr}}$ is connected) and using the fact that $\mathbf{x}_{\ell}w(\mathbf{x})$ is connected if and only if $(\mathbf{x}_{\ell})^{k}w(\mathbf{x})$ is connected for all $k \geq 1$, where w is any LDP.

To simplify notations, we will write $u_{\ell}, a_{\ell}, b_{\ell}$ instead of $u_{\ell}(\mathbf{Z}^{\ell}), a_{\ell}(\mathbf{Z}^{\ell}), b_{\ell}(\mathbf{Z}^{\ell})$ (i.e. we write u_{ℓ} to denote the value of the polynomial u_{ℓ} at \mathbf{Z}_{ℓ}). Introduce for $\ell \leq N-1$

$$v_{\ell}(z) = u_{\ell} + za_{\ell} + z^2 b_{\ell}$$

$$\xi_{\beta,c}(z) = \psi_{\beta,c}(v_{\ell}(z)).$$

It follows from (4.8), (4.9) that

$$\sum_{\ell=0}^{N-1} \left[\psi_{\beta,c} \left(\sum_{1 \le i < j \le n} r_{ij}^{\mathrm{Tr}}(\mathbf{Z}^{\ell+1}) \right) - \psi_{\beta,c} \left(\sum_{1 \le i < j \le n} r_{ij}^{\mathrm{Tr}}(\mathbf{Z}^{\ell}) \right) \right] = \sum_{\ell=0}^{N-1} \left[\xi_{\beta,c}(\mathbf{Y}_{\ell}) - \xi_{\beta,c}(\mathbf{X}_{\ell}) \right].$$
(4.12)

We will drop the subscripts in $\xi_{\beta,c}, \psi_{\beta,c}$ and instead simply write ξ, ψ . Using a 3rd order Taylor expansion, there exists $\mathbf{X}_{\ell}^* \in [-|\mathbf{X}_{\ell}|, |\mathbf{X}_{\ell}|]$ and $\mathbf{Y}_{\ell}^* \in [-|\mathbf{Y}_{\ell}|, |\mathbf{Y}_{\ell}|]$ such that

$$\begin{split} \xi(\mathbf{X}_{\ell}) &= \xi(0) + \mathbf{X}_{\ell} \xi^{(1)}(0) + \frac{\mathbf{X}_{\ell}^{2}}{2} \xi^{(2)}(0) + \frac{\mathbf{X}_{\ell}^{3}}{6} \xi^{(3)}(\mathbf{X}_{\ell}^{*}), \\ \xi(\mathbf{Y}_{\ell}) &= \xi(0) + \mathbf{Y}_{\ell} \xi^{(1)}(0) + \frac{\mathbf{Y}_{\ell}^{2}}{2} \xi^{(2)}(0) + \frac{\mathbf{Y}_{\ell}^{3}}{6} \xi^{(3)}(\mathbf{Y}_{\ell}^{*}). \end{split}$$

Taking the difference of the above and the expectation yields

$$\mathbb{E}\left[\xi(\mathbf{Y}_{\ell})\right] - \mathbb{E}\left[\xi(\mathbf{X}_{\ell})\right] = \frac{\mathbb{E}\left[\mathbf{Y}_{t}^{3}\xi^{(3)}(\mathbf{Y}_{\ell}^{*})\right] - \mathbb{E}\left[\mathbf{X}_{\ell}^{3}\xi^{(3)}(\mathbf{X}_{\ell}^{*})\right]}{6}.$$
(4.13)

We set $\beta = \log n$ in the definition of ψ for the remaining parts of the proof of Proposition 4.2. The next Lemma bounds (4.13).

Lemma 4.3. We have

$$\begin{split} |\mathbb{E}[\mathbf{X}_{\ell}^{3}\xi^{(3)}(\mathbf{X}_{\ell}^{*})]| &= O((\log n)^{2}n^{-\frac{7}{2}}), \\ |\mathbb{E}[\mathbf{Y}_{\ell}^{3}\xi^{(3)}(\mathbf{Y}_{\ell}^{*})]| &= O((\log n)^{2}n^{-3}d^{-\frac{1}{2}}), \end{split}$$

Proof of Lemma 4.3. Let $M_{\psi} \triangleq \|\psi^{(2)}\|_{\infty} + \|\psi^{(3)}\|_{\infty} + \|\psi^{(4)}\|_{\infty}$. Since $\beta = \log n$, it follows from item 2 of Lemma 3.3

$$M_{\psi} \le \left(\frac{\beta}{n}\right)^2 + \left(\frac{\beta}{n}\right)^3 + \left(\frac{\beta}{n}\right)^4 \lesssim \frac{(\log n)^2}{n^2}.$$

Introduce for $\ell \leq N-1$

$$v_{\ell}: \mathbb{R} \to \mathbb{R}, \quad z \mapsto u_{\ell} + za_{\ell} + z^2 b_{\ell},$$

so that $\xi(z) = \psi(v_{\ell}(z))$. Using Lemma 3.1 and the fact that $v_{\ell}^{(k)}(z) = 0, \forall k > 2$ we obtain

$$|\xi^{(3)}(z)| \le c' M_{\psi} \sum_{\substack{k_1+2k_2=3\\k_1,k_2\ge 0}} |a_{\ell}+2zb_{\ell}|^{k_1} |b_{\ell}|^{k_2} \le c' \frac{(\log n)^2}{n^2} \sum_{0\le j_1, j_2, j_3\le 3} |z|^{j_1} |a_{\ell}|^{j_2} |b_{\ell}|^{j_3} + \frac{1}{2} |b_{\ell}|^{j_3} |b_{\ell}|^{j_4} |b_{\ell}|^{j_4} \le c' \frac{(\log n)^2}{n^2} \sum_{0\le j_1, j_2, j_3\le 3} |z|^{j_1} |a_{\ell}|^{j_2} |b_{\ell}|^{j_3} + \frac{1}{2} |b_{\ell}|^{j_4} |b_{\ell}|^{j_4} |b_{\ell}|^{j_4} \le c' \frac{(\log n)^2}{n^2} \sum_{0\le j_1, j_2, j_3\le 3} |z|^{j_1} |a_{\ell}|^{j_2} |b_{\ell}|^{j_3} + \frac{1}{2} |b_{\ell}|^{j_4} |b_{\ell}|^{j$$

where c' denotes a different explicit constant in each inequality. Combining the above and $|\mathbf{X}_{\ell}^*| \leq |\mathbf{X}_{\ell}|$, we can bound $|\xi^{(3)}(\mathbf{X}_{\ell}^*)|$ by

$$\begin{aligned} |\xi^{(3)}(\mathbf{X}_{\ell}^{*})| &\leq c' \frac{(\log n)^{2}}{n^{2}} \sum_{0 \leq j_{1}, j_{2}, j_{3} \leq 3} |\mathbf{X}_{\ell}|^{j_{1}} |a_{\ell}|^{j_{2}} |b_{\ell}|^{j_{3}} \\ &\leq c' \frac{(\log n)^{2}}{n^{2}} \sum_{0 \leq j_{1}, j_{2}, j_{3} \leq 3} \left[(\mathbf{X}_{\ell})^{2j_{1}} (a_{\ell})^{2j_{2}} (b_{\ell})^{2j_{3}} + 1 \right], \end{aligned}$$

where we used the arithmetic geometric inequality in the last line. Therefore

$$\mathbb{E}\left[|\mathbf{X}_{\ell}|^{3}|\xi^{(3)}(\mathbf{X}_{\ell}^{*})|\right] \leq c' \frac{(\log n)^{2}}{n^{2}} \sum_{0 \leq j_{1}, j_{2}, j_{3} \leq 3} \left(\mathbb{E}\left[|\mathbf{X}_{\ell}|^{3+2j_{1}}(a_{\ell})^{2j_{2}}(b_{\ell})^{2j_{3}}\right] + \mathbb{E}\left[|\mathbf{X}_{\ell}|^{3}\right]\right)$$

Note that $\mathbf{X}_{\ell}^{3+2j_1}(a_{\ell})^{2j_2}(b_{\ell})^{2j_3}$ and \mathbf{X}_{ℓ}^3 are connected LDPs containing edge ℓ as the product of connected LDPs sharing edge ℓ . It follows from applying (3.12) in Lemma 3.12 with S the set of the endpoints of edge ℓ

$$\mathbb{E}\left[|\mathbf{X}_{\ell}|^{3}\right] = O(n^{-\frac{3}{2}}),$$
$$\mathbb{E}\left[|\mathbf{X}_{\ell}|^{3+2j_{1}}(a_{\ell})^{2j_{2}}(b_{\ell})^{2j_{3}}\right] = O(n^{-(2+\frac{3+2j_{1}}{2}-2)}) = O(n^{-\frac{3}{2}-j_{1}}) = O(n^{-\frac{3}{2}}),$$

where we used $j_1 \ge 0$. It follows readily,

$$\mathbb{E}[|\mathbf{X}_{\ell}|^{3}|\xi^{(3)}(\mathbf{X}_{\ell}^{*})|] = O((\log n)^{2}n^{-\frac{7}{2}}).$$
(4.14)

We bound $\mathbb{E}[|\mathbf{Y}_{\ell}|^3|\xi^{(3)}(\mathbf{Y}_{\ell}^*)|]|$ identically: the only difference lays in the application of Lemma 3.12, which yields for the disorder matrix \mathbf{Y} ,

$$\mathbb{E}\left[|\mathbf{Y}_{\ell}|^{3}\right] \leq O(n^{-1}d^{-\frac{1}{2}}),$$
$$\mathbb{E}\left[|\mathbf{Y}_{\ell}|^{3+2j_{1}}(a_{\ell})^{2j_{2}}(b_{\ell})^{2j_{3}}\right] \leq O(n^{-1}d^{-\frac{3+2j_{1}}{2}+1}) = O(n^{-1}d^{-\frac{1}{2}-j_{1}}) = O(n^{-1}d^{-\frac{1}{2}})$$

It follows that

$$\mathbb{E}[|\mathbf{Y}_{\ell}|^{3}|\xi^{(3)}(\mathbf{Y}_{\ell}^{*})|] = O((\log n)^{2}n^{-3}d^{-\frac{1}{2}}).$$
(4.15)

which ends the proof of Lemma 4.3.

Using Lemma 4.3 in (4.13), it follows that

$$|\mathbb{E}[\xi(\mathbf{Y}_{\ell})] - \mathbb{E}[\xi(\mathbf{X}_{\ell})]| = O((\log n)^2 n^{-3} d^{-\frac{1}{2}}).$$

Plugging the above in (4.12) and summing over $\ell \leq N - 1 \sim n^2/2$ yields

$$\left|\sum_{\ell=0}^{N-1} \mathbb{E}\left[\psi_{\beta,c}\left(\sum_{1\leq i< j\leq n} r_{ij}^{\mathrm{Tr}}(\mathbf{Z}^{\ell+1})\right)\right] - \mathbb{E}\left[\psi_{\beta,c}\left(\sum_{1\leq i< j\leq n} r_{ij}^{\mathrm{Tr}}(\mathbf{Z}^{\ell})\right)\right]\right| \leq O((\log n)^2 n^{-1} d^{-\frac{1}{2}}),$$

which implies

$$\left| \mathbb{E} \left[\psi_{\beta,c} \left(\sum_{1 \le i < j \le n} r_{ij}^{\mathrm{Tr}}(\mathbf{Y}) \right) \right] - \mathbb{E} \left[\psi_{\beta,c} \left(\sum_{1 \le i < j \le n} r_{ij}^{\mathrm{Tr}}(\mathbf{X}) \right) \right] \right| \le O((\log n)^2 n^{-1} d^{-\frac{1}{2}}).$$
(4.16)

We now show $\mathbb{P}\left(\frac{1}{n}\sum_{1\leq i< j\leq n}r_{ij}^{\mathrm{Tr}}(\mathbf{Y})\geq \mathcal{P}-\epsilon\right)=1-o(1)$. To ease notations, introduce

$$h: \mathbf{Z} \mapsto \sum_{1 \le i < j \le n} r_{ij}^{\mathrm{Tr}}(\mathbf{Z}).$$

Let $\epsilon' = \frac{\epsilon}{4}$ and set $c = \mathcal{P} - \frac{\epsilon}{2}$. Since $\psi_{\beta,c}$ is a nonnegative increasing function, we have

$$\mathbb{E}\left[\psi_{\beta,c}(h(\mathbf{X}))\right] \geq \mathbb{E}\left[\psi_{\beta,c}(h(\mathbf{X}))\mathbf{1}_{h(\mathbf{X})\geq n(\mathcal{P}-\epsilon')}\right] \\
\geq \psi_{\beta,c}\left(n(\mathcal{P}-\epsilon')\right)\mathbb{P}\left(h(\mathbf{X})\geq n(\mathcal{P}-\epsilon')\right) \\
= \phi\left(\frac{\beta\epsilon}{4}\right)\mathbb{P}\left(h(\mathbf{X})\geq n(\mathcal{P}-\epsilon')\right) \\
= (1-o(1))\mathbb{P}\left(h(\mathbf{X})\geq n(\mathcal{P}-\epsilon')\right) \\
= 1-o(1),$$
(4.17)

where (4.17) follows from $\beta \epsilon = \omega(1)$ due to $\beta = \log n$ and (4.18) follows from (4.2) in Lemma 4.1. Since $\psi_{\beta,c}$ is bounded above by 1, then (4.18) implies

$$\mathbb{E}\left[\psi_{\beta,c}(h(\mathbf{X}))\right] = 1 - o(1).$$

Combining the above with (4.16), it follows that

$$\mathbb{E}\left[\psi_{\beta,c}(h(\mathbf{Y}))\right] = 1 - o(1). \tag{4.19}$$

Since $\psi_{\beta,c} \leq 1$ and $\psi_{\beta,c}$ is increasing, we have

$$\mathbb{E}\left[\psi_{\beta,c}(h(\mathbf{Y}))\right] = \mathbb{E}\left[\psi_{\beta,c}(h(\mathbf{Y}))\mathbf{1}_{h(\mathbf{Y})< n(\mathcal{P}-\epsilon)}\right] + \mathbb{E}\left[\psi_{\beta,c}(h(\mathbf{Y}))\mathbf{1}_{h(\mathbf{Y})\geq n(\mathcal{P}-\epsilon)}\right] \\
\leq \psi_{\beta,c}\left(n(\mathcal{P}-\epsilon)\right) + \mathbb{P}\left(h(\mathbf{Y})\geq n(\mathcal{P}-\epsilon)\right) \\
= \phi\left(-\frac{\beta\epsilon}{2}\right) + \mathbb{P}\left(h(\mathbf{Y})\geq n(\mathcal{P}-\epsilon)\right) \\
= o(1) + \mathbb{P}\left(h(\mathbf{Y})\geq n(\mathcal{P}-\epsilon)\right),$$
(4.20)

where the last line follows from $-\beta \epsilon = -\omega(1)$. Using (4.20) and (4.19), it follows that

$$\mathbb{P}(h(\mathbf{Y}) \ge n(\mathcal{P} - \epsilon)) = 1 - o(1), \tag{4.21}$$

namely

$$\mathbb{P}\left(\frac{1}{n}\sum_{1\leq i< j\leq n}r_{ij}^{\mathrm{Tr}}(\mathbf{Y})\geq \mathcal{P}-\epsilon\right) = 1-o(1),\tag{4.22}$$

which concludes the proof of Proposition 4.2.

To conclude the proof of (2.5), let $q \triangleq \sum (r_{ij} - r_{ij}^{\text{Tr}})$ as in the proof of Lemma 4.1 and recall that we showed $(q^2)^{\text{Tr}} = 0$ and $||q^2|| = O(1)$. Applying (3.8) from Lemma 3.11 to q^2 with $S \triangleq \emptyset$, it follows that

$$\left|\mathbb{E}\left[q^{2}(\mathbf{Y})\right]\right| = \left|\mathbb{E}\left[q^{2}(\mathbf{Y}) - (q^{2})^{\mathrm{Tr}}(\mathbf{Y})\right]\right| = O(n^{2}d^{-1}).$$
(4.23)

Let $\epsilon > 0$. Using the above, it follows by Chebyshev's inequality

$$\mathbb{P}\left(\left|\sum_{1\leq i< j\leq n} \left[r_{ij}(\mathbf{Y}) - r_{ij}^{\mathrm{Tr}}(\mathbf{Y})\right]\right| \geq n\epsilon/2\right) = \mathbb{P}(q^2(\mathbf{Y}) \geq n^2\epsilon^2/4) \leq \frac{O(1)}{d\epsilon^2} = o(1)$$

Moreover, we have from Proposition 4.2 that $\frac{1}{n} \sum_{1 \le i < j \le n} r_{ij}^{\text{Tr}}(\mathbf{Y}) \ge \mathcal{P} - \epsilon/2$ whp. Therefore

$$\mathbb{P}\left(\frac{1}{n}\sum_{1\leq i< j\leq n}r_{ij}(\mathbf{Y})\geq \mathcal{P}-\epsilon\right)=1-o(1),$$

which completes the proof of (2.5) since $\frac{1}{2n}p(\mathbf{Y})^{\top}\mathbf{Y}p(\mathbf{Y}) = \frac{1}{n}\sum_{1 \le i < j \le n} r_{ij}(\mathbf{Y}).$

4.2 Proof of (2.6)

Proof. We use an interpolation argument from the Gaussian disorder **X** to the sparse disorder **Y**. For $c, \beta_1, \beta_2 > 0$ recall $\phi, \psi_{\beta_1,c}, \pi_{\beta_2}$ from Lemma 3.3. For convenience, we rewrite $\psi_{\beta_1,c}$ as

$$\psi_{\beta_1,c}(z) = \phi\left(-\beta_1\left(\frac{z}{\sqrt{n}} - c\right)\right)$$

Throughout this proof, we will assume the following

$$\beta_1 = \sqrt{\log \log d},$$

$$\beta_2 = \log \log d,$$
(4.24)

and we will also assume $\beta_2 \geq 2$. Denote by f the following function on $\mathbf{x} \in \mathbb{R}^{n \times n}$

$$f(\mathbf{x}) = \psi_{\beta_1,c} \left(\sqrt{\sum_{i=1}^n \pi_{\beta_2}(p_i(\mathbf{x}))^2} \right).$$

Our main technical result is the following Lemma.

Lemma 4.4. The following holds

$$|\mathbb{E}[f(\mathbf{Y})] - \mathbb{E}[f(\mathbf{X})]| = o(1). \tag{4.25}$$

Proof of Lemma 4.4. As in the proof of (2.5) we consider the interpolation path from **X** to **Y** given by $\mathbf{Z}^0, ..., \mathbf{Z}^N$ where $\mathbf{Z}^0 = \mathbf{X}, \mathbf{Z}^N = \mathbf{Y}$ and $N \triangleq \binom{n}{2}$. For $t \in [N]$ we will identify t with its corresponding (enumerated) edge (ij), and thus write $\mathbf{X}_t, \mathbf{Y}_t$ instead of $\mathbf{X}_{ij}, \mathbf{Y}_{ij}$. We have

$$\mathbb{E}[f(\mathbf{Y})] - \mathbb{E}[f(\mathbf{X})] = \sum_{\ell=0}^{N-1} \mathbb{E}[f(\mathbf{Z}^{\ell+1})] - \mathbb{E}[f(\mathbf{Z}^{\ell})].$$

Since $\forall i \in [n], p_i \in \mathcal{T}_{n,2}$, it follows that the monomials $\mathbf{Z}^{\boldsymbol{\alpha}}$ in $p_i(\mathbf{Z})$ satisfy $\max(\boldsymbol{\alpha}) \leq 2$. In particular, we can write for $\ell \leq N-1, i \in [n]$

$$p_i^{\mathrm{Tr}}(\mathbf{Z}^\ell) = u_{i,\ell}(\mathbf{Z}^\ell) + \mathbf{X}_\ell a_{i,\ell}(\mathbf{Z}^\ell) + \mathbf{X}_\ell^2 b_{i,\ell}(\mathbf{Z}^\ell), \qquad (4.26)$$

where $u_{i,\ell}(\mathbf{x}), a_{i,\ell}(\mathbf{x}), b_{i,\ell}(\mathbf{x})$ are LDPs in the variables $\mathbf{x} = (\mathbf{x}_{ij}, 1 \leq i < j \leq n)$ which do not depend on \mathbf{x}_{ℓ} . Furthermore, since we switch the distribution of $[\mathbf{Z}^{\ell}]_{\ell}$ from \mathbf{X}_{ℓ} to \mathbf{Y}_{ℓ} to obtain $\mathbf{Z}^{\ell+1}$ (and keep everything else unchanged), it follows that

$$p_i^{\rm Tr}(\mathbf{Z}^{\ell+1}) = u_{i,\ell}(\mathbf{Z}^{\ell}) + \mathbf{Y}_{\ell} a_{i,\ell}(\mathbf{Z}^{\ell}) + \mathbf{Y}_{\ell}^2 b_{i,\ell}(\mathbf{Z}^{\ell}).$$
(4.27)

Two important elementary properties of $u_{i,\ell}, a_{i,\ell}, b_{i,\ell}$ are

$$\forall i \in [n], \ell \in [0, N-1], \quad u_{i,\ell}(\mathbf{x}), \mathbf{x}_{\ell} a_{i,\ell}(\mathbf{x}), \mathbf{x}_{\ell} b_{i,\ell}(\mathbf{x}) \text{ are connected and contain node } i.$$

$$\max_{i \in [n], \ell \in [0, N-1]} \|u_{i,\ell}\| + \|a_{i,\ell}\| + \|b_{i,\ell}\| = O(1).$$

$$(4.29)$$

The above properties follow similarly to (4.10), (4.11). We introduce the following univariate functions

$$v_{i,\ell}(z) = u_{i,\ell} + za_{i,\ell} + z^2 b_{i,\ell},$$
$$g_{\ell}(z) = \psi_{\beta_1,c} \left(\sqrt{\sum_{i=1}^n \pi_{\beta_2}(v_{i,\ell}(z))^2} \right).$$

In particular, note that for $\ell \leq N-1$

$$f(\mathbf{Z}^{\ell+1}) - f(\mathbf{Z}^{\ell}) = g_{\ell}(\mathbf{Y}_{\ell}) - g_{\ell}(\mathbf{X}_{\ell}).$$

$$(4.30)$$

Using a 3rd order Taylor expansion, there exists $\mathbf{X}_{\ell}^* \in [-|\mathbf{X}_{\ell}|, |\mathbf{X}_{\ell}|]$ and $\mathbf{Y}_{\ell}^* \in [-|\mathbf{Y}_{\ell}|, |\mathbf{Y}_{\ell}|]$ such that

$$g_{\ell}(\mathbf{X}_{\ell}) = g_{\ell}(0) + \mathbf{X}_{\ell}g_{\ell}^{(1)}(0) + \frac{\mathbf{X}_{\ell}^{2}}{2}g_{\ell}^{(2)}(0) + \frac{\mathbf{X}_{\ell}^{3}}{6}g_{\ell}^{(3)}(\mathbf{X}_{\ell}^{*}),$$

$$g_{\ell}(\mathbf{Y}_{\ell}) = g_{\ell}(0) + \mathbf{Y}_{\ell}g_{\ell}^{(1)}(0) + \frac{\mathbf{Y}_{\ell}^{2}}{2}g_{\ell}^{(2)}(0) + \frac{\mathbf{Y}_{\ell}^{3}}{6}g_{\ell}^{(3)}(\mathbf{Y}_{\ell}^{*}).$$

Taking the difference of the above and the expectation yields

$$\mathbb{E}[g_{\ell}(\mathbf{Y}_{\ell})] - \mathbb{E}[g_{\ell}(\mathbf{X}_{\ell})] = \frac{\mathbb{E}\left[\mathbf{Y}_{\ell}^{3}g_{\ell}^{(3)}(\mathbf{Y}_{\ell}^{*})\right] - \mathbb{E}\left[\mathbf{X}_{\ell}^{3}g_{\ell}^{(3)}(\mathbf{X}_{\ell}^{*})\right]}{6}.$$
(4.31)

To simplify the representation of the derivatives of g_{ℓ} , we introduce the following two additional functions

$$k(z) = \sum_{i=1}^{n} \pi_{\beta_2} (v_{i,\ell}(z))^2,$$

$$h(z) = \sqrt{k(z)},$$

so that $g = \psi(h) = \psi(\sqrt{k})$. We first note the following two elementary properties of ψ, π from Lemma 3.3

$$\begin{aligned} \forall j \in [4], \quad \|\psi^{(j)}\|_{\infty} &\leq \left(\frac{\beta_1}{\sqrt{n}}\right)^j, \\ \forall z \in \mathbb{R}, \quad |\pi(z)| &\geq \frac{1}{2\beta_2 e^{\beta_2}}. \end{aligned}$$

Recalling $\beta_1 = \sqrt{\log \log d}, \beta_2 = \log \log d$, it follows that

$$\forall j \in [4], \quad \|\psi^{(j)}\|_{\infty} = \tilde{O}(n^{-\frac{j}{2}}), \tag{4.32}$$

$$\forall z \in \mathbb{R}, \quad |\pi(z)| \ge \frac{1}{2(\log \log d)(\log d)} = \tilde{\Omega}(1).$$
(4.33)

Note that (4.33) implies

$$\forall z \in \mathbb{R}, \quad \sqrt{k(z)} \ge \tilde{\Omega}(\sqrt{n}). \tag{4.34}$$

Using the above bound and Corollary 3.2, we have for all $z \in \mathbb{R}$, and $t_1, t_2, t_3 \in \mathbb{Z}_{\geq 0}$ satisfying $t_1 + 2t_2 + 3t_3 = 3$

$$\prod_{j=1}^{3} |h^{(j)}(z)|^{t_{j}} \lesssim \sum_{\substack{s_{1},m_{1},s_{2},m_{2} \in \mathbb{Z}_{\geq 0} \\ s_{1}m_{1}+s_{2}m_{2}=3}} \frac{|k^{(s_{1})}(z)|^{m_{1}}|k^{(s_{2})}(z)|^{m_{2}}}{(k(z))^{m_{1}+m_{2}-\frac{t_{1}+t_{2}+t_{3}}{2}}} \\
\lesssim \sum_{\substack{s_{1},m_{1},s_{2},m_{2} \in \mathbb{Z}_{\geq 0} \\ s_{1}m_{1}+s_{2}m_{2}=3}} |k^{(s_{1})}(z)|^{m_{1}}|k^{(s_{2})}(z)|^{m_{2}}n^{-m_{1}-m_{2}+\frac{t_{1}+t_{2}+t_{3}}{2}}.$$
(4.35)

We next bound $\left| \mathbb{E}[\mathbf{X}_{\ell}^{3}g_{\ell}^{(3)}(\mathbf{X}_{\ell}^{*})] \right|$. Using Lemma 3.1 and (4.32), it follows

$$\left| \mathbb{E}[\mathbf{X}_{\ell}^{3}g_{\ell}^{(3)}(\mathbf{X}_{\ell}^{*})] \right| \lesssim \sum_{\substack{t_{1}+2t_{2}+3t_{3}=3\\t_{1},t_{2},t_{3}\geq 0}} n^{-\frac{t_{1}+t_{2}+t_{3}}{2}} \mathbb{E}\left[|\mathbf{X}_{\ell}|^{3} \left(\prod_{j=1}^{3} |h^{(j)}(\mathbf{X}_{\ell}^{*})|^{t_{j}} \right) \right].$$

Combining with (4.35), it follows

$$\left| \mathbb{E}[\mathbf{X}_{\ell}^{3} g_{\ell}^{(3)}(\mathbf{X}_{\ell}^{*})] \right| \lesssim \sum_{\substack{s_{1}, m_{1}, s_{2}, m_{2} \in \mathbb{Z}_{\geq 0} \\ s_{1}m_{1}+s_{2}m_{2}=3}} n^{-m_{1}-m_{2}} \mathbb{E}\left[|\mathbf{X}_{\ell}|^{3} |k^{(s_{1})}(\mathbf{X}_{\ell}^{*})|^{m_{1}} |k^{(s_{2})}(\mathbf{X}_{\ell}^{*})|^{m_{2}} \right]$$

Combining the latter bound with (3.15) from Lemma 3.13 readily implies

$$\left| \mathbb{E}[\mathbf{X}_{\ell}^{3}g_{\ell}^{(3)}(\mathbf{X}_{\ell}^{*})] \right| \leq \tilde{O}(n^{-\frac{5}{2}}).$$

$$(4.36)$$

Following the same arguments above, we have with (3.16) from Lemma (3.13)

$$\left| \mathbb{E}[\mathbf{Y}_{\ell}^{3}g_{\ell}^{(3)}(\mathbf{Y}_{\ell}^{*})] \right| \leq \tilde{O}(n^{-2}d^{-\frac{1}{2}}).$$

$$(4.37)$$

Combining (4.36), (4.37) in (4.31), we have

$$|\mathbb{E}[g_{\ell}(\mathbf{Y}_{\ell})] - \mathbb{E}[g_{\ell}(\mathbf{X}_{\ell})]| \le \tilde{O}\left(\frac{1}{n^{\frac{5}{2}}} + \frac{1}{n^2\sqrt{d}}\right) = \tilde{O}(n^{-2}d^{-\frac{1}{2}}).$$

Summing the above over $\ell \leq N-1 \sim n^2/2$ and using (4.30) yields

$$|\mathbb{E}[f(\mathbf{Y})] - \mathbb{E}[f(\mathbf{X})]| \le \tilde{O}\left(\frac{1}{\sqrt{d}}\right) = o(1).$$
(4.38)

Which ends the proof of Lemma 4.4.

To end the proof of (2.6), we show the following Lemma.

Lemma 4.5. The following holds

$$\forall \epsilon > 0, \quad \mathbb{P}\left(d(p(\mathbf{Y}), [-1, 1]^n) \le \epsilon \sqrt{n}\right) = 1 - o(1).$$

Proof of Lemma 4.5. Fix $\epsilon > 0$. Set $c = \frac{\epsilon}{2}$ in $\psi_{\beta_1,c}$ and recall $\beta_1 = \sqrt{\log \log d}$. It follows

$$\mathbb{E}\left[\psi_{\beta_1,c}(d(p(\mathbf{X}), [-1,1]^n))\right] \ge \psi_{\beta_1,c}(\epsilon\sqrt{n}/4)\mathbb{P}\left(d(p(\mathbf{X}), [-1,1]^n) \le \epsilon\sqrt{n}/4\right)$$
(4.39)

$$=\phi(\beta_1 \epsilon/2)(1-o(1)) \tag{4.40}$$

$$= 1 - o(1), \tag{4.41}$$

where (4.39) follows from the decreasing property of $\psi_{\beta_1,c} \ge 0$, (4.40) from (2.4), and (4.41) from $\beta_1 \epsilon = \omega(1)$. Furthermore, since $\psi_{\beta_1,c} \le 1$ it follows

$$\mathbb{E}\left[\psi_{\beta_1,c}(d(p(\mathbf{X}), [-1,1]^n))\right] = 1 - o(1).$$
(4.42)

Using Lemma 3.4, we have

$$|\psi_{\beta_1,c}\left(d(p(\mathbf{X}), [-1,1]^n)\right) - f(\mathbf{X})| \le \frac{2\beta_1}{\beta_2} = o(1), \tag{4.43}$$

where the last equality follows from recalling $\beta_1 = \sqrt{\log \log d}$, $\beta_2 = \log \log d$. Similarly

$$|\psi_{\beta_1,c}\left(d(p(\mathbf{Y}), [-1,1]^n)\right) - f(\mathbf{Y})| = o(1).$$
(4.44)

Combining (4.42) and (4.43) it follows that

$$\mathbb{E}[f(\mathbf{X})] = 1 - o(1),$$

Combining the above and Lemma 4.4, it follows that

$$\mathbb{E}[f(\mathbf{Y})] = 1 - o(1)$$

Combining the above and (4.44), it follows that

$$\mathbb{E}[\psi_{\beta_1,c}\left(d(p(\mathbf{Y}), [-1,1]^n)\right)] = 1 - o(1).$$
(4.45)

Then

$$\mathbb{E}\left[\psi_{\beta_{1},c}(d(p(\mathbf{Y}),[-1,1]^{n}))\right] = \mathbb{E}\left[\psi_{\beta_{1},c}(d(p(\mathbf{Y}),[-1,1]^{n}))\left(\mathbf{1}_{d(p(\mathbf{Y}),[-1,1]^{n}) > \epsilon\sqrt{n}} + \mathbf{1}_{d(p(\mathbf{Y}),[-1,1]^{n}) \le \epsilon\sqrt{n}}\right)\right]$$

$$\leq \psi_{\beta_{1},c}(\epsilon\sqrt{n}) + \mathbb{P}\left(d(p(\mathbf{Y}),[-1,1]^{n}) \le \epsilon\sqrt{n}\right)$$

$$= \phi(-\beta_{1}\epsilon/2) + \mathbb{P}\left(d(p(\mathbf{Y}),[-1,1]^{n}) \le \epsilon\sqrt{n}\right)$$

$$= o(1) + \mathbb{P}\left(d(p(\mathbf{Y}),[-1,1]^{n}) \le \epsilon\sqrt{n}\right),$$
(4.47)

where (4.46) follows from the fact that $\psi_{\beta_{1,c}}$ is decreasing and $\|\psi_{\beta_{1,c}}\|_{\infty} = 1$, while (4.47) follows from $\beta_{1}\epsilon = \omega(1)$. Combining (4.47) and (4.45) yields

$$\mathbb{P}\left(d(p(\mathbf{Y}), [-1,1]^n) \le \epsilon \sqrt{n}\right) = 1 - o(1),\tag{4.48}$$

which concludes the proof of Lemma 4.5.

This concludes the proof of (2.6).

4.3 **Proof of (2.7)**

In this subsection, we extends a simple rounding procedure from [Mon19] to $p(\mathbf{Y})$ in order to obtain a point $\sigma(p(\mathbf{Y}), \mathbf{Y}) \in \{+1, -1\}^n$.

Proof. Recall that $\sigma(p(\mathbf{Y}), \mathbf{Y})$ is computed by first projecting the terms $p_i(\mathbf{Y})$ that lay outside [-1, 1] into [-1, 1], then sequentially setting the terms $p_i(\mathbf{Y})$ that lay in (-1, 1) to ± 1 while increasing the Hamiltonian objective value. Note in particular that $\sigma(p(\mathbf{Y}), \mathbf{Y})$ can be computed in polynomial time complexity from $p(\mathbf{Y})$.

Fix $\epsilon' > 0$ and let \mathbf{z} be the projection of $p(\mathbf{Y})$ onto the hypercube $[-1,1]^n$. From (2.6) it follows that $\|p(\mathbf{Y}) - \mathbf{z}\|_2 \le \epsilon' \sqrt{n}$ whp. We then have

$$\begin{aligned} \frac{1}{2n} |p(\mathbf{Y})^{\top} \mathbf{Y} p(\mathbf{Y}) - \mathbf{z}^{\top} \mathbf{Y} \mathbf{z}| &\leq \frac{1}{2n} |p(\mathbf{Y}) - \mathbf{z})^{\top} \mathbf{Y} (p(\mathbf{Y}) - \mathbf{z})| + \frac{1}{n} |\mathbf{z}^{\top} \mathbf{Y} (p(\mathbf{Y}) - \mathbf{z})| \\ &\leq \frac{1}{2n} ||\mathbf{Y}||_{\mathrm{op}} ||p(\mathbf{Y}) - \mathbf{z}||_{2}^{2} + \frac{1}{n} ||\mathbf{Y}||_{\mathrm{op}} ||p(\mathbf{Y}) - \mathbf{z}||_{2} ||\mathbf{z}||_{2} \\ &\leq \frac{c\epsilon'^{2}}{2} + c\epsilon', \end{aligned}$$

where we used $||z||_2 = \sqrt{n}$ and applied Lemma 3.10 in the last line, and c is a universal constant. Since the diagonal terms of \mathbf{Y} are null, it follows that the restriction of $\mathbf{z} \mapsto \mathbf{z}^{\top} \mathbf{Y} \mathbf{z}$ to any single entry \mathbf{z}_{ij} , $1 \le i < j \le$ is linear. In particular, if $\mathbf{z}_{ij} \in (-1, 1)$, we can increase the objective value $\mathbf{z}^{\top} \mathbf{Y} \mathbf{z}$ by moving \mathbf{z}_{ij} either to -1 or 1. Denote the new value of \mathbf{z}_{ij} by $\sigma(p(\mathbf{Y}), \mathbf{Y})$ and apply this rounding scheme iteratively for all $1 \le i < j \le n$. Since each rounding step improves the objective, it follows that

$$\begin{aligned} \frac{1}{2n}\sigma(p(\mathbf{Y}),\mathbf{Y})^{\top}\mathbf{Y}\sigma(p(\mathbf{Y}),\mathbf{Y}) &\geq \frac{1}{2n}\mathbf{z}^{\top}\mathbf{Y}\mathbf{z} \\ &\geq \frac{1}{2n}p(\mathbf{Y})^{\top}\mathbf{Y}p(\mathbf{Y}) - \frac{c\epsilon'^2}{2} - c\epsilon' \\ &\geq \mathcal{P} - \epsilon' - \frac{c\epsilon'^2}{2} - c\epsilon', \end{aligned}$$

where we used (2.5) in the last line. Taking ϵ' such that $\epsilon' + \frac{c\epsilon'^2}{2} + c\epsilon' \leq \epsilon$, yields the result of (2.7).

5 IAMP Representation and Proof of Theorem **2.2**

5.1 Approximation of IAMP by Polynomials

It is well known in the literature that AMP algorithms can be approximated by Low-Degree polynomials modulo some regularity assumptions on the denoiser functions f^t ([Mon22], [IS23]). However, this is mostly an informal statement and technical work must be done on a per-problem basis to rigorously justify the approximation.

We begin by restating the state evolution property satisfied by the AMP iterates.

Proposition 5.1 (Proposition 2.1 in [Mon21]). Assume the denoisers f^t are pseudo-Lipschitz of order m. Let $(U_j)_{j\geq 1}$ be a centered Gaussian process independent of U_0 with covariance $\mathbf{Q} = (\mathbf{Q}_{kj})_{k,j\geq 1}$ given defined by (2.9). Assume $\mathbf{Q}_{\leq k} \triangleq (\mathbf{Q}_{ij})_{1\leq i,j\leq k}$ is strictly positive definite for all $k \leq T$. Then for any $k, \ell \in \mathbb{Z}_{\geq 0}$, and any function $\psi : \mathbb{R}^{k+1} \to \mathbb{R}$ that is pseudo-Lipschitz of order ℓ , we have

$$\operatorname{p-lim}_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \psi(\mathbf{u}_i^0, ..., \mathbf{u}_i^k) = \mathbb{E}[\psi(U_0, ..., U_k)].$$

Next we restate a Lemma from [IS23] showing that general AMP can be well approximated by LDP under certain regularity assumptions stated below.

Lemma 5.2 (Lemma B.4 in [IS23]). Fix $\eta > 0$ and $T \in \mathbb{Z}_{>0}$. Suppose that

- $\{f^t : \mathbb{R}^{t+1} \to \mathbb{R}\}$ is a sequence of L-Lipschitz denoiser function which produce IAMP iterates $\mathbf{u}^0, ..., \mathbf{u}^T$.
- For each $i \in [n]$, $\frac{\partial f^t}{\partial \mathbf{u}_i}$ is either pseudo-Lipschitz or an indicator.
- The covariance matrix $\mathbf{Q}_{\leq t}$ for $t \in [T]$ satisfies $\mathbf{Q}_{\leq t} \succeq \mathbf{I}_t$ and $\|\mathbf{Q}_{\leq t}\|_{\infty} \leq 2$.

Then, there exists a sequence of polynomial denoisers $\{q^t : \mathbb{R}^{t+1} \to \mathbb{R}\}$ producing IAMP iterates $\hat{\mathbf{u}}^0, ..., \hat{\mathbf{u}}^T$ such that

$$\operatorname{p-lim}_{n \to \infty} \frac{1}{\sqrt{n}} \| \mathbf{u}^t - \hat{\mathbf{u}}^t \|_2 \le \eta,$$

for all $t \leq T$. Furthermore, there exists constants $\Delta = \Delta(\eta, T, L), c = c(\eta, T, L)$, such that the polynomials q^t have degree bounded by Δ , and coefficients bounded in absolute value by c.

As a Corollary of the above Lemma, we obtain a polynomial representation result for the IAMP algorithm described in [Mon19].

Corollary 5.3. Suppose $\mathbf{X} \stackrel{\mathsf{d}}{=} \operatorname{GOE}(n)$. Let $\eta > 0$ and $T \in \mathbb{Z}_{\geq 0}$. Consider the IAMP algorithm in [Mon19], and let its denoiser functions be $f^t : \mathbb{R}^{t+1} \to \mathbb{R}$, and the IAMP iterates be $\mathbf{u}^0, ..., \mathbf{u}^T$. Then, there exist $\Delta = \Delta(\eta, T), c = c(\eta, T)$, and a sequence of polynomial denoisers $\{q^t : \mathbb{R}^{t+1} \to \mathbb{R}\}$ satisfying $\operatorname{Deg}(q^t) \leq \Delta, \|q^t\| \leq c$, and producing AMP iterates $\hat{\mathbf{u}}^0, ..., \hat{\mathbf{u}}^T$ such that the following holds

$$\operatorname{p-lim}_{n \to \infty} \frac{\|\mathbf{u}^t - \hat{\mathbf{u}}^t\|_2}{\sqrt{n}} \le \eta,$$

for all $t \leq T$.

Proof. The proof is an application of Lemma 5.2 to the IAMP iterates in [Mon19]. The conditions required to apply the Lemma are: (1) f^t are Lipschitz functions. (2) f^t are weakly differentiable with pseudo-Lipschitz or indicator partial derivatives . (3) the covariance matrix $\mathbf{Q}_{\leq t}$ satisfies $\mathbf{Q}_{\leq t} \succeq \mathbf{I}_t$, $\|\mathbf{Q}_{\leq t}\|_{\infty} \leq 2$. The first condition is satisfied by the IAMP described in equations (9), (10) in [Mon19]: f^t are compositions/sums/products of bounded Lipschitz continuous functions. The second condition is only used to show state evolution for $b_{t,j}$ in the proof of Lemma 5.2 in [IS23] (see Claim B.7 in [IS23]), however, state evolution for $b_{t,j}$ is directly shown in Lemma II.3 in [Mon19]. Finally, the covariance matrix $\mathbf{Q}_{\leq t}$ is shown to be diagonal in Lemma II.2 in [Mon19] with entries upper/lower bounded by positive constants. We then claim that we can rescale the denoiser functions to satisfy condition (3) and detail this argument next. Given positive coefficients $a_0, ..., a_{T+1} \in \mathbb{R}_{>0}$, introduce for $t \leq T$

$$\tilde{f}^t : \mathbb{R}^{t+1} \to \mathbb{R}, \quad (x_0, ..., x_t) \mapsto a_{t+1} f^t (x_0/a_0, ..., x_t/a_t)$$

Let $\tilde{\mathbf{u}}^0 = a_0 \mathbf{u}^0$ and denote by $\tilde{\mathbf{u}}^t$ be the iterates of the IAMP with adjusted denoisers \tilde{f}^t and initialization $\tilde{\mathbf{u}}^0$. Furthermore denote by $\tilde{b}_{t,j}$ the adjusted $b_{t,j}$ coefficients for \tilde{f}^t . We first show by induction that $\forall t \leq T, \tilde{\mathbf{u}}^t = a_t \mathbf{u}^t$. If t = 0 the result follows by definition of $\tilde{\mathbf{u}}^0$. Assume the result for all $t' \in [t]$ for some t < T, then

$$\begin{split} \tilde{\mathbf{u}}^{t+1} &= \mathbf{X} \tilde{f}^{t}(\tilde{\mathbf{u}}^{0}, ..., \tilde{\mathbf{u}}^{t}) - \sum_{j=1}^{t} \tilde{b}_{t,j} \tilde{f}^{j-1}(\tilde{\mathbf{u}}^{0}, ..., \tilde{\mathbf{u}}^{j-1}) \\ &= a_{t+1} \mathbf{X} f^{t}(\tilde{\mathbf{u}}^{0}/a_{0}, ..., \tilde{\mathbf{u}}^{t}/a_{t}) - \sum_{j=1}^{t} \tilde{b}_{t,j} a_{j} f^{j-1}(\tilde{\mathbf{u}}^{0}/a_{0}, ..., \tilde{\mathbf{u}}^{j-1}/a_{j-1}) \\ &= a_{t+1} \left[\mathbf{X} f^{t}(\mathbf{u}^{0}, ..., \mathbf{u}^{t}) - \sum_{j=1}^{t} \frac{\tilde{b}_{t,j} a_{j}}{a_{t+1}} f^{j-1}(\mathbf{u}^{0}, ..., \mathbf{u}^{j-1}) \right]. \end{split}$$

For $j \in [t]$ we have

$$\begin{split} \tilde{b}_{t,j} &= \frac{1}{n} \sum_{i=1}^{n} \frac{\partial \tilde{f}^{t}}{\partial \tilde{\mathbf{u}}_{i}^{j}} (\tilde{\mathbf{u}}_{i}^{0}, ..., \tilde{\mathbf{u}}_{i}^{t}) \\ &= \frac{1}{n} \sum_{i=1}^{n} \frac{a_{t+1}}{a_{j}} \frac{\partial f^{t}}{\partial \mathbf{u}_{i}^{j}} (\tilde{\mathbf{u}}_{i}^{0}/a_{0}, ..., \tilde{\mathbf{u}}_{i}^{t}/a_{t}) \\ &= \frac{a_{t+1}}{a_{j}} b_{t,j}. \end{split}$$

Henceforth

$$\tilde{\mathbf{u}}^{t+1} = a_{t+1} \left[\mathbf{X} f^t(\mathbf{u}^0, ..., \mathbf{u}^t) - \sum_{j=1}^t b_{t,j} f^{j-1}(\mathbf{u}^0, ..., \mathbf{u}^{j-1}) \right]$$

= $a_{t+1} \mathbf{u}^{t+1}$,

which concludes the induction proof. Let $\mathbf{Q}, \tilde{\mathbf{Q}}$ be the covariance matrices associated with the IAMP obtained from denoisers f^t and \tilde{f}^t respectively, and given recursively by

$$\begin{aligned} \mathbf{Q}_{k+1,j+1} &= \mathbb{E}[f^k(U_0,...,U_k)f^j(U_0,...,U_t)],\\ \tilde{\mathbf{Q}}_{k+1,j+1} &= \mathbb{E}[\tilde{f}^k(\tilde{U}_0,...,\tilde{U}_k)\tilde{f}^j(\tilde{U}_0,...,\tilde{U}_t)], \end{aligned}$$

where $(U_j)_{j\geq 1}$ $((\tilde{U}_j)_{j\geq 1}$ resp.) is a centered Gaussian process independent of U_0 $(\tilde{U}_0$ resp.). As shown in Lemma II.2 in [Mon19], the matrix \mathbf{Q} is diagonal. Furthermore, the proof extends verbatim for $\tilde{\mathbf{Q}}$. We next show that $\tilde{\mathbf{Q}}_{t+1,t+1} = (a_{t+1})^2 \mathbf{Q}_{t+1,t+1}$ for all t < T. It is straighforward to see that we can take $\tilde{U}_t = a_t U_t$ for $t \leq T$, since $\tilde{U}_0 \stackrel{d}{=} a_0 U_0$ from our initialization $\tilde{\mathbf{u}}^0 = a_0 \mathbf{u}^0$. Therefore $(\tilde{U}_j)_{j\geq 1}$ is a centered Gaussian process independent of \tilde{U}_0 with diagonal covariance matrix, and we have for $0 \leq t < T$,

$$\begin{split} \tilde{\mathbf{Q}}_{t+1,t+1} &= \mathbb{E}[\tilde{f}^t(\tilde{U}_0, ..., \tilde{U}_t)\tilde{f}^t(\tilde{U}_0, ..., \tilde{U}_t)] \\ &= (a_{t+1})^2 \mathbb{E}[f^t(U_0, ..., U_t)f^t(U_0, ..., U_t)] \\ &= (a_{t+1})^2 \mathbf{Q}_{t+1,t+1}. \end{split}$$

Henceforth $\tilde{\mathbf{Q}}_{\leq T} = \text{Diag}((a_1)^2 \mathbf{Q}_{1,1}, ..., (a_T)^2 \mathbf{Q}_{T,T})$. It suffices then to pick $a_1, ..., a_T$ so that $\tilde{\mathbf{Q}}_{\leq T}$ is the identity matrix, and thus satisfies the two requirements $\tilde{\mathbf{Q}}_{\leq t} \succeq \mathbf{I}_t$ and $\|\tilde{\mathbf{Q}}_{\leq t}\|_{\infty} \leq 2$ for $t \leq T$, in order to apply Lemma 5.2. This concludes the proof.

The candidate for near optimum associated with the AMP with denoisers q^t from Lemma 5.2 is given similarly to (2.10) by

$$\hat{\mathbf{v}} = \sqrt{\delta} \sum_{k=1}^{\lfloor \bar{q}/\delta \rfloor} q^k(\hat{\mathbf{u}}^0, ..., \hat{\mathbf{u}}^k).$$

The next proposition shows that $\hat{\mathbf{v}}$ is close in ℓ_2 distance to \mathbf{v} given by (2.10), i.e. the candidate for near optimum associated with the IAMP algorithm in [Mon19].

Proposition 5.4. Fix $\theta, \delta > 0$ and $T \in \mathbb{Z}_{\geq 0}$. In the setting of Lemma 5.2, the polynomials q^t can be chosen to satisfy the following

$$\operatorname{p-lim}_{n \to \infty} \, \frac{\|\mathbf{v} - \hat{\mathbf{v}}\|_2}{\sqrt{n}} \le \theta$$

Proof. Let $\eta \in (0, 1)$. By Lemma 5.2, we can chose the polynomials q^t so that $\underset{n \to \infty}{\text{p-lim}} \frac{\|\mathbf{u}^t - \hat{\mathbf{u}}^t\|_2}{\sqrt{n}} \leq \eta, \forall t \leq T$. We have by the triangle inequality

$$\frac{\|\mathbf{v} - \hat{\mathbf{v}}\|_2}{\sqrt{n}} \le \sqrt{\delta} \sum_{k=1}^{\lfloor \bar{q}/\delta \rfloor} \frac{\|f^k(\mathbf{u}^0, ..., \mathbf{u}^k) - q^k(\hat{\mathbf{u}}^0, ..., \hat{\mathbf{u}}^k)\|_2}{\sqrt{n}}$$

therefore it suffices to show that each of the summands above is bounded by $O(\eta)$ whp and then take η small enough. Let $k \in [\lfloor \bar{q}/\delta \rfloor]$. We have

$$\|f^{k}(\mathbf{u}^{0},...,\mathbf{u}^{k}) - q^{k}(\hat{\mathbf{u}}^{0},...,\hat{\mathbf{u}}^{k})\|_{2} \leq \underbrace{\|f^{k}(\mathbf{u}^{0},...,\mathbf{u}^{k}) - f^{k}(\hat{\mathbf{u}}^{0},...,\hat{\mathbf{u}}^{k})\|_{2}}_{A} + \underbrace{\|f^{k}(\hat{\mathbf{u}}^{0},...,\hat{\mathbf{u}}^{k}) - q^{k}(\hat{\mathbf{u}}^{0},...,\hat{\mathbf{u}}^{k})\|_{2}}_{B}.$$

Let $\mathbf{U}, \hat{\mathbf{U}}$ be matrices in $\mathbb{R}^{n \times (k+1)}$ with columns $(\mathbf{u}^0, ..., \mathbf{u}^k)$, and $(\hat{\mathbf{u}}^0, ..., \hat{\mathbf{u}}^k)$ respectively. Since f^k is Lipschitz with constant Lipschitz coefficient L, it follows that

$$A \le L \|\mathbf{U} - \hat{\mathbf{U}}\|_{\text{fro}} = L \sqrt{\sum_{t=0}^{k} \|\mathbf{u}^t - \hat{\mathbf{u}}^t\|_2^2}.$$

From Lemma 5.2, we have $\|\mathbf{u}^t - \hat{\mathbf{u}}^t\|_2 \leq \eta \sqrt{n}, \forall t \in [T]$ whp as $n \to \infty$, it follows that $A \leq L\eta \sqrt{nk}$ whp. We now bound *B*. We have by Proposition 5.1 (using the fact that pseudo-Lipschitz functions are closed under sums/products)

$$p-\lim_{n \to \infty} \frac{B^2}{n} = p-\lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^n |f^k(\hat{\mathbf{u}}_i^0, ..., \hat{\mathbf{u}}_i^k) - q^k(\hat{\mathbf{u}}_i^0, ..., \hat{\mathbf{u}}_i^k)|^2$$

$$= \mathbb{E} \left[(f^k(\hat{U}^0, ..., \hat{U}^k) - q^k(\hat{U}^0, ..., \hat{U}^k))^2 \right]$$

$$< q_2(k)^2,$$

where g_2 is an increasing function introduced in Claim B.6 in [IS23] and satisfies $g_2(T) \leq c\eta^2$, where c is a constant depending on T, L. Hence

$$\operatorname{p-lim}_{n \to \infty} \frac{B^2}{n} \le c^2 \eta^4,$$

which readily implies $B \leq O(\eta\sqrt{n})$ whp. Taking η small enough, we conclude that $\Pr_{n\to\infty} \frac{\|\mathbf{v}-\hat{\mathbf{v}}\|_2}{\sqrt{n}} \leq \theta$, which ends the proof.

5.2 Approximation of AMP with Connected Tree-based Low-Degree Polynomials

We now state a general result on the representation of AMP iterations with polynomial denoisers. Following the statement of Theorem 2.2, we claim that we can assume that the initialization point \mathbf{u}^0 has entries bounded by a constant M > 0 independent of n. This assumption appears restrictive for IAMP, since [Mon19] requires a random initialization $\mathbf{u}^0 \sim \delta \mathcal{N}(0, \mathbf{I}_n)$ where δ is a fixed parameter driving the optimality gap of the IAMP output. Nonetheless this condition is implicitly satisfied by the IAMP algorithm as it includes a truncation step on the iterates, i.e. the denoisers depend on \mathbf{u}^0 only through $[\mathbf{u}^0]_M$ with $[u]_M \triangleq \max(-M, \min(u, M))$ for some fixed large enough M > 0. Indeed, let $\bar{\mathbf{u}}^k = [\mathbf{u}^k]_M, \forall k \in [T]$. Observe that for all $t \leq T$ the denoisers f^t given in (9), (10) in [Mon19] only depend on the iterates \mathbf{u}^k through $\bar{\mathbf{u}}^k$, i.e.

$$\forall t \leq T, f^t(\mathbf{u}^0, ..., \mathbf{u}^t) = f^t(\bar{\mathbf{u}}^0, ..., \bar{\mathbf{u}}^t),$$

thus switching the initialization **u** with $\bar{\mathbf{u}}^0$ has no effect on the IAMP iterations, and furthermore preserves the covariance matrix **Q** since it holds for $k, j \ge 0$

$$\mathbb{E}_{U_0,\dots,U_{\max(j,k)}}[f^k(U_0,\dots,U_k)f^j(U_0,\dots,U_j)] = \mathbb{E}_{U_0,\dots,U_{\max(j,k)}}[f^k([U_0]_M,\dots,U_k)f^j([U_0]_M,\dots,U_j)]$$
$$= \mathbb{E}_{[U_0]_M,\dots,U_{\max(j,k)}}[f^k([U_0]_M,\dots,U_k)f^j([U_0]_M,\dots,U_j)].$$

Adding the truncation step thus has the effect of adding a secondary limit $M \to \infty$ after the limits on n, d on all results mentioned in this paper.

Corollary 5.3 implies that the IAMP iterates can be approximated by LDP. In order to complete our second main result, Theorem 2.2 we need to show that LDP induced by denoisers q satisfy additional assumptions, namely, the connectivity, norm and degree bounds. This our next goal.

In the setting of Lemma 5.2 and Proposition 5.4, we modify the AMP iterations with polynomial denoisers q^t by replacing the coefficients $b_{t,j}$ by their respective limits in probability denoted $\bar{b}_{t,j} \triangleq \underset{n \to \infty}{\text{p-lim}} b_{t,j}$. i.e. consider the following iterations

$$\tilde{\mathbf{u}}^{t+1} = \mathbf{X}q^{t}(\tilde{\mathbf{u}}^{0}, ..., \tilde{\mathbf{u}}^{t}) - \sum_{j=1}^{t} \bar{b}_{t,j}q^{j-1}(\tilde{\mathbf{u}}^{0}, ..., \tilde{\mathbf{u}}^{j-1}),$$

$$b_{t,j} = \frac{1}{n} \sum_{i=1}^{n} \frac{\partial f^{t}}{\partial \hat{\mathbf{u}}_{i}^{j}} (\hat{\mathbf{u}}_{i}^{0}, ..., \hat{\mathbf{u}}_{i}^{t}),$$
(5.1)

where $\hat{\mathbf{u}}_{i}^{0}, ..., \hat{\mathbf{u}}_{i}^{T}$ are the iterates of the (standard) AMP with denoisers q^{t} . We note that the existence of the limits $\bar{b}_{t,j}$ is justified by applying Proposition 5.1 to the partial derivatives of the polynomial denoisers q^{t} . We show next that the above modification leads to effectively the same algorithm, and that the dynamics of $\tilde{\mathbf{u}}^{t}$ are roughly the same as $\hat{\mathbf{u}}^{t}$.

Proposition 5.5. Fix $\delta > 0, T \in \mathbb{Z}_{\geq 0}$. In the setting of Lemma 5.2, let $\hat{\mathbf{u}}^0, ..., \hat{\mathbf{u}}^T$ be the AMP iterates obtained with polynomial denoisers $q^t, t \in [T]$, and let $\tilde{\mathbf{u}}^0, ..., \tilde{\mathbf{u}}^T$ be the AMP iterates obtained by replacing $b_{t,j}$ with $\bar{b}_{t,j} \triangleq \underset{n \to \infty}{\text{p-lim}} b_{t,j}$ as shown in (5.1). Let the near-optimum candidates be given by

$$\hat{\mathbf{v}} = \sqrt{\delta} \sum_{k=1}^{\lfloor \bar{q}/\delta \rfloor} q^k(\hat{\mathbf{u}}^0, ..., \hat{\mathbf{u}}^k)$$
(5.2)

$$\tilde{\mathbf{v}} = \sqrt{\delta} \sum_{k=1}^{\lfloor \bar{q}/\delta \rfloor} q^k(\tilde{\mathbf{u}}^0, ..., \tilde{\mathbf{u}}^k),$$
(5.3)

Then

$$\operatorname{p-lim}_{n \to \infty} \frac{\|\hat{\mathbf{v}} - \tilde{\mathbf{v}}\|_2}{\sqrt{n}} = 0.$$

The proof of Proposition 5.5 is given in Section 5.4. Using Proposition 5.5, we next establish the key representation result we require.

Proposition 5.6. Let $\delta > 0, T \in \mathbb{Z}_{\geq 0}$ and consider the AMP iteration (5.1) with polynomial denoisers $q^t : \mathbb{R}^{t+1} \to \mathbb{R}$ with constant degrees Δ^t , and coefficients bounded in absolute value by a constant. Let $\tilde{\mathbf{u}}^0, ..., \tilde{\mathbf{u}}^T$ be the iterations produced where $\tilde{\mathbf{u}}^0$ will be assumed to satisfy $\|\tilde{\mathbf{u}}^0\|_{\infty} = O(1)$ whp. Then there exist LDPs w_i^t , and h_i such that for all $t \in [0, T], i \in [n]$

- 1. $\tilde{\mathbf{u}}_i^t = w_i^t(\mathbf{X}), \text{ and } \tilde{\mathbf{v}}_i = h_i(\mathbf{X}).$
- 2. $\max_i \|w_i^t\| = O(1)$, and $\max_i \|h_i\| = O(1)$.
- 3. w_i^t, h_i are a connected LDPs containing node *i*.

Proof. We first show the required properties for iterates $\tilde{\mathbf{u}}^t$, the claim for $\tilde{\mathbf{v}}$ will follow similarly. Note that by Proposition 5.1 and the fact that $\|\hat{\mathbf{u}}^0\|_{\infty} = O(1)$, it follows that $|\bar{b}_{t,j}| = O(1)$ for all $t \in [0,T]$ and $j \in [t]$, as the limit $\bar{b}_{t,j}$ is a constant. The proof proceeds by induction on $t \leq T$. If t = 0 then w_i^0 are constant polynomials and the graph $G_{\mathbf{0}_{\mathbb{Z}_{\geq 0}^{n \times n}}}$ is empty, thus it satisfies all claims in the proposition by definition. Assume the result for all $t' \in [0,t]$ where t < T is fixed. By symmetry, it suffices to check that it holds at t+1 for i = 1. We have

$$\tilde{\mathbf{u}}_{1}^{t+1} = \underbrace{\sum_{\ell=1}^{n} \mathbf{X}_{1\ell} q^{t}(\tilde{\mathbf{u}}_{\ell}^{0}, ..., \tilde{\mathbf{u}}_{\ell}^{t})}_{A_{1,t}} - \underbrace{\sum_{j=1}^{t} \bar{b}_{t,j} q^{j-1}(\tilde{\mathbf{u}}_{1}^{0}, ..., \tilde{\mathbf{u}}_{1}^{j-1})}_{O_{1,t}}.$$

We first show that $O_{1,t}$ is a connected LDP in **X** with constant coefficients. Each term $q^{j-1}(\tilde{\mathbf{u}}_1^0,...,\tilde{\mathbf{u}}_1^{j-1})$ is a polynomial in $\tilde{\mathbf{u}}_1^0,...,\tilde{\mathbf{u}}_1^t$ with constant degree and constant coefficients. Furthermore, by the inductive hypothesis, each of $\tilde{\mathbf{u}}_1^0,...,\tilde{\mathbf{u}}_1^t$ is a connected LDP in **X** with O(1) coefficients containing node 1. Therefore, it follows that $q^{j-1}(\tilde{\mathbf{u}}_1^0,...,\tilde{\mathbf{u}}_1^{j-1})$ is a LDP in **X** with O(1) coefficients. Moreover, the latter LDP is connected and contains node 1, since it is the product of connected monomials sharing node 1. As $\bar{b}_{t,j} = O(1)$, it follows that each summand in $O_{1,t}$ is a connected LDP with O(1) coefficients containing node 1. As $O_{1,t}$ is a finite sum of such LDPs, it readily follows that $O_{1,t}$ is a connected LDP in **X** with O(1) coefficients.

We now deal with the term $A_{1,t}$. Let $r_{\ell}(\mathbf{X}) = \mathbf{X}_{1\ell}q^t(\tilde{\mathbf{u}}_{\ell}^0, ..., \tilde{\mathbf{u}}_{\ell}^t)$ for $\ell \geq 2$ (we recall here that $\mathbf{X}_{\ell\ell} = 0, \forall \ell$). Note that $r_{\ell}, \ell \geq 2$ are LDPs and $\forall \ell \in [2, n], \{1, \ell\} \in r_{\ell}$. Furthermore, since each of $\tilde{\mathbf{u}}_{\ell}^0, ..., \tilde{\mathbf{u}}_{\ell}^t$ is a connected LDP in \mathbf{X} with O(1) coefficients containing node ℓ , it follows that $q^t(\tilde{\mathbf{u}}_{\ell}^0, ..., \tilde{\mathbf{u}}_{\ell}^t)$ is a connected LDP in \mathbf{X} with O(1) coefficients in **X** containing node ℓ . Therefore $r_{\ell}(\mathbf{X})$ is a connected LDP in **X** with O(1) coefficients containing nodes $\{1, \ell\}$ as the product of $\mathbf{X}_{1\ell}$ (an edge-tree with endpoints $1, \ell$) and $q^t(\tilde{\mathbf{u}}_{\ell}^0, ..., \tilde{\mathbf{u}}_{\ell}^t)$ (LDPs containing node 1). It follows readily that $A_{1,t}$ is a connected LDP in **X** containing node 1. It remains to show $||A_{1,t}|| = ||\sum_{2 \leq \ell \leq n} r_{\ell}|| = O(1)$. Using Lemma 3.6 with sets $S_{\ell} \triangleq \{1, \ell\}$ for $2 \leq \ell \leq n$ and $\gamma = 1, c = 1$, it follows that

$$||A_{1,t}|| \le (1 + \Delta^t) \max_{2 \le \ell \le n} ||r_\ell|| = O(1).$$

Therefore $\tilde{\mathbf{u}}_1^{t+1}$ is a connected LDP in **X** containing node 1 with O(1) coefficients. Finally, note that any bounds on the coefficients of the LDP $\tilde{\mathbf{u}}_1^{t+1}$ extend generically to the LDPs $\tilde{\mathbf{u}}_i^{t+1}$ for all $i \in [n]$, so that $\max_i \|\tilde{\mathbf{u}}_i^{t+1}\| = O(1)$. This completes the induction and ends the proof for $\tilde{\mathbf{u}}^t$.

We now show similarly the representation result on $\tilde{\mathbf{v}}$. By symmetry, it suffices to verify the result for i = 1. We have $\tilde{\mathbf{v}}_1 = \sqrt{\delta} \sum_{k=1}^{\lfloor \bar{q}/\delta \rfloor} q^k(\tilde{\mathbf{u}}_1^0, ..., \tilde{\mathbf{u}}_1^k)$. As shown above, we have that $\tilde{\mathbf{u}}_1^t, t \in [k]$ are connected LDPs in \mathbf{X} with constant coefficients and contain node 1. Since q^k are polynomials with constant coefficients, it follows readily that each of the summands $q^k(\tilde{\mathbf{u}}_1^0, ..., \tilde{\mathbf{u}}_1^k)$ is a LDP in \mathbf{X} with constant coefficients. Furthermore, since all the LDPs $\tilde{\mathbf{u}}_1^t, t \in [k]$ are connected and share node 1, it follows that all monomials in $q^k(\tilde{\mathbf{u}}_1^0, ..., \tilde{\mathbf{u}}_1^k)$ are connected and contain node 1. Therefore, the claims in items (1) and (3) in the proposition for h_1 hold. Finally, since $\tilde{\mathbf{v}}_1$ is a finite sum of LDPs with constant coefficients, it follows that $||h_1|| = O(1)$, and moreover $\max_i ||h_i|| = O(1)$ as the bound on $||h_1||$ extends to all h_i , which shows the claim of item (2) and concludes the proof.

5.3 Proof of Theorem 2.2

As a corollary of Proposition 5.6 and Proposition 5.5, for every fixed $\epsilon > 0$ we can construct an AMP algorithm with $T = T(\epsilon)$ iterations, and polynomial denoisers q^t satisfying the assumptions of Lemma 5.2, and we can encode the near optimum candidate as $h(\mathbf{X}) \triangleq (h_1(\mathbf{X}), ..., h_n(\mathbf{X}))$ where h_i satisfy the assumptions of Proposition 5.6. We next provide justification of the quality of the output of $p \triangleq h^{\mathrm{Tr}}$.

Proposition 5.7. Let $\mathbf{X} \stackrel{\mathsf{d}}{=} \operatorname{GOE}(n)$. Let $\epsilon > 0$. There exists $\delta = \delta(\epsilon) > 0$ and $T = T(\epsilon) \in \mathbb{Z}_{\geq 0}$ such that if $\tilde{\mathbf{v}} = h(\mathbf{X})$ in the setting of Proposition 5.6, and $p \triangleq h^{\operatorname{Tr}}$, then the following holds why as $n \to \infty$

$$\frac{1}{2n} |\mathbf{v}^{\top} \mathbf{X} \mathbf{v} - p(\mathbf{X})^{\top} \mathbf{X} p(\mathbf{X})| \le \epsilon,$$

$$\frac{d(p(\mathbf{X}), [-1, 1]^n)}{\sqrt{n}} \le \epsilon,$$
(5.4)

where \mathbf{v} is the output candidate of IAMP given in (2.10).

Proof. Let $\epsilon' > 0$. Let $\theta > 0$ as in Proposition 5.4, and $\hat{\mathbf{v}}$ be given in (5.2) Then, where $n \to \infty$

$$\frac{d(p(\mathbf{X}), [-1, 1]^n)}{\sqrt{n}} \leq \frac{d(p(\mathbf{X}), \tilde{\mathbf{v}})}{\sqrt{n}} + \frac{d(\tilde{\mathbf{v}}, \hat{\mathbf{v}})}{\sqrt{n}} + \frac{d(\hat{\mathbf{v}}, \mathbf{v})}{\sqrt{n}} + \frac{d(\mathbf{v}, [-1, 1]^n)}{\sqrt{n}} \\
\leq \frac{d(p(\mathbf{X}), \tilde{\mathbf{v}})}{\sqrt{n}} + \kappa_n + \theta + \epsilon',$$
(5.5)

where κ_n is a random variable s.t p-lim $\kappa_n = 0$, and the last line follows from applying Proposition 5.5, Proposition 5.4 and (2.12) respectively. We have

$$\mathbb{E}\left[d(p(\mathbf{X}), \tilde{\mathbf{v}})^2\right] = \sum_{i=1}^n \mathbb{E}\left[(h_i^{\mathrm{Tr}}(\mathbf{X}) - h_i(\mathbf{X}))^2\right].$$

Let $g_i \triangleq h_i^{\mathrm{Tr}} - h_i$. Since h_i is connected and $i \in h_i$ it follows that g_i is connected and $i \in g_i$. Since $g_i^{\mathrm{Tr}} = 0$ and g_i is connected, we have by Lemma 3.8 that $[(g_i)^2]^{\mathrm{Tr}} = 0$. Using Lemma 3.11 on g_i with $S \triangleq \{i\}$ yields

$$\mathbb{E}[g_i(\mathbf{X})^2] \le \frac{c \|h_i\|}{n},$$

where c is a constant. Therefore $\mathbb{E}\left[d(p(\mathbf{X}), \tilde{\mathbf{v}})^2\right] \leq c \sum_{i=1}^n \|h_i\|/n$. Since $\max_i \|h_i\| = O(1)$, it follows that $\mathbb{E}\left[d(p(\mathbf{X}), \tilde{\mathbf{v}})^2\right] = O(1)$. Using Chebyshev's inequality, we have for all $\eta > 0$

$$\mathbb{P}(d(p(\mathbf{X}), \tilde{\mathbf{v}}) \ge \eta \sqrt{n}) = \mathbb{P}(d(p(\mathbf{X}), \tilde{\mathbf{v}})^2 \ge \eta^2 n) \le \frac{O(1)}{\eta^2 n}.$$

Combining the above with (5.5), it holds whp

$$\frac{d(p(\mathbf{X}), [-1, 1]^n)}{\sqrt{n}} \le \eta + \kappa_n + \theta + \epsilon'.$$
(5.6)

Let $\mathbf{u} = p(\mathbf{X}) - \mathbf{v}$, so that $\|\mathbf{u}\|_2 / \sqrt{n} \le \eta + \kappa_n + \theta$ whp as shown above. We have

$$\begin{aligned} |p(\mathbf{X})^{\top} \mathbf{X} p(\mathbf{X}) - \mathbf{v}^{\top} \mathbf{X} \mathbf{v}| &= |2\mathbf{u}^{\top} \mathbf{X} \mathbf{v} + \mathbf{u}^{\top} \mathbf{X} \mathbf{u}| \\ &\leq \left(2 \|\mathbf{u}\|_2 \|\mathbf{v}\|_2 + \|\mathbf{u}\|_2^2\right) \|\mathbf{X}\|_{\text{op}}. \end{aligned}$$

From Proposition (2.12) it follows that $\|\mathbf{v}\|_2 \leq (1+\epsilon')\sqrt{n}$. Therefore $\frac{2\|\mathbf{u}\|_2\|\mathbf{v}\|_2+\|\mathbf{u}\|_2^2}{n} \leq 2(\eta+\kappa_n+\theta)(1+\epsilon')+(\eta+\kappa_n+\theta)^2$ whp. From standard random Gaussian matrix theory $\|\mathbf{X}\|_{\text{op}} = O(1)$ whp, therefore there exists a constant C > 0 such that

$$|p(\mathbf{X})^{\top}\mathbf{X}p(\mathbf{X}) - \mathbf{v}^{\top}\mathbf{X}\mathbf{v}| \le Cn\left(2(\eta + \kappa_n + \theta)(1 + \epsilon') + (\eta + \kappa_n + \theta)^2\right).$$
(5.7)

The claim of the proposition follows readily by taking η, θ, ϵ' small enough combined with $\underset{n \to \infty}{\text{p-lim}} \kappa_n = 0$ in (5.6) and (5.7).

This concludes the proof of Theorem 2.2.

5.4 Proof of Proposition 5.5

We first show the following Lemma on projection on conditional distribution of Gaussian matrices.

Lemma 5.8. Let $\mathbf{X} \stackrel{\mathsf{d}}{=} \operatorname{GOE}(n)$. Let $\mathbf{A} \in \mathbb{R}^{d \times n}$ and $\mathbf{D} \in \mathbb{R}^{d \times n}$. Suppose $d \leq n$ and $\operatorname{Rank}(\mathbf{A}) = d$. Let $\mathbf{X}' \stackrel{\mathsf{d}}{=} \operatorname{GOE}(n)$ be an independent copy of \mathbf{X} . Then, the distribution of \mathbf{X} conditional on $\mathbf{A}\mathbf{X} = \mathbf{D}$ is equal to the distribution of $\mathbf{A}^{\top}(\mathbf{A}\mathbf{A}^{\top})^{-1}\mathbf{D} + \operatorname{Proj}_{\{\mathbf{z} \in \mathbb{R}^n | \mathbf{A}\mathbf{z} = 0\}}(\mathbf{X}')$, where $\operatorname{Proj}_{\mathcal{V}}(\mathbf{Y})$ denotes the matrix obtained by projecting each column of \mathbf{Y} onto the vector space \mathcal{V} .

Proof. Let $\mathbf{B} \in \mathbb{R}^{(n-d) \times n}$ be a matrix with orthonormal rows, such that the rows of \mathbf{B} are also orthogonal to the rows of \mathbf{A} . In particular $[\mathbf{A}^{\top} | \mathbf{B}^{\top}]$ is invertible. Conditionally on $\mathbf{A}\mathbf{X} = \mathbf{D}$, we have

$$\begin{aligned} \mathbf{X} &= \mathbf{A}^{\top} (\mathbf{A} \mathbf{A}^{\top})^{-1} \mathbf{A} \mathbf{X} + \mathbf{B}^{\top} \mathbf{B} \mathbf{X} \\ &= \mathbf{A}^{\top} (\mathbf{A} \mathbf{A}^{\top})^{-1} \mathbf{D} + \mathbf{B}^{\top} \mathbf{B} \mathbf{X}, \end{aligned}$$

where we used $\mathbf{A}^{\top}(\mathbf{A}\mathbf{A}^{\top})^{-1}\mathbf{A} + \mathbf{B}^{\top}\mathbf{B} = \mathbf{I}_n$. Let \mathcal{V} be the vector space spanned by the rows of \mathbf{B} . Note that $\mathbf{B}^{\top}\mathbf{B}\mathbf{X} = \operatorname{Proj}_{\mathcal{V}}(\mathbf{X})$, and by Cochran's Theorem, it holds that conditionally on $\mathbf{A}\mathbf{X} = \mathbf{D}$, $\operatorname{Proj}_{\mathcal{V}}(\mathbf{X}) \stackrel{d}{=} \operatorname{Proj}_{\mathcal{V}}(\mathbf{X}')$. This concludes the proof.

Proof of Proposition 5.5. Given a sequence of random variables $X_i, i \ge 1$ and a constant C, we write p-limsup $X_n < C$ if $\lim_{n\to\infty} \mathbb{P}(X_n \ge C) = 0$. We first show the following Proposition.

Proposition 5.9. Let $\mathbf{X} \stackrel{\mathsf{d}}{=} \operatorname{GOE}(n)$. Let $q^t : \mathbb{R}^{t+1} \to \mathbb{R}, t = 0, .., T$ be multivariate polynomials with constant coefficients and degrees $\Delta_t = O(1)$. Let $\mathbf{u}^0 \in \mathbb{R}^n$ be an initialization independent of \mathbf{X} satisfying $\|\mathbf{u}^0\|_{\infty} \leq O(1)$ whp and $\forall \ell \in \mathbb{Z}_{\geq 0}, \hat{p}_{\mathbf{u}} \xrightarrow{W_{\ell}} p_{U_0}$ (where p_{U_0} is any distribution on \mathbb{R} with finite moments of all

orders, $\hat{p}_{\mathbf{u}}$ is the empirical distribution on $\{\mathbf{u}_1, ..., \mathbf{u}_n\}$, and the limit is in Wasserstein distance). Consider the iterates $\mathbf{u}^t, \tilde{\mathbf{u}}^t$ for $t \in [T]$ given recursively by

$$\mathbf{u}^{t+1} = \mathbf{X}q^{t}(\mathbf{u}^{0}, ..., \mathbf{u}^{t}) - \sum_{j=1}^{t} b_{t,j}q^{j-1}(\mathbf{u}^{0}, ..., \mathbf{u}^{j-1}),$$
$$\tilde{\mathbf{u}}^{t+1} = \mathbf{X}q^{t}(\tilde{\mathbf{u}}^{0}, ..., \tilde{\mathbf{u}}^{t}) - \sum_{j=1}^{t} \bar{b}_{t,j}q^{j-1}(\tilde{\mathbf{u}}^{0}, ..., \tilde{\mathbf{u}}^{j-1}).$$

where t < T, $\tilde{\mathbf{u}}^0 = \mathbf{u}^0$ and

$$b_{t,j} = \frac{1}{n} \sum_{i=1}^{n} \frac{\partial q^{t}}{\partial \mathbf{u}_{i}^{j}} (\mathbf{u}_{i}^{0}, ..., \mathbf{u}_{i}^{t}),$$

$$\bar{b}_{t,j} = \operatorname{p-lim}_{n \to \infty} b_{t,j},$$

where the existence of the limit points $\bar{b}_{t,j}$ is justified by Proposition 5.1. Let $(U_j)_{j\geq 1}$ be a centered Gaussian process independent of U_0 as defined in Proposition 5.1, and let \mathbf{Q} be its covariance matrix. Suppose

$$\exists c_1, c_2 > 0, \forall t \in [T], \mathbf{Q}_{\leq t} \succeq c_1 \mathbf{I}_t, and \|\mathbf{Q}\|_{\infty} \leq c_2.$$

$$(5.8)$$

Then, the following properties hold for all $t \in [0, T]$

• $\mathcal{A}_1(t)$: For all $\ell \in \mathbb{Z}_{>0}$, there exists constants $c_{t,\ell}$ such that

$$\operatorname{p-limsup}_{n \to \infty} \frac{1}{n} \| \mathbf{u}^t \|_{\ell}^{\ell} \le c_{t,\ell}, \quad \operatorname{p-limsup}_{n \to \infty} \frac{1}{n} \| \tilde{\mathbf{u}}^t \|_{\ell}^{\ell} \le c_{t,\ell}.$$

• $\mathcal{A}_2(t)$:

p-limsup
$$\frac{1}{\sqrt{n}} \| \mathbf{u}^t - \tilde{\mathbf{u}}^t \|_2 = 0$$

• $\mathcal{A}_3(t)$: For all $m \in \mathbb{Z}_{\geq 0}$ and any function $\psi : \mathbb{R}^{t+1} \mapsto \mathbb{R} \in \mathrm{PL}(m)$

$$\underset{n \to \infty}{\text{p-lim}} \left| \frac{1}{n} \sum_{i=1}^{n} \psi(\tilde{\mathbf{u}}_{i}^{0}, ..., \tilde{\mathbf{u}}_{i}^{t}) - \frac{1}{n} \sum_{i=1}^{n} \psi(\mathbf{u}_{i}^{0}, ..., \mathbf{u}_{i}^{t}) \right| = 0.$$

Proof. The proof is essentially identical to the proof of Proposition 2.1 in [Mon21] with minor adjustments. We provide a full proof here for completeness. Using the same notations as in [Mon21], we denote by $\mathbf{q}^t, \tilde{\mathbf{q}}^t$ the vectors $q^t(\mathbf{u}^0, ..., \mathbf{u}^t), q^t(\tilde{\mathbf{u}}^0, ..., \tilde{\mathbf{u}}^t)$ respectively. Let $\mathbf{F}_t, \tilde{\mathbf{F}}_t$ be the matrices in $\mathbb{R}^{n \times (t+1)}$ with columns $\mathbf{q}^0, ..., \mathbf{q}^t$, and $\tilde{\mathbf{q}}^0, ..., \tilde{\mathbf{q}}^t$ respectively. Let $\mathbf{P}_t, \tilde{\mathbf{P}}_t$ be the projection matrices on the column span of $\mathbf{F}_t, \tilde{\mathbf{F}}_t$ respectively, and let $\mathbf{P}_t^{\perp} = \mathbf{I}_n - \mathbf{P}_t$ and $\tilde{\mathbf{P}}_t^{\perp} = \mathbf{I}_n - \tilde{\mathbf{P}}_t$. Furthermore, let \mathbf{Q} be the covariance matrix associated with the AMP iterates \mathbf{u}^t . Applying Proposition 5.1 to the terms $\mathbf{q}^{j^{\top}}\mathbf{q}^i$, we have

$$\forall t \in [T], \text{ p-lim}_{n \to \infty} \frac{1}{n} \mathbf{F}_{t-1}^{\top} \mathbf{F}_{t-1} = \mathbf{Q}_{\leq t}.$$
(5.9)

We prove the three properties inductively on t. Claims $\mathcal{A}_2(0), \mathcal{A}_3(0)$ hold trivially as $\mathbf{u}^0 = \tilde{\mathbf{u}}^0$. Furthermore, $\mathcal{A}_{1}(0) \text{ holds as a direct application of Proposition 5.1, namely } \underset{n \to \infty}{\text{p-lim}} \frac{1}{n} \sum_{i=1}^{n} |\mathbf{u}_{i}^{0}|^{\ell} = \mathbb{E}[|U_{0}|^{\ell}] < \infty. \text{ Suppose that the properties } \mathcal{A}_{1}(t'), \mathcal{A}_{2}(t'), \mathcal{A}_{3}(t') \text{ hold for all } t' \in [0, t] \text{ for some } t < T.$ We first show $\mathcal{A}_{1}(t+1)$. For $\ell \geq 0$, let $\psi : \mathbb{R}^{t+2} \to \mathbb{R}, (u^{0}, ..., u^{t+1}) \mapsto |u^{t+1}|^{\ell}$. Clearly ψ is in $\text{PL}(O(\ell))$, thus by Proposition 5.1 $\underset{n \to \infty}{\text{p-lim}} \frac{1}{n} \sum_{i=1}^{n} \psi(\mathbf{u}_{i}^{0}, ..., \mathbf{u}_{i}^{t+1}) = \mathbb{E}[\psi(U_{0}, ..., U_{t+1})] < \infty.$ This proves

p-limsup $\frac{1}{n} \|\mathbf{u}^t\|_{\ell}^{\ell} < \infty$. Thus, we only need to show $\mathcal{A}_1(t+1)$ for the iterates $\tilde{\mathbf{u}}^t$. Let $\tilde{\mathbf{r}} = \sum_{j=1}^t \bar{b}_{t,j} \tilde{\mathbf{q}}^{j-1}$. Following [Mon21], we write

$$\begin{split} \tilde{\mathbf{u}}^{t+1} &= \mathbf{X}\tilde{\mathbf{q}}^t - \tilde{\mathbf{r}}^t \\ &= \tilde{\mathbf{P}}_{t-1}^{\perp} \mathbf{X} \tilde{\mathbf{P}}_{t-1}^{\perp} \tilde{\mathbf{q}}^t + \tilde{\mathbf{P}}_{t-1}^{\perp} \mathbf{X} \tilde{\mathbf{P}}_{t-1} \tilde{\mathbf{q}}^t + \tilde{\mathbf{P}}_{t-1} \mathbf{X} \tilde{\mathbf{q}}^t - \tilde{\mathbf{r}}^t \\ &\stackrel{\text{d}}{=} \tilde{\mathbf{P}}_{t-1}^{\perp} \hat{\mathbf{X}} \tilde{\mathbf{P}}_{t-1}^{\perp} \tilde{\mathbf{q}}^t + \tilde{\mathbf{P}}_{t-1}^{\perp} \mathbf{X} \tilde{\mathbf{P}}_{t-1} \tilde{\mathbf{q}}^t + \tilde{\mathbf{P}}_{t-1} \mathbf{X} \tilde{\mathbf{q}}^t - \tilde{\mathbf{r}}^t \\ &= \hat{\mathbf{X}} \tilde{\mathbf{P}}_{t-1}^{\perp} \tilde{\mathbf{q}}^t - \tilde{\mathbf{P}}_{t-1} \hat{\mathbf{X}} \tilde{\mathbf{P}}_{t-1}^{\perp} \tilde{\mathbf{q}}^t - \tilde{\mathbf{r}}^t + \tilde{\mathbf{P}}_{t-1}^{\perp} \mathbf{X} \tilde{\mathbf{P}}_{t-1} \tilde{\mathbf{q}}^t + \tilde{\mathbf{P}}_{t-1} \mathbf{X} \tilde{\mathbf{q}}^t \\ &= \mathbf{v}_1 - \mathbf{v}_2 - \mathbf{v}_3 + \mathbf{v}_4 + \mathbf{v}_5. \end{split}$$
(5.10)

where we used Lemma 5.8 in line (5.10) and $\hat{\mathbf{X}}$ is distributed as \mathbf{X} but independent of the σ -algebra $\sigma(\{\tilde{\mathbf{q}}^j, \tilde{\mathbf{u}}^j\}_{j \leq t})$. Furthermore, note that using $\mathcal{A}_3(j), j \leq t$, and (5.9)

$$\operatorname{p-lim}_{n \to \infty} \frac{1}{n} \tilde{\mathbf{F}}_{t-1}^{\top} \tilde{\mathbf{F}}_{t-1} = \operatorname{p-lim}_{n \to \infty} \frac{1}{n} \mathbf{F}_{t-1}^{\top} \mathbf{F}_{t-1} = \mathbf{Q}_{\leq t}.$$

Therefore $\tilde{\mathbf{F}}_{t-1}^{\top} \tilde{\mathbf{F}}_{t-1}$ is invertible whp and the following holds

$$\tilde{\mathbf{P}}_{t-1} = \tilde{\mathbf{F}}_{t-1} (\tilde{\mathbf{F}}_{t-1}^{\top} \tilde{\mathbf{F}}_{t-1})^{-1} \tilde{\mathbf{F}}_{t-1}^{\top}$$
(5.11)

We next bound p-limsup $\frac{1}{n^{1/\ell}} \| \mathbf{v}_k \|_{\ell}, k \in [5].$

1. We first deal with \mathbf{v}_1 . Denote $\mathbf{w} \triangleq \tilde{\mathbf{P}}_{t-1}^{\perp} \tilde{\mathbf{q}}^t$. Let $\mathbf{D} = \text{Diag}(z_1, ..., z_n)$ where z_i are independent normal $\mathcal{N}(0, 2/n)$, so that $\hat{\mathbf{X}} + \mathbf{D} \stackrel{d}{=} \text{GOE}(n)$, i.e the Gaussian Orthogonal Ensemble. Since $\hat{\mathbf{X}} + \mathbf{D}$ is independent of \mathbf{w} , it follows by simple verification that $(\hat{\mathbf{X}} + \mathbf{D})\mathbf{w} \stackrel{d}{=} (\|\mathbf{w}\|_2/\sqrt{n})\mathbf{g} + g_0\mathbf{w}/\sqrt{n}$ where $(g_0, \mathbf{g}) \stackrel{d}{=} \mathcal{N}(0, \mathbf{I}_{n+1})$. We then have

$$\begin{aligned} \|\mathbf{v}_1\|_{\ell} &= \|(\mathbf{X} + \mathbf{D})\mathbf{w} - \mathbf{D}\mathbf{w}\|_{\ell} \\ &\leq \|(\hat{\mathbf{X}} + \mathbf{D})\mathbf{w}\|_{\ell} + \|\mathbf{D}\mathbf{w}\|_{\ell} \\ &\leq \frac{\|\mathbf{w}\|_2}{\sqrt{n}} \|\mathbf{g}\|_{\ell} + \frac{|g_0|}{\sqrt{n}} \|\mathbf{w}\|_{\ell} + \|\mathbf{D}\mathbf{w}\|_{\ell}. \end{aligned}$$
(5.12)

To bound the p-limsup of the above, we first bound $\|\mathbf{w}\|_{\ell}$. Note that $\mathbf{w} = \tilde{\mathbf{q}}^t - \tilde{\mathbf{P}}_{t-1}\tilde{\mathbf{q}}^t$, and

$$\tilde{\mathbf{P}}_{t-1}\tilde{\mathbf{q}}^t = \sum_{j=0}^{t-1} a_{tj}\tilde{\mathbf{q}}^j, \quad a_{tj} \triangleq \sum_{k=0}^{t-1} (\tilde{\mathbf{F}}_{t-1}^\top \tilde{\mathbf{F}}_{t-1})_{jk}^{-1} \langle \tilde{\mathbf{q}}^k, \tilde{\mathbf{q}}^t \rangle.$$

Note that

$$|a_{tj}| \leq \sum_{k=0}^{t-1} \left| n(\tilde{\mathbf{F}}_{t-1}^{\top} \tilde{\mathbf{F}}_{t-1})_{jk}^{-1} \right| \left| \left\langle \frac{\tilde{\mathbf{q}}^k}{\sqrt{n}}, \frac{\tilde{\mathbf{q}}^t}{\sqrt{n}} \right\rangle \right|.$$

By $\mathcal{A}_1(k), k \leq t$ and using the fact that $\tilde{\mathbf{q}}^j, j \leq t$ are polynomials (of constant degrees/coefficients) in the iterates $\tilde{\mathbf{u}}^j$, it follows that p-limsup $|\langle \frac{\tilde{\mathbf{q}}^k}{\sqrt{n}}, \frac{\tilde{\mathbf{q}}^t}{\sqrt{n}} \rangle| = O(1)$. Furthermore, by (5.11) we have p-lim $n(\tilde{\mathbf{F}}_{t-1}^{\top}\tilde{\mathbf{F}}_{t-1})_{jk}^{-1} = (\mathbf{Q}_{\leq t})_{jk}^{-1} = O(1)$. Therefore, it follows that p-limsup $|a_{tj}| = O(1)$. Hence p-limsup $\frac{1}{n^{1/\ell}} \|\mathbf{w}\|_{\ell} \leq p$ -limsup $\frac{1}{n^{1/\ell}} \|\tilde{\mathbf{q}}^t\|_{\ell} + p$ -limsup $\frac{1}{n^{1/\ell}} \|\tilde{\mathbf{P}}_{t-1}\tilde{\mathbf{q}}^t\|_{\ell}$

$$\begin{split} \limsup_{n \to \infty} \frac{1}{n^{1/\ell}} \| \mathbf{w} \|_{\ell} &\leq \text{ p-limsup } \frac{1}{n^{1/\ell}} \| \tilde{\mathbf{q}}^t \|_{\ell} + \text{ p-limsup } \frac{1}{n^{1/\ell}} \| \mathbf{P}_{t-1} \tilde{\mathbf{q}}^t \| \\ &\leq O(1) + \text{ p-limsup } \sum_{j=0}^{t-1} \left(|a_{tj}| \frac{1}{n^{1/\ell}} \| \tilde{\mathbf{q}}^j \|_{\ell} \right) \\ &= O(1), \end{split}$$

where the last two inequalities follow from $\mathcal{A}_1(k), k \leq t$ combined with $p-\limsup_{n \to \infty} |a_{tj}| = O(1)$. Furthermore, we have

$$p-\limsup_{n \to \infty} \frac{1}{n^{1/\ell}} \|\mathbf{D}\mathbf{w}\|_{\ell} = p-\limsup_{n \to \infty} \frac{1}{n^{1/\ell}} \left(\sum_{i=1}^{n} |z_i|^{\ell} |\mathbf{w}_i|^{\ell} \right)^{\frac{1}{\ell}} \\ \leq p-\limsup_{n \to \infty} \max_{1 \le i \le n} |z_i| \frac{1}{n^{1/\ell}} \|\mathbf{w}\|_{\ell} \\ = 0.$$

where the last line follows from p-limsup $\frac{1}{n^{1/\ell}} \| \mathbf{w} \|_{\ell} = O(1)$ and the fact that $\max_{1 \le i \le n} |z_i| = \Theta(\sqrt{\log(n)}/\sqrt{n})$ with probability $1 - n^{-\Theta(1)}$. Putting everything together in (5.12) yields

$$\begin{aligned} \text{p-limsup} & \frac{1}{n^{1/\ell}} \| \mathbf{v}_1 \|_{\ell} \leq \text{ p-limsup} & \frac{1}{n^{1/\ell}} \frac{\| \mathbf{w} \|_2}{\sqrt{n}} \| \mathbf{g} \|_{\ell} + \text{ p-limsup} & \frac{1}{n^{1/\ell}} \frac{|g_0|}{\sqrt{n}} \| \mathbf{w} \|_{\ell} + \text{ p-limsup} & \frac{1}{n^{1/\ell}} \| \mathbf{D} \mathbf{w} \|_{\ell} \\ &= \text{ p-limsup} & \frac{\| \mathbf{w} \|_2}{\sqrt{n}} \frac{\| \mathbf{g} \|_{\ell}}{n^{1/\ell}} + \text{ p-limsup} & \frac{|g_0|}{\sqrt{n}} \frac{\| \mathbf{w} \|_{\ell}}{n^{1/\ell}}, \end{aligned}$$

note that p-limsup $\frac{\|\mathbf{w}\|_2}{\sqrt{n}} = O(1)$, p-limsup $\frac{\|\mathbf{w}\|_\ell}{n^{1/\ell}} = O(1)$ from previous arguments. Moreover, p-limsup $\frac{|g_0|}{n \to \infty} = 0$, and by virtue of CLT we have $\frac{\|\mathbf{g}\|_\ell}{n^{1/\ell}} = O(1)$. Henceforth p-limsup $\frac{1}{n^{1/\ell}} \|\mathbf{v}_1\|_\ell \le O(1)$. Which ends the analysis of $\|\mathbf{v}_1\|_\ell$.

2. We now deal with \mathbf{v}_2 similarly to the term \mathbf{v}_1 . Specifically, let $\mathbf{w} \triangleq \tilde{\mathbf{P}}_{t-1}^{\perp} \tilde{\mathbf{q}}^t$

$$\begin{aligned} \mathbf{v}_{2} &= \tilde{\mathbf{P}}_{t-1} \left((\hat{\mathbf{X}} + \mathbf{D}) \mathbf{w} - \mathbf{D} \mathbf{w} \right) \\ &= \tilde{\mathbf{P}}_{t-1} \left((\|\mathbf{w}\|_{2} / \sqrt{n}) \mathbf{g} + g_{0} \mathbf{w} / \sqrt{n} - \mathbf{D} \mathbf{w} \right) \\ &= (\|\mathbf{w}\|_{2} / \sqrt{n}) \tilde{\mathbf{P}}_{t-1} \mathbf{g} - \tilde{\mathbf{P}}_{t-1} \mathbf{D} \mathbf{w}, \end{aligned}$$

where we used $\tilde{\mathbf{P}}_{t-1}\mathbf{w} = 0$ in the last line as $\mathbf{w} = \tilde{\mathbf{P}}_{t-1}^{\perp}\tilde{\mathbf{q}}^t$. From the analysis of \mathbf{v}_1 , we have p-limsup $\|\mathbf{w}\|_2/\sqrt{n} = O(1)$. Furthermore

$$\tilde{\mathbf{P}}_{t-1}\mathbf{g} = \frac{1}{\sqrt{n}} \sum_{j=0}^{t-1} \tilde{g}_j \tilde{\mathbf{q}}^j,$$

where $(\tilde{g}_0, ..., \tilde{g}_{t-1})|_{\tilde{\mathbf{F}}_{t-1}} \stackrel{\mathsf{d}}{=} \mathcal{N}(0, (\tilde{\mathbf{F}}_{t-1}^\top \tilde{\mathbf{F}}_{t-1}/n)^{-1})$. By the induction hypothesis $\mathcal{A}_1(j), j \leq t$, we have p-limsup $\|\tilde{\mathbf{q}}^j\|_{\ell}/n^{1/\ell} = O(1)$. Therefore

$$p-\limsup_{n \to \infty} \frac{1}{n^{1/\ell}} \| \tilde{\mathbf{P}}_{t-1} \mathbf{g} \|_{\ell} \leq p-\limsup_{n \to \infty} \frac{O(1)}{\sqrt{n}} \sum_{j=0}^{t-1} | \tilde{g}_j | \\ \leq p-\limsup_{n \to \infty} \frac{O(t)}{\sqrt{n}} \| \tilde{\mathbf{g}} \|_2 \\ = 0,$$

where the last line follows from noting that $\mathbb{E}[\|\tilde{\mathbf{g}}\|_2^2|\tilde{\mathbf{F}}_{t-1}] = \operatorname{Tr}((\tilde{\mathbf{F}}_{t-1}^{\top}\tilde{\mathbf{F}}_{t-1}/n)^{-1}) \rightarrow_{n \to \infty} \operatorname{Tr}(\tilde{\mathbf{Q}}_{\leq t}^{-1}),$ which is bounded, then combining the latter with Markov inequality. Therefore

$$\operatorname{p-limsup}_{n \to \infty} \frac{1}{n^{1/\ell}} \| \mathbf{v}_2 \|_{\ell} \leq \operatorname{p-limsup}_{n \to \infty} \frac{1}{n^{1/\ell}} \| \tilde{\mathbf{P}}_{t-1} \mathbf{D} \mathbf{w} \|_{\ell}.$$

Note that

$$\tilde{\mathbf{P}}_{t-1}\mathbf{D}\mathbf{w} = \sum_{j=0}^{t-1} a_{tj}\tilde{\mathbf{q}}^j, \quad a_{tj} \triangleq \sum_{k=0}^{t-1} (\tilde{\mathbf{F}}_{t-1}^\top \tilde{\mathbf{F}}_{t-1})_{jk}^{-1} \langle \tilde{\mathbf{q}}^k, \mathbf{D}\mathbf{w} \rangle.$$

As shown in the analysis of \mathbf{v}_1 , it suffices to show $|a_{tj}| = O(1)$ to show p-limsup $\frac{1}{n^{1/\ell}} \|\tilde{\mathbf{P}}_{t-1}\mathbf{D}\mathbf{w}\|_{\ell} = O(1)$. We have

$$a_{tj} = \sum_{k=0}^{t-1} (\tilde{\mathbf{F}}_{t-1}^{\top} \tilde{\mathbf{F}}_{t-1}/n)_{jk}^{-1} \left\langle \frac{\tilde{\mathbf{q}}^k}{\sqrt{n}}, \frac{\mathbf{D}\mathbf{w}}{\sqrt{n}} \right\rangle.$$

As shown above $(\tilde{\mathbf{F}}_{t-1}^{\top}\tilde{\mathbf{F}}_{t-1}/n)_{jk}^{-1} = O(1)$, and by the induction hypothesis p-limsup $\|\tilde{\mathbf{q}}^k\|_2/\sqrt{n} = O(1)$, and as shown in the analysis of \mathbf{v}_1 , we have p-limsup $\|\mathbf{D}\mathbf{w}\|_2/\sqrt{n} = 0$. It follows that p-limsup $\|a_{t,j}\| = 0$. Henceforth, we have p-limsup $\frac{1}{n^{1/\ell}} \|\tilde{\mathbf{P}}_{t-1}\mathbf{D}\mathbf{w}\|_\ell = 0$, which readily yields p-limsup $\frac{1}{n^{1/\ell}} \|\mathbf{v}_2\|_\ell = O(1)$.

3. We now deal with \mathbf{v}_3 . We have

$$\operatorname{p-limsup}_{n \to \infty} \frac{1}{n^{1/\ell}} \| \mathbf{v}_3 \|_{\ell} \leq \operatorname{p-limsup}_{n \to \infty} \sum_{j=1}^t |\bar{b}_{t,j}| \frac{1}{n^{1/\ell}} \| \tilde{\mathbf{q}}^{j-1} \|_{\ell}.$$

We have $|\bar{b}_{tj}| = O(1)$, and by the inductive hypothesis p-limsup $n^{1/\ell} \|\tilde{\mathbf{q}}^{j-1}\|_{\ell} = O(1)$, therefore p-limsup $\frac{1}{n^{1/\ell}} \|\mathbf{v}_3\|_{\ell} = O(1)$.

4. We next deal with \mathbf{v}_4 . Let $\tilde{\mathbf{U}}_k \triangleq [\tilde{\mathbf{u}}^0|...|\tilde{\mathbf{u}}^{k+1}]$ and $\tilde{\mathbf{R}}_k \triangleq [0|\tilde{\mathbf{r}}^1|...|\tilde{\mathbf{r}}^k]$, and $\tilde{\mathbf{Y}}_k = \tilde{\mathbf{U}}_k + \tilde{\mathbf{R}}_k$ for $k \ge 0$. Note that $\mathbf{X}\tilde{\mathbf{F}}_{t-1} = \tilde{\mathbf{Y}}_{t-1}$. Introduce $\mathbf{b}^t \triangleq (\tilde{\mathbf{F}}_{t-1}^\top \tilde{\mathbf{F}}_{t-1})^{-1} \tilde{\mathbf{F}}_{t-1}^\top \tilde{\mathbf{q}}^t \in \mathbb{R}^t$. We have

$$\begin{split} \mathbf{v}_{4} &= \tilde{\mathbf{P}}_{t-1}^{\perp} \mathbf{X} \tilde{\mathbf{P}}_{t-1} \tilde{\mathbf{q}}^{t} \\ &= (\mathbf{I}_{n} - \tilde{\mathbf{P}}_{t-1}) \mathbf{X} \tilde{\mathbf{P}}_{t-1} \tilde{\mathbf{q}}^{t} \\ &= (\mathbf{I}_{n} - \tilde{\mathbf{P}}_{t-1}) \mathbf{X} \tilde{\mathbf{F}}_{t-1} (\tilde{\mathbf{F}}_{t-1}^{\top} \tilde{\mathbf{F}}_{t-1})^{-1} \tilde{\mathbf{F}}_{t-1}^{\top} \tilde{\mathbf{q}}^{t} \\ &= \tilde{\mathbf{Y}}_{t-1} \mathbf{b}^{t} - \tilde{\mathbf{P}}_{t-1} \tilde{\mathbf{Y}}_{t-1} \mathbf{b}^{t} \\ &\triangleq \mathbf{v}_{4,a} + \mathbf{v}_{4,b}. \end{split}$$

Using similar arguments to previous ones, we have p-limsup $\|\mathbf{b}^t\|_{\infty} = O(1)$. Furthermore, it follows from the inductive hypothesis $\mathcal{A}_1(j), j \leq t$ that p-limsup $\frac{1}{n^{1/\ell}} \|\tilde{\mathbf{u}}^j\|_{\ell} = O(1)$ and p-limsup $\frac{1}{n^{1/\ell}} \|\tilde{\mathbf{r}}^j\|_{\ell} = O(1)$ for all $j \leq t$. Therefore

$$p-\limsup_{n \to \infty} \frac{1}{n^{1/\ell}} \| \mathbf{v}_{4,a} \|_{\ell} \leq p-\limsup_{n \to \infty} \| \mathbf{b}^t \|_{\infty} \sum_{j=1}^t \frac{1}{n^{1/\ell}} \left(\| \tilde{\mathbf{u}}^j \|_{\ell} + \| \tilde{\mathbf{r}}^{j-1} \|_{\ell} \right)$$
$$= O(1).$$

We next bound $\mathbf{v}_{4,b}$. Note that

$$\mathbf{v}_{4,b} = \sum_{j=1}^{t} h_j \tilde{\mathbf{q}}^{j-1}, \quad (h_1, \dots, h_t) = (\tilde{\mathbf{F}}_{t-1}^{\top} \tilde{\mathbf{F}}_{t-1})^{-1} \tilde{\mathbf{F}}_{t-1}^{\top} \tilde{\mathbf{Y}}_{t-1} \mathbf{b}^t$$

Using the inductive hypothesis, we show similarly p-lim $\max_{1 \le j \le t} |h_j| = O(1)$, we then conclude using p-limsup $\frac{1}{n^{1/\ell}} \| \tilde{\mathbf{q}}^j \|_{\ell} = O(1)$ that p-limsup $\frac{1}{n^{1/\ell}} \| \mathbf{v}_{4,b} \| = O(1)$. Which ends the analysis of \mathbf{v}_4 .

5. The analysis of \mathbf{v}_5 is similar to \mathbf{v}_4 . Indeed, note that

$$\begin{aligned} \mathbf{v}_5 &= \mathbf{P}_{t-1} \mathbf{X} \tilde{\mathbf{q}}^t \\ &= \tilde{\mathbf{F}}_{t-1} (\tilde{\mathbf{F}}_{t-1}^\top \tilde{\mathbf{F}}_{t-1})^{-1} \tilde{\mathbf{F}}_{t-1}^\top \mathbf{X} \tilde{\mathbf{q}}^t \\ &= \tilde{\mathbf{F}}_{t-1} (\tilde{\mathbf{F}}_{t-1}^\top \tilde{\mathbf{F}}_{t-1})^{-1} (\mathbf{X} \tilde{\mathbf{F}}_{t-1})^\top \tilde{\mathbf{q}}^t \\ &= \tilde{\mathbf{F}}_{t-1} (\tilde{\mathbf{F}}_{t-1}^\top \tilde{\mathbf{F}}_{t-1})^{-1} \tilde{\mathbf{Y}}_{t-1}^\top \tilde{\mathbf{q}}^t. \end{aligned}$$

It suffices then to recall p-limsup $\|n(\tilde{\mathbf{F}}_{t-1}^{\top}\tilde{\mathbf{F}}_{t-1})^{-1}\|_{\infty} = O(1)$, and use the inductive hypothesis $\mathcal{A}_1(j), j \leq t$ to show p-limsup $\|\tilde{\mathbf{Y}}_{t-1}^{\top}\tilde{\mathbf{q}}^t\|_{\infty} = O(1)$, then conclude that p-limsup $\|\mathbf{v}_5\|_{\ell}/n^{1/\ell} = O(1)$.

This ends the proof of $\mathcal{A}_1(t+1)$. We next show $\mathcal{A}_2(t+1)$. We have

$$\frac{1}{\sqrt{n}} \|\tilde{\mathbf{u}}^{t+1} - \mathbf{u}^{t+1}\|_2 \le \|\mathbf{X}\|_{\text{op}} \frac{1}{\sqrt{n}} \|\tilde{\mathbf{q}}^t - \mathbf{q}^t\|_2 + \sum_{j=1}^t |b_{t,j} - \bar{b}_{t,j}| \frac{1}{\sqrt{n}} \|\mathbf{q}^{j-1}\|_2 + \sum_{j=1}^t |\bar{b}_{t,j}| \frac{1}{\sqrt{n}} \|\tilde{\mathbf{q}}^{j-1} - \mathbf{q}^{j-1}\|_2 \\ \triangleq a_1 + a_2 + a_3.$$

1. We first deal with a_1 . We have $\|\mathbf{X}\|_{\text{op}} \leq \Theta(1)$ with probability $1 - \exp(-\Theta(n))$. Therefore we have whp

$$a_1 \le O(1) \frac{1}{\sqrt{n}} \| \tilde{\mathbf{q}}^t - \mathbf{q}^t \|_2$$

Denote $\mathbf{s}_i = (\mathbf{u}_i^0, ..., \mathbf{u}_i^t)$ and $\tilde{\mathbf{s}}_i = (\tilde{\mathbf{u}}_i^0, ..., \tilde{\mathbf{u}}_i^t)$, so that

$$\frac{1}{n} \|\tilde{\mathbf{q}}^t - \mathbf{q}^t\|_2^2 = \frac{1}{n} \sum_{i=1}^n (q^t(\tilde{\mathbf{s}}_i) - q^t(\mathbf{s}_i))^2.$$

Since q^t is polynomial, it belongs to PL(m) for some $m \ge 0$. Therefore

$$\frac{1}{n}\sum_{i=1}^{n} (q^{t}(\tilde{\mathbf{s}}_{i}) - q^{t}(\mathbf{s}_{i}))^{2} \leq \frac{C}{n}\sum_{i=1}^{n} (1 + \|\tilde{\mathbf{s}}_{i}\|_{2}^{m-1} + \|\mathbf{s}_{i}\|_{2}^{m-1})^{2} \|\tilde{\mathbf{s}}_{i} - \mathbf{s}_{i}\|_{2}^{2} \\
\leq \frac{C}{n}\sum_{i=1}^{n} (1 + \|\tilde{\mathbf{s}}_{i}\|_{2} + \|\mathbf{s}_{i}\|_{2})^{2m-1} \|\tilde{\mathbf{s}}_{i} - \mathbf{s}_{i}\|_{2} \\
\leq C \left(\frac{1}{n}\sum_{i=1}^{n} (1 + \|\tilde{\mathbf{s}}_{i}\|_{2} + \|\mathbf{s}_{i}\|_{2})^{4m-2}\right)^{1/2} \left(\sum_{i=1}^{n} \frac{1}{n} \|\tilde{\mathbf{s}}_{i} - \mathbf{s}_{i}\|_{2}^{2}\right)^{1/2} \\
= C \left(\frac{1}{n}\sum_{i=1}^{n} (1 + \|\tilde{\mathbf{s}}_{i}\|_{2} + \|\mathbf{s}_{i}\|_{2})^{4m-2}\right)^{1/2} \left(\sum_{j=1}^{t} \frac{1}{n} \|\tilde{\mathbf{u}}^{j} - \mathbf{u}^{j}\|_{2}^{2}\right)^{1/2}.$$

Using the inductive hypothesis $\mathcal{A}_2(j), j \leq t$, we have p-limsup $\left(\sum_{j=1}^t \frac{1}{n} \|\tilde{\mathbf{u}}^j - \mathbf{u}^j\|_2^2\right)^{1/2} = 0$. Moreover, using $\mathcal{A}_1(j), j \leq t$, we have p-limsup $\left(\frac{1}{n}\sum_{i=1}^n (1+\|\tilde{\mathbf{s}}_i\|_2+\|\mathbf{s}_i\|_2)^{4m-2}\right)^{1/2} = O(1)$. Henceforth p-lim $\frac{1}{n}\sum_{i=1}^n (q^t(\tilde{\mathbf{s}}_i) - q^t(\mathbf{s}_i))^2 = 0$, which yields p-limsup $a_1 = 0$.

- 2. We now deal with a_2 . From the inductive hypothesis $\mathcal{A}_1(t)$, we have p-limsup $\frac{1}{\sqrt{n}} \|\mathbf{q}^t\|_2 = O(1)$, and by Proposition 5.1 we have p-limsup $b_{tj} = \bar{b}_{tj}$. Therefore p-limsup $a_2 = 0$.
- 3. The term a_3 is dealt with identically to a_1 and a_2 .

This concludes the proof of $\mathcal{A}_2(t+1)$. It remains to prove $\mathcal{A}_3(t+1)$, which we do next. Let $\psi \in PL(m)$, and use the same notations above. Then

$$\begin{aligned} \left| \frac{1}{n} \sum_{i=1}^{n} \psi(\tilde{\mathbf{s}}_{i}) - \frac{1}{n} \sum_{i=1}^{n} \psi(\mathbf{s}_{i}) \right| &\leq \frac{C}{n} \sum_{i=1}^{n} (1 + \|\tilde{\mathbf{s}}_{i}\|_{2} + \|\mathbf{s}_{i}\|_{2})^{m-1} \|\tilde{\mathbf{s}}_{i} - \mathbf{s}_{i}\|_{2} \\ &\leq C \left(\frac{1}{n} \sum_{i=1}^{n} (1 + \|\tilde{\mathbf{s}}_{i}\|_{2} + \|\mathbf{s}_{i}\|_{2})^{2m-2} \right)^{1/2} \left(\sum_{i=1}^{n} \frac{1}{n} \|\tilde{\mathbf{s}}_{i} - \mathbf{s}_{i}\|_{2}^{2} \right)^{1/2} \\ &= C \left(\frac{1}{n} \sum_{i=1}^{n} (1 + \|\tilde{\mathbf{s}}_{i}\|_{2} + \|\mathbf{s}_{i}\|_{2})^{2m-2} \right)^{1/2} \left(\sum_{j=1}^{t} \frac{1}{n} \|\tilde{\mathbf{u}}^{j} - \mathbf{u}^{j}\|_{2}^{2} \right)^{1/2}. \end{aligned}$$

Using the same arguments as in the proof of $\mathcal{A}_2(t+1)$, we conclude that $\operatorname{p-limsup}_{n\to\infty} \left| \frac{1}{n} \sum_{i=1}^n \psi(\tilde{\mathbf{s}}_i) - \frac{1}{n} \sum_{i=1}^n \psi(\mathbf{s}_i) \right| = 0$. Which ends the proof of $\mathcal{A}_3(t+1)$, and concludes the proof of Proposition 5.12.

We now turn to the proof of Proposition 5.5. Let \mathbf{Q}^{IAMP} be the covariance matrix associated with the IAMP algorithm in [Mon19]. Let $\hat{\mathbf{v}}, \tilde{\mathbf{v}}$ as in (5.2), (5.3) and let their associated iterates be $\hat{\mathbf{u}}^t, \tilde{\mathbf{u}}^t$, and let $\hat{\mathbf{Q}}$ be the covariance matrix of the AMP iterates $\hat{\mathbf{u}}^t$. In order to apply Proposition 5.9, on the iterates $\hat{\mathbf{u}}^t, \tilde{\mathbf{u}}^t$, we first need to verify (5.8) for $\hat{\mathbf{Q}}$, which we do next. Property (5.9) is shown to hold for \mathbf{Q}^{IAMP} in [Mon19], as the matrix \mathbf{Q}^{IAMP} is diagonal with positive diagonal terms bounded above and below by constants. Moreover, it is shown in [IS23](Claim B.5) that $\|\hat{\mathbf{Q}}_{\leq t} - \mathbf{Q}_{\leq t}^{\text{IAMP}}\|_{\text{fro}}$ can be taken arbitrarily small for any $t \in [T]$ and fixed number of iterations T. Hence, taking $\sup_{t \leq T} \|\hat{\mathbf{Q}}_{\leq t} - \mathbf{Q}_{\leq t}^{\text{IAMP}}\|_{\text{fro}}$ small enough, the matrix $\hat{\mathbf{Q}}$ satisfies (5.8). We then have

$$\frac{\|\hat{\mathbf{v}} - \tilde{\mathbf{v}}\|_2}{\sqrt{n}} \le \sqrt{\delta} \sum_{t=1}^{\lfloor \bar{q}/\delta \rfloor} \frac{\|q^t(\hat{\mathbf{u}}^0, ..., \hat{\mathbf{u}}^t) - q^t(\tilde{\mathbf{u}}^0, ..., \tilde{\mathbf{u}}^t)\|_2}{\sqrt{n}}.$$

Following the notations in the proof of Proposition 5.9, denote by $\hat{\mathbf{q}}^t$ the vector $q^t(\hat{\mathbf{u}}^0, ..., \hat{\mathbf{u}}^t)$, and let $\hat{\mathbf{s}}_i = (\hat{\mathbf{u}}_i^0, ..., \hat{\mathbf{u}}_i^t)$. Let $t \in [\lfloor \bar{q}/\delta \rfloor]$. Since q^t is a polynomial with constant degree and coefficients, we have $\mathbf{q}^t \in \mathrm{PL}(m)$ for some constant $m \ge 0$. We then have

$$\frac{\|q^{t}(\hat{\mathbf{u}}^{0},...,\hat{\mathbf{u}}^{t}) - q^{t}(\tilde{\mathbf{u}}^{0},...,\tilde{\mathbf{u}}^{t})\|_{2}^{2}}{n} \leq \frac{C}{n} \sum_{i=1}^{n} (1 + \|\hat{\mathbf{s}}_{i}\|_{2} + \|\tilde{\mathbf{s}}_{i}\|_{2})^{m-1} \|\hat{\mathbf{s}}_{i} - \tilde{\mathbf{s}}_{i}\|_{2}$$
$$\leq C \left(\frac{1}{n} \sum_{i=1}^{n} (1 + \|\hat{\mathbf{s}}_{i}\|_{2} + \|\tilde{\mathbf{s}}_{i}\|_{2})^{2m-2}\right)^{1/2} \left(\sum_{i=1}^{n} \frac{1}{n} \|\hat{\mathbf{s}}_{i} - \tilde{\mathbf{s}}_{i}\|_{2}^{2}\right)^{1/2}$$
$$= C \left(\frac{1}{n} \sum_{i=1}^{n} (1 + \|\hat{\mathbf{s}}_{i}\|_{2} + \|\tilde{\mathbf{s}}_{i}\|_{2})^{2m-2}\right)^{1/2} \left(\sum_{j=1}^{t} \frac{1}{n} \|\hat{\mathbf{u}}^{j} - \tilde{\mathbf{u}}^{j}\|_{2}^{2}\right)^{1/2}.$$

Applying $\mathcal{A}_1(j), \mathcal{A}_2(j), j \leq t$ from Proposition 5.9 to the above yields

$$\underset{n \to \infty}{\text{p-lim}} \frac{\|q^t(\hat{\mathbf{u}}^0, \dots, \hat{\mathbf{u}}^t) - q^t(\tilde{\mathbf{u}}^0, \dots, \tilde{\mathbf{u}}^t)\|_2^2}{n} = 0,$$

and therefore

$$\operatorname{p-lim}_{n \to \infty} \frac{\|\hat{\mathbf{v}} - \tilde{\mathbf{v}}\|_2}{\sqrt{n}} = 0$$

which ends the proof of Proposition 5.5.

References

- [AM23] Anurag Anshu and Tony Metger. "Concentration bounds for quantum states and limitations on the QAOA from polynomial approximations". In: *Quantum* 7 (2023), p. 999.
- [Arb] Louis François Antoine Arbogast. Du calcul des dérivations. Levrault, frères, 1800.
- [Bas22] Joao Basso et al. "Performance and limitations of the QAOA at constant levels on large sparse hypergraphs and spin glass models". In: 2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS). IEEE. 2022, pp. 335–343.
- [BMLM15] Mohsen Bayati, Marc Lelarge, and Andrea Montanari. "Universality in polytope phase transitions and message passing algorithms". In: (2015).
 - [Cha06] Sourav Chatterjee. "A generalization of the Lindeberg principle". In: Ann. Probab. 34.1 (2006), pp. 2061–2076.
 - [Dei14] Percy A Deift et al. "Universality in numerical computations with random data". In: *Proceedings* of the National Academy of Sciences 111.42 (2014), pp. 14973–14978.
 - [AMS17] Amir Dembo, Andrea Montanari, and Subhabrata Sen. "Extremal cuts of sparse random graphs". In: (2017).
 - [EMS23] Ahmed El Alaoui, Andrea Montanari, and Mark Sellke. "Local algorithms for maximum cut and minimum bisection on locally treelike regular graphs of large degree". In: *Random Structures* & Algorithms 63.3 (2023), pp. 689–715.
 - [Fen22] Oliver Y Feng et al. "A unifying tutorial on approximate message passing". In: Foundations and Trends® in Machine Learning 15.4 (2022), pp. 335–536.
 - [FK81] Zoltán Füredi and János Komlós. "The eigenvalues of random symmetric matrices". In: Combinatorica 1 (1981), pp. 233–241.
 - [IS23] Misha Ivkov and Tselil Schramm. "Semidefinite programs simulate approximate message passing robustly". In: arXiv preprint arXiv:2311.09017 (2023).
 - [Mon21] Andrea Montanari. "Optimization of the Sherrington–Kirkpatrick Hamiltonian". In: SIAM Journal on Computing 0 (2021), FOCS19–1.
 - [Mon19] Andrea Montanari. "Optimization of the Sherrington-Kirkpatrick Hamiltonian". In: 2019 IEEE 60th Annual Symposium on Foundations of Computer Science (FOCS). 2019.
 - [Mon22] Andrea Montanari and Alexander S Wein. "Equivalence of approximate message passing and low-degree polynomials in rank-one matrix estimation". In: *arXiv preprint arXiv:2212.06996* (2022).
 - [MOO10] Elchanan Mossel, Ryan O'Donnell, and Krzysztof Oleszkiewicz. "Noise stability of functions with low influences: Invariance and optimality". In: Annals of Mathematics (2010), pp. 295– 341.
 - [Pan13] Dmitry Panchenko. The Sherrington-Kirkpatrick model. Springer Science & Business Media, 2013.
 - [Par79] Giorgio Parisi. "Infinite number of order parameters for spin-glasses". In: Physical Review Letters 43.23 (1979), p. 1754.
 - [Sen18] Subhabrata Sen. "Optimization on sparse random hypergraphs and spin glasses". In: Random Structures & Algorithms 53.3 (2018), pp. 504–536.
 - [Tal06] Michel Talagrand. "The parisi formula". In: Annals of mathematics (2006), pp. 221–263.