

# Covariant LTB collapse in models of loop quantum gravity

Martin Bojowald<sup>1,\*</sup>, Erick I. Duque<sup>1†</sup> and Dennis Hartmann<sup>1,2‡</sup>

<sup>1</sup> Institute for Gravitation and the Cosmos,  
The Pennsylvania State University,  
104 Davey Lab, University Park, PA 16802, USA

<sup>2</sup> Department of Astronomy and Astrophysics,  
The Pennsylvania State University,  
525 Davey Lab, University Park, PA 16802, USA

## Abstract

Models of gravitational collapse provide important means to test whether non-classical space-time effects motivated for instance by quantum gravity can be realized in generic ways in physically relevant situations. Here, a detailed analysis of marginally bound Lemaître–Tolman–Bondi space-times is given in emergent modified gravity, which in particular includes a covariant formulation of holonomy modifications usually considered in models of loop quantum gravity. As a result, generic collapse in this setting is shown to imply a physical singularity that removes the bouncing behavior seen in vacuum space-times with the same type of modifications.

## 1 Introduction

Black holes present a fascinating problem in general relativity and possible extensions such as modified or quantum gravity. Being described by a small parameter space, they appear as rather simple objects from a classical perspective. However, it is often postulated that quantum effects should start to dominate near the singularity of a black hole, where curvature is large and divergences lead to undefined quantities in the classical theory. Various approaches to quantum gravity have therefore suggested modifications of the classical behavior of space-time with the main goal of resolving physical singularities.

Unraveling the mysteries of black holes requires an understanding of their formation. The conditions and mechanisms that cause the collapse of matter to a black hole are numerous. In this paper, we shall focus on the Lemaître–Tolman–Bondi (LTB) collapse model of dust matter within emergent modified gravity, extending the original analysis with canonical methods given in [1, 2]. We leave the critical collapse of scalar fields to a future paper. Both types of collapse models are well-established and have been studied from different perspectives. LTB collapse is of interest in the present paper in order to

---

\*e-mail address: [bojowald@psu.edu](mailto:bojowald@psu.edu)

†e-mail address: [eqd5272@psu.edu](mailto:eqd5272@psu.edu)

‡e-mail address: [dennis.hartmann@psu.edu](mailto:dennis.hartmann@psu.edu)

study a manifestly covariant model of matter collapse in which partial analytical results are available.

Emergent modified gravity [3, 4] is an approach to modified gravity crucially based on the canonical formulation of general relativity. Because canonical expressions do not require space-time integrals, they impose weaker assumptions on the basic setup than space-time scalars required for effective space-time actions. As a result, new theories of modified gravity are available, at least in spherical symmetry and polarized Gowdy systems [5], even if one works at the classical derivative order of equations of motion and does not include higher time derivatives or extra fields. The space-time geometries resulting from solutions of the theory are Riemannian, but the line element is not directly given by one of the fundamental fields and instead has to be derived from the field equations. Couplings to perfect fluids, scalar matter, and the electromagnetic field have been derived in [6, 7, 8].

Prior studies of black holes in emergent modified gravity or related approaches [9] have revealed interesting results, such as the potential for non-singular behavior [10, 11, 12] or, for a special set of modifications, dynamical signature change of metric components at classical scales [13] which demonstrates the non-standard nature of new space-time solutions. Collapse models have been analyzed in an Oppenheimer–Snyder version [6], and already in LTB models [14, 15]. The latter models were restricted to a subset of modification functions, in particular with a constant holonomy length, that are known to lead to large deviations from classical behavior even on small curvature scales. One of the main aims of our work is to provide a sufficiently general formulation and analysis of LTB models that also includes decreasing holonomy lengths.

Emergent modified gravity includes modification functions that clarify the appearance of holonomy modifications of loop quantum gravity [16]. It therefore provides a unique systematic way of turning preliminary constructions that implement loop effects in a fixed gauge or in a deparameterized theory into fully covariant space-time models. (Alternative constructions of a limited set of such models have been presented in [17]. However, as discussed in [18], they are special cases of emergent modified gravity and do not constitute a systematic derivation of modified gravity within a reliable setting of effective field theory.) In particular, emergent modified gravity can be used to identify problems in LTB models of loop quantum gravity, such as non-covariance in the recent [19, 20, 21, 22, 23], and amend them by including symmetry-restoring terms in the Hamiltonian constraint.

An issue that has been discussed extensively is the physical meaning of shell-crossing singularities and potential shock waves that may occur at a turning point of collapse, a possible cure of the classical curvature singularity. A careful analysis has been given in [21] where it was found important to compare results from different choices of coordinates and space-time slicings. However, the same paper pointed out that available models had not been shown to be covariant, which would be an important prerequisite to any physical analysis of space-time effects. Here, we complete the discussion by constructing a corresponding formulation that is explicitly shown to be covariant. Our main result is surprising: While covariance can be restored, we find that LTB models with modifications motivated by loop quantum gravity have a physical, rather than shell-crossing, singularity even though corresponding vacuum models are generically non-singular. A similar result

with dust [6] and scalar matter [7] had already been found in emergent modified gravity, and [14, 15] extended this effect to LTB collapse (with constant holonomy length). The latter papers then introduced a conformal transformation to a non-singular solution and argued that this should be the physical one. A second aim of the present paper is to critically examine this proposal, which we find inadmissible when seen from the perspective of a modified theory rather than a single modified solution.

The result of a generic re-appearance of physical singularities in LTB collapse raises important questions about the reliability of singularity resolution or other effects found in simple models of loop quantum gravity. It highlights the importance of constructing fully covariant space-time models whenever a space-time analysis based on effective line elements is performed: The singular behavior can be traced back to a crucial factor in the space-time metric that had been missed in non-covariant models. It is also important to include a generic set of modification functions in one's class of theories, such that all potential effects at the same order of derivatives can be included. Emergent modified gravity fulfills this condition from the point of view of effective field theory.

Our construction of LTB models in emergent modified gravity is presented in Section 2. We pay special attention to the non-fundamental nature of LTB conditions, which are motivated by convenient forms of solutions rather than fundamental principles such as symmetries. The procedures for choosing and implementing such conditions are therefore not as unique as, say, symmetry reduction. Moreover, the classical conditions are motivated by a specific form of the metric directly constructed from kinematical phase-space variables such as the spatial metric or triad, for which equations of motion are first formulated and then solved. Models of emergent modified gravity, and therefore covariant models of loop quantum gravity, are crucially different because their space-time metric in terms of phase-space variables is emergent and not known before equations of motion are imposed.

Our detailed discussion of these issues is followed in Section 3 by an analysis of various collapse solutions. In particular, in Section 3.3 we demonstrate that local covariant marginal LTB solutions with holonomy modifications are strictly reflection symmetric around a hypersurface of maximum curvature. This symmetry, which is explicitly demonstrated by specific solutions in Sections 3.5 and 3.6, implies that shock waves cannot form near maximum curvature unless they had already been present in the initial data of collapse. At the same time, we will see that the maximum curvature is infinite, implying a physical singularity even in the presence of modifications from loop quantum gravity. Our concluding Section 4 emphasizes several key advantages of embedding models of loop quantum gravity within emergent modified gravity that were crucial for our analysis.

## 2 LTB models in emergent modified gravity

Classical line elements of LTB type [24] are given by

$$ds^2 = -dt^2 + \frac{(R')^2}{1 + \kappa(x)} dx^2 + R^2(d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad (1)$$

where  $\kappa(x)$  is a non-dynamical radial function that distinguishes different models. The dynamical variable,  $R(t, x)$ , is a function of the time and radial coordinates subject to the differential equation

$$\dot{R}^2 = \kappa(x) + \frac{2m(x)}{R} \quad (2)$$

with another radial function,  $m(x)$ , that is determined by the initial mass distribution: The energy density of matter is given by

$$T_{tt} = \frac{m'}{4\pi GR^2 R'}. \quad (3)$$

Appendix A provides a more detailed review as well as several examples.

## 2.1 LTB models in canonical gravity

For an implementation in emergent modified gravity, the canonical formulation of an LTB reduction is crucial. Since LTB models are spherically symmetric, they can be expressed within the general class of line elements of the form

$$ds^2 = -N^2 dt^2 + \frac{(E^\varphi)^2}{E^x} (dx + N^x dt)^2 + E^x (d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad (4)$$

with free functions  $N(t, x)$ ,  $N^x(t, x)$ ,  $E^x(t, x)$  and  $E^\varphi(t, x)$ . In this form,  $N$  and  $N^x$  are the usual lapse function and radial shift vector, while  $E^x$  and  $E^\varphi$  are components of a densitized triad that may be used to parameterize spatial metric components. The orientation of the triad is given by the sign of  $E^x$ , which we will assume to be positive throughout this paper.

### 2.1.1 Vacuum properties

These variables are convenient in a canonical formulation, where, together with their momenta  $K_x$  and  $K_\varphi$ , they are subject to the Hamiltonian constraint [25, 26]

$$H^{\text{grav}}[N] = \int dx N \left[ \frac{((E^x)')^2}{8\sqrt{|E^x|}E^\varphi} - \frac{E^\varphi}{2\sqrt{|E^x|}} - \frac{E^\varphi K_\varphi^2}{2\sqrt{|E^x|}} - 2K_\varphi \sqrt{|E^x|} K_x - \frac{\sqrt{|E^x|}(E^x)'(E^\varphi)'}{2(E^\varphi)^2} + \frac{\sqrt{|E^x|}(E^x)''}{2E^\varphi} \right] = 0 \quad (5)$$

and the diffeomorphism constraint

$$H_x^{\text{grav}}[N^x] = \int dx N^x (K_\varphi' E^\varphi - K_x (E^x)') = 0 \quad (6)$$

in vacuum space-times. The LTB form of the line element is obtained with the gauge-fixing condition

$$N = 1 \quad , \quad N^x = 0 \quad (7)$$

and the LTB condition

$$E^\varphi = \frac{(E^x)'}{2\sqrt{1 + \kappa(x)}}. \quad (8)$$

The dynamical function  $R(t, x)$  is then identified with  $R = \sqrt{E^x}$ .

In a complete LTB solution, the LTB condition

$$L_1 \equiv E^\varphi - \frac{(E^x)'}{2\sqrt{1 + \kappa(x)}} = 0 \quad (9)$$

must be fulfilled at all times, and therefore  $L_1$  must Poisson commute with the Hamiltonian  $H[N] + H_x[N^x] = H[1]$  on-shell, when the constraints and equations of motion hold [1, 2]. As shown in more detail in App. A, this condition evaluates to

$$\{L_1, H[1]\} = \left( \frac{K_\varphi}{\sqrt{|E^x|}} + \frac{\sqrt{|E^x|}}{E^\varphi} 2K_x \right) L_1 - \frac{\sqrt{|E^x|}}{E^\varphi} \frac{H_x^{\text{grav}}}{\sqrt{1 + \kappa(x)}} \quad (10)$$

and therefore does vanish on-shell for vacuum LTB solutions, where  $H_x^{\text{grav}} = 0$ .

For further applications in the modified case, we note that our minimal implementation of the LTB condition, following [1, 2] but in contrast to [22], is suitable for a phenomenological reduction. Compared with symmetry reduction, for instance from the full theory to spherical symmetry, the LTB condition does not have a fundamental interpretation but rather imposes a restriction to a convenient but not physically characterized set of solutions. Such a phenomenological reduction is distinguished from a fundamental reduction in that the resulting solution space need not be equipped with a phase-space structure. Thus, we have minimally required that the LTB condition  $L_1$  is preserved in time, which led us to an on-shell condition that is automatically fulfilled for the classical dynamics. Since LTB models only implement a specific solutions space, an on-shell treatment is sufficient. It is not necessary to analyze our  $L_2$  off-shell and determine, for instance, whether the pair  $(L_1, L_2)$  is first class or second class and, in the latter case, derive the corresponding Dirac bracket. For phenomenological reductions, it is possible to implement a single condition,  $L_1 = 0$ . In a fundamental reduction, a single condition would always be first class and requires factoring out gauge transformations in order to obtain an even-dimensional phase space with a non-degenerate symplectic structure. A phenomenological reduction, by contrast, may easily work with an odd-dimensional solution space since no reduced phase-space structure is required. It is important to remember the distinction between phenomenological and fundamental reduction when the LTB condition is evaluated in modified gravity.

In what follows, we demonstrate that standard LTB solutions are obtained with this procedure when applied to general relativity. Any implementation of the same procedure in modified gravity in canonical form therefore provides valid modifications of the LTB solution space. It is, however, consistent to follow a more involved process such as the full implementation of the formal Dirac algorithm as performed in [22]. The availability of different methods for phenomenological reduction constitutes a source of ambiguities in models of modified gravity. For this reason, LTB models, in contrast, for instance,

to models of scalar collapse in spherical symmetry, do not provide conclusive answers to fundamental questions such as the fate of black-hole singularities or potential observational implications. Nevertheless, they play an important role as tractable testing grounds of various ideas in modified and quantum gravity.

### 2.1.2 Dust

Classical LTB models are compatible with dust-like matter as the source of the mass function  $m(x)$ . As an example of perfect fluids, dust is canonically described by two spatial scalars,  $T(x)$  and  $X(x)$ , that can be viewed as space-time coordinates of evolving fluid particles; see also [27]. Their momenta  $P_T$  and  $P_X$  are spatial densities and represent energy and momentum of the particles, which therefore have the radial velocity  $P_X/P_T$ . They appear in the matter contributions to the Hamiltonian constraint,

$$H_s^{\text{matter}} = \sqrt{sP_T^2 + \frac{E^x}{(E^\varphi)^2}(H_x^{\text{matter}})^2} - \sqrt{\det q} P(P_X^2/P_T^2), \quad (11)$$

and the diffeomorphism constraint,

$$H_x^{\text{matter}} = P_T T' + P_X X'. \quad (12)$$

In the first expression, the constant  $s$  takes the value  $s = 1$  for massive particles, and  $s = 0$  for massless ones. The pressure function  $P(P_X^2/P_T^2)$  vanishes in the case of dust, and  $\sqrt{\det q} = \sqrt{E^x E^\varphi}$  is the spatial volume element.

For dust, we have to evaluate the preservation of the LTB condition  $L_1$  with the Hamiltonian  $H[1] = H[1]^{\text{grav}} + H^{\text{matter}}[1]$ , which leads to

$$\{L_1, H[1]\} = \frac{\sqrt{E^x}}{E^\varphi} \frac{H_x^{\text{matter}}}{\sqrt{1 + \kappa(x)}} \quad (13)$$

on-shell. Stability of the LTB condition therefore requires  $H_x^{\text{matter}} = 0$  in the LTB gauge or slicing, which implies that the frame is co-moving with the fluid. We then fix  $s = 1$  because a physical co-moving frame exists only if the fluid particles are massive.

## 2.2 Emergent modified gravity

The spherically symmetric constraints have Poisson brackets

$$\{H_x[N^x], H_x[M^x]\} = -H_x[M^x(N^x)' - N^x(M^x)'], \quad (14)$$

$$\{H[N], H_x[M^x]\} = -H[M^x N'], \quad (15)$$

$$\{H[N], H[M]\} = -H_x[q^{xx}(MN' - NM')] \quad (16)$$

with the inverse radial metric  $q^{xx} = E^x/(E^\varphi)^2$ . They characterize canonical gauge transformations as hypersurface deformations. The gauge transformations of spatial metric

components (or the densitized triad) and their momenta, or any phase-space function depending on them, are directly obtained in canonical form as

$$\delta_\epsilon f(E, K) = \{f(E, K), H[\epsilon^0] + H_x[\epsilon^x]\} \quad (17)$$

with two components  $\epsilon^0$  and  $\epsilon^x$  of the gauge function  $\epsilon$ .

Gauge transformations of lapse and shift are more indirect because these functions do not have momenta in canonical gravity, but as metric components should nevertheless change when a spatial hypersurface is deformed in a new slicing. Their transformations,

$$\delta_\epsilon N = \dot{\epsilon}^0 + \epsilon^x N' - N^x (\epsilon^0)' \quad (18)$$

$$\delta_\epsilon N^x = \dot{\epsilon}^x + \epsilon^x (N^x)' - N^x (\epsilon^x)' + q^{xx} (\epsilon^0 N' - N (\epsilon^0)') \quad (19)$$

follow from an implementation of consistency of evolution generated by  $H[N] + H_x[N^x]$  with gauge transformations (17) generated by the same constraints [28, 29]. In particular, the structure function  $q^{xx}$  in (16) appears in the transformations (19).

The canonical theory is covariant because of a crucial on-shell equivalence of gauge transformations with space-time Lie derivatives of the metric, expressed as

$$\delta_\epsilon g_{\mu\nu}|_{\text{O.S.}} = \mathcal{L}_\xi g_{\mu\nu}|_{\text{O.S.}} \quad (20)$$

Here, ‘‘O.S.’’ refers to on-shell, requiring an evaluation of the condition on the solution space of the theory. In this way, it is possible to relate momenta in the canonical expressions on the left with time components of the Lie derivative on the right. Moreover, the space-time vector field  $\xi^\mu$  is related to the gauge functions by

$$\xi^\mu = \epsilon^0 n^\mu + \epsilon^x s^\mu = \xi^t t^\mu + \xi^x s^\mu \quad (21)$$

transforming the normal frame used for  $\epsilon$  to the coordinate frame used for  $\xi$ , with the time-evolution vector field  $t^\mu + N n^\mu + N^x s^\mu$  and a radial vector field  $s^\mu$  chosen as a basis for the tangent space on a radial hypersurface. The components are therefore related by

$$\xi^t = \frac{\epsilon^0}{N} \quad , \quad \xi^x = \epsilon^x - \frac{\epsilon^0}{N} N^x \quad (22)$$

This classical structure of space-time can be generalized if the original Hamiltonian constraint is modified such that the hypersurface-deformation brackets and the covariance condition are preserved. A theory of emergent modified gravity is then defined by a Hamiltonian constraint  $\tilde{H}^{\text{grav}}$  such that

$$\{H_x[N^x], H_x[M^x]\} = -H_x[M^x(N^x)' - N^x(M^x)'] \quad (23)$$

$$\{\tilde{H}[N], H_x[M^x]\} = -\tilde{H}[M^x N'] \quad (24)$$

$$\{\tilde{H}[N], \tilde{H}[M]\} = -H_x[\tilde{q}^{xx}(MN' - NM')] \quad (25)$$

with a phase-space function  $\tilde{q}^{xx}$  for which

$$\delta_\epsilon \tilde{g}_{\mu\nu}|_{\text{O.S.}} = \mathcal{L}_\xi \tilde{g}_{\mu\nu}|_{\text{O.S.}} \quad (26)$$

holds with a space-time metric  $\tilde{g}_{\mu\nu}$  derived from the emergent line element

$$ds^2 = -N^2 dt^2 + \tilde{q}_{xx}(dx + N^x dt)^2 + E^x d\Omega^2 \quad (27)$$

that makes use of the inverse  $\tilde{q}_{xx}$  of the new structure function  $\tilde{q}^{xx}$ . In contrast to classical general relativity,  $\tilde{q}^{xx}$  need not be defined a priori in terms of the fundamental fields. It must be derived from hypersurface-deformation brackets and the covariance condition and is therefore emergent. As the new structure function, the emergent spatial metric then appears in the gauge transformations

$$\delta_\epsilon N = \epsilon^0 + \epsilon^x N' - N^x (\epsilon^0)', \quad (28)$$

$$\delta_\epsilon N^x = \epsilon^x + \epsilon^x (N^x)' - N^x (\epsilon^x)' + \tilde{q}^{xx} (\epsilon^0 N' - N (\epsilon^0)') \quad (29)$$

of lapse and shift.

Hamiltonian constraints compatible with these conditions have been evaluated systematically in a derivative expansion that stops at the classical order of at most two spatial derivatives in each term. Even with this classical order, non-trivial modifications are possible because emergence of the metric is a new property that turns out to relax several implicit assumptions in action-based constructions of modified gravity. In the latter cases, a space-time volume element must be known before deriving equations of motion because it is required for the action integral.

In a canonical setting, the Hamiltonian constraint of a given theory is not uniquely defined but may always be subjected to canonical transformations. In contrast to classical gravity, there is no distinguished set of phase-space coordinates because it is not clear from the outset which functions will determine the space-time geometry. The freedom of performing canonical transformations must therefore be explored in a systematic derivation of modified canonical theories. In spherically symmetric models of emergent modified gravity up to second order in spatial derivatives, it turns out that one can fix the freedom of canonical transformations by requiring that the Hamiltonian constraint is strictly periodic in the momentum  $K_\varphi$ , which then appears with a constant coefficient  $\bar{\lambda}$  in trigonometric functions [4]. The set of covariant Hamiltonian constraints of this form is given by

$$\begin{aligned} \tilde{H}^{\text{grav}} = & -\bar{\lambda}_0 \frac{\sqrt{E^x}}{2} \left[ E^\varphi \left( -\bar{\lambda}_0 + \frac{\bar{\alpha}_0}{E^x} + 2 \frac{\sin^2(\bar{\lambda} K_\varphi)}{\bar{\lambda}^2} \frac{\partial c_f}{\partial E^x} + 4 \frac{\sin(2\bar{\lambda} K_\varphi)}{2\bar{\lambda}} \frac{\partial \bar{q}}{\partial E^x} \right. \right. \\ & + \frac{\bar{\alpha}_2}{E^x} \left( c_f \frac{\sin^2(\bar{\lambda} K_\varphi)}{\bar{\lambda}^2} + 2\bar{q} \frac{\sin(2\bar{\lambda} K_\varphi)}{2\bar{\lambda}} \right) \left. \right) + 4K_x \left( c_f \frac{\sin(2\bar{\lambda} K_\varphi)}{2\bar{\lambda}} + \bar{q} \cos(2\bar{\lambda} K_\varphi) \right) \\ & + \frac{((E^x)')^2}{E^\varphi} \left( -\frac{\bar{\alpha}_2}{4E^x} \cos^2(\bar{\lambda} K_\varphi) + \frac{K_x}{E^\varphi} \bar{\lambda}^2 \frac{\sin(2\bar{\lambda} K_\varphi)}{2\bar{\lambda}} \right) \\ & \left. + \left( \frac{(E^x)'(E^\varphi)'}{(E^\varphi)^2} - \frac{(E^x)''}{E^\varphi} \right) \cos^2(\bar{\lambda} K_\varphi) \right] \quad (30) \end{aligned}$$

such that the structure function equals

$$\tilde{q}^{xx} = \left( \left( c_f + \left( \frac{\bar{\lambda}(E^x)'}{2E^\varphi} \right)^2 \right) \cos^2(\bar{\lambda} K_\varphi) - 2\bar{q} \bar{\lambda}^2 \frac{\sin(2\bar{\lambda} K_\varphi)}{2\bar{\lambda}} \right) \bar{\lambda}_0^2 \frac{E^x}{(E^\varphi)^2}. \quad (31)$$



The parameter  $\bar{\lambda}$  is constant, while  $\bar{\lambda}_0$ ,  $\bar{\Lambda}_0$ ,  $\bar{\alpha}_0$ ,  $\bar{\alpha}_2$ ,  $c_f$  and  $\bar{q}$  are free functions of  $E^x$  that define a specific covariant modification of spherically symmetric general relativity. The functions  $\bar{\Lambda}_0$  and  $\bar{\alpha}_0$  can be combined to a dilaton potential and may also be realized in an action principle. There is no known action principle for the general set of modifications. Compared with non-covariant studies of LTB models with modifications from loop quantum gravity, the last two lines of (30) are new and crucial for covariance.

This form is useful as a mathematical condition to fix the freedom of gauge transformations, and also makes it possible to realize the Hamiltonian constraint as an operator in loop quantum gravity where periodic functions of  $K_\varphi$  can be related to basic holonomy operators. However, it is not always the best version in phenomenological evaluations of the theory. For this purpose, one may apply a canonical transformation to new phase-space variables such that  $\bar{\lambda}K_\varphi$  is replaced by  $\lambda(E^x)K_\varphi$  with a scale-dependent function  $\lambda(E^x)$ . A canonical transformation then transforms  $E^\varphi$  to  $(\bar{\lambda}/\lambda(E^x))E^\varphi$ , while  $E^x$  is unchanged and  $K_x$  is mapped to a specific linear combination of  $K_x$  and  $K_\varphi$  with triad-dependent coefficients. In this form, the nonperiodic modified constraint is

$$\begin{aligned}
\tilde{H}^{\text{grav}} = & -\sqrt{E^x} \frac{\lambda_0}{2} \left[ E^\varphi \left( -\Lambda_0 + \frac{\alpha_0}{E^x} + 2 \frac{\sin^2(\lambda K_\varphi)}{\lambda^2} \frac{\partial c_f}{\partial E^x} + 4 \frac{\sin(2\lambda K_\varphi)}{2\lambda} \frac{1}{\lambda} \frac{\partial(\lambda q)}{\partial E^x} \right. \right. \\
& + \left( \frac{\alpha_2}{E^x} - 2 \frac{\partial \ln \lambda^2}{\partial E^x} \right) \left( c_f \frac{\sin^2(\lambda K_\varphi)}{\lambda^2} + 2q \frac{\sin(2\lambda K_\varphi)}{2\lambda} \right) \\
& + 4 \left( \frac{K_x}{E^\varphi} + \frac{K_\varphi}{2} \frac{\partial \ln \lambda^2}{\partial E^x} \right) \left( c_f \frac{\sin(2\lambda K_\varphi)}{2\lambda} + q \cos(2\lambda K_\varphi) \right) \\
& + \frac{((E^x)')^2}{E^\varphi} \left( -\frac{\alpha_2}{4E^x} \cos^2(\lambda K_\varphi) + \left( \frac{K_x}{E^\varphi} + \frac{K_\varphi}{2} \frac{\partial \ln \lambda^2}{\partial E^x} \right) \lambda^2 \frac{\sin(2\lambda K_\varphi)}{2\lambda} \right) \\
& \left. + \left( \frac{(E^x)'(E^\varphi)'}{(E^\varphi)^2} - \frac{(E^x)''}{E^\varphi} \right) \cos^2(\lambda K_\varphi) \right] \quad (32)
\end{aligned}$$

with the associated structure function

$$\tilde{q}^{xx} = \left( \left( c_f + \lambda^2 \left( \frac{(E^x)'}{2E^\varphi} \right)^2 \right) \cos^2(\lambda K_\varphi) - 2q \lambda^2 \frac{\sin(2\lambda K_\varphi)}{2\lambda} \right) \lambda_0^2 \frac{E^x}{(E^\varphi)^2}. \quad (33)$$

The remaining modification functions are related by

$$\lambda_0 = \bar{\lambda}_0 \frac{\lambda}{\bar{\lambda}}, \quad q = \bar{q} \frac{\bar{\lambda}}{\lambda}, \quad \Lambda_0 = \frac{\bar{\lambda}^2}{\lambda^2} \bar{\Lambda}_0, \quad \alpha_0 = \frac{\bar{\lambda}^2}{\lambda^2} \bar{\alpha}_0, \quad \alpha_2 = \bar{\alpha}_2 + 4E^x \frac{\partial \ln \lambda}{\partial E^x}. \quad (34)$$

The constraint (30) has the weak observable

$$\begin{aligned}
\mathcal{M} = & d_0 + \frac{d_2}{2} \left( \exp \int dE^x \frac{\bar{\alpha}_2}{2E^x} \right) \left( c_f \frac{\sin^2(\bar{\lambda} K_\varphi)}{\bar{\lambda}^2} + 2\bar{q} \frac{\sin(2\bar{\lambda} K_\varphi)}{2\bar{\lambda}} - \cos^2(\bar{\lambda} K_\varphi) \left( \frac{(E^x)'}{2E^\varphi} \right)^2 \right) \\
& + \frac{d_2}{4} \int dE^x \left( \left( \bar{\Lambda}_0 + \frac{\bar{\alpha}_0}{E^x} \right) \exp \int dE^x \frac{\bar{\alpha}_2}{2E^x} \right) \quad (35)
\end{aligned}$$

which can be transformed canonically to the observable relevant for (32). These expressions can be related to the mass and are therefore useful in the context of LTB collapse.

Including perfect-fluid degrees of freedom, the general form of the matter contribution to the Hamiltonian constraint is given by [6]

$$\tilde{H}_s^{\text{matter}} = \sqrt{sP_T^2 + \tilde{q}^{xx}(H_x^{\text{matter}})^2} - \sqrt{\det \tilde{q}} P_q(E^x, P_X^2/P_T^2) - 4\pi\sqrt{E^x} E^\varphi \bar{\lambda}_0 P_0(E^x, P_X^2/P_T^2) \quad (36)$$

with the classical contribution (12) to the diffeomorphism constraint. In [6] only the pressure  $P_q$  was introduced, but it is straightforward to check that the alternative pressure function  $P_0$  preserves all the symmetry requirements to be imposed. A perfect fluid may therefore couple to the emergent spatial volume element,  $\sqrt{\tilde{q}}$ , or the classical volume element  $\sqrt{E^x} E^\varphi$ . This modified constraint is obtained from requiring anomaly freedom, normalization of the fluid's velocity co-vector

$$u_\mu dx^\mu = - \left( \dot{T} + \frac{P_X}{P_T} \dot{X} \right) dt - \left( T' + \frac{P_X}{P_T} X' \right) dx, \quad (37)$$

derived in (123), and the matter covariance condition

$$\delta_\epsilon u_\mu|_{\text{O.S.}} = \mathcal{L}_\xi u_\mu|_{\text{O.S.}}. \quad (38)$$

As shown in [10, 11, 12] for constant  $\lambda$  and in [16] for general  $\lambda$  in the vacuum case, a non-zero choice for this modification function implies that the classical singularity of spherically symmetric solutions is replaced by a transition between a black hole and a white hole. These two space-time regions are connected at a maximum-curvature hypersurface defined by  $\lambda K_\varphi = \pi/2$  at a local maximum of  $\sin(\lambda K_\varphi)$ . While  $K_\varphi$  could take greater values, this phase-space function no longer represents extrinsic curvature in a modified theory. Instead, the space-time curvature of the resulting emergent metric shows that a maximum is reached at the given value of  $\lambda K_\varphi$ .

This result has been extended to non-vacuum solutions for dust [6], electromagnetic fields [12, 8] and scalar fields [7]. In these cases, however, singularity resolution is not generic and therefore appears to be unstable under the inclusion of matter. In particular, the non-singular vacuum model may be used as a background for matter in a version of minimal coupling [9], but it re-introduces a physical singularity in collapse models [7]. (Additional modification functions in matter terms can be considered that may be preferred for various physical reasons, as discussed in detail in [7]. They can help to recover the nonsingular behavior, but their equations of motion differ from minimal coupling.) By extension, models of loop quantum gravity are also subject to the caveat that covariant matter couplings may undo singularity resolution observed in vacuum models. These results provide strong motivation for a detailed evaluation of LTB models in emergent modified gravity.

### 2.3 LTB conditions

Because the LTB conditions are phenomenological and not fundamental, since they merely restrict attention to a convenient set of solutions, there is no systematic derivation of new

conditions in a modified theory. The classical motivation for the specific condition relies on the resulting line element, but in emergent modified gravity (or any covariant model of loop quantum gravity), the complete space-time metric in terms of phase-space variables is available only after constraints and equations of motion have been analyzed in order to impose covariance. This feature results in a new ambiguity because the phase-space form of the LTB condition is no longer equivalent to the space-time form derived from the line element. In particular, if we maintain the phase-space condition (9), the resulting space-time solutions will not have a simple radial component given by  $(R')^2$ , but also contain an additional factor from  $\tilde{q}_{xx}$  that is in general not constant. Alternatively, one may try to derive a condition such that the new  $\tilde{q}_{xx}$  is of the simple form  $(R')^2$  (in the marginal case of  $\kappa(x) = 0$ ), which would then require a modified time dependence of  $R$  derived from the resulting equations of motion. However, this procedure would not only be more complicated, it might also be too restrictive because it is not guaranteed that such solutions exist for sufficiently general modification functions.

An additional subtlety is that the phase-space version of the LTB condition depends on the canonical variables in which the theory is evaluated. For instance, if we use the classical phase-space condition  $L_1$  with a choice of phase-space variables that implies the non-periodic version of the Hamiltonian constraint, the resulting solutions correspond to a periodic theory in which the transformed LTB condition

$$\tilde{L}_1 = \frac{\lambda(E^x)}{\bar{\lambda}} E^\varphi - \frac{(E^x)'}{2} \quad (39)$$

has been used. These two theories are equivalent, but using the periodic version with  $\tilde{L}_1$  is not the same as using the periodic version with the original  $L_1$ . We are not aware of any physical condition that could relate the choice of phase-space variables to a specific phase-space expression for  $L_1$ . Again, the specific implementation of LTB solutions within a covariant modified theory is mainly a matter of convenience.

Here, we maintain the LTB gauge (7) as well as the phase-space version (9) of the LTB condition in the non-periodic version of the modified Hamiltonian constraint, working with the original function  $L_1$ . This choice is appropriate because the non-periodic constraint and solutions in LTB form are motivated mainly by phenomenological rather than fundamental reasoning. We then have to re-evaluate the secondary condition that  $L_1$  be preserved under modified evolution. This condition may well place restrictions on the modification functions of the new constraint because there could be covariant modified theories that do not admit solutions in LTB form. The resulting equations become very complicated in the modified theory for non-trivial  $\lambda$ . However, the marginally bound case  $\kappa(x) = 0$  is tractable and considerable analytic progress is possible. We will therefore restrict our attention to this case in the following.

If we use the unmodified LTB condition (9), the secondary condition  $\tilde{L}_2 = \dot{L}_1 = \{L_1, \tilde{H}^{\text{grav}}[N] + H_x^{\text{grav}}[N^x]\}$  takes the form

$$\tilde{L}_2|_{\text{O.S., } N=1, N^x=0, L_1=0} = \gamma \left( \alpha_2 - 1 - 2E^x \frac{\partial \ln \lambda_0}{\partial E^x} \right), \quad (40)$$

where we have used the nonperiodic version of the Hamiltonian constraint in vacuum, given by (32), and defined

$$\gamma = \frac{\lambda_0(E^x)'}{2\sqrt{|E^x|}} \left( (c_f + \lambda^2) \frac{\sin(2\lambda K_\varphi)}{2\lambda} + 2q \cos(2\lambda K_\varphi) \right). \quad (41)$$

The gamma factor has the non-vanishing classical limit  $\gamma \rightarrow (E^x)'K_\varphi/(2\sqrt{|E^x|})$  and therefore the secondary condition implies the equation

$$\alpha_2 - 1 - 2E^x \frac{\partial \ln \lambda_0}{\partial E^x} = 0. \quad (42)$$

Thus, there is indeed a requirement on the modification functions for a consistent LTB form to exist. This condition can be solved for

$$\alpha_2 = 1 + 2E^x \frac{\partial \ln \lambda_0}{\partial E^x}, \quad (43)$$

which has the correct classical limit  $\alpha_2 \rightarrow 1$  as  $\partial \lambda_0 / \partial E^x \rightarrow 0$ . If this equation for the modification functions is satisfied, which would be the case, for instance, if we use the classical values  $\lambda_0 = 1 = \alpha_2$ , then the LTB condition is consistent with the equations of motion of the modified theory in vacuum too. There is no restriction on the other modification functions  $\lambda$ ,  $c_f$  and  $q$  that appear in the emergent metric.

As expected, implementing the LTB condition as a phase-space function does not produce a metric of the classical LTB form because space-time geometry is determined by a line element with the emergent metric (27). With the LTB condition, the emergent line element (33) simplifies slightly to

$$\tilde{q}^{xx} = \left( (c_f + \lambda^2) \cos^2(\lambda K_\varphi) - 2q\lambda^2 \frac{\sin(2\lambda K_\varphi)}{2\lambda} \right) \lambda_0^2 \frac{E^x}{(E^\varphi)^2}. \quad (44)$$

If all modification functions other than  $\lambda$  and  $\lambda_0$  take their classical values, the non-periodic version of the mass observable (35) can be used to solve for  $\cos^2(\lambda K_\varphi)$ , resulting in the line element

$$ds^2 = -dt^2 + \left( 1 + \lambda^2 \left( 1 - \frac{2\mathcal{M}}{R} \right) \right)^{-1} \frac{(R')^2}{\lambda_0^2} dx^2 + R^2 d\Omega^2. \quad (45)$$

We repeat the procedure described above, but this time, we include the perfect-fluid contributions. Taking the LTB gauge (7) and keeping an unmodified LTB condition (9), the secondary condition  $\tilde{L}_2 = \dot{L}_1 = \{L_1, \tilde{H}^{\text{grav}}[N] + \tilde{H}_s^{\text{matter}}[N] + H_x^{\text{grav}}[N^x] + H_x^{\text{matter}}[N^x]\}$  now takes the form

$$\tilde{L}_2|_{\text{O.S., } N=1, N^x=0, L_1=0} = \alpha H_x^{\text{matter}} + \beta P_q + \gamma (\alpha_2 - 1 - 2E^x (\ln \lambda_0)'). \quad (46)$$

where  $\gamma$  is again given by the expression (41), and

$$\beta = 4\pi \lambda_0 \gamma \sqrt{|E^x|} \frac{(E^x)^2}{(E^\varphi)^3} \lambda^4 (\tilde{q}_{xx})^{3/2}, \quad (47)$$

while  $\alpha$  is a more complicated expression. The secondary condition now requires the same vacuum condition (43), but also a vanishing pressure function  $P_q = 0$ . The remaining factor then simplifies to

$$\alpha = \frac{(E^x)^{3/2}}{\sqrt{E^x} \lambda_0 (E^\varphi)^2} \left( \left( \lambda^2 \lambda_0^3 q + \lambda^2 \bar{\lambda}_0 \tilde{q}^{xx} \frac{(E^\varphi)^2 \tan(\lambda K_\varphi)}{E^x} \frac{\lambda}{\lambda} \right) \frac{H_x^{\text{matter}}}{\tilde{H}^{\text{dust}}} + \tilde{q}^{xx} \left( \frac{(E^\varphi)^2}{E^x} \right)^{3/2} \frac{\cos(2\lambda K_\varphi)}{\cos^2(\lambda K_\varphi)} - 2\lambda^2 \lambda_0^2 q \frac{E^\varphi \tan(\lambda K_\varphi)}{\sqrt{E^x} \lambda} \right) \quad (48)$$

where

$$\tilde{H}^{\text{dust}} = \sqrt{s P_T^2 + \tilde{q}^{xx} (H_x^{\text{matter}})^2}. \quad (49)$$

Because  $\alpha$  does not vanish in the classical limit, the secondary condition is again satisfied only if  $H_x^{\text{matter}} = 0$ , which means that we must work in a slicing adapted to the fluid's frame and continue to use  $s = 1$ .

The energy density of the perfect fluid in the fluid's frame is given by

$$\rho = \frac{P_T}{\sqrt{\det \tilde{q}}}. \quad (50)$$

In LTB coordinates, we may define

$$\rho = \frac{1}{4\pi R^2} \frac{dm}{dR} = \frac{1}{4\pi R^2} \frac{m'}{R'}. \quad (51)$$

with some function  $m(x, t)$  we refer to as the mass parameter. Combining (50) and (51), we obtain

$$P_T = \lambda_0 \sqrt{1 + \lambda^2 \left( 1 - \frac{2\mathcal{M}}{R} \right) m'} \quad (52)$$

using  $\tilde{q}$  from (45).

### 3 Modified marginal LTB collapse

We are now ready to apply the general theory of emergent modified gravity in marginal LTB form to the problem of gravitational collapse. As a summary, the marginally bound LTB condition

$$E^\varphi = \frac{(E^x)'}{2} \quad (53)$$

in the gauge

$$N = 1, \quad N^x = 0, \quad (54)$$

can be consistently implemented in the modified theory provided the following secondary conditions are applied:

$$\alpha_2 = 1 + 2E^x \frac{\partial \ln \lambda_0}{\partial E^x} \quad (55)$$

for two of the gravitational modification functions,

$$P_q = 0 \tag{56}$$

for one of the possible pressure functions, and

$$P_X = \frac{P_T T'}{X'} \tag{57}$$

from  $H_x^{\text{matter}} = 0$  as in the classical case.

In the following we will take the classical values  $c_f = \alpha_0 = \alpha_2 = 1$ ,  $q = 0$ , and  $\Lambda_0 = \Lambda$ , and further assume a vanishing cosmological constant and pressure,  $\Lambda = 0$  and  $P_0 = 0$ , in order to study the collapse of dust in asymptotically flat space-times. Using these classical values, the condition (43) is satisfied for any constant  $\lambda_0 \neq 0$ , which we will keep in the following equations. Furthermore, relabeling  $E^x = R^2$  and imposing the LTB condition, the emergent metric (45) takes the form

$$ds^2 = -dt^2 + \left(1 + \lambda^2 \left(1 - \frac{2\mathcal{M}}{R}\right)\right)^{-1} \frac{(R')^2}{\lambda_0^2} dx^2 + R^2 d\Omega^2, \tag{58}$$

with  $\mathcal{M}$  being the vacuum mass observable (35) now taking the form

$$\mathcal{M} = \frac{R}{2} \frac{1 + \lambda^2}{\lambda^2} \sin^2(\lambda K_\varphi) \tag{59}$$

as a phase-space function.

The on-shell condition  $H_x = 0$  gives

$$K_x = \frac{K'_\varphi}{2}, \tag{60}$$

while  $\tilde{H} = 0$  implies

$$m' = \mathcal{M}', \tag{61}$$

where we have substituted  $P_T = \lambda_0 m'$ . We can therefore directly integrate (61), obtaining

$$m = \mathcal{M} + m_0(t), \tag{62}$$

where  $m_0$  is the constant of integration that we may set to zero such that  $m = \mathcal{M}$ . However, it is the latter that enters the space-time metric and therefore the actual value of  $m_0$  is not relevant for most purposes. Furthermore, it can be shown that

$$\dot{\mathcal{M}} = 0, \tag{63}$$

and hence the mass function is time independent in this gauge.

### 3.1 Equations of motion

The relevant equations of motion are

$$\dot{R} = \lambda_0(1 + \lambda^2) \frac{\sin(2\lambda K_\varphi)}{2\lambda}, \quad (64)$$

and

$$\begin{aligned} \left( \lambda_0(1 + \lambda^2) \frac{\sin(2\lambda K_\varphi)}{2\lambda} \right)^\bullet &= -\frac{\lambda_0^2(1 + \lambda^2)}{2R} \frac{\sin^2(\lambda K_\varphi)}{\lambda^2} \\ &\times \left( (1 + \lambda^2) \left( 1 - R \frac{\partial \ln \lambda}{\partial R} \right) \cos(2\lambda K_\varphi) - (\lambda^2 - 1) R \frac{\partial \ln \lambda}{\partial R} \right) \end{aligned} \quad (65)$$

for the gravitational variables, and

$$\dot{X} = 0 \quad , \quad \dot{T} = \text{sgn}(P_T) + \frac{T'}{X'} \dot{X} \quad , \quad \dot{P}_T = 0 \quad (66)$$

for the dust degrees of freedom, to which  $P_X$  is strictly related by (57).

The first two equations can be rewritten in terms of the mass function and  $R$ :

$$\dot{R} = s_{\sin} \lambda_0 \sqrt{\frac{2\mathcal{M}}{R} \left( 1 + \lambda^2 \left( 1 - \frac{2\mathcal{M}}{R} \right) \right)} \quad (67)$$

$$\ddot{R} = -\frac{\lambda_0^2 \mathcal{M}}{R^2} \left( 1 + \lambda^2 \left( 1 - \frac{4\mathcal{M}}{R} \right) - (R - 2\mathcal{M}) \frac{\partial \lambda^2}{\partial R} \right) \quad (68)$$

where  $s_{\sin} = \text{sgn}(\sin(2\lambda K_\varphi))$ . Taking a time derivative of (67) simply gives (68), and therefore only the former needs to be solved.

The Friedmann-like equation stemming from (67) is

$$\left( \frac{\dot{R}}{R} \right)^2 = \lambda_0^2 \left( \frac{2\mathcal{M}}{R^3} \left( 1 + \lambda^2 \left( 1 - \frac{2\mathcal{M}}{R} \right) \right) \right). \quad (69)$$

This result can be compared with the restriction to the marginal case of an equation derived in [19], which used a constant  $\Delta$  that, by comparing the equations, would be related to our function  $\lambda$  by

$$\lambda^2 = -\frac{2\mathcal{M}\Delta/R^3}{1 - 2\mathcal{M}/R} \quad (70)$$

provided  $\lambda_0 = 1$ . While the Friedmann-like equations might seem equivalent in these two cases, the  $\Delta$ -version would require a specific  $\lambda$ -function that does not agree with the form of holonomy modifications that motivated the original terms in [19]. The reason for this discrepancy can be seen in the fact that this reference did not implement covariance conditions. Furthermore,  $\lambda$  is a function of only  $E^x$  according to the covariance conditions for a constraint restricted to at most second-order spatial derivatives, while  $\mathcal{M}$  is a complicated phase space function of  $K_\varphi$  and  $E^\varphi$  and  $(E^x)'$ . (It is a simple function of  $x$  only on a given solution but not within a modified theory obtained after implementing the  $\Delta$ -terms.) The choice discussed in [19] therefore is not related to a valid covariant formulation.

### 3.2 Reflection-symmetry hypersurface

As discussed in [6, 16], the full constraints  $\tilde{H} = \tilde{H}^{\text{grav}} + \tilde{H}_s^{\text{matter}}$  and  $H_x = H_x^{\text{grav}} + H_x^{\text{matter}}$  are symmetric under the time reversal operation  $K_\varphi \rightarrow -K_\varphi$ ,  $K_x \rightarrow -K_x$ ,  $T \rightarrow -T$ ,  $P_X \rightarrow -P_X$  provided  $q = 0$ , which we assume here. Furthermore, the system possesses a similar symmetry at the phase-space hypersurface defined by  $K_\varphi = -\pi/(2\lambda)$  and  $K_x = T = P_X = 0$ : defining  $K_\varphi = -\pi/(2\lambda) - \delta$ , the Hamiltonian is symmetric under the operation  $\delta \rightarrow -\delta$ ,  $K_x \rightarrow -K_x$ ,  $T \rightarrow -T$ ,  $P_X \rightarrow -P_X$ . This transformation is a reflection symmetry in the sense that a given solution to the equations of motion in the region  $\delta < 0$  can also be used to describe a solution in the region  $\delta > 0$  by a simple reflection. By itself, this argument does not imply that a complete solution around the maximum-curvature hypersurface is strictly reflection symmetric; it only implies that the future part of the solution is the time reversal of some collapsing solution within the complete solutions space. However, in what follows we demonstrate by a detailed analysis of the specific equations to be solved that every single solution must be reflection symmetric.

Given its similarity to time-reversal symmetry, the solution in the region  $\delta > 0$  will look like a time-reversed solution of the region  $\delta < 0$  but it does describe a different region joined at the reflection-symmetry hypersurface. The identification and understanding of this symmetry will be useful in obtaining the global solution since our task is simplified by requiring only the dynamical solution on one side of the reflection symmetry hypersurface.

### 3.3 Generic formation of a singularity

Using (59), we find that at the maximum-curvature hypersurface given by  $K_\varphi = -\pi/(2\lambda)$  in space-time we have

$$R = 2\mathcal{M} \frac{\lambda^2}{1 + \lambda^2} \quad (71)$$

as a function of  $x$ . This hypersurface replaces  $R = 0$  in the classical theory, which is obtained in the limit of  $\lambda \rightarrow 0$ . Equation (59) also implies that, unless  $\lambda(R) = R/(c - R)$  with a constant  $c$  which would not be a valid large- $R$  behavior, decreasing  $R$  implies increasing  $\sin^2(\lambda K_\varphi)$ , and vice versa. If we extend  $K_\varphi$  across the value  $K_\varphi = -\pi/(2\lambda)$ , where the sine squared has a local maximum, we may use this variable as a local time coordinate around the reflection-symmetry hypersurface. Since we can simply reparametrize time along the trajectories of dust, this coordinate change does not affect the spatial coordinate  $x$ .

Therefore, we can take a spatial derivative of (71) at constant  $K_\varphi$ :

$$R' = \left( \frac{1 + \lambda^2}{2\lambda^2} - \frac{R}{2} \frac{\partial \ln \lambda^2}{\partial R} \right)^{-1} \mathcal{M}'. \quad (72)$$

This expression is non-zero for a non-constant mass profile, such that we avoid shell-crossing singularities. Equations (67) and (68) imply the time derivatives  $\dot{R} = 0$  and

$$\ddot{R} = \frac{\lambda_0^2 \mathcal{M}}{R^2} \left( 1 + \lambda^2 - 2\lambda^2 R \frac{\partial \ln \lambda^2}{\partial R} \right) \quad (73)$$



which allow us to compute the value of the Ricci scalar of the space-time metric (58) at the maximum-curvature hypersurface:

$$\begin{aligned}
\mathcal{R} = \Omega & \left[ \frac{\partial \lambda}{\partial R} \left[ -4\lambda^5 \mathcal{M}^3 R (\lambda^2 - 7) + 2\lambda^3 \mathcal{M}^2 R^2 (4\lambda^4 + 9\lambda^2 + 1) \right. \right. \\
& \left. \left. - \lambda \mathcal{M} R^2 (\lambda^2 + 1) (3\lambda^4 + 17\lambda^2 + 6) + 2\lambda R^4 (\lambda^2 + 1) \right] \right. \\
& \left. - 2\lambda^4 \mathcal{M} R^2 (R - 8\mathcal{M}) (R - 2\mathcal{M}) \left( \frac{\partial \lambda}{\partial R} \right)^2 - 2\lambda^2 \mathcal{M} R^3 (R - 6\mathcal{M}) \left( \frac{\partial \lambda}{\partial R} \right)^2 \right. \\
& \left. + (\lambda^2 + 1) \left[ 4\lambda^6 \mathcal{M}^3 - 19\lambda^4 \mathcal{M}^2 R (\lambda^2 + 1) + 2\lambda^2 \mathcal{M} R^2 (4\lambda^4 + 9\lambda^2 + 5) \right. \right. \\
& \left. \left. + 2\lambda^3 \mathcal{M} R^2 (2\mathcal{M} - R) (2\lambda^2 \mathcal{M} - (\lambda^2 + 1) R) \frac{\partial^2 \lambda}{(\partial R)^2} - (\lambda^2 + 1)^2 R^3 \right] + \frac{R^2}{\lambda_0^2} \right]
\end{aligned} \tag{74}$$

where

$$\Omega = \frac{2\lambda_0^2}{\lambda^2 (1 + \lambda^2) R^4 (R + \lambda^2 (R - 2\mathcal{M}))} \tag{75}$$

This expression is generically divergent at the maximum-curvature hypersurface for any  $\lambda$  because the overall denominator in the last term diverges,  $(R + \lambda^2(R - 2\mathcal{M}))^{-1} \rightarrow \infty$  according to (71).

Following the discussion of Subsection 3.2, the maximum-curvature hypersurface corresponds to the hypersurface of reflection symmetry. Therefore, the region beyond this hypersurface will be described by a family of reflected collapse solutions, one for each particle trajectory (at least locally, until a shell-crossing singularity is reached). Since the collapse of  $R$  is governed by a single first-order differential equation in time along a dust particle trajectory and the final value of collapse agrees with the initial value of subsequent expansion, a single solution is strictly reflection symmetric with expansion given precisely by the time reversal of collapse along the same trajectory. The global space-time can then be understood as a collapsing phase that bounces into an expanding phase, both joined at the singular maximum-curvature hypersurface defined by (71). This observation rules out the formation of shock waves if there are no shock profiles present in the initial data of collapse in a region around high curvature. (It is possible that shock waves may form at lower curvature before the region around reflection symmetry is reached. However, such effects would be unrelated to a potential bouncing property of the interior black-hole region.) While the collapsing phase describes the final stage of a black hole, the expanding phase describes the formation of a white hole. Unlike in vacuum solutions, these two regions are not causally connected but rather appear on two sides of a physical singularity. They may be connected only if one goes beyond an effective description and uses a microscopic theory of quantum space-time. However, possible results of such a transition are beyond the capabilities of current modeling in loop quantum gravity.

For a given space-time solution, it is possible to eliminate the singularity by choosing

a specific modification function

$$\lambda_0 \propto \sqrt{1 + \lambda^2(1 - 2\mathcal{M}/R)} \quad (76)$$

as suggested in [15]. (Alternatively, this choice could be implemented by applying a conformal transformation of the entire emergent metric by  $\omega^2 = 1 + \lambda^2(1 - 2\mathcal{M}/R)$  [14], but a conformal transformation changes the underlying physics if the new solution is interpreted in the observer's frame; see also [29] and App. B for additional details.) According to (43), a non-trivial choice would then be required for  $\alpha_2$ . While there may be a covariant modified theory that renders solutions with a given  $\mathcal{M}$  non-singular, solutions with other mass distributions would remain singular. It is not possible to use an  $\mathcal{M}$ -dependent (and thus solution-dependent) modification function  $\lambda_0$  in a single modified theory. Using the expression (59), the  $\mathcal{M}$ -dependence can be turned into a specific phase-space dependence on  $R$  and  $K_\varphi$  on the entire subspace of LTB solutions. However, if the LTB condition is not used, considering the full modified theory,  $\mathcal{M}$  depends also on  $(E^x)'$ , such that the required  $\lambda_0$  has a non-polynomial dependence on spatial derivatives of  $E^x$ . From the perspective of effective field theory, derivations within such a model would not be considered reliable because the remaining terms in the modified Hamiltonian constraint have been restricted to at most second order in spatial derivatives. It is then not clear how potential higher spatial derivatives in the other terms of the Hamiltonian constraint could affect properties of solutions.

We conclude that the covariant collapse of dust with holonomy modifications as suggested by loop quantum gravity is generically singular for solutions in marginal LTB form, which is the same conclusion that was reached in [6] for Oppenheimer-Snyder collapse.

### 3.4 Collapse in Gullstrand-Painlevé coordinates and causal structure

Even though solutions are generically singular at the maximum-curvature hypersurface, it is instructive to compare their causal structure to classical collapse. We have two cases of interest: 1) The simple case of constant holonomy parameter  $\lambda = \bar{\lambda}$ , and 2) the scale-dependent holonomy  $\lambda(E^x) = \sqrt{\Delta/E^x}$ . As already mentioned, it is possible to use a canonical transformation in order to map the  $K_\varphi$ -dependence of the Hamiltonian constraint with constant  $\bar{\lambda}$  to an equivalent version with non-constant  $\lambda(E^x)$ . However, this transformation, in general, changes the other modification functions according to (34); see [18] for a detailed discussion. If we make different choices for  $\lambda$  while keeping the other modification functions unchanged, we obtain physically inequivalent systems.

We will first transform the LTB gauge of space-time solutions with  $N^x = 0$  to a Gullstrand-Painlevé (GP) gauge which is useful for studying collapse across the point of horizon formation. The relevant equations can be derived for generic  $\lambda$  and will then be specialized and solved in the two main cases of interest.

To this end, it is convenient to choose  $R$  as our new radial coordinate, such that

$$dR = R'dx + \dot{R}dt$$

$$= R' dx + s_{\text{sin}} \lambda_0 \sqrt{\frac{2\mathcal{M}}{R} \left(1 + \lambda^2 \left(1 - \frac{2\mathcal{M}}{R}\right)\right)} dt \quad (77)$$

can be solved for

$$dx = \frac{dR}{R'} - s_{\text{sin}} \lambda_0 \sqrt{\frac{2\mathcal{M}}{R} \left(1 + \lambda^2 \left(1 - \frac{2\mathcal{M}}{R}\right)\right)} \frac{dt}{R'}. \quad (78)$$

The line element then takes the form

$$\begin{aligned} ds^2 &= -dt^2 + R^2 d\Omega^2 \\ &+ \lambda_0^{-2} \left(1 + \lambda^2 \left(1 - \frac{2\mathcal{M}}{R}\right)\right)^{-1} \left(dR - s_{\text{sin}} \lambda_0 \sqrt{\frac{2\mathcal{M}}{R} \left(1 + \lambda^2 \left(1 - \frac{2\mathcal{M}}{R}\right)\right)} dt\right)^2. \end{aligned} \quad (79)$$

In these coordinates we have a time-dependent mass distribution,  $\mathcal{M}(x) = \mathcal{M}(x(t, R))$ .

Gullstrand-Painlevé coordinates can be derived by using properties of infalling light rays. The lightlike condition on the tangent vector implies the velocity

$$\left. \frac{dR}{dt} \right|_{\text{null}} = \lambda_0 \sqrt{1 + \lambda^2 \left(1 - \frac{2\mathcal{M}}{R}\right)} \left(s_n + s_{\text{sin}} \sqrt{\frac{2\mathcal{M}}{R}}\right), \quad (80)$$

where  $s_n = \pm 1$  is the sign of the square root with  $s_n = 1$  being the outgoing ray and  $s_n = -1$  the ingoing ray. During the collapse stage we have  $s_{\text{sin}} = -1$  and hence the ingoing ray always has a negative radial velocity, whereas the outgoing ray has a vanishing velocity at the coordinates solving the equation

$$R = 2\mathcal{M}(t, R), \quad (81)$$

beyond which it becomes ingoing too. Both ingoing and outgoing null rays have a vanishing velocity at the maximum-curvature hypersurface.

Canonically, the metric (79) is defined by the GP gauge

$$N = 1 \quad , \quad E^x = R^2 \quad (82)$$

in terms of the radial coordinate  $R$ , together with the condition

$$E^\varphi = R^2, \quad (83)$$

which may be seen as an implication of the marginal LTB condition in the GP gauge. Assuming the classical values  $c_f = \alpha_0 = \alpha_2 = 1$ ,  $q = 0$ , and  $\Lambda_0 = \Lambda$ , and further choosing a vanishing cosmological constant and pressure,  $\Lambda = 0$  and  $P_0 = 0$ , the resulting equations of motion, together with the on-shell conditions  $H_x = 0 = \tilde{H}$ , imply

$$N^x = -\lambda_0 (1 + \lambda^2) \frac{\sin(2\lambda K_\varphi)}{2\lambda} = -s_{\text{sin}} \lambda_0 \sqrt{1 + \lambda^2} \sqrt{\frac{2\mathcal{M}}{R}} \sqrt{1 - \frac{2\mathcal{M}}{R} \frac{\lambda^2}{1 + \lambda^2}}. \quad (84)$$

In this gauge, the vacuum mass observable takes the form

$$\mathcal{M} = \frac{R}{2} \frac{1 + \lambda^2}{\lambda^2} \sin^2(\lambda K_\varphi) \quad (85)$$

and has the time derivative

$$\begin{aligned} \dot{\mathcal{M}} &= N^x \mathcal{M}' \\ &= -s_{\sin} \lambda_0 \sqrt{1 + \lambda^2} \sqrt{\frac{2\mathcal{M}}{R}} \sqrt{1 - \frac{2\mathcal{M}}{R} \frac{\lambda^2}{1 + \lambda^2}} \mathcal{M}'. \end{aligned} \quad (86)$$

### 3.5 Constant holonomy length

Since we use classical values for all modification functions except for  $\lambda$  and assume constant  $\lambda_0 = 1/\sqrt{1 + \lambda^2}$  (required for an asymptotically flat vacuum solution), a constant  $\lambda = \bar{\lambda}$  corresponds to a  $\mu_0$ -type scheme in the old language of loop quantum cosmology. (See [18] for drawbacks of the old classification of holonomy schemes and a covariant as well as unified treatment in the light of emergent modified gravity.)

If we insert a constant  $\lambda = \bar{\lambda}$  in equation (67) and use the initial condition  $R(t = 0, x) = x$  and  $s_{\sin} = -1$ , we obtain a unique real collapsing solution

$$R_- = x_{\bar{\lambda}} \left( \frac{\Xi^{1/3}}{2\bar{\lambda}^2 x_{\bar{\lambda}}^{2/3}} + \frac{2\bar{\lambda}^2 x_{\bar{\lambda}}^{2/3}}{\Xi^{1/3}} - 1 \right) \quad (87)$$

compatible with the classical limit. In this expression, we have the  $x$ -dependent function

$$x_{\bar{\lambda}} = 2\mathcal{M} \frac{\bar{\lambda}^2}{1 + \bar{\lambda}^2} \quad (88)$$

and the function

$$\begin{aligned} \Xi &= 4x_{\bar{\lambda}} \left( (a^2 + 2\bar{\lambda}^6) x_{\bar{\lambda}} - 3a\bar{\lambda}^2 \sqrt{1 + \bar{\lambda}^2} st \right) + 9\bar{\lambda}^4 (1 + \bar{\lambda}^2) t^2 \\ &\quad + \left[ 16a^4 x_{\bar{\lambda}}^4 - 96a^3 \bar{\lambda}^2 \sqrt{1 + \bar{\lambda}^2} st x_{\bar{\lambda}}^3 + 8a^2 \bar{\lambda}^4 x_{\bar{\lambda}}^2 (27(1 + \bar{\lambda}^2) t^2 + 8\bar{\lambda}^2 x_{\bar{\lambda}}^2) \right. \\ &\quad \left. - 24a \bar{\lambda}^6 x_{\bar{\lambda}} \sqrt{1 + \bar{\lambda}^2} st (9(1 + \bar{\lambda}^2) t^2 + 8\bar{\lambda}^2 x_{\bar{\lambda}}^2) \right. \\ &\quad \left. + 9(1 + \bar{\lambda}^2) \bar{\lambda}^8 t^2 (9(1 + \bar{\lambda}^2) t^2 + 16\bar{\lambda}^2 x_{\bar{\lambda}}^2) \right]^{1/2}. \end{aligned} \quad (89)$$

where  $s = \pm 1$ , as well as

$$a = \frac{A^3 - \lambda^6}{A^{3/2}} \quad (90)$$

and

$$A = \frac{\bar{\lambda}^2}{2x_{\bar{\lambda}}} \left( x + x_{\bar{\lambda}} + \sqrt{(x - x_{\bar{\lambda}})(x + 3x_{\bar{\lambda}})} \right). \quad (91)$$

This expression has the correct classical limit,

$$\lim_{\bar{\lambda} \rightarrow 0} R_- = \left( x^{3/2} - s \frac{3}{2} \sqrt{2\mathcal{M}(x)} t \right)^{2/3} \quad (92)$$

and is real only for

$$x > 2\mathcal{M}(x) \frac{\bar{\lambda}^2}{1 + \bar{\lambda}^2} \quad (93)$$

with a lower bound that in a region of constant  $\mathcal{M}$  is always less than the Schwarzschild radius.

Notice that in this solution, the sign parameter  $s = \pm 1$  always multiplies  $t$ , and, therefore, the solution with  $s = -1$  can be understood as the time-reversed version of  $s = +1$ . Hence, we may restrict ourselves to the latter solution. Furthermore, we note that Eq. (67) in the case  $s_{\text{sin}} = -1$ , is equivalent to the time-reversed version of the  $s_{\text{sin}} = +1$  case, differing only by the initial condition. In sight of this, it is more convenient to solve Eq. (67) with the initial condition  $R(t = 0) = x_{\bar{\lambda}}$ , such that we obtain the simpler, and manifestly time-reversal invariant, solution

$$R = x_{\bar{\lambda}} \left[ 2\bar{\lambda}^{2/3} x_{\bar{\lambda}}^{2/3} \left( 3|t| \sqrt{(1 + \bar{\lambda}^2) (9(1 + \bar{\lambda}^2)t^2 + 16\bar{\lambda}^2 x_{\bar{\lambda}}^2)} + 9(1 + \bar{\lambda}^2)t^2 + 8\bar{\lambda}^2 x_{\bar{\lambda}}^2 \right)^{-1/3} \right. \\ \left. + \frac{x_{\bar{\lambda}}^{-2/3}}{2\bar{\lambda}^{2/3}} \left( 3|t| \sqrt{(1 + \bar{\lambda}^2) (9(1 + \bar{\lambda}^2)t^2 + 16\bar{\lambda}^2 x_{\bar{\lambda}}^2)} + 9(1 + \bar{\lambda}^2)t^2 + 8\bar{\lambda}^2 x_{\bar{\lambda}}^2 \right)^{1/3} - 1 \right] \quad (94)$$

with classical limit

$$\lim_{\bar{\lambda} \rightarrow 0} R = \left( \frac{3}{2} \sqrt{2\mathcal{M}(x)} t \right)^{2/3}. \quad (95)$$

In this solution, the singularity and the reflection-symmetry hypersurface correspond to  $t = 0$ .

### 3.6 Scale-dependent holonomy function

A common scale-dependent choice for the holonomy function is given by  $\lambda = \sqrt{\Delta/E^x} = \sqrt{\Delta}/R$ , which provides a covariant formulation of the traditional  $\bar{\mu}$ -scheme of loop quantum cosmology. Asymptotic flatness of the vacuum solution then requires  $\lambda_0 = 1$  [16]. With this choice, Eq. (67) is hard to solve analytically. However, if we look at the initial stages of the collapse where  $\lambda \ll 1$ , we can approximate this equation to

$$\mathcal{M} \approx \frac{R}{4} \frac{1 - s_{\text{cos}} \sqrt{1 - 4\lambda^2 \dot{R}^2}}{\lambda^2}, \quad (96)$$

where  $s_{\text{cos}} = \text{sgn}(\cos(2\lambda K_{\varphi}))$ . Using the scale-dependent holonomy function, we obtain

$$R(t, x) \approx \left( x^3 \pm 3\sqrt{2\mathcal{M}t} \sqrt{x^3 - 2\Delta\mathcal{M}} + \frac{9\mathcal{M}t^2}{2} \right)^{1/3}, \quad (97)$$

where we have imposed the initial condition  $R(0) = x$  and used  $s_{\text{cos}} = +1$ , appropriate for the early stages of the collapse. This solution reduces to the classical one, (92), in the limit  $\Delta \rightarrow 0$  and is valid only for the region  $\sqrt{\Delta}/R \ll 1$ . However, for  $\Delta \neq 0$  the square root restricts this solution to the region  $x > (2\Delta\mathcal{M})^{1/3}$ . For  $\Delta$  of the order of a Planck area, this would be a very small distance and hence we may simply assume that all observable physical effects occur for  $x \gg (2\Delta\mathcal{M})^{1/3}$  and expand the term in the square root, simplifying it to

$$R(t, x) \approx \left( \left( x^{3/2} \pm \frac{3t}{2} \sqrt{2\mathcal{M}} \right)^2 \mp \frac{3\Delta t}{2} \left( \frac{2\mathcal{M}}{x} \right)^{3/2} \right)^{1/3}. \quad (98)$$

This approximation will be sufficient for exploring semiclassical effects at the early stages of the collapse at large distances. Taking the collapsing solution with the two sign choices in (98) given by  $(-, +)$ , we find that the modification term slows down the collapse.

## 4 Conclusions

We have constructed and analyzed models of marginal LTB collapse with modifications suggested by loop quantum gravity. Compared with previous proposals, we have included several key ingredients: We used emergent modified gravity in order to find space-time covariant theories that include the desired modifications. Different space-time gauges and slicings can then be analyzed without changing physical implications. The usual concepts of space-time curvature, geodesics, and horizons are then unambiguously defined and allow a meaningful analysis of black-hole properties.

Replacing previous proposals for LTB collapse in loop quantum gravity with covariant versions does not merely change a few coefficients in equations of motion. Instead, there are several conceptual differences that require a careful treatment of defining modified theories and implementing classes of solutions, for instance of LTB form. The underlying framework of emergent modified gravity shows that the crucial invariance condition given by general covariance can be implemented on the presence of holonomy modifications from loop quantum gravity only if the classical relationship between the space-time metric and the original phase-space variables is modified. The specific new dependence is strictly derived from the theory, specified by a modified Hamiltonian constraint compatible with covariance conditions, and does not require additional choices. However, the form of the new metric changes either the form of LTB conditions, or the form of the resulting space-time metric if the classical LTB conditions are used. Since an LTB reduction is phenomenological rather than fundamental, there is no preferred choice, except that some cases may be easier to solve than others. LTB models in loop quantum gravity are therefore more ambiguous than their classical versions.

The emergent nature of the space-time metric also means that there are no distinguished phase-space variables with a direct relationship to the spatial part of the metric. In this situation, it is important to make sure that the freedom of performing canonical

transformations is properly included in formulating distinct modified theories. Also here, the framework of emergent modified gravity has already provided a suitable classification of modified spherically symmetric theories, up to second order in spatial derivatives of the phase-space functions. A number of modification functions then characterizes physically distinct modifications, one of which is directly relevant for holonomy modifications.

The multitude of different modifications combined with the freedom of choosing a canonical phase-space representation implies that physical effects are best analyzed from the perspective of effective field theory. One should then consider theories on an equal footing that contain all possible terms of a certain derivative order and obey the required symmetries, given here by general covariance. In a canonical theory, the derivative order is determined by spatial derivatives only. General covariance then makes sure that they are accompanied by suitable time derivatives in the resulting equations of motion. Within this setting, one may choose specific modification functions (depending on phase-space degrees of freedom but not their spatial derivatives) in order to model certain effects such as holonomy modifications in models of loop quantum gravity.

Our analysis in this paper follows these principles. As a result, marginal LTB collapse in loop quantum gravity implies a physical singularity, rather than just a coordinate singularity from shell-crossing.

## Acknowledgements

This work was supported in part by NSF grant PHY-2206591.

## A LTB models in classical general relativity

We provide a brief review of the classical formulation of LTB models with an emphasis of canonical properties as required for emergent modified gravity.

### A.1 Geometric formulation

In classical general relativity, space-time solutions describing the collapse of inhomogeneous dust are given by the LTB metric

$$ds^2 = -dt^2 + \frac{(R')^2}{1 + \kappa(x)} dx^2 + R^2 d(\vartheta^2 + \sin^2 \vartheta d\varphi^2), \quad (99)$$

where the two functions  $R(t, x)$  and  $\kappa(x) > -1$  are related via Einstein's equation by

$$\dot{R}^2 = \kappa(x) + \frac{F(x)}{R}. \quad (100)$$

The new function  $F$  is implicitly given by the solution to

$$\frac{F'}{8\pi GR^2 R'} = T_{tt}. \quad (101)$$

If the energy density  $T_{tt}$  is written in terms of the Misner–Sharp mass  $m$ , it takes the form

$$T_{tt} = \frac{m'}{4\pi GR^2 R'} . \quad (102)$$

The last two equations then imply that

$$m = \frac{1}{2}F(x) \quad (103)$$

is a function of the radial coordinate only. We are therefore left with a single equation of motion,

$$\dot{R}^2 = \kappa(x) + \frac{2m(x)}{R} . \quad (104)$$

In the marginally bound case where  $\kappa(x) = 0$ , this equation can be directly integrated to yield

$$R(t, x) = \pm \left( \frac{3}{2} \sqrt{2m} (t - t_i(x)) \right)^{2/3} , \quad (105)$$

where  $t_i(x)$  is an arbitrary function specifying initial values. Choosing our initial values by  $R|_{t=0} = x$ , and the negative sign for collapse, we have

$$R(t, x) = \pm \left( x^{3/2} - \frac{3t}{2} \sqrt{2m(x)} \right)^{2/3} . \quad (106)$$

The Ricci scalar of (99) using (106) is given by

$$\mathcal{R} = \frac{6mm'}{R^{3/2} ((R^{3/2} - x^{3/2})m' + 3\sqrt{xm})} , \quad (107)$$

where  $m' = dm/dx$ . This spacetime has a geometric singularity at  $R = 0$  given by the surface

$$t(x) = \frac{2x^{3/2}}{3\sqrt{2m(x)}} . \quad (108)$$

There is a further singularity when the second term of the denominator in (107) vanishes. Solving for  $R$  in this case, we obtain

$$R(t, x) = \left( x^{3/2} - \frac{3\sqrt{xm(x)}}{m'(x)} \right)^{2/3} . \quad (109)$$

Setting this expression equal to (106) in the negative case, we derive a second singular surface given by

$$t(x) = -\frac{\sqrt{2xm(x)}}{m'(x)} . \quad (110)$$

Since this function is strictly negative for positive  $m'$ , it constitutes a past singularity. For a dust distribution of constant density,  $m \propto x^3$  and the singular  $t(x)$  is independent of  $x$ .



## A.2 Canonical formulation

Emergent modified gravity requires a canonical formulation of the gravitational field as well as matter ingredients.

### A.2.1 Vacuum LTB models

Using densitized-triad variables of Schwinger type [30], the spherically symmetric reduction of general relativity as derived in [25, 26] implies the spherical line element

$$ds^2 = -N^2 dt^2 + \frac{(E^\varphi)^2}{E^x} (dx + N^x dt)^2 + E^x (d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad (111)$$

with fields  $E^x$  and  $E^\varphi$ . We assume that  $E^x > 0$ , fixing the orientation of the spatial triad. The densitized-triad components  $E^x$  and  $E^\varphi$  as well as their momenta  $K_x$  and  $K_\varphi$  subject to the Hamiltonian constraint

$$H^{\text{grav}}[N] = \int dx N \left[ \frac{((E^x)')^2}{8\sqrt{|E^x|}E^\varphi} - \frac{E^\varphi}{2\sqrt{|E^x|}} - \frac{E^\varphi K_\varphi^2}{2\sqrt{|E^x|}} - 2K_\varphi \sqrt{|E^x|} K_x - \frac{\sqrt{|E^x|}(E^x)'(E^\varphi)'}{2(E^\varphi)^2} + \frac{\sqrt{|E^x|}(E^x)''}{2E^\varphi} \right] = 0, \quad (112)$$

and the diffeomorphism constraint

$$H_x^{\text{grav}}[N^x] = \int dx N^x (K_\varphi' E^\varphi - K_x (E^x)') = 0, \quad (113)$$

where the primes are radial derivatives. As usual [31, 32, 33], the lapse function  $N$  and radial shift vector  $N^x$  do not have momenta but rather can be used to specify the space-time gauge or slicing.

Comparing the two metrics (99) and (111) we find that the relevant LTB gauge is given by

$$N = 1 \quad , \quad N^x = 0. \quad (114)$$

Moreover, the phase space is further restricted by the LTB condition

$$E^\varphi = \frac{(E^x)'}{2\sqrt{1 + \kappa(x)}}. \quad (115)$$

We also identify  $R^2 = E^x$  without imposing a further constraint.

For a given  $\kappa(x)$ , we can directly enforce the LTB constraint

$$L_1 \equiv E^\varphi - \frac{(E^x)'}{2\sqrt{1 + \kappa(x)}}, \quad (116)$$

set to vanish dynamically. For a consistent implementation, this condition must be preserved under time evolution [1, 2], that is,  $L_2 \equiv \dot{L}_1 = \{L, H[N] + H_x[N^x]\} = 0$ , thereby

obtaining a secondary condition. (Recall that  $\kappa$  is a function of the radial coordinate only and hence  $\dot{\kappa} = 0$  so that the brackets do not act on it.)

We compute

$$\begin{aligned}
L_2 &= \dot{E}^\varphi - \frac{1}{2\sqrt{1+\kappa(x)}}(\dot{E}^x)' \\
&= -\frac{\delta H[N]}{\delta K_\varphi} + \frac{1}{2\sqrt{1+\kappa(x)}} \left( \frac{\delta H[N]}{\delta K_x} \right)' + \left( N^x E^\varphi - \frac{1}{2\sqrt{1+\kappa(x)}} N^x (E^x)' \right)' \\
&= -\frac{\delta H[1]}{\delta K_\varphi} + \frac{1}{2\sqrt{1+\kappa(x)}} \left( \frac{\delta H[1]}{\delta K_x} \right)' \\
&= -\frac{\partial H[1]}{\partial K_\varphi} + \left( \frac{\partial H[1]}{\partial K'_\varphi} \right)' - \left( \frac{\partial H[1]}{\partial K''_\varphi} \right)'' \\
&\quad + \frac{1}{2\sqrt{1+\kappa(x)}} \left( \left( \frac{\partial H[1]}{\partial K_x} \right)' - \left( \frac{\partial H[1]}{\partial K'_x} \right)'' + \left( \frac{\partial H[1]}{\partial K''_x} \right)''' \right). \tag{117}
\end{aligned}$$

Using (112) this expression equals

$$\begin{aligned}
L_2 &= \frac{E^\varphi K_\varphi}{\sqrt{|E^x|}} + 2\sqrt{|E^x|} K_x - \frac{(K_\varphi \sqrt{|E^x|})'}{\sqrt{1+\kappa(x)}} \\
&= \frac{K_\varphi}{\sqrt{|E^x|}} \left( E^\varphi - \frac{(E^x)'}{2\sqrt{1+\kappa(x)}} \right) - \frac{\sqrt{|E^x|}}{E^\varphi} \left( \frac{E^\varphi K'_\varphi}{\sqrt{1+\kappa(x)}} - 2K_x E^\varphi \right) \\
&= \left( \frac{K_\varphi}{\sqrt{|E^x|}} + \frac{\sqrt{|E^x|}}{E^\varphi} 2K_x \right) L_1 - \frac{\sqrt{|E^x|}}{E^\varphi} \frac{H_x^{\text{grav}}}{\sqrt{1+\kappa(x)}}. \tag{118}
\end{aligned}$$

We therefore find that in the given gauge the LTB condition  $L_1 = 0$  is consistent with the on-shell canonical dynamics, where  $H_x^{\text{grav}} = 0$ .

### A.2.2 Perfect-fluid LTB

The perfect fluid is a model of a collection of point particles in the space-time manifold which may locally be described by coordinate functions  $T(t, x)$  and  $X(t, x)$ . We interpret the time dependence  $(T(t, x), X(t, x))$  at fixed  $x$  as the event to which the particle at a position labeled by  $x$  has moved from the initial event  $(T(0, x), X(0, x))$ . The two functions  $T$  and  $X$  can therefore be used as basic fields of a dynamical formulation of perfect fluids. For a given action or set of canonical constraints, the dynamics will be valid in a region in which trajectories of individual particles do not cross. Boundaries of this region for a given theory are determined by shell-crossing singularities.

For a derivation of the canonical dynamics, we combine spatial fields  $T$  and  $X$  with canonical momenta  $P_T$  and  $P_X$  with units of energy and momentum, respectively. Both

momenta are scalar densities of weight one, which in one radial dimension transform like components of a co-vector. We have canonical Poisson brackets

$$\{T(x), P_T(y)\} = \{X(x), P_X(y)\} = \delta(x - y). \quad (119)$$

Since the two fields in spherical symmetry are spatial scalars, we immediately obtain the fluid contribution

$$H_x = P_T T' + P_X X' \quad (120)$$

to the diffeomorphism constraint.

The dual role of  $(T, X)$  as fields and space-time coordinates can be used to recognize  $H_x$  as the spatial component of the energy-momentum co-vector  $(P_T, P_X)$  transformed to a frame with the spatial coordinate  $x$  that appears in the derivatives. The corresponding time component in this frame is given by  $P_T \dot{T} + P_X \dot{X}$ . In general a general frame  $(t, x)$ , we therefore have the co-vector field

$$P_T dT + P_X dX = (P_T \dot{T} + P_X \dot{X}) dt + (P_T T' + P_X X') dx. \quad (121)$$

The left-hand side is applies in a co-moving frame where  $\dot{T} = 1 = X'$  and  $T' = 0 = \dot{X}$ .

In special relativity, the momentum co-vector  $(P_T, P_X)$  and the coordinate differential  $(dT, dX)$  are both expected to be proportional to the relativistic velocity  $u^\mu$  or its dual. The sum in (121) then takes the form of a scalar product

$$P_T dT + P_X dX = p_\mu dx^\mu \propto u_\mu \frac{dx^\mu}{d\tau} = g_{\mu\nu} u^\mu u^\nu = -s \quad (122)$$

where  $s = 1$  for massive particles and  $s = 0$  in the massless case. The proportionality factor in special relativity is given by the mass, but here we need a scalar of density weight one. The only available choice that can in general be assumed to be non-vanishing is given by  $P_T$ . We therefore define the velocity co-vector

$$u_\mu dx^\mu = -\frac{1}{P_T} (P_T dT + P_X dX) = -\left(\dot{T} + \frac{P_X}{P_T} \dot{X}\right) dt - \left(T' + \frac{P_X}{P_T} X'\right) dx. \quad (123)$$

The fluid's velocity (123) is normalized

$$g^{\mu\nu} u_\mu u_\nu = -s, \quad (124)$$

with  $s = 1$  or  $s = 0$  for a timelike or null fluid, respectively. Therefore,

$$\|(P_T, P_X)\|^2 = -s P_T^2. \quad (125)$$

The relation between the fluid's velocity and the configuration variables  $T$  and  $X$  imply that the latter are capable of locally coordinatizing the manifold with the simplest example given by  $T' = 0, \dot{T} = 1$  and  $X' = 1, \dot{X} = 0$  with  $P_X/P_T$  describing the velocity of the timelike fluid's particles.

The Hamiltonian-constraint contribution from the perfect fluid in its general covariant form is given by [6]

$$H_s^{\text{matter}} = \sqrt{sP_T^2 + \frac{E^x}{(E^\varphi)^2}(H_x^{\text{matter}})^2} - \sqrt{\det q} P (P_X^2/P_T^2) \quad (126)$$

with a free pressure function  $P$  depending on the velocity, and the diffeomorphism-constraint contribution by

$$H_x^{\text{matter}} = P_T T' + P_X X' \quad (127)$$

as already defined. The coefficient  $s = 1$  for a timelike fluid and  $s = 0$  for a null fluid appears in the general contribution to the Hamiltonian constraint. This square-root contribution represents the relativistic energy as the time component of the vector  $(H_s^{\text{matter}}, H_x^{\text{matter}})$  with norm squared  $-sP_T^2$ . Including the gravitational field, the complete Hamiltonian constraint is then given  $H = H^{\text{grav}} + H_s^{\text{matter}}$ , and the total diffeomorphism constraint by  $H_x = H_x^{\text{grav}} + H_x^{\text{matter}}$ . The full Hamiltonian generating the equations of motion is  $H_T[N, N^x] = H^{\text{grav}}[N] + H_s^{\text{matter}}[N] + H_x^{\text{grav}}[N^x] + H_x^{\text{matter}}[N^x]$ .

Because the full diffeomorphism constraint is now given by  $H_x = H_x^{\text{grav}} + H_x^{\text{matter}}$ , imposing the LTB condition, the secondary condition (118) receives the matter contribution

$$L_2|_{\text{O.S., } L_1=0} = \frac{\sqrt{E^x}}{E^\varphi} \frac{H_x^{\text{matter}}}{\sqrt{1 + \kappa(x)}}. \quad (128)$$

Therefore, the secondary condition is satisfied only if  $H_x^{\text{matter}} = 0$ , which means we must work in a slicing adapted to the fluid's frame, and therefore have

$$P_X = -\frac{P_T T'}{X'}, \quad (129)$$

and  $s = 1$ , since this is only possible for a timelike fluid.

The energy density of the perfect fluid in the fluid's frame is given by

$$\rho = \frac{P_T}{\sqrt{\det q}}. \quad (130)$$

In the LTB coordinates used in (99), we may define

$$\rho = \frac{1}{4\pi R^2} \frac{dm}{dR} = \frac{1}{4\pi R^2} \frac{m'}{R'}. \quad (131)$$

with some function  $m(x, t)$  we refer to as the mass parameter. Combining (130) and (131), we obtain

$$P_T = \frac{m'}{\sqrt{1 + \kappa(x)}}. \quad (132)$$

### A.3 LTB collapse

The on-shell condition  $H = 0$  together with (8) and  $R^2 = E^x$  implies

$$|P_T| = \frac{1}{2\sqrt{1+\kappa}} \left( (8\pi P - \Lambda) \frac{(R^3)'}{3} - (\kappa R)' + (RK_\varphi^2)' \right) \quad (133)$$

where  $P$  is the function introduced in (11). The relevant equations of motion are

$$\dot{R} = K_\varphi \quad (134)$$

and

$$\dot{K}_\varphi = \left( \frac{\Lambda}{2} - 4\pi P \right) R + \frac{\kappa}{2R} - \frac{K_\varphi^2}{2R} \quad (135)$$

for the unrestricted gravitational variables, as well as

$$\dot{X} = \frac{8\pi R^2}{\sqrt{1+\kappa}} R' \frac{T'}{X'} \frac{1}{P_T} \frac{\partial P}{\partial p} \quad (136)$$

and

$$\dot{T} = \text{sgn}(P_T) + \frac{T'}{X'} \dot{X} \quad (137)$$

for the dust coordinates, where  $p = P_X^2/P_T^2$ , and

$$\dot{P}_T = 0 \quad (138)$$

for the free dust momentum after using (129).

Using (132) and (134), the equations (133) and (135) can be rewritten as

$$m' = \frac{1}{2} \left( (8\pi P - \Lambda) \frac{(R^3)'}{3} + (R\dot{R}^2 - R\kappa)' \right), \quad (139)$$

$$\ddot{R} = \left( \frac{\Lambda}{2} - 4\pi P \right) R - \frac{\dot{R}^2 - \kappa}{2R}. \quad (140)$$

Equations (132) and (138) also imply that  $m(x)$  is independent of time.

For dust,  $P = 0$ , equation (139) can be integrated to yield

$$m(x) = \frac{R(\dot{R}^2 - \kappa)}{2} - \frac{\Lambda}{6} R^3 + m_0, \quad (141)$$

where  $m_0$  is the constant of integration. We may set this constant equal to zero for  $m|_{\dot{R}=0} = 0$ . Further setting  $\Lambda = 0$ , the equation can be exactly solved for  $R(t)$  in the marginally bound case  $\kappa = 0$ ,

$$R = \left( x^{3/2} \pm \frac{3t}{2} \sqrt{2m(x) - m_0} \right)^{2/3}, \quad (142)$$

where we have taken the initial condition  $R(t = 0, x) = x$  to fix the undetermined functions of  $x$  resulting from solving the equation. Direct substitution shows that this function solves (140) too. The solution (106) is precisely the geometric result with the negative sign chosen for the collapsing case.

The LTB metric (1) in this marginal case becomes

$$ds^2 = -dt^2 + \frac{x}{R} \left( 1 - \frac{t}{2} \sqrt{\frac{2m}{x}} (\ln m)' \right)^2 dx^2 + R^2 d\Omega^2. \quad (143)$$

This metric has a coordinate singularity at the time

$$t(x) = 2 \sqrt{\frac{x}{2m(x)} \frac{m(x)}{m'(x)}}. \quad (144)$$

### A.3.1 Collapse in Gullstrand-Painlevé coordinates

If we use  $R$  in (106) as a new radial coordinate we obtain

$$dx = \left( 1 - \sqrt{\frac{2m}{x}} \frac{m' t}{m} \right) \sqrt{\frac{R}{x}} \left( dR + \sqrt{\frac{2m}{R}} dt \right) =: \mathcal{W} \left( dR + \sqrt{\frac{2m}{R}} dt \right), \quad (145)$$

where  $m' = dm/dx$ . Substitution into the LTB metric (1) gives

$$ds^2 = -dt^2 + \left( dR + \sqrt{\frac{2m}{R}} dt \right)^2 + R^2 d\Omega^2, \quad (146)$$

which is the Gullstrand-Painlevé metric with non-constant  $m(t, R)$ . This form of the metric is useful to cross the event horizon. Taking an infalling/outgoing light ray, its velocity in the GP coordinates is given by

$$\frac{dR}{dt} = \pm 1 - \sqrt{\frac{2m}{R}}. \quad (147)$$

The horizon surface is given the vanishing of the outgoing rays's velocity, beyond which it becomes ingoing:

$$2m(t, R) = R. \quad (148)$$

### A.3.2 Uniform distribution

Consider a uniform mass distribution originally contained in a radius  $r$  in LTB coordinates, such that  $m(x) = Mx^3/r^3 = \rho_0 x^3$ , with total mass  $M < r/2$ . In LTB coordinates, the metric (143) with this distribution becomes

$$ds^2 = -dt^2 + \left( 1 - \frac{3\sqrt{\rho_0}}{\sqrt{2}} t \right)^{4/3} dx^2 + x^2 \left( 1 - \frac{3\sqrt{\rho_0}}{\sqrt{2}} t \right)^{4/3} d\Omega^2, \quad (149)$$

and the coordinate singularity (144) for this distribution is at the time

$$t = \frac{2}{3} \sqrt{\frac{r^3}{2M}} = \frac{\sqrt{2}}{3\sqrt{\rho_0}}. \quad (150)$$

The Ricci scalar (107) becomes

$$\mathcal{R} = \frac{6M}{r^3} \frac{x^3}{R^3} = \frac{6M}{r^3} \left(1 - \frac{3\sqrt{\rho_0}}{\sqrt{2}} t\right)^{-2}. \quad (151)$$

The coordinate singularity agrees with the geometric singularity.

With this distribution we obtain

$$\begin{aligned} R(t, x) &= x \left(1 - \frac{3t}{2r} \sqrt{\frac{2M}{r}}\right)^{2/3} \\ &=: a(t)x \end{aligned} \quad (152)$$

and the LTB metric (1) takes the form of a flat FLRW spacetime

$$ds^2 = -dt^2 + a(t)^2 (dx^2 + x^2 d\Omega^2) \quad (153)$$

with a matter-dominated scale factor

$$a(t) = \left(1 - \frac{3\sqrt{\rho_0}}{\sqrt{2}} t\right)^{2/3}. \quad (154)$$

We can then use (106) to get

$$dR = \left(1 - \frac{3\sqrt{\rho_0}}{\sqrt{2}} t\right)^{2/3} dx - R\sqrt{2\rho_0} \left(1 - \frac{3\sqrt{\rho_0}}{\sqrt{2}} t\right)^{-1} dt \quad (155)$$

and rewrite the metric in Gullstrand–Painlevé form (146):

$$ds^2 = -dt^2 + \left(dR + \frac{2R}{3} \left(\frac{\sqrt{2}}{3\sqrt{\rho_0}} - t\right)^{-1} dt\right)^2 + R^2 d\Omega^2. \quad (156)$$

Further redefining the time coordinate

$$\tau = t - \frac{\sqrt{2}}{3\sqrt{\rho_0}} \quad (157)$$

we obtain

$$ds^2 = -d\tau^2 + \left(dR + \frac{2R}{3\tau} d\tau\right)^2 + R^2 d\Omega^2, \quad (158)$$

which is the Oppenheimer-Snyder model in Gullstrand–Painlevé coordinates for a star collapsing from rest at spatial infinity.

The velocity of an infalling/outgoing light ray (147) is given by

$$\frac{dR}{dt} = \pm 1 - \sqrt{\frac{2M}{r}} \frac{R}{r} \left( 1 - \frac{3t}{2r} \sqrt{\frac{2M}{r}} \right)^{-1}. \quad (159)$$

The horizon surface (148) is given by

$$t = \frac{\sqrt{2}}{3\sqrt{\rho_0}} \left( 1 - \sqrt{\frac{2}{\rho_0}} R \right) \quad (160)$$

which differs from the coordinate singularity (150).

The geometric singularity (108) is

$$t(x) = \frac{\sqrt{2}}{3\sqrt{\rho_0}}, \quad (161)$$

which intersects the horizon at  $R = 0$ .

## B Conformal transformations of the emergent metric

Conformal transformations of space-time metrics are a useful tool for various aspects of space-time analysis. However, in such cases they do not change physical properties of the space-time geometry given by an explicit solution. Their application in recent studies of LTB collapse [14, 15] in order to remove physical singularities is conceptually different and requires further discussions, which we present in this appendix. We focus on the two-dimensional case relevant for the spherically symmetric reduced theory in these applications.

The study [14] defined a conformally modified two-dimensional emergent metric given by  $\tilde{\mathfrak{g}}_{\mu\nu} = \omega^2 \tilde{g}_{\mu\nu}$ , explicitly

$$\tilde{\mathfrak{g}}_{\mu\nu} dx^\mu dx^\nu = \omega^2 \left( -N^2 dt^2 + \tilde{q}_{xx} (dx + N^x dt)^2 \right). \quad (162)$$

If the covariance condition for the emergent metric (26) is satisfied, then the covariance condition for the conformally modified emergent metric

$$\delta_\epsilon \tilde{\mathfrak{g}}_{\mu\nu}|_{\text{o.s.}} = \mathcal{L}_\xi \tilde{\mathfrak{g}}_{\mu\nu}|_{\text{o.s.}}, \quad (163)$$

requires  $\omega$  be a space-time scalar, meaning its gauge transformation must give the Lie derivative  $\mathcal{L}_\xi \omega = \xi^\mu \partial_\mu \omega$  on-shell. Explicitly, the condition becomes

$$\delta_\epsilon \omega = \xi^t \dot{\omega} + \xi^x \omega'. \quad (164)$$



If this is the case, then (162) is obviously a covariant tensor. This is a feature not exclusive of emergent modified gravity. In general relativity, a covariant tensor of any rank multiplied by a scalar is still covariant and of the same rank. The question is whether (162) can be used as a space-time metric replacing the emergent one, since the space-time metric has several properties that single it out among other tensors, including off-shell properties.

As shown in [29], such a conformal transformation is possible and in accordance with the principles of emergent modified gravity (anomaly-freedom and covariance) only if the Hamiltonian constraint is rescaled to  $\bar{H}$  given by

$$\bar{H} = \omega^{-1} \tilde{H} \quad (165)$$

and the conformal factor obeys the off-shell gauge transformation

$$\{\omega, \tilde{H}[\epsilon^0] + H_x[\epsilon^x]\} = \Omega_0 \epsilon^0 + \omega' \epsilon^x \quad (166)$$

for some phase space function  $\Omega_0$ , with no derivatives of  $\epsilon^0$  or  $\epsilon^x$ , and only if  $\omega$  commutes with itself. In this particular system, if  $\omega$  is a function of only the vacuum gravitational observable  $\mathcal{M}$  and  $E^x$ , then these properties are satisfied. The algebra of  $\bar{H}$  with the diffeomorphism constraint then takes the hypersurface-deformation form

$$\{H_x[\bar{N}^x], H_x[\bar{M}^x]\} = -H_x[\bar{M}^x(\bar{N}^x)' - \bar{N}^x(\bar{M}^x)'] , \quad (167)$$

$$\{\bar{H}[\bar{N}], H_x[\bar{M}^x]\} = -\bar{H}[\bar{M}^x \bar{N}'] , \quad (168)$$

$$\{\bar{H}[\bar{N}], \bar{H}[\bar{M}]\} = -H_x[\bar{q}^{xx}(\bar{M} \bar{N}' - \bar{N} \bar{M}')] \quad (169)$$

with structure function

$$\bar{q}^{xx} = \omega^{-2} \tilde{q}^{xx} . \quad (170)$$

The emergent line-element compatible with the new algebra is given by

$$\begin{aligned} \bar{g}_{\mu\nu} &= -\bar{N}^2 dt^2 + \bar{q}_{xx} (dx + \bar{N}^x dt)^2 \\ &= -\bar{N}^2 dt^2 + \omega^2 \tilde{q}_{xx} (dx + \bar{N}^x dt)^2 . \end{aligned} \quad (171)$$

The conformally transformed geometry is therefore anomaly-free and covariant, in accordance with the principles of emergent modified gravity. However, there is an important conceptual distinction between the use of the tensors (162) and (171) as follows.

In the former case, the space-time geometry is given by the emergent metric (27), which is associated to the constraint  $\tilde{H} = \tilde{H}^{\text{grav}} + \tilde{H}_s^{\text{matter}}$  given by the expressions (32) and (36) that satisfy the hypersurface-deformation brackets (14)–(25); the dynamics are generated by the constraints  $\tilde{H}$  and  $H_x$  with lapse  $N$  and shift  $N^x$ ; and the expression (162) is a covariant tensor not strictly related to the space-time geometry.

In the latter case, the space-time geometry is given by the emergent metric (171), which is associated to the constraint  $\bar{H}$  given by the expression (165) that satisfies the hypersurface-deformation brackets (167)–(169); the dynamics are generated by the constraints  $\bar{H}$  and  $H_x$  with lapse  $\bar{N}$  and shift  $\bar{N}^x$ . Dynamics are, therefore, different in each

case: Setting  $N = 1$  will generate different equations of motion than  $\bar{N} = 1$ , the latter corresponding to  $N = \omega^{-1}$  in the original frame (using  $\tilde{H}$ ).

The conclusion that both the space-time geometry and the dynamics are different leads to further questions of which is the correct system for physical applications. While such a question can only be answered by experiments or observations, we may discuss theoretical or conceptual aspects favoring one over another. First, the fact that the space-time geometry is changed in the conformal case leads us to question whether the original matter coupling is still valid. While the system remains covariant, one would expect the correct theory to present the perfect fluid directly coupled to the space-time metric, in this case given by  $\bar{g}_{\mu\nu}$  instead of the original  $\tilde{g}_{\mu\nu}$ . If we amend this, following [6], the actual perfect fluid contribution to the Hamiltonian constraint should be given by

$$\bar{H}_s^{\text{matter}} = \sqrt{sP_T^2 + \bar{q}^{xx}(H_x^{\text{matter}})^2} - \sqrt{\det \bar{q}} P_q(E^x, P_X^2/P_T^2) - 4\pi\sqrt{E^x} E^\varphi \bar{\lambda}_0 P_0(E^x, P_X^2/P_T^2), \quad (172)$$

which differs from  $\omega^{-1}\tilde{H}_s^{\text{matter}}$ , with  $\tilde{H}_s^{\text{matter}}$  given by (36), by using  $\bar{q}^{xx}$  instead of  $\tilde{q}^{xx}$  and an overall factor of  $\omega^{-1}$ . The use of the amended matter constraint contribution (172) will, in turn, change the dynamics even further and, as a consequence, notions of the fluid frame will change, rendering the motivation of certain gauge choices more difficult.

Finally, we may expect canonical quantization to differ significantly between the different systems because the corresponding constraint, either  $\bar{H}$  or  $\tilde{H}$ , with its associated brackets must be used in the quantization procedure. In this application, it is the original system, with  $\tilde{H}$ , that is the more natural choice because it is the simpler constraint: The overall factor  $\omega^{-1}$  depends on the complicated phase-space function  $\mathcal{M}$ , which must be turned into a composite operator in the quantum theory, leading to ordering issues among others.

## References

- [1] M. Bojowald, T. Harada, and R. Tibrewala, Lemaitre-Tolman-Bondi collapse from the perspective of loop quantum gravity, *Phys. Rev. D* 78 (2008) 064057, [arXiv:0806.2593]
- [2] M. Bojowald, J. D. Reyes, and R. Tibrewala, Non-marginal LTB-like models with inverse triad corrections from loop quantum gravity, *Phys. Rev. D* 80 (2009) 084002, [arXiv:0906.4767]
- [3] M. Bojowald and E. I. Duque, Emergent modified gravity, *Class. Quantum Grav.* 41 (2024) 095008, [arXiv:2404.06375]
- [4] M. Bojowald and E. I. Duque, Emergent modified gravity: Covariance regained, *Phys. Rev. D* 108 (2023) 084066, [arXiv:2310.06798]
- [5] M. Bojowald and E. I. Duque, Emergent modified gravity: Polarized Gowdy model on a torus, *Phys. Rev. D* (2024) to appear, [arXiv:2407.13583]

- [6] E. I. Duque, Emergent modified gravity: The perfect fluid and gravitational collapse, *Phys. Rev. D* 109 (2024) 044014, [arXiv:2311.08616]
- [7] M. Bojowald and E. I. Duque, Emergent modified gravity coupled to scalar matter, *Phys. Rev. D* 109 (2024) 084006, [arXiv:2311.10693]
- [8] E. I. Duque, Emergent electromagnetism, *Phys. Rev. D* 110 (2024) 125006, [arXiv:2407.14954]
- [9] A. Alonso-Bardaj and D. Brizuela, Spacetime geometry from canonical spherical gravity, [arXiv:2310.12951]
- [10] A. Alonso-Bardaj, D. Brizuela, and R. Vera, An effective model for the quantum Schwarzschild black hole, *Phys. Lett. B* 829 (2022) 137075, [arXiv:2112.12110]
- [11] A. Alonso-Bardaj, D. Brizuela, and R. Vera, Nonsingular spherically symmetric black-hole model with holonomy corrections, *Phys. Rev. D* 106 (2022) 024035, [arXiv:2205.02098]
- [12] A. Alonso-Bardaj, D. Brizuela, and R. Vera, Singularity resolution by holonomy corrections: Spherical charged black holes in cosmological backgrounds, *Phys. Rev. D* 107 (2023) 064067, [arXiv:2302.10619]
- [13] M. Bojowald, E. I. Duque, and D. Hartmann, A new type of large-scale signature change in emergent modified gravity, *Phys. Rev. D* 109 (2024) 084001, [arXiv:2312.09217]
- [14] A. Alonso-Bardaj and D. Brizuela, Nonsingular collapse of a spherical dust cloud, *Phys. Rev. D* 109 (2024) 064023, [arXiv:2312.15505]
- [15] A. Alonso-Bardaj, Formation of nonsingular spherical black holes with holonomy corrections, [arXiv:2410.20529]
- [16] I. H. Belfaqih, M. Bojowald, S. Brahma, and E. I. Duque, Black holes in effective loop quantum gravity: Covariant holonomy modifications, [arXiv:2407.12087]
- [17] C. Zhang, J. Lewandowski, Y. Ma, and J. Yang, Black Holes and Covariance in Effective Quantum Gravity, [arXiv:2407.10168]
- [18] I. H. Belfaqih, M. Bojowald, S. Brahma, and E. I. Duque, Lessons for loop quantum gravity from emergent modified gravity, [arXiv:2411.17545]
- [19] F. Fazzini, V. Husain, and E. Wilson-Ewing, Shell-crossings and shock formation during gravitational collapse in effective loop quantum gravity, *Phys. Rev. D* 109 (2024) 084052, [arXiv:2312.02032]

- [20] J. G. Kelly, R. Santacruz, and E. Wilson-Ewing, Black hole collapse and bounce in effective loop quantum gravity, *Class. Quantum Grav.* 38 (2021) 04LT01, [arXiv:2006.09325]
- [21] V. Husain, J. G. Kelly, R. Santacruz, and E. Wilson-Ewing, Quantum gravity of dust collapse: shock waves from black holes, *Phys. Rev. Lett.* 128 (2022) 121301, [arXiv:2109.08667]
- [22] K. Giesel, H. Liu, E. Rullit, P. Singh, and S. A. Weigl, Embedding generalized LTB models in polymerized spherically symmetric spacetimes, [arXiv:2308.10949]
- [23] K. Giesel, H. Liu, P. Singh, and S. A. Weigl, Generalized analysis of a dust collapse in effective loop quantum gravity: fate of shocks and covariance, [arXiv:2308.10953]
- [24] K. Enqvist, Lemaitre-Tolman-Bondi model and accelerating expansion, *Gen. Rel. Grav.* 40 (2008) 451–466, [arXiv:0709.2044]
- [25] M. Bojowald, Spherically Symmetric Quantum Geometry: States and Basic Operators, *Class. Quantum Grav.* 21 (2004) 3733–3753, [gr-qc/0407017]
- [26] M. Bojowald and R. Swiderski, Spherically Symmetric Quantum Geometry: Hamiltonian Constraint, *Class. Quantum Grav.* 23 (2006) 2129–2154, [gr-qc/0511108]
- [27] J. D. Brown and K. V. Kuchař, Dust as a standard of space and time in canonical quantum gravity, *Phys. Rev. D* 51 (1995) 5600–5629
- [28] M. Bojowald, *Canonical Gravity and Applications: Cosmology, Black Holes, and Quantum Gravity*, Cambridge University Press, Cambridge, 2010
- [29] M. Bojowald, E. I. Duque, and A. Shah, Hypersurface deformations, [arXiv:2410.18807]
- [30] J. Schwinger, Quantized gravitational field, *Phys. Rev.* 130 (1962) 1253–1258
- [31] P. A. M. Dirac, The theory of gravitation in Hamiltonian form, *Proc. Roy. Soc. A* 246 (1958) 333–343
- [32] J. Katz, Les crochets de Poisson des contraintes du champ gravitationne, *Comptes Rendus Acad. Sci. Paris* 254 (1962) 1386–1387
- [33] R. Arnowitt, S. Deser, and C. W. Misner, The Dynamics of General Relativity, In L. Witten, editor, *Gravitation: An Introduction to Current Research*, Wiley, New York, 1962, Reprinted in [34]
- [34] R. Arnowitt, S. Deser, and C. W. Misner, The Dynamics of General Relativity, *Gen. Rel. Grav.* 40 (2008) 1997–2027