

LangYa: Revolutionizing Cross-Spatiotemporal Ocean Forecasting

Nan Yang^{1,2}, Chong Wang^{1,2}, Meihua Zhao^{1,2}, Zimeng Zhao^{1,2},
Huilin Zheng^{1,2}, Bin Zhang^{1,2}, Jianing Wang^{1,2}, Xiaofeng Li^{1,2*}

¹Key Laboratory of Ocean Observation and Forecasting, Qingdao, China.

²Key Laboratory of Ocean Circulation and Waves, Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China.

*Corresponding author(s). E-mail(s): lixf@qdio.ac.cn;

Abstract

Ocean forecasting is crucial for both scientific research and societal benefits. Currently, the most accurate forecasting systems are global ocean forecasting systems (GOFs), which represent the ocean state variables (OSVs) as discrete grids and solve partial differential equations (PDEs) governing the transitions of oceanic state variables using numerical methods. However, GOFs processes are computationally expensive and prone to cumulative errors. Recently, large artificial intelligence (AI)-based models significantly boosted forecasting speed and accuracy. Unfortunately, building a large AI ocean forecasting system that can be considered cross-spatiotemporal and air-sea coupled forecasts remains a significant challenge. Here, we introduce LangYa, a cross-spatiotemporal and air-sea coupled ocean forecasting system. Results demonstrate that the time embedding module in LangYa enables a single model to make forecasts with lead times ranging from 1 to 7 days. The air-sea coupled module effectively simulates air-sea interactions. The ocean self-attention module improves network stability and accelerates convergence during training, and the adaptive thermocline loss function improves the accuracy of thermocline forecasting. Compared to existing numerical and AI-based ocean forecasting systems, LangYa uses 27 years of global ocean data from the Global Ocean Reanalysis and Simulation version 12 (GLORYS12) for training and achieves more reliable deterministic forecasting results for OSVs. LangYa forecasting system provides global ocean researchers with access to a powerful software tool for accurate ocean forecasting and opens a new paradigm for ocean science.

Keywords: Global Ocean Forecasting, Large Model, AI for Science

1 Main

Ocean forecasting aims to forecast future ocean dynamic states and is crucial for understanding ocean activities. Numerical global ocean forecasting systems (GOFSS), such as the Mercator Ocean physical system (PSY4) [1] and the Real-Time Ocean Forecasting System (RTOFS) [2], rely on geophysical fluid dynamics and partial differential equations (PDEs) to describe future ocean conditions, and such systems remain widely used worldwide. However, improving the inference accuracy and speed of numerical forecast models is a significant challenge. First, numerical forecasting heavily depends on parameterizations and the extent of human knowledge regarding oceanic physical laws. Approximation functions can exacerbate forecast errors. In addition, even on supercomputers equipped with hundreds of computing nodes, it takes several hours to generate a single forecast. Large AI models are reshaping interactions between humans and technology through their massive learning parameters and powerful learning capabilities, e.g., ChatGPT and Sora. Such advances open the way for the development of geophysical system forecasting, including large AI models for weather (e.g., FourcastNet, PanGu-Weather) and ocean (e.g., XiHe [3], AI-GOMS [4]) forecasts. These large models are mainly data-driven, recasting air state variables (ASVs) and ocean state variables (OSVs) forecasting problems as autoregressive video generation, thereby avoiding much computational complexity and uncertainties inherent in physical processes. Moreover, specialized computing devices such as graphics processing units (GPUs) offer extremely fast inference speeds. By investigating and practicing ocean forecasting, we summarize two key barriers in current large ocean forecasting models: (1) Future OSVs are forecasted autoregressively in a step-by-step manner, with temporal information not incorporated into the training and inference processes. It neglects the time-dependent evolution of OSV fields and their interdependencies, making it impossible to achieve cross-timescale forecasts. (2) Air and ocean data are concatenated as model inputs to represent air-sea interactions, a simple way that cannot effectively simulate these interactions nor integrate their underlying mechanisms into the training and inference processes. Although researchers agree that large AI models hold tremendous potential for ocean forecasting, it should be acknowledged that revolutionary breakthroughs are still needed before an AI ocean forecasting system capable of cross-timescale, air-sea coupled forecasts can be realized. Fortunately, the breakthroughs have now come as expected. Here, we present the LangYa forecasting system (see the Methods section for "LangYa"), a powerful large AI-based model for cross-spatiotemporal, air-sea coupled ocean forecasting. LangYa delivers faster and more accurate forecasts for OSVs (temperature, salinity, and currents) than both traditional numerical weather prediction (NWP) systems and XiHe (an influential large AI-based model). Our contributions are fourfold: (1) We propose a large-language-model-based (LLM-based) Time Embedding Module that achieves high spatiotemporal resolution ($1/12^\circ$, 1 day) forecasts for OSVs. It enables cross-timescale forecasts (up to 7 days) without iterative steps, allowing training and inference for future 7-day OSVs with one-off. (2) We develop an asynchronous cross-iterative random sampling strategy to simulate the atmospheric's stochastic interaction with the ocean, improving the forecasting skill for variables with different rates of change, such as sea temperature and currents. (3) We introduce an ocean self-attention module

based on cosine attention, enabling more stable model training and faster convergence. With this module, a single epoch of distributed training on global OSVs from 1993 to 2019 can be completed in just 0.5 days. (4) We propose an adaptive loss function for thermocline, which significantly improves the accuracy of forecasts at a 100 m depth thermocline, yielding about a 50% improvement in flow forecast accuracy compared to other methods. Using GLORYS12 reanalysis data, we verify that LangYa can achieve stable training and convergence, and effectively yield real-time, accurate, multivariate forecasts across spatiotemporal scales, especially for the thermocline layer.

2 Cross-SpatioTemporal and Air-Sea Coupled Ocean Forecasting System

LangYa adopts the idea of data-driven that builds ocean forecasting as a video prediction task. The design contains training and testing phases by using oceanic and atmospheric reanalysis data as inputs and generating forecasts for Ocean State Variables (OSVs) — such as sea temperature, salinity, and currents — through single-pass inference over 1-7 days. The system uses 11,322 samples and mitigates overfitting risks by randomly shuffling sample order at the beginning of each training iteration. The training was conducted using a distributed data-parallel (DDP) strategy on a cluster of 16 NVIDIA A800 GPUs, completed in just 14 days. The core architecture of the LangYa system is shown in Figure 1a, which integrates multiple innovative components. The input state variables include ASVs, which are u- and v-components of wind speed, temperature, and relative humidity across 1000 to 200 hPa. Sea Surface State Variables (SSSVs): u- and v-components of 10-m wind speed, sea surface temperature, and mean sea level pressure. OSVs: Temperature, salinity, and currents spanning depths of 0 to -500 m. OSVs, ASVs, and SSSVs derived from the European Centre for Medium-Range Weather Forecasts (ECMWF) fifth-generation reanalysis dataset (ERA5) and GLORYS12 reanalysis datasets. The data is complemented by temporal information processed via a time-embedding module based on LLM (Figure 1b). The module extracts spatial features for individual time steps and temporal features across multiple time steps, and then integrates prior knowledge from the LLM to encode forecast durations. The unique fusion of spatiotemporal data supports highly efficient cross-spatiotemporal forecasting. A key innovation of LangYa is its Asynchronous Cross-Iterative Random Sampling Strategy (Figure 1c), which adaptively links ASVs, SSSVs, and OSVs to simulate air-sea interactions. Unlike traditional methods that rely on physics-based assumptions or simple data concatenation, this strategy learns spatiotemporal relationships and effectively captures air-sea variation patterns, significantly enhancing forecast accuracy without introducing additional physical constraints. The processed input data are passed through an encoder-decoder architecture equipped with an Ocean Self-Attention Module (Figure 1d). The module, based on cosine attention, ensures stable training and accelerates convergence by reducing gradient explosions common in high-dimensional data processing. Additionally, LangYa incorporates an Adaptive Thermocline Loss Function (Figure 1e) to address the challenges of forecasting thermocline mutations. The loss function is tailored to capture the steep vertical gradients and dynamic variability of the thermocline, improving




Fig. 1 Overview of the LangYa Forecasting System Architecture and Its Key Module. (A) Overall system pipeline integrating Time Embedding and deep feature extraction, (B) LLM-based Time Embedding module, (C) Asynchronous Cross-Iterative Random Sampling Strategy, (D) Ocean self-attention module based on cosine attention, and (E) Adaptive loss function for thermocline forecasts. W-MSA means window-based multi-head self-attention module. SW-MSA means shifted window-based multi-head self-attention.

forecast accuracy for this critical oceanic layer. By integrating these innovative components, LangYa has achieved several breakthroughs: It significantly reduces temporal distribution discrepancies and simplifies the complexity of air-sea coupling. It supports efficient, accurate, and stable forecasting across spatiotemporal scales. It yields state-of-the-art performance, particularly in challenging regions like thermoclines and areas with intense air-sea interactions. LangYa's comprehensive design represents a

major leap forward in AI-driven ocean forecasting, offering unprecedented efficiency and reliability for global ocean research and operational applications.

3 Experimental Setting and Main Results

The study trained the LangYa forecasting system using reanalysis data (GLORYS12 and ERA5) and evaluated it with both reanalysis (GLORYS12) and observational datasets. The observational datasets are derived from the GODAE Ocean View Inter-comparison and Validation Task Team (IV-TT) Class 4 framework, which provides observation datasets from Argo, Jason-1, Jason-2, and Envisat CLS AVISO level 3 satellite altimeters. 27 years of GLORYS12 data (1993-2019) were used for training LangYa, while data from 2020-2021 were reserved for testing. LangYa forecasts 128 variables, including temperature, salinity, and zonal and meridional ocean currents across 32 depth layers. Compared with reanalysis and observational data, LangYa achieved lower RMSEs for temperature and ocean currents (both zonal and meridional) than XiHe, XiHe-Autoregression (XiHe-AR), PSY4 [1], BLUElink OceanMAPS (BLK) [5], Global Ice Ocean Prediction System (GIOPS) [6], and Forecast Ocean Assimilation Model (FOAM) numerical models (Figure 2). Notably, the released XiHe version consists of 20 separate models, each designed to forecast OSVs for 1-10 days in either the upper or lower ocean layers. XiHe-AR refers to an autoregressive approach that generates multiday forecasts by sequentially chaining two 1-day XiHe models, resulting in higher cumulative errors. For salinity, LangYa’s forecast accuracy was comparable to XiHe and the numerical models but outperformed XiHe-AR.

In addition, LangYa achieved an inference time of 1 second for forecasts of any scale (1-7 days) on a single GPU, which is over 10,000 times faster than operational numerical models. Results from the 2020-2021 test data indicate that LangYa provided “perfect” quantitative forecasts (RMSE), particularly for challenging ocean phenomena such as thermoclines. Compared to the world’s leading OSVs forecasting systems (e.g., XiHe, PSY4, BLK, GIOPS, and FOAM numerical models), LangYa significantly reduced cumulative forecast errors and demonstrated remarkable improvements in thermocline forecasting.

Both quantitative and qualitative experimental results confirm that the AI-based LangYa forecasting system for OSVs holds tremendous potential for practical applications.

Figure 3 illustrates the RMSE of LangYa’s forecasts for temperature and salinity at 1, 4, and 7 days. Spatially, LangYa achieves RMSEs below 0.3°C for temperature forecasts across most ocean regions. Areas with higher RMSEs are concentrated in the western boundary currents (e.g., the Kuroshio Extension, Gulf Stream), eastern boundary currents (e.g., the Peru Current, California Current), and the Somali Current. These regions experience intense temperature variations, making them challenging to forecast accurately using numerical models or XiHe. Nevertheless, LangYa demonstrates significant advantages in these areas. For salinity, LangYa achieves RMSEs below 0.05 psu across most ocean regions, particularly in open oceans. Higher RMSEs are observed near the equator, western boundary currents, and the Indonesian Throughflow region, where salinity is heavily influenced by complex processes such




Fig. 2 Performance Comparison of LangYa, XiHe, XiHe-AR, and Numerical Models Across OSVs. A: RMSE of the u-component of ocean currents for 1-7 day forecasts. B: RMSE of the v-component of ocean currents for 1-7 day forecasts. C: RMSE of the ocean temperature for 1-7 day forecasts. D: RMSE of the salinity for 1-7 Day Forecasts.

as evaporation, precipitation, river runoff, and water mass exchange. Despite these complexities, LangYa delivers accurate salinity forecasts in these challenging regions. Furthermore, the spatial distribution of LangYa’s forecast errors for both temperature and salinity remains consistent as the forecast time increases, indicating the system’s high forecast stability.

Considering the significant differences in seawater characteristics across ocean basins, the global ocean was divided into eight basins (Northwest Pacific, Northeast Pacific, North Atlantic, North Indian Ocean, South Atlantic, South Pacific, South Indian Ocean, and Arctic Ocean) to evaluate LangYa’s forecasting performance. Figure 4 presents LangYa’s forecasting accuracy for four OSVs across these eight basins over 1-7 days. The results show substantial variations in RMSEs among the different ocean




Fig. 3 Global RMSE Distribution of LangYa Forecasts. Temperature and Salinity at 1, 4, and 7-day forecasts. A-C: RMSE distribution of temperature at 1, 4, and 7-day forecasts. D-F: RMSE distribution of salinity at 1, 4, and 7-day forecasts.

basins for forecasting ocean variables. Specifically, the Arctic Ocean exhibited the lowest RMSEs for U, V, and T, indicating that LangYa’s performance in this region is the most stable and accurate. This is due to the relatively stable hydrodynamic conditions and fewer drastic changes in the Arctic. Conversely, forecasting OSVs in the North Indian Ocean presented significant challenges, as monsoons, freshwater input, and complex oceanic dynamics influence the North Indian basin. The intricate physical processes and higher uncertainties posed difficulties for LangYa’s forecasts. In contrast, LangYa demonstrated more consistent forecasting performance in the other basins, showcasing the robustness of the LangYa forecasting system.

Notably, in salinity forecasts, there were significant differences in RMSEs between the North Indian Ocean and North Atlantic (relatively high values) and the two Southern Hemisphere basins (South Pacific and South Indian Ocean, relatively low values). Within the analysis in Figure 3, it can be inferred that the higher RMSEs in the North Atlantic are concentrated in the Gulf Stream extension region. The rapid salinity changes and spatial heterogeneity in this area are driven by freshwater input from the Gulf Stream and rivers. In the North Indian Ocean, the low accuracy of salinity forecasts is primarily due to the influence of complex monsoon systems and river runoff (e.g., Ganges and Brahmaputra Rivers). Additionally, the presence of coastal upwelling and the Indonesian Throughflow, which intensify water mass mixing, further complicates the forecasts in this region. These areas, characterized by complex physical processes and unique oceanic conditions, highlight the need for further development and refinement of the forecasting model.

4 Thermocline Forecast

The thermocline, a transitional layer in the ocean characterized by a sharp temperature gradient, is a critical interface for energy exchange, water mass mixing, and biogeochemical processes. Typically located at depths of 100-200 meter, the thermocline exhibits complex regional and seasonal variations due to its involvement in multiple coupled physical processes, such as ocean stratification, vertical mixing, and air-sea interactions. Accurate forecasting of the thermocline is essential for advancing




Fig. 4 RMSE of LangYa Forecasts for Temperature, Salinity, and Currents Across Eight Ocean Basins (1-7 Days). A-D: u-, v-component of ocean currents, temperature, and salinity forecast RMSE at different ocean basins.

the understanding of these processes and their implications for global climate systems, yet it has long been a challenge due to the steep gradients and non-linear dynamics in this layer.

LangYa uses a physics-guided adaptive loss function to address these challenges by enabling the model to focus on the thermocline’s complex temperature evolution. The advancement not only improves forecasting accuracy but also provides a deeper insight into the dynamics of thermocline formation, evolution, and interaction with surrounding layers.

Experimental results, shown in Figure 5, demonstrate that LangYa consistently outperforms numerical models and existing AI-based systems, such as XiHe, in temperature profile forecasting across all forecast horizons. The model’s superior performance is particularly evident near the thermocline, where traditional approaches struggle to capture the abrupt temperature changes. By minimizing RMSE near the thermocline, LangYa not only achieves unprecedented forecasting accuracy but also enhances the ability to simulate and analyze. In regions with intense air-sea interactions and surface mixing, LangYa’s performance remains robust, highlighting its ability to incorporate the stochastic and coupled nature of mixing processes. As depth increases and the model approaches the thermocline (identified by the pronounced rise in RMSE profiles), LangYa’s advantage becomes even more pronounced. The experimental




Fig. 5 Depth-Dependent RMSE Distributions of LangYa’s Temperature Forecasts at 1, 3, 5, and 7 Days. A: 1-day forecast result. B: 3-day forecast result. C: 5-day forecast result. D: 7-day forecast result.

result suggests that the adaptive loss function not only captures the vertical gradients but also enhances the model’s ability to generalize across varying oceanographic conditions.

The improved understanding and forecasting of thermocline dynamics enabled by LangYa provide significant implications for both scientific research and practical applications. By integrating advanced AI methods with physical knowledge, LangYa sets a new benchmark for forecasting accuracy and paves the way for more comprehensive studies on ocean stratification, climate variability, and marine resource management.

5 Discussion

LangYa forecasting system addresses the bottlenecks in cross-temporal, air-sea coupled, and thermocline forecasting. By integrating Time Embedding, air-sea interaction simulation, and adaptive loss function for thermocline, LangYa demonstrates significant advantages in forecasting OSVs across temporal scales. Compared with other forecasting systems (e.g., GOFs and XiHe), LangYa completes global OSVs inferencing within a short period and exhibits superior performance in accuracy, timeliness, and stability, offering feasibility for real-time ocean forecasting deployment.

Traditional ocean forecasting systems rely on iterative processes to estimate future OSVs, which are computationally expensive and prone to cumulative errors. LangYa overcomes the limitations by employing a Time Embedding module, enabling forecasts for up to 7 days without iterative steps. The innovation enhances its cross-temporal forecasting capabilities and improves forecasting flexibility and efficiency. In terms of simulating air-sea interactions, LangYa utilizes an asynchronous cross-iterative random sampling strategy, successfully overcoming the simplistic concatenation of air-sea

data used in existing AI-based ocean forecasting systems. The strategy effectively simulates the complex interactions between the atmosphere and ocean, significantly improving the forecasting accuracy of OSVs, such as temperature and currents. Moreover, LangYa’s training and convergence stability have been substantially enhanced through the incorporation of an ocean self-attention module. Ocean self-attention module optimizes the training process, accelerates convergence, and ensures stable forecasting outcomes.

A notable breakthrough of LangYa is its capability to forecast thermocline. The thermocline, a region in the ocean characterized by sharp temperature changes, has long been a challenging area in ocean science. By designing a loss function guided by physical knowledge, LangYa effectively captures the evolution characteristics of the thermocline, significantly improving forecasting accuracy. Experiments reveal that LangYa achieves lower RMSE in thermocline forecasts compared to existing AI-based ocean forecasting systems and numerical models, accurately reflecting the complex spatial and temporal variations of the thermocline.

LangYa’s experiments across eight global ocean basins further demonstrate its superior adaptability and stability. Whether in the relatively stable Arctic Ocean or the complex monsoon- and river-runoff-influenced ocean basins, LangYa maintains high forecasting accuracy and exhibits remarkable robustness across various ocean state variables. It indicates that LangYa can yield reliable forecasts even in dynamic and uncertain ocean environments, highlighting its broad application potential.

LangYa forecasting system provides a novel tool for oceanographic and Earth science research, advancing the integration of AI technologies with ocean science. Its innovative algorithm design and robust forecasting capabilities not only lay a theoretical foundation for real-time ocean forecasting systems but also offer precise and efficient support for global climate monitoring, marine resource management, and disaster early warning. With continuous technological advancements and data accumulation, LangYa is poised to become a core tool in global ocean science research, contributing significantly to the development of ocean science and its practical applications.

Despite its achievements, LangYa faces several limitations in practical applications: (1) LangYa is trained and tested on reanalysis data, whereas operational forecasting systems primarily rely on numerical assimilation data. The differences between reanalysis and numerical assimilation data may reduce model performance in applications, requiring further research into cross-data-source adaptability. (2) LangYa currently supports short-term forecasts of 1-7 days and has not yet been extended to medium- and long-term forecasts (e.g., monthly or seasonal scales). Addressing this challenge would require advancements in multi-scale coupling and capturing long-term dynamic changes. (3) LangYa focuses on forecasting OSVs and has not yet been extended to directly forecast ocean phenomena such as tropical cyclone tracks or El Niño events. The ocean phenomena hold greater scientific and practical value but require further breakthroughs in multivariable collaborative modeling. In the future, the integration of LangYa with numerical models will represent a novel approach to ocean forecasting research. The synergy will enable intelligent forecasting to expand from single-variable

predictions to complex phenomena, further enhancing the model’s applicability and scientific rigor.

References

- [1] Lellouche, J.-M., Greiner, E., Le Galloudec, O., Regnier, C., Benkiran, M., Testut, C.-E., Bourdalle-Badie, R., Drevillon, M., Garric, G., Drillet, Y.: Mercator ocean global high-resolution monitoring and forecasting system. *New Frontiers in Operational Oceanography*, 563–592 (2018)
- [2] Garraffo, Z.D., Cummings, J.A., Paturi, S., Iredell, D., Spindler, T., Balasubramanian, B., Rivin, I., Kim, H.-C., Mehra, A.: Rtofs-da: Real time ocean-sea ice coupled three dimensional variational global data assimilative ocean forecast system. *Research Activities in Earth System Modelling*, ed. E. Astakhova (Geneva: World Climate Research Programme) (2020)
- [3] Wang, X., Wang, R., Hu, N., Wang, P., Huo, P., Wang, G., Wang, H., Wang, S., Zhu, J., Xu, J., et al.: Xihe: A data-driven model for global ocean eddy-resolving forecasting. *arXiv preprint arXiv:2402.02995* (2024)
- [4] Xiong, W., Xiang, Y., Wu, H., Zhou, S., Sun, Y., Ma, M., Huang, X.: Ai-goms: Large ai-driven global ocean modeling system. *arXiv preprint arXiv:2308.03152* (2023)
- [5] Schiller, A., Brassington, G.B., Oke, P., Cahill, M., Divakaran, P., Entel, M., Freeman, J., Griffin, D., Herzfeld, M., Hoeke, R., et al.: Bluelink ocean forecasting australia: 15 years of operational ocean service delivery with societal, economic and environmental benefits. *Journal of Operational Oceanography* **13**(1), 1–18 (2020)
- [6] Smith, G.C., Roy, F., Reszka, M., Surcel Colan, D., He, Z., Deacu, D., Belanger, J.-M., Skachko, S., Liu, Y., Dupont, F., et al.: Sea ice forecast verification in the canadian global ice ocean prediction system. *Quarterly Journal of the Royal Meteorological Society* **142**(695), 659–671 (2016)
- [7] Jean-Michel, L., Eric, G., Romain, B.-B., Gilles, G., Angélique, M., Marie, D., Clément, B., Mathieu, H., Olivier, L.G., Charly, R., et al.: The copernicus global 1/12 oceanic and sea ice glorys12 reanalysis. *Frontiers in Earth Science* **9**, 698876 (2021)
- [8] Bell, B., Hersbach, H., Simmons, A., Berrisford, P., Dahlgren, P., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Radu, R., Schepers, D., et al.: The era5 global reanalysis: Preliminary extension to 1950. *Quarterly Journal of the Royal Meteorological Society* **147**(741), 4186–4227 (2021)
- [9] Ho, J., Jain, A., Abbeel, P.: Denoising diffusion probabilistic models. *Advances in neural information processing systems* **33**, 6840–6851 (2020)

- [10] Ryan, A., Regnier, C., Divakaran, P., Spindler, T., Mehra, A., Smith, G., Davidson, F., Hernandez, F., Maksymczuk, J., Liu, Y.: Godae oceanview class 4 forecast verification framework: global ocean inter-comparison. *Journal of Operational Oceanography* **8**(sup1), 98–111 (2015)

6 Methods

6.1 Dataset

The OSVs data used in LangYa are derived from the GLORYS12 reanalysis data [7]. GLORYS12 was developed and implemented under the framework of the Copernicus Marine Environment Monitoring Service (CMEMS). It has a horizontal resolution of $1/12^\circ$, 50 vertical layers, and a temporal spanning from January 1993 to December 2021. Due to its high resolution, temporal continuity, long coverage period, and excellent data quality, GLORYS12 has become the preferred dataset for large model development. Considering the important role of air-sea interactions in ocean dynamics, LangYa incorporates atmosphere state variables (ASVs, e.g. u- and v-components of wind speed, temperature, and relative humidity at 200, 500, 850, and 1000 hPa) and sea surface state variables (SSSVs, e.g. u- and v-components of 10-m wind speed, sea surface temperature, and mean sea level pressure) in the design. The ASVs and SSSVs are sourced from the ERA5 [8] produced by the European Centre for Medium Range Weather Forecasts (ECMWF). It has a spatial resolution of 0.25° and includes 37 vertical pressure levels.

6.2 Data Preprocessing

The input data for LangYa includes a total of 149 variables, categorized as follows: (1) 5 SSSVs: sea surface temperature, sea surface height, mean sea level pressure, u-, and v-components of 10-m wind speed. (2) 128 OSVs: These are distributed across 32 vertical depth layers (0.49, 1.54, 2.65, 3.82, 5.08, 6.44, 7.93, 9.57, 11.41, 13.47, 15.81, 18.50, 21.60, 25.21, 29.44, 34.43, 40.34, 47.37, 55.76, 65.81, 77.85, 92.32, 109.73, 130.67, 155.85, 186.13, 222.48, 266.04, 318.13, 380.21, 453.94, and 541.09 m), with four variables per depth layer: ocean temperature, salinity, u-, and v- components of ocean currents. (3) 6 ASVs: These are distributed across four pressure levels (200, 500, 850, and 1000 hPa), with four variables per level: air temperature, relative humidity, u-, and v-components of wind speed.

The study focuses on a global domain with longitude spanning -180° to 180° and latitude spanning -80° to 90° . The grid dimensions for OSVs are $H=4320$, $W=2041$. For consistent resolution, the ERA5 data were interpolated to a spatial resolution of $1/12^\circ$ to match the GLORYS12 reanalysis data using the nearest-neighbor interpolation method. The nearest-neighbor interpolation method was chosen for its simplicity and efficiency while preserving spatial consistency and spatiotemporal dependencies between grid points.

For each input channel, LangYa applies channel-wise Z-Score normalization to transform the data into a standard normal distribution, with a mean of 0 and a standard deviation of 1.

6.3 LangYa Forecasting System

LangYa was trained on 27 years of data from 1993 to 2019 and tested using data from 2020 and 2021. LangYa inputs OSVs, ASVs, and SSSVs, to forecast OSVs (channels, $C=128$) with a spatial resolution of $1/12^\circ$. As shown in Figure 1, the forecasting process can be expressed as:

$$\hat{X}_{t\tau} = \mathcal{F}_\theta (\mathcal{H} (X_t, A_{(t-7):t}) \mid E(t, \tau)) \quad (1)$$

where $X_t \in \mathbb{R}^{(D_x, H, W)}$ represents the OSVs at time t inputted to the forecasting system. It includes temperature, salinity, and ocean currents at 32 depth levels, which are combined along the channel dimension. $A_{(t-7):t} \in \mathbb{R}^{(D_A, H, W)}$ represents the historical ASVs over the previous 7 days, forming the contextual input for the forecasting system (The study primarily focuses on short-term forecasts of OSVs at a daily time resolution.). $\hat{X}_{t\tau}$ denotes the LangYa forecast for OSVs at time $\tau \in \{1, 2, \dots, K\}$. To standardize notation, $X_{t+\tau}^*$ represent the corresponding ground truth at the timestep $(t + \tau)$. $\mathcal{F}_\theta(\cdot)$ denotes LangYa’s ocean self-attention module. $\mathcal{H}(\cdot)$ represents LangYa’s modeling of air-sea interaction mechanisms, specifically implemented via the asynchronous cross-iterative random sampling strategy. $E(t, \tau)$ encodes the temporal information of LangYa’s initial and forecast horizons.

The specific forecasting steps of LangYa are as follows:

- (1) Coupling of OSVs, ASVs, and SSSVs: LangYa integrates OSVs, ASVs, and SSSVs using the asynchronous cross-iterative random sampling strategy within the air-sea coupled module. The process generates data with dimensions $2041 \times 4320 \times 133$. Subsequently, the Time Embedding module, based on LLMs, encodes the temporal information of the current and forecast times into a 96-dimensional feature vector. These two features are combined to form the input to LangYa.
- (2) Spatiotemporal Feature Extraction: LangYa employs an ocean self-attention module to extract spatiotemporal features, capturing the complex dependencies across spatial and temporal dimensions in the input data. For OSVs, LangYa utilizes a Patch Embedding module to perform dimensionality reduction by dividing the input grid into non-overlapping patches, with each patch encoded into a latent representation of dimension C . During experiments, the patch size is set to 4×4 , resulting in feature dimensions of $510 \times 1080 \times C$.
- (3) Deep Feature Extraction via Swin Transformer Blocks (ST Blocks): LangYa uses five ST blocks to extract deep features from the patch tokens. The ST blocks are organized hierarchically, where the number of tokens is progressively reduced through downsampling layers and then restored step by step through upsampling layers. Time Embeddings are incorporated into the ST blocks through tensor addition.
- (4) Upsampling for Forecast Generation: LangYa employs an upsampling module composed of interpolation and convolution operations to generate forecasted OSVs with dimensions $2041 \times 4320 \times 128$.

The technical details of the deep feature extraction architecture and the ST Block are as follows:

- (1) Deep Feature Extraction Architecture: For the first ST block, the input data has dimensions of $510 \times 1080 \times C$. In the subsequent three layers, the spatial dimensions are reduced by half, while the number of channels is doubled, resulting in data dimensions of $255 \times 540 \times 2C$. In the final ST block, the data dimensions are restored to $510 \times 1080 \times C$. During the downsampling process, 4 tokens are merged into 1, increasing the feature dimension from C to $4C$, and then reduced to $2C$ via a linear layer. During upsampling, the reverse operation is performed, restoring the data to its original dimensions.
- (2) Swin Transformer Block (ST Block): Each ST block consists of a Window-based Multi-Head Self-Attention (W-MSA) module followed by a Shifted Window-based Multi-Head Self-Attention (SW-MSA) module. Ocean data typically has extremely high spatial resolution, which imposes a significant computational burden on classical Transformers. To improve computational efficiency, the W-MSA module divides the input data into multiple small windows and applies the self-attention mechanism within each window for feature extraction. In our experiments, the window size is set to $[5, 12][5, 12][5, 12]$. However, window-based self-attention lacks connections between windows, which limits its modeling capacity. To introduce cross-window connections while maintaining the computational efficiency of non-overlapping windows, we employ the SW-MSA module, alternating W-MSA and SW-MSA modules within the ST block. In the ST block, the attention computation is performed only within each small window. Considering that the values of OSVs are closely related to the absolute geographic location on Earth, we incorporate Earth-specific positional encoding. The attention mechanism is computed as follows:

$$\text{Attention}(Q, K, V) = \text{SoftMax}\left(\frac{QK^T}{\|Q\|_2 \|K\|_2} + B\right)V \quad (2)$$

where Q represents the query, K the key, V the value, and B the Earth-specific positional bias.

6.4 Time Embedding Module

There are two methods for forecasting long-term time series of OSVs and ASVs: (1) autoregressive forecasting methods (e.g., Pangu-weather) and (2) building separate forecasting models for each forecast day (e.g., XiHe). The first method leads to significant cumulative errors, while the second entails high training costs due to the need for multiple models. Neither method satisfies the requirement of using a single model for long-term forecasting of OSVs, highlighting the urgent need to overcome this bottleneck. Inspired by recent progress in the denoising diffusion probabilistic model (DDPM) [9], LangYa employs time embedding as a carrier of temporal information to encode both the initial time and the forecast intervals. It enables a single model to achieve cross-temporal forecasting of OSVs for various initial times and forecast days. The periodic patterns exhibited by OSVs over years, seasons, and months are encoded into the initial time embedding. On the other hand, the nonlinear temporal variations over different forecast lead times are correspondingly encoded into

the interval time embedding. The combination of the above two items constitutes a complete time-embedded formalization:

$$E(t, \tau) \triangleq E(t_Y \oplus t_M \oplus t_D, \tau) = E_Y(t_Y) \oplus E_M(t_M) \oplus E_D(t_D) + E_\tau(\tau) \quad (3)$$

where Y , M , and D denote the values of year, month, and day, respectively. τ represents the forecast lead times. M and D are represented as discrete, learnable embedding vectors. Taking into account the fact that there is no clear periodic pattern in the numbering of years and that future years are not enumerable, we represent it as the linear combination of principal component embedding vectors:

$$E_Y(t_Y) \triangleq \sum_{i=1}^{N_Y} w_Y^{(i)} \cdot e_Y^{(i)} \quad (4)$$

where $\sum_{i=1}^{N_Y} w_Y^{(i)} = 1$ are the coefficients of the principal component vectors. These coefficients are derived from LLM with prior knowledge and are mapped to the required dimensions using a multilayer perceptron (MLP):

$$\left\{ w_Y^{(i)} \right\}_{i=1}^N = \text{MLP} [\text{LLM}(t_Y)] \quad (5)$$

In practice, the LLM is employed as the instruction-tuned Llama3.1-8B model. The following prompt is used for interaction with the LLM: "Global Ocean Status in the Year XXX." To obtain contextualized embeddings rather than word-wise embeddings, the hidden state embeddings from the last layer of the Llama model are utilized.

6.5 Air-Sea Coupling Module

The variations in OSVs do not occur in isolation but are driven by the complex interactions between the ocean and the atmosphere. These coupled relationships pose significant challenges for high-precision forecasting of OSVs, particularly under limited computational resources. Many existing models either fail to simultaneously handle ASVs, SSSVs, and OSVs data or inadequately account for their interactions, relying solely on simple data concatenation for modeling. Additionally, the inconsistency in spatial resolutions between ASVs, SSSVs, and OSVs data further complicates their integration within models.

To address the challenges, we propose a novel air-sea coupled training and sampling scheme (asynchronous cross-iterative random sampling strategy). The strategy simulates the interactions between the atmosphere and ocean by randomly sampling ASVs, SSSVs, and OSVs input data during the training process. The goal is to use ASVs data to achieve an accurate forecasting of OSVs and SSSVs. The core concept of the strategy is based on the fundamental fact that the temporal scale of atmospheric changes is significantly faster than that of oceanic changes. In other words, oceanic changes on a given day may be significantly influenced by atmospheric conditions from

several days prior. Accordingly, we define a temporal influence window of [3,7], indicating that ASVs influence the forecast of OSVs on a given day from the preceding 3 to 7 days. The process is formalized as:

$$H(X_t, A_{(t-7):t}) \triangleq [X_t \oplus SS_t] + H^{(A)} \left[\binom{1}{5} A_{(t-7):(t-3)} \right] \quad (6)$$

Here, SS refers to the SSSVs, while A denotes the ASVs. The operator $\binom{1}{5}$ represents the random selection of 1-day from the 5-day temporal window [3,7]. A linear interpolation operator $H^{(A)}$ is used to map 16-layer upper-air variables to the 133-layer OSVs and SSSVs. Specifically, the mechanism of the strategy works as follows: for the forecast of oceanic variables at the current time (t), OSVs may be influenced by ASVs between time t-7 and t-3. A specific time within the range is determined through random sampling. Similarly, at the current time t+1, the influence shifts to ASVs between time t-6 and t-2. The sliding-window-based random sampling mechanism effectively captures the stochasticity and complexity of air-sea interactions. The sampling strategy is integrated into the multi-node, multi-GPU training process and executed asynchronously. The design not only improves computational efficiency but also preserves the complex dynamic coupling relationships among the data. Therefore, it is called the Asynchronous Cross-Iterative Random Sampling Strategy.

The strategy not only provides a novel methodological perspective for modeling air-sea coupled but also offers significant technical support for high-resolution modeling under resource-constrained scenarios.

6.6 Ocean Self-Attention Module

OSVs, SSSVs, and ASVs often have very high resolutions, which makes gradient explosion (i.e., the rapid increase of deep activation values) more common during large model training. To address this issue, we propose a stability-enhancing module based on cosine attention. Two key modifications were made. First, we replaced the previous pre-normalization architecture with a post-normalization structure in the residual blocks. Thus, the output of each residual block is normalized before merging back into the main branch. The design ensures that the amplitude of the main branch does not accumulate as the network depth increases. The computation of each ST block is given by:

$$\begin{aligned} \hat{z}^1 &= \text{LN}(\text{W} - \text{MSA}(z^{1-1})) + z^{1-1} \\ z^1 &= \text{LN}(\text{MLP}(\hat{z}^1)) + \hat{z}^1 \\ \hat{z}^{1+1} &= \text{LN}(\text{SW} - \text{MSA}(z^1)) + z^1 \\ z^{1+1} &= \text{LN}(\text{MLP}(\hat{z}^{1+1})) + \hat{z}^{1+1} \end{aligned} \quad (7)$$

where z^{1-1} denotes the input to the ST block, z^{1+1} denotes the output of the block, "W-MSA" represents the window-based multi-head self-attention computation, SW-MSA represents the shifted-window multi-head self-attention computation, and LN denotes layer normalization.

In addition, ocean forecasting requires handling multidimensional input data, such as OSVs, ASVs, and SSSVs. The dot-product attention mechanism can be affected

by the scale variations of these high-dimensional inputs, leading to instability during training. To address it, we replaced the original dot-product attention with a scaled cosine attention mechanism (named ocean self-attention module), which ensures that the computation is independent of the input amplitude at each block.

6.7 Adaptive Thermocline Loss Function

The thermocline is a transition layer in the ocean where temperature changes drastically with depth. Its formation mechanism is complex, making accurate thermocline forecasting a significant challenge in ocean modeling. The difficulty of forecasting the thermocline lies in the sharp gradient changes in water temperature within this region, which are often difficult for traditional models to learn and capture effectively. To address this, we developed an adaptive thermocline loss function to enhance the model’s learning capability in the thermocline. The loss function is expressed as:

$$\mathcal{L}_T = \left| [X_T]_t^* - [\hat{X}_T]_t \right| \times \left[\text{Norm} \left(\frac{\partial [X_T]_t^*(x, y, z)}{\partial z} \right) + 1 \right] \quad (8)$$

where $[X_T]$ represents the grid for temperature, and $0 < x < W, 0 < y < H, 0 < z < C_T$ represents the grid coordinates for longitude, latitude, and depth. Norm denotes the min-max normalization, scaling the data to a range of $[0, 1]$. $[X_T]_t^*, [\hat{X}_T]$ represent the corresponding ground truth from the reanalysis and LangYa’s forecast values. The loss function enables the model to focus more on regions with significant vertical gradients, particularly the temperature variations near the thermocline, thereby significantly improving its ability to learn the complex evolution of the thermocline.

Based on the definition above, the overall training loss for LangYa is defined as:

$$\mathcal{L} = \left| X_t^* - \hat{X}_t \right| + \lambda \mathcal{L}_T \quad (9)$$

where $\lambda = 1.0$ is the weight for the thermocline loss.

6.8 Model Training and Metrics

LangYa was trained to forecast time ranging from 1 to 7 days. The air-sea coupled module considers atmospheric influences on the ocean for a period of 3 to 7 days. LangYa was trained for 100 epochs using the Adam optimizer. The training process was executed over 14 days on a cluster of 16 NVIDIA Tesla-A800 GPUs. The batch size was set to 16, and the initial learning rate was 0.001, which was gradually reduced to 0 during training using a cosine annealing strategy. To mitigate overfitting, the training dataset (1993-2019) was shuffled randomly within each epoch, while no random sampling was applied during the testing phase.

The study follows a recent work (XiHe) that adopted RMSE to measure the accuracy of forecasted OSVs compared to ground truth. To provide more objective experimental comparisons, MAE and PSNR are introduced as additional evaluation metrics. RMSE and MAE evaluate the global forecasting accuracy of variables from

a grid-level perspective, while PSNR captures the quality of the forecast data relative to the ground truth. The formula for calculating RMSE is defined as follows:

$$\begin{aligned}
\text{MAE} \left(X_t^*, \hat{X}_t \right) &= \frac{1}{C \cdot H \cdot W} \sum_{z,x,y}^{C,H,W} \left| X_t^*(z, x, y) - \hat{X}_t(z, x, y) \right| \\
\text{MSE} \left(X_t^*, \hat{X}_t \right) &= \frac{1}{C \cdot H \cdot W} \sum_{z,x,y}^{C,H,W} \left\| X_t^*(z, x, y) - \hat{X}_t(z, x, y) \right\|_2^2 \\
\text{RMSE} \left(X_t^*, \hat{X}_t \right) &= \sqrt{\text{MSE} \left(X_t^*, \hat{X}_t \right)} \\
\text{PSNR} \left(X_t^*, \hat{X}_t \right) &= 10 \cdot \log_{10} \frac{\max(X_t^*)^2}{\text{MSE} \left(X_t^*, \hat{X}_t \right)}
\end{aligned} \tag{10}$$

The results calculated using the above metrics are averaged across all time steps (testing set time range) and horizontal grid points to produce the mean RMSE, MAE, and PSNR values for the forecasted OSVs at a lead time. These metrics also remain robust in local regions, allowing for evaluation of RMSE, MAE, and PSNR for specific basins, such as Northwest Pacific, Northeast Pacific, etc.

6.9 Comparative Methods

Four numerical systems (PSY4, BLK, GIOPS, and FOAM) and one AI system (XiHe) that can be used for 1/125 numerical global thermosaline and current OSV forecasts are included in our comparison experiment.

The methods used to evaluate the model include the AI-based global OSVs forecasting system XiHe. XiHe achieves a high spatial resolution of approximately 1/12° and can complete a 10-day forecast within 0.36 seconds, which is thousands of times faster than traditional numerical Global Ocean Forecast Systems (GOFS). The evaluation also includes datasets from the IV-TT Class 4 framework, which provides observational datasets from Argo, Jason-1, Jason-2, and Envisat CLS AVISO level 3 satellite altimeters [10].

Additionally, the IV-TT Class 4 framework includes forecast results from numerical models (PSY4, BLK, GIOPS, and FOAM) at observational points for comparative validation. PSY4: Developed by Mercator Océan in France, this model features a high spatial resolution of approximately 1/12° and a forecast duration of 10 days. BLK: Developed by the Marine Hydrophysical Institute (MHI) in Russia, it offers a spatial resolution of 1/10° with a forecast duration of 7 days. GIOPS: Developed by Environment and Climate Change Canada (ECCC), this model has a spatial resolution of approximately 1/4° and a forecast duration of 10 days. FOAM: Developed by the Institute of Atmospheric Physics (IAP) of the Chinese Academy of Sciences, it features a spatial resolution of 0.5° with a forecast duration of 6 days.

7 Data availability

For training and testing LangYa, we downloaded the GLORYS12 dataset from https://data.marine.copernicus.eu/product/GLOBAL_MULTIYEAR_PHY_001_030/services and ERA5 dataset from <https://cds.climate.copernicus.eu/>. For comparison with other methods, we downloaded the IV-TT Class 4 framework from https://thredds.nci.org.au/thredds/catalog/rr6/intercomparison_files/catalog.html.

All these data are publicly available for research purposes. Source data will be provided in this paper.

8 Code availability

The code base of LangYa was established on PyTorch, a Pythonbased library for deep learning. The details of LangYa, including network architectures, modules, optimization tricks and hyperparameters, are available in the paper and the pseudocode. We released the trained models, inference code and the pseudocode of details to the public at a GitHub repository: https://github.com/iocaswolfteam/LangYa_v1.0. The trained models allow the researchers to explore LangYa’s ability on either GLORYS12 and ERA5 initial fields.