

# High-Rank Irreducible Cartesian Tensor Decomposition and Bases of Equivariant Spaces

**Shihao Shao**<sup>†</sup>

SHAOSHIHAO@PKU.EDU.CN

*State Key Lab of Vascular Homeostasis and Remodeling  
Department of Bioinformatics  
School of Basic Medical Sciences, Peking University*

**Yikang Li**

LIYK18@PKU.EDU.CN

*State Key Lab for General AI  
School of Intelligence Science and Technology  
Peking University, China*

**Zhouchen Lin**<sup>†</sup>

ZLIN@PKU.EDU.CN

*State Key Lab for General AI  
School of Intelligence Science and Technology, Peking University  
Pazhou Laboratory (Huangpu), China*

**Qinghua Cui**<sup>†</sup>

CUIQINGHUA@BJMU.EDU.CN

*State Key Lab of Vascular Homeostasis and Remodeling  
Department of Bioinformatics  
School of Basic Medical Sciences, Peking University*

## Abstract

Irreducible Cartesian tensors (ICTs) play a crucial role in the design of equivariant graph neural networks, as well as in theoretical chemistry and chemical physics. Meanwhile, the design space of available linear operations on tensors that preserve symmetry presents a significant challenge. The ICT decomposition and a basis of this equivariant space are difficult to obtain for high-order tensors. After decades of research, we recently achieve an explicit ICT decomposition for  $n = 5$  (Bonvicini, 2024) with factorial time/space complexity. This work, for the first time, obtains decomposition matrices for ICTs up to rank  $n = 9$  with reduced and affordable complexity, by constructing what we call path matrices. The path matrices are obtained via performing chain-like contraction with Clebsch-Gordan matrices following the parentage scheme. We prove and leverage that the concatenation of path matrices is an orthonormal change-of-basis matrix between the Cartesian tensor product space and the spherical direct sum spaces. Furthermore, we identify a complete orthogonal basis for the equivariant space, rather than a spanning set (Pearce-Crump, 2023b), through this path matrices technique. We further extend our result to the arbitrary tensor product and direct sum spaces, enabling free design between different spaces while keeping symmetry. The Python code is available in <https://github.com/ShihaoShao-GH/ICT-decomposition-and-equivariant-bases> where the  $n = 6, \dots, 9$  ICT decomposition matrices are obtained in 1s, 3s, 11s, and 4m32s, respectively.

**Keywords:** irreducible Cartesian tensor, Clebsch-Gordan coefficient, parentage scheme, equivariant graph neural networks, spherical harmonics

---

†. To whom correspondence should be addressed

## 1 Introduction

The deployment of equivariant neural networks in physics, chemistry, and robotics often requires certain equivariance or invariance to effectively incorporate inductive bias. Equivariant Graph Neural Networks (EGNNs) are well-suited for this purpose, as they can maintain equivariance while processing features within the graph (Zaverkin et al., 2024; Simeon and De Fabritiis, 2024; Batzner et al., 2022). One of the most promising approach to constructing equivariant layers involves leveraging irreducible representations. Depending on the basis of the space, two distinct types of irreducible representations are utilized in different EGNNs: spherical (spherical tensors) (Musaelian et al., 2023; Batatia et al., 2022) and Cartesian irreducible representations (Irreducible Cartesian Tensors, ICTs) (Zaverkin et al., 2024; Schütt et al., 2021) for  $O(3)$ . Spherical tensors serve as the starting point for EGNN based on irreducible representations. However, they require Clebsch-Gordan transforms with significant computational overhead to derive desired irreducible representations from existing ones during each inference, making them computationally intensive. In contrast, ICTs are more efficient, as high-order representations can be easily computed using tensor products (specifically, outer products in EGNNs), making them more computationally friendly for most deep learning frameworks. The ICTs are obtained through the ICT decomposition matrices  $H$ ,

$$\text{vec}\left(T^{(n;j;q)}\right) = H^{(n;j;q)}\text{vec}\left(T^{(n)}\right), \quad (1)$$

where  $T^{(n;j;q)}$  is an ICT,  $\text{vec}$  is the vectorization operation via flattening the tensor,  $T^{(n)}$  is a given tensor,  $n$  represents the rank of the tensor, and  $j$  is called the weight (the order of irreducible representations, which is distinct from the learnable weights in neural networks) of the irreducible representation, and  $q$  denotes the label (since the multiplicity for given  $n$  and  $j$  is sometimes greater than 1). The decomposition matrices  $H^{(n;j;q)}$  should be constructed such that  $T^{(n;j;q)}$  are irreducible. A formal definition is introduced in Section 3.1. The challenge lies in the fact that  $H$  of rank  $n > 5$  are unknown. The recent development of EGNNs research benefits from the irreducible decomposition matrices  $H$  obtained from the physics and chemistry communities. The decomposition matrices for  $n = 4$  have been established by Andrews and Ghoul (1982). However, only very recently have we obtained the decomposition matrices for  $n = 5$  (Bonvicini, 2024). Due to these limitations, state-of-the-art EGNNs cannot freely manipulate high-order ICTs or can only manipulate those whose weight equals their rank (Zaverkin et al., 2024).

It has been shown that  $l = 8$  irreducible representations can greatly benefit the model, but under the spherical basis (Liao et al., 2023). Therefore, the matrices for high-order orthogonal ICT decomposition are of great importance. Meanwhile, another crucial question is which linear operations applied to the Cartesian tensors are equivariant, as these operations constitute the design space of EGNNs, in other words, what is the vector space of linear mappings  $f$  such that

$$\rho(g)f(\text{vec}(T)) = f(\rho(g)\text{vec}(T)) \quad (2)$$

where  $g \in G$ , and  $\rho$  is the matrix representation. This vector space is called an equivariant space, also denoted as  $\text{End}_G\left(\left(\mathbb{R}^3\right)^{\otimes n}\right)$ . More generally, for different input and output

spaces, this equivariant space can be denoted as  $\text{Hom}_G\left(\left(\mathbb{R}^3\right)^{\otimes l}, \left(\mathbb{R}^3\right)^{\otimes k}\right)$ . Pearce-Crump (2023b) leverages Brauer (1937)’s theorem to find the spanning set of the equivariant space. However, as the rank increases, the number of elements in the spanning set becomes greater than the dimension of the space. Thus, it is also essential to identify a basis to reduce the number of parameters required.

In this work, one of our contributions is to efficiently obtain ICT decomposition matrices for  $6 \leq n \leq 9$ . Several previous methods (Andrews and Ghoul, 1982; Dinçkal, 2013; Bonvicini, 2024) rely on evaluating the full permutation of  $n$  (e.g., fundamental isotropic Cartesian tensors) and  $2n$  (e.g., natural projection tensors) indices, which requires factorial complexity, making it impractical for application to high-rank tensors. Bonvicini (2024) proposes a novel method to automate the procedure but does not reduce the complexity. The Reduced Row Echelon Form (RREF) algorithm is also required to run on an  $n! \times n!$  matrix. Instead, we leverage the fact that the Cartesian tensor product space is equivalent to a direct product of spherical spaces, up to a change of basis. In spherical space, it is straightforward to identify the desired irreducible representations, as they are diagonalized; then, the inverted change of basis can bring us back to Cartesian space. The challenge lies in finding such a change-of-basis matrix. We prove that one can construct this matrix by concatenating the results of sequential contractions and the outer products of Clebsch-Gordan matrices (CG matrices) according to the parentage scheme (Coope et al., 1965), which we refer to as path matrices. The change-of-basis matrix converts Cartesian tensors to the direct sum of spherical tensors.

If we treat tensors as vectors, we find that different spherical vectors are separated as distinct elements of the full-length vector. Thus, we can multiply it by a diagonalized matrix with only zeros and ones on the diagonals. This operation eliminates the other spherical vectors and retains only the desired one. The inverted change of basis can then convert the spherical direct product space back to the Cartesian tensor product space. The product of these three matrices forms the decomposition matrix, which decomposes the Cartesian tensor into the desired ICTs. This approach takes less than one second on CPUs to compute cases for  $n = 6$ , which is already faster than the current decomposition for  $n = 5$ . We also show that this process can be further accelerated. Specifically, we prove that path matrices have orthonormal columns up to multiplicative factors, which we leverage to demonstrate that a specific decomposition matrix can be generated via the product of the path matrix and its transpose.

Another challenge this work addresses is the construction of bases for the equivariant spaces. With the support of Schur’s lemma, we prove that the bases are formed by matrices that: 1) decompose a vector into its ICT, and 2) map an ICT to another ICT in an isomorphic space (i.e., with the same weight). These matrices are easy to construct in the spherical direct sum space, as their elements lie on the diagonal. Then, our path matrices can change the basis from the spherical space to the Cartesian tensor product space. We further extend our conclusions to linear equivariant operations between different direct sum and tensor product spaces. Additionally, we demonstrate the capability to construct bases for such general equivariant spaces, including spherical tensor product spaces with different parities. This enables the efficient construction of equivariant layers between different spaces, thereby expanding the design space of EGNNs.

Method	Rank	Orthogonality	Time/Space complexity
Coope et al. (1965)	3	No	calculated by hand
Andrews and Ghoul (1982)	4	No	calculated by hand
Dinçkal (2013)	3	Yes	calculated by hand
Bonvicini (2024)	5	No	factorial
Ours	6-9	Yes	exponential

Table 1: Comparison with previous works. The complexity is discussed in terms of rank. The complexity of this work will be polynomial if we focus on the dimensions.

This work has four main contributions:

- We propose a method to efficiently calculate high-rank ICT decomposition matrices and, for the first time, obtain the decomposition matrices for  $6 \leq n \leq 9$ , compared to predecessors, as shown in Table 1.
- We propose a general parentage scheme to extend ICT decomposition into a generalized ICT decomposition applicable to arbitrary tensor product spaces.
- We characterize the bases of equivariant spaces and propose a method to efficiently obtain these bases, overcoming the redundancy in Pearce-Crump (2023b)’s method.
- We extend our approach to efficiently obtain bases for equivariant spaces operating between different spaces.

These contributions enable EGNNs to process high-rank ICTs and flexibly design equivariant layers between desired spaces. Furthermore, this work benefits the physics and chemistry communities by facilitating theoretical analysis based on high-rank ICTs.

The following contents are organized as follows. In Section 1, we recall some basic ideas of representation theory, spherical harmonics, tensors, tensor product spaces, and we also introduce the Clebsch-Gordan coefficients and the parentage scheme. In Section 3, we formulate our problems: obtaining high-order ICT decomposition matrices and bases for arbitrary tensor product spaces. In Section 4, we introduce our theory and method to obtain high-order ICT decomposition matrices. In Section 5, we present the theory and method to obtain bases of equivariant spaces where the input and output spaces are the same. In Section 6, we extend the theory to the equivariant design spaces where the input and output spaces are different, and show the method to obtain bases for such general spaces. In Section 7, we review related works on designing EGNNs and calculating ICT decomposition matrices. In Section 8, we discuss the relation between Pearce-Crump (2023b) and this work, as well as the theory for  $SO(n)$ ,  $O(n)$ , and  $SU(n)$ . We provide concluding remarks in Section 9, and discuss the potential social impact in Section 10. We introduce the reproducibility details in Section 11. The acknowledgments are presented after the main text. The visualizations are

shown in the appendix. A list of mathematical symbols is provided in Table 2, and a mind map of this work is shown in Figure 1.

Table 2: Mathematical symbols and their meanings

Symbol	Meaning	Symbol	Meaning
$\ \cdot\ _{col}$	The $L_2$ -norm of an arbitrary column of $\cdot$	$G$	An arbitrary group
$SO(n)$	Special orthogonal group on $n$ -dimensional space	$SU(n)$	Special unitary group on $n$ -dimensional space
$O(n)$	Orthogonal group on $n$ -dimensional space	$U(n)$	Unitary group on $n$ -dimensional space
$\mathcal{S}$	A vector space	$\rho_{\mathcal{S}}$	A group representation
$\mathcal{S}^{\otimes n}$	A rank $n$ tensor product space from $\mathcal{S}$	$N^{(n;j)}$	The multiplicity of the ICT of rank $n$ and weight $j$
$T$	An arbitrary tensor $G \rightarrow \mathcal{S}$	$T^{(n)}$	A Cartesian tensor of rank $n$
$\hat{T}^{(n;j;q;p)}$	An ICT of rank $n$ , weight $j$ , index $q$ , and parity $p$ (sometimes omitted)	$Q_{\mathcal{S}}^{(l;p)}$	The set of all possible paths to decompose a tensor in space $\mathcal{S}$
$H^{(n;j;q)}$	An ICT decomposition matrix to obtain ICTs of rank $n$ , weight $j$ , and index $q$	$\hat{H}^{(\mathcal{S};j;q)}$	A general ICT decomposition matrix of space $\mathcal{S}$ , weight $j$ and index $q$
$C_{m_1 m_2 m_o}^{l_1 l_2 l_o}$	A CG coefficient with input weights $l_1, l_2$ , output weight $l_o$ , and the corresponding indices $m_1, m_2$ , and $m_o$ (the angular-momentum quantum numbers in physics)		
$C^{l_1 l_2 l_o}$	A CG tensor with input weights $l_1, l_2$ , and output weight $l_o$	$\hat{C}^{l_1 l_2 l_o}$	A CG matrix by merging the first two dimensions of the original CG tensor
$P^{(path)}$	A path matrix generated by $path$ , e.g., $(1 \rightarrow 3 \rightarrow 2)$	$\hat{P}^{(path)}$	A normalized path matrix generated by $path$
$D(\mathcal{S} \rightarrow \mathcal{S}')$	A change-of-basis matrix from space $\mathcal{S}$ to $\mathcal{S}'$	$v^{(l;p)}$	A representation weight $l$ and parity $p$ irreducible representation acts on
$\text{Hom}_G$	The set of $G$ -equivariant maps from one space to another	$\text{End}_G$	The set of $G$ -equivariant maps from one space to itself
irr	Returns all weights of irreducible representations in a decomposition	dim	The number of dimensions of a space
$(i_1 \xrightarrow{k_1} i_2 \xrightarrow{k_2} \dots)$	A path where $k_*$ are bridge numbers (omitted if all being 1)	$(l, p)$	An irreducible representation space of weight $l$ and parity $p$
$\otimes$	Tensor product	$\oplus$	Direct sum
$\odot$	Contraction	$\#$	Number of elements in a set

## 2 Preliminaries

### 2.1 Clebsch-Gordan coefficients

A review on irreducible representations and spherical spaces is in the appendix. The term *weights* in this paper refer to the order of irreducible representations. An important tool utilized in our work is the Clebsch-Gordan coefficients (CG coefficients). They are real numbers, and describe how to couple vectors from two spherical spaces into a new one. There are some rules for this coupling. First, given two vectors in weights  $l_1$  and  $l_2$  spherical spaces, the weights of the resulting vectors can only lie in  $|l_1 - l_2| \leq l_o \leq l_1 + l_2$ . For each index  $m_o$  of the obtained vector  $v_{l_o} \in (l = l_o)$ , it is calculated by homogeneous quadratic polynomials. Specifically, we have

$$v_{m_o}^{(l_o)} = \sum_{m_o=m_1+m_2} C_{m_1 m_2 m_o}^{l_1 l_2 l_o} v_{m_1}^{(l_1)} v_{m_2}^{(l_2)}, \quad (3)$$

and if we consider  $O(n)$  instead of  $SO(n)$ , parity  $p$  should also be considered. The parity is a property additional to the rank and weight, being 1 or  $-1$ . Having  $-1$  parity means that the representation will be inverse when the space is inverse, and *vice versa*. For example, the natural representations of  $O(n)$  always have  $p = -1$ . We will introduce the case that  $p = 1$  in the next section. Taking the parity into consideration, we can guarantee  $O(n)$  equivariance, and equation (3) becomes

$$v_{m_o}^{(l_o;p_o)} = \mathbb{1}_{(p_o=p_1 p_2)} \sum_{m_o=m_1+m_2} C_{m_1 m_2 m_o}^{l_1 l_2 l_o} v_{m_1}^{(l_1;p_1)} v_{m_2}^{(l_2;p_2)},$$

where  $\mathbb{1}_{(exp)}$  is the indicator function, outputting 1 if *exp* holds true, or 0 otherwise. Our discussion will be the same no matter we consider  $O(n)$  or  $SO(n)$ . But since Section 6, we need to take care of the parity which will play some roles in the further discussion. The CG coefficients are highly sparse, since  $C_{m_1 m_2 m_o}^{l_1 l_2 l_o} = 0$  if  $m_o \neq m_1 + m_2$ . The coefficients are identical to Wigner-3j symbols up to multiplicative factors, so we will adopt Wigner-3j for the implementation. The evaluation of CG coefficients is nearly  $\mathcal{O}(1)$  (if we do not consider the few calculations for the closed-form formula, which is cheap enough). To obtain the coefficients of  $O(3)/SO(3)$ , one often leverages the relation between  $SU(2)$  and  $SO(3)$ :  $SU(2)$  is a double cover of  $SO(3)$ , so the CG matrices of  $SU(2)$  and  $SO(3)$  are same up to a change-of-basis, from complex to real. The closed-form formulae for  $SU(2)$  CG coefficients are already known. Thus, this process can be efficiently evaluated without obvious overhead.

### 2.2 Tensors and tensor product spaces

To formally define a tensor, we need to consider vector space  $V$  and covector space  $V^*$ . A tensor product is a bilinear map  $\otimes : \mathcal{S} \times \mathcal{S}' \rightarrow \mathcal{S} \otimes \mathcal{S}'$ . The basis of  $\mathcal{S} \otimes \mathcal{S}'$  is the set of  $\hat{x}_i \otimes \hat{y}_j \in \mathcal{S} \otimes \mathcal{S}'$ , where  $\hat{x}_i$  and  $\hat{y}_j$  are basis vectors of  $\mathcal{S}$  and  $\mathcal{S}'$ , respectively. That is why one implements outer product as tensor product, as it meets the definition of tensor product. A tensor  $T : V^{\otimes r} \otimes V^{*\otimes s} \rightarrow F$ , where  $F$  is some field, is a mathematical object that converts things between different tensor product spaces. For example, the representation  $\rho(g)$  is a  $(r = 1, s = 1)$ -tensor that receives a vector  $v$  which is a  $(r = 1, s = 0)$ -tensor, and output another  $(r = 1, s = 0)$ -tensor  $\rho(g)v$ , as  $r$  and  $s$  vanish correspondingly. Tensor




Figure 1: The mind map of our method. The blocks in orange color are aligned with the three contributions of this work.

product can help us build up the ranks of tensors, e.g.,  $V \otimes V \otimes V$  is a  $(r = 3, s = 0)$  space. The representation space is accordingly  $(r = 3, s = 3)$ . Conversely, contraction can reduce the ranks of tensors. We express these operations via some index notations. For example,  $T_{i_1 i_2} = T'_{i_1} T''_{i_2}$  is a tensor product, and  $T_{i_1} = \sum_{j_1} T'_{i_1 j_1} T''_{j_1}$  is a contraction. Obviously, such operations can be mixed together, and we often omit the index of the output tensors and agree  $i_k$  will be placed at the  $k$ th index of the output tensor by default. We also agree that  $j_1$  and  $j_2$  will vanish due to the contraction. We will not use Einstein summation convention in this work to keep the conventions of machine learning community, and to avoid ambiguous during derivations. One should note that contractions can be expressed in term of matrix representation. Consider  $T = \sum_{j_1, j_2} T'_{i_1 j_1 j_2} T''_{j_1 j_2}$ , we have

$$\begin{bmatrix} T_1 \\ T_2 \end{bmatrix} = \begin{bmatrix} T'_{111} & T'_{112} & T'_{121} & T'_{122} \\ T'_{211} & T'_{212} & T'_{221} & T'_{222} \end{bmatrix} \begin{bmatrix} T''_{11} \\ T''_{12} \\ T''_{21} \\ T''_{22} \end{bmatrix}.$$

It is useful when we need to obtain the inverse of some linear operations. Also, contractions and tensor products are both  $O(n)$ -equivariant.

### 2.3 CG matrices and the parentage scheme

The  $O(3)$  group has its representations in tensor product space  $(\mathbb{R}^3)^{\otimes n}$ , which are some  $3^n \times 3^n$  matrices, or tensor product of  $n$   $(r = 1, s = 1)$ -tensors, which depends on one's view. If we interpret them as matrices, then the objects they act on are simply  $3^n$ -dimensional




Figure 2: The parentage scheme. The numbers in the circle represent multiplicity. The sum of each column equals equation (4).

vectors. The tensor product spaces  $(\mathbb{R}^3)^{\otimes n}$  for  $n > 1$  are generally reducible. Recall CG coefficients, two irreducible spherical vectors  $v$  and  $v'$  coupling can also be viewed as that they first form a rank 2 tensor, and then decompose into different new spherical vectors. These operations can also be viewed as a product of matrices and vectors. In this context, the CG coefficients  $\hat{C}^{l_1 l_2 l_o}$  form a  $(2l_1 + 1)(2l_2 + 1) \times (2l_o + 1)$  matrix, which we call a CG matrix. If we reshape it to a  $(2l_1 + 1) \times (2l_2 + 1) \times (2l_o + 1)$  tensor, then we instead call it an  $(l_1, l_2, l_o)$ -tensor  $C^{l_1 l_2 l_o}$ . Here are some important properties,

**Lemma 1 (Orthogonality)** *Given an  $(l_1, l_2, l_o)$ -CG tensor  $C^{l_1 l_2 l_o}$ , we have*

$$\sum_{j_1, j_2} C_{j_1 j_2 k}^{l_1 l_2 l_o} C_{j_1 j_2 k'}^{l_1 l_2 l_o} = \sum_{j_1, j_2} C_{j_1 k j_2}^{l_1 l_2 l_o} C_{j_1 k' j_2}^{l_1 l_2 l_o} = \sum_{j_1, j_2} C_{k j_1 j_2}^{l_1 l_2 l_o} C_{k' j_1 j_2}^{l_1 l_2 l_o} = 0$$

for  $k \neq k'$ . More generally, different CG tensors with the same  $l_1$  and  $l_2$ ,  $l_1$  and  $l_3$ , or  $l_2$  and  $l_3$  are also orthogonal in this way.

**Lemma 2 (Norm)** *Given an  $(l_1, l_2, l_o)$ -CG tensor  $C^{l_1 l_2 l_o}$ , we have*

$$\sum_{j_1, j_2} (C_{j_1 j_2 k}^{l_1 l_2 l_o})^2 = \sum_{j_1, j_2} (C_{j_1 j_2 k'}^{l_1 l_2 l_o})^2; \quad \sum_{j_1, j_2} (C_{j_1 k j_2}^{l_1 l_2 l_o})^2 = \sum_{j_1, j_2} (C_{j_1 k' j_2}^{l_1 l_2 l_o})^2; \quad \sum_{j_1, j_2} (C_{k j_1 j_2}^{l_1 l_2 l_o})^2 = \sum_{j_1, j_2} (C_{k' j_1 j_2}^{l_1 l_2 l_o})^2$$



for any pair of  $k$  and  $k'$ .

**Lemma 3 (Equivariance)** *An  $(l_1, l_2, l_o)$ -CG tensor  $C^{l_1 l_2 l_o}$  maintains the following equivari-  
ances,*

$$\begin{aligned}
& \rho_{l_1 \otimes l_2}(g) \odot C^{l_1 l_2 l_o} \\
&= \sum_{j_1, j_2} R_{i_1 j_1}^{l_1} R_{i_2 j_2}^{l_2} C_{j_1 j_2 i_3}^{l_1 l_2 l_o} \\
&= \sum_{j_1} C_{i_1 i_2 j_1}^{l_1 l_2 l_o} R_{j_1 i_3}^{l_3} \\
&= C^{l_1 l_2 l_o} \odot \rho_{l_o}(g),
\end{aligned}$$

where  $\odot$  denotes the tensor contraction,  $R^l$  is the matrix representation in  $l$  space. This equation holds for other permutations as Lemmas 1 and 2.

The above lemmas cover the CG matrices case. Since the columns of CG matrices have the same  $L_2$  norm, we denote as  $\|\cdot\|_{col}$  for shorthand, and we sometimes call the last index of the CG tensors as the index of their columns. Recall that CG matrices are non-zero if and only if  $|l_1 - l_2| \leq l_o \leq l_1 + l_2$ . If we start from an  $l = 1$  vector and couple an additional  $l = 1$  once a time, then this process has been drawn as a parentage scheme by Coope et al. (1965), as shown in Figure 2. Each time we move from left to closed right column we couple an additional  $l = 1$  vector. The parentage scheme also tells us which irreducible representations can a rank  $n$  tensor be decomposed into. Note that a rank  $n$  tensor can be decomposed into multiple spherical vectors, and the multiplicity for each weight given the rank is written in the scheme. Equivalently, it can also be decomposed into same numbers of ICTs, but in the Cartesian spaces. We also have a closed-form formula for the multiplicity,

**Proposition 4** *The multiplicities of the weight  $j$  ICTs of a rank  $n$  tensor is*

$$N^{(n;j)} = \sum_i^{\lfloor (n-j)/3 \rfloor} (-1)^i \frac{n!(2n-3i-j-2)!}{i!(n-i)!(n-2)!(n-3i-j)!}. \quad (4)$$

Note that the equation scales slower or equal to exponential, as it is smaller than  $3^n$ , the dimension of the tensor product space.

### 3 Problem formulation

We first formulate the task of obtaining ICT decomposition matrices. The properties derived during the process of obtaining these decomposition matrices are further leveraged to construct the bases for equivariant linear mappings between tensor product spaces.

#### 3.1 ICT decomposition

In this paper, one of our goal is to find an efficient way to obtain a set of matrices  $H$  to decompose a rank  $n$  tensor into ICTs with different weight  $l$  and index  $q$ , as shown in

equation (1). The resulting ICTs are in the Cartesian tensor product spaces, not in the spherical ones. Here, we precisely define what an ICT decomposition is. In other words, we specify the properties that the matrices  $H$  must satisfy:

**Definition 5 (ICT decomposition)** *Given a Cartesian tensor product space  $(\mathbb{R}^3)^{\otimes n}$  formed by tensor product of Cartesian spaces  $\mathbb{R}^3$ , an ICT decomposition is a set of decomposition matrices  $H^{(n;l;q)} \in \mathbb{R}^{3^n \times 3^n}$ , where  $l$  is the weight of ICTs, and  $q$  is the index of the multiplicities, such that:*

- 1)  $H^{(n;l;q)}$  are  $O(3)$ - $((\mathbb{R}^3)^{\otimes n}, (\mathbb{R}^3)^{\otimes n})$ -equivariant.
- 2) The summation of the the decomposition matrices is an identity matrix, i.e.,  $\sum_{l,q} H^{(n;l;q)} = I$ .
- 3)  $\{H^{(n;l;q)}\}$  is linearly independent.

Furthermore, if we have

$$\left(H^{(n;l;q)}v\right) \cdot \left(H^{(n;l';q')}v\right) = 0 \quad (5)$$

for any  $v \in (\mathbb{R}^3)^{\otimes n}$ , and  $l \neq l'$  or  $q \neq q'$ , then we further call it an orthogonal ICT decomposition, and the matrices orthogonal decomposition matrices.  $H^{(n;l;q)}v$  is called an ICT with rank  $n$ , weight  $l$ , and index  $q$ .

Note that ICT decomposition is not necessary to be orthogonal, but being orthogonal is the ultimate goal for ICT decomposition. To prove the orthogonality, we have

**Lemma 6** *Given matrices  $M : \mathcal{S} \rightarrow \mathcal{S}'$  and  $M' : \mathcal{S} \rightarrow \mathcal{S}'$  of the same shape, if  $M^T M' = 0$ , then*

$$(Mv) \cdot (M'v) = 0 \quad (6)$$

for any  $v \in \mathcal{S}$ .

**Proof** It holds true that

$$\begin{aligned} & (Mv) \cdot (M'v) \\ &= \sum_i \left( \sum_j M_{ij} v_j \right) \left( \sum_{j'} M'_{ij'} v_{j'} \right) \\ &= \sum_{j,j'} v_j v_{j'} \left( \sum_i M_{ij} M'_{ij'} \right). \end{aligned} \quad (7)$$

Assume we have  $M^T M' = 0$ , then for any  $j$  and  $j'$ , we have

$$\sum_i M_{ij} M'_{ij'} = 0. \quad (8)$$

Plug it back to equation (7), we obtain

$$(Mv) \cdot (M'v) = \sum_{j,j'} v_j v_{j'} \left( \sum_i M_{ij} M'_{ij'} \right) = 0. \quad (9)$$

This concludes the proof. ■

This work obtains orthogonal ICT decomposition matrices for tensor product spaces with rank  $6 \leq n \leq 9$ . A comparison with predecessors is shown in Table 1.

### 3.2 Basis of the equivariant design space

Given a tensor  $T \in (\mathbb{R}^3)^{\otimes n}$ , a critical challenge in the design of EGNNs is determining which linear operations acting on the tensor are equivariant, i.e., find the set of  $f$  in equation (2), denoted as  $\text{End}_{O(3)}((\mathbb{R}^3)^{\otimes n})$ . In real implementation, we flatten  $T$  to vectors  $v$ , and the linear mappings  $f$  are therefore matrices. Pearce-Crump (2023b) leverages Brauer (1937)'s theorem to obtain spanning sets for homomorphism spaces. However, the spanning set becomes increasingly redundant as the rank of tensors rises. In this work, we aim to efficiently find bases, instead of spanning sets, for equivariant spaces.

Additionally, we seek to find bases for the equivariant spaces  $\text{End}_{O(3)}(\mathcal{S}, \mathcal{S}')$ , where  $\mathcal{S}$  is not necessarily equal to  $\mathcal{S}'$ . Here,  $\mathcal{S}$  and  $\mathcal{S}'$  can be arbitrary direct sums of tensor product spherical spaces (which also include Cartesian spaces, recall Lemma 22). This approach enables us to design EGNNs more freely by making the domain and co-domain of equivariant layers steerable. In other words, given arbitrary domain and its co-domain, one can efficiently construct equivariant layers without redundant parameters. A discussion of the relationship between Pearce-Crump (2023b) and this work is provided in Section 8.1.

## 4 High-order orthogonal ICT decomposition

Our key to obtaining the orthogonal decomposition matrix is to find a change-of-basis matrix between irreducible spherical spaces and tensor product spaces. The first step is to construct what we call path matrices. We have first obtained the following lemmata:

**Lemma 7** *Given an  $(l_x \otimes l_y, l_2)$ -CG matrix  $\hat{C}^{l_x l_y l_2}$  and an  $(l_1, l_2, l_3)$ -CG tensor,  $C^{l_1 l_2 l_3}$ , if we let*

$$T = \sum_{j_1} \hat{C}_{i_2 j_1}^{l_x l_y l_2} C_{i_1 j_1 i_3}^{l_1 l_2 l_3} \quad (10)$$

*and we flatten  $T$  into a  $(2l_1 + 1)(2l_x + 1)(2l_y + 1) \times (2l_3 + 1)$  matrix, then the following statements hold true:*

- 1)  $T$  is  $O(3)$ - $(l_1 \otimes l_x \otimes l_y, l_3)$ -equivariant.
- 2) The columns of  $T$  are orthogonal.
- 3) The columns have the same  $L_2$ -norms.

**Proof** We first prove 1), given an arbitrary group element  $g \in O(3)$ ,

$$\begin{aligned}
 & \rho_{l_1 \otimes l_x \otimes l_y}(g) \odot T \\
 &= \rho_{l_1 \otimes l_x \otimes l_y}(g) \odot \left( \sum_{j_1} \hat{C}_{i_2 j_1}^{l_x l_y l_2} C_{i_1 j_1 i_3}^{l_1 l_2 l_3} \right) \\
 &= \sum_{i'_1, i'_2} R_{i_1 i'_1}^{l_1} R_{i_2 i'_2}^{l_x \otimes l_y} \left( \sum_{j_1} \hat{C}_{i'_2 j_1}^{l_x l_y l_2} C_{i'_1 j_1 i_3}^{l_1 l_2 l_3} \right) \\
 &= \sum_{j_1} \left( \sum_{i'_2} R_{i_2 i'_2}^{l_x \otimes l_y} \hat{C}_{i'_2 j_1}^{l_x l_y l_2} \right) \left( \sum_{i'_1} R_{i_1 i'_1}^{l_1} C_{i'_1 j_1 i_3}^{l_1 l_2 l_3} \right) \\
 &\stackrel{(a)}{=} \sum_{j_1} \left( \sum_{j'_1} \hat{C}_{i_2 j'_1}^{l_x l_y l_2} R_{j'_1 j_1}^{l_2} \right) \left( \sum_{i'_1} R_{i_1 i'_1}^{l_1} C_{i'_1 j_1 i_3}^{l_1 l_2 l_3} \right) \\
 &= \sum_{j_1} \hat{C}_{i_2 j_1}^{l_x l_y l_2} \left( \sum_{i'_1, j'_1} R_{j'_1 j_1}^{l_2} R_{i_1 i'_1}^{l_1} C_{i'_1 j'_1 i_3}^{l_1 l_2 l_3} \right) \\
 &\stackrel{(b)}{=} \sum_{j_1, i'_3} \hat{C}_{i_2 j_1}^{l_x l_y l_2} C_{i_1 j_1 i'_3}^{l_1 l_2 l_3} R_{i'_3 i_3}^{l_3} \\
 &= T \odot \rho_{l_3}(g),
 \end{aligned} \tag{11}$$

so  $T$  is  $O(3)$ - $(l_1 \otimes l_x \otimes l_y, l_3)$ -equivariant, where  $R^l$  is the matrix representation in  $l$  space, equalities (a) and (b) hold true because of Lemma 3.

Next we prove 2), for an arbitrary pair of the indexes of two columns  $k \neq k'$ , we have

$$\begin{aligned}
 & \sum_{j_1} T_{j_1 k} T_{j_1 k'} \\
 &= \sum_{j_1, j_3} \left( \sum_{j_2} \hat{C}_{j_1 j_2}^{l_x l_y l_2} C_{j_3 j_2 k}^{l_1 l_2 l_3} \right) \left( \sum_{j'_2} \hat{C}_{j_1 j'_2}^{l_x l_y l_2} C_{j_3 j'_2 k'}^{l_1 l_2 l_3} \right) \\
 &= \sum_{j_2, j'_2} \left( \sum_{j_1} \hat{C}_{j_1 j_2}^{l_x l_y l_2} \hat{C}_{j_1 j'_2}^{l_x l_y l_2} \right) \left( \sum_{j_3} C_{j_3 j_2 k}^{l_1 l_2 l_3} C_{j_3 j'_2 k'}^{l_1 l_2 l_3} \right) \\
 &\stackrel{(a)}{=} \sum_{j_1, j_2} \left( \hat{C}_{j_1 j_2}^{l_x l_y l_2} \right)^2 \left( \sum_{j_3} C_{j_3 j_2 k}^{l_1 l_2 l_3} C_{j_3 j_2 k'}^{l_1 l_2 l_3} \right) \\
 &= \|\hat{C}^{l_x l_y l_2}\|_{col}^2 \left( \sum_{j_2, j_3} C_{j_3 j_2 k}^{l_1 l_2 l_3} C_{j_3 j_2 k'}^{l_1 l_2 l_3} \right) \\
 &\stackrel{(b)}{=} 0,
 \end{aligned} \tag{12}$$

and thus the columns of  $T$  are orthogonal. Equalities (a) and (b) follow from Lemma 1 that  $\sum_{j_1} \hat{C}_{j_1 j_2}^{l_x l_y l_2} \hat{C}_{j_1 j_2'}^{l_x l_y l_2} = 0$  for  $j_2 \neq j_2'$ , and  $\sum_{j_2, j_3} C_{j_3 j_2 k}^{l_1 l_2 l_3} C_{j_3 j_2 k'}^{l_1 l_2 l_3} = 0$ .

Finally, we prove 3), for an arbitrary column with index  $k$ ,

$$\begin{aligned}
& \sqrt{\sum_{j_1} T_{j_1 k} T_{j_1 k}} \\
&= \sqrt{\sum_{j_1, j_3} \left( \sum_{j_2} \hat{C}_{j_1 j_2}^{l_x l_y l_2} C_{j_3 j_2 k}^{l_1 l_2 l_3} \right) \left( \sum_{j_2'} \hat{C}_{j_1 j_2'}^{l_x l_y l_2} C_{j_3 j_2' k}^{l_1 l_2 l_3} \right)} \\
&= \sqrt{\sum_{j_2, j_2'} \left( \sum_{j_1} \hat{C}_{j_1 j_2}^{l_x l_y l_2} \hat{C}_{j_1 j_2'}^{l_x l_y l_2} \right) \left( \sum_{j_3} C_{j_3 j_2 k}^{l_1 l_2 l_3} C_{j_3 j_2' k}^{l_1 l_2 l_3} \right)} \tag{13} \\
&= \sqrt{\sum_{j_2} \left( \sum_{j_1} \left( \hat{C}_{j_1 j_2}^{l_x l_y l_2} \right)^2 \sum_{j_3} \left( C_{j_3 j_2 k}^{l_1 l_2 l_3} \right)^2 \right)} \\
&= \|\hat{C}^{l_x l_y l_2}\|_{\text{col}} \sqrt{\sum_{j_2, j_3} \left( C_{j_3 j_2 k}^{l_1 l_2 l_3} \right)^2} \\
&= \|\hat{C}^{l_x l_y l_2}\|_{\text{col}} \|C^{l_1 l_2 l_3}\|_{\text{col}}.
\end{aligned}$$

From Lemma 2, we know that it takes the same value for different  $k$ , which satisfies 3). ■

Here is the strategy, we first let a  $(0 \otimes 0, 0)$ -CG matrix to contract with an  $(1, 0, 1)$ -CG tensor to obtain an  $(0 \otimes 1, 1)$ -matrix. This matrix can further contract with  $(1, 1, 2)$ ,  $(1, 1, 1)$ , and  $(1, 1, 0)$ -CG tensors to obtain the associated matrix. For example, we call the obtained  $(0 \otimes 1 \otimes 1, 2)$  a  $(0 \rightarrow 1 \rightarrow 2)$ -path matrix. The path matrices are constructed via the parentage scheme, which will decide what is the next CG tensor we are going to contract with, as shown in Figure 3. This gives us:

**Corollary 8** *Given a path matrix  $P^{(0 \rightarrow \dots \rightarrow l_2)}$  and a CG tensor  $C^{1, l_2 l_3}$ , if we let  $P^{(0 \rightarrow \dots \rightarrow l_2 \rightarrow l_3)} = \sum_{j_1} P_{i_2 j_1}^{(0 \rightarrow \dots \rightarrow l_2)} C_{i_1 j_1 i_3}^{1, l_2 l_3}$ , and we flatten  $P^{(0 \rightarrow \dots \rightarrow l_2 \rightarrow l_3)}$  as we did in Lemma 7, then we have the same statements hold as Lemma 7.*

**Proof** The path matrices have all the properties we used in Lemma 7. We prove this by a simple deduction. From the above lemma we know that  $(0 \rightarrow 1)$ -path matrix holds all the properties. This gives us the initial condition. Assume that a matrix holds the properties, then Lemma 7 tells us the next obtained path matrix also share them. ■

Next, we prove that




Figure 3: Construction of decomposition matrix. **Top:** The path matrices are formed via sequential contraction according to the path in the parentage scheme. **Bottom:** The decomposition matrix is obtained as the product of the path matrix and its transpose.

**Lemma 9** *The columns of different path matrices of the same path length are orthogonal.*

**Proof** We prove by deduction. The first to third columns of the parentage scheme generate  $(0 \rightarrow 1 \rightarrow 2)$ ,  $(0 \rightarrow 1 \rightarrow 1)$ , and  $(0 \rightarrow 1 \rightarrow 0)$ -path matrices. We first prove that the columns of these matrices are orthogonal. Since they come from the contraction of  $(0 \rightarrow 1)$ -path matrices and different CG matrices, we can prove a more general conclusion. That is, for any path matrix  $P$  and two different CG tensors  $C$  and  $C'$ , given two different column indexes

of there two tensors  $k$  and  $k'$ , consider the dot product of these two columns, we have

$$\begin{aligned}
& \sum_{j_1, j_3} \left( \sum_{j_2} P_{j_1 j_2} C_{j_3 j_2 k} \right) \left( \sum_{j'_2} P_{j_1 j'_2} C'_{j_3 j'_2 k'} \right) \\
&= \sum_{j_2, j'_2} \left( \sum_{j_1} P_{j_1 j_2} P_{j_1 j'_2} \right) \left( \sum_{j_3} C_{j_3 j_2 k} C'_{j_3 j'_2 k'} \right) \\
&\stackrel{a}{=} \|P\|_{col}^2 \underbrace{\left( \sum_{j_2, j_3} C_{j_3 j_2 k} C'_{j_3 j_2 k'} \right)}_0 \\
&\stackrel{b}{=} 0.
\end{aligned} \tag{14}$$

Equalities (a) and (b) follow from Lemma 1, the general cases. This general condition works for the columns of the length 3 path matrices as the initial condition. It also holds true for path matrices with arbitrary length (since the CG tensors must have orthogonal columns according to Lemma 1), even without the requirement for the assumption that the previous path matrices hold the orthogonality.  $\blacksquare$

Till now, it is adequate to show our solution to the problem:

**Theorem 10** *Given the set of all the paths of the same length  $n$   $\{p_i\}$  with  $\#\{p_i\} = K$  and path matrices  $\{P^{(p_i)}\}$ , the set of  $\{(\|P\|_{col}^{(p_i)})^{-2} P^{(p_i)} (P^{(p_i)})^T\}$  is a rank  $n - 1$  orthogonal ICT decomposition. The obtainment of this set is of exponential time and space complexity, in terms of rank  $n$ .*

**Proof** From Corollary 8, we know that each  $P^{(p_i)}$  has full rank (but not a square matrix), and has the same  $L_2$  norm of each column,  $\|P^{(p_i)}\|_{col}$ . Subsequently,  $\hat{P}^{(p_i)} = (\|P^{(p_i)}\|_{col})^{-1} P^{(p_i)}$  will make it with an unit  $L_2$  norm, as  $\|(\|P^{(p_i)}\|_{col})^{-1} P^{(p_i)}\|_{col} = 1$ . This does not affect the equivariance of the path matrices, as the  $L_2$ -norm is invariant. If we concatenate all the path matrices  $P$  along the last dimension,

$$D^{(\mathbb{R}^3)^{\otimes n-1} \rightarrow \text{irr}((\mathbb{R}^3)^{\otimes n-1})} = \left[ (\|P^{(p_1)}\|_{col})^{-1} P^{(p_1)} \quad \dots \quad (\|P^{(p_K)}\|_{col})^{-1} P^{(p_K)} \right] \tag{15}$$

where  $\text{irr}$  outputs the all possible irreducible representation spaces. The above equation forms a  $3^{n-1} \times 3^{n-1}$  matrix. In this proof, we will use  $D$  to represent  $D^{(\mathbb{R}^3)^{\otimes n-1} \rightarrow \text{irr}((l=1)^{\otimes n-1})}$  to save place. This is an orthogonal matrix,  $D^T D = D D^T = I$ , as the columns of it are all orthogonal and normalized to unit  $L_2$  norm as we discussed above, and  $K$  is the total multiplicity of the  $n$ th column of the parentage scheme. From Corollary 8, we know that  $D^T$  is  $O(3)$ - $((l=1)^{\otimes n-1}, \oplus_i^K (l^{(i)}))$ -equivariant, where  $l^{(i)} \in \text{irr}((l=1)^{\otimes n-1})$ . If we left multiply it with a selection matrix (a diagonal matrix with ones and zeros on the diagonal)  $I^{(j)}$  to select out the  $j$ th irreducible spherical representation, the resulting  $I^{(j)} P^T$  is  $O(3)$ - $((l=1)^{\otimes n-1}, l^{(j)})$ -equivariant. Similarly, the inverse  $P$  is  $O(3)$ - $(\oplus_i^K (l^{(i)}), (l=1)^{\otimes n-1})$ -equivariant. With the

---

**Algorithm 1** Decomposition Matrices Generation

---

**Input:** Target order  $n\_target$ ;  
 $n\_now, j\_now \leftarrow 0$ ;  
 $decomp\_list, path\_list, this\_path, pathmatrices\_list, ICT\_Decomp\_matrices \leftarrow$   
empty list;  
 $this\_pathmatrix \leftarrow Wigner\_3j(0, 0, 0)$ ;  $\triangleright$  Three arguments are  $j$ s in Wigner-3j symbols  
**function** PATHS\_GENERATOR( $n\_now, j\_now, this\_path, this\_wigner$ )  $\triangleright$  Generate the  
list of CG matrices and the list of corresponding paths  
  **if**  $n\_now \leq n\_target$  **then**  
     $this\_path.append(j\_now)$ ;  
    **for**  $j \in [j\_now + 1, j\_now, j\_now - 1]$  **do**  
      **if** NOT ( $j\_now = 0$  AND  $j \neq 1$ ) AND  $n\_now + 1 \leq n\_target$  **then**  
         $cgmatrix \leftarrow Wigner\_3j(1, j\_now, j)$ ;  
         $this\_pathmatrix_ \leftarrow contract(this\_pathmatrix, cgmatrix)$ ;  
        **call** Paths\_Generator( $n\_now + 1, j, this\_path, this\_pathmatrix$ );  
      **end if**  
    **end for**  
    **if**  $n\_now = n\_target$  **then**  
       $this\_pathmatrix \leftarrow normalize(this\_pathmatrix)$ ;  
       $pathmatrices\_list.append(this\_pathmatrix)$ ;  
       $path\_list.append(this\_path)$ ;  
    **end if**  
  **end if**  
**end function**  
**for**  $path\_matrix \in pathmatrices\_list$  **do**  
   $ICT\_Decomp\_matrx \leftarrow path\_matrix @ path\_matrix.T$ ;  $\triangleright$  @ denotes matrix multipli-  
cation  
   $ICT\_Decomp\_matrices.append(ICT\_Decomp\_matrx)$ ;  
**end for**  
**return**  $path\_list, ICT\_Decomp\_matrices$ ;

---



right multiplied  $I^{(j)}$ , it becomes  $O(3)$ - $(l^{(j)}, (l = 1)^{\otimes n-1})$ -equivariant. Together,  $PI^{(j)}P^T$  is  $O(3)$ - $((l = 1)^{\otimes n-1}, (l = 1)^{\otimes n-1})$ -equivariant. From Lemma 22, this relation is equivalent to  $O(3)$ - $(\mathbb{R}^3)^{\otimes n-1}, (\mathbb{R}^3)^{\otimes n-1}$ -equivariance in the  $n - 1$  rank Cartesian tensor product space. Meanwhile,

$$DI^{(j)}D^T = \hat{P}^{(p_j)}(\hat{P}^{(p_j)})^T, \quad (16)$$

which follows from the fact that  $D$  is an orthogonal matrix, and the concatenation of  $\hat{P}$  forms  $D$ . It therefore meets condition 1) of Definition 5.

Next we prove 2), it holds true because

$$\sum_j \hat{P}^{(p_j)} \left( \hat{P}^{(p_j)} \right)^T = \sum_j DI^{(j)}D^T = DD^T \sum_j I^{(j)} = I. \quad (17)$$

Finally, we prove the orthogonality, given  $\hat{P}^{(p_k)}(\hat{P}^{(p_k)})^T$  and  $\hat{P}^{(p_{k'})}(\hat{P}^{(p_{k'})})^T$  with different  $k \neq k'$ , consider dot product of an arbitrary pair of their columns  $c \neq c'$ , we have

$$\begin{aligned} & \left( \hat{P}^{(p_k)} \left( \hat{P}^{(p_k)} \right)^T \right)_{*,c} \cdot \left( \hat{P}^{(p_{k'})} \left( \hat{P}^{(p_{k'})} \right)^T \right)_{*,c'} \\ &= \sum_{j_1} \left( \sum_{j_2} \hat{P}_{j_1 j_2}^{(p_k)} \hat{P}_{c j_2}^{(p_k)} \right) \left( \sum_{j'_2} \hat{P}_{j_1 j'_2}^{(p_{k'})} \hat{P}_{c' j'_2}^{(p_{k'})} \right) \\ &= \sum_{j_2, j'_2} \hat{P}_{c j_2}^{(p_k)} \hat{P}_{c' j'_2}^{(p_{k'})} \underbrace{\left( \sum_{j_1} \hat{P}_{j_1 j_2}^{(p_k)} \hat{P}_{j_1 j'_2}^{(p_{k'})} \right)}_0 \\ &= 0. \end{aligned} \quad (18)$$

The last equality follows from Lemma 9. From Lemma 6, it satisfies the orthogonality requirement of Definition 5, which is stronger than condition 3), so 3) is implicitly guaranteed.

Time Complexity:

Note that Parentage Scheme is faster than completely traversing a ternary tree, which holds an exponential complexity. For algebraic complexity, there are several studies aiming to discover the complexity of tensor product and contraction (Kisil et al., 2021; Capelli et al., 2016). It can be concluded that our tensor operations will not be worse than exponential complexity. Finally, the dot product of path matrices  $\hat{P}^{(p_j)} \left( \hat{P}^{(p_j)} \right)^T$  are also with exponential complexity.

Space Complexity:

The largest matrices we maintain are the set of  $\hat{P}^{(p_i)}$  and the change-of-basis matrix  $D$ , of which the total size is  $3^{n-1} \times 3^{n-1}$ . ■

It is also noteworthy that the complexity cannot be lower than exponential in terms of rank  $n$ , as the ICT decomposition matrices themselves have an inherently exponential structure. Here we describe the process of constructing decomposition matrices using the parentage scheme. A path in the scheme uniquely corresponds to a path matrix. We start by creating a temporary path matrix, denoted as  $(0 \otimes 0, 0)$ , which will be updated in each round. This matrix is reshaped from a  $(0, 0, 0)$  CG tensor (merging the first two dimensions). The starting point is  $(0, 0)$  in the parentage scheme, where the first coordinate represents the rank, and the second represents the weight.

Next, we move from rank 0 to rank 1, i.e., from the first column to the second column. Observe that the only line connecting  $(0, 0)$  and  $(1, *)$  is the one between  $(0, 0)$  and  $(1, 1)$ . Therefore, the CG tensor to be contracted is  $(1, 0, 1)$ . The third 1 is determined by the second 1 in  $(1, 1)$  in the scheme. The second 0 is determined by the fact that our currently maintained temporary path matrix has 0 in the last position of  $(0 \otimes 0, 0)$ . The first 1 is fixed and remains unchanged regardless of which lines we are moving along (this behavior will change for the general parentage scheme, as introduced in Section 6).

The contraction is performed between the last dimension of the temporary path matrix and the second dimension of the CG tensor, as shown in Equation (10). By merging the first two dimensions of the resulting path matrices, we obtain  $(0 \rightarrow 1)$  path matrices, which can be represented as  $(0 \otimes 0 \otimes 1, 1)$  (we sometimes omit the  $0 \otimes 0$  for simplicity).

At this point, we are positioned at  $(1, 1)$ . The connected points in the next column are  $(2, 0)$ ,  $(2, 1)$ , and  $(2, 2)$ . Consequently, the CG tensors to be contracted are  $(1, 1, 0)$ ,  $(1, 1, 1)$ , and  $(1, 1, 2)$ , respectively. These contractions yield initial path matrices  $(0 \rightarrow 1 \rightarrow 2)$ ,  $(0 \rightarrow 1 \rightarrow 1)$ , and  $(0 \rightarrow 1 \rightarrow 0)$  with shapes  $(1 \otimes 1, 2)$ ,  $(1 \otimes 1, 1)$ , and  $(1 \otimes 1, 0)$ .

This process is repeated iteratively until we reach rank  $n$ . The process and an example are illustrated in Figure 3.

We also present the pseudocode in Algorithm 1. We also benchmark the speed of our algorithm for obtaining the set of all decomposition matrices for different ranks, as shown in Table 3. We run the experiments on 28-cores Intel<sup>®</sup> Xeon<sup>®</sup> Gold 6330 CPU @ 2.00GHz.

## 5 Basis of the equivariant design space

Recall Section 3.2 that one of our goal is to find the basis of the equivariant design space  $\{M \in \mathbb{R}^{3^n} \times \mathbb{R}^{3^n} \mid \rho(g)M = M\rho(g)\}$ . Here, we leverage Schur's lemma:

**Lemma 11 (Schur's lemma, special case)** *Given  $\rho$  a finite-dimensional irreducible representation of group  $G$ , and a matrix  $M$ . Then*

$$\rho(g)M = M\rho(g) \tag{19}$$

*holds for any  $g \in G$ , if and only if  $M = cI$ , where  $c$  is a scalar.*

Thus, for an endomorphism space from a single irreducible representation space to itself, the basis is simply an identity matrix  $I$  and it is a single dimension space. On the contrary, a rank- $n$  Cartesian tensor is a direct sum of several such irreducible representation spaces. In such situation, we obtain the following lemma:

**Lemma 12** *Given a representation  $\mathcal{S} = \oplus_i^n \mathcal{S}^{(i)}$  that can be written as the direct sum of several finite-dimensional irreducible representations  $\mathcal{S}^{(i)}$ . If they are isomorphic to each other  $\mathcal{S}^{(0)} \cong \dots \cong \mathcal{S}^{(n)}$ , then the number of dimensions of the equivariant design space is  $n^2$ , and the basis consists of one-dimensional linear projections with a scaling factor from  $\mathcal{S}^{(i)}$  to  $\mathcal{S}^{(j)}$  for any  $i$  and  $j$ .*

**Proof** First note that

$$(\mathcal{S}^{(i)} \oplus \mathcal{S}^{(j)}) \rightarrow \mathcal{S}' \cong (\mathcal{S}^{(i)} \rightarrow \mathcal{S}') + (\mathcal{S}^{(j)} \rightarrow \mathcal{S}'), \quad (20)$$

where  $+$  represents linear combinations, and conversely

$$\mathcal{S}' \rightarrow (\mathcal{S}^{(i)} \oplus \mathcal{S}^{(j)}) \cong (K' \rightarrow \mathcal{S}^{(i)}) \oplus (\mathcal{S}' \rightarrow \mathcal{S}^{(j)}). \quad (21)$$

Given a linear projection  $f : \oplus_i^n \mathcal{S}^{(i)} \rightarrow \oplus_i^n \mathcal{S}^{(i)}$ , we have

$$\begin{aligned} & \oplus_i^n \mathcal{S}^{(i)} \rightarrow \oplus_i^n \mathcal{S}^{(i)} \\ & \cong \oplus_j^n \left( \oplus_i^n \mathcal{S}^{(i)} \rightarrow \mathcal{S}^{(j)} \right) \\ & \cong \oplus_j^n \left( +_i^n \left( \mathcal{S}^{(i)} \rightarrow \mathcal{S}^{(j)} \right) \right). \end{aligned} \quad (22)$$

Note that  $(\mathcal{S}^{(i_1)} \rightarrow \mathcal{S}^{(j)})$  and  $(\mathcal{S}^{(i_2)} \rightarrow \mathcal{S}^{(j)})$  form a basis of  $(\mathcal{S}^{(i_1)} \rightarrow \mathcal{S}^{(j)}) + (\mathcal{S}^{(i_2)} \rightarrow \mathcal{S}^{(j)})$  if  $i_1 = i_2$ . Meanwhile,  $(\mathcal{S}^{(i_1)} \rightarrow \mathcal{S}^{(j_1)})$  and  $(\mathcal{S}^{(i_2)} \rightarrow \mathcal{S}^{(j_2)})$  must form a basis of  $(\mathcal{S}^{(i_1)} \rightarrow \mathcal{S}^{(j_1)}) \oplus (\mathcal{S}^{(i_2)} \rightarrow \mathcal{S}^{(j_2)})$  according to the definition of direct sum. Each mapping is of dimension one by Lemma 11. Thus,  $\oplus_i^n \mathcal{S}^{(i)} \rightarrow \oplus_i^n \mathcal{S}^{(i)}$  has totally  $n^2$  basis  $\mathcal{S}^{(i)} \rightarrow \mathcal{S}^{(j)}$ . ■

From the perspective of matrices, it is equivalent to say that the matrices which can *scale an irreducible representation and add it to another* (including itself) form a basis for the equivariant design space. Here, we provide an example,

$$\begin{aligned} & \begin{bmatrix} \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \\ \cdots & 0 & 0 & 0 & k & 0 & 0 & \cdots \\ \cdots & 0 & 0 & 0 & 0 & k & 0 & \cdots \\ \cdots & 0 & 0 & 0 & 0 & 0 & k & \cdots \\ \cdots & 0 & 0 & 0 & 0 & 0 & 0 & \cdots \\ \cdots & 0 & 0 & 0 & 0 & 0 & 0 & \cdots \\ \cdots & 0 & 0 & 0 & 0 & 0 & 0 & \cdots \\ \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} \vdots \\ v_{-1}^{(1;q)} \\ v_0^{(1;q)} \\ v_1^{(1;q)} \\ v_{-1}^{(1;q')} \\ v_0^{(1;q')} \\ v_1^{(1;q')} \\ \vdots \end{bmatrix} = \begin{bmatrix} \vdots \\ kv_{-1}^{(1;q')} \\ kv_0^{(1;q')} \\ kv_1^{(1;q')} \\ 0 \\ 0 \\ 0 \\ \vdots \end{bmatrix}, \quad (23) \\ & \quad \quad \quad A^{(l=1;q;q')} \end{aligned}$$

where  $A^{(1;q;q')}$  adds  $kv^{(1;q')}$  to the position of  $v^{(1;q)}$ , where  $v^{(1;q)}$  belongs to the  $(l = 1)$  spherical space, with index  $q$  corresponding to the multiplicity of the same weight, and  $k$  is a scalar. Note that  $A^{(1;q;q')}$  resides in the spherical space and is a  $(r = 1, s = 1)$ -tensor,

so we need to convert it back to Cartesian space via the change-of-basis matrices  $D$  and  $D^{-1} = D^T$ . We have that

$$\begin{aligned}
 & \begin{bmatrix} \vdots \\ \hat{P}_{*,0}^{(\cdots \rightarrow 2 \rightarrow 1)} \\ \hat{P}_{*,1}^{(\cdots \rightarrow 2 \rightarrow 1)} \\ \hat{P}_{*,2}^{(\cdots \rightarrow 2 \rightarrow 1)} \\ \hat{P}_{*,0}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ \hat{P}_{*,1}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ \hat{P}_{*,2}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ \vdots \end{bmatrix}^T \begin{bmatrix} \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \\ \cdots & 0 & 0 & 0 & k & 0 & 0 & \cdots \\ \cdots & 0 & 0 & 0 & 0 & k & 0 & \cdots \\ \cdots & 0 & 0 & 0 & 0 & 0 & k & \cdots \\ \cdots & 0 & 0 & 0 & 0 & 0 & 0 & \cdots \\ \cdots & 0 & 0 & 0 & 0 & 0 & 0 & \cdots \\ \cdots & 0 & 0 & 0 & 0 & 0 & 0 & \cdots \\ \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} \vdots \\ \hat{P}_{*,0}^{(\cdots \rightarrow 2 \rightarrow 1)} \\ \hat{P}_{*,1}^{(\cdots \rightarrow 2 \rightarrow 1)} \\ \hat{P}_{*,2}^{(\cdots \rightarrow 2 \rightarrow 1)} \\ \hat{P}_{*,0}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ \hat{P}_{*,1}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ \hat{P}_{*,2}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ \vdots \end{bmatrix} \\
 & \begin{matrix} P & P^T \end{matrix} \\
 & \begin{matrix} A^{(l;q;q')} \end{matrix} \\
 & = \begin{bmatrix} \vdots \\ \hat{P}_{*,0}^{(\cdots \rightarrow 2 \rightarrow 1)} \\ \hat{P}_{*,1}^{(\cdots \rightarrow 2 \rightarrow 1)} \\ \hat{P}_{*,2}^{(\cdots \rightarrow 2 \rightarrow 1)} \\ \hat{P}_{*,0}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ \hat{P}_{*,1}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ \hat{P}_{*,2}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ \vdots \end{bmatrix}^T \begin{bmatrix} \vdots \\ k\hat{P}_{*,0}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ k\hat{P}_{*,1}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ k\hat{P}_{*,2}^{(\cdots \rightarrow 1 \rightarrow 1)} \\ 0 \\ 0 \\ 0 \\ \vdots \end{bmatrix} \\
 & = k \sum_j \hat{P}_{i_1 j}^{(\cdots \rightarrow 2 \rightarrow 1)} \hat{P}_{i_2 j}^{(\cdots \rightarrow 1 \rightarrow 1)}.
 \end{aligned} \tag{24}$$

Thus, we can reduce the three square matrices multiplications to a series of products of sub-matrices. To conclude, we have

**Proposition 13** *Given a rank  $n$  Cartesian tensor product space.  $\hat{P}^{(p_i)} \left( \hat{P}^{(p_j)} \right)^T$  for all pair of paths  $p_i$  and  $p_j$  of which the last weights are the same, form an orthogonal basis of the equivariant design space with respect to the Cartesian space. The dimension of the equivariant space equals*

$$\sum_{n,0 \leq j \leq n} \left( \sum_i^{\lfloor (n-j)/3 \rfloor} (-1)^i \frac{n!(2n-3i-j-2)!}{i!(n-i)!(n-2)!(n-3i-j)!} \right)^2. \tag{25}$$

**Proof** We first prove the equivariance. For arbitrary  $\hat{P}^{(p_i)} \left( \hat{P}^{(p_j)} \right)^T$ , and group element  $g$ , we have

$$\begin{aligned}
 & \hat{P}^{(p_i)} \left( \hat{P}^{(p_j)} \right)^T \rho_{Cart}(g) = \hat{P}^{(p_i)} \left( \hat{P}^{(p_j)} \right)^{-1} \rho_{Cart}(g) \\
 & = \hat{P}^{(p_i)} \rho_{(l)}(g) \left( \hat{P}^{(p_j)} \right)^{-1} = \rho_{Cart}(g) \hat{P}^{(p_i)} \left( \hat{P}^{(p_j)} \right)^T.
 \end{aligned} \tag{26}$$

For the orthogonality, we have

$$\begin{aligned}
& \left( \hat{P}^{(p_i)} \left( \hat{P}^{(p_j)} \right)^T \right)^T \left( \hat{P}^{(p_{i'})} \left( \hat{P}^{(p_{j'})} \right)^T \right) \\
&= D \left( A^{(l;q;q')} \right)^T D^T D A^{(l;q'';q''')} D^T \\
&= D \underbrace{\left( \left( A^{(l;q;q')} \right)^T A^{(l;q'';q''')} \right)}_0 D^T \\
&= 0,
\end{aligned} \tag{27}$$

for arbitrary  $p_i \neq p_{i'}$  or  $p_j \neq p_{j'}$ , i.e.,  $q \neq q''$  or  $q' \neq q'''$ . The last equality follows from the fact that either  $A_{ij}^{(l;q;q')} = 0$  or  $A_{ji}^{(l;q'';q''')} = 0$  for  $q \neq q''$  or  $q' \neq q'''$ . This can form  $k^2$  orthogonal matrices for the irreducible spherical spaces with multiplicity  $k$ , so Lemma 12 guarantees they form a basis. Note that there is no linear mapping between the spaces that are not isomorphic. The dimension calculation is based on Proposition 4.  $\blacksquare$

Notice that the ICT decomposition matrices are also part of the basis of the equivariant space.

## 6 Extension to general spaces

### 6.1 General ICT decomposition

It is appealing to build equivariant layers between different desired spaces. The spherical tensor product spaces enable us to utilize high-order many-body information while remaining flexible in terms of memory cost and inference speed. However, it is difficult to construct such a layer. For example, if the input space is  $(l = 4 \otimes l = 5)$  and the output space is  $(l = 2 \otimes l = 6)$ , how can we parameterize the equivariant layer? More specifically, what is the basis of this layer? It is even not possible to find a spanning set by Pearce-Crump (2023b). Before answering this question, we first introduce the general problem of ICT decomposition and how to efficiently perform a general ICT decomposition on arbitrary input spaces.

**Definition 14** *Given a space  $\mathcal{S} = (l_{11} \otimes \dots \otimes l_{1i_1}) \oplus \dots \oplus (l_{k1} \otimes \dots \otimes l_{ki_k})$  with  $d$  dimension, a general irreducible decomposition is a series of matrices  $\hat{H}^{(\mathcal{S};l;q)} \in \mathbb{R}^{d \times d}$ , such that:*

- 1)  $\hat{H}^{(\mathcal{S};l;q)}$  is  $O(3)$ - $(\mathcal{S}, \mathcal{S})$ -equivariant.
- 2) The summation of the decomposition equals to identity matrix,  $\sum_i \hat{H}^{(\mathcal{S};l;q)} = I$ .
- 3)  $\{\hat{H}^{(\mathcal{S};l;q)}\}$  are linearly independent.

If  $H^{(i)}$  are orthogonal to each other, as defined in Definition 5, we further call it an orthogonal general irreducible decomposition.

Intuitively, an orthogonal general ICT decomposition can be obtained using a strategy similar to that of the classic ICT decomposition. However, we cannot use the parentage




Figure 4: Two methods for performing a general ICT decomposition.

scheme directly, as it only accepts a tensor product with an ( $l = 1$ ) one step at a time, proceeding from left to right. Here, we introduce a general parentage scheme. Unlike the classic scheme, this is not a fixed procedure.

For example, given a space  $(3 \otimes 4 \otimes 2) \oplus (2 \otimes 3)$ , we first separate the terms by the direct sum  $\oplus$ , resulting in  $(3 \otimes 4 \otimes 2)$  and  $(2 \otimes 3)$ . First, we use the general parentage scheme to decompose  $(3 \otimes 4 \otimes 2)$  into irreducible spherical spaces and construct the path matrices. We begin by writing a multiplicity of one in the weight-3 row, in the leftmost column. Then, we take the tensor product with ( $l = 4$ ), which results in multiplicities of one in  $(l = 7), \dots, (l = 1)$ . We also construct the path matrices  $(3 \xrightarrow{4} 7), \dots, (3 \xrightarrow{4} 1)$ . Note that we mark 4 on top of the arrow, which we refer to as the bridge number. Then, 2 is taken as the bridge number, and we accordingly write the multiplicities in the next column. Consequently, we apply the same procedure for  $(2 \otimes 3)$ . An important property is

**Corollary 15 (from Lemma 7)** *Given a tensor product space, the path matrices for **the same** general parentage scheme have orthogonal columns, and the  $L_2$  norms of columns of each path matrix are the same.*

**Proof** The proof for the is mostly identical to that of Lemma 7. ■

Here the words **the same** suggest us we need to fix our bridge number to obtain orthogonal path matrices. As our previous example,  $(3 \xrightarrow{4} 7 \xrightarrow{2} 6)$  and  $(4 \xrightarrow{3} 7 \xrightarrow{2} 6)$  will generate two different path matrices. But they are actually identical up to an index permutation. The result will be the same if we accordingly permute the index of our input vector. We can multiply the path matrices with their inverse to obtain decomposition matrices, as the classic ICT decomposition. Here we should note that the decomposition matrices of different sub schemes separated by the direct sum can not generally concatenated. We can, in general, concatenate some zeros to the path matrices and their transpose with short row and column, respectively. Or we can just leave the decomposition matrices of different shapes as they are, and multiply the matrices with different segments of the input vector. The two methods are shown in Figure 4.

## 6.2 Equivariant bases of arbitrary spaces

The general ICT decomposition gives us some hints. The constructed change-of-basis matrix  $D^T$  changes the basis from  $(3 \otimes 4 \otimes 2) \oplus (2 \otimes 3)$  to its decomposed irreducible spherical spaces, denoted as  $\text{irr}((3 \otimes 4 \otimes 2) \oplus (2 \otimes 3))$ , and its inverse  $D$  changes back the basis. An orthogonal basis from spaces  $\mathcal{S}_1$  to  $\mathcal{S}_2$  can be obtained via the change-of-basis matrices constructed based on the general ICT decompositions of these two spaces. But we need to take care of more when considering  $O(3)$ . The above discussion is identical for  $SO(3)$  and  $O(3)$ . Instead, here, the parity must get involved when we consider arbitrary spaces. First, one should note that a tensor product space, as irreducible spherical spaces, have parity 1 or  $-1$ . We have kept in mind that Cartesian space is of parity  $-1$ , which means that if we do the reflection, then the space will be reflected accordingly. It is equivalent to an  $(l = 1, p = -1)$  spherical space, from Lemma 22. Meanwhile, we should remember that there is also  $(l = 1, p = 1)$  spherical space, as a *cannot-be-reflected* version of the Cartesian space. The parity of the tensor product space is only decided by the number of the componential spaces with odd parities. If the number is odd, then the tensor product space has odd parity, and *vice versa*. Meanwhile,

**Proposition 16** *Given a tensor product space of rank  $n$ , the resulting spaces of the ICT decomposition are with the same parity as the tensor product space.*

**Proof** Suppose we have a rank  $n$  tensor  $T$  in the tensor product space, and assume that the parities of the ICTs and the original rank  $n$  tensor  $T$  are not the same. If a tensor  $T$  has an even parity, then reflection will not change  $T$ . Let  $\hat{T}_1^{(p=-1)}, \dots, \hat{T}_k^{(p=-1)}$  be the ICTs with odd parity. According to Definition 14, we should have

$$\begin{aligned} T &= \sum_i \hat{T}_i^{(p=-1)} + \sum_{i'} \hat{T}_{i'}^{(p=1)} \\ &= - \sum_i \hat{T}_i^{(p=-1)} + \sum_{i'} \hat{T}_{i'}^{(p=1)}, \end{aligned} \tag{28}$$

which gives us

$$\sum_i \hat{T}_i^{(p=-1)} = 0. \tag{29}$$

This contradicts with the definition that  $\hat{T}_i^{(p=-1)}$  are linearly independent. The same arguments can be given for the case that  $T$  has odd parity. ■

Now we are ready to introduce the procedure for a obtaining an orthogonal basis for different spaces. Given spaces  $\mathcal{S}^{in}$  and  $\mathcal{S}^{out}$ , we want to find a basis for the equivariant design space from  $\mathcal{S}^{in}$  to  $\mathcal{S}^{out}$ . We first find the irreducible spherical spaces of  $\mathcal{S}^{in}$  and  $\mathcal{S}^{out}$  via the general parentage scheme, denoted as  $\text{irr}(\mathcal{S}^{in})$  and  $\text{irr}(\mathcal{S}^{out})$ , and write down the paths. Then, the paths are grouped according to the last weight in them, as well as the parities of the original tensor product spaces, denoted as  $Q_{\mathcal{S}^{in}}^{(l;p)}$  and  $Q_{\mathcal{S}^{out}}^{(l;p)}$ . Let our equivariant space be  $\text{Hom}_{O(3)}(\mathcal{S}^{in}, \mathcal{S}^{out})$ , we have

---

**Algorithm 2** Equivariant basis generation

---

**Input:** Input and output spaces  $space\_in$ ,  $space\_out$ , CG possible decoupling weights generator  $CG\_decoup\_generator$ , group type  $group$ , path matrices generator  $path\_matrices\_generator$ ;  
 $paths\_comp$ ,  $paths\_decomp$   $\leftarrow$  empty list;  
**function** GENERAL\_SCHEME( $rank\_list$ ,  $this\_path$ ,  $rank\_now$ ,  $this\_wigner$ ,  $ind$ ,  $paths$ ,  $parity$ ,  $index$ )  
     $rank \leftarrow paths[ind]$   
    **for**  $i \in CG\_decoup\_generator(group, rank\_now, rank)$  **do**  
         $this\_path\_ \leftarrow this\_path.copy()$ ;  
         $this\_path\_ .append(rank, i)$ ;  
        **if**  $ind\_ \geq rank\_list.length()$  **then**  
             $path\_list.append(\{ 'path' : this\_path\_ , 'parity' : parity , 'index' : index \})$ ;  
        **else**  
            **call** General\_Scheme( $rank\_list, this\_path\_ , i, ind + 1, paths, parity, index$ );  
        **end if**  
    **end for**  
**end function**  
 $index \leftarrow 0$ ;  
**for**  $i \in space\_in[ 'space' ]$  **do**  
    General\_Scheme( $i, [[0, 0]], 0, 0, paths\_decomp, space\_in[ 'parity' ], index$ );  
     $index \leftarrow index + 1$ ;  
**end for**  
 $index \leftarrow 0$ ;  
**for**  $i \in space\_out[ 'space' ]$  **do**  
    General\_Scheme( $i, [[0, 0]], 0, 0, paths\_comp, space\_out[ 'parity' ], index$ );  
     $index \leftarrow index + 1$ ;  
**end for**  
 $equivariant\_basis, paths\_and\_spaces \leftarrow$  empty list;  
**for**  $path\_c \in paths\_comp$  **do**  
    **for**  $path\_d \in paths\_decomp$  **do**  
        **if**  $path\_d[ 'path' ][-1][-1] = path\_c[ 'path' ][-1][-1]$  and  $path\_d[ 'parity' ] = path\_c[ 'parity' ]$  **then**  
             $path\_list.append(\{ 'path' : this\_path\_ , 'parity' : parity , 'index' : index \})$ ;  
             $path\_matrix\_c \leftarrow path\_matrices\_generator(path\_c[ 'path' ], group)$ ;  
             $path\_matrix\_d \leftarrow path\_matrices\_generator(path\_d[ 'path' ], group)$ ;  
             $equivariant\_basis.append(path\_matrix\_c @ path\_matrix\_d.T)$ ;  
        **end if**  
    **end for**  
**end for**  
**return**  $paths\_and\_spaces, equivariant\_basis$ ;

---



---

**Algorithm 3** path matrices generator
 

---

**Input:** path  $path$ , group type  $group$ ;  
 $path\_matrix \leftarrow CG\_matrix(0, 0, 0, group)$ ;  $\triangleright$  generate group-specific CG matrix  
 $j\_now \leftarrow 0$ ;  
**for**  $bridge\_j, next\_j \in path$  **do**  
    $cg\_matrix \leftarrow CG\_matrix(bridge\_j, j\_now, next\_j, group)$ ;  
    $path\_matrix \leftarrow contract(path\_matrix, cg\_matrix)$ ;  
    $j\_now \leftarrow next\_j$   
**end for**  
 $path\_matrix \leftarrow \frac{path\_matrix}{\|path\_matrix\|_{col}}$ ;  
**return**  $path\_matrix$ ;  


---

$$\dim \text{Hom}_{O(3)}(\mathcal{S}^{in}, \mathcal{S}^{out}) = \sum_{l;p} \#Q_{\mathcal{S}^{in}}^{(l;p)} \#Q_{\mathcal{S}^{out}}^{(l;p)}, \quad (30)$$

where  $\#\cdot$  represents the number of elements in the set, which follows from Lemma 11 (Schur's lemma). We drop those paths sets that the last weight and the same parities do not show in both spaces. Then, we can generate path matrices based on those remained sets. Finally, we can conclude that

**Proposition 17** Given spaces  $\mathcal{S}^{in} = ((l_1^1, p_1^1) \otimes \cdots \otimes (l_{k_1}^1, p_{k_1}^1)) \oplus \cdots \oplus ((l_1^t, p_1^t) \otimes \cdots \otimes (l_{k_t}^t, p_{k_t}^t))$ , and  $\mathcal{S}^{out} = ((l_1^{l'}, p_1^{l'}) \otimes \cdots \otimes (l_{k_1}^{l'}, p_{k_1}^{l'})) \oplus \cdots \oplus ((l_1^{t'}, p_1^{t'}) \otimes \cdots \otimes (l_{k_{t'}}^{t'}, p_{k_{t'}}^{t'}))$ ,

$$\left\{ \hat{P}^{(out;l;p;q')} \left( \hat{P}^{(in;l;p;q)} \right)^T \mid p^{(in;l;p;q)} \in Q_{\mathcal{S}^{in}}^{(l;p)}, p^{(out;l;p;q')} \in Q_{\mathcal{S}^{out}}^{(l;p)}, \hat{p} \in \{-1, 1\} \right\}$$

forms an orthogonal basis of  $\text{Hom}_{O(3)}(\mathcal{S}^{in}, \mathcal{S}^{out})$ .

**Proof** Since  $\hat{P}^{(out;l;p;q')} \left( \hat{P}^{(in;l;p;q)} \right)^T$  are mappings between  $\mathcal{S}^{in}$  and  $\mathcal{S}^{out}$ , They can be represented as a  $\sum_{j=1}^t \left( \Pi_i^{k_j} 2l_i^j + 1 \right) \times \sum_{j=1}^{t'} \left( \Pi_i^{k'_j} 2l_i^{j'} + 1 \right)$  matrix. From the mapping relations, we know that it is a block matrix as

$$\begin{bmatrix} M_{11} & \cdots & \cdots & M_{1t} \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ M_{t'1} & \cdots & \cdots & M_{t't} \end{bmatrix}. \quad (31)$$

The sub-matrices are of shape  $\left( \Pi_i^{k'_{j'}} 2l_i^{j'} + 1 \right) \times \left( \Pi_i^{k_j} 2l_i^j + 1 \right)$  for some  $j$  and  $j'$ .  $\hat{P}^{(out;l;p;q')}$   $\left( \hat{P}^{(in;l;p;q)} \right)^T$  will be in the same column for those  $(l; p; q')$  belonging to the same direct




Figure 5: The general parentage scheme with the example of finding basis for  $((2 \otimes 2 \otimes 2), p = -1) \oplus ((1 \otimes 3), p = -1) \rightarrow ((3 \otimes 4), p = -1)$ . Compared to the classic parentage scheme, we need to additionally mark the bridge number in the scheme.

sum component of  $\mathcal{S}^{out}$ , and the same row for those  $(l; p; q)$  belonging to the same direct sum component of  $\mathcal{S}^{in}$ . When the non-zero values of  $\hat{P}^{(out; l; p; q')} \left( \hat{P}^{(in; l; p; q)} \right)^T$  appears in a block  $M$ , the rest of the sub-matrices will be zeros. Thus, given two elements in the basis, their corresponding block matrices must be orthogonal if the non-zero values do not

Rank	5	6	7	8	9
Time	<0.1s	1s	3s	11s	4m32s

Table 3: Benchmarking decomposition matrices generation speed.

show in the same block  $M$ . If they appear in the same  $M$ , then the question is equivalent to Proposition 13.  $\blacksquare$

The algorithm for generating equivariant bases is presented in Algorithm 2. Here, *CG\_decomp\_generator* receives the current weight and the bridge number as input and outputs the possible weights according to the selection rules of CG coefficients. These rules depend on the group type. For instance, in the case of  $O(3)$ ,  $l_1$  and  $l_2$  generate possible weights ranging from  $|l_1 - l_2|$  to  $l_1 + l_2$ . For  $SU(2)$ , an additional condition is imposed: the output weight must have the same parity (odd or even, though not the parity in  $O(3)$  used to determine whether it can be reflected) as the sum of the two input weights. The function *path\_matrices\_generator* in this algorithm is described in Algorithm 3, which also depends on the group type. It is elegant to generate equivariant bases as matrices, as they also enable efficient computations. The bases can also be efficiently obtained. However, as we test the spanning set method proposed by Pearce-Crump (2023b), when we assign a scalar parameter to each basis and combine these bases to form a single matrix, real-world neural network frameworks (e.g., PyTorch) may encounter memory issues. This is because a large computation graph needs to be maintained when the rank is very high. To address this issue in high-rank cases, it is preferable to first convert the tensor into the direct sum of different irreducible spherical tensors. Then, linear combinations of tensors with the same weight can be applied. Finally, the direct sum space is converted back into the original space. This approach also follows from the previous proofs and aligns with the core idea of this paper: performing operations in spherical spaces is simpler and more efficient. This process is detailed in Algorithm 4.

### 6.3 Multi-channel features, activation functions, normalizations, and biases

We have already introduced how to construct equivariant linear operations between some  $\mathcal{S}^{in}$  and  $\mathcal{S}^{out}$ . In a real implementation, additional considerations are required to make it an equivariant layer in a modern neural network. First, if we take vectors in  $\mathcal{S}^{in}$  and  $\mathcal{S}^{out}$  as features, it is both standard and beneficial for a neural network to have multi-channel features. Fortunately, this is straightforward in our framework. We can simply let  $(\mathcal{S}^{in})^{\oplus K^{in}}$  and  $(\mathcal{S}^{out})^{\oplus K^{out}}$  serve as our input and output spaces. In this case, the conclusion in Section 6.2 can seamlessly be applied. Notably, Proposition 17 already addresses the scenarios described here. Thus, for a space

$$\underbrace{((l_1^1, p_1^1) \otimes \cdots \otimes (l_{k_1}^1, p_{k_1}^1))}_{\mathcal{S}_{\oplus}^1} \oplus \cdots \oplus \underbrace{((l_1^t, p_1^t) \otimes \cdots \otimes (l_{k_t}^t, p_{k_t}^t))}_{\mathcal{S}_{\oplus}^t}, \quad (32)$$

---

**Algorithm 4** Equivariant basis generation – linear combinations in spherical spaces

---

**Input:** Input and output spaces  $space\_in$ ,  $space\_out$ , CG possible decoupling weights generator  $CG\_decoup\_generator$ , group type  $group$ , path matrices generator  $path\_matrices\_generator$ ;  
 $paths\_comp$ ,  $paths\_decomp$   $\leftarrow$  empty list;  
**define function** General\_Scheme( $rank\_list$ ,  $this\_path$ ,  $rank\_now$ ,  $this\_wigner$ ,  $ind$ ,  $paths$ ,  $parity$ ,  $index$ );  $\triangleright$  define function General\_Scheme as we did in Algorithm 2  
**define function** Spherical\_Multiplicities( $path$ );  $\triangleright$  return the number of dimensions, parity, and the multiplicity of the last point of each path  
 $index \leftarrow 0$ ;  
**for**  $i \in space\_in['space']$  **do**  
    General\_Scheme( $i$ ,  $[[0, 0]]$ ,  $0, 0, paths\_decomp, space\_in['parity'], index$ );  
     $index \leftarrow index + 1$ ;  
**end for**  
 $index \leftarrow 0$ ;  
**for**  $i \in space\_out['space']$  **do**  
    General\_Scheme( $i$ ,  $[[0, 0]]$ ,  $0, 0, paths\_comp, space\_out['parity'], index$ );  
     $index \leftarrow index + 1$ ;  
**end for**  
 $comp\_matrix \leftarrow concat(paths\_comp)$ ;  
 $decomp\_matrix \leftarrow concat(paths\_decomp)$ ;  
 $fc\_list \leftarrow$  empty list;  
**for**  $dim\_comp, p\_comp, multip\_comp \in Spherical\_Multiplicities(paths\_comp)$  **do**  
    **for**  $dim\_decomp, p\_decomp, multip\_decomp \in Spherical\_Multiplicities(paths\_decomp)$   
    **do**  
        **if**  $dim\_comp = dim\_decomp$  and  $p\_comp = p\_decomp$  **then**  
             $fc\_list.append(Learnable\_Matrix(multip\_comp, multip\_decomp))$ ;  
        **end if**  
    **end for**  
**end for**  
**return**  $fc\_list, decomp\_matrix.T, comp\_matrix$ ;

---

Rank	4	5	6	7	8
EMLP (Finzi et al., 2021)	2m52s	26m	OOM	OOM	OOM
Algorithm 2	<0.1s	1s	3s	10s	OOM
Algorithm 4	<0.1s	<0.1s	<0.1s	<0.1s	0.8s

Table 4: Comparison on the basis generation speed of  $\text{End}_{O(3)}\left(\left(\mathbb{R}^3\right)^{\otimes n}\right)$ . The out-of-memory error occurs when generating the bases using EMLP. In contrast, for Algorithm 2, the error occurs when multiplying the learnable parameters.

each direct sum component  $\mathcal{S}_{\oplus}^i$  is a single-channel feature in a multi-channel view, which can be different depending on the positions. This is more general than the regular multi-channel features.

On the other hand, activation functions introduce non-linearity to neural networks. As non-linear functions, their design space is very limited in order to preserve symmetry. However, it is completely safe if we can construct invariant linear mappings of the form  $f^{inv} : v^{out} \in \mathcal{S}^{out} \mapsto c \in \mathbb{R}$ . Then, we can apply an activation function to  $c$ , obtaining  $\phi(c)$ , and multiply  $\phi(c)$  with  $v^{out}$  to produce the activated result,  $\phi(f^{inv}(v^{out}))v^{out}$ . This process is equivariant, as

$$\begin{aligned}
& \phi\left(f^{inv}\left(\rho_{\mathcal{S}^{out}}(g) \odot v^{out}\right)\right) \rho_{\mathcal{S}^{out}}(g) \odot v^{out} \\
&= \phi\left(\underbrace{\rho_{\mathbb{R}}(g)}_I \odot f^{inv}\left(v^{out}\right)\right) \rho_{\mathcal{S}^{out}}(g) \odot v^{out} \\
&= \rho_{\mathcal{S}^{out}}(g) \odot \left(\phi\left(f^{inv}\left(v^{out}\right)\right) \odot v^{out}\right).
\end{aligned} \tag{33}$$

This process is similar to that of the spherical space-based methods (Batzner et al., 2022; Musaelian et al., 2023), but here the strategy is available for more general spaces.

For the normalizations, one can leverage the fact that the  $L_2$ -norm of the vectors are invariant to the rotation groups, in general spaces. This is because the matrix representations of  $O(n)$  always have an absolute determinant equal to one. Thus, for a vector  $v \in \mathcal{S}$ , we can simply calculate  $\|v\|_2$  to construct an equivariant normalization layer. The discussion on biases is mainly identical to Pearce-Crump (2023b). We can construct equivariant basis from  $\mathbb{R}$  to arbitrary  $\mathcal{S}^{out}$ . In our work, we can find an orthogonal basis through Proposition 17 instead of the spanning set. Here we should note that there are some situations that we cannot find such a mapping to  $\mathcal{S}^{out}$ . For example,  $\mathbb{R}$  with odd parity cannot map to  $\mathcal{S}^{out}$  with even parity, and *vice versa*. Also,  $(l=0, p=1)$  cannot map to  $(l=3, p=1) \otimes (l=2, p=1)$  either, as  $(l=0, p=1)$  is not in their irreducible decomposition set  $\text{irr}\left((l=3, p=1) \otimes (l=2, p=1)\right) = \{(l=5, p=1), (l=4, p=1), (l=3, p=1), (l=2, p=1), (l=1, p=1)\}$ . One can manually make the biases work by imaging the output space is  $((l=3, p=1) \otimes (l=2, p=1)) \oplus (l=0, p=1)$ , but with  $(l=0, p=1)$  being zero. The features can be well mixed in the next layer.

## 7 Related works

Research on equivariant neural networks is rooted in an abstract concept raised by Cohen and Welling (2016), where they extend the translation invariance of CNNs (LeCun et al., 1989) to a wide range of groups. Meanwhile, graph neural networks (Gori et al., 2005) are proven to be suitable tools for handling tasks involving point clouds. Their coupling, Equivariant Graph Neural Networks (EGNNs) has initiated an emerging area with very important applications to physics-informed tasks. Among the implementation, Message Passing Neural Networks (MPNNs) (Gilmer et al., 2017) are the most popular, where the constructed message is propagated to the neighboring points. Before that, Schütt et al. (2017) observed that group CNN frameworks require the points to lie on some grids and propose SchNet to flexibly handle atoms that randomly appear in space. (Gasteiger et al., 2020) further leveraged directional information to enhance the model expressivity. At this point, the features utilized were just scalars and vectors, and subsequent works began to use high-order irreducible representations to effectively encode many-body interactions. Thomas et al. (2018) proposed an abstract framework, the Tensor Field Network, to incorporate spherical irreducible representations into MPNNs. This abstract framework has been implemented with different designs to achieve SOTA performances on multiple popular benchmarks (Chmiela et al., 2017; Ruddigkeit et al., 2012; Fu et al., 2022). Gasteiger et al. (2021) demonstrated the universality of using such spherical irreducible representations. Batzner et al. (2022) implemented the Tensor Field Network with high order irreducible representations, showing strong performance. On the other hand, Drautz (2019) proposed a local descriptor-based method, Atomic Cluster Expansion (ACE), to encode the atomic information via polynomial basis functions. Batatia et al. (2022) further used high-order message-passing techniques to reduce the required number of layers utilizing spherical irreducible representations. Musaelian et al. (2023) proposes a strictly local EGNN, constructing high order spherical irreducible representations among edges. Recently, several studies have leveraged the fact that Cartesian tensor is more computationally efficient. Simeon and De Fabritiis (2024) first propose to use Cartesian tensor but only with rank-2 ICT decomposition, and they further apply it to charged molecules and spin states (Simeon et al., 2024). Cheng (2024) used high-rank Cartesian tensors but did not manipulate the irreducible components. Zaverkin et al. (2024) successfully incorporated high-weight ICT to MPNNs but only addressed the highest weight of a given Cartesian tensor, lacking a full treatment of all ICTs for a given Cartesian tensor.

Research on ICT decomposition has a long history and remains a highly active area in theoretical chemistry and chemistry physics communities, in addition to the recent strong interest from the machine learning community. The second-rank ICT decomposition is well known. Since mid-20th century, Coope et al. (1965) provided the explicit ICT decomposition for rank-3 Cartesian tensors. Seventeen years later, Andrews and Ghoul (1982) derived the rank-4 formula for this problem. Dinçkal (2013) further refined Coope et al. (1965)’s results to make them orthogonal. Very recently, Bonvicini (2024) used the reduced row echelon algorithm to handle the core part of finding linearly independent components in Andrews and Ghoul (1982)’s method, to successfully find rank-5 decomposition matrices. Zou et al. (2001) also proposed a general method to perform the decomposition in a recursive style. However, this approach requires recursively decomposing each tensor, rather than having a decomposition tensor that can be directly applied to arbitrary tensors. Moreover, previous

works have relied on finding the linearly independent components of isotropic tensors, which scales in a factorial level, limiting the efficiency. Snider (2017) summarized the development for ICT in his book.

Meanwhile, we aim to understand what operations can be performed on a Cartesian tensor to maintain equivariance. That is one of the reasons ICTs are significant, as linear combinations of them are naturally equivariant. Pearce-Crump (2023b) leveraged Brauer (1937)’s theorem to obtain a spanning set for the equivariant design space. Based on this, Pearce-Crump (2023a) further proposes an algorithm to generate the equivariant-preserving operation on the tensor product space. However, a direct construction of an orthogonal equivariant basis remains unknown. This work proposes a method to efficiently obtain orthogonal bases of equivariant design spaces.

## 8 Discussion

### 8.1 Relation to other works

The remarkable work done by Pearce-Crump (2023b) has successfully found spanning sets for the equivariant design spaces for various groups. Here, we discuss the relation to this work. First, we recall the theorem for  $O(n)$ :

**Theorem 18 (Pearce-Crump (2023b))**  $U_{i_1, j_1} U_{i_2, j_2} \dots U_{i_{\frac{l+k}{2}}, j_{\frac{l+k}{2}}}$  form a spanning set for  $\text{Hom}_{O(n)} \left( (\mathbb{R}^n)^{\otimes l}, (\mathbb{R}^n)^{\otimes k} \right)$ , where  $U$  are an unit matrices (also rank-2 tensors),  $i_i$  and  $j_i$  are different permutations of integer  $1, \dots, \frac{l+k}{2}$ .  $\text{Hom}_{O(n)} \left( (\mathbb{R}^n)^{\otimes l}, (\mathbb{R}^n)^{\otimes k} \right) = \emptyset$  when  $l+k$  is odd, so the above  $\frac{l+k}{2}$  is well defined.

Compared to his theorem, we identify orthogonal bases between arbitrary spaces where the input and output spaces can be any spherical tensor product spaces, not limited to the Cartesian tensor product spaces. For example, Pearce-Crump (2023b)’s theorem cannot handle  $((l = 3 \otimes l = 4), p = -1) \rightarrow ((l = 2 \otimes l = 2), p = -1)$ , but our method can well address it, with an orthogonal basis instead of the spanning set. As he has noted, there is no equivariant operation between Cartesian spaces of odd and even ranks, respectively, while through our method we can find an orthogonal basis if their parities are the same.

There is also a striking advance made by Finzi et al. (2021), who aim to find the basis of equivariant spaces. However, they adopt a numerical algorithm, EMLP, instead of our algorithm, which provides an analytical solution. As noted by Pearce-Crump (2023b), Finzi et al. (2021) cannot generate bases for continuous group tensor product spaces with higher ranks due to memory constraints and speed limitations. To further evaluate the performance of basis generation, we compare our method with that of Finzi et al. (2021) using their open-sourced library. The results are presented in Table 4, where EMLP is able to compute the basis of  $\text{End}_{O(3)} \left( (\mathbb{R}^{\otimes 5}) \right)$  in 26 minutes. In contrast, our method computes the same basis in less than one second and is capable of obtaining the basis of  $\text{End}_{O(3)} \left( (\mathbb{R}^{\otimes 8}) \right)$ .

It is worth noting that Finzi et al. (2021) claim polynomial complexity based on dimensions, whereas our complexity is determined by the ranks of the tensor product spaces.




Figure 6: The parentage scheme of decomposition of  $(l = 0.5)^{\otimes n}$  for  $SU(2)$ , where  $n$  is the rank. The numbers in the circle represent multiplicity.

## 8.2 Other groups

Although  $O(3)$  is the most popular and useful group considered in equivariant neural networks, our conclusion can also be extended to  $O(n)$ ,  $SO(n)$ , and  $SU(n)$ . The tensor product spaces can always be decomposed to smaller irreducible representation spaces, and we can always construct CG coefficients with orthogonality and normalization. The main challenge lies in obtaining the CG coefficients. Once they are obtained, the procedures outlined in this work can be repeated for those groups, and Proposition 17 will give us the equivariant basis. Fortunately, we already have a robust method for computing CG coefficients for  $SU(n)$ , proposed by Alex et al. (2011), who also provides an online service to freely compute the CG coefficients for  $SU(n)$ . One can generally follow the procedure of our work to extend the method to  $SU(n)$ . There are several relations between  $SU(n)$  and certain  $SO(n)$ . For example,  $SO(3)$  is double covered by  $SU(2)$ , which allows us to directly obtain CG coefficients for  $SO(3)$  from those for  $SU(2)$ . The relationship stems from the fact that  $SPIN(n)$  double covers  $SO(n)$ , and  $SU(2)$  is isomorphic to  $SPIN(3)$ . Further,  $SU(2) \otimes SU(2)$  is isomorphic to  $SPIN(4)$ , which double covers  $SO(4)$ . Similarly,  $SU(4)$  is isomorphic to  $SPIN(6)$ , and thus it double covers  $SO(6)$ . Consequently, the conclusion can also be extended to those  $SO(n)$  that are covered by  $SU(n')$  for high  $n$ . Further research



into the Lie algebra of  $SO(n)$  has the potential to directly discover the CG coefficients of  $SO(n)$ , rather than relying on the relationship with  $SU(n)$ . Additionally,  $SO(n)$  can be easily converting to  $O(n)$  by considering the parity. We additionally provide code for obtaining basis for equivariant spaces under  $U(2)/SU(2)$  in our code base. The parentage scheme is different from that of  $O(3)/SO(3)$ , as shown in Figure 6.

## 9 Conclusion

In this work, we leverage the relationship between spherical spaces and their tensor product spaces, constructing path matrices to propose methods to: 1) find high-order ICT decomposition matrices for  $6 \leq n \leq 9$ , 2) find orthogonal bases for equivariant layers where the input and output spaces are identical, and 3) generalize the conclusions to arbitrary pairs of spaces. The conclusions made in this work hold true for  $O(n)$ ,  $SO(n)$ , and  $SU(n)$ . However, currently, we can only efficiently calculate  $SU(n)$ ,  $O(3)$ , and  $SO(3)$  due to the availability of known CG coefficients. This also highlights our future work, which will focus on achieving efficient calculations or closed-form formulae for  $O(n)$  and  $SO(n)$ . The contributions of this work can be directly applied to EGNN designs as well as areas related to chemistry and physics.

## 10 Boarder social impact

There are several scenarios to which our theory can be applied. One of the most promising directions is the flexible design of EGNNs, which has already been shown to facilitate numerous applications in physics, chemistry, and broader biomedical fields. Examples include machine learning for force fields, drug design, blind docking, molecule generation, and many others. Additionally, there is a growing demand for equivariant methods in robotics (Wang et al., 2024). Thus, our work has the potential to positively impact these related fields. However, it is essential to establish strict regulations for methods that could potentially harm the health of people and animals.

## 11 Reproducibility statement

The code to efficiently implement the theory is provided in <https://github.com/ShihaoShao-GH/ICT-decomposition-and-equivariant-bases>. The only external dependency packages are PyTorch and e3nn. The e3nn package can be easily installed using the command `pip install --upgrade e3nn`. The code can run on any computer with Python version  $> 3.0.0$  and PyTorch version  $> 1.0$ . The code for generating the basis for  $SU(2)$  does not require the e3nn package. However, the code for generating Clebsch-Gordan (CG) coefficients is borrowed from <http://qutip.org/docs/3.1.0/modules/qutip/utilities.html>. The CG coefficients for  $SU(n)$  can be obtained from

<https://homepages.physik.uni-muenchen.de/~vondelфт/Papers/ClebschGordan/>, as provided by Alex et al. (2011). The code for generating these coefficients is detailed on pages 19–34 of their paper. The comparison with EMLP is based on their code base

<https://github.com/mfinzi/equivariant-MLP>.

## Acknowledgement

S. Shao acknowledges the support from the Principal's Scholarship of Peking University. Z. Lin was supported by National Key R&D Program of China (2022ZD0160300) and the NSF China (No. 62276004). Q. Cui was supported by grant from the National Natural Science Foundation of China (No. 62025102).

## Appendix A. Irreducible representations and spherical spaces

We first recall some basic facts. A thorough introduction to group theory can be found in Inui et al. (2012) and Bröcker and Tom Dieck (2013). We first focus on the  $O(3)$  group (3D rotation and reflection), but we also extend our conclusions to  $O(n)$ ,  $SO(n)$ , and  $SU(n)$  in Section 8. Consider a vector  $v \in V$ . The rotation and reflection operations should be able to act on it. Representation theory shows that  $O(3)$  can be represented by some real matrices in the Cartesian space via a group homomorphism  $\rho : G \rightarrow \mathbb{R}^{3 \times 3}$ , which is called the natural representation. It holds that

$$\rho(g_1 * g_2) = \rho(g_1) \cdot \rho(g_2)$$

for any  $g_1, g_2 \in G$ . The group element  $g$  acting on the vector  $v$  can therefore be represented as a matrix multiplication  $\rho(g) \cdot v$ . Sometimes, one may also refer to the space of  $v$  as a representation space. If the results of the group action remain in the same space, i.e.,  $\rho(g)v \in V$ . In this sense, this place is also called a  $G$ -invariant space or subspace. To represent  $O(3)$ , the matrix  $\rho(g)$  is constrained to be orthogonal. It is important to note that the co-domain of  $\rho$  can differ, meaning it can also be represented by matrices of other dimensions. In this paper, we denote the co-domain of  $\rho$  using a subscript when it should be made explicit, e.g.,  $\rho_{\mathbb{R}^3}$ . Now, it suffices to define equivariance. We claim that a function  $f : X \rightarrow Y$  is equivariant if and only if

$$\rho_Y f(x) = f(\rho_X(x)).$$

We first focus on the simpler case where  $X = Y$ . We will later generalize this to the case where  $X \neq Y$ , but  $Y$  is known and remains consistent across different  $M$ . When it is necessary to emphasize the change in space, we refer to a linear mapping  $M$  as  $G$ -( $X, Y$ )-equivariant.

We then introduce the definition of reducible and irreducible representation, in a matrix view:

**Definition 19** *A group homomorphism  $\rho \in \text{Hom}_G(\mathbb{R}^k)$  is said to be reducible, if there is an invertible matrix  $M$ , such that for every  $g \in G$ , we have*

$$M\rho(g)M^{-1} = \text{diag}(\rho_1(g), \rho_2(g))$$

for some non-trivial  $\rho_1$  and  $\rho_2$ .

The  $3 \times 3$  rotation and reflection matrices are irreducible. More generally,

**Proposition 20** *The natural representation  $\{M \in GL(n, \mathbb{R}) \mid MM^T = M^T M = I\}$  of  $O(n)$  is irreducible.*

If a representation is irreducible, then the vectors from the space it acts on is also said to be irreducible. Irreducible representations play a very important role in theoretical chemistry and physics, as they represent different symmetries. It is also one of the major building blocks for EGNNs in recent years. There are more irreducible representations than the natural one. Each irreducible representations space is spanned by the **real** spherical harmonics  $Y_{lm}$  as a basis of different degree  $l$  and order  $m$ . The complex spherical harmonics  $Y_l^m : S^2 \rightarrow \mathbb{C}$  are the solutions of Laplace’s equation, which are in the form of associated Legendre polynomials  $P_l^m$  taking spherical coordinates  $(\theta, \varphi)$  as variables.  $l$  can take values from  $(\frac{1}{2}, 1, \frac{3}{2}, \dots)$ . They are highly related to  $SU(2)$  group, which is of great importance for quantum physics. Leveraging the fact that  $SU(2)$  is a double-cover of  $SO(3)$ , analysis implies that the real form of spherical harmonics  $Y_{lm}$  of integer  $l$  form a basis for weight  $l$   $O(3)$  irreducible representation. In this case,  $l \in \{1, 2, \dots\}$ , and  $m \in \{-l, \dots, l\}$ . Thus  $O(3)$  has  $3 \times 3, 5 \times 5, \dots$  matrices as representations. We sometimes use  $(l = k)$  to denote a weight  $k$  spherical space. As said before, an important property of spherical harmonics is:

**Proposition 21** *The representation  $v$  in weight  $l$  spherical space with real spherical harmonics  $Y_{lm}$  as the basis,  $v \in \text{span}_{\mathbb{R}}(Y_{l,-l}, \dots, Y_{l,l})$ , is irreducible.*

There is also a connection between spherical and Cartesian spaces, which is a building block of our arguments:

**Lemma 22** *The weight one real spherical space is exactly the 3-d Cartesian space up to an orthogonal change-of-basis.*

This lemma is often utilized in engineering implementation (Thomas et al., 2018; Weiler et al., 2018; Kondor et al., 2018). In this work, it is a key connecting the spherical and Cartesian space. Also, we have the following properties:

**Proposition 23** *Real spherical harmonics are homogeneous polynomials,  $Y_{lm} \in P_l(x, y, z)$ , where  $l$  is the degree of the polynomials, and  $Y_{lm} \perp Y_{l'm'}$  for  $l' \neq l$  or  $m' \neq m$ .  $\{Y_{lm}\}$  span  $\binom{l+2}{2}$ -dimensional vector spaces, with unit  $L_2$ -norms.*

Appendix B. Visualization of decomposition matrices




Figure 7: ICT decomposition matrix with path  $(1 \rightarrow 1 \rightarrow 2 \rightarrow 3)$ .




Figure 8: ICT decomposition matrix with path  $(1 \rightarrow 1 \rightarrow 2 \rightarrow 3 \rightarrow 3)$ .




Figure 9: ICT decomposition matrix with path  $(1 \rightarrow 1 \rightarrow 0 \rightarrow 1 \rightarrow 2 \rightarrow 1)$ .




Figure 10: ICT decomposition matrix with path  $(1 \rightarrow 2 \rightarrow 3 \rightarrow 2 \rightarrow 2 \rightarrow 1 \rightarrow 1)$ .




Figure 11: ICT decomposition matrix with path  $(1 \rightarrow 1 \rightarrow 1 \rightarrow 2 \rightarrow 1 \rightarrow 2 \rightarrow 2)$ .




Figure 12: Equivariant basis matrix of  $\text{End}_{O(3)}\left(\left(\mathbb{R}^3\right)^{\otimes 6}\right)$  with paths  $(1 \rightarrow 2 \rightarrow 2 \rightarrow 3 \rightarrow 3 \rightarrow 3)$  and  $(1 \rightarrow 1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 3)$ .




Figure 13: Equivariant basis matrix of  $\text{Hom}_{O(3)}((l = 1 \otimes l = 2), (l = 1 \otimes l = 3))$  with paths  $(1 \xrightarrow{2} 2)$  and  $(1 \xrightarrow{3} 2)$ .




Figure 14: Equivariant basis matrix of  $\text{Hom}_{SU(2)}((0.5 \otimes 0.5 \otimes 1.5), (0.5 \otimes 0.5 \otimes 0.5))$  with paths  $(0.5 \xrightarrow{0.5} 0 \xrightarrow{1.5} 1.5)$  and  $(0.5 \xrightarrow{0.5} 1 \xrightarrow{0.5} 1.5)$ .



## References

- Arne Alex, Matthias Kalus, Alan Huckleberry, and Jan von Delft. A numerical algorithm for the explicit calculation of  $\text{su}(n)$  and  $\text{sl}(n, \mathbb{C})$  clebsch–gordan coefficients. *Journal of Mathematical Physics*, 52(2), 2011.
- DL Andrews and WA Ghoul. Irreducible fourth-rank cartesian tensors. *Physical Review A*, 25(5):2647, 1982.
- Ilyes Batatia, David P Kovacs, Gregor Simm, Christoph Ortner, and Gábor Csányi. Mace: Higher order equivariant message passing neural networks for fast and accurate force fields. *Advances in Neural Information Processing Systems*, 35:11423–11436, 2022.
- Simon Batzner, Albert Musaelian, Lixin Sun, Mario Geiger, Jonathan P Mailoa, Mordechai Kornbluth, Nicola Molinari, Tess E Smidt, and Boris Kozinsky. E (3)-equivariant graph neural networks for data-efficient and accurate interatomic potentials. *Nature communications*, 13(1):2453, 2022.
- Andrea Bonvicini. Irreducible cartesian tensor decomposition: A computational approach. *The Journal of Chemical Physics*, 160(22), 2024.
- Richard Brauer. On algebras which are connected with the semisimple continuous groups. *Annals of Mathematics*, 38(4):857–872, 1937.
- Theodor Bröcker and Tammo Tom Dieck. *Representations of compact Lie groups*, volume 98. Springer Science & Business Media, 2013.
- Florent Capelli, Arnaud Durand, and Stefan Mengel. The arithmetic complexity of tensor contraction. *Theory of Computing Systems*, 58:506–527, 2016.
- Bingqing Cheng. Cartesian atomic cluster expansion for machine learning interatomic potentials. *npj Computational Materials*, 10(1):157, 2024.
- Stefan Chmiela, Alexandre Tkatchenko, Huziel E Sauceda, Igor Poltavsky, Kristof T Schütt, and Klaus-Robert Müller. Machine learning of accurate energy-conserving molecular force fields. *Science advances*, 3(5):e1603015, 2017.
- Taco Cohen and Max Welling. Group equivariant convolutional networks. In *International conference on machine learning*, pages 2990–2999. PMLR, 2016.
- JAR Coope, RF Snider, and FR McCourt. Irreducible cartesian tensors. *The Journal of Chemical Physics*, 43(7):2269–2275, 1965.
- Çiğdem Dinçkal. Orthonormal decomposition of third rank tensors and applications. In *Proceedings of the World Congress on Engineering*, volume 1, 2013.
- Ralf Drautz. Atomic cluster expansion for accurate and transferable interatomic potentials. *Physical Review B*, 99(1):014104, 2019.

- Marc Finzi, Max Welling, and Andrew Gordon Wilson. A practical method for constructing equivariant multilayer perceptrons for arbitrary matrix groups. In *International conference on machine learning*, pages 3318–3328. PMLR, 2021.
- Xiang Fu, Zhenghao Wu, Wujie Wang, Tian Xie, Sinan Keten, Rafael Gomez-Bombarelli, and Tommi Jaakkola. Forces are not enough: Benchmark and critical evaluation for machine learning force fields with molecular simulations. *arXiv preprint arXiv:2210.07237*, 2022.
- Johannes Gasteiger, Janek Groß, and Stephan Günnemann. Directional message passing for molecular graphs. *arXiv preprint arXiv:2003.03123*, 2020.
- Johannes Gasteiger, Florian Becker, and Stephan Günnemann. Gemnet: Universal directional graph neural networks for molecules. *Advances in Neural Information Processing Systems*, 34:6790–6802, 2021.
- Justin Gilmer, Samuel S Schoenholz, Patrick F Riley, Oriol Vinyals, and George E Dahl. Neural message passing for quantum chemistry. In *International conference on machine learning*, pages 1263–1272. PMLR, 2017.
- Marco Gori, Gabriele Monfardini, and Franco Scarselli. A new model for learning in graph domains. In *Proceedings. 2005 IEEE international joint conference on neural networks, 2005.*, volume 2, pages 729–734. IEEE, 2005.
- Teturo Inui, Yukito Tanabe, and Yositaka Onodera. *Group theory and its applications in physics*, volume 78. Springer Science & Business Media, 2012.
- Ilya Kisil, Giuseppe G Calvi, Kriton Konstantinidis, Yao Lei Xu, and Danilo P Mandic. Reducing computational complexity of tensor contractions via tensor-train networks. *arXiv preprint arXiv:2109.00626*, 2021.
- Risi Kondor, Zhen Lin, and Shubhendu Trivedi. Clebsch–gordan nets: a fully fourier space spherical convolutional neural network. *Advances in Neural Information Processing Systems*, 31, 2018.
- Yann LeCun, Bernhard Boser, John S Denker, Donnie Henderson, Richard E Howard, Wayne Hubbard, and Lawrence D Jackel. Backpropagation applied to handwritten zip code recognition. *Neural computation*, 1(4):541–551, 1989.
- Yi-Lun Liao, Brandon Wood, Abhishek Das, and Tess Smidt. Equiformerv2: Improved equivariant transformer for scaling to higher-degree representations. *arXiv preprint arXiv:2306.12059*, 2023.
- Albert Musaelian, Simon Batzner, Anders Johansson, Lixin Sun, Cameron J Owen, Mordechai Kornbluth, and Boris Kozinsky. Learning local equivariant representations for large-scale atomistic dynamics. *Nature Communications*, 14(1):579, 2023.
- Edward Pearce-Crump. An algorithm for computing with brauer’s group equivariant neural network layers. *arXiv preprint arXiv:2304.14165*, 2023a.

- Edward Pearce-Crump. Brauer’s group equivariant neural networks. In *International Conference on Machine Learning*, pages 27461–27482. PMLR, 2023b.
- Lars Ruddigkeit, Ruud Van Deursen, Lorenz C Blum, and Jean-Louis Reymond. Enumeration of 166 billion organic small molecules in the chemical universe database gdb-17. *Journal of chemical information and modeling*, 52(11):2864–2875, 2012.
- Kristof Schütt, Pieter-Jan Kindermans, Huziel Enoc Saucedo Felix, Stefan Chmiela, Alexandre Tkatchenko, and Klaus-Robert Müller. Schnet: A continuous-filter convolutional neural network for modeling quantum interactions. *Advances in neural information processing systems*, 30, 2017.
- Kristof Schütt, Oliver Unke, and Michael Gastegger. Equivariant message passing for the prediction of tensorial properties and molecular spectra. In *International Conference on Machine Learning*, pages 9377–9388. PMLR, 2021.
- Guillem Simeon and Gianni De Fabritiis. Tensornet: Cartesian tensor representations for efficient learning of molecular potentials. *Advances in Neural Information Processing Systems*, 36, 2024.
- Guillem Simeon, Antonio Mirarchi, Raul P Pelaez, Raimondas Galvelis, and Gianni De Fabritiis. On the inclusion of charge and spin states in cartesian tensor neural network potentials. *arXiv preprint arXiv:2403.15073*, 2024.
- Robert F Snider. *Irreducible Cartesian Tensors*, volume 43. Walter de Gruyter GmbH & Co KG, 2017.
- Nathaniel Thomas, Tess Smidt, Steven Kearnes, Lusann Yang, Li Li, Kai Kohlhoff, and Patrick Riley. Tensor field networks: Rotation-and translation-equivariant neural networks for 3d point clouds. *arXiv preprint arXiv:1802.08219*, 2018.
- Dian Wang, Stephen Hart, David Surovik, Tarik Kelestemur, Haojie Huang, Haibo Zhao, Mark Yeatman, Jiuguang Wang, Robin Walters, and Robert Platt. Equivariant diffusion policy. In *8th Annual Conference on Robot Learning*, 2024.
- Maurice Weiler, Mario Geiger, Max Welling, Wouter Boomsma, and Taco S Cohen. 3d steerable cnns: Learning rotationally equivariant features in volumetric data. *Advances in Neural Information Processing Systems*, 31, 2018.
- Viktor Zaverkin, Francesco Alesiani, Takashi Maruyama, Federico Errica, Henrik Christiansen, Makoto Takamoto, Nicolas Weber, and Mathias Niepert. Higher-rank irreducible cartesian tensors for equivariant message passing. *arXiv preprint arXiv:2405.14253*, 2024.
- W-N Zou, Q-S Zheng, D-X Du, and J Rychlewski. Orthogonal irreducible decompositions of tensors of high orders. *Mathematics and Mechanics of Solids*, 6(3):249–267, 2001.