

# Using the Th III Ion for a Nuclear Clock and Searches for New Physics

V. A. Dzuba and V. V. Flambaum

*School of Physics, University of New South Wales, Sydney 2052, Australia*

The  $^{229}\text{Th}$  nucleus possesses a unique low-frequency transition at 8.4 eV, which is being considered for the development of an extremely accurate nuclear clock. We investigate an electronic bridge process in the Th III ion, where nuclear excitation occurs via electronic transitions, and demonstrate that a proper choice of laser frequencies can lead to a significant enhancement of this effect. Electrons also reduce the lifetime of the nuclear excited state.

Additionally, the electronic structure of the Th III ion exhibits features that make it particularly useful for probing new physics. Notably, it contains a metastable state connected to the ground state via a weak M2 transition, which can be utilized for quantum information processing, as well as searches for oscillating axion field, violation of local Lorentz invariance, test of the Einstein's equivalence principle, and measurement of nuclear weak quadrupole moment. The electronic states of the ion present a unique case of level crossing involving the  $5f$ ,  $6d$ , and  $7s$  single-electron states. This crossing renders the transition frequencies highly sensitive to potential time-variation of the fine-structure constant.

The nucleus of the  $^{229}\text{Th}$  isotope has a unique feature - a low-lying excited state [1] connected to the ground state via a nuclear M1 transition. It has been proposed to use this transition as the basis for a nuclear clock [2] with exceptionally high accuracy (up to  $\sim 10^{-19}$  [3]), which is also highly sensitive to new physics [4, 5].

The small size of the nucleus and the shielding effect of atomic electrons render the transition frequency largely insensitive to perturbations when the Th ion is in a "stretched" state, i.e., with maximal projections of both electronic and nuclear angular momenta [3]. However, the small nuclear transition energy (8.4 eV) arises from a strong cancellation between contributions from strong and electromagnetic nuclear forces. This makes the transition frequency highly sensitive to small changes in each of these contributions, which may result from a hypothetical variation of fundamental physical constants, such as the fine-structure constant  $\alpha$ , strong interaction constants, and quark masses [4].

Furthermore, this transition is highly sensitive to dark matter fields, which could induce variations in these constants [6, 7], as well as potential violations of Lorentz invariance and the Einstein's equivalence principle [5].

The measurement of the energy of the nuclear clock transition in  $^{229}\text{Th}$  has been an ongoing effort for many years [8–10], with significant recent progress [11–13]. The transition energy has been measured to be  $\omega_N = 8.355733(2)_{\text{stat}}(10)_{\text{sys}}$  eV ( $67393 \text{ cm}^{-1}$ ) [12] and  $\omega_N = 2,020,407,384,335(2)$  kHz [13] in Th atoms inside solids. A higher accuracy is expected in ion clocks.

The amplitude of the nuclear M1 transition is suppressed by five orders of magnitude compared to typical atomic E1 transitions. It is possible to enhance this transition using the electronic bridge (EB) process, in which electronic transitions induce nuclear transitions via hyperfine interaction and vice versa. The EB process has been explored in several Th II and Th IV ion calculations [14–17]. In particular, Ref. [16] proposed using the EB process for nuclear excitation in Th II.

In this work, we consider a similar EB process for the

Th III ion, which may offer several advantages over Th IV and Th II. The electronic spectrum of Th III is rich enough to enable strong enhancement of the EB effect through a two-step process, similar to that suggested in Ref. [16] for Th II. At the same time, its relatively simple electronic structure, consisting of two valence electrons above a closed-shell core, allows for more accurate calculations and a clearer interpretation of experimental results. The Th III ion also possesses low-energy E1 transitions, which can be used for ion cooling.

Beyond its role in nuclear clock development, the electronic structure of Th III has unique features that make it valuable for other applications. Its first excited state is a low-lying metastable state (excitation energy  $\approx 63 \text{ cm}^{-1}$ ) connected to the ground state via a weak M2 transition. This doublet of states can be used for quantum information processing (see similar proposal for highly charged ions in Ref. [20]) and for the search for oscillating axion fields. The axion field may interact directly with electrons or induce an oscillating nuclear magnetic quadrupole moment, which in turn stimulates the M2 transition. Unlike photon M2 transitions, axion-induced M2 transitions are not suppressed, leading to a significant relative reduction in background noise [21].

The ground state of Th III is also promising for testing local Lorentz invariance (LLI) violation. Its large total electron angular momentum and the presence of an electron in the open  $5f$  subshell ensure an enhancement of LLI violation effect - see detailed explanation in Ref. [22]. The M2 transition can further be used to search for violations of the Einstein's equivalence principle (EEP), which may manifest via an annual modulation of atomic frequencies due to the varying distance to Sun and corresponding variation of its gravitational potential - see e.g. [23–27].

Additionally, Th III exhibits a unique case of multiple level crossing. The energies of the  $5f$ ,  $6d$ , and  $7s$  single-electron states are approximately equal, making the transition frequencies particularly sensitive to potential time variation of the fine-structure constant  $\alpha$  - see

TABLE I. States of interest for electronic bridge process in Th III. Abbreviation GS stands for ground state. Other states are named in accordance with diagram of Fig. 1, i.e., states T1 and T2 appear on line  $t$ , states N1, N2, N3, N4 - on line  $n$ , states S1, S2, S3, S4 - on line  $s$ .

State name	Confi-guration	Term	$J$	NIST [34]		This work	
				Energy [cm <sup>-1</sup> ]	Landé	Energy [cm <sup>-1</sup> ]	Landé
GS	$5f6d$	$^3\text{H}^\circ$	4	0	0.888	0	0.885
T1	$5f^2$	$^1\text{G}$	4	25972	1.072	29323	1.11
T2	$5f7p$	$(7/2, 1/2)$	4	38580	1.105	38190	1.107
T3	$6d^2$	$^3\text{F}_2$	2	63	0.744	3056	0.7318
N1	$5f8s$	$(5/2, 1/2)^\circ$	3	74784		74030	1.05
N2	$5f7d$	$(5/2, 3/2)^\circ$	3	78328		78263	0.820
N3	$5f8s$		4	78417		78573	1.22
N4	$7s7p$	$^1\text{P}^\circ$	1	69001		71635	1.018
S1	$5f8s$		3	7501	1.027	6901	1.029
S2	$5f6d$	$^3\text{D}^\circ$	3	10741	1.22	13685	1.18
S3	$5f6d$	$^1\text{F}^\circ$	3	15453	1.07	18110	1.12
S4	$5f6d$	$^3\text{F}^\circ$	2	511	0.739	957	0.785

TABLE II. Electric dipole (E1) amplitudes (reduced matrix elements  $|\langle A||\mathbf{D}||B\rangle|$ , a.u.) between states of interest presented in Table I.

A \ B	GS	N1	N2	N3
T1	1.1628	0.2420	0.2245	0.4254
T2	3.520	0.9187	1.0049	3.0468

explanation in Refs. [31–33]. This sensitivity is especially pronounced for the aforementioned M2 transition, as the small value of its frequency  $\omega$  leads to an enhancement of the relative effect  $\delta\omega/\omega$ .

Finally, the ground state of Th III (with total electron angular momentum  $J = 4$  and negative parity) is mixed with the metastable  $J = 2$  positive-parity state at  $63 \text{ cm}^{-1}$  via the interaction of electrons with the nuclear weak quadrupole moment. The small  $63 \text{ cm}^{-1}$  energy denominator leads to an enhancement of corresponding parity-violating effects. Measuring these effects would enable, for the first time, the determination of the quadrupole moment of the neutron distribution in nuclei, which provides the dominant contribution to the weak quadrupole moment [28–30].

In this paper, we present accurate calculations of the discussed effects in the Th III ion. Our analysis includes the EB processes in the excitation and decay of the 8.4 eV nuclear clock state, the sensitivity of atomic frequencies to variations in the fine-structure constant  $\alpha$ , and other effects relevant to nuclear clock development and the search for new physics.

**Electronic bridge for nuclear excitation and decay in Th III.** Our approach follows the method used in Ref. [16] for the Th II ion. The decay of any atomic state with energy  $E_n > \omega_N$  may include nuclear excitation. Since the nuclear excitation energy  $\omega_N = 67393 \text{ cm}^{-1}$  lies outside the optical region, we consider a two-




FIG. 1. A diagram for the EB process in Th III. It is assumed that the atom is in initial electronic state  $t$ ;  $n$  and  $s$  are intermediate and final electronic states,  $g$  ground nuclear state,  $m$  is isomeric nuclear state.

step excitation of the electronic states. In the first step, the atom is excited from the ground state (GS) to an intermediate state  $t$  with energy  $\omega_1 \sim \omega_N/2$ . We have found two suitable states with electron angular momentum  $J = 4$ , referred to as T1 and T2 (see Table I), which connected to the GS by a strong electric dipole (E1) transition. We have not found suitable states with  $J = 5$ . We exclude states with  $J = 3$  to avoid leakage into the metastable state  $5f6d \ ^3\text{F}_2^\circ$  ( $E = 511 \text{ cm}^{-1}$ ).

In the second step, the ion is further excited by a second laser to satisfy the energy conservation condition for simultaneous nuclear excitation and excitation of the final electronic state,  $\omega_1 + \omega_2 = \omega_N + \omega_s$ . Note that the final state  $s$  is not the ground state but a low-lying excited state, as this configuration provides the largest EB amplitude. In this process, the intermediate electronic state with energy  $E_n$  is off-resonance, meaning it is virtually excited and subsequently decays, inducing nuclear excitation via hyperfine interaction (through magnetic dipole and electric quadrupole interactions). The diagram illustrating this process is shown in Fig. 1 (see also Ref. [16]).

The mathematical formulation of the EB process involves a summation over all intermediate states  $n$ . However, in this analysis, we focus on the dominant contributions, which come from specific intermediate states  $n$  and final states  $s$  providing a very small energy denominator in Eq. (3). Possible choices for the states  $t$ ,  $n$ , and  $s$  are presented in Table I.

The rate of an induced excitation from state  $b$  to state  $a$   $W_{ba}^{\text{in}}$  can be calculated using the rate of a spontaneous transition  $a \rightarrow b$   $W_{ab}$  [35]:

$$W_{ba}^{\text{in}} = W_{ab} \frac{4\pi^3 c^2}{\omega^3} I_\omega. \quad (1)$$

Here  $I_\omega$  is the intensity of isotropic and unpolarized incident radiation. Index  $b$  corresponds to nuclear ground state and electron excited state, index  $a$  corresponds to nuclear excited state and electronic ground or low energy excited state. The rate of a spontaneous transition via electronic bridge process is given by [14]

$$W_{ab} = \frac{4}{9} \left(\frac{\omega}{c}\right)^3 \frac{|\langle J_g || M_k || I_m \rangle|^2}{(2I_m + 1)(2J_t + 1)} G_2^{(k)}, \quad (2)$$




FIG. 2. A diagram for nuclear excitation by decaying of highly excited atomic state  $t$  and emitting photon.

Keeping in mind the relation (1), we assume that  $\omega$  in (2) is the frequency of second excitation ( $\omega \equiv \omega_2$ ) which is chosen to get into a resonance situation,  $\omega_2 = \epsilon_s + \omega_N - \omega_1$ . Factor  $G_2$  in (2) depends on electrons only. It corresponds to upper line of Fig. 1. In principle, it has summation over complete set of intermediate states  $n$ , see e.g. [14]). However, assuming resonance situation and keeping only one strongly dominating term, we have

$$G_2^{(k)} \approx \frac{1}{2J_n + 1} \left[ \frac{\langle s | T_k | n \rangle \langle n | D | t \rangle}{\omega_{ns} - \omega_N} \right]^2. \quad (3)$$

Here  $T_k$  is the electron part of the hyperfine interaction operator (magnetic dipole (M1) for  $k = 1$  and electric quadrupole (E2) for  $k = 2$ ),  $D$  is the electric dipole operator (E1). States  $n$  and  $s$  are chosen to get close to a resonance,  $\epsilon_n - \epsilon_s \equiv \omega_{ns} \approx \omega_N$ .

It is convenient to present the results in terms of dimensionless ratios ( $\beta$ ) of electronic transition rates to nuclear transition rates.

$$\beta_{M1} = \Gamma_{EB}^{(1)} / \Gamma_N(M1), \quad \beta_{E2} = \Gamma_{EB}^{(2)} / \Gamma_N(E2). \quad (4)$$

Here  $\Gamma^{(k)}$  is given by (2) ( $\Gamma^{(k)} \equiv W_{ab}$ ). Both parameters,  $\beta_{M1}$  and  $\beta_{E2}$  can be expressed via  $G_2$ , (Eq. (3)) [17]

$$\beta_{M1} = \left( \frac{\omega}{\omega_N} \right)^3 \frac{G_2^{(1)}}{3(2J_s + 1)}, \quad (5)$$

$$\beta_{E2} = \left( \frac{\omega}{\omega_N} \right)^3 \frac{4G_2^{(2)}}{k_n^2(2J_s + 1)}. \quad (6)$$

The  $M1$  and  $E2$  contributions can be combined into one effective parameter  $\tilde{\beta}$  using the known ratio of the widths of the nuclear  $M1$  and  $E2$  transitions,  $\gamma = \Gamma_\gamma(E2) / \Gamma_\gamma(M1) = 6.9 \times 10^{-10}$ , [38]. Then  $\tilde{\beta} = \beta_{M1}(1 + \rho)$ , where  $\rho = \gamma\beta_{E2} / \beta_{M1}$ .

The results of the calculations for a number of possible transitions are presented in Table III. We see that the probability of the nuclear excitation may be enhanced up to 295 times. This enhancement may be achieved by proper choices of laser frequencies in a two-step process of atomic excitation which is followed by the nuclear excitation.

There is another possibility of the nuclear excitation via EB process, see Fig. 2. If an atom is in excited state  $t$  with energy larger than the nuclear excitation energy,

$\epsilon_t > \omega_N$ , then the decay of this state may include a channel with the nuclear excitation via the hyperfine interaction while the excess of the energy is taken away by emitted photon. This contribution is strongly suppressed due to a small value of the photon frequency,  $\omega = \epsilon_t - \epsilon_s - \omega_N$ , since  $\beta \sim \omega^3$ , see Eqs. (5,6). For example, if  $t$  is the state with  $\epsilon_t = 83701 \text{ cm}^{-1}$  and  $J = 3$ , then  $\beta \approx 4 \times 10^{-3}$ .

The electronic bridge also decreases the lifetime of the nuclear excited state. There is no single dominant contribution in this case. The summation over all intermediate electron states (including continuum) and final electrons states gives  $\tilde{\beta} \approx 0.6$ .

**Th III ion cooling transition.** A cooling may be produced by the laser-induced E1 transition from the negative parity ground state with  $J = 4$  to the positive parity state with electron angular momentum  $J = 5$  and energy  $E = 17887 \text{ cm}^{-1}$ . The decay rate to GS,  $2.9 \times 10^5 \text{ 1/s}$ , is relatively high. Large difference of angular momenta,  $\Delta J = 3$ , helps to suppress leaking to metastable states with  $J = 2$ ,  $E = 511 \text{ cm}^{-1}$  and  $E = 63 \text{ cm}^{-1}$ .

**Applications of Th III ion not related to nuclear clock.** As it was mentioned above, Th III ion has very low lying ( $63 \text{ cm}^{-1}$ ) metastable state which has extremely large lifetime and can be considered as a second ground state. This doublet of states can be used for quantum information processing. In  $^{229}\text{Th}$  isotope the leading channel of decay is the E1 transition to the ground state mediated by the hyperfine interaction. It might be advantageous to use stable  $^{232}\text{Th}$  isotope instead. It has zero nuclear spin and thus no hyperfine structure. The metastable state is connected to the ground state by the extremely weak M2 transition with lifetime  $10^{10}$  years. The transition can be open by applying an external magnetic field. Then the transition amplitude is given by

$$A_{ab} = \left( \sum_n \frac{\langle b | M1 | n \rangle \langle n | E1 | a \rangle}{\epsilon_b - \epsilon_n} + \sum_n \frac{\langle b | E1 | n \rangle \langle n | M1 | a \rangle}{\epsilon_a - \epsilon_n} \right) B \quad (7)$$

Here  $B$  is an external magnetic field in. Using calculated E1 and M1 matrix elements for even and odd states with  $J = 3$  and experimental energies, we get the rate of spontaneous decay  $T_{ab} \approx 2 \times 10^{-5} \text{ s}^{-1} \text{ T}^{-2}$ . The rate of induced excitation can be estimated using Eq. (1). The decay rate is small due to the small transition frequency  $\omega = 63 \text{ cm}^{-1}$ . However, Eq. (1) contains this small frequency in denominator and the excitation probability is not suppressed. Transition between the doublet of the ground states may also be organised via E1 excitation and subsequent decay of higher states.

Ground state of the Th III ion can be used in search for the local Lorentz invariance (LLI) violation while the M2 transition can be used in search for the Einstein's equivalence principle (EEP) violation. Corresponding Hamil-

TABLE III. Versions of the EB process with resonance nuclear excitation. Notations  $t$ ,  $n$ ,  $s$  refer to the diagram on Fig. 1, State names Ti, Ni, Si ( $i=1,2,3$ ) correspond to states in Table I.  $\omega_1$  and  $\omega_2$  are the frequencies of first and second excitations respectively.  $\Delta$  is the energy denominator of the largest contribution.

	$t$	$n$	$s$	$\omega_1$ [cm <sup>-1</sup> ]	$\omega_2$ [cm <sup>-1</sup> ]	$\epsilon_s$ [cm <sup>-1</sup> ]	$\Delta$ [cm <sup>-1</sup> ]	$\beta_{M1}$	$\beta_{E2}$	$\rho$	$\tilde{\beta}$
1	T2	N1	S1	38580	36314	7501	-110	192	3.4[9]	0.01	194
2	T2	N2	S2	38580	39554	10741	193	1.64	1.5[8]	0.06	1.75
3	T2	N3	S2	38580	39554	10741	282	284	1.6[10]	0.04	295
4	T1	N1	S1	25972	48922	7501	-110	33	5.8[8]	0.01	33
5	T1	N3	S2	25972	52162	10741	282	13	7.0[8]	0.03	13

TABLE IV. Parameters of three low-lying states of Th III relevant to search for new physics.  $\alpha_0$  is static dipole polarizability needed for estimation of the BBR shift,  $Q$  is quadrupole moment of the state. Enhancement of two atomic frequencies of the transitions between different configurations due to variation of the fine structure constant.

	Conf.	$J^P$	$E_{\text{NIST}}$ [cm <sup>-1</sup> ]	$E_{\text{calc.}}$ [cm <sup>-1</sup> ]	$\alpha_0$ [a.u.]	$Q$ [a.u.]	$\langle a    T_0^{(2)}    a \rangle$ [a.u.]	$\langle a   H_K   a \rangle$ [a.u.]	$q$ [cm <sup>-1</sup> ]	$K$
GS	5f6d	4 <sup>-</sup>	0	0	13	-4.5	52	1.0	0	0
W1	6d <sup>2</sup>	2 <sup>+</sup>	63	3056	38	0.22	3.0	1.5	-33500	-1060
W2	5f7s	3 <sup>-</sup>	2527	2703					-24160	-19

tonian can be written as [22–25, 36]

$$\delta H = - \left( C_0^{(0)} - \frac{2U}{3c^2} c_{00} \right) \frac{\mathbf{p}^2}{2} - \frac{1}{6} C_0^{(2)} T_0^{(2)}, \quad (8)$$

where  $\mathbf{p}$  is electron momentum operator,  $c$  is speed of light,  $U$  is gravitation potential,  $C_0^{(0)}$ ,  $c_{00}$  and  $C_0^{(2)}$  are unknown constants to be found from measurements. First term in (8) violates the EEP via dependence of atomic frequencies on time of the year caused by varying Sun's gravitational potential. The change is periodical with minimum or maximum in January and July. To link the change to the unknown constant  $c_{00}$  one needs to perform the calculations of the matrix elements of the  $\mathbf{p}^2$  operator. The calculations must be relativistic since in the non-relativistic limit all atomic frequencies change at the same rate and the effect is unobservable [27]. The relativistic form of the operator of kinetic energy is  $H_K = c\gamma_0\gamma^j p_j$ . It is convenient to present the result in terms of the relativistic factor  $R$ , which describes the deviation of the energy shift caused by the kinetic energy operator from the value given by the virial theorem [27]

$$R = - \frac{\Delta E_a - \Delta E_b}{E_a - E_b}. \quad (9)$$

Then  $\Delta\omega/\omega = R\frac{2}{3}c_{00}\Delta U/c^2$ . The calculated values of  $\Delta E_a = \langle a | H_K | a \rangle$  and  $\Delta E_b = \langle b | H_K | b \rangle$  for the ground and first excited states of Th III are presented in Table IV. They lead to the large relativistic factor  $R \approx -1700$ . This is due to the small energy denominator in (9). Thus, the sensitivity of the frequency of the M2 transition to the change of the gravitation potential is strongly enhanced.

Second term in (8) causes LLI violation via dependence of the energy intervals between states with dif-

ferent projections  $M$  of the total atomic angular momentum  $J$  on the system orientation, e.g. due to the Earth rotation. The non-relativistic form of the tensor operator  $T_0^{(2)}$  is  $\mathbf{p}^2 - 3p_z^2$ , while the relativistic operator is  $c\gamma_0(\gamma^j p_j - 3\gamma^3 p_3)$ . It was formulated in Ref. [22] that there are at least two conditions for the effect to be large: (a) long living state, (b) large matrix element which can be found in states with open  $4f$  or  $5f$  shells. Both of these conditions are satisfied for the ground state of Th III. Moreover, using the ground state is an advantage compare e.g. to  $\text{Yb}^+$ , where the effect is zero for the ground state. Table IV presents the values of the reduced matrix elements of the tensor operator  $T_0^{(2)}$  for the ground and first excited state of Th III. Note that the value for the ground state is only 2 to 3 times smaller than the value of the matrix elements for excited states of  $\text{Yb}^+$ , which have holes in the  $4f$  subshell.

Finally, the M2 transition can be used in search for time variation of the fine structure constant  $\alpha$ . The sensitivity of the frequency of the transition to the variation of  $\alpha$  is strongly enhanced due to high  $Z$  and due to the fact that the transition is between states of different configurations. The latter can be explained in the following way. The relativistic energy shift of a single-electron state is given by [37]

$$\Delta_n \approx \frac{E_n(Z\alpha)^2}{\nu} \left[ \frac{1}{j+1/2} - 0.6 \right]. \quad (10)$$

Here  $\nu$  is the effective principal quantum number,  $\nu = 1/\sqrt{-2E_n}$ ,  $j$  is the total angular momentum of electron orbital. One can see from(10) that the maximum frequency shift due to  $\alpha$  variation ( $\delta\omega \approx \Delta_{n_1} - \Delta_{n_2}$ ) can be achieved for transitions with the largest  $\Delta j$ . The Th III is a unique atomic system in which the single-electron energies of the  $5f$ ,  $6d$  and  $7s$  states are very close. Therefore,

transition between low-lying states of Th III are usually either  $5f - 6d$  or  $6d - 7s$  transitions. To calculate the sensitivity of atomic frequencies to the variation of the fine structure constant we present them in a form

$$\omega(x) = \omega_0 + q[x - 1], \quad x \equiv \left(\frac{\alpha}{\alpha_0}\right)^2, \quad (11)$$

where  $\omega_0$  and  $\alpha_0$  are physical values of the frequency and fine structure constant respectively,  $q$  is sensitivity coefficient to be found from calculation by varying the value of  $\alpha$  in computer codes and calculating numerical derivative. Parameter  $q$  links variation of atomic frequency to the variation of  $\alpha$

$$\frac{\delta\omega}{\omega} = K \frac{\delta\alpha}{\alpha}. \quad (12)$$

The dimensionless factor  $K = 2q/\omega$  is called *enhancement factor*. The calculated values of  $q$  and  $K$  for two transitions between ground state (GS) and two excited states, marked as W1 and W2, are presented in Table IV. Note that all values of  $q$  and  $K$  are large and negative in nice agreement with (10). The value of  $K$  for the M2 transition ( $|K| \sim 10^3$ ) is one of the largest found in atomic systems (see, e.g. [39, 40]). It can be compared to recently estimated sensitivity of the nuclear transition to variation of  $\alpha$ ,  $K = 5900(2300)$  [41]. Note that if one frequency is measured against the other, the total sensitivity is further enhanced

$$\frac{\delta\omega_N}{\omega_N} - \frac{\delta\omega_{M2}}{\omega_{M2}} \approx 7000 \frac{\delta\alpha}{\alpha}. \quad (13)$$

For further estimations we have calculated static dipole polarizabilities and quadrupole moment for both clock states. The results are presented in Table IV. From

the values of polarizabilities we estimate the black body radiation shift being smaller than 1 Hz. The relative shift  $\sim 10^{-13}$ . The quadrupole moments for both clock states of Th III are smaller than those in clock states of Yb [42]. Note that corresponding frequency shift can be suppressed by averaging over projections of the total angular momentum [42].

This work was supported by the Australian Research Council Grant No. DP230101058.

## Appendix A: Method of calculation

We use the relativistic Hartree-Fock (RHF) method and the combination of the configuration interaction with the single-double coupled cluster (CI+SD) method [43] to calculate two-electron valence states of the Th III ion. To calculate transition amplitudes we use the time-dependent Hartree-Fock method [44] which is equivalent to the well-known random-phase approximation (RPA).

The calculations start from the closed-shell Th V ion. The single-electron basis states for valence electrons are calculated in the field of frozen core using the B-spline technique [45]. The SD equations are first solved for the core, then for valence states [43]. This leads to creation of the one- and two-electron correlation operators  $\Sigma_1$  and  $\Sigma_2$ , which are used in the CI calculations. Solving the RPA equations for valence states leads to the effective operators of external field, which are used to calculate matrix elements between valence states. The accuracy of the calculations is illustrated further in the text by comparing calculated energies and  $g$ -factors with experiment (see Table I).

Matrix elements of the electric dipole operator, which are used in the calculations, are presented in Table II.

- 
- [1] L. A. Kroger and C.W. Reich, Features of the low-energy level scheme of  $^{229}\text{Th}$  as observed in the  $\alpha$ -decay of  $^{233}\text{U}$ , Nuclear Physics A **259**, 29 (1976).
- [2] E. Peik and Chr. Tamm, Nuclear laser spectroscopy of the 3.5 eV transition in Th-229, Europh. Lett. **61**, 181 (2003).
- [3] C. J. Campbell, A. G. Radnaev, A. Kuzmich, V. A. Dzuba, V. V. Flambaum, and A. Derevianko, Single-Ion Nuclear Clock for Metrology at the 19th Decimal Place, Phys. Rev. Lett. **108**, 120802 (2012).
- [4] V. V. Flambaum, Enhanced Effect of Temporal Variation of the Fine Structure Constant and the Strong Interaction in  $^{229}\text{Th}$ , Phys. Rev. Lett. **97**, 092502 (2006).
- [5] V. V. Flambaum. Enhancing the effect of Lorentz invariance and Einstein equivalence principle violation in nuclei and atoms. Phys. Rev. Lett., **117**, 072501 (2016).
- [6] Asimina Arvanitaki, Junwu Huang, and Ken Van Tilburg, Searching for dilaton dark matter with atomic clocks. Phys. Rev. D **91**, 015015 (2015)
- [7] Y. V. Stadnik and V. V. Flambaum. Can dark matter induce cosmological evolution of the fundamental constants of nature? Phys. Rev. Lett., **115**, 201301 (2015).
- [8] C. W. Reich and R. G. Helmer, Energy Separation of the Doublet of Intrinsic States at the Ground State of  $^{229}\text{Th}$ , Phys. Rev. Lett. **64**, 271 (1990).
- [9] Z. O. Guimarães-Filho and O. Helen, Energy of the 3/2+ state of  $^{229}\text{Th}$  reexamined, Phys. Rev. C **71**, 044303 (2005).
- [10] B. R. Beck, J. A. Becker, P. Beiersdorfer, G. V. Brown, K. J. Moody, J. B. Wilhelmy, F. S. Porter, C. A. Kilbourne, and R. L. Kelley, Energy Splitting of the Ground-State Doublet in the Nucleus  $^{229}\text{Th}$ , Phys. Rev. Lett. **98**, 142501 (2007).
- [11] J. Tiedau, M. V. Okhapkin, K. Zhang, J. Thielking, G. Zitzler, E. Peik, F. Schaden, T. Pronebner, I. Morawetz, L. Toscani De Col, F. Schneider, A. Leitner, M. Pressler, G. A. Kazakov, K. Beeks, T. Sikorsky, and T. Schumm. Laser excitation of the th-229 nucleus. Phys. Rev. Lett., **132**, 182501, 2024.
- [12] R. Elwell, Christian Schneider, Justin Jeet, J. E. S. Ter-

- hune, H. W. T. Morgan, A. N. Alexandrova, H. B. Tran Tan, Andrei Derevianko, and Eric R. Hudson, Laser Excitation of the  $^{229}\text{Th}$  Nuclear Isomeric Transition in a Solid-State Host, *Phys. Rev. Lett.* **133**, 013201 (2024).
- [13] Zhang, C., Ooi, T., Higgins, J.S. et al. Frequency ratio of the  $^{229m}\text{Th}$  nuclear isomeric transition and the  $^{87}\text{Sr}$  atomic clock. *Nature* **633**, 63 (2024).
- [14] A. G. Porsev and V. V. Flambaum, Effect of atomic electrons on the 7.6-eV nuclear transition in  $^{229}\text{Th}^{3+}$ . *Phys. Rev. A* **81**, 032504 (2010).
- [15] S. G. Porsev and V. V. Flambaum, Electronic bridge process in  $^{229}\text{Th}^+$ , *Phys. Rev. A* **81**, 042516 (2010).
- [16] S. G. Porsev, V. V. Flambaum, E. Peik, and Chr. Tamm, Excitation of the Isomeric  $^{229m}\text{Th}$  Nuclear State via an Electronic Bridge Process in  $^{229}\text{Th}$ , *Phys. Rev. Lett.* **105**, 182501 (2010).
- [17] Pavlo V. Bilous, Nikolay Minkov, and Adriana Pálffy, Electric quadrupole channel of the 7.8 eV  $^{229}\text{Th}$  transition, *Phys. Rev. C* **97**, 044320 (2018).
- [18] Lin Li, Zi Li, Chen Wang, Wen-Ting Gan, Xia Hua, Xin Tong, Scheme for the excitation of thorium-229 nuclei based on electronic bridge excitation, *Nuclear Science and Techniques* 34:24 (2023). <https://doi.org/10.1007/s41365-023-01169-4>
- [19] Neng-Qiang Cai, Guo-Qiang Zhang, Chang-Bo Fu, Yu-Gang Ma, Populating  $^{229m}\text{Th}$  via two-photon electronic bridge mechanism *NUCL SCI TECH* (2021) 32:59 (0123456789(.,-volV)(0123456789(.,-volV) <https://doi.org/10.1007/s41365-021-00900-3>
- [20] M. S. Safronova, V. A. Dzuba, V. V. Flambaum, U. I. Safronova, S. G. Porsev, and M. G. Kozlov, Highly Charged Ions for Atomic Clocks, Quantum Information, and Search for  $\alpha$  variation, *Phys. Rev. Lett.* **113**, 030801 (2014).
- [21] V. V. Flambaum, H. B. Tran Tan, D. Budker, and A. Wickenbrock, Atomic and molecular transitions induced by axions via oscillating nuclear moments, *Phys. Rev. D* **101**, 073004 (2020).
- [22] V. A. Dzuba, V. V. Flambaum, M. S. Safronova, S. G. Porsev, T. Pruttivarasin, M. A. Hohensee, H. Häffner, Strongly enhanced effects of Lorentz symmetry violation in entangled  $\text{Yb}^+$  ions, DOI: 10.1038/nphys3610, *Nature Physics* 12, 465-468 (2016).
- [23] V. A. Kostelecky and J. D. Tasson, Matter-gravity couplings and Lorentz violation. *Phys. Rev. D* **83**, 016013 (2011).
- [24] V. A. Kostelecky and C. D. Lane, Constraints on Lorentz violation from clock-comparison experiments. *Phys. Rev. D* **60**, 116010 (1999).
- [25] V. A. Kostelecky and C. D. Lane, Nonrelativistic quantum Hamiltonian for Lorentz violation. *J. Math. Phys.* (N.Y.) **40**, 6245 (1999)
- [26] V.V. Flambaum, E.V. Shuryak, How changing physical constants and violation of local position invariance may occur? AIP conference proceedings, **995**, 1 (2008); arxiv: physics/0701220.
- [27] V. A. Dzuba and V. V. Flambaum, Limits on gravitational Einstein equivalence principle violation from monitoring atomic clock frequencies during a year, *Phys. Rev. D* **95**, 015019 (2017).
- [28] O.P. Sushkov, V.V.Flambaum. Effects of parity nonconservation in diatomic molecules. *Sov. Phys. JETP* **48**, 608, 1978 [*Zh. Exp. Teor. Fiz.* **75**, 1208 (1978)].
- [29] V.V. Flambaum, V.A. Dzuba, C. Harabati. Effect of nuclear quadrupole moment on parity nonconservation in atoms, *Phys. Rev. A* **96**, 012516 (2017) .
- [30] B.G.C. Lackenby, V.V. Flambaum, Weak quadrupole moments. *J. Phys. G* **45**, 075105 (2018).
- [31] V.V. Flambaum, S.G. Porsev. Enhanced sensitivity to the fine-structure constant variation in Th IV atomic clock transition, *Phys.Rev. A* **80**, 064502 (2009); arxiv: 0910.3459 physics.atom-ph
- [32] J. C. Berengut, V. A. Dzuba and V. V. Flambaum. Enhanced laboratory sensitivity to variation of the fine-structure constant using highly-charged ions, *Phys. Rev. Lett.* **105**, 120801 (2010),
- [33] J. C. Berengut, V. A. Dzuba, V. V. Flambaum, and A. Ong, Highly charged ions with E1, M1, and E2 transitions within laser range *Phys. Rev. A* **86**, 022517 (2012).
- [34] Kramida, A., Ralchenko, Yu., Reader, J., and NIST ASD Team (2024). NIST Atomic Spectra Database (ver. 5.12), [Online]. Available: <https://physics.nist.gov/asd> [2024, December 15]. National Institute of Standards and Technology, Gaithersburg, MD. DOI: <https://doi.org/10.18434/T4W30F>
- [35] I. I. Sobelman, *Atomic Spectra And Radiative Transitions*, (Springer-Verlag, Berlin, 1979).
- [36] M. A. Hohensee, N. Leefer, D. Budker, C. Harabati, V. A. Dzuba, and V. V. Flambaum, Limits on Violations of Lorentz Symmetry and the Einstein Equivalence Principle using Radio-Frequency Spectroscopy of Atomic Dysprosium, *Phys. Rev. Lett.* **111**, 050401 (2013).
- [37] V. A. Dzuba, V. V. Flambaum, and J. K. Webb, Space-Time Variation of Physical Constants and Relativistic Corrections in Atoms, *Phys. Rev. Lett.* **82**, 888 (1999).
- [38] Pavlo V. Bilous, Nikolay Minkov, and Adriana Pálffy, Electric quadrupole channel of the 7.8 eV  $^{229}\text{Th}$  transition, *Phys. Rev. C* **97**, 044320 (2018).
- [39] V. A. Dzuba and V. V. Flambaum, Atomic calculations and search for variation of the fine structure constant in quasar absorption spectra, *Can. J. Phys.* **87**, 15-23 (2009).
- [40] V. V. Flambaum and V. A. Dzuba, Search for variation of the fundamental constants in atomic, molecular and nuclear spectra, *Can. J. Phys.* **87**, 25-33 (2009).
- [41] K. Beeks, G. A. Kazakov, F. Schaden, I. Morawetz, L. T. de Col, T. Riebner, M. Bartokos, T. Sikorsky, T. Schumm, C. Zhang, T. Ooi, J. S. Higgins, J. F. Doyle, J. Ye, M. S. Safronova, Fine-structure constant sensitivity of the Th-229 nuclear clock transition, arXiv:2407.17300 (2024).
- [42] V. A. Dzuba, V. V. Flambaum, and S. Schiller, Testing physics beyond the standard model through additional clock transitions in neutral ytterbium, *Phys. Rev. A* **98**, 022501 (2018).
- [43] V. A. Dzuba, Combination of the single-double coupled cluster and the configuration interaction methods: application to barium, lutetium and their ions. *Phys. Rev. A* **90**, 012517 (2014).
- [44] V. A. Dzuba, V. V. Flambaum, P. G. Silvestrov, and O. P. Sushkov, Correlation potential method for the calculation of energy levels, hyperfine structure and E1 transition amplitudes in atoms with one unpaired electron, *J. Phys. B: At. Mol. Phys.*, **20**, 1399 (1987).
- [45] W. R. Johnson, S. A. Blundell, and J. Sapirstein, *Finite basis sets for the Dirac equation constructed from B splines*, *Phys. Rev. A* **37**, 307 (1988).