Efficiently Checking Separating Indeterminates

Bernhard Andraschko¹, Martin Kreuzer^{1*}, Le Ngoc Long²

^{1*}Fakultät für Informatik und Mathematik, Universität Passau, Innstraße 33, D-94032 Passau, Germany.

 $^{2^*}$ University of Education, Hue University, 34 Le Loi Street, Hue City, Vietnam.

*Corresponding author(s). E-mail(s): Martin.Kreuzer@uni-passau.de; Contributing authors: Bernhard.Andraschko@uni-passau.de; lelong@hueuni.edu.vn;

Abstract

In this paper we continue the development of a new technique for computing elimination ideals by substitution which has been called Z-separating re-embeddings. Given an ideal I in the polynomial ring $K[x_1, \ldots, x_n]$ over a field K, this method searches for tuples $Z = (z_1, \ldots, z_s)$ of indeterminates with the property that I contains polynomials of the form $f_i = z_i - h_i$ for $i = 1, \ldots, s$ such that no term in h_i is divisible by an indeterminate in Z. As there are frequently many candidate tuples Z, the task addressed by this paper is to efficiently check whether a given tuple Z has this property. We construct fast algorithms which check whether the vector space spanned by the generators of I or a somewhat enlarged vector space contain the desired polynomials f_i . We also extend these algorithms to Boolean polynomials and apply them to cryptoanalyse round reduced versions of the AES cryptosystem faster.

Keywords: elimination ideal, separating indeterminates, Groebner basis, Boolean polynomial, AES cryptosystem

2010: Primary 14Q20; Secondary 14R10 , 13E15 , 13P10

1 Introduction

Computing elimination ideals has long been one of the key tasks of computer algebra. In algebraic geometry, they define projections of varieties or schemes, in linear algebra

and number theory, they are applied to find minimal polynomials, and in numerous applications, they are used to help solve polynomial systems of equations. Traditionally, elimination ideals have been computed using resultants or using Gröbner bases with respect to elimination term orderings. As these techniques become impractical when the number of indeterminates is large, a new technique, called Z-separating reembeddings, and based on trying to perform elimination by substitution, has been developed in [12-14, 17]. This technique proceeds as follows.

Let $P = K[x_1, \ldots, x_n]$ be a polynomial ring over a field K, and let I be an ideal in P. Given a tuple of distinct indeterminates $Z = (z_1, \ldots, z_s)$ in $X = (x_1, \ldots, x_n)$, we say that a tuple of polynomials (f_1, \ldots, f_s) in I is Z-separating if there exists a term ordering σ such that $\operatorname{LT}_{\sigma}(f_i) = z_i$ for $i = 1, \ldots, s$. If we find such a tuple of polynomials, we can calculate a tuple of coherently Z-separating polynomials and then the elimination ideal $I \cap K[X \setminus Z]$ by potentially less costly interreductions and substitutions. Notice that in contrast to [12-14] we do not assume $I \subseteq \langle x_1, \ldots, x_n \rangle$ here, as this hypothesis is not needed for our algorithms to work.

Fast methods for determining good candidate tuples Z for which a Z-separating tuple of polynomials might exist in I were developed in [13, 14, 17]. Here we study the second phase of the method, namely to determine quickly whether the ideal I contains a Z-separating tuple of polynomials for a given Z. To answer this question in full generality, one would have to compute a Gröbner basis of I, the very task that we deem infeasible and want to avoid. Hence we will content ourselves with an algorithm which checks very quickly if the vector space spanned by the given generators of I, or a somewhat larger, easily calculated vector space, contains a Z-separating tuple. These algorithms are fast enough to allow us to scan many hundred candidate tuples Z in a matter of seconds, and are therefore suitable for applications involving larger polynomial systems. For instance, in the final section we apply them to speed up algebraic attacks on the AES-128 cryptosystem.

Now let us describe the contents of the paper in more detail. In Section 2 we recall the basic definitions of separating and coherently separating tuples of polynomials, and of separating re-embeddings. Then, in Section 3, we introduce the main algorithm of this paper (see Algorithm 3.2). It allows us to discover very efficiently when a given polynomial ideal contains a Z-separating tuple. However, it is allowed to fail in certain complicated cases. The idea of this algorithm is to construct a weight vector $W \in \mathbb{N}^n$ such that, for every term ordering σ which is compatible with the grading on P given by W, there exists a Z-separating tuple of polynomials (f_1, \ldots, f_s) in I with $\mathrm{LT}_{\sigma}(f_i) = z_i$ for $i = 1, \ldots, s$. Moreover, in Proposition 3.3, we prove that this algorithm succeeds exactly when the vector space spanned by the initially given system of generators of I contains a Z-separating tuple.

Section 4 introduces an optimization of our main algorithm. Algorithm 4.1 allows us to find Z-separating tuples of polynomials in I in more cases than Algorithm 3.2 at the cost of a small reduction in efficiency. The underlying idea is derived from the border basis algorithm (see [11]). We do not search for a Z-separating tuple in the vector space V spanned by the original generators of I, but in the larger vector space $V^+ = V + x_1V + \cdots + x_nV$. The loss in efficiency is mostly due to the necessity to perform some linear algebra operations (see Remark 4.3).

Now assume that one of these algorithms confirms the existence of a Z-separating tuple in I. It still remains to find an actual such tuple. This task is tackled in Section 5. In case the existence has been shown by Algorithm 3.2, a straightforward base change mechanism allows us to exhibit an actual Z-separating tuple of poylnomials in I (see Algorithm 5.1). If the existence has been shown using Algorithm 4.1, the situation is a bit trickier: in this case we have to mimick some linear algebra operations performed by the algorithm on simplified polynomials using the original generators instead (see Remark 5.3). We also explain how to pass from a Z-separating tuple to a coherently Z-separating tuple (see Remark 5.4).

If we want to apply these algorithms in situations involving Boolean polynomials, it suffices to adjust them slightly. In Section 6 we first provide the necessary background knowledge about canonical representatives, Boolean Gröbner bases, Zseparating and coherently Z-separating tuples of Boolean polynomials, as well as Boolean Z-separating re-embeddings. Based on the result that the canonical representatives of a tuple of Boolean polynomials are Z-separating if and only if the tuple itself is Z-separating (see Proposition 6.3), we show how to find Z-separating tuples of Boolean polynomials (see Proposition 6.4).

Finally, in Section 7, we apply the algorithms in order to improve algebraic attacks on the cryptosystem AES-128. These improvements are a consequence of better algebraic representations of the S-boxes of this cipher (see Example 7.1). They translate to more compact representations of r-round AES-128 in several logic normal forms such as CNF, CNF-XOR or XNF. For one round of AES-128, these better representations translate to meaningful speed-ups of state-of-the-art Boolean solvers (see Example 7.3), and for two rounds of AES, we get faster solutions in the case when we know some key bits (see Example 7.4).

All algorithms are illustrated by non-trival explicit examples. These examples were calculated using implementations of the algorithms in the computer algebra systems ApCoCoA(see [20]), CoCoA(see [1]), and SageMath (see [21]). The source code is available upon request from the authors. The general notation and definitions in this paper follow [15, 16].

2 Separating Tuples and Separating Re-Embeddings

In the following we let K be an arbitrary field, let $P = K[x_1, \ldots, x_n]$, let I be a proper ideal in P, and let R = P/I. Rings of this form are also called *affine K-algebras*, since they are the affine coordinate rings of closed subschemes of \mathbb{A}^n_K . The well-known task of re-embedding those subschemes into lower dimensional affine spaces can be phrased in terms of their coordinate rings as follows (see [12], [13]).

Definition 2.1. Let R = P/I be an affine K-algebra.

- (a) A K-algebra isomorphism $\Phi: R \longrightarrow P'/I'$, where P' is a polynomial ring over K
- and I' is an ideal in P', is called a **re-embedding** of I.
 (b) A re-embedding Φ : R → P'/I' of I is called **optimal**, if every K-algebra isomorphism R → P''/I'' with a polynomial ring P'' over K and an ideal I'' in P'' satisfies $\dim(P'') \ge \dim(P')$.

In a series of papers ([12], [13], [14]) methods were developed for finding good re-embeddings using the following technique. Let $X = (x_1, \ldots, x_n)$, and let $Z = (z_1, \ldots, z_s)$ be a tuple of pairwise distinct indeterminates in X.

Definition 2.2. Let I be a proper ideal in P.

- (a) The tuple Z is called a **separating tuple of indeterminates** for I, if there exists a term ordering σ and a tuple of polynomials $F = (f_1, \ldots, f_s)$ with $f_i \in I$ such that $LT_{\sigma}(f_i) = z_i$ for $i = 1, \ldots, s$.
- (b) In this case we say that F is a Z-separating tuple in I.
- (c) If, additionally, we have $f_i = z_i h_i$ with $h_i \in K[X \setminus Z]$ for i = 1, ..., s, then F is called a **coherently** Z-separating tuple in I.

Notice that, in contrast to [12], [13], and [14], we do not require $I \subseteq \mathfrak{M} = \langle x_1, \ldots, x_n \rangle$ here. Also note that the condition $\mathrm{LT}_{\sigma}(f_i) = z_i$ in this definition implies that no other term in $\mathrm{Supp}(f_i)$ is divisible by z_i . When we consider the subideal $J = \langle f_1, \ldots, f_s \rangle$ of I, the tuple F is a minimal monic σ -Gröbner basis for J. After interreducing the polynomials in F, we obtain the reduced σ -Gröbner basis $\widetilde{F} = (\widetilde{f}_1, \ldots, \widetilde{f}_s)$ of J, which is then a coherently Z-separating tuple.

Given a coherently Z-separating tuple in I, we can define a re-embedding of I as in the following proposition (cf. [12], [13]). Here we denote by $\operatorname{Lin}_{\mathfrak{M}}(I)$ the ideal $\langle \operatorname{Lin}(f) | f \in I \rangle$, where $\operatorname{Lin}(f)$ denotes the *linear part* of a polynomial $f \in I$, i.e., its homogeneous component of degree 1. Note that if I contains a polynomial with a non-zero constant term, then $\operatorname{Lin}_{\mathfrak{M}}(I) = \langle x_1, \ldots, x_n \rangle_K$. If I is contained in \mathfrak{M} then the ideal $\operatorname{Lin}_{\mathfrak{M}}(I)$ is generated by the linear parts of a system of generators of I (see [12], Prop. 1.9).

Proposition 2.3. (Z-Separating Re-Embeddings)

Let Z be a tuple of distinct indeterminate in X, let $Y = X \setminus Z$, let I be a proper ideal in P, and let $\tilde{F} = (\tilde{f}_1, \ldots, \tilde{f}_s)$ be a coherently Z-separating tuple of polynomials in I. For $i = 1, \ldots, s$, write $\tilde{f}_i = z_i - \tilde{h}_i$ with $\tilde{h}_i \in P$.

(a) The K-algebra homomorphism $\varphi : P \longrightarrow K[Y]$ defined by $z_i \mapsto \tilde{h}_i$ for $i = 1, \ldots, s$ and $\varphi(x_i) = x_i$ for $x_i \notin Z$ induces a K-algebra isomorphism

$$\Phi: P/I \longrightarrow K[Y]/(I \cap K[Y])$$

which is called a Z-separating re-embedding of I.

(b) Suppose that $I \subseteq \mathfrak{M} = \langle x_1, \ldots, x_s \rangle$ and $s = \dim_K(\operatorname{Lin}_{\mathfrak{M}}(I))$. Then we have $\operatorname{edim}(P/I) = n - s$, i.e., the Z-separating re-embedding of I is an optimal re-embedding.

In view of this proposition, the task of finding good, or even optimal, re-embeddings of a polynomial ideal can be divided into two steps:

(1) Find candidate tuples Z such that the ideal I has a reasonable chance of containing a Z-separating tuple of polynomials.

(2) Given a candidate tuple Z, check whether I does indeed contain a Z-separating tuple.

To perform the first step, for instance the following approaches have been proposed.

Remark 2.4. Let I be a proper ideal in P. Suppose we want to find a tuple Z of distinct indeterminates in X which is a separating tuple for I.

(a) If we can find a term ordering σ such that $Z \subseteq LT_{\sigma}(I)$ then Z is a separating tuple of indeterminates for I. For instance, we can take all indeterminates contained in $LT_{\sigma}(I)$.

Unfortunately, checking all leading term ideals $LT_{\sigma}(I)$ involves the computation of the Gröbner fan of I. This is known to be an extremely hard computation in general, as the Gröbner fan tends to contain many elements. One improvement can be to compute a *restricted Gröbner fan* (see [13]).

(b) If the ideal I is contained in $\mathfrak{M} = \langle x_1, \ldots, x_n \rangle$, it turns out that it is sufficient to compute the Gröbner fan of the linear part $\operatorname{Lin}_{\mathfrak{M}}(I)$ of I (see [14], Prop. 2.6). This task can be reduced to finding the minors of a matrix (see [14], Thm. 3.5 and Alg. 4.1). In general, this may still be quite demanding, but for instance when $\operatorname{Lin}_{\mathfrak{M}}(I)$ is a binomial linear ideal, it can be simplified further (see [14], Alg. 5.7).

In view of this remark we concentrate in the following on step (2) of the above procedure. In other words, we assume that we are given a tuple Z of indeterminates and want to check whether it is separating for I. Even if an ideal contains a Z-separating tuple of indeterminates, this may not be apparent by looking at its generators (see for instance [14], Example 2.4). In general, checking whether a given tuple Z of indeterminates is a separating tuple of indeterminates for I may require us to compute a Gröbner basis and thus be infeasible for large examples.

3 Checking Separating Tuples of Indeterminates

As before, we let I be a proper ideal in $P = K[x_1, \ldots, x_n]$, we let $X = (x_1, \ldots, x_n)$, and we let $Z = (z_1, \ldots, z_s)$ be a tuple of distinct indeterminates in X. Our goal in this section is to find fast methods for determining whether Z is a separating tuple of indeterminates for I, i.e., whether I contains a Z-separating tuple of polynomials. Since the calculation of a Gröbner basis of I with respect to an elimination term ordering for Z is, in general, too expensive, we construct procedures which are allowed to fail. We start by introducing the following algorithm, which is a part of Algorithm 3.2, the main algorithm of this paper.

Algorithm 3.1. (Linear Interreduction)

Let $Z = (z_1, \ldots, z_s)$ be a tuple of distinct indeterminates in X, and let σ be the lexicographic term ordering on P. Moreover, let $g_1, \ldots, g_r \in P$ such that $\dim_K(\langle \operatorname{Lin}(g_1), \ldots, \operatorname{Lin}(g_r) \rangle_K) = s$ and such that every term in the union of the supports of g_1, \ldots, g_r is divisible by at least one of the indeterminates in Z. Consider the following sequence of instructions.

- 1: Compute $T = (z_1, \ldots, z_s, t_1, \ldots, t_m)$ where $\{t_1, \ldots, t_m\}$ is the combined support of g_1, \ldots, g_r without z_1, \ldots, z_s , ordered such that $t_1 >_{\sigma} \cdots >_{\sigma} t_m$.
- 2: Form the coefficient matrix $M \in \operatorname{Mat}_{r,m+s}(K)$ of g_1, \ldots, g_r with respect to the terms in T.
- 3: Compute the matrix $N \in \operatorname{Mat}_{r',r}(K)$ such that $N \cdot M \in \operatorname{Mat}_{r',m+s}(K)$ is in reduced row echelon form and has no zero rows.
- 4: Compute $(g'_1, \ldots, g'_{r'})^{\operatorname{tr}} = N \cdot (g_1, \ldots, g_r)^{\operatorname{tr}}$ and return $g'_1, \ldots, g'_{r'}$.

This is an algorithm which computes a K-basis $g'_1, \ldots, g'_{r'}$ of $\langle g_1, \ldots, g_r \rangle_K$ such that the following conditions hold.

- (a) For i = 1, ..., s, the polynomial g'_i is of the form $g'_i = z_i h_i$, where $h_i \in P_{\geq 2}$.
- (b) Let u = r' s. For j = 1, ..., u, we get $g'_{s+j} = q_j \in P_{\geq 2}$ such that the polynomials $q_1, ..., q_u$ have distinct leading terms with respect to σ .
- (c) The polynomials q_1, \ldots, q_u form a K-basis of $\langle g_1, \ldots, g_r \rangle_K \cap P_{\geq 2}$.
- (d) For every $i \in \{1, \ldots, s\}$, we have $z_i \in \langle g_1, \ldots, g_r \rangle_K$ if and only if $g'_i = z_i$.

Proof. Let $U = \langle g_1, \ldots, g_r \rangle_K$. First observe that $N \cdot M$ is the coefficient matrix of $g'_1, \ldots, g'_{r'}$ with respect to the terms in T. Since the rows of $N \cdot M$ form a K-basis of the row space of M, the polynomials $g'_1, \ldots, g'_{r'}$ form a K-basis of U by the isomorphism between the row space of M and U.

To prove (a), we first note that $1 \notin \{t_1, \ldots, t_m\}$ since 1 is not divisible by any indeterminate in Z. Let $V = \langle \text{Lin}(g_1), \ldots, \text{Lin}(g_r) \rangle_K$. Since $\dim_K(V) = s$ and every term in the combined support is divisible by an indeterminate from Z, we have $V = \langle Z \rangle_K$. This shows that the upper left $s \times s$ submatrix of $N \cdot M$ is the identity matrix and hence $\text{Lin}(g'_i) = z_i$ for $i = 1, \ldots, s$ and we can write $g'_i = z_i - h_i$ where $h_i \in P_{\geq 2}$ since $1 \notin \text{Supp}(h_i)$.

To prove (b), we note that $\operatorname{Lin}(g'_{s+j}) = 0$ and $1 \notin \operatorname{Supp}(g'_{s+j})$ for $j = 1, \ldots, u$. This yields $g'_{s+j} \in P_{\geq 2}$. The remaining claim follows from the observations that the coefficient matrix $N \cdot M$ of $g'_1, \ldots, g'_{r'}$ is in reduced row-echelon form and that the terms t_1, \ldots, t_m are sorted decreasingly with respect to σ .

For the proof of (c), let $q \in U \cap P_{\geq 2}$. By (a), there exist $a_1, \ldots, a_s, b_1, \ldots, b_u \in K$ such that $q = a_1(z_1 - h_1) + \cdots + a_s(z_s - h_s) + b_1q_1 + \cdots + b_uq_u$. Then $0 = \text{Lin}(q) = a_1z_1 + \cdots + a_sz_s$ implies $a_1 = \cdots = a_s = 0$. Hence we get $q \in \langle q_1, \ldots, q_u \rangle_K$. This shows $\langle q_1, \ldots, q_u \rangle_K = U \cap P_{\geq 2}$. By (b), the polynomials q_1, \ldots, q_u are K-linearly independent, and therefore a K-basis of $U \cap P_{\geq 2}$.

Finally, we show (d). Let $i \in \{1, \ldots, s\}$ such that $z_i \in U$. Since $g'_i \in U$, we get $h_i = z_i - g'_i \in U \cap P_{\geq 2}$. Then (c) yields $h_i \in \langle q_1, \ldots, q_u \rangle_K$. Since $N \cdot M$ is in reduced row echelon form, the leading terms of q_1, \ldots, q_u do not appear in the support of h_i . However, since these polynomials have distinct leading terms according to (b), every nonzero element $q \in \langle q_1, \ldots, q_u \rangle_K$ satisfies $\mathrm{LT}_{\sigma}(q) = \mathrm{LT}_{\sigma}(q_j)$ for some j. This shows $h_i = 0$, and hence $g'_i = z_i$. For the other implication, note that if $g'_i = z_i$, then clearly $z_i = g'_i \in U$.

Now we are ready to introduce the main algorithm of this paper. Its basic idea is to try to find a tuple of weights $W = (w_1, \ldots, w_n)$ for the indeterminates in X such that for every term ordering σ compatible with the grading given by W there exist polynomials $f_i \in I$ with $LT_{\sigma}(f_i) = z_i$ for $i = 1, \ldots, s$.

Algorithm 3.2. (Checking Separating Indeterminates)

Let $g_1, \ldots, g_r \in P$, let $I = \langle g_1, \ldots, g_r \rangle$, and let $Z = (z_1, \ldots, z_s)$ be a tuple of distinct indeterminates in X. Consider the following sequence of instructions.

- 1: Set $w_i = 0$ for $i \in \{1, ..., n\}$, $\delta = \max\{\deg(g_1), \ldots, \deg(g_r)\}$, and d = 1.
- 2: Delete in g_1, \ldots, g_r every monomial which is not divisible by an indeterminate in Z and call the result g_1, \ldots, g_r again.
- 3: $if \dim_K((\operatorname{Lin}(g_1),\ldots,\operatorname{Lin}(g_r))_K) < \#Z$ then
- 4: return "Fail".
- 5: end if
- 6: repeat
- 7: Apply Algorithm 3.1 to g_1, \ldots, g_r and Z and call the result g_1, \ldots, g_r again.
- s: Set $\widetilde{Z} = Z \cap \{g_1, \dots, g_r\}.$
- 9: if $\widetilde{Z} = \emptyset$ then
- 10: return "Fail".
- 11: end if
- 12: for all $z_i \in \widetilde{Z}$ do
- 13: Let $k \in \{1, \ldots, n\}$ with $x_k = z_i$ and set $w_k = d$.
- 14: Remove z_i from Z.
- 15: end for
- 16: Delete in g_1, \ldots, g_r every monomial which is not divisible by an indeterminate in Z and call the result g_1, \ldots, g_r again.
- 17: Replace d by $\delta \cdot d + 1$.
- 18: until Z is empty.
- 19: **return** $W = (w_1, \ldots, w_n).$

This is an algorithm which returns either "Fail" or a tuple of non-negative weights $W = (w_1, \ldots, w_n) \in \mathbb{N}^n$. If it returns a tuple W, then there are $f_1, \ldots, f_s \in \langle g_1, \ldots, g_r \rangle_K$ such that (f_1, \ldots, f_s) is a Z-separating tuple of polynomials for I, and such that every term ordering σ which is compatible with the grading given by W satisfies $LT_{\sigma}(f_i) = z_i$ for $i = 1, \ldots, s$.

Proof. For the finiteness of the algorithm, it suffices to verify that the loop in Steps 6-18 terminates. In every iteration of the loop, the algorithm either stops in Step 10 or the number of elements in Z decreases in Step 14. This yields that the loop terminates after finitely many iterations.

Now suppose that we are at the start of an iteration of the loop in Steps 6-18. Let U be the K-vector space generated by the input polynomials and let T be the set of all terms deleted in Step 2 or in an execution of Step 16 in a previous iteration of the loop. We show that the following properties hold at the start of every iteration of the loop.

(1) The polynomials g_1, \ldots, g_r are a valid input for Algorithm 3.1, i.e.,

(1.1) every term in the support of g_1, \ldots, g_r is divisible by an indeterminate from Z, (1.2) $\dim_K((\operatorname{Lin}(g_1), \ldots, \operatorname{Lin}(g_r))_K) = \#Z.$

(2) We have $g_1, \ldots, g_r \in U + \langle T \rangle_K$.

(3) All terms in T have (w_1, \ldots, w_n) -weight smaller than d.

To prove (1.1), we note that every monomial not divisible by an indeterminate from Z was either deleted in Step 2 or in Step 16 in an earlier iteration of the loop, so the claim holds.

Next we show (1.2). In the first iteration, passing the condition in Steps 3-5 guarantees the claim. Assume that it holds at the start of an iteration of the loop that is not the last iteration. Then, after Step 7, Algorithm 3.1.a yields $h_1, \ldots, h_s, q_1, \ldots, q_r \in P_{\geq 2}$ such that $(g_1, \ldots, g_r) = (z_1 - h_1, \ldots, z_s - h_s, q_1, \ldots, q_r)$. Without loss of generality assume that $\widetilde{Z} = \{z_1, \ldots, z_k\}$ for k < s. Then, after the execution of Step 16 and therefore at the start of the next loop, we have $(g_1, \ldots, g_r) =$ $(0, \ldots, 0, z_{k+1} - \tilde{h}_{k+1}, \ldots, z_s - \tilde{h}_s, \tilde{q}_1, \ldots, \tilde{q}_r)$ where $\tilde{h}_{k+1}, \ldots, \tilde{h}_s, \tilde{q}_1, \ldots, \tilde{q}_r \in P_{\geq 2}$. Consequently, $\langle \text{Lin}(g_1), \ldots, \text{Lin}(g_r) \rangle_K = \langle z_{k+1}, \ldots, z_s \rangle_K$ has dimension s - k = #Z.

To show (2), we observe that it holds after the execution of Step 2, and thus at the start of the first iteration of the loop. Moreover, the property is preserved by K-linear interreductions and monomial deletions, which are the only operations performed on g_1, \ldots, g_r during the loop in Steps 6-18. This implies (2).

To prove (3), notice first that, if a term t is added to T at any point of the algorithm, then every later execution of Step 13 only changes the weight of an indeterminate in Z which does not divide t. Hence the weight of t does not change anymore. Moreover, if $t \in T$ is deleted in Step 2, then its weight is 0 which is smaller than d = 1. If it is deleted in Step 16, then every indeterminate dividing t has weight 0 or has been assigned a weight smaller than or equal to d. Hence $\deg(t) \leq \delta$ implies that the weight of t is smaller or equal to $\delta \cdot d < \delta \cdot d + 1$. This shows that, at the start of the next iteration of the loop, the weight of t is smaller than d.

Now assume that the algorithm successfully returned a weight tuple W. Then claim (2) yields that, for every $z_i \in Z$, there was an iteration of the loop in Steps 6-18 where we had $z_i \in U + \langle T \rangle_K$ in Step 8. Thus we can write $z_i = f_i + p_i$, where $f_i \in U$ and $p_i \in \langle T \rangle_K$. By claim (3), every term in T has W-weight smaller than d, while z_i has W-weight d. Hence every term ordering σ compatible with the grading given by W satisfies $\mathrm{LT}_{\sigma}(f_i) = \mathrm{LT}_{\sigma}(z_i - p_i) = z_i$. Notice that $f_i \in U = \langle g_1, \ldots, g_r \rangle_K$.

The next proposition characterizes when Algorithm 3.2 is successful.

Proposition 3.3. Let $g_1, \ldots, g_r \in P$, let $I = \langle g_1, \ldots, g_r \rangle$, and let $Z = (z_1, \ldots, z_s)$ be a tuple of distinct indeterminates in X. Then the following conditions are equivalent.

- (a) Algorithm 3.2 successfully computes a weight tuple W.
- (b) There exists a Z-separating tuple (f_1, \ldots, f_s) of polynomials in I such that $f_1, \ldots, f_s \in \langle g_1, \ldots, g_r \rangle_K$.

Proof. The correctness of Algorithm 3.2 shows that (a) implies (b). Thus we only need to show that (b) implies (a). Assume that there are $f_1, \ldots, f_s \in \langle g_1, \ldots, g_r \rangle_K$ such that (f_1, \ldots, f_s) is a Z-separating tuple of polynomials in I. Let τ be a term ordering on P such that $LT_{\tau}(f_i) = z_i$ for $i = 1, \ldots, s$.

First we show that the algorithm does not terminate in Step 4. For this, let f'_1, \ldots, f'_s be the result of deleting in f_1, \ldots, f_s every term which is not divisible by an indeterminate from Z. Similarly, for $i = 1, \ldots, r$, let g'_i be obtained from g_i in the same way. Then $f'_1, \ldots, f'_s \in \langle g'_1, \ldots, g'_r \rangle_K$. This shows $\operatorname{Lin}(f'_1), \ldots, \operatorname{Lin}(f'_s) \in$

 $(\operatorname{Lin}(g_1),\ldots,\operatorname{Lin}(g_r))_K$. Here $\operatorname{Lin}(f'_1),\ldots,\operatorname{Lin}(f'_s)$ are K-linearly independent, since $\operatorname{LT}_{\tau}(f'_i) = z_i$ yields $\operatorname{LT}_{\tau}(\operatorname{Lin}(f'_i)) = z_i$ for $i = 1,\ldots,s$. Hence the condition in Step 3 is false.

It remains to show that the algorithm does not fail in Step 10. For this, let $\hat{g}_1, \ldots, \hat{g}_{\hat{r}}$ be the original input polynomials of the algorithm, let $\hat{U} = \langle \hat{g}_1, \ldots, \hat{g}_{\hat{r}} \rangle_K$, and write $\hat{S} = \bigcup_{i=1}^{\hat{r}} \operatorname{Supp}(\hat{g}_i)$. At any state of the algorithm, let $U = \langle g_1, \ldots, g_r \rangle$, let $S = \bigcup_{i=1}^{r} \operatorname{Supp}(g_i)$, let $T = \hat{S} \setminus S$ be the set of deleted terms, and let $\varphi : \langle S' \rangle_K \to \langle S \rangle_K$ be the map representing the deletions, i.e., the K-linear map defined by $\varphi(t) = t$ for $t \in S$ and $\varphi(t) = 0$ for $t \in T$.

Let us show that $U = \varphi(\widehat{U})$ at every step of the algorithm. First notice that after the execution of Step 3, we have $r = \hat{r}$ and $g_i = \varphi(\widehat{g}_i)$ for $i = 1, \ldots, r$, so $U = \langle g_1, \ldots, g_r \rangle_K = \langle \varphi(\widehat{g}_1), \ldots, \varphi(\widehat{g}_{\widehat{r}}) \rangle_K = \varphi(\widehat{U})$. This still holds at the start of the first iteration of the loop in Steps 6-18. Since U is invariant under Step 7 in view of Algorithm 3.1, it is only left to show that the claim still holds after the execution of Step 16. For this, assume that we are after an execution of Step 16. Write $\overline{S}, \overline{\varphi}$, and \overline{U} for the values of S, φ , and U before the execution of Step 16, respectively. Write $\psi : \langle \overline{S} \rangle_K \to \langle S \rangle_K$ for the map representing the term deletion in this single execution of Step 16, i.e., the K-linear map with $\psi(t) = t$ for $t \in S$ and $\psi(t) = 0$ for $t \in \overline{S} \setminus S$. Then $\varphi = \psi \circ \overline{\varphi}$ and $U = \psi(\overline{U}) = \psi(\overline{\varphi}(\widehat{U})) = \varphi(\widehat{U})$. Therefore $U = \varphi(\widehat{U})$ holds after the execution of Step 16, and hence at the start of the next iteration of the loop.

Now we show that $\widetilde{Z} \neq \emptyset$ in every execution of Step 8. Assume that we are at the start of an iteration of the loop in Steps 6-18 and let $z_i \in Z$ be the minimal element of Z with respect to τ . Write $f_i = az_i + p_i$ with $a \in K \setminus \{0\}$ and $p_i \in P$ such that az_i is the leading monomial of f_i . Then none of the indeterminates in Z divides any of the terms in p_i , since any term divisible by an element of Z is larger than or equal to z_i with respect to τ . This shows $\operatorname{Supp}(p_i) \cap S = \emptyset$, which in turn implies $\varphi(p_i) = 0$, and hence $z_i = \varphi(f_i) \in \varphi(\widehat{U}) = U = \langle g_1, \ldots, g_r \rangle_K$. Finally, looking at Algorithm 3.1.d, we note that after the execution of Step 7 we have $g_j = z_i$ for some index $j \in \{1, \ldots, r\}$, and hence $g_j \in \widetilde{Z}$ in Step 8.

The following example illustrates Algorithm 3.2 at work. It will also be reconsidered in the next sections.

Example 3.4. Let $P = \mathbb{Q}[x_1, \ldots, x_{11}]$, and let *I* be the ideal in *P* generated by the polynomials $g_1, \ldots, g_9 \in P$ given by

$$\begin{split} g_1 &= x_1 x_4 x_8 x_{11} + x_5 x_6 x_8 + x_5 x_6 x_{10} + x_3 x_6 + x_7 x_8 + x_1 + x_4, \\ g_2 &= -x_1^2 x_3^4 - x_1 x_2 x_3 x_6 x_8 x_{11} + x_1^3 x_{10} + x_3 + x_7 + 1, \\ g_3 &= x_1 x_2 x_3^2 x_8^2 + x_6 x_7^2 x_8 + x_1 x_3 x_8^2 + x_7 x_8 + x_7 x_{10} + x_2 + x_5, \\ g_4 &= -x_1 x_7 x_9 + x_3 x_{11} + x_9, \\ g_5 &= x_1 x_4 x_8 x_{11} + x_1^2 x_6 + x_5 x_7, \\ g_6 &= x_8 x_{10}^2 + x_7 x_9, \\ g_7 &= x_1^2 x_3^4 + x_2^2 x_6^4 + x_1 x_2 x_3^2 x_8^2 + x_1 x_2 x_6^2 x_{10}^2 + x_1 x_2 x_3 x_6 x_8 x_{11}, \end{split}$$

$$g_8 = x_2^6 + x_1 x_6 x_8 - x_3 x_6,$$

$$g_9 = x_1^6 + x_1.$$

We apply Algorithm 3.2 to check whether $Z = (x_4, x_5, x_7)$ is a separating tuple of indeterminates for I.

- (1) In Step 1, we set $w_1 = \cdots = w_{11} = 0$, $\delta = 6$, and d = 1.
- (2) In Step 2, we delete all monomials in g_1, \ldots, g_9 not divisible by x_4, x_5 , or x_7 , and obtain

$$g_1 = x_1 x_4 x_8 x_{11} + x_5 x_6 x_8 + x_5 x_6 x_{10} + x_7 x_8 + x_4, \quad g_2 = x_7, \\ g_3 = x_6 x_7 x_8 + x_7 x_8 + x_7 x_{10} + x_5, \quad g_4 = -x_1 x_7 x_9, \\ g_5 = x_1 x_4 x_8 x_{11} + x_5 x_7, \quad g_6 = x_7 x_9, \quad g_7 = g_8 = g_9 = 0.$$

We have $\dim_{\mathbb{Q}}((\operatorname{Lin}(g_i) \mid i = 1, \ldots, 9)_{\mathbb{Q}}) = 3 = \#Z$, so the algorithm does not stop in Step 2.

(3) The result of applying Algorithm 3.1 in Step 7 is

 $\begin{array}{l} g_1 = x_4 - x_5 x_7 + x_5 x_6 x_8 + x_5 x_6 x_{10} + x_7 x_8, \\ g_2 = x_5 + x_6 x_7 x_8 + x_7 x_8 + x_7 x_{10}, \\ g_3 = x_7, \quad g_4 = x_1 x_4 x_8 x_{11} + x_5 x_7, \\ g_5 = -x_1 x_7 x_9, \quad g_6 = x_7 x_9. \end{array}$

- (4) In Step 8 we obtain $\widetilde{Z} = \{x_7\}.$
- (5) Since $Z \neq \emptyset$, Step 13 sets $w_7 = 1$ and Step 14 sets $Z = (x_4, x_5)$. Deleting all monomials not divisible by x_4 or x_5 in g_1, \ldots, g_6 in Step 16 yields

 $g_1 = x_4 - x_5 x_7 + x_5 x_6 x_8 + x_5 x_6 x_{10}, \quad g_2 = x_5, \quad g_3 = 0,$ $g_4 = x_1 x_4 x_8 x_{11} + x_5 x_7, \quad g_5 = g_6 = 0.$

Step 17 sets d to 7, and the loop repeats.

(6) Algorithm 3.1 returns

$$g_1 = x_4 - x_5 x_7 + x_5 x_6 x_8 + x_5 x_6 x_{10},$$

$$g_2 = x_5, \quad g_3 = x_1 x_4 x_8 x_{11} + x_5 x_7.$$

Step 8 finds $\widetilde{Z} = \{x_5\} \neq \emptyset$, so Step 13 sets $w_5 = 7$, and Step 14 sets $Z = (x_4)$. The deletions in Step 16 yield $g_1 = x_4$, $g_2 = 0$, and $g_3 = x_1 x_4 x_8 x_{11}$. Step 17 sets d = 43 and the loop repeats.

(7) Algorithm 3.1 returns $g_1 = x_4$ and $g_2 = x_1 x_4 x_8 x_{11}$. We have $\widetilde{Z} = \{x_4\}$ in Step 8, so Step 13 sets $w_4 = 43$ and Step 14 sets $Z = \emptyset$. After executing Steps 16 and 17, the loop terminates and the algorithm returns W = (0, 0, 0, 43, 7, 0, 1, 0, 0, 0, 0).

Hence there exists an (x_4, x_5, x_7) -separating tuple of polynomials in *I*.

It is worth noting that the computation of a τ -Gröbner basis of I from g_1, \ldots, g_9 for an elimination ordering τ for Z is quite hard and takes many hours using computer algebra systems such as CoCoA (see [1]), msolve (see [4]), or SageMath (see [21]). Algorithm 3.2, however, is done in less than a second.

The next example shows an application of Algorithm 3.2 to an ideal whose linear part is a binomial linear ideal which was studied in [14, Section 5]. This again shows that our method can work efficiently with rather large examples where any Gröbner basis calculation would be infeasible.

Example 3.5. In the polynomial ring $P = \mathbb{Q}[x_1, \ldots, x_{84}]$, consider the ideal I generated by nonzero entries of the pairwise commutators of the following three matrices

 $\begin{pmatrix} 0 & x_3 & x_5 & 0 & 0 & 0 & x_{12} \\ 0 & x_{15} & x_{17} & 0 & 0 & 0 & x_{24} \\ 0 & x_{27} & x_{29} & 0 & 0 & 0 & x_{36} \\ 1 & x_{39} & x_{41} & 0 & 0 & 0 & x_{48} \\ 0 & x_{51} & x_{53} & 1 & 0 & 0 & x_{60} \\ 0 & x_{63} & x_{65} & 0 & 1 & 0 & x_{72} \\ 0 & x_{75} & x_{77} & 0 & 0 & 1 & x_{84} \end{pmatrix} , \qquad , \qquad \begin{pmatrix} 0 & x_2 & x_4 & x_5 & x_7 & x_9 & x_{11} \\ 0 & x_{14} & x_{16} & x_{17} & x_{19} & x_{21} & x_{23} \\ 1 & x_{26} & x_{28} & x_{29} & x_{31} & x_{33} & x_{35} \\ 0 & x_{38} & x_{40} & x_{41} & x_{43} & x_{45} & x_{47} \\ 0 & x_{50} & x_{52} & x_{53} & x_{55} & x_{57} & x_{59} \\ 0 & x_{64} & x_{65} & x_{67} & x_{69} & x_{71} \\ 0 & x_{74} & x_{76} & x_{77} & x_{79} & x_{81} & x_{83} \end{pmatrix} , \qquad , \qquad ,$

Then a system of generators of I consists of 126 quadratic polynomials without constant, and so $I \subseteq \mathfrak{M}$. The linear part of I has dimension 57 and generates a binomial linear ideal. Using the method mentioned in Remark 2.4.b, we find that the following tuple Z of 57 indeterminates

 $(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10}, x_{11}, x_{12}, x_{18}, x_{19}, x_{20}, x_{21}, x_{22}, x_{23}, x_{30}, \\ x_{31}, x_{32}, x_{33}, x_{34}, x_{35}, x_{37}, x_{38}, x_{40}, x_{42}, x_{43}, x_{44}, x_{45}, x_{46}, x_{47}, x_{49}, x_{50}, x_{52}, x_{54}, x_{55}, \\ x_{56}, x_{57}, x_{58}, x_{59}, x_{61}, x_{62}, x_{64}, x_{66}, x_{67}, x_{68}, x_{69}, x_{70}, x_{71}, x_{78}, x_{79}, x_{80}, x_{81}, x_{82}, x_{83})$

is a possible candidate for a separating tuple of indeterminates for I. When we apply Algorithm 3.2, it shows that Z is indeed a separating tuple of indeterminates for Iand returns the weight tuple

(127, 127, 15, 127, 15, 31, 31, 31, 31, 31, 31, 15, 1, 1, 3, 3, 7, 7, 1, 1, 3, 3, 7, 7, 63, 63, 63, 15, 15, 31, 31, 31, 31, 31, 31, 31, 1, 1, 15, 15, 31, 31, 15, 15, 15, 1, 1, 3, 3, 15, 15, 1, 1, 3, 3, 7, 7).

Notice that we also have $\#Z = 57 = \dim_K(\operatorname{Lin}_{\mathfrak{M}}(I))$. Thus the Z-separating reembedding of I is an optimal re-embedding by [14, Corollary 2.8].

Moreover, the closed subscheme of $\mathbb{A}^{84}_{\mathbb{Q}}$ defined by I is the border basis scheme associated to a certain order ideal (see [3, Example 8]). Using techniques relying on Pommaret bases, it is shown in [3] that this scheme can be embedded into $\mathbb{A}^{42}_{\mathbb{Q}}$, while our method yields an optimal embedding into $\mathbb{A}^{27}_{\mathbb{Q}}$.

4 Optimized Checking of Separating Indeterminates

In this section we discuss an optimization of Algorithm 3.2. Instead of simply interreducing the polynomials g_1, \ldots, g_r linearly using Algorithm 3.1 in Step 7, we first extend them with some polynomials contained in $P_{\geq 2}$. This allows Algorithm 4.1 to succeed in cases where Algorithm 3.2 fails.

Algorithm 4.1. Let $I = \langle g_1, \ldots, g_r \rangle$ be a proper ideal in P, where $g_1, \ldots, g_r \in P \setminus \{0\}$, and let $Z = (z_1, \ldots, z_s)$ be a tuple of distinct indeterminates in X. Consider the following modifications of Algorithm 3.2.

(I) Replace Step 7 by the following sequence of steps.

 $\gamma_{a:}$ Form the set $H = \{x_i g_j \mid x_i \in X \setminus Z \text{ and } j \in \{1, \ldots, r\}\}.$

7b: Let τ be a degree-compatible term ordering. Compute a K-basis $\{h_1, \ldots, h_m\}$ of the space $\langle H \rangle_K$ such that $LT_{\tau}(h_i) \neq LT_{\tau}(h_j)$ for $i \neq j$.

7c: Compute the intersection $\{q_1, \ldots, q_u\} = \{h_1, \ldots, h_m\} \cap P_{\leq \delta}$.

7d: Apply Algorithm 3.1 to Z and $g_1, \ldots, g_r, q_1, \ldots, q_u$. Call the result g_1, \ldots, g_r again.

(II) Moreover, replace Step 17 by the following step:

17: Replace d by $2\delta d + 1$.

Then the resulting sequence of instructions is an algorithm which returns either "Fail" or a tuple of non-negative weights $W = (w_1, \ldots, w_n) \in \mathbb{N}^n$. If it returns a tuple W, then there is a Z-separating tuple of polynomials (f_1, \ldots, f_s) for I such that every term ordering σ which is compatible with the grading given by W satisfies $LT_{\sigma}(f_i) = z_i$ for $i = 1, \ldots, s$.

Proof. The finiteness of the algorithm follows in the same way as in the proof of Algorithm 3.2, so we only show its correctness whenever it returns a weight tuple $W = (w_1, \ldots, w_s)$. Also observe that Step 7b can be executed e.g. by executing Algorithm 3.1 on the input H and the empty tuple.

Let $\delta = \max\{\deg(g_1), \ldots, \deg(g_r)\}\$ as in Step 1 of the algorithm. For $k \in \mathbb{N}$, let $U_k = I_{\leq \delta+k}$ be the K-vector space of all polynomials in I of degree $\leq \delta + k$. At any state of the algorithm, let $W = (w_1, \ldots, w_n)$, and for every $\ell \geq 0$ let V_ℓ be the K-vector space generated by all terms in $K[X \setminus Z]_{\leq \delta+\ell}$ having W-weight $\leq \ell$. Notice that in every state of the algorithm, except between Steps 13 and 14, every indeterminate in $X \setminus Z$ has already been assigned a weight. So, if a term is in V_ℓ , then its W-weight does not change in a later step of the algorithm, even if some weights w_k are adjusted in a later execution of Step 13. Moreover, both sequences $(U_k)_{k\in\mathbb{N}}$ and $(V_\ell)_{\ell\in\mathbb{N}}$ are increasing sequences, i.e., $U_k \subseteq U_{k+1}$ and $V_\ell \subset V_{\ell+1}$ for all $k, \ell \in \mathbb{N}$.

Next we prove the following five claims.

- (1) At the start of the first iteration of the loop in Steps 6-18, we have $g_1, \ldots, g_r \in U_0 + V_0$.
- (2) If $g_1, \ldots, g_r \in U_m + V_{m'}$ for some $m, m' \in \mathbb{N}$ at the start of an iteration of the loop, then we have $g_1, \ldots, g_r \in U_{m+1} + V_{m'+(d-1)/2}$ after the execution of Steps 7a-7d.
- (3) If $g_1, \ldots, g_r \in U_m + V_{m'}$ for some $m, m' \in \mathbb{N}$ after an execution of Steps 7a-7d during an iteration of the loop, then we have $g_1, \ldots, g_r \in U_m + V_{m'} + V_{\delta d}$ after the execution of Step 16.
- (4) For every $k \in \mathbb{N}$ such that there exists a k-th iteration of the loop in Steps 6-18, we have $g_1, \ldots, g_r \in U_{k-1} + V_{(d-1)/2}$ at the start of the k-th iteration.
- (5) For every $k \in \mathbb{N}$ such that there exists a k-th iteration of the loop in Steps 6-18, we have $g_1, \ldots, g_r \in U_k + V_{d-1}$ after the execution of Steps 7a-7d in the k-th iteration of the loop.

To show (1), observe that all terms deleted in Step 2 are elements of $K[X \setminus Z]_{\leq \delta}$ and have W-weight 0. This shows $g_1, \ldots, g_r \in U_0 + V_0$ at the start of the first iteration of the loop in Steps 6-18.

For a proof of (2), assume that we are after the execution of Step 7c, and let $U = \langle H \rangle_K \cap P_{\leq \delta}$. We show the following.

(2.1) The polynomials q_1, \ldots, q_u computed in Step 7c form a K-basis of U.

(2.2) If $g_1, \ldots, g_r \in U_m + V_{m'}$ for some $m, m' \in \mathbb{N}$ before the execution of Step 7a, then we have $U \subseteq U_{m+1} + V_{m'+(d-1)/2}$ after the execution of Step 7c.

To verify (2.1), we note that $q_1, \ldots, q_u \in U$. Let $f \in U$ and suppose without loss of generality that $\operatorname{LT}_{\tau}(h_1) <_{\tau} \cdots <_{\tau} \operatorname{LT}_{\tau}(h_m)$ in Step 7b. Since τ is a degree-compatible term ordering, we have $\{q_1, \ldots, q_u\} = \{h_1, \ldots, h_u\}$. As $f \in \langle h_1, \ldots, h_m \rangle_K$ and the leading terms of h_1, \ldots, h_m are pairwise distinct, there is an index $j \in \{1, \ldots, m\}$ such that $\operatorname{LT}_{\tau}(f) = \operatorname{LT}_{\tau}(h_j)$. It follows that $f \in \langle h_1, \ldots, h_j \rangle_K$, because, for all i > j, we have $\deg(h_i) > \deg(f)$, and hence $\operatorname{LT}_{\tau}(h_i) >_{\tau} \operatorname{LT}_{\tau}(f)$. Moreover, $f \in P_{\leq \delta}$ implies $j \leq u$. This yields $f \in \langle h_1, \ldots, h_u \rangle_K = \langle q_1, \ldots, q_u \rangle_K$, and (2.1) follows.

For a proof of (2.2), it suffices to show that in Step 7a we have $x_i g_j \in U_{m+1} + V_{m'+(d-1)/2}$ for $x_i \in X \setminus Z$ and $j \in \{1, \ldots, r\}$. By the assumption, $g_j \in U_m + V_{m'}$, so there are polynomials $u_j \in U_m$ and $v_j \in V_{m'}$ with $g_j = u_j + v_j$. Every term in the support of v_j has W-weight $\leq m'$. Moreover, the W-weight of x_i was assigned in a previous iteration of the loop, so it is at most $(d-1)/2\delta$, and hence smaller than (d-1)/2. Altogether, the W-weight of every term in the support of $x_i v_j$ is at most m' + (d-1)/2 and we conclude that $x_i v_j \in V_{m'+(d-1)/2}$. It follows that

$$x_i g_j = x_i u_j + x_i v_j \in x_i \cdot U_m + V_{m'+(d-1)/2} \subseteq U_{m+1} + V_{m'+(d-1)/2}.$$

Hence we get $U \subseteq U_{m+1} + V_{m'+(d-1)/2}$. Now observe that after Step 7c we have

 $g_1, \ldots, g_r, q_1, \ldots, q_u \in (U_m + V_{m'}) \cup (U_{m+1} + V_{m'+(d-1)/2}) = U_{m+1} + V_{m'+(d-1)/2},$

so after the execution of Step 7d, we have $g_1, \ldots, g_r \in U_{m+1} + V_{m'+(d-1)/2}$. This shows Claim (2).

To show (3), suppose that $g_1, \ldots, g_r \in U_m + V_{m'}$ after an execution of Step 7d. During the execution of Steps 8-15, only $V_{m'}$ is extended. So, before the execution of Step 16, we still have $g_1, \ldots, g_r \in U_m + V_{m'}$. Let $j \in \{1, \ldots, r\}$, let c_1t_1, \ldots, c_et_e be the monomials deleted from g_j in Step 16, where $c_1, \ldots, c_e \in K$ and $t_1, \ldots, t_e \in \mathbb{T}^n$ are terms, and let $g'_j = g_j - c_1t_1 - \cdots - c_et_e$ be the resulting polynomial obtained from g_j after their deletion. Notice that, for $i = 1, \ldots, e$, we have $t_i \in K[X \setminus Z]$ and $\deg(t_i) \leq \delta$ by (2.1), so all t_1, \ldots, t_e have weight $\leq \delta d$. This shows $t_1, \ldots, t_e \in V_{\delta d}$. Hence $g_j \in U_m + V_{m'}$ implies $g'_j \in U_m + V_{m'} + V_{\delta d}$. Consequently, after the execution of Step 16, we have $g_j \in U_m + V_{m'} + V_{\delta d}$ for all $i \in \{1, \ldots, r\}$.

Next Claim (4) is shown inductively as follows. For k = 1, it holds by (1). Moreover, let $k \in \mathbb{N}$ such that there exist a k-th and a (k + 1)-st iteration of the loop, and suppose that $g_1, \ldots, g_r \in U_{k-1} + V_{(d-1)/2}$ at the start of the k-th iteration. Then (2) shows $g_1, \ldots, g_r \in U_k + V_{(d-1)/2+(d-1)/2} = U_k + V_{d-1}$ after the execution of Steps 7a-7d of the iteration, and (3) implies

$$g_1, \ldots, g_r \in U_k + V_{d-1} + V_{\delta d} = U_k + V_{\delta d}$$

after the execution of Step 16. Therefore, after the adjustment of d in Step 17 and at the start of the (k + 1)-st loop, we have $g_1, \ldots, g_r \in U_k + V_{(d-1)/2}$. This shows (4).

To prove (5), we note that at the start of the k-th iteration of the loop in Steps 6-18, we have $g_1, \ldots, g_r \in U_{k-1} + V_{(d-1)/2}$ by (4). Now (2) says that, after the execution of Steps 7a-7d, we have $g_1, \ldots, g_r \in U_k + V_{(d-1)/2+(d-1)/2} = U_k + V_{d-1}$.

Finally, we combine everything and finish the correctness proof of the algorithm. For every $z_i \in Z$, we eventually find $g_j = z_i$ for some index $j \in \{1, \ldots, r\}$, where we are in Steps 12-15 of the k-th iteration of the loop for some $k \ge 1$. Assume that the weight of z_i is set to d in this iteration. By Claim (5), we have $g_j \in U_k + V_{d-1}$ at this point, so there are $f_j \in U_k$ and $v_j \in V_{d-1}$ with $g_j = f_j + v_j$. Hence $f_j = g_j - v_j = z_i - v_j$, where every term in v_j has W-weight $\le d-1$, and where z_i has weight d. This shows $\mathrm{LT}_{\sigma}(f_j) = z_i$ for every term ordering σ that is compatible with the grading given by W. Consequently, (f_1, \ldots, f_s) is a Z-separating tuple of polynomials.

Algorithm 4.1 sometimes verifies separating tuples of indeterminates of larger lengths than Algorithm 3.2, as the following example shows.

Example 4.2. Let $P = \mathbb{Q}[x_1, \ldots, x_{11}]$ and let *I* be the ideal given in Example 3.4. There we used Algorithm 3.2 to verify that (x_4, x_5, x_7) is a separating tuple for *I*.

Now we apply Algorithm 4.1 to find a larger separating tuple of indeterminates for the ideal I. More precisely, consider $Z = (x_4, x_5, x_7, x_9)$.

After deleting all monomials not divisible by indeterminates from Z in Step 16, we are left with

$$g_1 = x_4 + x_1 x_4 x_8 x_{11} + x_5 x_6 x_8 + x_5 x_6 x_{10} + x_7 x_8, \quad g_2 = x_7$$

$$g_3 = x_5 + x_6 x_7 x_8 + x_7 x_8 + x_7 x_{10}, \quad g_4 = x_9 - x_1 x_7 x_9,$$

$$g_5 = x_1 x_4 x_8 x_{11} + x_5 x_7, \quad g_6 = x_7 x_9, \quad g_7 = g_8 = g_9 = 0.$$

At this point, Steps 7a-7d in Algorithm 4.1 enable us to reduce g_4 against x_1g_6 and get the new polynomial $g_4 = x_9$. Hence we can assign weight 1 to both x_7 and x_9 in the first iteration of the loop.

The remaining part of the algorithm operates similarly as in Example 3.4. However, in this case we find the weight tuple (0, 0, 0, 157, 13, 0, 1, 0, 1, 0, 0) and conclude that the tuple $Z = (x_4, x_5, x_7, x_9)$ is a separating tuple of indeterminates for I.

On the other hand, if we apply Algorithm 3.2 to g_1, \ldots, g_9 and Z, then we cannot eliminate the term $x_1x_7x_9$ in $\text{Supp}(g_4)$, which leads to a "Fail" in Algorithm 3.2.

Let us finish this discussion with a small optimization of Algorithm 4.1.

Remark 4.3. In the setting of Algorithm 4.1, to construct a K-basis $\{q_1, \ldots, q_u\}$ of the K-vector space $\langle H \rangle_K \cap P_{\leq \delta}$, as required in Steps 7b-7c, without introducing a degree-compatible term ordering τ , one can alternatively use the following instructions:

- (1) Write $H' = (h_1, \dots, h_m)$ where $H = \{h_1, \dots, h_m\}$ with m = (n-s)r. Compute the set $\{t_1, \dots, t_e\}$ of all terms of degree $> \delta$ in the support of one of the elements in H.
- (2) Write $h_j = h'_j + \sum_{k=1}^e a_{jk} t_k$, where $a_{jk} \in K$ and $\deg(h'_j) \leq \delta$ for all $j = 1, \ldots, m$, and form the matrix $A = (a_{kj}) \in \operatorname{Mat}_{e,m}(K)$.
- (3) Compute a K-basis $\{v_1, \ldots, v_u\}$ of Ker(A), and let $q_k = H \cdot v_k$ for $k = 1, \ldots, u$.

The advantage of this method is that the matrix A may be smaller than the cofficient matrix of h_1, \ldots, h_m . This may make the computation of this step more efficient than the computation in Step 7b.

5 Finding Z-Separating Tuples of Polynomials

The following algorithm provides a Z-separating tuple of polynomials in the ideal I in the case when Algorithm 3.2 is successful.

Algorithm 5.1. (Computing Z-Separating Polynomials)

Let $g_1, \ldots, g_r \in P$ be generators of a proper ideal I in P, and let Z be a tuple of distinct indeterminates in X such that Algorithm 3.2 returns a weight tuple $W \in \mathbb{N}^n$. Moreover, let σ be a term ordering on P which is compatible with the grading given by W. Consider the following sequence of instructions.

- 1: Compute $T = (t_1, \ldots, t_m)$ where $\{t_1, \ldots, t_m\}$ is the set of terms in the union of the supports of g_1, \ldots, g_r , ordered such that $t_1 >_{\sigma} \cdots >_{\sigma} t_m$.
- 2: Form the coefficient matrix $M \in \operatorname{Mat}_{r,m}(K)$ of g_1, \ldots, g_r with respect to terms in T.
- s: Compute the matrix $N \in \operatorname{Mat}_{r',r}(K)$ such that $N \cdot M \in \operatorname{Mat}_{r',m}(K)$ is in reduced row echelon form and has no zero rows.
- 4: Compute $(g'_1, \ldots, g'_{r'})^{\text{tr}} = N \cdot (g_1, \ldots, g_r)^{\text{tr}}$.
- 5: For $i = 1, \ldots, s$, let $f_i \in \{g'_1, \ldots, g'_{r'}\}$ be the element with $LT_{\sigma}(f_i) = z_i$. Return (f_1, \ldots, f_s) .

This is an algorithm which computes a Z-separating tuple of polynomials (f_1, \ldots, f_s) in I.

Proof. First we note that if the algorithm is executed correctly, then the resulting polynomials f_1, \ldots, f_s form indeed a Z-separating tuple of polynomials in I, as they are by construction elements of I which satisfy $LT_{\sigma}(f_i) = z_i$ for $i = 1, \ldots, s$. It remains to prove that the algorithm is always able to find f_1, \ldots, f_s as claimed in Step 4.

Let $z_i \in Z$. By Proposition 3.3, there is a polynomial $f'_i \in \langle g_1, \ldots, g_r \rangle_K$ with $LT_{\sigma}(f'_i) = z_i$. Since N is an invertible matrix, we also have $f'_i \in \langle g'_1, \ldots, g'_{r'} \rangle_K$, and since $N \cdot M$ is in reduced row echelon form, the leading terms of $g'_1, \ldots, g'_{r'}$ are pairwise distinct. Hence there exists an index $j \in \{1, \ldots, r'\}$ such that $LT_{\sigma}(g'_j) = LT_{\sigma}(f'_i) = z_i$. This shows that the algorithm correctly finds $f_i = g'_j$.

Example 5.2. Let g_1, \ldots, g_9 be the polynomials given in Example 3.4, and let $Z = (x_4, x_5, x_7)$. The application of Algorithm 3.2 in Example 3.4 yielded the weight tuple W = (0, 0, 0, 43, 7, 0, 1, 0, 0, 0, 0). As in [15, Definition 1.4.11], we let σ be the ordering on P represented by the matrix

Then σ is a term ordering on P which is compatible with the grading given by W. An application of Algorithm 5.1 to g_1, \ldots, g_9, Z and σ gives us the Z-separating tuple (f_1, f_2, f_3) of polynomials in I, where

$$f_1 = x_4 - x_5 x_7 + x_5 x_6 x_8 + x_5 x_6 x_{10} + x_7 x_8 - x_1^2 x_6 + x_3 x_6 + x_1,$$

$$f_2 = x_5 + x_6 x_7^2 x_8 + x_7 x_8 + x_7 x_{10} + x_1 x_2 x_3^2 x_8^2 + x_1 x_3 x_8^2 + x_2,$$

$$f_3 = x_7 + x_2^2 x_6^4 + x_1 x_2 x_3^2 x_8^2 + x_1 x_2 x_6^2 x_{10}^2 + x_1^3 x_{10} + x_3 + 1.$$

Notice that in the setting of Algorithm 4.1, it is somewhat trickier to find actual separating polynomials. Let us describe how to do this.

Remark 5.3. If Algorithm 3.2 is successful, then we can apply Algorithm 5.1 in order to find a Z-separating tuple of polynomials. However, if Algorithm 4.1 is successful, then it does not guarantee the existence of a Z-separating tuple of polynomials in the K-vector space $\langle g_1, \ldots, g_r \rangle_K$. Hence we can not apply Algorithm 5.1 to compute such a tuple. In this case, in order to compute a Z-separating tuple of polynomials in I, we can mimic all linear operations executed on g_1, \ldots, g_r on polynomials G_1, \ldots, G_r as follows.

At the start of the algorithm, let G_1, \ldots, G_r be copies of the input polynomials g_1, \ldots, g_r . In every execution of Steps 7a-7c, compute the matrix $A \in \operatorname{Mat}_{(n-s)r,u}(K)$ such that

$$(q_1, \ldots, q_u) = (x_i g_j \mid x_i \in X \setminus Z \text{ and } j \in 1, \ldots, r) \cdot A$$

and let

$$(Q_1,\ldots,Q_u) = (x_iG_j \mid x_i \in X \setminus Z \text{ and } j \in 1,\ldots,r) \cdot A.$$

Mimic the linear transformation performed on $g_1, \ldots, g_r, q_1, \ldots, q_u$ in Step 7d on $G_1, \ldots, G_r, Q_1, \ldots, Q_r$, and re-name the resulting polynomials as G_1, \ldots, G_r again. If $x_k = z_i$ in Step 13, let $f_i = G_k$.

Whenever Algorithm 4.1 successfully returns a weight tuple W, we obtain a Z-separating tuple (f_1, \ldots, f_s) of polynomials in I in this way.

The next remark shows how to obtain a coherently Z-separating tuple from a Z-separating tuple of polynomials.

Remark 5.4. Assume that (f_1, \ldots, f_s) is a Z-separating tuple of polynomials in I obtained by Algorithm 5.1 or Remark 5.3 after a successful run of Algorithm 3.2 or 4.1, respectively. Let W be the corresponding output of the latter algorithm and let σ be a term ordering on P which is compatible with the grading given by W.

If we renumber indices such that $z_1 <_{\sigma} z_2 <_{\sigma} \cdots <_{\sigma} z_s$, then f_i is of the form $f_i = z_i - p_i$, where $p_i \in K[X \setminus Z][z_1, \ldots, z_{i-1}] \leq_{\delta}$ for $i = 1, \ldots, s$. By substituting $z_1 \mapsto p_1$ in f_2 , we obtain a polynomial of the from $f'_2 = z_2 - p'_2$, where $p'_2 \in K[X \setminus Z] \leq_{\delta^2}$. Next we continue by substituting $z_1 \mapsto p_1$ and $z_2 \mapsto p'_2$ in f_3 and so on. In this way, we eventually get polynomials $f'_1 = z_1 - h'_1, \ldots, f'_s = z_s - h'_s$ with $h'_1, \ldots, h'_s \in K[X \setminus Z]$, i.e., a coherently Z-separating tuple (f'_1, \ldots, f'_s) of polynomials in I that defines a Z-separating re-embedding of I.

Let us apply this remark to our running example.

Example 5.5. Let P and I be defined as in Example 3.4, let $Z = (x_4, x_5, x_7, x_9)$, and let $Y = X \setminus Z$. In Example 4.2 we found out that Z is a separating tuple of indeterminates for I. Using Remark 5.3, we compute a Z-separating tuple (f_1, \ldots, f_4) of polynomials in I where

$$\begin{split} f_1 &= x_7 - x_1^2 x_3^4 - x_1 x_2 x_3 x_6 x_8 x_{11} + x_1^3 x_{10} + x_3 + 1, \\ f_2 &= x_9 + x_1 x_8 x_{10}^2 + x_3 x_{11}, \\ f_3 &= x_5 + x_1 x_2 x_3^2 x_8^2 + x_6 x_7^2 x_8 + x_1 x_3 x_8^2 + x_7 x_8 + x_7 x_{10} + x_2, \\ f_4 &= x_4 - x_1^2 x_6 + x_5 x_6 x_8 + x_5 x_6 x_{10} + x_3 x_6 - x_5 x_7 + x_7 x_8 + x_1. \end{split}$$

Write $p_1 = x_7 - f_1$ and $p_2 = x_9 - f_2$. If we substitute x_7 with p_1 and x_9 with p_2 in f_3 , we obtain

$$f_3' = x_5 - x_1 x_2 x_3^2 x_8^2 - x_6 x_8 p_1^2 - x_1 x_3 x_8^2 - x_8 p_1 - x_{10} p_1 + x_2.$$

Next, substituting x_7 , x_9 and x_5 by p_1 , p_2 and $p_3 = x_5 - f'_3$, respectively, into f_4 yields

$$f_4' = x_4 - x_1^2 x_6 + p_3 x_6 x_8 + p_3 x_6 x_{10} + x_3 x_6 - p_3 p_1 + p_1 x_8 + x_1.$$

Altogether, the tuple (f_1, f_2, f'_3, f'_4) is a coherently Z-separating tuple of polynomials in I which defines a Z-separating re-embedding of P/I into $K[Y]/(I \cap K[Y])$. If we now substitute x_4 , x_5 , x_7 and x_9 with p_1 , p_2 , p_3 , and $p_4 = x_4 - f'_4$, respectively, in

the generators g_1, \ldots, g_9 of I, we obtain polynomials $\overline{g}_1, \ldots, \overline{g}_9 \in K[Y]$ which are a generating set of $I \cap K[Y]$.

6 Checking Separating Boolean Indeterminates

Throughout this section we let $\mathbb{F}_2 = \mathbb{Z}/2\mathbb{Z}$, let $P = \mathbb{F}_2[X_1, \ldots, X_n]$ be the polynomial ring over \mathbb{F}_2 in the indeterminates X_1, \ldots, X_n , and let $\mathbb{B}_n = P/\mathbb{I}_n$ be the **ring of Boolean polynomials**. Here $\mathbb{I}_n = \langle X_1^2 - X_1, \ldots, X_n^2 - X_n \rangle$ is the **field ideal** of P. For $i \in \{1, \ldots, n\}$, we denote the residue class of X_i in \mathbb{B}_n by x_i . These residue classes x_1, \ldots, x_n are also referred to as **Boolean indeterminates**.

The goal of this section is to introduce (coherently) separating Boolean polynomials and to devise algorithms for checking whether given tuples of Boolean indeterminates are separating, and in the positive case for computing the corresponding separating Boolean polynomials. For basic facts about Boolean polynomials, we refer the reader to [6] or [9, Sec. 2.10].

It is well-known (see e.g. [6, Theorem 2.1.2]) that every Boolean polynomial $f \in \mathbb{B}_n$ has a unique representative $F \in P$ that is a sum of square-free terms, i.e., terms of the form $X_1^{\alpha_1} \cdots X_n^{\alpha_n}$ with $\alpha_i \in \{0, 1\}$. In this case, the polynomial F is called the **canonical representative** of f. We write deg $(f) = \deg(F)$ and $\operatorname{Supp}(f) =$ $\{t + \mathbb{I}_n \mid t \in \operatorname{Supp}(F)\}$. Moreover, given a term ordering σ on P, we let $\operatorname{LT}_{\sigma}(f) =$ $\operatorname{LT}_{\sigma}(F) + \mathbb{I}_n$ and call it the **leading term** of the Boolean polynomial f with respect to σ . We also write $\operatorname{NR}_{\sigma,G}(f) = \operatorname{NR}_{\sigma,(G_1,\ldots,G_r,X_1^2-X_1,\ldots,X_n^2-X_n)}(F) + \mathbb{I}_n$ for the normal remainder of f with respect to a tuple $G = (g_1,\ldots,g_r) \in \mathbb{B}_n^r$, where G_i is the canonical representative of g_i for $i = 1,\ldots,r$. This allows us also to define a **Boolean** σ -**Gröbner basis** of I as a finite set $G \subseteq I$ that satisfies $\langle \operatorname{LT}_{\sigma}(g) \mid g \in G \rangle = \operatorname{LT}_{\sigma}(I) =$ $\langle \operatorname{LT}_{\sigma}(f) \mid f \in I \rangle$.

In the following, let $X = (x_1, \ldots, x_n)$ be the tuple of all Boolean indeterminates in \mathbb{B}_n , and let $Z = (z_1, \ldots, z_s)$ be a tuple of distinct Boolean indeterminates from X. In analogy to Definition 2.2, the tuple Z is called a **separating tuple of Boolean indeterminates** for a proper ideal I in \mathbb{B}_n , if there exist a term ordering σ and a tuple $F = (f_1, \ldots, f_s)$ of Boolean polynomials in I such that $\mathrm{LT}_{\sigma}(f_i) = z_i$ for $i = 1, \ldots, s$. The tuple F is then called a Z-separating tuple of Boolean polynomials in I. Whenever such a tuple F satisfies the additional condition that none of the terms in the support of any f_i is divisible by z_j for $i, j \in \{1, \ldots, s\}$ with $i \neq j$, we say that Fis a **coherently** Z-separating tuple of Boolean polynomials in I.

The connection between Z-separating and coherently Z-separating tuples of Boolean polynomials is described by the following proposition. The analogous proposition for usual polynomials can be found as Proposition 2.2.d in [13].

Proposition 6.1. Let I be a proper ideal in \mathbb{B}_n , let (f_1, \ldots, f_s) be a Z-separating tuple of Boolean polynomials in I, and let σ be a term ordering on \mathbb{B}_n with $\mathrm{LT}_{\sigma}(f_i) = z_i$ for $i = 1, \ldots, s$. Moreover, write $f_i = z_i + h_i$ with $h_i \in \mathbb{B}_n$, and let $\tilde{h}_i = \mathrm{NR}_{\sigma,(f_1,\ldots,f_s)}(h_i)$ for $i = 1, \ldots, s$. Then the tuple $(z_1 - \tilde{h}_1, \ldots, z_s - \tilde{h}_s)$ is a coherently Z-separating tuple of Boolean polynomials in I.

Proof. For $i \in \{1, \ldots, s\}$, let F_i, Z_i, \tilde{H}_i be the canonical representatives of f_i, z_i, \tilde{h}_i , respectively. By the definition of the normal remainder, the polynomials \tilde{H}_i are fully reduced against F_1, \ldots, F_s . Thus no term in their support is divisible by any of the leading terms of F_1, \ldots, F_s . Since $\mathrm{LT}_{\sigma}(F_j) = Z_j$ for $j = 1, \ldots, s$, this implies that Z_j does not divide any of the terms in the support of \tilde{H}_i , and hence z_j does not divide any of the terms in the support of \tilde{h}_i for $i, j \in \{1, \ldots, s\}$. Consequently, none of the terms in the support of $z_i - \tilde{h}_i$ is divisible by z_j for $i, j \in \{1, \ldots, s\}$ with $i \neq j$. In addition, it is easy to see that $h_i - \tilde{h}_i \in \langle f_1, \ldots, f_s \rangle \subseteq I$, and so $z_i - \tilde{h}_i = f_i - (h_i - \tilde{h}_i) \in I$ for $i = 1, \ldots, s$. Therefore the tuple $(z_1 - \tilde{h}_1, \ldots, z_s - \tilde{h}_s)$ is a coherently Z-separating tuple of Boolean polynomials in I.

In order to construct separating re-embeddings of an ideal I in \mathbb{B}_n , one can apply the result below. Here a **Boolean re-embedding** of I is an \mathbb{F}_2 -algebra isomorphism

$$\Psi: \mathbb{B}_n/I \longrightarrow \mathbb{B}_m/I'$$

where $m \in \mathbb{N}$ and I' is an ideal of \mathbb{B}_m .

Proposition 6.2. (Boolean Z-Separating Re-Embeddings)

In the setting described above, let I be a proper ideal in \mathbb{B}_n , and let $\mathbb{B}_m = \mathbb{F}_2[X \setminus Z]$ with m = n - s. Assume that $F = (f_1, \ldots, f_s)$ is a coherently Z-separating tuple of Boolean polynomials in I with respect to a term ordering σ on \mathbb{B}_n . For $i = 1, \ldots, s$, write $f_i = z_i - h_i$ with $h_i \in \mathbb{B}_m$.

(a) The reduced Boolean σ -Gröbner basis of I is of the form

$$G = \{z_1 - \tilde{h}_1, \dots, z_s - \tilde{h}_s, g_1, \dots, g_r\}$$

where $\tilde{h}_1, \ldots, \tilde{h}_s, g_1, \ldots, g_r \in \mathbb{B}_m$.

(b) The \mathbb{F}_2 -algebra homomorphism $\Phi: \mathbb{B}_n/I \longrightarrow \mathbb{B}_m/(I \cap \mathbb{B}_m)$ given by

$$\Phi(x_i + I) = \begin{cases} x_i + (I \cap \mathbb{B}_m) & \text{for } x_i \notin Z \\ h_j + (I \cap \mathbb{B}_m) & \text{for } x_i = z_j \in Z \end{cases}$$

is an isomorphism of \mathbb{F}_2 -algebras. It is called a **Boolean** Z-separating reembedding of I.

(c) For every elimination ordering τ for Z, we have $\langle Z \rangle \subseteq LT_{\tau}(I)$.

Proof. Let G be the reduced Boolean σ -Gröbner basis of I, and let $h_i = \operatorname{NR}_{\sigma,G}(h_i)$ for $i = 1, \ldots, s$. By assumption, we have $z_i = \operatorname{LT}_{\sigma}(f_i) \in \operatorname{LT}_{\sigma}(I)$ and $h_i \in \mathbb{B}_m$. In particular, since $\operatorname{LT}_{\sigma}(I) \neq \langle 1 \rangle$, we have $z_i - \tilde{h}_i \in G$ and $\tilde{h}_i \in \mathbb{B}_m$ for $i = 1, \ldots, s$. Thus we can write $G = \{z_1 - \tilde{h}_1, \ldots, z_s - \tilde{h}_s, g_1, \ldots, g_r\}$, where $g_1, \ldots, g_r \in \mathbb{B}_n$. Moreover, since G is a reduced Gröbner basis, we know that z_i does not divide any term in $\operatorname{Supp}(g_j)$ for $i \in \{1, \ldots, s\}$ and $j \in \{1, \ldots, r\}$. Hence we get $g_1, \ldots, g_r \in \mathbb{B}_m$, and claim (a) follows.

To show (b), it suffices to prove that $I = \langle G \rangle$ is the kernel of the \mathbb{F}_2 -algebra epimorphism $\varphi : \mathbb{B}_n \longrightarrow \mathbb{B}_m / (I \cap \mathbb{B}_m)$ defined by $x_i \mapsto x_i + (I \cap \mathbb{B}_m)$ for $x_i \notin Z$ and by $x_i \mapsto h_j + (I \cap \mathbb{B}_m)$ for $x_i = z_j \in Z$.

Clearly, $G \subseteq \operatorname{Ker}(\varphi)$, and so $I = \langle G \rangle \subseteq \operatorname{Ker}(\varphi)$. For the other inclusion, let $h \in \operatorname{Ker}(\varphi)$ and write $\tilde{h} = \operatorname{NR}_{\sigma,G}(h)$. Then $\tilde{h} \in \mathbb{B}_m$ and $h = f + \tilde{h}$ for a Boolean polynomial $f \in I$. Since we already know $I \subseteq \operatorname{Ker}(\varphi)$, this implies $\varphi(\tilde{h}) = \varphi(h) = 0$, and so $\tilde{h} \in I \cap \mathbb{B}_m$. Hence we get $h = f + \tilde{h} \in I$.

Finally, claim (c) follows from the observation that $LT_{\tau}(f_i) = LT_{\tau}(z_i - h_i) = z_i \in LT_{\tau}(I)$ for $i = 1, \ldots, s$, because τ is an elimination ordering for Z.

The next proposition provides a connection between Z-separating Boolean polynomials and Z-separating polynomials in P.

Proposition 6.3. Let I' be a proper ideal in P that contains the field ideal \mathbb{I}_n , and let $I = I'/\mathbb{I}_n$. Let $Z = (z_1, \ldots, z_s)$ be a tuple of distinct Boolean indeterminates, and let $Z' = (Z_1, \ldots, Z_s)$ be the tuple of their canonical representatives, i.e., let $Z_i \in \{X_1, \ldots, X_n\}$ and $z_i = Z_i + \mathbb{I}_n$ for $i = 1, \ldots, s$. Furthermore, let $F_1, \ldots, F_s \in P$ and let $f_i = F_i + \mathbb{I}_n$ for $i = 1, \ldots, s$.

- (a) If (F_1, \ldots, F_s) is a Z'-separating tuple of polynomials in I', then (f_1, \ldots, f_s) is a Z-separating tuple of Boolean polynomials in I.
- (b) If (f_1, \ldots, f_s) is a Z-separating tuple in I and F_1, \ldots, F_s are the canonical representatives of f_1, \ldots, f_s , respectively, then (F_1, \ldots, F_s) is a Z'-separating tuple of polynomials in I'.
- (c) The tuple Z' is a separating tuple of indeterminates for I' if and only if the tuple Z is a separating tuple of Boolean indeterminates for I.

Proof. To show (a), let σ be a term ordering on P such that $\operatorname{LT}_{\sigma}(F_i) = Z_i$ for $i = 1, \ldots, s$. Fix $i \in \{1, \ldots, s\}$. Consider $t = x_{i_1} \cdots x_{i_k} \in \operatorname{Supp}(f_i) \setminus \{z_i\}$, where $k = \deg(t)$. Then there are $\alpha_1, \ldots, \alpha_k \in \mathbb{N}_+$ with $T = X_{i_1}^{\alpha_1} \cdots X_{i_k}^{\alpha_k} \in \operatorname{Supp}(F_i)$. It follows from $\operatorname{LT}_{\sigma}(F_i) = Z_i$ that $Z_i = \operatorname{LT}_{\sigma}(F_i) >_{\sigma} T \geq_{\sigma} X_{i_1} \cdots X_{i_k}$, and hence $z_i >_{\sigma} x_{i_1} \cdots x_{i_k} = t$. This implies that $\operatorname{LT}_{\sigma}(f_i) = z_i$. Therefore (f_1, \ldots, f_s) is a Z-separating tuple of Boolean polynomials in I.

To prove (b), let σ be a term ordering such that $LT_{\sigma}(f_i) = z_i$ for $i = 1, \ldots, s$. Then by the definition of leading terms of Boolean polynomials, we have $LT_{\sigma}(F_i) = Z_i$ for all $i = 1, \ldots, s$, and thus the tuple (F_1, \ldots, F_s) is a Z'-separating tuple of polynomials in I'.

By the definition of separating tuples of indeterminates, (c) follows immediately from (a) and (b). $\hfill \Box$

After these preparations, we are now ready to adapt our algorithms to check whether a tuple Z of Boolean indeterminates is a separating tuple for a given ideal in \mathbb{B}_n . The following result plays an important role for that.

Proposition 6.4 (Computation of Z-Separating Boolean Polynomials).

Let I be a proper ideal in \mathbb{B}_n , let $\{g_1, \ldots, g_r\} \subseteq \mathbb{B}_n$ be a system of generators of I, and let $G_1, \ldots, G_r \in P$ be the canonical representatives of g_1, \ldots, g_r . Furthermore, let Z =

 (z_1, \ldots, z_s) be a tuple of distinct Boolean indeterminates, and let $Z' = (Z_1, \ldots, Z_s)$ be the tuple of their canonical representatives.

- (a) If we apply Algorithm 3.2 to the input (G_1, \ldots, G_r) and Z', it successfully returns a weight tuple $W \in \mathbb{N}^n$ if and only if there exist $f_1, \ldots, f_s \in \langle g_1, \ldots, g_r \rangle_{\mathbb{F}_2}$ such that (f_1, \ldots, f_s) is a Z-separating tuple of Boolean polynomials in I.
- (b) In the case of (a), let σ be any term ordering which is compatible with the grading given by W. Then there exist $f'_1, \ldots, f'_s \in \langle g_1, \ldots, g_r \rangle_{\mathbb{F}_2}$ with $\mathrm{LT}_{\sigma}(f'_i) = z_i$ for $i = 1, \ldots, s$.

Proof. Let $U = \langle g_1, \ldots, g_r \rangle_{\mathbb{F}_2}$ and let $U' = \langle G_1, \ldots, G_r \rangle_{\mathbb{F}_2}$. Consider the \mathbb{F}_2 -linear map $\varphi : \mathbb{B}_n \to P$ that maps every Boolean polynomial to its canonical representative. Then the restriction map $\varphi|_U : U \to U'$ is an isomorphism of \mathbb{F}_2 -vector spaces. We first show that there is a Z-separating tuple of Boolean polynomials for I in U if and only if there is a Z'-separating tuple of polynomials for $I' = \langle G_1, \ldots, G_s \rangle + \mathbb{I}_n$ in U'.

Suppose that $(f_1, \ldots, f_s) \in U^s$ is a Z-separating tuple of Boolean polynomials for I. Then $(\varphi(f_1), \ldots, \varphi(f_s)) \in (U')^s$ is a Z'-separating tuple for I' by Proposition 6.3.b. Conversely, let $(F_1, \ldots, F_s) \in (U')^s$ be a Z'-separating tuple of polynomials for I', and let $f_i = F_i + \mathbb{I}_n$ for $i = 1, \ldots, s$. Then (f_1, \ldots, f_s) is a Z-separating tuple for I by Proposition 6.3.a. Since $F_1, \ldots, F_s \in U'$, every term in their support is in the support of one of the polynomials G_1, \ldots, G_r and thus square-free. It follows that every F_i is the canonical representative of f_i and therefore $f_i = \varphi^{-1}(F_i) \in \varphi^{-1}(U') = U$ for $i = 1, \ldots, s$.

In Proposition 3.3 it was shown that Algorithm 3.2 successfully returns a weight tuple W if and only if there exists a Z'-separating tuple of polynomials in U'. By the observation above, this is equivalent to the condition that there exists a Z-separating tuple of polynomials for I in U. This shows (a).

For (b), let σ be a term ordering which is compatible with the grading given by W. By Algorithm 3.2 and Proposition 3.3, there exists a Z'-separating tuple (F'_1, \ldots, F'_s) in U' such that $\operatorname{LT}_{\sigma}(F'_i) = Z_i$ for $i = 1, \ldots, s$. The canonical representative of the element $f'_i = F'_i + \mathbb{I}_n$ is F'_i for $i = 1, \ldots, s$. Therefore we conclude that $\operatorname{LT}_{\sigma}(f'_i) =$ $\operatorname{LT}_{\sigma}(F'_i) + \mathbb{I}_n = z_i$ for $i = 1, \ldots, s$.

Remark 6.5. Let $Z = (z_1, \ldots, z_s)$ be a tuple of distinct Boolean indeterminates in X, let $g_1, \ldots, g_r \in \mathbb{B}_n$, and let Z_i, G_j be the canonical representatives of z_i, g_j , respectively, where $i \in \{1, \ldots, s\}$ and $j \in \{1, \ldots, r\}$. The following methods can be applied to check whether Z is a separating tuple for the ideal $I = \langle g_1, \ldots, g_r \rangle$.

- (a) Apply Algorithm 3.2 to G_1, \ldots, G_r . By Proposition 6.4.a, Algorithm 3.2 successfully returns a weight tuple if and only if Z is a separating tuple of Boolean indeterminates for I and there is a Z-separating tuple of polynomials for I in $\langle g_1, \ldots, g_r \rangle_K$.
- (b) Suppose that Algorithm 3.2 successfully returns a weight tuple $W \in \mathbb{N}^n$, and let σ be a term ordering which is compatible with the grading given by W. Apply Algorithm 5.1 to (G_1, \ldots, G_r) and (Z_1, \ldots, Z_s) to obtain polynomials $F_1, \ldots, F_s \in P$. Then we have $F_1, \ldots, F_s \in \langle G_1, \ldots, G_r \rangle_{\mathbb{F}_2}$ with $\mathrm{LT}_{\sigma}(F_i) = Z_i$ for $i = 1, \ldots, s$.

Since all terms in the supports of G_1, \ldots, G_r are square-free, all terms in the supports of F_1, \ldots, F_s are also square-free. Thus F_i is the canonical representative of $f_i = F_i + \mathbb{I}_n$, and hence $\mathrm{LT}_{\sigma}(f_i) = \mathrm{LT}_{\sigma}(F_i) + \mathbb{I}_n = z_i$. This implies that (f_1, \ldots, f_s) is a Z-separating tuple of Boolean polynomials for I.

- (c) Let $\{H_1, \ldots, H_n\}$ be a set of generators of \mathbb{I}_n , let $I' = \langle G_1, \ldots, G_r \rangle + \mathbb{I}_n$, and let $Z' = (Z_1, \ldots, Z_s)$. According to Proposition 6.3.c, if Algorithm 4.1 applied to G_1, \ldots, G_r , H_1, \ldots, H_n and Z' verifies that Z' is a separating tuple of indeterminates for I', then the tuple Z is a separating tuple of Boolean indeterminates for I. In this case, a Z'-separating tuple (F_1, \ldots, F_s) of polynomials in I' can be computed by using Remark 5.3. Then Proposition 6.3.a yields that (f_1, \ldots, f_s) , where $f_i = F_i + \mathbb{I}_n$ for $i = 1, \ldots, s$, is a Z-separating tuple of polynomials in I.
- (d) Suppose that there exist $j \in \{1, ..., r\}$ and $i \in \{1, ..., s\}$ such that $g_j \in I$ has no constant term and $z_i \in \text{Supp}(g_j)$. Then the linear part of $z_i g_j$ is z_i . In some cases, it may be useful to add those polynomials $z_i g_j$ to the generators of the ideal to find separating tuples of Boolean indeterminates.

7 Application to the Cryptoanalysis of AES-128

The Advanced Encryption Standard (AES) was certified by the U.S. National Institute of Standards and Technology (NIST) in 2001 (see [19]) and is one of the most widely used symmetric encryption ciphers. In this section we apply our algorithms to polynomial equations describing AES-128 with the goal of improving algebraic attacks. We assume that the reader is familiar with the basic definition of AES, for instance as laid out in [19].

Example 7.1 (Polynomial Representation of the AES S-Box).

The only nonlinear part during the encryption process of AES-128 is the application of the Rijndael S-Box. This is a map $s : \mathbb{F}_2^8 \to \mathbb{F}_2^8$ which is detailed in Section 5.1.1 of [19]. We describe s by the vanishing ideal I_S of its graph, i.e., by the ideal I_S in \mathbb{B}_{16} in the Boolean indeterminates $x_1, \ldots, x_8, y_1, \ldots, y_8$ which is the vanishing ideal of the set $S = \{(a, s(a)) \mid a \in \mathbb{F}_2^8\} \subseteq \mathbb{F}_2^{16}$.

- (a) Using a computer algebra system such as CoCoA-5 (see [1]), we can start from the 256 points of S and use them to compute a generating set of I_S . In our case we used the function IdealOfPoints of CoCoA-5, which resulted in a set of 39 quadratic Boolean polynomials. We reference this set as $G_{\rm std}$ for the remaining part of this section.
- (b) Now we apply our implementation of Algorithm 3.2 in the computer algebra system SageMath (see [21]) to the polynomials in G_{std} and every $Z \subseteq \{x_1, \ldots, x_8\}$. The result is that Algorithm 3.2 is successful for every Z with $\#Z \leq 3$. We choose $Z' = (x_1, x_2, x_3)$ and use Algorithm 5.1 to compute a Z'-separating tuple G_{123} . The polynomials in G_{123} are of degree 2.
- (c) Proceeding as in (b), we also apply our implementation of Algorithm 4.1 to G_{std} and every $Z \subseteq \{x_1, \ldots, x_8\}$. It turns out to be successful for every tuple Z with $\#Z \leq 6$. We choose $Z'' = (x_1, x_2, x_3, x_4, x_5, x_6)$ and use Remark 5.3 to compute a Z''-separating tuple G_{1-6} in I_S . The polynomials in G_{1-6} are of degree 3.

- (d) If we strive to find a separating system of generators of I_S consisting of polynomials of lowest possible degree, we can take the three polynomials of degree 2 in G_{123} , three polynomials of degree 3 in G_{1-6} , and two polynomials of degrees 5 and 7, respectively, found using further rounds of Algorithm 4.1 or suitable elimination computations. The resulting tuple G_{1-8} is separating for (x_1, \ldots, x_8) and generates I_S .
- (e) Finally we can compute a lex-Gröbner basis of I_S and get a (x_1, \ldots, x_8) -separating tuple G_{lex} of polynomials of degree 7.

The sets G_{std} , $G_{\text{std}} \cup G_{123}$, $G_{\text{std}} \cup G_{1-6}$, G_{1-8} and G_{lex} generate I_S and allow us to eliminate 0, 3, 6, 8, and 8 Boolean indeterminates, respectively, from I_S . To perform the experiments below, we have to convert these Boolean polynomials to a suitable input for the solvers we consider. While Bosphorus and PolyBoRi work directly with algebraic normal forms and do not require any conversions, the solver Kissat needs CNF (conjunctive normal form) input, the solver CryptoMiniSat expects CNF-XOR, and the solver 2-Xornado uses XNF (xor-or-and normal form) formulas. Let us briefly collect the numbers of logical variables and clauses produced by the conversion tools described in [2].

Table 1 Logic Conversions

Generating set of I_S	# variables/clauses		
	CNF	CNF-XOR	XNF
$G_{\rm std}$	780/8722	256/519	136/399
G_{lex}	1062/6488	843/1424	555/1136
G_{1-8}	542/4061	346/606	213/473

Numbers of logical variables and clauses for the conversions of different systems of generators of I_S .

Of course, it does not make sense to include the numbers for $G_{\text{std}} \cup G_{123}$ and $G_{\text{std}} \cup G_{1-6}$ since these systems of generators properly contain G_{std} . In all three cases, the conversion of G_{lex} is the least efficient one, while most of the numbers for G_{std} and G_{1-8} are similar. However, the number of CNF clauses for G_{std} is large, because the conversion process creates long linear forms which blow up the CNF. An adjusted splitting technique (see [10]) could possibly improve this to some extent.

The question studied below is whether adding the polynomials in G_{123} and G_{1-6} speeds up algebraic attacks on AES-128. Let us describe the mathematical setting of general algebraic attacks first.

Remark 7.2 (Algebraic Attacks on *r*-AES-128).

The original cipher AES-128 is specified to have 10 rounds. For $r \in \{1, ..., 10\}$, we denote by r-AES-128 the version of AES-128 with r instead of 10 rounds, i.e., the number Nr in Figure 5 of [19] is set to r.

Consider the ring of Boolean polynomials \mathbb{B}_n where $n = 128 \cdot (4r + 3)$ whose indeterminates are \mathbf{k}_i , \mathbf{p}_i , \mathbf{c}_i , $\mathbf{rk}_{j,i}$, $\mathbf{so}_{j,i}$, and $\mathbf{rks}_{j,i}$ for $i = 1, \ldots, 128$ and

 $j = 1, \ldots, r$. Here the indeterminates \mathbf{k}_i , \mathbf{p}_i , \mathbf{c}_i , and $\mathbf{r}\mathbf{k}_{j,i}$ represent key, plaintext, ciphertext, and round key bits, respectively. The indeterminates $\mathbf{si}_{j,i}$ and $\mathbf{so}_{j,i}$ are additional variables for the inputs and outputs of the Subbytes transformations (see Section 5.1.1 in [19]) during the encryption process (see Figure 5 in [19]). Lastly, the indeterminates $\mathbf{rk}_{s,i}$ are additional variables for the outputs of the Subbytes transformations during the computation of the round keys (see Figure 11 in [19]).

Let $I_r \subseteq \mathbb{B}_n$ be the ideal modelling *r*-AES-128. In other words, a tuple $(\mathbf{k}, \mathbf{p}, \mathbf{c}, \mathbf{rk}, \mathbf{si}, \mathbf{so}, \mathbf{rks}) \in (\mathbb{F}_2^{128})^3 \times (\mathbb{F}_2^{128r})^4$ is a zero of I_r if and only if \mathbf{p} encrypted with *r*-AES-128 using the key \mathbf{k} yields \mathbf{c} , and during the encryption process, the round keys are \mathbf{rk} , the *S*-box inputs and outputs are \mathbf{si} and \mathbf{so} , respectively, and the outputs of the *S*-boxes during the round key computations are given by \mathbf{rks} . Using a generating set *G* for I_S as in Example 7.1, it is now possible to construct a generating set *H* of I_r that only contains linear polynomials and polynomials from *G* with their indeterminates substituted by the corresponding input and output indeterminates of the Subbytes steps.

The purpose of the next example is to verify that known-plaintext attacks on AES-128 based on finding the zeros of a specialization of the ideal I_r constructed in the preceding remark become easier if we enlarge the sets of generators by G_{123} or G_{1-6} to eliminate some indeterminates in the S-boxes. Currently, the most efficient tools to solve such systems are SAT solvers. In [2] it is described how systems of Boolean polynomials can be converted to XNF (xor-or-and normal form), to XOR-CNF, and to CNF such that they become suitable inputs for current SAT solvers.

Example 7.3 (Attacking One Round of AES-128).

For the following experiments, we generated Boolean polynomial systems H representing 1-AES-128. In order to use them to perform a known-plaintext attack, we proceeded as follows.

- (a) First we constructed a plaintext-ciphertext pair $(p, c) \in \mathbb{F}_2^{256}$ for 1-AES-128, where $p = (p_1, \ldots, p_{128})$ and $c = (c_1, \ldots, c_{128})$.
- (b) Then we specialized various systems of generators H of the ideal I_1 defining the cipher by letting $\mathbf{p}_i \mapsto p_i$ and $\mathbf{c}_i \mapsto c_i$ for $i = 1, \ldots, 128$. Here the various systems H differ by the system of generators G which is used to represent the S-box.
- (c) Next we transformed the specializations to suitable inputs for state-of-the-art Boolean system solvers.
- (d) Finally, we used these solvers to determine the unknown key bits k_i in 100 randomly generated instances and measured their average solving time.

In Table 2, we compare the running times of the XOR-CNF SAT solver Crypto-MiniSat, the CNF SAT solver Kissat (see [7]), and the Boolean polynomial solver Bosphorus (see [8]). For CryptoMiniSat and Kissat, we converted the Boolean polynomial systems H to XOR-CNF and CNF, respectively, using the methods described in [2].

As can be seen from this table, the Boolean polynomial system solver Bosphorus is significantly slower than the SAT solvers and is not effected much by the choice of the generating set of I_S . Both CryptoMiniSat and Kissat profit substantially from

Table 2 Experiments

Generating set of I_S	Average solving time		
	CryptoMiniSat	Kissat	Bosphorus
$G_{\rm std}$	40.23 s	$50.53~{\rm s}$	$244.74 \ s$
$G_{123} \cup G_{\mathrm{std}}$	$35.69 \mathrm{\ s}$	$27.36~{\rm s}$	257.01 s
$G_{1-6} \cup G_{\mathrm{std}}$	$21.47~\mathrm{s}$	$33.57~\mathrm{s}$	$252.78~\mathrm{s}$

Average solving times for 100 polynomial systems coming from known-plaintext attacks on one round of AES-128.

the addition of the separating tuples of polynomials G_{123} and G_{1-6} . More precisely, CryptoMiniSat performs best when using $G_{1-6} \cup G_{\text{std}}$ and Kissat performs best when using $G_{123} \cup G_{\text{std}}$ instead of G_{std} .

These Boolean polynomial systems were also given to the algebraic solver PolyBoRi (see [7]) and to the XNF-solver 2-Xornado (see [2]). However, neither of these solvers terminated on any of the instances in under one hour.

For our next experiment, we considered 2-AES-128 and assumed that a certain amount of bits of the secret key was known, as happens for instance in the case of some sidechannel attacks or for some types of guess-and-determine attacks.

Example 7.4 (Attacking Two Rounds of AES-128).

For this experiment, we created 220 instance of algebraic attacks to 2-AES-128. As above, we substituted randomly generated known plaintext-ciphertext pairs into the Boolean polynomials. Then we constructed 20 instances for each value $k \in \{70, \ldots, 80\}$, where we assume that we know the first k bits of the secret key. These values were also chosen randomly and substituted into the equations.

In the following cactus plot, we draw the solving times that Bosphorus and Cryp-toMiniSat needed for several systems of generators H of 2-AES-128 derived from the stated systems of generators G of the vanishing ideal I_S of the S-boxes.



Fig. 1 Running times of {solver}-{gens} for 220 satisfiable instances related to key-recovery attacks on 2-AES-128 with knowledge of the first $70 \le k \le 80$ key bits.

As one can see, the Boolean polynomial system based on the polynomials G_{1-8} constructed using Algorithms 3.2 and 4.1 is solved significantly faster by CryptoMiniSat (denoted by CMS5) than the system based on the original generators G_{std} . The solver Bosphorus did not terminate on the instances involving G_{lex} and G_{1-8} .

All experiments in this section were performed on an AMD Ryzen 5 7530U CPU with 16 GB of RAM under Manjaro Linux. We used CryptoMiniSat version 5.11.22, Kissat version 4.0.1, and Bosphorus version 3.0.0.

Conclusion

In recent years, a new technique for performing elimination on large-scale examples of polynomial ideals has emerged. It has been called elimination by substitution or the method of Z-separating re-embeddings. In this paper we made another important step in the development of this technique. Previous works discussed fast ways of finding tuples of indeterminates Z which are candidates for this type of elimination computation [13, 14, 17]. Frequently there is a large number of such candidate tuples. Here we provide several algorithms which allow us to check quickly whether a given tuple $Z = (z_1, \ldots, z_s)$ is suitable.

In other words, given an ideal I in a polynomial ring $K[X] = K[x_1, \ldots, x_n]$, we want to test whether $I \cap K[X \setminus Z]$ can be computed using elimination by substitution. For this, we want to check whether polynomials of the form $f_i = z_i - h_i$ exist in I, where h_i is not divisible by any indeterminate in Z. In full generality, this would require the calculation of a Gröbner basis of I which is infeasible in many cases. Therefore we search for the polynomials f_i in suitable subspaces of I.

The algorithms we develop are very fast and allow us to treat examples with dozens, even hundreds of indeterminates. The main idea is not to look for the polynomials f_i directly, but for a tuple of weights W such that, for any term ordering σ compatible with the grading given by W, there exist polynomials $f_i \in I$ with $LT_{\sigma}(f_i) = z_i$. Having found W, the actual polynomials defining the Z-separating re-embedding can then be calculated by simple interreduction.

Applications include examples of ideals for which traditional methods of elimination computation fail, such as ideals defining border basis schemes in Algebraic Geometry. Another area which may benefit from the new techniques is Cryptography. To apply the new method in this case, we developed a variant which is able to perform elimination by substitution for Boolean polynomials. For example, we show how Z-separating Boolean polynomials speed up several types of algebraic attacks on the cipher AES-128.

Acknowledgements

The first author gratefully acknowledges *Cusanuswerk e.V.* for financial support. The third author is supported by the *Vietnam Ministry of Education and Training* under the grant number B2025-DHH-02 and he thanks the University of Passau for its hospitality during part of the preparation of this paper.

References

- [1] J. Abbott, A.M. Bigatti, and L. Robbiano, CoCoA: a system for doing Computations in Commutative Algebra, available at https://cocoa.dima.unige.it.
- B. Andraschko, J. Danner, and M. Kreuzer, SAT solving using XOR-OR-AND normal forms, Math. Comput. Sci. 18 (2024), https://doi.org/10.1007/s11786-024-00594-x
- [3] C. Bertone, and F. Cioffi, The close relation between border and Pommaret marked bases, Collect. Math. 73 (2022), 181–201.
- [4] J. Berthomieu, C. Eder, and M. Safey El Din, msolve: a library for solving polynomial systems, in: *ISSAC '21: Proc. Int. Symp. on Symbolic and Algebraic Computation*, ACM Publications, New York, 2021, pp. 51–58.
- [5] A. Biere and M. Fleury, Gimsatul, IsaSAT and Kissat entering the SAT Competition 2022, in: Proc. SAT Competition 2022 - Solver and Benchmark Descriptions, University of Helsinki, Helsinki, 2022, pp. 10-11.
- [6] M. Brickenstein, Boolean Gröbner Bases: Theory, Algorithms and Applications, Springer-Verlag, Berlin, 2010.
- [7] M. Brickenstein and A. Dreyer, PolyBoRi: A framework for Gröbner-basis computations with Boolean polynomials, J. Symbolic Comput. 44 (2009), 1326–1345.

- [8] D. Choo, M. Soos, K. M. A. Chai, and K. S. Meel, Bosphorus: Bridging ANF and CNF solvers, in: *Proc. Design, Automation, and Test in Europe (DATE)*, Florence 2019, IEEE Xplore, pp. 468-473.
- [9] J. Horáček, Algebraic and logic solving methods for cryptanalysis, dissertation, University of Passau, Passau 2020.
- [10] P. Jovanovic and M. Kreuzer, Algebraic attacks using SAT solvers, Groups Complexity Cryptology 2 (2010), 247–259.
- [11] M. Kreuzer and A. Kehrein, Computing border bases, J. Pure Appl. Algebra 205 (2006), 279–295.
- [12] M. Kreuzer, L.N. Long, and L. Robbiano, Cotangent spaces and separating reembeddings, J. Algebra Appl. 21 (2022), 2250188.
- [13] M. Kreuzer, L.N. Long, and L. Robbiano, Restricted Gröbner fans and reembeddings of affine algebras, Sao Paulo J. Math. Sci. 17 (2023), 242–267.
- [14] M. Kreuzer, L.N. Long, and L. Robbiano, Re-embeddings of affine algebras via Gröbner fans of linear ideals, Beitr. Alg. Geom. 65 (2024), 827–851.
- [15] M. Kreuzer and L. Robbiano, Computational Commutative Algebra 1, Springer-Verlag, Berlin Heidelberg, 2000.
- [16] M. Kreuzer and L. Robbiano, Computational Commutative Algebra 2, Springer-Verlag, Berlin Heidelberg, 2005.
- [17] M. Kreuzer and L. Robbiano, Elimination by substitution, preprint 2024, 26 pages, available at arXiv:2403.06415 [math.AC]
- [18] A. Leventi-Peetz, O. Zendel, W. Lennartz, and K. Weber, CryptoMiniSat switches-optimization for solving cryptographic instances, in: *Proc. Pragmatics* of SAT 2015 and 2018, EPiC Series in Computing 59, EasyChair 2019, pp. 79-93.
- [19] National Institute of Standards and Technology (NIST), FIPS-197: Advanced Encryption Standard (AES), 2001, available at https://csrc.nist.gov/pubs/fips/197/final
- [20] The ApCoCoA Team, ApCoCoA: Applied Computations in Commutative Algebra, available at https://apcocoa.uni-passau.de.
- [21] The Sage Developers, SageMath, the Sage Mathematics Software System, Version 10.2, 2023, available at https://www.sagemath.org