# KP INTEGRABILITY OF NON-PERTURBATIVE DIFFERENTIALS

# A. ALEXANDROV, B. BYCHKOV, P. DUNIN-BARKOWSKI, M. KAZARIAN, AND S. SHADRIN

Abstract. We prove the KP integrability of non-perturbative topological recursion, which can be considered as a formal  $\hbar$ -deformation of the Krichever construction. This property goes back to a 2011 conjecture of Borot and Eynard.

## **CONTENTS**



## 1. INTRODUCTION

- <span id="page-0-1"></span><span id="page-0-0"></span>1.1. Notation. Throughout the paper we use the following notation:
	- Let  $\llbracket n \rrbracket$  be the set of indices  $\{1, \ldots, n\}$ . For each  $I \subset \llbracket n \rrbracket$  let  $z_I$  denote the set of formal variables / points on a Riemann surface / functions (respectively, depending on the context) indexed by  $i \in I$ .
	- The *n*-th symmetric group is denoted by  $\mathfrak{S}_n$ , and its subset of cycles of length *n* is denoted by  $C_n$ . Let  $\det^{\circ}(A_{ij})$  denote the connected determinant of a matrix  $(A_{ij})$ given in the case of an  $n \times n$  matrix by

(1) 
$$
\det^{\circ}(A_{ij}) \coloneqq (-1)^{n-1} \sum_{\sigma \in C_n} A_{i,\sigma(i)}.
$$

#### 2 A. ALEXANDROV, B. BYCHKOV, P. DUNIN-BARKOWSKI, M. KAZARIAN, AND S. SHADRIN

- The operator  $\bigcup_{z\to z'}$  is the operator of restriction of an argument z to the value z', that is,  $\bigcup_{z\to z'} f(z) = f(z')$ . When we write  $\bigcup_{z\to z}$  we mean that the argument of a function or a differential to which this operator applies is set to z, without specifying the notation for the former argument.
- In order to shorten the notation we often omit the summation over an index that runs from 1 to g. For instance,  $\theta_*\eta = \sum_{i=1}^g \theta_{*,i} \eta_i$ ,  $\int_{\mathfrak{B}} \partial_w := \sum_{i=1}^g \int_{\mathfrak{B}_i} \partial_{w_i}$ ,  $\eta(z)\partial_w\coloneqq\sum_{i=1}^g\eta_i(z)\partial_{w_i},\ k\mathcal{T} k=\sum_{i,j=1}^gk_i\mathcal{T}_{ij}k_j,\ bw=\sum_{i=1}^g\widetilde{b_i}w_i,\ \text{etc.}$ • Let  $\mathcal{S}(z)$  denote  $z^{-1}(e^{\frac{z}{2}}-e^{-\frac{z}{2}})$ .

<span id="page-1-0"></span>1.2. KP integrability for a system of differentials. We study KP integrability as a property of a system of differentials, see e.g. [\[KNTY88;](#page-20-0) [Oog90\]](#page-20-1).

Consider a Riemann surface  $\Sigma$ , not necessarily a compact one. Let  $\omega_n$ ,  $n \geq 1$ , be a system of symmetric meromorphic *n*-differentials such that  $\omega_n$  is regular on all diagonals for  $n > 2$ , and  $\omega_2$  has a pole of order two on the diagonal with biresidue 1. Define the bi-halfdifferential

<span id="page-1-1"></span>
$$
(2) \quad \Omega_1(z^+,z^-) = \frac{\sqrt{d\chi(z_1)}\sqrt{d\chi(z_2)}}{\chi(z_1)-\chi(z_2)} \, \exp\left(\sum_{m=1}^{\infty}\frac{1}{m!}\left(\int\limits_{z^-}^{z^+}\right)^m \left(\omega_m(z_{\llbracket n \rrbracket}) - \delta_{m,2}\frac{d\chi(z_1)d\chi(z_2)}{(\chi(z_1)-\chi(z_2))^2}\right)\right),
$$

where  $\chi$  is any local coordinate; the result is actually independent of its choice.

**Definition 1.1.** We say that  $\{\omega_n\}_{n\geq 1}$  is KP integrable, if the following determinantal identity holds:

(3) 
$$
\omega_n(z_{\llbracket n \rrbracket}) = \det^{\circ} \|\Omega_1(z_i, z_j)\|, \quad n \ge 2.
$$

The diagonal entries of the matrix  $\|\Omega_1(z_i, z_j)\|$  are not defined but the connected differential does not involve them for  $n \geq 2$ .

Remark 1.2. The differential  $\omega_1$  can also be recovered from  $\Omega_1$  by

(4) 
$$
\omega_1(z_1) = \bigcup_{z_2 \to z_1} \left( \Omega_1(z_1, z_2) - \frac{\sqrt{d_X(z_1)} \sqrt{d_X(z_2)}}{\chi(z_1) - \chi(z_2)} \right),
$$

for any choice of a local coordinate  $\chi$ , but this equality holds independently of KP integrability.

In fact, a choice of  $\omega_1$  does not affect KP integrability and we could put simply  $\omega_1 = 0$ and to drop the summand with  $m = 1$  in the formula for  $\Omega_1$ . With this modification of  $\Omega_1$ the determinantal formula still holds. Note, however, that it does influence the relation between the actually given  $\omega_1$  and  $\Omega_1$ , so we stick to the form of  $\Omega_1$  as above.

A point  $o \in \Sigma$  is called *regular* for the system of differentials  $\{\omega_n\}$  if  $\omega_n - \delta_{n,2} \frac{dz_1 dz_2}{(z_1 - z_2)}$  $\frac{1}{(z_1-z_2)^2}$ is regular at  $(o, \ldots, o) \in \Sigma^n$  for all  $n \in \mathbb{Z}_{>0}$ , where z is a local coordinate on  $\Sigma$  near o. To any regular point  $o \in \Sigma$  and a local coordinate z at this point we can associate a formal infinite power series  $\tau = \tau_{o,z}(t_1, t_2, \dots)$  defined by the power expansion in the local coordinate z of the following equality

<span id="page-1-2"></span>(5) 
$$
\omega_n(z_{\llbracket n \rrbracket}) = \Big|_{t \to 0} \delta_{z_1} \dots \delta_{z_n} \log(\tau) + \delta_{n,2} \frac{dz_1 dz_2}{(z_1 - z_2)^2}, \qquad \delta_z = \sum_{k=1}^{\infty} z^{k-1} dz \frac{\partial}{\partial t_k}.
$$

Then the KP integrability for a system of differentials  $\{\omega_n\}$  is equivalent to the condition that  $\tau$  is a KP tau function for at least one choice of regular o and z, and then it is automatically a KP tau function for any choice of regular  $o$  and  $z$  [\[ABDBKS23\]](#page-19-1).

While discussing KP integrability for a system of differentials  $\{\omega_n\}$ , it is useful to consider an extended set of  $n|n$  differentials (in fact, half-differentials)  $\Omega_n$  and  $\Omega_n^{\bullet}$  defined as follows. Let B be any symmetric bidifferential with a second order pole on the diagonal with biresidue one. Its choice is irrelevant for the following definition, and one can set  $B = \omega_2$  or simply  $B = \frac{dz_1dz_2}{(z_1-z_2)}$  $\frac{dz_1dz_2}{(z_1-z_2)^2}$  for some local coordinate z. In the global situation, when  $\Sigma$ is compact, one usually takes for  $B$  the Bergman kernel which is normalized by vanishing  $\mathfrak A$ -periods. Consider also the prime form E defined by

(6) 
$$
\frac{1}{E(z^+,z^-)} = \frac{\sqrt{d\chi(z^+)}\sqrt{d\chi(z^-)}}{\chi(z^+)-\chi(z^-)} \exp\left(\frac{1}{2}\int\limits_{z^-}^{z^+} \int\limits_{z^-}^{z^+} \left(B(z_1,z_2) - \frac{d\chi(z_1)d\chi(z_2)}{(\chi(z_1)-\chi(z_2))^2}\right)\right).
$$

In particular, the right hand side is independent of the choice of a local coordinate  $\chi$ . With this notation, the so called disconnected extended  $n|n$  differentials are defined by

<span id="page-2-0"></span>(7) 
$$
\Omega_n^{\bullet}(z_{\llbracket n \rrbracket}^+, z_{\llbracket n \rrbracket}^-) = \prod_{1 \le k < l \le n} \frac{E(z_k^+, z_l^+) E(z_k^-, z_l^-)}{E(z_k^+, z_l^-) E(z_k^-, z_l^+)} \prod_{i=1}^n \frac{1}{E(z_i^+, z_i^-)} \times \exp \left( \sum_{\substack{m_0, \dots, m_n \ge 0 \\ \sum_{i=1}^n m_i = m \ge 1}} \prod_{i=1}^n \frac{1}{m_i!} \left( \int \limits_{z_i^-}^{\cdot} \right)^{m_i} (\omega_m - \delta_{m,2} B) \right).
$$

Their connected counterparts are given by the inclusion-exclusion formula:

<span id="page-2-1"></span>(8) 
$$
\Omega_n(z_{\llbracket n \rrbracket}^+, z_{\llbracket n \rrbracket}^-) := \sum_{\ell=1}^n \frac{(-1)^{\ell-1}}{\ell} \sum_{I_1 \sqcup \dots \sqcup I_{\ell} = \llbracket n \rrbracket} \prod_{j=1}^{\ell} \Omega_{|I_j|}^{\bullet}(z_{I_j}^+, z_{I_j}^-).
$$

One can check that the definition of these symmetric bi-halfdifferentials does not depend on a choice of B. In particular, expressions [\(7\)](#page-2-0)–[\(8\)](#page-2-1) for  $\Omega_1 = \Omega_1^{\bullet}$  are equivalent to [\(2\)](#page-1-1). Notice that the pole on the diagonals  $z_i^+ = z_i^ \bar{i}$  cancel out for the connected  $n|n$  differentials for  $n \neq 1$ , and we have

(9) 
$$
\omega_n(z_{\llbracket n \rrbracket}) = \bigcup_{z_{\llbracket n \rrbracket}^+ \to z_{\llbracket n \rrbracket}} \bigcup_{z_{\llbracket n \rrbracket}^- \to z_{\llbracket n \rrbracket}} \Omega_n(z_{\llbracket n \rrbracket}^+, z_{\llbracket n \rrbracket}^-), \quad n \ge 2.
$$

The following statement is a well known reformulation of KP integrability (see [\[Kri77\]](#page-20-2), or, e.g., [\[EO24\]](#page-19-2)):

**Proposition 1.3.** A system of differentials  $\{\omega_n\}_{n>1}$  is KP integrable if and only if the following determinantal identities for extended differentials hold true:

(10) 
$$
\Omega_n^{\bullet}(z_{\llbracket n \rrbracket}^+, z_{\llbracket n \rrbracket}^-) = \det \| \Omega_1(z_i^+, z_j^-) \|,
$$

(11) 
$$
\Omega_n(z_{\llbracket n \rrbracket}^+, z_{\llbracket n \rrbracket}^-) = \det^{\circ} \|\Omega_1(z_i^+, z_j^-)\|.
$$

The following remarks regarding the definitions of  $\Omega_n^{\bullet}$  and  $\Omega_n$  are in order:

Remark 1.4. Equations [\(2\)](#page-1-1) and [\(7\)](#page-2-0) contain an infinite summation under the exponent. In order to avoid problems with convergence, we either have to interpret this formula as expansion in some suitable local coordinate, or assume some extra expansion of  $\omega_n$  in a formal parameter  $\hbar$  such that the coefficients of expansion in  $\hbar$  are finite expressions.

Remark 1.5. Working with half-differentials we have to choose some square roots of the canonical line bundle on  $\Sigma$  (this concerns both E and  $\Omega_n$ ). In the constructions below in terms of the Θ functions this choice is dictated by the choice of odd theta characteristics.

An example of a KP integrable system of differentials is given by the so-called Krichever differentials [\[Kri77\]](#page-20-2), which we briefly recall below. Another example of a KP integrable systems of differentials is any system of differentials produced by topological recursion on a genus 0 spectral curve (we recall the definition below), see [\[ABDBKS24b;](#page-19-3) [ABDBKS24c\]](#page-19-4).

#### 4 A. ALEXANDROV, B. BYCHKOV, P. DUNIN-BARKOWSKI, M. KAZARIAN, AND S. SHADRIN

Finally, our main result in this paper is that the former two constructions can be mixed into a system of the so-called non-perturbative differentials of topological recursion, proposed and further studied in [\[EM11;](#page-19-5) [BE12;](#page-19-6) [BE15;](#page-19-7) [EO24\]](#page-19-2), which we also prove to be KP integrable, as it was conjectured by Borot and Eynard in [\[BE12\]](#page-19-6).

We conclude this section with the following remark (not concerning KP integrability directly). For a given system of differentials  $\{\omega_n\}$ , along with the differentials  $\Omega_n^{\bullet}$  and  $\Omega_n$  we will use also their certain specializations. Namely, given additionally two functions x and y, we can rewrite the connected  $n|n$  half-differentials in different coordinates near the diagonal using the substitution  $z_i^{\pm} = e^{\pm \frac{uh}{2}\partial_{x_i}} z_i$ , where  $x_i := x(z_i)$ ,  $y_i := y(z_i)$ , simultaneously promoting them to the so-called extended  $n$ -differentials:

<span id="page-3-2"></span>(12) 
$$
\mathbb{W}_n(z_{\llbracket n \rrbracket}, u_{\llbracket n \rrbracket}) \coloneqq \Big( \prod_{i=1}^n e^{u_i \left( S(u_i \hbar \frac{d}{dx_i}) - 1 \right) y_i} \Big) \Big|_{z_i^{\pm} \to e^{\pm \frac{u_i \hbar}{2} \partial_{x_i}} z_i} \Omega_n(z_{\llbracket n \rrbracket}^+, z_{\llbracket n \rrbracket}^-),
$$

<span id="page-3-3"></span>(13) 
$$
\mathcal{W}_n(z, u; z_{\llbracket n \rrbracket}) \coloneqq \mathbb{W}_{n+1}(z, z_{\llbracket n \rrbracket}, u, u_{\llbracket n \rrbracket})\big|_{u_{\llbracket n \rrbracket} = 0}.
$$

Note that  $\omega_n$  is an obvious specialization of those:

(14) 
$$
\omega_n(z_{\llbracket n \rrbracket}) = \bigcup_{u_{\llbracket n \rrbracket} \to 0} (\mathbb{W}_n(z_{\llbracket n \rrbracket}, u_{\llbracket n \rrbracket}) - \delta_{n,1} \frac{dx_1}{u_1 \hbar}).
$$

The point is that there are explicit combinatorial formulas expressing  $\Omega_n$ , W<sub>n</sub>, and W<sub>n</sub> in terms of  $\omega$ -differentials with summation over certain graphs. We review and slightly revisit these formulas in Section [3.](#page-9-0) Here we mention just a formula for  $\mathcal{W}_n$ :

<span id="page-3-1"></span>(15) 
$$
\mathcal{W}_n(z, u; z_{\llbracket n \rrbracket}) = \frac{dx}{u\hbar} e^{\mathcal{T}_0(z, u)} \sum_{\llbracket n \rrbracket = \sqcup_{\alpha} J_{\alpha}} \prod_{\alpha} \mathcal{T}_{|J_{\alpha}|}(z, u; z_{J_{\alpha}}),
$$

where

$$
(16) \quad \mathcal{T}_n(z, u; z_{\llbracket n \rrbracket}) = \sum_{k=1}^{\infty} \frac{1}{k!} \prod_{i=1}^k \Big( \Big( \frac{1}{\tilde{z}_i \to z} u \hbar \mathcal{S}(u \hbar \frac{d}{d\tilde{x}_i}) \frac{1}{d\tilde{x}_i} \Big) \big( \omega_{k+n}(\tilde{z}_{\llbracket k \rrbracket}, z_{\llbracket n \rrbracket}) - \delta_{n,0} \delta_{k,2} \frac{d\tilde{x}_1 d\tilde{x}_2}{(\tilde{x}_1 - \tilde{x}_2)^2} \Big) + \delta_{n,0} u \big( \mathcal{S}(u \hbar \frac{d}{dx}) - 1 \big) y.
$$

<span id="page-3-0"></span>1.3. Topological recursion. Topological recursion of Chekhov–Eynard–Orantin [\[CEO06;](#page-19-8) [EO07\]](#page-19-9) associates a system of meromorphic differentials  $\omega_n^{(g)}$ ,  $g \ge 0$ ,  $n \ge 1$ ,  $2g - 2 + n \ge 0$ , to an input data that consists of a Riemann surface  $\Sigma$  and a finite set of points  $\mathcal{P} \subset \Sigma$ , two meromorphic functions x and y on  $\Sigma$  such that  $\bigcup_{\substack{q \to q}} dx = 0$  and  $\bigcup_{\substack{q \to q}} dy \neq 0$  for each  $q \in \mathcal{P}$  and a bi-differential B with the double pole on the diagonal with bi-residue 1. It has its origin in the computation of the cumulants of the matrix models, and by now it has multiple striking applications in algebraic geometry, enumerative combinatorics, and mathematical physics.

In the local setup [\[DBOSS14\]](#page-19-10) one can assume that  $\Sigma$  is just a union of disjoint discs around points in  $\mathcal{P}$ . In the global setup one assumes that  $\Sigma$  is a compact Riemann surface, and  $B$  is the Bergman kernel. But even in the last case it is still sufficient to have  $dx$ and dy defined in some neighborhood of the points in  $\mathcal{P}$ : the resulting differentials  $\omega_n^{(g)}$ of topological recursion are globally defined meromorphic.

We follow [\[ABDBKS24c\]](#page-19-4), though for the purposes of this paper we restrict ourselves to the case of a compact Riemann surface  $\Sigma$  with a fixed choice of  $\mathfrak A$  and  $\mathfrak B$  cycles and the Bergman kernel B normalized on  $\mathfrak A$  cycles. Let  $dx, dy$  be two meromorphic differentials on Σ. For each point  $q \in \Sigma$  we consider the local expansions of dx and dy in some local

coordinate z,

(17) 
$$
dx = a z^{r-1} (1 + O(z)) dz, \quad dy = b z^{s-1} (1 + O(z)) dz, \quad a, b \neq 0, r, s \in \mathbb{Z},
$$

and we say that the point q is non-special if either  $r = s = 1$  or  $r + s \leq 0$ , and special otherwise.

**Definition 1.6.** The initial data of generalized topological recursion is a tuple  $(\Sigma, dx, dy, \mathcal{P})$ , where  $\mathcal{P} \sqcup \mathcal{P}^{\vee}$  is an arbitrarily chosen split of the set of special points.

We call the points of  $P$  (respectively,  $P^{\vee}$ ) key-points (respectively,  $\vee$ -key-points).

**Definition 1.7.** The differentials of generalized topological recursion  $\omega_n^{(g)}$ ,  $g \ge 0$ ,  $n \ge 1$ ,  $2g - 2 + n \geq 0$ , for the initial data  $(\Sigma, dx, dy, \mathcal{P})$  are defined by  $\omega_2^{(0)} = B$  and for  $2q - 2 + n > 0$  they are given by

(18) 
$$
\omega_n^{(g)}(z, z_{\llbracket n-1 \rrbracket}) = \sum_{q \in \mathcal{P}} \text{res}_{\tilde{z}=q} \left( - \left( \sum_{z \to \tilde{z}} \sum_{r \ge 1} \left( -d \frac{1}{dy} \right)^r [u^r] \mathcal{W}_{n-1}^{(g)}(z, u; z_{\llbracket n-1 \rrbracket}) \right) \int \limits_0^{\tilde{z}} B(\cdot, z),
$$

where the differentials  $W_n = \sum_{g=0}^{\infty} \hbar^{2g-2+n} W_n^{(g)}$  are computed by [\(15\)](#page-3-1) for the system of differentials  $\omega_n = \sum_{(g,n)\neq(0,1)} g_{\geq 0}$  $\hbar^{2g-2+n}\omega_n^{(g)}, x=x(z), y=y(z).$ 

Remark 1.8. This definition might look intimidating without the context that led to it; we refer the reader to [\[ABDBKS24c\]](#page-19-4) for a full discussion. In particular, it indeed recursively defines  $\omega_n^{(g)}$ ,  $2g - 2 + n > 0$ , which turn out to be symmetric meromorphic differentials with the poles only at the key-points.

Yet another viewpoint to topological recursion is provided by the loop equations. They are applied if dy is holomorphic and non-vanishing at each key-point while  $dx$  gets simple zeros at these points. We refer to this situation as the *standard setup* of topological recursion.

**Definition 1.9.** We say that a system of differentials  $\{\omega_n^{(g)}\}, g \ge 0, n \ge 1, 2g-2+n \ge 0,$ satisfy the loop equations at the given point  $q \in \mathcal{P}$  if for any  $k \geq 0$  the following relation holds true:

(19) 
$$
[u^k]e^{uy}\mathcal{W}_n^{(g)}(z,u)\in \Xi_q,
$$

where the space  $\Xi_q$  is spanned by meromorphic differentials of the kind  $\left(d\frac{1}{dx}\right)^j \alpha$  where  $j \geq 0$  and  $\alpha$  is holomorphic.

Remark 1.10. In the standard setup it is sufficient to consider only  $k = 0$  and  $k = 1$  (the so-called linear and quadratic loop equations), see [\[ABDBKS22,](#page-19-11) Section 5]. However, we keep including all  $k \geq 0$  in the parts refereeing to the standard setup throughout the paper as it allows an immediate generalization of the relevant parts to the case when  $dx$ possibly has higher order zeros at key points.

Lemma 1.11 ([\[ABDBKS24c\]](#page-19-4)). In the standard setup the definition of the generalized topological recursion is equivalent to the following:

- $\bullet\ \omega_2^{(0)}=B;$
- for  $2g 2 + n > 0$  the differentials  $\omega_n^{(g)}$  have poles only at the key-points;
- $\{\omega_n^{(g)}\}$  satisfy the loop equations at the key-points;

### 6 A. ALEXANDROV, B. BYCHKOV, P. DUNIN-BARKOWSKI, M. KAZARIAN, AND S. SHADRIN

• they satisfy the projection property:

(20) 
$$
\omega_n(z, z_{\llbracket n-1 \rrbracket}) = \sum_{q \in \mathcal{P}} \operatorname{res}_{\tilde{z}=q} \omega_n(\tilde{z}, z_{\llbracket n-1 \rrbracket}) \int^{\tilde{z}} B(\cdot, z).
$$

In the global setup, that is, if  $\Sigma$  is compact, there is a canonical choice of B dictated by requirement of vanishing its  $\mathfrak A$  periods. For this choice of B, the projection property is also equivalent to the condition that the globally defined differentials  $\omega_n^{(g)}$  have also vanishing A periods.

If we assume that dx and dy are globally defined on a compact Riemann surface  $\Sigma$ , one can trace the effect of the swap of  $x$  and  $y$  in the initial data of topological recursion on the resulting differentials. It is a powerful technique developed in a number of papers [\[BCGFLS23;](#page-19-12) [Hoc24a;](#page-20-3) [Hoc23;](#page-20-4) [ABDBKS22\]](#page-19-11), and its natural development leads to a vast generalization and a revision of the definition of topological recursion, which was developed through a sequence of papers in [\[Hoc24b;](#page-20-5) [ABDBKS24a;](#page-19-13) [ABDBKS24d\]](#page-19-14) and got its final form in [\[ABDBKS24c\]](#page-19-4).

In this paper we use some techniques developed in [\[ABDBKS23;](#page-19-1) [ABDBKS24b;](#page-19-3) [AB-](#page-19-4)DBKS24c, and we refer the reader to the corresponding parts with these papers when appropriate:

- The  $x y$  swap action mentioned above and the fact that it preserves KP integrability of a system of differentials [\[ABDBKS23\]](#page-19-1).
- In the standard setup, there is a deformation formula that captures the effect of the infinitesimal change of  $dy$  near the key-points; it is proved that it preserves the KP integrability [\[ABDBKS24b\]](#page-19-3).
- Compatibility of the generalized topological recursion with particular type of limit behavior of the input data [\[ABDBKS24c\]](#page-19-4).

<span id="page-5-0"></span>1.4. Non-perturbative differentials. The following result was proved in [\[ABDBKS23;](#page-19-1) [ABDBKS24b\]](#page-19-3) for the standard setup of topological recursion and extended in [\[ABD-](#page-19-4)[BKS24c\]](#page-19-4) for generalized topological recursion:

**Proposition 1.12.** The system of differentials  $\omega_n$ ,  $n \geq 1$ , constructed by (generalized) topological recursion is KP integrable if and only if the underlying Riemann surface  $\Sigma$  has genus 0.

For the Riemann surfaces of higher genus one has to deal with the so-called nonperturbative differentials  $\omega_n^{\text{np}}$ ,  $n \geq 1$ , according to the conjecture of [\[BE12\]](#page-19-6). The nonperturbative differentials were initially introduced in [\[EM11\]](#page-19-5) in order to achieve the socalled background independence of the associated partition function. In [\[BE15\]](#page-19-7) they are related to the quantum invariants of knots.

The non-perturbative differentials can be expressed as formal power series in  $\hbar$  of non-topological type,  $\omega_n^{np} = \sum_{d=0}^{\infty} \omega_n^{np, \langle d \rangle}$ , with the leading terms given by the so-called Krichever differentials that are the cornerstones of the Krichever's construction of the finite-zone solutions of the KP equations.

*Remark* 1.13. Since we have this expansion in  $\hbar$ , the adjective "non-perturbative" might be a bit misleading, but it is by now the standard terminology in this area.

The non-perturbative differentials are the main objects of study in this paper. Their definition requires some preparation, so in Section [2](#page-6-1) we briefly summarize the standard facts on Θ functions and recall the construction of the Krichever differentials, and in Section [3.1](#page-9-1) we merge the construction of the Krichever differentials with the input data

of the generalized topological recursion discussed above in order to introduce the nonperturbative differentials and associated  $n|n$  half-differentials.

The main results about the non-perturbative differentials which are the main results of the paper are presented in the rest of Section [3](#page-9-0) and can be summarized as follows:

- In the standard setup, they can be described via loop equations (Theorem [3.7\)](#page-11-2) and a certain deformation of the projective property (Theorem [3.9\)](#page-11-3), which makes them the subject of a suitable generalization of the so-called blobbed topological recursion [\[BS17\]](#page-19-15).
- In the generalized setup, the non-perturbative differentials obey the standard formulas of the  $x - y$  swap (Theorem [3.11\)](#page-11-4) and dy-deformations (Theorem [3.12\)](#page-12-1). These formulas are known to preserve the KP integrability of a system of differentials.
- In the generalized setup, the previous two results (the former of them can be used in the generalized setup by taking limits) allow to prove the KP integrability of the system of non-perturbative differentials (Theorem [3.13\)](#page-12-2). We also present a version of this statement with some extra free parameters (Corollary [3.14\)](#page-12-3) that can be tuned to achieve the background independence and re-introduce the KP times that are otherwise omitted in our approach to KP integrability.

Remark 1.14. Theorem [3.13](#page-12-2) resolves the conjecture of Borot and Eynard in [\[BE12\]](#page-19-6).

<span id="page-6-3"></span>Remark 1.15. Theorems [3.7](#page-11-2) and [3.9](#page-11-3) mean that our results must be revisited in a much broader context of the so-called generalized blobbed topological recursion, a yet to be developed vast generalization and revision of the results of [\[BS17\]](#page-19-15) that should simultaneously feature the advantages of generalized topological recursion and allow non-topological expansions. This will be the subject of a forthcoming work.

<span id="page-6-0"></span>1.5. Acknowledgments. We thank G. Borot and B. Eynard for useful discussions and encouragement to apply our methods to this problem. Substantial part of this project was completed during the stay of the authors at the CGP IBS in Pohang, which we thank for hospitality.

A. A. was supported by the Institute for Basic Science (IBS-R003-D1). B. B. was supported by the ISF Grant 876/20. B. B., P. D.-B., and M. K. were supported by the Russian Science Foundation (grant No. 24-11-00366). S. S. was supported by the Dutch Research Council.

# 2. Necessary facts about Jacobians and Θ functions

<span id="page-6-2"></span><span id="page-6-1"></span>2.1. Θ functions. We fix a smooth algebraic genus q curve  $\Sigma$  and a system of  $\mathfrak{A}, \mathfrak{B}$ cycles on it. Let  $\eta_1, \ldots, \eta_g$  be the basis of holomorphic differentials on  $\Sigma$  normalized by the condition  $\oint_{\mathfrak{A}_i} \eta_j = \delta_{i,j}$  and B is the Bergman kernel normalized on the  $\mathfrak A$  cycles, that is  $\oint_{\mathfrak{A}_i} B = 0$  and  $\frac{1}{2\pi i} \int_{\mathfrak{B}_i} B = \eta_i$ ,  $i = 1, \ldots, g$ . The  $g \times g$  matrix  $\mathcal T$  of  $\mathfrak B$  periods  $\mathcal{T}_{i,j} = \oint_{\mathfrak{B}_i} \eta_j = \frac{1}{2\pi}$  $\frac{1}{2\pi i} \oint_{\mathfrak{B}_i} \oint_{\mathfrak{B}_j} B$  is symmetric and Im $\mathcal T$  is positively definite. The *Jacobian J* is defined as the quotient of  $\mathbb{C}^g$  by the 2g-dimensional integer lattice

(21) 
$$
J = \mathbb{C}^g / \{m + \mathcal{T}n\}, \quad m, n \in \mathbb{Z}^g.
$$

The lattice is chosen by the reason that the following *Abel map* is well defined:

(22) 
$$
\mathcal{A}: \Sigma \to J, \quad p \mapsto \left(\int_{q_0}^p \eta_1, \dots, \int_{q_0}^p \eta_g\right)
$$

with the same contours for all integrals and for some base point  $q_0$  fixed in advance. The Abel map extends to divisors,  $\mathcal{A}(\sum n_i p_i) = \sum n_i \mathcal{A}(p_i)$ , or even to Pic  $\Sigma$ , classes of linear equivalence of divisors: the values of  $A$  on equivalent divisors is the same.

The Riemann Θ function is defined by an explicit formula

(23) 
$$
\Theta : \mathbb{C}^{g} \to \mathbb{C},
$$

$$
\Theta(w) = \Theta(w | \mathcal{T}) = \sum_{k \in \mathbb{Z}^{g}} e^{2\pi i (\frac{1}{2}k \mathcal{T}k + kw)}
$$

The convergency follows from the positive definiteness of  $\text{Im }\mathcal{T}$ . A shift of the argument w by a lattice vector does not preserve Θ but multiplies it by an easily controlled factor. As a consequence, it does not define a function on the Jacobian but rather a section of certain line bundle over J. In particular, for any vector w, the "function" on  $\Sigma$  defined by

.

(24) 
$$
F(p) = \Theta(\mathcal{A}(p) + w)
$$

is holomorphic (has no poles) but multivalued. Nevertheless, the divisor of its zeroes (if F is not identically equal to zero) is well defined.

The function  $\Theta$  is even,  $\Theta(x) = \Theta(-x)$ . It is useful to consider also its odd analogue  $\Theta_*$ ,  $\Theta_*(-x) = -\Theta_*(x)$  defined as follows. Let  $\mu = (\mu_1, \ldots, \mu_g)$  and  $\nu = (\nu_1, \ldots, \nu_g)$  be two vectors (called *theta-characteristics*) with the entries in  $\{0, \frac{1}{2}\}$  $\frac{1}{2}$  such that the following congruence holds:  $4\langle \mu, \nu \rangle \equiv 1 \pmod{2}$ . Then, we set

(25) 
$$
\Theta_*(x) = \Theta_{\left[\begin{matrix} \mu \\ \nu \end{matrix}\right]}(x|\mathcal{T}) = \sum_{k \in \mathbb{Z}^g} e^{2\pi i \left(\frac{1}{2}(k+\mu)\mathcal{T}(k+\mu)+(k+\mu)(x+\nu)\right)}.
$$

This function does depend on a choice of  $\mu$  and  $\nu$ , but it is sufficient to fix one of these choices. In fact many relations also do not depend on this choice. For example, we have the following expression for the Bergman kernel:

(26) 
$$
B(z_1, z_2) = d_1 d_2 \log \Theta_* (\mathcal{A}(z_1 - z_2)),
$$

independently of the freedom in the definition of  $\Theta_*$ .

<span id="page-7-0"></span>2.2. Prime form. Let us expand  $\Theta_*$  at the origin,

(27) 
$$
\Theta_*(x) = \theta_* w + (\text{higher order terms}),
$$

and use the coefficients of linear terms in this expansion to define the holomorphic differential  $d\zeta = \theta_*\eta$ . Also, introduce the bi-half-differential

(28) 
$$
\frac{1}{E(z_1, z_2)} = \frac{\sqrt{d\zeta(z_1)}\sqrt{d\zeta(z_2)}}{\Theta_*(\mathcal{A}(z_1 - z_2))}.
$$

It is well defined in a neighborhood of the diagonal, but its global extension is multivalued. It should be considered as a section of a square root of the canonical bundle on  $\Sigma$  in each its variable, and the choice of the square root depends on the theta-characteristics.

Then, as we have already mentioned above in the Introduction, for any local coordinate, or just a meromorphic function  $\chi$  on  $\Sigma$ , we have:

<span id="page-7-1"></span>(29) 
$$
\frac{1}{E(z_1, z_2)} = \frac{\sqrt{d\chi_1}\sqrt{d\chi_2}}{\chi_1 - \chi_2} e^{\frac{1}{2}\int\limits_{z_2}^{z_1} \int\limits_{z_2}^{z_1} \left(B(\tilde{z}_1, \tilde{z}_2) - \frac{d\tilde{\chi}_1 d\tilde{\chi}_2}{(\tilde{\chi}_1 - \tilde{\chi}_2)^2}\right)}, \quad \chi_i = \chi(z_i).
$$

The integrals on the right hand side are not defined without regularization but it is easy to compute that the result is independent of the function z used for the regularization. This equality implies that  $1/E(z_1, z_2)$  has no singularities outside the diagonal (but is still multivalued with possible branchings along non-contractible loops on  $\Sigma$ ). The function  $\Theta_*(\mathcal{A}(z_1-z_2))$  regarded as a function in  $z_1$  has a zero at  $z_2$  and  $g-1$  more zeroes, denote them by  $r_1, \ldots, r_{q-1}$ . It follows that  $d\zeta$  gets double zeroes at  $r_1, \ldots, r_{q-1}$  and these points do not depend on  $z_2$ .

*Remark* 2.1. Notice the following useful identity for the prime form  $E(z_1, z_2)$  following from [\(29\)](#page-7-1) and appearing in some computations

<span id="page-8-4"></span>(30) 
$$
\exp\left(\int_{z_1^-}^{z_1^+} \int_{z_2^-}^{z_2^+} B\right) = \frac{E(z_1^+, z_2^+) E(z_1^-, z_2^-)}{E(z_1^+, z_2^-) E(z_1^-, z_2^+)}.
$$

<span id="page-8-0"></span>2.3. Krichever differentials. The Krichever construction associates a KP tau function to an arbitrary choice of a point  $q_{\infty} \in \Sigma$  and a local coordinate z at this point. It is given by an explicit formula

<span id="page-8-1"></span>(31) 
$$
\tau(t_1, t_2, \dots) = e^{-\frac{1}{2} \sum_{k,l \geq 1} b_{k,l} t_k t_l} \frac{\Theta_*(\sum_{k=1}^{\infty} t_k U_k + w)}{\Theta_*(w)},
$$

where the vectors  $U_k \in \mathbb{C}^g$  and the constants  $b_{k,l}$  are determined by the following expansions at the point  $q_{\infty}$ :

(32) 
$$
\eta(z) = \sum_{k=1}^{\infty} U_k z^{k-1} dz,
$$

(33) 
$$
B(z_1, z_2) = \frac{dz_1 dz_2}{(z_1 - z_2)^2} - \sum_{k,l \ge 1} b_{k,l} z_1^{k-1} dz_1 z_2^{l-1} dz_2.
$$

**Proposition 2.2** ([\[Kri77\]](#page-20-2)).  $\tau$  is a KP tau function for an arbitrary (generic) choice of the g-dimensional parameter w.

The *n*-point differentials  $\omega_n^{Kr}$  associated with this tau function according to [\(5\)](#page-1-2) extend globally and are given explicitly (see [\[KNTY88\]](#page-20-0)) by

<span id="page-8-2"></span>(34) 
$$
\omega_n^{\mathsf{Kr}} = \left( \prod_{i=1}^n \eta(z_i) \partial_w \right) \log \Theta_*(w | \mathcal{T}) + \delta_{n,2} B.
$$

Indeed, we see from [\(31\)](#page-8-1) that the loop insertion operator  $\delta_z$  acts on  $\log \tau$  by

(35) 
$$
\delta_z = \sum_{k=1}^{\infty} z^{k-1} dz \frac{\partial}{\partial t_k} = \sum_{k=1}^{\infty} z^{k-1} dz U_k \partial_w = \eta(z) \partial_w
$$

with a small correction implied by quadratic terms. This implies  $(34)$ . Similar computations provide also an explicit expression for the  $n|n$  half-differentials: (36)

$$
\Omega_n^{\mathsf{Kr},\bullet}(z^+_{[\![n]\!]},z^-_{[\![n]\!]}) = \prod_{1 \leq i < j \leq n} \frac{E(z^+_i,z^+_j) \, E(z^-_i,z^-_j)}{E(z^+_i,z^-_j) \, E(z^-_i,z^+_j)} \, \prod_{i=1}^n \frac{1}{E(z^+_i,z^-_i)} \frac{\Theta_*(\mathcal{A}(\sum_{i=1}^n (z^+_i - z^-_i)) + w)}{\Theta_*(w)}.
$$

The KP integrability means that these differentials obey determinantal identities

<span id="page-8-3"></span>(37) 
$$
\omega_n^{\text{Kr}}(z_{\llbracket n \rrbracket}^+, z_{\llbracket n \rrbracket}^-) = \det^{\circ}(\Omega_1^{\text{Kr}}(z_i, z_j)), \quad n \ge 2,
$$

(38) 
$$
\Omega_n^{\mathsf{Kr},\bullet}(z^+_{\llbracket n \rrbracket},z^-_{\llbracket n \rrbracket}) = \det(\Omega_1^{\mathsf{Kr}}(z^+_i,z^-_j)).
$$

The differential

(39) 
$$
\Omega_1^{\text{Kr}}(z^+, z^-) = \frac{\Theta_*(\mathcal{A}(z^+ - z^-) + w)}{E(z^+, z^-)\Theta_*(w)}
$$

appearing on the right hand side is known as Szegö kernel.

*Remark* 2.3. Equation [\(37\)](#page-8-3) for  $n = 2$  is a version of the so-called Fay trisecant identity. For example, we have

(40) 
$$
\omega_2^{\text{Kr}}(z_1, z_2) = -\Omega_1^{\text{Kr}}(z_1, z_2)\Omega_1^{\text{Kr}}(z_2, z_1) = \frac{\Theta(\mathcal{A}(z_1 - z_2) + w) \Theta(\mathcal{A}(z_2 - z_1) + w)}{E(z_1, z_2)^2 \Theta(w)^2}.
$$

Remark 2.4. Note that we get not just a single KP integrable system of differentials but rather a family of those parameterized by  $w$  and satisfying

(41) 
$$
\omega_{n+1}^{\mathsf{Kr}}(z, z_{\llbracket n \rrbracket}) - \delta_{n,1} B(z, z_1) = \eta(z) \, \partial_w \left( \omega_n^{\mathsf{Kr}}(z_{\llbracket n \rrbracket}) - \delta_{n,2} B(z_{\llbracket 2 \rrbracket}) \right).
$$

Also note that the freedom in a choice of the theta characteristics entering the definition of  $\Theta_*$  does not affect the KP integrability: a different choice  $\Theta_*$  leads to a shift of w, multiplication of  $\Theta_*$  by a constant, and by a factor whose logarithm is linear in w. All these transformations are obvious KP symmetries. For example, the last one affects the differential  $\omega_1^{Kr}$  only.

## 3. Basic constructions and properties

<span id="page-9-0"></span>The proofs of the statements in this section are relegated to Section [4.](#page-13-0)

<span id="page-9-1"></span>3.1. Definitions of non-perturbative objects. Let  $(\Sigma, dx, dy, \mathcal{P})$  be the initial data of (generalized) topological recursion, with the standard Bergman kernel, and arrange the differentials that it produces into the series

(42) 
$$
\omega_m \coloneqq \sum_{\substack{g \geq 0 \\ 2g-2+m>0}} \hbar^{2g-2+m} \omega_m^{(g)}.
$$

<span id="page-9-3"></span>**Definition 3.1** ([\[BE12\]](#page-19-6)). The non-perturbative disconnected  $n|n$  half-differentials  $\Omega_n^{\text{np},\bullet}$ are defined as

<span id="page-9-2"></span>
$$
(43) \ \ \Omega_n^{\text{np},\bullet}(z_{\llbracket n \rrbracket}^+,z_{\llbracket n \rrbracket}^-):=\prod_{1 \leq k < l \leq n} \frac{E(z_k^+,z_l^+)E(z_k^-,z_l^-)}{E(z_k^+,z_l^-)E(z_k^-,z_l^+)}\prod_{i=1}^n \frac{\exp\left(\frac{z_i^+}{\hbar}\int_{z_i^-}^{\omega_i^+}(u_j^0)\right)}{E(z_i^+,z_i^-)} \times \\ \frac{\exp\left(\sum\limits_{\substack{m_0,\ldots,m_n\geq 0 \\ \sum_{i=0}^n m_i=m\geq 1}}\frac{1}{\hbar}\left(\frac{1}{2\pi\hbar}\oint_{\mathfrak{B}}\partial_w\right)^{m_0}\prod_{i=1}^n\left(\frac{z_i^+}{z_i^-}\right)^{m_i}\omega_m\right)\Theta_*(w+\mathcal{A}\left(\sum\limits_{i=1}^n(z_i^+-z_i^-)\right)|\mathcal{T})}{\exp\left(\sum\limits_{m\geq 1}\frac{1}{m!}\left(\frac{1}{2\pi\hbar}\oint_{\mathfrak{B}}\partial_w\right)^m\omega_m\right)\Theta_*(w|\mathcal{T})}.
$$

The internal structure of this expression is described in the following Lemmas.

**Lemma 3.2.** The differentials  $\Omega_n^{np,\bullet}$  are extended n|n disconnected half-differentials associated with a certain family of symmetric n-differentials  $\omega_n^{np}$  by the following slight variation of formulas of Section [1.2:](#page-1-0)

<span id="page-10-1"></span>
$$
(44) \qquad \Omega_n^{\mathsf{np}}(z^+_{[\![n]\!]}, z^-_{[\![n]\!]}) \coloneqq \sum_{\ell=1}^n \frac{(-1)^{\ell-1}}{\ell} \sum_{I_1 \sqcup \dots \sqcup I_\ell = [\![n]\!]} \prod_{j=1}^\ell \Omega_n^{\mathsf{np}, \bullet}(z^+_{I_j}, z^-_{I_j}).
$$

(45) 
$$
\mathbb{W}_n^{\mathsf{np}}(z_{\llbracket n \rrbracket}, u_{\llbracket n \rrbracket}) := \Big( \prod_{i=1}^n e^{-u_i y_i} \Big) \Big|_{z_i^{\pm} \to e^{\pm \frac{u \hbar}{2} \partial_{x_i}} z_i} \Omega_n^{\mathsf{np}}(z_{\llbracket n \rrbracket}^+, z_{\llbracket n \rrbracket}^-),
$$

<span id="page-10-2"></span>(46) 
$$
\mathcal{W}_n^{\mathsf{np}}(z, u; z_{\llbracket n \rrbracket}) \coloneqq \mathbb{W}_{n+1}^{\mathsf{np}}(z, z_{\llbracket n \rrbracket}, u, u_{\llbracket n \rrbracket}) \Big|_{u_{\llbracket n \rrbracket} = 0},
$$

(47) 
$$
\omega_n^{\mathsf{np}}(z_{\llbracket n \rrbracket}) = \Big|_{u_{\llbracket n \rrbracket} \to 0} \big( \mathbb{W}_n^{\mathsf{np}}(z_{\llbracket n \rrbracket}, u_{\llbracket n \rrbracket}) - \delta_{n,1} \frac{dx_1}{u_1 \hbar} \Big),
$$

and  $\Omega_n^{\text{np},\bullet}$ ,  $n \geq 1$ , can be reconstructed from  $\omega_n^{\text{np}}$ ,  $n \geq 1$ , as

<span id="page-10-0"></span>(48) 
$$
\Omega_n^{\text{np},\bullet}(z_{\llbracket n \rrbracket}^+, z_{\llbracket n \rrbracket}^-) = \prod_{1 \le k < l \le n} \frac{E(z_k^+, z_l^+) E(z_k^-, z_l^-)}{E(z_k^+, z_l^-) E(z_k^-, z_l^+)} \prod_{i=1}^n \frac{\exp\left(\frac{1}{\hbar} \int_z^{z_i^+} \omega_1^{(0)}\right)}{E(z_i^+, z_i^-)} \\ \times \exp\left(\sum_{\substack{m_1, \ldots, m_n \ge 0 \\ \sum_{i=1}^{m_1, \ldots, m_n \ge 0}} \prod_{i=1}^n \frac{1}{m_i!} \left(\int_z^t \right)^{m_i} (\omega_m^{\text{np}} - \delta_{m,2} B)\right).
$$

The corresponding non-perturbative n-differentials  $\omega_n^{\text{np}}$  are formal power series in  $\hbar$ , whose coefficients are finite algebraic expressions in terms of the integrals of  $\omega_m^{(g)}$  and derivatives of  $\log \Theta_*(w|\mathcal{T})$ .

Remark 3.3. Note that the contribution of  $\omega_1^{(0)} = y dx$  is included to [\(43\)](#page-9-2), for the reason of consistency with the definitions in the other sources. This contribution does not affect the KP integrability but might be useful, for instance, for presentation of the loop equations. However, these terms are not included to  $\omega_n$  or  $\omega_n^{\text{np}}$ , according to our convention, in order to avoid appearance of negative powers of  $\hbar$ . By that reason, the relation between  $\Omega_n^{\text{np}}$ and  $\mathbb{W}_n^{\mathsf{np}}$  and  $\mathcal{W}_n^{\mathsf{np}}$  is slightly different from those of [\(12\)](#page-3-2)–[\(13\)](#page-3-3).

<span id="page-10-4"></span>**Lemma 3.4.** The leading term of the expansion of  $\omega_n^{\text{np}}$  in  $\hbar$  is the corresponding Krichever differential,

<span id="page-10-3"></span>(49) 
$$
\omega_n^{\mathsf{np}} = \omega_n^{\mathsf{Kr}} + O(\hbar).
$$

In particular,  $\bigcup_{\hbar \to 0} \omega_n^{\text{np}} - \delta_{n,2} B = \prod_{j=1}^n \eta(z_j) \partial_w \log \Theta_*(w|\mathcal{T})$  is holomorphic and  $\omega_n^{\text{np}} = \omega_n^{\text{Kr}}$ if  $\omega_m^{(g)} = 0, 2g - 2 + m > 0.$ 

The coefficient of any positive power of  $\hbar$  in  $\omega_n^{np}$  is a global symmetric meromorphic differential with the only poles in each variable at the points  $q \in \mathcal{P}$ .

Remark 3.5. Note that the expansions of the non-perturbative *n*-differentials in  $\hbar$  is not a topological expansion. But we still can define  $\omega_n^{\mathsf{np},\langle d \rangle}(z_{\llbracket n \rrbracket}) := [\hbar^d] \omega_n^{\mathsf{np}}(z_{\llbracket n \rrbracket}), d \geq 0$ . An existence of topological expansion would mean that these components could be nonzero only for the values of d of the form  $2g - 2 + n$ ,  $g \in \mathbb{Z}_{\geq 0}$ . But we have, in general, contributions for all nonnegative integers d.

Remark 3.6. Explicitly,  $\omega_n^{\text{np}}$  is defined in terms of  $\{\omega_m\}$  and  $\log \Theta_*$  by a closed combinatorial formula with summation over graphs. As a corollary, we obtain similar combinatorial formulas for all versions of extended differentials  $\Omega_n^{\text{np},\bullet}$ ,  $\Omega_n^{\text{np}}, \mathbb{W}_n^{\text{np}}$ ,  $\mathcal{W}_n^{\text{np}}$  associated to  $\{\omega_n^{\text{np}}\}$ by Equations [\(48\)](#page-10-0), [\(44\)](#page-10-1)–[\(46\)](#page-10-2). All these combinatorial expressions are reviewed in Sect. [4.](#page-13-0)

<span id="page-11-0"></span>3.2. Loop equations and projection property. Assume the standard setup of topological recursion. Define

(50) 
$$
\mathcal{W}_n^{\mathsf{np},\langle d\rangle} \coloneqq [\hbar^d] \mathcal{W}_n^{\mathsf{np}}(z,u; z_{\llbracket n \rrbracket}).
$$

<span id="page-11-2"></span>Theorem 3.7. The non-perturbative differentials satisfy the loop equations. Namely, for any key special point  $q \in \mathcal{P}$  such that both dx and dy are holomorphic at this point and  $dy \neq 0$  and for any  $k > 0$  we have:

(51) 
$$
[u^k]e^{uy}\mathcal{W}_n^{\mathsf{np},\langle d\rangle}(z,u)\in \Xi_q,
$$

where y is any local primitive of dy. This relation holds identically in  $z_1, \ldots, z_n$ .

Remark 3.8. As in the usual perturbative case [\[ABDBKS22;](#page-19-11) [ABDBKS24c\]](#page-19-4), an equivalent way to state the loop equations is to demand that

(52) 
$$
\sum_{r\geq 0} \left(-d\frac{1}{dy}\right)^r [u^r] \mathcal{W}_n^{\mathsf{np},\langle d\rangle}(z,u; z_{\llbracket n \rrbracket}) \text{ is holomorphic at } z = q \text{ for } q \in \mathcal{P}.
$$

Moreover, this relation holds as well in the general setting, with arbitrary orders of zeroes/poles of  $dx$  and  $dy$  at the key-points.

Theorem [3.7](#page-11-2) allows to compute the principal parts of  $\omega_n^{\mathsf{np},\langle d \rangle}$  in each variable once we know  $\omega_n^{np,(d')}$  with  $d' + n' < d + n$ . These principal parts allow us to reconstruct  $\omega_n^{np,(d)}$  for  $d \geq 1, n \geq 1$ , uniquely from  $\omega_n^{\mathsf{np},\langle 0 \rangle}, n \geq 1$ , using the following theorem

<span id="page-11-3"></span>**Theorem 3.9.** For any  $d \geq 0$ ,  $n \geq 0$ , we have the following projection property:

(53) 
$$
\oint_{z \in \mathfrak{A}} \left( \omega_{n+1}^{\mathsf{np}, \langle d \rangle}(z, z_{\llbracket n \rrbracket}) - \eta(z) \partial_w \omega_n^{\mathsf{np}, \langle d \rangle}(z_{\llbracket n \rrbracket}) \right) = 0.
$$

Remark 3.10. This setup is very close to the setup of the so-called blobbed topological recursion [\[BS17\]](#page-19-15). With this analogy the role of blobs in the present construction is played by  $\left(\prod_{i=1}^n \eta(z_i)\partial_w\right) \log \Theta_*(w|\mathcal{T})$ . Note, however, that the expansion in  $\hbar$  is non-topological. The coupling to blobs might also look a bit different at the first glance, but geometry of the underlying Riemann surface allows to reduce the  $\oint_{\mathfrak{B}}$  integrals to the residues at the points in P.

Let us mention that this setup can also be applied to generalized topological recursion, but it requires to develop a relevant piece of theory, see Remark [1.15.](#page-6-3)

<span id="page-11-1"></span>3.3. The  $x - y$  swap relation and dy deformations. Assume the generalized setup of topological recursion.

Consider the dual topological recursion with the initial data  $(\Sigma, dy, dx, \mathcal{P}^{\vee})$ . We use the differentials  $\omega_n^{\vee,(g)}$ , that it produces, to construct exactly the same sequence of nonperturbative objects, that is,  $\Omega_n^{\text{np},\vee,\bullet},\Omega_n^{\text{np},\vee},\mathbb{W}_n^{\text{np},\vee},\omega_n^{\text{np},\vee},$  and  $\mathcal{W}_n^{\text{np},\vee}$ . Then the  $x-y$  swap formula can be extended to the non-perturbative context:

<span id="page-11-4"></span>**Theorem 3.11.** The systems of non-perturbative multi-differentials for  $(\Sigma, dx, dy, \mathcal{P})$  and  $(\Sigma, dy, dx, \mathcal{P}^{\vee})$  are related by the following formulas:

(54) 
$$
\omega_n^{\mathsf{np},\vee}(z_{\llbracket n \rrbracket}) = (-1)^n \left( \prod_{i=1}^n \sum_{r=0}^{\infty} \left( -d_i \frac{1}{dy_i} \right)^r [u_i^r] \right) \mathbb{W}_n^{\mathsf{np}}(z_{\llbracket n \rrbracket}, u_{\llbracket n \rrbracket});
$$

(55) 
$$
\omega_n^{\mathsf{np}}(z_{\llbracket n \rrbracket}) = (-1)^n \left( \prod_{i=1}^n \sum_{r=0}^\infty \left( -d_i \frac{1}{dx_i} \right)^r [u_i^r] \right) \mathbb{W}_n^{\mathsf{np}}(z_{\llbracket n \rrbracket}, u_{\llbracket n \rrbracket}).
$$

A useful tool in the theory of topological recursion is the effect of the deformations of the initial data  $dx$  and  $dy$ . Our goal is to find the corresponding formulas in the nonperturbative case. Since we can use the  $x-y$  swap formula discussed above, it is sufficient to study the deformations of y.

Consider a deformation of dy given by  $dy + \epsilon d(\Delta y)$ , where  $\Delta y$  is some local primitive for the deformation of dy near the points in P. Assume that  $d(\Delta y)$  is regular at the points in P. Let  $\omega_n^{np} + \epsilon \Delta \omega_n^{np}$  be the corresponding deformations of the non-perturbative n-differentials.

## <span id="page-12-1"></span>Theorem 3.12. We have

<span id="page-12-4"></span>(56) 
$$
\Delta \omega_n^{\mathsf{np}}(z_{\llbracket n \rrbracket}) = \sum_{q \in \mathcal{P}} \operatorname{res}_{z=q} \omega_{n+1}^{\mathsf{np}}(z, z_{\llbracket n \rrbracket}) \int_q^z \Delta y \, dx.
$$

<span id="page-12-0"></span>3.4. KP integrability. Assume the generalized setup of topological recursion.

The system of non-perturbative  $n|n$  half-differentials and (extended) differentials satisfies the following KP integrability property expressed as determinantal formulas, which is the main theorem of the present paper.

<span id="page-12-2"></span>**Theorem 3.13.** The system of differentials  $\{\omega_n^{np}\}\$ is KP integrable, that is, we have:

(57) 
$$
\omega_n^{\mathsf{np}}(z_{\llbracket n \rrbracket}) = \det^{\circ}(\Omega_1^{\mathsf{np}}(z_i, z_j)), \quad n \ge 2.
$$

(58) 
$$
\Omega_n^{\mathsf{np},\bullet}(z^+_{\llbracket n \rrbracket}, z^-_{\llbracket n \rrbracket}) = \det(\Omega_1^{\mathsf{np}}(z^+_i, z^-_j)).
$$

The proof of Theorem [3.13](#page-12-2) is given in Section [4.6.](#page-18-0)

In order to incorporate slight changes in the choice of conventions in the definitions of the non-perturbative objects in [\[EM11;](#page-19-5) [BE12;](#page-19-6) [BE15;](#page-19-7) [EO24\]](#page-19-2), we also state the following slight variation of Theorem [3.13:](#page-12-2)

<span id="page-12-3"></span>**Corollary 3.14.** Let  $\{\tilde{\omega}_n^{\text{np}}\}$  be a system of non-perturbative differentials defined by the same formulas as in Definition 3.1, but for a different choice of theta characteristics same formulas as in Definition [3.1,](#page-9-3) but for a different choice of theta characteristics entering the definition of  $\Theta_*$ , or even where we replace  $\Theta_*(w|\mathcal{T})$  in Equation [\(43\)](#page-9-2) by  $\widetilde{\Theta}_*(w|\mathcal{T}) := ae^{bw}\Theta_*(w+c|\mathcal{T})$  for an arbitrary choice of constant a and vectors b, c such that  $\bigcup_{w\to 0} \widetilde{\Theta}_*(w+c|\mathcal{T}) \neq 0$ . Then  $\{\widetilde{\omega}_n^{\text{np}}\}$  is KP integrable.

Indeed, this modification of  $\Theta_*$  leads to a shift of w or to a modification of  $\omega_1^{Kr}$  in the Krichever construction, which obviously preserves KP integrability of Krichever differen-tials and the whole proof of Theorem [3.13](#page-12-2) works for a modified  $\Theta_*$ .

Remark 3.15. For instance, in [\[EM11\]](#page-19-5) the authors use the following choice of parameters

(59) 
$$
a = e^{\frac{1}{4\pi i\hbar^2} \int_{\mathfrak{A}} \omega_1^{(0)} \int_{\mathcal{T}\mathfrak{A}} \omega_1^{(0)} - 2\pi i \,\mu\nu}, \qquad b = \frac{1}{\hbar} \oint_{\mathcal{T}\mathfrak{A}} \omega_1^{(0)}, \qquad c = \frac{1}{2\pi i\hbar} \oint_{\mathfrak{B} - \mathcal{T}\mathfrak{A}} \omega_1^{(0)}
$$

in order to achieve the so-called background independence.

Remark 3.16. Corollary [3.14](#page-12-3) is also a tool to re-introduce the (multi-)KP times as it is done in [\[BE12;](#page-19-6) [EO24;](#page-19-2) [KZ23\]](#page-20-6), which we essentially suppressed everywhere so far for the clarity of exposition except for Equation [\(31\)](#page-8-1). To this end, we select a number of points  $p_1, \ldots, p_N \in \Sigma \setminus \mathcal{P}, N \geq 1$ , and consider the choices of the second kind differentials  $d\Upsilon_k^{\alpha}$  with the only pole at  $p_{\alpha}$  of order  $k+1, \alpha = 1, \ldots, N$ , normalized by the condition  $\oint_{\mathfrak{A}} d\Upsilon_k^{\alpha} = 0$ . Then the following choice of parameters introduces the multi-KP variables  $\check{t}_k^{\alpha}, \alpha = 1, \ldots, N, k \geq 1$ :

(60) 
$$
a = e^{\frac{1}{2}\sum_{\alpha,k}\sum_{\beta,l}\Upsilon^{\alpha\beta}_{kl}t^{\alpha\beta}_{k}t^{\beta}_{l}}, \qquad b = 0, \qquad c = \frac{1}{2\pi i}\oint_{\mathfrak{B}}\sum_{\alpha,k}t^{\alpha}_{k}d\Upsilon^{\alpha}_{k},
$$

where  $\Upsilon_{kl}^{\alpha\beta}$  is a bilinear form obtained by expanding the holomorphic parts of the primitives of  $d\Upsilon_k^{\alpha}$  at  $p_{\beta}$ . We refer to [\[KZ23,](#page-20-6) Sections 3, 4, 5] for a detailed exposition; note that this construction can be enhanced to include more variables of "discrete" kind, cf. also a review in [\[KS13,](#page-20-7) Examples 2 and 3].

## 4. Proofs

<span id="page-13-1"></span><span id="page-13-0"></span>4.1. Graphical formulas. The proofs are based on the graphical formula for  $\omega_n^{\text{np}}$  proposed in [\[BE12;](#page-19-6) [BE15\]](#page-19-7). More precisely, in [\[BE12;](#page-19-6) [BE15\]](#page-19-7) the authors give graphical formulas for the non-perturbative connected  $n|n$  half-differentials  $\Omega_n^{\text{np}}$ , and we further specialize them here in order to deal with  $\omega_n^{\text{np}}$  and  $\mathcal{W}_n^{\text{np}}$ .

In the rest of this section we use a variety of (different!) graphs that share a lot of common features. In each formula below we define a set  $G_{\star}$  of 3-level graphs  $\Gamma$  with some additional structure, and the common grounds are the following:

- The set of vertices  $V(\Gamma)$  splits into three subsets  $V(\Gamma) = V_{\ell}(\Gamma) \sqcup V_{\omega}(\Gamma) \sqcup V_{\theta}(\Gamma)$ , which we call (multi)leaves,  $\omega$ -vertices and  $\theta$ -vertices, respectively.
- Each edge in the set of edges  $E(\Gamma)$  connects an  $\omega$ -vertex to either a multileaf or a θ-vertex. Accordingly, the set  $E(\Gamma)$  splits as  $E(\Gamma) = E_{\ell-\omega}(\Gamma) \sqcup E_{\omega-\theta}(\Gamma)$ .
- Graph Γ must be connected.
- The set  $V_{\ell}(\Gamma)$  consists of n or, in some cases, of  $n+1$  ordered multileaves, depending on a particular problem that we address.
- Each edge  $e \in E_{\ell-\omega}(\Gamma)$  is decorated by an operator  $O(e)$  acting on the decoration of the  $\omega$ -vertex where e is attached. The operators have to be specified in each case.
- There is a map  $g: V_\omega(\Gamma) \to \mathbb{Z}_{\geq 0}$ . An  $\omega$ -vertex v of  $\Gamma$  with  $m(v)$  edges in  $E_{\ell-\omega}(\Gamma)$ and  $k(v)$  edges in  $E_{\omega-\theta}(\Gamma)$  attached to it and labeled by  $g(v)$  is called a  $(g(v), m(v), k(v))$ - $\omega$ -vertex.
- We require that  $2g(v) 2 + m(v) + k(v) \geq 0$  for each  $\omega$ -vertex v.
- The  $(0, 0, 2)$ - $\omega$ -vertices are not allowed.
- The  $(q, m, k)$ - $\omega$ -vertices v are decorated by some differentials  $\omega(v)$ . These decorations have to be specified depending on a particular problem that we address.
- Each  $\theta$ -vertex v is decorated by  $\theta(v) \coloneqq \log \Theta_*(w|\mathcal{T})$ .
- Each edge  $e \in E_{\omega-\theta}(\Gamma)$  is decorated by the bi-linear operator  $O(e) := \frac{1}{2\pi}$  $\frac{1}{2\pi i}\int_{\mathfrak{B}}\partial_w,$ where the  $\int_{\mathfrak{B}_i}$  operators act on the decoration of the attached  $\omega$ -vertex and  $\partial_{w_i}$ act on the decoration of the attached  $\theta$ -vertex.
- $|\text{Aut}(\Gamma)|$  denotes the order of the automorphisms group of a decorated graph  $\Gamma$ that preserves all decorations.
- We associate to a graph  $\Gamma$  its weight  $w_{\star}(\Gamma)$  defined as

(61) 
$$
\mathsf{w}_{\star}(\Gamma) \coloneqq \frac{1}{|\operatorname{Aut}(\Gamma)|} \prod_{e \in E(\Gamma)} O(e) \bigg( \prod_{v \in V_{\omega}(\Gamma)} \omega(v) \prod_{v \in V_{\log \theta}(\Gamma)} \theta(v) \bigg),
$$

where the action of the operators associated to leaves and edges on the decorations of the vertices is prescribed by the graph Γ.

This structure is specified in each particular case further by the choice of the labels of (multi)leaves and specific conditions on the index of these vertices (sometimes that are indeed of index 1, hence the name "leaves"), associated operators  $O(e)$  for  $e \in E_{\ell-\omega}(\Gamma)$ , and the decorations  $\omega(v)$  for the  $(q, m, k)$ - $\omega$ -vertices.

4.1.1. Borot–Eynard formula. Let  $G_{\Omega_n}$  be the set of 3-level graphs  $\Gamma$  with some additional structure described as follows:

- The set  $V_{\ell}(\Gamma)$  consists of n ordered multileaves  $\ell_1, \ldots, \ell_n$ , that is, there is a fixed bijection  $[\![n]\!] \to V_{\ell}(\Gamma)$  such that  $i \mapsto \ell_i$ .
- Each edge  $e \in E_{\ell-\omega}(\Gamma)$  attached to  $\ell_i$  is decorated by the operator  $O(e) \coloneqq \int_{z_i^-}^{z_i^+}$ .
- Each  $(g, m, k)$ -ω-vertex v is decorated by  $\omega(v) := \hbar^{2g-2+m+k} \omega_{m-k}^{(g)}$  $\lim_{m+k}$ , and the arguments of  $\omega_{m-l}^{(g)}$  $\binom{g}{m+k}$  are put in bijection with the edges attached to v.

<span id="page-14-0"></span>Proposition 4.1 ([\[BE12;](#page-19-6) [BE15\]](#page-19-7)). We have

<span id="page-14-1"></span>(62) 
$$
\Omega_n^{\mathsf{np}} = \prod_{i=1}^n \frac{\exp\left(\frac{1}{\hbar} \int\limits_{z_i^-}^{z_i^+} \omega_1^{(0)} + \int\limits_{z_i^-}^{z_i^+} \omega_1\right)}{E(z_i^+, z_i^-)} \sum_{\Gamma \in G_{\Omega_n}} \mathsf{w}_{\Omega_n}(\Gamma).
$$

<span id="page-14-2"></span>Remark 4.2. Proposition [4.1](#page-14-0) is already presented in [\[BE12;](#page-19-6) [BE15\]](#page-19-7); we just formalize their description in a way that is convenient for our proofs below. To this end, in addition to the standard Feynman graphs technique one has to use two additional observations. First of all, we use Equation [\(30\)](#page-8-4) in order to dissolve the first factor in [\(43\)](#page-9-2) into the  $(0, 2)$ - $\omega$ -vertices connected to two different multileaves. Moreover, we use the identity

(63) 
$$
\Theta_*(w + \mathcal{A}\left(\sum_{i=1}^n (z_i^+ - z_i^-)\right)|\mathcal{T}) = \exp\left(\sum_{i=1}^n \frac{1}{2\pi i} \int_{\mathfrak{B}} \int_{z_i^-}^{z_i^+} B \partial_w\right) \exp(\log \Theta_*(w|\mathcal{T}))
$$

to generate the  $(0, 2)$ - $\omega$ -vertices whose two edges are connected to a multileaf and to a  $\theta$ -vertex.

Remark 4.3. The right hand side of [\(62\)](#page-14-1) contains an infinite sum. However, both the left hand side and the right hand side of  $(62)$  can be expanded in  $\hbar$ . Then for each d there is still an infinite number of graphs  $\Gamma \in G_{\Omega_n}$  that contribute non-trivially to the coefficient of  $\hbar^d$  in the  $\hbar$ -expansion, but once we remove all  $(0, 2)$  vertices we get just a finite number of possible "core" graphs. Since the contribution of all possible configurations  $(0, 2)$ -vertices is of exponential type, see Remark [4.2,](#page-14-2) there is no problem with convergence.

Proposition [4.1](#page-14-0) can easily be specialized to the definitions of  $\omega_n^{np}$  and  $\mathcal{W}_n^{np}$ , let us describe the necessary modifications.

4.1.2. A formula for  $\omega_n^{np}$ . Let  $G_{\omega_n}$  be the set of 3-level graphs  $\Gamma$  with some additional structure described as follows:

- The set  $V_{\ell}(\Gamma)$  consists of n ordered leaves  $\ell_1, \ldots, \ell_n$ , that is, there is a fixed bijection  $[\![n]\!] \to V_{\ell}(\Gamma)$  such that  $i \mapsto \ell_i$ .<br>There is exactly one odge atter-
- There is exactly one edge attached to each  $\ell_i$ .
- The edge  $e \in E_{\ell-\omega}(\Gamma)$  attached to  $\ell_i$  is decorated by the operator  $O(e) := \bigcup_{\substack{\to \\ \to i}}$ .
- Each  $(g, m, k)$ -ω-vertex v is decorated by  $\omega(v) := \hbar^{2g-2+m+k} \omega_{m-k}^{(g)}$  $\lim_{m+k}$ , and the arguments of  $\omega_m^{(g)}$  are put in bijection with the edges attached to v.

A direct corollary of Proposition [4.1](#page-14-0) is the following

<span id="page-14-4"></span>Corollary 4.4. We have

<span id="page-14-3"></span>(64) 
$$
\omega_n^{\mathsf{np}} = \sum_{\Gamma \in G_{\omega_n}} \mathsf{w}_{\omega_n}(\Gamma).
$$

Remark 4.5. Equation [\(64\)](#page-14-3) is an equality of two formal power series in  $\hbar$ . For each d there is only a finite number of decorated graphs  $\Gamma \in G_{\omega_n}$  that contribute non-trivially to the coefficient of  $\hbar^d$  in the  $\hbar$ -expansion.

Remark 4.6. Note that the constant term in  $\hbar$  in Equation [\(64\)](#page-14-3) is indeed the one prescribed by Equation  $(49)$ .

Remark 4.7. Note also that feeding Equation [\(64\)](#page-14-3) into the right hand side of Equation [\(48\)](#page-10-0), and subsequently applying Equation [\(44\)](#page-10-1), we get back to Equation [\(62\)](#page-14-1) for  $\Omega_n^{\mathsf{np}}$ .

4.1.3. A formula for  $\mathcal{W}_n^{\text{np}}$ . Let  $G_{\mathcal{W}_n}$  be the set of 3-level graphs  $\Gamma$  with some additional structure described as follows:

- The set  $V_{\ell}(\Gamma)$  consists of  $n+1$  ordered (multi)leaves  $\ell_0, \ell_1, \ldots, \ell_n$ , that is, there is a fixed bijection  $\{0\} \cup [n] \rightarrow V_{\ell}(\Gamma)$  such that  $i \mapsto \ell_i$ .<br>There is exactly and edge attached to each  $\ell_i$  for  $i \in$
- There is exactly one edge attached to each  $\ell_i$  for  $i \in [\![n]\!]$ .<br>• The edge  $e \in F_i$  (F) attached to  $\ell_i$   $i \in [\![n]\!]$  is does
- The edge  $e \in E_{\ell-\omega}(\Gamma)$  attached to  $\ell_i, i \in [\![n]\!]$ , is decorated by the operator  $O(e) = |$  $O(e) \coloneqq |_{\dots}$
- Each edge  $e \in E_{\ell-\omega}(\Gamma)$  attached to  $\ell_0$  is decorated by the operator  $O(e)$  :=  $u\hbar\mathcal{S}(u\hbar\overset{\rightharpoonup}{\partial_{x(z)}}) \frac{1}{dx\beta}$  $\frac{1}{dx(z)}\Big|_{\rightarrow z}.$
- Each  $(g, m, k)$ -ω-vertex v is decorated by  $\omega(v) := \hbar^{2g-2+m+k} \omega_{m-k}^{(g)}$  $\lim_{m+k}^{(g)}$ , and the arguments of  $\omega_m^{(g)}$  are put in bijection with the edges attached to v.

A direct corollary of Proposition [4.1](#page-14-0) is the following

<span id="page-15-1"></span>Corollary 4.8. We have

<span id="page-15-2"></span>(65) 
$$
\mathcal{W}_n^{\mathsf{np}} = \frac{dx}{u\hbar} \exp\left(\Big|_{z' \to z} \Big|_{z'' \to z} u\hbar \mathcal{S}(u\hbar \partial_{x'}) u\hbar \mathcal{S}(u\hbar \partial_{x''}) \Big(\frac{B(z', z'')}{dx'dx''} - \frac{1}{(x' - x'')^2}\Big)\right) \times \exp\left(u\mathcal{S}(u\hbar \partial_x) y - uy\right) \sum_{\Gamma \in G_{\mathcal{W}_n}} \mathsf{w}_{\mathcal{W}_n}(\Gamma).
$$

Here  $x = x(z), x' = x(z'), x'' = x(z'').$ 

<span id="page-15-0"></span>4.2. Proof of the loop equations. In this Section we prove Theorem [3.7.](#page-11-2) To this end, we use Corollary [4.8.](#page-15-1) In the corresponding graph formula we see that we can recollect all prefactors in [\(65\)](#page-15-2) and the weights of all  $\omega$ -vertices attached to  $\ell_0$  and the operators on the edges incident to  $\ell_0$  into known graphical formulas for  $\mathcal{W}_m^{np}$  for some  $m \geq 0$ . Thus, we have the following lemma:

<span id="page-15-4"></span>Lemma 4.9. We have

<span id="page-15-3"></span>(66) 
$$
\mathcal{W}_n^{\mathsf{np}} = \sum_{\Gamma \in G_{\mathcal{W}_n'}} \mathsf{w}_{\mathcal{W}_n'}(\Gamma),
$$

where  $G_{\mathcal{W}'_n}$  is the set of 3-level graphs  $\Gamma$  with some additional structure described as follows:

- The set  $V_{\ell}(\Gamma)$  consists of  $n+1$  ordered leaves  $\ell_0, \ell_1, \ldots, \ell_n$ , that is, there is a fixed bijection  $\{0\} \cup [n] \rightarrow V_{\ell}(\Gamma)$  such that  $i \mapsto \ell_i$ .
- There is exactly one edge attached to each  $\ell_i$  for  $i \in \{0\} \cup [n]$ .
- The edge  $e \in E_{\ell-\omega}(\Gamma)$  attached to  $\ell_i, i \in [\![n]\!]$ , is decorated by the operator  $O(e) \coloneqq$  $\big\downarrow_{\rightarrow z_i}$ .
- The edge  $e \in E_{\ell-\omega}(\Gamma)$  attached to  $\ell_0$  is decorated by the operator  $O(e) \coloneqq \lfloor_{\rightarrow z}$ .
- Each  $(g, m, k)$ -w-vertex v not connected to  $\ell_0$  is decorated by  $\omega(v) \coloneqq \hbar^{2g-2+m+k} \omega_{m-k}^{(g)}$  $\binom{(g)}{m+k}$ and the arguments of  $\omega_m^{(g)}$  are put in bijection with the edges attached to v.
- The only  $(g, m + 1, k)$ -w-vertex v that is connected to  $\ell_0$  is decorated by  $\omega(v)$  :=  $\hbar^{2g-1+m+k}\mathcal{W}_{m+}^{(g)}$  $\mathcal{W}^{(g)}_{m+k},$  and the arguments of  $\mathcal{W}^{(g)}_k$  $\mathbf{F}_k^{(g)}$  are put in bijection with the edges

attached to v in such a way that the distinguished one corresponds to the edge that connects v to  $\ell_0$ .

According to the definition of the weight  $w_{W'_n}(\Gamma)$ , Equation [\(66\)](#page-15-3) represents  $\mathcal{W}_n^{\mathsf{np},\langle d\rangle}$  as a finite linear combination of  $\mathcal{W}_m^{(g)}(z, u; z'_1, \dots, z'_m)$  with  $2g + m \leq d$  with some operators (expressed as a finite some over graphs) in the variables  $z'_1, \ldots, z'_m$ . Note that each of these  $\mathcal{W}_m^{(g)}(z, u; z'_1, \ldots, z'_m)$  satisfies the loop equations, that is, for any key special point  $q \in \mathcal{P}$  such that both dx and dy are holomorphic at this point and  $dy \neq 0$  and for any  $k \geq 0$  we have:

<span id="page-16-2"></span>(67) 
$$
[u^k]e^{uy}\mathcal{W}_n^{(g)}(z,u)\in \Xi_q,
$$

where y is any local primitive of  $du$  (here we use the standard setup). Moreover, this relation holds identically in  $z_1, \ldots, z_n$ . Therefore, we have [\(67\)](#page-16-2) in z for any operator in in the variables  $z'_1, \ldots, z'_m$  applied to  $\mathcal{W}_m^{(g)}(z, u; z'_1, \ldots, z'_m)$ . Thus  $\mathcal{W}_n^{\mathsf{np}, \langle d \rangle}$  is expressed as a finite linear combination of elements of  $\Xi_q$ . Hence,  $[u^k]e^{uy}\mathcal{W}_n^{\mathsf{np},\langle d\rangle} \in \Xi_q$ . This completes the proof of Theorem [3.7.](#page-11-2)

<span id="page-16-0"></span>4.3. Proof of the projection formula. In this Section we prove Theorem [3.9.](#page-11-3) Consider Equation [\(64\)](#page-14-3), replacing n by  $n + 1$ . The sum over graphs in this expression can be split into two parts, depending on whether or not the leaf  $\ell_{n+1}$  is attached to a  $(0, 1, 1)$ - $\omega$ -vertex further attached to a  $\theta$ -vertex.

In the first case, since  $\frac{1}{2\pi i} \int_{\mathfrak{B}} B(z_{n+1}, \cdot) = \eta(z_{n+1})$ , we obtain the sum over graphs that is manifestly just

(68) 
$$
\eta(z_{n+1})\partial_w \sum_{\Gamma \in G_{\omega_n}} \mathsf{w}_{\omega}(\Gamma) = \eta(z_{n+1})\partial_w \omega_n^{\mathsf{np}}.
$$

The second sum for each  $\omega_{n+1}^{\mathsf{np},\langle d\rangle}$ ,  $d \geq 0$ ,  $n \geq 0$ , is a finite linear combination of the differentials  $\omega_{m+1}^{(g)}(z_1',\ldots,z_m',z_{n+1})$  with  $2g-1+m\geq 0$ , with some operators in  $z_1',\ldots,z_m'$ applied to them. Since each  $\omega_{m+1}^{(g)}(z_1',\ldots,z_m',z_{n+1})$  has vanishing  $\mathfrak{A}$ -periods in  $z_{n+1}$ , we conclude that the same holds for  $\omega_{n+1}^{\mathsf{np},\langle d \rangle} - \eta(z_{n+1}) \partial_w \omega_n^{\mathsf{np},\langle d \rangle}$ . This completes the proof of Theorem [3.9.](#page-11-3)

<span id="page-16-1"></span>4.4. Proof of the  $x - y$  swap relation. In this Section we prove Theorem [3.11.](#page-11-4) Recall that next to  $\omega_n^{(g)}$  and  $\omega_n^{\vee,(g)}$  we also considered in [\[ABDBKS22\]](#page-19-11) the two-index differentials  $\omega_{m,n}^{(g)}$  such that  $\omega_{m,0}^{(g)} = \omega_m^{(g)}$  and  $\omega_{0,n}^{(g)} = \omega_n^{\vee,(g)}$ . Since  $\omega_{m+1,n}^{(g)}(z) + \omega_{m,n}^{(g)}$  $\binom{g}{m,1+n}(z)$  is d-exact (here  $z$  is the argument that changes the side in the two-index differentials), we have

<span id="page-16-3"></span>(69) 
$$
\oint_{z \in \mathfrak{B}} \omega_{m+1,n}^{(g)} = \oint_{z \in \mathfrak{B}} \omega_{m,1+n}^{(g)}.
$$

Now, applying the same resummation as we did above in Lemma [4.9](#page-15-4) to the Borot– Eynard formula given in Proposition [4.1,](#page-14-0) we obtain the following formula for  $\mathbb{W}_n^{\mathsf{np}}$ :

# Lemma 4.10. We have

(70) 
$$
\mathbb{W}_n^{\mathsf{np}} = \sum_{\Gamma \in G_{\mathbb{W}_n}} \mathsf{w}_{\mathbb{W}_n}(\Gamma),
$$

where  $G_{W_n}$  is the set of 3-level graphs  $\Gamma$  with some additional structure described as follows:

- The set  $V_{\ell}(\Gamma)$  consists of n ordered leaves  $\ell_1, \ldots, \ell_n$ , that is, there is a fixed bijection  $\llbracket n \rrbracket \to V_\ell(\Gamma)$  such that  $i \mapsto \ell_i$ .
- There is exactly one edge attached to each  $\ell_i$ .
- The edge  $e \in E_{\ell-\omega}(\Gamma)$  attached to  $\ell_i$  is decorated by the operator  $O(e) \coloneqq \bigcup_{\substack{\to \\ \to (z_i, u_i)}}$ .

## 18 A. ALEXANDROV, B. BYCHKOV, P. DUNIN-BARKOWSKI, M. KAZARIAN, AND S. SHADRIN

• Each  $(g, m, k)$ -ω-vertex v is decorated by  $\omega(v) := \hbar^{2g-2+m+k} \prod_{i=1}^{k} \bigcup_{u_{m_i}\to 0} \mathbb{W}_{m-k}^{(g)}$  $\binom{g}{m+k}$ and the first m (respectively, last k) arguments of  $\mathbb{W}_{m}^{(g)}$  $\binom{(g)}{m+k}$  are put in bijection with the edges in  $E_{\ell-\omega}(\Gamma)$  (respectively, in  $E_{\omega-\theta}(\Gamma)$ ) attached to v.

Now, the theory of  $x - y$  duality in the standard perturbative case implies that

<span id="page-17-1"></span>(71) 
$$
(-1)^n \left( \prod_{i=1}^n \sum_{r=0}^\infty \left( -d_i \frac{1}{dy_i} \right)^r [u_i^r] \right) \sum_{\Gamma \in G_{W_n}} \mathsf{w}_{\mathbb{W}}(\Gamma)
$$

is given by exactly the same sum over decorated graphs as  $\mathbb{W}_n^{\mathsf{np}}$ , with the following two modifications:

- Now each  $(g, m, k)$ -ω-vertex v is decorated by  $\omega(v) \coloneqq \hbar^{2g-2+m+k} \omega_{k,m}^{(g)}$ , and the last m (respectively, the first k) arguments of  $\omega_{k,m}^{(g)}$  are put in bijection with the edges in  $E_{\ell-\omega}(\Gamma)$  (respectively, in  $E_{\omega-\theta}(\Gamma)$ ) attached to v.
- The edge  $e \in E_{\ell-\omega}(\Gamma)$  attached to  $\ell_i$  is decorated by the operator  $O(e) \coloneqq \bigcup_{\substack{\to z_i}}$ .

Using Equation [\(69\)](#page-16-3), we see that we can equivalently decorate each  $(g, m, k)$ - $\omega$ -vertex v by  $\hbar^{2g-2+m+k}\omega_{0,m+k}^{(g)} = \hbar^{2g-2+m+k}\omega_{m+k}^{\vee,(g)}$  $\binom{v,(g)}{m+k}$ . This turns the resulting formula for [\(71\)](#page-17-1) into exactly the same graphical expression as the one we have on the right hand side of Equa-tion [\(64\)](#page-14-3), with all  $\omega_m^{(g)}$ 's in the decorations of the  $\omega$ -vertices replaced by  $\omega_m^{\vee,(g)}$ 's. Thus, by Corollary [4.4,](#page-14-4) we have

(72) 
$$
(-1)^n \left( \prod_{i=1}^n \sum_{r=0}^\infty \left( -d_i \frac{1}{dy_i} \right)^r [u_i^r] \right) \sum_{\Gamma \in G_{\mathbb{W}_n}} \mathsf{w}_{\mathbb{W}_n}(\Gamma) = \omega_n^{\mathsf{np}, \vee}.
$$

This completes the proof of the first assertion of Theorem [3.11.](#page-11-4) Now, according to [\[AB-](#page-19-11)[DBKS22\]](#page-19-11), the second statement is a formal inverse of the first one, or, alternatively, one can prove if by exactly the same argument reversing the roles of  $dx$  and  $dy$ .

<span id="page-17-0"></span>4.5. Proof of the deformation formula. In this section we prove Theorem [3.12.](#page-12-1) We just substitute Equation [\(64\)](#page-14-3) on the left hand side and on the right hand side of Equation [\(56\)](#page-12-4).

On the left hand side, we can combine the known deformation formula for  $\omega_n^{(g)}$  and the Leibniz rule. This means that we change the decoration of one of the  $(g, m, k)$ - $\omega$ -vertices with  $2g - 2 + m + k > 0$  from  $\omega_{m}^{(g)}$  ${}_{m+k}^{(g)}$  to  $\sum_{q \in \mathcal{P}}$  res<sub> $z=q \omega_{m+k+1}^{(g)}(z) \int_q^z \Delta y dx$ .</sub>

On the right hand side, there are three possible case

- (1) First possible situation is that the leaf corresponding to the extra variable  $z$  is connected to a  $(q, m, k)$ -ω-vertex with  $2q - 2 + m + k > 1$ . The sum over all possible cases like this coincides with the left hand side.
- $(2)$  Second possibility is that the leaf corresponding to the extra variable z is connected to a  $(0, m, k)$ - $\omega$ -vertex with  $m+k=3$ . But then notice that  $\int_q^z \Delta y \, dx$  has a double zero at q and  $\omega_3^{(0)}$  $_3^{(0)}$  at most a double pole at q. This means that the sum of residues  $\sum_{q \in \mathcal{P}} \text{res}_{z=q} \,\omega_3^{(0)}$  $\int_3^{(0)}(z)\int_q^z \Delta y\,dx$  vanishes and these terms don't contribute.
- (3) The third possibility is that the leaf corresponding to the extra variable is connected to a  $(0,1,1)$ -ω- or  $(0,2,0)$ -ω-vertex. But again,  $\omega_2^{(0)} = B$  is holomorphic at each  $q \in \mathcal{P}$ , hence the sum of residues  $\sum_{q \in \mathcal{P}} res_{z=q} \omega_2^{(0)}$  $\int_{2}^{(0)}(z)\int_{q}^{z} \Delta y dx$  vanishes and these terms don't contribute.

Thus we prove that the left hand side and the right hand side of Equation [\(56\)](#page-12-4) are equal. This completes the proof of the theorem.

#### <span id="page-18-0"></span>4.6. Proof of the KP integrability. In this Section we prove Theorem [3.13.](#page-12-2)

4.6.1. *Standard setup.* We assume first that we are in the realm of the standard setup of topological recursion. That is, we assume that  $P$  is the set of zeros of dx and dy is regular and non-vanishing at each point  $q \in \mathcal{P}$ .

To this end, note that both the  $x-y$  swap and deformations of dy, such that dy is regular at zeros of  $dx$ , preserve the KP integrability of a system of differentials, see [\[ABDBKS23,](#page-19-1) Theorem 2.7] and [\[ABDBKS24b,](#page-19-3) Corollary 2.6], respectively. These statements can be applied to  $\{\omega_n^{np}\}\$ , since the formulas in Theorems [3.11](#page-11-4) and [3.12](#page-12-1) are exactly the same as in the case of the usual differentials of topological recursion. Combining these two operations, we are also allowed to deform  $dx$  in such a way that  $dx$  remains regular at the zeros of  $dy$ .

Thus, our strategy is to apply a sequence of deformations of  $dx$  and  $dy$  such that at the end we arrive to the situation that  $dx$  has poles of sufficiently high order at all zeros of  $dy$ . This would mean that the system of dual differentials  $\{\omega_n^{\vee,(g)}\}$  is computed by generalized topological recursion with  $\mathcal{P}^{\vee} = \emptyset$ , and all differentials in the stable range are equal to zero:  $\omega_n^{\vee,(g)} = 0$  for  $2g - 2 + n > 0$ . Hence, by Lemma [3.4,](#page-10-4)  $\omega_n^{\mathsf{np},\vee} = \omega_n^{\mathsf{Kr}}$ , and this system is integrable. Hence, its  $x - y$  dual system of differentials  $\{\omega_n^{np}\}\$ is also KP integrable, as well as any other system of non-perturbative differentials connected to it via KP integrable deformations, including the original system of non-perturbative differentials.

Let us describe a construction of the desired deformation. Denote the initial differentials by  $dx_0$  and  $dy_0$  and assume that  $dx_0$  is regular and non-vanishing at the zeros of  $dy_0$ . We need to choose some  $dx_1$  and  $dy_1$  satisfying the following conditions:

- $dy_0$  is regular and non-vanishing at the zeros of  $dx_1$ ;
- $dy_1$  is regular and non-vanishing at the zeros of  $dx_1$ ;
- $dx_1$  is regular and non-vanishing at the zeros of  $dy_0$ ;
- $dx_1$  has poles of sufficiently high order at the zeros of  $dy_1$ .

The choice is done in the following order. First, we choose some  $dy_1$  whose set of zeros is disjoint from the set of zeros of  $dy_0$ . Then we choose  $dx_1$  such that it has poles of sufficiently high order at the zeros of  $dy_1$  and no other poles, and its zeros are disjoint from the zeros and poles of  $dy_0$  and the poles of  $dy_1$ . Both choices are obviously possible.

Now we first deform  $(dx_0, dy_0)$  to  $(dx_t, dy_0)$ ,  $0 \le t \le 1$ , where  $dx_1$  is the one we have chosen, and in order to avoid any problems with the existence of the global differentials  $dx_t$  for  $0 < t < 1$  we demand only that they are defined locally at the zeros of  $dy_0$  (this relaxed setup is sufficient for the integrability properties, cf. [\[ABDBKS24b,](#page-19-3) Section 4]). Then we deform  $(dx_1, dy_0)$  to  $(dx_1, dy_s)$ ,  $0 \le s \le 1$ , where  $dy_1$  is the one we have chosen, and  $dy_s$  for  $0 < s < 1$  are defined only locally at the zeros od  $dx_1$ .

As we discussed above, this sequence of deformations preserves KP integrability, and the resulting system of differentials satisfies the conditions for KP integrability discussed above. This completes the proof of Theorem [3.13](#page-12-2) in the standard setup of topological recursion.

4.6.2. Extension to generalized setup. Now assume that the input data  $(\Sigma, dx, dy, \mathcal{P})$  corresponds to generalized topological recursion in the sense of [\[ABDBKS24c\]](#page-19-4). Then we can use exactly the same argument as in [\[ABDBKS24c,](#page-19-4) Proof of Theorem 6.4], which, in a nutshell, says that the system of differentials of generalized topological recursion can be realized as the limit of a one-parameter family of a system the differentials of the standard topological recursion. This extends without any effort to the non-perturbative systems of differentials. The observation that the KP integrability property is a closed condition completes the proof in this case.

#### <span id="page-19-0"></span>**REFERENCES**

- <span id="page-19-13"></span>[ABDBKS24a] A. Alexandrov, B. Bychkov, P. Dunin-Barkowski, M. Kazarian, and S. Shadrin. "Log Topological Recursion Through the Prism of  $x-y$  Swap". In: *Int. Math. Res. Not. IMRN* 21 (2024), pp. 13461–13487. DOI:  $10$ . [1093/imrn/rnae213](https://doi.org/10.1093/imrn/rnae213).
- <span id="page-19-11"></span>[ABDBKS22] A. Alexandrov, B. Bychkov, P. Dunin-Barkowski, M. Kazarian, and S. Shadrin. A universal formula for the  $x - y$  swap in topological recursion. 2022. arXiv: [2212.00320 \[math-ph\]](https://arxiv.org/abs/2212.00320). (to appear in J. Eur. Math. Soc.)
- <span id="page-19-1"></span>[ABDBKS23] A. Alexandrov, B. Bychkov, P. Dunin-Barkowski, M. Kazarian, and S. Shadrin. KP integrability through the  $x - y$  swap relation. 2023. arXiv: [2309.12176 \[math-ph\]](https://arxiv.org/abs/2309.12176).
- <span id="page-19-3"></span>[ABDBKS24b] A. Alexandrov, B. Bychkov, P. Dunin-Barkowski, M. Kazarian, and S. Shadrin. Any topological recursion on a rational spectral curve is KP integrable. 2024. arXiv: [2406.07391 \[math-ph\]](https://arxiv.org/abs/2406.07391).
- <span id="page-19-4"></span>[ABDBKS24c] A. Alexandrov, B. Bychkov, P. Dunin-Barkowski, M. Kazarian, and S. Shadrin. Degenerate and irregular topological recursion. 2024. arXiv: [24](https://arxiv.org/abs/2408.02608) [08.02608 \[math-ph\]](https://arxiv.org/abs/2408.02608).
- <span id="page-19-14"></span>[ABDBKS24d] A. Alexandrov, B. Bychkov, P. Dunin-Barkowski, M. Kazarian, and S. Shadrin. "Symplectic duality via log topological recursion". English. In: Commun. Number Theory Phys. 18.4 (2024), pp. 795–841. DOI: [10.4310/](https://doi.org/10.4310/CNTP.241203001416) [CNTP.241203001416](https://doi.org/10.4310/CNTP.241203001416).
- <span id="page-19-6"></span>[BE12] G. Borot and B. Eynard. "Geometry of spectral curves and all order dispersive integrable system". In: SIGMA Symmetry Integrability Geom. Methods Appl. 8 (2012), Paper 100, 53. DOI: [10.3842/SIGMA.2012.100](https://doi.org/10.3842/SIGMA.2012.100).
- <span id="page-19-7"></span>[BE15] G. Borot and B. Eynard. "All order asymptotics of hyperbolic knot invariants from non-perturbative topological recursion of A-polynomials". In: Quantum Topol. 6.1 (2015), pp. 39–138. DOI:  $10.4171/qT/60$ .
- <span id="page-19-15"></span>[BS17] G. Borot and S. Shadrin. "Blobbed topological recursion: properties and applications". In: Math. Proc. Cambridge Philos. Soc. 162.1 (2017), pp. 39– 87. doi: [10.1017/S0305004116000323](https://doi.org/10.1017/S0305004116000323).
- <span id="page-19-12"></span>[BCGFLS23] G. Borot, S. Charbonnier, E. Garcia-Failde, F. Leid, and S. Shadrin. Functional relations for higher-order free cumulants. 2023. arXiv: [2112.](https://arxiv.org/abs/2112.12184) [12184 \[math.OA\]](https://arxiv.org/abs/2112.12184).
- <span id="page-19-8"></span>[CEO06] L. Chekhov, B. Eynard, and N. Orantin. "Free energy topological expansion for the 2-matrix model". In: J. High Energy Phys. 12 (2006), pp. 053, 31. doi: [10.1088/1126-6708/2006/12/053](https://doi.org/10.1088/1126-6708/2006/12/053).
- <span id="page-19-10"></span>[DBOSS14] P. Dunin-Barkowski, N. Orantin, S. Shadrin, and L. Spitz. "Identification of the Givental formula with the spectral curve topological recursion procedure". In: *Comm. Math. Phys.* 328.2 (2014), pp. 669–700. DOI: [10.](https://doi.org/10.1007/s00220-014-1887-2) [1007/s00220-014-1887-2](https://doi.org/10.1007/s00220-014-1887-2).
- <span id="page-19-9"></span>[EO07] B. Eynard and N. Orantin. "Invariants of algebraic curves and topological expansion". In: Commun. Number Theory Phys. 1.2 (2007), pp. 347–452. doi: [10.4310/CNTP.2007.v1.n2.a4](https://doi.org/10.4310/CNTP.2007.v1.n2.a4).
- <span id="page-19-5"></span>[EM11] B. Eynard and M. Mariño. "A holomorphic and background independent partition function for matrix models and topological strings". In: J. Geom. Phys. 61.7 (2011), pp. 1181–1202. DOI: [10.1016/j.geomphys.](https://doi.org/10.1016/j.geomphys.2010.11.012) [2010.11.012](https://doi.org/10.1016/j.geomphys.2010.11.012).
- <span id="page-19-2"></span>[EO24] B. Eynard and S. Oukassi. Hirota, Fay and Geometry. 2024. arXiv: [2401.](https://arxiv.org/abs/2401.08317) [08317 \[math-ph\]](https://arxiv.org/abs/2401.08317).



<span id="page-20-7"></span><span id="page-20-5"></span><span id="page-20-4"></span><span id="page-20-3"></span><span id="page-20-0"></span>

<span id="page-20-6"></span><span id="page-20-2"></span><span id="page-20-1"></span>A. A.: CENTER FOR GEOMETRY AND PHYSICS, INSTITUTE FOR BASIC SCIENCE (IBS), POHANG 37673, Korea

Email address: alex@ibs.re.kr

B. B.: Department of Mathematics, University of Haifa, Mount Carmel, 3498838, Haifa, Israel

Email address: bbychkov@hse.ru

P. D.-B.: FACULTY OF MATHEMATICS, NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, USACHEVA 6, 119048 MOSCOW, RUSSIA; HSE-SKOLTECH INTERNATIONAL LABORAtory of Representation Theory and Mathematical Physics, Skoltech, Nobelya 1, 143026, Moscow, Russia; and ITEP, 117218 Moscow, Russia

Email address: ptdunin@hse.ru

M. K.: Faculty of Mathematics, National Research University Higher School of Economics, Usacheva 6, 119048 Moscow, Russia; and Center for Advanced Studies, Skoltech, Nobelya 1, 143026, Moscow, Russia

Email address: kazarian@mccme.ru

S. S.: Korteweg-de Vries Institute for Mathematics, University of Amsterdam, Postbus 94248, 1090GE Amsterdam, The Netherlands Email address: S.Shadrin@uva.nl