

Extension of the Standard Model with Chern-Simons type interaction

Volodymyr Gorkavenko¹, Ivan Hrynychak¹,
Oleksandr Khasai², Mariia Tsarenkova¹

¹ *Faculty of Physics, Taras Shevchenko National University of Kyiv,
64, Volodymyrs'ka str., Kyiv 01601, Ukraine*

² *Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine
14-b, Metrolohichna str., Kyiv 03143, Ukraine*

Abstract

Extension of the Standard Model with Chern-Simons type interaction contains a new vector massive boson (Chern-Simons boson) that couples to electroweak gauge bosons by the so-called effective Chern-Simons interaction. There is no direct interaction between the Chern-Simons bosons and SM fermions. We consider existing restrictions on the parameters of this SM extension, the effective loop interaction of a new vector boson with SM fermions, and the possibility of the manifestation of the long-lived GeV-scale Chern-Simons bosons in collider experiments.

1 Introduction

While the Standard Model (SM) has proven remarkably successful in describing collider experiments [1], there is compelling indirect evidence pointing toward the existence of new physics. Some examples of phenomena that SM can not explain are active neutrino oscillations [2–4], dark matter [5–8], and the baryon asymmetry of the Universe [9–11]. We can suggest the existence of new particles beyond SM (BSM particles) to solve SM's problems. However, there is also a possibility that there are new particles unrelated to solving these problems.

A natural question arises: if there are new particles, why have they not been detected in numerous experiments? There are two possible answers. First, the particles might be too heavy, and current accelerator energies are not sufficient to produce them. In this case, detection would require new, more powerful accelerators such as the FCC [12, 13]. Alternatively, the new particles could be light enough to be produced nowadays at the existing accelerators [14–17], but their interactions with SM particles are so feebly that they have not yet been observed. The search for such long-lived particles is already underway in the so-called intensity frontier experiments such as MATHUSLA [18], FACET [19], FASER [20, 21], SHiP [22, 23], NA62 [24–26], DUNE [27, 28], LHCb [29], etc.

To develop the phenomenology of these particles, we need to consider the different types they can be. Namely, they can be scalar [30–32], pseudoscalar (axionlike) [33–36],




Figure 1: Diagrams generating the Chern-Simons interaction. Heavy fermions, beyond the Standard Model, run in loop triangular diagrams.

fermion (heavy neutral leptons) [37–40], or vector (dark photons) [41–44] particles, see detail, e.g., in reviews [18,23]. Each of these possibilities results in a theory that introduces different new terms in the SM Lagrangian, often referred to as portals. This work focuses on a portal involving a new massive vector boson with Chern-Simons-like interaction.

Chern-Simons interactions are known to arise in a variety of theoretical models, including those with extra dimensions and string theory frameworks [45–50].

The idea of creating a Chern-Simons portal is based on the phenomenon of chiral anomaly cancellation in physical theories. Interest in the Lagrangian terms generated by the chiral anomaly is explained by the fact that their appearance does not depend on the mass of the particle, which runs along the internal triangular loop. This means that if it exists a very heavy particle that cannot be born in a collider experiments and detected directly, then this heavy particle can manifest itself by generating the corresponding chiral term of the interaction. Actually, it is the only way heavy fermions with a mass that can exceed the capabilities of modern accelerators by many orders of magnitude can manifest themselves in low-energy experiments. For example, if assume that the capacities of modern accelerators are not sufficient for the discovery of the heavy t -quark, then it would manifest itself as a necessary additional term for cancellation of the chiral anomalies in SM.

Following [51], we will be interested in the possibility of the manifestation of the interaction of the new vector massive gauge field X_μ , for example, of the group $U_1(X)$, with SM particles that are generated by the condition of cancellation of the chiral anomaly. Assume that there are new heavy fermions charged with respect to the gauge group of the SM $U_Y(1)$ and certain additional group $U_X(1)$. At the same time, SM fermions are considered as uncharged with respect to the $U_X(1)$ group and, accordingly, do not directly interact with the X_μ field. Heavy fermions cannot manifest themselves at modern accelerators; accordingly, it would seem that the X_μ field cannot manifest itself at low energies either. In [51], SM is modified to a theory with symmetry $SU_C(3) \times SU_W(2) \times U_Y(1) \times U_X(1)$, in which the X_μ boson manifests itself in the interaction with SM vector fields thanks to the diagrams presented in Fig.1.

As a result of the chiral anomaly effect the interaction between the new vector bosons (henceforth we will call them Chern-Simons or CS bosons) and SM particles is induced in the form of gauge-invariant Lagrangian described by operators with minimum dimension 6 [23,51]:

$$\mathcal{L}_1 = \frac{C_Y}{\Lambda_Y^2} \cdot X_\mu (\mathfrak{D}_\nu H)^\dagger H B_{\lambda\rho} \cdot \epsilon^{\mu\nu\lambda\rho} + h.c., \quad (1)$$

$$\mathcal{L}_2 = \frac{C_{SU(2)}}{\Lambda_{SU(2)}^2} \cdot X_\mu (\mathfrak{D}_\nu H)^\dagger F_{\lambda\rho} H \cdot \epsilon^{\mu\nu\lambda\rho} + h.c.. \quad (2)$$

Here, Λ_Y and $\Lambda_{SU(2)}$ represent new scales of the theory, while C_Y and $C_{SU(2)}$ are dimensionless coupling constants. $\epsilon^{\mu\nu\lambda\rho}$ stands for the Levi-Civita tensor ($\epsilon^{0123} = +1$), and X_μ is a CS vector boson. Please note that X_μ is a Stueckelberg field [52, 53], which ensures the gauge invariance of the Lagrangians (1) and (2). The Higgs doublet scalar field is denoted by H , and $B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$ and $F_{\mu\nu} = -ig \sum_{i=1}^3 \frac{\tau^i}{2} V_{\mu\nu}^i$ refer to the field strength tensors of the $U_Y(1)$ and $SU_W(2)$ gauge fields of the SM, respectively.

After electroweak symmetry breaking, the Lagrangians (1) and (2) produce many terms, including three-field interactions written as four-dimensional operators

$$\mathcal{L}_{CS} = c_z \epsilon^{\mu\nu\lambda\rho} X_\mu Z_\nu \partial_\lambda Z_\rho + c_\gamma \epsilon^{\mu\nu\lambda\rho} X_\mu Z_\nu \partial_\lambda A_\rho + \{c_w \epsilon^{\mu\nu\lambda\rho} X_\mu W_\nu^- \partial_\lambda W_\rho^+ + h.c.\}, \quad (3)$$

where A_μ is the electromagnetic field, and W_μ^\pm and Z_μ are the fields associated with weak interactions. The coefficients c_z , c_γ , and c_w are independent and dimensionless. Both c_z and c_γ are real, but c_w can be complex $c_w = \Theta_{W1} + i\Theta_{W2}$. Importantly, the CS vector boson X_μ does not directly interact with SM fermions.

In this paper, we consider existing restrictions on the parameters of this SM extension for the case of light CS bosons, the effective loop interaction of a new vector boson with SM fermions and the possibility of the manifestation of the long-lived GeV-scale CS bosons in collider experiments.

2 Constrains from interaction with vector fields of SM

In the case $M_X < M_W, M_Z$ interaction in the form (3) leads to the additional channel of W^\pm - and Z -boson decay. Namely, the following process are allowed: $Z_\mu \rightarrow X_\mu + \gamma$, $W_\mu^\pm \rightarrow X_\mu + q_n + \bar{q}_m$, $W_\mu^\pm \rightarrow X_\mu + \ell_n + \bar{\nu}_m$.

The corresponding decay width of Z -boson can be easily calculated:

$$\Gamma(Z \rightarrow X\gamma) = c_\gamma^2 \frac{M_Z}{96\pi} \left(1 - \frac{M_X^2}{M_Z^2}\right)^3 \left(1 + \frac{M_Z^2}{M_X^2}\right). \quad (4)$$

In the limit of small M_X , $M_X/M_Z \ll 1$ we have

$$\Gamma_{ZX} = c_\gamma^2 \frac{M_Z}{96\pi} \left(\frac{M_Z}{M_X}\right)^2. \quad (5)$$

The corresponding decay width of the W^\pm boson can also be analytically calculated in the approximation of massless quarks and leptons and without taking into account hadronization effects

$$\Gamma(W^+ \rightarrow Xu\bar{d}) = \frac{N_C M_W^3 G_F |V_{ud}|^2}{3456\sqrt{2}\pi^3} (\Theta_{W1}^2 F_1(x) + \Theta_{W2}^2 F_2(x)), \quad (6)$$

where $x = M_W/M_X$, $N_C = 3$ – number of quark colors and F_1, F_2 are dimensionless functions

$$F_1(x) = \frac{4}{x^2} - 6 \ln x (24 - 108x^2 + 20x^4 - x^6) - 392 + 639x^2 - 274x^4 + 23x^6 +$$

$$3x(14-x^2)(4-x^2)^{3/2} \left(\pi - 2 \arctan \frac{x(3-x^2)}{\sqrt{4-x^2(1-x^2)}} \right),$$

$$F_2(x) = 4 + 6x^6 \ln x - 45x^2 + 18x^4 + 23x^6 +$$

$$\frac{3x^3(20+2x^2-x^4)}{\sqrt{4-x^2}} \left(\pi - 2 \arctan \frac{x(3-x^2)}{\sqrt{4-x^2(1-x^2)}} \right).$$

It should be noted that in (6) there is no terms $\sim \Theta_{W1}\Theta_{W2}$. So contributions from real and imaginary parts of coupling c_w do not interfere with each other.

In the limit of small M_X , $M_X/M_W \ll 1$ we have

$$\Gamma(W^+ \rightarrow X u \bar{d}) = \Theta_{W1}^2 \frac{N_C M_W^3 G_F |V_{ud}|^2}{864 \sqrt{2} \pi^3} \left(\frac{M_W}{M_X} \right)^2. \quad (7)$$

Taking into account channels of decay W -boson into X -boson and quarks (main contribution is from decay into u, \bar{d} and c, \bar{s} because of diagonal elements of the CKM matrix) and into three generations of the charged lepton and corresponding neutrino also. So the full decay width of the W^+ -boson with the production of X -boson for the case $M_X/M_W \ll 1$ can be written as

$$\Gamma_{WX} = \frac{\Gamma(W^+ \rightarrow X u \bar{d})}{|V_{ud}|^2} (|V_{ud}|^2 + |V_{cs}|^2 + 3/N_C) \approx \Theta_{W1}^2 \frac{M_W^3 G_F}{96 \sqrt{2} \pi^3} \frac{M_W^2}{M_X^2}. \quad (8)$$

Current measurements of the decay width of the W^\pm, Z bosons are in agreement with SM predictions. However experimental measurements have some uncertainties [54], namely we have $\Gamma_W = 2.085 \pm 0.042$ GeV and $\Gamma_Z = 2.4955 \pm 0.0023$ GeV. So, new possible channels of the W^\pm, Z decay into CS boson can exist only if $\Gamma_{ZX} < 2\Delta\Gamma_Z$ and $\Gamma_{WX} < 2\Delta\Gamma_W$, where $\Delta\Gamma_Z = 0.0023$ GeV, $\Delta\Gamma_W = 0.042$. Therefore we can roughly estimate from above the magnitude of couplings [23]

$$c_\gamma^2 \lesssim 10^{-6} \left(\frac{M_X}{1 \text{ GeV}} \right)^2, \quad [Re c_w]^2 \lesssim 10^{-2} \left(\frac{M_X}{1 \text{ GeV}} \right)^2. \quad (9)$$

It should be noted, that in the case of c_γ a significantly stronger bound comes from the measuring of the single photon events at LEP [55]. There the branching at the level $Br = \Gamma_{ZX}/\Gamma_{Z,total} < 10^{-6}$ was established for photons with energy above 15 GeV. This leads to the stronger bound $c_\gamma^2 \lesssim 10^{-9} \left(\frac{M_X}{1 \text{ GeV}} \right)^2$. In the case of light CS bosons ($M_X/M_W \ll 1$), we expect $Z \rightarrow X\gamma$ decays with photon energy $E \lesssim 45$ GeV.

3 Effective loop interaction with fermions of the different flavour

In addition to interaction with SM vector fields, the CS boson can interact effectively with SM fermions due to loop diagrams. Such interaction of the CS bosons with quarks




Figure 2: CS boson loop interactions with two quarks of different flavours.

of different flavours, see Fig.2, was considered in [56–58]. In this case, effective interaction occurs only due to the interaction of the CS boson with W^\pm boson via coupling $c_w = \Theta_{W1} + i\Theta_{W2}$.

It was shown that the divergent part of the loop diagrams is proportional to a non-diagonal element of the unity matrix V^+V (V is the Cabibbo–Kobayashi–Maskawa matrix) and is removed. It allows one to construct an effective Lagrangian of the interaction of the CS bosons with quarks of different flavours. If $\Theta_{W1} \neq 0$ than the dominant terms of this Lagrangian have form [58]

$$\mathcal{L}_{quarks}^{CS} = \sum_{m < n} \Theta_{W1} \left(C_{mn} \bar{d}_m \gamma^\mu \hat{P}_L d_n X_\mu + C_{nm}^+ \bar{d}_n \gamma^\mu \hat{P}_L d_m X_\mu \right), \quad (10)$$

where d_n is down-type quark, the summation occurs over the quark generations,

$$C_{mn} = \frac{3a}{2\sqrt{2}\pi^2} G_F m_t^2 V_{d_m t}^+ V_{td_n}, \quad (11)$$

and

$$\begin{aligned} a &= 0.13, & |C_{sb}| &= 1.97 \cdot 10^{-4}, \\ |C_{db}| &= 4.43 \cdot 10^{-5}, & |C_{ds}| &= 1.77 \cdot 10^{-6}. \end{aligned} \quad (12)$$

As one can see the effective Lagrangian depends only on one of two unknown couplings (Θ_{W1}) of the CS boson interaction with W boson. It was shown that the interaction of the GeV-scale CS boson with up-type quarks can be neglected.

The obtained effective Lagrangian (10) allows us to compute dominant production channels of GeV-scale CS bosons in decays of mesons due to reactions $b \rightarrow s + X$, $b \rightarrow d + X$, $s \rightarrow d + X$. These reactions correspond to the production of the CS bosons in the following decays of charged and neutral mesons: $B \rightarrow K + X$, $B \rightarrow \pi + X$, $K \rightarrow \pi + X$, where final mesons may also be in excited states, see details in [58].

4 Effective loop interaction with fermions of the same flavour

Unlike the case of effective loop interaction of the CS bosons with quarks of the different flavours, where interaction is defined only by diagrams with W -bosons, see Fig.2 and by coupling c_w , in the case of interaction of the CS boson with quarks of the same flavours or with leptons the diagrams with Z -bosons and photons are also important, see Fig.3. Thus the interaction with quarks of the same flavours or leptons depends on c_w , c_γ and c_Z couplings of Lagrangian (3).




Figure 3: Diagrams of the CS boson's decay into leptons in the unitary gauge. The interaction depends on c_w , c_γ and c_Z couplings.

We would like the divergences in the loop diagrams of the interaction of the CS bosons with fermions of the same flavours to be automatically removed as well otherwise we will have a problem. These divergences cannot be removed via counterterms because the initial Lagrangian (3) does not contain terms of the interaction of the CS boson and SM fermions.

As was shown in [58], for effective loop interaction of CS bosons with quarks of the same flavours or with leptons, divergences in loop diagrams with vertex XWW are not automatically removed during calculations. In [59] this interaction was considered in the unitary gauge taking into account all corresponding diagrams, see Fig.3. It was hoped that the sum of the differences of all diagrams could be cancelled for a certain relation between c_w , c_γ and c_Z couplings. Unfortunately, it was concluded that using Lagrangian (3), we can not eliminate the divergences in the effective interaction of the CS bosons with fermions of the same flavours.

We can only hope that, perhaps, further consideration of this problem in non-unitary gauge will help solve the problem of divergences. Otherwise, it will mean that the interaction of the CS bosons with fermions of the same flavours must be considered within the framework of an effective field theory, namely with the help of corresponding effective operators with a set of new couplings. Or maybe it will be necessary to supplement the initial Lagrangian (3) with additional terms, which will allow us to remove differences via corresponding counterterms. In both cases, additional couplings appear in addition to c_w , c_γ and c_Z couplings, which complicates the task of finding manifestations of the CS bosons in experiments.

5 Discussion

In this paper, we considered an extension of the SM including a new light massive vector boson with the Chern-Simons interactions. We considered the low energy Lagrangian of the CS boson interaction with vector fields of SM (3) and we have given the known limits on the parameters of this Lagrangian c_w, c_Z, c_γ . Also, we considered the effective loop interaction of the CS bosons with fermions. While the interaction with quarks of different flavours is well defined, the interaction with fermions of the same flavours contains divergences that have not yet been removed [59].

Even though there is no hope of seeing the direct interaction of the BSM particles with collider detectors, the BSM long-lived particles can still be searched in collider experiments. The main idea of such experiments is not to directly search for the BSM particles but to search for products of its decay into the SM particles. To do this, it is necessary to produce as many BSM particles as possible as a result of SM reactions, e.g., in proton-proton collisions. The produced BSM particles must be further isolated from the SM particles to avoid background events, and then look for very rare decay events of the BSM particles. This is the idea of the intensity frontier experiments mentioned in the Introduction.

As is well known, before the experimental search of a BSM particle at intensity frontier experiments one has to compute the sensitivity region of these experiments, namely the parameter region of the new particle (mass and coupling of interaction with SM particles) where the particle will manifest itself in the experiment. The procedure of computation sensitivity region is well known, see e.g. [60]. It should be noted, that for computation of sensitivity region, one needs to know the technical parameters of the experiment facility and all dominant channels of the BSM particle production and decays.

Let us now consider the possibility of experimentally searching for the CS boson in the intensity frontier experiments.

Regarding production channels, only the production from heavy mesons' decay into the lighter mesons is currently calculated [56–58] due to obtained effective Lagrangian (10) of the interaction the CS boson with quarks of different flavours. But the CS boson can be produced and via interaction the CS boson with quarks of the same flavours. For example, in reactions, e.g, $\pi^0 \rightarrow X\gamma$, $\omega \rightarrow \eta X$, $\phi \rightarrow \eta X$ or due to bremsstrahlung, or deep inelastic scattering in proton-proton interactions.

As for the decay channels, we can only consider decays of the CS bosons into quarks of different flavours followed by hadronization, but even the decays into lepton pairs are not yet available for calculation.

As one can see, the question of the effective interaction of the CS bosons with fermions of the same flavour is crucial for the experimental search for the Chern-Simons bosons. This is what should be the main focus of further research.

6 Acknowledgments

The authors would like to thank the Organizers of the conference "New Trends in High-Energy and Low-x Physics" in Sfantu Gheorghe (Romania) for their kind hospitality during this extremely interesting and inspiring meeting. The work of V.G., I.H., and O.Kh. was supported by the National Research Foundation of Ukraine under project No. 2023.03/0149.

References

- [1] W.N. Cottingham, D.A. Greenwood. An Introduction to the Standard Model of Particle Physics, (Cambridge University Press, 2023) [ISBN: 978-1-00-940168-5, 978-1-00-940172-2, 978-1-00-940170-8, 978-0-511-27136-6, 978-0-521-85249-4].
- [2] S.M. Bilenky, S.T. Petcov. Massive Neutrinos and Neutrino Oscillations. *Rev. Mod. Phys.*, **59**, 671 (1987) [DOI: 10.1103/RevModPhys.59.671] [Erratum: Rev.Mod.Phys. 61, 169 (1989), Erratum: Rev.Mod.Phys. 60, 575–575 (1988)].
- [3] A. Strumia, F. Vissani. Neutrino masses and mixings and... arXiv:hep-ph/0606054 (2006).
- [4] P.F. de Salas, D.V. Forero, C.A. Ternes, M. Tortola, J.W.F. Valle. Status of neutrino oscillations 2018: 3σ hint for normal mass ordering and improved CP sensitivity. *Phys. Lett. B*, **782**, 633 (2018) [DOI: 10.1016/j.physletb.2018.06.019].
- [5] P.J.E. Peebles. Dark Matter. *Proc. Nat. Acad. Sci.*, **112**, 2246 (2015) [DOI: 10.1073/pnas.1308786111].
- [6] V. Lukovic, P. Cabella, N. Vittorio. Dark matter in cosmology. *Int. J. Mod. Phys. A*, **29**, 1443001 (2014) [DOI: 10.1142/S0217751X14430015].
- [7] G. Bertone, D. Hooper. History of dark matter. *Rev. Mod. Phys.*, **90**, 045002 (2018) [DOI: 10.1103/RevModPhys.90.045002].
- [8] M. Cirelli, A. Strumia, J. Zupan. Dark matter. arXiv:hep-ph/2406.01705 (2024).
- [9] G. Steigman. Observational tests of antimatter cosmologies. *Ann. Rev. Astron. Astrophys.*, **14**, 339 (1976) [DOI: 10.1146/annurev.aa.14.090176.002011].
- [10] A. Riotto, M. Trodden. Recent progress in baryogenesis. *Ann. Rev. Nucl. Part. Sci.*, **49**, 35 (1999) [DOI: 10.1146/annurev.nucl.49.1.35].
- [11] L. Canetti, M. Drewes, M. Shaposhnikov. Matter and Antimatter in the Universe. *New J. Phys.*, **14**, 095012 (2012) [DOI: 10.1088/1367-2630/14/9/095012].
- [12] T. Golling et al. Physics at a 100 TeV pp collider: beyond the Standard Model phenomena. arXiv:1606.00947 (2016).
- [13] A. Abada et al. FCC Physics Opportunities: Future Circular Collider Conceptual Design Report Volume 1. *Eur. Phys. J. C*, **79**, 474 (2019) [DOI: 10.1140/epjc/s10052-019-6904-3].
- [14] V.M. Gorkavenko. Search for Hidden Particles in Intensity Frontier Experiment SHiP. *Ukr. J. Phys.*, **64**, 689 (2019) [DOI: 10.15407/ujpe64.8.689].
- [15] J. Beacham et al. Physics Beyond Colliders at CERN: Beyond the Standard Model Working Group Report. *J. Phys. G*, **47**, 010501 (2020) [DOI: 10.1088/1361-6471/ab4cd2].

- [16] G. Lanfranchi, M. Pospelov, P. Schuster. The Search for Feebly Interacting Particles. *Ann. Rev. Nucl. Part. Sci.*, **71**, 279 (2021) [DOI: 10.1146/annurev-nucl-102419-055056].
- [17] C. Antel et al. Feebly Interacting Particles: FIPs 2022 workshop report. *Eur. Phys. J. C*, **83**, 1122 (2023) [DOI: 10.1140/epjc/s10052-023-12168-5].
- [18] D. Curtin et al. Long-Lived Particles at the Energy Frontier: The MATH-USLA Physics Case. *Rept. Prog. Phys.*, **82**, 116201 (2019) [DOI: 10.1088/1361-6633/ab28d6].
- [19] S. Cerci et al. FACET: A new long-lived particle detector in the very forward region of the CMS experiment. arXiv:2201.00019 (2021).
- [20] A. Ariga et al. Letter of Intent for FASER: ForWArD Search ExpeRiment at the LHC. arXiv:1811.10243 (2018) [REPORT NUMBER: CERN-LHCC-2018-030, LHCC-I-032, UCI-TR-2018-18, KYUSHU-RCAPP-2018-05].
- [21] A. Ariga et al. FASER’s physics reach for long-lived particles. *Phys. Rev. D*, **99**, 095011 (2019) [DOI: 10.1103/PhysRevD.99.095011].
- [22] M. Anelli et al. A facility to Search for Hidden Particles (SHiP) at the CERN SPS. arXiv:1504.04956 (2015).
- [23] S. Alekhin et al. A facility to Search for Hidden Particles at the CERN SPS: the SHiP physics case. *Rept. Prog. Phys.*, **79**, 124201 (2016) [DOI: 10.1088/0034-4885/79/12/124201].
- [24] P. Mermod. Prospects of the SHiP and NA62 experiments at CERN for hidden sector searches. *PoS, NuFact2017*, 139 (2017) [DOI: 10.22323/1.295.0139].
- [25] E. Cortina Gil et al. Search for heavy neutral lepton production in K^+ decays. *Phys. Lett. B*, **778**, 137 (2018) [DOI: 10.1016/j.physletb.2018.01.031].
- [26] M. Drewes, J. Hajer, J. Klaric, G. Lanfranchi. NA62 sensitivity to heavy neutral leptons in the low scale seesaw model. *JHEP*, **07**, 105 (2018) [DOI: 10.1007/JHEP07(2018)105].
- [27] R. Acciarri et al. Long-Baseline Neutrino Facility (LBNF) and Deep Underground Neutrino Experiment (DUNE): Conceptual Design Report, Volume 2: The Physics Program for DUNE at LBNF. arXiv:1512.06148 (2015) [REPORT NUMBER: FERMILAB-DESIGN-2016-02].
- [28] B. Abi et al. Prospects for beyond the Standard Model physics searches at the Deep Underground Neutrino Experiment. *Eur. Phys. J. C*, **81**, 322 (2021) [DOI: 10.1140/epjc/s10052-021-09007-w].
- [29] V. Gorkavenko, B.K. Jashal, V. Kholoimov, Y. Kyselov, D. Mendoza, M. Ovchynnikov et al. LHCb potential to discover long-lived new physics particles with lifetimes above 100 ps. *Eur. Phys. J. C* **84**, 608 (2024) [DOI: 10.1140/epjc/s10052-024-12906-3].

- [30] B. Patt, F. Wilczek. Higgs-field portal into hidden sectors. arXiv:hep-ph/0605188 (2006).
- [31] F. Bezrukov, D. Gorbunov. Light inflaton Hunter’s Guide. *JHEP*, **05**, 010 (2010) [DOI: 10.1007/JHEP05(2010)010].
- [32] I. Boiarska, K. Bondarenko, A. Boyarsky, V. Gorkavenko, M. Ovchinnikov, A. Sokolenko. Phenomenology of GeV-scale scalar portal. *JHEP*, **11**, 162 (2019) [DOI: 10.1007/JHEP11(2019)162].
- [33] R.D. Peccei, H.R. Quinn. CP Conservation in the Presence of Instantons. *Phys. Rev. Lett.*, **38**, 1440 (1977) [DOI: 10.1103/PhysRevLett.38.1440].
- [34] S. Weinberg. A New Light Boson? *Phys. Rev. Lett.*, **40**, 223 (1978) [DOI: 10.1103/PhysRevLett.40.223].
- [35] F. Wilczek. Problem of Strong P and T Invariance in the Presence of Instantons. *Phys. Rev. Lett.*, **40**, 279 (1978) [DOI: 10.1103/PhysRevLett.40.279].
- [36] K. Choi, S.H. Im, C.S. Shin. Recent progress in physics of axions or axion-like particles. arXiv:2012.05029 (2020).
- [37] T. Asaka, M. Shaposhnikov. The ν MSM, dark matter and baryon asymmetry of the universe. *Phys. Lett. B*, **620**, 17 (2005) [DOI: 10.1016/j.physletb.2005.06.020].
- [38] T. Asaka, S. Blanchet, M. Shaposhnikov. The nuMSM, dark matter and neutrino masses. *Phys. Lett. B*, **631**, 151 (2005) [DOI: 10.1016/j.physletb.2005.09.070].
- [39] K. Bondarenko, A. Boyarsky, D. Gorbunov, O. Ruchayskiy. Phenomenology of GeV-scale Heavy Neutral Leptons. *JHEP*, **11**, 032 (2018) [DOI: 10.1007/JHEP11(2018)032].
- [40] A. Boyarsky, M. Drewes, T. Lasserre, S. Mertens, O. Ruchayskiy. Sterile neutrino Dark Matter. *Prog. Part. Nucl. Phys.*, **104**, 1 (2019) [DOI: 10.1016/j.pnpnp.2018.07.004].
- [41] L.B. Okun. LIMITS OF ELECTRODYNAMICS: PARAPHOTONS? *Sov. Phys. JETP*, **56**, 502 (1982).
- [42] B. Holdom. Two U(1)’s and Epsilon Charge Shifts. *Phys. Lett. B*, **166**, 196 (1986) [DOI: 10.1016/0370-2693(86)91377-8].
- [43] P. Langacker. The Physics of Heavy Z' Gauge Bosons. *Rev. Mod. Phys.*, **81**, 1199 (2009) [DOI: 10.1103/RevModPhys.81.1199].
- [44] P. Ilten, Y. Soreq, M. Williams, Wei Xue. Seren10.1007/JHEP06(2018)004dipity in dark photon searches. *JHEP*, **06**, 004 (2018) [DOI: 10.1007/JHEP06(2018)004].
- [45] I. Antoniadis, E. Kiritsis, T.N. Tomaras. A D-brane alternative to unification. *Phys. Lett. B*, **486**, 186 (2000) [DOI: 10.1016/S0370-2693(00)00733-4].
- [46] C. Coriano, N. Irges, E. Kiritsis. On the effective theory of low scale orientifold string vacua. *Nucl. Phys. B*, **746**, 77 (2006) [DOI: 10.1016/j.nuclphysb.2006.04.009].

- [47] P. Anastasopoulos, M. Bianchi, E. Dudas, E. Kiritsis. Anomalies, anomalous U(1)'s and generalized Chern-Simons terms. *JHEP*, **11**, 057 (2006) [DOI: 10.1088/1126-6708/2006/11/057].
- [48] J.A. Harvey, C.T. Hill, R.J. Hill. Standard Model Gauging of the Wess-Zumino-Witten Term: Anomalies, Global Currents and pseudo-Chern-Simons Interactions. *Phys. Rev. D*, **77**, 085017 (2008) [DOI: 10.1103/PhysRevD.77.085017].
- [49] P. Anastasopoulos, F. Fucito, A. Lionetto, G. Pradisi, A. Racioppi, Y.S. Stanev. Minimal Anomalous U(1)-prime Extension of the MSSM. *Phys. Rev. D*, **78**, 085014 (2008) [DOI: 10.1103/PhysRevD.78.085014].
- [50] J. Kumar, A. Rajaraman, J.D. Wells. Probing the Green-Schwarz Mechanism at the Large Hadron Collider. *Phys. Rev. D*, **77**, 066011 (2008) [DOI: 10.1103/PhysRevD.77.066011].
- [51] I. Antoniadis, A. Boyarsky, S. Espahbodi, O. Ruchayskiy, J.D. Wells. Anomaly driven signatures of new invisible physics at the Large Hadron Collider. *Nucl. Phys. B*, **824**, 296 (2010) [DOI: 10.1016/j.nuclphysb.2009.09.009].
- [52] H. Ruegg and M. Ruiz-Altaba. The Stueckelberg field. *Int. J. Mod. Phys. A* **19**, 3265 (2004) [DOI: 10.1142/S0217751X04019755].
- [53] G.D. Kribs, G. Lee and A. Martin. Effective field theory of Stückelberg vector bosons. *Phys. Rev. D* **106**, 055020 (2022) [DOI: 10.1103/PhysRevD.106.055020].
- [54] S. Navas et al. (Particle Data Group), *Phys. Rev. D* **110**, 030001 (2024) [DOI: 10.1103/PhysRevD.110.030001].
- [55] M. Acciarri et al. Search for new physics in energetic single photon production in e^+e^- annihilation at the Z resonance. *Phys. Lett. B* **412**, 201 (1997) [DOI: 10.1016/S0370-2693(97)01003-4].
- [56] J.A. Dror, R. Lasenby, M. Pospelov. New constraints on light vectors coupled to anomalous currents. *Phys. Rev. Lett.*, **119**, 141803 (2017) [DOI: 10.1103/PhysRevLett.119.141803].
- [57] J.A. Dror, R. Lasenby, M. Pospelov. Dark forces coupled to nonconserved currents. *Phys. Rev. D*, **96**, 075036 (2017) [DOI: 10.1103/PhysRevD.96.075036].
- [58] Y. Borysenkova, P. Kashko, M. Tsarenkova, K. Bondarenko, V. Gorkavenko. Production of Chern-Simons bosons in decays of mesons. *J. Phys. G*, **49**, 085003 (2022) [DOI: 10.1088/1361-6471/ac77a7].
- [59] Y. Borysenkova, V. Gorkavenko, I. Hrynychak, O. Khasai, M. Tsarenkova, Divergences in the effective loop interaction of the Chern-Simons bosons with leptons. The unitary gauge case. *Ukr. J. Phys.*, **69**, 897 (2024). [DOI: 10.15407/ujpe69.12.897].
- [60] M. Ovchinnikov, J.-L. Tastet, O. Mikulenko, K. Bondarenko. Sensitivities to feebly interacting particles: public and unified calculations. *Phys. Rev. D* **108**, 075028 (2023). [DOI: 10.1103/PhysRevD.108.075028].