FORMAL SERIES OF JACOBI FORMS

HIROKI AOKI, TOMOYOSHI IBUKIYAMA, CRIS POOR

ABSTRACT. We prove for general paramodular level that formal series of scalar Jacobi forms with an involution condition necessarily converge and are therefore the Fourier-Jacobi expansions at the standard 1-cusp of paramodular Fricke eigenforms.

CONTENTS

1. Introduction.	1
2. Notation.	4
3. Formal Series.	5
4. Vanishing Order of Jacobi Forms.	9
5. Invariance under subgroups of finite	index in $\Gamma^0(N)$. 13
6. Specialization.	17
7. Locally Bounded.	23
8. Holomorphicity.	27
9. Main Theorem.	33
10. Concluding Remarks.	39
References	40

1. INTRODUCTION.

A parmodular form is a Siegel modular form of degree two for the following discrete group

$$K(N) = \operatorname{Sp}(4, \mathbb{Q}) \cap \begin{pmatrix} \mathbb{Z} & N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & \mathbb{Z} & \mathbb{Z} & \frac{1}{N}\mathbb{Z} \\ \mathbb{Z} & N\mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ N\mathbb{Z} & N\mathbb{Z} & N\mathbb{Z} & \mathbb{Z} \end{pmatrix} = \operatorname{Stab}_{\operatorname{Sp}(4, \mathbb{Q})} \begin{pmatrix} \mathbb{Z} \\ \mathbb{Z} \\ \mathbb{N} \\ \mathbb{Z} \end{pmatrix}.$$

Paramodular forms $M_k(K(N))$ are a natural generalization of elliptic modular forms $M_k(\Gamma_0(N))$ and are interesting in many ways. Roberts and Schmidt [35, 36] gave a sophisticated theory of local and global

Date: December 30, 2024.

²⁰²⁰ Mathematics Subject Classification. Primary: 11F46, 11F50.

Key words and phrases. Formal series, Jacobi form, Paramodular form.

AOKI, IBUKIYAMA, POOR

paramodular newforms. Paramodular newforms have applications to modularity; weight two to abelian surfaces [8, 9], and weight three to nonrigid Calabi-Yau threefolds [15] of Hodge type (1, 1, 1, 1). Conjectures of Ibukiyama [22, 23], and of Ibukiyama and Kitayama [25], connecting paramodular forms to algebraic modular forms on a compact twist of GSp(4), motivated mainly by independent calculations of dimension formulae, have recently been proven [39, 37]. In [10], this connection was generalized to a correspondence with Fricke eigenspaces. Utilizing results of [10], dimension formulae for Fricke plus and minus spaces of paramodular forms for prime level were computed in [24]. For weights $k \geq 3$, these recent proofs allow paramodular Hecke eigensystems to also be computed using orthogonal modular forms [10, 3]. Finally, Gritsenko lifts and Borcherds products provide concrete examples of paramodular forms [19].

Paramodular forms $f \in M_k(K(N))$ have Fourier expansions in three variables but the natural generalization of the Fourier series of an elliptic modular form is perhaps the Fourier-Jacobi expansion of a paramodular form, which, using the components $\Omega = \begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} \in \mathcal{H}_2$, recollects the Fourier series in powers of $e(\omega) = e^{2\pi i \omega}$,

$$f\left(\begin{smallmatrix} \tau & z \\ z & \omega \end{smallmatrix}\right) = \sum_{m=0}^{\infty} \phi_m(\tau, z) e\left(Nm\omega\right).$$

Each coefficient is a Jacobi form, $\phi_m \in J_{k,Nm}$, and has its own Fourier expansion $\phi_m(\tau, z) = \sum_{n \ge 0, r \in \mathbb{Z}} c(n, r; \phi_m) e(n\tau + rz)$. The Fourier-Jacobi expansion thus defines a map to formal series of Jacobi forms FJ : $M_k(K(N)) \to \mathbb{M}(k, N) = \prod_{m=0}^{\infty} J_{k,Nm}$. This map is not surjective because we cannot freely select a sequence of Jacobi forms and obtain the convergent Fourier-Jacobi expansion of a paramodular form. One source of consistency conditions among the Fourier-Jacobi coefficients (ϕ_m) arises from a normalizing involution of K(N), the paramodular Fricke involution $\mu_N = \begin{pmatrix} {}^{t}F_N^{-1} & 0 \\ 0 & F_N \end{pmatrix}$, where $F_N = \frac{1}{\sqrt{N}} \begin{pmatrix} 0 & 1 \\ -N & 0 \end{pmatrix}$ is the Fricke involution on $\Gamma_0(N)$. This involution splits $M_k(K(N))$ into plus and minus forms, $M_k(K(N)) = M_k(K(N))^+ \oplus M_k(K(N))^-$. The block diagonal form of μ_N gives a simple action on the Fourier series and consequently gives the following *involution condition* on the Fourier-Jacobi coefficients of any $f \in M_k(K(N))^\epsilon$, $\epsilon \in \{\pm 1\}$:

(1) For all semidefinite
$$\binom{n-r/2}{r/2}$$
 with $n, r, m \in \mathbb{Z}$,
 $c(n, r; \phi_m) = (-1)^k \epsilon c(m, r; \phi_n).$

Let $\mathbb{M}(k, N, \epsilon)$ be the subspace of formal series of Jacobi forms satisfying the involution condition:

$$\mathbb{M}(k, N, \epsilon) = \{ \mathfrak{f} \in \mathbb{M}(k, N) : \mathfrak{f} \text{ satisfies condition } (1) \}.$$

The Fourier-Jacobi expansion gives $FJ: M_k(K(N))^{\epsilon} \to \mathbb{M}(k, N, \epsilon)$. It seems a bit audacious to hope that the involution condition alone forces the convergence of a formal series to a paramodular Fricke eigenform, but theoretical results for low level N and computed examples suggest that this is true, and we prove it here.

Theorem 1.1 (Main Theorem). Let $N \in \mathbb{N}$, $k \in \mathbb{N}_0$, and $\epsilon \in \{\pm 1\}$. The map $\mathrm{FJ} : M_k (K(N))^{\epsilon} \to \mathbb{M}(k, N, \epsilon)$ is an isomorphism.

The case k = 0 is not difficult, and neither is the case k = 1, when both the domain and codomain are trivial by a result of Skoruppa [38], but k = 2 already has important examples. The case N = 1 of Theorem 1.1 was proven by the first author [1], the cases $2 \le N \le 4$ by Yuen and the second and third authors [26]. These results raised the question, made explicit in [26], of whether the involution condition alone implies the convergence of a formal series of Jacobi forms. Theorem 1.1 resolves this question in the affirmative for the first time.

Bruinier [6], Raum [40], and Bruinier and Raum [7, 5] have approached the theory of formal series of Jacobi forms in a more general setting but with a different set of hypotheses. Bruinier [6] and Raum [40] independently extended Aoki's result for $Sp(4, \mathbb{Z})$ to vector valued formal series of Jacobi forms, and thereby resolved a conjecture of Kudla [27, 28, 29] concerning generating series of special cycles for degree n = 2. In [7], Bruinier and Raum proved that formal series of Jacobi forms for $\operatorname{Sp}(2n,\mathbb{Z})$ converge under the assumption of all $\operatorname{GL}(n,\mathbb{Z})$ symmetries, thereby proving Kudla's modularity conjecture for general n. Pollack [32] has also given a proof of automatic convergence for cuspidal automorphic forms on Sp(2n) that takes Jacobi and Levi symmetries as hypotheses. Bruinier and Raum [5] have recently developed a theory of formal series for subgroups commensurable with $\operatorname{Sp}(2n,\mathbb{Z})$; they prove that compatible formal series of Jacobi forms at every 1-cusp and Levi symmetries imply convergence. They interpret formal series as sections of a line bundle over a formal complex space. Our Main Theorem 1.1 for paramodular groups in degree two and levels N > 3 is not a consequence of any of these results. We only use formal series of Jacobi forms at the standard 1-cusp and only assume symmetry under a single involution.

Our main result has applications to computations. Jacobi restriction [26, 4, 33] is a method that attempts to rigorously compute the space $\operatorname{FJ}(S_k(K(N))^{\epsilon})$ by imposing necessary linear relations on the space of Jacobi forms $\prod_{m=1}^{d} J_{k,Nm}^{\operatorname{cusp}}$. Jacobi restriction has provided rigorous upper bounds for dim $S_k(K(N))^{\epsilon}$, even though the authors could not guarantee in advance that the method would work. The following corollary proves that any schema for computing spaces of paramodular forms that spans spaces of Jacobi forms and imposes the involution condition is in principle sound.

Corollary 1.2. For $d \in \mathbb{N}$, define the \mathbb{C} -vector space $\mathbb{M}(k, N, \epsilon)[d]$ as $\{(\phi_m) \in \prod_{m=0}^d J_{k,Nm} : (\phi_m) \text{ satisfies } (1) \text{ for all } n, m \leq d \}$. The sequence $\dim_{\mathbb{C}} \mathbb{M}(k, N, \epsilon)[d]$ is monotonically decreasing for $d \geq \frac{1}{6}Nk$, and we have $\lim_{d\to+\infty} \dim_{\mathbb{C}} \mathbb{M}(k, N, \epsilon)[d] = \dim_{\mathbb{C}} M_k (K(N))^{\epsilon}$.

In particular, we have $\dim_{\mathbb{C}} M_k (K(N))^{\epsilon} \leq \dim_{\mathbb{C}} \mathbb{M}(k, N, \epsilon)[d]$ for $d > \frac{1}{6}Nk$, and we have equality for sufficiently large d.

The first author was supported by JSPS KAKENHI Grant Numbers JP19K03429 and JP23K03039. The second author was supported by the following grants: JSPS KAKENHI Grant Number JP19K03424, JP20H00115 and JP23K03031. The authors thank the American Institute of Mathematics for its critical support of this research.

2. NOTATION.

We denote the natural numbers by $\mathbb{N} = \{1, 2, 3, ...\}$ and the whole numbers by $\mathbb{N}_0 = \{0, 1, 2, ...\}$. Throughout this article $N \in \mathbb{N}$ denotes a level, $k \in \mathbb{N}_0$ a weight, and $\epsilon \in \{\pm 1\}$ a sign. We write σ' for the transpose of a matrix σ , and σ^* for the transpose inverse. Let $t[\sigma] = \sigma' t\sigma$ for compatibly sized matrices, and $\langle a, b \rangle = tr(ab)$ for $a, b \in M_{n \times n}^{\text{sym}}(\mathbb{C})$. For $z \in \mathbb{C}$, set $e(z) = e^{2\pi i z}$.

For the theory of Jacobi forms see [11]. The upper half plane is $\mathcal{H}_1 = \{\tau \in \mathbb{C} : \operatorname{Im}(\tau) > 0\}$. For a Jacobi form $\phi \in J_{k,m}$ of weight k and index $m \in \mathbb{N}_0$ with $\phi : \mathcal{H}_1 \times \mathbb{C} \to \mathbb{C}$ we write the Fourier expansion as $\phi(\tau, z) = \sum_{n \in \mathbb{N}_0, r \in \mathbb{Z}} c(n, r; \phi) q^n \zeta^r$ with $q = e(\tau)$ and $\zeta = e(z)$. The order of a nonzero ϕ is $\operatorname{ord} \phi = \min\{n \in \mathbb{N}_0 : \exists r \in \mathbb{Z} : c(n, r; \phi) \neq 0\}$. For $\nu \in \mathbb{N}_0$, we set $J_{k,m}(\nu) = \{\phi \in J_{k,m} : \operatorname{ord} \phi \geq \nu\}$.

For the theory of Siegel modular forms we refer to [12]. For a ring $R \subseteq \mathbb{R}$ define the positive definite cone with entries in R by $\mathcal{P}_n(R) = \{s \in M_{n \times n}^{\text{sym}}(R) : s > 0\}$, and the positive semidefinite cone by $\overline{\mathcal{P}}_n(R) = \{s \in M_{n \times n}^{\text{sym}}(R) : s \ge 0\}$. The Siegel upper half space is $\mathcal{H}_n = \{\Omega \in M_{n \times n}^{\text{sym}}(\mathbb{C}) : \text{Im}(\Omega) \in \mathcal{P}_n(\mathbb{R})\}$. An element $\sigma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in the real symplectic group $\text{Sp}(2n, \mathbb{R})$ acts on the symmetric space \mathcal{H}_n by $\sigma \langle \Omega \rangle = (a\Omega + b)(c\Omega + d)^{-1}$, and the Siegel factor of automorphy $j : \text{Sp}(2n, \mathbb{R}) \times \mathcal{H}_n \to \mathbb{C}^{\times}$ is $j(\sigma, \Omega) = \det(c\Omega + d)$. For a function

 $f: \mathcal{H}_n \to \mathbb{C}$ the slash k-action $(f|_k \sigma)(\Omega) = j(\sigma, \Omega)^{-k} f(\sigma \langle \Omega \rangle)$ defines another function $f|_k \sigma : \mathcal{H}_n \to \mathbb{C}$. For a discrete group $\Gamma \subseteq \operatorname{Sp}(2n, \mathbb{R})$ commensurable with $\operatorname{Sp}(2n, \mathbb{Z})$, write $M_k(\Gamma)$ for the \mathbb{C} -vector space of Siegel modular forms of weight k, and $S_k(\Gamma)$ for the subspace of cusp forms.

For the paramodular group K(N) in degree n = 2, write the Fourier expansion for $f \in M_k(K(N))$ as $f(\Omega) = \sum_{t \in \bar{\mathcal{X}}(N)} a(t; f)e(\langle \Omega, t \rangle)$, where $\bar{\mathcal{X}}(N) = \{ \begin{pmatrix} n & r/2 \\ r/2 & Nm \end{pmatrix} \in \bar{\mathcal{P}}_2(\mathbb{Q}) : n, r, m \in \mathbb{Z} \}$. For the Fourier-Jacobi expansion of f, we write $\Omega = \begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix}$ and collect the Fourier expansion in $e(\omega)$ to obtain $f\begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} = \sum_{m=0}^{\infty} \phi_m(\tau, z)e(Nm\omega)$ with Fourier-Jacobi coefficients $\phi_m \in J_{k,Nm}$. The paramodular group has a normalizing involution, the paramodular Fricke involution, given by $\mu_N = \begin{pmatrix} {}^t F_N^{-1} & 0 \\ 0 & F_N \end{pmatrix} \in \operatorname{Sp}(4, \mathbb{R})$, with $F_N = \frac{1}{\sqrt{N}} \begin{pmatrix} 0 & 1 \\ -N & 0 \end{pmatrix} \in \operatorname{SL}(2, \mathbb{R})$. Define the Fricke paramodular group $K(N)^+ = \langle K(N), \mu_N \rangle$. For $\epsilon \in \{\pm 1\}$, let $M_k(K(N))^{\epsilon} = \{f \in M_k(K(N)) : f|_k \mu_N = \epsilon f\}$ be the Fricke eigenspaces. There are graded rings, $M(K(N)) = \bigoplus_{k \in \mathbb{N}_0} M_k(K(N))$, $M_{\pm}(K(N)) = \bigoplus_{k \in \mathbb{N}_0, \epsilon \in \{\pm 1\}} M_k(K(N))^{\epsilon}$, as well as the graded ring of Fricke plus forms $M(K(N)^+) = \bigoplus_{k \in \mathbb{N}_0} M_k(K(N)^+)$.

•
$$N_{\infty}(\eta) = \{\tau \in \mathcal{H}_1 : \operatorname{Im}(\tau) > \eta\} \text{ for } \eta \ge 0.$$

•
$$P_{2,1}(\mathbb{Z}) = Sp(4,\mathbb{Z}) \cap \begin{pmatrix} \mathbb{Z} & 0 & \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & \mathbb{Z} & \mathbb{Z} & \mathbb{Z} \\ \mathbb{Z} & 0 & \mathbb{Z} & \mathbb{Z} \\ 0 & 0 & 0 & \mathbb{Z} \end{pmatrix}$$
.

3. Formal Series.

A paramodular form $f \in M_k(K(N))$ has a Fourier series $f(\Omega) = \sum_{t \in \bar{\mathcal{X}}(N)} a(t; f) e(\langle \Omega, t \rangle)$ that converges absolutely and uniformly on compact subsets of \mathcal{H}_2 . The absolute convergence allows rearrangement into a Fourier-Jacobi series $f(\Omega) = \sum_{m=0}^{\infty} \phi_m(\tau, z) e(Nm\omega)$ where each Fourier-Jacobi coefficient is a Jacobi form $\phi_m \in J_{k,Nm}$. This follows from the fact that the Fourier-Jacobi expansion is term-by-term invariant under $P_{2,1}(\mathbb{Z}) \subseteq K(N)$, see [18]. We define the formal Fourier-Jacobi series FJ(f) of a paramodular form f by

$$FJ: M_k(K(N)) \to \mathbb{M}(k, N) = \prod_{m=0}^{\infty} J_{k, Nm}$$
$$f \mapsto \sum_{m=0}^{\infty} \phi_m \xi^{Nm},$$

AOKI, IBUKIYAMA, POOR

where ξ is a place-holding variable. The Fourier coefficients of a paramodular form also satisfy symmetries determined by the following group:

$$\{\sigma \in \operatorname{GL}(2,\mathbb{R}) : \left(\begin{smallmatrix} \sigma & 0 \\ 0 & \sigma^* \end{smallmatrix}\right) \in K(N)\} = \langle \Gamma^0(N), \operatorname{diag}(1,-1) \rangle = \Gamma^0_{\pm}(N).$$

We call equation (2) below the $\Gamma^0(N)$ -symmetries,

(2)
$$\forall \sigma \in \Gamma^0_{\pm}(N), \forall t \in \bar{\mathcal{X}}(N), a(t[\sigma]; f) = \det(\sigma)^k a(t; f).$$

The Fourier coefficients of f are related to the Fourier coefficients of the Jacobi forms ϕ_m in the Fourier-Jacobi expansion of f by

$$a\left(\left(\begin{smallmatrix}n&r/2\\r/2&Nm\end{smallmatrix}\right);f\right)=c(n,r;\phi_m).$$

Accordingly, we define a subspace of $\mathbb{M}(k, N)$ that satisfies corresponding symmetries. Let $\Gamma \subseteq \Gamma^0_{\pm}(N)$ be a subgroup.

$$\mathbb{M}(k,N;\Gamma) = \{ \mathfrak{f} = \sum_{m=0}^{\infty} \phi_m \xi^{Nm} \in \mathbb{M}(k,N) : \mathfrak{f} \text{ satisfies equation (3)} \},$$

(3)
$$\forall \sigma \in \Gamma, \forall t_1 = \begin{pmatrix} n_1 & r_1/2 \\ r_1/2 & Nm_1 \end{pmatrix}, t_2 = \begin{pmatrix} n_2 & r_2/2 \\ r_2/2 & Nm_2 \end{pmatrix} \in \mathcal{X}(N),$$

if $t_1[\sigma] = t_2$ then $c(n_2, r_2; \phi_{m_2}) = \det(\sigma)^k c(n_1, r_1; \phi_{m_1}).$

Thus we have $FJ: M_k(K(N)) \to \mathbb{M}(k, N; \Gamma)$.

We write $\phi_m = \phi_m(\mathfrak{f})$ if we need to indicate the dependence of this Jacobi form on the formal series \mathfrak{f} . It is often helpful to reformulate the definition of $\mathbb{M}(k, N; \Gamma)$ by using the notation $a(t; \mathfrak{f}) = c(n, r; \phi_m(\mathfrak{f}))$ for $t = \binom{n - r/2}{r/2 - Nm} \in \bar{\mathcal{X}}(N)$. In this way equation (3) shortens to

(4)
$$\forall \sigma \in \Gamma, \forall t \in \bar{\mathcal{X}}(N), a(t[\sigma]; \mathfrak{f}) = \det(\sigma)^k a(t; \mathfrak{f})$$

We may sometimes avoid tracking boundary conditions in summations by setting $a(t; \mathfrak{f}) = 0$ for $t \in M_{2 \times 2}^{\text{sym}}(\mathbb{Q}) \setminus \overline{\mathcal{X}}(N)$.

Lemma 3.1. Let $\Gamma \subseteq \Gamma^0_{\pm}(N)$ be a subgroup. The Cauchy product gives $\mathbb{M}(N) = \bigoplus_{k=0}^{\infty} \mathbb{M}(k, N)$ the structure of a graded ring, and $\mathbb{M}(N; \Gamma) = \bigoplus_{k=0}^{\infty} \mathbb{M}(k, N; \Gamma)$ a graded subring. In particular, $\mathbb{M}(k_1, N)\mathbb{M}(k_2, N) \subseteq \mathbb{M}(k_1 + k_2, N)$, and $\mathbb{M}(k_1, N; \Gamma)\mathbb{M}(k_2, N; \Gamma) \subseteq \mathbb{M}(k_1 + k_2, N; \Gamma)$. The sets $\mathbb{S}(N) = \bigoplus_{k=0}^{\infty} \prod_{m=1}^{\infty} J_{k,Nm}^{\text{cusp}}$, and $\mathbb{S}(N; \Gamma) = \mathbb{M}(N; \Gamma) \cap \mathbb{S}(N)$ are graded ideals in $\mathbb{M}(N)$, and $\mathbb{M}(N; \Gamma)$, respectively. The Fourier-Jacobi map FJ : $M(K(N)) \to \mathbb{M}(N; \Gamma)$ is an injective ring homomorphism of graded rings that sends S(K(N)) to $\mathbb{S}(N; \Gamma)$, and the inverse image of $\mathbb{S}(N; \Gamma)$ is S(K(N)).

Proof. When the formal series $\mathfrak{f}_j \in \mathbb{M}(k_j, N)$, for j = 1, 2, are written as $\mathfrak{f}_j = \sum_{m=0}^{\infty} \phi_m(\mathfrak{f}_j) \xi^{Nm}$ then by the definition of the Cauchy product we have $\mathfrak{f}_1 \mathfrak{f}_2 = \sum_{m=0}^{\infty} \phi_m(\mathfrak{f}_1 \mathfrak{f}_2) \xi^{Nm}$ where

$$\phi_m(\mathfrak{f}_1\mathfrak{f}_2) = \sum_{m_1,m_2 \in \mathbb{N}_0: m_1+m_2=m} \phi_{m_1}(\mathfrak{f}_1)\phi_{m_2}(\mathfrak{f}_2).$$

Thus $\mathbb{M}(k_1, N)\mathbb{M}(k_2, N) \subseteq \mathbb{M}(k_1 + k_2, N)$ follows from the grading $J_{k_1,Nm_1}J_{k_2,Nm_2} \subseteq J_{k_1+k_2,N(m_1+m_2)}$ on Jacobi forms, which shows that $\mathbb{M}(N)$ is a graded ring. To derive a similar result for $\mathbb{M}(N;\Gamma)$ we reformulate the Cauchy product on $\mathbb{M}(N)$ as

(5)
$$\forall t \in \overline{\mathcal{X}}(N), \ a(t; \mathfrak{f}_1 \mathfrak{f}_2) = \sum_{t_1, t_2 \in \overline{\mathcal{X}}(N): \ t_1 + t_2 = t} a(t_1; \mathfrak{f}_1) a(t_2; \mathfrak{f}_2).$$

To see that (5) follows from the Cauchy product, note the following four equalities.

$$a(t;\mathfrak{f}_{1}\mathfrak{f}_{2}) = c(n,r;\phi_{m}(\mathfrak{f}_{1}\mathfrak{f}_{2})) = \sum_{\substack{m_{1}+m_{2}=m\\m_{1}+m_{2}=m}} c(n,r;\phi_{m_{1}}(\mathfrak{f}_{1})\phi_{m_{2}}(\mathfrak{f}_{2}))$$
$$= \sum_{\substack{m_{1},m_{2}\in\mathbb{N}_{0}:\\m_{1}+m_{2}=m\\m_{1}+m_{2}=m}} \sum_{\substack{n_{1},n_{2}\in\mathbb{N}_{0}:\\n_{1}+n_{2}=n\\m_{1}+n_{2}=n}} c(n_{1},r_{1};\phi_{m_{1}}(\mathfrak{f}_{1})) c(n_{2},r_{2};\phi_{m_{2}}(\mathfrak{f}_{2}))$$
$$= \sum_{\substack{t_{1}+t_{2}=t\\t_{1}+t_{2}=t}} a(t_{1};\mathfrak{f}_{1})a(t_{2};\mathfrak{f}_{2}).$$

Conversely, equation (5) implies the second equality, which is equivalent to the Cauchy product. If we assume $\mathfrak{f}_j \in \mathbb{M}(k_j, N; \Gamma)$ for j = 1, 2, then, for any $\sigma \in \Gamma$, $a(t[\sigma]; \mathfrak{f}_1 \mathfrak{f}_2) = \sum_{t_1+t_2=t[\sigma]} a(t_1; \mathfrak{f}_1)a(t_2; \mathfrak{f}_2)$. We use $\{(t_1, t_2) \in \overline{\mathcal{X}}(N)^2 : t_1+t_2=t[\sigma]\} = \{(s_1[\sigma], s_2[\sigma]) \in \overline{\mathcal{X}}(N)^2 : s_1+s_2=t\}$ to change the indices of summation,

$$a(t[\sigma]; \mathfrak{f}_{1}\mathfrak{f}_{2}) = \sum_{s_{1}+s_{2}=t} a(s_{1}[\sigma]; \mathfrak{f}_{1})a(s_{2}[\sigma]; \mathfrak{f}_{2})$$

$$= \sum_{s_{1}+s_{2}=t} \det(\sigma)^{k_{1}}a(s_{1}; \mathfrak{f}_{1})\det(\sigma)^{k_{2}}a(s_{2}; \mathfrak{f}_{2})$$

$$= \det(\sigma)^{k_{1}+k_{2}}a(t; \mathfrak{f}_{1}\mathfrak{f}_{2}).$$

Thus $\mathbb{M}(k_1, N; \Gamma)\mathbb{M}(k_2, N; \Gamma) \subseteq \mathbb{M}(k_1 + k_2, N; \Gamma)$ and $\mathbb{M}(N; \Gamma)$ is a graded ring, noting that a sum of paramodular forms of distinct weights is zero as a holomorphic function if and only if each summand is zero as a holomorphic function. The sets $\mathbb{S}(N)$, and $\mathbb{S}(N; \Gamma)$ are graded ideals simply because $J_{k_1,Nm_1}^{\text{cusp}} J_{k_2,Nm_2} \subseteq J_{k_1+k_2,N(m_1+m_2)}^{\text{cusp}}$. The convergence of the Fourier-Jacobi expansion shows that FJ is injective on

each graded piece and hence on M(K(N)). The absolute convergence of the Fourier-Jacobi expansion for $f_j \in M_{k_j}(K(N))$ proves that FJ : $M(K(N)) \to \mathbb{M}(N;\Gamma)$ is a homomorphism because for $\Omega = \begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} \in \mathcal{H}_2$,

$$f_1(\Omega)f_2(\Omega) = \sum_{m=0}^{\infty} \left(\sum_{m_1+m_2=m} \phi_{m_1}(\mathrm{FJ}(f_1))(\tau, z) \phi_{m_2}(\mathrm{FJ}(f_2))(\tau, z)\right) e(Nm\omega),$$

and therefore

$$FJ(f_1f_2) = \sum_{m=0}^{\infty} \left(\sum_{m_1+m_2=m} \phi_{m_1}(FJ(f_1)) \phi_{m_2}(FJ(f_2)) \right) \xi^{Nm}$$
$$= FJ(f_1)FJ(f_2).$$

For a cusp form $f \in S(K(N))$, the Siegel Φ map gives the containment $\operatorname{supp}(f) = \{t \in \overline{\mathcal{X}}(N) : a(t; f) \neq 0\} \subseteq \mathcal{X}(N)$. By examining the support of ϕ_m ,

$$supp(\phi_m) = \{(n,r) \in \mathbb{N}_0 \times \mathbb{Z} : c(n,r;\phi_m) \neq 0\}$$
$$= \{(n,r) \in \mathbb{N}_0 \times \mathbb{Z} : a\left(\binom{n}{r/2} Nm}{r/2}; f\right) \neq 0\}$$
$$\subseteq \{(n,r) \in \mathbb{N} \times \mathbb{Z} : 4Nmn - r^2 > 0\},$$

we see that $\phi_m \in J_{k,Nm}^{\text{cusp}}$, and that $\operatorname{FJ}(f) \in \mathbb{S}(N)$, from which $\operatorname{FJ}(f) \in \mathbb{S}(N; \Gamma)$ follows. The assertion that $\{f \in M_k(K(N)) : \operatorname{FJ}(f) \in \mathbb{S}(N; \Gamma)\} = S_k(K(N))$ relies on special properties of the paramodular group. For general subgroups commensurable with $\operatorname{Sp}(4, \mathbb{Z})$ it is not true that the Fourier-Jacobi expansion of a Siegel modular form with coefficients that are all cusp forms necessarily comes from a Siegel modular form because Reefschläger's double coset decomposition [34] of $K(N) \setminus \operatorname{Sp}(4, \mathbb{Q}) / \operatorname{P}_{2,1}(\mathbb{Q})$ has representatives of the form $\begin{pmatrix} u & 0 \\ 0 & u^* \end{pmatrix}$ with $u \in \operatorname{GL}(2, \mathbb{Q})$; see also Corollary 2.5 of Gritsenko [18].

The involution $\mu_N = \begin{pmatrix} F_N^* & 0 \\ 0 & F_N \end{pmatrix}$ decomposes paramodular forms into plus and minus forms $M_k(K(N)) = M_k(K(N))^+ \oplus M_k(K(N))^-$ where $M_k(K(N))^{\epsilon} = \{f \in M_k(K(N)) : f|_k \mu_N = \epsilon f\}$. The action of F_N^* on $\bar{\mathcal{X}}(N)$ is $F_N^{-1}tF_N^* = \begin{pmatrix} m & -r/2 \\ -r/2 & Nn \end{pmatrix}$ if $t = \begin{pmatrix} n & r/2 \\ r/2 & Nm \end{pmatrix} \in \bar{\mathcal{X}}(N)$. Therefore the Fourier coefficients of a Fricke eigenform $f \in M_k(K(N))^{\epsilon}$ satisfy the *involution condition*

(6)
$$\forall t \in \bar{\mathcal{X}}(N), \ a\left(F_N^{-1}tF_N^*; f\right) = \epsilon \ a\left(t; f\right).$$

Accordingly, we define a subspace of formal series satisfying the corresponding involution condition

$$\mathbb{M}(k, N, \epsilon) = \{ \mathfrak{f} = \sum_{m=0}^{\infty} \phi_m \xi^{Nm} \in \mathbb{M}(k, N) : \mathfrak{f} \text{ satisfies equation (7)} \},\$$

$$(7) \qquad \forall \begin{pmatrix} n & r/2 \\ r/2 & Nm \end{pmatrix} \in \bar{\mathcal{X}}(N), \ c(n, r; \phi_m) = (-1)^k \epsilon \, c(m, r; \phi_n).$$

The map FJ : $M_k(K(N))^{\epsilon} \to \mathbb{M}(k, N, \epsilon)$ injects. It is helpful to rewrite equation (7) as $\forall t \in \overline{\mathcal{X}}(N)$, $a(t[F_N^*]; \mathfrak{f}) = (-1)^k \epsilon a(t; \mathfrak{f})$. If we write $\mathbb{M}(k, N, \epsilon; \Gamma) = \mathbb{M}(k, N; \Gamma) \cap \mathbb{M}(k, N, \epsilon)$ we also have FJ : $M_k(K(N))^{\epsilon} \to \mathbb{M}(k, N, \epsilon; \Gamma)$. We set $\mathbb{S}(k, N, \epsilon) = \mathbb{M}(k, N, \epsilon) \cap \mathbb{S}(k, N)$, and the more frequently used $\mathbb{S}(k, N, \epsilon; \Gamma) = \mathbb{M}(k, N, \epsilon; \Gamma) \cap \mathbb{S}(k, N)$.

Lemma 3.2. Let $\Gamma \subseteq \Gamma^0_{\pm}(N)$ be a subgroup. The Cauchy product gives $\mathbb{M}_{\pm}(N) = \bigoplus_{k,\epsilon} \mathbb{M}(k, N, \epsilon)$, $\mathbb{M}_{\pm}(N; \Gamma) = \bigoplus_{k,\epsilon} \mathbb{M}(k, N, \epsilon; \Gamma)$, and $\mathbb{M}(N, +) = \bigoplus_{k=0}^{\infty} \mathbb{M}(k, N, +)$ the structure of graded rings. In particular, we have $\mathbb{M}(k_1, N, \epsilon_1)\mathbb{M}(k_2, N, \epsilon_2) \subseteq \mathbb{M}(k_1 + k_2, N, \epsilon_1\epsilon_2)$, and $\mathbb{M}(k_1, N, \epsilon_1; \Gamma)\mathbb{M}(k_2, N, \epsilon_2; \Gamma) \subseteq \mathbb{M}(k_1 + k_2, N, \epsilon_1\epsilon_2; \Gamma)$. The subsets $\mathbb{S}_{\pm}(N) = \bigoplus_{k,\epsilon} \mathbb{S}(k, N, \epsilon)$, and $\mathbb{S}_{\pm}(N; \Gamma) = \mathbb{M}_{\pm}(N; \Gamma) \cap \mathbb{S}_{\pm}(N)$ are graded ideals in $\mathbb{M}_{\pm}(N)$ and $\mathbb{M}_{\pm}(N; \Gamma)$, respectively. The Fourier-Jacobi expansion map FJ : $M_{\pm}(K(N)) \to \mathbb{M}_{\pm}(N; \Gamma)$ is an injective homomorphism of graded rings that sends $S_{\pm}(K(N))$ to $\mathbb{S}_{\pm}(N; \Gamma)$, and the inverse image of $\mathbb{S}_{\pm}(N; \Gamma)$ is $S_{\pm}(K(N))$.

Proof. The proof is very similar to that of Lemma 3.1.

Lemma 3.3. Let $N \in \mathbb{N}$. The graded ring of formal series of Jacobi forms, $\mathbb{M}(N) = \bigoplus_{k=0}^{\infty} \mathbb{M}(k, N)$, is an integral domain.

Proof. An element in $\mathbb{M}(N)$ is zero precisely when each graded piece is zero. If a product of nonzero elements from $\mathbb{M}(N)$ is zero then the product of the nonzero graded pieces of highest weight in each factor must be zero. Each nonzero element in each graded piece has a leading term $\phi_m \in J_{k,Nm} \subseteq \mathcal{O}(\mathcal{H}_1 \times \mathbb{C})$. The product of these leading terms cannot be zero because the ring $\mathcal{O}(\mathcal{H}_1 \times \mathbb{C})$ is an integral domain. \Box

4. VANISHING ORDER OF JACOBI FORMS.

In Corollary 3.1 of [2], Aoki significantly improved the known bounds on vanishing orders of Jacobi forms [11, 16], which were O(m) in the index m for a fixed weight k. He proved that a nontrivial Jacobi form of weight k and index m cannot have a vanishing order ν greater than $\frac{k+1}{6}(\sqrt{2m+1}+1)$. As a consequence the \mathbb{C} -vector spaces of formal series $\mathbb{M}(k, N, \epsilon)$ are finite dimensional. In Theorem 4.3 we state Aoki's main theorem from [2] in the case of even weights and use it to improve the estimate concerning when $J_{k,Nm}(m) = \{0\}$ enough to get the desired bound on the growth in k for dimensions of spaces of formal series of Jacobi forms, namely, dim $\mathbb{M}(k, N, \epsilon) \in O(N^3 k^3)$ for $N, k \in \mathbb{N}$.

Definition 4.1. For $j \in \mathbb{N}$ we define $\psi_j : \mathbb{N} \to \mathbb{Q}^+$ by

$$\psi_j(u) = \begin{cases} 1 & \text{if } u = 1, \\ \prod_{p \mid u: p \text{ prime}} \left(1 - \frac{1}{p^j} \right) & \text{if } u \ge 2. \end{cases}$$

It is clear that $a \mid t$ implies $\psi_j(a) \ge \psi_j(t)$.

Definition 4.2. We define $\psi : \mathbb{N} \setminus \{1\} \to \mathbb{Q}^+$ by

$$\psi(t) = \begin{cases} t^2 \psi_2(t) = 3 & \text{if } t = 2, \\ \frac{1}{2} t^2 \psi_2(t) = \frac{1}{2} t^2 \prod_{p|t: \ p \text{ prime}} \left(1 - \frac{1}{p^2} \right) & \text{if } t \ge 3. \end{cases}$$

It is easy to check that $\psi(t) \ge 3$ for $t \ge 2$.

Theorem 4.3 ([2]). Let $k \in 2\mathbb{N}$, $m \in \mathbb{N}$. Let $\phi \in J_{k,m}$ have $\operatorname{ord}(\phi) = \mu$. We have

$$\min\left(m, m - 6\mu + \frac{k}{2}\right) \ge \sum_{t:(\clubsuit)} \psi(t),$$

where t runs over all natural numbers satisfying the condition

$$t \neq 1 \text{ and } \sum_{c=0}^{t-1} \psi_1\left(\gcd(t,c)\right) \max\left(\mu - \frac{mc(t-c)}{t^2}, 0\right) > \frac{kt}{12}\psi_2(t).$$
 (\blacklozenge)

Lemma 4.4. Let $t, \mu, m \in \mathbb{N}$. If $\mu < m$, we have

$$\sum_{c=0}^{t-1} \max\left(\mu - \frac{mc(t-c)}{t^2}, 0\right) > \frac{\mu^2 t}{2m}$$

Proof. The condition $\mu < m$ is used in the first equality.

$$\sum_{c=0}^{t-1} \max\left(\mu - \frac{mc(t-c)}{t^2}, 0\right) \ge \sum_{c=0}^{t-1} \max\left(\mu - \frac{mc}{t}, 0\right)$$
$$= \sum_{c=0}^{\operatorname{floor}\left(\frac{\mu t}{m}\right)} \left(\mu - \frac{mc}{t}\right) = \left(\mu - \frac{m}{2t}\operatorname{floor}\left(\frac{\mu t}{m}\right)\right) \left(\operatorname{floor}\left(\frac{\mu t}{m}\right) + 1\right)$$
$$\ge \left(\mu - \frac{\mu}{2}\right) \left(\operatorname{floor}\left(\frac{\mu t}{m}\right) + 1\right) = \frac{\mu}{2} \left(\operatorname{floor}\left(\frac{\mu t}{m}\right) + 1\right) > \frac{\mu^2 t}{2m}.$$

Proposition 4.5. Let $k, N, \nu \in \mathbb{N}$. If $\nu > \frac{1}{6}Nk$ then $J_{k,N\nu}(\nu) = \{0\}$.

Proof. The case of odd k follows from that of even k. If $\phi \in J_{k,N\nu}(\nu)$ then $\phi^2 \in J_{2k,N(2\nu)}(2\nu)$. Since $\nu > \frac{1}{6}Nk$ we have $2\nu > \frac{1}{6}N(2k)$ and so $\phi^2 = 0$ assuming the result for even weights; hence $\phi = 0$.

Suppose that there is a nontrivial $\varphi \in J_{k,N\nu}(\nu)$ with vanishing order $\mu = \operatorname{ord} \varphi \geq \nu > \frac{1}{6}Nk$ and k even. We will obtain a contradiction to Theorem 4.3. For $m = N\nu$, we will contradict

$$\min\left(m, m - 6\mu + \frac{k}{2}\right) \ge \sum_{t:(\spadesuit)} \psi(t).$$

Since ψ has a positive minimum, it suffices to show that the set of positive integers satisfying (\blacklozenge) is infinite.

We show that all sufficiently large primes p satisfy

(**(**)
$$\sum_{c=0}^{p-1} \psi_1 \left(\gcd(p,c) \right) \max \left(\mu - \frac{mc(p-c)}{p^2}, 0 \right) > \frac{kp}{12} \psi_2(p).$$

We know that $a \mid t$ implies $\psi_1(a) \ge \psi_1(t)$. Since $gcd(p,c) \mid p$, we have $\psi_1(gcd(p,c)) \ge \psi_1(p)$. Therefore

$$\sum_{c=0}^{p-1} \psi_1\left(\gcd(p,c)\right) \max\left(\mu - \frac{mc(p-c)}{p^2}, 0\right)$$
$$\geq \psi_1(p) \sum_{c=0}^{p-1} \max\left(\mu - \frac{mc(p-c)}{p^2}, 0\right).$$

Lemma 4.4 says that for $t, m, \mu \in \mathbb{N}$ with $\mu < m$ we have

$$\sum_{c=0}^{t-1} \max\left(\mu - \frac{mc(t-c)}{t^2}, 0\right) > \frac{\mu^2 t}{2m}.$$

We use the linear bound $\operatorname{ord} \varphi \leq \frac{k+2m}{12}$ of [16], Proposition 3.2, to check the hypothesis $\mu < m$:

$$\mu = \operatorname{ord} \varphi \le \frac{k + 2m}{12} < \frac{\frac{6\nu}{N} + 2m}{12} = \frac{\frac{6}{N^2} + 2}{12} m < m.$$

Thus by Lemma 4.4 we have

$$\sum_{c=0}^{p-1} \psi_1\left(\gcd(p,c)\right) \max\left(\mu - \frac{mc(p-c)}{p^2}, 0\right)$$
$$> \psi_1(p) \frac{\mu^2 p}{2m} \ge \psi_1(p) \frac{\nu^2 p}{2m} = \psi_1(p) \frac{\nu p}{2N}.$$

Thus a sufficient condition for a prime p to satisfy (\spadesuit) is

$$\psi_1(p)\frac{\nu p}{2N} > \frac{kp}{12}\psi_2(p).$$

This simplifies to $\nu > \frac{Nk}{6} \left(1 + \frac{1}{p}\right)$, or $\nu - \frac{1}{6}Nk > \frac{Nk}{6}\frac{1}{p}$, which is true for all sufficiently large p because $\nu - \frac{1}{6}Nk$ is positive.

Corollary 4.6. For $k, N \in \mathbb{N}$ we have dim $\mathbb{M}(k, N, \epsilon) \in O(N^3 k^3)$.

Proof. We have dim $\mathbb{M}(k, N, \epsilon) \leq \sum_{j=0}^{\infty} \dim J_{k,Nj}(j+\delta)$, where δ is 0 or 1 as $(-1)^k \epsilon$ is 1 or -1, by Lemma 3.2 of [26]. By Proposition 4.5 we may cap the summation at floor $(\frac{1}{6}Nk)$ and, by Theorem 2.3 in [11], the codimension of $J_{k,m}^{\text{cusp}}$ in $J_{k,m}$ is at most floor(b/2) + 1 where b is the largest integer with $b^2 \mid m$. Therefore

$$\dim \mathbb{M}(k, N, \epsilon) \leq \sum_{j=0}^{\text{floor}(\frac{1}{6}Nk)} \dim J_{k,Nj}(j) \leq \sum_{j=0}^{\text{floor}(\frac{1}{6}Nk)} \dim J_{k,Nj}$$
$$\leq \dim J_{k,0} + \sum_{j=1}^{\text{floor}(\frac{1}{6}Nk)} \dim J_{k,Nj}^{\text{cusp}} + \frac{1}{2}\sqrt{Nj} + 1$$
$$\leq \frac{k+12}{12} + \frac{2}{3} \left(\frac{1}{6}Nk + 1\right)^{3/2} + \frac{1}{6}Nk + \sum_{j=1}^{\text{floor}(\frac{1}{6}Nk)} \dim J_{k,Nj}^{\text{cusp}}$$

It is known that for $k, m \in \mathbb{N}$, dim $J_{k,m}^{\text{cusp}} \in O(km)$. An easy estimate from the dimension formula [11] is dim $J_{k,m}^{\text{cusp}} \leq \frac{m+1}{24}(2k+35)$. From dim $J_{k,Nj}^{\text{cusp}} \in O(kNj)$ we have $\sum_{j=1}^{\text{floor}(\frac{1}{6}Nk)} \dim J_{k,Nj}^{\text{cusp}} \in O(N^3k^3)$.

Proposition 4.7. The integral domain $\mathbb{M}(N, +) = \bigoplus_{k=0}^{\infty} \mathbb{M}(k, N, +)$ is algebraic over its subring FJ $(M(K(N)^+)) = \bigoplus_{k=0}^{\infty} \text{FJ}(M_k(K(N)^+))$. For $k \in \mathbb{N}_0$, each $\mathfrak{f} \in \mathbb{M}(k, N, +)$ satisfies a polynomial relation of the type

$$\mathrm{FJ}(f_0)\mathfrak{f}^d + \mathrm{FJ}(f_1)\mathfrak{f}^{d-1} + \dots + \mathrm{FJ}(f_j)\mathfrak{f}^{d-j} + \dots + \mathrm{FJ}(f_d) = 0,$$

for some $d \in \mathbb{N}$, $k_0 \in \mathbb{N}_0$, and some $f_j \in M_{k_0+jk}(K(N)^+)$ with f_0 not identically zero.

Proof. It suffices to prove the second statement. The group $\Gamma = K(N)^+$ is commensurable with Sp(4, Z), so by Theorem 6.11 in Freitag [12], the homogeneous quotient field of $M(\Gamma)$, $\mathcal{K}(\Gamma)$, has transcendence degree three. Take three meromorphic functions $f_1, f_2, f_3 \in \mathcal{K}(\Gamma)$ that are algebraically independent over \mathbb{C} and select a common denominator $D \in M_{k^*}(\Gamma)$. We must have $k^* \in \mathbb{N}$ because the f_j are not

constant. We obtain four paramodular forms $g_1 = Df_1$, $g_2 = Df_2$, $g_3 = Df_3$, and $g_4 = D$ in $M_{k^*}(\Gamma)$ that are algebraically independent over \mathbb{C} . This follows because we may reduce to the case where any putative polynomial relation is homogeneous in the g_j .

Take $\mathfrak{f} \in \mathbb{M}(k, N, +)$. We may assume that \mathfrak{f} is nontrivial because otherwise $f^1 = 0$ satisfies the conclusion with $k_0 = 0, f_0 = 1 \in M_0(\Gamma)$, and $f_1 = 0 \in M_k(\Gamma)$. Similarly, if k = 0 then $\mathfrak{f} = c$ is constant and $\mathfrak{f}^1 - c = 0$ satisfies the conclusion with $k_0 = 0, f_0 = 1 \in M_0(\Gamma)$, and $f_1 = -c \in M_0(\Gamma)$. So we assume k > 0 as well. The four formal series $\mathrm{FJ}(g_i^k) \in \mathbb{M}(k^*k, N, +)$ are algebraically independent over \mathbb{C} because FJ is a monomorphism. Consider the list of five formal series $\mathrm{FJ}(g_i^k), \, \mathfrak{f}^{k^*} \in \mathbb{M}(k^*k, N, +).$ For any $\mu \in \mathbb{N}$, there are $\binom{\mu+4}{4}$ distinct monomials $x_1^{i_1} \cdots x_5^{i_5}$ in five variables with $i_1 + \cdots + i_5 = \mu$. By substitution of the five formal series into these monomials, we have $\binom{\mu+4}{4}$ elements $\operatorname{FJ}(g_1^{ki_1})\cdots\operatorname{FJ}(g_4^{ki_4})\mathfrak{f}^{k^*i_5} \in \mathbb{M}(\mu k^*k, N, +)$. By Corollary 4.6, however, we have dim $\mathbb{M}(\mu k^* k, N, +) \in O((Nk^*k)^3 \mu^3)$, so for sufficiently large μ there is a nontrivial \mathbb{C} -linear dependence relation among these $\binom{\mu+4}{4}$ elements. At least one supported monomial in the dependence relation must contain a positive power of f because the remaining four formal series are algebraically independent. For the same reason, when a positive power of f is supported then its coefficient, after collecting like terms in powers of \mathfrak{f} , must be nontrivial. If $d \in \mathbb{N}$ is the highest power of \mathfrak{f} that is supported in the dependence relation, then we may write this relation as $\sum_{j=0}^{d} \operatorname{FJ}(f_j) f^{d-j} = 0$ where the f_j are \mathbb{C} linear combinations of monomials in the four g_1, \ldots, g_4 . Since f^d is supported, we have f_0 nontrivial. Let the weight of f_0 be $k_0 \in \mathbb{N}_0$ so that $f_0 \in M_{k_0}(\Gamma)$. The terms $\mathrm{FJ}(f_j)\mathfrak{f}^{d-j}$ all have the same weight, $k_0 + kd$, which is the weight of $\mathrm{FJ}(f_0)\mathfrak{f}^d$. Therefore we may take $f_j \in M_{k_0+k_j}(\Gamma)$, as required. \square

5. Invariance under subgroups of finite index in $\Gamma^0(N)$.

In Proposition 4.7 of the previous section we saw that a formal series of Jacobi forms \mathfrak{f} possessing the involution condition for $\epsilon = +1$ satisfies a polynomial P(X) = 0 whose coefficients are formal Fourier-Jacobi expansions of paramodular forms. As a consequence of this polynomial relation we will show in this section that \mathfrak{f} is invariant under a subgroup Γ of finite index in $\Gamma^0(N)$. These arguments best take place inside the ring of formal Fourier series.

The ring structure on the ring of formal Fourier series, $\mathbb{C}^{\bar{\mathcal{X}}(N)}$, is defined by the Cauchy product, noting that for every $t \in \bar{\mathcal{X}}(N)$ the set

 $\{(t_1, t_2) \in \overline{\mathcal{X}}(N) \times \overline{\mathcal{X}}(N) : t_1 + t_2 = t\}$ is finite. We accordingly use a place-holding variable q to write an element $\psi \in \mathbb{C}^{\overline{\mathcal{X}}(N)}$ as

$$\psi = \sum_{t \in \bar{\mathcal{X}}(N)} a(t; \psi) q^t$$

The ring of formal Fourier series, $\mathbb{C}^{\bar{\mathcal{X}}(N)}$ is also an integral domain, which will be proven in Corollary 5.2. The Fourier expansion of a paramodular form defines a map

$$FS: M_k(K(N)) \to \mathbb{C}^{\mathcal{X}(N)}$$
$$f \mapsto \sum_{t \in \bar{\mathcal{X}}(N)} a(t; f) q^t.$$

Given a formal series of Jacobi forms $\mathfrak{f} \in \mathbb{M}(k, N)$, we may define the associated formal Fourier series, $AFS(\mathfrak{f}) \in \mathbb{C}^{\bar{\mathcal{X}}(N)}$, by

$$AFS: \mathbb{M}(k, N) \to \mathbb{C}^{\mathcal{X}(N)}$$
$$\mathfrak{f} = \sum_{m=0}^{\infty} \phi_m \xi^{Nm} \mapsto \sum_{t = \binom{n-r/2}{r/2} \in \bar{\mathcal{X}}(N)} c(n, r; \phi_m) q^t = \sum_{t \in \bar{\mathcal{X}}(N)} a(t; \mathfrak{f}) q^t.$$

Extending by linearity we have a map AFS : $\mathbb{M}(N) \to \mathbb{C}^{\bar{\mathcal{X}}(N)}$ from the ring $\mathbb{M}(N) = \bigoplus_{k \in \mathbb{N}_0} \mathbb{M}(k, N)$. That AFS is a ring homomorphism is an exercise using the Cauchy product similar to the computations demonstrating equation (5). For $f \in M_k(K(N))$ we have the compatibility AFS (FJ(f)) = FS(f).

Formal Fourier series share some properties with formal power series $\mathbb{C}[[a, b, c]]$ in three variables due to the following monomorphism that sends q^t to the monomial $a^{\langle s_1, t \rangle} b^{\langle s_2, t \rangle} c^{\langle s_3, t \rangle}$.

Lemma 5.1. Set
$$s_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$
, $s_2 = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$, $s_3 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$. The map
 $\iota : \mathbb{C}^{\bar{\mathcal{X}}(N)} \to \mathbb{C}[[a, b, c]]$
 $\psi = \sum_{t \in \bar{\mathcal{X}}(N)} a(t; \psi) q^t \mapsto \sum_{i, j, k \in \mathbb{N}_0} \left(\sum_{\substack{t \in \bar{\mathcal{X}}(N):\\\langle s_1, t \rangle = i, \langle s_2, t \rangle = j, \langle s_3, t \rangle = k}} a(t; \psi) \right) a^i b^j c^k$

is a ring homomorphism satisfying the following properties.

(1) $\iota(\psi) = \sum_{i,j,k\in\mathbb{N}_0} a\left(\left(\frac{i}{\frac{j-i-k}{2}},\psi\right);\psi\right) a^i b^j c^k,$ (2) $\iota(\psi) = \sum_{t\in\bar{\mathcal{X}}(N)} a\left(t;\psi\right) a^{\langle s_1,t\rangle} b^{\langle s_2,t\rangle} c^{\langle s_3,t\rangle},$ (3) ι is injective.

Proof. Setting $t = \binom{n-r/2}{r/2 Nm} \in \bar{\mathcal{X}}(N)$ and solving the system of linear equations $\langle s_1, t \rangle = i$, $\langle s_2, t \rangle = j$, and $\langle s_3, t \rangle = k$, the unique solution is $n = i, r = \frac{j-i-k}{2}$, and Nm = k. This proves formula (1) if we understand that $a(t; \psi) = 0$ for $t \notin \bar{\mathcal{X}}(N)$. Formula (2) follows from (1) since rearrangements of formal power series are equal. The ring homomorphism property is then formal because the $\langle s_\ell, t \rangle$ are linear in t. The injectivity (3) follows from formula (1).

Corollary 5.2. The ring $\mathbb{C}^{\bar{\mathcal{X}}(N)}$ is an integral domain.

Proof. A ring with a monomorphism to an integral domain is an integral domain. The monomorphism here is $\iota : \mathbb{C}^{\bar{\mathcal{X}}(N)} \to \mathbb{C}[[a, b, c]]$. \Box

There is a copy of $\Gamma^0(N)$ inside the group of automorphisms of formal Fourier series.

Lemma 5.3. For $\sigma \in \Gamma^0(N)$ define

$$\mathbf{j}(\sigma): \mathbb{C}^{\bar{\mathcal{X}}(N)} \to \mathbb{C}^{\bar{\mathcal{X}}(N)}$$
$$\sum_{t \in \bar{\mathcal{X}}(N)} a(t; \psi) q^t \mapsto \sum_{t \in \bar{\mathcal{X}}(N)} a(t[\sigma]; \psi) q^t.$$

The map $j(\sigma)$ is an automorphism of $\mathbb{C}^{\bar{\mathcal{X}}(N)}$ and the map

$$\mathbf{j}: \Gamma^0(N) \to \operatorname{Aut}\left(\mathbb{C}^{\bar{\mathcal{X}}(N)}\right)$$

 $\sigma \mapsto \mathbf{j}(\sigma)$

is a homomorphism. For $\mathfrak{f} \in \mathbb{M}(k, N)$, we have $\mathfrak{j}(\sigma)AFS(\mathfrak{f}) = AFS(\mathfrak{f})$ if and only if $a(t[\sigma]; \mathfrak{f}) = a(t; \mathfrak{f})$ for all $t \in \overline{\mathcal{X}}(N)$. For $f \in M_k(K(N))$ we have $\mathfrak{j}(\sigma)FS(f) = FS(f)$ for all $\sigma \in \Gamma^0(N)$.

Proof. We show $j(\sigma)$ is an automorphism. The map $j(\sigma)$ has $j(\sigma^{-1})$ as an inverse and the additivity of $j(\sigma)$ is clear so that it suffices to prove $j(\sigma) (\psi_1 \psi_2) = (j(\sigma)\psi_1) (j(\sigma)\psi_2)$. We have

$$j(\sigma) (\psi_1 \psi_2) = j(\sigma) \sum_{t \in \bar{\mathcal{X}}(N)} \left(\sum_{t_1, t_2 \in \bar{\mathcal{X}}(N): t_1 + t_2 = t} a(t_1; \psi_1) a(t_2; \psi_2) \right) q^t$$
$$= \sum_{t \in \bar{\mathcal{X}}(N)} \left(\sum_{t_1, t_2 \in \bar{\mathcal{X}}(N): t_1 + t_2 = t[\sigma]} a(t_1; \psi_1) a(t_2; \psi_2) \right) q^t.$$

We use the equality $\{(t_1, t_2) \in \overline{\mathcal{X}}(N) \times \overline{\mathcal{X}}(N) : t_1 + t_2 = t[\sigma]\} = \{(s_1[\sigma], s_2[\sigma]) \in \overline{\mathcal{X}}(N) \times \overline{\mathcal{X}}(N) : s_1 + s_2 = t\}$ to change the index of

summation.

$$j(\sigma) (\psi_1 \psi_2) = \sum_{t \in \bar{\mathcal{X}}(N)} \left(\sum_{s_1, s_2 \in \bar{\mathcal{X}}(N): s_1 + s_2 = t} a(s_1[\sigma]; \psi_1) a(s_2[\sigma]; \psi_2) \right) q^t$$
$$= \sum_{s_1 \in \bar{\mathcal{X}}(N)} a(s_1[\sigma]; \psi_1) q^{s_1} \sum_{s_2 \in \bar{\mathcal{X}}(N)} a(s_2[\sigma]; \psi_2) q^{s_2}$$
$$= (j(\sigma)\psi_1) (j(\sigma)\psi_2).$$

We show that $j : \Gamma^0(N) \to \operatorname{Aut} \left(\mathbb{C}^{\bar{\mathcal{X}}(N)}\right)$ is a homomorphism. Take $\sigma_1, \sigma_2 \in \Gamma^0(N)$ and $\psi = \sum_{t \in \bar{\mathcal{X}}(N)} a(t; \psi) q^t \in \mathbb{C}^{\bar{\mathcal{X}}(N)}$. We have

$$\mathbf{j}(\sigma_2)\psi = \sum_t a(t[\sigma_2];\psi)q^t$$
$$(\mathbf{j}(\sigma_1)\mathbf{j}(\sigma_2))\psi = \mathbf{j}(\sigma_1)(\mathbf{j}(\sigma_2)\psi) = \sum_t a(t[\sigma_1][\sigma_2];\psi)q^t = \mathbf{j}(\sigma_1\sigma_2)\psi.$$

For $\mathfrak{f} = \sum_{m=0}^{\infty} \phi_m \xi^{Nm} \in \mathbb{M}(k, N)$ we have $\operatorname{AFS}(\mathfrak{f}) = \sum_t a(t; \mathfrak{f})q^t$ where $a(t; \mathfrak{f}) = c(n, r; \phi_m)$ for $t = \binom{n r/2}{r/2 Nm} \in \overline{\mathcal{X}}(N)$. We then have $\mathfrak{j}(\sigma)\operatorname{AFS}(\mathfrak{f}) = \sum_t a(t[\sigma]; \mathfrak{f})q^t$ and, by definition of formal series, this equals $\operatorname{AFS}(\mathfrak{f})$ if and only if $a(t[\sigma]; \mathfrak{f}) = a(t; \mathfrak{f})$ for all $t \in \overline{\mathcal{X}}(N)$. The final assertion, $\mathfrak{j}(\sigma)\operatorname{FS}(f) = \operatorname{FS}(f)$ for all $\sigma \in \Gamma^0(N)$, follows from the $\Gamma^0(N)$ -symmetries of $f \in M_k(K(N))$ in equation (2). \Box

The next proposition shows that a formal series of Jacobi forms satisfying the involution condition necessarily has additional symmetries.

Proposition 5.4. Let $\mathfrak{f} \in \mathbb{M}(k, N, +)$. There is a subgroup Γ of finite index in $\Gamma^0(N)$ such that $\mathfrak{f} \in \mathbb{M}(k, N, +; \Gamma)$.

Proof. By Proposition 4.7, \mathfrak{f} satisfies a polynomial relation of the type

$$\mathrm{FJ}(f_0)\mathfrak{f}^d + \dots + \mathrm{FJ}(f_j)\mathfrak{f}^{d-j} + \dots + \mathrm{FJ}(f_d) = 0,$$

for some $d \in \mathbb{N}$, $k_0 \in \mathbb{N}_0$, and some $f_j \in M_{k_0+jk}(K(N)^+)$ with f_0 not identically zero. Apply the monomorphism AFS : $\mathbb{M}(N) \to \mathbb{C}^{\bar{\mathcal{X}}(N)}$ to obtain

$$FS(f_0)(AFS(\mathfrak{f}))^d + \dots + FS(f_j)(AFS(\mathfrak{f}))^{d-j} + \dots + FS(f_d) = 0.$$

The polynomial $\sum_{j=0}^{d} \operatorname{FS}(f_j) X^{d-j} \in \mathbb{C}^{\bar{\mathcal{X}}(N)}[X]$ has a root $\operatorname{AFS}(\mathfrak{f})$. Noting by Lemma 5.3 that $\mathfrak{j}(\sigma) \operatorname{FS}(f_j) = \operatorname{FS}(f_j)$ for every $\sigma \in \Gamma^0(N)$, each element of the orbit $\mathfrak{j}(\Gamma^0(N)) \operatorname{AFS}(\mathfrak{f})$ is a root. However, the ring $\mathbb{C}^{\bar{\mathcal{X}}(N)}$ is an integral domain and a polynomial of positive degree d over an integral domain has at most d roots. If we want to be definite we can find $u_1, \ldots, u_{d_1} \in \Gamma^0(N)$ with $d_1 \leq d$ such that $\mathfrak{j}(\Gamma^0(N)) \operatorname{AFS}(\mathfrak{f}) =$

 $\{j(u_1)AFS(\mathfrak{f}),\ldots,j(u_{d_1})AFS(\mathfrak{f})\}$. The natural homomorphism ρ from $\Gamma^0(N)$ to permutations of the orbit $j(\Gamma^0(N))AFS(\mathfrak{f})$ is specified in this labeling by $j(\sigma)j(u_i)AFS(\mathfrak{f}) = j(u_{\rho(\sigma)i})AFS(\mathfrak{f})$ for $i = 1,\ldots,d_1$. Let $\Gamma = \ker(\rho)$ be the kernel of $\rho : \Gamma^0(N) \to S_{d_1}$. Then Γ is a normal subgroup of $\Gamma^0(N)$ of index at most d_1 !. For every $\sigma \in \Gamma$ we have $j(\sigma)AFS(\mathfrak{f}) = AFS(\mathfrak{f})$ so that, by Lemma 5.3, $a(t[\sigma];\mathfrak{f}) = a(t;\mathfrak{f})$ for all $t \in \overline{\mathcal{X}}(N)$. Thus $\mathfrak{f} \in \mathbb{M}(k, N, +) \cap \mathbb{M}(k, N; \Gamma) = \mathbb{M}(k, N, +; \Gamma)$ as claimed.

6. Specialization.

Let $\Gamma \subseteq \Gamma^0(N)$ be a subgroup of finite index. A formal series of Jacobi forms $\mathfrak{f} = \sum_{m=1}^{\infty} \phi_m \xi^{Nm} \in \mathbb{S}(k, N; \Gamma)$ has the defining symmetries $a(t[\sigma]; \mathfrak{f}) = a(t; \mathfrak{f})$ for all $\sigma \in \Gamma$ and all $t \in \overline{\mathcal{X}}(N)$. We will construct a dense subset $\mathcal{W}_1(\Gamma)$ of $\mathcal{H}_1 \times \mathbb{C}$ where the formal series \mathfrak{f} specializes to a holomorphic function of one variable. More precisely, for each $(\tau_1, z_1) \in \mathcal{W}_1(\Gamma)$, the series $\sum_{m=1}^{\infty} \phi_m(\tau_1, z_1) e(Nm\omega)$ will converge to a holomorphic function $H(\tau_1, z_1, \mathfrak{f})(\omega)$ on the neighborhood of infinity $\{\omega \in \mathcal{H}_1 : (\frac{\tau_1}{z_1} \frac{z_1}{\omega}) \in \mathcal{H}_2\}$. The formal series \mathfrak{f} thus converges on a dense subset of \mathcal{H}_2 .

Definition 6.1. For $x \in \mathbb{Q}$, let $\text{Denom}(x) = \min\{n \in \mathbb{N} : nx \in \mathbb{Z}\}$ be the minimal positive denominator of x.

Definition 6.2. Let $\Gamma \subseteq SL(2,\mathbb{Z})$ be a subgroup of finite index. Let $U = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in GL(2,\mathbb{Z})$. Define

 $\mathcal{W}_0(\Gamma) = \{(x, y) \in \mathbb{Q}^2 : x \in (U\Gamma U) \text{-}orbit(\infty) \text{ and } y \operatorname{Denom}(x) \in \mathbb{Z}\},\$ $\mathcal{W}_1(\Gamma) = \{(\tau, z) \in \mathcal{H}_1 \times \mathbb{C} : \exists (x, y) \in \mathcal{W}_0(\Gamma) : z = x\tau + y\},\$ $\mathcal{W}(\Gamma) = \{(\begin{smallmatrix} \tau & z \\ z & \omega \end{smallmatrix}) \in \mathcal{H}_2 : (\tau, z) \in \mathcal{W}_1(\Gamma)\}.$

We follow Lehner [30] for the theory of Fuchsian groups. We view the Riemann sphere $\mathbb{P}^1(\mathbb{C}) = \mathbb{C} \cup \{\infty\}$ as the extended complex plane. The Riemann sphere is the disjoint union $\mathbb{P}^1(\mathbb{C}) = \mathbb{P}^1(\mathbb{R}) \amalg \mathcal{H}_1 \amalg \overline{\mathcal{H}}_1$ of the extended real numbers, $\mathbb{P}^1(\mathbb{R}) = \mathbb{R} \cup \{\infty\}$, and the upper and lower half planes. The groups $\mathrm{SL}(2,\mathbb{R})$ and $\mathrm{PSL}(2,\mathbb{R}) = \mathrm{SL}(2,\mathbb{R})/\{\pm I\}$ act on $\mathbb{P}^1(\mathbb{C})$ by Möbius transformations and preserve this disjoint union. We will only consider subgroups $\Gamma \subseteq \mathrm{SL}(2,\mathbb{R})$ and the corresponding transformation groups $\overline{\Gamma} = \langle \Gamma, -I \rangle / \{\pm I\} \subseteq \mathrm{PSL}(2,\mathbb{R})$.

Definition 6.3. Let $\Gamma \subseteq SL(2, \mathbb{R})$. The limit set $\Lambda(\Gamma)$ of Γ is the set of $z \in \mathbb{P}^1(\mathbb{C})$ such that there exists a $w \in \mathbb{P}^1(\mathbb{C})$ and a sequence of distinct $\gamma_n \in \Gamma$ with $\lim_{n \to +\infty} \gamma_n \langle w \rangle = z$.

For subgroups $\Gamma_1, \Gamma_2 \subseteq SL(2, \mathbb{R})$, if Γ_1 has finite index in Γ_2 then $\Lambda(\Gamma_1) = \Lambda(\Gamma_2)$; this is the theorem in section 2C of [30], page 11. The corollary in section 2F, page 14, is that either $\Lambda(\Gamma) = \mathbb{P}^1(\mathbb{C})$ or $\Lambda(\Gamma) \subseteq \mathbb{P}^1(\mathbb{R})$. We mention some terminology to assist readers who use a different reference. A group $\overline{\Gamma} \subseteq PSL(2, \mathbb{R})$ is *Fuchsian* when $\Lambda(\Gamma) \subseteq \mathbb{P}^1(\mathbb{R})$. The theorem in section 2F of [30], page 13, characterizes Fuchsian groups as the discrete subgroups of $PSL(2, \mathbb{R})$. For us, the salient result is Theorem 3 in section 3E, page 21.

Theorem 6.4 ([30]). Let $\Gamma \subseteq SL(2, \mathbb{R})$ be a subgroup. If $F \subseteq \mathbb{P}^1(\mathbb{C})$ is a closed set containing at least two points, such that $\Gamma F \subseteq F$, then $F \supseteq \Lambda(\Gamma)$.

As Lehner comments on page 21, this theorem may be rephrased: when $\Lambda(\Gamma)$ has more than one point, $\Lambda(\Gamma)$ is the smallest closed Γ invariant set containing at least two points.

An example of a Fuchsian group is $\mathrm{PSL}(2,\mathbb{Z})$. The orbit of ∞ is $\mathbb{P}^1(\mathbb{Q})$ and $\Lambda(\mathrm{SL}(2,\mathbb{Z})) = \mathbb{P}^1(\mathbb{R})$. This implies that any subgroup $\Gamma \subseteq \mathrm{SL}(2,\mathbb{Z})$ of finite index also has $\Lambda(\Gamma) = \mathbb{P}^1(\mathbb{R})$. For a ring $R \subseteq \mathbb{R}$, set $\mathcal{P}_{10}(R) = \{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2, R) : c = 0 \}$. The stabilizer of ∞ in $\mathrm{SL}(2, \mathbb{R})$ is $\mathcal{P}_{10}(\mathbb{R})$ and we have an orbit-stabilizer bijection $\mathrm{SL}(2,\mathbb{Z})/\mathcal{P}_{10}(\mathbb{Z}) \leftrightarrow \mathbb{P}^1(\mathbb{Q})$ given by sending $\gamma \mathcal{P}_{10}(\mathbb{Z}) \mapsto \gamma \langle \infty \rangle$. This bijection shows that a subgroup of finite index in $\mathrm{SL}(2,\mathbb{Z})$ cannot stabilize ∞ . We require the following ergodic corollary of Theorem 6.4.

Lemma 6.5. Let $\Gamma \subseteq SL(2,\mathbb{Z})$ be a subgroup of finite index. The Γ -orbit of ∞ is dense in $\Lambda(\Gamma) = \mathbb{P}^1(\mathbb{R})$.

Proof. We have $\Lambda(\Gamma) = \Lambda(\mathrm{SL}(2,\mathbb{Z})) = \mathbb{P}^1(\mathbb{R})$ because Γ has finite index in $\mathrm{SL}(2,\mathbb{Z})$. Furthermore, Γ cannot stabilize ∞ for the same reason. Therefore, the Γ -orbit of ∞ has at least two points, as does its closure $F = \overline{\Gamma}$ -orbit $(\infty) \subseteq \mathbb{P}^1(\mathbb{R})$. We check that the closure F remains Γ invariant. Take $z \in F$ and $\gamma \in \Gamma$. We have $z = \lim_{n \to +\infty} \gamma_n \langle \infty \rangle$ for some $\gamma_n \in \Gamma$, so that $\gamma \langle z \rangle = \lim_{n \to +\infty} (\gamma \gamma_n) \langle \infty \rangle$ for $\gamma \gamma_n \in \Gamma$. Hence $\gamma \langle z \rangle \in F$ and F is Γ -invariant. Thus, by Theorem 6.4 we have $F = \Lambda(\Gamma) = \mathbb{P}^1(\mathbb{R})$. The Γ -orbit of ∞ is thus dense in $\mathbb{P}^1(\mathbb{R})$. \Box

Lemma 6.6. Let $\Gamma \subseteq SL(2,\mathbb{Z})$ be a subgroup of finite index. The sets $\mathcal{W}_0(\Gamma)$, $\mathcal{W}_1(\Gamma)$, and $\mathcal{W}(\Gamma)$ are dense in \mathbb{R}^2 , $\mathcal{H}_1 \times \mathbb{C}$, and \mathcal{H}_2 , respectively.

Proof. It suffices to prove $\mathcal{W}_0(\Gamma)$ is dense in \mathbb{R}^2 . Take $(\xi, \eta) \in \mathbb{R}^2$ and any neighborhoods X of ξ and Y of η . We just need to find an element of $\mathcal{W}_0(\Gamma)$ in $X \times Y$. Pick an irrational number $\xi_o \in X$.

The group $U\Gamma U$ has finite index in $SL(2,\mathbb{Z})$ and so $\Lambda(U\Gamma U) = \mathbb{P}^1(\mathbb{R})$. By Lemma 6.5, the $(U\Gamma U)$ -orbit (∞) is dense in $\Lambda(U\Gamma U) = \mathbb{P}^1(\mathbb{R})$, and hence $\mathbb{Q} \cap (U\Gamma U)$ -orbit $(\infty) = (U\Gamma U)$ -orbit $(\infty) \setminus \{\infty\}$ is dense in \mathbb{R} . Accordingly there is a sequence $x_j \in \mathbb{Q} \cap (U\Gamma U)$ -orbit (∞) with $\lim_j x_j = \xi_o$, and we have $x_j \in X$ for all sufficiently large j. Set $D_j = \text{Denom}(x_j)$. Since ξ_o is irrational we have $\lim_j D_j = +\infty$. Set $y_j = \frac{1}{D_j} \text{floor}(\eta D_j)$ so that $(x_j, y_j) \in \mathcal{W}_0(\Gamma)$ and $\lim_j y_j = \eta$. For all sufficiently large j we have $(x_j, y_j) \in X \times Y$, which shows that $\mathcal{W}_0(\Gamma) \cap (X \times Y)$ is nonempty, and hence that $\mathcal{W}_0(\Gamma)$ is dense in \mathbb{R}^2 . \Box

Let Γ be a subgroup of $\mathrm{SL}(2,\mathbb{Z})$ with finite index. For $f \in S_k(\Gamma)$, define the *Hecke bound* $\mathrm{HB}(f) = \sup_{\tau \in \mathcal{H}_1}(\mathrm{Im}(\tau))^{k/2}|f(\tau)|$. The Hecke bound is a norm on the vector space $S_k(\Gamma)$ and the Fourier coefficients $a(t, \cdot) : S_k(\Gamma) \to \mathbb{C}$, given by $f \mapsto a(t; f)$, become bounded linear functionals. Hence the Fourier coefficients are continuous in f in the topology on $S_k(\Gamma)$ induced by this norm. For Jacobi cusp forms $\phi \in J_{k,m}^{\mathrm{cusp}}$ the Hecke bound [11], compare page 27, is given by

$$\operatorname{HB}(\phi) = \sup_{(\tau,z)\in\mathcal{H}_1\times\mathbb{C}} (\operatorname{Im}(\tau))^{k/2} e^{-2\pi m \frac{(\operatorname{Im}(z))^2}{\operatorname{Im}(\tau)}} |\phi(\tau,z)|.$$

Lemma 6.7. Let $\phi \in J_{k,m}^{\text{cusp}}$ with $m \in \mathbb{N}$. For all $(n,r) \in \mathbb{N} \times \mathbb{Z}$ we have the bound $|c(n,r;\phi)| \leq \text{HB}(\phi) \left(\frac{e\pi}{mk} |4mn - r^2|\right)^{k/2}$.

Proof. Let $\phi(\tau, z) = \sum_{n \in \mathbb{N}, r \in \mathbb{Z}} c(n, r; \phi) e(n\tau + rz)$ be the Fourier series with $\tau = u + iv \in \mathcal{H}_1$ and $z = x + iy \in \mathbb{C}$. By the familiar formula for the Fourier coefficients, we have

$$\begin{aligned} c(n,r;\phi) &= \iint_{[0,1]^2} \phi(\tau,z) e(-n\tau - rz) \, du \, dx \\ |c(n,r;\phi)| &\leq \iint_{[0,1]^2} |\phi(\tau,z)| \, e^{2\pi n v} e^{2\pi r y} \, du \, dx \\ &\leq \iint_{[0,1]^2} \operatorname{HB}(\phi) v^{-k/2} e^{2\pi m \frac{y^2}{v}} e^{2\pi (nv+ry)} \, du \, dx \\ &= \operatorname{HB}(\phi) v^{-k/2} e^{2\pi (m \frac{y^2}{v} + ry + nv)}. \end{aligned}$$

Choosing $y = -\frac{rv}{2m}$ we have $m\frac{y^2}{v} + ry + nv = \frac{4mn-r^2}{4m}v$. Thus we have $|c(n,r;\phi)| \leq \text{HB}(\phi)v^{-k/2}e^{2\pi\frac{4mn-r^2}{4m}v}$. For $4mn-r^2 > 0$ this is minimized by $v = \frac{km}{\pi(4mn-r^2)}$ proving $|c(n,r;\phi)| \leq \text{HB}(\phi)\left(\frac{e\pi}{mk}(4mn-r^2)\right)^{k/2}$. This proves the result when $4mn-r^2 > 0$, and when $4mn-r^2 \leq 0$ we have $c(n,r;\phi) = 0$ so the Lemma's conclusion holds as well. \Box

The following proposition gives a specialization of a cuspidal formal series $\mathfrak{f} \in \mathbb{S}(k, N; \Gamma)$ for each $(x, y) \in \mathcal{W}_0(\Gamma)$. Each specialization depends only upon a finite number of the Fourier-Jacobi coefficients of \mathfrak{f} , although this finite number may increase with the denominator of x.

Proposition 6.8. Let $\Gamma \subseteq \Gamma^0(N)$ be a subgroup of finite index. Let $\mathfrak{f} = \sum_{m=1}^{\infty} \phi_m \xi^{Nm} \in \mathbb{S}(k, N; \Gamma)$. For $(x, y) \in \mathcal{W}_0(\Gamma)$, set D = Denom(x). We have $f_m(\tau) := e(Nmx^2\tau) \phi_m(\tau, x\tau + y) \in S_k(\Gamma(D^2))$. The sequence of Hecke bounds satisfies

$$\operatorname{HB}(f_m) \in O\left(m^{\frac{k+1}{2}}\right),$$

where the implied constant depends only on f and x.

Proof. Take $\mathfrak{f} = \sum_{m=1}^{\infty} \phi_m \xi^{Nm} \in \mathbb{S}(k, N; \Gamma)$. For each $m \in \mathbb{N}$ there is a positive constant K_m such that $|c(n, \ell; \phi_m)| \leq K_m \left(\frac{4Nmn-\ell^2}{4Nm}\right)^{k/2}$ for all $(n, \ell) \in \operatorname{supp}(\phi_m)$. Here we use the assumption that each ϕ_m is a cusp form and Lemma 6.7. By Proposition 2.2 of [20] or Theorem 4.2 of [2], for example, we have $f_m \in S_k(\Gamma(D^2))$ because $Dx, Dy \in \mathbb{Z}$. This elliptic modular cusp form has the Fourier expansion

$$f_m(\tau) = \sum_{v > 0: \ D^2 v \in \mathbb{Z}} a(v; f_m) q^v; \ a(v; f_m) = \sum_{(n,\ell) \in \Lambda_{Nm,x,v}} e(y\ell) c(n,\ell;\phi_m),$$

for $\Lambda_{Nm,x,v} = \{(n,\ell) \in \mathbb{N} \times \mathbb{Z} : Nmx^2 + x\ell + n = v, 4Nmn > \ell^2\}$. Next we check $|\Lambda_{Nm,x,v}| \leq 4\sqrt{Nmv} + 1$. For any $(n,\ell) \in \Lambda_{Nm,x,v}$ we first note that n is determined by ℓ , and that ℓ satisfies

$$(2Nmx + \ell)^2 + 4Nmn - \ell^2 = 4Nm(Nmx^2 + \ell x + n) = 4Nmv.$$

Thus $|2Nmx + \ell| \leq 2\sqrt{Nmv}$ and the number of integers ℓ in this interval is at most the length plus one, showing $|\Lambda_{Nm,x,v}| \leq 4\sqrt{Nmv} + 1$. Pick an $L \in \mathbb{N}$ such that the Fourier coefficients a(v; f) for $v \leq L$ determine $f \in S_k(\Gamma(D^2))$. By virtue of the valence inequality, the choice $L = L(k, D) = \text{floor}\left(\frac{k}{12}[\text{SL}(2,\mathbb{Z}):\Gamma(D^2)]\right)$ will work. The following supremum exists because the denominator is nonzero, the quotient is continuous in f, and the supremum may be taken over a compact sphere.

$$C(k,D) = \sup_{f \in S_k(\Gamma(D^2)) \setminus \{0\}} \frac{\operatorname{HB}(f)}{\sum_{0 < v \le L(k,D)} |a(v;f)|}$$

Therefore we have

$$\operatorname{HB}(f_m) \le C(k, D) \sum_{0 < v \le L(k, D)} |a(v; f_m)|$$

$$\leq C(k,D) \sum_{0 < v \leq L(k,D)} \sum_{(n,\ell) \in \Lambda_{Nm,x,v}} |c(n,\ell;\phi_m)|$$
$$\leq C(k,D) \sum_{(n,\ell) \in \tilde{\Lambda}_{Nm,x,L}} |c(n,\ell;\phi_m)|$$

for $\tilde{\Lambda}_{Nm,x,L} = \{(n,\ell) \in \mathbb{N} \times \mathbb{Z} : Nmx^2 + x\ell + n \leq L, 4Nmn > \ell^2\}$ or, equivalently, $\tilde{\Lambda}_{Nm,x,L} = \bigcup_{0 < v \leq L} \Lambda_{Nm,x,v}$. Each $\Lambda_{Nm,x,v}$ has at most $4\sqrt{Nmv} + 1$ elements and there are at most D^2L of them so we have $|\tilde{\Lambda}_{Nm,x,L}| \leq D^2L(4\sqrt{NmL}+1)$. Note that $(n,\ell) \in \tilde{\Lambda}_{Nm,x,L}$ implies that $4Nmn - \ell^2 \leq 4NmL$. This is because

$$4Nmn - \ell^2 \le (2Nmx + \ell)^2 + 4Nmn - \ell^2 = 4Nm(Nmx^2 + \ell x + n) \le 4NmL$$

We now come to the main point where we use the Γ -conditions for \mathfrak{f} . Our assumption $(x, y) \in \mathcal{W}_0(\Gamma)$ gives us $x \in (U\Gamma U)$ -orbit (∞) . Accordingly there exists a matrix $\begin{pmatrix} a & \eta \\ c & \xi \end{pmatrix} \in U\Gamma U$ with x = a/c. Letting D = Denom(x), the conditions $\gcd(a, c) = 1$ and x = a/c imply $c = \pm D$ and a = cx. Moving this matrix to Γ we define

$$\sigma = \begin{pmatrix} \xi & c \\ \eta & a \end{pmatrix} = U \begin{pmatrix} a & \eta \\ c & \xi \end{pmatrix} U \in \Gamma.$$

We note that $\Gamma \subseteq \Gamma^0(N)$ implies $N \mid D$. For $\binom{n \ \ell/2}{\ell/2 \ Nm} \in \mathcal{X}(N)$, compute $\sigma' \binom{n \ \ell/2}{\ell/2 \ Nm} \sigma = \binom{n_1 \ \ell_1/2}{\ell_1/2 \ Nm_1} \in \mathcal{X}(N)$, where $n_1 = Nm\eta^2 + \ell\xi\eta + n\xi^2$,

$$n_1 = l \cdot m_\eta + l \cdot \zeta \eta + n \zeta \eta,$$

$$\ell_1 = c \left(2n\xi + \ell(\eta + \xi x) + 2Nmx\eta\right),$$

$$m_1 = \frac{D}{N}D \left(Nmx^2 + \ell x + n\right).$$

,

For $(n, \ell) \in \tilde{\Lambda}_{Nm,x,L}$ we have $Nmx^2 + \ell x + n \leq L$ so that $m_1 \leq \frac{D^2}{N}L$, as well as $\binom{n \ell/2}{\ell/2 Nm}$ and $\binom{n_1 \ell_1/2}{\ell_1/2 Nm_1} \in \mathcal{X}(N)$. By the Γ -conditions of equation (3) for \mathfrak{f} we see

$$|c(n,\ell;\phi_m)| = |c(n_1,\ell_1;\phi_{m_1})| \le K_{m_1} \left(\frac{4Nm_1n_1 - \ell_1^2}{4Nm_1}\right)^{k/2}$$
$$= K_{m_1} \left(\frac{4Nmn - \ell^2}{4Nm_1}\right)^{k/2} \le K_{m_1} \left(\frac{4NmL}{4Nm_1}\right)^{k/2} \le K_{m_1} \left(Lm\right)^{k/2}$$

if we remember that $(n, \ell) \in \tilde{\Lambda}_{Nm,x,L}$ implies $0 < 4Nmn - \ell^2 \leq 4NmL$ and note $m_1 \geq 1$. Therefore, for $(n, \ell) \in \tilde{\Lambda}_{Nm,x,L}$, we have

$$|c(n, \ell; \phi_m)| \le K_{m_1} (Lm)^{k/2} \le \left(\max_{1 \le j \le \frac{D^2}{N}L} K_j\right) L^{k/2} m^{k/2}$$

AOKI, IBUKIYAMA, POOR

$$= K_{D,N,L} L^{k/2} m^{k/2}$$

since $m_1 \leq \frac{D^2}{N}L$. The constant $K_{D,N,L}$ grows with the denominator of $(x, y) \in \mathcal{W}_0(\Gamma)$ but in each individual case controls the growth of all HB (f_m) in terms of the finite number of HB (ϕ_j) with $j \leq \frac{D^2}{N}L$. Therefore we have

$$HB(f_m) \leq C(k, D) \sum_{(n,\ell)\in\tilde{\Lambda}_{Nm,x,L}} |c(n,\ell;\phi_m)| \\ \leq C(k, D) \sum_{(n,\ell)\in\tilde{\Lambda}_{Nm,x,L}} K_{D,N,L} L^{k/2} m^{k/2} \\ = C(k, D) |\tilde{\Lambda}_{Nm,x,L}| K_{D,N,L} L^{k/2} m^{k/2} \\ \leq C(k, D) K_{D,N,L} L^{k/2} D^2 L (4\sqrt{NmL} + 1) m^{k/2} \\ \leq C_2(k, D, N, L) (4\sqrt{NmL} + 1) m^{k/2}.$$

Thus $\operatorname{HB}(f_m) \in O\left(m^{\frac{k+1}{2}}\right)$ as was to be shown.

Consider a formal series $\mathfrak{f} = \sum_{m=0}^{\infty} \phi_m \xi^{Nm} \in \mathbb{M}(k, N)$. For a point $\Omega_o = \begin{pmatrix} \tau_o & z_o \\ z_o & \omega_o \end{pmatrix} \in \mathcal{H}_2$, if the series $\sum_{m=0}^{\infty} \phi_m(\tau_o, z_o) e(Nm\omega_o)$ converges absolutely, we say that the formal series \mathfrak{f} converges absolutely at Ω_o .

Corollary 6.9. Let $\Gamma \subseteq \Gamma^0(N)$ be a subgroup of finite index. Let $\mathfrak{f} = \sum_{m=1}^{\infty} \phi_m \xi^{Nm} \in \mathbb{S}(k, N; \Gamma)$. The formal series \mathfrak{f} converges absolutely on $\mathcal{W}(\Gamma)$. For $(\tau_1, z_1) \in \mathcal{W}_1(\Gamma)$ set $\eta_1 = (\mathrm{Im}(z_1))^2/\mathrm{Im}(\tau_1)$. The function $H(\tau_1, z_1, \mathfrak{f}) : N_{\infty}(\eta_1) \to \mathbb{C}$ defined by

$$H(\tau_1, z_1, \mathfrak{f})(\omega) = \sum_{m=1}^{\infty} \phi_m(\tau_1, z_1) e(Nm\omega)$$

is holomorphic on $N_{\infty}(\eta_1)$.

Proof. Pick $\Omega = \begin{pmatrix} \tau & x\tau+y \\ x\tau+y & \omega \end{pmatrix} \in \mathcal{W}(\Gamma)$ with $(x,y) \in \mathcal{W}_0(\Gamma)$. By Proposition 6.8, $\operatorname{HB}(f_m) \leq C(\mathfrak{f}, x)m^{\frac{k+1}{2}}$ for some constant $C(\mathfrak{f}, x)$ and the elliptic forms $f_m \in S_k(\Gamma(D^2))$ given by $f_m(\tau) = e(Nmx^2\tau) \phi_m(\tau, x\tau+y)$. From $(\operatorname{Im}(\tau))^{k/2}|f_m(\tau)| \leq \operatorname{HB}(f_m) \leq C(\mathfrak{f}, x)m^{\frac{k+1}{2}}$ we infer

$$|\phi_m(\tau, x\tau + y)| \le C(\mathfrak{f}, x)(\operatorname{Im}(\tau))^{-k/2} e^{2\pi Nmx^2(\operatorname{Im}(\tau))} m^{\frac{k+1}{2}},$$
$$\limsup_{m \to +\infty} |\phi_m(\tau, x\tau + y)|^{\frac{1}{m}} \le e^{2\pi Nx^2(\operatorname{Im}(\tau))}.$$

The radius of convergence R of the series $\sum_{m\in\mathbb{N}} \phi_m(\tau, x\tau + y)\xi^{Nm}$ thus satisfies $R \ge e^{-2\pi N x^2(\operatorname{Im}(\tau))}$, and so this series converges absolutely for $|\xi|^N < e^{-2\pi N x^2(\operatorname{Im}(\tau))^2}$. In particular the series converges at Ω if $\operatorname{Im}(\omega) >$

 $x^{2}(\operatorname{Im}(\tau)) = \frac{(\operatorname{Im}(x\tau+y))^{2}}{\operatorname{Im}(\tau)}.$ The condition $\operatorname{Im}(\omega)\operatorname{Im}(\tau) > (\operatorname{Im}(x\tau+y))^{2},$ however, is just $\Omega = \begin{pmatrix} \tau & x\tau+y \\ x\tau+y & \omega \end{pmatrix} \in \mathcal{H}_{2}.$

Consider $(\tau_1, z_1) \in \mathcal{W}_1(\Gamma)$ so that $(\tau_1, z_1) = (\tau_1, x\tau_1 + y)$ for some $(x, y) \in \mathcal{W}_0(\Gamma)$. We have seen that the power series in $e(\omega)$,

(8)
$$\sum_{m\in\mathbb{N}}\phi_m(\tau_1, z_1)e\left(Nm\omega\right),$$

converges when $\begin{pmatrix} \tau_1 & z_1 \\ z_1 & \omega \end{pmatrix} \in \mathcal{H}_2$. A power series that converges on an open set is holomorphic there. Thus $H(\tau_1, z_1, \mathfrak{f})$ is holomorphic on the open set $\{\omega \in \mathcal{H}_1 : \begin{pmatrix} \tau_1 & z_1 \\ z_1 & \omega \end{pmatrix} \in \mathcal{H}_2\} = N_{\infty}(\eta_1)$.

7. LOCALLY BOUNDED.

Locally bounded families of holomorphic functions possess remarkable convergence properties. In Theorem 7.4 we show that, for subgroups $\Gamma \subseteq \Gamma^0(N)$ of finite index, the partial sums of the formal series $\mathfrak{g} = \sum_{m=1}^{\infty} \phi_m \xi^{Nm} \in \mathbb{S}(k, N, \epsilon; \Gamma)$ are locally bounded on \mathcal{H}_2 if \mathfrak{g} is integral over FJ ($\mathbb{M}_{\pm}(N; \Gamma)$). The following homomorphism respects the ring structure but forgets the grading on $\mathbb{M}_{\pm}(N)$.

Lemma 7.1. For each $(\tau_1, z_1) \in \mathcal{H}_1 \times \mathbb{C}$ there is a ring homomorphism $A(\tau_1, z_1) : \mathbb{M}_{\pm}(N) \to \mathbb{C}[[\xi]]$ defined by sending $\mathfrak{f} = \sum_{m=0}^{\infty} \phi_m \xi^{Nm} \in \mathbb{M}(k, N, \epsilon)$ to $A(\tau_1, z_1)\mathfrak{f} = \sum_{m=0}^{\infty} \phi_m(\tau_1, z_1)\xi^{Nm} \in \mathbb{C}[[\xi]]$ and extending additively to $\mathbb{M}_{\pm}(N) = \bigoplus_{k,\epsilon} \mathbb{M}(k, N, \epsilon)$.

Proof. The ring structures are defined by the Cauchy product rule, so substitution is a homomorphism. \Box

Let $D = \{z \in \mathbb{C} : |z| < r\}$ be the open disk about the origin in \mathbb{C} with positive radius r. Let $\mathcal{S}(D) = \{w \in \mathbb{C}[[\xi]] : w$ converges on $D\}$ be the ring of power series that converge on D. The map $T_0 : \mathcal{O}(D) \to \mathcal{S}(D)$ that sends a holomorphic function on D to its Taylor series about the origin is a ring isomomorphism. For $\xi_o \in D$ and $w = \sum_{n=0}^{\infty} w_n \xi^n \in$ $\mathcal{S}(D)$, let $w[\xi_o] = \sum_{n=0}^{\infty} w_n \xi_o^n \in \mathbb{C}$ be the sum of w at $\xi := \xi_o$. The evaluation map $[\xi_o] : \mathcal{S}(D) \to \mathbb{C}$ sending w to $w[\xi_o]$ is a ring homomorphism with the property that $(T_0^{-1}w)(\xi_o) = w[\xi_o]$.

Lemma 7.2. For each $(\tau_1, z_1) \in \mathcal{H}_1 \times \mathbb{C}$, define $\eta_1 = \frac{(\operatorname{Im}(z_1))^2}{\operatorname{Im}(\tau_1)}$, a radius $r_1 = e^{-2\pi\eta_1}$, and a disk $D_1 = \{\xi \in \mathbb{C} : |\xi| < r_1\}$. The ring homomorphism $A(\tau_1, z_1) : \mathbb{M}_{\pm}(N) \to \mathbb{C}[[\xi]]$ sends $\operatorname{FJ}(M_{\pm}(N))$ into $\mathcal{S}(D_1)$. For $\omega \in N_{\infty}(\eta_1)$, set $\Omega_1(\omega) = \begin{pmatrix} \tau_1 & z_1 \\ z_1 & \omega \end{pmatrix} \in \mathcal{H}_2$. For $f \in M_k(K(N))^{\epsilon}$ and $\omega \in N_{\infty}(\eta_1)$, we have $(A(\tau_1, z_1)\operatorname{FJ}(f))[e(\omega)] = f(\Omega_1(\omega))$. Let $\Gamma \subseteq \Gamma^0(N)$ be a subgroup of finite index. For $(\tau_1, z_1) \in \mathcal{W}_1(\Gamma)$, $A(\tau_1, z_1)$ sends $\mathbb{S}_{\pm}(N; \Gamma)$ into $\mathcal{S}(D_1)$. For $\mathfrak{f} \in \mathbb{S}(k, N, \epsilon; \Gamma)$ and $\omega \in N_{\infty}(\eta_1)$ we have $(A(\tau_1, z_1)\mathfrak{f})[e(\omega)] = H(\tau_1, z_1, \mathfrak{f})(\omega)$ for the holomorphic function $H(\tau_1, z_1, \mathfrak{f})$ defined in Corollary 6.9. The map $H(\tau_1, z_1) :$ $\mathbb{S}_{\pm}(N; \Gamma) \to \mathcal{O}(N_{\infty}(\eta_1))$ that sends \mathfrak{f} to $H(\tau_1, z_1, \mathfrak{f})$ is linear and multiplicative.

Proof. We know that $A(\tau_1, z_1)$ is a ring homomorphism from Lemma 7.1. We need to check that the relevant power series converge in D_1 and correctly label their values. For $f \in M_k(K(N))^{\epsilon}$ the Fourier-Jacobi expansion $f(\Omega) = \sum_{m=0}^{\infty} \phi_m(\tau, z) e(Nm\omega)$ converges for $\Omega = \begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} \in \mathcal{H}_2$. The formal series is $\mathrm{FJ}(f) = \sum_{m=0}^{\infty} \phi_m \xi^{Nm}$ and the value of $A(\tau_1, z_1)$ on it is $A(\tau_1, z_1)\mathrm{FJ}(f) = \sum_{m=0}^{\infty} \phi_m(\tau_1, z_1)\xi^{Nm} \in \mathbb{C}[[\xi]]$. For any $\xi_o \in D_1$, there is an $\omega_o \in N_{\infty}(\eta_1)$ with $\xi_o = e(\omega_o)$ and $\Omega_1(\omega_o) \in \mathcal{H}_2$. Accordingly, $f(\Omega_1(\omega_o)) = \sum_{m=0}^{\infty} \phi_m(\tau_1, z_1)\xi_o^{Nm}$ is convergent and, since $\xi_o \in D_1$ was arbitrary, $A(\tau_1, z_1)\mathrm{FJ}(f) \in \mathcal{S}(D_1)$. For any $\omega \in N_{\infty}(\eta_1)$ we have $e(\omega) \in D_1$ and $(A(\tau_1, z_1)\mathrm{FJ}(f))[e(\omega)] = \sum_{m=0}^{\infty} \phi_m(\tau_1, z_1)e(Nm\omega) =$ $f(\Omega_1(\omega))$.

Assume that $(\tau_1, z_1) \in \mathcal{W}_1(\Gamma)$. Since $\mathfrak{f} \in \mathbb{S}(k, N, \epsilon; \Gamma) \subseteq \mathbb{S}(k, N; \Gamma)$, we can apply Corollary 6.9 to assure the convergence of the series $H(\tau_1, z_1, \mathfrak{f})(\omega) = \sum_{m=1}^{\infty} \phi_m(\tau_1, z_1) e(Nm\omega)$ for $\omega \in N_{\infty}(\eta_1)$. Any $\xi \in D_1$ may be written $\xi = e(\omega)$ for $\omega \in N_{\infty}(\eta_1)$, so the power series $A(\tau_1, z_1)\mathfrak{f} = \sum_{m=1}^{\infty} \phi_m(\tau_1, z_1)\xi^{Nm}$ converges on D_1 and so $A(\tau_1, z_1)\mathfrak{f} \in \mathcal{S}(D_1)$. The value of the convergent series at $e(\omega) \in D_1$ is given by $(A(\tau_1, z_1)\mathfrak{f})[e(\omega)] = \sum_{m=1}^{\infty} \phi_m(\tau_1, z_1)e(Nm\omega) = H(\tau_1, z_1, \mathfrak{f})(\omega).$

Both $A(\tau_1, z_1)$ and the evaluation map $[e(\omega)]$ are ring homomorphisms so their composition, restricted to the ideal $\mathbb{S}_{\pm}(N; \Gamma)$ is linear and multiplicative. This shows, for all $\omega \in N_{\infty}(\eta_1)$, that we have

$$\begin{aligned} H(\tau_1, z_1, \mathfrak{f}_1 \mathfrak{f}_2)(\omega) &= (A(\tau_1, z_1) \mathfrak{f}_1 \mathfrak{f}_2) \left[e(\omega) \right] = \\ (A(\tau_1, z_1) \mathfrak{f}_1) \left[e(\omega) \right] (A(\tau_1, z_1) \mathfrak{f}_2) \left[e(\omega) \right] &= H(\tau_1, z_1, \mathfrak{f}_1)(\omega) H(\tau_1, z_1, \mathfrak{f}_2)(\omega), \\ \text{which is what } H(\tau_1, z_1, \mathfrak{f}_1 \mathfrak{f}_2) &= H(\tau_1, z_1, \mathfrak{f}_1) H(\tau_1, z_1, \mathfrak{f}_2) \text{ means.} \end{aligned}$$

Lemma 7.3. Let $d \in \mathbb{N}$ and $z \in \mathbb{C}$ be given. Let a monic polynomial $P \in \mathbb{C}[X]$ of degree d be given by $P(X) = X^d + \sum_{j=1}^d a_j X^{d-j}$ for $a_1, \ldots, a_d \in \mathbb{C}$. If P(z) = 0 then $|z| \leq 1 + \sum_{j=1}^d |a_j|$.

Proof. Apply the Triangle Inequality to the case $|z| \ge 1$.

Theorem 7.4. Let $N, k, d \in \mathbb{N}$. Let $\epsilon \in \{\pm 1\}$. Let $\Gamma \subseteq \Gamma^0(N)$ be a subgroup of finite index. Let $\mathfrak{g} = \sum_{m=1}^{\infty} \phi_m \xi^{Nm} \in \mathbb{S}(k, N, \epsilon; \Gamma)$ satisfy the monic polynomial relation $P(\mathfrak{g}) = 0$ in $\mathbb{M}(kd, N, \epsilon^d; \Gamma)$ for

$$P(X) = X^d + \dots + FJ(g_j)X^{d-j} + \dots + FJ(g_d)$$

with $g_j \in M_{kj} (K(N))^{\epsilon^j}$ for $j = 1, \ldots, d$.

Then the sequence of partial sums $\sum_{m=1}^{M} \phi_m(\tau, z) e(Nm\omega)$ for $M \in \mathbb{N}$ is locally bounded on $\begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} \in \mathcal{H}_2$.

Proof. Pick $\Omega_o \in \mathcal{H}_2$. We need to make a neighborhood B_1 of Ω_o and a positive constant A_1 such that for all $M \in \mathbb{N}$ and all $\begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} \in B_1$ we have $|\sum_{m=1}^{M} \phi_m(\tau, z)e(Nm\omega)| \leq A_1$. Let B_1 be an open Euclidean ball centered at Ω_o with closure $\overline{B}_1 \subseteq \mathcal{H}_2$. We can push the closed ball \overline{B}_1 down by $-i\begin{pmatrix} 0 & 0 \\ 0 & \epsilon \end{pmatrix}$ so that this translation is still contained in \mathcal{H}_2 . We do this in detail. For any $\epsilon > 0$, the translated ball lies in the space $L = \{\begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} \in M_{2 \times 2}^{\text{sym}}(\mathbb{C}) : \tau \in \mathcal{H}_1\}$. Define the function $h : L \to \mathbb{R}$ for any $\Omega = \begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} \in L$ by

$$h(\Omega) = \frac{\det(\operatorname{Im}(\Omega))}{\langle \operatorname{Im}(\Omega), \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \rangle} = \operatorname{Im}(\omega) - \frac{(\operatorname{Im}(z))^2}{\operatorname{Im}(\tau)}$$

The function h is continuous on L and positive on $\mathcal{H}_2 \subseteq L$ and so has a positive minimum on \overline{B}_1 . Set $\epsilon = \frac{1}{2} \min_{\Omega \in \overline{B}_1} h(\Omega) > 0$. We remark that ϵ only depends on the choice of the ball B_1 . Translate the ball B_1 by $-i\begin{pmatrix} 0 & 0 \\ 0 & \epsilon \end{pmatrix}$ to obtain the ball $B_2 = \{\Omega - i\begin{pmatrix} 0 & 0 \\ 0 & \epsilon \end{pmatrix} \in L : \Omega \in B_1\}$ with closure in L given by $\overline{B}_2 = \{\Omega - i\begin{pmatrix} 0 & 0 \\ 0 & \epsilon \end{pmatrix} \in L : \Omega \in \overline{B}_1\}$. We have

$$\inf_{\Omega \in \overline{B}_2} h(\Omega) = \inf_{\Omega \in \overline{B}_1} (h(\Omega) - \epsilon) = 2\epsilon - \epsilon = \epsilon > 0.$$

Therefore det(Im(Ω)) > 0 for $\Omega \in \overline{B}_2$ and $\overline{B}_2 \subseteq \mathcal{H}_2$. By this process, B_2 is completely determined by B_1 . Next define a compact set $K_2 = \{\Omega + \begin{pmatrix} 0 & 0 \\ 0 & \mu \end{pmatrix} \in \mathcal{H}_2 : \Omega \in \overline{B}_2, 0 \leq \mu \leq 1\}$ that contains \overline{B}_2 . Use the continuity of the g_j to define $A_0 = A_0(B_1, P) = \sup_{\Omega \in K_2} \left(1 + \sum_{j=1}^d |g_j(\Omega)|\right)$. We will show that $A_1 = \frac{A_0}{e^{2\pi N\epsilon} - 1}$ works as the local bound at B_1 .

The main step will be to show $|\sum_{m=1}^{M} \phi_m(\tau_1, z_1)e(Nm\omega_1)| \leq A_1$ for every $M \in \mathbb{N}$ and every $\Omega_1 = \begin{pmatrix} \tau_1 & z_1 \\ z_1 & \omega_1 \end{pmatrix} \in B_1 \cap \mathcal{W}(\Gamma)$. The conclusion $|\sum_{m=1}^{M} \phi_m(\tau, z)e(Nm\omega)| \leq A_1$ on the neighborhood B_1 then follows because $\sum_{m=1}^{M} \phi_m(\tau, z)e(Nm\omega)$ is continuous on B_1 and $B_1 \cap \mathcal{W}(\Gamma)$ is dense in B_1 . The assumption $\Omega_1 \in B_1 \cap \mathcal{W}(\Gamma)$ implies $(\tau_1, z_1) \in \mathcal{W}_1(\Gamma)$. We may rewrite the hypothesis

$$\mathfrak{g}^d + \cdots + \mathrm{FJ}(g_j)\mathfrak{g}^{d-j} + \cdots + \mathrm{FJ}(g_d) = 0$$

as $\sum_{j=0}^{d} \operatorname{FJ}(g_j)\mathfrak{g}^{d-j} = 0$ where $g_0 = 1 \in M_0(K(N))^+$. Apply the ring homomorphism $A(\tau_1, z_1) : \mathbb{M}_{\pm}(N) \to \mathbb{C}[[\xi]]$ to the given relation $\sum_{j=0}^{d} \operatorname{FJ}(g_j)\mathfrak{g}^{d-j} = 0$ in $\mathbb{M}(kd, N, \epsilon^d; \Gamma)$ to obtain a relation among

formal power series

(9)
$$\sum_{j=0}^{d} \left(A(\tau_1, z_1) \mathrm{FJ}(g_j) \right) \left(A(\tau_1, z_1) \mathfrak{g} \right)^{d-j} = 0.$$

By Lemma 7.2, we know $A(\tau_1, z_1)\mathfrak{g} \in \mathcal{S}(D_1)$ for $\eta_1 = \frac{(\operatorname{Im}(z_1))^2}{\operatorname{Im}(\tau_1)}$, the radius $r_1 = e^{-2\pi\eta_1} \leq 1$, and the disk $D_1 = \{z \in \mathbb{C} : |z| < r_1\}$. Moreover, the power series $A(\tau_1, z_1) \operatorname{FJ}(g_j)$ converges in D_1 . Therefore the formal power series in equation (9) is in the subring $\mathcal{S}(D_1)$ of *convergent* power series on D_1 .

For $\omega \in N_{\infty}(\eta_1)$, we have $\xi_1 = e(\omega) \in D_1$ and so we may use the evaluation homomorphism $[\xi_1] : \mathcal{S}(D_1) \to \mathbb{C}$ to obtain a relation among complex numbers

(10)
$$\sum_{j=0}^{d} \left(A(\tau_1, z_1) \mathrm{FJ}(g_j) \right) \left[\xi_1 \right] \left(\left(A(\tau_1, z_1) \mathfrak{g} \right) \left[\xi_1 \right] \right)^{d-j} = 0.$$

For $\omega \in N_{\infty}(\eta_1)$, define $\Omega_1(\omega) = \begin{pmatrix} \tau_1 & z_1 \\ z_1 & \omega \end{pmatrix} \in \mathcal{H}_2$. By Lemma 7.2, for any $\omega \in N_{\infty}(\eta_1)$, we have both $(A(\tau_1, z_1) \mathrm{FJ}(g_j))[\xi_1] = g_j(\Omega_1(\omega))$ and $(A(\tau_1, z_1)\mathfrak{g})[\xi_1] = H(\tau_1, z_1, \mathfrak{g})(\omega)$. We rewrite equation (10) as

$$\left(H(\tau_1, z_1, \mathfrak{g})(\omega)\right)^d + \sum_{j=1}^d g_j\left(\Omega_1(\omega)\right) \left(H(\tau_1, z_1, \mathfrak{g})(\omega)\right)^{d-j} = 0.$$

By Lemma 7.3 we have

(11)
$$|H(\tau_1, z_1, \mathfrak{g})(\omega)| \le 1 + \sum_{j=1}^d |g_j(\Omega_1(\omega))|$$

By the definition of $A_0 = \sup_{\Omega \in K_2} \left(1 + \sum_{j=1}^d |g_j(\Omega)| \right)$, we have

(12)
$$|H(\tau_1, z_1, \mathfrak{g})(\omega)| \le A_0, \text{ if } \Omega_1(\omega) \in K_2.$$

We have $\omega_1 - i\epsilon \in N_{\infty}(\eta_1) = \{\omega \in \mathcal{H}_1 : \Omega_1(\omega) \in \mathcal{H}_2\}$ because $\Omega_1(\omega_1 - i\epsilon) \in B_2 \subseteq \mathcal{H}_2$. Furthermore, for $0 \leq \mu \leq 1$, we have $\omega_1 - i\epsilon + \mu \in N_{\infty}(\eta_1)$ because $\Omega_1(\omega_1 - i\epsilon + \mu) \in K_2 \subseteq \mathcal{H}_2$, and the function $H(\tau_1, z_1, \mathfrak{g})$ is holomorphic on $\omega_1 - i\epsilon + [0, 1] \subseteq N_{\infty}(\eta_1)$. Thus we may compute the Fourier coefficients of

$$H(\tau_1, z_1, \mathfrak{g})(\omega) = \sum_{m=1}^{\infty} \phi_m(\tau_1, z_1) e\left(Nm\omega\right)$$

by the familiar formula

$$\phi_m(\tau_1, z_1) = \int_0^1 H(\tau_1, z_1, \mathfrak{g})(\omega_1 - i\epsilon + \mu)e\left(-Nm(\omega_1 - i\epsilon + \mu)\right) d\mu.$$

We calculate, for any $M \in \mathbb{N}$,

$$\begin{aligned} &|\sum_{m=1}^{M} \phi_m(\tau_1, z_1) e\left(Nm\omega_1\right)| = \\ &|\sum_{m=1}^{M} \left(\int_0^1 H(\tau_1, z_1, \mathfrak{g})(\omega_1 - i\epsilon + \mu) e\left(-Nm(\omega_1 - i\epsilon + \mu)\right) d\mu\right) e\left(Nm\omega_1\right) \\ &\leq \sum_{m=1}^{M} e^{-2\pi Nm \operatorname{Im}(\omega_1)} \int_0^1 |H(\tau_1, z_1, \mathfrak{g})(\omega_1 - i\epsilon + \mu)| e^{2\pi Nm(\operatorname{Im}(\omega_1) - \epsilon)} d\mu \\ &= \sum_{m=1}^{M} e^{-2\pi Nm\epsilon} \int_0^1 |H(\tau_1, z_1, \mathfrak{g})(\omega_1 - i\epsilon + \mu)| d\mu. \end{aligned}$$

As mentioned above, we have $\Omega_1(\omega_1 - i\epsilon + \mu) = \begin{pmatrix} \tau_1 & z_1 \\ z_1 & \omega_1 - i\epsilon + \mu \end{pmatrix} \in K_2$ for $0 \leq \mu \leq 1$, so that $|H(\tau_1, z_1, \mathfrak{g})(\omega_1 - i\epsilon + \mu)| \leq A_0$ by equation (12). Thus

$$\left|\sum_{m=1}^{M} \phi_{m}(\tau_{1}, z_{1}) e\left(Nm\omega_{1}\right)\right|$$

$$\leq A_{0} \sum_{m=1}^{M} e^{-2\pi Nm\epsilon} < \frac{A_{0}e^{-2\pi N\epsilon}}{1 - e^{-2\pi N\epsilon}} = \frac{A_{0}}{e^{2\pi N\epsilon} - 1} = A_{1}.$$

This completes the main step and, as explained, consequently shows that $|\sum_{m=1}^{M} \phi_m(\tau, z) e(Nm\omega)| \leq A_1$ on the neighborhood B_1 of Ω_o for all $M \in \mathbb{N}$. Since $\Omega_o \in \mathcal{H}_2$ was arbitrary, the sequence of partial sums $\sum_{m=1}^{M} \phi_m(\tau, z) e(Nm\omega)$ is locally bounded on \mathcal{H}_2 .

8. HOLOMORPHICITY.

In Theorem 8.2 we show that the divisibility of paramodular Fricke eigenforms is implied by the cuspidality of the quotient of their formal series. Later, after proving our main result, Theorem 10.1 improves Theorem 8.2 by dropping the cuspidality assumption.

We review divisor theory following Gunning [21]. Let $G \subseteq \mathbb{C}^n$ be a domain. For open $U \subseteq G$, let $\mathcal{O}(U)$ be the ring of holomorphic functions on U. For $p \in G$, let \mathcal{O}_p be the ring of germs of holomorphic functions, and \mathcal{M}_p the field of germs of meromorphic functions at p. A holomorphic subvariety is a subset $V \subseteq G$ such that each point $p \in V$ has a neighborhood U where $U \cap V$ is the zero set of finitely many functions holomorphic on U. A holomorphic subvariety V is *irreducible* if $V = V_1 \cup V_2$ for holomorphic subvarieties V_1 , V_2 , implies $V_1 = V$ or $V_2 = V$. The *regular points* of V, $\mathcal{R}(V)$, are the points $p \in V$ where $U \cap V$ is a complex manifold for some neighborhood U of p. The regular points are an open dense subset of V, locally finitely connected, and $\mathcal{R}(V)$ is connected precisely when V is irreducible. The dimension, dim V, of an irreducible V is the dimension of the complex manifold $\mathcal{R}(V)$, and the codimension of V is $n - \dim V$.

We let $\mathcal{E}(G)$ be the set of irreducible, codimension one holomorphic subvarieties of G. For a function $\Delta : \mathcal{E}(G) \to \mathbb{Z}$, the support of Δ is $\operatorname{supp}(\Delta) = \{V \in \mathcal{E}(G) : \Delta(V) \neq 0\}$. We say Δ is locally finite if, for all open $U \subseteq G$ with $\overline{U} \subseteq G$, the set $\{V \in \operatorname{supp}(\Delta) : U \cap V \neq \emptyset\}$ is finite. The group of divisors in G, \mathcal{D}_G , is the abelian group of locally finite functions $\Delta : \mathcal{E}(G) \to \mathbb{Z}$. Since $\operatorname{supp}(\Delta)$ is countable, we often write a global divisor as $\Delta = \sum_{j \in \mathbb{N}} \nu_j V_j$, for $\nu_j = \Delta(V_j) \in \mathbb{Z}$ and distinct $V_j \in \operatorname{supp}(\Delta)$. The semigroup of effective divisors, $\mathcal{D}_G^+ \subseteq \mathcal{D}_G$, is defined by $\nu_j \geq 0$.

For the germ of a holomorphic subvariety at p, V_p , the dimension, dim V_p , is the maximal dimension of the finitely many connected components of $\mathcal{R}(V)$ whose closure contains p. We say V_p is pure dimensionsal when all these connected components have the same dimension. The definition of irreducibility for V_p is as before. Let $\mathcal{E}(p)$ be the set of germs of holomorphic subvarieties at p that are irreducible and have codimension one. A function $\delta : \mathcal{E}(p) \to \mathbb{Z}$ has support $\operatorname{supp}(\delta) = \{V_p \in \mathcal{E}(p) : \delta(V_p) \neq 0\}$. The group of local divisors at p, \mathcal{D}_p , is the free abelian group on $\mathcal{E}(p)$. We write local divisors $\delta \in \mathcal{D}_p$ as $\delta = \sum_{j=1}^{\ell} \nu_j V_j$ with $\nu_j = \delta(V_j) \in \mathbb{Z}$ and distinct $V_j \in \mathcal{E}(p)$ but, fundamentally, a local divisor at p is a function $\delta : \mathcal{E}(p) \to \mathbb{Z}$ with finite support. The semigroup of effective local divisors, $\mathcal{D}_p^+ \subseteq \mathcal{D}_p$, is defined by $\nu_j \geq 0$. If $p \in \mathcal{R}(V)$ then V_p is irreducible. For $V \in \mathcal{E}(G)$ and general $p \in V$, $V_p = V'_p \cup V''_p \cup \cdots$ decomposes into a finite union of distinct $V'_p, V''_p, \ldots \in \mathcal{E}(p)$.

The ring \mathcal{O}_p is a unique factorization domain and noetherian. The fundamental correspondence of algebraic geometry holds between germs of holomorphic subvarieties and germs of holomorphic functions at p. The ideal $\mathrm{id}(V_p) \subseteq \mathcal{O}_p$ consists of the germs that vanish on V_p and, for an ideal $\mathfrak{a} \subseteq \mathcal{O}_p$, $\mathrm{loc}(\mathfrak{a})$ is the germ of the holomorphic subvariety defined by the simultaneous vanishing of the elements of \mathfrak{a} . The germ V_p is irreducible precisely when $\mathrm{id}(V_p)$ is prime, and V_p has pure codimension one precisely when $\mathrm{id}(V_p)$ is principal. Therefore, for $V \in \mathcal{E}(G)$ and $p \in V$, there is a neigborhood U of p and a $\varpi \in \mathcal{O}(U)$, with ϖ_p prime in \mathcal{O}_p , such that $V \cap U = \{Z \in U : \varpi(Z) = 0\}$. We call this a local equation at p and refer to ϖ as a uniformizer. As a consequence of Cartan's Theorem, see Theorem F6 of [21], there is a neighborhood

 $U'' \subseteq U$ of p such that ϖ_q generates id V_q at all points $q \in U'' \cap V$. Cartan's Theorem asserts the existence of a neighborhood $U' \subseteq U$ of pand of functions $h_1, \ldots, h_\ell \in \mathcal{O}(U')$ such that id $V_q = \mathcal{O}_q(h_{1q}, \ldots, h_{\ell q})$ at all points $q \in U' \cap V$. Since ϖ_p divides each h_{jp} there is a neighborhood $U'' \subseteq U'$ of p where id $V_q = \mathcal{O}_q(h_{1q}, \ldots, h_{\ell q}) \subseteq \mathcal{O}_q \varpi_q$ at all points $q \in U'' \cap V$. Hence id $V_q = \mathcal{O}_q \varpi_q$ and ϖ_q is prime for all $q \in U'' \cap V$.

For a ring R, let R^{\times} be the group of units and $R^* = R \setminus \{0\}$. The sequence of semigroups $0 \to (\mathcal{O}_p^{\times}, \cdot) \to (\mathcal{O}_p^{*}, \cdot) \xrightarrow{\operatorname{div}_p} (\mathcal{D}_p^+, +) \to 0$ is exact, where $\operatorname{div}_p(f_p) = \sum_{j=1}^{\ell} \nu_j \operatorname{loc}(\mathcal{O}_p f_j)$ is the semigroup homomorphism determined by the factorization $f_p = (\text{unit})f_1^{\nu_1}\cdots f_{\ell}^{\nu_{\ell}}$ in \mathcal{O}_p into powers of nonassociate irreducibles f_j , unique up to order and units. The rule $\operatorname{div}_p(f_p/g_p) = \operatorname{div}_p(f_p) - \operatorname{div}_p(g_p)$ defines a group homomorphism $\operatorname{div}_p : (\mathcal{M}_p^*, \cdot) \to (\mathcal{D}_p, +)$. The local divisor $\operatorname{div}_p(f_p)$ is effective precisely when $f_p \in \mathcal{O}_p^* \subseteq \mathcal{M}_p^*$. To each global divisor $\Delta \in \mathcal{D}_G$ and point $p \in G$, we associate a local divisor, $\operatorname{germ}_p(\Delta) =$ $\sum_{V \in \text{supp}(\Delta): p \in V} \Delta(V) \left(V'_p + V''_p + \cdots \right)$, where $V_p = V'_p \cup V''_p \cup \cdots$ is the finite decomposition of V_p into irreducibles. For each $f \in \mathcal{O}(G)^*$, the global divisor div(f) is the unique $\Delta \in \mathcal{D}_G^+$ such that germ_p $(\Delta) =$ $\operatorname{div}_p(f_p)$ for all $p \in G$. The rule $\operatorname{div}(f/g) = \operatorname{div}(f) - \operatorname{div}(g)$ extends div to a group homomorphism div : $(\mathcal{M}(G)^*, \cdot) \to (\mathcal{D}_G, +),$ where the field of meromorphic functions $\mathcal{M}(G)$ is the quotient field of $\mathcal{O}(G)$. The global assertion that div(f) is effective precisely when $f \in \mathcal{O}(G)^* \subseteq \mathcal{M}(G)^*$ follows from the corresponding local assertion.

For $V \in \mathcal{E}(G)$ and $f \in \mathcal{O}(G)^*$, a direct way to compute $\nu = \operatorname{ord}_V(f) = (\operatorname{div}(f))(V)$ is to take a regular point $p \in \mathcal{R}(V)$ and a local equation at $p, V \cap U = \{Z \in U : \varpi(Z) = 0\}$, so that V_p is irreducible and ϖ_p is prime with $\mathcal{O}_p \varpi_p = \operatorname{id}(V_p)$. The factor ϖ_p^{ν} in the unique factorization of f_p in \mathcal{O}_p defines $\nu \in \mathbb{N}_0$. This ν is independent of the choice of the regular point p and of the uniformizer ϖ .

We use $c_N = \begin{pmatrix} -1 & -N \\ 0 & 1 \end{pmatrix} \in GL(2, \mathbb{Z})$ and $C_N = \begin{pmatrix} c_N & 0 \\ 0 & c_N^* \end{pmatrix} \in K(N)$ in the following lemma. The usefulness of the following lemma is that a point on the $K(N)^+$ -orbit of a codimension one irreducible holomorphic subvariety of \mathcal{H}_2 can be found where the uniformizer ϖ is regular in ω at that point. This allows us to use the Weierstrauss preparation theorem on ϖ in the proof of Theorem 8.2. In Lemma 8.1, the term "regular" regettably has two distinct meanings. A holomorphic ϖ is regular in ω at (τ_o, z_o, ω_o) means $\varpi(\tau_o, z_o, \omega) \neq 0$ in all neighborhoods of ω_o , whereas a point p of a holomorphic subvariety V is regular if $V \cap U$ is a complex manifold for some neighborhood U of p. **Lemma 8.1.** Let $k, N \in \mathbb{N}$, and $\epsilon \in \{\pm 1\}$. Let $f \in M_k(K(N))^{\epsilon}$. Let $V \subseteq \mathcal{H}_2$ be an irreducible component of div(f). For any $g \in K(N)^+$, $V_g = g\langle V \rangle$ is also an irreducible component of div(f), and we have $\operatorname{ord}_V f = \operatorname{ord}_{V_a} f$.

There is a $g \in \{I, \mu_N, C_N\} \subseteq K(N)^+$, a regular point $p \in V_g$, a neighborhood $U \subseteq \mathcal{H}_2$ of p, and a function $\varpi \in \mathcal{O}(U)$ with an irreducible germ $\varpi_p \in \mathcal{O}_p$, and with $\varpi \begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix}$ regular in ω at p, such that $V_g \cap U = \{Z \in U : \varpi(Z) = 0\}.$

Proof. Each $g \in K(N)^+$ is biholomorphic on \mathcal{H}_2 so V is an irreducible holomorphic subvariety of codimension one if and only if V_g is. The automorphy of f shows that V is supported in div(f) if and only if V_g is, although we remark that V and V_g might be equal. Since g is biholomorphic, we have $\operatorname{ord}_V f = \operatorname{ord}_{g\langle V \rangle} f \circ g^{-1}$. The automorphy of fgives us $(f \circ g^{-1})(\Omega) = j(g^{-1}, \Omega)^k f(\Omega)$ so that f and $f \circ g^{-1}$ differ by a multiplicative unit and $\operatorname{ord}_{g\langle V \rangle} f \circ g^{-1} = \operatorname{ord}_{V_g} f$.

Every regular point $p \in V$ has a local defining equation for V given by $V \cap U = \{Z \in U : \varpi(Z) = 0\}$ for some neighborhood U of p, and some uniformizer $\varpi \in \mathcal{O}(U)$ with an irreducible germ ϖ_p represented by (ϖ, U) . We may select U so that every point of $U \cap V$ is also a regular point of V because V is a complex manifold at p. By Cartan's Theorem, we may also assume that a single function element (ϖ, U) satisfies id $V_q = \mathcal{O}_q \varpi_q$ for all $q \in U \cap V$. For the proof of the second part, we break the discussion into two mutually exclusive cases. Case I. Some regular point $p \in V$ has $\varpi \begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix}$ regular in ω or τ at p. If $\varpi \begin{pmatrix} \tau_0 & z_0 \\ z_0 & \omega \end{pmatrix} \neq 0$ in ω for some regular $p = \begin{pmatrix} \tau_0 & z_0 \\ z_0 & \omega_0 \end{pmatrix} \in V$ then we take g = I to satisfy our conclusion. Whereas if $\varpi \begin{pmatrix} \tau & z_0 \\ z_0 & \omega_0 \end{pmatrix} \neq 0$ in τ we take $g = \mu_N$ and consider $q = \begin{pmatrix} \tau_1 & z_1 \\ z_1 & \omega_1 \end{pmatrix} = \mu_N \langle p \rangle = \begin{pmatrix} \omega_0/N & -z_0 \\ -z_0 & N\tau_0 \end{pmatrix} \in V_g$. The point $q \in g \langle U \rangle$ is a regular point of V_g because p is a regular point of V and g is biholomorphic. The local uniformizer at $q \in V_g$ may be represented by $(\varpi \circ g^{-1}, g \langle U \rangle)$. In this case

$$\left(\varpi \circ g^{-1}\right)\left(\begin{smallmatrix} \tau_1 & z_1 \\ z_1 & \omega \end{smallmatrix}\right) = \varpi \left(\begin{smallmatrix} \omega/N & -z_1 \\ -z_1 & N\tau_1 \end{smallmatrix}\right) = \varpi \left(\begin{smallmatrix} \omega/N & z_o \\ z_o & \omega_o \end{smallmatrix}\right)$$

is regular in ω at q. From $V \cap U = \{Z \in U : \varpi(Z) = 0\}$ we deduce $V_g \cap g\langle U \rangle = \{Z \in g\langle U \rangle : (\varpi \circ g^{-1})(Z) = 0\}$ so that q and $g\langle U \rangle$ play the role of p and U in the statement of the lemma. Case II. Every regular point $p \in V$ has $\varpi \begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix}$ not regular in ω and not regular in τ at p. Consider any regular point $p_o = \begin{pmatrix} \tau_o & z_o \\ z_o & \omega_o \end{pmatrix} \in V$. Take a polydisk $\Delta = \Delta_{11} \times \Delta_{12} \times \Delta_{22} \subseteq U$ about p_o and use the function element (ϖ, Δ) . We will ultimately show that Case II is very special and that $V \cap \Delta = \{Z \in \Delta : z - z_o = 0\}$. We claim that $\varpi \begin{pmatrix} \tau & z_o \\ z_o & \omega \end{pmatrix} = 0$ for all $\begin{pmatrix} \tau & z_o \\ z_o & \omega \end{pmatrix} \in \Delta$.

If not there is some $r_o = \begin{pmatrix} \tau_1 & z_o \\ z_o & \omega_1 \end{pmatrix} \in \Delta$ with $\varpi(r_o) \neq 0$. Since ϖ is not regular in ω at p_o we have $\varpi\left(\begin{smallmatrix} \tau_o & z_o \\ z_o & \omega \end{smallmatrix}\right) \equiv 0$ for ω in some neighborhood of ω_o , hence necessarily for all $\omega \in \Delta_{22}$. Therefore $\varpi \begin{pmatrix} \tau_o & z_o \\ z_o & \omega_1 \end{pmatrix} = 0$. The setting of Case II applies equally well to $r = \begin{pmatrix} \tau_o & z_o \\ z_o & \omega_1 \end{pmatrix} \in V \cap \Delta$, which is still a regular point of V, so that ϖ is not regular in τ at r, and $\varpi\left(\begin{smallmatrix} \tau & z_o \\ z_o & \omega_1 \end{smallmatrix}\right) \equiv 0$ for τ in a neighborhood of τ_o , and hence for all $\tau \in \Delta_{11}$. Thus $\varpi(r_o) = \varpi\left(\begin{smallmatrix} \tau_1 & z_o \\ z_o & \omega_1 \end{smallmatrix}\right) = 0$, contrary to our supposition. Thus $\varpi \begin{pmatrix} \tau & z_o \\ z_o & \omega \end{pmatrix} = 0$ for all $\begin{pmatrix} \tau & z_o \\ z_o & \omega \end{pmatrix} \in \Delta$. Therefore ϖ vanishes on $\{Z \in Z \in Z \}$ $\Delta : z - z_o = 0$ and so the irreducible $z - z_o$ divides ϖ in Δ ; however, the germ ϖ_{p_o} of ϖ at p_o is irreducible and so ϖ and $z - z_o$ differ by a multiplicative unit in Δ , after possibly shrinking Δ further. Therefore in Case II we have the possible but special circumstance $V \cap \Delta = \{Z \in Z\}$ $\Delta: z - z_o = 0$. Without loss of generality we may adjust the selection of ϖ by a unit and assume that $\varpi \begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} = z - z_o$ in Δ . Now set $g = C_N$ and consider $q = \begin{pmatrix} \tau_1 & z_1 \\ z_1 & \omega_1 \end{pmatrix} = C_N \langle p_o \rangle = \begin{pmatrix} \tau_o + 2Nz_o + N^2\omega_o & z_o + N\omega_o \\ z_o + N\omega_o & \omega_o \end{pmatrix} \in V_g$. The local uniformizer at $q \in V_g$ may be represented by $(\varpi \circ g^{-1}, g \langle \Delta \rangle)$. For this choice we have

$$(\varpi \circ g^{-1}) \begin{pmatrix} \tau_1 & z_1 \\ z_1 & \omega \end{pmatrix} = \varpi \begin{pmatrix} \tau_1 - 2Nz_1 + N^2\omega & z_1 - N\omega \\ z_1 - N\omega & \omega \end{pmatrix}$$
$$= \varpi \begin{pmatrix} \tau_o + N^2(\omega - \omega_o) & z_o + N(\omega_o - \omega) \\ z_o + N(\omega_o - \omega) & \omega \end{pmatrix} = N(\omega_o - \omega)$$

which is regular in ω of order 1 at q. Thus the conclusion of the lemma holds for q and $g\langle\Delta\rangle$ in the role of p and U.

Theorem 8.2. Let $k, k_0, N \in \mathbb{N}$ and $\epsilon, \epsilon_0 \in \{\pm 1\}$. Let $\Gamma \subseteq \Gamma^0(N)$ be a subgroup of finite index. Let $\mathfrak{f} \in \mathbb{S}(k, N, \epsilon; \Gamma)$. Let $f_0 \in M_{k_0}(K(N))^{\epsilon_0}$ be nontrivial and $G \in M_{k_0+k}(K(N))^{\epsilon_0\epsilon}$. Let $V \subseteq \mathcal{H}_2$ be an irreducible component of div (f_0) .

If $\mathrm{FJ}(G) = \mathfrak{f} \mathrm{FJ}(f_0)$ in $\mathbb{S}(k_0 + k, N, \epsilon_0 \epsilon; \Gamma)$ then $\mathrm{ord}_V f_0 \leq \mathrm{ord}_V G$. The meromorphic G/f_0 is holomorphic and $G/f_0 \in M_k(K(N))^{\epsilon}$.

Proof. By Lemma 8.1 we can find a point $p \in V_g$, for some $g \in K(N)^+$, where the local defining equation $V_g \cap U = \{Z \in U : \varpi(Z) = 0\}$ has the following nice properties. The point p is a regular point of V_g , the uniformizer ϖ is regular in ω at p, and the germ ϖ_p is prime in \mathcal{O}_p . By Lemma 8.1, in order to prove $\operatorname{ord}_V f_0 \leq \operatorname{ord}_V G$ it suffices to prove $\operatorname{ord}_{V_g} f_0 \leq \operatorname{ord}_{V_g} G$, so we may rename V_g as V without loss of generality.

In this paragraph we outline the remainder of the proof. We will find a neighborhood U_1 of p such that every point $q \in U_1 \cap V$ has the same nice properties that p does. We use this to twice move to nearby points. First we select $p_1 \in U_1 \cap V$ so that ϖ_{p_1} is the only irreducible factor of G_{p_1} that vanishes at p_1 . Then we use the Weierstrauss preparation theorem on ϖ_{p_1} to find a $p_2 \in \mathcal{W}(\Gamma) \cap V \cap U_1$, and a holomorphic curve Ω_2 inside $\mathcal{W}(\Gamma) \cap V$ and passing through p_2 , which guarantees that the formal series \mathfrak{f} converges on $\Omega_2(\omega)$. The conclusion about the orders of vanishing readily follows from the convergence of \mathfrak{f} on the holomorphic curve.

The set of regular points $\mathcal{R}(V)$ is open in V. Regularity in ω at p is a local condition in p because regularity of order less than or equal to ν at p is implied by the nonvanishing of a partial derivative $\partial_{\omega}^{\nu} \varpi(p)$. By Cartan's Theorem, there is a neighborhood $U'' \subseteq U$ of p such that ϖ_q generates id V_q at all points $q \in U'' \cap V$. Accordingly there is a smaller neighborhood U_1 of p where every $q \in U_1 \cap V$ has the nice properties $q \in \mathcal{R}(V)$, ϖ regular in ω at q, and ϖ_q prime in \mathcal{O}_q .

In the local ring \mathcal{O}_p , factor $(f_0)_p = (\Delta_1)_p \varpi_p^{\nu(f_0)}$ and $G_p = (\Delta_2)_p \varpi_p^{\nu(G)}$, where $(\Delta_1)_p, (\Delta_2)_p$ are finite products of irreducibles that are not associate to ϖ_p , and where $\nu(f_0) = \operatorname{ord}_V f_0$ and $\nu(G) = \operatorname{ord}_V G$ by the direct way of computing the vanishing order on V. We select a neighborhood $N_1 \subseteq U_1$ where the germs of the above factors all have representative function elements. If $\Delta_2(q) = 0$ for all $q \in N_1 \cap V$ then $(\Delta_2)_p$ vanishes on V_p and ϖ_p divides $(\Delta_2)_p$ in \mathcal{O}_p because ϖ_p is prime, contradicting the fact that $(\Delta_2)_p$ is a finite product of irreducibles that are not associate to ϖ_p in \mathcal{O}_p . Hence there is a point $p_1 \in V \cap N_1$ such that $\Delta_2(p_1) \neq 0$. By shrinking N_1 , we may assume that Δ_2 is nonzero on N_1 .

We apply the Weierstrauss preparation theorem to ϖ , which is regular in ω at $p_1 = \begin{pmatrix} \tau_1 & z_1 \\ z_1 & \omega_1 \end{pmatrix}$. There exist neighborhoods U_2 of (τ_1, z_1) and W_2 of ω_1 such that, $N_2 = U_2 \times W_2 \subseteq N_1$ and

$$\{Z = \begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} \in N_2 : \varpi(Z) = 0\} = \{Z = \begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} \in N_2 : W(\tau, z, \omega) := \sum_{j=0}^t h_j(\tau, z)(\omega - \omega_1)^{t-j} = 0\},\$$

for some $t \in \mathbb{N}$, $h_j \in \mathcal{O}(U_2)$ for $j = 0, \ldots, t$, and $h_0 = 1$. The set $\mathcal{W}_1(\Gamma)$ is dense in U_2 by Lemma 6.6, so by taking $(\tau_2, z_2) \in \mathcal{W}_1(\Gamma)$ close enough to $(\tau_1, z_1) \in U_2$ so that we may choose a root ω_2 of $W(\tau_2, z_2, \omega)$ close enough to ω_1 , we have

$$p_2 = \begin{pmatrix} \tau_2 & z_2 \\ z_2 & \omega_2 \end{pmatrix} \in V \cap N_2 \cap \mathcal{W}(\Gamma).$$

From $N_2 \subseteq N_1$ we inherit the representative function elements f_0 , G, Δ_1 , Δ_2 , and ϖ , so that $f_0 = \Delta_1 \varpi^{\nu(f_0)}$ and $G = \Delta_2 \varpi^{\nu(G)}$ in N_2 , and Δ_2 is nonzero on N_2 .

The formal series of G is given by $FJ(G) = \mathfrak{f} FJ(f_0)$. Recalling that $(\tau_2, z_2) \in \mathcal{W}_1(\Gamma)$, and using Lemma 7.2 to apply $A(\tau_2, z_2)$, we have the equality $A(\tau_2, z_2)FJ(G) = (A(\tau_2, z_2)\mathfrak{f})(A(\tau_2, z_2)FJ(f_0))$ of convergent

power series on $D_2 = \{\xi \in \mathbb{C} : |\xi| < r_2\}$ for $r_2 = e^{-2\pi\eta_2}$ and $\eta_2 = (\operatorname{Im}(z_2))^2/\operatorname{Im}(\tau_2)$. Also set $\Omega_2(\omega) = \begin{pmatrix} \tau_2 & z_2 \\ z_2 & \omega \end{pmatrix} \in \mathcal{H}_2$ for $\omega \in N_{\infty}(\eta_2)$. For $\omega \in N_{\infty}(\eta_2)$ we have $e(\omega) \in D_2$ and may apply the evaluation homomorphism $[e(\omega)] : \mathcal{S}(D_2) \to \mathbb{C}$ to obtain

$$(A(\tau_2, z_2)\mathrm{FJ}(G))[e(\omega)] = (A(\tau_2, z_2)\mathfrak{f})[e(\omega)] (A(\tau_2, z_2)\mathrm{FJ}(f_0))[e(\omega)].$$

Making use of Lemma 7.2, this may be written on $W_2 \cap N_{\infty}(\eta_2)$, which contains a neighborhood of ω_2 , as

$$G(\Omega_2(\omega)) = H(\tau_2, z_2, \mathfrak{f})(\omega) f_0(\Omega_2(\omega)).$$

By specializing the factorizations of G and f_0 in $\mathcal{O}(N_2)$ to the holomorphic curve $\Omega_2(\omega) = \begin{pmatrix} \tau_2 & z_2 \\ z_2 & \omega \end{pmatrix}$ we have

$$\Delta_2(\Omega_2(\omega)) \varpi (\Omega_2(\omega))^{\nu(G)} = H(\tau_2, z_2, \mathfrak{f})(\omega) \Delta_1(\Omega_2(\omega)) \varpi (\Omega_2(\omega))^{\nu(f_0)}$$

Note that $\varpi(\Omega_2(\omega))$ has at most t zeros on W_2 and is hence nontrivial. Of the two cases $\nu(G) \geq \nu(f_0)$, and $\nu(G) < \nu(f_0)$, the first is our conclusion, so we will conclude the proof by showing that the second does not occur. In the second case, by cancelling powers of the nontrivial $\varpi(\Omega_2(\omega))$, we have

$$\Delta_2(\Omega_2(\omega)) = H(\tau_2, z_2, \mathfrak{f})(\omega) \,\Delta_1(\Omega_2(\omega)) \,\varpi \left(\Omega_2(\omega)\right)^{\nu(f_0) - \nu(G)}$$

We evaluate these at $\omega_2 \in W_2 \cap N_{\infty}(\eta_2)$ to obtain

$$\Delta_2(p_2) = H(\tau_2, z_2, \mathfrak{f})(\omega_2) \,\Delta_1(p_2) \left(0^{\nu(f_0) - \nu(G)} \right) = 0,$$

which contradicts $\Delta_2(p_2) \neq 0$. Thus we have $\nu(G) \geq \nu(f_0)$.

A meromorphic function with an effective divisor is holomorphic. Therefore G/f_0 is holomorphic and continuous on \mathcal{H}_2 . There is an open dense subset of \mathcal{H}_2 where G/f_0 transforms like an element of $M_k(K(N))^{\epsilon}$. By the continuity of G/f_0 and of the factor of automorphy, G/f_0 transforms like an element of $M_k(K(N))^{\epsilon}$ on \mathcal{H}_2 and hence $G/f_0 \in M_k(K(N))^{\epsilon}$.

9. MAIN THEOREM.

Theorem 9.1. Let $U \subseteq \mathbb{C}^d$ be open. Let $\{f_j\}$ be a locally bounded sequence of holomorphic functions on U that converges on a dense subset of U. Then the sequence $\{f_j\}$ converges on U and uniformly on compact subsets of U.

Proof. This is Exercise 4a of section 4 in Chapter 1 of [14].

The proof given here imitates the proof of Lemma IV.4.8 in [13] for the one-dimensional case. It suffices to show that $\{f_j\}$ is uniformly Cauchy on compact subsets of Ω . Pick a compact $K \subseteq \Omega$ and an $\epsilon > 0$. There is no loss of generality in assuming that K is a closed ball. We will construct an $M \in \mathbb{N}$ from the given data $\{f_j\}, K$, and ϵ .

The family $\{f_j\}$ is equicontinuous on K because it is locally bounded, see pages 11-12 of [31]. Select $\delta > 0$ to enforce this equicontinuity on K for $\frac{1}{3}\epsilon$.

(13)
$$\forall z, w \in K, \forall n \in \mathbb{N}, |z - w| < \delta \implies |f_n(z) - f_n(w)| < \frac{1}{3}\epsilon.$$

Cover K with open balls inside Ω at each point of K with radii that are less than $\frac{1}{2}\delta$. By the compactness of K we have $K \subseteq \bigcup_{i=1}^{\ell} B(z_i, \delta_i) \subseteq \Omega$ for some $z_i \in K$, and $0 < \delta_i < \frac{1}{2}\delta$. Since S is dense in K, we can pick an $s_i \in K \cap S \cap B(z_i, \delta_i)$; this uses that K is a closed ball. Note that for every point $z \in B(z_i, \delta_i)$ we have $|z - s_i| < \delta$ and therefore $|f_n(z) - f_n(s_i)| < \frac{1}{3}\epsilon$ by (13). By hypothesis each sequence $f_j(s_i)$ converges and so is Cauchy. Select $M_i > 0$ so that

(14)
$$\forall m, n \in \mathbb{N}, m, n > M_i \implies |f_m(s_i) - f_n(s_i)| < \frac{1}{3}\epsilon$$

Let $M = \max_{1 \le i \le \ell} M_i$ be the promised natural number. We have

$$|f_m(z) - f_n(z)| \le |f_m(z) - f_m(s_i)| + |f_m(s_i) - f_n(s_i)| + |f_n(s_i) - f_n(z)|$$

for any $m, n \in \mathbb{N}, z \in K$, and $1 \leq i \leq \ell$. If m, n > M, then $m, n > M_i$ and so $|f_m(s_i) - f_n(s_i)| < \frac{1}{3}\epsilon$ by (14). Choose *i* so that $z \in B(z_i, \delta_i)$; then $|f_j(z) - f_j(s_i)| < \frac{1}{3}\epsilon$ for all $j \in \mathbb{N}$ by (13). Therefore, for all $m, n \in \mathbb{N}$ with m, n > M, and all $z \in K$, we have $|f_m(z) - f_n(z)| < \epsilon$, and the sequence $\{f_j\}$ is uniformly Cauchy on K. \Box

The essence of the proof of the main result, Theorem 9.6, lies in the argument for the following special case of Fricke plus cusp forms. The general case will be reduced to Theorem 9.2.

Theorem 9.2. The map $FJ : S_k(K(N)^+) \to S(k, N, +)$ is an isomorphism for $N, k \in \mathbb{N}$.

Proof. Since the map FJ is injective, the only issue is surjectivity. Take a nontrivial $\mathfrak{f} \in \mathbb{S}(k, N, +)$. By Proposition 5.4, there is a subgroup Γ of finite index in $\Gamma^0(N)$ such that $\mathfrak{f} \in \mathbb{M}(k, N, +; \Gamma)$, which implies $\mathfrak{f} \in \mathbb{M}(k, N, +; \Gamma) \cap \mathbb{S}(k, N, +) = \mathbb{S}(k, N, +; \Gamma)$. Completely separately, by Proposition 4.7, \mathfrak{f} satisfies a polynomial relation in $\mathbb{M}(k_0+dk, N, +)$,

$$\mathrm{FJ}(f_0)\mathfrak{f}^d + \mathrm{FJ}(f_1)\mathfrak{f}^{d-1} + \dots + \mathrm{FJ}(f_j)\mathfrak{f}^{d-j} + \dots + \mathrm{FJ}(f_d) = 0,$$

for some $d \in \mathbb{N}$, $k_0 \in \mathbb{N}_0$, and some $f_j \in M_{k_0+jk}(K(N)^+)$ with f_0 not identically zero. Since $\mathrm{FJ}(f_j)$, $\mathfrak{f} \in \mathbb{M}(N, +; \Gamma)$, this polynomial relation is actually in $\mathbb{M}(k_0 + dk, N, +; \Gamma)$. Since $\mathbb{S}(N, +; \Gamma)$ is a graded ideal

in $\mathbb{M}(N, +; \Gamma)$, set

$$\mathfrak{g} = \mathrm{FJ}(f_0)\mathfrak{f} = \sum_{m=1}^{\infty} \phi_m \xi^{Nm} \in \mathbb{S}(k+k_0, N, +; \Gamma).$$

Then \mathfrak{g} is integral over FJ $(M(K(N)^+))$ and satisfies a monic polynomial relation in $\mathbb{M}((k+k_0)d, N, +; \Gamma)$ given by

$$\mathfrak{g}^d + \mathrm{FJ}(g_1)\mathfrak{g}^{d-1} + \cdots + \mathrm{FJ}(g_j)\mathfrak{g}^{d-j} + \cdots + \mathrm{FJ}(g_d) = 0$$

with $g_j = f_0^{j-1} f_j \in M_{(k+k_0)j}(K(N)^+)$ for $j = 1, \dots, d$.

By Theorem 7.4, the sequence of partial sums $\sum_{m=1}^{M} \phi_m(\tau, z) e(Nm\omega)$ is locally bounded for $\begin{pmatrix} \tau & z \\ z & \omega \end{pmatrix} \in \mathcal{H}_2$. By Corollary 6.9, we also know that the formal series \mathfrak{g} converges on the dense subset $\mathcal{W}(\Gamma)$ of \mathcal{H}_2 , noting $\mathfrak{g} \in \mathfrak{S}(k+k_0, N, +; \Gamma) \subseteq \mathfrak{S}(k+k_0, N; \Gamma)$. Therefore, by Theorem 9.1, the sequence of partial sums converges uniformly on compact sets of \mathcal{H}_2 . The limiting function $G \in \mathcal{O}(\mathcal{H}_2)$ is given by $G(\Omega) =$ $\sum_{m=1}^{\infty} \phi_m(\tau, z) e(Nm\omega)$ and satisfies $G|_{k+k_0}g = G$ for all $g \in \mathcal{P}_{2,1}(\mathbb{Z})$ because each term $\phi_m(\tau, z)e(Nm\omega)$ is so invariant. The holomorphic function G is hence periodic with respect to the translation lattice $\{ \begin{pmatrix} I & s \\ 0 & I \end{pmatrix} \in \mathcal{P}_{2,1}(\mathbb{Z}) : s \in M^{\text{sym}}_{2 \times 2}(\mathbb{Z}) \}$ and has a Fourier series that converges absolutly and uniformly on compact sets of \mathcal{H}_2 . This absolute convergence of the Fourier series of G is the crux of the matter because it will allow us to rearrange the order of summation and thereby deduce the crucial invariance property $G|_{k+k_0}\mu_N = G$ from the involution condition $c(n,r;\phi_m) = (-1)^k \epsilon c(m,r;\phi_n)$ on \mathfrak{g} . Since the convergence of the sequence of partial sums is uniform on compact subsets, the Fourier coefficients of G are supported on $\overline{\mathcal{X}}(N) \subseteq M^{\text{sym}}_{2\times 2}(\mathbb{Q})$ and, for $t = \begin{pmatrix} n & r/2 \\ r/2 & Nm \end{pmatrix} \in \bar{\mathcal{X}}(N)$, satisfy $a(t;G) = c(n,r;\phi_m)$. For \mathfrak{g} in this proof we have $\epsilon = +1$. Recall that $t \mapsto F_N^{-1} t F_N^*$ is an involution of $\bar{\mathcal{X}}(N)$ satisfying $F_N^{-1} \begin{pmatrix} n & r/2 \\ r/2 & Nm \end{pmatrix} F_N^* = \begin{pmatrix} m & -r/2 \\ -r/2 & Nn \end{pmatrix}$, and that absolutely convergent summations over $t \in \overline{\mathcal{X}}(N)$ may instead be rearranged to be taken over $F_N^{-1}tF_N^*$.

$$\begin{aligned} (G|\mu_N)\left(\Omega\right) &= G(F_N^*\Omega F_N^{-1}) = \sum_{t\in\bar{\mathcal{X}}(N)} a(t;G)e\left(\langle F_N^*\Omega F_N^{-1},t\rangle\right) \\ &= \sum_t a(t;G)e\left(\langle\Omega,F_N^{-1}tF_N^*\rangle\right) = \sum_t a(F_N^{-1}tF_N^*;G)e\left(\langle\Omega,t\rangle\right) \\ &= \sum_{t=\binom{n-r/2}{r/2}\in\bar{\mathcal{X}}(N)} c(m,-r;\phi_n)e\left(\langle\Omega,t\rangle\right) = \sum_t c(n,r;\phi_m)e\left(\langle\Omega,t\rangle\right) \end{aligned}$$

$$= \sum_t a(t;G) e\left(\langle \Omega,t\rangle \right) = G(\Omega).$$

Combining these automorphic properties of G we obtain $G|\sigma = G$ for all $\sigma \in \langle P_{2,1}(\mathbb{Z}), \mu_N \rangle = \langle K(N), \mu_N \rangle = K(N)^+$, see Gritsenko [17, 18]. Therefore $G \in M_{k+k_0}(K(N)^+)$ and $FJ(G) = \mathfrak{g} \in \mathbb{S}(k + k_0, N, +; \Gamma)$. Together these imply that $G \in S_{k+k_0}(K(N)^+)$ by Reefschläger's decomposition [34], as in Lemma 3.1.

We have $\operatorname{FJ}(G) = \mathfrak{g} = \operatorname{FJ}(f_0)\mathfrak{f}$ for the formal series $\mathfrak{f} \in \mathbb{S}(k, N, +; \Gamma)$, and the paramodular forms $f_0 \in M_{k_0}(K(N)^+)$, $G \in S_{k+k_0}(K(N)^+)$. By Theorem 8.2, $f := G/f_0 \in M_k(K(N)^+)$. From $f_0f = G$ we see that $\operatorname{FJ}(f_0)\operatorname{FJ}(f) = \operatorname{FJ}(G)$. Combining this with $\operatorname{FJ}(f_0)\mathfrak{f} = \operatorname{FJ}(G)$, we have $\operatorname{FJ}(f_0)(\operatorname{FJ}(f) - \mathfrak{f}) = 0$, which implies $\operatorname{FJ}(f) = \mathfrak{f}$ since $\mathbb{M}(N)$ is an integral domain and f_0 is nontrivial. By Reefschläger's result, $\operatorname{FJ}(f) =$ $\mathfrak{f} \in \mathbb{S}(k, N, +; \Gamma)$ and $f \in M_k(K(N)^+)$ imply $f \in S_k(K(N)^+)$, which proves that FJ is surjective. \Box

Lemma 9.3. There is a nontrivial cusp form $\chi_N \in S_{11}(K(N))^-$ for N > 1.

Proof. There are a number of possible approaches to proving the existence of nontrivial minus cusp forms. We might use an asymptotic trace formula or the oldform theory of Roberts and Schmidt [35, 36]. Perhaps the briefest is to use the injectivity of the Gritsenko lift [18], Grit : $J_{k,m}^{\text{cusp}} \to S_k (K(m))^{\epsilon}$ for $\epsilon = (-1)^k$. An estimate from the dimension formula [11] is dim $J_{11,m}^{\text{cusp}} \geq \frac{7}{4}$ floor $(\frac{m-3}{6}) + \frac{7}{12}$ for $m \in \mathbb{N}$. Considering dim $J_{11,2}^{\text{cusp}} = 1$, we see dim $J_{11,m}^{\text{cusp}} \geq 1$ for $m \geq 2$. Thus there is a nontrivial Gritsenko lift χ_N in $S_{11} (K(N))^-$ for N > 1.

Theorem 9.4. The map $FJ : S_k(K(N))^{\epsilon} \to \mathbb{S}(k, N, \epsilon)$ is an isomorphism for $N, k \in \mathbb{N}$, and $\epsilon \in \{\pm 1\}$.

Proof. The case $\epsilon = +1$ is Theorem 9.2, so assume $\epsilon = -1$ and N > 1. Take $\mathfrak{f} \in \mathbb{S}(k, N, -)$. There is a nontrivial cusp form $\chi_N \in S_{11}(K(N))^$ by Lemma 9.3. Multiply \mathfrak{f} by $\mathrm{FJ}(\chi_N)$ to obtain a new formal series $\mathfrak{g} = \mathrm{FJ}(\chi_N)\mathfrak{f} \in \mathbb{S}(k + 11, N, +)$. By Theorem 9.2, there exists a $G \in S_{k+11}(K(N))^+$ such that $\mathrm{FJ}(G) = \mathfrak{g} = \mathrm{FJ}(\chi_N)\mathfrak{f}$. Both $\mathrm{FJ}(G)$ and $\mathrm{FJ}(\chi_N)$ have $\Gamma^0(N)$ -symmetries and we will show that \mathfrak{f} does too. Apply the ring homomorphism AFS : $\mathbb{M}(N) \to \mathbb{C}^{\bar{\mathcal{X}}(N)}$ of section 5 to $\mathrm{FJ}(G) = \mathrm{FJ}(\chi_N)\mathfrak{f}$ to obtain $\mathrm{FS}(G) = \mathrm{FS}(\chi_N)\mathrm{AFS}(\mathfrak{f})$. Now, for $\sigma \in \Gamma^0(N)$, apply the automorphism $\mathfrak{j}(\sigma)$ of Lemma 5.3, which fixes $\mathrm{FS}(G)$ and $\mathrm{FS}(\chi_N)$ to obtain $0 = \mathrm{FS}(\chi_N)(\mathrm{AFS}(\mathfrak{f}) - \mathfrak{j}(\sigma)\mathrm{AFS}(\mathfrak{f}))$. Since $\mathbb{C}^{\bar{\mathcal{X}}(N)}$ is an integral domain and $\mathrm{FS}(\chi_N)$ is nontrivial, we have

 $j(\sigma)AFS(\mathfrak{f}) = AFS(\mathfrak{f})$ for all $\sigma \in \Gamma^0(N)$. By Lemma 5.3, we conclude that $\mathfrak{f} \in \mathbb{S}(k, N, -; \Gamma^0(N))$. By Theorem 8.2 with $\Gamma = \Gamma^0(N)$, the equation $FJ(G) = FJ(\chi_N)\mathfrak{f}$ implies that $f:=G/\chi_N$ is holomorphic, and, consequently, that $f \in M_k(K(N))^-$. From $FJ(G) = FJ(\chi_N)\mathfrak{f}$ and $FJ(G) = FJ(\chi_N)FJ(f)$ we obtain $0 = FJ(\chi_N)(\mathfrak{f} - FJ(f))$. Since $\mathbb{M}(N)$ is an integral domain and $FJ(\chi_N)$ is nontrivial, we have $FJ(f) = \mathfrak{f}$. By Reefschläger's result and $\mathfrak{f} \in \mathbb{S}(k, N, -; \Gamma)$ we have $f \in S_k(K(N))^-$. Thus FJ is surjective.

Lemma 9.5. Let $N \in \mathbb{N}$ and $I = [K(N)^+ : K(N)^+ \cap K(1)]$. Let $p \in \mathcal{H}_2$. There is a cusp form $\chi_p \in S_{60I}(K(N)^+)$ such that $\chi_p(p) \neq 0$.

Proof. The divisor of $\chi_{10} \in S_{10}(K(1))$ is the reducible locus R,

 $R = \bigcup_{\sigma \in K(1)} \{ \sigma \langle \begin{pmatrix} \tau & 0 \\ 0 & \omega \end{pmatrix} \rangle \in \mathcal{H}_2 : \tau, \omega \in \mathcal{H}_1 \}.$

For $\psi_{12} \in S_{12}(K(1))$, we have $\psi_{12} \begin{pmatrix} \tau & 0 \\ 0 & \omega \end{pmatrix} = \Delta(\tau)\Delta(\omega)$ for the nonvanishing $\Delta \in S_{12}(\mathrm{SL}(2,\mathbb{Z}))$. Thus χ_{10} and ψ_{12} have no common zeros on \mathcal{H}_2 . Pick *I* coset representatives σ for $K(N)^+ = \coprod_{\sigma} (K(N)^+ \cap K(1))\sigma$. For any $(\alpha, \beta) \in \mathbb{C}^2 \setminus \{(0, 0)\}$, consider the nontrivial norm

$$H^{\alpha,\beta} = \prod_{[\sigma] \in (K(N)^+ \cap K(1)) \setminus K(N)^+} \left(\alpha \chi_{10}^6 + \beta \psi_{12}^5 \right) |_{60} \sigma \in S_{60I} \left(K(N)^+ \right).$$

We have $H^{\alpha,\beta}(p) = (\prod_{\sigma} j(\sigma, p)^{-60}) \prod_{\sigma} (\alpha \chi_{10}^6(\sigma \langle p \rangle) + \beta \psi_{12}^5(\sigma \langle p \rangle))$. Pick any $(\alpha_o, \beta_o) \in \mathbb{C}^2$ that is not on any of the finitely many codimension one complex lines through the origin $\{(z_1, z_2) \in \mathbb{C}^2 : z_1 \chi_{10}^6(\sigma \langle p \rangle) + z_2 \psi_{12}^5(\sigma \langle p \rangle) = 0\}$. The cusp form $\chi_p = H^{\alpha_o,\beta_o}$ is as claimed. \Box

Theorem 9.6 (Main Theorem). Let $N \in \mathbb{N}$, $k \in \mathbb{N}_0$, and $\epsilon \in \{\pm 1\}$. The map $\mathrm{FJ} : M_k(K(N))^{\epsilon} \to \mathbb{M}(k, N, \epsilon)$ is an isomorphism. The map $\mathrm{FJ} : S_k(K(N))^{\epsilon} \to \mathbb{S}(k, N, \epsilon)$ is also an isomorphism.

Proof. The case k = 0 is easy so assume $k \in \mathbb{N}$. Take $\mathfrak{f} \in \mathbb{M}(k, N, \epsilon)$. For any point $p \in \mathcal{H}_2$ there is a cusp form $\chi_p \in S_{60I}(K(N)^+)$ such that $\chi_p(p) \neq 0$, as in Lemma 9.5. Define a corresponding product of formal series $\mathfrak{g}_p = \mathrm{FJ}(\chi_p)\mathfrak{f} \in \mathbb{S}(k + 60I, N, \epsilon)$. By Theorem 9.4, there exists a $G_p \in S_{k+60I}(K(N))^{\epsilon}$ such that $\mathrm{FJ}(G_p) = \mathfrak{g}_p = \mathrm{FJ}(\chi_p)\mathfrak{f}$. Note that $S_{k+60I}(K(N))^{\epsilon} = S_{k+60I}(K(N)^+, \lambda_{\epsilon})$ where $\lambda_{\epsilon} : K(N)^+ \to \{\pm 1\}$ is the character defined by $\lambda_{\epsilon}(\mu_N) = \epsilon$ and $\lambda_{\epsilon} = 1$ on K(N).

In the same way, for any point $q \in \mathcal{H}_2$ distinct from p we also have $\mathrm{FJ}(G_q) = \mathrm{FJ}(\chi_q)\mathfrak{f}$. Since $\mathbb{M}(N)$ is an integral domain we necessarily have $\mathrm{FJ}(G_p)\mathrm{FJ}(\chi_q) - \mathrm{FJ}(G_q)\mathrm{FJ}(\chi_p) = 0$, or $G_p\chi_q = G_q\chi_p$. Therefore we have $\frac{G_p}{\chi_p} = \frac{G_q}{\chi_q}$ whenever both quotients are defined. We may define $f \in \mathcal{O}(\mathcal{H}_2)$ by setting $f = \frac{G_p}{\chi_p}$ in neighborhoods where the denominator

AOKI, IBUKIYAMA, POOR

does not vanish, and by noting this is independent of the choice of p. Therefore we have $\chi_p f = G_p$, $\chi_p |\sigma f| \sigma = G_p |\sigma$, and $\chi_p f |\sigma = \lambda_{\epsilon}(\sigma) G_p$, for any point p and any $\sigma \in K(N)^+$. Thus $\chi_p (f |\sigma - \lambda_{\epsilon}(\sigma) f) = 0$. Since M(K(N)) is an integral domain and χ_p is nontrivial, we have $f |\sigma = \lambda_{\epsilon}(\sigma) f$ and $f \in M_k(K(N)^+, \lambda_{\epsilon}) = M_k(K(N))^{\epsilon}$. It remains to show $FJ(f) = \mathfrak{f}$. This follows from $FJ(G_p) = FJ(\chi_p)\mathfrak{f}$, $FJ(G_p) =$ $FJ(\chi_p)FJ(f)$, and the fact that $\mathfrak{M}(N)$ is an integral domain and χ_p is nontrivial. The result for cusp forms follows from the general result because, by Lemma 3.2, the inverse image of $\mathfrak{S}(k, N, \epsilon)$ under FJ is $S_k (K(N))^{\epsilon}$. \Box

To help prove Corollary 1.2 we use a lemma from linear algebra.

Lemma 9.7. Let U, V, and W be finite dimensional vector spaces. Let $L': U \to W$ and $L'': V \to W$ be linear maps. Define a subspace $X \subseteq U \times V$ by $X = \{(u, v) \in U \times V : L'u = L''v\}$. Define the projection $\pi_U: X \to U$ by $\pi_U(u, v) = u$. If L'' is injective then π_U is injective, and conversely.

Proof. We have $\ker(\pi_U) = \{(0, v) \in U \times V : 0 = L''v\} = \{0\} \times \ker(L'')$. Therefore $\ker(\pi_U) = \{(0, 0)\}$ if and only if $\ker(L'') = \{0\}$. \Box

We now prove Corollary 1.2 from the Introduction.

Proof of Corollary 1.2. To prove that the sequence dim $\mathbb{M}(k, N, \epsilon)[d]$ is monotonically decreasing for $d \geq \frac{1}{6}Nk$, it suffices to prove that the projection $\pi_d : \mathbb{M}(k, N, \epsilon)[d+1] \to \mathbb{M}(k, N, \epsilon)[d]$, sending $(\phi_m)_{m=0}^{d+1}$ to $(\phi_m)_{m=0}^d$, injects. To employ the notation of Lemma 9.7, we set $U = \mathbb{M}(k, N, \epsilon)[d]$ and $V = J_{k,N(d+1)}$. To define W, first define the finite set $I = \{(m, r) \in \mathbb{Z}^2 : 0 \leq m \leq d, 4N(d+1)m \geq r^2\}$ and set $W = \mathbb{C}^{|I|}$. Define a linear map $L' : U \to W$ by $L'\left((\phi_m)_{m=0}^d\right) =$ $(c(d+1, r; \phi_m))_{(m,r)\in I}$, and a linear map $L'' : V \to W$ by $L''(\phi_{d+1}) =$ $\left((-1)^k \epsilon c(m, r; \phi_{d+1})\right)_{(m,r)\in I}$. From the definition of L'' we see that the kernel of L'' consists of the Jacobi forms vanishing to order d + 1, so that $\ker(L'') = J_{k,N(d+1)}(d+1)$. From equation (1) we see that $\mathbb{M}(k, N, \epsilon)[d] \subseteq X$ for

$$X = \{ ((\phi_m)_{m=0}^d, \phi_{d+1}) \in U \times V : L' \left((\phi_m)_{m=0}^d \right) = L''(\phi_{d+1}) \}.$$

Since $d+1 > \frac{1}{6}Nk$, Proposition 4.5 implies $\ker(L'') = J_{k,N(d+1)}(d+1) = \{0\}$, so that $\pi_U : X \to U$ injects by Lemma 9.7. Since π_d is the restriction of π_U to $\mathbb{M}(k, N, \epsilon)[d+1]$, we have shown that the projection $\pi_d : \mathbb{M}(k, N, \epsilon)[d+1] \to \mathbb{M}(k, N, \epsilon)[d]$ injects for $d \geq \frac{1}{6}Nk$.

For the inverse system $\pi_d : \mathbb{M}(k, N, \epsilon)[d+1] \to \mathbb{M}(k, N, \epsilon)[d]$ we have $\mathbb{M}(k, N, \epsilon) = \varprojlim_d \mathbb{M}(k, N, \epsilon)[d]$. The reason for this is that any equation $c(n, r; \phi_m) = (-1)^k \epsilon c(m, r; \phi_m)$ imposed by equation (1) on $\mathfrak{f} = (\phi_m)_{m=0}^{\infty} \in \mathbb{M}(k, N, \epsilon)$ is also imposed on $\pi_d \mathfrak{f} = (\phi_m)_{m=0}^d \in \mathbb{M}(k, N, \epsilon)[d]$ for all $d \ge \max(m, n)$.

Since the sequence dim $\mathbb{M}(k, N, \epsilon)[d] \in \mathbb{N}_0$ is eventually monotonically decreasing, it is eventually constant, and the injective π_d are eventually isomorphisms. Thus dim $\mathbb{M}(k, N, \epsilon) = \lim_d \dim \mathbb{M}(k, N, \epsilon)[d]$. From Theorem 9.6 we have dim $M_k(K(N))^{\epsilon} = \dim \mathbb{M}(k, N, \epsilon)$ and this completes the proof.

10. Concluding Remarks.

The Main Theorem 9.6 gives an quick proof, without needing to check convergence, of the existence of the Gritsenko lift [18], Grit : $J_{k,N} \to M_k (K(N))^{\epsilon}$, for $\epsilon = (-1)^k$. For $\ell \in \mathbb{N}$, let $V_{\ell} : J_{k,m} \to J_{k,m\ell}$ be the index raising operator from [11]. For $\phi \in J_{k,N}$, the formal series $\mathfrak{f} = c(0,0;\phi)\frac{1}{2}\zeta(1-k)E_k + \sum_{m\in\mathbb{N}} (\phi|V_m) \xi^{Nm}$ is directly checked to be in $\mathbb{M}(k, N, \epsilon)$. Here, the first term with the elliptic Eisenstein series E_k is zero unless $k \geq 4$ is even. By the Main Theorem there is a $\operatorname{Grit}(\phi) \in M_k (K(N))^{\epsilon}$ with FJ ($\operatorname{Grit}(\phi)$) = \mathfrak{f} .

The Main Theorem 9.6 can be used to construct the global level raising operators in the paramodular newform theory of Roberts and Schmidt [35, 36]. For a prime ℓ , the three level raising operators $\theta_{\ell}, \theta'_{\ell} : S_k(K(N)) \to S_k(K(N\ell))$, and $\eta_{\ell} : S_k(K(N)) \to S_k(K(N\ell^2))$, are used to create the oldforms. We can directly define the global level raising operators by giving their action on Fourier-Jacobi expansions. For $\ell \in \mathbb{N}$, let $U_{\ell} : J_{k,m} \to J_{k,m\ell^2}$ be the index raising operator from [11]. The simplest to define is η_{ℓ} . For $f \in S_k(K(N))^{\epsilon}$ with $\mathrm{FJ}(f) = \sum_{m=1}^{\infty} \phi_m \xi^{Nm}$, define $A(f) = \sum_{m=1}^{\infty} (\phi_m | U_{\ell}) \xi^{N\ell^2 m} \in$ $\mathbb{S}(k, N\ell^2)$. One directly checks $A(f) \in \mathbb{S}(k, N\ell^2, \epsilon)$ so that by the main result there is an $\eta_{\ell} f \in S_k(K(N\ell^2))^{\epsilon}$ with $\mathrm{FJ}(\eta_{\ell} f) = A(f)$. This defines $\eta_{\ell} : S_k(K(N)) \to S_k(K(N\ell^2))$ since the entire space is the direct sum of the plus and minus forms.

In order to define θ_{ℓ} and θ'_{ℓ} , set $\Xi_{\ell}f = \sum_{m=1}^{\infty} (\phi_m | V_{\ell}) \xi^{N\ell m} \in \mathbb{S}(k, N\ell)$, and $\Xi'_{\ell}f = \sum_{m=1}^{\infty} \left(\sum_{\delta | \gcd(\ell,m)} \delta^{k-1} \phi_{\frac{\ell m}{\delta^2}} | U_{\delta} \right) \xi^{N\ell m} \in \mathbb{S}(k, N\ell)$. One directly checks $B^{\pm}(f) = \Xi_{\ell}(f) \pm \Xi'_{\ell}(f | \mu_N) \in \mathbb{S}(k, N\ell, \pm 1)$ so that by the main result there are $b^{\pm}(f) \in S_k (K(N\ell))^{\pm}$ with FJ $(b^{\pm}(f)) = B^{\pm}(f)$. Defining $\theta_{\ell}f = \frac{1}{2}(b^+(f) + b^-(f)) \in S_k (K(N\ell))$ and correspondingly $\theta'_{\ell}f = \frac{1}{2}(b^+(f | \mu_N) - b^-(f | \mu_N)) \in S_k (K(N\ell))$, we have $\theta_{\ell}, \theta'_{\ell} : S_k (K(N)) \to S_k (K(N\ell))$ with FJ $(\theta_{\ell}f) = \Xi_{\ell}f$, FJ $(\theta'_{\ell}f) = \Xi'_{\ell}f$, and $(\theta_{\ell} f)|\mu_{N\ell} = \theta'_{\ell}(f|\mu_N)$. Since these constructions work for $\ell \in \mathbb{N}$, this gives a generalization of the global level raising operators.

The theory of formal series for arithmetic subgroups developed in [5] frames different hypotheses than we do in the case of the paramodular groups. Still, in their Theorem 4.8, Bruinier and Raum considered the implications of their general theory for paramodular groups when a single formal series at the standard 1-cusp is given. In our notation, they proved the following. For squarefree N, let $\Gamma_0(N)^*$ be the extension of $\Gamma_0(N)$ by all the Atkin-Lehner involutions. Let $\chi : \Gamma_0(N)^* \to \{\pm 1\}$ be a character trivial on $\Gamma_0(N)$. If a formal series $\mathfrak{f} \in \mathbb{M}(k, N)$ satisfies $\mathfrak{j}(\sigma^*) \operatorname{AFS}(\mathfrak{f}) = \chi(\sigma) \operatorname{AFS}(\mathfrak{f})$ for all $\sigma \in \Gamma_0(N)^*$, then \mathfrak{f} converges on \mathcal{H}_2 . Using simple generators for $\Gamma_0(N)^*$, this reproves the cases N = 2, 3from [26]. In comparison, our main result allows general N and assumes a single symmetry under the paramodular Fricke involution.

Theorem 8.2 only needed to be proven for cuspidal quotients \mathfrak{f} because the Main Theorem 9.6 was reduced to the case of formal series of Jacobi cusp forms. With the Main Theorem in hand, however, we may improve Theorem 8.2 to obtain a general criterion for the divisibility of paramodular Fricke eigenforms that is interesting in its own right because it avoids all discussion of divisors.

Theorem 10.1. Let $k, k_0, N \in \mathbb{N}$ and $\epsilon, \epsilon_0 \in \{\pm 1\}$. Let $\mathfrak{f} \in \mathbb{M}(k, N)$, $f_0 \in M_{k_0}(K(N))^{\epsilon_0}$ be nontrivial, and $G \in M_{k_0+k}(K(N))^{\epsilon_0\epsilon}$. If we have $\mathrm{FJ}(G) = \mathrm{FJ}(f_0)\mathfrak{f}$ in $\mathbb{M}(k_0+k, N)$ then the meromorphic G/f_0 is holomorphic and $G/f_0 \in M_k(K(N))^{\epsilon}$.

Proof. We have $\mathfrak{f} \in \mathbb{M}(k, N, \epsilon)$ if and only if $\mathfrak{j}(F_N^*)AFS(\mathfrak{f}) = \epsilon AFS(\mathfrak{f})$. Applying AFS to $FJ(G) = FJ(f_0)\mathfrak{f}$ we obtain $FS(G) = FS(f_0)AFS(\mathfrak{f})$. Applying $\mathfrak{j}(F_N^*)$ we have $\epsilon_0\epsilon FS(G) = \epsilon_0FS(f_0)\mathfrak{j}(F_N^*)AFS(\mathfrak{f})$. Since $\mathbb{C}^{\overline{\mathcal{X}}(N)}$ is an integral domain and the element $FS(f_0)$ is nontrivial, we have $\mathfrak{j}(F_N^*)AFS(\mathfrak{f}) = \epsilon AFS(\mathfrak{f})$, and thus $\mathfrak{f} \in \mathbb{M}(k, N, \epsilon)$. By the Main Theorem 9.6 there is an $f \in M_k(K(N))^{\epsilon}$ such that $FJ(f) = \mathfrak{f}$. Thus the quotient $G/f_0 = f \in M_k(K(N))^{\epsilon}$ is holomorphic. \Box

References

- Hiroki Aoki. Estimating Siegel modular forms of genus 2 using Jacobi forms. J. Math. Kyoto Univ., 40(3):581–588, 2000.
- [2] Hiroki Aoki. On the upper bound of the orders of Jacobi forms. Internat. J. Math., 33(8):Paper No. 2250056, 14, 2022.
- [3] Eran Assaf, Watson Ladd, Gustavo Rama, Gonzalo Tornaría, and John Voight. A database of paramodular forms from quinary orthogonal modular forms. In LuCaNT: LMFDB, computation, and number theory, volume 796 of Contemp. Math., pages 243–259. Amer. Math. Soc., [Providence], RI, [2024] ©2024.

- [4] Jeffery Breeding, II, Cris Poor, and David S. Yuen. Computations of spaces of paramodular forms of general level. J. Korean Math. Soc., 53(3):645–689, 2016.
- [5] Jan Bruinier and Martin Raum. Formal Siegel modular forms for arithmetic subgroups. Trans. Amer. Math. Soc. Ser. B, 11:1394–1434, 2024.
- [6] Jan Hendrik Bruinier. Vector valued formal Fourier-Jacobi series. Proc. Amer. Math. Soc., 143(2):505–512, 2015.
- [7] Jan Hendrik Bruinier and Martin Westerholt-Raum. Kudla's modularity conjecture and formal Fourier-Jacobi series. Forum Math. Pi, 3:7, 30, 2015.
- [8] Armand Brumer and Kenneth Kramer. Paramodular abelian varieties of odd conductor. Trans. Amer. Math. Soc., 366(5):2463-2516, 2014.
- [9] Armand Brumer and Kenneth Kramer. Paramodular abelian varieties of odd conductor. arXiv:1004.4699, 2018.
- [10] N. Dummigan, A. Pacetti, G. Rama, and G. Tornaría. Quinary forms and paramodular forms. *Math. Comp.*, 93(348):1805–1858, 2024.
- [11] Martin Eichler and Don Zagier. The theory of Jacobi forms, volume 55 of Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, 1985.
- [12] E. Freitag. Siegelsche Modulfunktionen, volume 254 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1983.
- [13] Eberhard Freitag and Rolf Busam. Complex analysis. Universitext. Springer-Verlag, Berlin, second edition, 2009.
- [14] Klaus Fritzsche and Hans Grauert. From holomorphic functions to complex manifolds, volume 213 of Graduate Texts in Mathematics. Springer-Verlag, New York, 2002.
- [15] Vasily Golyshev and Duco van Straten. Congruences via fibered motives. Pure Appl. Math. Q., 19(1):233–265, 2023.
- [16] V. Gritsenko and K. Hulek. Commutator coverings of Siegel threefolds. Duke Math. J., 94(3):509–542, 1998.
- [17] Valeri Gritsenko. Arithmetical lifting and its applications. In Number theory (Paris, 1992–1993), volume 215 of London Math. Soc. Lecture Note Ser., pages 103–126. Cambridge Univ. Press, Cambridge, 1995.
- [18] Valeri Gritsenko. Irrationality of the moduli spaces of polarized abelian surfaces. In Abelian varieties (Egloffstein, 1993), pages 63–84. de Gruyter, Berlin, 1995. With an appendix by the author and K. Hulek.
- [19] Valeri A. Gritsenko and Viacheslav V. Nikulin. Automorphic forms and Lorentzian Kac-Moody algebras. II. Internat. J. Math., 9(2):201–275, 1998.
- [20] Valery Gritsenko and Haowu Wang. Powers of Jacobi triple product, Cohen's numbers and the Ramanujan Δ-function. Eur. J. Math., 4(2):561–584, 2018.
- [21] Robert C. Gunning. Introduction to holomorphic functions of several variables, Vol. II: Local theory. The Wadsworth & Brooks/Cole Mathematics Series. Wadsworth & Brooks/Cole Advanced Books & Software, Monterey, CA, 1990.
- [22] Tomoyoshi Ibukiyama. On relations of dimensions of automorphic forms of Sp(2, **R**) and its compact twist Sp(2). I. In Automorphic forms and number theory (Sendai, 1983), volume 7 of Adv. Stud. Pure Math., pages 7–30. North-Holland, Amsterdam, 1985.

- [23] Tomoyoshi Ibukiyama. Paramodular forms and compact twist. In Automorphic forms on GSp(4): Proceedings of the 9th Autumn Workshop on Number Theory, pages 37–48, 2007.
- [24] Tomoyoshi Ibukiyama. Dimensions of paramodular forms and compact twist modular forms with involutions. arXiv:2208.13578v2, 2024.
- [25] Tomoyoshi Ibukiyama and Hidetaka Kitayama. Dimension formulas of paramodular forms of squarefree level and comparison with inner twist. J. Math. Soc. Japan, 69(2):597–671, 2017.
- [26] Tomoyoshi Ibukiyama, Cris Poor, and David S. Yuen. Jacobi forms that characterize paramodular forms. Abh. Math. Semin. Univ. Hambg., 83(1):111–128, 2013.
- [27] Stephen S. Kudla. Algebraic cycles on Shimura varieties of orthogonal type. Duke Math. J., 86(1):39–78, 1997.
- [28] Stephen S. Kudla. Integrals of Borcherds forms. Compositio Math., 137(3):293– 349, 2003.
- [29] Stephen S. Kudla. Special cycles and derivatives of Eisenstein series. In *Heegner points and Rankin L-series*, volume 49 of *Math. Sci. Res. Inst. Publ.*, pages 243–270. Cambridge Univ. Press, Cambridge, 2004.
- [30] Joseph Lehner. A short course in automorphic functions. Holt, Rinehart and Winston, New York-Toronto-London, 1966.
- [31] Junjiro Noguchi. Analytic function theory of several variables. Springer, Singapore, 2016. Elements of Oka's coherence.
- [32] Aaron Pollack. Automatic convergence for Siegel modular forms. arXiv:2408.16392v1, 2024.
- [33] Cris Poor, Jerry Shurman, and David S. Yuen. Siegel paramodular forms of weight 2 and squarefree level. Int. J. Number Theory, 13(10):2627–2652, 2017.
- [34] Helmut Reefschläger. Berechnung der Anzahl der 1-Spitzen der Paramodularen Gruppen 2-ten Grades. PhD thesis, Georg-August-Universität zu Göttingen, 1973.
- [35] Brooks Roberts and Ralf Schmidt. On modular forms for the paramodular groups. In Automorphic forms and zeta functions, pages 334–364. World Sci. Publ., Hackensack, NJ, 2006.
- [36] Brooks Roberts and Ralf Schmidt. Local Newforms for GSp(4), volume 1918 of Lecture Notes in Mathematics. Springer, Berlin, 2007.
- [37] Mirko Rösner and Rainer Weissauer. Global liftings between inner forms of GSp(4). J. Number Theory, 263:79–138, 2024.
- [38] Nils-Peter Skoruppa. Uber den Zusammenhang zwischen Jacobiformen und Modulformen halbganzen Gewichts, volume 159 of Bonner Mathematische Schriften [Bonn Mathematical Publications]. Universität Bonn, Mathematisches Institut, Bonn, 1985. Dissertation, Rheinische Friedrich-Wilhelms-Universität, Bonn, 1984.
- [39] Pol van Hoften. A geometric Jacquet-Langlands correspondence for paramodular Siegel threefolds. *Math. Z.*, 299(3-4):2029–2061, 2021.
- [40] Martin Westerholt-Raum. Formal Fourier Jacobi expansions and special cycles of codimension two. Compos. Math., 151(12):2187–2211, 2015.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND TECHNOLOGY, TOKYO UNIVERSITY OF SCIENCE, NODA, CHIBA, 278-8510 JAPAN *Email address*: aoki_hiroki_math@nifty.com

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF MATHEMATICS, OS-AKA UNIVERSITY, MACHIKANEYAMA 1-1, TOYONAKA, OSAKA, 560-0043 JAPAN *Email address*: ibukiyam@math.sci.osaka-u.ac.jp

Department of Mathematics, Fordham University, Bronx, NY 10458 $\rm USA$

Email address: poor@fordham.edu