ARC-TRANSITIVE MAPS WITH EDGE NUMBER COPRIME TO THE EULER CHARACTERISTIC – I

CAI HENG LI AND LUYI LIU

ABSTRACT. This is one of a series of papers which aim towards a classification of edge-transitive maps of which the Euler characteristic and the edge number are coprime. This one establishes a framework and carries out the classification work for arc-transitive maps with solvable automorphism groups, which illustrates how the edge number impacts on the Euler characteristic for maps. The classification is involved with the constructions of various new and interesting arc-regular maps.

Key words: arc-regular, maps, Euler characteristic

1. INTRODUCTION

A map is a 2-cell embedding of a graph into a closed surface. Throughout the paper, we denote by $\mathcal{M} = (V, E, F)$ a map with vertex set V, edge set E, and face set F. The underlying graph (V, E) of \mathcal{M} is denoted by Γ , and the supporting surface of \mathcal{M} is denoted by \mathcal{S} . The Euler characteristic of a map $\mathcal{M} = (V, E, F)$ is

$$\chi(\mathcal{M}) = |V| - |E| + |F|.$$

Clearly, the number of edges |E| directly affects $\chi(\mathcal{M})$. In this paper, we investigate highly symmetric maps where $gcd(|E|, \chi(\mathcal{M})) = 1$. Moreover, we always assume that \mathcal{M} has no loops.

Let $\mathcal{M} = (V, E, F)$ be a map. An *arc* of \mathcal{M} is an ordered adjacent pair (α, e) of a vertex v and an edge e, while a *flag* of \mathcal{M} is an adjacent triple (α, e, f) of a vertex α , an edge e, and a face f. Let e be an edge with endpoints α and β . By the definition of a map, there are exactly two faces f and f' adjacent to e. The edge e is adjacent to two arcs (α, e) and (β, e) , and four flags: $(\alpha, e, f), (\alpha, e, f'),$ $(\beta, e, f), \text{ and } (\beta, e, f')$. The arc set and flag set of \mathcal{M} are denoted by \mathcal{A} and \mathcal{F} , respectively, and satisfy $|\mathcal{F}| = 2|\mathcal{A}| = 4|E|$. We denote by E(v) and E(f) the sets of edges incident to vertex α and face f, respectively. The vertex valency of α is defined as $|E(\alpha)|$, and the face length of f is defined as |E(f)|.

An automorphism of \mathcal{M} is a permutation σ of its flags that preserves all incidence relations. The set of all automorphisms of \mathcal{M} forms a group, denoted by Aut(\mathcal{M}). If an automorphism $\sigma \in \text{Aut}(\mathcal{M})$ fixes a flag, then it must fix all flags, and therefore σ is the identity element. This implies that Aut(\mathcal{M}) acts semiregularly on the flag set \mathcal{F} of \mathcal{M} . If Aut(\mathcal{M}) acts transitively on \mathcal{F} , then \mathcal{M} is called a *regular map*. If Aut(\mathcal{M}) acts transitively on the arc set \mathcal{A} but intransitively on \mathcal{F} , then it acts regularly on \mathcal{A} , and \mathcal{M} is termed an *arc-regular map*. Similarly, \mathcal{M} is called *edgeregular* if Aut(\mathcal{M}) acts regularly on the edge set E. The classification problem of certain arc-transitive maps has been extensively studied; see [2, 5, 6, 16, 17, 18] and references therein.

This work was partially supported by NSFC grant 11931005.

A map is called *locally finite* if all vertex valencies and face lengths are finite. Locally finite edge-transitive maps and their automorphism groups are classified by Graver and Watkins [9] into fourteen types based on the local actions of their automorphism groups. Among these types, five are arc-transitive which are explain as follows.

Let $G \leq \operatorname{Aut}(\mathcal{M})$ act arc-transitively on \mathcal{M} . Then G is transitive on the vertex set V, and the valency $\ell = |E(\alpha)|$ is constant for all $\alpha \in V$, called the *valency* of \mathcal{M} . Furthermore, there exists an involution $z \in G$ that swaps the two arcs (α, e) and (β, e) adjacent to an edge $e \in E$.

First, suppose that G acts transitively on the flag set of \mathcal{M} , then $G = \operatorname{Aut}(\mathcal{M})$, and \mathcal{M} is a regular map. Furthermore, there exist involutions $x, y \in G$ such that

$$G = \langle x, y, z \rangle, \quad G_v = \langle x, y \rangle = \mathcal{D}_{2\ell}, \quad G_e = \langle x, z \rangle = \mathcal{D}_4.$$

The triple (x, y, z) is called a *regular triple* of G.

Now assume that G is not transitive on the flag set. Then G is regular on the arc set of \mathcal{M} , and \mathcal{M} is a G-arc-regular map. In this case, G_v is cyclic or dihedral of order equal to ℓ . If $G_{\alpha} = \langle a \rangle = \mathbb{Z}_{\ell}$, then \mathcal{M} is said to be G-vertex-rotary, and (a, z)is called a rotary pair for G. On the other hand, if $G_{\alpha} = \langle x, y \rangle = D_{\ell}$, then \mathcal{M} is said to be G-vertex-reversing, and (x, y, z) is called a reversing triple for G. Further, according to the action of z on $\{f, f'\}$, vertex-rotary maps are divided into rotary maps and bi-rotary maps, and vertex-reversing maps are divided into reversing maps and bi-reversing maps, as shown in the table below.

TABLE 1. Five types of arc-transitive maps

Type	G_{α}	$(f,f')^z$	Name
1	$\langle x, y \rangle = \mathcal{D}_{2k}$	(f, f') or (f', f)	regular
2^{*}	$\langle x, y \rangle = \mathcal{D}_k$	(f,f')	reversing
2^{P}	$\langle x, y \rangle = \mathbf{D}_k$	(f',f)	bi-reversing
$2^{*} ex$	$\langle a \rangle = \mathbb{Z}_k$	(f', f)	rotary (orientably regular)
$2^{\mathrm{P}}\mathrm{ex}$	$\langle a \rangle = \mathbb{Z}_k$	(f, f')	bi-rotary

We classify arc-transitive maps $\mathcal{M} = (V, E, F)$ with $gcd(|E|, \chi(\mathcal{M})) = 1$ in two papers. This one presents a classification with $Aut(\mathcal{M})$ solvable, and the second one [14] completes the work for the case with $Aut(\mathcal{M})$ nonsolvable.

Noticing that arc-transitive embeddings of multicycles are determined in [10], and arc-transitive embeddings of complete multigraphs are classified in [17]. For the orientably-regular embeddings, see [11]. The main result of this paper is stated as follows.

Theorem 1.1. Let $\mathcal{M} = (V, E, F)$ be a map with $gcd(\chi(\mathcal{M}), |E|) = 1$. Assume that $G \leq Aut(\mathcal{M})$ is indecomposable, solvable and arc-transitive on \mathcal{M} , and the underlying graph of \mathcal{M} is neither $K_4^{(\lambda)}$ nor $C_n^{(\lambda)}$ with $n \geq 2$. Then one of the following holds:

(1) $G = \langle g \rangle : \langle h \rangle$, where $C_{\langle h \rangle}(g) \leq \Phi(\langle h \rangle)$, |h| = 2m and $g^{h^m} = g^{-1}$; the map \mathcal{M} is G-vertex-rotary with a rotary pair (a, z) is equivalent to (gh^i, h^m) or (gh^{2i}, h^m) ,

where gcd(i, |h|) = 1; the number of inequivalent rotary pairs for G is $\frac{2\varphi(|h|)}{|\tau_{h_0}|}$ if $G_2 \cong D_{2^e}$ with $e \ge 3$; and $\varphi(|h|)/|\tau_{h_0}|$ otherwise, where $\langle h_0 \rangle = C_{\langle h \rangle}(g)$.

In the following, Sylow 2-subgroups of G are dihedral. Let u, v be two involutions such that $\langle u, v \rangle = G_2$ and w = uv.

(2) $G = (\langle g \rangle \times \langle g_v \rangle):(\langle u, v \rangle \times \langle h \rangle)$, where $|g|, |g_v|$ and |h| are pairwise coprime, $C_{\langle h \rangle}(gg_v) \leq \Phi(\langle h \rangle)$, and $\langle u, v \rangle = D_{2^e}$ with $e \geq 2$ such that

 $(g, g_v)^v = (g, g_v^{-1}) \text{ and } (g, g_v)^u = (g, g_v),$

where $g_v \neq 1$; and either

(i) \mathcal{M} is G-vertex-reversing, $G = \langle g_v \rangle : \langle u, v \rangle$, relabeling if necessary, a reversing triple (x, y, z) is equivalent to

 $(u, g_v w^{2j} v, v)$, or $(v, g_v w, u)$ when |w| = 2,

and there are $6 \cdot 2^{e-2}$ inequivalent reversing triples of G; or

(ii) \mathcal{M} is G-vertex-rotary, and a rotary pair (a, z) is equivalent to

 (gg_vwh^j, v) or (gg_vvh^j, v) ,

where gcd(j, |h|) = 1, and there are $2\varphi(|h|)/|\tau_{h_0}|$ inequivalent rotary pairs of G.

(3) $G = (\langle g_u \rangle \times \langle g_v \rangle \times \langle g \rangle):(\langle u, v \rangle \times \langle h \rangle), \text{ where } |g_u|, |g_v|, |g| \text{ and } |h| \text{ are pairwise coprime, } C_{\langle h \rangle}(g_u g_v g) \leqslant \Phi(\langle h \rangle), \text{ and } \langle u, v \rangle = D_{2^e} \text{ with } e \geqslant 2 \text{ such that}$

 $(g_u, g_v, g)^v = (g_u, g_v^{-1}, g^{-1})$ and $(g_u, g_v, g)^u = (g_u^{-1}, g_v, g^{-1}),$

where if $g_u = 1$, then $g_v \neq 1$ and $g \neq 1$; suppose that $g_v = 1$, then either

(1) \mathcal{M} is G-vertex-reversing, h = 1, and relabeling if necessary, a reversing triple (x, y, z) is equivalent to

 $(g_u g^{k_1} w^{2i} u, g^{k_2} v, u),$

where $gcd(k_1, k_2) = 1$, ℓ_1 is odd, and there are

$$6 \cdot 2^{e-3} \cdot \frac{|\{(k_1, k_2) \mid \gcd(k_1, k_2) = 1\}|}{\varphi(|g|)}$$

inequivalent reversing triples of G;

(2) \mathcal{M} is G-vertex-rotary, and a rotary pair (a, z) is equivalent to

 $(g_u g w h^{\ell}, u)$ or $(g_u g w^2 v h^{\ell}, u)$,

and there are $2\varphi(|h|)/|\tau_{h_0}|$ inequivalent rotary pairs in G.

(4) $G = (\langle g \rangle \times \langle w^2, v \rangle):((\langle h \rangle: \langle u \rangle) \times \langle c \rangle), \text{ where } \langle u, v \rangle = D_8 \text{ such that } h^u = h^{-1} \text{ and } g^u = g^{-1}, \langle h \rangle \text{ is a Sylow 3-subgroup such that } (w^2v, v, w^2)^h = (v, w^2, w^2v) \text{ and } [g, h] = 1 \text{ and } \langle g \rangle \text{ and } \langle c \rangle \text{ are Hall subgroups such that } C_{\langle c \rangle}(g) \leqslant \Phi(\langle c \rangle); \text{ and } either$

(i) \mathcal{M} is G-vertex-rotary with a rotary pair (a, z) being equivalent to

$$(gth^{j}c^{k}, u) \text{ or } (gth^{j}uc^{k}, u),$$

where $gcd(j, 3) = gcd(k, |c|) = 1$, and $t \in \{v, w^{2}\}$, and there are
 $\varphi(|h|)\varphi(|c|)/2|\tau_{h^{3}}||\tau_{c_{0}}|$

inequivalent rotary pairs of G.

(ii) *M* is G-vertex-reversing, c = 1; let tt' ∈ ⟨w², v⟩, relabeling if necessary, a reversing triple (x, y, z) is equivalent to one of the following triples:
(i) (t(gh)ⁱu, t', u) with t' ≠ 1,

(*ii*)
$$(t(gh)^i u, g^{i'}h^{3j_0}w^2 u, u),$$

(*iii*) $(t(gh)^{i}u, g^{i'}t'h^{j'}u, u)$ with $(t, t') \neq (1, 1)$,

where gcd(i, |gh|) = 1.

2. Preliminaries

In this section, we collect some basic definitions and preliminary results which will be used in the ensuing sections.

Let $\mathcal{M} = (V, E, F)$ be a map, and $G \leq \operatorname{Aut}(\mathcal{M})$ be arc-regular on \mathcal{M} . Then the stabilizer G_e of an edge e is generated by an involution z say, explicitly,

$$G_e = \langle z \rangle \cong \mathbb{Z}_2$$

and $(\alpha, e)^z = (\beta, e)$. Let $f, f' \in F$ be incident with e. Then z either fixes f and f', or interchanges f and f', so that

$$z: (f, f') \mapsto (f, f') \text{ or } (f', f).$$

Since G acts regularly on the arc set, the vertex stabilizer G_{α} is regular on $E(\alpha)$. The pair (G_{α}, z) defines an arc-regular coset graph $\Gamma = (V, E)$, where

$$V = [G:G_{\alpha}] = \{G_{\alpha}g \mid g \in G\}, \quad E = [G:\langle z \rangle] = \{\langle z \rangle h \mid h \in G\}$$

with incidence defined by non-empty intersection: a vertex $G_{\alpha}g$ and an edge $\langle z \rangle h$ are incident if and only if $G_{\alpha}g \cap \langle z \rangle h \neq \emptyset$. This coset graph, denoted by $\mathsf{Cos}(G, G_{\alpha}, \langle z \rangle)$, may have multiple edges; see [16] for details.

Assume that $G_{\alpha} = \langle a \rangle$. Then the rotary pair (a, z) determines two vertex-rotary embeddings of Γ , namely, a rotary map or a bi-rotary map, defined below.

Definition 2.1. Let G be a group generated by a, z with z being an involution, and let $\Gamma = \text{Cos}(G, \langle a \rangle, \langle z \rangle)$. Let $V = [G : G_{\alpha}], E = [G : \langle z \rangle]$, and let

- (i) RotaMap(G, a, z) be the embedding of Γ with face set $[G : \langle az \rangle]$, and
- (ii) BiRotaMap(G, a, z) be the embedding of Γ with face set $[G : \langle z, z^a \rangle]$,

with incidences defined by non-empty intersection. Then $\mathsf{RotaMap}(G, a, z)$ is a rotary map, and $\mathsf{BiRotaMap}(G, a, z)$ is a bi-rotary map.

Now let $G_{\alpha} = \langle x, y \rangle$ with x and y being involutions. Then the reversing triple (x, y, z) determines two vertex-reversing embeddings of Γ , namely, a reversing map or a bi-reversing map, defined below.

Definition 2.2. Let G be a group generated by involutions x, y, z, and let $\Gamma = Cos(G, \langle x, y \rangle, \langle z \rangle)$. Let

- (i) RevMap(G, x, y, z) be the embedding of Γ with $F = [G:\langle x, z \rangle] \cup [G:\langle y, z \rangle]$,
- (ii) BiRevMap(G, x, y, z) be the embedding of Γ with $F = [G:\langle x, y^z \rangle]$,

with incidence defined by non-empty intersection. Then $\operatorname{RevMap}(G, x, y, z)$ is a reversing map, and $\operatorname{BiRevMap}(G, x, y, z)$ is a bi-reversing map.

To end of this section, we provide some definitions for symbols used in this paper. For an integer n, let n_p be the p-part, which means that $n = n_p m$ such that n_p is a p-power and gcd(p, m) = 1. Let $\pi(n)$ be the set of prime divisors of the integer n.

For a group G, let $\pi(G) = \pi(|G|)$, the set of prime divisors of the order |G|. For a subset π of $\pi(G)$, denote by G_{π} a Hall π -subgroup, and $G_{\pi'}$ a Hall π' -subgroup of G, where $\pi' = \pi(G) \setminus \pi$. If $\pi = \{p\}$, then $G_{\pi} = G_p$ is a Sylow p-subgroup of G. For a subgroup $H \leq G$, let $N_G(H)$ and $C_G(H)$ be the normaliser and the centralizer of H in G, respectively. Let $\Phi(G)$ be the Frattini subgroup of G, which consists of all non-generators of G. By \mathbb{Z}_n , we mean a cyclic group of order n, and D_{2n} is a dihedral group of order 2n.

3. The edge number and the Euler characteristic

In this section, we investigate the effect of the Euler characteristic on edge numbers and automorphism groups of maps. Also, we provided a characterization of automorphism groups of maps such that the Euler characteristic and the edge number are coprime.

Lemma 3.1. Assume that \mathcal{M} has constant vertex valency ℓ and constant face length k, and $gcd(\chi(\mathcal{M}), |E|) = 1$. Then |E| divides $k\ell$, |V| divides 2k and |F| divides 2ℓ .

Proof. Since each edge is adjacent to two vertices and two faces, there are

$$|V| \cdot \ell = 2|E|$$
 and $|F| \cdot k = 2|E|$.

Combining these with the Euler characteristic formula $\chi(\mathcal{M}) = |V| - |E| + |F|$, we substitute

$$|V| = \frac{2|E|}{\ell}, \quad |F| = \frac{2|E|}{k}$$

to obtain

$$k\ell\chi(\mathcal{M}) = (2\ell - k\ell + 2k)|E|.$$

It follows from $gcd(\chi(\mathcal{M}), |E|) = 1$ that |E| divides $k\ell$. Consequently, $|V| = \frac{2|E|}{\ell}$ divides $\frac{2k\ell}{\ell} = 2k$, and $|F| = \frac{2|E|}{k}$ divides $\frac{2k\ell}{k} = 2\ell$.

This lemma shows that, if $gcd(\chi, |E|) = 1$ then the edge number |E| is small relative to the valency and the face length.

The next lemma characterizes Sylow subgroups of the automorphism group of a map whose Euler characteristic is coprime to the number of edges.

Lemma 3.2. Let $\mathcal{M} = (V, E, F)$ be a map, and G be a subgroup of Aut (\mathcal{M}) . Assume that $gcd(\chi(\mathcal{M}), |E|) = 1$. Then the following statements are true:

- (1) $gcd(\chi(\mathcal{M}), |G|)$ divides 4, and $gcd(\chi(\mathcal{M}), |G|) \neq 1$ only if the edge number |E| is odd;
- (2) each Sylow subgroup of G is a cyclic or dihedral subgroup;
- (3) $|G| = \operatorname{lcm}\{|G_{\omega}| : \omega \in V \cup E \cup F\}.$

Proof. Recall that Aut(\mathcal{M}) acts semiregular on the flag set \mathcal{F} . So |G| divides $|\mathcal{F}|$. Let $\chi = \chi(\mathcal{M})$ for convenience. Since $|\mathcal{F}| = 4|E|$ and $gcd(\chi, |E|) = 1$, $gcd(\chi, |G|)$ divides $gcd(\chi, |\mathcal{F}|) = gcd(\chi, 4|E|) = gcd(\chi, 4)$. Suppose that $gcd(\chi, |G|) \neq 1$. Then χ is even, and so the edge number |E| is odd, as in part (1).

Let p be a prime divisor of |G|. First, suppose that $p \mid \chi$. It follows from statement (1) that p = 2 and |E| is odd. Since |G| divides 4|E|, the Sylow 2-subgroups of G have order dividing 4. Explicitly, Sylow 2-subgroups are cyclic or dihedral.

Now, suppose that $p \nmid \chi$. We claim that exists $\omega \in V \cup E \cup F$ such that $p \nmid \frac{|G|}{|G_{\omega}|}$. Otherwise, p divides the length of each G-orbit on $V \cup E \cup F$, and so p divides $|V| - |E| + |F| = \chi$ which is the sum all G-orbits length. Contradiction comes to $p \nmid \gcd(\chi, |G|)$, and so the claim is proved. Therefore Sylow p-subgroups of G are isomorphic to a subgroup of G_{ω} by Sylow's theorem. So they are cyclic or dihedral as G_{ω} is cyclic or dihedral for each $\omega \in V \cup E \cup F$, as in statement (2).

Let $\ell = \operatorname{lcm}(|G_{\omega}| \mid \omega \in V \cup E \cup F)$ and $d = \operatorname{gcd}(|G|/|G_{\omega}| \mid \omega \in V \cup E \cup F)$. Then $|G| = \ell d$, and the greatest common divisor d divides $\operatorname{gcd}(\chi, |G|)$.

Assume that $d \neq 1$. By statement (1), $gcd(\chi, |G|)$ is 2 or 4, and |G| = 2|E| or 4|E| where the edge number |E| is odd. There follows two cases as below.

- (a) If |G| = 4|E|, then \mathcal{M} is a regular map which implies that edge stabilizers in G are isomorphic to D₄. Hence 4 divides ℓ , and so d = 1 which is a contradiction.
- (b) If |G| = 2|E|, then d = 2 and $\ell = |E|$ is odd. Therefore $|G_{\omega}|$ is odd for each $\omega \in V \cup E \cup F$. A contradiction comes to that |E| is the sum of all lengths of G-orbits on E, which are all even.

Above all, d = 1 and so $G = \ell$ as in statement (3).

Regular maps on orientable surface of genus p + 1, where p is prime, have been classified[4]. In particular, automorphism groups of regular maps \mathcal{M} with order coprime to $\chi/2$ are determined. Here by part (1) of Lemma 3.2, if a *G*-arc-transitive map \mathcal{M} satisfies $gcd(\chi, |E|) = 1$ and $gcd(\chi/2, |G|) = 1$, the edge number |E| of \mathcal{M} is odd. To avoid overlaps, we will not discuss on the above case.

4. Almost Sylow-cyclic groups

A group is called an *almost Sylow-cyclic group* if its odd order Sylow subgroups are cyclic and each Sylow 2-subgroup contains an index 2 cyclic subgroup, refer to [3]. Non-solvable almost Sylow-cyclic groups were completely classified by Suzuki [19] and Wong [20], and solvable almost Sylow-cyclic groups were characterized by Zassenhaus [21].

The authors of this paper presents an explicitly classification of solvable almost Sylow-cyclic groups, extending Theorem 4.1 of [3].

Theorem 4.1 ([15]). Let G be an indecomposable solvable finite group of which each Sylow subgroup is cyclic or dihedral. Then G is one of the following five types:

(I) $G = (\langle g \rangle : \langle h \rangle), \text{ where } C_{\langle h \rangle}(g) \leq \Phi(\langle h \rangle).$

 $\mathbf{6}$

In the following, Sylow 2-subgroups of G are dihedral. Let u, v be two involutions such that $\langle u, v \rangle = G_2$ and w = uv.

(II) $G = (\langle g \rangle \times \langle g_v \rangle):(\langle u, v \rangle \times \langle h \rangle)$, where $|g|, |g_v|$ and |h| are pairwise coprime, $C_{\langle h \rangle}(gg_v) \leq \Phi(\langle h \rangle)$, and $\langle u, v \rangle = D_{2^e}$ with $e \geq 2$ such that

$$(g, g_v)^v = (g, g_v^{-1}) and (g, g_v)^u = (g, g_v),$$

where $g_v \neq 1$;

(III) $G = (\langle g_u \rangle \times \langle g_v \rangle \times \langle g \rangle):(\langle u, v \rangle \times \langle h \rangle), \text{ where } |g_u|, |g_v|, |g| \text{ and } |h| \text{ are pairwise coprime, } C_{\langle h \rangle}(g_u g_v g) \leqslant \Phi(\langle h \rangle), \text{ and } \langle u, v \rangle = D_{2^e} \text{ with } e \ge 2 \text{ such that}$

$$(g_u, g_v, g)^v = (g_u, g_v^{-1}, g^{-1}) \text{ and } (g_u, g_v, g)^u = (g_u^{-1}, g_v, g^{-1}),$$

where if $g_u = 1$, then $g_v \neq 1$ and $g \neq 1$;

- (IV) $G = (\langle g \rangle \times \langle u, v \rangle) : \langle h \rangle$, where |g|, |h| are coprime, |h| is divisible by 3, $C_{\langle h^3 \rangle}(g) \leq \Phi(\langle h^3 \rangle)$, and $\langle u, v \rangle \cong D_4$ such that $(u, v, w)^h = (v, w, u)$;
- (V) $G = (\langle g \rangle \times \langle w^2 v, v \rangle):((\langle h \rangle: \langle u \rangle) \times \langle c \rangle), \text{ where } \langle h \rangle \text{ is a Sylow 3-subgroup, } |g|, |h|, |c| \text{ are pairwise coprime, } C_{\langle c \rangle}(g) \leq \Phi(\langle c \rangle), \text{ and } \langle u, v \rangle = D_8 \text{ such that}$

$$(w^2v, v, w)^h = (v, w, w^2v)$$
 and $h^u = h^{-1}$.

Automorphism groups of cyclic groups and dihedral groups are well known. For convenience, we provide the definitions below.

Definition 4.2. For a cyclic group $L = \langle l \rangle$ and an element $l_0 \in \Phi(L)$, define an automorphism of L:

$$\tau_{l_0}: l \to l^{m \frac{|l|}{|l_0|}+1}$$

where *m* is an integer such that $gcd(m|l_0| + 1, |l|) = 1$. Then $\langle \tau_{l_0} \rangle = Aut(L)_{(L/\langle l_0 \rangle)}$.

Definition 4.3. Let $Q = \langle w \rangle : \langle u \rangle = D_{2m}$ with $m \ge 3$. Define automorphisms of Q(1) $\xi : (w, w^i u) \mapsto (w, w^{i+1}u), i = 1, ..., m$, and

(2) $\mu : wu^j \mapsto w^i u^j$, where *i* is a primitive divisor of m-1 and j=1,2. Then $\operatorname{Aut}(Q) = \langle \xi \rangle : \langle \mu \rangle \cong \operatorname{Hol}(\mathbb{Z}_m)$.

Automorphism groups of such group are determined, as quoted below.

Theorem 4.4. Let G be an indecomposable solvable finite group of which each Sylow subgroup is cyclic or dihedral. Then G = H:L where H, L are Hall subgroups, and

$$\operatorname{Aut}(G) = H: \mathcal{H} \times \mathcal{L},$$

where $\operatorname{Aut}(H) \cong \mathcal{H} \leq \operatorname{Aut}(G)_{(L)}$ and $\operatorname{Aut}(L)_{(L/C)} \cong \mathcal{L} \leq \operatorname{Aut}(G)_L \cap \operatorname{Aut}(G)_{(H)}$. Following notations in Theorem 4.1, one of the following statements holds:

- (I) $H = \langle g_{2'} \rangle$, $L = \langle g_2 \rangle : \langle h \rangle$, and $\mathcal{L} \cong \operatorname{Hol}(\langle g_2 \rangle) \times \langle \tau_{h_0} \rangle$;
- (II) $H = \langle gg_v \rangle, L = \langle u, v \rangle \times \langle h \rangle, and \mathcal{L} \cong \langle \xi^2 \rangle : \langle \mu \rangle \times \langle \tau_{h_0} \rangle;$
- (III) $H = \langle g_u g_v g \rangle, \ L = \langle u, v \rangle \times \langle h \rangle, \ and \ \mathcal{L} \cong \langle \xi^2 \rangle : \langle \mu \rangle \times \langle \tau_{h_0} \rangle;$

(IV)
$$H = \langle g \rangle$$
, $L = \langle u, v \rangle$: $\langle h \rangle$, and \mathcal{L} is isomorphic to
 $(\langle \tilde{u}, \tilde{v} \rangle \times \langle \tau_{h^3} \rangle)$: $(\langle \tilde{h} \rangle$: $\langle \mu_r \rangle)$, if $[h, g] = 1$;
 $\langle \tilde{u}, \tilde{v} \rangle$: $\langle \tilde{h} \rangle \times \langle \tau_{h^3} \rangle$, if $[h^3, g] = 1$.
(V) $H = \langle g \rangle$, $L = \langle w^2, v \rangle$: $(\langle h \rangle$: $\langle u \rangle) \times \langle c \rangle$, and \mathcal{L} is isomorphic to
 $(\langle \tilde{u}, \tilde{v} \rangle \times \langle \tau_{h^3} \rangle)$: $(\langle \tilde{h} \rangle$: $\langle \mu_r \rangle) \times \langle \tau_{c_0} \rangle$, if $[h, g] = 1$;
 $\langle \tilde{u}, \tilde{v} \rangle$: $\langle \tilde{h} \rangle \times \langle \tau_{h^3} \rangle \times \langle \tau_{c_0} \rangle$, if $[h^3, g] = 1$,
where $\langle c_0 \rangle = C_{\langle c \rangle}(g)$.

5. Type I: Metacyclic groups

In this section, we always assume that G is metacyclic, that is, G is a group of type I.

Lemma 5.1. Assume that $G = \langle g \rangle : \langle h \rangle = D_{2m}$ is a dihedral group, and \mathcal{M} is a *G*-arc-regular map with underlying graph Γ .

Then either

- (1) \mathcal{M} is G-vertex-rotary with rotary pair (a, z), where the pair is equivalent to $(g, g^{j}h)$ or (h, gh), and Γ is $K_{2}^{(m)}$ or C_{m} , respectively; or
- (2) \mathcal{M} is G-vertex-reversing with reversing triple (x, y, z), where the triple is equivalent to $\{h, gh, g^{i}h\}$ or $\{h, gh, g^{m/2}\}$, and Γ is a multiloop or $C_{m/2}^{(2)}$.

Proof. Suppose that \mathcal{M} is *G*-vertex-rotary with a rotary pair (a, z). Then $G = \langle a, z \rangle$, and so $a \neq 1$. It follows that $a = g^i$ or $a = g^i h$ for some integer *i*. Consequently, (a, z) is of the form $(g^i, g^j h)$ with gcd(i, m) = 1, or $(h, g^j h)$ with gcd(j, m) = 1. By equivalence under Aut(G), the pair (a, z) is either $(g, g^j h)$ or (h, gh).

The subgroup $\langle a \rangle$ corresponds to a vertex stabilizer in G, and its order equals the vertex valency of Γ . When |a| = m, this implies $\Gamma \cong K_2^{(m)}$; if |a| = 2, then $\Gamma \cong C_m$. This aligns with case (1).

Now assume \mathcal{M} is *G*-vertex-reversing with a reversing triple (x, y, z). In the dihedral group *G*, any two non-commuting involutions generate *G*; let these be *x* and *y*. Without loss of generality, we may set (x, y) = (h, gh). The third involution *z* can be chosen arbitrarily.

After possible relabeling, the triple $\{x, y, z\}$ is equivalent to:

- $\{h, gh, g^ih\}$ for some *i*, yielding a multiloop graph Γ , or
- $\{h, gh, g^{m/2}\}$ when m is even, resulting in $\Gamma \cong C_{m/2}^{(2)}$.

Up to duality, vertex stabilizers in G are isomorphic to either G or D_4 , determining the graph structure as above.

We now consider non-dihedral groups G and characterize necessary conditions for the existence of rotary pairs in G.

Lemma 5.2. Let \mathcal{M} be a G-arc-transitive map. Assume that $G = \langle g \rangle : \langle h \rangle$, where $C_{\langle h \rangle}(g) \leq \Phi(\langle h \rangle)$ and $|h| \geq 3$. Then the following statements hold:

- (1) The map \mathcal{M} is G-vertex-rotary.
- (2) The group satisfies $G = \langle g \rangle : \langle h \rangle$ with |h| = 2m and $g^{h^m} = g^{-1}$.
- (3) A rotary pair (a, z) is equivalent to (gh^i, h^m) or (gh^{2i}, h^m) , where gcd(i, |h|) = 1. The latter case occurs only if $G_2 \cong D_{2^e}$ with $e \ge 3$.
- (4) Let $\langle h_0 \rangle = C_{\langle h \rangle}(g)$. The number of inequivalent rotary pairs for G is $\frac{2\varphi(|h|)}{|\tau_{h_0}|}$ if $G_2 \cong D_{2^e}$ with $e \ge 3$, and $\frac{\varphi(|h|)}{|\tau_{h_0}|}$ otherwise.

Proof. The group G has no reversing triples. Suppose otherwise, then $G/\langle g \rangle \cong \langle h \rangle$ with $|h| \ge 3$ would be generated by involutions, leading to a contradiction. This establishes part (1).

It remains to classify the rotary pairs of G. Assume first that |h| is odd. Since $\langle g \rangle : \langle h \rangle$ is indecomposable and non-cyclic, |g| is also odd. Consequently, there exist no involutions in G.

Hence, $\langle h \rangle$ have even order 2m, containing a unique involution $u = h^m$. If u centralizes a Sylow *p*-subgroup $\langle g \rangle_p$ of $\langle g \rangle$, then $\langle g \rangle_{p'}: \langle u \rangle$ would be normal in G, making the quotient $G/(\langle g \rangle_{p'}: \langle u \rangle)$ cyclic. This would split G as $\langle g \rangle_p \times G_{p'}$, contradicting the assumption. Therefore, u centralizes no Sylow subgroup of $\langle g \rangle$, and $g^u = g^{-1}$ follows as in part (2).

We now classify rotary pairs of G. A involution z is equivalent to u under $\operatorname{Aut}(G)$, so set $z = u = h^m$. Write $a = g^i h^j$ for integers i, j. From $G = \langle a, z \rangle = \langle g^i h^j, u \rangle \leqslant \langle g^i, h^j, u \rangle$, we deduce $\operatorname{gcd}(i, |g|) = 1$ and $\operatorname{gcd}(j, |h|) \mid 2$. Note $\operatorname{gcd}(j, |h|) = 2$ requires $|h|_2 = 2$. Theorem 4.1 further implies a is equivalent to gh^j or gh^{2j} where $\operatorname{gcd}(j, |h|) = 1$.

Still, we need to check that $\langle a, z \rangle = G$. It follows from $g^u = g^{-1}$ that $[z, a] = a^z a^{-1} = g^{-2}$, and thus

$$\langle a, z \rangle = \langle g_{2'}, a_2, a_{\pi(|h|)_{2'}}, z \rangle.$$

Suppose that $|h_2| \ge 4$. Then $g_2 = 1$, so $\langle a_2 \rangle = \langle h_2 \rangle$ is a Sylow 2-subgroup of G. On the other hand, $\langle a_2, z \rangle$ is also a Sylow 2-subgroup of G when $|h_2| = 2$, since $h_2 = u$ and $\langle a_2, z \rangle = \langle g_2, u \rangle$. In either case, $\langle a, z \rangle = G$ holds, proving part (3).

Next, we show part (4): There are |g| involutions in $G \setminus Z(G)$. By part (3), there are $\varphi(|g|)\varphi(|h|)$ rotary pairs (a, z) for each $z \in G \setminus Z(G)$. Since $\langle a, z \rangle = G$, Aut(G) acts semiregularly on the set of rotary pairs of G. By Theorem 4.1, $|\operatorname{Aut}(G)| = |g|\varphi(|g|)|\tau_{h_0}|$ where $\langle h_0 \rangle = C_{\langle h \rangle}(g)$. Thus, there are $2\varphi(|h|)/|\tau_{h_0}|$ inequivalent rotary pairs for G when $G_2 \cong D_{2^e}$ with $e \ge 3$, and $\varphi(|h|)/|\tau_{h_0}|$ otherwise. This completes the proof of part (4).

6. Groups of types (II)-(III): 2-Nilpotent groups

In this section, we determine rotary pairs and reversing triples in the 2-nilpotent groups G given in parts (II)-(III) of Proposition 4.1. The first lemma treats the groups in (II) of Proposition 4.1.

Lemma 6.1. Let $G = (\langle g \rangle \times \langle g_v \rangle):(\langle u, v \rangle \times \langle h \rangle)$, where $|g|, |g_v|$ and |h| are pairwise coprime, $C_{\langle h \rangle}(gg_v) \leq \Phi(\langle h \rangle)$, and $\langle u, v \rangle = D_{2^e}$ with $e \geq 2$ such that

$$(g, g_v)^v = (g, g_v^{-1}) \text{ and } (g, g_v)^u = (g, g_v),$$

where $g_v \neq 1$.

Let \mathcal{M} be a G-arc-transitive map. Assume that $G \leq \operatorname{Aut}(\mathcal{M})$. Then either

(1) \mathcal{M} is G-vertex-reversing, $G = \langle g_v \rangle : \langle u, v \rangle$, relabeling if necessary, a reversing triple (x, y, z) is equivalent to

$$(u, g_v w^{2j} v, v)$$
, or $(v, g_v w, u)$ when $|w| = 2$,

and there are $6 \cdot 2^{e-2}$ inequivalent reversing triples of G; or

(2) \mathcal{M} is G-vertex-rotary, and a rotary pair (a, z) is equivalent to

 (gg_vwh^j, v) or (gg_vvh^j, v) ,

where gcd(j, |h|) = 1, and there are $2\varphi(|h|)/|\tau_{h_0}|$ inequivalent rotary pairs of G.

Proof. Note that $C_{\langle u,v\rangle}(g) = \langle \rho^2, u \rangle = D_{2^{e-1}}$ since $g^u = g$ and $g^v = g^{-1}$.

Case 1. Suppose that (x, y, z) is a reversing triple of G.

Firstly, we claim that h = g = 1, and

$$G = \langle g_v \rangle : \langle u, v \rangle = (\langle g_v \rangle \times \langle w^2, u \rangle) : \langle v \rangle$$

By definition, $C_G(g) \ge \langle g, g_v \rangle : \langle u, v \rangle$, and |h|, |g| are odd. It follows that h = 1, as $\langle h \rangle \cong G/(\langle g \rangle : G_2)$ is generated by involutions, and so g = 1, as $G = \langle g \rangle \times \langle g \rangle : \langle u, v \rangle$. The claim is thereby proved.

Next, we determine the triple (x, y, z). The quotient group $G/\langle g \rangle$ is dihedral, hence is generated by two involutions of x, y and z, say x and z without loss of generality. Hence $\langle x, y \rangle$ is a Sylow 2-subgroup of G, and so we may assume that $x, z \in \langle u, v \rangle$ by Sylow's theorem. By Lemma 4.1, $\operatorname{Aut}(G) = \langle \tilde{g_v} \rangle$: $\operatorname{Aut}(g_v) \times \langle \xi^2 \rangle$: $\langle \mu \rangle$, where $\langle \xi^2 \rangle$: $\langle \mu \rangle \leq \operatorname{Aut}(G)_{g_v} \cap \operatorname{Aut}(G)_{\langle u,v \rangle}$. Hence $\{x, y\}$ is equivalent to $\{u, v\}$. Now we have that $y = g_v^i t$, where $t \in \langle u, v \rangle$, and the integer i satisfies $\operatorname{gcd}(i, |g_v|) = 1$. Otherwise $\langle x, y, z \rangle \leq \langle g_v^i, u, v, w^{2j}v \rangle < G$. Also, the involution t is either $w^{2j}v$ or wwith |w| = 2.

Consequently, the triple (x, y, z) is equivalent to

$$(u, g_v w^{2j} v, v)$$
, or $(v, g_v w, u)$ when $|w| = 2$,

permuting x, y if necessary. Therefore, the group G has 2^{e-1} inequivalent reversing triples, as in statement (1).

Case 2. Suppose that (a, z) be a rotary pair of G. By Sylow's theorem, we may assume that $z \in \langle u, v \rangle$. Furthermore, $z \notin \langle w^2, u \rangle$ since $G/C_{\langle u,v \rangle}(gg_v)$ is not cyclic. Hence, we may take

z = v.

(Under $\langle \mu \rangle$). Without loss of generality, there is

$$a = (gg_v)^i th^j$$

where i, j are integers and $t \in \langle u, v \rangle$. Integers i and j satisfy $gcd(i, |gg_v|) = 1$ and gcd(j, |h|) = 1, respectively, and $\langle t, v \rangle \cong \langle u, v \rangle$, since $G = \langle a, z \rangle \leq \langle (gg_v)^i, t, v, h^j \rangle$. Therefore, $t = w^{2k+1}$ or $w^{2k}u$. Finally, by Theorem 4.1, pairs $((gg_v)^i w^{2k+1} h^j, v)$ Next, we consider the groups listed in (III) of Proposition 4.1.

Lemma 6.2. Let $G = (\langle g_u \rangle \times \langle g_v \rangle \times \langle g \rangle):(\langle u, v \rangle \times \langle h \rangle)$, where $|g_u|, |g_v|, |g|$ and |h| are pairwise coprime, $C_{\langle h \rangle}(g_u g_v g) \leq \Phi(\langle h \rangle)$, and $\langle u, v \rangle = D_{2^e}$ with $e \geq 2$ such that

$$(g_u, g_v, g)^v = (g_u, g_v^{-1}, g^{-1}) \text{ and } (g_u, g_v, g)^u = (g_u^{-1}, g_v, g^{-1}),$$

where if $g_u = 1$, then $g_v \neq 1$ and $g \neq 1$;

Let \mathcal{M} be a G-arc-transitive map. Assume that $G \leq \operatorname{Aut}(\mathcal{M})$. Then g_v or $g_u = 1$. Suppose that $g_v = 1$. Then either

(1) \mathcal{M} is G-vertex-reversing, $G = \langle g, g_u \rangle : \langle u, v \rangle$, and relabeling if necessary, a reversing triple (x, y, z) is equivalent to

$$(g_u g^{k_1} w^{2i} u, g^{k_2} v, u)$$

where $gcd(k_1, k_2) = 1$, ℓ_1 is odd, and there are

$$6 \cdot 2^{e-3} \cdot \frac{|\{(k_1, k_2) \mid \gcd(k_1, k_2) = 1\}|}{\varphi(|g|)}$$

inequivalent reversing triples of G;

(2) \mathcal{M} is G-vertex-rotary, and a rotary pair (a, z) is equivalent to

 $(g_u gwh^{\ell}, u)$ or $(g_u gw^2 vh^{\ell}, u),$

and there are $2\varphi(|h|)/|\tau_{h_0}|$ inequivalent rotary pairs in G.

Proof. Case 1. Suppose that (x, y, z) is a reversing triple of G. Then the element h of odd order must satisfy h = 1, because the quotient group $\langle h \rangle \cong G/\langle g_1, g_2, g_3, u, v \rangle$ is generated by involutions. This implies

$$G = (\langle g_1 \rangle \times \langle g_2 \rangle \times \langle g_3 \rangle) : \langle u, v \rangle,$$

and we proceed to classify the involutions of G. Without loss of generality, let $t = g_u^i g_v^j g^k w^{\ell} u$ be an involution for integers i, j, k, ℓ . If ℓ is odd, then conjugation by $w^{\ell} u$ maps (g_u, g_v, g) to (g_u, g_v^{-1}, g^{-1}) . Therefore,

$$t^{2} = g_{u}^{i}g_{v}^{j}g^{k}w^{\ell}ug_{u}^{i}g_{v}^{j}g^{k}w^{\ell}u = g_{u}^{i}g_{v}^{j}g^{k} \cdot g_{u}^{i}g_{v}^{-j}g^{-k} = g_{u}^{2i} = 1.$$

Since $|g_u|$ is odd, this forces $g_u^i = 1$. Thus, $t = g_v^j g^k w^\ell u$. Similarly, if ℓ is even, we derive $g_v^j = 1$ and $t = g_u^j g^k w^\ell u$. Additionally, the involution $w^{2^{e-2}}$ cannot belong to any reversing triple.

By Sylow's theorem, we may assume $z \in \langle u, v \rangle$, so

$$z = w^m u$$

for some integer m. Let

$$x = g_u^{i_1} g_v^{j_1} g^{k_1} w^{\ell_1} u, \ y = g_u^{i_2} g_v^{j_2} g^{k_2} w^{\ell_2} u,$$

where $i_1, j_1, k_1, \ell_1, i_2, j_2, k_2, \ell_2$ are integers. If both ℓ_1 and ℓ_2 are even, then $g_v^{j_1} = g_v^{j_2} = 1$. However, this leads to $\langle x, y, z \rangle \leq \langle g_u, g, u, v \rangle < G$, a contradiction. Similarly, both ℓ_1 and ℓ_2 being odd also yields a contradiction. Hence, after relabeling, we may take ℓ_1 even and ℓ_2 odd. This implies $g_v^{j_1} = 1$ and $g_u^{i_2} = 1$, simplifying to:

$$x = g_u^{i_1} g^{k_1} w^{\ell_1} u, \quad y = g_v^{j_2} g^{k_2} w^{\ell_2} u.$$

Since $\langle x, y, z \rangle \leqslant \langle g_u^{i_1}, g^{k_1}, w^{\ell_1}, u, g_v^{j_2}, g^{k_2}, w^{\ell_2}u, w^m u \rangle = G$, we require

$$gcd(i_1, |g_u|) = gcd(j_2, |g_v|) = gcd(k_1, k_2) = 1.$$

By Theorem 4.1,

$$\operatorname{Aut}(G)_{(L)} \ge \operatorname{Aut}(H) \text{ and } \operatorname{Aut}(G)_L \cap \operatorname{Aut}(G)_{(H)} \ge \langle \xi^2 \rangle : \langle \mu \rangle.$$

Consequently, the triple (x, y, z) is equivalent to either

$$(g_u g^{k_1} w^{\ell_1} u, g_v g^{k_2} v, u)$$
 or $(g_u g^{k_1} w^{\ell_1} u, g_v g^{k_2} v, v),$

as stated in part (i).

Now we verify whether the above triples generate G. First, consider the triple $(x, y, z) = (g_u g^{k_1} w^{\ell_1} u, g_v g^{k_2} v, u)$, where ℓ_1 is even. Using the conjugation relation $(g_u, g_v, g)^w = (g_u^{-1}, g_v^{-1}, g)$ and the condition $gcd(k_1, k_2) = 1$, we compute:

$$\langle x, y, z \rangle = \langle g^{k_1}, g^{k_2}, g_u, w^{\ell_1}, g_v w^{-1}, u \rangle$$

= $\langle g, g_u, w^{\ell_1}, g_v w^{-1}, u \rangle.$

Observe that the quotient group

$$G = \langle x, y, z \rangle / \langle g, g_u \rangle \cong \langle w^{\ell_1}, g_u w^{-1}, u \rangle \cong \langle u, v \rangle$$

forces $g_v = 1$. A similar argument shows $g_u = 1$ for the cases that (x, y, z) is equivalent to $(g_u g^{k_1} w^{\ell_1} u, g_v g^{k_2} v, v)$.

Conversely, if $g_v = 1$ (respectively $g_u = 1$), the triple $(g_u g^{k_1} w^{\ell_1} u, g^{k_2} v, u)$ (respectively $(g_u g^{k_1} w^{\ell_1} u, g^{k_2} v, v)$) indeed generates G, by the argument above.

Thus, G has a reversing triple (x, y, z) if and only if one of the following holds:

(1) $g_v = 1$, and the triple is equivalent to $(g_u g^{k_1} w^{\ell_1} u, g^{k_2} v, u);$

(2) $g_u = 1$, and the triple is equivalent to $(g^{k_1}w^{\ell_1}u, g_vg^{k_2}v, v)$,

where $gcd(k_1, k_2) = 1$, $\ell_1 = 2i$ is even.

The number of inequivalent reversing triples is given by:

$$\frac{\{(k_1, k_2) \mid \gcd(k_1, k_2) = 1\}|}{\varphi(|g|)} \cdot 2^{e-2}.$$

Case 2. Suppose that (a, z) is a rotary pair of G. By Sylow's theorem, we may assume that $z \in \langle u, v \rangle$, so that

$$z = w^m u,$$

for some integer m. Without loss of generality,

$$a = g_u^i g_v^j g^k t h^\ell,$$

where $t \in \langle u, v \rangle$, and i, j, k, ℓ are integers. These integers i, j, k, ℓ and the element t satisfy

$$\gcd(i, |g_u|) = \gcd(j, |g_v|) = \gcd(k, |g|) = \gcd(\ell, |h|) = 1 \text{ and } \langle t, \rho^m u \rangle = \langle u, v \rangle,$$

so that $\langle a, z \rangle \leq \langle g_u^i, g_v^j, g^k, t, w^m u, h^\ell \rangle = G$. Here by Theorem 4.1,

 $\operatorname{Aut}(G)_{(L)} \ge \operatorname{Aut}(H) \text{ and } \operatorname{Aut}(G)_L \cap \operatorname{Aut}(G)_{(H)} \ge \langle \xi^2 \rangle : \langle \mu \rangle,$

where $H = \langle g_u g_v g \rangle$ and $L = \langle u, v \rangle \times \langle h \rangle$. The pair (a, z) is equivalent to

 $(g_u g_v gth^\ell, z),$

where $gcd(\ell, |h|) = 1$ and $(t, z) \in \{(w^i, u), (w^i, v), (w^{2i}v, u), (w^{2i}u, v)\}$ with i being an odd integer.

Next, we verify whether these pairs generate G. The key observation is that $G' = \langle [a, z] \rangle$, which holds because

$$\langle [a, z] \rangle$$
 char $\langle g_u g_v g \rangle \times \langle w^2 \rangle \lhd G$.

Suppose $(a, z) = (g_u g_v g w^i h^{\ell}, u)$ with *i* odd. Then $G' = \langle [a, z] \rangle = \langle g_u^2 g^2 w^2 \rangle$, and the abelian quotient

 $G/G' \cong \langle g_v \rangle \times \langle u, v \rangle / \langle w^2 \rangle \times \langle h \rangle.$

This implies $g_v^v = g_v$ since $gcd(|G'|, |g_v|) = 1$. From the definition of g_v , it follows that $g_v = 1$. Similarly, if z = v, then $g_u = 1$ must hold.

Conversely, if $g_u = 1$ (respectively $g_v = 1$), the pairs $(g_v g w^i h^\ell, u)$ and $(g_v g w^{2i} v h^\ell, u)$ (respectively $(g_u g w^i h^\ell, v)$ and $(g_u g w^{2i} u h^\ell, v)$) generate G by the argument above.

In summary, G admits a rotary pair (a, z) if and only if one of the following holds:

(1) $g_v = 1$, and (a, z) is equivalent to $(g_u g w h^{\ell}, u)$ or $(g_u g w^2 v h^{\ell}, u)$;

(2) $g_u = 1$, and (a, z) is equivalent to $(g_v g w h^{\ell}, v)$ or $(g_v g w^2 u h^{\ell}, v)$;

Furthermore, there are $\frac{2\varphi(|h|)}{|\tau_{h_0}|}$ inequivalent rotary pairs in G.

7. Groups of types (IV)-(V)

In this section, we assume that G is a group of type (IV) or type (V). We analyse the two cases separately.

7.1. Groups of type (IV). The first lemma treats groups of type IV.

Lemma 7.1. Let $G = (\langle g \rangle \times \langle u, v \rangle): \langle h \rangle = (\mathbb{Z}_m \times \mathbb{Z}_2^2):\mathbb{Z}_{3^{f_n}}$ with $G/O_{2'}(G) = A_4$, where m, n, 6 are pairwise coprime. Then \mathcal{M} is G-vertex-rotary with a rotary pair (a, z), and

(1) $G = \langle u, v \rangle : \langle h \rangle = \mathbb{Z}_2^2 : \mathbb{Z}_{3\lambda} = \mathbb{Z}_{\lambda} . A_4, and$

(2) $a = h^i$ with $gcd(i, 3\lambda) = 1$, and $z \in \{u, v, uv\}$, and

(3) the underlying graph of \mathcal{M} is $\Gamma = K_4^{(\lambda)}$.

Proof. Let $N = \langle u, v \rangle = \mathbb{Z}_2^2$, and let $C = C_G(N)$. Then $G/C \leq \operatorname{Aut}(N)$, and hence $G/C = \mathbb{Z}_3$. It follows that \mathcal{M} is G-vertex-rotary, and $\langle g \rangle : \langle h \rangle \cong G/N$ is of odd order and generated by a single element. Thus g = 1, and $G = N: G_{2'} = \langle u, v \rangle : \langle h \rangle \cong \mathbb{Z}_{\lambda}.$ A₄, with $\lambda = |h|/3$, as in part (1).

Let (a, z) be a rotary pair for G. Then $z \in \{u, v, uv\}$, and a is conjugate to a generator of $\langle h \rangle$. Without loss of generality, we may let $a = h^i$, with gcd(i, |h|) = 1, as in part (2).

Finally, the vertex stabilizer $G_{\alpha} = \langle a \rangle \cong \mathbb{Z}_{3\lambda}$, and $|G : G_{\alpha}| = 4$. Thus |V| = 4, and the kernel $K = G_{(V)} = \mathbb{Z}_{\lambda}$, and so $\Gamma = \mathrm{K}_{4}^{(\lambda)}$, as in part (3).

7.2. Groups of type (V).

Lemma 7.2. Let $G = (\langle g \rangle \times \langle w^2, v \rangle):((\langle h \rangle: \langle u \rangle) \times \langle c \rangle)$, where $\langle u, v \rangle = D_8$, $\langle h \rangle$ is a Sylow 3-subgroup, both $\langle g \rangle$ and $\langle c \rangle$ are Hall subgroups such that $C_{\langle c \rangle}(g) \leq \Phi(\langle c \rangle)$, $(w^2v, v, w^2)^h = (v, w^2, w^2v)$ and $h^u = h^{-1}$.

Assume that G is indecomposable, and G has a rotary pair or a reversing triple. Then $g^u = g^{-1}$ and [g, h] = 1.

Proof. Suppose that $\langle g_p \rangle$ be a Sylow *p*-subgroup of $\langle g \rangle$ such that $[g_p, u] = 1$. Then $\langle g_p \rangle : \langle c \rangle$ is non-abelian, otherwise, $G = \langle g_p \rangle \times G_{p'}$, contradiction G being indecomposable. Thus the odd order factor group

$$\overline{G} := G/\langle g_{p'}, u, v, h \rangle \cong \langle g_p \rangle : \langle c \rangle$$

is non-abelian.

Let (a, z) be a rotary pair and (x, y, z) be a reversing triple, and so either $G = \langle a, z \rangle$ or $G = \langle x, y, z \rangle$. By Sylow's theorem, we assume that $z \in \langle u, v \rangle$. It follows that the factor group \overline{G} is either cyclic or dihedral, contradiction. Therefore, $g_p^w = g_p^{-1}$ for each $p \in \pi(|g|)$, and so $g^u = g^{-1}$. Let $C = C_G(g)$. The abelian group

$$G/C \ge \langle uC \rangle.$$

Hence $h \in C$, equivalently [g, h] = 1, otherwise $\langle hC, uC \rangle$ is a non-abelian subgroup of G/C.

Before treating the groups of type (V), we give a typical construction for rotary pairs in this case, which is built upon the symmetric group $G = \text{Sym}\{1, 2, 3, 4\} = S_4$, and a rotary pair ((123), (14)).

Example 7.3. We construct rotary pairs in some groups of type (V).

(1) Let $G = \text{Sym}\{1, 2, 3, 4\} = \langle u, v \rangle : (\langle h_0 \rangle : \langle w \rangle) = \mathbb{Z}_2^2 : S_3 = S_4$, where u = (12)(34), v = (13)(24), w = (14) and $h_0 = (124)$. Let

$$(a_1, z_1) = ((123), (14)).$$

(2) Let $X = \langle u, v \rangle$: $(\langle h \rangle : \langle w \rangle) = \mathbb{Z}_2^2$: $\mathbb{D}_{2,3^f} = \langle h^3 \rangle$. $G = \mathbb{Z}_{3^{f-1}}$. \mathbb{S}_4 , where $\mathbb{Z}_2^2 = \langle u, v \rangle < \langle u, w \rangle = \langle v, w \rangle = \mathbb{D}_8$, $|h| = 3^f$, and $(u, v, uv)^h = (v, uv, u)$, and $h^w = h^{-1}$. Let $(a_2, z_2) = (h, w^u)$.

Noticing that $h^u = uhu = hvu$ and $(h^u)^w (h^u)^{-1} = h^{-2}$, we have that $h \in \langle h^u, w \rangle$, and hence $\langle h^u, w \rangle = X$. So $(a_2, z_2) = (h^u, w)^u$ is a rotary pair for X.

(3) Let $Y = \langle g \rangle : X = (\langle g \rangle \times \langle u, v \rangle) : (\langle h \rangle : \langle w \rangle) = (\mathbb{Z}_m \times \mathbb{Z}_2^2) : \mathbb{D}_{2.3^f}$, where |g| = m is coprime to 3, gh = hg and $g^w = g^{-1}$. Let

 $(a_3, z_3) = (ga_2, z_2) = (gh, w^u).$

Then $\langle a_3, z_3 \rangle = \langle gh, z_2 \rangle = \langle g, h, z_2 \rangle = \langle g \rangle : \langle h, z_2 \rangle = Y$, so that (a_3, z_3) is a rotary pair for the group Y.

(4) Let $Z = Y: \langle c \rangle = (\langle g \rangle \times \langle u, v \rangle): ((\langle h \rangle: \langle w \rangle) \times \langle c \rangle) = (\mathbb{Z}_m \times \mathbb{Z}_2^2): (\mathbb{D}_{2.3^f} \times \mathbb{Z}_n)$, where [h, Y] = 1 and $[\langle g \rangle, \langle c \rangle] \leq \Phi(\langle g \rangle)$. Let $(a_4, z_4) = (a_3 c, z_3) = (ghc, w^u).$ Then $\langle a_4, z_4 \rangle = \langle c, gh, z_3 \rangle = Y: \langle c \rangle = Z$, so (a_4, z_4) is a rotary pair for Z. \Box

The following lemma determines rotary pairs and reversing triples in the groups of type (V). Let

$$G = (\langle g \rangle \times \langle w^2, v \rangle) : ((\langle h \rangle : \langle u \rangle) \times \langle c \rangle),$$

a group of type (V). We first consider the subgroup $\langle w^2, v \rangle : \langle h, u \rangle$ of G.

Lemma 7.4. Following notations above, $t \in \langle w^2, v \rangle$ is such that $\langle th, u \rangle = \langle w^2, v \rangle$: $\langle h, u \rangle$ if and only if t = v or w^2 .

Proof. From $(w^2v, v, w^2)^h = (v, w^2, w^2v)$ and $h^u = h^{-1}$, we have: $\langle th, u \rangle = D_{4\cdot 3^e} \iff (th)^u = (th)^{-1} \iff t^u h^{-1} = h^{-1}t \iff t^u = t^h \iff t \in \{1, w^2v\}.$ Since $D_{4\cdot 3^e}$ is a maximal subgroup of $\langle w^2, v \rangle : \langle h, w \rangle \cong \mathbb{Z}_{3^{e-1}}.S_4$, we have that $\langle th, u \rangle = \langle w^2, v \rangle : \langle h, w \rangle$ if and only if

$$t \in \langle u, v \rangle \backslash \{1, w^2 v\} = \{v, w^2\}.$$

Suppose that $t = w^2$. Then $(w^2h)^2 u^{w^2h} = w^2 v h^2 h^{-2} u = w$, and so $\langle w^2h, u \rangle = \langle w^2, v \rangle : \langle h, u \rangle$. Similarly, it is straightforward to check that $\langle vh, u \rangle = \langle w^2, v \rangle : \langle h, u \rangle$. \Box

Now, we are ready to classify rotary pairs and reversing triples of groups of type V.

Lemma 7.5. Following notations in Lemma 7.2. Assume that \mathcal{M} is G-vertexreversing with a reversing triple (x, y, z), and let $t, t' \in \langle u, v \rangle$, i be an integer such that gcd(i, |gh|) = 1. Then c = 1, and relabeling if necessary, the reversing triple (x, y, z) is equivalent to one of the following triples:

- (1) $(t(gh)^{i}u, t', u)$, where $t' \neq 1$;
- (2) $(t(gh)^{i}u, g^{i'}h^{3j_0}w^2u, u);$
- (3) $(t(gh)^{i}u, g^{i'}t'h^{j'}u, u)$, where $(t, t') \neq (1, 1)$

Proof. First, c = 1, since every factor group of G is generated by involutions. By Lemma 7.2, we have that [g, h] = 1. Thus $\overline{G} := G/\langle w^2, v \rangle \cong \langle gh \rangle : \langle u \rangle$ is a dihedral group.

For each $l \in G$, let \overline{l} denote the image of l in \overline{G} . Among the three involutions $\overline{x}, \overline{y}$, and \overline{z} , two generate the dihedral group \overline{G} . Without loss of generality, assume these are \overline{x} and \overline{z} . It follows that $z \notin \langle w^2, v \rangle$, and therefore we may assume

$$z = u$$

by Sylow's theorem. Write

$$x = t(gh)^i u$$

where $t \in \langle w^2, v \rangle$ and gcd(i, |gh|) = 1. Since x is an involution, the element t satisfies $t^u = t^{h^i}$.

implying $t \in \{w^2v, 1\}$ if $i \equiv 1 \pmod{3}$, or $t \in \{v, 1\}$ if $i \equiv 2 \pmod{3}$. The remaining involution y is either in $\langle w^2, v \rangle$ or of the following:

$$\begin{split} g^i w^2 v h^j u &= g^i h^j v u \quad (j \equiv 1 \pmod{3}); \\ g^i v h^j u &= g^i h^j w \qquad (j \equiv 2 \pmod{3}); \\ g^i h^{3j_0} w^2 u \text{ or } g^i h^j u. \end{split}$$

Suppose $y \in \{w^2v, v, w^2\}$. Then $\{w^2v, v, w^2\} = y^{\langle xz \rangle}$, and thus $\langle x, y, z \rangle = \langle t(gh)^i u, t', u \rangle = G$, as in case (1).

Suppose $y = g^{i'}h^{3j_0}w^2u$. Then $\langle y, z \rangle = (\langle w^2 \rangle \times \langle g^i, h^{3j_0} \rangle): \langle u \rangle$, which implies $w^2 \in \langle x, y, z \rangle$. Since $\{w^2v, v, w^2\} = (w^2)^{\langle xz \rangle}$, it follows that $\langle x, y, z \rangle = G$, as in case (2).

For the remaining choices of y, direct verification shows $\langle x, y, z \rangle = G$ except when t = 1 and $y = (gh)^{i'}u$, as described in case (3).

Lemma 7.6. Following notations in Lemma 7.2. Assume that \mathcal{M} is G-vertex-rotary with a rotary pair (a, z). Then (a, z) is equivalent to

 $(gth^{j}c^{k}, u) \text{ or } (gth^{j}uc^{k}, u),$ where gcd(j, 3) = gcd(k, |c|) = 1, and $t \in \{v, w^{2}\}$. There are $\varphi(|h|)\varphi(|c|)/2|\tau_{h^{3}}||\tau_{c_{0}}|$

inequivalent rotary pairs of G.

Proof. By Sylow's theorem, we may assume that the involution $z \in \langle u, v \rangle$. Suppose that $z \in \langle w^2, v \rangle$. Then $G/\langle u, v \rangle \cong \langle g \rangle : ((\langle h \rangle : \langle u \rangle) \times \langle c \rangle)$ is isomorphic to a subgroup of $\langle a \rangle$. This is not possible since $\langle g \rangle : ((\langle h \rangle : \langle u \rangle) \times \langle c \rangle)$ is non-abelian. Hence $z \in \langle u, v \rangle \setminus \langle w^2, v \rangle = \{u, w^2u\}$, and we may further let

z = u,

as $(w^2 u)^v = w$.

Since $G = (\langle g \rangle \times \langle w^2, v \rangle):((\langle h \rangle: \langle u \rangle) \times \langle c \rangle)$, the element *a* has two possibilities:

either
$$a = g^i t h^j u c^k$$
, or $a = g^i t h^j c^k$

where i, j, k are integers and $t \in \langle w^2, v \rangle$. In both cases, one always have that

$$G = \langle a, z \rangle \leqslant \langle g^i, t, h^j, w, c^k \rangle,$$

which implies that gcd(i, |g|) = gcd(j, 3) = gcd(k, |c|) = 1, and $t \neq 1$. Again in either cases, one have that $\langle w^2, v \rangle : \langle h, u \rangle \times \langle c \rangle \cong \langle a, z \rangle / \langle g \rangle \cong \langle th^j c^k, u \rangle$. Hence t = v or w^2 by Lemma 7.4.

Conversely, we show explicitly that the above pairs generate G. Without loss of generality, let z = u and $a = gthu^{\ell}c$, where $t \in \{v, w^2\}$ and $\ell \in \{0, 1\}$. Then $\langle a, z \rangle = \langle gthc, u \rangle$, and we let a' = gthc. Then

$$a'^{2} = gg^{c^{-1}}tt^{h^{-1}}h^{2}c^{2}$$
 and $z^{a'}z = (g^{2})^{ch^{-1}}(t^{u}t)^{h^{-1}}h^{2}$.

16

Since gcd(|g|, 6) = 1 and $[g, \langle w^2, v \rangle : \langle h \rangle] = 1$, we have that

$$g \in \langle z^a \ z \rangle.$$

So $\langle tt^{h^{-1}}h^2c^2 \rangle \leqslant \langle g, a'^2 \rangle.$ As $gcd(|c|, 6) = 1$ and $[c, \langle w^2, v \rangle: \langle h \rangle] = 1,$
 $c \in \langle g, a'^2 \rangle.$
It follows from Lemma 7.4 that $\langle g, t \rangle = \langle g, ath, v \rangle = C$ on $t \in \{w^2, v\}$.

It follows from Lemma 7.4 that $\langle a, z \rangle = \langle g, c, th, u \rangle = G$ as $t \in \{w^2, v\}$.

8. Proof of Theorem 1.1

Let $\mathcal{M} = (V, E, F)$ be a *G*-arc-transitive map with $G \leq \operatorname{Aut}(\mathcal{M})$ and underlying graph $\Gamma = (V, E)$. Assume that $\operatorname{gcd}(\chi(\mathcal{M}), |E|) = 1$. Then each Sylow subgroup of *G* is cyclic or dihedral by Lemma 3.2, and then *G* is divided into five types in Proposition 4.1. The groups of type (I) are treated in Lemma 5.1-5.2, which show that \mathcal{M} and *G* satisfy part (1) of Theorem 1.1. Then the groups of types (II)-(III) are dealt with by Lemmas 6.1 and 6.2, which show that \mathcal{M} and *G* satisfy parts (2)-(3) of Theorem 1.1. Finally, the groups of types (IV)-(V) are treated in Lemmas 7.1, 7.6 and 7.5. This completes the proof of Theorem 1.1. \Box

References

- [1] J. N. Bray, D. F. Holt, C. M. Roney-Dougal, The maximal subgroups of the low-dimensional finite classical groups. *London Mathematical Society Lecture Note Series*, **407** (2013), xiv+438.
- [2] J. Y. Chen, Edge-transitive maps with few vertices and faces, in preparation.
- [3] M. Conder, P. Potočnik, J. Širáň, Regular maps with almost Sylow-cyclic automorphism groups, and classification of regular maps with Euler characteristic $-p^2$. J. Algebra, **324** (2010), 2620–2635.
- [4] M. Conder, J. Širáň, T. W. Tucke, The genera, reflexibility and simplicity of regular maps. J. Eur. Math. Soc., 12 (2010), 343–364.
- [5] A. B. d'Azevedo, D.A. Catalano, J. Širáň, Bi-rotary maps of negative prime characteristic. Ann. Comb. 23 (2019), 27-50.
- [6] A. B. d'Azevedo, R. Nedela, J. Širáň, Classification of regular maps of negative prime Euler characteristic. Trans. Amer. Math. Soc., 357 (2004), 4175–4190.
- [7] Gill, Nick, Orientably regular maps with Euler characteristic divisible by few primes. J. Lond. Math. Soc. (2) 88 (2013), 118-136.
- [8] D. Gorenstein, J. H. Walter, The characterization of finite groups with dihedral Sylow 2subgroups. I. J. Algebra, 2 (1965), 218–270.
- [9] J. E. Graver, M. E. Watkins, Locally finite, planar, edge-transitive graphs. Mem. Amer. Math. Soc, 126 (1997), vi+75.
- [10] Zheng Guo and Luyi Liu, Arc-transitive embeddings of multicycles, in preparation.
- [11] S. Gyurki, S. Pavlikova and J. Siran, Orientably-regular embeddings of complete multigraphs, preprint, arXiv:2311.096666v1 [math.CO] 16 Nov 2023.
- [12] A. W. Joseph, Spaces of constant curvature. AMS Chelsea, (2011), xviii+424.
- [13] G. A. Jones, Automorphism groups of edge-transitive maps. Acta Math. Univ. Comenian. (N.S.), 88 (2019), 841–847
- [14] C. H. Li and Luyi Liu, Arc-transitive maps with edge number coprime to the Euler characteristic – II, preprint.
- [15] C. H. Li and Luyi Liu, Automorphism groups of finite almost Sylow-cyclic groups, preprint.
- [16] C. H. Li, C. E. Praeger, S. J. Song, Locally finite vertex-rotary maps and coset graphs with finite valency and finite edge multiplicity. *J. Combin. Theory Ser. B*, **169** (2024), 1–44.

[17] C. H. Li, C. E. Praeger, S. J. Song, A new characterisation of the five types of arc-transitive maps, in preparation.

[18] J. Širáň, How symmetric can maps on surfaces be? London Math. Soc. Lecture Note Ser, 409 (2013), 161–238

[19] M. Suzuki, On finite groups with cyclic Sylow subgroups for all odd primes. Amer. J. Math., 77 (1955), 657–691.

[20] W. J. Wong, On finite groups with semi-dihedral Sylow 2-subgroups. J. Algebra, 4 (1966), 52–63.

[21] H. Zassenhaus, Über endliche Fastkörper, Abh. Math. Sem. Univ. Hamburg 11 (1935), 187-220.

SUSTECH INTERNATIONAL CENTER FOR MATHEMATICS, DEPARTMENT OF MATHEMATICS, SOUTHERN UNIVERSITY OF SCIENCE AND TECHNOLOGY, SHENZHEN, GUANGDONG 518055, P. R. CHINA

Email address: lich@sustech.edu.cn

SHENZHEN INTERNATIONAL CENTER FOR MATHEMATICS, SOUTHERN UNIVERSITY OF SCI-ENCE AND TECHNOLOGY, SHENZHEN, GUANGDONG 518055, P. R. CHINA

Email address: 12031108@mail.sustech.edu.cn