

SOME PROPERTIES OF OVERPARTITIONS INTO NONMULTIPLES OF TWO INTEGERS

ABDULAZIZ M. ALANAZI, AUGUSTINE O. MUNAGI, AND MANJIL P. SAIKIA

ABSTRACT. We consider properties of overpartitions that are simultaneously ℓ -regular and μ -regular, where ℓ and μ are positive relatively prime integers. We prove a seven-way combinatorial identity related to these overpartitions. We also prove several congruence properties satisfied by this class of partitions (and a further related class) using both generating functions and modular forms with Radu's Algorithm.

1. INTRODUCTION

A partition of a positive integer n is a non-increasing sequence of positive integers $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ that sum to n . The λ_i 's are called parts of the partition. An overpartition of n is a partition of n where the first occurrence of each part size may be overlined. We denote the number of overpartitions of n by $\overline{p}(n)$, with $\overline{p}(0) = 1$. For example, $\overline{p}(3) = 8$, which enumerates the following overpartitions

$$(3), (\overline{3}), (2, 1), (\overline{2}, 1), (2, \overline{1}), (\overline{2}, \overline{1}), (1, 1, 1), (\overline{1}, 1, 1).$$

The three overpartitions with no overlined parts are the ordinary partitions of 3. We will also use the alternative notation of a partition: $\lambda = (c_1^{u_1}, c_2^{u_2}, \dots, c_r^{u_r})$, where $c_1 > c_2 > \dots > c_r > 0$ and $r \leq k$.

Given a positive integer ℓ a partition λ is called ℓ -regular if no part of λ is a multiple of ℓ . Munagi and Sellers [9] studied combinatorial and arithmetic properties of overpartitions of n with ℓ -regular overlined parts, denoted by $A_\ell(n)$. Alanazi and Munagi [2] investigated the combinatorial identities for ℓ -regular overpartitions of n , denoted by $\overline{R}_\ell(n)$. In addition, Alanazi et. al. [1] studied combinatorial and arithmetic properties of the overpartitions of n with ℓ -regular non-overlined parts. We also note that Alanazi et. al. [3] and Shen [17] discussed various arithmetic properties of $\overline{R}_\ell(n)$. We also recall the generating function from [1]

$$(1) \quad \sum_{n \geq 0} \overline{R}_\ell^*(n) q^n = \prod_{n=1}^{\infty} \frac{(1 - q^{n\ell})(1 + q^n)}{1 - q^n} = \frac{f_2 f_\ell}{f_1^2},$$

where $\overline{R}_\ell^*(n)$ counts overpartitions of n wherein non-overlined parts are ℓ -regular and there are no restrictions on the overlined parts. Alanazi et. al. [1] proved several interesting results related to this function, some of which were recently extended by Sellers [16]. The second equality in (1) follows from the notation f_n^k is defined by $f_n^k := \prod_{i \geq 1} (1 - q^{in})^k$ for all integers n, k with $n > 0$.

In this paper, we extend the previous studies and consider properties of overpartitions that are simultaneously ℓ -regular and μ -regular, where ℓ and μ are positive relatively prime integers. Let $\overline{R}_{\ell, \mu}(n)$ be the number of overpartitions of n where no parts are divisible by ℓ or μ , where $\gcd(\ell, \mu) = 1$. We will also call this function the number of (ℓ, μ) -regular overpartitions of n . It is easy to see that the generating function is given by

$$(2) \quad \sum_{n \geq 0} \overline{R}_{\ell, \mu}(n) q^n = \prod_{n=1}^{\infty} \frac{(1 + q^n)(1 - q^{\ell n})(1 - q^{\mu n})(1 + q^{\ell \mu n})}{(1 - q^n)(1 + q^{\ell n})(1 + q^{\mu n})(1 - q^{\ell \mu n})} = \frac{f_2 f_\ell^2 f_\mu^2 f_{2\mu\ell}}{f_1^2 f_{2\ell} f_{2\mu} f_{\mu\ell}^2}.$$

The paper is organized as follows. In Section 2 we prove a seven-way partition identity using generating functions and combinatorial techniques thus establishing the equivalence of several sets of partitions. In Section 3 we prove some congruences modulo small powers of 2 satisfied by the functions $\overline{R}_{\ell, \mu}(n)$ and

2020 *Mathematics Subject Classification.* 05A17, 11P81, 11P83.

Key words and phrases. Overpartition, theta function, congruence, Modular Form, Radu's Algorithm.

The work of the last author is partially supported by an Ahmedabad University Start-Up Grant (Reference No. URBSASI24A5).

$\overline{R}_\ell^*(n)$. Then in Section 4 we prove further congruences for $\overline{R}_{\ell,\mu}(n)$ and $\overline{R}_\ell^*(n)$ using modular forms, and end the paper in Section 5 with concluding remarks.

2. A GENERAL PARTITION THEOREM

Our first result is the following seven-way identity.

Theorem 2.1. *Let*

- $A_{\ell,\mu}(\ell n)$ denote the number of μ -regular partitions of ℓn where parts that are non-multiples of ℓ appear ℓ times and parts that are multiples of ℓ appear less than ℓ times,
- $B_{\ell,\mu}(2n)$ denote the number of (ℓ, μ) -regular partitions of $2n$ in which ℓ and μ are odd and odd parts occur with even multiplicities,
- $C_{\ell,\mu}(2n)$ denote the number of μ -regular partitions of $2n$ in which odd parts appear with multiplicities $2, 4, \dots, 2(\ell - 2)$ or $2(\ell - 1)$ and even parts appear less than ℓ times, where $\ell < \mu$ and ℓ and μ are odd,
- $D_{\ell,\mu}(2n)$ denote the number of $(\ell, 2\mu)$ -regular partitions of $2n$ in which odd parts occur with even multiplicities and parts $\equiv \mu \pmod{2\mu}$ appear at most once, where μ is even,
- $E_{\ell,\mu}(2n)$ denote the number of (2μ) -regular partitions of $2n$ in which odd parts occur with multiplicities $2, 4, \dots, 2(\ell - 2)$ or $2(\ell - 1)$, even parts appear less than ℓ times and parts $\equiv \mu \pmod{2\mu}$ appear at most once, where $\ell < \mu$ and μ is even, and
- $F_{\ell,\mu}(\ell n)$ denote the number of (ℓ^2, μ) -regular partitions of ℓn in which parts that are non-multiples of ℓ appear either 0 or ℓ times.

Then

$$A_{\ell,\mu}(\ell n) = B_{\ell,\mu}(2n) = C_{\ell,\mu}(2n) = D_{\ell,\mu}(2n) = E_{\ell,\mu}(2n) = F_{\ell,\mu}(\ell n) = \overline{R}_{\ell,\mu}(n).$$

Proof. The generating function for $A_{\ell,\mu}(\ell n)$ is

$$\begin{aligned}
 \sum_{n=0}^{\infty} A_{\ell,\mu}(\ell n) q^{\ell n} &= \prod_{n=1}^{\infty} \frac{(1 + q^{\ell(\ell n - 1)}) \cdots (1 + q^{\ell(\ell n - (\ell - 1))}) (1 + q^{\ell n} + \cdots + q^{(\ell - 1)\ell n})}{(1 + q^{\mu \ell(\ell n - 1)}) \cdots (1 + q^{\mu \ell(\ell n - (\ell - 1))}) (1 + q^{\mu \ell n} + \cdots + q^{(\ell - 1)\mu \ell n})} \\
 &= \prod_{n=1}^{\infty} \frac{(1 + q^{\ell(\ell n - 1)}) \cdots (1 + q^{\ell(\ell n - (\ell - 1))}) (1 + q^{\ell n} + \cdots + q^{(\ell - 1)\ell n})}{(1 + q^{\mu \ell(\ell n - 1)}) \cdots (1 + q^{\mu \ell(\ell n - (\ell - 1))}) (1 + q^{\mu \ell n} + \cdots + q^{(\ell - 1)\mu \ell n})} \\
 (3) \quad &\quad \times \frac{(1 + q^{\ell^2 n})(1 + q^{\mu \ell^2 n})}{(1 + q^{\ell^2 n})(1 + q^{\mu \ell^2 n})} \\
 &= \prod_{n=1}^{\infty} \frac{(1 + q^{\ell n})(1 + q^{\ell n} + \cdots + q^{(\ell - 1)\ell n})(1 + q^{\mu \ell^2 n})}{(1 + q^{\mu \ell n})(1 + q^{\mu \ell n} + \cdots + q^{(\ell - 1)\mu \ell n})(1 + q^{\ell^2 n})} \\
 &= \prod_{n=1}^{\infty} \frac{(1 + q^{\ell n})(1 - q^{\ell^2 n})(1 - q^{\mu \ell n})(1 + q^{\mu \ell^2 n})}{(1 + q^{\mu \ell n})(1 - q^{\ell n})(1 - q^{\mu \ell^2 n})(1 + q^{\ell^2 n})}.
 \end{aligned}$$

Replacing q^ℓ by q yields the generating function for $\overline{R}_{\ell,\mu}(n)$, that is, (2).

$$\begin{aligned}
 \sum_{n=0}^{\infty} B_{\ell,\mu}(2n) q^{2n} &= \prod_{n=1}^{\infty} \frac{(1 - q^{2\ell n})(1 - q^{2\mu n})(1 - q^{2\ell(2n - 1)})(1 - q^{2\mu(2n - 1)})}{(1 - q^{2n})(1 - q^{2(2n - 1)})(1 - q^{2\ell\mu n})(1 - q^{2\mu(2n - 1)})} \\
 &= \prod_{n=1}^{\infty} \frac{(1 + q^{2n})(1 - q^{2\ell n})(1 - q^{2\mu n})(1 + q^{2\ell\mu n})}{(1 - q^{2n})(1 + q^{2\ell n})(1 + q^{2\mu n})(1 - q^{2\ell\mu n})}.
 \end{aligned}$$

Then replacing q^2 by q yields (2).

$$\begin{aligned}
\sum_{n=0}^{\infty} C_{\ell,\mu}(2n)q^{2n} &= \prod_{n=1}^{\infty} \frac{(1+q^{2n}+\dots+q^{(\ell-1)2n})(1+q^{2(2n-1)}+\dots+q^{2(\ell-1)(2n-1)})}{(1+q^{2\mu n}+\dots+q^{(\ell-1)2\mu n})(1+q^{2\mu(2n-1)}+\dots+q^{2(\ell-1)\mu(2n-1)})} \\
&= \prod_{n=1}^{\infty} \frac{(1-q^{2\ell n})(1-q^{2\ell(2n-1)})(1-q^{2\mu n})(1-q^{2\mu(2n-1)})}{(1-q^{2n})(1-q^{2(2n-1)})(1-q^{2\ell\mu n})(1-q^{2\ell\mu(2n-1)})} \\
(4) \qquad &= \sum_{n=0}^{\infty} B_{\ell,\mu}(2n)q^{2n}.
\end{aligned}$$

The generating function of 2μ -regular partitions of n in which odd parts occur with even multiplicities and each part $\equiv \mu \pmod{2\mu}$ appears at most once is

$$\prod_{n=1}^{\infty} \frac{1-q^{2\mu n}}{(1-q^{2n})(1+q^{2\mu n})(1-q^{2(2n-1)})}.$$

Thus, the generating function for $D_{\ell,\mu}(2n)$ is

$$\begin{aligned}
\sum_{n=0}^{\infty} D_{\ell,\mu}(2n)q^{2n} &= \prod_{n=1}^{\infty} \frac{1-q^{2\mu n}}{(1-q^{2n})(1+q^{2\mu n})(1-q^{2(2n-1)})} \\
&\quad \times \frac{(1-q^{2\ell n})(1+q^{2\ell\mu n})(1-q^{2\ell(2n-1)})}{1-q^{2\ell\mu n}} \\
&= \sum_{n=0}^{\infty} B_{\ell,\mu}(2n)q^{2n}.
\end{aligned}$$

Next, we have

$$\begin{aligned}
\sum_{n=0}^{\infty} E_{\ell,\mu}(2n)q^{2n} &= \prod_{n=1}^{\infty} \frac{(1-q^{2\ell n})(1-q^{2\ell(2n-1)})}{(1-q^{2n})(1-q^{2(2n-1)})} \times \frac{(1-q^{2\mu n})}{(1-q^{2\ell\mu n})} \times \frac{1+q^{2\ell\mu n}}{1+q^{2\mu n}} \\
&= \prod_{n=1}^{\infty} \frac{(1+q^{2n})(1-q^{2\ell n})(1-q^{2\mu n})(1+q^{2\ell\mu n})}{(1-q^{2n})(1+q^{2\ell n})(1+q^{2\mu n})(1-q^{2\ell\mu n})}.
\end{aligned}$$

Then replacing q^2 by q gives (2).

Lastly,

$$\begin{aligned}
\sum_{n=0}^{\infty} F_{\ell,\mu}(\ell n)q^{\ell n} &= \prod_{n=1}^{\infty} \frac{(1+q^{\ell(\ell n-1)})\dots(1+q^{\ell(\ell n-(\ell-1))})(1-q^{\ell^2 n})(1-q^{\mu\ell n})}{(1+q^{\mu\ell(\ell n-1)})\dots(1+q^{\mu\ell(\ell n-(\ell-1))})(1-q^{\ell n})(1-q^{\mu\ell^2 n})} \\
&= \prod_{n=1}^{\infty} \frac{(1+q^{\ell n})(1+q^{\mu\ell^2 n})(1-q^{\ell^2 n})(1-q^{\mu\ell n})}{(1+q^{\mu\ell n})(1+q^{\ell^2 n})(1-q^{\ell n})(1-q^{\mu\ell^2 n})}.
\end{aligned}$$

Then replacing q^ℓ by q yields (2) as desired. This completes the proof. \blacksquare

3. CONGRUENCE PROPERTIES VIA ANALYTIC & COMBINATORIAL TECHNIQUES

Before stating and proving our results, we state the following lemmas, the proofs of which may be found in [9]. We recall the definition of Ramanujan's theta function

$$\varphi(q) := \sum_{n=-\infty}^{\infty} q^{n^2} = \prod_{n=1}^{\infty} (1+q^{2n-1})^2(1-q^{2n}).$$

Lemma 3.1. *We have*

$$\varphi(-q^2)^2 = \varphi(q)\varphi(-q).$$

Lemma 3.2. *We have*

$$\frac{1}{\varphi(-q)} = \varphi(q)\varphi(q^2)^2\varphi(q^4)^4\dots$$

We can rewrite the generating function of $\overline{R_{\ell,\mu}}(n)$ in terms of Ramanujan's theta function $\varphi(q)$ as follows

$$(5) \quad \sum_{n \geq 0} \overline{R_{\ell,\mu}}(n)q^n = \frac{\varphi(-q^\ell)\varphi(-q^\mu)}{\varphi(-q)\varphi(-q^{\ell\mu})}.$$

Incorporating the results of Lemmas 3.1 and 3.2, we can re-write

$$(6) \quad \sum_{n \geq 0} \overline{R_{\ell,\mu}}(n)q^n = \frac{\varphi(q)\varphi(q^2)^2\varphi(q^4)^4 \cdots \varphi(q^{\ell\mu})\varphi(q^{2\ell\mu})^2\varphi(q^{4\ell\mu})^4 \cdots}{\varphi(q^\ell)\varphi(q^{2\ell})^2\varphi(q^{4\ell})^4 \cdots \varphi(q^\mu)\varphi(q^{2\mu})^2\varphi(q^{4\mu})^4 \cdots}.$$

The following easy corollary now follows.

Corollary 3.3. *For all $n \geq 1$, we have $\overline{R_{\ell,\mu}}(n) \equiv 0 \pmod{2}$.*

Proof. Since $\varphi(q) = 1 + 2 \sum_{n \geq 1} q^{n^2}$, we know that $\varphi(q) \equiv 1 \pmod{2}$. So (6) gives us

$$\sum_{n \geq 0} \overline{R_{\ell,\mu}}(n)q^n \equiv 1 \pmod{2}.$$

This gives an analytic proof. ■

We give a combinatorial proof as well.

Combinatorial Proof of Corollary 3.3. We can obtain an overpartition by overlining the first occurrence of any distinct part of an ordinary partition $\lambda = (c_1^{u_1}, c_2^{u_2}, \dots, c_r^{u_r})$ with $c_1 > c_2 > \cdots > c_r$. Since we may choose to either overline a part or not, the number of overpartitions obtainable from λ is $\overline{p}(\lambda) := 2^r$. Thus for all $n \geq 1$, $\overline{p}(n) \equiv 0 \pmod{2}$. Analogously, if we consider only partitions λ that are simultaneously ℓ - and μ -regular, we would obtain an even number of (ℓ, μ) -regular overpartitions. Thus For all $n \geq 1$, we have $\overline{R_{\ell,\mu}}(n) \equiv 0 \pmod{2}$. ■

We now give a complete modulo 4 characterization of $\overline{R_{\ell,\mu}}(n)$.

Theorem 3.4. *For all $n \geq 1$,*

(i) *if ℓ is a square and μ is not, then*

$$\overline{R_{\ell,\mu}}(n) \equiv \begin{cases} 2 \pmod{4} & \text{if } n = k^2 \text{ or } n = \mu k^2, \text{ where } \ell \text{ and } k \text{ are relatively prime;} \\ 0 \pmod{4} & \text{otherwise,} \end{cases}$$

(ii) *if ℓ and μ are both squares, then*

$$\overline{R_{\ell,\mu}}(n) \equiv \begin{cases} 2 \pmod{4} & \text{if } n = k^2, \text{ where } k, \mu \text{ and } \ell \text{ are relatively prime;} \\ 0 \pmod{4} & \text{otherwise,} \end{cases}$$

(iii) *if neither ℓ nor μ is a square, then*

$$\overline{R_{\ell,\mu}}(n) \equiv \begin{cases} 2 \pmod{4} & \text{if } n = k^2, n = \ell k^2, n = \mu k^2 \text{ or } n = \ell \mu k^2; \\ 0 \pmod{4} & \text{otherwise.} \end{cases}$$

Analytic Proof of Theorem 3.4. Using (6) we have

$$\sum_{n \geq 0} \overline{R_{\ell,\mu}}(n)q^n \equiv \frac{\varphi(q)\varphi(q^{\ell\mu})}{\varphi(q^\ell)\varphi(q^\mu)} \pmod{4},$$

since $\varphi(q^i)^j \equiv 1 \pmod{4}$ for any $j \geq 2$. Next, we deduce from Lemma 3.1, that

$$\varphi(q) = \frac{\varphi(-q^2)^2}{\varphi(-q)}.$$

Thus

$$\begin{aligned} \overline{R_{\ell,\mu}}(n)q^n &\equiv \frac{\varphi(q)\varphi(q^{\ell\mu})}{\varphi(q^\ell)\varphi(q^\mu)} \equiv \frac{\varphi(q)\varphi(q^\ell\mu)\varphi(-q^\ell)\varphi(-q^\mu)}{\varphi(-q^{2\ell})^2\varphi(-q^{2\mu})^2} \pmod{4} \\ &\equiv \varphi(q)\varphi(q^\ell\mu)\varphi(-q^\ell)\varphi(-q^\mu) \pmod{4}, \end{aligned}$$

since $\varphi(-q^{2\ell})^2 \equiv 1 \pmod{4}$.

Hence,

$$\begin{aligned} \sum_{n \geq 0} \overline{R_{\ell, \mu}}(n)q^n &\equiv \varphi(q)\varphi(q^\ell \mu)\varphi(-q^\ell)\varphi(-q^\mu) \pmod{4} \\ &= (1 + 2 \sum_{n \geq 1} q^{n^2})(1 + 2 \sum_{n \geq 1} (q^{\ell\mu})^{n^2})(1 + 2 \sum_{n \geq 1} (-q^\ell)^{n^2})(1 + 2 \sum_{n \geq 1} (-q^\mu)^{n^2}) \\ &\equiv 1 + 2 \sum_{n \geq 1} q^{n^2} + 2 \sum_{n \geq 1} (q^{\ell\mu})^{n^2} + 2 \sum_{n \geq 1} (-q^\ell)^{n^2} + 2 \sum_{n \geq 1} (-q^\mu)^{n^2} \pmod{4} \\ &\equiv 1 + 2 \sum_{n \geq 1} q^{n^2} + 2 \sum_{n \geq 1} q^{\ell\mu n^2} + 2 \sum_{n \geq 1} q^{\ell n^2} + 2 \sum_{n \geq 1} q^{\mu n^2} \pmod{4}. \end{aligned}$$

A straightforward interpretation of the last congruence gives the three results as required. \blacksquare

We give a combinatorial proof as well.

Combinatorial Proof of Theorem 3.4. Let $m(\ell|n)$ be the number of multiples ℓ dividing n and define $\Delta(n, \ell, \mu) := \tau(n) - (m(\ell|n) + m(\mu|n) - m(\ell\mu|n))$, where $\tau(n)$ is the number of divisors of n . We claim that

$$(7) \quad \overline{R_{\ell, \mu}}(n) \equiv \begin{cases} 2 \pmod{4} & \text{if } \Delta(n, \ell, \mu) \text{ is odd;} \\ 0 \pmod{4} & \text{otherwise.} \end{cases}$$

We decompose (ℓ, μ) -regular overpartitions into two classes: those containing a unique part-size and those containing two or more different part-sizes. By the proof of Corollary 3.3 the latter class has cardinality of the form $m2^r, m > 0, r > 1$ which is divisible by 4. However, partitions with a single part-size arise from divisors of n . Each divisor d of n (excluding ℓ, μ and $\ell\mu$) gives the ordinary partition $(d^{n/d})$ which in turn produces two (ℓ, μ) -regular overpartitions. Thus $\Delta(n, \ell, \mu) \equiv 1 \pmod{2}$ if and only if divisors of n contribute an odd number of pairs of (ℓ, μ) -regular overpartitions. Hence (7) follows.

We will use the easily proved relation $m(\ell|n) = \tau(n/\ell)$.

We consider part (i). In view (7) it will suffice to find the parity of $\Delta(n, \ell, \mu)$ under each constraint. If $n = k^2$ and $\ell \nmid n$, then $\tau(n)$ is odd and $m(\ell|n) = 0$. Since both $m(\mu|n)$ and $m(\ell\mu|n)$ are even, it follows that $\Delta(n, \ell, \mu)$ is odd. Similarly, the case $\mu \nmid n$ implies that $\Delta(n, \ell, \mu)$ is odd. If $n = k^2$ and $\mu | n$, then $m(\mu|n)$ is even (since μ is not a square). Hence $\Delta(n, \ell, \mu)$ is odd. But if $n = \mu k^2$, then $\tau(n)$ is even and $m(\mu|n) = \tau(k^2)$ which is odd. So $\Delta(n, \ell, \mu)$ is odd. The proof of the first line of part (i) is complete.

For the second line we consider the the following negations, given that ℓ is a square and μ is not: (a) $n = k^2$ and $\ell | n$, and (b) $n \neq k^2$ and $n \neq \mu k^2$. In (a) we find that both $\tau(n)$ and $m(\ell|n)$ are odd. Since $m(\mu|n)$ and $m(\ell\mu|n)$ are clearly even, it follows that $\Delta(n, \ell, \mu)$ is even. In (b) it is clear that all the relevant functions are even. This completes the proof of part (i).

The other parts may be proved in a similar manner. For example, the first line of part (iii) may be established by noting that exactly one member of the set $\{\tau(n), m(\ell|n), m(\mu|n), m(\ell\mu|n)\}$ is odd at a time with all the others being even. \blacksquare

We close this section by proving a general mod 8 congruence for the $\overline{R_\ell^*}(n)$ function. We will need the following lemma.

Lemma 3.5. *We have*

$$\sum_{n \geq 0} \overline{R_6^*}(3n + 2)q^n \equiv 4f_6^3 \pmod{4}.$$

Proof. We need the following identity [6, Theorem 1]

$$(8) \quad \frac{f_2}{f_1^2} = \frac{f_6^4 f_9^6}{f_3^8 f_{18}^3} + 2q \frac{f_6^3 f_9^3}{f_3^7} + 4q^2 \frac{f_6^2 f_{18}^3}{f_3^6}.$$

Using (8) in (1) we have the following

$$\sum_{n \geq 0} \overline{R}_6^*(3n+2)q^n = 4 \frac{f_2^3 f_6^3}{f_1^6} \equiv 4 \frac{f_1^6 f_6^3}{f_1^6} \equiv 4f_6^3 \pmod{4}.$$

■

Theorem 3.6. *Let $p \geq 5$ be a prime and let r with $1 \leq r \leq p-1$ be such that $\text{inv}(3, p) \cdot 4 \cdot r + 1$ is a quadratic nonresidue modulo p , where $\text{inv}(3, p)$ is the inverse of 3 modulo p . Then, for all $n \geq 0$ we have $\overline{R}_6^*(3(pn+r)+2) \equiv 0 \pmod{8}$.*

Proof. We need Jacobi's triple product identity

$$(9) \quad f_1^3 = \sum_{j=0}^{\infty} (-1)^j (2j+1) q^{j(j+1)/2}.$$

From Lemma 3.5 and (9) we have

$$\sum_{n \geq 0} \overline{R}_6^*(3n+2)q^n \equiv 4 \sum_{j=0}^{\infty} (-1)^j (2j+1) q^{3j(j+1)} \pmod{8}.$$

We are interested in values of the form $\overline{R}_6^*(3(pn+r)+2)$, and we want to know whether we can have $pn+r = 3j(j+1)$, for some nonnegative integer j . If such a representation for $pn+r$ exists, then $r \equiv 3j(j+1) \pmod{p}$. Since $p \geq 3$, so this is equivalent to $\text{inv}(3, p) \cdot r \equiv j(j+1) \pmod{p}$, which in turn is equivalent to $\text{inv}(3, p) \cdot 4 \cdot r + 1 \equiv (2j+1)^2 \pmod{p}$. From our assumption that $\text{inv}(3, p) \cdot 4 \cdot r + 1$ is a quadratic nonresidue modulo p , the result now follows. ■

4. CONGRUENCE PROPERTIES VIA MODULAR FORMS

In this section, we will use the theory of modular forms to find several congruences. The first result is given below.

Theorem 4.1. *For all $n \geq 0$, we have*

$$\begin{aligned}
(10) \quad & \overline{R_{2,3}}(9n+6) \equiv 0 \pmod{6}, \\
(11) \quad & \overline{R_{4,3}}(6n+3) \equiv 0 \pmod{6}, \\
(12) \quad & \overline{R_{4,3}}(6n+5) \equiv 0 \pmod{12}, \\
(13) \quad & \overline{R_{4,3}}(9n+3) \equiv 0 \pmod{6}, \\
(14) \quad & \overline{R_{4,3}}(12n+7) \equiv 0 \pmod{24}, \\
(15) \quad & \overline{R_{4,3}}(12n+11) \equiv 0 \pmod{72}, \\
(16) \quad & \overline{R_{4,9}}(8n+4) \equiv 0 \pmod{12}, \\
(17) \quad & \overline{R_{4,9}}(12n+4) \equiv 0 \pmod{12}, \\
(18) \quad & \overline{R_{4,9}}(12n+8) \equiv 0 \pmod{72}, \\
(19) \quad & \overline{R_{4,9}}(16n+8) \equiv 0 \pmod{24}, \\
(20) \quad & \overline{R_{4,9}}(18n+12) \equiv 0 \pmod{96}, \\
(21) \quad & \overline{R_{4,9}}(24n+20) \equiv 0 \pmod{216}, \\
(22) \quad & \overline{R_{4,9}}(18n+15) \equiv 0 \pmod{48}, \\
(23) \quad & \overline{R_{4,9}}(96n+80) \equiv 0 \pmod{864}, \\
(24) \quad & \overline{R_{8,27}}(36n+15) \equiv 0 \pmod{24}, \\
(25) \quad & \overline{R_{8,27}}(36n+21) \equiv 0 \pmod{96}, \\
(26) \quad & \overline{R_{8,27}}(36n+24) \equiv 0 \pmod{12}, \\
(27) \quad & \overline{R_{8,27}}(36n+27) \equiv 0 \pmod{6}, \\
(28) \quad & \overline{R_{16,81}}(36n+33) \equiv 0 \pmod{48}, \\
(29) \quad & \overline{R_{16,81}}(72n+60) \equiv 0 \pmod{48}.
\end{aligned}$$

Theorem 4.2. *For all $n \geq 0$, we have*

$$\begin{aligned}
(30) \quad & \overline{R_3^*}(9n+4) \equiv 0 \pmod{12}, \\
(31) \quad & \overline{R_3^*}(9n+7) \equiv 0 \pmod{48}, \\
(32) \quad & \overline{R_6^*}(9n+5) \equiv 0 \pmod{24}, \\
(33) \quad & \overline{R_6^*}(9n+8) \equiv 0 \pmod{96}.
\end{aligned}$$

Remark 4.3. *Alanazi et. al. [1] had obtained*

$$\overline{R_3^*}(9n+4) \equiv \overline{R_3^*}(9n+7) \equiv 0 \pmod{3}.$$

While Sellers [16] had very recently obtained

$$\overline{R_3^*}(9n+4) \equiv \overline{R_3^*}(9n+7) \equiv 0 \pmod{4}.$$

Our proof of the first congruence in Theorem 4.2 is independent of the techniques used in the proofs of the results of Alanazi et. al. and Sellers.

We can also get congruences for finer arithmetic progressions, we list only two of them without proof here.

Theorem 4.4. *For all $n \geq 0$, we have*

$$\begin{aligned}
(34) \quad & \overline{R_6^*}(27n+11) \equiv 0 \pmod{64}, \\
(35) \quad & \overline{R_6^*}(81n+47) \equiv 0 \pmod{24}.
\end{aligned}$$

Theorems 4.1 and 4.2 are proved in Section 4.4 using an algorithmic approach, due to Smoot [18].

The following two congruences will be proved using a manual implementation of an algorithm of Radu [13, 12], due to computational difficulties.

Theorem 4.5. *For all $n \geq 0$, we have*

$$(36) \quad \overline{R_{3,5}}(9n + 3) \equiv 0 \pmod{6},$$

$$(37) \quad \overline{R_{2,5}}(18n + 9) \equiv 0 \pmod{6}.$$

We prove Theorem 4.5 in Section 4.5.

4.1. Preliminaries on the Algorithmic Approach. In this section we describe our methods for proving Theorems 4.1, 4.2 and 4.5. Theorems 4.1 and 4.2 are proved using a Mathematica implementation of an algorithm of Radu [13, 12] due to Smoot [18], while Theorem 4.5 is proved using Radu's algorithm manually. We first describe Radu's algorithm in Subsection 4.2 and then Smoot's implementation in Subsection 4.3.

4.2. Radu's Algorithm. For a positive integer N , we define the following matrix groups:

$$\Gamma := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : a, b, c, d \in \mathbb{Z}, ad - bc = 1 \right\},$$

$$\Gamma_\infty := \left\{ \begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix} \in \Gamma : n \in \mathbb{Z} \right\}.$$

$$\Gamma_0(N) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma : c \equiv 0 \pmod{N} \right\},$$

and

$$[\Gamma : \Gamma_0(N)] := N \prod_{\ell|N} \left(1 + \frac{1}{\ell}\right),$$

where ℓ is a prime.

We need some preliminary results, which describe an algorithmic approach to proving partition concurrences, developed by Radu [12, 13]. For integers $M \geq 1$, suppose that $R(M)$ is the set of all the integer sequences

$$(r_\delta) := (r_{\delta_1}, r_{\delta_2}, r_{\delta_3}, \dots, r_{\delta_k})$$

indexed by all the positive divisors δ of M , where $1 = \delta_1 < \delta_2 < \dots < \delta_k = M$. For integers $m \geq 1$, $(r_\delta) \in R(M)$, and $t \in \{0, 1, 2, \dots, m-1\}$, we define the set $P(t)$ as

$$P(t) := \left\{ t' \in \{0, 1, 2, \dots, m-1\} : t' \equiv ts + \frac{s-1}{24} \sum_{\delta|M} \delta r_\delta \pmod{m} \right.$$

$$(38) \quad \left. \text{for some } [s]_{24m} \in \mathbb{S}_{24m} \right\},$$

where $[x]_m$ denote the residue class of x , \mathbb{Z}_m^* denote the set of the invertible elements of \mathbb{Z}_m , and \mathbb{S}_m denote the set of the squares of \mathbb{Z}_m^* .

For integers $N \geq 1$, $\gamma := \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$, $(r_\delta) \in R(M)$, and $(r'_\delta) \in R(N)$, we also define

$$p(\gamma) := \min_{\lambda \in \{0, 1, \dots, m-1\}} \frac{1}{24} \sum_{\delta|M} r_\delta \frac{\gcd(\delta(a + k\lambda c), mc)^2}{\delta m},$$

$$p'(\gamma) := \frac{1}{24} \sum_{\delta|N} r'_\delta \frac{\gcd(\delta, c)^2}{\delta}.$$

For integers $m \geq 1$; $M \geq 1$, $N \geq 1$, $t \in \{0, 1, 2, \dots, m-1\}$, $k := \gcd(m^2 - 1, 24)$, and $(r_\delta) \in R(M)$, define Δ^* to be the set of all tuples $(m, M, N, t, (r_\delta))$ such that all of the following conditions are satisfied

1. Prime divisors of m are also prime divisors of N ;
2. If $\delta | M$, then $\delta | mN$ for all $\delta \geq 1$ with $r_\delta \neq 0$;
3. $24 | kN \sum_{\delta|M} \frac{r_\delta m N}{\delta}$;

$$4. 8 \mid kN \sum_{\delta \mid M} r_\delta;$$

$$5. \frac{24m}{\left(-24kt - k \sum_{\delta \mid M} \delta r_\delta, 24m\right)} \mid N;$$

6. If $2 \mid m$ then either $4 \mid kN$ and $8 \mid \delta N$ or $2 \mid s$ and $8 \mid (1-j)N$, where $\prod_{\delta \mid M} \delta^{|r_\delta|} = 2^s \cdot j$.

We now state a result of Radu [13], which we use in completing the proof of Theorem 4.1.

Lemma 4.6. [13, Lemma 4.5] *Suppose that $(m, M, N, t, (r_\delta)) \in \Delta^*$, $(r'_\delta) := (r'_\delta)_{\delta \mid N} \in R(N)$, $\{\gamma_1, \gamma_2, \dots, \gamma_n\} \subseteq \Gamma$ is a complete set of representatives of the double cosets of $\Gamma_0(N) \backslash \Gamma / \Gamma_\infty$, and $t_{\min} := \min_{t' \in P(t)} t'$,*

$$(39) \quad \nu := \frac{1}{24} \left(\left(\sum_{\delta \mid M} r_\delta + \sum_{\delta \mid N} r'_\delta \right) [\Gamma : \Gamma_0(N)] - \sum_{\delta \mid N} \delta r'_\delta - \frac{1}{m} \sum_{\delta \mid M} \delta r_\delta \right) - \frac{t_{\min}}{m},$$

$p(\gamma_j) + p'(\gamma_j) \geq 0$ for all $1 \leq j \leq n$, and $\sum_{n=0}^{\infty} A(n)q^n := \prod_{\delta \mid M} f_\delta^{r_\delta}$. If for some integers $u \geq 1$, all $t' \in P(t)$,

and $0 \leq n \leq \lfloor \nu \rfloor$, $A(mn + t') \equiv 0 \pmod{u}$ is true, then for integers $n \geq 0$ and all $t' \in P(t)$, we have $A(mn + t') \equiv 0 \pmod{u}$.

The following lemma supports Lemma 4.6 in the proof of Theorem 4.1.

Lemma 4.7. [13, Lemma 2.6] *Let N or $\frac{N}{2}$ be a square-free integer, then we have*

$$\bigcup_{\delta \mid N} \Gamma_0(N) \begin{pmatrix} 1 & 0 \\ \delta & 1 \end{pmatrix} \Gamma_\infty = \Gamma.$$

4.3. Smoot's Implementation of Radu's Algorithm. We will also use Smoot's [18] implementation of Radu's algorithm [13, 12], which can be used to prove Ramanujan type congruences. The algorithm takes as an input the generating function

$$\sum_{n=0}^{\infty} a_r(n)q^n = \prod_{\delta \mid M} \prod_{n=1}^{\infty} (1 - q^{\delta n})^{r_\delta},$$

and positive integers m and N , with M another positive integer and $(r_\delta)_{\delta \mid M}$ is a sequence indexed by the positive divisors δ of M . With this input, Radu's algorithm tries to produce a set $P_{m,j}(j) \subseteq \{0, 1, \dots, m-1\}$ which contains j and is uniquely defined by $m, (r_\delta)_{\delta \mid M}$ and j . Then, it decides if there exists a sequence $(s_\delta)_{\delta \mid N}$ such that

$$q^\alpha \prod_{\delta \mid M} \prod_{n=1}^{\infty} (1 - q^{\delta n})^{s_\delta} \cdot \prod_{j' \in P_{m,j}(j)} \sum_{n=0}^{\infty} a(mn + j')q^n,$$

is a modular function with certain restrictions on its behaviour on the boundary of \mathbb{H} .

Smoot [18] implemented this algorithm in Mathematica and we use his `RaduRK` package, which requires the software package `4ti2`. Documentation on how to install and use these packages are available from Smoot [18]. We use this implemented `RaduRK` algorithm to prove Theorem 4.1 in the next section.

It is natural to guess that $N = m$ (which corresponds to the congruence subgroup $\Gamma_0(N)$), but this is not always the case, although they are usually closely related to one another. The determination of the correct value of N is an important problem for the usage of `RaduRK` and it depends on the Δ^* criterion described in the previous subsection. It is easy to check the minimum N which satisfies this criterion by running `minN[M, r, m, j]`.

4.4. Proofs of Theorems 4.1 and 4.2. In this section we prove Theorems 4.1 and 4.2 using Smoot's implementation described in Subsection 4.3.

Proof of Theorem 4.1. Since the proof of all congruences listed in Theorem 4.1 are similar, we only prove (10) here. The rest of the output can be obtained by visiting <https://manjilsaikia.in/publ/mathematica/smoot>. We use the procedure call

$$\text{RK}[12, 12, \{-2, 3, 2, -1, -3, 1\}, 9, 6]$$

which gives a straight proof of (10). Here we give the output of RK.

$$\text{In}[1] := \text{RaduRK}[12, 12, \{-2, 3, 2, -1, -3, 1\}, 9, 6]$$

$$\prod_{\delta|M} (q^\delta; q^\delta)_\infty^{r_\delta} = \sum_{n=0}^{\infty} a(n) q^n$$

$$\boxed{f_1(q) \cdot \prod_{j' \in P_{m,r}(j)} \sum_{n=0}^{\infty} a(mn + j') q^n = \sum_{g \in AB} g \cdot p_g(t)}$$

Modular Curve: $X_0(N)$

Out [2] =

N:	12
$\{M, (r_\delta)_{\delta M}\}$:	$\{12, \{-2, 3, 2, -1, -3, 1\}\}$
m:	9
$P_{m,r}(j)$:	$\{6\}$
$f_1(q)$:	$\frac{(q; q)_\infty^5 (q^4; q^4)_\infty^2 (q^6; q^6)_\infty^9}{q^3 (q^2; q^2)_\infty (q^3; q^3)_\infty^3 (q^{12}; q^{12})_\infty^{12}}$
t:	$\frac{(q^4; q^4)_\infty^4 (q^6; q^6)_\infty^2}{q (q^2; q^2)_\infty^2 (q^{12}; q^{12})_\infty^4}$
AB:	$\{1\}$
$\{p_g(t) : g \in AB\}$	$\{6t^3 + 6t^2 - 6t - 6\}$
Common Factor:	6

The interpretation of this output is as follows.

The first entry in the procedure call $\text{RK}[12, 12, \{-2, 3, 2, -1, -3, 1\}, 9, 6]$ corresponds to specifying $N = 12$, which fixes the space of modular functions

$$M(\Gamma_0(N)) := \text{the algebra of modular functions for } \Gamma_0(N).$$

The second and third entry of the procedure call $\text{RK}[12, 12, \{-2, 3, 2, -1, -3, 1\}, 9, 6]$ gives the assignment $\{M, (r_\delta)_{\delta|M}\} = \{12, (-2, 3, 2, -1, -3, 1)\}$, which corresponds to specifying $(r_\delta)_{\delta|M} = (r_1, r_2, r_4) = (-2, 3, 2, -1, -3, 1)$, so that

$$\sum_{n \geq 0} \bar{R}_{2,3}(n) q^n = \prod_{\delta|M} (q^\delta; q^\delta)_\infty^{r_\delta} = \frac{f_2^3 f_3^2 f_{12}}{f_1^2 f_4 f_6^3}.$$

The last two entries of the procedure call $\text{RK}[12, 12, \{-2, 3, 2, -1, -3, 1\}, 9, 6]$ corresponds to the assignment $m = 8$ and $j = 7$, which means that we want the generating function

$$\sum_{n \geq 0} \bar{R}_{2,3}(n)(mn + j)q^n = \sum_{n \geq 0} \bar{R}_{2,3}(n)(9n + 6)q^n.$$

So, $P_{m,r}(j) = P_{9,r}(6)$ with $r = (-2, 3, 2, -1, -3, 1)$.

The output $P_{m,r}(j) := P_{9,(-2,3,2,-1,-3,1)}(6) = \{6\}$ means that there exists an infinite product

$$f_1(q) = \frac{(q; q)_\infty^5 (q^4; q^4)_\infty^2 (q^6; q^6)_\infty^9}{q^3 (q^2; q^2)_\infty (q^3; q^3)_\infty^3 (q^{12}; q^{12})_\infty^{12}},$$

such that

$$f_1(q) \sum_{n \geq 0} \overline{R}_{2,3}(n)(9n+6)q^n \in M(\Gamma_0(12)).$$

Finally, the output

$$t = \frac{(q^4; q^4)_\infty^4 (q^6; q^6)_\infty^2}{q (q^2; q^2)_\infty^2 (q^{12}; q^{12})_\infty^4}, \quad AB = \{1\}, \quad \text{and} \quad \{p_g(t): g \in AB\},$$

presents a solution to the question of finding a modular function $t \in M(\Gamma_0(12))$ and polynomials $p_g(t)$ such that

$$f_1(q) \sum_{n \geq 0} \overline{R}_{2,3}(n)(9n+6)q^n = \sum_{g \in AB} p_g(t) \cdot g$$

In this specific case, we see that the singleton entry in the set $\{p_g(t): g \in AB\}$ has the common factor 6, thus proving equation (10). \blacksquare

Remark 4.8. *The interested reader can refer to [11] or [4] for some more recent applications of the method.*

Proof of Theorem 4.2. Since the proof of Theorem 4.2 is similar to the proof of Theorem 4.1, we just refer the reader to the output file available at https://manjilsaikia.in/publ/mathematica/smoot_rl.nb \blacksquare

4.5. Proof of Theorem 4.5. We use the material in Subsection 4.2 without commentary. We give the full details for the proof of (36), and mention the parameters in Radu's algorithm for (37).

For the purposes of (36), it is enough to take

$$(m, M, N, t, (r_\delta)) = (9, 30, 30, 3, (-2, 1, 2, 2, -1, -1, -2, 1)).$$

It is routine to check that this choice satisfies the Δ^* conditions. By equation (38) we see that $P(t) = \{3\}$. For the choice of $(r'_\delta) = (12, 0, 0, 0, 0, 0, 0, 0, 0)$ we see that

$$p \left(\begin{pmatrix} 1 & 0 \\ \delta & 1 \end{pmatrix} \right) + p' \left(\begin{pmatrix} 1 & 0 \\ \delta & 1 \end{pmatrix} \right) \geq 0 \quad \text{for all } \delta \mid N,$$

and $\lfloor \nu \rfloor = 101$ for $t = 3$. So we need to just check the validity of (36) for all $n \leq \lfloor \nu \rfloor$ and then by Lemmas 4.6 and 4.7 we would have proved our result. This can be checked using Mathematica and hence the result follows.

For (37), we take

$$(m, M, N, t, (r_\delta)) = (18, 20, 60, 9, (-2, 3, -1, 2, -3, 1)),$$

which will give us

$$P(t) = \{9\}, \quad (r'_\delta) = (54, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0) \quad \text{and} \quad \lfloor \nu \rfloor = 1322.$$

We can then verify the congruence up to $n = 1322$ via Mathematica from which the result follows. \blacksquare

5. CONCLUDING REMARKS

- (1) It is possible to give a combinatorial proof of Theorem 2.1.
- (2) A modulo 8 characterization in the same vein as the modulo 4 characterization in Section 3 is also possible. We leave it as an open problem.
- (3) It is desirable to have elementary proofs of the results in Section 4.
- (4) A further study of congruences is also desired. To that end, we make the following conjecture.

Conjecture 5.1. *For all $n \geq 0$, $\ell \geq 2$, and $1 \leq k \leq \ell$, we have*

$$\overline{R}_{4,9}(4\ell n + 4k) \equiv 0 \pmod{6}.$$

REFERENCES

- [1] Abdulaziz Alanazi, Bashair Alenazi, William Keith, and Augustine Munagi. Refining overpartitions by properties of non-overlined parts. *Contrib. Discrete Math.*, 17(2):96–111, 2022.
- [2] Abdulaziz M. Alanazi and Augustine O. Munagi, Combinatorial identities for ℓ -regular overpartitions, *Ars Combin.* **CXXX** (2017), 55–66.
- [3] Abdulaziz M. Alanazi, Augustine O. Munagi and James A. Sellers, An infinite family of congruences for ℓ -regular overpartitions, *Integers* **16** (2016) 8pp, #A37.
- [4] George E. Andrews and Peter Paule. MacMahon’s partition analysis. XIV: Partitions with n copies of n . *J. Comb. Theory, Ser. A*, 203:34, 2024. Id/No 105836.
- [5] Bruce C. Berndt, *Number Theory in the Spirit of Ramanujan*, American Mathematical Society, Providence (2006).
- [6] Michael D. Hirschhorn and James A. Sellers. Arithmetic relations for overpartitions. *J. Comb. Math. Comb. Comput.*, 53:65–73, 2005.
- [7] Michael D. Hirschhorn and James A. Sellers, Arithmetic properties of overpartitions into odd parts, *Ann. Comb.* **10** (2006), 353–367.
- [8] William J. Keith, Partitions into parts simultaneously regular, distinct and/or flat, *Proceedings of CANT 2016*, https://link.springer.com/chapter/10.1007/978-3-319-68032-3_10
- [9] Augustine O. Munagi and James A. Sellers. Refining overlined parts in overpartitions via residue classes: bijections, generating functions, and congruences, *Util. Math.* **95** (2014), pp. 33–49
- [10] M. S. M. Naika and C. Shivashankar, Arithmetic properties of ℓ -regular overpartition pairs, *Turk J Math*, 41 (2017), 756–774.
- [11] Manjil P. Saikia. Some missed congruences modulo powers of 2 for t -colored overpartitions. *Bol. Soc. Mat. Mex.*, III. Ser., 29(1):10, 2023. Id/No 15.
- [12] Cristian-Silviu Radu. An algorithmic approach to Ramanujan-Kolberg identities. *J. Symbolic Comput.*, 68(1):225–253, 2015.
- [13] Silviu Radu and James A. Sellers. Congruence properties modulo 5 and 7 for the pod function. *Int. J. Number Theory*, 7(8):2249–2259, 2011.
- [14] D. Ranganatha, On some new congruences for ℓ -regular overpartitions, *Palestine Journal of Mathematics*, **7** (2018), 345–362.
- [15] Nipen Saikia, and Chayanika Boruah, Congruences for ℓ -regular overpartition for $\ell \in \{5, 6, 8\}$, *Indian J Pure Appl Math* **48** (2017), 295–308.
- [16] James A. Sellers. Extending congruences for overpartitions with l -regular non-overlined parts. *Bull. Aust. Math. Soc.*, accepted, 2024.
- [17] E. Y. Y. Shen, Arithmetic Properties of ℓ -regular Overpartitions, *Int. J. Number Theory* **12** (2016), 841–852.
- [18] Nicolas Allen Smoot. On the computation of identities relating partition numbers in arithmetic progressions with eta quotients: an implementation of Radu’s algorithm. *J. Symbolic Comput.*, 104:276–311, 2021.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, UNIVERSITY OF TABUK, P.O.BOX 741, TABUK 71491, SAUDI ARABIA

Email address: `am.alenezi@ut.edu.sa`

SCHOOL OF MATHEMATICS, UNIVERSITY OF THE WITWATERSRAND, JOHANNESBURG, SOUTH AFRICA

Email address: `augustine.munagi@wits.ac.za`

MATHEMATICAL AND PHYSICAL SCIENCES DIVISION, SCHOOL OF ARTS AND SCIENCES, AHMEDABAD UNIVERSITY, AHMEDABAD 380009, GUJARAT, INDIA

Email address: `manjil@saikia.in`