

FACTORIZATION OF SPREAD POLYNOMIALS

JOHANN CIGLER AND HANS-CHRISTIAN HERBIG

ABSTRACT. We present a proof of a conjecture of Goh and Wildberger on the factorization of the spread polynomials. We indicate how the factors can be effectively calculated and exhibit a connection to the factorization of Fibonacci numbers into primitive parts.

CONTENTS


1.	Introduction	1
2.	Some well-known facts about Lucas polynomials	2
3.	Factorization of the polynomials $L_n(x) - 2$	3
4.	Some properties of $Z_n(x)$	4
5.	Proof of Theorem 1.1	5
6.	How to calculate the $\phi_k(x)$	6
6.1.	Odd index	6
6.2.	The case indexed by powers of 2	6
6.3.	General case	7
7.	Final remarks	8
	References	8

1. INTRODUCTION

The sequence of *spread polynomials* $(S_n(x))_{n \geq 1}$ of Norman J. Wildberger is uniquely defined by the requirement

$$(1.1) \quad S_n(\sin^2 \theta) = \sin^2(n\theta).$$

The spread polynomials play a central role in Wildberger's *rational trigonometry* [8]. In order to make plane geometry free of transcendental expressions and square roots he suggests to replace lengths by their squares, which are referred to as *quadrances*, and angles by *spreads*. In the right angled triangle with sides a, b and c




the spread $s \in [0, 1]$ is defined as the ratio $s = b^2/c^2$ of the quadrance of the opposite leg b by the quadrance of the hypotenuse c . The geometric property of the spread polynomials that makes them play a central role in rational trigonometry is the relation

$$(1.2) \quad s_n = S_n(s)$$

for an arrangement of $n + 1$ lines which all meet in a single point and whose spreads of neighboring lines all coincide. Here s_n is the spread between the extremal lines of the arrangement.

2020 *Mathematics Subject Classification*. primary 11B83, secondary 11B39, 97G40.

Key words and phrases. spread polynomials, cyclotomic polynomials, Fibonacci numbers.



The spread polynomials can be written in terms of the Lucas polynomials $L_n(x)$

$$(1.3) \quad S_n(x) = \frac{2 - L_n(2 - 4x)}{4}.$$

Following [3] we consider the closely related *zspread polynomials*

$$(1.4) \quad Z_n(x) = 4S_n\left(\frac{x}{4}\right) = 2 - L_n(2 - x).$$

A conjecture of Wildberger's honours student Shuxiang Goh (see [9]), stated in terms of $Z_n(x)$, claims the following.

Theorem 1.1. *There is a sequence of polynomials $(\Phi_d(x))_{d \geq 1}$, $\Phi_d(x) \in \mathbb{Z}[x]$, such that $Z_n(x) = \prod_{d|n} \Phi_d(x)$ and $\deg(\Phi_d) = \varphi(d)$, where $\varphi(d)$ is Euler's totient function.*

The aim of this note is to prove the theorem as a consequence of (1.4) and known results about the Chebyshev polynomials obtained in [1], [2] and [7]. Moreover, we obtain some concrete facts about the polynomials $\Phi_d(x)$, some of which had already been conjectured in [3].

Acknowledgements. We would like to thank Tri Nguyen for generously sharing the idea [4] with us.

2. SOME WELL-KNOWN FACTS ABOUT LUCAS POLYNOMIALS

The Lucas polynomials $L_n(x)$ are defined by the recursion

$$(2.1) \quad L_n(x) = xL_{n-1}(x) - L_{n-2}(x)$$

for $n \geq 2$ with initial values $L_0(x) = 2$ and $L_1(x) = x$.¹ The first terms are (cf. [6, entry A034807]):

n	0	1	2	3	4	5	6	...
$L_n(x)$	2	x	$-2 + x^2$	$-3x + x^3$	$2 - 4x^2 + x^4$	$5x - 5x^3 + x^5$	$-2 + 9x^2 - 6x^4 + x^6$...

Binet's formula gives

$$(2.2) \quad L_n(x) = (\alpha(x))^n + (\beta(x))^n$$

with $\alpha(x) = \frac{x + \sqrt{x^2 - 4}}{2}$ and $\beta(x) = \frac{1}{\alpha(x)} = \frac{x - \sqrt{x^2 - 4}}{2}$. The Lucas polynomials are related to the Chebyshev polynomials of the first kind $T_n(x)$ by

$$(2.3) \quad L_n(x) = 2T_n\left(\frac{x}{2}\right)$$

and are characterized by

$$(2.4) \quad L_n\left(z + \frac{1}{z}\right) = z^n + \frac{1}{z^n}$$

for $n \geq 1$ and $z \neq 0$ because $z^n + \frac{1}{z^n} = \left(z + \frac{1}{z}\right)\left(z^{n-1} + \frac{1}{z^{n-1}}\right) - \left(z^{n-2} + \frac{1}{z^{n-2}}\right)$. For $z = e^{\sqrt{-1}\theta}$ Equation (2.4) implies

$$(2.5) \quad L_n(2 \cos \theta) = 2 \cos(n\theta).$$

Equation (2.4) also implies

$$(2.6) \quad L_{mn}(x) = L_m(L_n(x))$$

for all $m, n \geq 0$ since $z^{mn} + \frac{1}{z^{mn}} = (z^n)^m + \left(\frac{1}{z^n}\right)^m$.

¹Up to the index $n = 0$ the sequence of Lucas polynomials coincides with the sequence of the pyramidal polynomials from [3], the latter coming from a Riordan array.

Let us also mention that $z^{2n} + \frac{1}{z^{2n}} - 2 = (z^n + \frac{1}{z^n} - 2)(z^n + \frac{1}{z^n} + 2)$ implies

$$(2.7) \quad L_{2n}(x) - 2 = (L_n(x) - 2)(L_n(x) + 2)$$

and that

$$(2.8) \quad L_{2n}(x) + 2 = (L_n(x))^2$$

because $z^{2n} + \frac{1}{z^{2n}} + 2 = (z^n + \frac{1}{z^n})^2$.

3. FACTORIZATION OF THE POLYNOMIALS $L_n(x) - 2$

Proposition 3.1. *For $n \geq 1$ the roots of $f_n(x) := L_n(x) - 2$ are $\nu_k = 2 \cos(\frac{2\pi k}{n})$ with $0 \leq k \leq \frac{n}{2}$. For odd $n = 2m + 1$ the roots ν_k with $1 \leq k \leq m$ are double roots and $\nu_0 = 2$ is a simple root. For even $n = 2m$ the roots ν_k with $1 \leq k < m$ are double roots and ν_0, ν_m are simple roots.*

Proof. Since $f_n(2 \cos(\frac{2\pi k}{n})) = L_n(2 \cos(\frac{2\pi k}{n})) - 2 = 2 \cos(2\pi k) - 2 = 0$ each $\nu_k = 2 \cos(\frac{2\pi k}{n})$ for $k \in \mathbb{Z}$ is a root of $f_n(x)$. All different roots of this form are given by $0 \leq k \leq \frac{n}{2}$.

We claim that in this way we obtain all the roots of $f_n(x)$ because the number of roots above, counted with multiplicity, in fact already amounts to n .

To show the claim for $n = 2m + 1$ it suffices to show that

$$(3.1) \quad (L_{2m+1}(x) - 2)(x - 2) = (L_{m+1}(x) - L_m(x))^2,$$

which is a square of a polynomial. Identity (3.1) in turn can be shown by setting $x = z + \frac{1}{z}$ and evaluating

$$\begin{aligned} \left(z^{2m+1} + \frac{1}{z^{2m+1}} - 2\right) \left(z + \frac{1}{z} - 2\right) &= \frac{(z^{2(2m+1)} - 2z^{2m+1} + 1)(z^2 - 2z + 1)}{z^{2m+2}} \\ &= \left(\frac{(z^{2m+1} - 1)(z - 1)}{z^{m+1}}\right)^2 = \left(\left(z^{m+1} + \frac{1}{z^{m+1}}\right) - \left(z^m + \frac{1}{z^m}\right)\right)^2 \end{aligned}$$

For $n = 2m$ in turn it suffices to show the identity

$$(3.2) \quad (L_{2m}(x) - 2)(x - 2)(x + 2) = (L_{m+1}(x) - L_{m-1}(x))^2.$$

This, however, follows from

$$\frac{z^{2m} + \frac{1}{z^{2m}} - 2}{\left(z + \frac{1}{z} - 2\right)\left(z + \frac{1}{z} + 2\right)} = \left(\frac{(z^{2m} - 1)}{z^{m-1}(z^2 - 1)}\right)^2$$

and $(z + \frac{1}{z} - 2)(z + \frac{1}{z} + 2) = \left(\frac{(z^2 - 1)}{z}\right)^2$ because

$$\begin{aligned} \left(z^{2m} + \frac{1}{z^{2m}} - 2\right) \left(\frac{(z^2 - 1)}{z}\right)^2 &= \left(\frac{(z^{2m} - 1)}{z^{m-1}(z^2 - 1)} \left(\frac{(z^2 - 1)}{z}\right)^2\right)^2 = \left(\frac{(z^{2m} - 1)(z^2 - 1)}{z^{m+1}}\right)^2 \\ &= \left(z^{m+1} - \frac{1}{z^{m-1}} - z^{m-1} + \frac{1}{z^{m+1}}\right)^2. \end{aligned}$$

□

Next we use the fact that a polynomial $p(x)$ with an algebraic number a as a root has the minimal polynomial of a as a factor.

The minimal polynomial $\psi_n(x)$ of $2 \cos(\frac{2\pi}{n})$ is

$$(3.3) \quad \psi_n(x) = \prod_{\substack{\gcd(j,n)=1, \\ 0 < j < n/2}} \left(x - 2 \cos \frac{2\pi j}{n}\right)$$

for $n > 2$ with $\psi_1(x) = x - 2$ and $\psi_2(x) = x + 2$. The first examples are:

n	1	2	3	4	5	6	7	8	9	...
$\psi_n(x)$	$-2 + x$	$2 + x$	$1 + x$	x	$-1 + x + x^2$	$-1 + x$	$-1 - 2x + x^2 + x^3$	$-2 + x^2$	$1 - 3x + x^3$...

Theorem 3.2. Let $\alpha(x) = \frac{x + \sqrt{x^2 - 4}}{2}$ and $C_n(x)$ be the n th cyclotomic polynomial. Then for $n \geq 3$ the minimal polynomial of $2 \cos\left(\frac{2\pi}{n}\right)$ is

$$(3.4) \quad \psi_n(x) = \frac{C_n(\alpha(x))}{(\alpha(x))^{\varphi(n)/2}}.$$

Proof. For $n \geq 3$ the cyclotomic polynomial $C_n(x)$ is a symmetric polynomial with integer coefficients of even degree $\varphi(n)$. For example:

n	3	4	5	6	7	8	9	...
$C_n(x)$	$1 + x + x^2$	$1 + x^2$	$1 + x + x^2 + x^3 + x^4$	$1 - x + x^2$	$1 + x + \dots + x^6$	$1 + x^4$	$1 + x^3 + x^6$...

Therefore $\frac{C_n(x)}{x^{\varphi(n)/2}}$ is a sum of terms of the form $x^k + \frac{1}{x^k}$ with integer coefficients. Since $L_k = \alpha^k + \frac{1}{\alpha^k}$ we can write

$$\frac{C_n(\alpha(x))}{(\alpha(x))^{\varphi(n)/2}} = \sum_{k=0}^{\varphi(n)/2} c_k L_k(x) \in \mathbb{Z}[x]$$

with integer coefficients c_k and note that this polynomial is of degree $\varphi(n)/2$. Since $\alpha(2 \cos \theta) = \cos \theta + \sqrt{\cos^2 \theta - 1} = \cos \theta + \sqrt{-\sin^2 \theta} = e^{\sqrt{-1}\theta}$ we get

$$C_n\left(\alpha\left(2 \cos \frac{2\pi}{n}\right)\right) = C_n\left(e^{2\sqrt{-1}\pi/n}\right) = 0.$$

Therefore $\frac{C_n(\alpha(x))}{(\alpha(x))^{\varphi(n)/2}}$ is a monic polynomial of degree $\varphi(n)/2$ with integer coefficients and root $2 \cos \frac{2\pi}{n}$. Consequently it must coincide with the minimal polynomial $\psi_n(x)$. \square

For example, $\frac{C_9(x)}{x^3} = 1 + x^3 + \frac{1}{x^3}$ gives $\frac{C_9(\alpha(x))}{(\alpha(x))^3} = 1 + (\alpha(x))^3 + \frac{1}{(\alpha(x))^3} = 1 + L_3(x) = 1 - 3x + x^3 = \psi_9(x)$.

Corollary 3.3. For $n \geq 1$ we have $\psi_{2n+2}(x) = L_{2n}(x)$.

Proof. Since $C_{2n+2}(x) = x^{2n+1} + 1$ we get $\frac{C_{2n+2}(x)}{x^{2n}} = x^2 + \frac{1}{x^{2n}}$ and $(\alpha(x))^{2n} + \frac{1}{(\alpha(x))^{2n}} = L_{2n}(x)$. \square

Proposition 3.4. [1, Proposition 2.2] For $n \geq 1$ we have $L_n(x) - 2 = \psi_1(x) (\psi_2(x))^{e_n} \prod_{k|n, k \neq 1, 2} \psi_k^2(x)$ with $e_n = (1 + (-1)^n)/2$.

Proof. This follows from the multiplicities of the roots of $L_n(x) - 2$ and the fact that for $\gcd(j, n) = d$ the minimal polynomial of $2 \cos(2\pi j/n)$ is $\psi_{n/d}(x)$. \square

4. SOME PROPERTIES OF $Z_n(x)$

The formula $Z_n(x) = 2 - L_n(2 - x)$ shows that $Z_n(x)$ is closely related to the Lucas polynomials $L_n(x)$. The first terms are:

n	1	2	3	4	5	...
$Z_n(x)$	x	$4x - x^2$	$9x - 6x^2 + x^3$	$16x - 20x^2 + 8x^3 - x^4$	$25x - 50x^2 + 35x^3 - 10x^4 + x^5$...

Note that $-Z_n(-x)$ is a monic polynomial of degree n with positive coefficients. Since $L_n(2) = 2$ we have $Z_n(0) = 0$.

Let $c(n, k) = [x^k] L_n(2 + x)$. From $L_n(2 + x) = (2 + x)L_{n-1}(2 + x) - L_{n-2}(2 + x)$ we get $c(n, k) = 2c(n-1, k) + c(n-1, k-1) - c(n-2, k)$. Since $c(0, 1) = 0$ and $c(1, 1) = 1$ we get with induction that $c(n, 1) = n^2$. It is easily verified that $c(n, k) = \binom{n+k-1}{n-k} \frac{n}{k}$. Therefore, we get

$$(4.1) \quad Z_n(x) = \sum_{k=1}^n (-1)^{k-1} \binom{n+k-1}{n-k} \frac{n}{k} x^k.$$

With $\lambda(x) = \alpha(2 - x) = \frac{2-x+\sqrt{x^2-4x}}{2}$ we get

$$(4.2) \quad Z_n(x) = \frac{(\lambda(x)^n - 1)^2}{\lambda(x)^n}$$

because $Z_n(x) = 2 - L_n(2 - x) = 2 - \left((\lambda(x))^n + \frac{1}{(\lambda(x))^n} \right) = -\frac{((\lambda(x))^n - 1)^2}{(\lambda(x))^n}$.

Another connection with the Lucas polynomials is the following.

Remark 4.1.

$$\begin{aligned} Z_{2n+1}(x^2) &= (L_{2n+1}(x))^2, \\ Z_{2n}(x^2) &= 4 - (L_{2n}(x))^2. \end{aligned}$$

Proof. For odd m we have $Z_m(x^2) = 2 - L_m(2 - x^2) = 2 - L_m(-L_2(x)) = 2 + L_{2m}(x) = (L_m(x))^2$ because of Equation (2.8). The second identity follows from

$$\begin{aligned} Z_{2n}(x^2) &= 2 - L_{2n}(2 - x^2) = 2 - L_{2n}\left(2 - \left(z + \frac{1}{z}\right)^2\right) = 2 - L_{2n}\left(z^2 + \frac{1}{z^2}\right) = 2 - \left(z^{4n} + \frac{1}{z^{4n}}\right) \\ &= 4 - \left(z^{2n} + \frac{1}{z^{2n}}\right)^2 = 4 - (L_{2n}(x))^2. \end{aligned}$$

□

Proposition 4.2. *For arbitrary $u \neq 0$ we have*

$$(4.3) \quad Z_n\left(-\left(u - \frac{1}{u}\right)^2\right) = -\left(u^n - \frac{1}{u^n}\right)^2,$$

$$(4.4) \quad Z_{mn}(x) = Z_m(Z_n(x)).$$

Proof. Note that $Z_n\left(-\left(u - \frac{1}{u}\right)^2\right) = 2 - L_n\left(2 + \left(u - \frac{1}{u}\right)^2\right) = 2 - L_n\left(u^2 + \frac{1}{u^2}\right) = 2 - \left(u^{2n} + \frac{1}{u^{2n}}\right) = -\left(u^n - \frac{1}{u^n}\right)^2$. On the other hand, $Z_{mn}(x) = 2 - L_{mn}(2 - x) = 2 - L_m(L_n(2 - x)) = 2 - L_m(2 - Z_n(x)) = Z_m(Z_n(x))$. □

5. PROOF OF THEOREM 1.1

Let $z_n(x) = (-1)^{n-1}Z_n(x) = (-1)^{n-1}(2 - L_n(2 - x))$ be the monic versions of the $Z_n(x)$. The first terms are:

n	1	2	3	4	5	...
$z_n(x)$	x	$-4x + x^2$	$9x - 6x^2 + x^3$	$-16x + 20x^2 - 8x^3 + x^4$	$25x - 50x^2 + 35x^3 - 10x^4 + x^5$...

Proposition 3.4 gives

$$(5.1) \quad z_n(x) = \psi_1(2 - x)(\psi_2(2 - x))^{e_n} \prod_{k|n, k \neq 1, 2} (\psi_k(2 - x))^2.$$

Since all the $z_n(x)$ are monic we can replace all the polynomials of the right hand side by their monic versions. This gives

$$(5.2) \quad z_n(x) = \phi_1(x)(\phi_2(x))^{e_n} \prod_{k|n, k \neq 1, 2} (\phi_k(x))^2$$

with $\phi_1(x) = x$, $\phi_2(x) = x - 4$, $\phi_n(x) = (-1)^{\varphi(n)/2}\psi_n(2 - x)$.

Observing that $2 - 2 \cos\left(\frac{2\pi k}{n}\right) = 4 \sin^2\left(\frac{k\pi}{n}\right)$ we get from Equation (3.3) by changing $x \mapsto 2 - x$

$$(5.3) \quad \phi_n(x) = \prod_{\gcd(k,n)=1, 0 < k < n/2} \left(x - 4 \sin^2\left(\frac{k\pi}{n}\right)\right) = (-1)^{\varphi(n)/2} \frac{C_n(\lambda(x))}{(\lambda(x))^{\varphi(n)/2}}$$

for $n \geq 3$ with $\lambda(x) = \alpha(2-x) = \frac{2-x+\sqrt{x^2-4x}}{2}$. Thus $\phi_n(x)$ is the minimal polynomial of $4\sin^2\left(\frac{\pi}{n}\right)$. The first terms are:

n	1	2	3	4	5	6	7	
$\phi_n(x)$	x	$-4+x$	$-3+x$	$-2+x$	$5-5x+x^2$	$-1+x$	$-7+14x-7x^2+x^3$	
						8	9	\dots
						$2-4x+x^2$	$-3+9x-6x^2+x^3$	\dots

(see [6, entry A232633]).

Let

$$(5.4) \quad \begin{aligned} \Phi_n(x) &:= (\phi_n(x))^2 \text{ for } n \geq 3, \\ \Phi_1(x) &:= x, \quad \Phi_2(x) := 4-x = -\phi_2(x) \end{aligned}$$

such that $\phi_1(x)(\phi_2(x))^{e_n} = (-1)^{n-1}\Phi_1(x)(\Phi_2(x))^{e_n}$. Then $\Phi_n(x) \in \mathbb{Z}[x]$ with $\deg \Phi_n = \varphi(n)$.

With this the proof of Theorem 1.1 is a consequence of Equation (5.2).

6. HOW TO CALCULATE THE $\phi_k(x)$

6.1. Odd index. For odd m we know from Remark 4.1 that $Z_m(x^2) = (L_m(x))^2$. This implies

$$(6.1) \quad L_m(x) = x \prod_{d|m, d>1} \phi_d(x^2).$$

For example,

$$\begin{aligned} L_1(x) &= x, \quad L_3(x) = x(x^2-3) = x\phi_3(x^2), \quad L_5(x) = x(x^4-5x+5) = x\phi_5(x^2), \\ L_7(x) &= x(x^6-7x^4+14x^2-7) = x\phi_7(x^2), \quad L_9(x) = x(x^2-3)(x^6-6x^4+9x^2-3) = x\phi_3(x^2)\phi_9(x^2). \end{aligned}$$

Thus, for odd m the polynomial $\phi_m(x)$ can be easily calculated in terms of the Lucas polynomials.

6.2. The case indexed by powers of 2. Observing that $Z_{2^k}(x) = \prod_{j=0}^k \Phi_{2^j}(x)$ and $\Phi_{2^k}(x) = (\phi_{2^k}(x))^2$ we get for $k \geq 2$ that

$$(6.2) \quad \Phi_{2^k}(x) = \frac{Z_{2^k}(x)}{Z_{2^{k-1}}(x)} = \frac{1}{(\lambda(x))^{2^{k-1}}} \left(\frac{(\lambda(x))^{2^k} - 1}{(\lambda(x))^{2^{k-1}} - 1} \right)^2 = \frac{(\lambda(x))^{2^{k-1}} + 1}{(\lambda(x))^{2^{k-1}}} = \left((\lambda(x))^{2^{k-2}} + \frac{1}{\lambda(x)^{2^{k-2}}} \right)^2$$

implies

$$(6.3) \quad \phi_{2^k}(x) = (\lambda(x))^{2^{k-2}} + \frac{1}{(\lambda(x))^{2^{k-2}}} = \left((\lambda(x))^{2^{k-3}} + \frac{1}{(\lambda(x))^{2^{k-3}}} \right) - 2 = (\phi_{2^{k-1}}(x))^2 - 2$$

for $k \geq 3$. The first terms are:

n	1	2	3	4	5	\dots
$\phi_{2^n}(x)$	x	$-4+x$	$-2+x$	$2-4x+x^2$	$2-16x+20x^2-8x^3+x^4$	\dots

The same consideration for $L_{2^n}(z + \frac{1}{z}) = z^{2^n} + \frac{1}{z^{2^n}}$ together with $L_2(x) = x^2 - 2$ and $\phi_{2^2}(x^2) = x^2 - 2$ gives

$$(6.4) \quad \phi_{2^{n+1}}(x^2) = L_{2^n}(x)$$

for $n \geq 1$.

6.3. General case.

Theorem 6.1. *For odd $m \geq 3$ and $k \geq 2$*

$$(6.5) \quad \phi_{2m}(x) = (-1)^{\varphi(m)/2} \phi_m(4-x) = (-1)^{\varphi(m)/2} (\phi_m \circ \Phi_2)(x),$$

$$(6.6) \quad \phi_{2^k m}(x) = \phi_m((\phi_{2^k}(x))^2) = (\phi_m \circ \Phi_{2^k})(x).$$

Proof. We have $Z_{2m}(x) = Z_m(x)Z_m(4-x)$ because $Z_m(4-x) = 2 - L_m(2-4+x) = 2 - L_m(x-2) = 2 + L_m(2-x)$ and $Z_{2m}(x) = 2 - L_{2m}(2-x) = (2 - L_m(2-x))(2 + L_m(2-x))$ (see Equation (2.7)).

Comparing $Z_{2m}(x) = \prod_{d|2m} \Phi_d(x) = \prod_{d|m} \Phi_d(x)\Phi_{2d}(x)$ and

$$Z_{2m}(x) = Z_m(x)Z_m(4-x) = \prod_{d|m} \Phi_d(x)\Phi_d(4-x)$$

gives $\Phi_{2m}(x) = \Phi_m(4-x)$ for odd $m \geq 3$. Since $\Phi_m(x) = (\phi_m(x))^2$ and ϕ_m is monic of degree $\varphi(m)/2$ we get Equation (6.5).

We reproduce here the proof of Tri Nguyen [4] of Equation (6.6). It is based on the fact that $\phi_n(x)$ is the minimal polynomial of $4 \sin^2(\frac{\pi}{n})$ and has degree $\deg \phi_n = \frac{\varphi(n)}{2}$. It is sufficient to show that the root $\gamma_k = 4 \sin^2(\frac{\pi}{2^k m})$ of $\phi_{2^k m}(x)$ is also a root of $\phi_m((\phi_{2^k}(x))^2)$ as $\phi_{2^k m}(x)$ is irreducible and $\deg(\phi_m \circ (\phi_{2^k}(x))^2) = \frac{\varphi(m)}{2} 2^{k-1} = \frac{\varphi(2^k m)}{2} = \deg \phi_{2^k m}$.

Let $f^{(1)}(x) = f(x) = x^2 - 2$ and $f^{(n)}(x) = f \circ f^{(n-1)}(x)$ for $n \geq 2$. By Equation (6.3) $f(\phi_{2^{k-1}}(x)) = \phi_{2^k}(x)$, and therefore we have for $1 \leq j < k$

$$f^{(k-j)}(\phi_{2^j}(x)) = \phi_{2^k}(x).$$

Let us first show by induction that for any $\theta \in \mathbb{R}$

$$(6.7) \quad f^{(n)}(4 \sin^2(\theta) - 2) = 2 - 4 \sin^2(2^n \theta).$$

This is true for $n = 1$ since

$$f(4 \sin^2(\theta) - 2) = (4 \sin^2(\theta) - 2)^2 - 2 = (2 \cos^2(\theta) - 2 \sin^2(\theta))^2 - 2 = 4 \cos^2(2\theta) - 2 = 2 - 4 \sin^2(2\theta).$$

Equation (6.7) follows now from

$$f^{(n)}(4 \sin^2(\theta) - 2) = f\left(f^{(n-1)}(4 \sin^2(\theta) - 2)\right) = f(2 - 4 \sin^2(2^{n-1}\theta)) = 2 - 4 \sin^2(2^n \theta).$$

Now let us compute $\phi_{2^k}(\gamma_k)$. For $k = 2$ we find

$$\phi_4(\gamma_2) = \gamma_2 - 2 = 4 \sin^2\left(\frac{\pi}{4m}\right) - 2 = 2 \sin^2\left(\frac{\pi}{4m}\right) - 2 \cos^2\left(\frac{\pi}{4m}\right) = -2 \cos\left(\frac{\pi}{2m}\right),$$

and for $k > 2$

$$\begin{aligned} \phi_{2^k}(\gamma_k) &= f^{(k-2)}(\phi_4(\gamma_k)) = f^{(k-2)}(\gamma_k - 2) = f^{(k-2)}\left(4 \sin^2\left(\frac{\pi}{2^k m}\right) - 1\right) = 2 - 4 \sin^2\left(\frac{2^{k-2}\pi}{2^k m}\right) \\ &= 2 - 4 \sin^2\left(\frac{\pi}{4m}\right) = 2 \cos\left(\frac{\pi}{2m}\right). \end{aligned}$$

Finally, using Equation (6.5) we get

$$(\phi_m \circ \phi_{2^k}^2)(\gamma_k) = \phi_m\left(4 \cos^2\left(\frac{\pi}{2m}\right)\right) = (-1)^{\varphi(m)/2} \phi_{2m}\left(4 \sin^2\left(\frac{\pi}{2m}\right)\right) = 0.$$

□

Remark 6.2. The relation $Z_m(\Phi_{2^n}(x)) = \Phi_{2^n}(Z_m(x))$ holds for $n \geq 2$.

Proof. For $n = 2$ we have $\Phi_4(x) = (x-2)^2$. This gives $Z_m(\Phi_4(x)) = Z_m((x-2)^2) = (L_m(x-2))^2 = (L_m(2-x))^2 = (2 - Z_m(x))^2 = \Phi_4(Z_m(x))$. By induction we get

$$\begin{aligned} Z_m(\Phi_{2^{n+1}}(x)) &= Z_m\left(\left((\phi_{2^n}(x))^2 - 2\right)^2\right) = (L_m\left(\left(\phi_{2^n}(x)\right)^2 - 2\right))^2 = (2 - Z_m\left(\left(\phi_{2^n}(x)\right)^2\right))^2 \\ &= (2 - Z_m(\Phi_{2^n}(x)))^2 = (2 - \Phi_{2^n}(Z_m(x)))^2 = \left(\left(\phi_{2^n}(Z_m(x))\right)^2 - 2\right)^2 \\ &= \left(\phi_{2^{n+1}}(Z_m(x))\right)^2 = \Phi_{2^{n+1}}(Z_m(x)), \end{aligned}$$

where we have been using Remark 4.1 and Equation (6.3). □

7. FINAL REMARKS

Let us finally show some connection with the Fibonacci numbers $(F_n)_{n \geq 0} = (0, 1, 1, 2, 3, 5, 8, 13, 21, \dots)$. They satisfy $F_n = F_{n-1} + F_{n-2}$ for $n \geq 2$ with $F_0 = 0$, $F_1 = 1$ and Binet's Formula gives

$$(7.1) \quad F_n = \frac{\rho^n - \sigma^n}{\rho - \sigma}$$

with $\rho = \frac{1+\sqrt{5}}{2}$ and $\sigma = \frac{-1}{\rho} = \frac{1-\sqrt{5}}{2}$. It is known (see [5]) that the Fibonacci numbers form what is called a divisibility sequence.

Since $\lambda(5) = \frac{2-5+\sqrt{25-20}}{2} = \frac{-3+\sqrt{5}}{2} = -\left(\frac{1-\sqrt{5}}{2}\right)^2 = -\sigma^2$ we get $(\lambda(5))^n + \frac{1}{(\lambda(5))^n} = (-1)^n (\sigma^{2n} + \rho^{2n}) = (-1)^n \left(5 \left(\frac{\rho^n - \sigma^n}{\sqrt{5}}\right) + 2(-1)^n\right) = 5(-1)^n F_n^2 + 2$, which implies

$$(7.2) \quad Z_n(5) = (-1)^{n-1} 5 F_n^2.$$

Theorem 1.1 gives then the factorization of the Fibonacci numbers into primitive parts (cf. [6, entry A061446] and [5])

$$F_n = \prod_{d|n} p_d$$

with $p_1 = 1$ and $p_n = |\phi_n(5)|$ for $n \geq 2$. We get:

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	
$\phi_n(5)$	5	1	-2	-3	5	-4	-13	7	-17	11	-89	6	233	-29	61	47	...

For example

n	1	2	3	4	5	6	7	8	9
$F_n = \prod_{d n} p_d$	1	1 · 1	1 · 2	1 · 1 · 3	1 · 5	1 · 1 · 2 · 4	1 · 13	1 · 1 · 3 · 7	1 · 2 · 17

REFERENCES

1. Daniel J. Grubb, *Factoring Chebyshev polynomials*, Fibonacci Q. **52** (2014), no. 4, 360–366.
2. Yusuf Z. Gürtaş, *Chebyshev polynomials and the minimal polynomial of $\cos(2\pi/n)$* , Am. Math. Mon. **124** (2017), no. 1, 74–78.
3. Hans-Christian Herbig and Mateus de Jesus Gonçalves, *On the numerology of trigonometric polynomials*, 2023, arXiv:2311.13604.
4. Tri Nguyen, math.stackexchange.com, question 5005043.
5. Andrzej Nowicki, *Strong divisibility and LCM-sequences*, Am. Math. Mon. **122** (2015), no. 10, 958–966.
6. OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, 2023, Published electronically at <http://oeis.org>.
7. William Watkins and Joel Zeitlin, *The minimal polynomial of $\cos(2\pi/n)$* , Am. Math. Mon. **100** (1993), no. 5, 471–474.
8. Norman J. Wildberger, *Divine proportions. Rational trigonometry to universal geometry.*, Kingsford: Wild Egg, 2005.
9. Norman J. Wildberger, *Advice for maths exploration — Chebyshev and spread polynomials: the remarkable Goh factorization*, https://www.youtube.com/watch?v=P5bcwA_0ye0, May 2022.

DEPARTMENT OF MATHEMATICS, OSKAR-MORGENSTERN-PLATZ 1, 1090 VIENNA, AUSTRIA
 Email address: johann.cigler@univie.ac.at

DEPARTAMENTO DE MATEMÁTICA APLICADA, UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, AV. ATHOS DA SILVEIRA RAMOS 149, CENTRO DE TECNOLOGIA - BLOCO C, CEP: 21941-909 - RIO DE JANEIRO, BRAZIL
 Email address: herbighc@gmail.com