Rotating Traversable wormholes with two cosmic strings at the throat.

Vedant Subhash[∗](#page-1-0)

HEPCOS, Physics Department, State University of New York at Buffalo. (Dated: December 30, 2024)

Abstract

This paper investigates how rotating traversable wormholes incorporate two cosmic strings at the throat of the wormhole. Building on Edward Teo's framework for rotating traversable wormholes and Sung-Won Kim's analysis of scalar perturbations, a modified metric that accounts for these cosmic string perturbations is presented. This paper evaluates the Null Energy Condition in the presence of these string perturbations and focuses mainly on the scalar field dynamics in this modified spacetime. Using localized Gaussian perturbations to model cosmic string effects, this paper discusses the impact of angular and radial distributions from the cosmic strings on the spacetime geometry. Regularization has been applied to the shape function and energy condition violations at the wormhole throat, helping to eliminate divergences. The paper concludes with the result that the presence of cosmic strings induces unique anisotropic contributions. Future work could explore gravitational wave signatures associated with similar systems.

I. INTRODUCTION

The study of wormhole physics and cosmic strings represents an intersection of general relativity and quantum field theory, which I have tried to do the same in this paper (first calculating the NEC) and calculating scalar wave equation. In this introduction, I will provide a review of the literature on wormholes and cosmic strings, starting from the history of wormholes, exploring various configurations, and examining their combination.

Origin of Wormholes: Einstein-Rosen Bridge

The concept of wormholes, as solutions to Einstein's field equations, originated with the work of Albert Einstein and Nathan Rosen in 1935. Their seminal paper, [\[1\]](#page-27-0) introduced what is now known as the Einstein-Rosen Bridge. Initially, this mathematical construct was proposed as a way to model elementary particles, describing a "bridge" connecting two distinct regions of spacetime. The metric for the Einstein-Rosen Bridge is derived from the Schwarzschild solution and can be expressed as:

$$
ds^{2} = -\left(1 - \frac{2GM}{r}\right)dt^{2} + \left(1 - \frac{2GM}{r}\right)^{-1}dr^{2} + r^{2}d\Omega^{2},
$$
\n(1)

[∗] Also at HEPCOS, Physics Department, State University of New York at Buffalo.

where G is the gravitational constant, M the mass, and $d\Omega^2$ represents the angular components. This solution describe a non-traversable structure. In order to avoid the singularities, this investigation model matter and charge proposing a two flat spacetime connected by "bridge" as a representation of "particles". The paper addresses the issue of singularities in classical solutions like the Schwarzschild and Reissner-Nordström metrics, where the authors propose a modification to the field equations and this results in some kind of a "bridge-like" structure.[\[1\]](#page-27-0)

FIG. 1. Schemetic representation of an Einstein Rosen bridge. This is a black hole in disguise.[\[3\]](#page-27-1)

The potential role of connecting very distant regions of the universe was the main motivation behind describing this shortcut using the term "womrhole" in the 1950s. Different wormhole configurations are considered.

• Static Spherically Symmetric Wormholes: These wormholes, first studied in detail by Kip Thorne and Michael Morris in their 1988 paper, are described by the Morris-Thorne metric [\[2\]](#page-27-2)

$$
ds^{2} = -e^{2\Phi(r)}dt^{2} + \frac{dr^{2}}{1 - b(r)/r} + r^{2}(d\theta^{2} + \sin^{2}\theta d\phi^{2}),
$$
\n(2)

where $\Phi(r)$ is the redshift function and $b(r)$ the shape function. The throat of the wormhole is $b(r) = r$, and the structure is maintained by exotic matter that violates the Null Energy Condition (NEC).

• Rotating Wormholes: This rotating wormhole incorporates the effects of framedragging and off-diagnol metric components, as inspired by Kerr solution. [\[5\]](#page-27-3) [\[6\]](#page-27-4)

$$
ds^{2} = -e^{2\Phi(r)}dt^{2} + \frac{dr^{2}}{1 - b(r)/r} + r^{2} \left(d\theta^{2} + \sin^{2}\theta d\phi^{2}\right) - 2\omega r^{2} \sin^{2}\theta dt d\phi \qquad (3)
$$

where ω represents the angular velocity.

• Traversable Wormholes: Traversable wormholes are configurations where a traveler can pass through the wormhole without encountering singularities or horizons. Stability analyses of these wormholes often focus on perturbation theory and energy condition violations.

Progress in Traversable Wormholes and Rotating Traversable Wormholes: Morris and Thorne introduced traversable wormholes as viable solutions which require exotic matter for the throat of the wormhole to stay open. Subsequent studies by Matt Visser, Sayan Kar, and others have extensively explored the stability of these wormholes and the minimal energy violations needed to sustain them. In the context of rotating wormholes, Edward Teo's work introduced a rotating traversable wormhole model, highlighting frame-dragging effects. These rotating wormholes are different in the thin shell formalism of wormholes as in the former ones, I have a continuous spacetime, and in the thin shell structure there is a discontinuity. Theories of modified gravity, such as $f(R)$ gravity and scalar-tensor theories, have expanded the parameter space for traversable wormholes, reducing or eliminating the need for exotic matter.[\[12\]](#page-28-0)

Traversable Wormholes with Cosmic Strings

Cosmic strings are hypothesized as topological defects formed during symmetry-breaking phase transitions. These linear defects induce a conical spacetime geometry characterized by a deficit angle $\Delta \phi = 8\pi G\mu$, where μ is the string tension.[\[11\]](#page-28-1)

The exploration of wormholes, cosmic strings, and their associated theoretical frameworks, represents a convergence of geometry, topology, and high-energy physics. This compilation of works spans diverse approaches. The works by Dzhunushaliev et al.[\[15\]](#page-28-2) and Kuhfittig [\[14\]](#page-28-3)explore the dynamics of rotating wormholes, emphasizing their stability and traversability under exotic matter conditions. These studies extend classical Morris-Thorne solutions, incorporating angular momentum considerations, and provide insights into energy condition violations.

Radhakrishnan et al. [\[12\]](#page-28-0) review wormhole solutions within extended gravity frameworks, such as f(R) and Einstein-Gauss-Bonnet theories. These models attempt to minimize exotic matter requirements, contributing to singularity problems.

The interplay between cosmic strings and wormholes is highlighted in works by $[20][18][19]$ $[20][18][19]$ $[20][18][19]$ Clement, Aros, and others. These studies link cylindrical geometries with energy condition violations, suggesting cosmic strings can act as natural candidates for the wormhole throat.[\[14\]](#page-28-3) Their application in observational cosmology is also discussed. Bejarano et al. and related studies utilize the thin-shell formalism to construct and analyze wormholes within cylindrical and cosmic string frameworks.^{[\[21\]](#page-28-7)}[\[17\]](#page-28-8) This approach not only clarifies the conditions under which wormholes can exist but also examines their stability under perturbations using Israel junction methods.[\[22\]](#page-28-9) From stability analyses to the incorporation of quantum effects and modified gravity, traversable wormholes challenge conventional notions of causality, while cosmic string solutions may provide observable signatures. Tracing back to the Einstein-Rosen bridge, through Morris-Thorne wormholes[\[13\]](#page-28-10), to modern investigations into higher-dimensional and modified theories, this body of research reflects a consistent expansion of theoretical boundaries. In [\[25\]](#page-28-11) it was shown that the brane tension can counteract gravity without exotic matter in the brane world scenario. Also, in [\[26\]](#page-28-12) a solution describing a wormhole connecting two black holes in the same universe was constructed and it was shown that the repulsive force of the cosmological constant in de Sitter space can counteract gravity. Related to potential observational signature of wormholes, precision of astrophysical observations in near future will be sufficient for very sensitive searches for wormholes in astrophysical settings, as it was shown in [\[28\]](#page-29-0)[\[29\]](#page-29-1)[\[27\]](#page-28-13).

This project aims to explore the interplay between rotating traversable wormholes and cosmic strings, focusing on configurations that combine rotation and effects from strings. The primary aim of this research is to explore the impact of incorporating two cosmic strings at the throat of a rotating traversable wormhole. This work extends the foundations laid by E. Teo in his study on rotating traversable wormholes [\[6\]](#page-27-4) and Sung-Won Kim in his examination of scalar perturbations in rotating wormholes [\[9\]](#page-27-5).

The modifications in this study focus on introducing cosmic string perturbations into the wormhole metric and analyzing their implications for the Null Energy Condition (NEC) and scalar field dynamics. The project is structured as follows:

- 1. Metric Modification: Building upon Teo's metric, I introduced cosmic strings into the geometry. This requires careful adjustments to the angular metric components to accommodate the presence of the strings.
- 2. NEC Violation Analysis: The methodology for evaluating NEC violations is similar to Teo's analysis of null vectors and the stress-energy tensor components at the wormhole throat.
- 3. Scalar Perturbation Analysis: Following Sung-Won Kim's approach, I derived and simplified the scalar wave equation within the context of the modified metric. This process involved the use of a toy model to decouple and solve the equation.

THE MODIFIED METRIC

The spacetime metric, incorporating the effects of cosmic strings, is expressed as:

$$
ds^{2} = -N(r,\theta)^{2}dt^{2} + \frac{r}{r-b(r)}dr^{2} + r^{2}K(r,\theta)^{2} \left(d\theta^{2} + g(r,\theta)^{2}\sin^{2}\theta(d\phi - \omega(r,\theta)dt)^{2}\right)
$$
 (4)

where:

- $N(r, \theta)$: Gravitational redshift factor,
- $b(r)$: Shape function satisfying the flare-out condition at the throat,
- $K(r, \theta)$: Angular metric factor,
- $f(r, \theta) = K(r, \theta)g(r, \theta)$: Cosmic string perturbation,
- $g(r, \theta) = 1 \epsilon h(r, \theta)$, where $h(r, \theta)$ introduces the cosmic string deformation,
- $\bullet \hspace{0.1cm} \omega(r,\theta) \colon$ Rotational function describing frame-dragging effects.

ENERGY CONDITION ANALYSIS

By defining appropriate null vectors, I analyzed the violation of the NEC in the presence of cosmic strings. This step aligns with the principles outlined in Teo's paper [\[6\]](#page-27-4), where the NEC is evaluated along specific trajectories. The cosmic strings modify the stress-energy tensor, introducing unique terms that alter the geometric behavior at the throat.

SCALAR PERTURBATION DYNAMICS

Incorporating scalar fields, I extended the analysis to wave propagation within this modified wormhole geometry. Using Sung-Won Kim's framework [\[9\]](#page-27-5), I derived the wave equation for a scalar field and separated it into radial and angular components. The simplified toy model allows for the explicit solution of the decoupled equations and use the Frobenius series solution to solve for the Radial equation.

This research opens avenues for studying more complex perturbative effects, such as gravitational waves induced by cosmic string-wormhole interactions or the dynamics of multiplestring configurations. Additionally, exploring the semi-classical regime, as suggested by Teo, could yield insights into quantum-scale wormholes [\[6\]](#page-27-4).

II. MATHEMATICAL FORMULATION

The spacetime metric, incorporating the effects of cosmic strings, is expressed as:

$$
ds^{2} = -N(r,\theta)^{2}dt^{2} + \frac{r}{r - b(r)}dr^{2} + r^{2}K(r,\theta)^{2} \left(d\theta^{2} + g(r,\theta)^{2}\sin^{2}\theta(d\phi - \omega(r,\theta)dt)^{2}\right) (5)
$$

For simplification, the following relationship is used:

$$
f(r,\theta) = K(r,\theta)g(r,\theta)
$$
\n(6)

where the effects of the cosmic strings are encapsulated in $g(r, \theta)$, defined as:

$$
g(r, \theta) = 1 - \epsilon h(r, \theta) \tag{7}
$$

Here, ϵ represents the deficit angle parameter introduced by the cosmic strings, and $h(r, \theta)$ is a function describing the radial and angular distribution of the cosmic string's influence

The metric tensor components in spherical coordinates (t, r, θ, ϕ) are given by:

$$
g_{\mu\nu} = \begin{bmatrix} -N(r,\theta) + r^2 f(r,\theta)^2 \sin^2 \theta \, \omega(r,\theta)^2 & 0 & 0 & -r^2 f(r,\theta)^2 \sin^2 \theta \, \omega(r,\theta) \\ 0 & \frac{1}{1 - \frac{b(r,\theta)}{r}} & 0 & 0 \\ 0 & 0 & r^2 K(r,\theta)^2 & 0 \\ -r^2 f(r,\theta)^2 \sin^2 \theta \, \omega(r,\theta) & 0 & 0 & r^2 f(r,\theta)^2 \sin^2 \theta \end{bmatrix}
$$
(8)

A. Localized Perturbations from Cosmic Strings

The function $h[r, \theta]$, which models the perturbations introduced by the cosmic strings, is expressed as:

$$
h[r,\theta] = \exp\left(-\frac{(r-r_0)^2}{\sigma_r^2}\right) \left[\exp\left(-\frac{(\theta-\pi/4)^2}{\sigma_\theta^2}\right) + \exp\left(-\frac{(\theta-3\pi/4)^2}{\sigma_\theta^2}\right)\right],\qquad(9)
$$

where:

- r_0 is the radial position of the throat.
- σ_r controls the radial spread of the perturbation.
- σ_{θ} governs the angular spread of the perturbation, $\theta = \pi/4$ and $\theta = 3\pi/4$.
- The two Gaussian terms correspond to the localized effects of the two cosmic strings, symmetrically placed at $\theta = \pi/4$ (45°) and $\theta = 3\pi/4$ (135°).

The Gaussian form ensures that the perturbations are smooth, rapidly decaying, and concentrated around specific regions in both radial and angular directions.

1. Visualization of Perturbations

Figure [2](#page-8-0) illustrates the perturbation $h[r, \theta]$ due to the two cosmic strings. The plot is presented in polar coordinates, highlighting the localization of the perturbations around $\theta = \pi/4$ and $\theta = 3\pi/4$.

2. Interpretation of the Plot

The plot demonstrates several key features of the perturbation function:

- Radial Localization: The perturbation magnitude peaks around r_0 , with a rapid falloff determined by σ_r . This ensures that the cosmic string effects are localized near the wormhole throat.
- Angular Localization: The perturbation is symmetrically distributed around $\theta =$ $\pi/4$ and $\theta = 3\pi/4$, with the spread governed by σ_{θ} . This corresponds to the two cosmic strings' influence.

FIG. 2. Localized perturbations $h[r, \theta]$ due to two cosmic strings. The color scale represents the magnitude of the perturbation $\epsilon h[r, \theta]$, with brighter regions indicating stronger effects. The perturbations are concentrated around $\theta = \pi/4$ and $\theta = 3\pi/4$, with radial localization controlled by σ_r .

• Magnitude Variations: The color gradient illustrates the relative strength of the perturbation, with the highest values concentrated near the intersection of $r \approx r_0$ and $\theta = \pi/4, 3\pi/4.$

These localized perturbations significantly affect the anisotropy of the wormhole geometry. The next section will analyze how these perturbations affect the Null Energy Condition (NEC) and provides a detailed analysis of the Null Energy Condition (NEC) for a rotating traversable wormhole, incorporating the effects of two cosmic strings at the throat. The NEC is evaluated based on the spacetime metric components and the Ricci tensor components, with emphasis on the physical contributions from frame-dragging due to rotation and cosmic string perturbations.

B. Verification of the Null Condition

The null vector k^{μ} for our spacetime metric is defined as:

$$
k^{\mu} = \left(\frac{\alpha(r,\theta)}{N(r,\theta)}, \sqrt{\frac{1-b(r)}{r}}, \frac{\beta(r,\theta)}{rK(r,\theta)}, \frac{\omega(r,\theta)(1-\epsilon h(r,\theta))}{N(r,\theta)}\right)
$$
(10)

The spacetime metric is given by:

$$
ds^{2} = -N(r,\theta)^{2}dt^{2} + \frac{r}{r-b(r)}dr^{2} + r^{2}K(r,\theta)^{2} \left(d\theta^{2} + g(r,\theta)^{2}\sin^{2}\theta(d\phi - \omega(r,\theta)dt)^{2}\right)
$$
 (11)

where:

- $N(r, \theta)$: Gravitational redshift factor,
- $b(r)$: Shape function,
- $K(r, \theta)$: Angular metric factor,
- $f(r, \theta) = K(r, \theta)g(r, \theta)$: Cosmic string perturbation,
- $g(r, \theta) = (1 \epsilon h(r, \theta))$
- $\omega(r, \theta)$: Rotational function.

III. VIOLATION OF THE NULL ENERGY CONDITION

In general relativity, the Null Energy Condition (NEC) states that the contraction of the Ricci tensor with a null vector k^{μ} must satisfy:

$$
R_{\mu\nu}k^{\mu}k^{\nu}\geq 0.
$$

The NEC is central to many classical theorems, including the singularity theorems and topological censorship.[\[3\]](#page-27-1) In this work, we analyze the NEC violation in a spacetime influenced by two cosmic strings and rotation. This analysis builds on methods previously explored in [\[6\]](#page-27-4), where NEC violation is explicitly demonstrated in traversable wormhole geometries.

A. Setup: Stationary Axisymmetric Metric with Cosmic Strings and Rotation

The spacetime metric is stationary and axisymmetric, given by:

$$
ds^2 = -N(r,\theta)^2 dt^2 + \frac{r}{r-b(r)} dr^2 + r^2 K(r,\theta)^2 \left(d\theta^2 + f(r,\theta)^2 \sin^2\theta (d\phi - \omega(r,\theta)dt)^2\right),
$$

where:

- $N(r, \theta)$: Gravitational redshift factor,
- $b(r)$: Shape function, encoding the throat geometry,
- $K(r, \theta)$: Angular metric factor,
- $f(r, \theta) = K(r, \theta)(1 \epsilon h(r, \theta))$: Cosmic string perturbation,
- $\omega(r, \theta)$: Rotational function.

B. Null Vector and NEC Expression

The null vector k^{μ} for the spacetime is constructed as:

$$
k^{\mu} = \left(\frac{\alpha(r,\theta)}{N(r,\theta)}, \sqrt{\frac{1-b(r)}{r}}, \frac{\beta(r,\theta)}{rK(r,\theta)}, \frac{\omega(r,\theta)(1-\epsilon h(r,\theta))}{N(r,\theta)}\right).
$$

ANALYSIS OF NEC NEGATIVITY AT THE THROAT

To ensure the Null Energy Condition (NEC) is negative at the throat $(r = r_0)$, we strategically selected specific forms for the parameters $b(r)$, $f(r, \theta)$, $K(r, \theta)$, $\omega(r, \theta)$, and $N(r, \theta)$. This approach ensures dominant negative contributions from critical terms.

Parameter Selection

$$
b(r) = r_0 - \epsilon (r - r_0)^2,
$$

\n
$$
f(r, \theta) = K(r, \theta)g(r, \theta), \text{ where } g(r, \theta) = 1 - \epsilon h(r, \theta),
$$

\n
$$
K(r, \theta) = 1 + \gamma \sin^2(\theta),
$$

\n
$$
f(r, \theta) = (1 + \gamma \sin^2(\theta))(1 - \epsilon h(r, \theta)).
$$

\n
$$
\omega(r, \theta) = \alpha e^{-\lambda(r - r_0)^2} \cos(\theta),
$$

\n
$$
N(r) = \sqrt{1 - \frac{b(r)}{r}}.
$$

The components $h(r, \theta)$ represent localized perturbations introduced by cosmic strings and are typically expressed as:

$$
h(r,\theta) = \exp\left(-\frac{(r-r_0)^2}{\sigma_r^2}\right) \left[\exp\left(-\frac{(\theta-\pi/4)^2}{\sigma_\theta^2}\right) + \exp\left(-\frac{(\theta-3\pi/4)^2}{\sigma_\theta^2}\right)\right].\tag{12}
$$

The remaining parameters are as follows:

$$
b(r) = r_0 - \epsilon (r - r_0)^2, \quad \epsilon > 0,
$$
\n(13)

$$
\omega(r,\theta) = \alpha e^{-\lambda(r-r_0)^2} \cos(\theta), \quad \alpha, \lambda > 0,
$$
\n(14)

$$
N(r) = \sqrt{1 - \frac{b(r)}{r}}.\tag{15}
$$

Now, we have computed all the 30 terms of the NEC in Appendix A, we will check whether the NEC is violated at the throat of the wormhole. Substituting the parameters values in the NEC equation, we see that the 3rd term vanishes $N(r_0)$ tends to zero and the term extends to negative infinity. In order to remove this divergence, I will regularize two parameters $b(r)$ and $N(r)$. The parameter $b(r)$ is a really interesting one to follow as it must be differentiable and physically regular (no singularities) across the spacetime. $b(r)$ < 1 ensures the flare-out condition is met. $b(r)$ tends to zero as r tends to infinity, ensures the wormhole asymptotically matches a flat or nearly-flat spacetime.

1. Regularized Parameter Forms

To avoid divergence at the throat, $b(r)$ should approach r_0 smoothly while ensuring $r - b(r) \neq 0$ for small deviations from r₀. A regularized form is:

$$
b(r) = r_0 \left(1 - \epsilon \frac{(r - r_0)^2}{r_0^2} \right),
$$

where $\epsilon > 0$. This ensures:

$$
b(r_0) = r_0, \quad \frac{\partial b(r)}{\partial r} = -2\epsilon \frac{(r - r_0)}{r_0^2}, \quad \frac{\partial^2 b(r)}{\partial r^2} = -\frac{2\epsilon}{r_0^2}.
$$

Cosmic String Factor $f(r, \theta)$:

To avoid $f(r, \theta) \to 0$, we regularize $g(r, \theta)$ to ensure positivity:

$$
g(r, \theta) = \max(1 - \epsilon h(r, \theta), \delta),
$$

where $\delta > 0$ is a small positive constant ensuring $f(r, \theta) > 0$ everywhere. The regularized $f(r, \theta)$ becomes:

$$
f(r,\theta) = (1 + \gamma \sin^2(\theta)) \max(1 - \epsilon h(r,\theta), \delta).
$$

Summary of NEC Analysis

The Null Energy Condition (NEC) is computed as the sum of 30 terms evaluated at the throat of the wormhole $(r = r_0)$, where $b(r_0) = r_0$. Using the revised parameters:

$$
b(r) = r_0 - \epsilon (r - r_0)^2, \quad f(r, \theta) = (1 + \gamma \sin^2(\theta))(1 - \epsilon h(r, \theta)),
$$

$$
K(r, \theta) = 1 + \gamma \sin^2(\theta), \quad \omega(r, \theta) = \alpha e^{-\lambda (r - r_0)^2} \cos(\theta), \quad N(r) = \sqrt{1 - \frac{b(r)}{r}},
$$

we analyze the sign of the NEC at the throat.

Key Findings

- 1. **Regularized Contributions:** Terms involving $N(r)$, such as Term 3, are regularized by the choice $b(r) = r_0 - \epsilon (r - r_0)^2$, ensuring all terms are finite. For example, $N(r)^2 \sim \epsilon (r - r_0)^2$ at the throat, eliminating divergences.
- 2. Negative NEC Contributions: Several terms contribute negatively due to their dependence on derivatives of $f(r, \theta)$, $h(r, \theta)$, and $\omega(r, \theta)$. Notably:

Term 17:
$$
\frac{-8f(r_0, \theta) (1 - \epsilon h(r_0, \theta))^2 \omega(r_0, \theta)^2 \sin(2\theta) \frac{\partial f(r_0, \theta)}{\partial \theta}}{K(r_0, \theta)^2 N(r_0)^2},
$$
Term 20:
$$
\frac{2(r_0 - b(r_0))\omega(r_0, \theta) \sin^3(\theta) (-2\lambda \alpha \cos(\theta))}{N(r_0)^2}.
$$

At $r = r_0$, the regularized $N(r_0)$ amplifies these negative contributions.

3. Angular and Localized Effects: The perturbation function $h(r, \theta)$, defined as:

$$
h(r,\theta) = \exp\left(-\frac{(r-r_0)^2}{\sigma_r^2}\right) \left[\exp\left(-\frac{(\theta-\pi/4)^2}{\sigma_\theta^2}\right) + \exp\left(-\frac{(\theta-3\pi/4)^2}{\sigma_\theta^2}\right)\right] \tag{16}
$$

contributes significantly to the angular variations. This is evident in terms like Term 11 and Term 17, where derivatives of $h(r, \theta)$ combine with sinusoidal terms, producing localized negative energy density contributions.

- 4. **Parameter-Driven Dominance:** The choice of parameters such as $\epsilon > 0, \gamma > 0, \alpha > 1$ $0, \lambda > 0$ ensures:
	- Smooth regularization near the throat with $r b(r) \sim \epsilon (r r_0)^2$.
	- Enhanced angular contributions from $K(r, \theta) = 1 + \gamma \sin^2(\theta)$ and $\omega(r, \theta)$, ensuring finite but negative terms.

After carefully calculating and analyzing the terms and certain conditions under which the terms are positive and negative, the overall NEC comes to be negative. Finite nature of then NEC was achieved through regularization, and negative dominance is achieved through carefully choosing the parameters which are physically consistent.

Dominance of Negative Terms

The overall sign of the NEC is negative, dominated by terms like Term 17 and Term 20. These terms involve:

- The quadratic form of $b(r)$, ensuring $(r b(r)) \rightarrow 0$ smoothly, amplifying negative factors in $N(r)$ terms.
- Sinusoidal variations from $\omega(r, \theta)$, contributing negatively through $\cos(\theta)$ and $\sin(2\theta)$ factors.
- Localized perturbations from $h(r, \theta)$, which add angular dependence to derivatives of $f(r, \theta)$ and $\omega(r, \theta)$.

Conclusion

This analysis confirms that the NEC is negative at the throat under the chosen parameter values, demonstrating the presence of exotic matter required to sustain the wormhole structure. The analysis done in this study closely aligns with the one done here.

IV. SCALAR PERTURBATIONS OF ROTATING WORMHOLE INCORPORAT-ING TWO COSMIC STRINGS AT THE THROAT

The scalar wave equation has the following form:[\[7\]](#page-27-6)

$$
\nabla_{\mu}\nabla^{\mu}\Phi = \frac{1}{\sqrt{-g}}\partial_{\mu}\left(g^{\mu\nu}\sqrt{-g}\,\partial_{\nu}\Phi\right) = 0.
$$
 (17)

Now, with the given modified metric in eq[4], we can solve for every components and then combine together as shown below:

$$
\Box[t, r, \theta, \phi] = \frac{e^{i(-tw+m\phi)}}{2r^2(r-b[r])f[r, \theta]^2K[r, \theta]^3N[r, \theta]} \Big(
$$

\n
$$
2m^2(-r+b[r])\csc^2(\theta)K[r, \theta]^3N[r, \theta]R[r]S[\theta]
$$

\n
$$
+ 2(r-b[r])f[r, \theta]K[r, \theta]N[r, \theta] \Big(r(r-b[r])K[r, \theta]^2S[\theta]\partial_r f[r, \theta]\partial_r R[r]
$$

\n
$$
+ R[r]\partial_{\theta}f[r, \theta]\partial_{\theta}S[\theta]\Big)
$$

\n
$$
+ f[r, \theta]^2 \Big(2r(r-b[r])^2K[r, \theta]^2N[r, \theta]S[\theta]\partial_r K[r, \theta]\partial_r R[r]
$$

\n
$$
+ (r-b[r])K[r, \theta]^3S[\theta] \Big(2r^2R[r](w-m\omega[r, \theta])^2
$$

\n
$$
+ r(r-b[r])\partial_r N[r, \theta]\partial_r R[r]
$$

\n
$$
+ N[r, \theta] \Big(-(3b[r] + r(-4 + \partial_r b[r]))\partial_r R[r]
$$

\n
$$
+ 2r(r-b[r])\partial_{r^2}R[r]\Big)\Big)
$$

\n
$$
+ 2(-r+b[r])N[r, \theta]R[r]\partial_{\theta}K_1[r, \theta]\partial_{\theta}S[\theta]
$$

\n
$$
+ K[r, \theta]R[r] \Big((r-b[r])\partial_{\theta}N[r, \theta]\partial_{\theta}S[\theta] + \partial_{\theta}S[\theta]\Big)
$$

\n
$$
+ N[r, \theta] \Big(2(r-b[r]) \Big(\cot(\theta)\partial_{\theta}S[\theta] + \partial_{\theta}S[\theta]\Big)\Big)\Big)\Big)\Big)
$$

The wave equation is highly complex to separate them, so certain assumptions are taken to simplify this wave equation in a compact form so that separation of variables can be applied. In the study, I have defined the perturbation from the two cosmic strings in a more localized manner: The perturbation $h[r, \theta]$ is localized and modeled as:

$$
h[r,\theta] = \exp\left(-\frac{(r-r_0)^2}{\sigma_r^2}\right) \left[\exp\left(-\frac{(\theta-\pi/4)^2}{\sigma_\theta^2}\right) + \exp\left(-\frac{(\theta-3\pi/4)^2}{\sigma_\theta^2}\right)\right],\qquad(19)
$$

where:

- r_0 is the radial position of the throat.
- \bullet σ_r and σ_θ control the spread of the perturbation.
- The two Gaussian terms correspond to the localized effects of the two cosmic strings at $\theta = \pi/4$ and $\theta = 3\pi/4$.

Given the new definitions: $f(r, \theta) = K(r, \theta)g(r, \theta)$ and $g(r, \theta) = 1 - \epsilon h(r, \theta)$. The function $b(r)$ is given by: $b(r) = \frac{r_0^2}{r}$, with $K(r, \theta) = 1$, the simplifications yield: $f(r, \theta) = 1 - \epsilon h(r, \theta)$. The terms involving $f(r, \theta)$, $\frac{\partial f(r, \theta)}{\partial r}$, and $\frac{\partial f(r, \theta)}{\partial \theta}$ now become:

$$
f(r,\theta) = 1 - \epsilon h(r,\theta) \tag{20}
$$

$$
\frac{\partial f(r,\theta)}{\partial r} = -\epsilon \frac{\partial h(r,\theta)}{\partial r}
$$
\n(21)

$$
\frac{\partial f(r,\theta)}{\partial \theta} = -\epsilon \frac{\partial h(r,\theta)}{\partial \theta} \tag{22}
$$

For $h(r, \theta)$: $\frac{\partial h(r, \theta)}{\partial r} = -\frac{2(r-r_0)}{\sigma_r^2}$ $\frac{\dot{r}-r_0}{\sigma_r^2}h(r,\theta).$

$$
\frac{\partial h(r,\theta)}{\partial \theta} = \exp\left(-\frac{(r-r_0)^2}{\sigma_r^2}\right) \left[-\frac{2(\theta-\frac{\pi}{4})}{\sigma_\theta^2} \exp\left(-\frac{(\theta-\frac{\pi}{4})^2}{\sigma_\theta^2}\right) - \frac{2(\theta-\frac{3\pi}{4})}{\sigma_\theta^2} \exp\left(-\frac{(\theta-\frac{3\pi}{4})^2}{\sigma_\theta^2}\right) \right]
$$
(23)

Under these new definitions and the subsequent simplification using the above expressions, we get the scalar equation as:

$$
\Box[t,r,\theta,\phi] = \frac{e^{i(-tw+m\phi)}}{2r^2 \left(r - \frac{r_0^2}{r}\right) \left(1 - \frac{\epsilon h(r,\theta)}{e^{(r-r_0)^2/\sigma_r^2}}\right)^2} \left\{ 2m^2 \left(r - \frac{r_0^2}{r}\right) \csc^2(\theta)R(r)S(\theta) \right.\n+ \left(1 - \frac{\epsilon h(r,\theta)}{e^{(r-r_0)^2/\sigma_r^2}}\right)^2 \left[-\frac{2r_0^2}{r}\left(r - \frac{r_0^2}{r}\right)^2 S(\theta)\frac{\partial R(r)}{\partial r} \right.\n+ \left(r - \frac{r_0^2}{r}\right)S(\theta)\left(2r^2(w - m\omega(r,\theta))^2 R(r) \right.\n- \frac{3r_0^2}{r}\frac{\partial R(r)}{\partial r} + r\left(r - \frac{r_0^2}{r}\right)\frac{\partial R(r)}{\partial r} \right.\n+ 2r\left(r - \frac{r_0^2}{r}\right)\frac{\partial^2 R(r)}{\partial r^2}\right)\left.\left.\right\|_{\theta=\theta} = \epsilon R(r)\frac{\partial S(\theta)}{\partial \theta}\frac{\partial h(r,\theta)}{\partial \theta} \right.\n- \epsilon r\left(r - \frac{r_0^2}{r}\right)S(\theta)\frac{\partial R(r)}{\partial r}\frac{\partial h(r,\theta)}{\partial r}\right)\left.\right\}.
$$
\n(24)

The newly defined functions are: $N = 1$, $K = 1$, $b(r) = \frac{r_0^2}{r}$, $f(r, \theta) = K(r, \theta)g(r, \theta), \quad g(r, \theta) = 1 - \epsilon h(r, \theta),$

$$
h(r,\theta) = \exp\left(-\frac{(r-r_0)^2}{\sigma_r^2}\right) \left[\exp\left(-\frac{(\theta-\pi/4)^2}{\sigma_\theta^2}\right) + \exp\left(-\frac{(\theta-3\pi/4)^2}{\sigma_\theta^2}\right)\right] \tag{25}
$$

Using these definitions, we analyze the scalar equation by separating variables into radial $(r$ -dependent) and angular (θ -dependent) parts. To separate variables in the simplified scalar equation, we aim to write it in a form where r and θ dependent components are isolated. The following methods involves systematic inspection and grouping of terms based on their dependence on r and θ

The equation contains terms involving:

- *r*-dependent functions: $R(r)$, $b(r) = \frac{r_0^2}{r}$, and their derivatives.
- θ -dependent functions: $S(\theta)$, $h(r, \theta)$, and their derivatives.
- Mixed terms: Products of r and θ -dependent functions, e.g.,

$$
R(r)S(\theta), \quad \frac{\partial f(r,\theta)}{\partial r} \frac{\partial f(r,\theta)}{\partial \theta} \tag{26}
$$

By grouping and dividing appropriately, we identify terms that must individually satisfy separate differential equations. Let's work through this step by step. Now in such a highly complex model, we will display the r, and θ dependent terms separately.

r-Dependent Terms

- $R(r) \cdot (r \frac{r_0^2}{r})$
- $R(r) \cdot (r \frac{r_0^2}{r})$

$$
\bullet \ -2\cdot \tfrac{r_0^2}{r}\cdot (r-\tfrac{r_0^2}{r})^2\cdot \tfrac{\partial R(r)}{\partial r}
$$

- $(r \frac{r_0^2}{r}) \cdot 2r^2 \cdot (w m\omega(r, \theta))^2 \cdot R(r)$
- \bullet $(r-\frac{r_0^2}{r})\cdot \left(-3\cdot\frac{r_0^2}{r}\right)\cdot \frac{\partial R(r)}{\partial r}$ ∂r

$$
\bullet \ (r - \tfrac{r_0^2}{r})^2 \cdot 2r \cdot \tfrac{\partial^2 R(r)}{\partial r^2} \]
$$

•
$$
2 \cdot (1 - \epsilon h(r, \theta) e^{-(r-r_0)^2/\sigma_r^2}) \cdot \left(r - \frac{r_0^2}{r}\right) \cdot R(r)
$$

θ-Dependent Terms

- $S(\theta) \cdot \csc^2(\theta)$
- $S(\theta)$
- \bullet $-\epsilon \cdot \frac{\partial S(\theta)}{\partial \theta} \cdot \frac{\partial h(r,\theta)}{\partial \theta}$ ∂θ

Now that we have a clear picture of all the r - dependent terms and the θ dependent terms, we can work on Radial, and Angular Equations separately. The former one looks more complex than the latter, as we shall see in the following sections. Combine all r dependent terms into:

$$
R(r) = R(r) \left(m^2 \cdot \left(r - \frac{r_0^2}{r} \right) + 2r^2 \cdot (w - m\omega(r, \theta))^2 \right) \tag{27}
$$

$$
-2 \cdot \frac{r_0^2}{r} \cdot \left(r - \frac{r_0^2}{r}\right)^2 \cdot \frac{\partial R(r)}{\partial r}
$$
\n(28)

$$
-3 \cdot \frac{r_0^2}{r} \cdot \left(r - \frac{r_0^2}{r}\right) \cdot \frac{\partial R(r)}{\partial r} \tag{29}
$$

$$
+ 2 \cdot r \cdot \left(r - \frac{r_0^2}{r}\right)^2 \cdot \frac{\partial^2 R(r)}{\partial r^2}.
$$
\n(30)

This becomes a second-order differential equation for $R(r)$. Now, group similar terms in the expression and substituting $b(r) = \frac{(r_0)^2}{r}$ $\frac{p_0^2}{r}$ and using $r - b(r) = r - \frac{r_0^2}{r}$, we simplify the combined expression to get a second-order linear differential equation. Write the equation as:

$$
2r \cdot (r - b(r))^{2} \cdot \frac{\partial^{2} R(r)}{\partial r^{2}} - \frac{\partial R(r)}{\partial r} \cdot \left[2b(r) \cdot (r - b(r))^{2} + 3b(r) \cdot (r - b(r)) \right] + R(r) \cdot \left[m^{2} \cdot (r - b(r)) + 2r^{2} \cdot (w - m\omega(r, \theta))^{2} \right] = 0.
$$

In order to solve a second-order linear differential equation, introduce a substitution of $y(r) = R(r)$. Then I solve the equation using Forbenius series.[\[24\]](#page-28-14)

After substituting $b(r) = \frac{r_0^2}{r}$ into the equation, leads to $r - b(r) = r - \frac{r_0^2}{r}$ Thus, the terms involving $b(r)$ expand as follows:

$$
(r - b(r))^{2} = \left(r - \frac{r_0^{2}}{r}\right)^{2}
$$
\n(31)

$$
b(r) \cdot (r - b(r)) = \frac{r_0^2}{r} \cdot (r - \frac{r_0^2}{r})
$$
\n(32)

Rewrite the equation, grouping terms by derivatives of $R(r)$. In order to solve this system of equations, I assume the solution for $R(r)$ is:

$$
R(r) = \sum_{n=0}^{\infty} a_n r^{n+c}
$$
\n(33)

where c is the indicial exponent. Substitute this into the differential equation. For the first and the second derivative term:

$$
\frac{\partial R(r)}{\partial r} = \sum_{n=0}^{\infty} a_n (n+c)r^{n+c-1}
$$
\n(34)

$$
\frac{\partial^2 R(r)}{\partial r^2} = \sum_{n=0}^{\infty} a_n (n+c)(n+c-1)r^{n+c-2}
$$
\n(35)

We solve the radial equation using a Frobenius series:

$$
R(r) = \sum_{n=0}^{\infty} a_n r^{n+c}.
$$
\n(36)

The indicial equation, derived from the lowest power of r, gives: $c(c-1) = 0 \implies c =$ 0 or $c = 1$. For finite behavior near $r = 0$, we choose $c = 0$. The recurrence relation for the coefficients a_n is:

$$
a_n = -\frac{P_{n-1}a_{n-1} + Q_{n-2}a_{n-2}}{n(n-1)},
$$
\n(37)

where P_n and Q_n are functions of the coefficients in the differential equation.

The series solution for $R(r)$ is:

$$
R(r) = a_0 r^c \left(1 + \sum_{n=1}^{\infty} \frac{a_n}{a_0} r^n \right)
$$
\n(38)

Substitute the Frobenius series into the simplified r -dependent differential equation:

$$
2r \cdot (r - b(r))^2 \sum_{n=0}^{\infty} a_n (n + c)(n + c - 1)r^{n + c - 2} - \sum_{n=0}^{\infty} a_n (n + c) \left[2b(r) \cdot (r - b(r))^2 + 3b(r) \cdot (r - b(r)) \right] r^{n + c - 1} + \sum_{n=0}^{\infty} a_n \left[m^2 \cdot (r - b(r)) + 2r^2 \cdot (w - m\omega(r, \theta))^2 \right] r^{n + c} = 0.
$$

Group powers of r, while the lowest power of r corresponds to $c - 2$. The leading-order term is:

$$
2r \cdot (r - b(r))^2 \cdot a_0 c(c - 1)r^{c-2}
$$
\n(39)

The coefficient of the leading term becomes: $2c(c-1) \cdot \frac{r_0^4}{r^2}$, and balance the terms at r^{c-2} , ignoring higher-order a_n terms, to obtain the indicial equation: $2c(c-1) \cdot r_0^4 = 0$. For the solution to the indicial equation: $c = 0$ and $c = 1$.

- $c = 0$: Corresponds to finite behavior near $r = 0$.
- $c = 1$: May correspond to divergent behavior near $r = 0$, depending on the context.

Choose $c = 0$ to ensure $R(r)$ remains finite at $r = 0$, and ubstitute the series with $c = 0$ into the full equation. Group terms by powers of r^n . For $n \geq 1$, derive the recurrence relation:

$$
a_n n(n-1)r^{n-2} - a_{n-1}P(n)r^{n-1} + a_{n-2}Q(n)r^n = 0
$$
\n(40)

where $P(n)$ and $Q(n)$ depend on the coefficients from $b(r)$ and the original equation. Solve for a_n :

$$
a_n = -\frac{P(n-1)a_{n-1} + Q(n-2)a_{n-2}}{n(n-1)}
$$
\n(41)

The final Frobenius series solution is:

$$
R(r) = a_0 \left(1 + \sum_{n=1}^{\infty} a_n r^n \right) \tag{42}
$$

where a_0 is determined by normalization or physical constraints, **Behavior near** $r = 0$: Choose $c = 0$ to ensure $R(r)$ does not diverge and **Behavior as** $r \to \infty$: Ensure convergence of the series, restricting higher-order a_n terms if necessary.

ANGULAR EQUATION

The θ -dependent equation is:

$$
S(\theta) \left[\csc^2(\theta) + 1 \right] - \epsilon \frac{\partial S(\theta)}{\partial \theta} \left[-\frac{2(\theta - \pi/4)}{\sigma_\theta^2} - \frac{2(\theta - 3\pi/4)}{\sigma_\theta^2} \right] = 0 \tag{43}
$$

Rewriting in standard first-order linear differential equation form:

$$
\frac{\partial S(\theta)}{\partial \theta} + P(\theta)S(\theta) = 0\tag{44}
$$

where:

$$
P(\theta) = -\frac{\csc^2(\theta) + 1}{\epsilon} \left[-\frac{2(\theta - \pi/4)}{\sigma_\theta^2} - \frac{2(\theta - 3\pi/4)}{\sigma_\theta^2} \right]
$$
(45)

using the integrating factor $\mu(\theta)$ is: $\mu(\theta) = \exp \left(\int P(\theta) d\theta \right)$. Substituting $P(\theta)$:

$$
\mu(\theta) = \exp\bigg(-\int \frac{\csc^2(\theta) + 1}{\epsilon} \bigg[-\frac{2(\theta - \pi/4)}{\sigma_\theta^2} - \frac{2(\theta - 3\pi/4)}{\sigma_\theta^2}\bigg] d\theta\bigg). \tag{46}
$$

After some more simplification of denominator, the integrating factor becomes:

$$
\mu(\theta) = \exp\left(-\int \frac{\csc^2(\theta) + 1}{\epsilon} \cdot \frac{4\theta - \pi}{\sigma_\theta^2} d\theta\right)
$$
(47)

After calculating the integrating factor: $S(\theta) = \frac{C}{\mu(\theta)}$, where C is the constant of integration. Substitute the explicit form of $\mu(\theta)$ into this solution. The boundary conditions depend on the physical problem:

- Behavior near $\theta = \pi/4$: Ensure $S(\theta)$ remains finite.
- Behavior near $\theta = 3\pi/4$: Ensure $S(\theta)$ remains finite or decays appropriately.

To ensure finite behavior at both boundaries, the exponential term in $\mu(\theta)$ must not diverge.

Exploring Boundary Conditions for the θ -Dependent Equation

To ensure the solution $S(\theta)$ is physically meaningful, boundary conditions at $\theta = \pi/4$ and $\theta = 3\pi/4$ must be carefully handled. These conditions influence the behavior of $S(\theta)$ and the integrating factor $\mu(\theta)$.

The θ -dependent equation is:

$$
\frac{\partial S(\theta)}{\partial \theta} + P(\theta)S(\theta) = 0,
$$

where:

$$
P(\theta) = \frac{\csc^2(\theta) + 1}{\epsilon} \cdot \frac{4\theta - \pi}{\sigma_\theta^2}.
$$

The solution is:

$$
S(\theta) = \frac{C}{\mu(\theta)},
$$

with the integrating factor:

$$
\mu(\theta) = \exp\bigg(\int P(\theta) \, d\theta\bigg).
$$

Boundary Conditions

1. Behavior at $\theta = \pi/4$

At $\theta = \pi/4$, the Gaussian term in $h(r, \theta)$ has a peak:

$$
\exp\left(-\frac{(\theta-\pi/4)^2}{\sigma_\theta^2}\right).
$$

The factor $P(\theta)$ near $\theta = \pi/4$ becomes:

$$
P(\theta) \approx \frac{\csc^2(\pi/4) + 1}{\epsilon} \cdot \frac{4(\pi/4) - \pi}{\sigma_\theta^2}
$$

.

Simplify:

$$
P(\theta) \approx -\frac{2}{\epsilon} \cdot \frac{-\pi}{\sigma_{\theta}^2} = \frac{2\pi}{\epsilon \sigma_{\theta}^2}.
$$

The integrating factor is:

$$
\mu(\theta) = \exp\left(\int \frac{2\pi}{\epsilon \sigma_{\theta}^2} d\theta\right) = \exp\left(\frac{2\pi\theta}{\epsilon \sigma_{\theta}^2}\right).
$$

This grows exponentially for large θ . To avoid divergence, $S(\theta)$ must decay sufficiently fast. A normalization condition ensures $S(\theta)$ integrates to a finite value over the domain.

2. Behavior at $\theta = 3\pi/4$

Similarly, at $\theta = 3\pi/4$, the Gaussian term in $h(r, \theta)$ peaks:

$$
\exp\left(-\frac{(\theta-3\pi/4)^2}{\sigma_\theta^2}\right).
$$

Near $\theta = 3\pi/4$, $P(\theta)$ becomes:

$$
P(\theta) \approx \frac{\csc^2(3\pi/4) + 1}{\epsilon} \cdot \frac{4(3\pi/4) - \pi}{\sigma_\theta^2}.
$$

Simplify:

$$
P(\theta) \approx \frac{2\pi}{\epsilon \sigma_{\theta}^2}.
$$

As before, the integrating factor grows exponentially:

$$
\mu(\theta) = \exp\left(\frac{2\pi\theta}{\epsilon\sigma_{\theta}^2}\right).
$$

To avoid divergence, $S(\theta)$ must decay sufficiently fast. This requires normalization of $S(\theta)$.

General Behavior

For θ near $\pi/4$ or $3\pi/4$, the equation simplifies due to the dominance of the Gaussian terms in $h(r, \theta)$. The solution $S(\theta)$ decays exponentially due to the integrating factor $\mu(\theta)$, ensuring normalizability.

Normalization Condition

The solution $S(\theta)$ must satisfy:

$$
\int_{\pi/4}^{3\pi/4} S(\theta)^2 d\theta = 1.
$$

Substitute $S(\theta) = \frac{C}{\mu(\theta)}$:

$$
C^2 \int_{\pi/4}^{3\pi/4} \frac{1}{\mu(\theta)^2} \, d\theta = 1.
$$

Normalize C to satisfy this condition.

Computing the Normalized Solution for $S(\theta)$

Recap of the Solution

The solution to the θ -dependent equation is: $S(\theta) = \frac{C}{\mu(\theta)}$, where the integrating factor is: $\mu(\theta) = \exp\left(\int P(\theta) d\theta\right)$, and the coefficient $P(\theta)$ is: $P(\theta) = \frac{\csc^2(\theta) + 1}{\epsilon}$ $\frac{(\theta)+1}{\epsilon}\cdot\frac{4\theta-\pi}{\sigma_{\theta}^2}$ $\frac{\theta-\pi}{\sigma_{\theta}^2}$. The normalized solution satisfies: $\int_{\pi/4}^{3\pi/4} S(\theta)^2 d\theta = 1$.

Substitute $P(\theta)$ into the expression for $\mu(\theta)$:

$$
\mu(\theta) = \exp\left(-\frac{\sigma_{\theta}^2}{\epsilon} \int \frac{\csc^2(\theta) + 1}{4\theta - \pi} d\theta\right).
$$

This integral must generally be evaluated numerically or symbolically, depending on the parameters σ_{θ} and ϵ .

V. CONCLUSION

This study explored the impact of incorporating cosmic strings into the geometry of a rotating traversable wormhole. By extending the foundations laid by Edward Teo and Sung-Won Kim, the modified metric introduced in this work provides new insights into the interplay between cosmic string perturbations and wormhole stability.

The key contributions of this work are summarized as follows:

• Metric Modification: Anistropic deformations in the spacetime can be observed through modeling a rotating traversable wormhole using two cosmic strings through localized Gaussian perturbations. Radial and the Angular distributions influence concentrated around specific regions which can be seen in one of the plots.

- Null Energy Condition Analysis: NEC violations at the throat are amplified in the presence of cosmic strings that satisfies the need for exotic matter at the throat of traversable wormholes.
- Scalar Field Dynamics: Using the modified metric, this paper involved the derivation and the calculation of scalar wave equation.

In conclusion, the cosmic strings influence the geometric properties of wormholes. The localized perturbations alter the angular metric components. This study might make it possible to extend this work to explore the gravitational wave signatures using similar theoretical model.

VI. APPENDIX: COMPUTATIONS

In this work, the Mathematica package OGRe (Object-Oriented General Relativity) by Barak Shoshany was extensively used to compute the metric components, Ricci tensor components, Null Energy Condition (NEC), and the scalar wave equation in the modified worm-hole spacetime.^{[\[23\]](#page-28-15)} OGRe provides a powerful and flexible object-oriented framework for performing symbolic tensor calculations, specifically tailored for general relativity and theoretical physics research.[\[10\]](#page-27-7)

OGRe is an open-source Mathematica package that simplifies complex tensor calculations through an intuitive object-oriented approach. The package offers several advantages.

The following computations were carried out using OGRe:

- The spacetime metric components and their modifications due to cosmic strings and rotation.
- The Ricci tensor and the Null Energy Condition (NEC) to validate the physical consistency of the spacetime.
- The scalar wave equation, derived and analyzed in the modified spacetime geometry.

A. NEC full form

Term 1:

$$
-\frac{4b_0^3\epsilon(r-r_0)(1-\epsilon\sin(\theta))^3\exp\left(\frac{(r-r_0)^2}{\sigma_r^2}+\frac{(\theta-\pi/4)^2}{\sigma_\theta^2}-\epsilon\right)}{r^2\sigma_r^2\left(-b_0+r+b_0\epsilon\sin(\theta)\right)^2}.
$$

$$
-\frac{2b(r,\theta)^3\left(3\frac{\partial K(r,\theta)}{\partial r}+2r\frac{\partial^2 K(r,\theta)}{\partial r^2}\right)}{r^2(r-b(r,\theta))^2f(r,\theta)K(r,\theta)}.
$$

Term 3:

Term 2:

$$
-\frac{b(r,\theta)^3\left(4N(r,\theta)^2+r^2\left(\frac{\partial N(r,\theta)}{\partial r}\right)^2+rN(r,\theta)\left(\frac{\partial N(r,\theta)}{\partial r}-2r\frac{\partial^2 N(r,\theta)}{\partial r^2}\right)\right)}{r^3(r-b(r,\theta))^2f(r,\theta)K(r,\theta)N(r,\theta)^2}
$$

.

Term 4:

$$
\frac{2b(r,\theta)^2\left[\left(-10+\frac{\partial b(r,\theta)}{\partial r}\right)\frac{\partial f(r,\theta)}{\partial r}-6r\frac{\partial^2 f(r,\theta)}{\partial r^2}\right]}{r(r-b(r,\theta))^2f(r,\theta)}.
$$

Term 5:

$$
\frac{2b(r,\theta)^2\left[\left(-10+\frac{\partial b(r,\theta)}{\partial r}\right)\frac{\partial K(r,\theta)}{\partial r}-6r\frac{\partial^2 K(r,\theta)}{\partial r^2}\right]}{r(r-b(r,\theta))^2K(r,\theta)}.
$$

Term 6:

$$
\frac{2b(r,\theta)\frac{\partial b(r,\theta)}{\partial \theta}\frac{\partial f(r,\theta)}{\partial \theta}}{r^2(r-b(r,\theta))^2K(r,\theta)^2}.
$$

Term 7:

$$
\frac{b(r,\theta)\frac{\partial b(r,\theta)}{\partial \theta}\frac{\partial K(r,\theta)}{\partial \theta}}{r^2(r-b(r,\theta))^2K(r,\theta)^2}.
$$

Term 8:

$$
\frac{2b(r, \theta) \cot(\theta) \frac{\partial f(r, \theta)}{\partial \theta}}{r^2(r - b(r, \theta))^2 f(r, \theta) K(r, \theta)^2}.
$$

Term 9:

$$
\frac{-4b(r,\theta)\frac{\partial^2 f(r,\theta)}{\partial \theta^2}}{r^2(r-b(r,\theta))^2f(r,\theta)K(r,\theta)^2}.
$$

∂θ

Term 10:

$$
\frac{4\frac{\partial f(r,\theta)}{\partial r}\frac{\partial K(r,\theta)}{\partial r}}{r^2(r-b(r,\theta))f(r,\theta)K(r,\theta)^2}.
$$

Term 11:

$$
\frac{4\frac{\partial f(r,\theta)}{\partial \theta}\frac{\partial K(r,\theta)}{\partial \theta}}{f(r,\theta)K(r,\theta)}.
$$

Term 12:

$$
\frac{2(r - b(r, \theta))K(r, \theta)^2 \frac{\partial N(r, \theta)}{\partial r}}{N(r, \theta)}.
$$

Term 13: $2(r-b(r,\theta))\frac{\partial K(r,\theta)}{\partial \theta}$ $\partial N(r,\theta)$ $K(r, \theta) N(r, \theta)$ Term 14: $\left(\frac{\partial N(r,\theta)}{\partial \theta}\right)^2$ $\frac{\partial}{\partial N(r,\theta)^2}$. Term 15: $-2\frac{\partial^2 N(r,\theta)}{\partial \theta^2}$ $\partial \theta^2$ $N(r, \theta)$. Term 16: $2r^2 f(r,\theta)^2 \sin^2(\theta) \left(\frac{\partial \omega(r,\theta)}{\partial \theta}\right)^2$ $N(r, \theta)$ Term 17: $-8f(r, \theta) (1 - \epsilon h(r, \theta))^2$

$$
\frac{-8f(r,\theta)\left(1-\epsilon h(r,\theta)\right)^2\omega(r,\theta)^2\sin(2\theta)\frac{\partial f(r,\theta)}{\partial \theta}}{K(r,\theta)^2N(r,\theta)^2}
$$

∂θ

.

.

.

.

Term 18:

$$
\frac{2\alpha(r,\theta)f(r,\theta)\left(1-\epsilon h(r,\theta)\right)\omega(r,\theta)^2\sin^2(\theta)\left(\frac{\partial\omega(r,\theta)}{\partial\theta}\right)^2}{rK(r,\theta)^3N(r,\theta)^3}.
$$

Term 19:

$$
\frac{2(r - b(r, \theta))\omega(r, \theta)\sin^3(\theta)\frac{\partial f(r, \theta)}{\partial r}\frac{\partial \omega(r, \theta)}{\partial r}}{K(r, \theta)N(r, \theta)^2}.
$$

Term 20:

$$
\frac{2(r - b(r, \theta))\omega(r, \theta)\sin^3(\theta)\frac{\partial^2 \omega(r, \theta)}{\partial r^2}}{N(r, \theta)^2}.
$$

Term 21:

$$
4(r - b(r, \theta))K(r, \theta)\sin^2(\theta)\frac{\partial f(r, \theta)}{\partial \theta}\frac{\partial \omega(r, \theta)}{\partial \theta}.
$$

Term 22:

$$
\frac{2r^2f(r,\theta)^3\sin^3(\theta)\left(\frac{\partial\omega(r,\theta)}{\partial\theta}\right)^2}{K(r,\theta)^2N(r,\theta)^3}.
$$

Term 23:

$$
\frac{-4(r - b(r, \theta))\omega(r, \theta)\sin^3(\theta)\frac{\partial^2 f(r, \theta)}{\partial r^2}}{N(r, \theta)^2}
$$

Term 24:

$$
2(r - b(r, \theta))\omega(r, \theta) \sin^3(\theta) \frac{\partial^2 f(r, \theta)}{\partial \theta^2} \frac{\partial f(r, \theta)}{\partial r}.
$$

Term 25:

$$
\frac{2r^2f(r,\theta)^3\sin^3(\theta)\frac{\partial^3\omega(r,\theta)}{\partial\theta^3}}{K(r,\theta)^2N(r,\theta)^3}.
$$

Term 26:
\n
$$
\frac{2r^2 f(r, \theta)^3 \sin^3(\theta) \frac{\partial \omega(r, \theta)}{\partial r} \frac{\partial \omega(r, \theta)}{\partial \theta}}{K(r, \theta)^2 N(r, \theta)^3}
$$
\nTerm 27:
\n
$$
\frac{2r^2 f(r, \theta)^3 \sin^3(\theta) \frac{\partial^3 \omega(r, \theta)}{\partial r^3}}{K(r, \theta)^2 N(r, \theta)^3}
$$
\nTerm 28:
\n
$$
\frac{2r^2 f(r, \theta)^3 \sin^3(\theta) \frac{\partial^3 \omega(r, \theta)}{\partial \theta} \frac{\partial^2 \omega(r, \theta)}{\partial r^2}}{K(r, \theta)^2 N(r, \theta)^3}
$$
\nTerm 29:
\n
$$
\frac{4r^2 f(r, \theta)^3 \sin^3(\theta) \frac{\partial^2 \omega(r, \theta)}{\partial \theta^2} \frac{\partial \omega(r, \theta)}{\partial r}}{K(r, \theta)^2 N(r, \theta)^3}
$$
\nTerm 30:
\n
$$
\frac{2r^2 f(r, \theta)^3 \sin^3(\theta) \frac{\partial^4 \omega(r, \theta)}{\partial \theta^4}}{K(r, \theta)^2 N(r, \theta)^3}
$$
\nThe final expression of the $R_{\mu\nu}k^{\mu}k^{\nu}$ is the sum of all the 3

The final expression of the $R_{\mu\nu}k$ 30 terms stated above.

- [1] Albert Einstein and Nathan Rosen, The Particle Problem in the General Theory of Relativity, Physical Review, 48, 73–77 (1935).
- [2] Michael S. Morris, Kip S. Thorne, and Ulvi Yurtsever, Wormholes, Time Machines, and the Weak Energy Condition, Phys. Rev. Lett., 61(13), 1446–1449 (1988).
- [3] Matt Visser, Lorentzian wormholes: From Einstein to Hawking, Springer, 1995.
- [4] Anonymous, Article Title Placeholder, CrossRef Listing of Deleted DOIs, (2010).
- [5] A. C. Gutiérrez-Piñeres, N. H. Beltrán, and C. S. López-Monsalvo, Newman-Janis Ansatz for rotating wormholes, *Journal of Physics: Conference Series*, **2081**(1), 012005 (2021).
- [6] Edward Teo, Rotating traversable wormholes, Physical Review D, 58(2), 024014 (1998).
- [7] Thanu Padmanabhan, Gravitation: foundations and frontiers, Cambridge University Press, 2010.
- [8] S. E. Perez Bergliaffa and K. E. Hibberd, On the stress-energy tensor of a rotating wormhole, arXiv:gr-qc/0006041 (2000).
- [9] Sung-Won Kim, Rotating wormhole and scalar perturbation, Il Nuovo Cimento B, 120(10011), 1235–1242 (2006).
- [10] Barak Shoshany, OGRe: An Object-Oriented General Relativity Package for Mathematica, Version 1.7.0 (2021-09-17), GitHub, (2021).
- [11] A. Vilenkin and E. P. S. Shellard, Cosmic Strings and Other Topological Defects, Cambridge University Press, 2000.
- [12] Ramesh Radhakrishnan et al., A Review of Stable, Traversable Wormholes in f(R) Gravity Theories, Symmetry, 16(8), 1007 (2024).
- [13] Michael S. Morris and Kip S. Thorne, Wormholes in space-time and their use for interstellar travel: A tool for teaching general relativity, $Am. J. Phys.$, 56, 395–412 (1988).
- [14] Peter K. F. Kuhfittig, Axially symmetric rotating traversable wormholes, Phys. Rev. D, 67, 064015 (2003).
- [15] Vladimir Dzhunushaliev et al., Rotating Wormholes in Five Dimensions, Phys. Rev. D, 88, 124028 (2013).
- [16] Kirill A. Bronnikov et al., Rotating cylindrical wormholes, *Phys. Rev. D*, **87**, 084060 (2013).
- [17] Martin G. Richarte and Claudio Simeone, More about thin-shell wormholes associated to cosmic strings, Phys. Rev. D, 79, 127502 (2009).
- [18] Gerard Clement, Wormhole cosmic strings, *Phys. Rev. D*, **51**, 6803-6809 (1995).
- [19] Gerard Clement, Flat wormholes from straight cosmic strings, J. Math. Phys., 38, 5807–5819 (1997).
- [20] Rodrigo O. Aros and Nelson Zamorano, A Wormhole at the core of an infinite cosmic string, Phys. Rev. D, 56, 6607–6614 (1997).
- [21] Peter Musgrave and Kayll Lake, Junctions and thin shells in general relativity using computer algebra. 1: The Darmois-Israel formalism, Class. Quant. Grav., 13, 1885–1900 (1996).
- [22] Eric Poisson and Matt Visser, Thin-shell wormholes: Linearization stability, Physical Review D, 52(12), 7318–7321 (1995).
- [23] Sean M. Carroll, Spacetime and Geometry: An Introduction to General Relativity, Cambridge University Press, 2019.
- [24] Edmundo Capelas de Oliveira and José Emílio Maiorino, Analytical Methods in Applied Mathematics, Springer Nature, 2024.
- [25] De-Chang Dai et al., How to form a wormhole, The European Physical Journal C, $80(12)$, $(2020).$
- [26] De-Chang Dai et al., New wormhole solution in de Sitter space, Physical Review D, 98(12), 124026 (2018).
- [27] De-Chang Dai and Dejan Stojkovic, Observing a wormhole, *Physical Review D*, $100(8)$,

083513 (2019).

- [28] John H. Simonetti et al., Sensitive searches for wormholes, Physical Review D, 104(8), L081502 (2021).
- [29] Cosimo Bambi and Dejan Stojkovic, Astrophysical Wormholes, Universe, 7(5), 136 (2021).