ON THE CENTER OF TWO-PARAMETER (v, t)-QUANTUM GROUPS

ZHAOBING FAN AND ZIQI XIN

ABSTRACT. This paper mainly considers the centre of two-parameter quantum group $U_{v,t}$ [FL] of finite type via an analogue of the Harish-Chandra homomorphism. Through combining the connection between one-parameter quantum group case and two-parameter quantum group case, we get the description of centre in the sense of Harish-Chandra homomorphism.

CONTENTS

1.	Introduction	1
2.	The two-parameter quantum algebra $U_{v,t}$	2
3.	The ad-invariant bilinear form	6
4.	The centre of $U_{v,t}$	11
5.	The image of the Harish-Chandra homomorphism of $U_{v,t}(g)$	17
References		24

1. INTRODUCTION

In the one-parameter quantum group case, much work has been done on the centre of quantum groups for the finite-dimensional simple Lie algebras, and also for generalized Kac-Moody algebras. The approach taken in many of these papers is to define a nondegenerate bilinear form on the quantum group which is invariant under the adjoint action. The next step involves constructing an algebra homomorphism ξ called the Harish-Chandra homomorphism and proving the map ξ is injective. The main difficulty lies in determining the image of ξ . In the two-parameter case for type A_n [BKL], a new phenomenon arises: the n odd and even cases behave differently. Additional central elements arise when n is odd. This phenomenon complicates the description in two-parameter quantum groups.

In [FL], the first author and Li provided a noval presentation of the two-parameter quantum algebra $U_{v,t}(\mathfrak{g})$ by a geometric approach, where both parameters v and thave geometric meaning. Moreover, this presentation unifies the various quantum

Date: December 30, 2024.

Key words and phrases. quantum algebra, centre, Harish-Chandra homomorphism.

Z. FAN AND Z. XIN

algebras in literature. By various specialization, one can obtain one-parameter quantum algebras [L], two-parameter quantum algebras [BW], quantum superalgebras [CFYW] and multi-parameter quantum algebras [HPR].

In [FX], they defined a skew-Hopf pairing on the deformed quantum algebra which unifies various quantum algebras in literatures, such as one-parameter quantum algebras [L], two-parameter quantum algebras [BW], quantum superalgebras [CFYW] and multi-parameter quantum algebras [HPR]. This provided a tool to construct the ad-invariant bilinear form on two-parameter quantum groups.

In this paper, first we show the Harish-Chandra homomorphism on $U_{v,t}(g)$ is injective. We adjust the prove in [BKL] to make it doesn't rely on the weight module representation. Then we construct some central elements based on weight representation theory and a bilinear form which is invariant under the adjoint action. Next we give a necessary and sufficient condition to show if there are some other central elements. Different from other's work, we consider the result from the perspective of grading on $U_{v,t}$. Furthermore, we define U_J , a set of subalgebras of $U_{v,t}$ and show the image of Harish-Chandra homomorphism on these subalgebras contains the image of centre on the whole algebra. Finally, we show the existence of central elements on some certain degrees to make the consequence more accurate.

This paper is organized as follows. In Section 2, we recall the definition of twoparameter quantum algebra $U_{v,t}$ from [FL]. Besides, we prove some related properties of Verma module of $U_{v,t}$. In Section 3, we construct the nondegenerate bilinear form on $U_{v,t}$ which is invariant under the adjoint action. By using this bilinear form, we construct some central elements using the weight representation. In Section 4, we define the Harish-Chandra homomorphism ξ on $U_{v,t}$ and show ξ is injective. In section 5, we analyse the centre of two-parameter quantum groups via the connection between one-parameter quantum groups and two parameter quantum groups. By the analysis, we give a necessary and sufficient condition to show if there exist some other central elements. Then we define a set of subalgebras of $U_{v,t}$ and show the image of Harish-Chandra homomorphism on these subalgebras contains the image of centre of the whole algebra and construct some central elements to make the consequence more accurate.

2. The two-parameter quantum algebra $U_{v,t}$

We briefly review the definition of the two-parameter quantum algebra $U_{v,t}$ in [FL]. Given a Cartan datum (I, \cdot) , let $\Omega = (\Omega_{ij})_{i,j \in I}$ be an integer matrix satisfying that

- (a) $\Omega_{ii} \in \mathbb{Z}_{>0}, \ \Omega_{ij} \in \mathbb{Z}_{\leq 0}$ for all $i \neq j \in I$; (b) $\frac{\Omega_{ij} + \Omega_{ji}}{\Omega_{ii}} \in \mathbb{Z}_{\leq 0}$ for all $i \neq j \in I$;
- (c) the greatest common divisor of all Ω_{ii} is equal to 1.

To Ω , we associate the following three bilinear forms on $\mathbb{Z}[I]$.

$$\begin{array}{lll} \langle i,j\rangle &=& \Omega_{ij}, & \forall i,j\in I. \\ [i,j] &=& 2\delta_{ij}\Omega_{ii} - \Omega_{ij}, & \forall i,j\in I. \end{array}$$

$$i \cdot j = \langle i, j \rangle + \langle j, i \rangle, \quad \forall i, j \in I.$$

The matrix Ω is called of symmetric type if $\Omega_{ii} = 1$, $\forall i \in I$. In this case, the associated Cartan datum is of symmetric type. For simplicity, we assume that Ω is of symmetric type.

2.1. The free algebra 'f. For indeterminates v and t, we set $v_i = v^{i \cdot i/2}$ and $t_i = t^{i \cdot i/2}$. Denoted by $v_{\nu} = \prod_i v_i^{\nu_i}$, $t_{\nu} = \prod_i t_i^{\nu_i}$ and $\operatorname{tr}(\nu) = \sum_{i \in I} \nu_i \in \mathbb{N}$, for any $\nu = \sum \nu_i i \in \mathbb{N}[I]$.

Let \mathfrak{f} be the free unital associative algebra over $\mathbb{Q}(v,t)$ generated by the symbols $\theta_i, \forall i \in I$. By setting the degree of the generator θ_i to be i, the algebra \mathfrak{f} becomes an $\mathbb{N}[I]$ -graded algebra. For any $\nu \in \mathbb{N}[I]$, we denote by \mathfrak{f}_{ν} the subspace of all homogenous elements of degree ν . We have $\mathfrak{f} = \bigoplus_{\nu \in \mathbb{N}[I]} \mathfrak{f}_{\nu}$ and denote by |x| the degree of a homogenous element $x \in \mathfrak{f}$.

2.1.1. The tensor product $\mathfrak{f} \otimes \mathfrak{f}$. On the tensor product $\mathfrak{f} \otimes \mathfrak{f}$, we define an associative $\mathbb{Q}(v, t)$ -algebra structure by

$$(x_1 \otimes x_2)(y_1 \otimes y_2) = v^{|y_1| \cdot |x_2|} t^{\langle |y_1|, |x_2| \rangle - \langle |x_2|, |y_1| \rangle} x_1 y_1 \otimes x_2 y_2,$$

for homogeneous elements x_1, x_2, y_1 and y_2 in \mathcal{F} . It is associative since the forms \langle, \rangle and " \cdot " are bilinear.

Let $r: \mathfrak{f} \to \mathfrak{f} \otimes \mathfrak{f}$ be the $\mathbb{Q}(v, t)$ -algebra homomorphism such that

$$r(\theta_i) = \theta_i \otimes 1 + 1 \otimes \theta_i$$
, for all $i \in I$.

Proposition 2.1. [FL, Proposition 13] There is a unique symmetric bilinear form (,) on \mathfrak{f} with values in $\mathbb{Q}(v,t)$ such that

(a) (1,1) = 1;(b) $(\theta_i, \theta_j) = \delta_{ij} \frac{1}{1-v_i^{-2}}, \text{ for all } i, j \in I;$ (c) $(x, y'y'') = (r(x), y' \otimes y''), \text{ for all } x, y', y'' \in \mathfrak{f};$ (d) $(x'x'', y) = (x' \otimes x'', r(y)), \text{ for all } x', x'', y \in \mathfrak{f}.$

Here the bilinear form on $\mathfrak{f} \otimes \mathfrak{f}$ is defined by

$$(x_1 \otimes x_2, y_1 \otimes y_2) = t^{2[|x_1|, |x_2|]}(x_1, y_1)(x_2, y_2).$$

2.1.2. The maps r_i and ir. For any $i \in I$, let $r_i(\text{resp. }_ir) : \mathfrak{f} \to \mathfrak{f}$ be the unique linear map satisfying the following properties.

$$\begin{aligned} r_i(1) &= 0, \ r_i(\theta_j) = \delta_{ij}, \ \forall j \in I \text{ and } r_i(xy) = v^{i \cdot |y|} t^{\langle |y|,i\rangle - \langle i,|y| \rangle} r_i(x)y + xr_i(y); \\ ir(1) &= 0, \ ir(\theta_j) = \delta_{ij}, \ \forall j \in I \text{ and } ir(xy) = ir(x)y + v^{i \cdot |x|} t^{\langle i,|x| \rangle - \langle |x|,i\rangle} x_i r(y). \end{aligned}$$

By an induction on |x|, we can show that $r(x) = r_i(x) \otimes \theta_i$ (resp. $r(x) = \theta_i \otimes i r(x)$) plus other terms.

2.1.3. Quantum serre relations. Let \mathfrak{J} be the radical of the bilinear form (-, -). It is clear that \mathfrak{J} is a two-sided ideal of \mathfrak{f} . Denote the quotient algebra of \mathfrak{f} by

 $\mathfrak{f} = \mathfrak{f}/\mathfrak{J}.$

Recall the quantum integers from [FL]. For any $n \in \mathbb{N}$, we have

$$[n]_{v,t} = \frac{(vt)^n - (vt^{-1})^{-n}}{vt - (vt^{-1})^{-1}}, \qquad [n]_{v,t}^! = \prod_{k=1}^n [k]_{v,t}.$$

Denote by

$$\theta_i^{(n)} = \frac{\theta_i^n}{[n]_{v_i, t_i}^!}.$$

Proposition 2.2. [FL, Proposition 14] The generators θ_i of \mathfrak{f} satisfy the following identities.

$$\sum_{\substack{p+p'=1-2\frac{i\cdot j}{i\cdot i}}} (-1)^p t_i^{-p(p'-2\frac{\langle i,j\rangle}{i\cdot i}+2\frac{\langle j,i\rangle}{i\cdot i})} \theta_i^{(p)} \theta_j \theta_i^{(p')} = 0, \quad \forall i \neq j \in I.$$

2.2. The two-parameter quantum algebra $U_{v,t}$. By Drinfeld double construction, we get the following presentation of the entire two-parameter quantum algebra $U_{v,t}$, generated by symbols $E_i, F_i, K_i^{\pm 1}, K_i^{\prime \pm 1}, \forall i \in I$, and subjects to the following relations.

$$\begin{array}{ll} (R1) & K_{i}^{\pm 1}K_{i}^{\mp 1} = K_{i}^{\prime \pm 1}K_{i}^{\prime \mp 1} = 1. \\ (R2) & K_{i}E_{j}K_{i}^{-1} = v^{i\cdot j}t^{\langle j,i\rangle - \langle i,j\rangle}E_{j}, \quad K_{i}^{\prime}E_{j}K_{i}^{\prime - 1} = v^{-i\cdot j}t^{\langle j,i\rangle - \langle i,j\rangle}E_{j}, \\ & K_{i}^{\prime}F_{j}K_{i}^{\prime - 1} = v^{i\cdot j}t^{\langle i,j\rangle - \langle j,i\rangle}F_{j}, \quad K_{i}F_{j}K_{i}^{-1} = v^{-i\cdot j}t^{\langle i,j\rangle - \langle j,i\rangle}F_{j}. \\ (R3) & E_{i}F_{j} - F_{j}E_{i} = \delta_{ij}\frac{K_{i} - K_{i}^{\prime}}{v_{i} - v_{i}^{-1}}. \\ (R4) & \sum_{p+p^{\prime}=1-2\frac{i\cdot j}{i\cdot i}}(-1)^{p}t_{i}^{-p(p^{\prime}-2\frac{\langle i,j\rangle}{i\cdot i}+2\frac{\langle j,i\rangle}{i\cdot i})}E_{i}^{(p)}E_{j}E_{i}^{(p^{\prime})} = 0, \quad \text{if } i \neq j, \\ & \sum_{p+p^{\prime}=1-2\frac{i\cdot j}{i\cdot i}}(-1)^{p}t_{i}^{-p(p^{\prime}-2\frac{\langle i,j\rangle}{i\cdot i}+2\frac{\langle j,i\rangle}{i\cdot i})}F_{i}^{(p^{\prime})}F_{j}F_{i}^{(p)} = 0, \quad \text{if } i \neq j. \end{array}$$

The algebra $U_{v,t}$ has a Hopf algebra structure with the comultiplication Δ , the counit ε and the antipode S given as follows.

$$\begin{split} \Delta(K_i^{\pm 1}) &= K_i^{\pm 1} \otimes K_i^{\pm 1}, \qquad \Delta(K_i'^{\pm 1}) = K_i'^{\pm 1} \otimes K_i'^{\pm 1}, \\ \Delta(E_i) &= E_i \otimes 1 + K_i \otimes E_i, \quad \Delta(F_i) = 1 \otimes F_i + F_i \otimes K_i', \\ \varepsilon(K_i^{\pm 1}) &= \varepsilon(K_i'^{\pm 1}) = 1, \qquad \varepsilon(E_i) = \varepsilon(F_i) = 0, \qquad S(K_i^{\pm 1}) = K_i^{\mp 1}, \\ S(K_i'^{\pm 1}) &= K_i'^{\mp 1}, \qquad S(E_i) = -K_i^{-1}E_i, \qquad S(F_i) = -F_iK_i'^{-1}. \end{split}$$

Let $U_{v,t}^+$ (resp. $U_{v,t}^-$) be a subalgebra of $U_{v,t}$ generated by E_i (resp. F_i).

2.3. The module of $U_{v,t}$. A $U_{v,t}$ -module M is called weight module if it admits a decomposition $M = \bigoplus_{\lambda \in \mathbb{N}[I]} M_{\lambda}$ of vector spaces such that

$$M_{\lambda} = \{ m \in M | K_i m = v^{i \cdot \lambda} c_{i,\lambda} m, K'_i m = v^{-i \cdot \lambda} c_{i,\lambda} m, \forall i \in I \},\$$

where

$$c_{i,\lambda} = t^{\langle \lambda, i \rangle - \langle i, \lambda \rangle}$$

let Λ and Q be the weight lattice and the root lattice, respectively. And Let Λ^+ denote the dominant weights.

Fix $\lambda \in \Lambda$ and assume that $J_{v,t}(\lambda)$ is the left ideal of $U_{v,t}(g)$ generated by E_i , $K_i - v^{i \cdot \lambda} t^{\langle \lambda, i \rangle - \langle i, \lambda \rangle}$, and $K'_i - v^{-i \cdot \lambda} t^{\langle \lambda, i \rangle - \langle i, \lambda \rangle}$ for all $i \in I$. Denote by $M_{v,t}(\lambda) = U_{v,t}(g)/J_{v,t}(\lambda)$ the Verma module of two-parameter quantum group. This is a $U_{v,t}$ module by left multiplication and the highest weight vector $v_{\lambda} = 1 + J_{v,t}(\lambda)$.

By analogue with the one-parameter quantum group case, $M_{v,t}(g)$ has the following proposition.

Proposition 2.3. $M_{v,t}(\lambda)$ satisfies:

- (1) As a $U_{v,t}$ -module, $M_{v,t}(\lambda)$ is free of rank 1, generated by the highest weight vector $v_{\lambda} = 1 + J_{v,t}(\lambda)$;
- (2) Every highest weight $U_{v,t}$ -module with highest weight λ is a homomorphic image of $M_{v,t}(g)$;
- (3) The Verma module $M_{v,t}(g)$ has a unique maximal submodule.

Proof. The proof is almost the same with the one parameter quantum group case. \Box

Theorem 2.4. For $\lambda \in \Lambda^+$, $i \in I$, set $n = \lambda \cdot i + 1$, then there is a homomorphism of $U_{v,t}$ -module

$$f: M_{v,t}(\lambda - ni) \mapsto M_{v,t}(\lambda)$$

with $f(v_{\lambda-ni}) = F_i^n v_{\lambda}$.

Proof. From the universal property of $M_{v,t}(g)$, it is enough to show that $E_j F_i^n v_\lambda = 0$ for all $j \in I$. If $j \neq i$, $E_j F_i^n v_\lambda = 0$ is obvious. We only have to check the case when j = i. The communication relation of E_i and F_i^n is as follows

$$E_i F_i^n - F_i^n E_i = \frac{\sum_{k=0}^{n-1} (F_i^k K_i F_i^{n-1-k} - F_i^k K_i' F_i^{n-1-k})}{v_i - v_i^{-1}}$$
$$= \frac{\sum_{k=0}^{n-1} (v^{-(n-1-k)i \cdot i} F_i^{n-1} K_i - v^{(n-1-k)i \cdot i} F_i^{n-1} K_i')}{v_i - v_i^{-1}}.$$

Since $E_i v_{\lambda} = 0$, we have

$$E_i F_i^n v_{\lambda} = (v_i - v_i^{-1})^{-1} F_i^{n-1} \left(\frac{1 - v_i^{-2n}}{1 - v_i^{-2}} K_i - \frac{1 - v_i^{2n}}{1 - v_i^2} K_i'\right) v_{\lambda}.$$

So $E_i v_{\lambda} = 0$ is equivalent to say

$$\left(\frac{1-v_i^{-2n}}{1-v_i^{-2}}K_i - \frac{1-v_i^{2n}}{1-v_i^2}K_i'\right)v_{\lambda} = 0.$$

From the above equation, we get $n = \lambda \cdot i + 1$.

3. The ad-invariant bilinear form

Proposition 3.1. [FX] There exists a unique bilinear form $(\cdot, \cdot) : U_{v,t}^{\leq} \times U_{v,t}^{\geq} \to \mathbb{Q}(v,t)$ such that for all $x, x' \in U_{v,t}^{\leq}, y, y' \in U_{v,t}^{\geq}, \mu, \nu \in \mathbb{Z}[I]$ and $i, j \in I$, we have

$$(x, yy') = (\Delta(x), y \otimes y'), \ (xx', y) = (x' \otimes x, \Delta(y)), \ (F_i, E_i) = \delta_{ij}(v_i^{-1} - v_i)^{-1}, (K'_{\mu}, K_{\nu}) = v^{\mu \cdot \nu} t^{\langle \nu, \mu \rangle - \langle \mu, \nu \rangle}, \ (K'_{\mu}, E_i) = (F_i, K_{\mu}) = 0,$$

where $K_{\mu} = \prod_{i \in I} K_i^{\mu_i}$, $K'_{\mu} \prod_{i \in I} K_i^{\mu'_i}$ for each $\mu = \sum_{i \in I} \mu_i \alpha_i \in Q$.

The above bilinear form is the skew-Hopf pair of $U_{v,t}(g)$. Since we will frequently use the commutation relation between the positive part and negative part, we need the following lemma.

Lemma 3.2. For each $i \in I$, there exist two linear maps p_i and p'_i on $U^+_{v,t}$, such that for any $x \in U^+_{v,t}$

$$xF_i - F_i x = \frac{p_i(x)K_i - K'_i p'_i(x)}{v_i - v_i^{-1}}$$

and

$$\Delta(x) = x \otimes 1 + \sum_{i \in I} p_i(x) K_i \otimes E_i + M_1$$
$$= K_\mu \otimes x + \sum_{i \in I} E_i K_{\mu-i} \otimes p'_i(x) + M_2$$

where M_1 and M_2 denote the sums of the other terms.

Proof. Consider the definition of co-product, we can get

$$\Delta(U_{\xi}^{+}) = \bigoplus_{0 \le \nu \le \xi} U_{\xi-\nu}^{+} K_{\nu} \otimes U_{\nu}^{+},$$

where \leq is the usual partial order on Q^+ : $\nu \leq \mu$ if $\nu - \mu \in Q^+$. Thus, for each $i \in I$, we can define two linear maps p_i and p'_i on $U^+_{v,t}$, satisfying the following equation

$$\Delta(x) = x \otimes 1 + \sum_{i \in I} p_i(x) K_i \otimes E_i + M_1$$
$$= K_\mu \otimes x + \sum_{i \in I} E_i K_{\mu-i} \otimes p'_i(x) + M_2$$

where M_1 and M_2 denote the sums of the other terms.

On the other hand, for $x \in U_{v,t}^+$ and $i \in I$, we have

$$xF_i - F_i x = \frac{b_i(x)K_i - K'_i b'_i(x)}{v_i - v_i^{-1}},$$

where b_i and b'_i are linear maps on $U^+_{v,t}$.

We have to show the map b_i is equal to p_i and the map b'_i is equal to p'_i .

Assume x_1 and x_2 are two homogeneous elements in $U_{v,t}^+$, we have

$$\frac{x_1 x_2 F_i - F_i x_1 x_2 =}{(x_1 b_i(x_2) + v^{i \cdot |x_2|} t^{\langle i, |x_2| \rangle - \langle |x_2|, i \rangle} b_i(x_1) x_2) K_i - K'_i(v^{i \cdot |x_1|} t^{\langle |x_1|, i \rangle - \langle i, |x_1| \rangle} x_1 b'_i(x_2) + b'_i(x_1) x_2)}{v_i - v_i^{-1}}$$

From the above equation, we get the following relations:

$$b_i(x_1x_2) = x_1b_i(x_2) + v^{i \cdot |x_2|} t^{\langle i, |x_2| \rangle - \langle |x_2|, i \rangle} b_i(x_1)x_2,$$

$$b_i'(x_1x_2) = v^{i \cdot |x_1|} t^{\langle |x_1|, i \rangle - \langle i, |x_1| \rangle} x_1 b_i'(x_2) + b_i'(x_1)x_2.$$

On the other hand,

$$\Delta(x_1 x_2) = ((v^{i \cdot |x_2|} t^{\langle i, |x_2| \rangle - \langle |x_2|, i \rangle} p_i(x_1) x_2 + x_1 p_i(x_2)) K_i \otimes E_i) \oplus M,$$

where M denotes the sum of the other terms.

So we have the following relation:

$$p_i(x_1x_2) = x_1p_i(x_2) + v^{i \cdot |x_2|} t^{\langle i, |x_2| \rangle - \langle |x_2|, i \rangle} p_i(x_1) x_2.$$

Comparing $b_i(x_1x_2)$ with $p_i(x_1x_2)$, we conclude $b_i(x) = p_i(x)$ for all $x \in U_{v,t}^+$. Similarly,

$$\Delta(x_1 x_2) = (E_i K_{|x_1| + |x_2| - i} \otimes v^{i \cdot |x_1|} t^{\langle |x_1|, i \rangle - \langle i, |x_1| \rangle} x_1 p'_i(x_2) + p'_i(x_1) x_2) \oplus M',$$

where M' denotes the sum of the other terms.

So we have the following relation:

$$p'_i(x_1x_2) = v^{i \cdot |x_1|} t^{\langle |x_1|, i \rangle - \langle i, |x_1| \rangle} x_1 p'_i(x_2) + p'_i(x_1) x_2.$$

Comparing $b'_i(x_1x_2)$ with $p'_i(x_1x_2)$, we conclude $b'_i(x) = p'_i(x)$ for all $x \in U^+_{v,t}$.

After the above discussion, we have the map b_i is equal to p_i and the map b'_i is equal to p'_i .

Parallel to the positive part, there are also two linear maps defined on the negative part. For any $i \in I$, by analogue with $U_{v,t}^+$, denote by a_i and a'_i the linear maps on $U_{v,t}^-$, such that

$$x^{-}E_{i} - E_{i}x^{-} = \frac{a_{i}(x)K_{i}' - K_{i}a_{i}'(x)}{v_{i} - v_{i}^{-1}}$$

for any $x^- \in U_{v,t}^-$.

Besides, the maps a_i and a'_i also satisfy

$$\Delta(x^{-}) = 1 \otimes x^{-} + \sum_{i \in I} F_i \otimes a_i(x^{-})K'_i + M'_1$$

= $x^{-} \otimes K'_{\mu} + \sum_{i \in I} a'_i(x^{-}) \otimes F_iK'_{\mu-i} + M'_2.$

where M'_1 and M'_2 denote the sums of the other terms.

It is well-known that $U_{v,t}$ can act on itself by adjoint representation. Recall the adjoint representation as follows,

$$ad(u)m = \sum_{(u)} u_{(1)}mS(u_{(2)})$$

where $u, m \in U_{v,t}$ and $\Delta(u) = \Sigma_{(u)} u_{(1)} \bigotimes u_{(2)}$.

It is evident from the triangular decomposition that $U_{v,t}$ has the decomposition

$$\bigoplus_{\mu,\nu\in Q^+} (U^-_{-\nu}K'_{-\nu})\otimes (U^0)\otimes (U^+_{\mu})\to U_{v,t}.$$

Recall the skew-Hopf pair (,) of $U_{v,t}$ and the above decomposition allows us to define the following bilinear form.

Definition 3.3. Set ρ the half sum of positive roots. Define

$$\langle yK'_{-\nu}K'_{\eta}K_{\phi}x, y_1K'_{-\nu_1}K'_{\eta_1}K_{\phi_1}x_1 \rangle = (y, x_1)(y_1, x)(K'_{\eta}, K_{\phi_1})(K'_{\eta_1}, K_{\phi})v^{2\rho \cdot \nu}$$

for all $x \in U^+_{\mu}$, $x_1 \in U^+_{\mu_1}$, $y \in U^-_{\nu}$, $y_1 \in U^-_{\nu_1}$, $\nu, \nu_1, \mu, \mu_1 \in Q^+$ and all $\eta, \eta_1, \phi, \phi_1 \in Q$. Extend it to the whole algebra $U_{v,t}$, we get the bilinear form $\langle \cdot | \cdot \rangle \colon U_{v,t} \times U_{v,t} \to \mathbb{Q}(v,t)$.

Form the above definition and the non-degeneracy of skew-Hopf pair, we can conclude the following lemma.

Lemma 3.4. Let $\mu, \mu_1, \nu, \nu_1 \in Q^+$. Then

$$\langle U^{-}_{-\nu}U^{0}U^{+}_{\mu}|U^{-}_{-\nu_{1}}U^{0}U^{+}_{\mu_{1}}\rangle = 0$$

unless $\nu = \mu_1$ and $\mu = \nu_1$.

Proposition 3.5. The bilinear form is ad-invariant on $U_{v,t}$. That is, we have

$$\langle ad(u)u_1|u_2\rangle = \langle u_1|ad(S(u))u_2\rangle$$

for all $u, u_1, u_2 \in U_{v,t}$.

Proof. We only have to prove that the bilinear form is ad-invariant when u is one of the generators E_i, F_i, K_i, K'_i for all $i \in I$. And for convenience, assume u_1 and u_2 have the following form,

$$u_{1} = yK_{\nu}^{-1}K_{\eta}'K_{\phi}x$$
$$u_{2} = y_{1}K_{\nu_{1}}^{-1}K_{\eta_{1}}'K_{\phi_{1}}x_{1}$$

where $|y| = -\nu$, $|y_1| = -\nu_1$, $|x| = \mu$ and $|x_1| = \mu_1$.

When $u = K_i$, we have

$$ad(K_i)u_1 = K_i u_1 K_i^{-1} = v^{i \cdot (\mu - \nu)} t^{\langle \nu - \mu, i \rangle - \langle i, \nu - \mu \rangle} u_1$$

So we get

$$\langle ad(K_i)u_1|u_2\rangle = v^{i\cdot(\mu-\nu)}t^{\langle\nu-\mu,i\rangle-\langle i,\nu-\mu\rangle}\langle u_1|u_2\rangle.$$

Besides, we have

$$ad(S(K_i))u_2 = K_i^{-1}u_2K_i = v^{i \cdot (-\mu_1 + \nu_1)}t^{\langle -\nu_1 + \mu_1, i \rangle - \langle i, -\nu_1 + \mu_1 \rangle}u_2,$$

so we get

$$\langle u_1 | ad(S(K_i))u_2 \rangle = v^{i \cdot (-\mu_1 + \nu_1)} t^{\langle -\nu_1 + \mu_1, i \rangle - \langle i, -\nu_1 + \mu_1 \rangle} \langle u_1 | u_2 \rangle$$

If $\langle u_1 | u_2 \rangle = 0$, it is obvious to get the conclusion.

If $\langle u_1|u_2\rangle \neq 0$, we have $\nu - \mu = \mu_1 - \nu_1$ from the lemma 3.4. So we can conclude $\langle ad(K_i)u_1|u_2\rangle = \langle u_1|ad(S(K_i))u_2\rangle$.

When $u = F_i$, we have

$$ad(F_i)u_1 = -u_1F_iK_i^{\prime-1} + F_iu_1K_i^{\prime-1}.$$

From Lemma 3.2, we get

$$xF_i - F_i x = \frac{p_i(x)K_i - K'_i p'_i(x)}{v_i - v_i^{-1}}$$

for any $x \in U_{v,t}^+$. Then

$$\begin{split} ad(F_{i})u_{1} &= u_{1}S(F_{i}) + F_{i}u_{1}S(K_{i}') = -u_{1}F_{i}K_{i}'^{-1} + F_{i}u_{1}K_{i}'^{-1} \\ &= -yK_{\nu}'^{-1}K_{\eta}'K_{\phi}xF_{i}K_{i}'^{-1} + F_{i}yK_{\nu}'^{-1}K_{\eta}'K_{\phi}xK_{i}'^{-1} \\ &= -yK_{\nu}'^{-1}K_{\eta}'K_{\phi}(F_{i}x + (v_{i} - v_{i}^{-1})^{-1}(p_{i}(x)K_{i} - K_{i}'p_{i}'(x)))K_{i}'^{-1} + F_{i}yK_{\nu}'^{-1}K_{\eta}'K_{\phi}xK_{i}'^{-1} \\ &= -yK_{\nu}'^{-1}K_{\eta}'K_{\phi}F_{i}xK_{i}'^{-1} + (v_{i} - v_{i}^{-1})^{-1}(yK_{\nu}'^{-1}K_{\eta}'K_{\phi}K_{i}'p_{i}'(x)K_{i}'^{-1}) \\ &- (v_{i} - v_{i}^{-1})^{-1}(yK_{\nu}'^{-1}K_{\eta}'K_{\phi}p_{i}(x)K_{i}K_{i}'^{-1}) + F_{i}yK_{\nu}'^{-1}K_{\eta}'K_{\phi}xK_{i}'^{-1} \\ &= -v^{(-i)\cdot(\mu+\phi+\nu-\eta)}t^{\langle i,\mu+\phi+\eta-\nu\rangle-\langle \mu+\phi+\eta-\nu,i\rangle}yF_{i}K_{\nu+i}'K_{\eta}'K_{\phi}x \\ &+ (v_{i} - v_{i}^{-1})^{-1}v^{(-i)\cdot(\mu-i)}t^{\langle i,\mu-i\rangle-\langle \mu-i,i\rangle}yK_{\nu}'^{-1}K_{\eta}'K_{\phi}p_{i}'(x) \\ &- (v_{i} - v_{i}^{-1})^{-1}v^{(-2i)\cdot(\mu-i)}yK_{\nu}'^{-1}K_{\eta-i}'K_{\phi+i}p_{i}(x) \\ &+ v^{-i\cdot\mu}t^{\langle i,\mu\rangle-\langle \mu,i\rangle}F_{i}yK_{\nu+i}'K_{\eta}'K_{\phi}x \end{split}$$

Now

$$ad(S(F_i))u_2 = ad(-F_iK_i'^{-1})u_2 = -v^{i \cdot (\mu_1 - \nu_1)}t^{\langle i, \nu_1 - \mu_1 \rangle - \langle \nu_1 - \mu_1, i \rangle}ad(F_i)u_2.$$

Similarly, repeat the previous calculation with u_1 replaced by u_2 , we have

$$ad(S(F_{i}))u_{2} = v^{-i \cdot (\phi_{1}+2\nu_{1}-\eta_{1})} t^{\langle i,\phi_{1}+\eta_{1}\rangle-\langle\phi_{1}+\eta_{1},i\rangle} y_{1}F_{i}K_{\nu_{1}+i}^{\prime-1}K_{\eta_{1}}^{\prime}K_{\phi_{1}}x_{1}$$

$$- (v_{i} - v_{i}^{-1})^{-1} v^{-i \cdot (\nu_{1}-i)} t^{\langle i,\nu_{1}-i\rangle-\langle\nu_{1}-i,i\rangle} y_{1}K_{\nu_{1}}^{\prime-1}K_{\eta_{1}}^{\prime}K_{\phi_{1}}p_{i}^{\prime}(x_{1})$$

$$+ (v_{i} - v_{i}^{-1})^{-1} v^{i \cdot (2i-\mu_{1}-\nu_{1})} t^{\langle i,\nu_{1}-\mu_{1}\rangle-\langle\nu_{1}-\mu_{1},i\rangle} y_{1}K_{\nu_{1}}^{\prime-1}K_{\eta_{1}-i}^{\prime}K_{\phi_{1}+i}p_{i}(x_{1})$$

$$- v^{-i \cdot \nu_{1}} t^{\langle i,\nu_{1}\rangle-\langle\nu_{1},i\rangle} F_{i}y_{1}K_{\nu_{1}+i}^{\prime-1}K_{\eta_{1}}^{\prime}K_{\phi_{1}}x_{1}$$

Note that $\langle U_{-\nu}^{-}U^{0}U_{\mu}^{+}|U_{-\nu_{1}}^{-}U^{0}U_{\mu_{1}}^{+}\rangle=0$ unless $\mu_{1}=\nu$ and $\mu=\nu_{1}$. Then $\langle ad(F_{i})u_{1}|u_{2}\rangle$ and $\langle u_{1}|ad(S(F_{i}))u_{2}\rangle$ are non-zero only when either (a) $\nu + i = \mu_{1}$ and $\mu = \nu_{1}$, or (b) $\nu = \mu_{1}$ and $\nu_{1} = \mu - i$. Here, we only prove (a) since (b) can be proved in a similar way.

When $\nu + i = \mu_1$ and $\mu = \nu_1$. From the definition of $\langle \cdot | \cdot \rangle$, we have

$$\begin{aligned} \langle ad(F_{i})u_{1}|u_{2}\rangle &= \langle -v^{(-i)\cdot(\mu+\phi+\nu-\eta)}t^{\langle i,\mu+\phi+\eta-\nu\rangle-\langle\mu+\phi+\eta-\nu,i\rangle}yF_{i}K_{\nu+i}^{\prime-1}K_{\eta}^{\prime}K_{\phi}x|y_{1}K_{\nu_{1}}^{\prime-1}K_{\eta_{1}}^{\prime}K_{\phi_{1}}x_{1}\rangle \\ &+ \langle v^{-i\cdot\mu}t^{\langle i,\mu\rangle-\langle\mu,i\rangle}F_{i}yK_{\nu+i}^{\prime-1}K_{\eta}^{\prime}K_{\phi}x|y_{1}K_{\nu_{1}}^{\prime-1}K_{\eta_{1}}^{\prime}K_{\phi_{1}}x_{1}\rangle \\ &= -v^{(-i)\cdot(\mu+\phi+\nu-\eta)}t^{\langle i,\mu+\phi+\eta-\nu\rangle-\langle\mu+\phi+\eta-\nu,i\rangle} \\ &\times (yF_{i},x_{1})(y_{1},x)(K_{\eta}^{\prime},K_{\phi_{1}})(K_{\eta_{1}}^{\prime},K_{\phi})v^{2\rho\cdot(\nu+i)} \\ &+ v^{-i\cdot\mu}t^{\langle i,\mu\rangle-\langle\mu,i\rangle}(F_{i}y,x)(y_{1},x)(K_{\eta}^{\prime},K_{\phi_{1}})(K_{\eta_{1}}^{\prime},K_{\phi})v^{2\rho\cdot(\nu+i)} \\ &= A \times (x,y_{1})(K_{\phi},K_{\eta_{1}}^{\prime})(K_{\phi_{1}},K_{\eta}^{\prime})v^{2\rho\cdot(\nu+i)}, \end{aligned}$$

where

 $A = -v^{(-i)\cdot(\mu+\phi+\nu-\eta)}t^{\langle i,\mu+\phi+\eta-\nu\rangle-\langle\mu+\phi+\eta-\nu,i\rangle}(yF_i,x) + v^{-i\cdot\mu}t^{\langle i,\mu\rangle-\langle\mu,i\rangle}(F_iy,x).$ Similarly,

$$\langle yK'_{-\nu}K'_{\eta}K_{\phi}x|ad(S(F_i))u_2\rangle = B \times (x,y_1)(K_{\phi},K'_{\eta_1})(K_{\phi_1},K'_{\eta})v^{2\rho\cdot\nu},$$

where

$$B = -(v_i - v_i^{-1})^{-1} v^{(-i) \cdot (\nu_1 - i)} t^{\langle i, \nu_1 - i \rangle - \langle \nu_1 - i, i \rangle} (p'_i(x), y) + (v_i - v_i^{-1})^{-1} v^{(i) \cdot (2i - \mu_1 - \nu_1)} t^{\langle i, \nu_1 - \mu_1 \rangle + \langle \nu_1 - \mu_1, i \rangle} (p_i(x_1, y))$$

Comparing $\langle ad(F_i)u_1|u_2\rangle$ with $\langle u_1|ad(S(F_i))u_2\rangle$, we have $\langle ad(F_i)u_1|u_2\rangle = \langle u_1|ad(S(F_i))u_2\rangle$ for all $i \in I$.

When $u = E_i$. This case is similar to the case when $u = F_i$.

In the rest of this paper, the Cartan matrix (a_{ij}) , where $a_{ij} = i \cdot j$, for $i, j \in I$, is of finite type.

For $\eta, \phi \in Q \times Q$, we define $\chi_{\eta,\phi} : Q \times Q \to \mathbb{Q}(v,t)$ by

$$\chi_{\eta,\phi}(\eta_1,\phi_1) = (K'_{\eta},K_{\phi_1})(K'_{\eta_1},K_{\phi}), \text{ for all } (\eta_1,\phi_1) \in Q \times Q$$

Lemma 3.6. If $\chi_{\eta,\phi} = \chi_{\eta',\phi'}$, then $(\eta,\phi) = (\eta',\phi')$.

Proof. By the definition, we get that for all $i \in I$

$$\chi_{\eta,\phi}(0,\alpha_i) = (K'_{\eta}, K_i) = v^{\eta \cdot i} t^{\langle i,\eta \rangle - \langle \eta,i \rangle},$$

$$\chi_{\eta',\phi'}(0,\alpha_i) = (K'_{\eta'}, K_i) = v^{\eta' \cdot i} t^{\langle i,\eta' \rangle - \langle \eta',i \rangle},$$

$$\chi_{\eta,\phi}(\alpha_i, 0) = (K'_i, K_{\phi}) = v^{i \cdot \phi} t^{\langle \phi,i \rangle - \langle i,\phi \rangle},$$

$$\chi_{\eta',\phi'}(\alpha_i, 0) = (K'_i, K_{\phi'}) = v^{i \cdot \phi'} t^{\langle \phi',i \rangle - \langle i,\phi' \rangle}.$$

Since the Cartan matrix (a_{ij}) , where $a_{ij} = i \cdot j$, for $i, j \in I$, is non-degenerate, we have $\eta = \eta_1$ and $\phi = \phi_1$.

Proposition 3.7. The ad-invariant bilinear form $\langle \cdot | \cdot \rangle$ is non-degenerate on $U_{v,t}$.

Proof. Since for $\mu, \mu_1, \nu, \nu_1 \in Q^+$, $\langle U^-_{-\nu} U^0 U^+_{\mu} | U^-_{-\nu_1} U^0 U^+_{\mu_1} \rangle = 0$ unless $\nu = \mu_1$ and $\mu = \nu_1$, it is sufficient to argue that if $u \in U^-_{-\mu} U^0 U^+_{\nu}$ and $\langle u | v \rangle = 0$ for all $v \in U^-_{-\mu} U^0 U^+_{\nu}$, then u = 0.

For each $\mu \in Q^+$, choose a basis $u_1^{\mu}, u_2^{\mu}, \ldots, u_{d_{\mu}}^{\mu}$, where $d_{\mu} = \dim(U_{\mu}^+)$. Then we can take a dual basis $v_1^{\mu}, v_2^{\mu}, \ldots, v_{d_{\mu}}^{\mu}$ of $U_{-\mu}^-$, i.e. $(v_i^{\mu}, u_j^{\mu}) = \delta_{i,j}$. Then we get a basis of $U_{-\nu}^- U^0 U_{\mu}^+$ as

$$\{(v_i^{\nu} K_{\nu}^{\prime-1}) K_{\eta}^{\prime} K_{\phi} u_j^{\mu} | 1 \le i \le d_{\nu}, \ 1 \le j \le d_{\mu}, \text{ and } \eta, \phi \in Q\}.$$

From the definition of the bilinear form $\langle \cdot | \cdot \rangle$, we obtain

$$\langle (v_i^{\nu} K_{\nu}^{\prime-1}) K_{\eta}^{\prime} K_{\phi} u_j^{\mu} \mid (v_k^{\mu} K_{\mu}^{\prime-1}) K_{\eta_1}^{\prime} K_{\phi_1} u_l^{\nu} \rangle$$

= $(v_i^{\nu}, u_l^{\nu}) (v_k^{\mu}, u_j^{\mu}) (K_{\eta}^{\prime}, K_{\phi_1}) (K_{\eta_1}^{\prime}, K_{\phi}) v^{2\rho \cdot \nu}$
= $\delta_{i,l} \delta_{j,k} (K_{\eta}^{\prime}, K_{\phi_1}) (K_{\eta_1}^{\prime}, K_{\phi}) v^{2\rho \cdot \nu} .$

Now write $u = \sum_{i,j,\eta,\phi} \theta_{i,j,\eta,\phi} (v_i^{\nu} K_{\nu}^{\prime-1}) K_{\eta}^{\prime} K_{\phi} u_j^{\mu}$, and $v = (v_k^{\mu} K_{\mu}^{\prime-1}) K_{\eta_1}^{\prime} K_{\phi_1} u_l^{\nu}$ with $1 \le k \le d_{\mu}$ and $1 \le l \le d_{\nu}$. From the assumption $\langle u | v \rangle = 0$, we have

$$\sum_{\eta,\phi} \theta_{i,j,\eta,\phi}(K'_{\eta}, K_{\phi_1})(K'_{\eta_1}, K_{\phi})v^{2\rho \cdot \nu} = 0,$$

for all $1 \leq k \leq d_{\mu}$ and $1 \leq l \leq d_{\nu}$. That is,

$$\sum_{\eta,\phi} \theta_{i,j,\eta,\phi} \chi_{\eta,\phi} v^{2\rho \cdot \nu} = 0$$

It follows from lemma 3.6 and linear independence of different characters that $\theta_{i,j,\eta,\phi} = 0$. Hence we get $\langle \cdot | \cdot \rangle$ is non-degenerate.

4. The centre of $U_{v,t}$

We denote the centre of $U_{v,t}$ by $Z(U_{v,t})$. And since the set $Z(U_{v,t})$ commutes with K_i , K'_i for all $i \in I$, we have

$$Z(U_{v,t}) \subseteq U_0 = U^0 \bigoplus_{v>0} U^-_{-v} U^0 U^+_v.$$

Besides, we define an algebra automorphism $\gamma^{-\rho}: U^0 \to U^0$ as

$$\gamma^{-\rho}(K_{\eta}K_{\phi}') = v^{-\rho \cdot (\eta-\phi)} t^{\langle \eta+\phi,\rho\rangle-\langle\rho,\eta+\phi\rangle} K_{\eta}K_{\phi}',$$

specially, we have $\gamma^{-\rho}(K_{\eta}K'_{-\eta}) = v^{-2\rho\cdot\eta}K_{\eta}K'_{-\eta}$.

The Harish-Chandra map $\xi : Z \to U^0$ is the composition of the two following maps:

$$\gamma^{-\rho} \circ \pi : U_0 \to U^0 \to U^0$$

where π is the canonical projection.

Theorem 4.1. The Harish-Chandra map $\xi : Z \to U^0$ is an injective algebra homomorphism.

Proof. Let $U_0 = U^0 \oplus K$, where $K = \bigoplus_{\nu>0} U^-_{-\nu} U^0 U^+_{\nu}$. Since K is the two-sided ideal in U_0 which is the kernel of π , and hence of ξ , we get ξ is an algebra homomorphism. Assume $z \in Z(U_{v,t})$ and $\xi(z) = 0$, to prove π is injective, it is enough to show z must be zero. Since $z \in Z(U_{v,t})$ and $\xi(z) = 0$, z must have the form $z = \sum_{\nu>0} z_{\nu}$ with $z_{\nu} \in U^-_{-\nu} U^0 U^+_{\nu}$. If z is not zero, we can choose one $\nu \in Q^+/0$ minimal with the property that $z_{\nu} \neq 0$. Fix $\{x_l\}$ as bases of U^+_{ν} . We can write $z_{\nu} = \bigoplus_{\{x_l\}} \bigoplus_{\eta, \phi \in Q} y_{\eta, \phi, l} K_{\eta} K'_{\phi} x_l$, where $y_{\mu, \phi, l} \in U^-_{-\nu}$. From the fact $\{K_{\eta} K'_{\phi}\}_{\eta, \phi \in Q}$ can form a basis of U^0 and $\{x_l\}$ is a basis of U^+_{ν} , there must exist some x_l and η' , ϕ' such that

$$y_{\eta',\phi',l'}K_{\eta'}K_{\phi'}'x_{l'} \neq 0$$

and ϕ' is the maximal one among the nonzero terms. Since $y_{\eta',\phi',l'} \neq 0$, referring to [FL], lemma 9, there exists some $i \in I$, such that $a_i(y_{\eta',\phi',l'}) \neq 0$. Write

$$0 = E_i z - zE_i = \sum_{\gamma \neq \nu} (E_i z_\gamma - z_\gamma E_i) + E_i z_\nu - z_\nu E_i.$$

Consider all the terms belonging to $U^-_{-(\nu-i)}U^0U^+_{\nu}$ and note that $E_i z_{\nu} - z_{\nu}E_i$ is the only term in $E_i z - zE_i$ involving $U^-_{-(\nu-i)}U^0U^+_{\nu}$. We write

$$E_i z_{\nu} - z_{\nu} E_i$$

= $E_i (\bigoplus_{\{x_l\}} \bigoplus_{\eta, \phi \in Q} y_{\eta, \phi, l} K_\eta K'_\phi x_l) - (\bigoplus_{\{x_l\}} \bigoplus_{\eta, \phi \in Q} y_{\eta, \phi, l} K_\eta K'_\phi x_l) E_i.$

Substitute $x^-E_i - E_i x^- = \frac{a_i(x)K'_i - K_i a'_i(x)}{v_i - v_i^{-1}}$ into the above formula, then we must have the conclusion

$$(v_i - v_i^{-1})^{-1} \bigoplus_{\{x_l\}} \bigoplus_{\eta, \phi \in Q} (a_i(y_{\eta, \phi, l})K'_i - K_i a'_i(y_{\eta, \phi, l}))K_\eta K'_\phi x_l = 0.$$

Since $\{K_{\eta}K'_{\phi}\}_{\eta,\phi\in Q}$ can form a basis of U^0 and $\{x_l\}$ is a basis of U^+_{ν} , we must have $a_i(y_{\eta',\phi',l})K_{\eta'}K'_{\phi'+i}x_{l'}=0$. So $a_i(y_{\eta',\phi',l'})=0$, which is a contradiction.

Define an algebra homomorphism ρ^{λ} : $U^0 \to \mathbb{Q}(v, t)$ by

$$\varrho^{\lambda}(K'_{\eta}K_{\phi}) = v^{\lambda \cdot \phi - \eta} t^{\langle \phi + \eta, \lambda \rangle - \langle \lambda, \phi + \eta \rangle}$$

for all $\lambda \in \Lambda$ and $\eta \times \phi \in Q \times Q$.

Specially, when $\lambda \in Q$, we have

$$\varrho^{\lambda}(K_{i}) = (K_{i}, K_{\lambda}') = v^{\lambda \cdot i} t^{\langle i, \lambda \rangle - \langle \lambda, i \rangle},$$
$$\varrho^{\lambda}(K_{i}') = (K_{\lambda}, K_{i}')^{-1} = v^{-i \cdot \lambda} t^{\langle i, \lambda \rangle - \langle \lambda, i \rangle}.$$

Define an algebra homomorphism $\varrho^{0,\lambda}$: $U^0 \to \mathbb{Q}(v,t)$ by

$$\varrho^{0,\lambda}(K_i'K_j) = v^{2(i+j)\cdot\lambda},$$

for all $\lambda \in \Lambda$ and $i, j \in I$.

We also define an algebra homomorphism $\rho^{\lambda,\mu}: U^0 \to \mathbb{Q}(v,t)$ by

$$\varrho^{\lambda,\mu}(K'_{\eta}K_{\phi}) = \varrho^{\lambda}(K'_{\eta}K_{\phi})\varrho^{0,\mu}(K'_{\eta}K_{\phi}),$$

for all $\lambda \times \mu \in \Lambda \times \Lambda$ and $\eta \times \phi \in Q \times Q$.

Lemma 4.2. Let $u = K'_{\eta}K_{\phi}$, $\eta, \phi \in Q$. If $\varrho^{\lambda,\mu}(u) = 1$ for all $\lambda, \mu \in \Lambda$, then u = 1.

Proof. If $\rho^{0,\lambda}(K'_{\eta}K_{\phi}) = 0$ for all $\lambda \in \Lambda$, then $\eta + \phi = 0$ from the definition of $\rho^{0,\lambda}$. If $\rho^{\lambda,0}(K'_{\eta}K_{\phi}) = 0$ for all $\lambda \in \Lambda$, consider

$$\varrho^{\lambda}(K'_{\eta}K_{\phi}) = v^{\lambda \cdot \phi - \eta} t^{\langle \phi + \eta, \lambda \rangle - \langle \lambda, \phi + \eta \rangle}$$

since $\lambda \cdot (\eta - \phi) = 0$ for all $\lambda \in Q$ and the Cartan matrix (a_{ij}) , where $a_{ij} = i \cdot j$ for $i, j \in I$, is non-degenerate, we can conclude $\eta - \phi = 0$. Remembering $\eta + \phi = 0$, $\eta = \phi = 0$ is concluded.

Lemma 4.3. If $u \in U^0$, $\rho^{\lambda,\mu}(u) = 0$, for all $(\lambda,\mu) \in \Lambda \times \Lambda$, then u = 0.

Proof. Corresponding to each $(\eta, \phi) \in Q \times Q$ is the character on the group $\Lambda \times \Lambda$ defined by

$$(\lambda,\mu)\mapsto \varrho^{\lambda,\mu}(K'_{\eta}K_{\phi}).$$

It follows from the lemma 4.2 that different (η, ϕ) gives to different characters.

Suppose now that $u = \sum \theta_{\eta,\phi} K'_{\eta} K_{\phi}$, where $\theta_{\eta,\phi} \in \mathbb{Q}(v,t)$. By assumption,

$$\sum \theta_{\eta,\phi} \varrho^{\lambda,\mu} (K'_{\eta} K_{\phi}) = 0,$$

for all $(\lambda, \mu) \in \Lambda \times \Lambda$. By the linear independence of different characters, $\theta_{\eta,\phi} = 0$ for all $\eta \times \phi \in Q \times Q$, so u = 0.

Define the Weyl group action on $U^0_{\flat} = \bigoplus_{\eta \in Q} \mathbb{Q}(v, t) K'_{\eta} K_{-\eta}$ as follows:

$$\sigma(K'_{\eta}K_{-\eta}) = K'_{\sigma(\eta)}K_{\sigma(-\eta)}.$$

Theorem 4.4. For $U_{v,t}$, $\varrho^{\sigma(\lambda),\mu}(u) = \varrho^{\lambda,\mu}(\sigma^{-1}(u))$, where $u \in U^0_{\flat}$, $\sigma \in W$ and $\lambda, \mu \in \Lambda$.

Proof. Assume that $u = K'_{\eta}K_{-\eta}$. We only need to check the theorem for each σ_i . Note that $\varrho^{0,\mu}(K'_{\eta}K_{-\eta}) = \varrho^{0,\mu}(\sigma^{-1}(K'_{\eta}K_{-\eta})) = 1$ from the definition of $\varrho^{0,\mu}$. So, we get $\varrho^{\sigma(\lambda),\mu}(u) = \varrho^{\sigma(\lambda),0}(K'_{\eta}K_{-\eta})\varrho^{0,\mu}(K'_{\eta}K_{-\eta}) = \varrho^{\lambda,0}(\sigma^{-1}(K'_{\eta}K_{-\eta}))\varrho^{0,\mu}(\sigma^{-1}(K'_{\eta}K_{-\eta})) = \varrho^{\lambda,\mu}(\sigma^{-1}(u)).$

Lemma 4.5. We set ι as the unit map of $U_{v,t}$ and ϵ the co-unit map of $U_{v,t}$. Then $z \in Z(U_{v,t})$ if and only if $ad(x)z = (\iota \circ \epsilon)(x)z$ for all $u \in U_{v,t}$.

Proof. Let $z \in Z(U_{v,t})$. Then for all $x \in U_{v,t}$, we have

$$ad(x)z = \sum_{(x)} x_{(1)}zS(x_{(2)}) = z\sum_{(x)} x_{(1)}S(x_{(2)}) = (\iota \circ \varepsilon)(x)z.$$

Conversely, assume that $ad(x)z = (\iota \circ \varepsilon)(x)z$ for all $x \in U_{v,t}$. We can consider it on the generators of $U_{v,t}$. Then

$$K_i z K_i^{-1} = ad(K_i) z = (\iota \circ \varepsilon)(K_i) z = z.$$

Similarly, we have $K'_i z K'^{-1}_i = z$. Furthermore, we have

$$0 = (\iota \circ \varepsilon)(E_i)z = ad(E_i)z = E_i z - K_i z K_i^{-1} E_i = E_i z - z E_i$$

and

$$0 = (\iota \circ \varepsilon)(F_i)z = ad(F_i)z = -zF_i(K'_i)^{-1} + F_iz(K'_i)^{-1} = (-zF_i + F_iz)(K'_i)^{-1}.$$

Hence, $z \in Z(U_{v,t})$.

Lemma 4.6. Assume that $\Psi: U^-_{-\mu} \times U^+_{\nu} \to \mathbb{Q}(v,t)$ is a linear map, and let $(\eta, \phi) \in Q \times Q$. Then there exists $u \in U^-_{-v} U^0 U^+_{\mu}$ such that

$$\langle u | (yK_{\mu}^{\prime-1})K_{\eta_1}^{\prime}K_{\mu_1}x \rangle = (K_{\eta_1}^{\prime}, K_{\phi})(K_{\eta}^{\prime}, K_{\phi_1})\Psi(y, x)$$

for all $x \in U_{\nu}^+$, $y \in U_{-\mu}^-$ and $(\eta_1, \phi_1) \in Q \times Q$.

Proof. For each $\mu \in Q^+$, we choose a basis $u_1^{\mu}, u_2^{\mu}, \ldots, u_{d_{\mu}}^{\mu}$ $(d_{\mu} = \dim(U_{\mu}^+))$ of U_{μ}^+ and a dual basis $v_1^{\mu}, v_2^{\mu}, \ldots, v_{d_{\mu}}^{\mu}$ of $U_{-\mu}^-$ such that $(v_i^{\mu}, u_j^{\mu}) = \delta_{i,j}$. If we set

$$u = \sum_{(i,j)} \Psi(v_j^{\mu}, u_i^{\nu}) v_i^{\nu} (K_{\nu}')^{-1} K_{\eta}' K_{\phi} u_j^{\mu} v^{-2\rho \cdot \nu},$$

then it is straightforward to verify that u satisfies the requirement.

We define a $U_{v,t}$ -module structure on the dual space $U_{v,t}^*$ by setting (x.f)(v) = f(S(x)v), for any $u, v \in U_{v,t}, f \in U_{v,t}^*$. We also define a map $\beta : U_{v,t} \to U_{v,t}^*$ by setting

$$\beta(u)(v) = \langle u | v \rangle \quad \forall u, v \in U_{v,t}.$$

Considering that $U_{v,t}$ -module structure on $U_{v,t}$ is given by the adjoint representation. We get β is an injective $U_{v,t}$ -module homomorphism because of the adinvariance and the non-degeneracy of the bilinear form \langle , \rangle .

Definition 4.7. Let M be a finite dimensional module of $U_{v,t}$. For each $m \in M$ and $f \in M^*$, we define an element $C_{f,m} \in U^*_{v,t}$ by $C_{f,m}(v) = f(v.m)$, for any $v \in U_{v,t}$.

Proposition 4.8. Assume that M is a finite $U_{v,t}$ -module, $M = \bigoplus_{\lambda \in wt(M)} M_{\lambda}$. Suppose that the weight set wt(M) of M satisfies: $wt(M) \in Q$. For each $f \in M^*$, $m \in M$, there exist a unique $u \in U_{v,t}$ such that

$$C_{f,m}(v) = \langle u | v \rangle$$
 for all $v \in U_{v,t}$

Proof. Let $m \in M_{\lambda}$, $v = (yK'_{\mu})K'_{\eta_1}K_{\phi_1}x$, where $x \in U^+_{\nu}$, $y \in U^-_{-\mu}$, $(\eta_1, \phi_1) \in Q \times Q$. Then for each $f \in M^*$, we have

$$C_{f,m}(v) = C_{f,m}((yK'_{\mu})K'_{\eta_1}K_{\phi_1}x) = f((yK'_{\mu})K'_{\eta_1}K_{\phi_1}x.m)$$
$$= \varrho^{\nu+\lambda}(K'_{\eta_1}K_{\phi_1})f((yK'_{\mu})x.m).$$

Note that $(y, x) \mapsto f(yx.m)$ is bilinear, and $\rho^{\nu+\lambda}(K'_{\eta_1}) = (K'_{\eta_1}, K_{-\nu-\lambda}), \rho^{\nu+\lambda}(K_{\phi_1}) = (K'_{\nu+\lambda}, K_{\phi_1})$. By lemma 4.6, there exist a unique $u_{\nu\mu}$ such that $C_{f,m}(v) = \langle u_{\nu\mu}, v \rangle$ for all $v \in U^-_{-\mu} U^0 U^+_{\nu}$. Now for arbitrary $v \in U_{v,t}$, we write $v = \sum_{\mu,\nu} v_{\mu\nu}$ where $v_{\mu\nu} \in U^-_{-\mu} U^0 U^+_{\nu}$. Since M is finite-dimensional, there exists a finite set Υ of pair $Q \times Q$, such that

$$C_{f,m}(v) = C_{f,m}(\sum_{(\mu,\nu)\in\Upsilon} v_{\mu\nu}).$$

Setting $u = \sum_{\mu,\nu} u_{\mu\nu}$, we have

$$C_{f,m}(v) = C_{f,m}\left(\sum_{(\mu,\nu)\in\Upsilon} v_{\mu\nu}\right) = \sum_{(\mu,\nu)\in\Upsilon} C_{f,m}(v_{\mu\nu})$$
$$= \sum_{(\mu,\nu)\in\Upsilon} \langle u_{\mu\nu} | v_{\mu\nu} \rangle = \langle u | v \rangle.$$

The uniqueness follows from proposition 4.7.

Assume M is a finite dimensional $U_{v,t}$ -module, and we define a linear map Θ on M by

$$\Theta(m_{\lambda}) = v^{-2\rho \cdot \lambda} m_{\lambda},$$

for all $m \in M_{\lambda}, \lambda \in \Lambda$.

Lemma 4.9. $\Theta u = S^2(u)\Theta$, for all $u \in U_{v,t}$.

Proof. We have only to check it holds for generators E_i, F_i, K_i and K'_i . Also we fix one $m_{\lambda} \in M_{\lambda}$:

 $\begin{array}{l} (1) \text{When } u = K_i. \ S^2(K_i)\Theta(m_{\lambda}) = K_i\Theta(m_{\lambda}) = \Theta K_i(m_{\lambda}). \\ (2) \text{When } u = K'_i. \ S^2(K'_i)\Theta(m_{\lambda}) = K'_i\Theta(m_{\lambda}) = \Theta K'_i(m_{\lambda}). \\ (3) \text{When } u = E_i. \ S^2(E_i)\Theta(m_{\lambda}) = K_i^{-1}E_iK_i\Theta(m_{\lambda}) = v^{i\cdot i}E_i\Theta(m_{\lambda}) = v^{-2\rho\cdot(\lambda+i)}E_im_{\lambda} \\ = \Theta(E_i.m_{\lambda}). \\ (4) \text{When } u = F_i. \ S^2(F_i)\Theta(m_{\lambda}) = K'_iF_iK'_i^{-1}\Theta(m_{\lambda}) = v^{i\cdot i}F_i\Theta(m_{\lambda}) = v^{-2\rho\cdot(\lambda-i)}F_im_{\lambda} \\ = \Theta(F_i.m_{\lambda}). \end{array}$

For $\lambda \in \Lambda$, define $f_{\lambda} \in U_{v,t}^*$ by $f_{\lambda}(u) = tr_{L(\lambda)}(u\Theta)$ for each $u \in U_{v,t}$.

Lemma 4.10. If $\lambda \in \Lambda^+ \cap Q$, then $f_{\lambda} \in Im(\beta)$.

Proof. We have only to find an element $u \in U_{v,t}$ such that $\langle u|v \rangle = f_{\lambda}(v)$, for any $v \in U_{v,t}$. Let $\lambda \in \Lambda^+ \cap Q$ and $k = \dim(L_{\lambda})$. Denote by $\{m_i\}_{i=1,2,\dots,k}$ a basis of L_{λ} and $\{f_i\}_{i=1,2,\dots,k}$ the dual basis of L_{λ}^* . Then by definition we have

$$f_{\lambda}(v) = tr_{L(\lambda)}(v\Theta).$$

Since Θ is invertible, $\{\Theta m_i\}_{i=1,2,\dots,k}$ form a basis of L_{λ}

$$tr_{L(\lambda)}(v\Theta) = \sum_{i=1}^{k} f_i(v\Theta m_i) = \sum_{i=1}^{k} C_{f_i,\Theta m_i}(v) = \sum_{i=1}^{k} \langle u_i | v \rangle = \langle \sum_{i=1}^{k} u_i | v \rangle.$$

Set $u = \sum_{i=1}^{k} u_i$, u is the element we want.

Theorem 4.11. For $\lambda \in \Lambda^+ \cap Q$, $z_{\lambda} = \beta^{-1}(f_{\lambda}) \in Z(U_{v,t})$.

Proof. For all $x \in U_{v,t}$, we have

$$(S^{-1}(x)f_{\lambda})(u) = f_{\lambda}(ad(x)u)$$

= $tr_{L(\lambda)}(\sum_{(x)} x_{(1)}uS(x_{(2)})\Theta)$
= $tr_{L(\lambda)}(u\sum_{(x)} S(x_2)\Theta x_{(1)})$
= $tr_{L(\lambda)}(u\sum_{(x)} S(x_2)S^2(x_{(1)})\Theta)$
= $tr_{L(\lambda)}(uS(\sum_{(x)} S(x_{(1)})x_{(2)})\Theta)$
= $(\iota \circ \epsilon)(x)tr_{L(\lambda)}(u\Theta)$
= $(\iota \circ \epsilon)(x)f_{\lambda}(u).$

Substituting x for $S^{-1}(x)$ in the above and noticing that $\varepsilon \circ S = \varepsilon$, we can conclude

$$x.f_{\lambda} = (\iota \circ \epsilon)(x)f_{\lambda}$$

Now let

$$xf_{\lambda} = x\beta(\beta^{-1}(f_{\lambda})) = \beta(ad(S(x))\beta^{-1}(f_{\lambda})),$$
$$(\iota \circ \epsilon)(x)f_{\lambda} = (\iota \circ \epsilon)(x)\beta(\beta^{-1}(f_{\lambda})) = \beta((\iota \circ \epsilon)(x)\beta^{-1}f_{\lambda}).$$

Since β is an injective map, for all $x \in U_{v,t}$

$$ad(x)(\beta^{-1}(f_{\lambda})) = (\iota \circ \epsilon)(x)\beta^{-1}(f_{\lambda})$$

then we have $\beta^{-1}(f_{\lambda}) \in Z(U_{v,t})$.

Recall that $U^0_{\flat} = \bigoplus_{\eta \in Q} \mathbb{Q}(v, t) K'_{\eta} K_{-\eta}$ and define $(U^0_{\flat})^W = \{ u \in U^0_{\flat} | \sigma(u) = u, \forall \sigma \in W \}$

Theorem 4.12. For $U_{v,t}$, $(U_{\flat}^0)^W \subseteq \xi(Z(U_{v,t}))$.

Proof. Set $z_{\lambda} = \beta^{-1}(f_{\lambda})$ for $\lambda \in \Lambda^+ \cap Q$ and write

$$z_{\lambda} = \sum_{v \ge 0} z_{\lambda,v},$$

and specially,

$$z_{\lambda,0} = \sum_{\eta,\phi\in Q\times Q} \theta_{\eta,\phi} K'_{\eta} K_{\phi},$$

where $z_{\lambda,v} \in U^-_{-v} U^0 U^+_v$ and $\theta_{\eta,\phi} \in \mathbb{Q}(v,t)$. Now for $(\eta_1,\phi_1) \in Q \times Q$, we have

$$\langle z_{\lambda} | K'_{\eta_1} K_{\phi_1} \rangle = \langle z_{\lambda,0} | K'_{\eta_1} K_{\phi_1} \rangle = \sum_{(\eta,\phi)} \theta_{\eta,\phi}(K'_{\eta_1}, K_{\phi})(K'_{\eta}, K_{\phi_1}).$$

On the other hand, from the definition of z_{λ} , we also have

$$\begin{aligned} \langle z_{\lambda} | K'_{\eta_1} K_{\phi_1} \rangle &= \beta(z_{\lambda}) (K'_{\eta_1} K_{\phi_1}) = f_{\lambda} (K'_{\eta_1} K_{\phi_1}) = tr_{L(\lambda)} (K'_{\eta_1} K_{\phi_1} \Theta) \\ &= \sum_{\mu \leq \lambda} v^{-2\rho \cdot \mu} \dim(L(\lambda)_{\mu}) \varrho^{\mu} (K'_{\eta_1} K_{\phi_1}) \\ &= \sum_{\mu \leq \lambda} v^{-2\rho \cdot \mu} \dim(L(\lambda)_{\mu}) (K'_{\eta_1}, K_{-\mu}) (K'_{\mu}, K_{\phi_1}). \end{aligned}$$

Now we get the following equation,

$$\sum_{(\eta,\phi)} \theta_{\eta,\phi} \chi_{\eta,\phi} = \sum_{\mu \le \upsilon} v^{-2\rho \cdot \lambda} \dim(L(\lambda)_{\mu}) \chi_{\mu,-\mu}.$$

Since these different characters are linearly independent, we have (1) $\theta_{\eta,\phi} = \dim(L(\lambda))v^{-2\rho\cdot\lambda}m_{\lambda}$, when $(\eta,\phi) = (\mu,-\mu)$. (2) $\theta_{\eta,\phi} = 0$, otherwise.

So, we have

$$z_{\lambda,0} = \sum_{\mu \le \lambda} v^{-2\rho \cdot \mu} m_{\lambda} \dim(L(\lambda)_{\mu}) K'_{\mu} K_{-\mu}$$

Then, we obtain

$$\xi(z_{\lambda}) = \varrho^{-\rho}(z_{\lambda,0}) = \sum_{\mu \le \lambda} \dim(L(\lambda)_{\mu}) K'_{\mu} K_{-\mu}$$

For $\lambda \in \Lambda^+ \cap Q$, we define

$$av(\lambda) = \frac{1}{|W|} \sum_{\sigma \in W} \sigma(K'_{\lambda}K_{-\lambda}) = \frac{1}{|W|} \sum_{\sigma \in W} K'_{\sigma(\lambda)}K_{-\sigma(\lambda)}$$

For each $\eta \in Q$, there exists a unique $\sigma \in W$ such that $\sigma(\eta) \in \Lambda^+ \cap Q$, so $\{av(\lambda) | \lambda \in \Lambda^+ \cap Q\}$ forms a basis of $(U^0_{\flat})^W$. So we only have to show $av(\lambda) \in \text{Im}(\xi)$. By induction on the height of λ , if $\lambda = 0$, $av(\lambda) = 1 = \xi(1)$. Assume that $\lambda > 0$. Since $\dim L(\lambda)_{\mu} = \dim L(\lambda)_{\sigma(\mu)}$, for all $\sigma \in W$, and $\dim(L(\lambda)_{\lambda}) = 1$, we can obtain

$$\xi(z_{\lambda}) = |W|av(\lambda) + |W| \sum \dim(L(\lambda)_{\mu})av(\mu),$$

where the sum is over μ such that $\mu < \lambda$ and $\mu \in \Lambda^+ \cap Q$. By the induction hypothesis, we get $av(\lambda) \in \text{Im}(\xi)$. So $(U_b^0)^W \subseteq \xi(Z(U_{v,t}))$.

5. The image of the Harish-Chandra homomorphism of $U_{v,t}(g)$

Let us recall the connection between the one-parameter quantum group and the two-parameter quantum group in [FL].

For any $\gamma = (\gamma_1, \gamma_2), \eta = (\eta_1, \eta_2) \in \mathbb{Z}^I \times \mathbb{Z}^I$, we define a bilinear form on $\mathbb{Z}^I \times \mathbb{Z}^I$ by

$$[\gamma, \eta]' = [\gamma_2, \eta_2] - [\gamma_1, \eta_1].$$

The algebra $U_{v,t}$ admits a $\mathbb{Z}^I \times \mathbb{Z}^I$ -grading by defining the degrees of generators as follows,

$$\deg(E_i) = (i, 0), \deg(K_i) = (i, i) = \deg(K'_i)$$
$$\deg(F_i) = (0, i), \deg(K_i^{-1}) = (-i, -i) = \deg(K'_i^{-1}).$$

And in [FL], the authors define a new multiplication "*" on $U_{v,t}$ by

$$x * y = t^{-\lfloor |x|, |y|\rfloor'} xy$$

for any homogeneous elements $x, y \in U_{v,t}$. Since [,]' is a bilinear form, $(U_{v,t}, *)$ is an associative algebra over $\mathbb{Q}(v, t)$. Define the multiplication, denoted by *, on $U_{v,t} \otimes U_{v,t}$ by

$$(x \otimes y) * (x' \otimes y') = x * x' \otimes y * y'.$$

This gives a algebra structure on $U_{v,t} \otimes U_{v,t}$. $(U_{v,t}, *)$ has a Hopf algebra structure with the comultiplication Δ^* , the counit ε^* and the antipode S^* . The image of generators E_i, F_i, K_i and K'_i under the map Δ^* (resp. ε^* and S^*) are the same as the ones under the map Δ (resp. ε and S).

Theorem 5.1. [FL, Theorem 4] If (I, \cdot) is the Cartan datum associated to Ω , then there is a Hopf-algebra isomorphism

$$(U_{v,t}, *, \triangle^*, \varepsilon^*, S^*) \simeq (U_{v,t}(I, \cdot), \cdot, \triangle_1, \varepsilon_1, S_1)$$

sending the generators in $U_{v,t}$ to the respective generators in $U_{v,t}(I, \cdot)$, where $\mathbb{Q}(v, t) \otimes U_v(I, \cdot) \cong (U_{v,t}(I, \cdot), *)$, and $U_v(I, \cdot)$ is the one-parameter case with Cartan datum (I, \cdot) .

In the rest of this section, grading on $U_{v,t}$ refers to the $\mathbb{Z}^I \times \mathbb{Z}^I$ -grading (\cdot, \cdot) . In this case, homogeneous central elements can be defined with degree (η, η) for some $\eta \in \mathbb{Z}^I$ from the fact that central elements commute with K_i and K'_i for all $i \in I$. This grading allows us to consider homogeneous central elements for convenience.

Theorem 5.2. The two following propositions are equivalent :

- (a) There doesn't exist any nonzero $\eta \in Q$, such that $[\eta, i] [i, \eta] = 0$ for all $i \in I$.
- (b) $\xi(Z(U_{v,t})) = (U_{\flat}^{0})^{W}$.

Proof. Assume (b) holds. If there is some nonzero $\eta' \in \mathbb{Z}^I$, such that $[\eta', i] - [i, \eta'] = 0$ for all $i \in I$, then consider the commutation relation of the element $K'_{\eta'}K_{\eta'}$. For all $i \in I$,

$$K'_{\eta'}K_{\eta'}E_{i} = t^{\langle 2\eta',i\rangle - \langle i,2\eta'\rangle}E_{i}K'_{\eta'}K_{\eta'} = t^{[i,2\eta']-[2\eta',i]}E_{i}K'_{\eta'}K_{\eta'} = E_{i}K'_{\eta'}K_{\eta'},$$

$$K'_{\eta'}K_{\eta'}F_{i} = t^{\langle i,2\eta'\rangle - \langle 2\eta',i\rangle}F_{i}K'_{\eta'}K_{\eta'} = t^{[2\eta',i]-[i,2\eta']}F_{i}K'_{\eta'}K_{\eta'} = F_{i}K'_{\eta'}K_{\eta'}.$$

So we have $K'_{n'}K_{n'} \in Z(U_{v,t})$, which is contradictory to (b).

Conversely, assume (a) holds. Since we have proved $(U_{\flat}^{0})^{W} \subseteq \xi(Z(U_{v,t}))$ in theorem 4.12, then $\xi(Z(U_{v,t})) = (U_{\flat}^{0})^{W}$ is equivalent to $(U_{\flat}^{0})^{W} \supseteq \xi(Z(U_{v,t}))$. Parallel to the one parameter case, the images of Harish-Chandra homomorphism of the central elements with degree (0,0) are in $(U_{\flat}^{0})^{W}$. So, to show $(U_{\flat}^{0})^{W} \supseteq \xi(Z(U_{v,t}))$, it is enough to show there is no central element z with degree (η, η) , where $\eta \neq 0$ and η is under the assumption (a). If there exists some homogeneous central element u with degree $|u| = (\eta, \eta), \eta \in \mathbb{Z}^I \setminus \{0\}$, we have

$$u \in \bigoplus_{\alpha \in Q^+} U^-_{(0,\alpha)} U^0_{(\eta - \alpha, \eta - \alpha)} U^+_{(\alpha,0)}.$$

Choose $\gamma \in Q^+$ as the maximal one in u such that $U^-_{(0,\gamma)}U^0_{(\eta-\gamma,\eta-\gamma)}U^+_{(\gamma,0)}$ are nonzero. Also choose a basis $\{x_1^{\gamma}, x_2^{\gamma}, \ldots, x_{d_{\gamma}}^{\gamma}\}$ of $U^-_{(0,\gamma)}$ where d_{γ} denote the dimension of $U^-_{(0,\gamma)}$. From the triangle decomposition, we have

$$u = \sum_{j=1}^{d_{\gamma}} \theta_j x_j^{\gamma} (\sum_{a+b=\eta-\gamma} K_a' K_b U_{(\gamma,0)}^+) + \sum_{\alpha \neq \gamma} U_{(0,\alpha)}^- U_{(\eta-\alpha,\eta-\alpha)}^0 U_{(\alpha,0)}^+$$

where $\theta_j \in \mathbb{Q}(v, t)$. Under the assumption (a), there is some $i \in I$ satisfying $[\eta, i] - [i, \eta] \neq 0$. Then $E_i u = u E_i$ can be expressed as

$$E_{i}\left(\sum_{j=1}^{d_{\gamma}}\theta_{j}x_{j}^{\gamma}\left(\sum_{a+b=\eta-\gamma}K_{a}'K_{b}U_{(\gamma,0)}^{+}\right)+\sum_{\alpha\neq\gamma}U_{(0,\alpha)}^{-}U_{(\eta-\alpha,\eta-\alpha)}^{0}U_{(\alpha,0)}^{+}\right)$$
$$=\left(\sum_{j=1}^{d_{\gamma}}\theta_{j}x_{j}^{\gamma}\left(\sum_{a+b=\eta-\gamma}K_{a}'K_{b}U_{(\gamma,0)}^{+}\right)+\sum_{\alpha\neq\gamma}U_{(0,\alpha)}^{-}U_{(\eta-\alpha,\eta-\alpha)}^{0}U_{(\alpha,0)}^{+}\right)E_{i}.$$

Note that $E_i U^-_{(0,\gamma)} U^0_{(\eta-\gamma,\eta-\gamma)} U^+_{(\gamma,0)}$ is the only term of $E_i u$ involving $U^-_{(0,\gamma)} U^0_{(\eta-\gamma,\eta-\gamma)} U^+_{(\gamma+i,0)}$ because of the maximum of γ . For the same reason, $U^-_{(0,\gamma)} U^0_{(\eta-\gamma,\eta-\gamma)} U^+_{(\gamma,0)} E_i$ is the only term of uE_i involving $U^-_{(0,\gamma)} U^0_{(\eta-\gamma,\eta-\gamma)} U^+_{(\gamma+i,0)}$.

And

$$E_{i}\left(\sum_{j=1}^{d_{\gamma}}\theta_{j}x_{j}^{\gamma}\left(\sum_{a+b=\eta-\gamma}K_{a}'K_{b}U_{(\gamma,0)}^{+}\right)\right)$$

$$=\sum_{j=1}^{d_{\gamma}}\theta_{j}x_{j}^{\gamma}E_{i}\left(\sum_{a+b=\eta-\gamma}K_{a}'K_{b}U_{(\gamma,0)}^{+}\right)+U_{(0,\gamma-i)}U^{0}U_{(\gamma,0)}$$

$$=\sum_{j=1}^{d_{\gamma}}v^{(b-a)\cdot i}t^{\langle i,\eta-\gamma\rangle-\langle \eta-\gamma,i\rangle}\theta_{j}x_{j}^{\gamma}\left(\sum_{a+b=\eta-\gamma}K_{a}'K_{b}E_{i}U_{(\gamma,0)}^{+}\right)+U_{(0,\gamma-i)}U^{0}U_{(\gamma,0)}.$$

Choose some $\theta_k \neq 0$, $K_{a'}$ and $K'_{b'}$, since $E_i u = u E_i$, we must have

$$v^{(b'-a')\cdot i}t^{\langle i,\eta-\gamma\rangle-\langle \eta-\gamma,i\rangle}\theta_k x_k^{\gamma}K_{a'}'K_{b'}E_iU_{(\gamma,0)}^+ = \theta_k x_k^{\gamma}K_{a'}'K_{b'}U_{(\gamma,0)}^+E_i.$$

Since $\{x_1^{\gamma}, x_2^{\gamma}, \dots, x_{d_{\gamma}}^{\gamma}\}$ is a basis of $U_{(0,\gamma)}^-$, we obtain

$$v^{(b'-a')\cdot i}t^{\langle i,\eta-\gamma\rangle-\langle\eta-\gamma,i\rangle}E_iU^+_{(\gamma,0)}=U^+_{(\gamma,0)}E_i.$$

Rewrite the above equation under the multiplication *, we have

$$v^{(b'-a')\cdot i}t^{\langle i,\eta\rangle-\langle \eta,i\rangle}E_i * U^+_{(\gamma,0)} = U^+_{(\gamma,0)} * E_i.$$

Z. FAN AND Z. XIN

Here the element $U^+_{(\gamma,0)}$ can be expressed as the finite sum $\theta' \sum_{i \in \mathbb{Z}} t^i \otimes x_i$ where $\theta' \in \mathbb{Q}(v,t)$ and $x_i \in U_v$, and note that only finite x_i are nonzero. From the equation

$$v^{(b'-a')\cdot i}t^{\langle i,\eta\rangle - \langle \eta,i\rangle}E_i * U^+_{(\gamma,0)} = U^+_{(\gamma,0)} * E_i,$$

we know $\langle i, \eta \rangle - \langle \eta, i \rangle$ must be 0. It is a contradiction.

Now we would like to consider some subalgebras of $U_{v,t}(g)$ and compare the images of Harish-Chandra homomorphism. For $J \subseteq I$, let U_J be the subalgebra of $U_J =$ $\langle U_{v,t}^0, E_i, F_i | i \in J \rangle$ generated by $U_{v,t}^0$ and E_i, F_i with $i \in J$. We denote by Z_J the centre of U_J and ξ_J the Harish-Chandra homomorphism for U_J . Then we would like to show $\operatorname{Im}(\xi) \subset \operatorname{Im}(\xi_J)$. Denote by U_J^+ the subalgebra of U_J generated by E_i with $i \in J$ and U_J^- the subalgebra of U_J generated by F_i with $i \in J$. Recall the skew-Hopf pair $(\cdot|\cdot)$ on $U_{v,t}$. Set

$$R_J^+ = \{ x \in U_{v,t}^+ | (x|U_J^-) = 0 \} = \{ x \in U_{v,t}^+ | (x|U_J^-U_{v,t}^0) = 0 \},$$

$$R_J^- = \{ y \in U_{v,t}^- | (U_j^+|y) = 0 \} = \{ y \in U_{v,t}^- | (U_{v,t}^0U_J^+|y) = 0 \},$$

$$R_J = R_J^-U_{v,t}^0U_{v,t}^+ + U_{v,t}^-U_{v,t}^0R_J^+.$$

Lemma 5.3. We have

(1)
$$U_{v,t} = U_J \oplus R_J,$$

(2) $U_J R_J U_J \subset R_J,$
(3) $(\epsilon \otimes 1 \otimes \epsilon)(R_J) = 0.$

Proof. (1). It suffices to show $U_{\gamma}^+ = U_{J,\gamma}^+ \oplus R_{J,\gamma}^+$ for any $\gamma \in Q^+$. From the definition of R_{I}^{+} , we get the following equation,

$$R_{J,\gamma}^+ = \operatorname{Ker}(U_{\gamma}^+ \to (U_{-\gamma}^-)^* \to (U_{J,-\gamma}^-)^*),$$

where $U_{\gamma}^+ \to (U_{-\gamma}^-)^*$ is an algebra isomorphism because of the fact that skew-Hopf pair is nondegenerate on $U_{\gamma}^+ \times U_{-\gamma}^-$ and $(U_{-\gamma}^-)^* \to (U_{J,-\gamma}^-)^*$ is the canonical projection.

From the above equation and the epimorphism of $U^+_{\gamma} \to (U^-_{-\gamma})^*$, we have the following dimensional equation,

$$\dim R^+_{J,\gamma} = \dim U^+_{\gamma} - \dim U^-_{J,\gamma} = \dim U^+_{\gamma} - \dim U^+_{J,\gamma}$$

Since $(\cdot|\cdot)$ is nondegenerate on $U_{J,\gamma}^+ \times U_{J,-\gamma}^-$, we have $R_{J,\gamma}^+ \cap U_{J,\gamma}^+ = \{0\}$.

(2). For one $x_J^- \in U_J^-$, from proposition 3.1 and lemma 3.2, we get

$$(E_i x | x_J^-) = (E_i | F_i K'_{|x_J^-|-i})(x | a'_i(x_J^-)), \text{ for all } x \in R_J^+ \text{ and } i \in I$$

Since $a'_i(x_J^-) \in U_J^-$, we obtain $(E_i x | x_J^-) = 0$.

Similarly,

$$(xE_i|x_J^-) = (E_i|F_i)(x|a_i(x_J^-)K_i'), \text{ for all } x \in R_J^+$$

Since $a_i(x_J^-) \in U_J^-$, we obtain $(xE_i|x_J^-) = 0$.

20

$$U_J^+ R_J^- \subset R_J^- U_{v,t}, \qquad R_J^+ U_J^- \subset U_{v,t} R_J^+.$$

We show $U_J^+ R_J^- \subset R_J^- U_{v,t}$ for example, and the other case can be proved in a similar way.

Set any $y \in R_J^-$ and $j \in J$, from lemma 3.2, we can get

$$yE_j - E_j y = \frac{a_j(y)K'_j - K_j a'_j(y)}{v_j - v_j^{-1}}$$

Since $(E_j K'_{|y|-j} U_J^+|y) = (E_j K_{|y|-j}|F_j K'_{|y|-j})(U_J^+|a'_j(y))$ and $E_j U_J^+ \in U_J^+$, we have $(E_i U_I^+|y) = 0$. Then $(U_I^+|a_i'(y)) = 0$ because $(E_i|F_i) \neq 0$. That is to say, $a_i'(y)$ belongs to R_I^- . Similarly, $a_i(y)$ belongs to R_I^- . (2) is proved.

(3). Clear.

Proposition 5.4. $Im(\xi) \subseteq Im(\xi_J)$

Proof. Let $z \in Z(U_{v,t})$ and write $z = z_1 + z_2$ with $z_1 \in U_J$, $z_2 \in R_J$. lemma 5.3 (2), $z_1 \in Z(U_J)$, and hence by By lemma 5.3 (3), $\xi(z) = \xi_J(z_1) \in Im(\xi_J)$.

With different choices of the set J, we shall get different results. Now consider a special case when J = i for one $i \in I$, then the algebra U_{J_i} is generated by E_i , F_i for this fixed i and $K_j^{\pm 1}$, $K_j^{\prime \pm 1}$ for all $j \in I$. And let $U_{v,t}(g_i)$ be the subalgebra generated by $E_i, F_i, K_i^{\pm 1}, K_i^{\pm 1}$ for this fixed $i \in I$.

The centre of $U_{v,t}(g_i)$ has a basis of monomials $X^m Y^n$, $m \in \mathbb{Z}$, $n \in \mathbb{Z}_{\geq 0}$, where $X = K_i K'_i$ and Y is the Casimir element,

$$Y = E_i F_i + \frac{v^{-1} K_i + v K'_i}{(v - v^{-1})^2} = F_i E_i + \frac{v K_i + v^{-1} K'_i}{(v - v^{-1})^2}.$$

The following theorem gives the description of the centre of U_{J_i} for any fixed $i \in I$.

Theorem 5.5. For any $i \in I$, denote by Q' the \mathbb{Z} -lattice spanned by $I/\{i\}$. Then any $u \in Z(U_{J_i})$ can be expressed as the sum

$$u = \bigoplus_{x \in Q', y \in Q'} \mathbb{Q}(v, t) K_x K'_y L$$

where $L \in U_{v,t}(g_i)$ and K_x , K'_y and L satisfy the following two conditions:

(1) we require K_x and K'_y to meet the condition: $E_i K_x K'_y = v^{-2k} K_x K'_y E_i$ for some $k \in \mathbb{Z}$.

(2) $L = \{K_i^k(\Sigma_{m,n\in\mathbb{Z}}\mathbb{Q}(v,t)X^mY^n)\}, \text{ where } X = K_iK_i', Y = E_iF_i + \frac{v^{-1}K_i + vK_i'}{(v-v^{-1})^2} \text{ and } K_i'$ k is in (1).

Proof. For any $u \in U_{J_i}$, u can be expressed as the sum

$$u = \bigoplus_{x \in Q', y \in Q'} K_x K'_y L,$$

where $L \in U_{v,t}(g_i)$.

To determine the centre of U_{J_i} , it is enough to find the central elements with the form $K_x K'_y L$. That is to say, we have to check the equation

$$u(K_x K'_y L) = (K_x K'_y L)u,$$

for all $u \in U_{J_i}$.

(1) When $u = K_j$ for all $j \in I$. L has the form $L = \bigoplus_{a \in \mathbb{N}; b, c \in \mathbb{Z}} E_i^a K_{bi} K'_{ci} F_i^a, i$ is the fixed one.

(2) When $u = K'_i$ for all $j \in I$. The result is the same as in case (1).

(3) When $u = E_i$, where *i* is the fixed one. From the equation $E_i K_x K'_y L = K_x K'_y L E_i$, we may consider $K_x K'_y (E_i L) = K_x K'_y (\theta L E_i) \pounds \neg$ where θ is a constant in $\mathbb{Q}(v, t)$ determined by E_i and $K_x K'_y$. Assume *L* has the form as in (1):

$$L = \bigoplus_{a \in \mathbb{N}} E_i^a t_a F_i^a,$$

here $t_a \in \mathbb{Q}(v, t)[K_i^{\pm}, K_i'^{\pm}].$

Denote by m the maximal number such that $E_i^m t_m F_i^m \neq 0$. Since we have $E_i L = \theta L E_i$, then

$$E_i \bigoplus_{a \in \mathbb{N}} E_i^a t_a F_i^a = \theta \bigoplus_{a \in \mathbb{N}} E_i^a t_a F_i^a E_i,$$
$$E_i^{m+1} t_m F_i^m + E_i \bigoplus_{a \in \mathbb{N}, a < m} E_i^a t_a F_i^a = \theta E_i^m t_m F_i^m E_i + \theta \bigoplus_{a \in \mathbb{N}, a < m} E_i^a t_a F_i^a E_i$$

It is obvious that $\theta E_i^m t_m F_i^m E_i$ is the only term on the right including $E_i^{m+1} t_m F_i^m$. From the commutative relation of $U_{v,t}(g_i)$, we conclude $\theta \in v^{2n}$, where $n \in \mathbb{Z}$ determined by E_i and t_m .

The equation $E_i K_x K'_y L = K_x K'_y L E_i$ is equivalent to say $E_i L = \theta L E_i$, where θ determined by $K_x K'_y$ and E_i . Form the above argument, we know θ is not an arbitrary element in $\mathbb{Q}(v, t)$ but belongs to v^{2n} . That is to say, $K_x K'_y$ should satisfy

$$v^{2n}E_iK_xK'_u = K_xK'_uE_i, n \in \mathbb{Z}.$$

The final problem is to find all elements $L \in U_{v,t}(g_i)$ satisfying

$$E_i L = v^{2n} L E_i, n \in \mathbb{Z}.$$

For some $k \in \mathbb{Z}$, set $\Gamma_k = \{L | L \in U_{v,t}(g_i), E_i L = v^{2k} L E_i\}$. In fact, for arbitrary $k \in \mathbb{Z}$, we have $E_i K_i^k Z(U_{v,t}(g_i)) = v^{2k} K_i^k Z(U_{v,t}(g_i)) E_i$, and since K_i is invertible, we conclude

$$\Gamma_k = \{ K_i^k (\Sigma_{m,n \in \mathbb{Z}} \mathbb{Q}(v,t) X^m Y^n) \mid X = K_i K_i', \ Y = E_i F_i + \frac{v^{-1} K_i + v K_i'}{(v-v^{-1})^2} \},$$

for the fixed i and k.

(4) When $u = F_i$, where *i* is the fixed one. The result is the same as in case (3).

Recall that theorem 5.2 gives a criteria to justify when the extra central elements appear. Besides, to make the proposition 5.4 more accurate, we calculate the centre of U_{J_i} for each fixed $i \in I$ in theorem 5.5. Finally, we get the main theorem as follows.

Theorem 5.6. We have

(1) If there exists $\eta \in Q$ such that $[\eta, i] - [i, \eta] = 0$ for all $i \in I$, then the centre $Z(U_{v,t})$ is isomorphic under ξ to a subalgebra satisfying $\mathbb{Q}(v,t)[K'_{\eta}K_{\eta},K'_{\eta}^{-1}K_{\eta}^{-1}] \otimes (U^0_{\mathfrak{b}})^W \subseteq \xi(Z(U_{v,t})) \subseteq \bigcap_{i \in I} \xi(Z(U_{J_i}))$, where $Z(U_{J_i})$ is described in theorem 5.5. (2) Otherwise we have $\xi : Z(U_{v,t}) \to (U^0_{\mathfrak{b}})^W$ is an isomorphism.

Recall in theorem 5.2 we show that the nonzero homogeneous central elements only exist in the degree (η, η) with $\eta \in Q$ satisfying $[\eta, i] - [i, \eta] = 0$, $\forall i \in I$. So far, there are two situations described in the sense of Harish-Chandra homomorphism:

(1) Denote by Z_0 the set of central elements with degree (0,0), they are described via the Harish-Chandra homomorphism $\xi : Z_0 \to (U_{\flat}^0)^W$.

(2) Denote by Z_{ev} the set of central elements with nonzero degree $(2\eta, 2\eta)$ with $\eta \in Q$ satisfying $[\eta, i] - [i, \eta] = 0$, $\forall i \in I$. Since $K'_{\eta}K_{\eta}$ is invertible central elements, Z_{ev} are described via the Harish-Chandra homomorphism $\xi : Z_{ev} \to \mathbb{Q}(v, t)[K'_{\eta}K_{\eta}, K'^{-1}_{\eta}K^{-1}_{\eta}] \otimes (U^0_{\flat})^W$.

Then it is natural to ask the question : besides the central elements whose images of Harish-Chandra homomorphism are in $\mathbb{Q}(v,t)[K'_{\eta}K_{\eta},K'^{-1}K^{-1}_{\eta}]\otimes (U^0_{\flat})^W$, are there any other extra central elements? Consider the $U_{v,t}(sl_2)$ case, the Casimir element $EF + \frac{v^{-1}K+vK'}{(v-v^{-1})^2}$ hasn't been described in $\mathbb{Q}(v,t)[K'_{\eta}K_{\eta},K'^{-1}K^{-1}_{\eta}]\otimes (U^0_{\flat})^W$ under the Harish-Chandra homomorphism. To answer the question in more general cases, we have the following theorem.

Theorem 5.7. The homogeneous central elements exist in and only in grading (η, η) for all $\eta \in Q$ satisfying $[\eta, i] - [i, \eta] = 0$, $\forall i \in I$.

Proof. It is sufficient to show the existence. Set $\{\alpha_i\}_{i\in I}$ as the set of simple roots. For one arbitrary $\eta \in Q$ satisfying $[\eta, i] - [i, \eta] = 0$, $\forall i \in I$. We express η as the sum $x_1\alpha_1 + x_2\alpha_2 + \ldots + x_n\alpha_n$ where $x_1, x_2, \ldots, x_n \in \mathbb{Z}$ and $\alpha_1, \alpha_2, \ldots, \alpha_n \in \{\alpha_i\}_{i\in I}$.

Recall the definition of multiplication *, it is equivelent to show the existence of central elements with grading (η, η) under the multiplication *.

For finite type, a direct calculation shows if $\eta = x_1\alpha_1 + x_2\alpha_2 + \ldots + x_n\alpha_n$ satisfying $[\eta, i] - [i, \eta] = 0 \ \forall i \in I$, then there always exists $y_1, y_2, \ldots, y_n \in \mathbb{Z}$ such that for every $i = 1, 2, \ldots, n, x_i$ and y_i have the same parity and $\nu = y_1\alpha_1 + y_2\alpha_2 + \ldots + y_n\alpha_n \in Q \cap 2\Lambda$.

For any $\nu \in Q \cap 2\Lambda$, from the conclusion of Harish-Chandra isomorphism in one parameter case, we know there exist one unique central element whose image of Harish-Chandra isomorphism is $(K_{\nu})^W$. We can denote this central element as Z_{ν} and express it as $Z_{\nu} = \bigoplus_{\nu'>0} U_{\nu'}^{-} U^0 U_{\nu'}^+$. And for every $\nu' > 0$, denote by $\{x_{\nu'}\}$ and $\{y_{\nu'}\}$ the basis of $U_{\nu'}^-$ and $U_{\nu'}^+$ respectively. Then Z_{ν} can be expressed as

$$Z_{\nu} = \bigoplus_{\nu' \ge 0} \bigoplus_{x_{\nu'} \in \{x_{\nu'}\}, y_{\nu'} \in \{y_{\nu'}\}} x_{\nu'} U^0 y_{\nu'},$$

where U^0 denotes the polynomial algebra in Cartan part.

Fix some $\nu' \in Q$, $x_{\nu'}$ and $y_{\nu'}$, then write $x_{\nu'}U^0y_{\nu'}$ as $\bigoplus_{\varsigma\in Q} \mathbb{Q}(v,t)x_{\nu'}K_{\varsigma}y_{\nu'}$. Corresponding to every $x_{\nu'}U^0y_{\nu'}$, we define one term $x_{\nu'}\tilde{U}^0y_{\nu'}$ in $((U_{v,t},(I,\cdot)),*)$ as $\bigoplus_{\varsigma\in Q} \mathbb{Q}(v,t)x_{\nu'}K_{\frac{\eta-\nu'+\varsigma}{2}}K'_{\frac{\eta-\nu'-\varsigma}{2}}y_{\nu'}$.

Then we can define one element Z_{ν} as

$$\tilde{Z}_{\nu} = \bigoplus_{\nu' \ge 0} \bigoplus_{x_{\nu'} \in \{x_{\nu'}\}, y_{\nu'} \in \{y_{\nu'}\}} \mathbb{Q}(v) x_{\nu'} \tilde{U}^0 y_{\nu'},$$

comparing the communicative relation, since the only difference exists in Cartan part, we know \tilde{Z}_{ν} is in the centre of $U_{v,t}(I, \cdot)$.

References

- [BW] G. Benkart and S. Witherspoon, *Two-parameter quantum groups and Drinfel'd doubles*. Algebras & Representation Theory, 2004, 7(3):261-286.
- [C] S. Clark, Odd knot invariants from quantum covering groups. Algebraic & Geometric Topology, 2017, 17(5):2961-3005.
- [CFYW] S. Clark, Z. Fan, and Y. Li, elt, Quantum supergroups III. Twistors. Communications in Mathematical Physics, 2014, 332(1): 415-436.
- [FL] Z. Fan and Y. Li, Two-parameter quantum algebras, canonical bases and categorifications. International Mathematics Research Notices, 2015, 2015(16):7016-7062.
- [FX] Z. Fan and J. Xing, Drinfeld double of deformed quantum algebras. Journal of Algebra, 2019, 534:358-383.
- [HPR] N. Hu, Y. Pei, and M. Rosso, Multi-parameter quantum groups and quantum shuffles. I. Quantum Affine Algebras, Extended Affine Lie Algebras, and their Applications, 145-71. Contemp. Math. 506. Providence, RI: American Mathematical Society, 2010.
- [Jan] J.C. Jantzen, *Lectures on quantum groups*. American Mathematical Society, Providence, 1996.
- [Kas] C. Kassel, *Quantum groups*. Springer-Verlag, New York, 1995.
- [K11] L. H. Kauffman, Remarks on Khovanov homology and the Potts model. Perspectives in Analysis, Geometry, and Topology, 2011:237-262.
- [K16] L. H. Kauffman, Knot logic and topological quantum computing with Majorana fermions. Logic and Algebraic Structures in Quantum Computing, 2016:223-336.
- [L] G. Lusztig, *Introduction to quantum groups*. Modern Birkhäuser Classics, Birkhäuser/Springer, New York, 2010.
- [NSS] C. Nayak, S. H. Simon and A. Stern, elt, Non-Abelian anyons and topological quantum computation. Reviews of Modern Physics, 2008, 80:1083-1159.
- [RT] N. Y. Reshetikhin and V. G. Turaev, *Ribbon graphs and their invariants derived from quantum groups*. Communications in Mathematical Physics, 1990, 127:1-26.
- [Tur90] V G. Turaev, Operator invariants of tangles, and R-matrices. Mathematics of the USSR-Izvestiya, 1990, 35(2):411-444.

- [W] E. Witten, *Quantum field theory and the jones polynomial*. Communications in Mathematical Physics, 1989, 121(3): 351-399.
- [X97] J. Xiao, Drinfeld double and Ringel-Green Theory of Hall algebras. Journal of Algebra, 1997, 190(1):100-144.
- [BKL] G. Benkart, S.J. Kang, K.H. Lee, On the centre of two-parameter quantum groups. Proc. Roy. Soc. Edinburgh Sect. A 136(3)(2006)445-472.

College of Mathematical Sciences, Harbin Engineering University, Harbin, China, 150001

Email address: fanzhaobing@hrbeu.edu.cn (Fan)

College of Mathematical Sciences, Harbin Engineering University, Harbin, China, 150001

Email address: 2016@hrbeu.edu.cn (Xin)