

# A robust $C^0$ interior penalty method for a gradient-elastic Kirchhoff plate model

Mingqing Chen<sup>a</sup>, Jianguo Huang<sup>b,1</sup>, Xuehai Huang<sup>c,2</sup>

<sup>a</sup>*School of Mathematical Sciences, Zhejiang Normal University, Zhejiang 321004, China*

<sup>b</sup>*School of Mathematical Sciences, and MOE-LSC, Shanghai Jiao Tong University, Shanghai 200240, China*

<sup>c</sup>*School of Mathematics, Shanghai University of Finance and Economics, Shanghai 200433, China*

---

## Abstract

This paper is devoted to proposing and analyzing a robust  $C^0$  interior penalty method for a gradient-elastic Kirchhoff plate (GEKP) model over a convex polygon. The numerical method is obtained by combining the triangular Hermite element and a  $C^0$  interior penalty method, which can avoid the use of higher order shape functions or macroelements. Next, a robust regularity estimate is established for the GEKP model based on our earlier result for a triharmonic equation on a convex polygon. Furthermore, some local lower bound estimates of the a posteriori error analysis are established. These together with an enriching operator and its error estimates lead to a Céa-like lemma. Thereby, the optimal error estimates are achieved, which are also robust with respect to the small size parameter. In addition, it is proved that this numerical method is convergent without any additional regularity assumption for the exact solution. Some numerical experiments are performed to verify the theoretical findings.

*Keywords:* gradient-elastic Kirchhoff plate model; regularity;  $C^0$  interior penalty method; robustness; error estimates;

---

## 1. Introduction

To account for small-scale phenomena in micro-scale or even nano-scale structures, higher-order continuum theories were developed systematically by many researchers. The key feature is that the corresponding constitutive relations require to depend on additional size parameters to describe micro/nano scale properties of structures. We refer the reader to the references [16, 50, 38, 34, 37, 1, 2, 47] for details along this line. Most of the underlying mathematical models involve two or more size parameters. Aifantis and his collaborators proposed in [2, 47] a one-parameter simplified strain gradient elasticity (SGE) theory for material deformation at the micro/nano scale, which is effective in simulating the deformation of elastic plastic materials. Its constitutive relation reads (cf. [41])

$$\tilde{\sigma}_{ij} = C_{ijkl}(\varepsilon_{kl} - \iota^2 \varepsilon_{kl,mm}), \quad (1.1)$$

where,  $\varepsilon = (\varepsilon_{ij})$  is the strain tensor in elasticity,  $\tilde{\sigma} = (\tilde{\sigma}_{ij})$  is the stress tensor of the SGE model,  $C_{ijkl}$  is a fourth-order tensor and  $\iota$  denotes the size parameter describing the

---

*Email addresses:* chenmq@zjnu.edu.cn (Mingqing Chen), jghuang@sjtu.edu.cn (Jianguo Huang), huang.xuehai@sufe.edu.cn (Xuehai Huang)

<sup>1</sup>Corresponding author. The work of this author was partially supported by NSFC (Grant No. 12071289) and the National Key Research and Development Project (2020YFA0709800).

<sup>2</sup>The work of this author was partially supported by NSFC (Grant No. 12171300 and 12071289).

length scale of the microstructure of the material under discussion. Here and below, we use Einstein's summation convention. Besides, for linear elastic problem,

$$C_{ijkl} = \lambda \delta_{ij} \delta_{kl} + \mu \delta_{ik} \delta_{jl} + \mu \delta_{il} \delta_{jk}, \quad (1.2)$$

where  $\delta$  is the Kronecker delta function, and  $\lambda$  and  $\mu$  are the Lamé constants.

Given a thin plate occupying a three dimensional region  $\mathcal{P} = \Omega \times (-\frac{t}{2}, \frac{t}{2})$ , denote the displacement by  $\mathbf{u} = (u_i) : \mathcal{P} \rightarrow \mathbb{R}^3$ , where  $\Omega$  is the midsurface of the plate and  $t \ll \text{diam}(\Omega)$  is the thickness of the plate. Under the Kirchhoff-Love hypothesis, the displacement field can be approximated as

$$u_1 = -x_3 \frac{\partial w(x_1, x_2)}{\partial x_1}, \quad u_2 = -x_3 \frac{\partial w(x_1, x_2)}{\partial x_2}, \quad u_3 = w(x_1, x_2),$$

where  $w : \Omega \rightarrow \mathbb{R}$  is the transverse deflection at the midsurface  $\Omega \times \{0\}$ . Then, the classical strains in (1.1) can be simplified to the plane strains  $\varepsilon_{ij} = -x_3 \frac{\partial^2 w}{\partial x_i \partial x_j}$  with  $i, j = 1, 2$ . Next, we introduce a curvature tensor  $\boldsymbol{\kappa} = \boldsymbol{\kappa}(\nabla w) = (\frac{\partial^2 w}{\partial x_i \partial x_j}) : \Omega \rightarrow \mathbb{R}^{2 \times 2}$  as a restriction of the classic strain tensor field to the midsurface. Then, we can obtain the classic bending moment tensor  $\mathbf{M} = \mathcal{D}((1 - \nu)\boldsymbol{\kappa} + \nu \text{tr}(\boldsymbol{\kappa})\mathbf{I})$  and the classical transverse force vector  $\mathbf{Q} = \text{div } \mathbf{M}$ . Here,  $\mathcal{D} = \frac{Et^3}{12(1-\nu^2)}$  is the bending rigidity related to Poisson's ratio  $\nu \in (0, 1/2)$  and Young's module  $E$ . Then, the combination of the Kirchhoff plate bending equation, (1.1) and (1.2), one can deduce a sixth-order governing equation for the gradient-elastic Kirchhoff plate (GEKP) model

$$\text{div}((1 - \iota^2 \Delta)\mathbf{Q}) = \mathcal{D}(\Delta^2 w - \iota^2 \Delta^3 w) = f, \quad (1.3)$$

where  $\iota \in (0, 1)$  denotes the size parameter, and  $f \in L^2(\Omega)$  is the transversal load. We refer to [41] for details of the above model.

It is easy to observe that the mathematical model (1.3) is a singularly perturbed problem with respect to the size parameter  $\iota$ . In fact, if  $\iota = 0$ , then the original problem will degenerate to a fourth-order elliptic problem. This implies that the solution of the reduced problem may not meet all of the boundary conditions of problem (3.1) and this would lead to the phenomenon of boundary layer as  $\iota$  goes to zero. Thus, it is extremely necessary to develop a robust numerical method for approximating the GEKP model with respect to  $\iota$ .

As far as we know, there are some works on numerically solving (3.1), such as the analytical methods [41, 44, 56, 18, 58], the finite element methods (FEM) [45, 7], the extended Kantorovich method [6, 52], the differential quadrature method [39, 32, 33, 35], the machine learning approach [57], the isogeometric analysis method [40], etc. However, most of these works mainly pay attention to numerical simulation using the underlying methods. In other words, there are few works on the construction of a robust numerical method for the GEKP model and developing the corresponding robust error analysis.

Hence, one of the main motivations of our study is to develop a robust numerical method for solving the GEKP model (3.1). We can refer to [42, 54, 15, 9] and references therein for details about some robust nonconforming FEMs for approximating singularly perturbed fourth-order problems. However, these numerical methods only possess the sharp half-order convergence rate due to the effect of boundary layer. Inspired by the Nitsche's technique [43] and the penalty technique [5, 10], the authors in [24, 30] devised some robust nonconforming finite element methods by imposing one of boundary conditions weakly for a singularly perturbed fourth-order problem. See also the work in [54].

Furthermore, there has developed a few of FEMs that approximate triharmonic equations, including the conforming FEM [26, 14], the decoupled mixed FEM [48, 3, 19], the nonconforming FEM [55, 28, 27, 53, 36], the  $C^0$  interior penalty method [23, 12], the virtual

element method [17, 11, 4, 13, 29], and hybrid high-order method [46]. It is well-known that the conforming finite element methods for approaching high-order partial differential equations are hard to be constructed and implemented due to the quite a number of degrees of freedom (DoFs) and the existence of supersmooth DoFs. Therefore we prefer to apply less smooth finite element. In particular, the authors in [23, 12] proposed some  $C^0$  interior penalty methods for a third-order Laplace equation using the cubic Lagrange finite element space. This element only has 10 DoFs. Making use of the a posteriori analysis techniques, some Céa-like lemmas have been established with minimal regularity assumptions on the exact solution. This kind of methods has an advantage of using a small number of degrees of freedom. Thus, we are tempted to develop an interior penalty method to approximate such singularly perturbed high-order problems.

Another purpose of our study is to build up the regularity theory for the GEKP problem (3.1) and further to deduce robust error analysis of the underlying numerical method. In [42], the authors derived a uniform regularity estimate for a singularly perturbed fourth-order problem with the help of the regularity result for a biharmonic equation. Recall that the regularity theory of a triharmonic equation on a convex polygon has been established in [15]. This together with the regularity theory of the reduced problem (3.3) in [20] can lead to the following robust regularity estimate

$$|w - w_0|_2 + \iota \|w\|_3 + \iota^2 \|w\|_4 \lesssim \iota^{1/2} \|f\|_0.$$

The main results of our work in this paper can be summarized as follows. Borrowing the ideas in [43, 5, 10, 23], we design a  $C^0$  interior penalty method to discretize problem (3.1). The element we used here is the triangular Hermite element having 10 DoFs, which is easy to implement numerically. Using the similar approach as in [23, 21, 22, 30], we first introduce a quasi interpolation operator and an enriching operator connecting the Hermite element space to a  $C^2$  element space. Next, we derive some local lower bound estimates of a posteriori error analysis for the resulting numerical method, which will be exploited to achieve a Céa-like lemma. Thereby, the optimal and robust error estimates with respect to the size parameter can be obtained. Furthermore, it can be shown that the numerical method we proposed is convergent for any  $w \in H_0^3(\Omega)$ .

The rest of this paper is arranged as follows. In Sect. 2, we introduce some notation and existing results for further analysis. In Sect. 3, we present the variational formulation of problem (3.1) and develop the underlying regularity estimate relying on Theorem 3.5 in [15]. In Sect. 4, we design a  $C^0$  interior penalty method of problem (3.1) and prove its well-posedness. In Sect. 5, we define two interpolation operators and derive their error estimates. In Sect. 5.3, a Céa-like lemma can be derived with the aid of some interpolation error estimates and the a posteriori analysis techniques. Then, we are able to analyse the convergence properties of the method. In the last section, the numerical experiments are performed to verify the theoretical findings.

## 2. Preliminaries

Throughout this paper, let  $\Omega$  be a bounded convex domain, which is partitioned into a family of shape regular triangles  $\mathcal{T}_h = \{K\}$ . That is to say that there exists a constant  $\gamma_0 \geq 1$  such that

$$\frac{h_K}{\rho_K} \leq \gamma_0 \quad \forall K \in \mathcal{T}_h,$$

where  $h_K = \text{diam}(K)$  and  $h = \max_{K \in \mathcal{T}_h} h_K$ . Use  $\mathcal{V}(\mathcal{T}_h)$  and  $\mathcal{V}^i(\mathcal{T}_h)$  (resp.  $\mathcal{E}(\mathcal{T}_h)$  and  $\mathcal{E}^i(\mathcal{T}_h)$ ) to stand for the sets of all and interior vertices (resp. edges) of  $\mathcal{T}_h$ , respectively. We then introduce the patch of vertices and edges for further use. For any vertex  $\delta \in \mathcal{V}(\mathcal{T}_h)$  and edge  $e \in \mathcal{E}(\mathcal{T}_h)$ , write  $\omega_\delta$  and  $\omega_e$  to be respectively the unions of all triangles in  $\mathcal{T}_\delta$

and  $\mathcal{T}_e$ , where  $\mathcal{T}_\delta$  and  $\mathcal{T}_e$  is the set of all triangles in  $\mathcal{T}_h$  sharing common vertex  $\delta$  and edge  $e$ , respectively. In addition, for a finite set  $A$ , denote by  $\#A$  its cardinality. For each  $K \in \mathcal{T}_h$ , denote by  $\mathbf{n}_K$  the unit outward normal to  $\partial K$  and denote the unit tangential direction of  $\partial K$  by  $\mathbf{t}_K$  obtained by rotating  $\mathbf{n}_K$  90° counterclockwise. Without causing any confusion, abbreviate  $\mathbf{n}_K$  and  $\mathbf{t}_K$  as  $\mathbf{n}$  and  $\mathbf{t}$  for simplicity. For all  $e \in \mathcal{E}^i(\mathcal{T}_h)$  sharing by two triangles  $K^+$  and  $K^-$ , we preset the unit normal vector of  $e$  by  $\mathbf{n}_e := \mathbf{n}_+$ , where  $\mathbf{n}_+$  is the unit outward normal to  $e$  of  $K^+$  and denote the unit tangential direction of  $e$  by  $\mathbf{t}_e$  obtained by rotating  $\mathbf{n}_e$  90° counterclockwise. Define the jump and average on  $e$  as follows:

$$\begin{aligned} \llbracket \partial_{\mathbf{a}} v \rrbracket|_e &:= \partial_{\mathbf{a}_e} v|_{K^+} - \partial_{\mathbf{a}_e} v|_{K^-}, & \{\{\partial_{\mathbf{a}} v\}\}|_e &:= \frac{\partial_{\mathbf{a}_e} v|_{K^+} + \partial_{\mathbf{a}_e} v|_{K^-}}{2}, \\ \llbracket \partial_{\mathbf{ab}} v \rrbracket|_e &:= \partial_{\mathbf{a}_e \mathbf{b}_e} v|_{K^+} - \partial_{\mathbf{a}_e \mathbf{b}_e} v|_{K^-}, & \{\{\partial_{\mathbf{ab}} v\}\}|_e &:= \frac{\partial_{\mathbf{a}_e \mathbf{b}_e} v|_{K^+} + \partial_{\mathbf{a}_e \mathbf{b}_e} v|_{K^-}}{2}, \\ \llbracket \partial_{\mathbf{abc}} v \rrbracket|_e &:= \partial_{\mathbf{a}_e \mathbf{b}_e \mathbf{c}_e} v|_{K^+} - \partial_{\mathbf{a}_e \mathbf{b}_e \mathbf{c}_e} v|_{K^-}, & \{\{\partial_{\mathbf{abc}} v\}\}|_e &:= \frac{\partial_{\mathbf{a}_e \mathbf{b}_e \mathbf{c}_e} v|_{K^+} + \partial_{\mathbf{n}_e \mathbf{n}_e \mathbf{t}_e} v|_{K^-}}{2}. \end{aligned}$$

where  $\mathbf{a}, \mathbf{b}, \mathbf{c} = \mathbf{n}, \mathbf{t}$ . For any  $e \in \mathcal{E}^\partial(\mathcal{T}_h)$ , the associated jump and average are given by

$$\begin{aligned} \llbracket \partial_{\mathbf{a}} v \rrbracket|_e &:= \partial_{\mathbf{a}_e} v, & \{\{\partial_{\mathbf{a}} v\}\}|_e &:= \partial_{\mathbf{a}_e} v, \\ \llbracket \partial_{\mathbf{ab}} v \rrbracket|_e &:= \partial_{\mathbf{a}_e \mathbf{b}_e} v, & \{\{\partial_{\mathbf{ab}} v\}\}|_e &:= \partial_{\mathbf{a}_e \mathbf{b}_e} v, \\ \llbracket \partial_{\mathbf{abc}} v \rrbracket|_e &:= \partial_{\mathbf{a}_e \mathbf{b}_e \mathbf{c}_e} v, & \{\{\partial_{\mathbf{abc}} v\}\}|_e &:= \partial_{\mathbf{a}_e \mathbf{b}_e \mathbf{c}_e} v. \end{aligned}$$

Clearly,

$$\llbracket uv \rrbracket = \{\{u\}\} \llbracket v \rrbracket + \llbracket u \rrbracket \{\{v\}\}.$$

Given a bounded domain  $D$  and any integers  $m, k \geq 0$ , denote by  $H^m(D)$  the standard Sobolev spaces on  $D$  with norm  $\|\cdot\|_{m,D}$  and semi-norm  $|\cdot|_{m,D}$ , and  $H_0^m(D)$  the closure of  $C_0^\infty(D)$  with respect to  $\|\cdot\|_{m,D}$ . The notation  $(\cdot, \cdot)_D$  symbolizes the standard inner product on  $D$ . For  $D = \Omega$ , we abbreviate  $\|\cdot\|_{m,\Omega}$ ,  $|\cdot|_{m,\Omega}$  and  $(\cdot, \cdot)_D$  as  $\|\cdot\|_m$ ,  $|\cdot|_m$  and  $(\cdot, \cdot)$ , respectively. Let  $\mathbb{P}_k(D)$  be the set of all polynomials on  $D$  with the total degree up to  $k$ . Denote by  $\Pi_D^k$  the standard  $L^2$  projection operator from  $L^2(D)$  to  $\mathbb{P}_k(D)$ .

For simplicity of expression, we use  $\lesssim$  to stand for  $\leq C$ , where  $C$  may be a generic positive constant independent of the mesh size  $h$  and the material parameter  $\iota$ . And  $a \approx b$  indicates  $a \lesssim b \lesssim a$ . Here  $\nabla_h$  (resp.  $\Delta_h$ ) is the elementwise version of  $\nabla$  (resp.  $\Delta$ ) with respect to  $\mathcal{T}_h$ .

For forthcoming theoretical analysis, recall some basic results. As given in [20, Theorem 1.5.1.10], the following multiplicative trace inequality holds

$$\|v\|_{0,\partial D}^2 \lesssim \|v\|_{0,D} \|v\|_{1,D} \quad \forall v \in H^1(D),$$

where  $D$  is a bounded domain with Lipschitz boundary. And by the scaling argument, there exists a positive constant  $C(\varepsilon)$  for any  $\varepsilon > 0$ , such that

$$\|v\|_{0,\partial K} \lesssim \varepsilon h_K^{1/2} |v|_{1,K} + C(\varepsilon) h_K^{-1/2} \|v\|_{0,K} \quad \forall v \in H^1(K), K \in \mathcal{T}_h. \quad (2.1)$$

Moreover, by the inverse inequality of polynomials, one has readily

$$\|\nabla^m p\|_{0,\partial K} \lesssim h_K^{-1/2} \|\nabla^m p\|_{0,K} \quad \forall p \in \mathbb{P}_k(K), K \in \mathcal{T}_h. \quad (2.2)$$

### 3. Regularity results

Let us consider the following boundary value problem with the Dirichlet boundary conditions:

$$\begin{cases} \mathcal{D}(\Delta^2 w - \iota^2 \Delta^3 w) = f & \text{in } \Omega, \\ w = \partial_{\mathbf{n}} w = \partial_{\mathbf{nn}} w = 0 & \text{on } \partial\Omega. \end{cases} \quad (3.1)$$

Without loss of generality, set  $\mathcal{D} = 1$  throughout this paper. The weak formulation of problem (3.1) is to find  $w \in V := H_0^3(\Omega)$  such that

$$\iota^2 a(w, v) + b(w, v) = (f, v) \quad \forall v \in V, \quad (3.2)$$

where

$$a(w, v) = (\nabla^3 w, \nabla^3 v) \quad \text{and} \quad b(w, v) = (\nabla^2 w, \nabla^2 v).$$

Set  $\iota = 0$ , problem (3.1) will reduce to a fourth-order elliptic problem:

$$\begin{cases} \Delta^2 w_0 = f & \text{in } \Omega, \\ w_0 = \partial_n w_0 = 0 & \text{on } \partial\Omega. \end{cases} \quad (3.3)$$

Its weak formulation is to find  $w_0 \in H_0^2(\Omega)$  such that

$$b(w_0, v) = (f, v) \quad \forall v \in H_0^2(\Omega). \quad (3.4)$$

It's worth noting that for any convex polygon  $\Omega$ , the operator  $\Delta^2$  is an isomorphism from  $H^3(\Omega) \cap H_0^2(\Omega)$  to  $H^{-1}(\Omega)$  (cf. [20, Corollary 7.3.2.5]). Then,  $w_0 \in H^3(\Omega)$  and there exists a positive constant  $C$  such that

$$\|w_0\|_3 \leq C \|f\|_0. \quad (3.5)$$

For the regularity estimate of (3.1), we recall a useful result of the triharmonic equation.

**Theorem 3.1.** ([15, Theorem 3.5]) *Let  $\Omega$  be a convex polygon. For any  $w \in H_0^3(\Omega)$  satisfying*

$$-\Delta^3 w = f \in H^{-2}(\Omega),$$

*it holds  $w \in H^4(\Omega)$  and*

$$\|w\|_4 \lesssim \|f\|_{-2}.$$

Following the argument in [42, Lemma 5.1], we have the following regularity result.

**Theorem 3.2.** *Assume  $\Omega$  be a convex polygon. Let  $w \in H_0^3(\Omega)$  and  $w_0 \in H_0^2(\Omega)$  be the solutions of problems (3.2) and (3.4), respectively. Then it holds  $w \in H^4(\Omega)$  and*

$$|w - w_0|_2 + \iota \|w\|_3 + \iota^2 \|w\|_4 \lesssim \iota^{1/2} \|f\|_0. \quad (3.6)$$

*Here, the hidden constant is independent of the size parameter  $\iota$ .*

*Proof.* The first equation of problem (3.1) can be reformulated as

$$-\Delta^3 w = \iota^{-2} \Delta^2 (w_0 - w) \in H^{-2}(\Omega),$$

which along with Theorem 3.1 yields  $w \in H^4(\Omega)$  and

$$\|w\|_4 \lesssim \|\iota^{-2} \Delta^2 (w_0 - w)\|_{-2} \lesssim \iota^{-2} |w_0 - w|_2. \quad (3.7)$$

Thus, it suffices to prove

$$|w - w_0|_2 + \iota \|w\|_3 \lesssim \iota^{1/2} \|f\|_0.$$

For any  $v \in H_0^2(\Omega) \cap H^3(\Omega)$ , it holds

$$v|_{\partial\Omega} = \partial_n v|_{\partial\Omega} = \partial_{nt} v|_{\partial\Omega} = 0.$$

Then, by (3.2) and (3.4), we have

$$\iota^2(\nabla^3 w, \nabla^3 v) + (\nabla^2 w, \nabla^2 v) - \iota^2(\partial_{\mathbf{nnn}} w, \partial_{\mathbf{nn}} v)_{\partial\Omega} = (\nabla^2 w_0, \nabla^2 v). \quad (3.8)$$

Choosing  $v$  as  $w - w_0$  in (3.8), we have by integration by parts that

$$\iota^2|w|_3^2 + |w - w_0|_2^2 = \iota^2(\nabla^3 w, \nabla^3 w_0) - \iota^2(\partial_{\mathbf{nnn}} w, \partial_{\mathbf{nn}} w_0)_{\partial\Omega}. \quad (3.9)$$

By the multiplicative trace inequality, (3.5) and (3.7),

$$\begin{aligned} -\iota^2(\partial_{\mathbf{nnn}} w, \partial_{\mathbf{nn}} w_0)_{\partial\Omega} &\lesssim \iota^2 \|w\|_3^{1/2} \|w\|_4^{1/2} \|w_0\|_2^{1/2} \|w_0\|_3^{1/2} \\ &\lesssim \iota \|w\|_3^{1/2} |w - w_0|_2^{1/2} \|f\|_0. \end{aligned}$$

From (3.5) we get

$$\iota^2(\nabla^3 w, \nabla^3 w_0) \leq \iota^2 |w|_3 |w_0|_3 \lesssim \iota^2 |w|_3 \|f\|_0.$$

Combining (3.9) and the last two inequalities gives

$$\begin{aligned} \iota^2|w|_3^2 + |w - w_0|_2^2 &\lesssim (\iota^{1/2} \|w\|_3^{1/2} |w - w_0|_2^{1/2} + \iota^{3/2} |w|_3) \iota^{1/2} \|f\|_0 \\ &\lesssim (\iota \|w\|_3 + |w - w_0|_2) \iota^{1/2} \|f\|_0, \end{aligned}$$

which along with the absorbing technique yields the desired result.  $\square$

#### 4. $H^3$ nonconforming finite element method

For a triangle  $K \in \mathcal{T}_h$ , let  $\mathcal{V}(K)$  and  $\mathcal{E}(K)$  be the set of all vertices and edges of  $K$ , respectively.

##### 4.1. Finite element space

The Hermite element in a triangle takes  $V(K) := \mathbb{P}_3(K)$  as the local shape function space. The corresponding DoFs are

$$\begin{cases} v(\zeta) & \forall \zeta \in \mathcal{V}(K), \\ \nabla v(\zeta) & \forall \zeta \in \mathcal{V}(K), \\ v(\zeta_K). \end{cases} \quad (4.1)$$

where  $\zeta_K$  is the barycenter of  $K$ . Let

$$V_h := \{v \in H^1(\Omega) : v|_K \in V(K) \text{ for each } K \in \mathcal{T}_h, v \text{ and } \nabla v \text{ are continuous at all vertices of triangulation } \mathcal{T}_h \text{ and all DoFs (4.1) vanish on boundary}\}.$$

##### 4.2. Discrete formulation

Assume  $w$  is sufficiently smooth. For any  $v \in V_h \subset H_0^1(\Omega)$ , it follows from integration by parts that

$$\begin{aligned} -(\Delta^3 w, v) &= (\nabla \Delta^2 w, \nabla v) \\ &= \sum_{K \in \mathcal{T}_h} \left( -(\nabla^2 \Delta w, \nabla^2 v)_K + (\partial_{\mathbf{n}} \nabla \Delta w, \nabla v)_{\partial K} \right) \\ &= \sum_{K \in \mathcal{T}_h} \left( (\nabla^3 w, \nabla^3 v)_K - (\partial_{\mathbf{n}} \nabla^2 w, \nabla^2 v)_{\partial K} + (\partial_{\mathbf{n}} \nabla \Delta w, \nabla v)_{\partial K} \right) \end{aligned} \quad (4.2)$$

and

$$(\Delta^2 w, v) = \sum_{K \in \mathcal{T}_h} \left( (\nabla^2 w, \nabla^2 v)_K - (\partial_{\mathbf{n}} \nabla w, \nabla v)_{\partial K} \right). \quad (4.3)$$

Note that

$$\partial_i v = n_i \partial_{\mathbf{n}} v + t_i \partial_{\mathbf{t}} v \quad \forall v \in H^1(K), \quad i = 1, 2, \quad (4.4)$$

with any  $\mathbf{n} = (n_1, n_2)^\top \in \mathbb{R}^2$  and  $\mathbf{t} = (t_1, t_2)^\top \in \mathbb{R}^2$ . Then we get easily that

$$\nabla^2 v = \mathbf{n} \partial_{\mathbf{n} \mathbf{t}} v \mathbf{t}^\top + \mathbf{n} \partial_{\mathbf{n} \mathbf{n}} v \mathbf{n}^\top + \mathbf{t} \partial_{\mathbf{t} \mathbf{n}} v \mathbf{n}^\top + \mathbf{t} \partial_{\mathbf{t} \mathbf{t}} v \mathbf{t}^\top, \quad (4.5)$$

$$\partial_{\mathbf{n}} \nabla \Delta w = \partial_{\mathbf{n} \mathbf{n}} \Delta w \mathbf{n}^\top + \partial_{\mathbf{n} \mathbf{t}} \Delta w \mathbf{t}^\top, \quad (4.6)$$

$$\partial_{\mathbf{n}} \nabla^2 w = \mathbf{n} \partial_{\mathbf{n} \mathbf{n} \mathbf{n}} w \mathbf{n}^\top + \mathbf{n} \partial_{\mathbf{n} \mathbf{n} \mathbf{t}} w \mathbf{n}^\top + \mathbf{t} \partial_{\mathbf{n} \mathbf{n} \mathbf{t}} w \mathbf{n}^\top + \mathbf{t} \partial_{\mathbf{n} \mathbf{t} \mathbf{t}} w \mathbf{t}^\top, \quad (4.7)$$

$$\partial_{\mathbf{n}} \nabla w = \partial_{\mathbf{n} \mathbf{n}} w \mathbf{n}^\top + \partial_{\mathbf{n} \mathbf{t}} w \mathbf{t}^\top. \quad (4.8)$$

Combining (4.4)-(4.7) and (4.2), we find

$$\begin{aligned} -(\Delta^3 w, v) &= \sum_{K \in \mathcal{T}_h} \left( (\nabla^3 w, \nabla^3 v)_K + (\partial_{\mathbf{n} \mathbf{n}} \Delta w, \partial_{\mathbf{n}} v)_{\partial K} + (\partial_{\mathbf{n} \mathbf{t}} \Delta w, \partial_{\mathbf{t}} v)_{\partial K} \right) \\ &\quad - \sum_{K \in \mathcal{T}_h} \left( (\partial_{\mathbf{n} \mathbf{n} \mathbf{n}} w, \partial_{\mathbf{n} \mathbf{n}} v)_{\partial K} + 2(\partial_{\mathbf{n} \mathbf{n} \mathbf{t}} w, \partial_{\mathbf{n} \mathbf{t}} v)_{\partial K} + (\partial_{\mathbf{n} \mathbf{t} \mathbf{t}} w, \partial_{\mathbf{t} \mathbf{t}} v)_{\partial K} \right). \end{aligned} \quad (4.9)$$

Due to the fact that  $v \in H_0^1(\Omega) \cap C^0(\bar{\Omega})$  and  $w \in H_0^3(\Omega)$ , it holds

$$\begin{aligned} -(\Delta^3 w, v) &= \sum_{K \in \mathcal{T}_h} (\nabla^3 w, \nabla^3 v)_K \\ &\quad + \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\{\{\partial_{\mathbf{n} \mathbf{n}} \Delta w\}\}, [\partial_{\mathbf{n}} v])_e + \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{\mathbf{n}} w], \{\{\partial_{\mathbf{n} \mathbf{n}} \Delta v\}\})_e \\ &\quad - \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\{\{\partial_{\mathbf{n} \mathbf{n} \mathbf{n}} w\}\}, [\partial_{\mathbf{n} \mathbf{n}} v])_e - \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{\mathbf{n} \mathbf{n}} w], \{\{\partial_{\mathbf{n} \mathbf{n} \mathbf{n}} v\}\})_e \\ &\quad - 2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\{\{\partial_{\mathbf{n} \mathbf{n} \mathbf{t}} w\}\}, [\partial_{\mathbf{n} \mathbf{t}} v])_e - 2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{\mathbf{n} \mathbf{t}} w], \{\{\partial_{\mathbf{n} \mathbf{n} \mathbf{t}} v\}\})_e \\ &\quad + \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} ([\partial_{\mathbf{n} \mathbf{n}} w], [\partial_{\mathbf{n} \mathbf{n}} v])_e + \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-3} ([\partial_{\mathbf{n}} w], [\partial_{\mathbf{n}} v])_e. \end{aligned} \quad (4.10)$$

Here,  $\eta > 0$  is a sufficiently large constant.

Besides, from (4.4), (4.8) and (4.3), one has

$$(\Delta^2 w, v) = \sum_{K \in \mathcal{T}_h} (\nabla^2 w, \nabla^2 v)_K - \sum_{K \in \mathcal{T}_h} (\partial_{\mathbf{n} \mathbf{n}} w, \partial_{\mathbf{n}} v)_{\partial K} - \sum_{K \in \mathcal{T}_h} (\partial_{\mathbf{n} \mathbf{t}} w, \partial_{\mathbf{t}} v)_{\partial K}. \quad (4.11)$$

Then, it holds

$$\begin{aligned} (\Delta^2 w, v) &= \sum_{K \in \mathcal{T}_h} (\nabla^2 w, \nabla^2 v)_K - \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\{\{\partial_{\mathbf{n} \mathbf{n}} w\}\}, [\partial_{\mathbf{n}} v])_e \\ &\quad - \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{\mathbf{n}} w], \{\{\partial_{\mathbf{n} \mathbf{n}} v\}\})_e + \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} ([\partial_{\mathbf{n}} w], [\partial_{\mathbf{n}} v])_e. \end{aligned}$$

For any  $w, v \in V_h + H^3(\Omega)$ , define

$$\begin{aligned}
a_h(w, v) &= \sum_{K \in \mathcal{T}_h} (\nabla^3 w, \nabla^3 v)_K \\
&\quad - \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\{\{\partial_{nnn} w\}\}, [\partial_{nn} v])_e - \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{nn} w], \{\{\partial_{nnn} v\}\})_e \\
&\quad - 2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\{\{\partial_{nnt} w\}\}, [\partial_{nt} v])_e - 2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{nt} w], \{\{\partial_{nnt} v\}\})_e \\
&\quad + \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} ([\partial_{nn} w], [\partial_{nn} v])_e + \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-3} ([\partial_n w], [\partial_n v])_e, \\
b_h(w, v) &= \sum_{K \in \mathcal{T}_h} (\nabla^2 w, \nabla^2 v)_K - \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\{\{\partial_{nn} w\}\}, [\partial_n v])_e \\
&\quad - \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_n w], \{\{\partial_{nn} v\}\})_e + \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} ([\partial_n w], [\partial_n v])_e.
\end{aligned}$$

Here, the positive penalty parameter  $\eta$  is independent of  $h$  and  $\iota$ . Then the discretization formulation of (3.2) is to find  $w_h \in V_h$  such that

$$\iota^2 a_h(w_h, v_h) + b_h(w_h, v_h) = (f, v_h) \quad \forall v_h \in V_h. \quad (4.12)$$

Define the weighted norm on  $V_h$  by

$$\|v_h\|_{\iota, h}^2 := \|v_h\|_{2, h}^2 + \iota^2 \|v_h\|_{3, h}^2,$$

where,

$$\begin{aligned}
\|v_h\|_{2, h}^2 &:= |v_h|_{2, h}^2 + \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} \|[\partial_n v_h]\|_{0, e}^2, \\
\|v_h\|_{3, h}^2 &:= |v_h|_{3, h}^2 + \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} \|[\partial_{nn} v_h]\|_{0, e}^2 + \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-3} \|[\partial_n v_h]\|_{0, e}^2, \\
|v_h|_{2, h}^2 &:= \sum_{K \in \mathcal{T}_h} |v_h|_{2, K}^2, \quad |v_h|_{3, h}^2 := \sum_{K \in \mathcal{T}_h} |v_h|_{3, K}^2.
\end{aligned}$$

**Lemma 4.1.** *The discrete bilinear form  $\iota^2 a_h(\cdot, \cdot) + b_h(\cdot, \cdot)$  is continuous on  $V_h \times V_h$  and the discrete coercivity holds:*

$$\iota^2 a_h(v, v) + b_h(v, v) \geq \frac{1}{2} \|v\|_{\iota, h}^2 \quad \forall v \in V_h. \quad (4.13)$$

*Proof.* Using the inverse inequality, one has

$$\|[\partial_{nt} v]\|_{0, e} \lesssim h_e^{-1} \|[\partial_n v]\|_{0, e}. \quad (4.14)$$

By the inverse inequality, (2.2) and (4.14), there exists a generic constant  $C_1 > 0$  dependent only on  $\gamma_0$  such that

$$\begin{aligned}
& - 2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\{\{\partial_{nnn} v\}\}, [\partial_{nn} v])_e - 4 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\{\{\partial_{nnt} w\}\}, [\partial_{nt} v])_e \\
& \leq 2 \left( \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e \| \{\{\partial_{nnn} v\}\} \|_{0, e}^2 \right)^{1/2} \left( \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} \| [\partial_{nn} v] \|_{0, e}^2 \right)^{1/2} \\
& \quad + 4 \left( \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e \| \{\{\partial_{nnt} v\}\} \|_{0, e}^2 \right)^{1/2} \left( \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} \| [\partial_{nt} v] \|_{0, e}^2 \right)^{1/2} \\
& \leq \frac{1}{2} |v|_{3, h}^2 + C_1 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} \left( h_e^{-1} \| [\partial_{nn} v] \|_{0, e}^2 + h_e^{-3} \| [\partial_n v] \|_{0, e}^2 \right). \quad (4.15)
\end{aligned}$$

Similarly, it follows from (2.2) and the Cauchy-Schwarz inequality that

$$-2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\{\{\partial_{nn}v\}\}, [\partial_n v])_e \leq \frac{1}{2} |v|_{2,h}^2 + C_2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} \|[\partial_n v]\|_{0,e}^2, \quad (4.16)$$

where  $C_2 > 0$  is a generic constant dependent only on  $\gamma_0$ . Combining with (4.15)-(4.16), we have

$$\begin{aligned} \iota^2 a_h(v, v) + b_h(v, v) &\geq \frac{1}{2} \iota^2 |v|_{3,h}^2 + \iota^2 (\eta - C_1) \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} \|[\partial_{nn}v]\|_{0,e}^2 \\ &\quad + \iota^2 (\eta - C_1) \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-3} \|[\partial_n v]\|_{0,e}^2 \\ &\quad + \frac{1}{2} |v|_{2,h}^2 + (\eta - C_2) \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} \|[\partial_n v]\|_{0,e}^2. \end{aligned}$$

The discrete coercivity is derived by setting  $\eta$  as  $\frac{1}{2} + \max\{C_1, C_2\}$  in the last inequality.

We omit the proof of the continuity of  $\iota^2 a_h(\cdot, \cdot) + b_h(\cdot, \cdot)$ , which is easy to check by (4.14), the inverse inequality and the Cauchy-Schwarz inequality.  $\square$

**Remark 4.2.** As shown in [25, 31], we count the global DOFs in view of the relation:

$$\lim_{h \rightarrow 0} \frac{N_e}{N_v} = 3, \quad \text{and} \quad \lim_{h \rightarrow 0} \frac{N_T}{N_v} = 2.$$

where  $N_e = \#\mathcal{E}(\mathcal{T}_h)$ ,  $N_v = \#\mathcal{V}(\mathcal{T}_h)$ ,  $N_T = \#\mathcal{T}_h$ . In [23], Gudi use the cubic Lagrange element to approximate the sixth-order problem, which has the same number of the local DoFs as the Hermite element in a triangle. However, the number of the global DoFs for the cubic Lagrange element space is  $9N_v$ , while one for  $V_h$  is  $5N_v$ .

## 5. Error analysis

### 5.1. The quasi interpolation operator

In order to derive some robust and optimal error estimates, we first define an weak interpolation operator, which is motivated by the definition of the average projection operators in [49, Sect. 3.5]. Denote the weak interpolation operator from  $H_0^2(\Omega)$  to  $V_h$  by  $I_h$ . Then, for any  $v \in H_0^2(\Omega)$ , the DoFs of  $I_h v$  are given as:

$$I_h v(\zeta) = \frac{1}{\#\mathcal{T}_\zeta} \sum_{K' \in \mathcal{T}_\zeta} (\Pi_{K'}^3 v)(\zeta), \quad (5.1)$$

$$\nabla I_h v(\zeta) = \frac{1}{\#\mathcal{T}_\zeta} \sum_{K' \in \mathcal{T}_\zeta} (\nabla \Pi_{K'}^3 v)(\zeta), \quad (5.2)$$

$$I_h v(\zeta_K) = v(\zeta_K), \quad (5.3)$$

where  $\zeta \in \mathcal{V}^i(\mathcal{T}_h)$ ,  $K \in \mathcal{T}_h$  and DoFs (5.1)-(5.2) vanish on boundary. Following the ideas for proving Theorem 3.5.3 in [49], we have

$$|v - I_h v|_{m,h} \lesssim h^{s-m} |v|_s \quad \forall v \in H^s(\Omega) \cap H_0^2(\Omega), \quad 0 \leq m \leq s, \quad 2 \leq s \leq 4. \quad (5.4)$$

**Lemma 5.1.** Let  $w \in V \cap H^4(\Omega)$  and  $w_0 \in H_0^2(\Omega) \cap H^s(\Omega)$  ( $s = 3, 4$ ) be the solutions of problems (3.2) and (3.4), respectively. Then, it holds

$$\|w - I_h w\|_{\iota,h} \lesssim \iota^{1/2} \|f\|_0 + h^{s-2} |w_0|_s, \quad (5.5)$$

$$\|w - I_h w\|_{\iota,h} \lesssim (\iota h^{s-3} + h^{s-2}) |w|_s. \quad (5.6)$$

*Proof.* By the trace inequality (2.1), we have

$$\begin{aligned} & \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} \|[\partial_{\mathbf{nn}}(w - I_h w)]\|_e^2 + \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-3} \|[\partial_{\mathbf{n}}(w - I_h w)]\|_e^2 \\ & \lesssim \sum_{K \in \mathcal{T}_h} (h_K^{-4} |w - I_h w|_{1,K}^2 + h_K^{-2} |w - I_h w|_{2,K}^2 + |w - I_h w|_{3,K}^2), \end{aligned}$$

which combined with (5.4) and (3.6) yields

$$\iota^2 \| \|w - I_h w\|_{3,h}^2 \lesssim \iota^2 |w|_3^2 \lesssim \iota \|f\|_0^2, \quad (5.7)$$

$$\iota^2 \| \|w - I_h w\|_{3,h}^2 \lesssim \iota^2 h^{2s-6} |w|_s^2. \quad (5.8)$$

Similarly, employing the trace inequality (2.1) and (5.4), one gets

$$\| \|w - I_h w\|_{2,h}^2 \lesssim \sum_{K \in \mathcal{T}_h} (h_K^{-2} |w - I_h w|_{1,K}^2 + |w - I_h w|_{2,K}^2) \lesssim h^{2s-4} |w|_4^2. \quad (5.9)$$

On the other hand, combining the trace inequality (2.1), (5.4) and (3.6) leads to

$$\begin{aligned} \| \|w - I_h w\|_{2,h}^2 & \lesssim \sum_{K \in \mathcal{T}_h} (h_K^{-2} |w - I_h w|_{1,K}^2 + |w - I_h w|_{2,K}^2) \\ & \lesssim \sum_{K \in \mathcal{T}_h} (h_K^{-2} |w - w_0 - I_h(w - w_0)|_{1,K}^2 + |w - w_0 - I_h(w - w_0)|_{2,K}^2) \\ & \quad + \sum_{K \in \mathcal{T}_h} (h_K^{-2} |w_0 - I_h w_0|_{1,K}^2 + |w_0 - I_h w_0|_{2,K}^2) \\ & \lesssim |w - w_0|_2^2 + h^{2s-4} |w_0|_s^2 \lesssim \iota \|f\|_0^2 + h^{2s-4} |w_0|_s^2. \end{aligned} \quad (5.10)$$

Finally, the estimate (5.5) can be derived by (5.7) and (5.10), while the estimate (5.6) is derived by (5.8) and (5.9).  $\square$

## 5.2. The enriching operator

To obtain the error estimate under minimal regularity of  $w$ , we also need to construct an enriching operator connecting  $V_h$  to a  $H^3$  conforming finite element space. Hence, we first introduce a  $C^2$ -finite element (cf. [8, 51]). Its shape function space is chosen as  $V^c(K) := \mathbb{P}_9(K)$  and the DoFs consist of:

$$\begin{cases} \nabla^\alpha v(\zeta) & \forall \zeta \in \mathcal{V}(K), \alpha = 0, 1, 2, 3, 4, \\ \partial_{\mathbf{nn}} v(\zeta_{e,1}), \partial_{\mathbf{nn}} v(\zeta_{e,2}) & \forall e \in \mathcal{E}(K), \\ \partial_{\mathbf{n}} v(\zeta_e) & \forall e \in \mathcal{E}(K) \\ v(\zeta_K). \end{cases} \quad (5.11)$$

Here,  $\zeta_K$  is the barycenter of  $K$  and for each edge  $e = \overline{a_1^e a_2^e} \in \mathcal{E}(K)$  with  $a_1^e, a_2^e \in \mathcal{V}(K)$ ,  $\zeta_e = (a_1^e + a_2^e)/2$  is the midpoint of  $e$ ,  $\zeta_{e,1} = (a_1^e + 2a_2^e)/3$ ,  $\zeta_{e,2} = (2a_1^e + a_2^e)/3$ . Set

$$W_h^c = \{v \in H^3(\Omega) : v|_K \in V^c(K) \text{ for all } K \in \mathcal{T}_h, \nabla^\alpha v \text{ is continuous at all vertices of triangulation } \mathcal{T}_h \text{ with } \alpha = 0, 1, 2, 3, 4, \partial_{\mathbf{n}} v \text{ is continuous at the midpoints of } e \in \mathcal{E}(K), \partial_{\mathbf{nn}} v \text{ is continuous at the points of trisection of } e \in \mathcal{E}(K) \text{ and all DoFs (5.11) vanish on boundary}\}.$$

Then, it holds  $W_h^c \subset H_0^3(\Omega)$  and

$$[\partial_{\mathbf{nn}} v]|_e = [\partial_{\mathbf{n}} v]|_e = [\partial_{\mathbf{nt}} v]|_e = [\partial_{\mathbf{tt}} v]|_e = 0 \quad \forall v \in W_h^c, e \in \mathcal{E}(\mathcal{T}_h). \quad (5.12)$$

The enriching operator  $E_h^c : V_h \rightarrow W_h^c$  is defined as follows:

$$\nabla^\alpha E_h^c v(\zeta) = \frac{1}{\#\mathcal{T}_\zeta} \sum_{K' \in \mathcal{T}_\zeta} (\nabla^\alpha v|_{K'}) (\zeta), \quad \alpha = 0, 1, 2, 3, 4, \quad (5.13)$$

$$\partial_{nn} E_h^c v(\zeta_{e,i}) = \frac{1}{\#\mathcal{T}_e} \sum_{K' \in \mathcal{T}_e} \partial_{nn}(v|_{K'}) (\zeta_{e,i}), \quad i = 1, 2, \quad (5.14)$$

$$\partial_n E_h^c v(\zeta_e) = \frac{1}{\#\mathcal{T}_e} \sum_{K' \in \mathcal{T}_e} \partial_n(v|_{K'}) (\zeta_e), \quad (5.15)$$

$$E_h^c v(\zeta_K) = v(\zeta_K), \quad (5.16)$$

where,  $K \in \mathcal{T}_h$ ,  $e \in \mathcal{E}^i(\mathcal{T}_h)$  and  $\zeta \in \mathcal{V}^i(\mathcal{T}_h)$ . Noted that the DoFs of  $E_h^c v$  vanish on boundary  $\partial\Omega$ .

**Lemma 5.2.** *For any  $v \in V_h$ , it holds*

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} h_K^{-4} \|v - E_h^c v\|_{0,K}^2 + \sum_{K \in \mathcal{T}_h} h_K \|\partial_{nn}(v - E_h^c v)\|_{0,\partial K}^2 \\ + \sum_{K \in \mathcal{T}_h} h_K^{-1} \|\partial_n(v - E_h^c v)\|_{0,\partial K}^2 \lesssim \|v\|_{2,h}^2, \end{aligned} \quad (5.17)$$

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} h_K^{-6} \|v - E_h^c v\|_{0,K}^2 + \sum_{K \in \mathcal{T}_h} h_K \|\partial_{nnn}(v - E_h^c v)\|_{0,\partial K}^2 \\ + \sum_{K \in \mathcal{T}_h} h_K^{-1} \|\partial_{nn}(v - E_h^c v)\|_{0,\partial K}^2 + \sum_{K \in \mathcal{T}_h} h_K^{-3} \|\partial_n(v - E_h^c v)\|_{0,\partial K}^2 \lesssim \|v\|_{3,h}^2, \end{aligned} \quad (5.18)$$

$$\|E_h^c v\|_{l,h} \lesssim \|v\|_{l,h}. \quad (5.19)$$

*Proof.* By (5.13)-(5.16) and scaling argument, we have

$$\begin{aligned} \|v - E_h^c v\|_{0,K}^2 &\lesssim h_K^{2+2\alpha} \sum_{\alpha=2}^4 \sum_{\zeta \in \mathcal{V}(K)} |\nabla^\alpha v|_K - \nabla^\alpha E_h^c v|^2(\zeta) + h_K^4 \sum_{e \in \mathcal{E}(K)} |\partial_n v|_K - \partial_n E_h^c v|^2(\zeta_e) \\ &\quad + h_K^6 \sum_{i=1}^2 \sum_{e \in \mathcal{E}(K)} |\partial_{nn} v|_K - \partial_{nn} E_h^c v|^2(\zeta_{e,i}) \\ &\lesssim h_K^{1+2\alpha} \sum_{\alpha=2}^4 \sum_{\zeta \in \mathcal{V}(K)} \sum_{e \in \mathcal{E}(\mathcal{T}_h), \zeta \in \partial e} \|[\nabla^\alpha v]\|_{0,e}^2 + h_K^3 \sum_{e \in \mathcal{E}(K)} \|[\partial_n v]\|_{0,e}^2 \\ &\quad + h_K^5 \sum_{e \in \mathcal{E}(K)} \|[\partial_{nn} v]\|_{0,e}^2, \end{aligned} \quad (5.20)$$

Applying the inverse inequality, (5.20) and (2.2), we can get

$$\begin{aligned} \|v - E_h^c v\|_{0,K}^2 &\lesssim h_K^{2\alpha} \sum_{\alpha=2}^4 \sum_{\zeta \in \mathcal{V}(K)} \sum_{K' \in \mathcal{T}_\zeta} \|\nabla^\alpha v\|_{0,K'}^2 + h_K^3 \sum_{e \in \mathcal{E}(K)} \|[\partial_n v]\|_{0,e}^2 \\ &\lesssim h_K^4 \sum_{\zeta \in \mathcal{V}(K)} \sum_{K' \in \mathcal{T}_\zeta} |v|_{2,K'}^2 + h_K^3 \sum_{e \in \mathcal{E}(K)} \|[\partial_n v]\|_{0,e}^2. \end{aligned} \quad (5.21)$$

On the other hand, due to the fact that  $V_h \in H_0^1(\Omega) \cap C(\bar{\Omega})$  and using (4.5) and the inverse inequality, we can obtain

$$\|[\nabla^2 v]\|_{0,e} \lesssim \|[\partial_{nn} v]\|_{0,e} + \|[\partial_{nt} v]\|_{0,e} \lesssim \|[\partial_{nn} v]\|_{0,e} + h_e^{-1} \|[\partial_n v]\|_{0,e},$$

and

$$\|\nabla^4 v\|_{0,K} \lesssim h_K^{-1} \|\nabla^3 v\|_{0,K},$$

which together with (5.20) yields

$$\begin{aligned} \|v - E_h^c v\|_{0,K}^2 &\lesssim h_K^6 \sum_{\zeta \in \mathcal{V}(K)} \sum_{K' \in \mathcal{T}_\zeta} |v|_{3,K'}^2 + h_K^3 \sum_{e \in \mathcal{E}(K)} \|[\partial_{\mathbf{n}} v]\|_{0,e}^2 \\ &\quad + h_K^5 \sum_{e \in \mathcal{E}(K)} \|[\partial_{\mathbf{nn}} v]\|_{0,e}^2. \end{aligned} \quad (5.22)$$

Summing over all triangles in  $\mathcal{T}_h$  and employing (2.2) and the inverse inequality, we can derive the estimate (5.17) from (5.21), and we can derive the estimate (5.18) from (5.22). Moreover, the inequality (5.19) can be obtained by (5.17), (5.18) and the inverse inequality.  $\square$

### 5.3. Error estimates

**Lemma 5.3.** *For any  $e \in \mathcal{E}^i(\mathcal{T}_h)$  and  $v_h \in V_h$ , it holds*

$$\|f\|_{0,K} \lesssim \|f - \Pi_K^0 f\|_{0,K} + \iota^2 h_K^{-3} |w - v_h|_{3,K} + h_K^{-2} |w - v_h|_{2,K}, \quad (5.23)$$

$$\begin{aligned} \iota^4 \|[\partial_{\mathbf{nnn}} v_h]\|_{0,e}^2 &\lesssim h_e^5 \sum_{K \in \mathcal{T}_e} \|f - \Pi_K^0 f\|_{0,K}^2 + \iota^4 h_e^{-1} \sum_{K \in \mathcal{T}_e} |w - v_h|_{3,K}^2 \\ &\quad + h_e \sum_{K \in \mathcal{T}_e} |w - v_h|_{2,K}^2, \end{aligned} \quad (5.24)$$

$$\begin{aligned} \|[\partial_{\mathbf{nn}} v_h]\|_{0,e}^2 &\lesssim h_e^3 \sum_{K \in \mathcal{T}_e} \|f - \Pi_K^0 f\|_{0,K}^2 + \iota^4 h_e^{-3} \sum_{K \in \mathcal{T}_e} |w - v_h|_{3,K}^2 \\ &\quad + h_e^{-1} \sum_{K \in \mathcal{T}_e} |w - v_h|_{2,K}^2. \end{aligned} \quad (5.25)$$

*Proof.* Set

$$\phi_K = \begin{cases} b_K^3 \Pi_K^0 f & \text{in } K, \\ 0 & \text{in } \Omega \setminus K. \end{cases}$$

Obviously,  $\phi_K|_K \in H_0^3(K)$  and we have from scaling argument that

$$\|\phi_K\|_{0,K}^2 \lesssim \|\Pi_K^0 f\|_0^2 \lesssim (\Pi_K^0 f, \phi_K)_K. \quad (5.26)$$

By integration by parts and (3.2), it holds for any  $v_h \in V_h$  that

$$\begin{aligned} \|\Pi_K^0 f\|_{0,K}^2 &\lesssim (\Pi_K^0 f - f, \phi_K)_K + (f - \Delta^2 v_h + \iota^2 \Delta^3 v_h, \phi_K)_K \\ &= (\Pi_K^0 f - f, \phi_K)_K + \iota^2 (\nabla^3 w - \nabla^3 v_h, \nabla^3 \phi_K)_K + (\nabla^2 w - \nabla^2 v_h, \nabla^2 \phi_K)_K. \end{aligned}$$

Then, using the inverse inequality and the Cauchy-Schwarz inequality, we can obtain

$$\|\Pi_K^0 f\|_{0,K}^2 \lesssim (\|f - \Pi_K^0 f\|_{0,K} + \iota^2 h_K^{-3} |w - v_h|_{3,K} + h_K^{-2} |w - v_h|_{2,K}) \|\phi_K\|_{0,K}.$$

We get (5.23) from (5.26) and the triangle inequality readily.

For each interior edge  $e \in \mathcal{E}^i(\mathcal{T}_h)$ , there exist two adjacent triangles,  $K_1$  and  $K_2$  in  $\mathcal{T}_h$  sharing a common edge  $e$  and define a bubble function as follow:

$$\varrho_e = \begin{cases} \lambda_{K_1,1} \lambda_{K_1,2} \lambda_{K_2,1} \lambda_{K_2,2} & \text{in } \omega_e, \\ 0 & \text{in } \Omega \setminus \omega_e, \end{cases}$$

where,  $\lambda_{K_1,i}$  and  $\lambda_{K_2,i}$  are barycentric coordinates of  $K_1$  and  $K_2$  related to the vertices of  $e$ ,  $i = 1, 2$ . Besides, the line where  $e$  lying in can be represented by  $\mathbf{l}_e = \mathbf{n}_e \cdot \mathbf{x} + C_e$  with a constant  $C_e$ . A simple calculation shows  $\mathbf{l}_e|_e = 0$ ,  $\partial_{\mathbf{n}_e} \mathbf{l}_e|_e = 1$  and

$$|\mathbf{l}_e|_{\omega_e} = h_e \left| \frac{(\mathbf{n}_e \cdot \mathbf{x} + C_e)|_{\omega_e}}{h_e} \right| \lesssim h_e. \quad (5.27)$$

We now prove (5.24). Define  $J_{1,\omega_e}$  by extending the jump  $[[\partial_{nnn}v_h]]|_e$  to  $\omega_e$  constantly along the normal to  $e$ . Take  $\phi_e = \varrho_e^3 J_{1,\omega_e} \mathbf{l}_e^2$ . Then,  $\phi_e \in H_0^3(\omega_e)$ ,  $\phi_e|_e = \partial_{\mathbf{n}_e} \phi_e|_e = 0$  and

$$\partial_{nn} \phi_e|_e = (\varrho_e^3 J_{1,\omega_e} \partial_{nn} \mathbf{l}_e^2)|_e = 2\varrho_e^3 [[\partial_{nnn}v_h]]|_e.$$

By integrating by parts, (3.1), scaling argument and the similar arguments for obtaining (4.9), it holds for any  $e \in \mathcal{E}^i(\mathcal{T}_h)$  that

$$\begin{aligned} \iota^2 \|[[\partial_{nnn}v_h]]\|_{0,e}^2 &\lesssim \iota^2 ([[ \partial_{nnn}v_h ], \partial_{nn} \phi_e)_e = \iota^2 \sum_{K \in \mathcal{T}_e} (\partial_{nnn}v_h, \partial_{nn} \phi_e)_{\partial K} \\ &= \iota^2 \sum_{K \in \mathcal{T}_e} (\nabla^3(v_h - w), \nabla^3 \phi_e)_K + (f, \phi_e)_{\omega_e} - (\Delta^2 w, \phi_e)_{\omega_e} \\ &= \iota^2 \sum_{K \in \mathcal{T}_e} (\nabla^3(v_h - w), \nabla^3 \phi_e)_K + (f, \phi_e)_{\omega_e} \\ &\quad - \sum_{K \in \mathcal{T}_e} (\nabla^2 v_h, \nabla^2 \phi_e)_K + \sum_{K \in \mathcal{T}_e} (\nabla^2(v_h - w), \nabla^2 \phi_e)_K, \end{aligned} \quad (5.28)$$

Noted that  $v_h|_K \in \mathbb{P}_3(K)$ ,  $\phi_e \in H_0^3(\Omega)$  and  $\phi_e|_e = \partial_{\mathbf{n}_e} \phi_e|_e = 0$ . Using integration by parts, we have

$$\sum_{K \in \mathcal{T}_e} (\nabla^2 v_h, \nabla^2 \phi_e)_K = \sum_{K \in \mathcal{T}_e} (\Delta^2 v_h, \phi_e)_K + \sum_{\hat{e} \in \partial K, K \in \mathcal{T}_e} ((\nabla^2 v_h, \partial_n \phi_e)_{\hat{e}} - (\partial_n \Delta v_h, \phi_e)_{\hat{e}}) = 0,$$

which combined with (5.28) and the inverse inequality implies

$$\begin{aligned} &\iota^2 \|[[\partial_{nnn}v_h]]\|_{0,e}^2 \\ &\lesssim \iota^2 \sum_{K \in \mathcal{T}_e} |v_h - w|_{3,K} |\phi_e|_{3,K} + \sum_{K \in \mathcal{T}_e} |v_h - w|_{2,K} |\phi_e|_{2,K} + \|f\|_{0,\omega_e} \|\phi_e\|_{0,\omega_e} \\ &\lesssim (\iota^4 h_e^{-6} \sum_{K \in \mathcal{T}_e} |v_h - w|_{3,K}^2 + h_e^{-4} \sum_{K \in \mathcal{T}_e} |v_h - w|_{2,K}^2 + \|f\|_{0,\omega_e}^2)^{1/2} \|\phi_e\|_{0,\omega_e}. \end{aligned} \quad (5.29)$$

From (5.27) and the inverse inequality, one has

$$\|\phi_e\|_{0,\omega_e} \lesssim h_e^2 \|J_{1,\omega_e}\|_{0,\omega_e} \lesssim h_e^3 \|[[\partial_{nnn}v_h]]\|_{\infty,e} \lesssim h_e^{5/2} \|[[\partial_{nnn}v_h]]\|_{0,e}. \quad (5.30)$$

Therefore, (5.24) can be achieved by (5.23), (5.29) and (5.30).

Finally, we are going to prove (5.25). For any  $e \in \mathcal{E}^i(\mathcal{T}_h)$ , define  $J_{2,\omega_e}$  by extending the jump  $[[\partial_{nn}v_h]]|_e$  to  $\omega_e$  constantly along the normal to  $e$ . Set  $\varphi_e = \varrho_{e,2}^3 J_{2,\omega_e} \mathbf{l}_e \in H_0^3(\omega_e)$  and it's easy to know that

$$\partial_{\mathbf{n}_e} \varphi_e|_e = \varrho_{e,2}^3 [[\partial_{nn}v_h]]|_e.$$

We get from the inverse inequality and (5.27) that

$$\|\varphi_e\|_{0,\omega_e} \lesssim h_e \|J_{2,\omega_e}\|_{0,\omega_e} \lesssim h_e^2 \|[[\partial_{nn}v_h]]\|_{\infty,e} \lesssim h_e^{3/2} \|[[\partial_{nn}v_h]]\|_{0,e}. \quad (5.31)$$

By scaling argument and the similar arguments for obtaining (4.11),

$$\|[[\partial_{nn}v_h]]\|_{0,e}^2 \lesssim ([[ \partial_{nn}v_h ], \partial_{\mathbf{n}_e} \varphi_e)_e = \sum_{K \in \mathcal{T}_e} (\partial_{nn}v_h, \partial_{\mathbf{n}_e} \varphi_e)_{\partial K} = \sum_{K \in \mathcal{T}_e} (\nabla^2 v_h, \nabla^2 \varphi_e)_K.$$

Then, using integration by parts and (3.1), it holds

$$\begin{aligned} \|\llbracket \partial_{\mathbf{nn}} v_h \rrbracket\|_{0,e}^2 &\lesssim (f, \varphi_e)_{\omega_e} + \sum_{K \in \mathcal{T}_e} (\nabla^2(v_h - w), \nabla^2 \varphi_e)_K \\ &\quad + \iota^2 \sum_{K \in \mathcal{T}_e} (\nabla^3(v_h - w), \nabla^3 \varphi_e)_K - \iota^2 \sum_{K \in \mathcal{T}_e} (\nabla^3 v_h, \nabla^3 \varphi_e)_K. \end{aligned} \quad (5.32)$$

Since  $\varphi_e \in H_0^3(\omega_e)$ , we get from the similar arguments for deriving (4.9) that

$$\iota^2 (\nabla^3 v_h, \nabla^3 \varphi_e)_{\omega_e} = \iota^2 \sum_{K \in \mathcal{T}_e} (\partial_{\mathbf{nnn}} v_h, \partial_{\mathbf{nn}} \varphi_e)_{\partial K} = \iota^2 (\llbracket \partial_{\mathbf{nnn}} v_h \rrbracket, \partial_{\mathbf{nn}} \varphi_e)_e. \quad (5.33)$$

By (5.32)-(5.33) and the inverse inequality, it holds

$$\begin{aligned} \|\llbracket \partial_{\mathbf{nn}} v_h \rrbracket\|_{0,e}^2 &\lesssim \|f\|_{0,\omega_e} \|\varphi_e\|_{0,\omega_e} + \sum_{K \in \mathcal{T}_e} |v_h - w|_{2,K} |\varphi_e|_{2,K} \\ &\quad + \iota^2 \sum_{K \in \mathcal{T}_e} |v_h - w|_{3,K} |\varphi_e|_{3,K} + \iota^2 \|\llbracket \partial_{\mathbf{nnn}} v_h \rrbracket\|_{0,e} \|\partial_{\mathbf{nn}} \varphi_e\|_{0,e} \\ &\lesssim (\|f\|_{0,\omega_e} + \iota^2 h_e^{-5/2} \|\llbracket \partial_{\mathbf{nnn}} v_h \rrbracket\|_{0,e}) \|\varphi_e\|_{0,\omega_e} \\ &\quad + \left( h_e^{-4} \sum_{K \in \mathcal{T}_e} |v_h - w|_{2,K}^2 + \iota^4 h_e^{-6} \sum_{K \in \mathcal{T}_e} |v_h - w|_{3,K}^2 \right)^{1/2} \|\varphi_e\|_{0,\omega_e}, \end{aligned}$$

which together with (5.23), (5.24) and (5.31) yields (5.25) immediately.  $\square$

**Lemma 5.4.** *For any  $w_I, v_h \in V_h$ , it holds*

$$\iota^2 a_h(w_I, v_h - E_h^c v_h) \lesssim (\text{Osc}_h(f) + \|w - w_I\|_{L,h}) \|v_h\|_{L,h},$$

where

$$\text{Osc}_h(f)^2 = \sum_{K \in \mathcal{T}_h} h_K^4 \|f - \Pi_K^0 f\|_{0,K}^2. \quad (5.34)$$

*Proof.* Let  $\beta_{v_h} = v_h - E_h^c v_h$ . Employing the same arguments for proving (4.9), one gets

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} (\nabla^3 w_I, \nabla^3 \beta_{v_h})_K &= \sum_{K \in \mathcal{T}_h} (\partial_{\mathbf{nnn}} w_I, \partial_{\mathbf{nn}} \beta_{v_h})_{\partial K} + 2 \sum_{K \in \mathcal{T}_h} (\partial_{\mathbf{ntt}} w_I, \partial_{\mathbf{nt}} \beta_{v_h})_{\partial K} \\ &\quad + \sum_{K \in \mathcal{T}_h} (\partial_{\mathbf{ntt}} w_I, \partial_{\mathbf{tt}} \beta_{v_h})_{\partial K}. \end{aligned}$$

Then, recalling the definition of the bilinear form  $a_h(\cdot, \cdot)$ , we get

$$\begin{aligned} &a_h(w_I, \beta_{v_h}) \\ &= \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} (\llbracket \partial_{\mathbf{nnn}} w_I \rrbracket, \{\{\partial_{\mathbf{nn}} \beta_{v_h}\}\})_e + 2 \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} (\llbracket \partial_{\mathbf{ntt}} w_I \rrbracket, \{\{\partial_{\mathbf{nt}} \beta_{v_h}\}\})_e \\ &\quad + \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} (\llbracket \partial_{\mathbf{ntt}} w_I \rrbracket, \{\{\partial_{\mathbf{tt}} \beta_{v_h}\}\})_e - \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\llbracket \partial_{\mathbf{nn}} w_I \rrbracket, \{\{\partial_{\mathbf{nnn}} \beta_{v_h}\}\})_e \\ &\quad - 2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\llbracket \partial_{\mathbf{nt}} w_I \rrbracket, \{\{\partial_{\mathbf{ntt}} \beta_{v_h}\}\})_e + \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} (\llbracket \partial_{\mathbf{nn}} w_I \rrbracket, \llbracket \partial_{\mathbf{nn}} \beta_{v_h} \rrbracket)_e \\ &\quad + \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-3} (\llbracket \partial_{\mathbf{n}} w_I \rrbracket, \llbracket \partial_{\mathbf{n}} \beta_{v_h} \rrbracket)_e. \end{aligned} \quad (5.35)$$

Combining the Cauchy-Schwarz inequality, (5.24) and (5.17)-(5.18), we find

$$\begin{aligned}
& \iota^2 \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} ([\partial_{nnn} w_I], \{\{\partial_{nn} \beta_{v_h}\}\})_e \\
& \lesssim \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} \left( h_e^4 \|f - \Pi_K^0 f\|_{0,\omega_e}^2 + \sum_{K \in \mathcal{T}_e} |w - v_h|_{2,\omega_e}^2 \right)^{1/2} h_e^{1/2} \|\{\{\partial_{nn} \beta_{v_h}\}\}\|_{0,e} \\
& \quad + \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} \iota^2 h_e^{-1/2} \left( \sum_{K \in \mathcal{T}_e} |w - v_h|_{3,K}^2 \right)^{1/2} \|\{\{\partial_{nn} \beta_{v_h}\}\}\|_{0,e} \\
& \lesssim (Osc_h(f) + |w - v_h|_{2,h}) \|v_h\|_{2,h} + \iota^2 |w - v_h|_{3,h} \|v_h\|_{3,h}. \tag{5.36}
\end{aligned}$$

By the Cauchy-Schwarz inequality and the inverse inequality, it holds

$$\begin{aligned}
& 2\iota^2 \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} ([\partial_{nnt} w_I], \{\{\partial_{nt} \beta_{v_h}\}\})_e - 2\iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{nt} w_I], \{\{\partial_{nnt} \beta_{v_h}\}\})_e \\
& \lesssim \iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} \left( h_e^{-2} \|[\partial_{nn} w_I]\|_{0,e} \|\{\{\partial_n \beta_{v_h}\}\}\|_{0,e} + h_e^{-2} \|[\partial_n w_I]\|_{0,e} \|\{\{\partial_{nn} \beta_{v_h}\}\}\|_{0,e} \right) \\
& = \iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-2} \|[\partial_{nn}(w_I - w)]\|_{0,e} \|\{\{\partial_n \beta_{v_h}\}\}\|_{0,e} \\
& \quad + \iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-2} \|[\partial_n(w_I - w)]\|_{0,e} \|\{\{\partial_{nn} \beta_{v_h}\}\}\|_{0,e}. \tag{5.37}
\end{aligned}$$

Here we have used the fact that

$$[\partial_{nn} w]|_e = [\partial_n w]|_e = 0 \quad \forall e \in \mathcal{E}(\mathcal{T}_h), \tag{5.38}$$

which also implies

$$\begin{aligned}
& -\iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{nnn} w_I], \{\{\partial_{nnn} \beta_{v_h}\}\})_e \\
& \lesssim \iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} \|[\partial_{nn}(w - w_I)]\|_{0,e} \|\{\{\partial_{nnn} \beta_{v_h}\}\}\|_{0,e}. \tag{5.39}
\end{aligned}$$

Similarly, we get from (5.38), the inverse inequality and (2.2) that

$$\begin{aligned}
\iota^2 \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} ([\partial_{ntt} w_I], \{\{\partial_{tt} \beta_{v_h}\}\})_e & \lesssim \iota^2 \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} h_e^{-4} \|[\partial_n w_I]\|_{0,e} \|\{\{\beta_{v_h}\}\}\|_{0,e} \\
& \lesssim \iota^2 \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} h_e^{-9/2} \|[\partial_n(w_I - w)]\|_{0,e} \|\beta_{v_h}\|_{0,\omega_e}. \tag{5.40}
\end{aligned}$$

Using the Cauchy-Schwarz inequality, (5.12) and (5.38), one has

$$\begin{aligned}
& \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} ([\partial_{nn} w_I], [\partial_{nn} \beta_{v_h}])_e + \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-3} ([\partial_n w_I], [\partial_n \beta_{v_h}])_e \\
& = \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} ([\partial_{nn}(w_I - w)], [\partial_{nn} \beta_{v_h}])_e + \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-3} ([\partial_n(w_I - w)], [\partial_n \beta_{v_h}])_e \\
& \lesssim \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} \|[\partial_{nn}(w_I - w)]\|_{0,e} \|[\partial_{nn} \beta_{v_h}]\|_{0,e} \\
& \quad + \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-3} \|[\partial_n(w_I - w)]\|_{0,e} \|[\partial_n \beta_{v_h}]\|_{0,e}. \tag{5.41}
\end{aligned}$$

Finally, we end the proof by combining (5.35)-(5.41) and (5.18).  $\square$

**Lemma 5.5.** For any  $w_I, v_h \in V_h$ , it holds

$$(f, v_h - E_h^c v_h) - b_h(w_I, v_h - E_h^c v_h) \lesssim (\text{Osc}_h(f) + \|w - w_I\|_{l,h}) \|v_h\|_{l,h},$$

where,  $\text{Osc}_h(f)$  is determined by (5.34).

*Proof.* Using the similar arguments for deriving (4.11), we have

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} (\nabla^2 w_I, \nabla^2 \beta_{v_h})_K &= \sum_{K \in \mathcal{T}_h} (\partial_{nn} w_I, \partial_n \beta_{v_h})_{\partial K} + \sum_{K \in \mathcal{T}_h} (\partial_{nt} w_I, \partial_t \beta_{v_h})_{\partial K} \\ &= \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} ([\partial_{nn} w_I], \{\{\partial_n \beta_{v_h}\}\})_e + \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\{\{\partial_{nn} w_I\}\}, [\partial_n \beta_{v_h}])_e \\ &\quad + \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} ([\partial_{nt} w_I], \{\{\partial_t \beta_{v_h}\}\})_e. \end{aligned}$$

Then, it holds

$$\begin{aligned} (f, \beta_{v_h}) - b_h(w_I, \beta_{v_h}) &= (f, \beta_{v_h}) - \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} ([\partial_{nn} w_I], \{\{\partial_n \beta_{v_h}\}\})_e \\ &\quad - \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} ([\partial_{nt} w_I], \{\{\partial_t \beta_{v_h}\}\})_e + \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_n w_I], \{\{\partial_{nn} \beta_{v_h}\}\})_e \\ &\quad - \eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} ([\partial_n w_I], [\partial_n \beta_{v_h}])_e. \end{aligned} \tag{5.42}$$

Applying the Cauchy-Schwarz inequality and (5.23), we get

$$\begin{aligned} (f, \beta_{v_h}) &\lesssim \sum_{K \in \mathcal{T}_h} \|f\|_{0,K} \|\beta_{v_h}\|_{0,K} \\ &\lesssim \sum_{K \in \mathcal{T}_h} (h_K^2 \|f - \Pi_K^0 f\|_{0,K} + |w - v_h|_{2,K}) h_K^{-2} \|\beta_{v_h}\|_{0,K} \\ &\quad + \sum_{K \in \mathcal{T}_h} \iota^2 h_K^{-3} |w - v_h|_{3,K} \|\beta_{v_h}\|_{0,K}, \end{aligned}$$

which together with (5.17) and (5.18) yields

$$\begin{aligned} (f, \beta_{v_h}) &\lesssim (\text{Osc}_h(f) + |w - w_I|_{2,h}) \| \|v_h\| \|_{2,h} + \iota^2 |w - w_I|_{3,h} \| \|v_h\| \|_{3,h} \\ &\lesssim (\text{Osc}_h(f) + \|w - w_I\|_{l,h}) \|v_h\|_{l,h}. \end{aligned} \tag{5.43}$$

Similarly, by (5.25), (5.17) and (5.18),

$$\begin{aligned} &- \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} ([\partial_{nn} w_I], \{\{\partial_n \beta_{v_h}\}\})_e \\ &\lesssim \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} \|[\partial_{nn} w_I]\|_{0,e} \|\{\{\partial_n \beta_{v_h}\}\}\|_{0,e} \\ &\lesssim \sum_{e \in \mathcal{E}(\mathcal{T}_h)} \sum_{K \in \mathcal{T}_e} (h_K^4 \|f - \Pi_K^0 f\|_{0,K}^2 + |w - w_I|_{2,K}^2)^{1/2} h_e^{-1/2} \|\{\{\partial_n \beta_{v_h}\}\}\|_{0,e} \\ &\quad + \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} \iota^2 h_e^{-3/2} \left( \sum_{K \in \mathcal{T}_e} |w - w_I|_{3,K}^2 \right)^{1/2} \|\{\{\partial_n \beta_{v_h}\}\}\|_{0,e} \\ &\lesssim (\text{Osc}_h(f) + \|w - w_I\|_{l,h}) \|v_h\|_{l,h}. \end{aligned} \tag{5.44}$$

Similarly, we get from (5.38), the inverse inequality, (2.2) and (5.17) that

$$\begin{aligned}
& - \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} ([\partial_{\mathbf{n}t} w_I], \{\{\partial_t \beta_{v_h}\}\})_e \tag{5.45} \\
& \lesssim \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-2} \|[\partial_{\mathbf{n}} w_I]\|_{0,e} \|\{\{\beta_{v_h}\}\}\|_{0,e} \\
& \lesssim \left( \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} h_e^{-1} \|[\partial_{\mathbf{n}}(w_I - w)]\|_{0,e}^2 \right)^{1/2} \left( \sum_{e \in \mathcal{E}^i(\mathcal{T}_h)} h_e^{-4} \|\beta_{v_h}\|_{0,\omega_e}^2 \right)^{1/2} \\
& \lesssim \|w_I - w\|_{2,h} \|v_h\|_{2,h}. \tag{5.46}
\end{aligned}$$

By (5.12), (5.38) and (5.17), we have

$$\begin{aligned}
& \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{\mathbf{n}} w_I], \{\{\partial_{\mathbf{n}n} \beta_{v_h}\}\})_e \\
& = \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{\mathbf{n}}(w_I - w)], \{\{\partial_{\mathbf{n}n} \beta_{v_h}\}\})_e \\
& \lesssim \left( \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} \|[\partial_{\mathbf{n}}(w_I - w)]\|_{0,e}^2 \right)^{1/2} \left( \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e \|\{\{\partial_{\mathbf{n}n} \beta_{v_h}\}\}\|_{0,e}^2 \right)^{1/2} \\
& \lesssim \|w_I - w\|_{2,h} \|v_h\|_{2,h} \tag{5.47}
\end{aligned}$$

and

$$\begin{aligned}
-\eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} ([\partial_{\mathbf{n}} w_I], [\partial_{\mathbf{n}} \beta_{v_h}])_e & = -\eta \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} ([\partial_{\mathbf{n}}(w_I - w)], [\partial_{\mathbf{n}} v_h])_e \\
& \lesssim \|w_I - w\|_{2,h} \|v_h\|_{2,h}. \tag{5.48}
\end{aligned}$$

Hence, we obtain the desired result by (5.42)-(5.48).  $\square$

**Lemma 5.6.** *Let  $w \in V$  be the solution of problem (3.2). Then for any  $w_I, v_h \in V_h$ , it holds*

$$\iota^2 a_h(w - w_I, E_h^c v_h) + b_h(w - w_I, E_h^c v_h) \lesssim \|w - w_I\|_{\iota,h} \|v_h\|_{\iota,h}. \tag{5.49}$$

*Proof.* For simplicity, let  $\beta_w = w - w_I$ . By (5.12), we can rewrite the discrete bilinear form as

$$\begin{aligned}
& \iota^2 a_h(\beta_w, E_h^c v_h) + b_h(\beta_w, E_h^c v_h) \\
& = \iota^2 \sum_{K \in \mathcal{T}_h} (\nabla^3 \beta_w, \nabla^3 E_h^c v)_K - \iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{\mathbf{n}n} \beta_w], \{\{\partial_{\mathbf{n}n} E_h^c v\}\})_e \\
& \quad - 2\iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{\mathbf{n}t} \beta_w], \{\{\partial_{\mathbf{n}nt} E_h^c v\}\})_e \\
& \quad + \sum_{K \in \mathcal{T}_h} (\nabla^2 \beta_w, \nabla^2 E_h^c v)_K - \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{\mathbf{n}} \beta_w], \{\{\partial_{\mathbf{n}n} E_h^c v\}\})_e. \tag{5.50}
\end{aligned}$$

By (2.2) and the Cauchy-Schwarz inequality, one has

$$\begin{aligned}
& -\iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{\mathbf{n}n} \beta_w], \{\{\partial_{\mathbf{n}n} E_h^c v\}\})_e - \sum_{e \in \mathcal{E}(\mathcal{T}_h)} ([\partial_{\mathbf{n}} \beta_w], \{\{\partial_{\mathbf{n}n} E_h^c v\}\})_e \\
& \lesssim \iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1/2} \|[\partial_{\mathbf{n}n} \beta_w]\|_{0,e} |E_h^c v|_{3,\omega_e} + \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1/2} \|[\partial_{\mathbf{n}} \beta_w]\|_{0,e} |E_h^c v|_{2,\omega_e} \\
& \lesssim \|\beta_w\|_{\iota,h} \|E_h^c v\|_{\iota,h}. \tag{5.51}
\end{aligned}$$

Using the Cauchy-Schwarz inequality again, we have from (5.38) and the inverse inequality that

$$\begin{aligned}
& -2\iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\llbracket \partial_{\mathbf{nt}} \beta_w \rrbracket, \{\{\partial_{\mathbf{nn}t} E_h^c v\}\})_e \\
& = 2\iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} (\llbracket \partial_{\mathbf{nt}} w_I \rrbracket, \{\{\partial_{\mathbf{nn}t} E_h^c v\}\})_e \\
& \lesssim \iota^2 \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-2} \|\llbracket \partial_{\mathbf{n}} w_I \rrbracket\|_{0,e} \|\{\{\partial_{\mathbf{nn}} E_h^c v\}\}\|_{0,e} \\
& \lesssim \iota^2 \left( \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-3} \|\llbracket \partial_{\mathbf{n}} \beta_w \rrbracket\|_{0,e}^2 \right)^{1/2} \left( \sum_{e \in \mathcal{E}(\mathcal{T}_h)} h_e^{-1} \|\partial_{\mathbf{nn}} E_h^c v\|_{0,e}^2 \right)^{1/2} \\
& \lesssim \|\beta_w\|_{\iota,h} \|E_h^c v\|_{\iota,h}. \tag{5.52}
\end{aligned}$$

The inequalities (5.50)-(5.52) implies

$$\iota^2 a_h(\beta_w, E_h^c v_h) + b_h(\beta_w, E_h^c v_h) \lesssim \|w - w_I\|_{\iota,h} \|E_h^c v\|_{\iota,h}.$$

Applying (5.19) to the last inequality leads to the desired result immediately.  $\square$

**Lemma 5.7.** *Let  $w \in V$  and  $w_h \in V_h$  be the solutions of problems (3.2) and (4.12), respectively. Then it holds*

$$\|w - w_h\|_{\iota,h} \lesssim \text{Osc}_h(f) + (1 + \eta) \inf_{w_I \in V_h} \|w - w_I\|_{\iota,h}, \tag{5.53}$$

where,  $\text{Osc}_h(f)$  is determined by (5.34).

*Proof.* For any  $w_I \in V_h$ , let  $\tilde{w}_h = w_h - w_I$ . By Lemma 4.1, (4.12) and (3.2), it follows that

$$\begin{aligned}
\|w_h - w_I\|_{\iota,h}^2 & \lesssim (f, \tilde{w}_h) - \iota^2 a_h(w_I, \tilde{w}_h) - b_h(w_I, \tilde{w}_h) \\
& = (f, \tilde{w}_h - E_h^c \tilde{w}_h) - b_h(w_I, \tilde{w}_h - E_h^c \tilde{w}_h) - \iota^2 a_h(w_I, \tilde{w}_h - E_h^c \tilde{w}_h) \\
& \quad + \iota^2 a_h(w - w_I, E_h^c \tilde{w}_h) + b_h(w - w_I, E_h^c \tilde{w}_h).
\end{aligned}$$

Then, applying Lemma 5.4, Lemma 5.5 and Lemma 5.6 with  $v_h = \tilde{w}_h$  leads to

$$\|w_h - w_I\|_{\iota,h}^2 \lesssim (\text{Osc}_h(f) + (1 + \eta) \|w - w_I\|_{\iota,h}) \|w_h - w_I\|_{\iota,h}. \tag{5.54}$$

The proof is completed by the triangle inequality and (5.54).  $\square$

**Theorem 5.8.** *Let  $w \in V \cap H^4(\Omega)$ ,  $w_0 \in H_0^2(\Omega) \cap H^s(\Omega)$  and  $w_h \in V_h$  be the solutions of problems (3.2), (3.4) and (4.12), respectively. Then the following error estimates hold:*

$$\|w - w_h\|_{\iota,h} \lesssim \text{Osc}_h(f) + (h^2 + \iota h) |w|_4, \tag{5.55}$$

$$\|w - w_h\|_{\iota,h} \lesssim \text{Osc}_h(f) + \iota^{1/2} \|f\|_0 + h^{s-2} |w_0|_s, \tag{5.56}$$

where  $s = 3, 4$  and  $\text{Osc}_h(f)$  is determined by (5.34).

*Proof.* Setting  $w_I$  as  $I_h w$  in (5.53). Then the inequality (5.55) is derived by (5.6) and the inequality (5.56) is obtained by (5.5).  $\square$

**Theorem 5.9.** *Let  $w_0 \in H_0^2(\Omega) \cap H^s(\Omega)$  and  $w_h \in V_h$  be the solutions of problems (3.4) and (4.12), respectively. Then it holds*

$$|w_0 - w_h|_{2,h} + \iota |w_0 - w_h|_{3,h} \lesssim \text{Osc}_h(f) + \iota^{1/2} \|f\|_0 + h^{s-2} |w_0|_s,$$

where  $s = 3, 4$  and  $\text{Osc}_h(f)$  is determined by (5.34).

Table 1: The error  $\|w - w_h\|_{\iota, h}$  for Example 6.1 with  $\eta = 10$ .

$\iota$	$h$					rate
	1/4	1/8	1/16	1/32	1/64	
1	9.133e+01	5.226e+01	2.699e+01	1.361e+01	6.818e+00	1.00
$10^{-2}$	1.090e+01	5.996e+00	2.696e+00	1.326e+00	6.729e-01	0.98
$10^{-4}$	4.513e+00	1.583e+00	4.929e-01	1.711e-01	7.269e-02	1.24
$10^{-6}$	4.398e+00	1.446e+00	3.736e-01	8.883e-02	2.224e-02	2.00
0	4.397e+00	1.444e+00	3.722e-01	8.761e-02	2.113e-02	2.05

*Proof.* Using (3.6) and (3.5), one has

$$|w_0 - w|_{2,h} + \iota |w_0 - w|_{3,h} \lesssim \iota^{1/2} \|f\|_0. \quad (5.57)$$

Applying the triangle inequality and (5.56) we obtain the inequality as required.  $\square$

Lastly, we are going to demonstrate that the method (4.12) is convergent without any additional regularity assumption for the exact solution  $w$ .

**Theorem 5.10.** *Let  $w \in V$  and  $w_h \in V_h$  be the solutions of problems (3.2) and (4.12), respectively. Then we have*

$$\|w - w_h\|_{\iota, h} \rightarrow 0, \quad h \rightarrow 0. \quad (5.58)$$

*Proof.* As is known to all,  $C^\infty(\bar{\Omega}) \cap H^3(\Omega)$  is dense in  $H^3(\Omega)$ . Thus, for any  $\epsilon > 0$ , there exists  $\tilde{w} \in C^\infty(\bar{\Omega})$  such that  $\|w - \tilde{w}\|_3 < \epsilon$  and it holds

$$\begin{aligned} \|w - \tilde{w}\|_{\iota, h}^2 &= |w - \tilde{w}|_2^2 + \iota^2 |w - \tilde{w}|_3^2 \\ &< (1 + \iota^2) \epsilon^2. \end{aligned} \quad (5.59)$$

Notice that  $Osc(f) = O(h^2)$ . Setting  $w_I$  as  $I_h \tilde{w}$  in (5.53) and employing the similar arguments for deriving (5.6), one finds

$$\begin{aligned} \|w - w_h\|_{\iota, h} &\lesssim (1 + \eta) \|w - I_h \tilde{w}\|_{\iota, h} + O(h^2) \\ &\lesssim (1 + \eta) (\|w - \tilde{w}\|_{\iota, h} + \|\tilde{w} - I_h \tilde{w}\|_{\iota, h}) + O(h^2) \\ &\lesssim (1 + \eta) ((1 + \iota) \epsilon + (h + \iota) h |\tilde{w}|_4) + O(h^2), \end{aligned} \quad (5.60)$$

Hence, it holds

$$\limsup_{h \rightarrow 0} \|w - w_h\|_{\iota, h} \lesssim \epsilon, \quad (5.61)$$

and the proof is completed by the arbitrariness of  $\epsilon$  and  $\|w - w_h\|_{\iota, h} \geq 0$ .  $\square$

## 6. Numerical experiments

**Example 6.1.** The exact solution of (3.1) is chosen as

$$w = \sin^3(\pi x) \sin^3(\pi y),$$

and the right side function is determined by (3.1) and  $w$ . The example is used to verify the estimate (5.55). Let  $\eta = 10$ . We display the error  $\|w - w_h\|_{\iota, h}$  in Tab. 1 with different mesh size  $h$  and size parameter  $\iota$ . Besides, we test the robustness of the numerical method with respect to the penalization parameter  $\eta$  in Tab. 2.

Table 2: The performance for Example 6.1.

$\iota$	$\eta$	$h$					rate
		1/4	1/8	1/16	1/32	1/64	
$10^{-8}$	$10^{-4}$	4.200e+00	1.231e+00	3.203e-01	7.985e-02	1.989e-02	2.01
	$10^{-6}$	4.200e+00	1.231e+00	3.203e-01	7.985e-02	1.989e-02	2.01
	1	4.230e+00	1.266e+00	3.376e-01	8.538e-02	2.137e-02	2.00
	$10^4$	7.303e+00	3.523e+00	1.663e+00	7.591e-01	3.132e-01	1.28
	$10^6$	7.323e+00	3.565e+00	1.731e+00	8.575e-01	4.259e-01	1.01

Table 3: The performance for Example 6.2 with  $\iota = 10^{-6}$  and  $\eta = 10$ .

$h$	1/4	1/8	1/16	1/32	1/64
$\ w_0 - w_h\ _{\iota,h}$	2.941e+00	8.607e-01	2.098e-01	4.997e-02	1.256e-02
rate	-	1.77	2.04	2.07	1.99
$ w_0 - w_h _1$	1.806e-01	2.513e-02	2.958e-03	3.457e-04	4.206e-05
rate	-	2.84	3.09	3.10	3.04
$ w_0 - w_h _{2,h}$	2.840e+00	8.113e-01	1.888e-01	4.286e-02	1.027e-02
rate	-	1.81	2.10	2.14	2.06
$ w_0 - w_h _{3,h}$	5.073e+01	2.837e+01	1.381e+01	6.609e+00	3.243e+00
rate	-	0.84	1.04	1.06	1.03
$\ w_0 - w_h\ _0$	2.723e-02	3.063e-03	2.545e-04	1.760e-05	1.138e-06
rate	-	3.15	3.59	3.85	3.95

As shown in Tab. 1, we observe that the convergence rate of  $\|w - w_h\|_{\iota,h}$  for  $\iota = 1, 10^{-2}, 10^{-4}$  is close to  $O(h)$ , while one for  $\iota = 0, 10^{-6}$  would reach to  $O(h^2)$  nearly. Since  $O_{sc_h}(f) = O(h^2)$ , it can be shown that the numerical results are in coincidence with the theoretical predication in (5.55). Moreover, when  $\iota = 10^{-8}$ , it can be seen from Tab. 2 that the numerical method (4.12) behaves well for a large range of  $\eta$ .

**Example 6.2.** The exact solution of the reduced problem (3.3) is set to be a function in the form

$$w_0 = \sin^2(\pi x) \sin^2(\pi y).$$

Set  $f$  computed from (3.3) be the right side term of (3.1). Then, the exact solution of (3.1) possesses strong boundary layer. Besides, It's easy to check that  $f$  is independent of  $\iota$  and the exact solution of (3.1) is unknown. We use this example to demonstrate the robustness of the proposed method (4.12) with respect to size parameter  $\iota$ . We display the error  $w_0 - w_h$  in Tab. 3 and Tab. 4 when  $\iota = 10^{-6}, 10^{-8}$  and  $\eta = 10$ .

Table 4: The performance for Example 6.2 with  $\iota = 10^{-8}$  and  $\eta = 10$ .

$h$	1/4	1/8	1/16	1/32	1/64
$\ w_0 - w_h\ _{\iota,h}$	2.940e+00	8.600e-01	2.092e-01	4.943e-02	1.207e-02
rate	-	1.77	2.04	2.08	2.03
$ w_0 - w_h _1$	1.806e-01	2.513e-02	2.958e-03	3.457e-04	4.206e-05
rate	-	2.84	3.09	3.10	3.04
$ w_0 - w_h _{2,h}$	2.840e+00	8.113e-01	1.888e-01	4.286e-02	1.027e-02
rate	-	1.81	2.10	2.14	2.06
$ w_0 - w_h _{3,h}$	5.073e+01	2.837e+01	1.381e+01	6.609e+00	3.243e+00
rate	-	0.84	1.04	1.06	1.03
$\ w_0 - w_h\ _0$	2.723e-02	3.063e-03	2.545e-04	1.760e-05	1.136e-06
rate	-	3.15	3.59	3.85	3.95

By (5.57) and  $Osc_h(f) = O(h^2)$ , it suffices to observe the behavior of error  $w_0 - w_h$  to verify (5.56). We can observe from Tab. 3 and Tab. 4 that when  $\iota = 10^{-6}$ ,  $\|w_0 - w_h\|_{\iota,h} = O(h^{1.99})$  and when  $\iota = 10^{-8}$ ,  $\|w_0 - w_h\|_{\iota,h} = O(h^{2.03})$ . It can be seen that the numerical results are in agreement with Theorem 5.9, which implies that the proposed method (4.12) is robust with respect to the size parameter  $\iota$ . Besides, it can be seen that  $|w_0 - w_h|_{3,h} = O(h^{1.03})$ ,  $|w_0 - w_h|_{2,h} = O(h^{1.99})$ ,  $|w_0 - w_h|_1 = O(h^{3.04})$ ,  $\|w_0 - w_h\|_0 = O(h^{3.95})$ .

## 7. Declarations

1. **Conflicts of interest/Competing interests.** The authors have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this manuscript.
2. **Availability of data and material.** The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.
3. **Code availability.** The codes during the current study are available from the corresponding author on reasonable request.
4. **Authors' contributions.** All authors contributed equally to this manuscript.

## Reference

- [1] E. C. Aifantis. On the microstructural origin of certain inelastic models. *J. Eng. Mater. Technol.*, 106(4):326–330, 1984.
- [2] S. B. Altan and E. C. Aifantis. On the structure of the mode III crack-tip in gradient elasticity. *Scr. Metall. Mater.*, 26(2):319–324, 1992.
- [3] Q. An, X. Huang, and C. Zhang. A decoupled finite element method for the triharmonic equation. *Appl. Math. Lett.*, 147:108843, 8, 2024.
- [4] P. F. Antonietti, G. Manzini, and M. Verani. The conforming virtual element method for polyharmonic problems. *Comput. Math. Appl.*, 79(7):2021–2034, 2020.
- [5] D. N. Arnold. An interior penalty finite element method with discontinuous elements. *SIAM. J. Numer. Anal.*, 19(4):742–760, 1982.
- [6] A. Ashoori Movassagh and M. Mahmoodi. A micro-scale modeling of Kirchhoff plate based on modified strain-gradient elasticity theory. *Eur. J. Mech. A-Solids*, 40:50–59, 2013.
- [7] B. Babu and B. Patel. A new computationally efficient finite element formulation for nanoplates using second-order strain gradient Kirchhoff's plate theory. *Composites Part B: Engineering*, 168:302–311, 2019.
- [8] J. H. Bramble and M. Zlámal. Triangular elements in the finite element method. *Math. Comp.*, 24:809–820, 1970.
- [9] S. C. Brenner and M. Neilan. A  $C^0$  interior penalty method for a fourth order elliptic singular perturbation problem. *SIAM J. Numer. Anal.*, 49(2):869–892, 2011.
- [10] S. C. Brenner and L.-Y. Sung.  $C^0$  interior penalty methods for fourth order elliptic boundary value problems on polygonal domains. *J. Sci. Comput.*, 22/23:83–118, 2005.

- [11] C. Chen, X. Huang, and H. Wei.  $H^m$ -conforming virtual elements in arbitrary dimension. *SIAM J. Numer. Anal.*, 60(6):3099–3123, 2022.
- [12] H. Chen, J. Li, and W. Qiu. A  $C^0$  interior penalty method for  $m$ th-Laplace equation. *ESAIM Math. Model. Numer. Anal.*, 56(6):2081–2103, 2022.
- [13] L. Chen and X. Huang. Nonconforming virtual element method for  $2m$ th order partial differential equations in  $\mathbb{R}^n$ . *Math. Comp.*, 89(324):1711–1744, 2020.
- [14] L. Chen and X. Huang. Geometric decompositions of the simplicial lattice and smooth finite elements in arbitrary dimension, 2021, arXiv:2111.10712.
- [15] M. Chen, J. Huang, and X. Huang. A robust lower-order mixed finite element method for a strain gradient elastic model. *SIAM J. Numer. Anal.*, 61(5):2237–2260, 2023.
- [16] E. Cosserat and F. Cosserat. *Theorie des corp deformables*. Herman, Paris, 1909.
- [17] F. Dassi, D. Mora, C. Reales, and I. Velásquez. Mixed variational formulations of virtual elements for the polyharmonic operator  $(-\Delta)^n$ . *Comput. Math. Appl.*, 158:150–166, 2024.
- [18] G. Fu, S. Zhou, and L. Qi. On the strain gradient elasticity theory for isotropic materials. *Internat. J. Engrg. Sci.*, 154:103348, 24, 2020.
- [19] D. Gallistl. Stable splitting of polyharmonic operators by generalized Stokes systems. *Math. Comp.*, 86(308):2555–2577, 2017.
- [20] P. Grisvard. *Elliptic Problems in Nonsmooth Domains*. Pitman, Boston, MA, 1985.
- [21] T. Gudi. A new error analysis for discontinuous finite element methods for linear elliptic problems. *Math. Comp.*, 79(272):2169–2189, 2010.
- [22] T. Gudi. Some nonstandard error analysis of discontinuous Galerkin methods for elliptic problems. *Calcolo*, 47(4):239–261, 2010.
- [23] T. Gudi and M. Neilan. An interior penalty method for a sixth-order elliptic equation. *IMA J. Numer. Anal.*, 31(4):1734–1753, 2011.
- [24] J. Guzmán, D. Leykekhman, and M. Neilan. A family of non-conforming elements and the analysis of Nitsche’s method for a singularly perturbed fourth order problem. *Calcolo*, 49(2):95–125, 2012.
- [25] O. y. Hjelle and M. Dæ hlen. *Triangulations and applications*. Mathematics and Visualization. Springer-Verlag, Berlin, 2006.
- [26] J. Hu, T. Lin, and Q. Wu. A construction of  $C^r$  conforming finite element spaces in any dimension. *Found. Comput. Math.*, 24:1941–1977, 2024.
- [27] J. Hu and S. Zhang. A canonical construction of  $H^m$ -nonconforming triangular finite elements. *Ann. Appl. Math.*, 33(3):266–288, 2017.
- [28] J. Hu and S. Zhang. A cubic  $H^3$ -nonconforming finite element. *Commun. Appl. Math. Comput.*, 1(1):81–100, 2019.
- [29] X. Huang. Nonconforming virtual element method for  $2m$ th order partial differential equations in  $\mathbb{R}^n$  with  $m > n$ . *Calcolo*, 57(4):42, 2020.
- [30] X. Huang, Y. Shi, and W. Wang. A Morley-Wang-Xu element method for a fourth order elliptic singular perturbation problem. *J. Sci. Comput.*, 87(3):84, 2021.

- [31] X. H. Huang, J. J. Lai, and W. Q. Wang. A modified Argyris element method for Kirchhoff plates bending problems. *J. Shanghai Jiaotong Univ. (Chinese Ed.)*, 47(2):203–209, 2013.
- [32] M. Ishaquddin and S. Gopalakrishnan. A novel weak form quadrature element for gradient elastic beam theories. *Appl. Math. Model.*, 77:1–16, 2020.
- [33] M. Ishaquddin and S. Gopalakrishnan. Differential quadrature-based solution for non-classical Euler-Bernoulli beam theory. *Eur. J. Mech. A Solids*, 86:104135, 26, 2021.
- [34] W. T. Koiter. Couple-stresses in the theory of elasticity, I and II. *Nederl. Akad. Wetensch. Proc. Ser. B*, 67:17–44, 1964.
- [35] A. Li, X. Ji, S. Zhou, L. Wang, J. Chen, and P. Liu. Nonlinear axisymmetric bending analysis of strain gradient thin circular plate. *Appl. Math. Model.*, 89:363–380, 2021.
- [36] J. Li and S. Wu. A construction of canonical nonconforming finite element spaces for elliptic equations of any order in any dimension, 2024, arXiv:2409.06134.
- [37] R. D. Mindlin. Micro-structure in linear elasticity. *Arch. Ration. Mech. Anal.*, 16(1):51–78, 1964.
- [38] R. D. Mindlin and H. F. Tiersten. Effects of couple-stresses in linear elasticity. *Arch. Ration. Mech. Anal.*, 11(1):415–448, 1962.
- [39] S. M. Mousavi, J. Niiranen, and A. H. Niemi. Differential cubature method for gradient-elastic Kirchhoff plates. *J. Structural Mech.*, 48(3):164–180, 2015.
- [40] J. Niiranen, J. Kiendl, A. H. Niemi, and A. Reali. Isogeometric analysis for sixth-order boundary value problems of gradient-elastic Kirchhoff plates. *Comput. Methods Appl. Mech. Engrg.*, 316:328–348, 2017. Special Issue on Isogeometric Analysis: Progress and Challenges.
- [41] J. Niiranen and A. H. Niemi. Variational formulations and general boundary conditions for sixth-order boundary value problems of gradient-elastic Kirchhoff plates. *Eur. J. Mech. A-Solids*, 61:164–179, 2017.
- [42] T. Nilssen, X.-C. Tai, and R. Winther. A robust nonconforming  $H^2$ -element. *Math. Comput.*, 70(234):489–505, 2001.
- [43] J. Nitsche. Über ein Variationsprinzip zur Lösung von Dirichlet-Problemen bei Verwendung von Teilräumen, die keinen Randbedingungen unterworfen sind. *Abh. Math. Sem. Univ. Hamburg*, 36:9–15, 1971.
- [44] S. Papargyri-Beskou, A. Giannakopoulos, and D. Beskos. Variational analysis of gradient elastic flexural plates under static loading. *Internat. J. Solids Structures*, 47(20):2755–2766, 2010.
- [45] I. P. Pegios, S. Papargyri-Beskou, and D. E. Besko. Finite element static and stability analysis of gradient elastic beam structures. *Acta Mech.*, 226(3):745–768, 2015.
- [46] Y. Ren and D. Liu. The hybrid high-order method for  $p$ -Laplace problem. *Math. Numer. Sin.*, 46(4):397–408, 2024.
- [47] C. Q. Ru and E. C. Aifantis. A simple approach to solve boundary-value problems in gradient elasticity. *Acta Mech.*, 101(1-4):59–68, 1993.

- [48] M. Schedensack. A new discretization for  $m$ th-Laplace equations with arbitrary polynomial degrees. *SIAM J. Numer. Anal.*, 54(4):2138–2162, 2016.
- [49] Z. Shi and M. Wang. *Finite Element Methods*. Academic Press, Beijing, 2013.
- [50] R. A. Toupin. Elastic materials with couple-stresses. *Arch. Ration. Mech. Anal.*, 11(1):385–414, 1962.
- [51] A. Ženišek. Interpolation polynomials on the triangle. *Numer. Math.*, 15:283–296, 1970.
- [52] B. Wang, S. Huang, J. Zhao, and S. Zhou. Reconsiderations on boundary conditions of Kirchhoff micro-plate model based on a strain gradient elasticity theory. *Appl. Math. Model.*, 40(15):7303–7317, 2016.
- [53] M. Wang and J. Xu. Minimal finite element spaces for  $2m$ -th-order partial differential equations in  $R^n$ . *Math. Comp.*, 82(281):25–43, 2013.
- [54] W. Wang, X. Huang, K. Tang, and R. Zhou. Morley-Wang-Xu element methods with penalty for a fourth order elliptic singular perturbation problem. *Adv. Comput. Math.*, 44(4):1041–1061, 2018.
- [55] S. Wu and J. Xu. Nonconforming finite element spaces for  $2m$ th order partial differential equations on  $\mathbb{R}^n$  simplicial grids when  $m = n + 1$ . *Math. Comp.*, 88(316):531–551, 2019.
- [56] X. Xu, Z. Deng, J. Meng, and K. Zhang. Bending and vibration analysis of generalized gradient elastic plates. *Acta Mech.*, 225(12):3463–3482, 2014.
- [57] C. A. Yan, R. Vescovini, and N. Fantuzzi. A neural network-based approach for bending analysis of strain gradient nanoplates. *Eng. Anal. Bound. Elem.*, 146:517–530, 2023.
- [58] Y. Zhou and K. Huang. On simplified deformation gradient theory of modified gradient elastic Kirchhoff-Love plate. *Eur. J. Mech. A Solids*, 100:105014, 2023.