HOM COMPLEXES OF GRAPHS WHOSE CODOMAINS ARE SQUARE-FREE

TAKAHIRO MATSUSHITA

ABSTRACT. The Hom complex Hom(G, H) of graphs is a simplicial complex associated to a pair of graphs G and H, and its homotopy type is of interest in the graph coloring problem and the homomorphism reconfiguration problem. In this paper, we show that if G is a connected graph and H is a square-free connected graph, then every connected component of Hom(G, H) is homotopy equivalent to a point, a circle, H or a connected double cover over H. We also obtain a certain relation between the fundamental group of Hom(G, H) and realizable walks studied in the homomorphism reconfiguration problem.

1. INTRODUCTION

1.1. Background and the main result. Lovász's proof [18] of the Kneser conjecture is widely known as one of the most significant achievements in the application of algebraic topology to combinatorics. He introduced the neighborhood complex $\mathcal{N}(G)$, which is a simplicial complex associated to a graph G, and showed that there is a profound relation between the chromatic number of G and the homotopy type of $\mathcal{N}(G)$. The neighborhood complex is a central object of topological combinatorics, and has been extensively studied (see [4, 19, 24, 25, 31, 33, 35] for example).

The Hom complex $\operatorname{Hom}(G, H)$ of graphs is a simplicial complex associated to a pair of graphs G and H, which is a generalization of the neighborhood complex $\mathcal{N}(H)$ of H. As is the case of the neighborhood complex, the Hom complex has also been studied in the context of the graph coloring problem. For example, when G belongs to a certain class of graphs, called homotopy test graphs, there is a close relation between the chromatic number of H and the homotopy type of $\operatorname{Hom}(G, H)$. Lovász's bound for the chromatic number [18] implies that K_2 is a homotopy test graph. Babson and Kozlov [1, 2] showed that complete graphs and cycle graphs having at least three vertices are homotopy test graphs. In such a context of the graph coloring problem, the homotopy types of Hom complexes have been studied by many authors (see [10, 20, 23, 27, 28] for example). Recently, Hom complexes are also of interest in the study of the homomorphism reconfiguration (see [7, 11, 34]).

The purpose of this paper is to investigate the homotopy type of Hom(G, H) whose codomain H is square-free. Since our main theorem (Theorem 1.2) is a generalization of the case that H is a cycle graph, we recall the previous work on this case. The homotopy types of Hom complexes between cycles were determined by Čukić and Kozlov [6]. Recently, Fujii et al. [12] studied the homotopy type of Hom(G, H) when H is a cycle graph, and proved the following theorem:

²⁰²⁰ Mathematics Subject Classification. Primary 05C15; Secondary 55U10.

Key words and phrases. Hom complex; square free graph; homomorphism reconfiguration.

Theorem 1.1 (Fujii et al. [12]). Let G be a finite connected graph and n an integer at least 3. Then every component of $\text{Hom}(G, C_n)$ is contractible or homotopy equivalent to a circle.

By the famous fold theorem (see [17] for example), one can show that every connected component of $\text{Hom}(G, C_4)$ is contractible. Hence their contribution is to show the case $n \neq 4$. Our main theorem is a generalization of this case:

Theorem 1.2. Let G be a finite connected graph and H a finite square-free connected graph. Then every component of $\operatorname{Hom}(G, H)$ is homotopy equivalent to one of the following spaces: One point space, a circle, the graph H or a connected double cover of H. Moreover, if G is non-bipartite, then every component of $\operatorname{Hom}(G, H)$ is contractible or homotopy equivalent to a circle.

Remark 1.3. Soichiro Fujii, Kei Kimura and Yuta Nozaki [13] independently proved the above theorem. Since their approach is different from that of this paper, we decided to announce them as separate papers.

Theorem 1.2 follows from Theorem 5.8 and Corollary 5.9 (for our conventions, see Subsection 2.8). Note that the homotopy type of a connected double cover of a connected graph H does not depend on the choice of the connected double cover since the homotopy type of a connected graph is determined by its Euler characteristic. Furthermore, when H is neither contractible nor homotopy equivalent to a circle, we determine the number of components homotopy equivalent to H or its connected double cover (see Corollary 5.9 and Proposition 5.10).

Here we discuss the universality problem, which is a fundamental problem in topological combinatorics described as follows: Given a construction K(G) of a simplicial complex from a graph G, determine the class of homotopy or topological types of spaces that can arise as K(G) for some graph G. For instance, Csorba [5] showed that for every finite free $\mathbb{Z}/2$ -simplicial complex X, there exists a graph G such that $\operatorname{Hom}(K_2, G) \simeq_{\mathbb{Z}/2} X$. Furthermore, Dochtermann [9] proved that if a connected graph T contains at least one edge, then for any finite simplicial complex X, there exists a (reflexive) graph G such that $X \simeq \operatorname{Hom}(T, G)$.

On the other hand, Theorem 1.2 implies that if H is square-free, the class of spaces that can appear as Hom(G, H) is highly restricted. For instance, when H is square-free, there does not exist a graph G such that $\text{Hom}(G, H) \simeq S^n$ for any $n \ge 2$.

1.2. Construction of the universal covering. The proof of Theorem 1.2 is divided into three parts. First, we construct the universal covering of a connected component of $\operatorname{Hom}(G, H)$. Secondly, we show that the universal covering is contractible. Finally, we determine the fundamental group of each component of $\operatorname{Hom}(G, H)$. Such a rough outline of the proof is common to [12].

When we consider the universal covering of a space of mappings between topological spaces X and Y, it can sometimes be described as a space of equivariant maps between their universal coverings. For example, it is usual to describe the universal covering of the space Homeo⁺(S^1) of orientation preserving homeomorphisms of a circle S^1 as follows:

$$\widetilde{\text{Homeo}^+}(S^1) = \{ f \in \text{Homeo}(\mathbb{R}) \mid f(x+1) = f(x) + 1 \text{ for every } x \in \mathbb{R} \}.$$

Our construction of the universal covering of a connected component of Hom(G, H) is similar to the geometric intuition above. However, we do not employ the usual covering map theory of graphs, since the usual covering maps are not so compatible with the topology of Hom complexes. Instead, we use the theory of 2-covering maps, studied in [22, 30, 32].

Here, we briefly describe our construction of the covering of a component of $\operatorname{Hom}(G, H)$. In [22], the author introduced the 2-fundamental group $\pi_1^2(G)$ of graphs and 2-covering maps (see Section 2). Then, as is the case of the covering space theory (see [14, Section 1.3]) in classical topology, there are the 2-coverings $p_G \colon \tilde{G} \to G$ and $p_H \colon \tilde{H} \to H$ which are *universal*, that is, $\pi_1^2(\tilde{G})$ and $\pi_1^2(\tilde{H})$ are trivial. Let $f \colon G \to H$ be a graph homomorphism. Then the group $\Gamma = \pi_1^2(G)$ acts on \tilde{G} and \tilde{H} , and we have a Γ -equivariant lift $\tilde{f} \colon \tilde{G} \to \tilde{H}$ of f. We consider the Γ -equivariant Hom complex $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$, which consists of Γ -equivariant multi-homomorphisms. There is a natural surjective map $\pi \colon \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H}) \to \operatorname{Hom}(G, H)$, which turns out to be a covering map (see Proposition 3.1).

Our main task of the proof is to show that if H is square-free then this map π gives a contractible covering of each component of $\operatorname{Hom}(G, H)$. Let $\operatorname{Hom}(G, H)_f$ denote the component of $\operatorname{Hom}(G, H)$ containing f, and $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$ the component of $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$ containing \tilde{f} . The following theorem is deduced from Theorem 4.1.

Theorem 1.4. If H is square-free, then $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$ is contractible.

1.3. Fundamental groups and realizable walks. The homotopy type of a connected simplicial complex X with a contractible universal covering is determined by its fundamental group $\pi_1(X)$ (see [14, Theorem 1B.8]). Therefore, when H is square-free, Theorem 1.4 implies that the homotopy type of $\text{Hom}(G, H)_f$ is determined by its fundamental group. Thus, our next task is to study the fundamental group of $\text{Hom}(G, H)_f$.

For the moment, we do not assume that H is square-free. Then, the covering $\operatorname{Hom}^{\Gamma}(G, H)_{\tilde{f}}$ is not necessarily simply connected (see Remark 5.5). Nonetheless, we can describe a remarkable connection in the general situation between this covering $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$ and the group $\Pi(f, v)$ of realizable elements in $\pi_1^2(H, f(v))$, which is a key concept in Wrochna's work [34] on the homomorphism reconfiguration problem.

We briefly recall his work. While Wrochna originally considered only the case where H is square-free, it is straightforward to extend his concept to the general setting in terms of 2-fundamental groups. Let v be a non-isolated vertex of G. Consider a \times -homotopy $h: G \times I_n \to H$ from f to f (see Subsection 2.3). Then there is a closed walk

$$(h(v,0), w_1, h(v,1), w_2, \dots, w_n, h(v,n))$$

at f(v) with even length in H. Although this closed walk depends on the choice of w_k , its 2-homotopy class does not. Thus, it defines an element of $\pi_1^2(H, f(v))$. We call an element of $\pi_1^2(H, f(v))$ obtained in this manner *realizable*. The realizable elements form a subgroup of $\pi_1^2(H, f(v))$, which we denote by $\Pi(f, v)$.

We are now ready to state our next result, which reveals a relation among $\pi_1(\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}})$, $\pi_1(\operatorname{Hom}(G, H)_f)$ and $\Pi(f, v)$.

Theorem 1.5. Let G and H be connected graphs. Then there is a following short exact sequence of groups:

$$1 \to \pi_1(\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}, \tilde{f}) \to \pi_1(\operatorname{Hom}(G, H)_f, f) \to \Pi(f, v) \to 1.$$

When H is square-free, Theorem 1.4 implies that $\text{Hom}(G, H)_f$ is the $K(\Pi(f, v), 1)$ -space. Wrochna [34] determined the isomorphism types of $\Pi(f, v)$ in this case (see Theorem 5.7), which completes the proof of Theorem 1.2.

The rest of this paper is organized as follows: In Section 2, we review necessary definitions and facts. In Section 3 we construct the covering $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$ and show that the natural map π : $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H}) \to \operatorname{Hom}(G, H)$ is a covering map (Proposition 3.1). In Section 4, we show Theorem 1.4. In Section 5, we prove Theorem 1.5, and complete the proof of Theorem 1.2. In Appendix, we provide an alternative proof of Wrochna's result (Theorem 5.7) required for the proof of Theorem 1.2.

2. Preliminaries

In this section, we review necessary definitions and facts, following [14, 15, 16, 22].

2.1. **Graphs.** A graph is a pair G = (V(G), E(G)) consisting of a set V(G) together with a symmetric subset E(G) of $V(G) \times V(G)$, i.e., $(x, y) \in E(G)$ implies $(y, x) \in E(G)$. Hence our graphs may have loops but no multiple edges. A vertex v is *looped* if $(v, v) \in E(G)$. If G has no looped vertices, then G is said to be *simple*. For $x, y \in V(G)$, we write $x \sim_G y$ or $x \sim y$ to mean that x and y are adjacent.

For $n \geq 3$, the *n*-cycle graph C_n is defined by $V(C_n) = \mathbb{Z}/n$ and $E(C_n) = \{(x, x \pm 1) \mid x \in \mathbb{Z}/n\}$. A square-free graph is a simple graph which has no subgraph isomorphic to C_4 .

For graphs G and H, a graph homomorphism from G to H is a map $f: V(G) \to V(H)$ such that $(v, w) \in E(G)$ implies $(f(v), f(w)) \in E(H)$.

Let P_n be the path graph with n + 1 vertices. Namely, $V(P_n) = \{0, 1, \ldots, n\}$ and $E(P_n) = \{(x, y) \mid |x - y| = 1\}$. A walk in a graph G with length n is a graph homomorphism $\gamma: P_n \to G$, and set length $(\gamma) = n$. A walk γ is called a path when γ is injective as a set map. We sometimes write $(\gamma(0), \cdots, \gamma(n))$ to indicate the walk γ . For a pair of walks $\gamma: P_n \to G$ and $\delta: P_m \to G$ such that $\gamma(n) = \delta(0)$, the concatenation $\gamma \cdot \delta$ of γ and δ is defined by the walk $(\gamma(0), \cdots, \gamma(n), \delta(1), \cdots, \delta(m))$ with length m + n.

Remark 2.1. The order of concatenation of walks aligns with that in [14, 30, 32] but is the reverse of the order used in [22]. While this does not entail any essential change, it reverses the roles of the starting and terminal points of walks. Furthermore, whereas the action of the 2-fundamental group on the universal 2-covering is from the right in [22], this action is from the left in this paper (see Subsection 2.6).

Let γ be a walk (x_0, \ldots, x_n) , and suppose $x \sim x_0$ and $x_n \sim y$. We denote the walk (x, x_0, \ldots, x_n) by $x - \gamma$, the walk (x_0, \ldots, x_n, y) by $\gamma - y$, and the walk (x_0, \ldots, x_{n-1}) by $\gamma \setminus x_n$.

A based graph is a pair (G, v) consisting of a graph G equipped with a vertex v of G. For based graphs (G, v) and (H, w), a based graph homomorphism from (G, v) to (H, w) is a graph homomorphism from G to H sending v to w. A closed walk of a based graph (G, v)is a walk from v to v. The categorical product $G \times H$ of graphs G and H is defined by

$$V(G \times H) = V(G) \times V(H), \quad E(G \times H) = \{((x, y), (x', y')) \mid (x, x') \in E(G), (y, y') \in E(H)\}.$$

2.2. **Poset topology.** Let P and Q be (possibly infinite) posets. A chain in P is a subset σ of P such that any two elements of σ are comparable. The order complex $\Delta(P)$ of P is the simplicial complex whose simplices are finite chains in P. The classifying space of P is the geometric realization of $\Delta(P)$, and denoted by |P|. We apply topological terms to posets, using classifying spaces. For example, a subposet Q of P is said to be a deformation retract of P if |Q| is a deformation retract of |P|.

A map $f: P \to Q$ is said to be *monotone* or *order preserving* if for any two elements x, y of $P, x \leq y$ implies $f(x) \leq f(y)$. Then a monotone map induces a continuous map $|f|: |P| \to |Q|$ between their classifying spaces.

An ascending (or descending) closure operator of P is a monotone map $c: P \to P$ such that $c(x) \ge x$ ($c(x) \le x$, respectively) for every $x \in P$ and $c^2 = c$.

Theorem 2.2 ([16, Theorem 13.12]). Let $c: P \to P$ be an ascending or descending closure operator. Then c(P) is a deformation retract of P.

Definition 2.3. A monotone map $f: P \to Q$ is a *poset covering map* if for every pair y and y' of elements of Q with $y \leq y'$, both of the following two conditions are satisfied:

(1) For every $x \in P$ with f(x) = y, there is unique $x' \in P$ such that $x \leq x'$ and f(x') = y'.

(2) For every $x' \in P$ with f(x') = y', there is unique $x \in P$ such that $x \leq x'$ and f(x) = y.

Proposition 2.4 (see [3, Proposition 2.9 and Theorem 3.2] for example). A poset covering map $f: P \to Q$ induces a covering map $|f|: |P| \to |Q|$ between their classifying spaces.

2.3. Hom complexes. For a set X, let $\mathcal{P}(X)$ denote the power set of X. For graphs G and H, a multi-homomorphism from G to H is a map $\eta: V(G) \to \mathcal{P}(V(H)) \setminus \{\emptyset\}$ such that $(x,y) \in E(G)$ implies $\eta(x) \times \eta(y) \subset E(H)$, i.e., every element of $\eta(x)$ and every element of $\eta(y)$ are adjacent. Let $\operatorname{Hom}(G, H)$ be the set of multi-homomorphisms from G to H. We define a partial order on $\operatorname{Hom}(G, H)$ as follows: For $\eta, \eta' \in \operatorname{Hom}(G, H), \eta \leq \eta'$ if and only if $\eta(x) \subset \eta'(x)$ for every $x \in V(G)$. A map $f: V(G) \to V(H)$ is a graph homomorphism if and only if the map $x \mapsto \{f(x)\}$ is a multi-homomorphism. From this point of view, we regard a graph homomorphism as an element of $\operatorname{Hom}(G, H)$.

A graph homomorphism $p: H_1 \to H_2$ induces a monotone map $p_*: \text{Hom}(G, H_1) \to \text{Hom}(G, H_2)$ defined by $p_*(\eta)(x) = p(\eta(x))$.

Remark 2.5. If G has no edge, then any map from V(G) to $\mathcal{P}(V(H))$ is a multi-homomorphism. Thus, $\operatorname{Hom}(G, H)$ is empty or contractible, and Theorem 1.2 is trivial. This is an exceptional case, and we assume that G and H have at least one edge in the subsequent sections (see Subsection 2.8).

Now we recall a bit of ×-homotopy theory, studied by Dochtermann [7]. For a nonnegative integer n, let I_n be the graph defined by $V(I_n) = \{0, 1, \dots, n\}, E(I_n) = \{(i, j) \mid i, j \in V(I_n), |i-j| \leq 1\}$. Let $f, g: G \to H$ be graph homomorphisms. A ×-homotopy from f to g is a graph homomorphism $h: G \times I_n \to H$ such that h(x, 0) = f(x) and h(x, n) = g(x) for every $x \in V(G)$. Two graph homomorphisms f and g are said to be ×-homotopic if there is a ×-homotopy from f to g. The proof of the following lemma is straightforward.

Lemma 2.6. For graph homomorphisms $f, g: G \to H$, the following are equivalent:

- (1) The map $h: V(G \times I_1) \to V(H)$ defined by h(x,0) = f(x) and h(x,1) = g(x) is a graph homomorphism from $G \times I_1$ to H.
- (2) The map $f \cup g: V(G) \to \mathcal{P}(V(H)) \setminus \{\emptyset\}, x \mapsto \{f(x), g(x)\}$ is a multi-homomorphism.
- (3) There is a multi-homomorphism $\eta \in \text{Hom}(G, H)$ such that $f \leq \eta$ and $g \leq \eta$.

This implies the following:

Proposition 2.7 ([7, Proposition 4.7]). Let f and g be graph homomorphisms from G to H. Then f and g are \times -homotopic if and only if f and g belong to the same connected component of Hom(G, H).

2.4. 2-fundamental groups. The theory of 2-fundamental groups and 2-covering maps was first studied in [22], and applied to multiplicative graphs in [30, 32]. Another type of combinatorial fundamental groups related to Hom complexes was studied in [8].

Let $v, v' \in V(G)$, and let $\Omega(G, v, v')$ be the set of walks joining v to v'. Let \simeq_2 be the equivalence relation on $\Omega(G, v, v')$ generated by the following two relations (1) and (2): For walks (x_0, \dots, x_m) and (y_0, \dots, y_n) joining v to v',

- (A) n = m + 2 and there is $k \in \{0, 1, ..., m\}$ such that $x_i = y_i$ for every $i \le k$ and $x_i = y_{i+2}$ for every $i \ge k$.
- (B) m = n and $\#\{i \mid x_i \neq y_i\} \le 1$.

Let $\pi_1^2(G, v, v')$ be the quotient set $\Omega(G, v, v')/\simeq_2$. We call the equivalence class containing $\varphi \in \Omega(G, v, v')$ the 2-homotopy class of φ . We write $\pi_1^2(G, v)$ instead of $\pi_1^2(G, v, v)$. The concatenation of walks defines a group operation on $\pi_1^2(G, v)$. We call this group the 2-fundamental group of (G, v). Note that a based graph homomorphism $f \colon (G, v) \to (H, w)$ induces a group homomorphism $f_* \colon \pi_1^2(G, v) \to \pi_1^2(H, f(v))$.

Let \simeq_1 be the equivalence relation on $\Omega(G, v)$ generated by the condition (A) and set $\pi_1^1(G, v) = \Omega(G, v) / \simeq_1$. Similarly, by the concatenation of closed walks, $\pi_1^1(G, v)$ becomes a group. We call this group the 1-fundamental group of (G, v). By the definition there is a natural quotient map $\pi_1^1(G, v) \to \pi_1^2(G, v)$. When G is simple, then the 1-fundamental group is isomorphic to the edge path group (see [29]) of (G, v), and hence is isomorphic to the usual fundamental group. Thus, in this case we write $\pi_1(G, v)$ instead of $\pi_1^1(G, v)$.

If G is square-free, then (B) is obtained from (A). Thus we have the following:

Lemma 2.8. If G is square-free and $v \in V(G)$, then the natural quotient map $\pi_1(G, v) \rightarrow \pi_1^2(G, v)$ is an isomorphism.

By the definition of the 2-fundamental group $\pi_1^2(G, v)$, there is a well-defined group homomorphism

$$\pi_1^2(G, v) \to \mathbb{Z}/2, \quad [\gamma] \mapsto \text{length}(\gamma) \mod 2.$$

The kernel of this group homomorphism is called the *even part of* $\pi_1^2(G, v)$ and is denoted by $\pi_1^2(G, v)_{ev}$. If the component of G containing v is bipartite, then $\pi_1^2(G, v)_{ev}$ coincides with $\pi_1^2(G, v)$. If the component of G containing v is non-bipartite, then $\pi_1^2(G, v)_{ev}$ is a subgroup of $\pi_1^2(G, v)$ whose index is 2.

Finally, we recall the following close relationship between the 2-fundamental groups and the fundamental groups of neighborhood complexes. **Theorem 2.9** (Theorem 1.1 of [22]). Let (G, v) be a based graph, and assume that v is not isolated. Then there is a natural isomorphism

$$\pi_1^2(G, v)_{ev} \xrightarrow{\cong} \pi_1(\mathcal{N}(G), v).$$

2.5. 2-covering maps.

Definition 2.10. Let G be a graph and x a vertex of G. Define $N_G(x)$ and $N_G^2(x)$ by

$$N_G(x) = \{y \in V(G) \mid (x, y) \in E(G)\}, \quad N_G^2(x) = \bigcup_{y \in N_G(x)} N_G(y).$$

We write N(x) (or $N^2(x)$) instead of $N_G(x)$ (or $N_G^2(x)$, respectively).

Definition 2.11. A covering map or a 1-covering map is a graph homomorphism $p: G \to H$ such that $p|_{N(v)}: N(v) \to N(p(v))$ is bijective for every $v \in V(G)$. A 2-covering map is a graph homomorphism $p: G \to H$ such that for every $v \in V(G)$, the maps $p|_{N(v)}: N(v) \to$ N(p(v)) and $p|_{N^2(v)}: N^2(v) \to N^2(p(v))$ are bijective.

Theorem 2.12 ([22, Proposition 6.9]). Let (G, v) be a based graph. Then there is a connected based 2-covering map $p: (\tilde{G}, \tilde{v}) \to (G, v)$ such that $\pi_1^2(\tilde{G}, \tilde{v})$ is trivial. Moreover, such a based 2-covering over (G, v) is unique up to isomorphisms.

We call this graph \tilde{G} the universal 2-covering of G.

Remark 2.13. The universal 2-covering \tilde{G} of G is always simple and bipartite. Indeed, $\pi_1^2(\tilde{G}, \tilde{v}) \cong 1$ implies that there is no odd element of $\pi_1^2(\tilde{G}, \tilde{v})$, and this means that \tilde{G} has neither loops nor odd cycles.

We will frequently use the following analogy to the lifting criterion of covering spaces [14, Proposition 1.33].

Lemma 2.14 (lifting criterion [22, Lemma 6.8]). Let $p: (G, v) \to (H, w)$ be a based 2covering map, and $f: (T, x) \to (H, w)$ be a based graph homomorphism. Assume that T is connected. Then the following are equivalent:

- (1) $f_*(\pi_1^2(T,x)) \subset p_*(\pi_1^2(G,v)).$
- (2) There is a based graph homomorphism $\tilde{f}: (T, x) \to (G, v)$ such that $p \circ \tilde{f} = f$.
- (3) There is a unique based graph homomorphism $\tilde{f}: (T, x) \to (G, v)$ such that $p \circ \tilde{f} = f$.

2.6. 2-covering actions.

Definition 2.15. Let Γ be a group and G a graph. A Γ -action on G is a Γ -action on V(G) such that for every $\gamma \in \Gamma$, the map $x \mapsto \gamma x$ is a graph homomorphism from G to G.

For a Γ -action on a graph G, define the quotient graph G/Γ as follows: The vertex set of G/Γ is the orbit set $V(G)/\Gamma$. The edge set of G/Γ is defined by

$$E(G/\Gamma) = \{ (\alpha, \beta) \in V(G/\Gamma) \times V(G/\Gamma) \mid (\alpha \times \beta) \cap E(G) \neq \emptyset \}.$$

In other words, $(\alpha, \beta) \in E(G/\Gamma)$ if and only if there are $x \in \alpha$ and $y \in \beta$ satisfying $(x, y) \in E(G)$.

It is easy to see that the projection $p_G: G \to G/\Gamma$ is a graph homomorphism.

Definition 2.16 (the sentence before Proposition 6.10 of [22]). A Γ -action on a graph G is a 2-covering action if $N^2(x) \cap N^2(\gamma x) = \emptyset$ for every $x \in V(G)$ and $\gamma \in \Gamma \setminus \{1_{\Gamma}\}$.

Then the following equivalence holds:

Proposition 2.17 (see [22, Proposition 6.10]). Let G be a Γ -graph, and assume that the Γ -action is free as a set action. Then the following are equivalent:

- (1) The projection $G \to G/\Gamma$ is a 2-covering map.
- (2) The Γ -action on G is a 2-covering action.

Let (G, v) be a connected based graph and (\tilde{G}, \tilde{v}) the universal 2-covering over (G, v). Then $\pi_1^2(G, v)$ acts on \tilde{G} as follows: Let $\tilde{x} \in V(\tilde{G})$, and let φ be a walk joining \tilde{v} to \tilde{x} . Let $\alpha \in \pi_1^2(G, v)$ and γ a representative of α . Then there is a walk ψ on \tilde{G} such that $p_G \circ \psi = \gamma \cdot (p_G \circ \varphi)$ starting from \tilde{v} . Define $\alpha \tilde{x}$ to be the terminal point of ψ . This is the description of the 2-covering action of $\pi_1^2(G, v)$ on \tilde{G} (see the proof of [22, Theorem 6.11]). In [22], the author considered a right action, which is due to the fact that the order of concatenation of walks is the reverse of that in this paper (see Remark 2.1).

This description of the $\pi_1^2(G, v)$ -action on \tilde{G} implies the following:

Lemma 2.18. Let $p_G: (\tilde{G}, \tilde{v}) \to (G, v)$ be the universal 2-covering of (G, v), and let $g \in \pi_1^2(G, v)$. Let φ be a walk of \tilde{G} joining \tilde{v} to $g\tilde{v}$. Then the 2-homotopy class of the closed walk $p_G \circ \varphi$ of (G, v) coincides with g.

Let $p_G: (\tilde{G}, \tilde{v}) \to (G, v)$ and $p_H: (\tilde{H}, \tilde{w}) \to (H, w)$ be the universal 2-coverings. Let $f: (G, v) \to (H, w)$ be a based graph homomorphism. Then Lemma 2.14 implies that there is a based graph homomorphism $\tilde{f}: (\tilde{G}, \tilde{v}) \to (\tilde{H}, \tilde{w})$ satisfying $p_H \circ \tilde{f} = f \circ p_G$. Set $\Gamma = \pi_1^2(G, v)$. Then Γ acts on \tilde{G} as above. Moreover, Γ acts on \tilde{H} by $g\tilde{w} = f_*(g)\tilde{y}$ for $g \in \Gamma$ and $\tilde{y} \in V(\tilde{H})$. By the above description of the actions on the universal 2-coverings, the graph homomorphism $\tilde{f}: \tilde{G} \to \tilde{H}$ is Γ -equivariant.

2.7. A bit of algebraic topology.

Lemma 2.19 ([26, Example 2 and Theorem A in Appendix]). Let X be a simplicial complex and $(X_n)_{n \in \mathbb{N}}$ an increasing sequence of subcomplexes of X. Suppose that X_i is a deformation retract of X_{i+1} for each i and $X = \bigcup_{n \in \mathbb{N}} X_n$. Then the inclusion $X_0 \hookrightarrow X$ is a homotopy equivalence.

Theorem 2.20 (see [14, Proposition 1.39] and its proof). Let (X, x_0) be a based simplicial complex and $p: (Y, y_0) \to (X, x_0)$ a connected covering space. Let Γ_0 be a group, and suppose that Γ_0 acts on Y freely and $p(\gamma y) = p(y)$ for every $\gamma \in \Gamma_0$ and for every $x \in X$. If

$$p^{-1}(x_0) = \{\gamma y_0 \mid \gamma \in \Gamma_0\},\$$

then there is a following short exact sequence of groups:

$$1 \to \pi_1(Y, y_0) \to \pi_1(X, x_0) \xrightarrow{\Psi} \Gamma_0 \to 1.$$

Moreover, the group homomorphism $\Psi: \pi_1(X, x_0) \to \Gamma_0$ is described as follows: Let $[\gamma] \in \pi_1(X, x_0)$. Then there is a unique $\tilde{\gamma}: [0, 1] \to Y$ such that $\tilde{\gamma}(0) = y_0$ and $p \circ \tilde{\gamma} = \gamma$. Then define $\Psi([\gamma])$ by $\gamma(1) = \Psi([\gamma]) \cdot y_0$.

2.8. Notation and conventions. For the rest of this paper, G and H are assumed to be finite non-empty connected graphs having at least one edge (see Remark 2.5). The graph His not assumed to be square-free or even simple unless it is specified. Our results are valid for graphs having loops if it is not specified. Let $f: G \to H$ denote a graph homomorphism. Let $v \in V(G)$ and set w = f(v). Let Γ denote the 2-fundamental group $\pi_1^2(G, v)$ of (G, v). Let $p_G: (\tilde{G}, \tilde{v}) \to (G, v), p_H: (\tilde{H}, \tilde{w}) \to (H, w)$ be the universal 2-coverings of (G, v) and (H, w), respectively. Let $\tilde{f}: \tilde{G} \to \tilde{H}$ be a lift of f, and suppose $\tilde{f}(\tilde{v}) = \tilde{w}$.

3. Construction of the covering

As was described in Subsection 2.6, $\Gamma = \pi_1^2(G, v)$ acts on \tilde{G} and \tilde{H} , and the lift $\tilde{f} : \tilde{G} \to \tilde{H}$ of $f : G \to H$ is Γ -equivariant. A multi-homomorphism $\eta \in \operatorname{Hom}(\tilde{G}, \tilde{H})$ is said to be Γ equivariant if $f_*(\gamma)\eta(\tilde{x}) = \eta(\gamma \tilde{x})$ for every $\gamma \in \Gamma$ and $\tilde{x} \in V(\tilde{G})$. Then we have $\tilde{f} \in$ $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$, and let $\operatorname{Hom}(\tilde{G}, \tilde{H})_{\tilde{f}}$ denote the component of $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$ containing \tilde{f} .

Define the map

 $\pi \colon \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H}) \to \operatorname{Hom}(G, H)$

by $\pi(\eta)(p_G(\tilde{x})) = p_H(\eta(\tilde{x}))$. It is straightforward to see that π is a well-defined monotone map. The goal of this section is to prove that the map π is a poset covering map.

Proposition 3.1. The map $\pi \colon \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H}) \to \operatorname{Hom}(G, H)$ is a poset covering map.

Proof. Let $\eta, \eta' \in \text{Hom}(G, H)$ and suppose $\eta \leq \eta'$. By the definition of poset covering maps (Definition 2.3), it suffices to show the following:

- (1) For every $\tilde{\eta} \in \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$ with $\pi(\tilde{\eta}) = \eta$, there is unique $\tilde{\eta}' \in \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$ such that $\tilde{\eta} \leq \tilde{\eta}'$ and $\pi(\tilde{\eta}') = \eta'$.
- (2) For every $\tilde{\eta}' \in \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$ with $\pi(\tilde{\eta}') = \eta'$, there is unique $\tilde{\eta} \in \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$ such that $\tilde{\eta} \leq \tilde{\eta}'$ and $\pi(\tilde{\eta}) = \eta$.

We first show (1). Choose $v_x \in \tilde{\eta}(x) \subset V(\tilde{H})$ for each $x \in V(\tilde{G})$ so that $v_{\gamma x} = \gamma v_x$. Then set $\tilde{\eta}'(x) = (p_H|_{N_{\tilde{H}}^2(v_x)})^{-1}(\eta'(x))$, where $p_H|_{N_{\tilde{H}}^2(v_x)} \colon N_{\tilde{H}}^2(v_x) \to N_{H}^2(p_H(v_x))$. Since \tilde{G} has no isolated vertices (see Subsection 2.8), we have $\tilde{\eta}(x) \subset N_{\tilde{H}}^2(v_x)$ and $p_H(\tilde{\eta}(x)) = \eta(x) \subset \eta'(x)$. Thus we have $\tilde{\eta}(x) \subset \tilde{\eta}'(x)$. Now we check the Γ -equivariance of $\tilde{\eta}'$. Since $\gamma v_x = v_{\gamma x}$, we have $\gamma \tilde{\eta}'(x) \subset \gamma N_{\tilde{H}}^2(v_x) \subset N_{\tilde{H}}^2(v_{\gamma x})$. Since $p_H(\gamma \tilde{\eta}'(x)) = \eta'(p_G(x)) = \eta'(p_G(\gamma x)) = p_H(\tilde{\eta}'(\gamma x))$, we have $\gamma \tilde{\eta}'(x) = \tilde{\eta}'(\gamma x)$. Hence $\tilde{\eta}'$ is Γ -equivariant.

We show that $\tilde{\eta}'$ is a multi-homomorphism. Let $(x, y) \in E(\tilde{G})$, and let $a \in \tilde{\eta}'(x)$ and $b \in \tilde{\eta}'(y)$. It suffices to show that $a \sim_{\tilde{H}} b$. Since $p_H(a) \in \eta'(x) \subset N_H(p_H(v_y))$, there is $a' \in N_{\tilde{H}}(v_y)$ such that $p_H(a) = p_H(a')$. Since $v_x \sim_{\tilde{H}} v_y \sim_{\tilde{H}} a'$, we have $a' \in N_{\tilde{H}}^2(v_x)$. Since $a, a' \in N_{\tilde{H}}^2(v_x)$, we have a = a' from the injectivity of $p_H|_{N_{\tilde{H}}^2(v_x)}$. Therefore we have $a = a' \sim_{\tilde{H}} v_y$. Similarly, we have $v_x \sim_{\tilde{H}} b$. Hence we have

$$a \sim_{\tilde{H}} v_y \sim_{\tilde{H}} v_x \sim_{\tilde{H}} b.$$

Note that $p_H(a) \in \eta'(x)$, $p_H(b) \in \eta'(y)$ and η' is a multi-homomorphism. Hence we have $p_H(a) \sim_H p_H(b)$. Then there is $b' \in N_{\tilde{H}}(a)$ such that $p_H(b') = p_H(b)$. Since $b, b' \in N_{\tilde{H}}^2(v_y)$, we have b = b'. Therefore we conclude that $a \sim b' = b$ and $\tilde{\eta}'$ is a multi-homomorphism.

Finally, we mention the uniqueness of $\tilde{\eta}'$. Suppose that $\tilde{\eta}'' \in \text{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$ such that $\eta \leq \tilde{\eta}''$ and $\pi(\tilde{\eta}'') = \eta'$. Since $\tilde{\eta}''$ is a multi-homomorphism, we have $\tilde{\eta}''(x) \subset N^2_{\tilde{H}}(v_x)$ for every $x \in V(\tilde{G})$. Since $p_H|_{N^2_{\tilde{H}}(v_x)}$ is bijective and $p_H(\tilde{\eta}''(x)) = \eta'(p_G(x)) = p_H(\tilde{\eta}'(x))$, we have $\tilde{\eta}'(x) = \tilde{\eta}''(x)$. This completes the proof of (1).

The proof of (2) is easier. Indeed, for every $x \in V(\tilde{G})$, let $v_x \in \tilde{\eta}'(x)$ so that $\gamma v_x = v_{\gamma x}$. Define $\tilde{\eta}(x)$ by $\tilde{\eta}(x) = (p_H|_{N^2(v_x)})^{-1}(\eta(x))$. Then $\tilde{\eta}(x)$ is clearly a multi-homomorphism since $\tilde{\eta}(x) \subset \tilde{\eta}'(x)$ and $\tilde{\eta}'$ is a multi-homomorphism. The proofs for verifying the other conditions are the same as that of (1). This completes the proof. \Box

Remark 3.2. In the above proof, the hypothesis used in most parts is that $p_H: H \to H$ is a 2-covering map. Hence, other variations of this proposition, such as the following, can also be shown by the same argument: If $p: H_1 \to H_2$ be a 2-covering map, then the map $p_*: \operatorname{Hom}(G, H_1) \to \operatorname{Hom}(G, H_2)$ is a poset covering map. In fact, this version appeared in an earlier draft of [22] (see [21, Lemma 7.1]).

4. Contractibility

It is straightforward to see that if H is square-free then H is a tree (see [30] for example). Thus, to prove Theorem 1.4, it suffices to show the following theorem.

Theorem 4.1. If \tilde{H} is a tree, then $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$ is contractible.

The goal of this section is to prove Theorem 4.1. Throughout this section, we assume that \tilde{H} is a tree.

4.1. Outline of the proof. Here we outline the proof of Theorem 4.1. We construct the filtration of $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$ in three stages as follows:

• For each $n \ge 0$, we construct a subposet X_n such that

$$\{\tilde{f}\} = X_0 \subset X_1 \subset X_2 \cdots, \quad \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}} = \bigcup_{n \ge 0} X_n.$$

• Fix $n \ge 0$. For each $m \ge -1$, we construct a subposet $X_{n,m}$ such that

$$X_{n-1} = X_{n,-1}, \quad X_{n,-1} \subset X_{n,0} \subset X_{n,1} \subset \cdots,$$

and $X_{n,m} = X_n$ for sufficiently large m.

• Fix $n \ge 0$ and $m \ge -1$. For each $i \ge 0$, we construct a subposet $X_{n,m,i}$ such that

 $X_{n,m,0} = X_{n,m}, \quad X_{n,m,0} \subset X_{n,m,1} \subset X_{n,m,2} \subset \cdots,$

and $X_{n,m,i} = X_{n,m}$ for sufficiently large *i*.

Finally, we show that $X_{n,m,i-1}$ is a deformation retract of $X_{n,m,i}$ (Proposition 4.10), which concludes the proof of Theorem 4.1 by Lemma 2.19.

4.2. Subposet X_n . In this subsection we define the subposet X_n of $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$ for each $n \geq 0$. For $\eta \in \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$ and for $\tilde{x} \in V(\tilde{G})$, set

$$u_{\eta}(\tilde{x}) = \max\{d_{\tilde{H}}(f(\tilde{x}), \tilde{y}) \mid \tilde{y} \in \eta(\tilde{x})\}.$$

Here $d_{\tilde{H}}$ denotes the path metric of \tilde{H} .

Lemma 4.2. For every $\eta \in \text{Hom}(\tilde{G}, \tilde{H})$ and for every $\tilde{x} \in V(\tilde{G})$, the integer $u_n(\tilde{x})$ is even.

Proof. Since \tilde{H} is bipartite (see Remark 2.13), there is a graph homomorphism $\varepsilon \colon \tilde{H} \to K_2$. Then ε induces a map $\varepsilon_* \colon \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}} \to \operatorname{Hom}(\tilde{G}, K_2) \cong S^0$. Since $u_{\eta}(\tilde{x})$ is odd, $\eta(\tilde{x})$ has an element \tilde{y} such that $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y})$ is odd. Thus, $\varepsilon_*(\eta)(\tilde{x})$ contains an element different from $\varepsilon_*\tilde{f}(\tilde{x}) = \varepsilon(\tilde{f}(\tilde{x}))$. Hence we have $\varepsilon_*(\eta) \neq \tilde{f}$. This is a contradiction since $\operatorname{Hom}(\tilde{G}, K_2) \cong S^0$ and $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$ is connected. \Box

Since η is Γ -equivariant, for every $\gamma \in \Gamma$ and $\tilde{x} \in V(\tilde{G})$, we have $u_{\eta}(\gamma \tilde{x}) = u_{\eta}(\tilde{x})$. Hence u_{η} induces a function \bar{u}_{η} on V(G). Since G is finite, \bar{u}_{η} is bounded, which implies that u_{η} is also bounded. Define the subposet X_n of $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$ by

$$X_n = \{ \eta \in \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}} \mid u_{\eta}(\tilde{x}) \le 2n \text{ for every } \tilde{x} \in V(\tilde{G}) \}$$

Since u_{η} is bounded, $\eta \in X_n$ holds for some n. Thus we have

$$\{\tilde{f}\} = X_0 \subset X_1 \subset X_2 \subset \cdots, \quad \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}} = \bigcup_{n \ge 0} X_n.$$

4.3. Directed subgraphs $\tilde{G}(\eta)$ and $G(\eta)$. Our next task is to construct the filtration $X_{n,m}$ of X_n . To define the filtration, we need to introduce a directed subgraph $\tilde{G}(\eta)$ of \tilde{G} and a directed subgraph $G(\eta)$ of G for each $\eta \in X_n$.

First we define $G(\eta)$ and $G(\eta)$ as undirected graphs. Let $\eta \in X_n$. Define the subgraph $\tilde{G}(\eta)$ of \tilde{G} to be the induced subgraph of \tilde{G} whose vertex set is $u_{\eta}^{-1}(2n)$. Similarly, define the subgraph $G(\eta)$ of G to be the induced subgraph of G whose vertex set is $\bar{u}_{\eta}^{-1}(2n)$. Next, we consider the direction of each edge of $\tilde{G}(\eta)$ and $G(\eta)$. Here we recall that \tilde{H} is a tree, and that for any two vertices of \tilde{H} there is a unique path joining them.

Proposition 4.3. Let $(\tilde{x}, \tilde{x}') \in G(\eta)$. Then one of the following two conditions holds, and the other does not.

- (1) The set $\eta(\tilde{x})$ consists of one vertex \tilde{y} of \tilde{H} and for every element $\tilde{y}' \in \eta(\tilde{x}')$ with $d_{\tilde{H}}(\tilde{f}(\tilde{x}'), \tilde{y}') = 2n$, the path joining $\tilde{f}(\tilde{x}')$ to \tilde{y}' cointains \tilde{y} .
- (2) The set $\eta(\tilde{x}')$ consists of one vertex \tilde{y}' of \tilde{H} and for every element $\tilde{y} \in \eta(\tilde{x})$ with $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y}) = 2n$, the path joining $\tilde{f}(\tilde{x})$ to \tilde{y} cointains \tilde{y}' .

Proof. Since $\tilde{x}' \in V(\tilde{G}(\eta))$, we have $u_{\eta}(\tilde{x}') = 2n$. Hence there is $\tilde{y}' \in \eta(\tilde{x}')$ such that $d_{\tilde{H}}(\tilde{f}(\tilde{x}'), \tilde{y}') = 2n$. Similarly, there is $\tilde{y} \in \eta(\tilde{x})$ such that $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y}) = 2n$. Let P be the path in \tilde{H} joining $\tilde{f}(\tilde{x})$ to \tilde{y} .

Suppose that \tilde{y}' is not contained in P. Then the path P can be extended to the path $P \cdot \tilde{y}'$. Since \tilde{H} is a tree, we have $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y}') = 2n + 1$. Let $\tilde{y}_0 \in \eta(\tilde{x})$. Since $\eta \in X_n$,

 $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y}_0) \leq 2n \text{ and } \tilde{y}_0 \text{ is adjacent to } \tilde{y}'$. Since \tilde{H} is a tree, this implies that \tilde{y}_0 coincides with \tilde{y} . If $\tilde{f}(\tilde{x}')$ is not contained in P, then $\tilde{f}(\tilde{x}') \cdot P \cdot \tilde{y}'$ is a path joining $\tilde{f}(\tilde{y})$ to \tilde{y}' with length 2n + 2. This means that $d_{\tilde{H}}(\tilde{f}(\tilde{x}'), \tilde{y}') = 2n + 2$, and this is a contradiction. Hence $\tilde{f}(\tilde{x}')$ is contained in the path P, and is the next point from the starting point $\tilde{f}(\tilde{x})$. Hence the path joining $\tilde{f}(\tilde{x}')$ to \tilde{y}' contains \tilde{y} , and this means that (1) holds but (2) does not hold.

On the other hand, suppose that \tilde{y}' is contained in P. Since \tilde{H} is a tree, \tilde{y}' is the previous point of the terminal point \tilde{y} of P. Since $d_{\tilde{H}}(\tilde{f}(\tilde{x}'), \tilde{y}') = 2n$, $\tilde{f}(\tilde{x}')$ is not contained in P. Then $Q = \tilde{f}(\tilde{x}') \cdot P \setminus \tilde{y}$ is the path joining $\tilde{f}(\tilde{x}')$ to \tilde{y}' . Thus, applying the above paragraph to Q instead of P, we conclude that (2) holds but (1) does not. \Box

We define the direction on an edge of $\tilde{G}(\eta)$ as follows: If $(\tilde{x}, \tilde{x}') \in E(\tilde{G}(\eta))$ satisfies (1) of Proposition 4.3, then we consider the source of e is \tilde{x} and the target of e is \tilde{x}' , and if the other holds, then we consider the source of e is \tilde{x}' and the target of e is \tilde{x} . Set

 $\vec{E}(\tilde{G}(\eta)) = \{(\tilde{x}, \tilde{x}') \in E(\tilde{G}(\eta)) \mid \text{the source of the edge is } \tilde{x} \text{ and the target of the edge is } \tilde{x}'\}$ This direction on $\tilde{G}(\eta)$ is Γ -equivariant, i.e., for every edge $(\tilde{x}, \tilde{x}') \in \vec{E}(\tilde{G}(\eta)), (\gamma \tilde{x}, \gamma \tilde{y}) \in \vec{E}(\tilde{G}(\eta))$. Therefore, the direction on $\tilde{G}(\eta)$ induces the direction on $G(\eta)$. We define the set $\vec{E}(G(\eta))$ in the same way.

Lemma 4.4. Let $e = (\tilde{x}, \tilde{x}') \in E(\tilde{G}(\eta))$. Then the following are equivalent:

- (1) $(\tilde{x}, \tilde{x}') \in \vec{E}(\tilde{G}(\eta)).$
- (2) There are $\tilde{y} \in \eta(\tilde{x})$ and $\tilde{y}' \in \eta(\tilde{x}')$ such that $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y}) = d_{\tilde{H}}(\tilde{f}(\tilde{x}'), \tilde{y}') = 2n$ and the path joining $\tilde{f}(\tilde{x}')$ to \tilde{y}' contains \tilde{y} .

Proof. By the definition of the direction of edges, (1) clearly implies (2). Next, suppose (2). It suffices to show that (2) of Proposition 4.3 does not hold. By (2) of this lemma, the path P in \tilde{H} joining $\tilde{f}(\tilde{x}')$ to \tilde{y}' contains \tilde{y} . Then \tilde{y} is a previous point of \tilde{y}' in P. Since $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y}) = 2n, \tilde{f}(\tilde{x})$ is not contained in P, and $Q = \tilde{f}(\tilde{x}) - P \setminus \tilde{y}'$ is the path joining $\tilde{f}(\tilde{x})$ to \tilde{y} . Then Q does not contain \tilde{y}' and hence (2) in Proposition 4.3 does not hold. This completes the proof.

Definition 4.5. A vertex \tilde{x} of $\tilde{G}(\eta)$ is a *sink* if there is no edge of $\tilde{G}(\eta)$ whose source is \tilde{x} . Similarly, a vertex x of $G(\eta)$ is a *sink* if there is no edge of $G(\eta)$ whose source is x.

Lemma 4.6. Let $\eta, \eta' \in X_n$ such that $\eta \leq \eta'$. Then the following hold:

- (1) The directed graph $\tilde{G}(\eta)$ is a directed subgraph of $\tilde{G}(\eta')$.
- (2) Suppose that $e \in \vec{E}(\tilde{G}(\eta'))$ and the target of e is not a sink of η' . Then $e \in \vec{E}(\tilde{G}(\eta))$.

Proof. We first show (1). Then we have $V(\tilde{G}(\eta)) = u_{\eta}^{-1}(2n) \subset u_{\eta'}^{-1}(2n) = V(\tilde{G}(\eta'))$. To complete the proof of (1), it suffices to show that $e \in \vec{E}(\tilde{G}(\eta))$ implies $\vec{E}(\tilde{G}(\eta))$, i.e., e has the same direction in $\tilde{G}(\eta)$ and $\tilde{G}(\eta')$. This clearly follows from Lemma 4.4.

Next we show (2). Let $e = (\tilde{x}, \tilde{x}') \in \vec{E}(\tilde{G}(\eta'))$. By the assumption, there is an edge of $\tilde{G}(\eta')$ whose source is \tilde{x}' . This means that $\eta'(\tilde{x})$ and $\eta'(\tilde{x}')$ are one point sets (see Proposition 4.3). Thus we have $\eta(\tilde{x}) = \eta'(\tilde{x})$ and $\eta(\tilde{x}') = \eta'(\tilde{x}')$, and e has the same direction as that in $\tilde{G}(\eta)$.

Lemma 4.7. The directed graph $G(\eta)$ has no directed cycle.

Proof. Suppose that $G(\eta)$ has a directed cycle $\gamma = (x_0, \ldots, x_m)$ with $x_0 = x_m$, and let $\tilde{\gamma} = (\tilde{x}_0, \ldots, \tilde{x}_m)$ be a lift of γ on \tilde{G} . Then there is $g \in \Gamma$ such that $\tilde{x}_m = g\tilde{x}_0$. Then we have a directed walk

(4.1)
$$g^{-1}\tilde{x}_{m-1} \to \tilde{x}_0 \to \tilde{x}_1 \to \dots \to \tilde{x}_m \to g\tilde{x}_1$$

in G. Proposition 4.3 implies that $\eta(\tilde{x}_0), \dots, \eta(\tilde{x}_m)$ are one point sets. Let $\eta' \in \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$. If $\eta' \leq \eta$, then $\eta'(\tilde{x}_i) = \eta(\tilde{x}_i)$ for every *i*. If $\eta \leq \eta'$, then Lemma 4.6 implies that $\tilde{G}(\eta')$ contains the above directed path (4.1). Hence $\eta'(\tilde{x}_0), \dots, \eta'(\tilde{x}_m)$ are also one point sets and hence we have $\eta(\tilde{x}_i) = \eta'(\tilde{x}_i)$. This means that for every $\eta'' \in \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}, \eta''(\tilde{x}_i) = \eta(\tilde{x}_i)$. This is a contradiction since $\tilde{f} \in \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$ and $d_{\tilde{H}}(\tilde{f}(\tilde{x}_i), \eta(\tilde{x}_i)) = 2n > 0$.

Since G is a finite graph, Lemma 4.7 immediately deduces the following corollary:

Corollary 4.8. The directed graph $G(\eta)$ has a sink, and so does $G(\eta)$.

Lemma 4.9. Let $e = (\tilde{x}, \tilde{x}') \in E(\tilde{G})$ and suppose $u_{\eta}(\tilde{x}) = 2n$. Let $\tilde{y} \in \eta(\tilde{x})$ such that $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y}) = 2n$. Suppose that $\tilde{x}' \notin V(\tilde{G}(\eta))$ or $(\tilde{x}', \tilde{x}) \in \vec{E}(\tilde{G}(\eta))$. Then $\eta(\tilde{x}')$ consists of one point \tilde{y}' and \tilde{y}' is contained in the path P joining $\tilde{f}(\tilde{x})$ to \tilde{y} . Moreover, \tilde{y}' is the previous point of the terminal point \tilde{y} of P.

Proof. Let $\tilde{y}' \in \eta(\tilde{x}')$. Suppose that \tilde{y}' is not contained in P. Then $P - \tilde{y}'$ is the path with length 2n + 1 joining $\tilde{f}(\tilde{x})$ to \tilde{y}' . Since $\tilde{f}(\tilde{x}')$ is adjacent to $\tilde{f}(\tilde{x})$ and $\eta \in X_n$, we have $d_{\tilde{H}}(\tilde{f}(\tilde{x}'), \tilde{y}') = 2n$ and $\tilde{x}' \in V(\tilde{G}(\eta))$. Since P does not contain \tilde{y}' , we have $(\tilde{x}, \tilde{x}') \in \vec{E}(\tilde{G}(\eta))$. This contradicts the hypothesis.

Hence \tilde{y}' is contained in *P*. Since \tilde{y} and \tilde{y}' are adjacent, \tilde{y}' is the previous point of \tilde{y} . Since such a point is unique, the set $\eta(\tilde{x}')$ consists of one point \tilde{y}' .

4.4. Subposets $X_{n,m}$ and $X_{n,m,i}$. We are now ready to introduce the filtrations $X_{n,m}$ and $X_{n,m,i}$ of X_n . Let m be an integer at least -1. Define the induced subposet $X_{n,m}$ of X_n as follows: $\eta \in X_{n,m}$ if and only if $G(\eta)$ has no directed path having the length greater than m. Here we consider that a vertex is a directed path with length 0. Then we have $X_{n-1} = X_{n,-1}$ and $X_{n,m-1} \subset X_{n,m}$ for $m \ge 0$. Since G is a finite graph, we have $X_n = X_{n,m}$ for sufficiently large m.

Next we define the subposet $X_{n,m,i}$. Let P_1, \ldots, P_l be the directed paths with length m in G. For $i \ge 0$, let $X_{n,m,i}$ be the induced subposet of $X_{n,m}$ consisting of multi-homomorphisms $\eta \in X_{n,m}$ such that $G(\eta)$ does not contain P_{i+1}, \ldots, P_l . Then $X_{n,m,0} = X_{n,m-1}$ and $X_{n,m,i} = X_{n,m}$ for every $i \ge l$. These are the desired conditions described in Subsection 4.1.

4.5. **Deformation retracts.** Thus, to complete the proof of Theorem 4.1, it suffices to show the following (see Lemma 2.19):

Proposition 4.10. For $i \ge 1$, the poset $X_{n,m,i-1}$ is a deformation retract of $X_{n,m,i}$

The rest of this section is devoted to the proof of Proposition 4.10. We define an ascending closure operator $U: X_{n,m,i} \to X_{n,m,i}$ and a descending closure operator $D: U(X_{n,m,i}) \to U(X_{n,m,i})$ such that $DU(X_{n,m,i}) = X_{n,m,i-1}$.

First we define the map $U: X_{n,m,i} \to X_{n,m,i}$ as follows: Let $\eta \in X_{n,m,i}$. If $\eta \in X_{n,m,i-1}$, then we set $U(\eta) = \eta$. Suppose $\eta \notin X_{n,m,i-1}$. Then $G(\eta)$ contains P_i .

Let x be the terminal point of P_i . Let \tilde{x} be a vertex of $V(G(\eta))$ such that $p_G(\tilde{x}) = x$. Then there is $\tilde{y} \in \eta(\tilde{x})$ such that $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y}) = 2n$. Let P be the path joining $\tilde{f}(\tilde{x})$ to \tilde{y} . Let \tilde{x}_{η} be the vertex on P such that $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{x}_{\eta}) = 2n - 2$. Then \tilde{x}_{η} does not depend on the choice of \tilde{y} . Indeed, let \tilde{x}' be a vertex adjacent to \tilde{x} . Lemma 4.9 implies that $\eta(\tilde{x}')$ consists of one vertex \tilde{y}' , and \tilde{y}' is the previous vertex of the terminal point \tilde{y} of P. Then \tilde{x}_{η} is the previous vertex of the terminal point \tilde{y} does not depend on the choice of $\tilde{y} \in \eta(\tilde{x}), \tilde{x}_{\eta}$ does not depend on the choice of \tilde{y} .

Now we define the map $U(\eta) \colon V(\tilde{G}) \to \mathcal{P}(V(\tilde{G})) \setminus \{\emptyset\}$ as follows:

$$U(\eta)(\tilde{z}) = \begin{cases} \eta(\tilde{z}) \cup \{\tilde{z}_{\eta}\} & (p_G(\tilde{z}) = x) \\ \eta(\tilde{z}) & (p_G(\tilde{z}) \neq x). \end{cases}$$

Then $U(\eta)$ is clearly Γ -equivariant and $U^2 = U$. We would like to show the following:

- (1) $U(\eta)$ is a multi-homomorphism.
- (2) $G(\eta) = G(U(\eta))$. Similarly, $G(\eta) = G(U(\eta))$. In particular, $U(\eta) \in X_{n,m,i}$.
- (3) U is monotone.

First we show that $U(\eta)$ is a multi-homomorphism. Since $U(\eta)$ and η coincide except for $p_G^{-1}(x)$, it suffices to observe edges connecting with a vertex contained in $p_G^{-1}(x)$. Let $\tilde{x} \in p_G^{-1}(x)$ and \tilde{x}' a vertex adjacent to \tilde{x} . Let $\tilde{y}' \in \eta(\tilde{x}')$. It suffices to check that \tilde{x}_{η} and \tilde{y}' are adjacent. Let $\tilde{y} \in \eta(\tilde{x})$ satisfying $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y}) = 2n$, and P the path joining $\tilde{f}(\tilde{x})$ to \tilde{y} . Then it follows from Lemma 4.9 that \tilde{y}' is the previous point of the terminal point \tilde{y} of P. Since \tilde{x}_{η} is the vertex of P with $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{x}_{\eta}) = 2n - 2$, \tilde{x}_{η} and \tilde{y}' are adjacent. This completes the proof of (1).

Next, we show (2). It suffices to show only $\tilde{G}(\eta) = \tilde{G}(U(\eta))$. Since for every $\tilde{z} \in V(\tilde{G})$, $U(\eta)(\tilde{z})$ is just adding a vertex at distance 2n-2 from $\tilde{f}(\tilde{z})$, we have $V(\tilde{G}(\eta)) = V(\tilde{G}(U(\eta)))$ and the direction of each edge coincides (see Lemma 4.4). This completes the proof of (2).

Finally, we show (3). Let $\eta, \eta' \in X_{n,m,i}$ and $\eta \leq \eta'$. If $U(\eta) = \eta$, then $U(\eta) = \eta \leq \eta' \leq U(\eta')$. Hence we consider the case that $\eta < U(\eta)$. Then $\eta \notin X_{n,m,i-1}$. Let x be the terminal point of P_i and $\tilde{x} \in p_G^{-1}(x)$. Since $U(\eta) \neq \eta$, $\eta(\tilde{x})$ does not contain \tilde{x}_{η} . Since $\eta \leq \eta'$, Lemma 4.6 implies that $G(\eta')$ also contains P_i . If x is not a sink in $G(\eta')$, then $G(\eta')$ contains a directed path whose length is greater than m. This contradicts the assumption that $\eta' \in X_{n,m}$. Hence x is a sink of $G(\eta')$ and $\tilde{x}_{\eta'} = \tilde{x}_{\eta}$. Hence we have

$$U(\eta')(\tilde{x}) = \eta'(\tilde{x}) \cup \{\tilde{x}_{\eta'}\} \supset \eta(\tilde{x}) \cup \{\tilde{x}_{\eta}\} = U(\eta)(x).$$

This completes the proof of (3).

Next we define the map $D: U(X_{n,m,i}) \to U(X_{n,m,i})$. If $\eta \in X_{n,m,i-1}$, then set $D(\eta) = \eta$. Suppose $\eta \notin X_{n,m,i-1}$. Then $G(\eta)$ contains P_i . Let x be the terminal point of P_i and let $\tilde{x} \in p_G^{-1}(x)$. Since $\eta \in U(X_{n,m,i})$ and $U^2 = U$, $\eta(\tilde{x})$ contains \tilde{x}_{η} . Hence we define $D(\eta) \in X_{n,m,i}$ by

$$D(\eta)(\tilde{z}) = \begin{cases} \{\tilde{z}_{\eta}\} & (p_G(\tilde{z}) = x) \\ \eta(\tilde{z}) & (p_G(\tilde{z}) \neq x). \end{cases}$$

14

Since $\emptyset \neq D(\eta)(\tilde{z}) \subset \eta(\tilde{z})$ for each $\tilde{z} \in V(G)$ and η is a multi-homomorphism, $D(\eta)$ is a multi-homomorphism. It is clear that $D(\eta)$ is Γ -equivariant. Note that $D(\eta)(\tilde{x})$ does not contain a vertex \tilde{y} such that $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y}) = 2n$. This means $\tilde{x} \notin V(\tilde{G}(D(\eta)))$. By Lemma 4.6, we have $D(\eta) \in X_{n,m,i-1} \subset U(X_{n,m,i})$.

We show that $D: U(X_{n,m,i}) \to U(X_{n,m,i})$ is monotone. Let $\eta, \eta' \in U(X_{n,m,i})$. We only need to consider the case $D(\eta') < \eta'$. Let x be the terminal point of P_i . Suppose that $D(\eta) \leq D(\eta')$ does not hold. Then $\eta \leq D(\eta')$ does not hold and $\eta \leq \eta'$. Hence there is $\tilde{x} \in p_G^{-1}(x)$ and $\tilde{y} \in \eta(\tilde{x}) \subset \tilde{\eta}'(\tilde{x})$ such that $d_{\tilde{H}}(\tilde{f}(\tilde{x}), \tilde{y}) = 2n$. By Lemma 4.6, $G(\eta)$ contains P_i . Since $\eta \in U(X_{n,m,i})$, we have $U(\eta) = \eta$, and hence we have that $\eta(\tilde{x})$ contains \tilde{x}_{η} . This means that $\tilde{y} \notin D(\eta)(\tilde{x})$, this is a contradiction. This completes the proof that D is monotone.

We have shown that $U: X_{n,m,i} \to X_{n,m,i}$ and $D: U(X_{n,m,i}) \to U(X_{n,m,i})$ are closure operators and $DU(X_{n,m,i}) = X_{n,m,i-1}$. By Theorem 2.2, $X_{n,m,i-1}$ is a deformation retract of $X_{n,m,i}$. This completes the proof of Proposition 4.10.

5. Fundamental group of $\operatorname{Hom}(G, H)_f$

In this section, we study the fundamental group of $\text{Hom}(G, H)_f$. In Subsection 5.1, we establish a relationship between the fundamental group of $\text{Hom}(G, H)_f$ and realizable walks studied in Wrochna [34] (Theorem 1.5). In Subsection 5.2, we complete the proof of Theorem 1.2, using a fact (Theorem 5.7) proved in [34].

5.1. **Realizable elements.** In his study of the homomorphism reconfiguration problem, Wrochna [34] considered the group $\Pi(f, v)$ for the case that H is square-free. While his work deals only with the case where H is square-free, it is straightforward to generalize it in terms of the 2-homotopy classes of walks (see also [30]).

For the moment, suppose that H is an arbitrary connected graph. Let h be a \times -homotopy from f to f (see Subsection 2.3), i.e., a graph homomorphism $h: G \times I_n \to H$ such that h(x,0) = h(x,n) = f(x). Let $v' \in N_G(v)$. Then we have a closed walk

$$f(v) = h(v, 0) \sim h(v', 0) \sim h(v, 1) \sim \dots \sim h(v', n-1) \sim h(v, n) = f(v)$$

of H with even length. Although this closed walk depends on the choice of $v' \in N_G(v)$, its 2-homotopy class α does not. In this case, we say that α is realized by h. We call an element of $\pi_1^2(H, w)_{ev}$ realizable if it is realized by a \times -homotopy. The realizable elements form a subgroup of $\pi_1^2(H, w)$, and we denote this group by $\Pi(f, v)$.

We now provide a characterization of the realizability in terms of $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$. In the following theorem, recall that $\pi_1^2(H, w)$ acts on the universal 2-covering \tilde{H} (Subsection 2.6).

Theorem 5.1. Let ϖ be an element of $\pi_1^2(H, w)$. Then the following are equivalent:

(1) ϖ is realizable, i.e., $\varpi \in \Pi(f, v)$.

(2) $\varpi \tilde{f}$ belongs to Hom^{Γ} $(\tilde{G}, \tilde{H})_{\tilde{f}}$. Here $\varpi \tilde{f}$ is a graph homomorphism sending \tilde{x} to $\varpi(\tilde{f}(\tilde{x}))$.

Proof. We first prove (2) \Rightarrow (1). If (2) holds, by the same argument to show Proposition 2.7, we have a \times -homotopy $\tilde{h}: \tilde{G} \times I_n \to \tilde{H}$ from \tilde{f} to $\varpi \cdot \tilde{f}$ such that for every $t \in \{0, 1, \dots, n\}$,

the map $V(\tilde{G}) \to V(\tilde{H}), \tilde{x} \mapsto \tilde{h}(\tilde{x}, t)$ is Γ -equivariant. Let $\tilde{v}' \in N_{\tilde{G}}(\tilde{v})$ and consider the walk $\varphi = (\tilde{h}(\tilde{v}, 0), \tilde{h}(\tilde{v}', 0), \tilde{h}(\tilde{v}, 1), \tilde{h}(\tilde{v}', 1), \cdots \tilde{h}(\tilde{v}, n))$

in \tilde{G} . Then φ joins \tilde{w} to $\varpi \cdot \tilde{w}$. By Lemma 2.18, the 2-homotopy class of $p_H \circ \varphi$ is ϖ .

On the other hand, \tilde{h} induces a ×-homotopy $h: G \times I_n \to H$ such that $h(p_G(\tilde{x}), t) = p_H(\tilde{h}(\tilde{x}, t))$. Then the 2-homotopy class realized by h coincides with the 2-homotopy class of $p_H \circ \varphi$, which is ϖ . This completes the proof of $(2) \Rightarrow (1)$.

Next we show (1) \Rightarrow (2). Suppose that ϖ is realizable. Let $h: G \times I_n \to H$ be a ×homotopy from f to f such that the 2-homotopy class realized by h is ϖ . For $i = 0, 1, \dots, n$, define the graph homomorphism $f_i: G \to H$ by $f_i(x) = h(x, i)$. Then Lemma 2.6 implies that there is a sequence

$$f = f_0 \le \eta_0 \ge f_1 \le \dots \ge f_n = f$$

in Hom(G, H). By Proposition 3.1, there is a sequence

$$\tilde{f} = \tilde{f}_0 \le \tilde{\eta}_0 \ge \tilde{f}_1 \le \dots \ge \tilde{f}_n$$

in $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})$ such that $\pi(\tilde{f}_i) = f_i$ and $\pi(\tilde{\eta}_i) = \eta_i$. Then we have a ×-homotopy $\tilde{h} \colon \tilde{G} \times I_n \to \tilde{H}$ defined by $\tilde{h}(\tilde{x}, i) = \tilde{f}_i(\tilde{x})$. Note that the 2-homotopy class of walks realized by \tilde{h} is a lift of that of h. Hence, by the description of the action on \tilde{H} (see Subsection 2.6), the 2-homotopy class of walks realized by \tilde{h} joins $\tilde{f}(\tilde{v})$ to $\varpi \tilde{f}(\tilde{v})$. This means that $\tilde{f}_n(\tilde{v}) = \varpi \cdot \tilde{f}(\tilde{v})$ and $p_H \circ \tilde{f}_n = p_H \circ (\varpi \tilde{f})$. Thus, the uniqueness of the lifting criterion (Lemma 2.14) implies that $\varpi \tilde{f} = \tilde{f}_n \in \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$.

The following corollary is known for the case that H is square-free (see [34, Lemma 4.1]).

Corollary 5.2. Let $\alpha \in \pi_1^2(G, v)$ and $\varpi \in \Pi(f, v)$. Then $f_*(\alpha) \varpi = \varpi f_*(\alpha)$ in $\pi_1^2(H, w)$.

Proof. It follows from Theorem 5.1 that \tilde{f} and $\varpi \tilde{f}$ are Γ -equivariant. Hence we have

$$(f_*(\alpha)\varpi) \cdot \tilde{f}(\tilde{v}) = (\varpi\tilde{f})(\alpha\tilde{v}) = (\varpi f_*(\alpha))(\tilde{f}(\tilde{v})).$$

Since the action of $\pi_1^2(H, w)$ on $V(\tilde{H})$ is free, we conclude $\varpi f_*(\alpha) = f_*(\alpha) \varpi$.

Lemma 5.3. Let $\tilde{g} \in \operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$ and suppose $\pi(\tilde{g}) = f$ (for the definition of π : $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H}) \to \operatorname{Hom}(G, H)$, see Section 3). Then there is $\varpi \in \pi_1^2(H, w)$ such that $\tilde{g} = \varpi \tilde{f}$.

Proof. Since $p_H(\tilde{g}(\tilde{v})) = \pi(\tilde{g})(v) = f(v) = p_H(\tilde{f}(\tilde{v}))$, there is $\varpi \in \pi_1^2(H, w)$ such that $\tilde{g}(\tilde{v}) = \varpi \tilde{f}(\tilde{v})$. By the uniqueness of the lifting criterion (Lemma 2.14), we have $\tilde{g} = \varpi \tilde{f}$. \Box

We are now ready to prove Theorem 1.5. For the reader's convenience, we restate the theorem below.

Theorem 1.5. There is a following short exact sequence of groups:

$$1 \to \pi_1(\operatorname{Hom}^{\Gamma}(G,H)_{\tilde{f}},f) \to \pi_1(\operatorname{Hom}(G,H)_f,f) \to \Pi(f,v) \to 1.$$

Proof of Theorem 1.5. This follows from Theorem 2.20, Theorem 5.1 and Lemma 5.3. \Box

The following corollary is a direct consequence of Theorem 1.5.

Corollary 5.4. If $\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}}$ is simply connected, then $\pi_1(\operatorname{Hom}(G, H)_f, f)$ and $\Pi(f, v)$ are isomorphic.

Remark 5.5. In general, $\operatorname{Hom}^{\Gamma}(G, H)_{\tilde{f}}$ is not necessarily simply connected. Indeed, consider the case that $G = K_n$, $H = K_{n+1}$ for $n \geq 3$, and $f: G \to H$ is any graph homomorphism. Since $\pi_1^2(H, f(v))_{ev} \cong \pi_1(\mathcal{N}(K_{n+1}), f(v)) \cong 1$ (Theorem 2.9). Since $\Pi(f, v) \subset \pi_1^2(H, f(v))_{ev}$, $\Pi(f, v)$ is trivial. Hence Theorem 1.5 implies that $\pi_1(\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})) \cong \pi_1(\operatorname{Hom}(G, H)_f)$. Since $\operatorname{Hom}(G, H)$ is a wedge of circles and is not contractible (see [1, Section 4]), $\pi_1(\operatorname{Hom}^{\Gamma}(\tilde{G}, \tilde{H})_{\tilde{f}})$ is non-trivial in this case.

Recall that if the universal covering of a simplicial complex X is contractible, then the homotopy type of X is determined by its fundamental group (see [14, Theorem 1B.8]). Thus for a group Π , we denote by $K(\Pi, 1)$ the simplicial complex such that its universal covering is contractible and $\pi_1(K(\Pi, 1)) \cong \Pi$. Note that if Π is trivial then $K(\Pi, 1)$ is contractible, and that if Π is an infinite cyclic group then $K(\Pi, 1)$ is a circle.

Theorem 4.1 and Corollary 5.4 imply the following:

Corollary 5.6. If H is a tree, then $\operatorname{Hom}(G, H)_f$ is $K(\Pi(f, v), 1)$.

We conclude this subsection by providing a direct description of the group homomorphism $\Psi \colon \pi_1(\operatorname{Hom}(G,H)_f, f) \to \Pi(f,v)$ in Theorem 1.5. Let $\alpha \in \pi_1(\operatorname{Hom}(G,H)_f, f)$. Then there is an edge path (see [29])

$$f = f_0 \le \eta_1 \ge \eta'_1 \le \eta_2 \ge \dots \le \eta_n \ge f_n = f$$

in Hom $(G, H)_f$ representing α . For each $i = 1, \dots, n-1$, let f_i be a graph homomorphism such that $f_i \leq \eta'_i$. Then the above edge path is homotopic to the edge path

$$f = f_0 \le \eta_1 \ge f_1 \le \eta_2 \ge \dots \le \eta_n \ge f_n.$$

Then Lemma 2.6 implies that we have a \times -homotopy $h: G \times I_n \to H$ from f to f. The element realized by h is $\Psi(\alpha)$.

To see this, by the same argument of the proof of Theorem 5.1, we can take

$$\tilde{f} = \tilde{f}_0 \le \tilde{\eta}_1 \ge \tilde{f}_1 \le \tilde{\eta}_2 \ge \dots \le \tilde{\eta}_n \ge \tilde{f}_n$$

a sequence of $\operatorname{Hom}^{\Gamma}(\tilde{f}, \tilde{g})_{\tilde{f}}$ such that $\pi(\tilde{f}_i) = f_i$ and $\pi(\tilde{\eta}_i) = \eta_i$. By Theorem 2.20 we have $\tilde{f}_n = \Psi(\alpha) \cdot \tilde{f}$. Define the Γ -equivariant \times -homotopy $\tilde{h} \colon \tilde{G} \times I_n \to \tilde{H}$ by $\tilde{h}(\tilde{x}, i) = \tilde{f}_i(\tilde{x})$. The 2-homotopy class of walks realized by \tilde{h} joins \tilde{v} to $\Psi(\alpha)\tilde{v}$, and $p_H \colon \tilde{H} \to H$ sends the 2-homotopy class realized by \tilde{h} to the 2-homotopy class realized by h. Thus Lemma 2.18 implies that the 2-homotopy class realized by h is $\Psi(\alpha)$. This completes the proof.

5.2. **Proof of Theorem 1.2.** In this subsection, we complete the proof of Theorem 1.2.

When H is square-free, Wrochna provided an algorithm to determine the isomorphism type of $\Pi(f, v)$ (see [34, Theorem 8.1]). However, to prove Theorem 1.2, it suffices to rely on a theorem of the following form. Here, recall that $\pi_1(H, w)$ is a free group, and every subgroup of a free group is also a free group (see [14, Proposition 1A.2 and Theorem 1A.4]).

Theorem 5.7 (Wrochna [34]). Assume that H is square-free. Let K be the image of the group homomorphism $f_*: \pi_1(G, v) \to \pi_1(H, w)$. Then the following hold:

(1) If K is non-abelian free, then $\Pi(f, v)$ is trivial.

- (2) If K is an infinite cyclic group, then $\Pi(f, v)$ is trivial or an infinite cyclic group.
- (3) If K is trivial, then $\Pi(f, v)$ is $\pi_1(H, f(v))_{ev}$.

In Appendix, we provide an alternative proof of Theorem 5.7. In this section, we complete the proof of the main theorem (Theorem 1.2) by Theorem 5.7. The following is the main part of Theorem 1.2.

Theorem 5.8. Let H a square-free graph. Let K be the image of the group homomorphism

$$f_* \colon \pi_1(G, v) \to \pi_1(H, w).$$

Then the following hold:

- (1) If K is a non-abelian free group, then $\operatorname{Hom}(G, H)_f$ is contractible.
- (2) If K is an infinite cyclic group, then $\operatorname{Hom}(G, H)_f$ is contractible or homotopy equivalent to a circle.
- (3) If K is trivial and H is bipartite, then $\operatorname{Hom}(G,H)_f$ is homotopy equivalent to H.
- (4) If K is trivial and H is non-bipartite, then $\operatorname{Hom}(G, H)_f$ is homotopy equivalent to a connected double cover of H.

Proof of Theorem 5.8. Suppose (1). Then Theorem 5.7 implies that $\pi_1(\operatorname{Hom}(G,H)_f) \cong \Pi(f,v)$ is trivial, and hence $\operatorname{Hom}(G,H)_f = K(\Pi(f,v),1)$ (Corollary 5.6) is contractible. The proof of (2) is similar and is omitted. To see (3) and (4), suppose that K is trivial. Then, by Theorem 5.7 and Corollary 5.6, we have $\operatorname{Hom}(G,H)_f = K(\pi_1(H,v)_{ev},1)$. If H is bipartite, we have $\pi_1(H,v)_{ev} = \pi_1(H,v)$, hence $\operatorname{Hom}(G,H)_f = K(\pi_1(H),1) \simeq H$. If H is non-bipartite, then $\pi_1(H,v)_{ev}$ is a subgroup of $\pi_1(H,v)$ index 2. Hence there is a connected double cover H^* of H such that $\pi_1(H^*)$ is isomorphic to $\pi_1(H,v)_{ev}$. This shows that $\operatorname{Hom}(G,H)_f = K(\pi_1(H^*),1)$ is homotopy equivalent to H^* . This completes the proof. \Box

In particular, when G is non-bipartite, we have the following corollary:

Corollary 5.9. Let G be a non-bipartite graph, and H a square-free graph. Then every component of Hom(G, H) is contractible or homotopy equivalent to S^1 .

Proof. Let $f: G \to H$ be a graph homomorphism. Since G is non-bipartite, $\pi_1(G)$ has an odd element α . Then $f_*(\alpha) \in \pi_1(H)$ is also an odd element, and hence is non-trivial. Therefore $f_*: \pi_1(G) \to \pi_1(H)$ is non-trivial, and Theorem 5.8 completes the proof. \Box

We are now ready to complete the proof of Theorem 1.2.

Proof of Theorem 1.2. This follows from Theorem 5.8 and Corollary 5.9. \Box

Next we consider the case where G is bipartite. Then there is a graph homomorphism $f: G \to H$ such that $f_*: \pi_1(G) \to \pi_1(H)$ is trivial.

Proposition 5.10. Let G be a bipartite graph, and H a square-free graph which is neither contractible nor homotopy equivalent to S^1 . Then the following hold:

(1) If H is bipartite, then the number of components of Hom(G, H) homotopy equivalent to H is 2.

18

(2) If H is non-bipartite, then the number of components of Hom(G, H) homotopy equivalent to a connected double cover of H is 1.

Proposition 5.10 follows from Lemma 5.11 below and Theorem 5.8. Although Lemma 5.11 is likely a folklore fact, we provide the proof here for the reader's convenience.

Lemma 5.11. Let G be a bipartite graph. Then the following hold:

- (1) If H is bipartite, then the number of components of Hom(G, H) containing a graph homomorphism f which induces a trivial map $\pi_1(G) \to \pi_1(H)$ is 2.
- (2) If H is non-bipartite, then the number of components of Hom(G, H) containing a graph homomorphism f which induces a trivial map $\pi_1(G) \to \pi_1(H)$ is 1.

The following lemma seems to be well-known, and easily follows from the fold theorem (see [17, Theorem 3.3] for example) and Lemma 2.19.

Lemma 5.12. Let T be a (possibly infinite) tree having at least one edge. Then we have $\operatorname{Hom}(G,T) \simeq S^0$. Moreover, for two graph homomorphisms f_0 , $f_1: G \to T$, f_0 and f_1 belong to the same connected component of $\operatorname{Hom}(G,T)$ if and only if for some (or every) $x \in V(G)$, the distance $d_T(f_0(x), f_1(x))$ is even.

Proof of Lemma 5.11. In this proof, we call a graph homomorphism f null-homotopic if the map $f_*: \pi_1(G) \to \pi_1(H)$ induced by f is trivial.

Let $p: T \to H$ be the usual universal covering. Suppose that $f: G \to H$ is null-homotopic. By the usual lifting criterion [14, Proposition 1.33], there is a graph homomorphism $f': G \to T$ such that $p \circ f' = f$. Hence, f is contained in the image of $p_*: \text{Hom}(G, T) \to \text{Hom}(G, H)$. Since $\text{Hom}(G, T) \simeq S^0$ (Lemma 5.12), the number of components of Hom(G, H) containing null-homotopic graph homomorphisms is at most 2.

We now show (1). Suppose that H is bipartite and let $\varepsilon \colon H \to K_2$ be a graph homomorphism. Then there are two null-homotopic graph homomorphisms $f_0, f_1 \colon G \to H$ such that $\varepsilon \circ f_0 \neq \varepsilon \circ f_1$. Since $\operatorname{Hom}(G, K_2) = S^0$, f_0 and f_1 belong to distinct components of $\operatorname{Hom}(G, H)$. Hence, the number of components containing null-homotopic graph homomorphisms is 2. This completes the proof of (1).

Next we show (2). Suppose that H is non-bipartite. Then there is an odd element $\alpha \in \pi_1(H)$. Then f' and $\alpha f'$ belong to different components of Hom(G, T) since the length of path joining f'(x) to $\alpha f'(x)$ is odd for every $x \in V(G)$ (see Lemma 5.12). Since $p_H \circ f' = p_H \circ (\alpha f')$, p_* sends two components of Hom(G, T) to the same component of Hom(G, H). Thus the number of components containing null-homotopic graph homomorphisms is 1. \Box

Appendix A. Alternative proof of Theorem 5.7

In Subsection 5.2, we proved Theorem 1.2, by using Theorem 5.7. In this appendix, we present an alternative proof of Theorem 5.7 from the perspective of 2-covering theory.

For a group Γ_0 and for a subset S of Γ_0 , let $Z_{\Gamma_0}(S)$ be the subgroup of Γ_0 consisting of elements of Γ_0 that commute with every element of S.

Lemma A.1. Let F be a free group and K a subgroup of F. Then the following hold: (1) If K is a non-abelian free group, then $Z_F(K)$ is trivial.

(2) If K is an infinite cyclic group, then $Z_F(K)$ is an infinite cyclic group.

Proof. We first show (1). Let $x \in Z_F(K)$ and consider the subgroup K' of F generated by $K \cup \{x\}$. Then, K' is a non-abelian free group, and x belongs to the center of K'. Since a non-abelian free group has the trivial center, we conclude that x is trivial.

Next we show (2). Let $x \in K$ be a generator of K. Then $x \in Z_F(K)$ and hence $Z_F(K)$ is a non-trivial subgroup of F. If $Z_F(K)$ is not an infinite cyclic group, then $Z_F(K)$ is a non-abelian free group and $x \in Z_F(Z_F(K))$. Since x is non-trivial, this contradicts (1). \Box

Alternative proof of Theorem 5.7. Consider the following commutative diagram:

$$\begin{array}{c} \pi_1(G,v) \xrightarrow{f_*} \pi_1(H,w) \\ \downarrow \qquad \qquad \downarrow \cong \\ \pi_1^2(G,v) \xrightarrow{f_*} \pi_1^2(H,w). \end{array}$$

The vertical arrows are the natural quotient maps (see Subsection 2.4). Since H is squarefree, the right vertical arrow is an isomorphism (Lemma 2.8). Hence $K = f_*(\pi_1(G, v))$ coincides with $f_*(\pi_1^2(G, v))$. Corollary 5.2 implies $\Pi(f, v) \subset Z_{\pi_1^2(H,w)}(K)$. Thus, (1) and (2) of Theorem 5.7 follows from Lemma A.1.

Next we show (3). Since $K = f_*(\pi_1^2(G, v))$ is trivial, Lemma 2.14 implies that there is a graph homomorphism $f': G \to \tilde{H}$ such that $p_H \circ f' = f$. Since H is square-free, \tilde{H} is a tree. Let $\alpha \in \pi_1^2(H, w)_{ev}$. It follows from Lemma 5.12 that f' and $\alpha f'$ belong to the same connected component of $\operatorname{Hom}(G, \tilde{H})$. By Proposition 2.7, there is a ×-homotopy $h: G \times I_n \to \tilde{H}$ from f' to $\alpha f'$. Then $p_H \circ h$ is a ×-homotopy from f to f. By Lemma 2.18, the 2-homotopy class realized by $p_H \circ h$ is α . Thus we have $\pi_1^2(H, w)_{ev} \subset \Pi(f, v)$. By the definition of $\Pi(f, v)$, it is clear that $\Pi(f, v) \subset \pi_1^2(H, w)_{ev}$. This completes the proof of (3).

Acknowledgement

The author was supported in part by JSPS KAKENHI Grant Numbers JP23K12975. The author thanks Soichiro Fujii, Kei Kimura and Yuta Nozaki for fruitful discussion. In particular, the discussions with them were important for coming up with the formulation of Theorem 1.2. The author also thanks Shun Wakatsuki for useful comments.

References

- E. Babson and D. N. Kozlov. Complexes of graph homomorphisms. Israel J. Math., 152:285–312, 2006. doi:10.1007/BF02771988.
- [2] E. Babson and D. N. Kozlov. Proof of the Lovász conjecture. Ann. of Math. (2), 165(3):965–1007, 2007. doi:10.4007/annals.2007.165.965.
- [3] J. A. Barmak and E. G. Minian. A note on coverings of posets, A-spaces and polyhedra. Homology Homotopy Appl., 18(1):143-150, 2016. doi:10.4310/HHA.2016.v18.n1.a8.
- [4] P. Csorba. Homotopy types of box complexes. Combinatorica, 27(6):669-682, 2007. doi:10.1007/s00493-007-2204-x.
- [5] P. Csorba. On the simple Z₂-homotopy types of graph complexes and their simple Z₂-universality. Canad. Math. Bull., 51(4):535-544, 2008. doi:10.4153/CMB-2008-053-9.

20

- [6] S. L. Čukić and D. N. Kozlov. The homotopy type of complexes of graph homomorphisms between cycles. Discrete Comput. Geom., 36(2):313–329, 2006. doi:10.1007/s00454-006-1245-z.
- [7] A. Dochtermann. Hom complexes and homotopy theory in the category of graphs. European J. Combin., 30(2):490-509, 2009. doi:10.1016/j.ejc.2008.04.009.
- [8] A. Dochtermann. Homotopy groups of Hom complexes of graphs. J. Combin. Theory Ser. A, 116(1):180–194, 2009. doi:10.1016/j.jcta.2008.06.001.
- [9] A. Dochtermann. The universality of Hom complexes of graphs. Combinatorica, 29(4):433-448, 2009. doi:10.1007/s00493-009-2376-7.
- [10] A. Dochtermann and C. Schultz. Topology of Hom complexes and test graphs for bounding chromatic number. Israel J. Math., 187:371–417, 2012. doi:10.1007/s11856-011-0085-6.
- [11] A. Dochtermann and A. Singh. Homomorphism complexes, reconfiguration, and homotopy for directed graphs. European J. Combin., 110:Paper No. 103704, 31, 2023. doi:10.1016/j.ejc.2023.103704.
- [12] S. Fujii, Y. Iwamasa, K. Kimura, Y. Nozaki, and A. Suzuki. Homotopy types of Hom complexes of graph homomorphisms whose codomains are cycles, 2024. URL: https://arxiv.org/abs/2408.04802, arXiv:2408.04802.
- [13] S. Fujii, K. Kimura, and Y. Nozaki. Homotopy types of Hom complexes of graph homomorphisms whose codomains are square-free, in preparation, 2024. URL: https://arxiv.org/abs/2412.19039, arXiv:2412.19039.
- [14] A. Hatcher. Algebraic topology. Cambridge University Press, Cambridge, 2002.
- [15] P. Hell and J. Nešetřil. Graphs and homomorphisms, volume 28 of Oxford Lecture Series in Mathematics and its Applications. Oxford University Press, Oxford, 2004. doi:10.1093/acprof:oso/9780198528173.001.0001.
- [16] D. Kozlov. Combinatorial algebraic topology, volume 21 of Algorithms and Computation in Mathematics. Springer, Berlin, 2008. doi:10.1007/978-3-540-71962-5.
- [17] D. N. Kozlov. A simple proof for folds on both sides in complexes of graph homomorphisms. Proc. Amer. Math. Soc., 134(5):1265–1270, 2006. doi:10.1090/S0002-9939-05-08105-0.
- [18] L. Lovász. Kneser's conjecture, chromatic number, and homotopy. J. Combin. Theory Ser. A, 25(3):319– 324, 1978. doi:10.1016/0097-3165(78)90022-5.
- [19] J. Matoušek and G. M. Ziegler. Topological lower bounds for the chromatic number: a hierarchy. Jahresber. Deutsch. Math.-Verein., 106(2):71–90, 2004.
- [20] T. Matsushita. Answers to some problems about graph coloring test graphs. European J. Combin., 45:59-64, 2015. doi:10.1016/j.ejc.2014.10.006.
- [21] T. Matsushita. Fundamental groups of neighborhood complexes, 2015. URL: https://arxiv.org/abs/1210.2803, arXiv:1210.2803.v2.
- [22] T. Matsushita. Fundamental groups of neighborhood complexes. J. Math. Sci. Univ. Tokyo, 24(3):321– 353, 2017.
- [23] T. Matsushita. Homotopy types of the Hom complexes of graphs. European J. Combin., 63:216-226, 2017. doi:10.1016/j.ejc.2017.03.009.
- [24] T. Matsushita. Z₂-indices and Hedetniemi's conjecture. Discrete Comput. Geom., 62(3):662-673, 2019. doi:10.1007/s00454-019-00090-1.
- [25] T. Matsushita and S. Wakatsuki. Dominance complexes, neighborhood complexes and combinatorial Alexander duals. J. Combin. Theory Ser. A, 211:Paper No. 105978, 2025. doi:10.1016/j.jcta.2024.105978.
- [26] J. Milnor. Morse theory, volume No. 51 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 1963. Based on lecture notes by M. Spivak and R. Wells.
- [27] C. Schultz. Graph colorings, spaces of edges and spaces of circuits. Adv. Math., 221(6):1733–1756, 2009.
 doi:10.1016/j.aim.2009.03.006.
- [28] C. Schultz. The equivariant topology of stable Kneser graphs. J. Combin. Theory Ser. A, 118(8):2291– 2318, 2011. doi:10.1016/j.jcta.2011.04.009.
- [29] E. H. Spanier. Algebraic topology. McGraw-Hill Book Co., New York-Toronto-London, 1966.
- [30] C. Tardif and M. Wrochna. Hedetniemi's conjecture and strongly multiplicative graphs. SIAM J. Discrete Math., 33(4):2218–2250, 2019. doi:10.1137/19M1245013.

- [31] J. W. Walker. From graphs to ortholattices and equivariant maps. J. Combin. Theory Ser. B, 35(2):171– 192, 1983. doi:10.1016/0095-8956(83)90070-9.
- [32] M. Wrochna. Square-free graphs are multiplicative. J. Combin. Theory Ser. B, 122:479–507, 2017. doi:10.1016/j.jctb.2016.07.007.
- [33] M. Wrochna. On inverse powers of graphs and topological implications of Hedetniemi's conjecture. J. Combin. Theory Ser. B, 139:267-295, 2019. doi:10.1016/j.jctb.2019.02.008.
- [34] M. Wrochna. Homomorphism reconfiguration via homotopy. SIAM J. Discrete Math., 34(1):328–350, 2020. doi:10.1137/17M1122578.
- [35] R. T. Živaljević. WI-posets, graph complexes and Z₂-equivalences. J. Combin. Theory Ser. A, 111(2):204-223, 2005. doi:10.1016/j.jcta.2004.12.002.

DEPARTMENT OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCE, SHINSHU UNIVERSITY, MATSUMOTO, NAGANO 390-8621, JAPAN

Email address: matsushita@shinshu-u.ac.jp