Fluid-particle interactions and fluctuation-dissipation relations III - Correlated fluctuations, regularity and added mass

Massimiliano Giona,^{1,*} Giuseppe Procopio,^{1,[†](#page-0-1)} and Chiara Pezzotti^{1,[‡](#page-0-2)}

¹Dipartimento di Ingegneria Chimica Materiali Ambiente Facoltà di Ingegneria Industriale, La Sapienza Università di Roma, via Eudossiana 18, 00184, Roma, Italy

(Dated: December 30, 2024)

Abstract

The fluctuation-dissipation theory is grounded on the Langevin condition expressing the local independence between the thermal force and the particle velocity history. Upon hydrodynamic grounds, it is reasonable to relax this condition in order to account for the correlated fluid fluctuations, especially in the case of liquids, consistently with the inclusion of acoustic effects and with the finite speed of propagation of internal shear stresses. We show that the introduction of correlated stochastic processes in the basic fluctuational patterns defined in Giona et al. (2024), preserves the global fluctuation-dissipation relation, connecting diffusivity to the global friction factor, and the resulting velocity fluctuations become almost everywhere smooth functions of time. Moreover, a fluctuational added mass arises as a consequence of correlations. This leads to a fluctuation-inertia relation, connecting the fluctuational added mass at microscale to its occurrence for macroscopic objects.

[∗] massimiliano.giona@uniroma1.it

[†] giuseppe.procopio@uniroma1.it

[‡] chiara.pezzotti@uniroma1.it

I. INTRODUCTION

In [\[1\]](#page-34-0) and [\[2\]](#page-35-0), henceforth referred to as part I and part II, respectively, we have followed the classical Kubo approach to fluctuation-dissipation relations, assuming that the thermal force $\mathbf{R}(t)$ at time t is uncorrelated to particle velocity at previous time instants,

$$
\langle \mathbf{R}(t) \otimes \mathbf{v}(0) \rangle = 0, \qquad t \ge 0 \tag{1}
$$

where \otimes indicates the tensor product. This is indeed the Langevin condition [\[3](#page-35-1), [4](#page-35-2)], that undoubtedly displays analytical advantages, as it provides straightforwardly the evolution equation for the velocity autocorrelation function, and suggests the white-noise nature of the stochastic forcings entering the various fluctuation-dissipation patterns introduced in part I, typically regarded as distributional derivatives of independent Wiener processes.

The Langevin condition eq. [\(1\)](#page-1-0) and the equipartition relation for the squared variance of the velocity entries at equilibrium

$$
\langle v_i^2 \rangle = \frac{k_B T}{m}, \qquad i = 1, 2, 3 \tag{2}
$$

where m is the particle mass and T the constant temperature of the fluid medium (coinciding with that of the particle), are the two foundative properties deriving from statistical physics, defining particle velocity fluctuations and characterizing the Kubo fluctuation-dissipation theory.

In recent years, stort-time and high-resolution experiments on Brownian particles both in liquids and gases [\[5](#page-35-3)[–8](#page-35-4)] have qualitatively demostrated the correctness of the scaling of the velocity autocorrelation function with time predicted by the Kubo theory, supplementing it with the hydromechanic analysis of the force exterted by the fluid onto the particle including also the fluid-inertial effects, i.e. the Basset force [\[9](#page-35-5)[–12\]](#page-35-6). On the other hand, these experiments have shown that Brownian motion trajectories and velocities at short time scales in Newtonian fluids (water, acetone) are smooth functions of time and more regular than what expected at the Einstein's times $[8, 13]$ $[8, 13]$ $[8, 13]$. Moreover, deviations from eq. (2) , i.e. from the equipartition result, have been reported [\[14](#page-35-8)].

The latter phenomenon can be attributed hydrodynamically to the added mass effect, that occurs as an instantaneous fluid-inertial contribution in incompressible fluids, proportional to the actual value of the particle acceleration [\[15](#page-35-9), [16](#page-35-10)]. However, as correctly observed

by Zwanzig [\[17\]](#page-36-0), this instantaneous effect could be the consequence of the simplified and unphysical assumption of the infinite sound velocity subsumed in the assumption of incompressibility. The analysis performed by Chow and Hermans [\[18](#page-36-1)], considering a compressible time-dependent Stokes equation for the fluid (i.e., a linear proportionality between pressure and density) supports this thesis. Therefore, the possible occurrence of a non-vanishing added-mass influencing the squared velocity variance, i.e.,

$$
\langle v_i^2 \rangle = \frac{k_B T}{m + m_a}, \qquad i = 1, 2, 3
$$
\n⁽³⁾

with $m_a > 0$, instead of eq. [\(2\)](#page-1-1), has become a crucial issue both theoretically and experimentally [\[19\]](#page-36-2), with relavant implications in the modeling of hydrodynamics at microscale [\[20\]](#page-36-3), and in the foundations of statistical physics.

The inclusion of the "tiny acoustic effects" (i.e. compressibility) to explain the statistical properties of micrometric particles in still fluids (Brownian motion) clearly indicates that the removal of the unphysical assumption of infinite sound velocity determines significant changes in the qualitative statistical properties of the particle velocity fluctuations. In turn, it necessarily implies the occurrence of correlated pressure fluctuations possessing a bounded velocity of propagation. It is important to underline that the results by Chow and Hermans are grounded on the classical approach to fluctuation-dissipation relations, namely on the validity of eq. [\(1\)](#page-1-0) and on the representation of the stochastic forcings as white-noise processes.

Assuming, as physically correct and statistically relevant, this "regularity ansatz" of the acoustic effects, an equivalent regularity ansatz should be extended, by consistency, to the other contributions involved in the fluctuation-dissipation characterization of particle dynamics. Specifically, the simplification of δ-correlated stochastic forcing (white-noise processes), should be removed on equal footing. And this, ultimately questions the physical validity of the Langevin condition eq. [\(1\)](#page-1-0), as an acceptable approximation at least for liquids.

The inclusion of correlated and more regular stochastic forcings, replacing the δ -correlated distributional derivatives of Wiener processes, in particle hydromechanic equations is also consistent with the hydrodynamic picture of fluid-inertial effects, giving rise to coherent and correlated motion of fluid elements in the neighbourhood of the moving solid object (the Darwin paths) [\[21](#page-36-4)]. It is also consistent with a principle of mutual coupling between physical systems: the thermal force $\mathbf{R}(t)$ originates also from the velocity fluctuations occurring in the fluid medium due to the motion of the colloidal particle, and therefore it is expected, especially in liquids, that $\mathbf{R}(t)$ should be correlated with the previous velocity history of the particle itself. In this regard, the Langevin condition eq. [\(1\)](#page-1-0) should be viewed as a long-term approximation for time-scales much larger than the characteristic dissipation and internal stress propagation times.

However, the inclusion of correlated forcings in the particle equation of motion and ultimately in fluctuation-dissipation theory, should be performed with great caution, and should be compatible with some fundamental form of fluctuation-dissipation condition confirmed by the experiments. This condition, at equilibrium, is given by the global fluctuationdissipation relation, that enabled Einstein to extract from the analysis of Brownian motion trajectories, the Avogadro number, subsequently confirmed experimentally by Perrin and by many others [\[22](#page-36-5)[–24\]](#page-36-6).

In this article we show that, introducing correlated stochastic forcings within the basic fluctuational patterns defined in part I, the global fluctuation-dissipation relation is preserved under very loose assumptions (essentially related to a normalization condition) in any complex fluid medium for which particle diffusion is asymptotically regular (i.e., the particle mean square displacement is asymptotically a linear function of time). The effect of correlated stochastic forcings within the fluctuational patterns is twofold: (i) compared to the white-noise driven counterparts, a transition is observed from fractal to regular velocity fluctuations; (ii) correlations induce the occurrence of an added mass in any fluid dynamic conditions (including the case of compressible hydrodynamics). These results are indeed in agreement with the recent experimental results on highly resolved Brownian motion in liquids [\[8,](#page-35-4) [13,](#page-35-7) [14](#page-35-8)].

Within the extension of fluctuation-dissipation theory developed in this article, the added mass is a consequence of the correlated hydrodynamic fluctuations in a macroscopically still fluid phase, induced by the high-frequency erratic motion of micrometric particles. A delicate issue arises in providing the connection between the fluctuation-induced added mass at microscale (i.e. for micrometric particles) and the "instantaneous" added mass observed for macroscopic objects (from millimetric particles to submarines and transoceanic ships), as consequence of fluid inertia [\[25](#page-36-7)[–27](#page-36-8)]. This connection is addressed in Section [VI,](#page-27-0) providing a new form of constraint referred to as the fluctuation-inertia relation.

The article is organized as follows. Section [II](#page-4-0) introduces the hydrodynamics of the problems and the importance of removing all the approximations associated with infinite propagation velocity of hydrodynamic fields in the analysis of microparticle velocity fluctuations. Section [III](#page-8-0) addresses the inclusion of correlated stochastic forcings within the basic fluctuational patterns introduced in part I, and derives for this class of models the validity of the global fluctuation-dissipation relation followed, as an aftermath, by the occurrence of a fluctuational added mass. Section [IV](#page-14-0) discusses some prototypical examples. The regularity of velocity fluctuations emerging from the correlated fluctuational patterns is the main focus of Section [V.](#page-22-0) More precisely, the presence of fluid inertia necessarily induces in the classical fluctuation-dissipation paradigm (based on white-noise stochastic forcings) an almost-everywhere singular (fractal) structure of velocity fluctuations that becomes regularized whenever correlated forcings are accounted for in these patterns. Section [VI](#page-27-0) addresses the connection between the fluctuational added mass deriving from the present theory and the instataneous added mass of incompressible hydrodynamics referred to macroscopic objects accelerating in a fluid. The concluding Section addresses some open issues and the experimental scrutiny of the theory.

II. STATEMENT OF THE PROBLEM: HYDRODYNAMIC CONSIDERATIONS

The advances in the theory of Brownian motion, as the paradigmatic problem involving the interaction between hydrodynamics and thermal fluctuations leading to the formulation of the fluctuation-dissipation theory, have been marked by a progressive regularization of the hydrodynamic description of the fluid medium, removing the paradoxes associated with the infinite propagation speed of the hydrodynamic fields.

Owing to linearity, particle hydromechanics in the free space (i.e. in the absence of constraints deriving from confinement, see part I) can be always expressed as

$$
m\frac{d\mathbf{v}(t)}{dt} = \mathbf{F}_{f\rightarrow p}[\mathbf{v}(t)] + \mathbf{R}(t)
$$
\n(4)

where $\mathbf{F}_{f\rightarrow p}[\mathbf{v}(t)]$ is the mean field force exerted by the fluid onto the particle, deriving from the hydrodynamic equations, and $\mathbf{R}(t)$ the thermal force possessing zero mean, $\langle \mathbf{R}(t) \rangle = 0$. The term $\mathbf{F}_{f\rightarrow p}[\mathbf{v}(t)]$ is a linear functional of the history of particle velocity from the initial time to the actual time t.

These advances towards an increasingly accurate hydrodynamic description of the fluid rheology and of the fluid-particle interactions correspond to the complexification of the fluctuational patterns discussed in part I, say from $\mathbb{S}_{m,E}^{(1,1,0)}$, where $E = k_B T$, to $\mathbb{D} \mathbb{I}_{m,E}^{(3,N_d,N_i)}$, i.e. a three-dimensional tensor model including dissipative and fluid-inertial memory effects via N_d and N_i modes, respectively, in which either the complex geometry of the object, the generic rheology of the fluid medium, and the time-dependent hydrodynamics of the fluid velocity field are taken into account.

The Einstein-Langevin picture refers to an incompressible Stokes hydrodynamics in which the reaction of the fluid to the perturbation associated with the motion of a solid object immersed in it is instantaneous (infinite speed of propagation of velocity perturbations). Thus, for a spherical particle of radius R_p , the force exerted by the fluid on the particle,

$$
\mathbf{F}_{f \to p}[\mathbf{v}(t)] = -6 \pi \mu R_p \mathbf{v}(t)
$$
\n(5)

is a function of the actual velocity value $\mathbf{v}(t)$, where μ is the fluid viscosity.

The hydrodynamic theory of Brownian motion developed in the '70 of the last century focused and improved the paradoxes (in the meaning by Birkhoff [\[28\]](#page-36-9)) of this representation. The instantaneous Stokes description was replaced by the time-dependent incompressible Stokes hydrodynamics. This leads for a Newtonian fluid to the following expression for the force $\mathbf{F}_{f\rightarrow p}[\mathbf{v}(t)]$ at time t

$$
\mathbf{F}_{f \to p}[\mathbf{v}(t)] = -6 \pi \mu R_p \mathbf{v}(t) - 6\sqrt{\pi \mu \rho_0} R_p^2 \int_0^t \frac{1}{\sqrt{t - \tau}} \left(\frac{d\mathbf{v}(\tau)}{d\tau} + \mathbf{v}(0) \delta(\tau) \right) d\tau
$$

\n
$$
- \frac{2}{3} \pi R_p^3 \rho_0 \frac{d\mathbf{v}(t)}{dt}
$$

\n
$$
= \frac{2}{3} \pi R_p^3 \rho_0 \frac{d\mathbf{v}(t)}{dt}
$$
 (6)

where ρ_0 is the density of the fluid, in which the Basset force emerges as a memory integral over the particle velocity history and the instantaneous added-mass term arises, proportional to the actual value of the particle acceleration. Expressed via a memory kernel, the added-mass term corresponds to an impulsive fluid-inertial kernel, proportional to a Dirac δ-distribution centered at $t = 0$.

Also the instantaneous added-mass term is a consequence of the paradox of infinite propagation velocity of the pressure field, intrinsic to the assumption of incompressibility. Incompressibility overlooks acoustic effects and corresponds to an infinite speed of sound in the fluid [\[15\]](#page-35-9).

The removal of the latter infinite-velocity paradox (and consequently the inclusion of acoustics within the particle hydromechanics) has been considered by Chow and Hermans [\[18\]](#page-36-1), expressing compressibility effects and finite sound speed in the form of a linear relation between pressure P and density ρ (of the fluid), that in the present case is no longer constant,

$$
\frac{dP}{d\rho} = c_s^2 \tag{7}
$$

where c_s is the sound velocity, considering the linearized hydrodynamics for the fluid velocity field $\mathbf{u}(\mathbf{x},t)$,

$$
\rho_0 \frac{\partial \mathbf{u}}{\partial t} = \mu \nabla^2 \mathbf{u} + \left(\frac{\mu}{3} + \kappa\right) \nabla (\nabla \cdot \mathbf{u}) - \nabla P
$$

\n
$$
\frac{\partial \rho}{\partial t} = -\rho_0 \nabla \cdot \mathbf{u}
$$
\n(8)

where ρ_0 is the constant density of the fluid at rest, and κ the bulk (compression) viscosity.

Solving the hydrodynamic problem in the free space equipped with no-slip conditions at the external surface S_p of the spherical particle, $\mathbf{u}(\mathbf{x}, t)|_{\mathbf{x} \in S_p} = \mathbf{v}(t)$, and evaluating the force exerted by the fluid on the particle, the added-mass effect disappears as an instantaneous contribution. More precisely, the hydromechanics induced by fluid-inertial effects is described by a convolution integral with respect to the particle acceleration defined by a smooth continuous kernel $k(t)$ not containing any impulsive (Dirac- δ) contribution.

The hydrodynamic pressure P accounts for compressible internal stresses in the fluid, that superimpose to the shear stresses. Including finite sound velocity and compressibility implies, by consistency, that also the paradox of infinite propagation velocity should be removed from the shear stress dynamics, which resides in the Newtonian constitutive equation of direct (and instantaneous) proportionality between the shear-stress tensor and the deformation tensor. The simplest way for achieving this goal is to consider a linear viscoelastic constitutive equation characterized by a single relaxation rate (Maxwell fluid) [\[29\]](#page-36-10).

In point of fact, the Newtonian constitutive equations are an approximation corresponding to the instantaneous response of the internal shear stresses to deformations. Even water at room temperature admits a viscoelastic behavior characterized by a relaxation rate order of 1 ps [\[30\]](#page-36-11).

Performing this extension (in the incompressible case) [\[31\]](#page-36-12), the Basset inertial contribution becomes a memory integral,

$$
\mathbf{F}_{f \to p}[\mathbf{v}(t)]|_{\text{Basset}} = -\int_0^t k(t - \tau) \left(\frac{d\mathbf{v}(\tau)}{d\tau} + \mathbf{v}(0) \,\delta(\tau) \right) d\tau \tag{9}
$$

where

$$
k(t) = \frac{6\sqrt{\pi \mu \rho_0} R_p^2}{\sqrt{\tau_c}} e^{-t/2\tau_c} I_0\left(\frac{t}{2\tau_c}\right)
$$
 (10)

where $I_0(\xi)$ is the modified Bessel function the first kind of zeroth order. and the singularity at $t = 0$ disappears. The regularization of the Basset kernel via the inclusion of viscoelastic constitutive equations, is the typical manifestation of the removal of the paradoxes and infinities associated with the infinite speed of propagation that characterizes all the parabolic transport schemes [\[32](#page-37-0), [33](#page-37-1)].

Starting from this hydrodynamic premise, it is consequential to extend the regularity issue to all the contribution entering the fluctuation-dissipation patterns, removing in this way the paradoxes of infinite propagation velocity and their unphysical consequences. According to this program, it is more realistic, as supported by the above mentioned hydrodynamic observations, to formulate a fluctuation-dissipation theory in which the stochastic forcing terms entering the fluctuational patterns would possess some colored characters (i.e. continuous and almost everywhere smooth correlation functions), instead of white-noise processes, such as the distributional derivatives of Wiener processes. As a consequence of this extension, the thermal force $\mathbf{R}(t)$ may be correlated with $\mathbf{v}(\tau)$ for $\tau \leq t$, i.e.

$$
\langle \mathbf{R}(t) \otimes \mathbf{v}(0) \rangle \neq 0, \qquad t \ge 0 \tag{11}
$$

corresponding to the breaking of the Langevin condition eq. [\(1\)](#page-1-0). Again, the hydrodynamic phenomenology supports this generalization: inertial effects and the finite sound velocity of acoustics in a fluid medium, push for the adoption of a corresponding property (of bounded propagation speed) as regards the stochastic forcings acting either on $\mathbf{v}(t)$ or on the auxiliary variables characterizing the local representation of the memory dynamics. However, the choice of the stochastic forcing $\mathbf{R}(t)$ could not be arbitrary, as it should fulfill some basic properties characterizing thermal equilibrium, expressed by the global fluctuation-dissipation relation. This can be achieved by resorting to the basic fluctuational patterns introduced in part I, substituting the white-noise processes with correlated ones.

III. CORRELATED FLUCTUATIONS IN BASIC PATTERNS: MAIN PROPER-TIES

In extending fluctuation-dissipation theory beyond the Langevin condition, the founding principle is the validity of the Einsteinian global fluctuation-dissipation relation for a spherical particle

$$
D\,\eta = k_B\,T\tag{12}
$$

relating the particle diffusion coefficient D to the Stokes friction factor η in a Newtonian fluid at constant temperature T.

Consider the case of a generic fluid characterized by linear rheological constitutive equation and a particle of arbitrary shape. The force acting on the particle is expressed by

$$
\mathbf{F}_{f \to p}[\mathbf{v}(t)] = -\mathbf{h}(t) * \mathbf{v}(t) - \mathbf{k}(t) * \left(\frac{d\mathbf{v}(t)}{dt} + \mathbf{v}(0)\,\delta(t)\right)
$$
(13)

where in general the dissipative $h(t)$ and the fluid-inertial $k(t)$ kernels have a tensorial character. As a byproduct, also particle motion becomes anisotropic and characterized by a diffusivity tensor D. Henceforth "∗" will indicate convolution.

The fluid inertial kernel does not contribute to dissipation, surely in the case of regularly diffusive case, and the extension of the Stokes-Einstein relation for the particle diffusion tensor D becomes

$$
\mathbf{D}\,\boldsymbol{\eta}_{\infty} = k_B\,T\,\mathbf{I} \tag{14}
$$

where I is the 3×3 identity matrix and the long-term friction factor η_{∞} is expressed by

$$
\eta_{\infty} = \int_0^{\infty} \mathbf{h}(t) dt
$$
 (15)

If η_{∞} is bounded, particle diffusion in the long-term is regular (i.e. characterized by a linear scaling of the mean-square displacement with time), otherwise anomalies (subdiffusive motion) may occur as discussed in part II. In the remainder, we consider exclusively the regular case. Enforcing the Green-Kubo relation [\[4\]](#page-35-2), eq. [\(14\)](#page-8-1) expresses a global condition on the properties of the particle velocity autocorrelation tensor $\mathbf{C}_{vv}(t) = \langle \mathbf{v}(t) \otimes \mathbf{v}(0) \rangle_{\text{eq}}$ (where the expected value $\langle \cdot \rangle_{\text{eq}}$, is referred to the probability mesure of velocity fluctuations at equilibrium),

$$
\int_0^\infty \mathbf{C}_{vv}(t) dt = k_B T \, \pmb{\eta}_{\infty}^{-1} \tag{16}
$$

This global condition is assumed as the fundamental constraint in the extension of fluctuation-dissipation theory to correlated perturbations.

Let $\widehat{\mathbf{C}}_{vv}(s) = \int_0^\infty e^{-st} \mathbf{C}_{vv}(t) dt$ be the Laplace transform of the velocity autocorrelation tensor. Equation [\(14\)](#page-8-1) corresponds to a constraint on the value of $\hat{\mathbf{C}}_{vv}(s)$ at $s = 0$,

$$
\widehat{\mathbf{C}}_{vv}(0) = k_B T \,\boldsymbol{\eta}_{\infty}^{-1} \tag{17}
$$

All the fluctuational patterns introduced in part I fulfill this condition.

In dealing with the Fourier extension of the Laplace transform of autocorrelation functions, say $\mathbf{C}_{vv}(t)$, an even extension to negative time is adopted, i.e. $\mathbf{C}_{vv}(-t) = \mathbf{C}_{vv}(t)$, $t > 0$.

In order to generalize the theory, it is convenient to consider the functional structures of all the fluctuation patterns discussed in part I. These patterns involve white-noise perturbations (either in the form of distributional derivatives of Wiener processes as in part I, or distributional derivatives of compound counting processes as in part II) that are mutually independent of each other. Let $\boldsymbol{\xi}(t) = (\xi_1(t), \ldots, \xi_N(t))$ be the vector collecting all these processes, i.e. in the most general case of $\mathbb{DI}_{m,E}^{(3,N_d,N_i)}(\boldsymbol{\xi}^{(d)}(t),\boldsymbol{\xi}^{(i)}(t)), \boldsymbol{\xi}(t) = (\boldsymbol{\xi}^{(d)}(t),\boldsymbol{\xi}^{(i)}(t)),$ where $\boldsymbol{\xi}^{(d)}(t)$ and $\boldsymbol{\xi}^{(i)}(t)$ refer to the elementary white-noise forcings acting on the internal dissipative and fluid-inertial variables (and, due to inertial effects, also directly to $\mathbf{v}(t)$, at least as regards $\xi^{(i)}(t)$) The dimension N of the vector $\xi(t)$ is $3(N_d+N_i)$, as we consider the general three-dimensional case. Independence and white noise nature imply

$$
\langle \xi_i(t) \xi_j(t') \rangle = \delta_{ij} \, \delta(t - t') \,, \qquad i, j = 1, \dots, N \tag{18}
$$

Owing to linearity and stationarity of the dynamics associated with the fluctuational patterns, the velocity entry $v_h(t)$ can be expressed as the sum of convolutions of these impulsive stochastic contributions

$$
v_h(t) = \sum_{i=1}^{N} M_{hi}(t) * \xi_i(t), \qquad h = 1, 2, 3
$$
 (19)

where $M_{hi}(t)$ are the response functions to the elementary stochastic forcings. Transforming this equation in the Laplace domain, and setting $\hat{v}_h[\omega] = \hat{v}_h(s)|_{s=i\omega}$ for the Fourier-Laplace transform of $v_h(t)$ (and similarly for the other quantties), we have

$$
\widehat{v}_{h}[\omega] = \sum_{i=1}^{N} \widehat{M}_{hi}[\omega] \,\widehat{\xi}_{i}[\omega] \tag{20}
$$

Owing to the mutual independence of the elementary stochastic forcings eq. [\(18\)](#page-9-0), we get for the velocity power spectral density tensor $P_{hk}^{(v)}(\omega) = \langle \hat{v}_h[\omega] \hat{v}_k^*[\omega] \rangle$, $(z^*$ is the complex conjugate of the complex number z).

$$
P_{hk}^{(v)}(\omega) = \sum_{i=1}^{N} \sum_{j=1}^{N} \widehat{M}_{h,i}[\omega] \widehat{M}_{k,j}^*[\omega] P_{ij}^{(\xi)}(\omega)
$$
 (21)

where the power spectral density of the elementary stochastic forcings equal $P_{ij}^{(\xi)}(\omega) = \delta_{ij}$ by definition of independent white-noise processes. Thus, eq. [\(21\)](#page-10-0) reduces to

$$
P_{hk}^{(v)}(\omega) = \sum_{i=1}^{N} \widehat{M}_{h,i}[\omega] \widehat{M}_{k,i}^{*}[\omega]
$$
\n(22)

or in matrix form

$$
\mathbf{P}^{(v)}(\omega) = \widehat{\mathbf{M}}[\omega] \widehat{\mathbf{M}}^{*T}[\omega]
$$
\n(23)

where the superscript " T " indicates transposition.

As developed in part I, the fluctuational patterns satisfy, by definition, the Kubo fluctuation-dissipation relations of the first and second kind, extended also to the presence of fluid inertial effects. Therefore, provided that the long-term friction factor is bounded, they satisfy a-fortiori the global fluctuation-dissipation relation eq. [\(17\)](#page-9-1). Enforcing the Wiener-Khinchin theorem, the velocity power-spectral density is the Fourier transform of the velocity autocorrelation function (extended in an even way over the entire real line, as discussed above). For this reason we have

$$
\widehat{\mathbf{C}}_{vv}(0) = \frac{\mathbf{P}^{(v)}(0)}{2} \tag{24}
$$

and the global fluctuation-dissipation relation for the fluctuational patterns becomes

$$
\mathbf{D} = \widehat{\mathbf{C}}_{vv}(0) = \frac{1}{2} \widehat{\mathbf{M}}[\omega] \widehat{\mathbf{M}}^{*T}[\omega]
$$
 (25)

Next, consider a generalization of the fluctuational patterns in which each white-noise elementary stochastic forcing $\xi_i(t)$ is replaced by a correlated stationary stochastic process $q_i(t)$, and these processes are independent of each other, i.e.

$$
\langle q_i(t) q_j(t') \rangle = \delta_{ij} C_{q_i q_j}(|t - t'|), \qquad i, j = 1, \dots, N
$$
\n(26)

t, $t' \in \mathbb{R}$, where $C_{q_i q_i}(|t-t'|)$, $i = 1, ..., N$ are Lipshitz-continuous functions of their argument. More precisely $C_{q_i q_i}(t)$ are smooth for any $|t| > 0$, with the only cusp singularity at $t = 0$, where both left and right derivatives exist. The fluctuation patterns extended in this way to correlated elementary forcings will be indicated with a letter "C" preceding the symbolic code of the pattern, so that in the simplest scalar Stokes case $\mathbb{CS}_{m,E}^{(1,1,0)}(q(t))$ is the same pattern as $\mathbb{S}_{m,E}^{(1,1,0)}(\xi(t))$ with a correlated process $q(t)$ substituting $\xi(t)$.

Since the functional structure of a fluctuational pattern remains unchanged, we still have in the correlated case

$$
v_h(t) = \sum_{i=1}^{N} M_{hi}(t) * q_i(t)
$$
\n(27)

Enforcing the mutual independence of the $q_i(t)$'s, and repeating the same calculation developed above for the white-noise case, we arrive to

$$
P_{hk}^{(v)}(\omega) = \sum_{i=1}^{N} \widehat{M}_{hi}[\omega] \widehat{M}_{k,i}^{*}[\omega] P_{ii}^{(q)}[\omega]
$$
\n(28)

Since $P_{ii}^{(q)}[\omega] = \hat{C}_{q_i q_i}(0)/2$, the effective diffusivity tensor associated with the correlated pattern is given by

$$
D_{hk} = \sum_{i=1} \widehat{M}_{hi}[0] \widehat{M}_{ki}^*[0] \widehat{C}_{q_i q_i}(0) \tag{29}
$$

where $C_{q_i q_i}(0)$ is the value of the Laplace transform of the autocorrelation function $C_{q_i q_i}(t) =$ $\langle q_i(t) q_i(0) \rangle_{\text{eq}}$ of $q_i(t)$. We have the following result:

Theorem I - If the correlated stationary processes $q_i(t)$, $i = 1, \ldots, N$, are mutually independent, and are characterized by the properties

$$
\int_0^\infty C_{q_i q_i}(t) dt = \widehat{C}_{q_i q_i}(0) = \frac{1}{2}, \qquad i = 1, ..., N
$$
\n(30)

then the correlated fluctuational pattern built upon $\{q_i(t)\}_{i=1}^N$ satisfies the global fluctuation dissipation relation eq. [\(14\)](#page-8-1).

Proof - Under the condition of mutual independence, eq. [\(29\)](#page-11-0) holds. If the autocorrelation functions of the $q_i(t)$ -processes satisfy eq. [\(30\)](#page-11-1) then eqs. [\(29\)](#page-11-0) coincide with eq. [\(25\)](#page-10-1) for the original fluctuational pattern in the presence of elementary forcings that cor-responds to the global fluctuation-dissipation relation eq. [\(14\)](#page-8-1). This proves the statement. \diamond

The condition expressed by eq. [\(30\)](#page-11-1) admits a simple physical explanation. Consider a correlated process $q(t)$ satisfying eq. [\(30\)](#page-11-1) and characterized by a finite correlation rate ν , and the diffusive dynamics generated by it,

$$
\frac{dx_q(t)}{dt} = q(t) \tag{31}
$$

compared with the diffusive dynamics generated by the distributional derivative $\xi(t)$ of a Wiener process

$$
\frac{dx(t)}{dt} = \xi(t) \tag{32}
$$

starting from $x_q(0) = x(0) = 0$. From eq. [\(30\)](#page-11-1) it follows that in the long-term $\langle x_q^2(t) \rangle \simeq$ $2 D_q t$, where the diffusivity is $D_q = \widehat{C}_{qq}(0) = 1/2$, and coincides with that of eq. [\(32\)](#page-12-0) driven by a Wiener process. Moreover, in the long term the central limit theorem applies, and $x_q(t)$ approaches the statistical properties of $x(t)$ defined by eq. [\(32\)](#page-12-0). This justifies why the substitution of distributional derivatives of Wiener processes with correlated processes $q_i(t)$ within the structure of a fluctuational pattern does not modify the global fluctuation-dissipation relation, as the two dynamics possess in the long term the same statistical properties.

To complete the analysis, the behavior of the velocity autocorrelation function at $t = 0$ should be investigated, as this corresponds to the equilibrium values of the second-order velocity moments. To simplify the analysis consider the scalar case for a velocity $v(t)$ as the extension to the vectorial problem is straightforward. Let $C_{vv}(t)$ be the velocity autocorrelation function for any fluctuational pattern driven by white-noise elementary forcings, and $C_{vv,corr}(t)$ the same function for the corresponding correlated fluctuational pattern, i.e. for the same hydrodynamic conditions but with the elementary forcings expresses as correlated processes satisfying the conditions of Theorem I. We have

$$
C_{vv}(t) = \langle v^2 \rangle_{\text{eq}} \phi(t) , \qquad \langle v^2 \rangle_{\text{eq}} = \frac{k_B T}{m}
$$
 (33)

and

$$
C_{\rm vv,corr}(t) = \langle v^2 \rangle_{\rm eq,corr} \phi_{\rm corr}(t)
$$
\n(34)

where $\phi(t)$ and $\phi_{\text{corr}}(t)$ are the normalized autocorrelation functions, $\phi(0) = \phi_{\text{corr}}(0) = 1$. Since, by Theorem I the integrals of $C_{vv}(t)$ and $C_{vv,corr}(t)$ over $t \in [0, \infty)$ coincide, we have

$$
\langle v^2 \rangle_{\text{eq,corr}} = \frac{k_B T}{m} \frac{\int_0^\infty \phi(t) \, dt}{\int_0^\infty \phi_{\text{corr}}(t) \, dt} \tag{35}
$$

For one and the same structure of the fluctuational pattern, the normalized velocity autocorrelation function should be greater in the presence of correlated forcings than in the case of δ -correlated ones, i.e. $\phi_{\text{corr}}(t) > \phi(t)$, and thus from eq. [\(35\)](#page-12-1) we have

$$
\langle v^2 \rangle_{\text{eq,corr}} \le \frac{k_B T}{m} \tag{36}
$$

A proof of the latter statement is provided below.

Theorem II - For any correlated fluctuational pattern built upon the correlated processes $\{q_i(t)\}_{i=1}^N$, independent of each other and satisfying the conditions of Theorem I, eq. [\(36\)](#page-13-0) applies.

Proof - Consider a generic velocity entry $v_h(t)$. In the case of a Wiener-driven fluctuational patter we have from eqs. $(18)-(19)$ $(18)-(19)$

$$
\langle v_h(t) \rangle = \sum_{i=1}^N \sum_{j=1}^N \int_0^t M_{hi}(\tau) d\tau \int_0^t M_{hj}(\tau') \langle \xi_i(t-\tau) \xi_j(t-\tau') \rangle d\tau'
$$

$$
= \sum_{i=1}^N \int_0^t M_{hi}^2(\tau) d\tau
$$
(37)

the equilibrium value corresponds to the limit for $t \to \infty$, and fulfills eq. [\(2\)](#page-1-1),

$$
\langle v_h^2 \rangle_{\text{eq}} = \sum_{i=1}^N \int_0^\infty M_{hi}^2(\tau) \, d\tau = \frac{k_B \, T}{m} \tag{38}
$$

Next, consider its correlated counterpart substituting $\{\xi_i(t)\}_{i=1}^N$ with independent correlated processes $\{q_i(t)\}_{i=1}^N$. Enforcing their mutual independence we have

$$
\langle v_h^2(t) \rangle_{\text{corr}} = \sum_{i=1}^N \int_0^t M_{hi}(\tau) d\tau \int_0^t M_{hi}(\tau') C_{q_i q_i}(|\tau - \tau'|) d\tau'
$$

=
$$
2 \sum_{i=1}^N \int_0^t M_{hi}(\tau') d\tau' \int_{\tau'}^t M_{hi}(\tau) C_{q_i q_i}(\tau - \tau') d\tau
$$
(39)

In the limit for $t \to \infty$ the equilibrium value is achieved,

$$
\langle v_h^2(t) \rangle_{\text{eq,corr}} = 2 \sum_{i=1}^N \int_0^\infty M_{hi}(\tau') d\tau' \int_{\tau'}^\infty M_{hi}(\tau) C_{q_i q_i}(\tau - \tau') d\tau
$$

$$
= 2 \sum_{i=1}^N \int_0^\infty M_{hi}(\tau') d\tau' \int_0^\infty M_{hi}(\theta + \tau') C_{q_i q_i}(\theta) d\theta
$$

$$
= 2 \sum_{i=1}^N \int_0^\infty C_{q_i q_i}(\theta) d\theta \int_0^\infty M_{hi}(\tau') M_{hi}(\theta + \tau') d\tau'
$$
(40)

In the last integral, the Schwartz inequality can be applied, namely

$$
\left| \int_0^\infty M_{hi}(\tau') M_{hi}(\theta + \tau') d\tau' \right| \le \int_0^\infty M_{h,i}^2(\tau') d\tau' \tag{41}
$$

Thus,

$$
\langle v_h^2 \rangle_{\text{eq,corr}} \leq \sum_{i=1}^N \left| 2 \int_0^\infty C_{q_i q_i}(\theta) \, d\theta \, \middle| \, \left| \int_0^\infty M_{hi}^2(\tau') \, d\tau' \right| \right.
$$

$$
\leq \sum_{i=1}^N \int_0^\infty M_{hi}^2(\tau') \, d\tau' = \frac{k_B T}{m} \tag{42}
$$

where we have used eq. (30) and eq. (38) . \diamond .

The proof of Theorem II is grounded on the Schwarz inequality eq. [\(41\)](#page-14-1),in which the equal sign applies only in the limit case $q_i(t)$ tending towards $\xi_i(t)$. This is interesting as it indicates that a non-vanishing added mass, no matter how small, characterized any correlated fluctuational patterns. If the elementary stochastic forcings in a fluctuational pattern possess some characteristic correlation time, either the global fluctuation-dissipation relation (as in the present formulation of the theory) or the equipartition result can be fulfilled, but never both at the same time. The only case in which these two properties are simultaneously satisfied occurs for the classical fluctuational patterns discussed in part I and II driven by white-noise processes.

Eq. [\(36\)](#page-13-0) indicates that an added mass arises as an emergent property of correlated fluctuations defining the elementary stochastic forcing of a correlated fluctuational pattern (the C-patterns).

In a physical perspective, this results applies on equal footing to incompressible and compressible flows, as it is not a property of the hydromechanic kernels, but a consequence of the regular nature of the hydrodynamic fluctuations. A connection between this fluctuational property and the fluid-particle hydromechanics is addressed in Section [VI](#page-27-0) via the fluctuationinertia relation. Some prototypical examples of application of Theorem II are discussed in the next Section.

IV. EXAMPLES

This Section discusses several prototypical examples, considering for notational simplicity the scalar case, i.e. a scalar velocity $v(t)$, corresponding to the isotropic dynamics of a spherical object.

Before analyzing some case studies, let us consider how to construct admissible correlated processes, i.e. stochastic processes $q(t)$ fulfilling eq. [\(30\)](#page-11-1). There are two simple ways of defining them: either through successive low-pass filterings of distributional derivatives of Wiener processes or by considering discontinuous correlated processes, such as the Poisson-Kac processes [\[34](#page-37-2)[–36](#page-37-3)].

Consider the first case. The simplest example is the output $q_{\nu}(t)$ of a low-pass filtering of $\xi(t)$, i.e. of a distributional derivative of a Wiener process through a filter with characteristic time ν^{-1} under the condition that, for $\nu \to \infty$ $q_{\nu}(t)$, converges stochastically to $\xi(t)$. For notational simplicity we set $q(t) = q_{\nu}(t)$. The process is defined by the dynamic equation

$$
\frac{dq(t)}{dt} = -\nu \left(q(t) - \xi(t) \right) \tag{43}
$$

which is characterized by the equilibrium (long-term) properties: $\langle q^2 \rangle_{\text{eq}} = \nu/2$ and

$$
C_{qq}(t) = \frac{\nu e^{-\nu t}}{2} \tag{44}
$$

where $C_{qq}(t) = \langle q(t) q(0) \rangle_{\text{eq}}$ is its autocorrelation function. Therefore, $\hat{C}_{qq}(0) = 1/2$, consistently with the condition expressed by Theorem I.

Similarly, this mechanism can be generalized to a cascade of low-pass filters of different characteristic times. For example, in the case of two filters we have for $q(t)$

$$
\frac{dq(t)}{dt} = -\nu(q(t) - q_1(t))
$$
\n
$$
\frac{dq_1(t)}{dt} = -\nu_1(q_1(t) - \xi(t))
$$
\n(45)

and it is characterized by the steady-state moments

$$
m_{q_1q_1}^* = \langle q_1^2 \rangle_{\text{eq}} = \frac{\nu_1}{2}, \qquad m_{qq_1}^* = \langle q \, q_1 \rangle_{\text{eq}} = m_{qq}^* = \langle q^2 \rangle_{\text{eq}} = \frac{\nu \, \nu_1}{2(\nu + \nu_1)} \tag{46}
$$

Its autocorrelation function $C_{qq}(t)$ is the solution of the equations

$$
\frac{dC_{qq}(t)}{dt} = -\nu(C_{qq}(t) - C_{q_1q}(t))
$$
\n
$$
\frac{dC_{q_1q}(t)}{dt} = -\nu_1 C_{q_1q}(t)
$$
\n(47)

equipped with the initial conditions $C_{qq}(0) = m_{qq}^*$, $C_{q_1q}(0) = m_{qq_1}^*$, the solution of which reads in the Laplace domain

$$
\widehat{C}_{qq}(s) = \frac{m_{qq}^*}{s + \nu} + \frac{\nu m_{qq_1}^*}{(s + \nu)(s + \nu_1)}\tag{48}
$$

Substituting the values of the stationary moments eq. [\(46\)](#page-15-0) into eq. [\(48\)](#page-15-1), $\hat{C}_{qq}(0) = 1/2$ follows, confirming its admissibility.

Alternatively, one may define $q(t)$ as a Poisson-Kac process or its generalizations [\[34](#page-37-2)[–36\]](#page-37-3). The simplest example is provided by the process

$$
q(t) = c(-1)^{\chi(t,\nu)}\tag{49}
$$

where $\chi(t,\nu)$ is a Poisson process with rate ν , and c a constant to be determined. Since, for $t \geq 0$, we have

$$
C_{qq}(t) = c^2 \langle (-1)^{\chi(t,\nu)} (-1)^{\chi(0,\nu)} \rangle = c^2 e^{-2\nu t} \Rightarrow \int_0^\infty C_{qq}(t) dt = \frac{c^2}{2\nu} \tag{50}
$$

the admissibility of this process, i.e. the condition of Theorem I, implies $c = \sqrt{\nu}$.

Next, consider the application of correlated forcings to the prototypical fluctuational patterns. The process $q(t)$ defined by eq. [\(43\)](#page-15-2) will be explicitly considered.

$\mathbf A.\quad \mathbb{CS}^{(1,1,0)}_{m,E}(q(t))$

For the scalar Stokesian pattern we have $a = \sqrt{k_B T \eta}$, and the dynamics of $\mathbb{CS}_{m,R}^{(1,1,0)}(q(t))$ is thus defined by

$$
m\frac{dv(t)}{dt} = -\eta v(t) + \sqrt{2} a q(t)
$$

$$
\frac{dq(t)}{dt} = -\nu(q(t) - \xi(t))
$$
(51)

Setting $\alpha = \eta/m$, $\beta = \sqrt{2}a/m$, the second-order moments $m_{vv}(t) = \langle v^2(t) \rangle$, $m_{vv}(t) =$ $\langle v(t)q(t)\rangle$, $m_{qq}(t) = \langle q^2(t)\rangle$ evolve according to the dynamics

$$
\frac{dm_{vv}(t)}{dt} = -2 \alpha m_{vv}(t) + 2\beta m_{vq}(t)
$$
\n
$$
\frac{dm_{vq}(t)}{dt} = -\alpha m_{vq}(t) + \beta m_{qq}(t) - \nu m_{vq}(t)
$$
\n
$$
\frac{dm_{qq}(t)}{dt} = -2 \nu m_{qq}(t) + \nu^2
$$
\n(52)

providing the steady-state values

$$
m_{vv}^* = \frac{\beta^2 \nu}{2 \alpha (\alpha + \nu)}, \qquad m_{vq}^* = \frac{\beta \nu}{2 (\alpha + \nu)}, \quad m_{qq}^* = \frac{\nu}{2}
$$
 (53)

The squared velocity variance at equilibrium is given by

$$
\langle v^2 \rangle_{\text{eq}} = \frac{k_B T}{m \left(1 + \tau_c^*\right)}\tag{54}
$$

where τ_c^* is the dimensionless correlation time equal to the ratio of the correlation time $1/\nu$ to the dissipation time m/η ,

$$
\tau_c^* = \frac{1}{\nu^*} = \frac{\eta}{m\,\nu} \tag{55}
$$

where $\nu^* = \nu/\alpha$. It follows from eq. [\(54\)](#page-17-0) that the nondimensional added mass of this pattern is given by

$$
\frac{m_a}{m} = \tau_c^* \tag{56}
$$

As regards the velocity autocorrelation function $C_{vv}(t)$, we have

$$
\frac{dC_{vv}(t)}{dt} = -\alpha C_{vv}(t) + \beta C_{qv}(t)
$$
\n
$$
\frac{dC_{qv}(t)}{dt} = -\mu C_{qv}(t)
$$
\n(57)

equipped with the initial conditions $C_{vv}(0) = m^*_{vv}, C_{qv}(0) = m^*_{vq}$, The Laplace trasform of $C_{vv}(t)$, solution of eq. [\(57\)](#page-17-1), is

$$
\widehat{C}_{vv}(s) = \frac{m_{vv}^*}{s+\alpha} + \frac{\beta m_{vq}^*}{(s+\alpha)(s+\nu)}\tag{58}
$$

from which, substituting the equilibrium values for the moments eq. [\(53\)](#page-16-0), the Stokes-Einstein relation $\widehat{C}_{vv}(0) = k_B T/m \eta$ is recovered. The explicit expression for the nondimensional velocity autocorrelation $C_{\text{nd}}(t') = m C_{vv}(t)/k_B T|_{t=t'/\alpha}$ reads for $t > 0$

$$
C_{\rm nd}(t') = \begin{cases} \frac{1}{1+1/\nu^*} e^{-t'} + \frac{\nu^*}{1-\nu^{*2}} \left[e^{-\nu^* t'} - e^{-t'} \right] & \nu^* \neq 1\\ \frac{1}{1+1/\nu^*} e^{-t'} + \frac{\nu^*}{1+\nu^*} t' e^{-t'} & \nu^* = 1 \end{cases}
$$
(59)

where $t' = \alpha t$ is the nondimensional time rescaled with respect to the dissipation time. As a realistic situation, consider the case where the fluctational added mass coincides with the hydrodynamic added mass for an object possessing the same density of the solvent fluid. In this case $m_a/m = 1/2$ $m_a/m = 1/2$ $m_a/m = 1/2$, and this corresponds to $\nu^* = 2$. Figure 1 depicts the comparison of the normalized velocity autocorrelation functions for the patterns $\mathbb{S}_{m,E}^{(1,1,0)}(\xi(t))$ and $\mathbb{CS}_{m,E}^{(1,1,0)}(q(t))$ at $\nu^* = 2$. A slight difference can be observed between the δ-correlated and the correlated patterns.

FIG. 1. Normalized velocity autocorrelation functions $C_{\text{nd}}(t')$ vs the nondimensional time t' . Line (a) refers to $\mathbb{S}_{m,E}^{(1,1,0)}(\xi(t))$, line (b) to $\mathbb{CS}_{m,E}^{(1,1,0)}(q(t))$ at a value of $\nu^* = 2$.

As a final observation, consider the Langevin condition. In this model, $R(t)$ is proportional to $q(t)$, and we have $(t \geq 0)$

$$
\langle R(t) v(0) \rangle_{\text{eq}} = \sqrt{2 k_B T \eta} \langle q(t) v(0) \rangle_{\text{eq}} = \frac{k_B T \nu}{1 + \nu m / \eta} e^{-\nu t}
$$
(60)

Thus, for any finite ν , the Langevin condition is not satisfied, as intuitively expected. In the limit for $\nu \to \infty$, i.e. when $q(t)$ approaches $\xi(t)$, one gets from eq. [\(60\)](#page-18-1)

$$
\lim_{\nu \to \infty} \langle R(t) \, v(0) \rangle_{\text{eq}} = \begin{cases} k_B \, T \, \eta/m & t = 0 \\ 0 & t > 0 \end{cases} \tag{61}
$$

The correlation between $R(t)$ and $v(0)$ is vanishing for any $t > 0$, and attains at $t = 0$ a finite value. So for any practical purpose the limit eq. (61) converges in the L²-norm to the Langevin condition, i.e. to zero.

B. $CD_{m,E}^{(1,1,0)}(q(t))$

The next pattern in order of complexity is the correlated dissipative pattern $\mathbb{CD}_{m,E}^{(1,1,0)}(q(t))$ associated with the pattern $\mathbb{D}_{m,E}^{(1,1,0)}(\xi(t))$, in the presence of a single exponential memory term. This correlated pattern is defined by the dynamics

$$
\begin{aligned}\n\frac{dv(t)}{dt} - \alpha \theta(t) \\
\frac{d\theta(t)}{dt} &= -\lambda \theta(t) + \lambda v(t) + \sqrt{2} b \lambda q(t) \\
\frac{dq(t)}{dt} &= -\nu(q(t) - \xi(t))\n\end{aligned} \tag{62}
$$

where $b = \sqrt{k_B T / \eta}$. The steady-state moments are given by $m_{v\theta}^* = 0$, $m_{qq}^* = \nu/2$,

$$
m_{\theta q}^* = \frac{b\,\lambda\,\nu}{\sqrt{2}\left(\lambda + \nu + \alpha\,\lambda/\nu\right)}, \qquad m_{\nu q}^* = -\frac{\alpha\,b\,\lambda}{\sqrt{2}\left(\lambda + \nu + \alpha\,\lambda/\nu\right)}, \quad m_{\theta \theta}^* = \frac{b^2\,\lambda\,\nu}{\left(\lambda + \nu + \alpha\,\lambda/\nu\right)}\tag{63}
$$

and

$$
m_{vv}^* = \langle v^2 \rangle_{\text{eq}} = \frac{b^2 \alpha \left(\lambda + \nu \right)}{\left(\lambda + \nu + \alpha \lambda / \nu \right)} = \frac{k_B T}{m} \left(1 + \frac{\alpha \lambda}{\nu \left(\lambda + \nu \right)} \right)^{-1} \tag{64}
$$

and this corresponds to a fluctuational added mass

$$
\frac{m_a}{m} = \frac{\eta \lambda}{m \nu (\lambda + \nu)}\tag{65}
$$

Also for this pattern, consider the case $m_a/m = 1/2$. This corresponds to a dimensionless correlation rate $\nu^* = \nu/\alpha$, depending on the dimensionless relaxation rate $\lambda^* = \lambda/\alpha$, given by

$$
\nu^* = \frac{\lambda^*}{2} \left[\sqrt{1 + \frac{8}{\lambda^*}} - 1 \right] \tag{66}
$$

Figure [2](#page-20-0) depicts the behaviour of ν^* , providing an added mass equal half of the particle mass, as a function of λ^* .

For very large λ^* , one recovers the Stokesian result discussed above $\nu^* \simeq 2$. For $\lambda^* \ll 1$, $\nu^* \sim \sqrt{2\lambda^*}$. This result is not surprising. In a viscoelastic fluids, the correlation rate ν to achieve a given added mass depends on the viscoelastic relaxation rate λ , and if $\lambda \ll \alpha$, $\nu \simeq \sqrt{2 \lambda \alpha}$, i.e. it is proportional to the geometric mean of the dissipation and relaxation rates.

Next, consider the velocity autocorrrelation function. Setting $C_{nd}(t') = m C_{vv}(t')/k_B T$ where $t' = \alpha t$ is the nondimensional time, and $\tilde{C}_{\theta v}(t') = C_{\theta v}(t')/b^2 \alpha$, $\tilde{C}_{qv}(t') = \sqrt{2} C_{qv}(t')/b \alpha$, the nondimensional velocity autocorrelation function is the solution of this system of equa-

FIG. 2. ν^* vs λ^* (line a) for the $\mathbb{CD}_{m,E}^{(1,1,0)}(q(t))$ pattern providing $m_a/m = 1/2$. Line (b) refers to the Stokesian limit $\nu^* = 2$, line (c) to the low-relaxation rate limit $\nu^* = \sqrt{2\lambda^*}$.

tions

$$
\frac{dC_{nd}(t')}{dt'} = -\widetilde{C}_{\theta v}(t')
$$
\n
$$
\frac{d\widetilde{C}_{\theta v}(t')}{dt'} = -\lambda^* \widetilde{C}_{\theta v}(t') + \lambda^* C_{nd}(t') + \lambda^* \widetilde{C}_{qv}(t')
$$
\n
$$
\frac{d\widetilde{C}_{qv}(t')}{dt'} = -\nu^* \widetilde{C}_{qv}(t')
$$
\n(67)

equipped with the initial conditions $C_{nd}(0) = (1 + \lambda^*/\nu^*(\lambda^* + \nu^*))^{-1}$, $C_{\theta\nu}(0) = 0$, $C_{qv}(0) =$ $-\lambda^*/(\lambda^* + \nu^* + \lambda^*/\nu^*)$. Figure [3](#page-21-0) depicts the behaviour of the normalized velocity autocorrelation function at $\lambda^* = 1$, for values of ν^* providing $m_a/m = 1/2$ and $m_a/m = 1/6$ respectively. λ^* .

To conclude, consider the Langevin condition. Since $R(t) = -m\sqrt{2} \alpha b \lambda e^{-\lambda t} * q(t)$ we have

$$
\langle R(t) v(0) \rangle_{\text{eq}} = -m \sqrt{2} \alpha b \lambda \left(e^{-\lambda t} * C_{qv}(t) \right) = -m \sqrt{2} \alpha b \lambda m_{vq} \left(e^{-\lambda t} * e^{-\nu t} \right) \tag{68}
$$

Expressing the Langevin term with respect to the dimensionless time t' , eq. [\(68\)](#page-20-1) becomes

$$
\langle R(t')\, v(0)\rangle_{\text{eq}} = \frac{k_B\,T\,\eta}{m} \frac{(\lambda^*)^2}{\lambda^* + \nu^* + \lambda^*/\nu^*} \begin{cases} \frac{e^{-\nu^*t'} - e^{-\lambda^*t'}}{\lambda^* - \nu^*} & \nu \neq \lambda\\ t' \, e^{-\lambda^*t'} & \nu = \lambda \end{cases} \tag{69}
$$

In this case, $\langle R(0) v(0) \rangle_{\text{eq}} = 0$, while $\langle R(t') v(0) \rangle_{\text{eq}} > 0$ for $t' > 0$.

FIG. 3. $C_{nd}(t')$ vs t' for the CD-patterns at $\lambda^* = 1$. Line (a) refers to the corresponding D-pattern. Line (b) refers to $\nu^* = 1$ corresponding to $m_a/m = 1/2$, line (c) to $\nu^* = 2$ corresponding to $m/m_a = 1/6.$

C. $\mathbb{C} \mathbb{I}_{m,E}^{(1,1,1)}(q(t),q_1(t))$

The next prototypical model of increasing hydrodynamic complexity involves the occurrence of a fluid inertial contribution superimposed to a Stokesian dissipative term, i.e. $\mathbb{I}_{m,E}^{(1,1,1)}(\xi(t),\xi_1(t))$. In the presence of correlated forcing it becomes $\mathbb{CI}_{m,E}^{(1,1,1)}(q(t),q_1(t))$ where $q(t)$ and $q_1(t)$ are the low-pass filtering of the white-noise independent processes $\xi(t)$ and $\xi_1(t)$. The dynamics of this correlated pattern reads

$$
\frac{dv(t)}{dt} = -(\gamma + \alpha) v(t) + \gamma z(t) + \sqrt{2}a q(t) + \sqrt{2} d q_1(t)
$$
\n
$$
\frac{dz(t)}{dt} = -\mu z(t) + \mu v(t) + \sqrt{2} c q_1(t)
$$
\n
$$
\frac{dq(t)}{dt} = -\nu(q(t) - \xi(t))
$$
\n
$$
\frac{dq_1(t)}{dt} = -\nu_1(q(t) - \xi_1(t))
$$
\n(70)

where $\alpha = \eta/m$, $\gamma = g/m$ and (see part I)

$$
a = \delta = \frac{\sqrt{k_B T \eta}}{m}, \quad d = \sqrt{\frac{\gamma}{\alpha}} \delta, \quad c = -\frac{\delta}{\sqrt{\alpha \gamma}}
$$
(71)

From the analysis of the moments at steady state, one immediately gets $m_{qq}^* = \nu/2$, $m_{q_1q_1}^* =$ $\nu_1/2, m_{qq_1}^* = 0$. The remaining seven moments satisfy a linear system, the solution of which

FIG. 4. Normalized squared velocity variance $\langle v^2 \rangle_{\text{eq}}^*$ vs ν^* at $\mu = 1, \gamma = 1$, for different values of the rate ν_1^* . Line (a) refers to $\nu_1 = \nu^*$. The remaining lines indicated by an arrow refer to $\nu = 1$ for increasing values of $\nu_1^* = 1$, 10, 100. The horizontal line depicts the limit behaviour $\langle v^2 \rangle_{\text{eq}}^* = 1$.

provides the value for $m_{vv}^* = \langle v^2 \rangle_{\text{eq}}$, see the Appendix. For finite values of ν and ν_1 , we have $\langle v^2 \rangle_{\text{eq}} < k_B T/m$, and

$$
\lim_{\nu \to \infty} \lim_{\nu_1 \to \infty} \frac{m \langle v^2 \rangle_{\text{eq}}}{k_B T} = 1
$$
\n(72)

consistently with the limit of $q(t)$ and $q_1(t)$ tending to white-noise processes. From the analysis of the velocity autocorrelation function it follows that the global fluctuation-dissipation relation is satisfied for any choice of ν and ν_1 , as expected from Theorem I. Figure [4](#page-22-1) depicts the behaviour of the normalized squared velocity variance $\langle v^2 \rangle_{\text{eq}}^* = m \langle v^2 \rangle_{\text{eq}} / k_B T$ obtained from $\mathbb{C}[\Pi_{m,E}^{(1,1,1)}(q(t), q_1(t))]$ patterns as a function of $\nu^* = \nu/\alpha$, for different settings of $\nu_1^* = \nu_1/\alpha$.

V. REGULARITY OF THE VELOCITY FLUCTUATIONS

There is another general qualitative property that arises as a consequence of fluid-inertial interactions, related to the regularity of velocity fluctuations, in the meaning of their Hölder continuity [\[37](#page-37-4)]. Consider a single realization of the stochastic velocity process $v(t)$ in some time interval $I = [a, b], a > 0, b > a$. The process is Hölder continuous in I with exponent H if, for any generic realization $v(t)$, and for any $t_1, t_2 \in I$, $t_2 > t_1$, , there exist constants $C_1, C_2 > 0, C_2 > C_1$ such that

$$
C_1 |t_2 - t_1|^H \le |v(t_2) - v(t_1)| \le C_2 |t_2 - t_1|^H \tag{73}
$$

The exponent H is referred to as the Hölder exponent. If $H = 1$, $v(t)$ is Lipshitz continuous (something more than continuity, something less than differentiability). Values of H less than 1 implies that the function $v(t)$, (regarding a realization of the process as a function of time t), displays a singular behaviour almost everywhere in I , in the meaning that it cannot be defined a tangent vector at almost any $t \in I$. This follows readily for the simple observation that, setting $t_2 = t_1 + \Delta t$, $\Delta t > 0$, eq. [\(73\)](#page-23-0) implies

$$
\frac{C_1}{\Delta t^{1-H}} \le \frac{|v(t_1 + \Delta t) - v(t_1)|}{\Delta t} \le \frac{C_2}{\Delta t^{1-H}}
$$
\n(74)

Therefore, in the limit for $\Delta t \to 0$, the absolute value of the derivative of $v(t)$ at t_1 diverges. In the case of a realization of a Wiener process $w(t)$, it follows from its definition [\[38\]](#page-37-5) that $w(t_2) - w(t_1) = \sqrt{t_2 - t_1} r$ where $r \in \mathcal{N}(0, 1)$ is a random variable normally distributed, and thus $H = 1/2$ [\[39\]](#page-37-6).

An Hölder exponent less than 1 is the manifestation of the fractal nature of ageneric realization $v(t)$. More precisely, the fractal dimension d_v of $v(t)$, characterized by the Hölder exponent H , is given by [\[37\]](#page-37-4)

$$
d_v = 2 - H \tag{75}
$$

For a Wiener process, its fractal dimension equals $3/2$. Eq. [\(75\)](#page-23-1) implies that the estimate of H can be conveniently obtained by considering the scaling of the length $L(\Delta t)$ of $v(t)$ in the interval I as a function of the temporal resolution Δt , since

$$
L(\Delta t) \sim \Delta t^{1-H} \tag{76}
$$

The length-scale analysis associated with eq. [\(76\)](#page-23-2) is experimentally feasible and computationally very simple as it requires solely a single, statistically representative, realization of the particle velocity, i.e. a single time series.

Next, consider the different hydromechanic mechanisms and their influence as regards the regularity properties of the particle velocity $v(t)$. The structure of the different fluctuational patterns introduced in part I as regards the action of the stochastic forcings is particularly useful in this analysis. Without loss of generality we consider the onedimensional scalar case (spherical particle of radius R_p) and the nondimensional formulation adopted in part I,

FIG. 5. Length-resolution analysis, $L(\Delta t)$ vs Δt , for a realization of the $\mathbb{S}_{1,1}^{(1,1,0)}$ $_{1,1}^{(1,1,0)}(\xi(t))$ pattern in the presence of a harmonic potential with $k = 0.2$. Symbols (\bullet) are the results of the stochastic simulations, the solid line represents the scaling $L(\Delta t) \sim \Delta t^{-1/2}$.

rescaling time with respect to the characteristic dissipation time and velocity so that $\langle v^2 \rangle_{\text{eq}} =$ 1. In order to consider a model closely connected with the high-resolution experiments [\[5](#page-35-3)[–8\]](#page-35-4), we include also the influence of a harmonic potential ensuring in the experiments particle localization due to the action of an optical trap.

To begin with, consider the Stokes approximation, i.e. the pattern $\mathbb{S}^{(1,1,0)}_{1,1}$ $_{1,1}^{(1,1,0)}(\xi(t))$, the dynamics of which, in the presence of a quadratic potential, reads

$$
\dot{x}(t) = v(t) \tag{77}
$$

$$
\dot{v}(t) = -v(t) - k x(t) + \sqrt{2} \xi(t) \tag{78}
$$

where $\dot{x}(t) = dx(t)/dt$ and similarly for the other variables, and k is the nondimensional spring constant of the trap. It is a typical feature of the Stokesian pattern that $\dot{v}(t)$ is proportional to the distributional derivative $\xi(t) = dw(t)/dt$ of a Wiener process $w(t)$ and thus $v(t) \sim w(t)$. Consequently, the regularity properties of $v(t)$ stemming from $\mathbb{S}^{(1,1,0)}_{1,1}$ $_{1,1}^{(1,1,0)}(\xi(t))$ are those of a Wiener process, implying the fractal nature of the velocity realizations with a Hölder exponent $H = 1/2$. Figure [5](#page-24-0) depicts the scaling of the length $L(\Delta t)$ vs the temporal resolution Δt for a velocity realization of this pattern.

The subsequent pattern in order of hydrodynamic complexity is $\mathbb{D}_{1,1}^{(1,1,0)}$ $1,1^{(1,1,0)}(\xi^{(d)}(t))$ characterized by the presence of a dissipative memory dynamics. We consider for simplicity the case

FIG. 6. Length-resolution analysis, $L(\Delta t)$ vs Δt , for a realization of the $\mathbb{D}_{1,1}^{(1,1,0)}$ $_{1,1}^{(1,1,0)}(\xi^{(d)}(t))$ patterns in the presence of a harmonic potential with $k = 0.2$ at different values of λ . Symbols are the results of the stochastic simulations: (\Box) and line (a): $\lambda = 1$, (\circ) and line (b): $\lambda = 10$, (\bullet) and line (c): $\lambda = 100$. Line (d) represents the Wiener scaling $L(\Delta t) \sim \Delta t^{-1/2}$.

of a single mode, but the analysis applies to a generic superposition of dissipative modes. The equations of motion of this pattern are

$$
\begin{aligned}\n\dot{x}(t) &= v(t) \\
\dot{v}(t) &= -\lambda \theta(t) - k \, x(t) \\
\dot{\theta}(t) &= -\lambda \theta(t) + v(t) + \sqrt{2} \, \xi^{(d)}(t)\n\end{aligned} \tag{79}
$$

where $\xi^{(d)}(t) = dw^{(d)}(t)/dt$, and $w^{(d)}(t)$ is a Wiener process. The characteristic structure of the D-patterns is that the Wiener forcings, in the present case $\xi^{(d)}(t)$ do not act directly on the velocity dynamics, but solely on the auxiliary θ -variables. Since $\theta(t)$ is continuous and almost everywhere bounded with probabiliy 1, (as its equilibrium statistics is Gaussian, see part II), the velocity $v(t)$ is a Hölder continuous function with exponent $H = 1$ almost everwhere with probability 1.

This is indeed a generic property of dissipative memory dynamics with finite relaxation rates λ , in the absence of fluid-inertial effects. In other words, the presence of a linear viscoelastic behaviour not only solves the paradox of the infinite speed of propagation of the shear stresses (subsumed by the Newtonian constitutive equations), but also regularizes the local structure of the velocity fluctuations of a colloidal particle immersed in the fluid. Figure

[6](#page-25-0) depicts the scaling of the length $L(\Delta t)$ vs Δt for realizations of this pattern at different values of λ . The value $\lambda = 1$ corresponds to a relaxation time equal to the dissipation time $t_{\text{diss}},$ and $\lambda = 10$, 100 correspond to short relaxation time, a tenth and a hundredth smaller than t_{diss} , respectively. But even in this rather extreme cases, the saturation towards a constant value of $L(\Delta t)$ is clearly evident starting from $\Delta t \sim 1/\lambda$ and reducing further Δt .

To conclude, consider the case of $\mathbb{DI}_{1,1}^{(1,1,1)}(\xi^{(d)}(t),\xi^{(i)}(t))$, namely the coupling of fluid inertial effects with a dissipative memory dynamics. A single mode for each hydrodynamic mechanism is considered. For this pattern (see part I), the dynamic equations read

$$
\begin{aligned}\n\dot{x}(t) &= v(t) \\
\dot{v}(t) &= -\lambda \theta(t) - \gamma v(t) + \gamma \mu \, z(t) - k \, x(t) + \sqrt{2 \gamma} \, \xi^{(i)}(t) \\
\dot{\theta}(t) &= -\lambda \theta(t) + v(t) + \sqrt{2} \, \xi^{(d)}(t) \\
\dot{z}(t) &= -\mu \, z(t) + v(t) - \sqrt{\frac{2}{\gamma}} \, \xi^{(i)}(t)\n\end{aligned} \tag{80}
$$

with $\xi^{(\beta)}(t) = dw^{(\beta)}(t)/dt$, $\beta = d, i$, where $w^{(d)}(t)$ and $w^{(i)}(t)$ are Wiener processes independent of each other.

As can be observed from the structure of eqs. [\(80\)](#page-26-0), and even more clearly from the blockdiagram of the stochastic couplings of this pattern depicted in part I, the presence of fluid inertia determines a direct connection (proportionality) between the particle acceleration $\dot{v}(t)$ and $\xi^{(i)}(t)$. Consequently, $v(t) \sim w^{(i)}(t)$, from which it follows that the regularity properties of the particle velocity should be those of a Wiener process. This implies that in DI-patterns we would expect a fractal velocity signal characterized by a Hölder exponent $H = 1/2$, as in the case of the Stokesian $\mathbb{S}^{(1,1,0)}_{1,1}$ $1,1,1,0)$ ($\xi(t)$) pattern, indepedently of the presence of a dissipative memory dynamics. This result is generic as regards the number of modes considered, and it is valid a fortiori if we replace the dissipative memory dynamics with an instantaneous Stokesian friction. This result is indeed a direct consequence of the presence of an impulsive contribution in the autocorrelation function of the stochastic force $R(t)$ associated with the effects of fluid-inertial interactions, eqs. (63) and (64) in part I.

We can therefore conclude that, independently of the fluid-dynamic model adopted in determining the fluid inertial contribution, stemming either from an incompressible model, or including the effects of compressibility, as in the Chow and Hermans analysis, we would observe from the time-series of particle velocities in DI-patterns an almost everywhere singular behaviour with a fractal dimension $d_v = 3/2$. The DI-pattern is the most general fluctuation-dissipation scheme deriving from the assumption of the validity of the Langevin condition. Its correlated generalization, namely the pattern $\mathbb{CDI}_{m,E}^{(n,N_d,N_i)}(\xi^{(d)}(t),\xi^{(i)}(t))$ removes this singularity due to the presence of a correlated stochastic forcing $q^{(i)}(t)$ in place of $\xi^{(i)}(t)$, acting exactly in the same way as a viscoelastic memory term, and restores a regular behaviour of the particle velocity characterized by the Hölder exponent $H = 1$.

Figure [7](#page-28-0) panels (A) and (B) depict the fractal length-resolution analysis of velocity realizations arising from this pattern. As predicted, a singular scaling characterized by the fractal dimension $d_v = 3/2$ of the Wiener processes is observed over the whole range of values of the inertial parameters μ and γ . The effect of μ is practically negligible as can be observed from the data of panel (B), almost collapsing into a single curve. The effect of γ (panel A) is much more sensible and, as γ decreases, the resolution required to observe a fractal behaviour is below $\Delta t = 0.1$, i.e. order of a tenth of the dissipation time.

Therefore, the possible occurrence of regular velocity dynamics in Brownian motion experiments, (this has been reported in [\[6](#page-35-11), [13\]](#page-35-7), albeit it should be further checked experimentally in order to obtain a definitive validation of it), strongly supports the thesis developed in this article of correlated forcings, at least as regards the inertial terms $\boldsymbol{\xi}^{(i)}(t)$. But, as discussed in the previous Sections, this implies also the occurrence of a non vanishing added mass stemming from correlated stochastic forcings.

Back to the hydrodynamic reasoning, this result is not surprising if not intuitive: the inclusion in particle hydromechanics of the effects of a finite speed of sound, i.e. compressibility, represents solely a partial and incomplete refinement of the theory, as it keeps in classical fluctuational pattern theory, an infinite propagation velocity of the stochastic forcings arising a consequence of the hydrodynamic fluctuations in the fluid medium. The complete regularization of the hydrodynamic interactions, removing also the singularity of Wiener forcings in fluctuation-dissipation models, not only ensures the regularity of $v(t)$, but, as an aftermath, determines the emergence of the added-mass effect in a way that is consistent with the global fluctuation-dissipation relation.

VI. FLUCTUATION-INERTIA RELATION

In the previous Sections we have outlined a fluctuational theory of the added-mass effect consistent with the global fluctuation-dissipation relation and associated with correlated

FIG. 7. Length-resolution analysis, $L(\Delta t)$ vs Δt , for a realization of the $\mathbb{DI}_{1,1}^{(1,1,1)}(\xi^{(d)}(t),\xi^{(i)}(t))$ patterns in the presence of a harmonic potential with $k = 0.2$ and $\lambda = 1$, at different values of the parameters characterizing the inertial term. Symbols are the results of the stochastic simulations. Panel (A): $\mu = 1$. Symbols (\square) and line (a) refer to $\gamma = 0.1$, (\circ) and line (b) to $\gamma = 1$, (\bullet) and line (c) to $\gamma = 10$. The solid line (d) represents the scaling $L(\Delta t) \sim \Delta t^{-1/2}$. Panel (B): $\gamma = 0.1$. Symbols (\square) refer to $\mu = 0.1$, (\circ) to $\mu = 1$, (\bullet) to $\mu = 10$. The solid line (a) represents the scaling $L(\Delta t) \sim \Delta t^{-1/2}.$

stochastic forcings acting within the structure of the fluctuational patterns.

But the phenomenon of added mass is also a hydrodynamic property occurring in the accelerated motion of macroscopic objects in a fluid medium of non-negligible density with respect to their densities. It is therefore important to address the connection between the fluctuational added mass and its macroscopic hydrodynamic counterpart. For simplicty, consider the case of a spherical particle, amenable to a scalar formalism.

The starting point is the structure of the force exerted by the fluid on the particle in a Newtonian incompressible fluid eq. [\(6\)](#page-5-0). Apart from the specific instantaneous nature of the dissipative kernel and of the component of the fluid-inertial kernel pertaining to the instantaneous added mass, the dependence of the hydromecanic kernels on the physical parameters characterizing the fluid (density ρ_0 and viscosity μ) reveals:

• the presence of a dissipative kernel $h(t)$ proportional to the fluid viscosity μ ,

$$
h(t) \sim \mu \tag{81}
$$

and consequenly vanishing in the limit $\mu \to \infty$ of an inviscid fluid;

• the decomposition of the fluid inertial kernel into two contributions,

$$
k(t) = k_{\mu}(t) + k_0(t)
$$
\n(82)

where $k_{\mu}(t)$ depends both on ρ_0 and μ

$$
k_{\mu}(t) \sim \sqrt{\mu \,\rho_0} \tag{83}
$$

vanishing for $\mu \to \infty$ and responsible, in eq. [\(6\)](#page-5-0), for the Basset force, and $k_0(t)$ proportional exclusively to the fluid density,

$$
k_0(t) \sim \rho_0 \tag{84}
$$

that in the Newtonian incompressible case accounts for the instantaneous added mass.

This classification can be extended to generic fluids, substituting the Newtonian viscosity with the long-term viscosity as the integral of the viscosity memory kernel $|40|$. By the linearity of the hydrodynamic regimes considered, the decomposition eq. [\(82\)](#page-29-0) depends on the splitting of the overall shear-stress tensor into dissipative shear stresses associated with $k_{\mu}(t)$, and compressible pressure forces determining $k_0(t)$. The latter contribution can be thus uniquely defined starting from the overall fluid inertial kernel $k(t)$ as

$$
k_0(t) = \lim_{\mu \to 0} k(t) \tag{85}
$$

In the most accurate representation of fluid-particle interactions in which acoustic effects are also accounted for, $k_0(t)$ is a continuous function of its argument, rapidly falling to zero for timescales comparable with the characteristic dissipation time $t_{\text{diss}} = m/(6 \pi \mu R_p)$ (order of 10^{-6} -10⁻⁷ s, for a micrometric particle in water), as the acoustic time scale $t_{\text{acou}} \simeq R_p/c_s$, is order of 10[−]⁹ s in the same particle-fluid system. Introducting the decomposition eq. [\(82\)](#page-29-0) into particle hydromechanics, we have

$$
m\frac{dv(t)}{dt} = -\int_0^t h(t-\tau)v(\tau) d\tau - \int_0^t k_\mu(t-\tau) \left(\frac{dv(\tau)}{d\tau} + v(0)\delta(\tau)\right) d\tau
$$

+
$$
\int_0^t k_0(t-\tau) \left(\frac{dv(\tau)}{d\tau} + v(0)\delta(\tau)\right) d\tau
$$
(86)

Next, consider a macroscopic object slowly accelerating in a fluid. For the time-scales at which $k_0(t)$ is significantly different from zero, $dv(t)/dt$ is practically a constant acceleration, and consequently the last term in eq. [\(86\)](#page-30-0), upon a change of variables in the convolution integral, can be expressed as

$$
\int_0^t k_0(\tau) \left(\frac{dv(t-\tau)}{d\tau} + v(0)\delta(t-\tau) \right) d\tau \simeq \frac{dv(t)}{dt} \int_0^t k_0(\tau) d\tau \simeq \frac{dv(t)}{dt} m_a^{(\text{hydro})} \tag{87}
$$

where the hydrodynamic added mass $m_a^{(\text{hydro})}$ equals the integral of the inviscid kernel $k_0(t)$,

$$
m_a^{\text{(hydro)}} = \int_0^\infty k_0(t) \, dt \tag{88}
$$

From the above analysis it may seem that $m_a^{(\text{hydro})}$ and the fluctuational added mass entering the statistical properties of velocity fluctuations of micrometric particles eq. [\(3\)](#page-2-0) refer to two different hydrodynamic phenomenologies. But the origin of the added-mass effect, either at microscale or for macroscopic objects, should stem from one and the same physical mechanism. This is further confirmed qualitatively (yet not fully quantitatively, thus a further careful experimental scrutiny is required) by the experimental results reported in [\[14](#page-35-8)].

This observation leads to a principle of equivalence of the added masses, that can be referred to as the *fluctuation-inertia* relation. This is stated by the following conjecture:

Conjecture - Fluctuation-inertia relation - The fluctuational added mass m_a , defined by eq. [\(3\)](#page-2-0) and the hydrodynamic added mass defined by eq. [\(88\)](#page-30-1), for the same object in the same hydrodynamic conditions, coincide.

Given the validity of the above principle, it remains to determine the structure of the correlated stochastic forcings. If we assume that each $q_i(t)$ is characterized by the same relaxation rate ν then, for any specific fluctuational pattern, the fluctuational added mass m_a is a monotonically decreasing functon of ν , $m_a = \Phi(\nu)$ such that $\lim_{\nu \to \infty} \Phi(\nu) = 0$. Making use of the fluctuation-inertia principle, ν is uniquely defined by the relation

$$
\nu = \Phi^{-1} \left(m_a^{\text{(hydro)}} \right) \tag{89}
$$

Of course, it is important to stress that the velocity autocorrelation function determined by the correlated patterns enforcing eq. [\(89\)](#page-31-0) is different from the Kubo FD1k relation based on the validity of the Langevin condition. However, once the theory is applied to realistic Prony expansions of hydromechanic kernels, it may happen that the difference in the two autocorrelation functions may be tiny and difficult to detect from experimental measurements.

Eq. [\(89\)](#page-31-0) represents a first closure of a correlated theory of hydromechanics velocity fluctuations. Further theoretical work is needed in order to improve this formulation in the form of fluctuation-inertia relations analogous to the Kubo first and second fluctuationdissipation relations, defining the fine structure of the correlated fluctuations and relating them to functional structure of $k_0(t)$, and not only to its integral.

The fluctuation-inertia relation unifies the inertial phenomenology at micro- and macroscales. It physically indicates that the characteristic correlation time associated with fluid fluctuations is related to the microscopic added mass, and vice versa. For this reason, it can be regarded as the inertial counterpart of the global fluctuation-dissipation relation eq. [\(12\)](#page-8-2). Further experiments on micrometric particles in liquids should be performed in order to verify this conjecture. On the other hand, the proposed conjecture stresses also the theoretical hydrodynamic modelling of a compressible liquid phase as, the compressible approach proposed by Chow and Hermans should be extended to a more realistic description of a compressible liquid phase. To this purpose, the theory proposed in [\[20\]](#page-36-3) is an interesting starting point.

VII. CONCLUDING REMARKS

This third, and final article of the trilogy on fluctuation-dissipation relations has addressed a new theoretical formulation of particle hydromechanics grounded on the overcoming of the Langevin condition, within the constraint imposed by the global fluctuationdissipation relation. Technically, this is made possible by considering correlated stochastic forcing within the basic fluctuational patterns introduced in part I. This generalization follows the research line towards a progressively more accurate description of the hydrodynamics at microscale, removing the unphysical approximations, such e.g. the assumption of incompressibility, thus accounting for the finite speed of propagation of any hydrodynamic field (pressure/density, internal stresses).

While the inclusion of acoustic effects, still imbedded within a Wiener-based theory of fluctuations, such e.g. in the Chow-Hermans work, has "eliminated " the occurrence of the added-mass effect, restoring the energy equipartition for the squared velocity variance of colloidal particles in a still liquid, in virtue of the regularity (finite speed of sound) of the acoustic propagation, the consistent, and almost due extension of the same regularity principle, applied to stochastic forcings entering the equations of motion, restored the addedmass effect as a generic property of correlated elementary fluctuations. This restoration of the added-mass is altogeher different from the same hydrodynamic counterpart occuring due to incompressibility, as it occurs in generic compressible flows due to the correlated properties of thermal/hydrodynamic fluctuations.

In the development of the theory, the role of the fluctuational patterns introduced in part I should be stressed, as the basic structural form for describing in a compact way FD3k in different hydrodynamic conditions of increasing complexity. The structure of these patterns has been exploited in part II in order to analyze the properties of white-noise impulsive stochastic forcings, and in the present article to extend the theory to correlated fluctuations maintaining the validity of the global fluctuation-dissipation relation.

Moreover, we have linked the occurrence of the added-mass effect, associated with fluctuational properties, to the regularity of the velocity signal $v(t)$. And also this property clearly emerges from the structure of the fluctuational patterns accounting for fluid-inertial effects, i.e. $\mathbb{DI}_{m,E}^{(n,N_d,N_i)}$. While velocity signals in $\mathbb{DI}_{m,E}^{(n,N_d,N_i)}$ displays a fractal character (almost everywhere not differentiable), this property disappear in their correlated counterparts, i.e.

 $\mathbb{CDI}_{m,E}^{(n,N_d,N_i)}$, possessing Lipshitzian velocity signals.

The regularity of velocity signals (i.e. their almost everywhere Lipshitz continuity) can be "easily" checked experimentally starting from the analysis of velocity time series in highly resolved Brownian experiments. It constitutes the local (in time) counterpart on the cumulative statistical effect expressed by a non-vanishing fluctuational added mass.

A major thoretical effort is still to be performed in order to derive a finer characterization of fluctuation-inertia relations conceptually similar to the FD1k and FD2k relations of the classical dissipative Kubo theory grounded on the Langevin condition.

A general fluctuation-inertia principle has also been outlined, connecting the fluctuational added mass occurring at microscales to the hydrodynamic added mass characterizing the inertial properties of macroscopic objects accelerating in a liquid.

The proposed theory is potentially rich of new theoretical implications as regards either statistical physics or hydrodynamics of liquid phases. These implications will be thoroughly addressed in a forthcoming communication.

Acknowledgment - This research received financial support from ICSC—Centro Nazionale di Ricerca in High Performance Computing, Big Data and Quantum Computing, funded by European Union—NextGenerationEU. One of the authors (M.G.) is grateful to M. G. Raizen for precious discussions.

Appendix A: Moment analysis of the $\mathbb{CI}^{(1,1,1)}_{m,E}$ pattern

Consider the second-order moments associated with the dynamics variables of the correlated pattern $\mathbb{CI}_{m,E}^{(1,1,1)}(q(t),q_1(t))$ analyzed in paragraph [IV C.](#page-21-1) Set

$$
\mathbf{X} = (m_{vv}^*, m_{vz}^*, m_{vq}^*, m_{vq_1}^*, m_{zz}^*, m_{z1}^*, m_{zq1}^*)^T
$$
\n(A1)

The vector X of the steady-state (equilibrium) values is the solution of the linear system

$$
A X = b \tag{A2}
$$

where

$$
\mathbf{A} = \begin{pmatrix}\n-(\gamma + \alpha) & \gamma & \sqrt{2}a & \sqrt{2}d & 0 & 0 & 0 \\
0 & \mu & 0 & 0 & -\mu & 0 & \sqrt{2}c \\
\mu & -(\gamma + \alpha + \mu) & 0 & \sqrt{2}c & \gamma & \sqrt{2}a & \sqrt{2}d \\
0 & 0 & -(\gamma + \alpha + \nu) & 0 & 0 & \gamma & 0 \\
0 & 0 & 0 & -(\gamma + \alpha + \nu_1) & 0 & 0 & \gamma \\
0 & 0 & \mu & 0 & 0 & -(\mu + \nu) & 0 \\
0 & 0 & 0 & \mu & 0 & 0 & -(\mu + \nu_1)\n\end{pmatrix}
$$
\n(A3)

and

$$
\mathbf{b} = \left(0, 0, 0, -\frac{a\nu}{\sqrt{2}}, -\frac{d\nu_1}{\sqrt{2}}, 0, -\frac{c\nu_1}{\sqrt{2}}\right)^T
$$
(A4)

Solving this linear system the expression for $m_{vv}^* = \langle v^2 \rangle_{eq}$ takes the form

$$
\frac{m\langle v^2 \rangle_{\text{eq}}}{k_B T} = \frac{N_0 + N_1 \alpha + N_2 \alpha^2}{D_0 + D_1 \alpha + D_2 \alpha^2 + D_3 \alpha^3} \tag{A5}
$$

$$
N_0 = \nu \nu_1 (\gamma + \mu) (\mu + \nu_1) (\gamma + \mu + \nu)
$$

\n
$$
N_1 = \nu (\mu + \nu) (\mu + \nu_1)^2 + \gamma [\mu^2 \nu + \nu_1^2 (\mu + \nu) + \nu \nu_1 (2 \mu + \nu)]
$$

\n
$$
N_2 = \nu (\mu + \nu) (\mu + \nu_1)
$$

\n
$$
D_0 = \nu \nu_1 (\gamma + \mu) (\gamma + \mu + \nu) (\gamma + \mu + \nu_1)
$$

\n
$$
D_1 = \nu_1 (\gamma + \mu) (\mu + \nu) (\gamma + \mu + \nu_1) + \nu (\gamma + \mu) (\mu + \nu_1) (\gamma + \mu + \nu)
$$

\n
$$
+ \nu \nu_1 (\gamma + \mu + \nu) (\gamma + \mu + \nu_1)
$$

\n
$$
D_2 = (\gamma + \mu) (\mu + \nu) (\mu + \nu_1) + \nu_1 (\mu + \nu) (\gamma + \mu + \nu_1) + \nu (\mu + \nu_1) (\gamma + \mu + \nu)
$$

\n
$$
D_3 = (\mu + \nu) (\mu + \nu_1)
$$

Set $\mathbf{p} = (\gamma, \mu, \nu, \nu_1)$ and let $\mathbf{p}^* = (\gamma^*, \mu^*, \nu^*, \nu_1^*)$ be the corresponding nondimensional set of rates rescaled by $1/\alpha$, i.e. $\gamma^* = \gamma/\alpha$ and similarly for the other entries. The squared velocity variance at equilibrium can be expressed as

$$
\frac{m\langle v^2 \rangle_{\text{eq}}}{k_B T} = \frac{N_0 + N_1 + N_2}{D_0 + D_1 + D_2 + D_3} \bigg|_{\mathbf{p} = \mathbf{p}^*}
$$
(A7)

[1] M. Giona, G. Procopio and C. Pezzotti, Fluid-particle interactions and fluctuation-dissipation relations I - General linear theory and basic fluctuation patterns, submitted to arXiv (2024).

- [2] C. Pezzotti, M. Giona and G. Procopio, Fluid-particle interactions and fluctuation-dissipation relations II - Gaussianity and Gaussianity breaking, submitted to arXiv (2024).
- [3] P. Langevin, Sur la theorie du mouvement brownien. C. R. Acad. Sci. (Paris) 146, 530-533 (1908), see also the English translation in Am. J. Phys. 65, 1079-1081 (1997).
- [4] R. Kubo, M. Toda and N. Hashitsune, Statistical Physics II Nonequilibrium Statistical Mechanics (Springer-Verlag, Berlin, 1991).
- [5] R. Huang, I. Chavez, K. M. Taute, B. Lukic, S. Jeney, M. G. Raizen and E. L. Florin, Direct observation of the full transition from ballistic to diffusive Brownian motion in a liquid. Nature Phys. 7, 576-580 (2011).
- [6] T. Franosch, M. Grimm, M. Belushkin, F. M. Mor, G. Foffi, L. Forró and S. Jeney, Resonances arising from hydrodynamic memory in Brownian motion, Nature 478, 85-88 (2011).
- [7] P. N. Pusey, Brownian motion goes ballistic, Science 332, 802-803 (2011).
- [8] S. Kheifets, A. Simha, K. Melin, T. Li and M. G. Raizen, Observation of Brownian motion in liquids at short times: instantaneous velocity and memory loss, Science 343, 1493-1496 (2014).
- [9] A. Widom, Velocity Fluctuations of a Hard-Core Brownian Particle, Phys. Rev. A 3, 1394- 1396 (1971).
- [10] R. M. Mazo, Theory of Brownian Motion. IV. A Hydrodynamic Model for the Friction Factor, J. Chem. Phys. 54 3712-3713 (1971).
- [11] R. E. Burgess, Brownian motion and the equipartition theorem, Phys. Lett. 42A, 395-396 (1973).
- [12] H. J. H. Clercx and P. P. J. M. Schram, Brownian particles in shear flow and harmonic potentials: A study of long-time tails, Phys. Rev. A 46 1942-1950 (1992).
- [13] M. G. Raizen and T. Li, The measurement Einstein deemed impossible, Physics Today 68, 56-57 (2016).
- [14] J. Mo, A. Simha, S. Kheifets and M. G. Raizen, Testing the Maxwell-Boltzmann distribution using Brownian particles, Optics Exp. 23 1888-1893 (2015).
- [15] L. D. Landau and E. M. Lifshitz, Fluid Mechanics (Pergamon Press, Oxford, 1959).
- [16] S. Kim and S. J. Karrila, Microhydrodynamics - Principles and Selected Applications (Dover Publ., Mineola USA, 2005).
- [17] R. Zwanzig and M. Bixon, Compressibility effects in the hydrodynamic theory of Brownian motion, J. Fluid Mech. 69, 21-25 (1975).
- [18] T. S. Chow and J. J. Hermans, Brownian motion of a spherical particle in a compressible fluid, Physica 65, 156-162 (1973).
- [19] L. M. Cheung and C. Jarzynski, Added mass effect in coupled Brownian particles, [arXiv:2405.01696](http://arxiv.org/abs/2405.01696) (2024).
- [20] M. Giona, G. Procopio, A. Adrover and R. Mauri, New formulation of the Navier–Stokes equations for liquid flows, J. Non-Equilibrium Thermodyn. 48, 207-228 (2023).
- [21] C. Darwin, Note on hydrodynamics, Math. Proc. Cambridge Phil. Soc. 49, 342-354 (1953).
- [22] A. Einstein, Investigations on the Theory of Brownian Movement (Dover Publ., Mineola USA, 1956).
- [23] J. B. Perrin, Le mouvement brownien et la réalité moléulaire, Ann. Chem. Phys. 18, 5-114 (1909).
- [24] R. Newburgh, J. Peidle and W. Rueckner, Einstein, Perrin, and the reality of atoms: 1905 revisited, Am. J. Phys. 74, 478-481 (2006).
- [25] C. E. Brennen, A Review of Added Mass and Fluid Inertial Forces, Technical Report–CR82.010; Naval Civil Engineering Laboratory: Port Hueneme, CA, USA, 1982.
- [26] D. T. Sen and T. C. Vinh, Determination of added mass and inertia moment of marine ships moving in 6 degrees of freedom, Int. J. Transp. Eng. Technol. 2 8-14 (2016).
- [27] K. McKee and A. Czarnecki, Acceleration due to buoyancy and mass renormalization, Am. J. Phys. 87 165-170 (2019).
- [28] G. Birkhoff, *Hydrodynamics. A study in logic, fact and similitude*, Revised Edition, (Princeton University Press, Princeton, 1960).
- [29] M. Grimm, S. Jeney and T. Franosch, Brownian motion in a Maxwell fluid, Soft Matter 7, 2076-2084 (2011).
- [30] A. Cunsolo, G. Ruocco, F. Sette, C. Masciovecchio, A. Mermet, G. Monaco, M. Sampoli and R. Verbeni, Experimental determination of the structural relaxation in liquid water, Phys. Rev. Lett. 82, 775-778 (1999).
- [31] G. Procopio and M. Giona, Modal representation of inertial effects in fluid-particle interactions and the regularity of the memory kernels, Fluids 8, 84 (2023).
- [32] D. Jou, J. Casa-Vázquez and G. Lebon, *Extended Irreversible Thermodynamics* (Springer-Verlag, Berlin, 2001).
- [33] M. Giona and L. Pucci, Hyperbolic heat/mass transport and stochastic modelling-Three simple problems, Mathematics in Engineering 1, 224-251 (2019).
- [34] M. Kac, A stochastic model related to the telegrapher's equation, Rocky Mountain J. Math. 4, 497-509 (1974).
- [35] M. Giona, A. Brasiello and S. Crescitelli, Stochastic foundations of undulatory transport phenomena: Generalized Poisson–Kac processes - Part I basic theory, J. Phys. A 50, 335002 (2017).
- [36] M. Giona, A. Cairoli and R. Klages, Extended Poisson-Kac theory: A unifying framework for stochastic processes with finite propagation velocity, Phys. Rev. X 12, 021004 (2022).
- [37] C. Tricot, Curves and Fractal Dimension (Springer-Verlag, New York, 1995).
- [38] A. Lasota and M. C. Mackey, Chaos, Fractals, and Nois (Springer-Vwelag, New York, 1994).
- [39] K. Falconer, *Fractal Geometry* (J. Wiley $\%$ Sons, Chichester, 2014).
- [40] C. W. Macosko, Rheology - Principles, Measurements, and Applications (Wiley-VCH, New York, 1994).