PRIORS FOR SECOND-ORDER UNBIASED BAYES ESTIMATORS

MANA SAKAI*[†] TAKERU MATSUDA*[‡] TATSUYA KUBOKAWA*

*The University of Tokyo

†RIKEN Center for Advanced Intelligence Project

‡RIKEN Center for Brain Science

ABSTRACT. Asymptotically unbiased priors, introduced by Hartigan (1965), are designed to achieve second-order unbiasedness of Bayes estimators. This paper extends Hartigan's framework to non-i.i.d. models by deriving a system of partial differential equations that characterizes asymptotically unbiased priors. Furthermore, we establish a necessary and sufficient condition for the existence of such priors and propose a simple procedure for constructing them.

The proposed method is applied to several examples, including the linear regression model and the nested error regression (NER) model (also known as the random effects model). Simulation studies evaluate the frequentist properties of the Bayes estimator under the asymptotically unbiased prior for the NER model, highlighting its effectiveness in small-sample settings.

1. Introduction

1.1. **Background.** In Bayesian inference, the choice of prior distribution is a crucial consideration. When researchers have prior knowledge about the parameter of interest, they can incorporate this information into the prior. In the absence of prior knowledge, however, it is common to use so-called objective priors. Well-known examples of objective priors include Jeffreys' prior (Jeffreys, 1961) and reference priors (Bernardo, 1979; Berger and Bernardo, 1992), among others.

In addition to these objective priors, Hartigan (1965) introduced the concept of asymptotically unbiased priors that are designed to achieve second-order unbiasedness of Bayes estimators. It should be noted that Hartigan (1965) approached this topic within the framework of decision theory, considering the general loss functions and Bayes estimators. Additionally, the definition of bias in his work differs slightly from the standard one. In this paper, we focus on the squared error loss and the posterior mean as the Bayes estimator, referring to bias as the difference between the expected value of the estimator and the true parameter value.¹

Hartigan (1965) showed that, in general, the posterior mean has a bias of $O(n^{-1})$, where n is the sample size. He defined the asymptotically unbiased prior as one that reduces this bias to $o(n^{-1})$, achieving second-order unbiasedness. Hartigan (1965) considered i.i.d.

Date: December 30, 2024.

Email: mana.sakai.77@gmail.com, matsuda@mist.i.u-tokyo.ac.jp, tatsuya@e.u-tokyo.ac.jp

¹Note that for the squared error loss, the definition of bias in Hartigan (1965) corresponds to the square of the bias in the standard (or our) definition.

models and derived conditions that asymptotically unbiased priors should satisfy. For onedimensional parameter models, the asymptotically unbiased priors are proportional to the Fisher information of the models. For multi-parameter models, such priors are characterized as solutions to a system of partial differential equations.

The utility of asymptotically unbiased priors lies in their ability to reduce bias of the Bayes estimator, which is particularly important when the sample size is small and bias tends to be large. In this sense, asymptotically unbiased priors are similar to the bias reduction approach for maximum likelihood estimators (MLE) introduced by Firth (1993).

1.2. **Motivation and contributions.** The results of Hartigan (1965) are significant, but the restriction to i.i.d. models often proves too limiting in practice. For instance, in Bayesian regression problems, it is customary to treat the response variables as random while assuming the explanatory variables are fixed. In such cases, the data are not i.i.d. and the results of Hartigan (1965) do not directly apply.

In this paper, we extend the findings of Hartigan (1965) to non-i.i.d. models, deriving a system of partial differential equations that characterizes asymptotically unbiased priors. The generalization broadens the applicability of asymptotically unbiased priors to a wider range of models.

Moreover, we establish a necessary and sufficient condition for the existence of such priors and present a simple procedure for constructing them. These results are particularly valuable, as it provides a unified framework for constructing asymptotically unbiased priors across various models. We note that this condition and the construction method are also applicable to other classes of priors $\pi(\theta)$ for a p-dimensional parameter θ , which are defined as solutions to a system of partial differential equations of the form

$$\frac{\partial}{\partial \theta_r} \log \pi(\theta) = A_r(\theta), \quad r = 1, \dots, p,$$

such as the moment matching priors proposed by Ghosh and Liu (2011).

To demonstrate the usefulness of our construction method, we apply it to several examples, including the linear regression model, the nested error regression (NER) model (also known as the random effects model), and other simpler models. In the linear regression model, we show that the asymptotically unbiased prior we derive leads to an exactly unbiased estimator for both the regression coefficients and the error variance. For the NER model, although an analytical form of the posterior mean with the asymptotically unbiased prior is not available, we evaluate its performance through simulation studies. The prior we construct for the NER model, to the best of our knowledge, has not been explored in the existing literature. Our simulation studies highlight its effectiveness in small samples (i.e., small number of areas/groups), particularly in terms of bias reduction.

Our contributions can be summarized as follows.

- 1. We derive the system of partial differential equations that asymptotically unbiased priors should satisfy in a general setting where the model is not necessarily i.i.d. This extends the work of Hartigan (1965) under the squared error loss.
- 2. We establish a necessary and sufficient condition for the existence of asymptotically unbiased priors and propose a simple procedure for constructing such priors.

- 3. We propose a novel asymptotically unbiased prior for the NER/random effects model. Simulation studies demonstrate that the proposed prior reduces bias effectively in small samples.
- 1.3. Organization of the paper. The remainder of the paper is organized as follows. In Section 2, we present the main theoretical results. In Section 3, we apply these results to several examples, including the linear regression model and the NER model. Section 4 focuses on simulation studies to evaluate the performance of the asymptotically unbiased prior for the NER model. In Section 5, we provide proofs of the main results.

2. Main results

In this section, we present the main theoretical results of this paper. Section 2.1 begins with the derivation of the second-order bias of Bayes estimators under simple regularity conditions (Corollary 2). Building on this result, with additional assumptions, we further simplify the evaluation of bias (Corollaries 3 and 4). In Section 2.2, we establish the necessary and sufficient condition for the existence of an asymptotically unbiased prior (Theorem 5) and propose a practical construction method for such priors (Corollary 7). Finally, we demonstrate that an asymptotically unbiased prior always exists for one-parameter models.

2.1. Second-order bias of Bayes estimators. Suppose (X_1, \ldots, X_n) has a joint density function $f_n(x_1, \ldots, x_n | \theta)$. Here, θ is an interior point of the parameter space Θ , which is a rectangular subspace of \mathbb{R}^p . The log-likelihood function is denoted by

$$\ell_n(\theta) = \log f_n(x_1, \dots, x_n | \theta).$$

To find the maximum likelihood estimator (MLE) $\hat{\theta}^{ML}$, we need to solve the equation

$$\left. \frac{\partial}{\partial \theta} \ell_n(\theta) \right|_{\theta = \hat{\theta}^{ML}} = 0. \tag{1}$$

Let $\hat{\theta}^B$ denote the posterior mean, which we call the (generalized) Bayes estimator, corresponding to a prior density π . $\hat{\theta}^B$ can be calculated as

$$\hat{\theta}^{B} = \frac{\int_{\Theta} \theta \pi(\theta) f_{n}(x_{1}, \dots, x_{n} | \theta) d\theta}{\int_{\Theta} \pi(\theta) f_{n}(x_{1}, \dots, x_{n} | \theta) d\theta} = \frac{\int_{\Theta} \theta \pi(\theta) \exp(\ell_{n}(\theta)) d\theta}{\int_{\Theta} \pi(\theta) \exp(\ell_{n}(\theta)) d\theta}.$$
 (2)

The negative second-order derivative of the log-likelihood is denoted by

$$H_n(\theta) = -\frac{1}{n} \frac{\partial^2}{\partial \theta \partial \theta^{\top}} \ell_n(\theta).$$

We assume the following regularity conditions.

Assumption 1. Assume that the following conditions are satisfied:

- (i) $\hat{\theta}^{ML} \theta = O_p(n^{-1/2}).$
- (ii) $\ell_n(\theta) = O_p(n)$.
- (iii) $\ell_n(\theta)$ is three times continuously differentiable.
- (iv) The prior density $\pi(\theta)$ is differentiable.
- (v) $H_n(\theta)$ is invertible. $H_n(\hat{\theta}^{ML})$ is positive definite.

Note that (i) and (ii) imply $\frac{\partial}{\partial \theta} \ell_n(\theta) = O_p(n^{1/2})$. For notational simplicity, we write

$$I_{n}(\theta) = \frac{1}{n} \left(\frac{\partial}{\partial \theta} \ell_{n}(\theta) \right) \left(\frac{\partial}{\partial \theta} \ell_{n}(\theta) \right)^{\top},$$

$$J_{n,rs,t}(\theta) = \frac{1}{n} \left(\frac{\partial^{2}}{\partial \theta_{r} \partial \theta_{s}} \ell_{n}(\theta) \right) \left(\frac{\partial}{\partial \theta_{t}} \ell_{n}(\theta) \right),$$

$$K_{n,rst}(\theta) = \frac{1}{n} \frac{\partial^{3}}{\partial \theta_{r} \partial \theta_{s} \partial \theta_{t}} \ell_{n}(\theta).$$

Additionally, we denote the (r, s)-th element of any matrix M as M_{rs} and the (r, s)-th element of its inverse (if it exists) as M^{rs} .

Theorem 1. Suppose the model satisfies Assumption 1. Then, the Bayes estimator can be decomposed as

$$\hat{\theta}^B = \theta + \frac{1}{n} H_n^{-1}(\theta) \left(\frac{\partial}{\partial \theta} \left(\log \pi(\theta) + 2\ell_n(\theta) \right) + \frac{1}{2} \sum_{r=1}^p \sum_{s=1}^p H_n^{rs}(\theta) A_{n,rs}(\theta) \right) + o_p \left(\frac{1}{n} \right), \quad (3)$$

where $A_{n,rs}(\theta)$ for each $r,s \in \{1,\ldots,p\}$ is given by

$$A_{n,rs}(\theta) = \begin{bmatrix} K_{n,1rs}(\theta) + 2J_{n,1r,s}(\theta) \\ \vdots \\ K_{n,prs}(\theta) + 2J_{n,pr,s}(\theta) \end{bmatrix} + \sum_{t=1}^{p} \sum_{u=1}^{p} H_n^{tu}(\theta) I_{n,su}(\theta) \begin{bmatrix} K_{n,rt1}(\theta) \\ \vdots \\ K_{n,rtp}(\theta) \end{bmatrix}.$$

The proof of Theorem 1 is provided in Section 5.1.1. Assuming that the expectation of the remainder term $o_p(n^{-1})$ in (3) is $o(n^{-1})$, we can evaluate the bias of the Bayes estimator as follows.

Corollary 2. Suppose the model satisfies Assumptions 1. Then, the bias of the Bayes estimator is given by

$$\mathbb{E}(\hat{\theta}^B) - \theta = \frac{1}{n} \mathbb{E} \left[H_n^{-1}(\theta) \left(\frac{\partial}{\partial \theta} \left(\log \pi(\theta) + 2\ell_n(\theta) \right) + \frac{1}{2} \sum_{r=1}^p \sum_{s=1}^p H_n^{rs}(\theta) A_{n,rs}(\theta) \right) \right] + o\left(\frac{1}{n}\right). \tag{4}$$

If, in particular, $H_n(\theta)$ is non-stochastic (and so is $K_{n,rst}$), the bias of the Bayes estimator can be expressed as

$$\mathbb{E}(\hat{\theta}^B) - \theta = \frac{1}{n} H_n^{-1}(\theta) \left(\frac{\partial}{\partial \theta} \log \pi(\theta) + \frac{1}{2} \sum_{r=1}^p \sum_{s=1}^p H_n^{rs}(\theta) \mathbb{E} \left[A_{n,rs}(\theta) \right] \right) + o\left(\frac{1}{n}\right).$$

Thus, the Bayes estimator is second-order unbiased if the prior satisfies

$$\frac{\partial}{\partial \theta} \log \pi(\theta) = \phi(\theta) := -\frac{1}{2} \sum_{r=1}^{p} \sum_{s=1}^{p} H_n^{rs}(\theta) \mathbb{E} \left[A_{n,rs}(\theta) \right]. \tag{5}$$

Note that in this case, $\mathbb{E}[A_{n,rs}(\theta)]$ has the form of

$$\mathbb{E}\left[A_{n,rs}(\theta)\right] = \begin{bmatrix} K_{n,1rs}(\theta) + 2\mathbb{E}\left[J_{n,1r,s}(\theta)\right] \\ \vdots \\ K_{n,prs}(\theta) + 2\mathbb{E}\left[J_{n,pr,s}(\theta)\right] \end{bmatrix} + \sum_{t=1}^{p} \sum_{u=1}^{p} H_n^{tu}(\theta)\mathbb{E}\left[I_{n,su}(\theta)\right] \begin{bmatrix} K_{n,rt1}(\theta) \\ \vdots \\ K_{n,rtp}(\theta) \end{bmatrix}.$$

It should be noted that the prior satisfying (5) depends on the sample size n.

It seems relatively easier to find a prior that ensures second-order unbiasedness of the Bayes estimator if $H_n(\theta)$ is non-stochastic. However, in general, $H_n(\theta)$ is a random matrix. In such cases, we need alternative strategies to find a prior that leads a second-order unbiased Bayes estimator. One possible approach is to consider the asymptotic properties of elements such as $H_n(\theta)$ and $K_{n,rst}(\theta)$. With some additional assumptions, we can simplify the evaluation of the bias of the Bayes estimator given in (4).

Assumption 2. Assume that the following conditions are satisfied:

(i) There exists a non-singular $p \times p$ constant matrix $H(\theta)$ that satisfies

$$H_n(\theta) = H(\theta) + O_p(n^{-1/2})$$

and consequently,

$$H_n^{-1}(\theta) = H^{-1}(\theta) + O_p(n^{-1/2}).$$

(ii) For each $r, s, t \in \{1, ..., p\}$, there exists a constant $K_{rst}(\theta)$ that satisfies

$$K_{n,rst}(\theta) = K_{rst}(\theta) + o_p(1).$$

(iii) There exists a $p \times p$ matrix $I(\theta)$ that satisfies

$$\mathbb{E}\left[I_n(\theta)\right] = I(\theta) + o(1).$$

(iv) For each $r, s, t \in \{1, ..., p\}$, there exists $J_{rs,t}(\theta)$ that satisfies

$$\mathbb{E}\left[J_{n,rs,t}\right] = J_{rs,t}(\theta) + o(1).$$

With Assumption 2 in place, we can now proceed to simplify the evaluation of the bias of the Bayes estimator, taking advantage of the asymptotic behavior of the matrices involved. Specifically, these conditions allow us to express the bias in a form that does not depend on the sample size n.

Corollary 3. Suppose the model satisfies Assumptions 1 and 2. Then, the bias of the Bayes estimator is given by

$$\mathbb{E}(\hat{\theta}^B) - \theta = \frac{1}{n} H^{-1}(\theta) \left(\frac{\partial}{\partial \theta} \log \pi(\theta) + \frac{1}{2} \sum_{r=1}^p \sum_{s=1}^p H^{rs}(\theta) A_{rs}(\theta) \right) + o\left(\frac{1}{n}\right),$$

where $A_{rs}(\theta)$ for each $r, s \in \{1, ..., p\}$ is defined as

$$A_{rs}(\theta) = \begin{bmatrix} K_{1rs}(\theta) + 2J_{1r,s}(\theta) \\ \vdots \\ K_{prs}(\theta) + 2J_{pr,s}(\theta) \end{bmatrix} + \sum_{t=1}^{p} \sum_{u=1}^{p} H^{tu}(\theta)I_{su}(\theta) \begin{bmatrix} K_{rt1}(\theta) \\ \vdots \\ K_{rtp}(\theta) \end{bmatrix}.$$

This implies that the Bayes estimator is asymptotically unbiased if the prior $\pi(\theta)$ satisfies

$$\frac{\partial}{\partial \theta} \log \pi(\theta) = \phi(\theta) := -\frac{1}{2} \sum_{r=1}^{p} \sum_{s=1}^{p} H^{rs}(\theta) A_{rs}(\theta). \tag{6}$$

The proof of Corollary 3 is provided in Section 5.1.2.

Remark 1. It can be seen that when $H(\theta) = I(\theta)$ holds, we have

$$A_{rs}(\theta) = 2 \begin{bmatrix} K_{1rs}(\theta) + J_{1r,s}(\theta) \\ \vdots \\ K_{prs}(\theta) + J_{pr,s}(\theta) \end{bmatrix}.$$

Hence, the definition of $\phi(\theta)$ in (6) simplifies to

$$\phi_t(\theta) = -\sum_{r=1}^p \sum_{s=1}^p H^{rs}(\theta)(K_{trs}(\theta) + J_{tr,s}(\theta)), \quad t = 1, \dots, p.$$
 (7)

2.1.1. Application to models with i.i.d. random variables. Suppose X_1, \ldots, X_n is a sequence of i.i.d. random variables with density function $f_n(x_1, \ldots, x_n | \theta) = \prod_{i=1}^n f(x_i | \theta)$. Under some regularity conditions, it is well known that $\sqrt{n}(\hat{\theta}^{ML} - \theta)$ converges in distribution to $N(0, I^{-1}(\theta))$, where $I(\theta)$ is the Fisher information matrix

$$I(\theta) = \mathbb{E}\left[\left(\frac{\partial}{\partial \theta} \log f(X|\theta)\right) \left(\frac{\partial}{\partial \theta} \log f(X|\theta)\right)^{\top}\right] = -\mathbb{E}\left[\frac{\partial^2}{\partial \theta \partial \theta^{\top}} \log f(X|\theta)\right]. \tag{8}$$

This means condition (i) of Assumption 1 is satisfied. We compute the limits defined in Assumption 2 in this situation. Since X_1, \ldots, X_n are i.i.d., applying the central limit theorem yields

$$\sqrt{n}(H_n(\theta) - H(\theta)) \xrightarrow{d} N(0, \Sigma),$$

where $H(\theta)$ equals $I(\theta)$ and Σ is some covariance function. Thus, condition (i) of Assumption 2 is satisfied with $H(\theta) = I(\theta)$. Applying the law of large numbers, we can show that condition (ii) of Assumption 2 is also satisfied with

$$K_{rst}(\theta) = \mathbb{E}\left[\frac{\partial^3}{\partial \theta_r \partial \theta_s \partial \theta_t} \log f(X|\theta)\right].$$

Conditions (iii) and (iv) of Assumption 2 are straightforward; we can take $I(\theta)$ in condition (iii) as the Fisher information matrix in (8), and $J_{rs,t}(\theta)$ in condition (iv) as

$$J_{rs,t}(\theta) = \mathbb{E}\left[\left(\frac{\partial^2}{\partial \theta_r \partial \theta_s} \log f(X|\theta)\right) \left(\frac{\partial}{\partial \theta_t} \log f(X|\theta)\right)\right].$$

Since the matrix $H(\theta)$ is identical to the Fisher information matrix $I(\theta)$, by Remark 1, $\phi(\theta)$ is given by (7). Noting that the relation

$$\frac{\partial}{\partial \theta_t} I_{rs}(\theta) + J_{rs,t}(\theta) + K_{rst}(\theta) = 0 \quad (r, s, t \in \{1, \dots, p\})$$
(9)

holds in general, we conclude the following:

Corollary 4. Given the assumptions previously discussed, the bias of the Bayes estimator is given by

$$\mathbb{E}(\hat{\theta}^B) - \theta = n^{-1}I^{-1}(\theta) \left(\frac{\partial}{\partial \theta} \log \pi(\theta) - \sum_{r=1}^p \sum_{s=1}^p I^{rs}(\theta) \begin{bmatrix} \frac{\partial}{\partial \theta_s} I_{1r}(\theta) \\ \vdots \\ \frac{\partial}{\partial \theta_s} I_{pr}(\theta) \end{bmatrix} \right) + o(n^{-1}),$$

where $I(\theta)$ is the Fisher information matrix defined in (8). Consequently, the Bayes estimator is asymptotically unbiased if the prior $\pi(\theta)$ satisfies

$$\frac{\partial}{\partial \theta} \log \pi(\theta) = \phi(\theta) := \sum_{r=1}^{p} \sum_{s=1}^{p} I^{rs}(\theta) \begin{bmatrix} \frac{\partial}{\partial \theta_{s}} I_{1r}(\theta) \\ \vdots \\ \frac{\partial}{\partial \theta_{s}} I_{pr}(\theta) \end{bmatrix}. \tag{10}$$

Remark 2. The above result for i.i.d. models is consistent with the existing result; see Section 7 of Hartigan (1965).

Remark 3. There is an interesting connection between the asymptotically unbiased prior for i.i.d. models, Firth's method (Firth, 1993), and the moment matching prior (Ghosh and Liu, 2011). Firth's method reduces the bias of the MLE to $o(n^{-1})$ by modifying the score equation (1) as

$$\frac{\partial}{\partial \theta} \ell_n(\theta) = -\sum_{r=1}^p \sum_{s=1}^p I^{rs}(\theta) \begin{bmatrix} J_{1r,s}(\theta) + \frac{1}{2} K_{1rs}(\theta) \\ \vdots \\ J_{pr,s}(\theta) + \frac{1}{2} K_{prs}(\theta) \end{bmatrix}. \tag{11}$$

In contrast, the moment matching prior $\pi^M(\theta)$ is defined such that the posterior mean matches the MLE up to $o_p(n^{-1})$. This prior satisfies the equation

$$\frac{\partial}{\partial \theta} \log \pi^{M}(\theta) = -\frac{1}{2} \sum_{r=1}^{p} \sum_{s=1}^{p} I^{rs}(\theta) \begin{bmatrix} K_{1rs}(\theta) \\ \vdots \\ K_{prs}(\theta) \end{bmatrix}.$$
 (12)

Interestingly, by considering the general relation (9), it can be found that the sum of the right-hand sides of (11) and (12) corresponds to the right-hand side of (10). This observation suggests that the asymptotically unbiased prior can be interpreted as a bias-reduced version of the moment matching prior, where the bias reduction is achieved using Firth's method.

- Remark 4. Corollaries 2, 3 and 4 demonstrate a trade-off between the restrictiveness of the assumptions and the simplicity of the derived expressions for the bias. Under minimal regularity conditions, we obtain a general but complex result (Corollary 2), which may have limited practical applicability. By introducing additional assumptions, or by assuming i.i.d. observations, the results become simpler, enabling more practical application (Corollaries 3 and 4). This highlights how stronger assumptions can facilitate application of the results, albeit at the expense of reduced generality. In Section 3, we illustrate the application of Corollaries 3 and 4 to specific models.
- 2.2. Existence of asymptotically unbiased priors. An asymptotically unbiased prior, if it exists, can be obtained by solving (5) (assuming $H_n(\theta)$ is non-stochastic), (6) (assuming the model satisfies Assumption 2), or (10) (for i.i.d. data). However, there is no guarantee that such priors exist for all models. To address this issue, we examine the necessary and sufficient condition for these equations to have a solution. The following theorem applies a classical result from the theory of partial differential equations.

Theorem 5. Let $\phi: \Theta \to \mathbb{R}^p$ be a differentiable vector-valued function,² and assume that the order of integration and differentiation in ϕ can be interchanged, i.e.,

$$\int \left(\frac{\partial}{\partial \theta}\phi(\theta)\right) d\theta = \frac{\partial}{\partial \theta} \left(\int \phi(\theta) d\theta\right).$$

Then, a twice-differentiable prior density $\pi(\theta)$ satisfying

$$\frac{\partial}{\partial \theta} \log \pi(\theta) = \phi(\theta) \tag{13}$$

exists if and only if the following holds:

$$\frac{\partial}{\partial \theta_u} \phi_t(\theta) = \frac{\partial}{\partial \theta_t} \phi_u(\theta) \quad \text{for all } t, u \in \{1, \dots, p\}.$$
 (14)

We refer to (14) as the integrability condition.

The proof of Theorem 5 is provided in Section 5.2.

Remark 5. It is worth noting that the asymptotically unbiased prior that is independent of the sample size n is unique up to a constant multiple, provided it exists. Indeed, if two priors $\pi_1(\theta)$ and $\pi_2(\theta)$ satisfy (13), then for any $r = 1, \ldots, p$, we have

$$\frac{\partial}{\partial \theta_r} \log \left(\frac{\pi_1(\theta)}{\pi_2(\theta)} \right) = 0.$$

This implies that the ratio $\pi_1(\theta)/\pi_2(\theta)$ is a constant.

Remark 6. Meng and Zaslavsky (2002) proposed the single observation unbiased prior (SOUP), which is a prior that ensures unbiasedness of the Bayes estimator for a single observation. Our result is closely related to the concept of SOUP in that the SOUP is always second-order unbiased when considering repeated sampling from the same distribution. Consequently, the necessary and sufficient condition for the existence of an asymptotically unbiased prior also serves as a necessary condition for the existence of a SOUP. In fact, the asymptotically unbiased priors we derived often result in Bayes estimators that are exactly unbiased (see Examples 2, 3, and 5).

If the model is i.i.d. and the Fisher information matrix $I(\theta)$ is diagonal, a simpler sufficient condition for (14) can be considered.

Proposition 6. Suppose the model satisfies the assumptions of Theorem 5. Suppose further that the model is i.i.d. and has a diagonal Fisher information matrix $I(\theta)$ that is twice continuously differentiable. Then, there exists an asymptotically unbiased prior if for any $t, u \in \{1, \ldots, p\}$, the ratio $I_{tt}(\theta)/I_{uu}(\theta)$ can be expressed as the product of the following two functions.

- (a) A twice continuously differentiable function that does not depend on θ_t (but may depend on $\{\theta_r : r \neq t\}$).
- (b) A twice continuously differentiable function that does not depend on θ_u (but may depend on $\{\theta_r : r \neq u\}$).

 $^{^{2}\}phi:\Theta\to\mathbb{R}^{p}$ corresponds to the right-hand side of either (5), (6), or (10).

Proof. Fix $t, u \in \{1, ..., p\}$. By assumption, the ratio $I_{tt}(\theta)/I_{uu}(\theta)$ can be expressed as

$$\frac{I_{tt}(\theta)}{I_{uu}(\theta)} = k_t(\theta_{-t})k_u(\theta_{-u})$$

where $k_t(\theta_{-t})$ and $k_u(\theta_{-u})$ satisfy conditions (a) and (b), respectively. Taking the logarithm of both sides, we have

$$\log I_{tt}(\theta) - \log k_t(\theta_{-t}) = \log I_{uu}(\theta) + \log k_u(\theta_{-u}).$$

Furthermore, differentiating both sides with respect to θ_t and θ_u , we obtain

$$\frac{\partial^2}{\partial \theta_u \partial \theta_t} \left(\log I_{tt}(\theta) - \log k_t(\theta_{-t}) \right) = \frac{\partial^2}{\partial \theta_t \partial \theta_u} \left(\log I_{uu}(\theta) + \log k_u(\theta_{-u}) \right),$$

that is,

$$\frac{\partial}{\partial \theta_u} \left(\frac{1}{I_{tt}(\theta)} \frac{\partial}{\partial \theta_t} I_{tt}(\theta) \right) = \frac{\partial}{\partial \theta_t} \left(\frac{1}{I_{uu}(\theta)} \frac{\partial}{\partial \theta_u} I_{uu}(\theta) \right).$$

Since the Fisher information is diagonal, we have $I^{rr}(\theta) = 1/I_{rr}(\theta)$ for each r, and $I_{rs}(\theta) = I^{rs}(\theta) = 0$ for $r \neq s$. This implies that the above equation is equivalent to (14) with $\phi(\theta)$ defined as in (10).

The following corollary offers a practical method for constructing an asymptotically unbiased prior. This builds on the sufficiency part of the proof of Theorem 5, presented in Section 5.2.

Corollary 7. Suppose the model satisfies the assumptions of Theorem 5. If the integrability condition (14) holds, a prior constructed using the following procedure satisfies (13).

- (i) Fix a constant $(c_1, \ldots, c_p) \in \Theta$ arbitrarily.
- (ii) Define

$$\psi_t(\theta_t, \dots, \theta_n) = \phi_t(c_1, \dots, c_{t-1}, \theta_t, \dots, \theta_n)$$

and compute

$$\tilde{\pi}(\theta) = \exp\left(\sum_{t=1}^{p} \int_{c_t}^{\theta_t} \psi_t(z, \theta_{t+1}, \dots, \theta_p) dz\right).$$

(iii) Take the prior as $\pi(\theta) \propto \tilde{\pi}(\theta)$.

The prior constructed using the procedure in Corollary 7 may often be improper, meaning that its integral over the parameter space diverges to infinity. This does not invalidate the posterior mean, as long as the posterior distribution is proper (i.e., integrable).

In the following section, we present examples of models where an asymptotically unbiased prior can be constructed using the procedure described above.

3. Examples

In this section, we provide examples of asymptotically unbiased priors for various models. We begin with one-parameter models, for which we demonstrate that the integrability condition is always satisfied. Next, we present several simple multi-parameter models, which often have analytical posterior distributions and enable us to verify the exact unbiasedness

of the Bayes estimator. Finally, we consider a more complex model, the nested error regression (NER)/random effects model, which is widely used in many fields, including small area estimation.

3.1. One-parameter models. Suppose the model satisfies the assumptions of Theorem 5 with p=1. In this one-parameter case, since $\phi(\theta)$ is a scalar, the integrability condition is automatically satisfied, ensuring the existence of an asymptotically unbiased prior. Specifically, if the model is i.i.d., $\phi(\theta)$ defined in Corollary 4 is written as

$$\phi(\theta) = \phi_1(\theta) = I^{-1}(\theta)I'(\theta),$$

where $I'(\theta)$ is the derivative of the Fisher information. An asymptotically unbiased prior can be constructed using the method in Corollary 7 as follows:

- (i) Fix a constant $c \in \Theta$ arbitrarily.
- (ii) Define $\psi_1(\theta) = \phi_1(\theta) = I^{-1}(\theta)I'(\theta)$ and compute

$$\tilde{\pi}(\theta) = \exp\left(\int_{c}^{\theta} \psi_{1}(z)dz\right) = \exp\left(\left[\log I(z)\right]_{z=c}^{z=\theta}\right) = \frac{I(\theta)}{I(c)}.$$

(iii) Take the prior as

$$\pi(\theta) \propto I(\theta)$$
.

Thus, when the model is i.i.d., the asymptotically unbiased prior is proportional to the Fisher information itself.³

Example 1 (Binomial distribution). Suppose X_1, \ldots, X_n are i.i.d. random variables following a Bernoulli distribution with parameter $\theta \in (0,1)$. By the previous argument, the asymptotically unbiased prior for θ is given by

$$\pi(\theta) \propto I(\theta) = \frac{1}{\theta(1-\theta)}.$$

Let $\bar{X} = n^{-1} \sum_{i=1}^{n} X_i$ denote the sample mean of the *n* Bernoulli random variables. The posterior density is computed as

$$\pi(\theta|X_1,\ldots,X_n) \propto \theta^{n\bar{X}-1}(1-\theta)^{n(1-\bar{X})-1},$$

which is a beta distribution with parameters $(n\bar{X}, n(1-\bar{X}))$. Thus, the posterior mean of θ is given by

$$\hat{\theta}^B = \bar{X}.$$

which is an unbiased estimator of θ .

3.2. Simple multi-parameter models. In this section, we illustrate the construction of asymptotically unbiased priors for multi-parameter models. We show that, in certain cases, the priors constructed using this method result in Bayes estimators that are exactly unbiased. Furthermore, we provide an example where an asymptotically unbiased prior does not exist.

Example 2 (Mean and variance parameters of normal distribution). Suppose X_1, \ldots, X_n are i.i.d. $N(\mu, \sigma^2)$ random variables with density function

$$f(x|\theta) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right),$$

³Recall that Jeffreys' prior is proportional to the square root of the Fisher information.

where $\theta = (\mu, \sigma^2)$ is the parameter of interest. We construct an asymptotically unbiased prior based on the result of Corollary 4. The Fisher information $I(\theta)$ is given by

$$I(\theta) = \begin{bmatrix} \sigma^{-2} & 0 \\ 0 & \sigma^{-4}/2 \end{bmatrix} = \begin{bmatrix} \theta_2^{-1} & 0 \\ 0 & \theta_2^{-2}/2 \end{bmatrix}.$$

Observe that the Fisher information matrix is diagonal with $I_{11}(\theta)/I_{22}(\theta) = 2\theta_2$. Thus, by Proposition 6 and Theorem 5, the existence of an asymptotically unbiased prior is guaranteed. We can also compute the inverse of the Fisher information matrix and its partial derivatives as

$$I^{-1}(\theta) = \begin{bmatrix} \sigma^2 & 0 \\ 0 & 2\sigma^4 \end{bmatrix}, \quad \frac{\partial}{\partial \theta_1} I(\theta) = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad \text{and} \quad \frac{\partial}{\partial \theta_2} I(\theta) = \begin{bmatrix} -\sigma^{-4} & 0 \\ 0 & -\sigma^{-6} \end{bmatrix},$$

respectively. According to Corollary 4, we compute $\phi_1(\theta)$ and $\phi_2(\theta)$ as

$$\phi_1(\theta) = \sum_{r=1}^2 \sum_{s=1}^2 I^{rs}(\theta) \frac{\partial}{\partial \theta_s} I_{1r}(\theta) = 0 \quad \text{and} \quad \phi_2(\theta) = \sum_{r=1}^2 \sum_{s=1}^2 I^{rs}(\theta) \frac{\partial}{\partial \theta_s} I_{2r}(\theta) = -2\sigma^{-2}.$$

We follow the procedure of Corollary 7 to construct an asymptotically unbiased prior.

- (i) Fix an arbitrary constant c > 0.
- (ii) Define $\psi_1(\theta_1, \theta_2) = 0$ and $\psi_2(\theta_2) = \psi_2(\sigma^2) = -2\sigma^{-2}$. Compute

$$\tilde{\pi}(\theta) := \exp\left(\int_{c}^{\sigma^2} \psi_2(z) dz\right) = \exp\left(-2[\log z]_{z=c}^{z=\sigma^2}\right) = \frac{c^2}{\sigma^4}.$$

(iii) Take $\pi(\theta) \propto \sigma^{-4}$.

Note that the obtained prior is different from Jeffreys' prior since Jeffreys' prior for this model is $\pi_J(\theta) \propto (\det I(\theta))^{1/2} \propto \sigma^{-3}$. Next, we calculate the posterior mean $\hat{\theta}^B$ corresponding to the prior $\pi(\theta) = \pi(\mu, \sigma^2) \propto \sigma^{-4}$. Define

$$\bar{X} = \frac{1}{n} \sum_{i=1}^{n} X_i$$
 and $S = \left(\sum_{i=1}^{n} (X_i - \bar{X})^2\right)^{\frac{1}{2}}$.

Then, the posterior density is written as

$$\pi(\mu, \sigma^2 | X_1, \dots, X_n) = \sqrt{\frac{n}{2\pi}} \frac{1}{\Gamma(\frac{n+1}{2})} \left(\frac{S^2}{2}\right)^{\frac{n+1}{2}} \sigma^{-n-4} \exp\left(-\frac{S^2 + n(\bar{X} - \mu)^2}{2\sigma^2}\right).$$

Therefore, we obtain the posterior mean $\hat{\theta}^B = (\hat{\mu}^B, \hat{\sigma}^{2,B})$ as

$$\hat{\mu}^B = \int_0^\infty \int_{-\infty}^\infty \mu \pi(\mu, \sigma^2 | X_1, \dots, X_n) d\mu d\sigma^2 = \bar{X},$$

$$\hat{\sigma}^{2,B} = \int_0^\infty \int_{-\infty}^\infty \sigma^2 \pi(\mu, \sigma^2 | X_1, \dots, X_n) d\mu d\sigma^2 = \frac{S^2}{n-1}.$$

This estimator is an exactly unbiased estimator of $\theta = (\mu, \sigma^2)$.

Example 3 (Location and scale parameters of normal distribution). We consider the same model as in Example 2 but we are interested in the estimation of $\theta = (\mu, \sigma)$ instead of (μ, σ^2) . In this case, the Fisher information $I(\theta)$ is given by

$$I(\theta) = \begin{bmatrix} \sigma^{-2} & 0 \\ 0 & 2\sigma^{-2} \end{bmatrix} = \theta_2^{-2} \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}.$$

Since the Fisher information matrix is diagonal with $I_{11}(\theta)/I_{22}(\theta) = 1/2$, the existence of an asymptotically unbiased prior is guaranteed by Proposition 6 and Theorem 5. Following Corollary 7 as in the previous example, we obtain the asymptotically unbiased prior

$$\pi(\theta) = \pi(\mu, \sigma) \propto \sigma^{-2}$$
.

Note that the asymptotically unbiased prior we constructed above corresponds to Jeffreys' prior, unlike the parametrization of Example 2. The posterior density is given by

$$\pi(\mu, \sigma | X_1, \dots, X_n) = \sqrt{\frac{2n}{\pi}} \frac{1}{\Gamma\left(\frac{n}{2}\right)} \left(\frac{S^2}{2}\right)^{\frac{n}{2}} \sigma^{-n-2} \exp\left(-\frac{S^2 + n(\bar{X} - \mu)^2}{2\sigma^2}\right),$$

and the posterior mean of $\theta = (\mu, \sigma)$ is computed as

$$\hat{\mu}^{B} = \int_{0}^{\infty} \int_{-\infty}^{\infty} \mu \pi(\mu, \sigma^{2} | X_{1}, \dots, X_{n}) d\mu d\sigma^{2} = \bar{X},$$

$$\hat{\sigma}^{B} = \int_{0}^{\infty} \int_{-\infty}^{\infty} \sigma^{2} \pi(\mu, \sigma^{2} | X_{1}, \dots, X_{n}) d\mu d\sigma^{2} = \frac{\Gamma\left(\frac{n-1}{2}\right)}{\sqrt{2}\Gamma\left(\frac{n}{2}\right)} S.$$

Again, these are the exact unbiased estimators of μ and σ .

Example 4 (Location and scale parameters of gamma distribution). Suppose X_1, \ldots, X_n are i.i.d. $\Gamma(\theta_1, \theta_2)$ random variables with density function

$$f(x|\theta) = \frac{1}{\Gamma(\theta_1)\theta_2^{\theta_1}} x^{\theta_1 - 1} \exp\left(-\frac{x}{\theta_2}\right),$$

where $\theta = (\theta_1, \theta_2)$ is the parameter of interest. We look for a prior that satisfies (10) in Corollary 4. Let Λ denote the derivative of the digamma function, that is, $\Lambda(\theta_1) = \frac{d^2}{d\theta_1^2} \log \Gamma(\theta_1)$. Then, we can write the Fisher information $I(\theta)$ as

$$I(\theta) = \begin{bmatrix} \Lambda(\theta_1) & \theta_2^{-1} \\ \theta_2^{-1} & \theta_1 \theta_2^{-2} \end{bmatrix}.$$

The inverse of the Fisher information matrix and its partial derivatives can be computed as

$$\begin{split} I^{-1}(\theta) &= \frac{1}{\theta_1 \Lambda(\theta_1) - 1} \begin{bmatrix} \theta_1 & -\theta_2 \\ -\theta_2 & \theta_2^2 \Lambda(\theta_1) \end{bmatrix}, \\ \frac{\partial}{\partial \theta_1} I(\theta) &= \begin{bmatrix} \frac{d}{d\theta_1} \Lambda(\theta_1) & 0 \\ 0 & \theta_2^{-2} \end{bmatrix}, \quad \frac{\partial}{\partial \theta_2} I(\theta) &= \begin{bmatrix} 0 & -\theta_2^{-2} \\ -\theta_2^{-2} & -2\theta_1\theta_2^{-3} \end{bmatrix}, \end{split}$$

respectively. Therefore, according to Corollary 4, we compute $\phi(\theta) = (\phi_1(\theta), \phi_2(\theta))$ as

$$\phi_1(\theta) = \sum_{r=1}^2 \sum_{s=1}^2 I^{rs}(\theta) \frac{\partial}{\partial \theta_s} I_{1r}(\theta) = (\theta_1 \Lambda(\theta_1) - 1)^{-1} \left(\theta_1 \frac{d}{d\theta_1} \Lambda(\theta_1) - \Lambda(\theta_1) \right),$$

$$\phi_2(\theta) = \sum_{r=1}^2 \sum_{s=1}^2 I^{rs}(\theta) \frac{\partial}{\partial \theta_s} I_{2r}(\theta) = (\theta_1 \Lambda(\theta_1) - 1)^{-1} (-2\theta_1 \theta_2^{-1} \Lambda(\theta_1)).$$

While the derivative of ϕ_1 with respect to θ_2 is zero, the derivative of ϕ_2 with respect to θ_1 is not zero in general. This implies that the integrability condition is not satisfied. Hence, an asymptotically unbiased prior independent of n does not exist for this model, according to Theorem 5 and Remark 5.

Example 5 (Linear regression model). Consider the model

$$y|X \sim N(X\beta, \sigma^2 I_n),$$

where $y = [y_1 \dots y_n]^{\top}$ is an *n*-dimensional random vector of response variables, $X = [x_1 \dots x_n]^{\top}$ is an $n \times p$ non-random matrix of explanatory variables and $\beta \in \mathbb{R}^p$ is the coefficient vector. We consider the estimation problem of the parameter $\theta = (\theta_1, \dots, \theta_p, \theta_{p+1}) = (\beta, \sigma^2) \in \mathbb{R}^p \times (0, \infty)$. Since the model is not i.i.d., we cannot apply the result of Hartigan (1965) given in Corollary 4. Hence, we apply Corollary 3 to obtain an asymptotically unbiased prior.

The log-likelihood is given by

$$\ell_n(\theta) = -\frac{n}{2}\log\sigma^2 - \frac{1}{2\sigma^2}(y - X\beta)^{\top}(y - X\beta) + (const.).$$

We compute $H(\theta)$ and $I(\theta)$, that are found to be identical in this case, as

$$H(\theta) = I(\theta) = \begin{bmatrix} \sigma^{-2} \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} x_i x_i^{\top} & 0\\ 0 & \sigma^{-4}/2 \end{bmatrix}.$$

To obtain $\phi(\theta)$ in Corollary 3, we first compute $J_{tr,s}(\theta)$. For any $r, s \leq p$ and $t \leq p+1$, we get

$$J_{rs,t}(\theta) = 0$$

and for other cases, we have

$$J_{r(p+1),s}(\theta) = \begin{cases} -\sigma^{-4} \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} x_{ir} x_{is}, & r, s \le p, \\ 0, & r \le p, s = p+1, \\ 3\sigma^{-4} \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} x_{ir}, & r = p+1, s \le p, \\ -\sigma^{-6}, & r = s = p+1. \end{cases}$$

Next, we compute $K_{rst}(\theta)$. We get

$$K_{rst}(\theta) = 0, r, s, t \le p,$$

$$K_{rs(p+1)}(\theta) = \sigma^{-4} \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} x_{ir} x_{is}, r, s \le p,$$

$$K_{r(p+1)(p+1)}(\theta) = 0, r \le p,$$

$$K_{(p+1)(p+1)(p+1)}(\theta) = 2\sigma^{-6}.$$

By substituting these values into the definition of ϕ_t given in Corollary 3, we obtain

$$\phi(\theta) = \begin{bmatrix} 0 & \cdots & 0 & -2\sigma^{-2} \end{bmatrix}^{\top}.$$

It is easy to verify that $\phi_t(\theta)$ is zero for all $t \leq p$, and hence, the integrability condition is satisfied. Thus, we can construct an asymptotically unbiased prior using the construction method in Corollary 7 as

$$\pi(\theta) \propto \exp\left(-\int_{c}^{\sigma^{2}} (-2)z^{-1}dz\right) \propto \sigma^{-4}.$$

The corresponding posterior distribution can be described as

$$\beta | \sigma^2, y \sim N\left(\hat{\beta}^{OLS}, \sigma^2(X^\top X)^{-1}\right),$$

$$\sigma^2 | y \sim \text{InverseGamma}\left(\frac{n-p}{2} + 1, \frac{1}{2}y^\top (I_n - X(X^\top X)^{-1}X^\top)y\right),$$

where $\hat{\beta}^{OLS} = (X^{\top}X)^{-1}X^{\top}y$ is the ordinary least squares (OLS) estimator. We can compute the posterior mean of β as

$$\hat{\beta}^B = \hat{\beta}^{OLS},$$

and the posterior mean of σ^2 as

$$\hat{\sigma}^{2,B} = \frac{1}{n-p} \left(y^{\top} (I_n - X(X^{\top}X)^{-1}X^{\top}) y \right)$$

when n > p. It can be shown that $(\hat{\beta}^B, \hat{\sigma}^{2,B})$ is an unbiased estimator of $\theta = (\beta, \sigma^2)$.

3.3. Nested error regression (NER) model. We now consider the NER/random effects model, which is widely used for analyzing hierarchically structured data in many fields, such as small area estimation. In this model, the response variable y_i is expressed as a linear function of the covariates x_i , a random effect v_i , and error terms ϵ_{ij} for $j \leq n_i$. The random effect is shared within each group i, while the error terms ϵ_{ij} are independent across units j within the same area/group. For more details and applications of the NER model in small area estimation, please refer to, for example, Sugasawa and Kubokawa (2023) and Rao and Molina (2015).

Example 6 (NER model). We consider the NER model

$$y_{ij} = x_{ij}^{\top} \beta + v_i + \epsilon_{ij}, \quad i = 1, ..., m, \quad j = 1, ..., n_i,$$

 $v_i \sim N(0, \tau^2), \quad \epsilon_{ij} \sim N(0, \sigma^2), \quad v_i \perp \epsilon_{ij},$

where y_{ij} is the response variable for each unit j in area i, x_{ij} is a p-dimensional non-random covariate vector, $\beta \in \mathbb{R}^p$ is the coefficient vector, v_i represents the random effect for area i, and ϵ_{ij} is the error term. If we define

$$y_i = \begin{bmatrix} y_{i1} \\ \vdots \\ y_{in_i} \end{bmatrix}, \ x_i = \begin{bmatrix} x_{i1}^\top \\ \vdots \\ x_{in_i}^\top \end{bmatrix}, \ \epsilon_i = \begin{bmatrix} \epsilon_{i1} \\ \vdots \\ \epsilon_{in_i} \end{bmatrix},$$

then we can express the model in vector form as

$$y_i = x_i \beta + v_i \iota_{n_i} + \epsilon_i, \quad i = 1, \dots, m,$$

where ι_{n_i} is an n_i -dimensional column vector of ones. The data are assumed to be independent across areas i. We consider the asymptotic setting where $m \to \infty$, with n_i fixed. The parameter of interest is

$$\theta = (\theta_1, \dots, \theta_p, \theta_{p+1}, \theta_{p+2}) = (\beta, \tau^2, \sigma^2) \in \Theta = \mathbb{R}^p \times (0, \infty)^2.$$

If we write

$$V_i = V_i(\theta) = \tau^2 \iota_{n_i} \iota_{n_i}^\top + \sigma^2 I_{n_i}$$

the likelihood function is given by

$$f_m(y_1, \dots, y_m | x_1, \dots, x_m, \theta) = \prod_{i=1}^m \left[(2\pi)^{-\frac{n_i}{2}} \det(V_i)^{-\frac{1}{2}} \exp\left(-\frac{1}{2} (y_i - x_i \beta)^\top V_i^{-1} (y_i - x_i \beta)\right) \right].$$

Thus, the log-likelihood function is written as

$$\ell_m(\theta) = -\frac{1}{2} \sum_{i=1}^m \left[\log \det(V_i) + (y_i - x_i \beta)^\top V_i^{-1} (y_i - x_i \beta) \right] + (const.).$$

We obtain an asymptotically unbiased prior using the result of Corollary 3. The first step is to compute $\phi(\theta)$. The expression for $\phi(\theta)$ in Corollary 3 is given by

$$\phi_t(\theta) = -\frac{1}{2} \sum_{r=1}^{p+2} \sum_{s=1}^{p+2} H^{rs}(\theta) A_{rst}(\theta) \quad (t = 1, \dots, p+2),$$

where $A_{rs}(\theta)$ is defined as

$$A_{rs}(\theta) = \begin{bmatrix} K_{1rs}(\theta) + 2J_{1r,s}(\theta) \\ \vdots \\ K_{prs}(\theta) + 2J_{pr,s}(\theta) \end{bmatrix} + \sum_{t=1}^{p+2} \sum_{u=1}^{p+2} H^{tu}(\theta) I_{su}(\theta) \begin{bmatrix} K_{rt1}(\theta) \\ \vdots \\ K_{rtp}(\theta) \end{bmatrix}$$

and $H(\theta), I(\theta), J(\theta)$ and $K(\theta)$ are defined such that Assumption 2 is satisfied. By a simple calculation, we get $H(\theta)$ and $I(\theta)$ as

$$H(\theta) = I(\theta) = \lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} \begin{bmatrix} x_i^{\top} V_i^{-1} x_i & 0 & 0 \\ 0 & 2^{-1} n_i^2 (\sigma^2 + n_i \tau)^{-2} & 2^{-1} n_i (\sigma^2 + n_i \tau)^{-2} \\ 0 & 2^{-1} n_i (\sigma^2 + n_i \tau)^{-2} & 2^{-1} \text{Tr}(V_i^{-2}) \end{bmatrix}.$$

This means that the inverse of this matrix is represented as

$$H^{-1}(\theta) = I^{-1}(\theta) = \begin{bmatrix} \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} x_i^{\top} V_i^{-1} x_i \right)^{-1} & 0\\ 0 & W(\theta) \end{bmatrix},$$

where

$$W(\theta) = w(\theta) \lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} \begin{bmatrix} \text{Tr}(V_i^{-2}) & -n_i(\sigma^2 + n_i\tau)^{-2} \\ -n_i(\sigma^2 + n_i\tau)^{-2} & n_i^2(\sigma^2 + n_i\tau)^{-2} \end{bmatrix}$$

and

$$w^{-1}(\theta) = 2^{-1} \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} n_i^2 (\sigma^2 + n_i \tau^2)^{-2} \right) \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} \operatorname{Tr}(V_i^{-2}) \right)$$
$$- 2^{-1} \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} n_i (\sigma^2 + n_i \tau^2)^{-2} \right)^2.$$

Since $H(\theta)$ and $I(\theta)$ are identical, by Remark 1, we have

$$\phi_t(\theta) = -\sum_{r=1}^{p+2} \sum_{s=1}^{p+2} H^{rs}(\theta) (K_{trs}(\theta) + J_{tr,s}(\theta)), \quad t = 1, \dots, p+2.$$

Furthermore, since $H^{-1}(\theta)$ is block diagonal, we have

$$\phi_t(\theta) = -\sum_{r=1}^p \sum_{s=1}^p H^{rs}(\theta) (K_{trs}(\theta) + J_{tr,s}(\theta)) - \sum_{r=p+1}^{p+2} \sum_{s=p+1}^{p+2} H^{rs}(\theta) (K_{trs}(\theta) + J_{tr,s}(\theta))$$
 (15)

for t = 1, ..., p + 2. Thus, we only need to compute $K_{rst}(\theta)$, $J_{tr,s}(\theta)$ for all t = 1, ..., p + 2 and $(r,s) \in \{1,...,p\}^2$, or $(r,s) \in \{p+1,p+2\}^2$. This can be done by a simple calculation; for $r,s \leq p$, we get

$$J_{tr,s}(\theta) = \begin{cases} 0, & t \leq p, \\ -\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^2 + n_i \tau^2)^{-2} \iota_{n_i}^{\top} x_{i \cdot r} x_{i \cdot s}^{\top} \iota_{n_i}, & t = p+1, \\ -\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} x_{i \cdot r}^{\top} V_i^{-2} x_{i \cdot s}^{\top}, & t = p+2, \end{cases}$$

and

$$K_{rst}(\theta) = -J_{tr,s}(\theta),$$

where $x_{i,r}$ is the r-th column vector of x_i . For $r, s \in \{p+1, p+2\}$, we have

$$J_{t(p+1),p+1}(\theta) = \begin{cases} 0, & t \leq p, \\ -\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^2 + n_i \tau^2)^{-3} n_i^3, & t = p+1, \\ -\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^2 + n_i \tau^2)^{-3} n_i^2, & t = p+2, \end{cases}$$

$$K_{(p+1)(p+1)t}(\theta) = -2J_{t(p+1),p+1}(\theta),$$

$$J_{t(p+1),p+2}(\theta) = \begin{cases} 0, & t \leq p, \\ -\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^2 + n_i \tau^2)^{-3} n_i^2, & t = p+1, \\ -\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^2 + n_i \tau^2)^{-3} n_i, & t = p+2, \end{cases}$$

$$K_{(p+1)(p+2)t}(\theta) = -2J_{t(p+1),p+2}(\theta),$$

$$J_{t(p+2),p+1}(\theta) = J_{t(p+1),p+2}(\theta),$$

$$K_{(p+2)(p+1)t}(\theta) = K_{(p+1)(p+2)t}(\theta),$$

$$J_{t(p+2),p+2}(\theta) = \begin{cases} 0, & t \leq p, \\ -\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^2 + n_i \tau^2)^{-3} n_i, & t = p+1, \\ -\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^2 + n_i \tau^2)^{-3} n_i, & t = p+1, \end{cases}$$

$$-\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} \mathrm{Tr}(V_i^{-3}), & t = p+2, \end{cases}$$

$$K_{(p+2)(p+2)t}(\theta) = -2J_{t(p+2),p+2}(\theta).$$

Substituting these values into (15) yields $\phi_t(\theta) = 0$ for $t \leq p$ and

$$-w^{-1}(\theta)\phi_{p+1}(\theta) = \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} \text{Tr}(V_{i}^{-2})\right) \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-3} n_{i}^{3}\right)$$

$$-2 \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-3} n_{i}^{2}\right) \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-2} n_{i}\right)$$

$$+ \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-3} n_{i}\right) \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-2} n_{i}^{2}\right),$$

$$-w^{-1}(\theta)\phi_{p+2}(\theta) = \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} \text{Tr}(V_{i}^{-2})\right) \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-3} n_{i}^{2}\right)$$

$$-2 \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-3} n_{i}\right) \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-2} n_{i}\right)$$

$$+ \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} \text{Tr}(V_{i}^{-3})\right) \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-2} n_{i}^{2}\right).$$

It can be verified that $\phi_{p+1}(\theta)$ and $\phi_{p+2}(\theta)$ are expressed as

$$\phi_{p+1}(\theta) = \frac{\partial}{\partial \tau^2} \log g(\tau^2, \sigma^2)$$
 and $\phi_{p+2}(\theta) = \frac{\partial}{\partial \sigma^2} \log(\sigma^{-4}g(\tau^2, \sigma^2))$

where $g(\tau^2, \sigma^2)$ is given by

$$g(\tau^{2}, \sigma^{2}) = \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-2} n_{i}\right) \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-2} n_{i} (n_{i} - 1)\right) \sigma^{4}$$

$$+ 2 \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-2} n_{i}^{2}\right) \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-2} n_{i} (n_{i} - 1)\right) \sigma^{2} \tau^{2}$$

$$+ \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-2} n_{i}^{2}\right) \left(\lim_{m \to \infty} \frac{1}{m} \sum_{i=1}^{m} (\sigma^{2} + n_{i}\tau^{2})^{-2} n_{i}^{2} (n_{i} - 1)\right) \tau^{4}.$$

This representation of $\phi(\theta)$ implies that the integrability condition (14) is satisfied, and thus an asymptotically unbiased prior exists.

Finally, we compute the asymptotically unbiased prior according to Corollary 7. For arbitrary constants $c_1, c_2 > 0$, we define

$$\psi_{p+1}(\theta_{p+1}) = \frac{\partial}{\partial \tau^2} \log g(\tau^2, \sigma^2), \quad \psi_{p+2}(\theta_{p+2}) = \frac{\partial}{\partial \sigma^2} \log(\sigma^{-4}g(c_1, \sigma^2)).$$

Then, the asymptotically unbiased prior is obtained as

$$\pi(\theta) \propto \exp\left(\int_{c_1}^{\tau^2} \psi_{p+1}(z)dz + \int_{c_2}^{\sigma^2} \psi_{p+2}(z)dz\right) \propto \sigma^{-4}g(\tau^2, \sigma^2).$$
 (16)

Example 7 (Balanced NER model). We consider the case where $n_i \equiv n$ for all i in the previous example. In this case, the calculation of the limit terms in $g(\tau^2, \sigma^2)$ is simplified, and $g(\tau^2, \sigma^2)$ reduces to

$$g(\tau^2, \sigma^2) = \frac{n^2(n-1)}{(\sigma^2 + n\tau^2)^4} \sigma^4 + 2 \frac{n^3(n-1)}{(\sigma^2 + n\tau^2)^4} \sigma^2 \tau^2 + \frac{n^4(n-1)}{(\sigma^2 + n\tau^2)^4} \tau^2$$
$$= \frac{n^2(n-1)}{(\sigma^2 + n\tau^2)^2}.$$

Therefore, the asymptotically unbiased prior in (16) simplifies to

$$\pi(\theta) \propto \sigma^{-4}(\sigma^2 + n\tau^2)^{-2}.\tag{17}$$

The corresponding posterior distribution of θ is given by

$$\pi(\theta|\mathcal{D}) \propto (\sigma^2)^{-2 - \frac{(n-1)m}{2}} (\sigma^2 + n\tau^2)^{-2 - \frac{m}{2}} \exp\left(-\frac{1}{2} \sum_{i=1}^m (y_i - x_i \beta)^\top V^{-1} (y_i - x_i \beta)\right), \quad (18)$$

where $V = \tau^2 \iota_n \iota_n^\top + \sigma^2 I_n$ is the covariance matrix of $y_i - x_i \beta$ and $\mathcal{D} = (y_i, x_i)_{1 \leq i \leq m}$ represents the data. The posterior propriety of the prior in (17) is discussed in Appendix A.

We demonstrate an MCMC algorithm to obtain samples from the posterior distribution in (18). Instead of directly sampling from $\pi(\theta|\mathcal{D})$, we consider a transformation of θ defined as

$$\bar{\theta} = (\beta, \rho, \sigma^2), \quad \rho = \frac{\sigma^2}{\sigma^2 + n\tau^2} \in (0, 1).$$

The corresponding posterior distribution of $\bar{\theta}$ is given by

$$\bar{\pi}(\bar{\theta}|\mathcal{D}) \propto (\sigma^2)^{-3-\frac{nm}{2}} \rho^{\frac{m}{2}} \exp\left(-\frac{1}{2} \sum_{i=1}^{m} (y_i - x_i \beta)^{\top} \bar{V}^{-1} (y_i - x_i \beta)\right),$$
 (19)

where \bar{V}^{-1} is defined as $\bar{V}^{-1} = \sigma^{-2}(I_n - (1 - \rho)\iota_n\iota_n^{\top}/n)$. For posterior inference, we first sample $\bar{\theta}$ from the distribution $\bar{\pi}(\bar{\theta}|\mathcal{D})$, and subsequently recover θ .

Sampling from the posterior distribution (19) is performed using a Gibbs sampler. We can show that the full conditional distributions of β , ρ , and σ^2 are given by

$$\beta | \rho, \sigma^{2}, \mathcal{D} \sim N \left(\left(\sum_{i=1}^{m} x_{i}^{\top} \bar{V}^{-1} x_{i} \right)^{-1} \left(\sum_{i=1}^{m} x_{i}^{\top} \bar{V}^{-1} y_{i} \right), \left(\sum_{i=1}^{m} x_{i}^{\top} \bar{V}^{-1} x_{i} \right)^{-1} \right),$$

$$\rho | \beta, \sigma^{2}, \mathcal{D} \sim \text{Truncated Gamma} \left(\frac{m}{2} + 1, \frac{1}{2n\sigma^{2}} \sum_{i=1}^{m} (y_{i} - x_{i}\beta)^{\top} \iota_{n} \iota_{n}^{\top} (y_{i} - x_{i}\beta) \right)$$

$$\text{truncated on } (0, 1),$$

$$\sigma^{2} | \beta, \rho, \mathcal{D} \sim \text{Inverse Gamma} \left(\frac{nm}{2} + 2, \frac{1}{2} \sum_{i=1}^{m} (y_{i} - x_{i}\beta)^{\top} \left(I_{n} - \frac{1 - \rho}{n} \iota_{n} \iota_{n}^{\top} \right) (y_{i} - x_{i}\beta) \right).$$

In the simulation studies presented in Section 4, we utilize the inverse transform sampler to draw samples of ρ from the truncated gamma distribution.




FIGURE 1. Bias of the posterior mean when the true parameter is $\beta_1 = \beta_2 =$ $\tau^2 = \sigma^2 = 1.$

4. Simulation studies on the NER model

Most of the examples presented in Section 3 yield analytical posterior distributions and exactly unbiased Bayes estimators. However, for the NER model, deriving an analytical posterior distribution is not feasible, and consequently, the properties of the Bayes estimator remain unknown. To address this issue, we conduct simulation studies to evaluate the bias of the Bayes estimator under the asymptotically unbiased prior we developed. This evaluation is particularly critical when the sample size (that is, the number of areas/groups) is small, which is a common scenario in practical applications of small area estimation. We compare the proposed prior with several alternative priors and assess whether the proposed prior results in smaller bias when the sample size is limited. All simulation codes are available at https://github.com/manasakai/second-order-unbiased-NER.

We consider the balanced NER model described in Example 7. For simplicity, we assume that β is two-dimensional. The true parameter values are set under the following two scenarios:

•
$$\beta_1 = \beta_2 = \tau^2 = \sigma^2 = 1$$
.

•
$$\beta_1 = \beta_2 = \tau^2 = \sigma^2 = 1$$
.
• $\beta_1 = \beta_2 = 1, \tau^2 = 0.5, \sigma^2 = 4$.

As will be demonstrated, the simulation results depend on the true parameter settings. Further details on the simulation settings are provided in Appendix B.

We compare the following three priors.

• [AU] The asymptotically unbiased prior:

$$\pi(\theta) \propto [\sigma^2(\sigma^2 + n\tau^2)]^{-2}.$$




FIGURE 2. Bias of the posterior mean when the true parameter is $\beta_1 = \beta_2 = 1$, $\tau^2 = 0.5$, $\sigma^2 = 4$.

• [JF] Jeffreys' prior for the variance components with a flat prior for the regression coefficients:⁴

$$\pi(\theta) \propto [\sigma^2(\sigma^2 + n\tau^2)]^{-1}.$$

• [DG] Inverse-gamma priors for the variance components with a flat prior for the regression coefficients:⁵

$$\pi(\beta) \propto 1$$
, $\tau^2 \sim \mathrm{IG}(a_\tau, b_\tau)$, $\sigma^2 \sim \mathrm{IG}(a_\sigma, b_\sigma)$.

Here, we set $a_{\tau} = b_{\tau} = a_{\sigma} = b_{\sigma} = 5$.

Figures 1 and 2 illustrate the computed absolute bias of the posterior mean under the two parameter settings. As shown in these figures, the bias of β remains relatively small and stable across priors and sample sizes. However, for τ^2 and σ^2 , the bias is highly sensitive to the choice of prior, particularly when the sample size is small (m = 10, 32). For instance, the proposed prior [AU] exhibits smaller bias for τ^2 under both parameter settings, while [JF] shows an advantage in estimating σ^2 in the second setting $(\tau^2 = 0.5, \sigma^2 = 4)$. These findings suggest that while the [AU] prior provides robust performance for variance components in general, alternative priors, such as [JF], may excel in specific parameter settings.

⁴A similar composition of prior is employed in Tiao and Tan (1965), where the random effects model of the form $y_{ij} = \mu + \alpha_i + \epsilon_{ij}$ is considered. In their approach, a flat prior is used for μ , while Jeffreys' prior was used for the variance components.

⁵This prior was proposed by Datta and Ghosh (1991).

The evaluation of the mean squared error (MSE) and the coverage probability can be found in Appendix B. These additional metrics provide further insights into the performance of the proposed prior and its comparisons.

5. Proofs

In this section, we present the remaining proofs of the results we introduced in Section 2.

5.1. Proofs of Theorem 1 and Corollary 3.

5.1.1. Proof of Theorem 1. The proof strategy is to evaluate $\hat{\theta}^B - \hat{\theta}^{ML}$ and $\hat{\theta}^{ML} - \theta$ separately, and combine them to obtain the decomposition (3).

We sometimes include 'dots' instead of indices $r, s, t \in \{1, ..., p\}$ to denote a vector or a matrix composed of $K_{n,rst}$. For example, we write

$$K_{n,r\cdot t}(\theta) = \begin{bmatrix} K_{n,r1t}(\theta) \\ \vdots \\ K_{n,rpt}(\theta) \end{bmatrix} \text{ and } K_{n,r\cdot t}(\theta) = \begin{bmatrix} K_{n,r11}(\theta) & \cdots & K_{n,r1p}(\theta) \\ \vdots & \ddots & \vdots \\ K_{n,rp1}(\theta) & \cdots & K_{n,rpp}(\theta) \end{bmatrix}.$$

We first evaluate $\hat{\theta}^B - \hat{\theta}^{ML}$ using Laplace's method.

Lemma 8. Suppose the model satisfies Assumption 1. Then, we can decompose the difference between the Bayes estimator and the MLE as

$$\hat{\theta}^B - \hat{\theta}^{ML} = \frac{1}{n} H_n^{-1}(\theta) \left(\frac{\partial}{\partial \theta} \log \pi(\theta) + \frac{1}{2} \sum_{r=1}^p \sum_{s=1}^p H_n^{rs}(\theta) K_{n,rs}(\theta) \right) + o_p \left(\frac{1}{n} \right). \tag{20}$$

Proof. Noting that $\pi(\theta)$ is null if $\theta \notin \Theta$, we can write the Bayes estimator given in (2) as

$$\hat{\theta}^B = \frac{\int_{\mathbb{R}^p} \theta \pi(\theta) \exp(\ell_n(\theta)) d\theta}{\int_{\mathbb{R}^p} \pi(\theta) \exp(\ell_n(\theta)) d\theta}.$$

Here, we can expand $\pi(\theta)$ around $\hat{\theta}^{ML}$ as

$$\pi(\theta) = \pi(\hat{\theta}^{ML}) + \dot{\pi}(\hat{\theta}^{ML})^{\top}(\theta - \hat{\theta}^{ML}) + O_p(n^{-1}),$$

where $\dot{\pi}(\hat{\theta}^{ML}) = \frac{\partial}{\partial \theta} \pi(\theta) \Big|_{\theta = \hat{\theta}^{ML}}$ is the first order derivative of $\pi(\theta)$ at $\hat{\theta}^{ML}$. Similarly, we can expand $\ell_n(\theta)$ around $\hat{\theta}^{ML}$ as

$$\ell_n(\theta) = \ell_n(\hat{\theta}^{ML}) - \frac{n}{2}(\theta - \hat{\theta}^{ML})^{\top} H_n(\hat{\theta}^{ML}) (\theta - \hat{\theta}^{ML}) + \frac{1}{6} \left(\sum_{r=1}^p (\theta_r - \tilde{\theta}_r) \frac{\partial}{\partial \theta_r} \right)^3 \ell_n(\theta) \Big|_{\theta = \tilde{\theta}} + O_p \left(\frac{1}{n} \right),$$

where we used (1) to eliminate the first order term. By the positive definiteness of $\hat{H}_n := H_n(\hat{\theta}^{ML})$, there exists a symmetric and positive definite matrix $\hat{G}_n := G_n(\hat{\theta}^{ML})$ that satisfies $n\hat{H}_n = \hat{G}_n^2$. Define $\eta = \hat{G}_n(\theta - \hat{\theta}^{ML})$, and for simplicity, we write $\pi(\hat{\theta}^{ML})$ as $\hat{\pi}$. Then, we can evaluate the Bayes estimator as

$$\hat{\theta}^B - \hat{\theta}^{ML}$$

$$= \hat{G}_n^{-1} \frac{\int_{\mathbb{R}^p} \eta \left(\hat{\pi} + \dot{\pi} (\hat{\theta}^{ML})^\top \hat{G}_n^{-1} \eta + O_p(n^{-1}) \right) \exp \left(-\frac{1}{2} \eta^\top \eta + g(\eta) + O_p(n^{-1}) \right) d\eta}{\int_{\mathbb{R}^p} \left(\hat{\pi} + \dot{\pi} (\hat{\theta}^{ML})^\top \hat{G}_n^{-1} \eta + O_p(n^{-1}) \right) \exp \left(-\frac{1}{2} \eta^\top \eta + g(\eta) + O_p(n^{-1}) \right) d\eta}, \quad (21)$$

where we defined $g(\eta) = \frac{1}{6} \left(\sum_{r=1}^p \sum_{s=1}^p \hat{G}_n^{rs} \eta_s \frac{\partial}{\partial \theta_r} \right)^3 \ell_n(\theta) \Big|_{\theta = \hat{\theta}^{ML}}$. Noting that expanding $\exp(g(\eta) + O_p(n^{-1}))$ yields

$$\exp(g(\eta) + O_p(n^{-1})) = 1 + g(\eta) + O_p(n^{-1}),$$

we have

$$(\hat{\pi} + \dot{\pi}(\hat{\theta}^{ML})^{\top} \hat{G}_n^{-1} \eta + O_p(n^{-1})) \exp(g(\eta) + O_p(n^{-1}))$$

$$= \hat{\pi} + \dot{\pi}(\hat{\theta}^{ML})^{\top} \hat{G}_n^{-1} \eta + \hat{\pi}g(\eta) + O_p(n^{-1}).$$

Thus, (21) can be written as

$$\hat{\theta}^B - \hat{\theta}^{ML}$$

$$= \hat{G}_{n}^{-1} \frac{\int_{\mathbb{R}^{p}} \eta \exp\left(-\frac{1}{2}\eta^{\top}\eta\right) \left(\dot{\pi}(\hat{\theta}^{ML})^{\top} \hat{G}_{n}^{-1} \eta + \hat{\pi}g(\eta) + O_{p}(n^{-1})\right) d\eta}{(2\pi)^{p/2}\hat{\pi} + O_{p}(n^{-1})}$$

$$= \frac{\hat{G}_{n}^{-1}}{\hat{\pi}} \int_{\mathbb{R}^{p}} \frac{\eta \exp\left(-\frac{1}{2}\eta^{\top}\eta\right)}{(2\pi)^{p/2}} \left(\dot{\pi}(\hat{\theta}^{ML})^{\top} \hat{G}_{n}^{-1} \eta + \hat{\pi}g(\eta) + O_{p}(n^{-1})\right) d\eta + o_{p}\left(\frac{1}{n}\right)$$

$$= \frac{\hat{G}_{n}^{-1}}{\hat{\pi}} \left(\int_{\mathbb{R}^{p}} \frac{\eta \eta^{\top} \exp\left(-\frac{1}{2}\eta^{\top}\eta\right)}{(2\pi)^{p/2}} d\eta\right) \hat{G}_{n}^{-1} \dot{\pi}(\hat{\theta}^{ML})$$

$$+ \hat{G}_{n}^{-1} \int_{\mathbb{R}^{p}} \frac{\eta \exp\left(-\frac{1}{2}\eta^{\top}\eta\right)}{(2\pi)^{p/2}} g(\eta) d\eta + o_{p}\left(\frac{1}{n}\right)$$

$$= \frac{1}{n} \hat{H}_{n}^{-1} \frac{\partial}{\partial \theta} \log \pi(\theta) \Big|_{\theta = \hat{\theta}^{ML}} + \hat{G}_{n}^{-1} \int_{\mathbb{R}^{p}} \frac{\eta \exp\left(-\frac{1}{2}\eta^{\top}\eta\right)}{(2\pi)^{p/2}} g(\eta) d\eta + o_{p}\left(\frac{1}{n}\right). \tag{22}$$

The third equality follows from the symmetry of \hat{G}_n^{-1} , and the last equality follows from $\int_{\mathbb{R}^p} (2\pi)^{-p/2} \eta \eta^{\top} \exp\left(-\frac{1}{2}\eta^{\top}\eta\right) d\eta$ being the identity matrix. The integral in the second term of (22) is calculated as

$$\frac{1}{2}\hat{G}_n^{-1} \sum_{r=1}^p \sum_{s=1}^p \hat{H}_n^{rs} K_{n,rs}(\hat{\theta}^{ML})$$

by the definition of $g(\eta)$. Thus, we obtain

$$\hat{\theta}^B - \hat{\theta}^{ML} = \frac{1}{n} \hat{H}_n^{-1} \left(\frac{\partial}{\partial \theta} \log \pi(\theta) + \frac{1}{2} \sum_{r=1}^p \sum_{s=1}^p \hat{H}_n^{rs} K_{n,rs}(\hat{\theta}^{ML}) \right) + o_p \left(\frac{1}{n} \right). \tag{23}$$

By conditions (i) and (ii) of Assumption 1, \hat{H}_n can be approximated as

$$\hat{H}_n = H_n(\theta) + O_n(n^{-1/2})$$

and consequently by condition (ii) of Assumption 1, we have

$$\hat{H}_n^{-1} = H_n^{-1}(\theta) + O_p(n^{-1/2}).$$

Similarly, by conditions (i) and (ii) of Assumption 1, we can approximate $K_{n,rs}(\hat{\theta}^{ML})$ as

$$K_{n,rs}(\hat{\theta}^{ML}) = K_{n,rs}(\theta) + O_p(n^{-1/2}).$$

Substituting these approximations into (23), we obtain (20).

Next, we evaluate $\hat{\theta}^{ML} - \theta$.

Lemma 9. Under Assumption 1, the difference between the MLE and the true parameter can be approximated as

$$\hat{\theta}^{ML} - \theta = \frac{1}{n} H_n^{-1}(\theta) \left(2 \frac{\partial}{\partial \theta} \ell_n(\theta) + \frac{1}{2} \sum_{r=1}^p \sum_{s=1}^p H_n^{rs}(\theta) B_{n,rs}(\theta) \right) + o_p \left(\frac{1}{n} \right),$$

where $B_{n,rs}(\theta)$ is defined as

$$B_{n,rs}(\theta) = 2 \begin{bmatrix} J_{n,1r,s}(\theta) \\ \vdots \\ J_{n,pr,s}(\theta) \end{bmatrix} + \sum_{t=1}^{p} \sum_{u=1}^{p} H_n^{tu}(\theta) I_{su}(\theta) K_{n,r\cdot t}(\theta).$$

Proof. By expanding (1) around the true parameter, we obtain

$$0 = \frac{\partial}{\partial \theta} \ell_n(\theta) - n \left(H_n(\theta) - \frac{1}{2} \sum_{r=1}^p (\hat{\theta}_r^{ML} - \theta_r) K_{n,r..}(\theta) \right) (\hat{\theta}^{ML} - \theta) + o_p(1), \tag{24}$$

or equivalently,

$$\hat{\theta}^{ML} - \theta = \frac{1}{n} \left(H_n(\theta) - \frac{1}{2} \sum_{r=1}^p (\hat{\theta}_r^{ML} - \theta_r) K_{n,r}(\theta) \right)^{-1} \left(\frac{\partial}{\partial \theta} \ell_n(\theta) + o_p(1) \right). \tag{25}$$

Note that by conditions (i) and (ii) of Assumption 1, we have

$$H_{n}(\theta) \left(H_{n}(\theta) - \frac{1}{2} \sum_{r=1}^{p} (\hat{\theta}_{r}^{ML} - \theta_{r}) K_{n,r..}(\theta) \right)^{-1}$$

$$= \left(I_{p} - \frac{1}{2} \sum_{r=1}^{p} (\hat{\theta}_{r}^{ML} - \theta_{r}) K_{n,r..}(\theta) H_{n}^{-1}(\theta) \right)^{-1}$$

$$= I_{p} + \frac{1}{2} \sum_{r=1}^{p} (\hat{\theta}_{r}^{ML} - \theta_{r}) K_{n,r..}(\theta) H_{n}^{-1}(\theta) + O_{p} \left(\frac{1}{n} \right)$$

$$= 2I_{p} - H_{n}(\theta) H_{n}^{-1}(\theta) + \frac{1}{2} \sum_{r=1}^{p} (\hat{\theta}_{r}^{ML} - \theta_{r}) K_{n,r..}(\theta) H_{n}^{-1}(\theta) + O_{p} \left(\frac{1}{n} \right). \tag{26}$$

Furthermore, conditions (i) and (ii) of Assumption 1 and (24) imply

$$0 = \frac{\partial}{\partial \theta} \ell_n(\theta) - nH_n(\theta)(\hat{\theta}^{ML} - \theta) + O_p(1).$$

This can be rewritten as

$$\hat{\theta}^{ML} - \theta = \frac{1}{n} H_n^{-1}(\theta) \frac{\partial}{\partial \theta} \ell_n(\theta) + O_p\left(\frac{1}{n}\right),$$

or equivalently, for each $r \in \{1, \dots, p\}$,

$$\hat{\theta}_r^{ML} - \theta_r = \frac{1}{n} \sum_{s=1}^p H_n^{rs}(\theta) \frac{\partial}{\partial \theta_s} \ell_n(\theta) + O_p\left(\frac{1}{n}\right). \tag{27}$$

Combining (27) with (25) and (26), we have

$$nH_n(\theta)(\hat{\theta}^{ML} - \theta) = 2\frac{\partial}{\partial \theta}\ell_n(\theta) + \frac{1}{2}\sum_{r=1}^p \sum_{s=1}^p H_n^{rs}(\theta)B_{n,rs}(\theta) + o_p(1).$$

This completes the proof.

5.1.2. Proof of Corollary 3. By applying Assumption 2 to Lemma 8, we can straightforwardly evaluate the difference between the Bayes estimator and the MLE as

$$\hat{\theta}^B - \hat{\theta}^{ML} = \frac{1}{n} H^{-1}(\theta) \left(\frac{\partial}{\partial \theta} \log \pi(\theta) + \frac{1}{2} \sum_{r=1}^p \sum_{s=1}^p H^{rs}(\theta) K_{rs}(\theta) \right) + o_p \left(\frac{1}{n} \right).$$

It remains to evaluate the difference between the MLE and the true parameter. First, we replace (26) with a slightly different expression, which is

$$H(\theta) \left(H_n(\theta) - \frac{1}{2} \sum_{r=1}^{p} (\hat{\theta}_r^{ML} - \theta_r) K_{n,r}..(\theta) \right)^{-1}$$

$$= \left(I_p - (H(\theta) - H_n(\theta)) H^{-1}(\theta) - \frac{1}{2} \sum_{r=1}^{p} (\hat{\theta}_r^{ML} - \theta_r) K_{n,r}..(\theta) H^{-1}(\theta) \right)^{-1}$$

$$= I_p + (H(\theta) - H_n(\theta)) H^{-1}(\theta) + \frac{1}{2} \sum_{r=1}^{p} (\hat{\theta}_r^{ML} - \theta_r) K_{n,r}..(\theta) H^{-1}(\theta) + O_p \left(\frac{1}{n} \right)$$

$$= 2I_p - H_n(\theta) H^{-1}(\theta) + \frac{1}{2} \sum_{r=1}^{p} (\hat{\theta}_r^{ML} - \theta_r) K_{n,r}..(\theta) H^{-1}(\theta) + O_p \left(\frac{1}{n} \right).$$

This equation, combined with (25) and (27), gives us

$$\begin{split} nH_n(\theta)(\hat{\theta}^{ML} - \theta) \\ &= 2\frac{\partial}{\partial \theta}\ell_n(\theta) + \sum_{r=1}^p \sum_{s=1}^p H^{rs}(\theta) \begin{bmatrix} J_{n,1r,s}(\theta) \\ \vdots \\ J_{n,pr,s}(\theta) \end{bmatrix} \\ &+ \frac{1}{2} \sum_{r=1}^p \sum_{s=1}^p H_n^{rs}(\theta) \sum_{t=1}^p \sum_{u=1}^p H^{tu} K_{n,r\cdot t}(\theta) I_{n,su}(\theta) + o_p(1) \\ &= 2\frac{\partial}{\partial \theta}\ell_n(\theta) + \sum_{r=1}^p \sum_{s=1}^p H^{rs}(\theta) \begin{bmatrix} J_{n,1r,s}(\theta) \\ \vdots \\ J_{n,pr,s}(\theta) \end{bmatrix} + \frac{1}{2} \sum_{t=1}^p \sum_{u=1}^p H^{tu}(\theta) K_{r\cdot t}(\theta) I_{n,su}(\theta) + o_p(1). \end{split}$$

Thus, we have

$$\hat{\theta}^B - \theta = \frac{1}{n} H^{-1}(\theta) \left(\frac{\partial}{\partial \theta} \left(\log \pi(\theta) + 2\ell_n(\theta) \right) + \frac{1}{2} \sum_{r=1}^p \sum_{s=1}^p H^{rs}(\theta) \bar{A}_{n,rs}(\theta) \right) + o_p \left(\frac{1}{n} \right)$$

where

$$\bar{A}_{n,rs}(\theta) = \begin{bmatrix} K_{1rs}(\theta) + 2J_{n,1r,s}(\theta) \\ \vdots \\ K_{prs}(\theta) + 2J_{n,pr,s}(\theta) \end{bmatrix} + \frac{1}{2} \sum_{t=1}^{p} \sum_{u=1}^{p} H^{tu}(\theta) K_{r\cdot t}(\theta) I_{n,su}(\theta)$$

Taking the expectation of the above equation, we obtain the desired result.

5.2. **Proof of Theorem 5.** We prove the following general result.

Lemma 10. Suppose Θ is a rectangular subspace of \mathbb{R}^p . Let $\phi: \Theta \to \mathbb{R}^p$ be a vector-valued function that is first-order differentiable. We refer to the r-th component of $\phi(\theta)$ as $\phi_r(\theta)$. Consider a system of partial differential equations

$$\frac{\partial}{\partial \theta} \xi(\theta) = \phi(\theta) \quad \text{for all inner points } \theta \in \Theta$$
 (28)

for some function $\xi: \Theta \to \mathbb{R}$. Suppose we can interchange the order of integration and differentiation of ϕ , i.e.,

$$\int \left(\frac{\partial}{\partial \theta}\phi(\theta)\right) d\theta = \frac{\partial}{\partial \theta} \left(\int \phi(\theta) d\theta\right).$$

Then, (28) has a twice continuously differentiable solution $\xi:\theta\to\mathbb{R}$ if and only if

$$\frac{\partial}{\partial \theta_s} \phi_r(\theta) = \frac{\partial}{\partial \theta_r} \phi_s(\theta) \tag{29}$$

holds for all $r, s = 1, \ldots, p$.

Proof. (\Longrightarrow): Differentiating both sides of (28) with respect to θ , we have

$$\frac{\partial}{\partial \theta \partial \theta^{\top}} \xi(\theta) = \frac{\partial}{\partial \theta} \phi(\theta).$$

Observe that the left-hand side of the above equation is a symmetric matrix since ξ is twice continuously differentiable. Therefore, the condition (29) holds.

 (\Leftarrow) : Fix an arbitrary constant vector $(c_1,\ldots,c_p)\in\Theta$. For each $r=1,\ldots,p$, define a function $\psi_r:\mathbb{R}^{p-r+1}\to\mathbb{R}$ by

$$\psi_r(\theta_r,\ldots,\theta_p) = \phi_r(c_1,\ldots,c_{r-1},\theta_r,\ldots,\theta_p).$$

Then, for an arbitrary constant C, the function

$$\xi(\theta) = \sum_{r=1}^{p} \int_{c_r}^{\theta_r} \psi_r(z, \theta_{r+1}, \dots, \theta_p) dz + C$$
(30)

is a solution to (28). Indeed, by (29), for s > r, we have

$$\frac{\partial}{\partial \theta_s} \psi_r(\theta_r, \dots, \theta_p) = \frac{\partial}{\partial \theta_s} \phi_r(\theta) \Big|_{(\theta_1, \dots, \theta_{r-1}) = (c_1, \dots, c_{r-1})} = \frac{\partial}{\partial \theta_r} \phi_s(\theta) \Big|_{(\theta_1, \dots, \theta_{r-1}) = (c_1, \dots, c_{r-1})}.$$
(31)

Thus, for $s = 1, \ldots, p$, we obtain

 $=\phi_s(\theta_1,\ldots,\theta_n),$

$$\frac{\partial}{\partial \theta_s} \left(\sum_{r=1}^p \int_{c_r}^{\theta_r} \psi_r(z, \theta_{r+1}, \dots, \theta_p) dz + C \right)$$

$$= \sum_{r=1}^{s-1} \int_{c_r}^{\theta_r} \left(\frac{\partial}{\partial \theta_s} \psi_r(z, \theta_{r+1}, \dots, \theta_p) \right) dz + \psi_s(\theta_s, \dots, \theta_p)$$

$$= \sum_{r=1}^{s-1} \int_{c_r}^{\theta_r} \left(\frac{\partial}{\partial z} \phi_s(\theta_1, \dots, \theta_{r-1}, z, \theta_{r+1}, \dots, \theta_p) \Big|_{(\theta_1, \dots, \theta_{r-1}) = (c_1, \dots, c_{r-1})} \right) dz + \psi_s(\theta_s, \dots, \theta_p)$$

$$= \sum_{r=1}^{s-1} \left(\phi_s(c_1, \dots, c_{r-1}, \theta_r, \dots, \theta_p) - \phi_s(c_1, \dots, c_r, \theta_{r+1}, \dots, \theta_p) \right) + \phi_s(c_1, \dots, c_{s-1}, \theta_s, \dots, \theta_p)$$

where the second equality follows from (31) and the third equality follows from the fundamental theorem of calculus. Hence, (30) is a solution to (28).

ACKNOWLEDGEMENTS

The authors would like to thank Kaoru Irie, Shonosuke Sugasawa, and Shintaro Hashimoto for their valuable comments on the content of this article. The authors also thank Tomoya Wakayama for his assistance in improving the simulation code. Takeru Matsuda was supported by JSPS KAKENHI Grant Numbers 21H05205, 22K17865, 24K02951 and JST Moonshot Grant Number JPMJMS2024. Tatsuya Kubokawa was supported by JSPS KAKENHI Grant Number 22K11928.

References

Berger, J. O. and J. M. Bernardo (1992). On the development of reference priors. In *Bayesian Statistics 4: Proceedings of the Fourth Valencia International Meeting, Dedicated to the memory of Morris H. DeGroot, 1931-1989.* Oxford University Press.

Bernardo, J. M. (1979). Reference posterior distributions for Bayesian inference. *Journal of the Royal Statistical Society: Series B (Methodological)* 41(2), 113–128.

Datta, G. S. and M. Ghosh (1991). Bayesian prediction in linear models: Applications to small area estimation. *The Annals of Statistics* 19(4), 1748–1770.

Firth, D. (1993). Bias reduction of maximum likelihood estimates. *Biometrika* 80(1), 27–38. Ghosh, M. and R. Liu (2011). Moment matching priors. *Sankhyā: The Indian Journal of Statistics*, *Series A* (2008-) 73(2), 185–201.

Hartigan, J. A. (1965). The asymptotically unbiased prior distribution. The Annals of Mathematical Statistics 36(4), 1137–1152.

Jeffreys, H. (1961). The Theory of Probability (Third ed.). Oxford University Press.

Meng, X.-L. and A. M. Zaslavsky (2002). Single observation unbiased priors. The Annals of Statistics 30(5), 1345 - 1375.

Rao, J. N. and I. Molina (2015). Small area estimation (Second ed.). John Wiley & Sons. Sugasawa, S. and T. Kubokawa (2023). Mixed-Effects Models and Small Area Estimation. SpringerBriefs in Statistics. Springer Singapore.

Tiao, G. C. and W. Y. Tan (1965). Bayesian analysis of random-effect models in the analysis of variance. I. Posterior distribution of variance-components. *Biometrika* 52(1/2), 37-53.

Appendix A. Posterior propriety of the asymptotically unbiased prior for the balanced NER model

In this section, we prove that the posterior distribution corresponding to the prior given in (17) is proper.

As discussed earlier in Example 7, we introduce a transformation of the parameters $\theta = (\beta, \tau^2, \sigma^2)$ given by

$$\bar{\theta} = (\beta, \rho, \sigma^2) \in \bar{\Theta} = \mathbb{R}^p \times (0, 1) \times (0, \infty), \quad \rho = \frac{\sigma^2}{\sigma^2 + n\tau^2}.$$

The posterior distribution in terms of $\bar{\theta}$, as shown in (19), is expressed as

$$\bar{\pi}(\bar{\theta}|\mathcal{D}) \propto (\sigma^2)^{-3-\frac{nm}{2}} \rho^{\frac{m}{2}} \exp\left(-\frac{1}{2} \sum_{i=1}^m (y_i - x_i \beta)^\top \bar{V}^{-1} (y_i - x_i \beta)\right) =: \bar{\pi}(\bar{\theta}|\mathcal{D}),$$

where \bar{V}^{-1} is defined as $\bar{V}^{-1} = \sigma^{-2}(I_n - (1 - \rho)\iota_n\iota_n^\top/n)$. To establish posterior propriety, we need to show that the integral of $\bar{\pi}(\bar{\theta}|\mathcal{D})$ over $\bar{\Theta}$ is finite.

Assumption 3. Assume that the following conditions are satisfied:

- (i) There exists $i \in \{1, ..., m\}$ such that x_i has full column rank.
- (ii) There exists $i \in \{1, ..., m\}$ such that $[y_i \ x_i]$ has full column rank.
- (iii) There exists $i \in \{1, ..., m\}$ such that the sample variance covariance matrix

$$\frac{1}{n} \sum_{j=1}^{n} \left(x_{ij} - \frac{1}{n} \sum_{j=1}^{n} x_{ij} \right) \left(x_{ij} - \frac{1}{n} \sum_{j=1}^{n} x_{ij} \right)^{\top}$$

is positive semidefinite.

(iv) There exists $i \in \{1, ..., m\}$ such that the sample variance covariance matrix

$$\frac{1}{n} \sum_{j=1}^{n} \left(\begin{bmatrix} y_{ij} \\ x_{ij} \end{bmatrix} - \frac{1}{n} \sum_{j=1}^{n} \begin{bmatrix} y_{ij} \\ x_{ij} \end{bmatrix} \right) \left(\begin{bmatrix} y_{ij} \\ x_{ij} \end{bmatrix} - \frac{1}{n} \sum_{j=1}^{n} \begin{bmatrix} y_{ij} \\ x_{ij} \end{bmatrix} \right)^{\top}$$

is positive semidefinite.

Proposition 11. If Assumption 3 is satisfied, we have

$$\int_{\bar{\Theta}} \bar{\pi}(\bar{\theta}|\mathcal{D}) d\bar{\theta} < \infty.$$

Proof. For notational simplicity, we define

$$y = \begin{bmatrix} y_1 \\ \vdots \\ y_m \end{bmatrix}, \quad X = \begin{bmatrix} x_1 \\ \vdots \\ x_m \end{bmatrix}.$$

Using this notation, the posterior distribution $\bar{\pi}(\bar{\theta}|\mathcal{D})$ can expressed as

$$\bar{\pi}(\bar{\theta}|\mathcal{D}) = (\sigma^2)^{-3 - \frac{nm}{2}} \rho^{\frac{m}{2}} \exp\left(-\frac{1}{2\sigma^2} (y - X\beta)^{\top} \bar{Q}(\rho) (y - X\beta)\right),$$

where $\bar{Q}(\rho)$ is a block diagonal matrix with m blocks, each given by

$$I_n - \frac{1-\rho}{n} \iota_n \iota_n^{\mathsf{T}}.$$

Since $X^{\top}\bar{Q}(\rho)X$ is positive definite for $\rho \in [0,1]$ by Lemmas 12 and 13, we can integrate out β as

$$\begin{split} &\int_{\mathbb{R}^p} \exp\left(-\frac{1}{2\sigma^2}(y - X\beta)^\top \bar{Q}(\rho)(y - X\beta)\right) d\beta \\ &= (2\pi\sigma^2)^{p/2} \det(X^\top \bar{Q}(\rho)X)^{-1/2} \exp\left(-\frac{1}{2\sigma^2}y^\top \bar{Q}(\rho)y - y^\top \bar{Q}(\rho)X(X^\top \bar{Q}(\rho)X)^{-1}X^\top \bar{Q}(\rho)y\right) \\ &= (2\pi\sigma^2)^{p/2} \det(X^\top \bar{Q}(\rho)X)^{-1/2} \exp\left(-\frac{1}{2\sigma^2}h(\rho)\right), \end{split}$$

where $h(\rho)$ is given by

$$h(\rho) = \det(X^{\top} \bar{Q}(\rho) X)^{-1} \det\left(\begin{bmatrix} y^{\top} \\ X^{\top} \end{bmatrix} \bar{Q}(\rho) \begin{bmatrix} y & X \end{bmatrix}\right).$$

Since Lemma 12 also implies

$$\det(X^\top \bar{Q}(\rho)X) > 0 \quad \text{and} \quad \det\left(\begin{bmatrix} y^\top \\ X^\top \end{bmatrix} \bar{Q}(\rho) \begin{bmatrix} y & X \end{bmatrix}\right) > 0,$$

we can express the integral as

$$\int_{\bar{\Theta}} \bar{\pi}(\bar{\theta}|\mathcal{D}) d\bar{\theta}
= (2\pi)^{p/2} \int_{0}^{1} \int_{0}^{\infty} (\sigma^{2})^{-3 - \frac{nm-p}{2}} \rho^{\frac{m}{2}} \det(X^{\top} \bar{Q}(\rho) X)^{-1/2} \exp\left(-\frac{1}{2\sigma^{2}} h(\rho)\right) d\sigma^{2} d\rho
= (2\pi)^{p/2} 2^{2 + \frac{nm-p}{2}} \Gamma\left(2 + \frac{nm-p}{2}\right) \int_{0}^{1} \rho^{\frac{m}{2}} \det(X^{\top} \bar{Q}(\rho) X)^{-1/2} h(\rho)^{-\left(2 + \frac{nm-p}{2}\right)} d\rho.$$

Since the integrand is continuous and the domain of integration is bounded in the last equation, the integral is finite. Therefore, we can conclude that the posterior distribution is proper. \Box

Lemma 12. Suppose conditions (i) and (ii) of Assumption 3 hold. Then, for any $\rho > 0$, we have

$$X^{\top} \bar{Q}(\rho) X > 0$$
 and $\begin{bmatrix} y^{\top} \\ X^{\top} \end{bmatrix} \bar{Q}(\rho) \begin{bmatrix} y & X \end{bmatrix} > 0.$

Proof. We first show the positive definiteness of $I_n - \frac{1-\rho}{n} \iota_n \iota_n^{\top}$. Indeed, for any non-zero vector $b \in \mathbb{R}^n \setminus \{0\}$, by the Cauchy–Schwarz inequality, we have

$$b^{\top} \left(I_n - \frac{1 - \rho}{n} \iota_n \iota_n^{\top} \right) b = \|b\|^2 - \frac{1 - \rho}{n} (\iota_n^{\top} b)^2 \ge \|b\|^2 - \frac{1 - \rho}{n} \|\iota_n\|^2 \|b\|^2 = \rho \|b\|^2 > 0.$$

Next, we note that the matrices $X^{\top}\bar{Q}(\rho)X$ and $[y\ X]^{\top}\bar{Q}(\rho)[y\ X]$ can be expressed as

$$X^{\top} \bar{Q}(\rho) X = \sum_{i=1}^{m} x_{i}^{\top} \left(I_{n} - \frac{1 - \rho}{n} \iota_{n} \iota_{n}^{\top} \right) x_{i},$$
$$\begin{bmatrix} y^{\top} \\ X^{\top} \end{bmatrix} \bar{Q}(\rho) \begin{bmatrix} y & X \end{bmatrix} = \sum_{i=1}^{m} \begin{bmatrix} y_{i}^{\top} \\ x_{i}^{\top} \end{bmatrix} \left(I_{n} - \frac{1 - \rho}{n} \iota_{n} \iota_{n}^{\top} \right) \begin{bmatrix} y_{i} & x_{i} \end{bmatrix}.$$

Observe that $x_i^{\top} \left(I_n - \frac{1-\rho}{n} \iota_n \iota_n^{\top} \right) x_i$ and $[y_i \ x_i]^{\top} \left(I_n - \frac{1-\rho}{n} \iota_n \iota_n^{\top} \right) [y_i \ x_i]$ are positive semidefinite for each i. Furthermore, by conditions (i) and (ii) of Assumption 3, there exists i, i' such that

$$x_i^{\top} \left(I_n - \frac{1-\rho}{n} \iota_n \iota_n^{\top} \right) x_i > 0 \quad \text{and} \quad \begin{bmatrix} y_{i'}^{\top} \\ x_{i'}^{\top} \end{bmatrix} \left(I_n - \frac{1-\rho}{n} \iota_n \iota_n^{\top} \right) \begin{bmatrix} y_{i'} & x_{i'} \end{bmatrix} > 0$$

hold. Therefore, we conclude that both $X^{\top}\bar{Q}(\rho)X$ and $[y\ X]^{\top}\bar{Q}(\rho)[y\ X]$ are positive definite.

Lemma 13. Suppose Assumption 3 holds. Then for $\rho = 0$, we have

$$X^{\top}\bar{Q}(0)X>0 \quad and \quad \begin{bmatrix} y^{\top} \\ X^{\top} \end{bmatrix} \bar{Q}(0) \begin{bmatrix} y & X \end{bmatrix}>0.$$

Proof. By the same argument as in the previous lemma, we can show that $I_n - \frac{1}{n}\iota_n\iota_n$ is positive semidefinite. Thus, $x_i^{\top} \left(I_n - \frac{1}{n}\iota_n\iota_n^{\top}\right) x_i \geq 0$ and $[y_i \ x_i]^{\top} \left(I_n - \frac{1}{n}\iota_n\iota_n^{\top}\right) [y_i \ x_i] \geq 0$ hold for each i. Furthermore, by condition (iii) of Assumption 3, there exists i that satisfies

$$x_i^{\top} \left(I_n - \frac{1}{n} \iota_n \iota_n^{\top} \right) x_i = \sum_{j=1}^n \left(x_{ij} - \frac{1}{n} \sum_{j=1}^n x_{ij} \right) \left(x_{ij} - \frac{1}{n} \sum_{j=1}^n x_{ij} \right)^{\top} > 0.$$

Therefore, we have

$$X^{\top} \bar{Q}(0) X = \sum_{i=1}^{m} x_i^{\top} \left(I_n - \frac{1}{n} \iota_n \iota_n^{\top} \right) x_i > 0.$$

By a similar argument, we also obtain

$$\begin{bmatrix} y^{\top} \\ X^{\top} \end{bmatrix} \bar{Q}(0) \begin{bmatrix} y & X \end{bmatrix} = \sum_{i=1}^{m} \begin{bmatrix} y_i^{\top} \\ x_i^{\top} \end{bmatrix} \left(I_n - \frac{1}{n} \iota_n \iota_n^{\top} \right) \begin{bmatrix} y_i & x_i \end{bmatrix} > 0$$

as required.

APPENDIX B. SIMULATION DETAILS

In this section, we provide the details of the simulation studies in Section 4.

For each area i, the number of units is set to n = 5, and the sample size is varied across $m \in \{10, 32, 100, 316, 1000\}$. The covariates are generated as $x_{ij} \sim N(\mu, \Sigma)$, where

$$\mu = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad \Sigma = \begin{bmatrix} 4 & 1 \\ 1 & 1 \end{bmatrix}.$$

For [AU] and [JF], the MCMC sample size is set to N=2000 with a warm-up size of warmup=100, while for [DG], N=20000 with warmup=1000. For [DG], a larger sample size was chosen due to the higher autocorrelation of the chain compared to the

other priors. All generated chains have at least 300 effective sample sizes, indicating good convergence and reliable posterior inference.

To examine the frequentist properties of Bayes estimators, we simulate 10000 independent datasets of $\mathcal{D} = (y_i, x_i)_{1 \leq i \leq m}$ for each sample size m. For each dataset, we compute the posterior mean $\hat{\theta}^B$, the bias, the mean squared error (MSE), and the coverage probability of the 95% credible interval for each parameter.

For each prior, Gibbs sampler is applied to sample from the posterior distribution. Sampling from the posterior distribution corresponding to $[\mathbf{AU}]$ is explained in Example 7, and a similar transformation of the parameter is used for $[\mathbf{JF}]$. For $[\mathbf{DG}]$, we treat the model as a hierarchical model and sample $\beta, \tau^2, \sigma^2, \{v_i : i \leq m\}$ by Gibbs sampler.

Figures 3 and 4 show the log-MSE of the posterior mean under the two parameter settings. Figures 5 and 6 show the coverage probability of the 95% credible interval, each corresponding to the two parameter settings. From these results, it can be observed that [AU] and [JF] exhibit comparable performance in terms of MSE and coverage probability. In contrast, the performance of [DG] is more sensitive to the true parameter settings, which might be attributed to the choice of hyperparameters in the prior.




FIGURE 3. Log-MSE of the posterior mean when the true parameter is $\beta_1 = \beta_2 = \tau^2 = \sigma^2 = 1$.




FIGURE 4. Log-MSE of the posterior mean when the true parameter is $\beta_1=\beta_2=1, \tau^2=0.5, \sigma^2=4.$




FIGURE 5. Coverage probability of the posterior mean when the true parameter is $\beta_1 = \beta_2 = \tau^2 = \sigma^2 = 1$.




FIGURE 6. Coverage probability of the posterior mean when the true parameter is $\beta_1 = \beta_2 = 1, \tau^2 = 0.5, \sigma^2 = 4$.