

Implications for the non-Gaussianity of primordial gravitational waves from pulsar timing arrays

Zhi-Zhang Peng^{1,*}, You Wu^{2,†} and Lang Liu^{3,‡}

¹*School of Physics and Astronomy, Beijing Normal University, Beijing 100875, People's Republic of China*

²*College of Mathematics and Physics, Hunan University of Arts and Science, Changde, 415000, China*

³*Faculty of Arts and Sciences, Beijing Normal University, Zhuhai 519087, China*

The detection of a stochastic signal by recent pulsar timing array (PTA) collaborations, including NANOGrav, PPTA, EPTA+InPTA, CPTA and MPTA, has opened a new window to explore gravitational waves (GWs) at nanohertz frequencies. Motivated by the possibility that such a signal could arise from primordial gravitational waves (PGWs), we investigate the implications of tensor non-Gaussianity for the PGW power spectrum. Utilizing PTA data sets, we provide constraints on local-type tensor non-Gaussianity parameter \mathcal{F}_{NL} . We find $|\mathcal{F}_{\text{NL}}| \lesssim 7.97$ for a log-normal PGW power spectrum. Our analysis reveals that even moderate tensor non-Gaussianity can lead to significant deviations from standard predictions, thereby offering a novel means to test inflationary scenarios and probe the underlying dynamics of the early Universe. Future multi-band GW observatories, such as LISA, Taiji, and TianQin, will be instrumental in complementing these efforts and further refining our understanding of tensor non-Gaussianity.

Introduction. Primordial gravitational waves (PGWs), as one of the key predictions of inflationary cosmology, offer an exceptional opportunity to probe the early Universe [1, 2]. The standard inflationary paradigm predicts that PGWs follow a nearly Gaussian distribution due to the linearity of quantum field perturbations. However, certain extensions to the inflationary framework introduce non-Gaussianities in PGWs, manifesting as deviations from Gaussian statistics in correlation functions. Investigating non-Gaussian features provides an opportunity to explore inflationary dynamics beyond the simplest models, offering deeper insights into the fundamental physics underlying the early Universe.

Non-Gaussianities in PGWs arise in various inflationary scenarios, such as extra fields [3–10], non-attractor phase for tensor fluctuations [11], massive gravity theory [12, 13], generalized G-inflation [14–17], non-standard vacuum states [18–21], higher-derivative Weyl terms [22–24], Chern-Simons interactions [24–31], and more generally, effective field theory [32–35]. Constraints on tensor non-Gaussianity can be derived from measurements of the three-point correlation function of the cosmic microwave background (CMB) temperature anisotropies and polarization. Up to now, the theoretical investigation of different non-Gaussianity shapes and their detectability has been conducted. Moreover, some of these shapes have already been evaluated using CMB data [36–41]. However, large-scale non-Gaussianity constraints alone are insufficient to distinguish between different inflationary models. With the upcoming launch of space-based GW interferometers, such as LISA [42], Taiji [43], and TianQin [44], it is crucial to probe small-scale tensor non-Gaussianity to complete the puzzle of full-scale tensor non-Gaussianity.

Usually, tensor non-Gaussianity primarily impacts the PGW power spectrum by introducing nontrivial spectral tilts, which can enhance or suppress the GW across different frequency ranges. Additionally, non-Gaussian corrections may shift the peak frequency of the PGW, pushing it to either higher or lower frequencies. This sensitivity to tensor non-Gaussianity offers a pathway to distinguish between various inflationary models and to probe the fundamental dynamics of inflation. Recently, multiple pulsar timing array (PTA) collaborations—such as the Chinese PTA (CPTA) [45], the European PTA (EPTA) in collaboration with the Indian PTA (InPTA) [46, 47], the Parkes PTA (PPTA) [48, 49], the North American Nanohertz Observatory for GWs (NANOGrav) [50, 51], and the MeerKAT PTA (MPTA) [52, 53] have independently unveiled robust evidence suggesting the presence of a stochastic gravitational-wave background (SGWB). If the observed signal can be attributed to PGWs, its power spectrum may exhibit observable effects induced by non-Gaussianity. PTA data analysis can therefore constrain non-Gaussianity parameters associated with PGWs.

In this letter, we consider the impact of tensor non-Gaussianity on the power spectrum of PGWs. As the first step, we only focus on the local-type non-Gaussianities, parameterized by a tensor non-Gaussianity parameter \mathcal{F}_{NL} . Using the latest PTA datasets, including the NANOGrav 15-year data release, PPTA DR3, and EPTA DR2, we perform a joint constraints on \mathcal{F}_{NL} . We find $-7.97 \lesssim \mathcal{F}_{\text{NL}} \lesssim 7.97$ when primordial tensor perturbation follows a log-normal form. This result establishes a robust framework for testing inflationary models and predictions of the statistical properties of PGWs.

Recent studies suggest that second-order scalar-induced GWs could also account for the PTA signals [54–77]. In this context, GWs induced by non-Gaussian curvature perturbations have attracted considerable attention [78–83], and one can use PTA data to constrain

* pengzhizhang@bnu.edu.cn

† Corresponding author: youwuphy@gmail.com

‡ Corresponding author: liulang@bnu.edu.cn

the non-Gaussianity of curvature perturbations on small scales [56, 57, 59, 84]. We emphasize that although scalar-induced GWs also exhibit non-Gaussianity, this non-Gaussianity arises from the curvature perturbations themselves, rather than from the tensor non-Gaussianity inherent to the GWs. Our work serves as an important supplement to the study of scalar non-Gaussianity and will provide valuable insights into exploring the physics of inflation.

Primordial gravitational waves. To investigate the impact of tensor non-Gaussianity on the power spectrum of PGWs, we consider parameterizing the non-Gaussianity of PGWs. A straightforward approach is to mimic the local expansion used in characterizing primordial scalar perturbations. Specially, we assume that the deviation from Gaussianity is small, leading to tensor perturbations in real space expressed as the sum of a Gaussian component and its quadratic term [85–87]

$$h_{ij}(\mathbf{x}) = h_{ij,G}(\mathbf{x}) + \mathcal{F}_{\text{NL}}(h_{ij,G}^2(\mathbf{x}) - \langle h_{ij,G}^2(\mathbf{x}) \rangle), \quad (1)$$

where $h_{ij,G}(\mathbf{x})$ follows Gaussian statistics and \mathcal{F}_{NL} is the dimensionless non-Gaussian parameter. Here, we want to reminder the reader that the value of local-type non-Gaussianities of primordial tensor perturbations can be different from the value of local-type non-Gaussianities of primordial scalar perturbations. In principle, \mathcal{F}_{NL} is a quantity that depends on the GW polarization. For the sake of simplicity, however, we treat it as a constant independent of the polarization. Note that in this expansion, we omit temporal dependence as the tensor non-Gaussianity is evaluated at the end of inflation, and then PGWs imprinted with non-Gaussian features evolve with the expansion of the Universe. The variance term has been subtracted from the quadratic part in order that the expectation value satisfies $\langle h_{ij} \rangle = 0$.

The Fourier modes are written as

$$h_{ij}(\mathbf{k}) = h_{ij,G}(\mathbf{k}) + \mathcal{F}_{\text{NL}} \int \frac{d^3 p}{(2\pi)^3} h_{ij,G}(\mathbf{p}) h_{ij,G}(\mathbf{k} - \mathbf{p}). \quad (2)$$

Note that $h_{ij}(\mathbf{k})$ consists of two polarization components of GWs, i.e., $h_{ij}(\mathbf{k}) = h_{\mathbf{k}}^+ e_{ij}^+(\mathbf{k}) + h_{\mathbf{k}}^\times e_{ij}^\times(\mathbf{k})$, where $e_{ij}^+(\mathbf{k})$ and $e_{ij}^\times(\mathbf{k})$ are polarization tensors which satisfy $e_{ij}^\lambda(\mathbf{k}) e_{ij}^{\lambda'}(-\mathbf{k}) = \delta^{\lambda\lambda'}$, $h_{\mathbf{k}}^+$ and $h_{\mathbf{k}}^\times$ denote tensor perturbations with each polarization. We focus on the two-point correlation of PGWs, and the contributions deviating from the Gaussian state can be extracted. The explicit expression is given by

$$\begin{aligned} & \langle h_{ij}(\mathbf{k}) h_{ij}(\mathbf{k}') \rangle \\ &= \langle h_{ij,G}(\mathbf{k}) h_{ij,G}(\mathbf{k}') \rangle + \int \frac{d^3 p}{(2\pi)^3} \int \frac{d^3 q}{(2\pi)^3} \\ & \times 2\mathcal{F}_{\text{NL}}^2 \langle h_{ij,G}(\mathbf{p}) h_{ij,G}(\mathbf{q}) \rangle \langle h_{ij,G}(\mathbf{k} - \mathbf{p}) h_{ij,G}(\mathbf{k}' - \mathbf{q}) \rangle, \end{aligned} \quad (3)$$

where the four-point correlation function has been decomposed into the product of two-point correlation functions and the disconnected term vanishes.

The dimensionless power spectrum of PGWs, $\mathcal{P}_h(k)$, is defined as

$$\langle h_{ij}(\mathbf{k}) h_{ij}(\mathbf{k}') \rangle = \frac{2\pi^2}{k^3} \mathcal{P}_h(k) (2\pi)^3 \delta^{(3)}(\mathbf{k} + \mathbf{k}'). \quad (4)$$

Note that $\mathcal{P}_h(k)$ represents the total power spectrum including contributions from both PGW polarization modes. We shall decompose the power spectrum into Gaussian contribution and non-Gaussian one

$$\mathcal{P}_h(k) = \mathcal{P}_h^G(k) + \mathcal{P}_h^{\text{NG}}(k). \quad (5)$$

Combining Eqs. (3), (4) and (5), we obtain

$$\mathcal{P}_h^{\text{NG}}(k) = \frac{k^3 \mathcal{F}_{\text{NL}}^2}{2\pi} \int d^3 p \frac{\mathcal{P}_h^G(p) \mathcal{P}_h^G(|\mathbf{k} - \mathbf{p}|)}{p^3 |\mathbf{k} - \mathbf{p}|^3}. \quad (6)$$

After introducing two dimensionless variables $u = |\mathbf{k} - \mathbf{p}|/k$ and $v = p/k$, the non-Gaussian correction can be rewritten as

$$\mathcal{P}_h^{\text{NG}}(k) = \mathcal{F}_{\text{NL}}^2 \int_0^\infty dv \int_{|1-v|}^{1+v} du \frac{\mathcal{P}_h^G(uk) \mathcal{P}_h^G(vk)}{u^2 v^2}. \quad (7)$$

The spectrum of GW energy density per logarithmic frequency interval today can be conveniently normalized as [88]

$$\Omega_{\text{GW}}(k, \tau_0) = \frac{1}{\rho_c} \frac{d\rho_{\text{GW}}}{d \ln k} = \frac{k^2}{12a_0^2 H_0^2} \mathcal{P}_h(k) \overline{T^2(k, \tau_0)}, \quad (8)$$

where $\rho_c = 3H_0^2/(8\pi G)$, $\tau_0 = 1.41 \times 10^4$ Mpc, $a_0 = 1$, $H_0 = 67.4$ km s⁻¹ Mpc⁻¹, and the overline denotes the oscillation average. The transfer function $T(k, \tau_0)$ can be obtained by numerically solving the post-inflationary evolution of PGWs. A good fit to the transfer function is [89]

$$T(k, \tau_0) = \frac{3\Omega_m j_1(k\tau_0)}{k\tau_0} \sqrt{1.0 + 1.36 \frac{k}{k_{\text{eq}}} + 2.50 \left(\frac{k}{k_{\text{eq}}}\right)^2}, \quad (9)$$

where $k_{\text{eq}} = 0.073 \Omega_m h^2$ Mpc⁻¹ is that of the perturbation mode that entered the horizon at the equality of matter and radiation.

The evolution of PGW strongly depends on the thermal history of the Universe, including factors such as dark energy with time-varying equation of state, tensor anisotropic stress due to free-streaming relativistic particles in the early Universe, deviations from the standard equation of state parameter $w = 1/3$ in the radiation era, and so on [89–92]. Since our focus is on the impact of tensor non-Gaussianity on the GW power spectrum, we neglect the detailed corrections during the propagation process.

To step forward, we shall specify the k dependence of $\mathcal{P}_h(k)$. We consider the following log-normal parametrization for the Gaussian power spectrum

$$\mathcal{P}_h^G(k) = \frac{A}{\sqrt{2\pi\Delta^2}} \exp\left(-\frac{\ln^2(k/k_*)}{2\Delta^2}\right), \quad (10)$$




FIG. 1. Energy density spectrum of PGWs predicted by the non-Gaussian model \mathcal{M}_{NG} . The olive solid line shows the median prediction, with the surrounding shaded region indicating the 90% credible interval. The violin plots display the energy density spectra derived from PTA observations: NANOGrav 15-year data (red), PPTA DR3 (green), and EPTA DR2 (blue). For comparison, we show the power-law integrated sensitivity curves for future space-based GW detectors: LISA (solid black), Taiji (dashed black), and TianQin (dash-dotted black).

Parameter	A	f_*/Hz	Δ	$ \mathcal{F}_{\text{NL}} $
Prior	$\log\mathcal{U}(-1, 0)$	$\log\mathcal{U}(-8, -6.4)$	$\mathcal{U}(0.05, 3.5)$	$\log\mathcal{U}(-5, 3)$
Result for \mathcal{M}_G	$0.60_{-0.28}^{+0.29}$	$2.36_{-1.41}^{+1.43} \times 10^{-7}$	$1.58_{-0.33}^{+0.18}$	—
Result for \mathcal{M}_{NG}	$0.39_{-0.27}^{+0.43}$	$2.37_{-1.44}^{+1.40} \times 10^{-7}$	$1.67_{-0.36}^{+0.41}$	$\lesssim 7.97$

TABLE I. Prior choices and posterior constraints for parameters in both the non-Gaussian model (\mathcal{M}_{NG}) and Gaussian model (\mathcal{M}_G). Prior distributions are specified using uniform (\mathcal{U}) and log-uniform ($\log\mathcal{U}$) distributions. For each parameter, we report the median value and symmetric 90% credible interval derived from the posterior distributions.

where A denotes the amplitude of the power spectrum, k_* is the pivot scale corresponding to the peak location, and Δ characterizes the width of the spectrum.

Methodology and result. To investigate non-Gaussianity of PGWs, we perform a joint analysis of data from three major PTA collaborations: the NANOGrav 15-year dataset [51], PPTA DR3 [48], and EPTA DR2 [46]. These long-term observational campaigns represent the culmination of over a decade of precise pulsar timing measurements, with each collaboration independently reporting evidence for a SGWB showing the characteristic Hellings-Downs [93] spatial correlations. The datasets encompass observations from com-

plementary pulsar observations: NANOGrav monitors 68 pulsars over 16.03 years [51], PPTA tracks 32 pulsars for up to 18 years [48], and EPTA observes 25 pulsars across 24.7 years [46]. The detection of consistent signals [94] across these independent arrays strengthens the case for a GW origin, motivating our joint analysis approach to maximize parameter estimation precision.

Our analysis utilizes the free spectral representation of the SGWB derived by each collaboration using Hellings-Downs correlations. The frequency coverage is determined by each array's observational timespan T_{obs} , with a minimum frequency $f_{\min} = 1/T_{\text{obs}}$. The spectral analysis encompasses 14 frequency bins from NANOGrav,

28 from PPTA, and 24 from EPTA, yielding a combined spectrum of 66 frequency components spanning 1.28–49.1nHz (visualized in Fig. 1). To maintain consistency with cosmological constraints, we incorporate bounds from CMB and baryon acoustic oscillation measurements [95]. Specifically, we impose the integrated energy density constraint that $\int_{k_{\min}}^{\infty} h^2 \Omega_{\text{GW},0}(k) d \ln k \lesssim 2.9 \times 10^{-7}$, where $h = 0.674$ is the dimensionless Hubble parameter [95, 96].

Beginning with the time delay data $d(f)$ released by each collaboration, we first reconstruct the power spectral density through

$$S(f) = d^2(f) T_{\text{obs}}, \quad (11)$$

where T_{obs} is the observational timespan. The GW characteristic strain spectrum $h_c(f)$ is then derived via

$$h_c(f) = \sqrt{12\pi^2 f^3 S(f)}. \quad (12)$$

We further compute the dimensionless GW energy density spectrum as

$$\hat{\Omega}_{\text{GW}}(f) = \frac{2\pi^2}{3H_0^2} f^2 h_c^2(f) = \frac{8\pi^4}{H_0^2} T_{\text{obs}} f^5 d^2(f). \quad (13)$$

To construct the likelihood function, we estimate the kernel density \mathcal{L}_i from the posterior distributions of $\hat{\Omega}_{\text{GW}}(f_i)$ at each frequency bin. The total likelihood for our parameter set $\Lambda = \{A, \Delta, f_*, |\mathcal{F}_{\text{NL}}|\}$ takes the form of

$$\mathcal{L}(\Lambda) = \prod_{i=1}^{66} \mathcal{L}_i(\Omega_{\text{GW}}(f_i, \Lambda)), \quad (14)$$

accounting for correlations across the entire frequency range of our analysis.

For parameter estimation, we employ the nested sampling algorithm implemented in `dynesty` [97], interfaced through the `Bilby` [98, 99] package. This approach efficiently explores the multi-dimensional parameter space while providing robust evidence calculations. The prior distributions and ranges for all model parameters are detailed in Table I.

Our analysis compares two competing models for the PGW background: a standard Gaussian model (\mathcal{M}_G) and a non-Gaussian model (\mathcal{M}_{NG}) incorporating primordial tensor non-Gaussianity. The resulting parameter constraints, displayed in Fig. 2, reveal a significant degeneracy between the amplitude parameter A and the non-Gaussianity parameter \mathcal{F}_{NL} in the \mathcal{M}_{NG} analysis, arising from their similar effects on the overall signal strength near the spectrum's peak. Despite this degeneracy, our joint analysis yields constraints, with the amplitude bounded to $A = 0.39^{+0.43}_{-0.27}$ (90% C.L.) and non-Gaussianity parameter constrained to $|\mathcal{F}_{\text{NL}}| \lesssim 7.97$ (90% C.L.). Due to the symmetry in how non-Gaussian corrections enter our model, we find equivalent constraints for positive and negative values, leading to the bilateral




FIG. 2. Joint posterior distributions of model parameters, showing both one-dimensional marginalized distributions and two-dimensional correlation contours. Results are presented for both the Gaussian model \mathcal{M}_G (red) and non-Gaussian model \mathcal{M}_{NG} (blue), derived from the combined analysis of NANOGrav 15-year, PPTA DR3, and EPTA DR2 datasets. Two-dimensional contours represent 68.3%, 95.4%, and 99.7% credible regions, respectively.

bound $-7.97 \lesssim \mathcal{F}_{\text{NL}} \lesssim 7.97$, with complete parameter estimates and their uncertainties presented in Table I.

Conclusion and discussion. Inflation is a cornerstone of modern cosmology, providing insights into the dynamics of the early Universe and aligning well with observational data. However, key questions about the inflationary mechanism remain unresolved, such as the energy scale of inflation and its particle content. The most promising observables for addressing these questions are the power spectrum and bispectrum of primordial perturbations, both in the scalar and tensor sectors. Current data impose stringent constraints on the inflationary scalar sector at CMB scales. In contrast, the tensor sector may provide vital insights into the mass and spin of extra fields and is directly connected to modified gravity. Thus, exploring tensor non-Gaussianity complements the scalar one, helping to test various inflationary models.

In this letter, assuming the signal detected by the PTA collaborations originates from PGWs, we jointly utilize the NANOGrav 15-yr data set, PPTA DR3, and EPTA DR2 to constrain the PGWs with non-Gaussianity. For the first time, we place a limit on the non-Gaussianity parameter $-7.97 \lesssim \mathcal{F}_{\text{NL}} \lesssim 7.97$ for a log-normal power spectrum of PGWs. Moreover, as illustrated in Figure 1, the energy density spectrum of PGWs can typically be extended to the frequency range detectable by space-

based GW observatories. Consequently, the combined multi-band observations from PTAs and the upcoming space-borne GW detectors, including LISA [42], Taiji [43], and TianQin [44], will offer a complementary approach to studying tensor non-Gaussianity.

For simplicity, we focus on the tensor non-Gaussianity up to second-order, characterized by non-Gaussianity parameter \mathcal{F}_{NL} . A complete one-loop correction shall include the third-order corrections, akin to \mathcal{G}_{NL} order in the study of non-Gaussianity of curvature perturbations [100, 101]. For curvature perturbation case, the full one-loop contribution from the scalar sector significantly affects both the abundance of primordial black holes and the energy spectrum of scalar-induced GWs, while the tensor sector primarily influences the power spectrum of PGWs. We will address these details in future work.

In the search of Planck, they concentrate on tensor non-Gaussianity of the equilateral type, and place constraints on two specific inflationary models, including either a U(1)-axion coupling or an SU(2)-axion one. Taking U(1)-axion model for example, the constraints on $\mathcal{F}_{\text{NL}}^{\text{equil}}$ is $\mathcal{F}_{\text{NL}}^{\text{equil}} \approx 6.4 \times 10^{11} \mathcal{P}^3 \epsilon^3 e^{6\pi\xi} / \xi^9$, where \mathcal{P} is the curvature perturbation power spectrum, ϵ is a slow-roll parameter of the inflaton field, and ξ expresses the strength of the U(1)-axion coupling [102, 103]. Then, the upper bound on ξ is derived as $\xi < 3.3$ (95 % C.L.) [39]. The results presented above provide constraints on tensor non-Gaussianity on large scales. If the GWs generated in axion models can also explain the PTA signals, we can obtain constraints on tensor non-Gaussianity on small scales, allowing for further restrictions on the parameter ξ . This would enable a more thorough examination of the viability of the axion model as a candidate for inflationary scenarios. We leave the further investigation for future work.

It is pointed out that the tensor bispectrum is largely suppressed at interferometer scales due to decoherence of the phase and the finite observation time [104–106]. However, this does not completely preclude the detection of tensor non-Gaussianity on small scales. Several potential strategies have been proposed to circumvent this challenge. The most straightforward approach is to ex-

amine the angular three-point function of the SGWB energy density, which remains unaffected by the fast phases [107]. Additionally, one could consider special forms of the tensor bispectrum, such as stationary graviton non-Gaussianity [108] and ultra-squeezed configuration [109], where phase differences from source to detection are eliminated.

Finally, although we emphasize in the present work that the tensor non-Gaussianity of PGWs and the non-Gaussianity of scalar-induced GWs are fundamentally distinct, it is noteworthy that both can be related in certain inflationary models. For instance, in a single-field inflationary model with a periodic structure on the potential [110] or with sound speed resonance [78, 111], where the inflaton field perturbations can be significantly enhanced, both scalar-induced GWs and observable PGWs are produced. Previous investigation found that the tensor bispectrum mediated by enhanced inflaton perturbations peaks in the squeezed configuration [112]. Once the tensor non-Gaussianity is properly assessed, combined with PTA data, we can place constraints on the amplitude of the field perturbations. It was found that non-Gaussianities from different sectors (i.e., tensor and scalar) can share the same origin (i.e., the enhanced field perturbations), even though they enter the energy spectrum of GWs at different epochs: during inflation and after inflation, respectively. Therefore, by combining the tensor and scalar non-Gaussianities on small scales, the parameter space of such models can be significantly constrained.

ACKNOWLEDGMENTS

This work has been supported by the National Key Research and Development Program of China (No. 2023YFC2206704). ZZP is supported by the National Natural Science Foundation of China under Grant No. 12447166. YW is supported by the National Natural Science Foundation of China under Grant No. 12405057. LL is supported by the National Natural Science Foundation of China Grant under Grant No. 12433001.

-
- [1] Alexei A. Starobinsky, “Spectrum of relict gravitational radiation and the early state of the universe,” *JETP Lett.* **30**, 682–685 (1979).
 - [2] V. A. Rubakov, M. V. Sazhin, and A. V. Veryaskin, “Graviton Creation in the Inflationary Universe and the Grand Unification Scale,” *Phys. Lett. B* **115**, 189–192 (1982).
 - [3] Maresuke Shiraishi, Daisuke Nitta, Shuichiro Yokoyama, Kiyotomo Ichiki, and Keitaro Takahashi, “Cosmic microwave background bispectrum of tensor passive modes induced from primordial magnetic fields,” *Phys. Rev. D* **83**, 123003 (2011), arXiv:1103.4103 [astro-ph.CO].
 - [4] Maresuke Shiraishi, “Parity violation of primordial magnetic fields in the CMB bispectrum,” *JCAP* **06**, 015 (2012), arXiv:1202.2847 [astro-ph.CO].
 - [5] Aniket Agrawal, Tomohiro Fujita, and Eiichiro Komatsu, “Large tensor non-Gaussianity from axion-gauge field dynamics,” *Phys. Rev. D* **97**, 103526 (2018), arXiv:1707.03023 [astro-ph.CO].
 - [6] Emanuela Dimastrogiovanni, Matteo Fasiello, Giandomaso Tasinato, and David Wands, “Tensor non-Gaussianities from Non-minimal Coupling to the Inflaton,” *JCAP* **02**, 008 (2019), arXiv:1810.08866 [astro-ph.CO].
 - [7] Garrett Goon, Kurt Hinterbichler, Austin Joyce, and

- Mark Trodden, “Shapes of gravity: Tensor non-Gaussianity and massive spin-2 fields,” *JHEP* **10**, 182 (2019), arXiv:1812.07571 [hep-th].
- [8] Aniket Agrawal, Tomohiro Fujita, and Eiichiro Komatsu, “Tensor Non-Gaussianity from Axion-Gauge-Fields Dynamics : Parameter Search,” *JCAP* **06**, 027 (2018), arXiv:1802.09284 [astro-ph.CO].
- [9] Mohammad Ali Gorji, Seyed Ali Hosseini Mansoori, and Hassan Firouzjahi, “Inflation with multiple vector fields and non-Gaussianities,” *JCAP* **11**, 041 (2020), arXiv:2008.08195 [astro-ph.CO].
- [10] Laura Iacconi, Matteo Fasiello, Hooshyar Assadullahi, and David Wands, “Small-scale Tests of Inflation,” *JCAP* **12**, 005 (2020), arXiv:2008.00452 [astro-ph.CO].
- [11] Ogan Ozsoy, Maria Mylova, Susha Parameswaran, Cari Powell, Gianmassimo Tasinato, and Ivonne Zavala, “Squeezed tensor non-Gaussianity in non-attractor inflation,” *JCAP* **09**, 036 (2019), arXiv:1902.04976 [hep-th].
- [12] Guillem Domènech, Takashi Hiramatsu, Chunshan Lin, Misao Sasaki, Maresuke Shiraishi, and Yi Wang, “CMB Scale Dependent Non-Gaussianity from Massive Gravity during Inflation,” *JCAP* **05**, 034 (2017), arXiv:1701.05554 [astro-ph.CO].
- [13] Tomohiro Fujita, Shuntaro Mizuno, and Shinji Mukohyama, “Primordial Tensor Non-Gaussianity from Massive Gravity,” *JCAP* **01**, 023 (2020), arXiv:1909.07563 [astro-ph.CO].
- [14] Xian Gao, Tsutomu Kobayashi, Masahide Yamaguchi, and Jun’ichi Yokoyama, “Primordial non-Gaussianities of gravitational waves in the most general single-field inflation model,” *Phys. Rev. Lett.* **107**, 211301 (2011), arXiv:1108.3513 [astro-ph.CO].
- [15] Xian Gao, Tsutomu Kobayashi, Maresuke Shiraishi, Masahide Yamaguchi, Jun’ichi Yokoyama, and Shuichiro Yokoyama, “Full bispectra from primordial scalar and tensor perturbations in the most general single-field inflation model,” *PTEP* **2013**, 053E03 (2013), arXiv:1207.0588 [astro-ph.CO].
- [16] Sayantan Choudhury and Supratik Pal, “Primordial non-Gaussian features from DBI Galileon inflation,” *Eur. Phys. J. C* **75**, 241 (2015), arXiv:1210.4478 [hep-th].
- [17] Yongqing Huang, Anzhong Wang, Razieh Yousefi, and Tao Zhu, “Primordial non-Gaussianity of gravitational waves in Hořava-Lifshitz gravity,” *Phys. Rev. D* **88**, 023523 (2013), arXiv:1304.1556 [hep-th].
- [18] Shingo Akama, Shin’ichi Hirano, and Tsutomu Kobayashi, “Primordial tensor non-Gaussianities from general single-field inflation with non-Bunch-Davies initial states,” *Phys. Rev. D* **102**, 023513 (2020), arXiv:2003.10686 [gr-qc].
- [19] Abhishek Naskar and Supratik Pal, “Generic 3-point statistics with tensor modes in light of Swampland and Trans-Planckian Censorship Conjecture,” *Eur. Phys. J. C* **82**, 900 (2022), arXiv:2003.14066 [astro-ph.CO].
- [20] Sugumi Kanno and Misao Sasaki, “Graviton non-gaussianity in α -vacuum,” *JHEP* **08**, 210 (2022), arXiv:2206.03667 [hep-th].
- [21] Jinn-Ouk Gong, Maria Mylova, and Misao Sasaki, “New shape of parity-violating graviton non-Gaussianity,” *JHEP* **10**, 140 (2023), arXiv:2303.05178 [hep-th].
- [22] Valerio De Luca, Gabriele Franciolini, Alex Kehagias, Antonio Riotto, and Maresuke Shiraishi, “Constraining graviton non-Gaussianity through the CMB bispectra,” *Phys. Rev. D* **100**, 063535 (2019), arXiv:1908.00366 [astro-ph.CO].
- [23] Juan M. Maldacena and Guilherme L. Pimentel, “On graviton non-Gaussianities during inflation,” *JHEP* **09**, 045 (2011), arXiv:1104.2846 [hep-th].
- [24] Maresuke Shiraishi, Daisuke Nitta, and Shuichiro Yokoyama, “Parity Violation of Gravitons in the CMB Bispectrum,” *Prog. Theor. Phys.* **126**, 937–959 (2011), arXiv:1108.0175 [astro-ph.CO].
- [25] Arthur Lue, Li-Min Wang, and Marc Kamionkowski, “Cosmological signature of new parity violating interactions,” *Phys. Rev. Lett.* **83**, 1506–1509 (1999), arXiv:astro-ph/9812088.
- [26] Stephon Alexander and Jerome Martin, “Birefringent gravitational waves and the consistency check of inflation,” *Phys. Rev. D* **71**, 063526 (2005), arXiv:hep-th/0410230.
- [27] Jiro Soda, Hideo Kodama, and Masato Nozawa, “Parity Violation in Graviton Non-gaussianity,” *JHEP* **08**, 067 (2011), arXiv:1106.3228 [hep-th].
- [28] Nicola Bartolo and Giorgio Orlando, “Parity breaking signatures from a Chern-Simons coupling during inflation: the case of non-Gaussian gravitational waves,” *JCAP* **07**, 034 (2017), arXiv:1706.04627 [astro-ph.CO].
- [29] Nicola Bartolo, Giorgio Orlando, and Maresuke Shiraishi, “Measuring chiral gravitational waves in Chern-Simons gravity with CMB bispectra,” *JCAP* **01**, 050 (2019), arXiv:1809.11170 [astro-ph.CO].
- [30] Maria Mylova, “Chiral primordial gravitational waves in extended theories of Scalar-Tensor gravity,” (2019), arXiv:1912.00800 [gr-qc].
- [31] Nicola Bartolo, Luca Caloni, Giorgio Orlando, and Angelo Ricciardone, “Tensor non-Gaussianity in chiral scalar-tensor theories of gravity,” *JCAP* **03**, 073 (2021), arXiv:2008.01715 [astro-ph.CO].
- [32] Abhishek Naskar and Supratik Pal, “Non-Gaussian Features of Primordial Gravitational Waves,” *Phys. Rev. D* **98**, 083520 (2018), arXiv:1806.08178 [astro-ph.CO].
- [33] Abhishek Naskar and Supratik Pal, “Enhanced tensor non-Gaussianities in presence of a source,” *Eur. Phys. J. C* **80**, 1158 (2020), arXiv:1906.08558 [astro-ph.CO].
- [34] Lorenzo Bordin and Giovanni Cabass, “Graviton non-Gaussianities and Parity Violation in the EFT of Inflation,” *JCAP* **07**, 014 (2020), arXiv:2004.00619 [astro-ph.CO].
- [35] Giovanni Cabass, David Stefanyszyn, Jakub Supel, and Ayngaran Thavanesan, “On graviton non-Gaussianities in the Effective Field Theory of Inflation,” *JHEP* **10**, 154 (2022), arXiv:2209.00677 [hep-th].
- [36] Maresuke Shiraishi and Toyokazu Sekiguchi, “First observational constraints on tensor non-Gaussianity sourced by primordial magnetic fields from cosmic microwave background,” *Phys. Rev. D* **90**, 103002 (2014), arXiv:1304.7277 [astro-ph.CO].
- [37] Maresuke Shiraishi, Michele Liguori, and James R. Fergusson, “Observed parity-odd CMB temperature bispectrum,” *JCAP* **01**, 007 (2015), arXiv:1409.0265 [astro-ph.CO].
- [38] Maresuke Shiraishi, Michele Liguori, and James R. Fergusson, “CMB bounds on tensor-scalar-scalar inflationary correlations,” *JCAP* **01**, 016 (2018), arXiv:1710.06778 [astro-ph.CO].

- [39] Y. Akrami et al. (Planck), “Planck 2018 results. IX. Constraints on primordial non-Gaussianity,” *Astron. Astrophys.* **641**, A9 (2020), arXiv:1905.05697 [astro-ph.CO].
- [40] Maresuke Shiraishi, “Tensor Non-Gaussianity Search: Current Status and Future Prospects,” *Front. Astron. Space Sci.* **6**, 49 (2019), arXiv:1905.12485 [astro-ph.CO].
- [41] Oliver H. E. Philcox and Maresuke Shiraishi, “Non-Gaussianity Beyond the Scalar Sector: A Search for Tensor and Mixed Tensor-Scalar Bispectra with Planck Data,” (2024), arXiv:2409.10595 [astro-ph.CO].
- [42] Pau Amaro-Seoane et al. (LISA), “Laser Interferometer Space Antenna,” (2017), arXiv:1702.00786 [astro-ph.IM].
- [43] Wen-Hong Ruan, Zong-Kuan Guo, Rong-Gen Cai, and Yuan-Zhong Zhang, “Taiji program: Gravitational-wave sources,” *Int. J. Mod. Phys. A* **35**, 2050075 (2020), arXiv:1807.09495 [gr-qc].
- [44] Jun Luo et al. (TianQin), “TianQin: a space-borne gravitational wave detector,” *Class. Quant. Grav.* **33**, 035010 (2016), arXiv:1512.02076 [astro-ph.IM].
- [45] Heng Xu et al., “Searching for the Nano-Hertz Stochastic Gravitational Wave Background with the Chinese Pulsar Timing Array Data Release I,” *Res. Astron. Astrophys.* **23**, 075024 (2023), arXiv:2306.16216 [astro-ph.HE].
- [46] J. Antoniadis et al. (EPTA), “The second data release from the European Pulsar Timing Array - I. The dataset and timing analysis,” *Astron. Astrophys.* **678**, A48 (2023), arXiv:2306.16224 [astro-ph.HE].
- [47] J. Antoniadis et al. (EPTA, InPTA:), “The second data release from the European Pulsar Timing Array - III. Search for gravitational wave signals,” *Astron. Astrophys.* **678**, A50 (2023), arXiv:2306.16214 [astro-ph.HE].
- [48] Andrew Zic et al., “The Parkes Pulsar Timing Array third data release,” *Publ. Astron. Soc. Austral.* **40**, e049 (2023), arXiv:2306.16230 [astro-ph.HE].
- [49] Daniel J. Reardon et al., “Search for an Isotropic Gravitational-wave Background with the Parkes Pulsar Timing Array,” *Astrophys. J. Lett.* **951**, L6 (2023), arXiv:2306.16215 [astro-ph.HE].
- [50] Gabriella Agazie et al. (NANOGrav), “The NANOGrav 15 yr Data Set: Evidence for a Gravitational-wave Background,” *Astrophys. J. Lett.* **951**, L8 (2023), arXiv:2306.16213 [astro-ph.HE].
- [51] Gabriella Agazie et al. (NANOGrav), “The NANOGrav 15 yr Data Set: Observations and Timing of 68 Millisecond Pulsars,” *Astrophys. J. Lett.* **951**, L9 (2023), arXiv:2306.16217 [astro-ph.HE].
- [52] Matthew T. Miles et al., “The MeerKAT Pulsar Timing Array: The 4.5-year data release and the noise and stochastic signals of the millisecond pulsar population,” (2024), 10.1093/mnras/stae2572, arXiv:2412.01148 [astro-ph.HE].
- [53] Matthew T. Miles et al., “The MeerKAT Pulsar Timing Array: The first search for gravitational waves with the MeerKAT radio telescope,” (2024), 10.1093/mnras/stae2571, arXiv:2412.01153 [astro-ph.HE].
- [54] Keisuke Inomata, Kazunori Kohri, and Takahiro Terada, “Detected stochastic gravitational waves and subsolar-mass primordial black holes,” *Phys. Rev. D* **109**, 063506 (2024), arXiv:2306.17834 [astro-ph.CO].
- [55] Zu-Cheng Chen, Chen Yuan, and Qing-Guo Huang, “Pulsar Timing Array Constraints on Primordial Black Holes with NANOGrav 11-Year Dataset,” *Phys. Rev. Lett.* **124**, 251101 (2020), arXiv:1910.12239 [astro-ph.CO].
- [56] Lang Liu, Zu-Cheng Chen, and Qing-Guo Huang, “Implications for the non-Gaussianity of curvature perturbation from pulsar timing arrays,” *Phys. Rev. D* **109**, L061301 (2024), arXiv:2307.01102 [astro-ph.CO].
- [57] Gabriele Franciolini, Antonio Iovino, Junior., Ville Vaskonen, and Hardi Veermae, “Recent Gravitational Wave Observation by Pulsar Timing Arrays and Primordial Black Holes: The Importance of Non-Gaussianities,” *Phys. Rev. Lett.* **131**, 201401 (2023), arXiv:2306.17149 [astro-ph.CO].
- [58] Seyed Ali Hosseini Mansoori, Fereshteh Felegray, Alireza Talebian, and Mohammad Sami, “PBHs and GWs from T^2 -inflation and NANOGrav 15-year data,” *JCAP* **08**, 067 (2023), arXiv:2307.06757 [astro-ph.CO].
- [59] Sai Wang, Zhi-Chao Zhao, Jun-Peng Li, and Qing-Hua Zhu, “Implications of pulsar timing array data for scalar-induced gravitational waves and primordial black holes: Primordial non-Gaussianity fNL considered,” *Phys. Rev. Res.* **6**, L012060 (2024), arXiv:2307.00572 [astro-ph.CO].
- [60] Jia-Heng Jin, Zu-Cheng Chen, Zhu Yi, Zhi-Qiang You, Lang Liu, and You Wu, “Confronting sound speed resonance with pulsar timing arrays,” *JCAP* **09**, 016 (2023), arXiv:2307.08687 [astro-ph.CO].
- [61] Lang Liu, Zu-Cheng Chen, and Qing-Guo Huang, “Probing the equation of state of the early Universe with pulsar timing arrays,” *JCAP* **11**, 071 (2023), arXiv:2307.14911 [astro-ph.CO].
- [62] Qing-Hua Zhu, Zhi-Chao Zhao, Sai Wang, and Xin Zhang, “Unraveling the early universe’s equation of state and primordial black hole production with PTA, BBN, and CMB observations*,” *Chin. Phys. C* **48**, 125105 (2024), arXiv:2307.13574 [astro-ph.CO].
- [63] Zhu Yi, Zhi-Qiang You, You Wu, Zu-Cheng Chen, and Lang Liu, “Exploring the NANOGrav signal and planet-mass primordial black holes through Higgs inflation,” *JCAP* **06**, 043 (2024), arXiv:2308.14688 [astro-ph.CO].
- [64] Keisuke Harigaya, Keisuke Inomata, and Takahiro Terada, “Induced gravitational waves with kination era for recent pulsar timing array signals,” *Phys. Rev. D* **108**, 123538 (2023), arXiv:2309.00228 [astro-ph.CO].
- [65] Shyam Balaji, Guillem Domènech, and Gabriele Franciolini, “Scalar-induced gravitational wave interpretation of PTA data: the role of scalar fluctuation propagation speed,” *JCAP* **10**, 041 (2023), arXiv:2307.08552 [gr-qc].
- [66] Zhu Yi, Zhi-Qiang You, and You Wu, “Model-independent reconstruction of the primordial curvature power spectrum from PTA data,” *JCAP* **01**, 066 (2024), arXiv:2308.05632 [astro-ph.CO].
- [67] Zhi-Qiang You, Zhu Yi, and You Wu, “Constraints on primordial curvature power spectrum with pulsar timing arrays,” *JCAP* **11**, 065 (2023), arXiv:2307.04419 [gr-qc].
- [68] Lang Liu, You Wu, and Zu-Cheng Chen, “Simultaneously probing the sound speed and equation of state of the early Universe with pulsar timing arrays,” *JCAP* **04**, 011 (2024), arXiv:2310.16500 [astro-ph.CO].
- [69] Sayantan Choudhury, Kritartha Dey, Ahaskar Karde, Sudhakar Panda, and M. Sami, “Primordial non-Gaussianity as a saviour for PBH overproduction in SIGWs generated by pulsar timing arrays for

- Galileon inflation,” *Phys. Lett. B* **856**, 138925 (2024), arXiv:2310.11034 [astro-ph.CO].
- [70] Sayantan Choudhury, Kritartha Dey, and Ahaskar Karde, “Untangling PBH overproduction in w -SIGWs generated by Pulsar Timing Arrays for MST-EFT of single field inflation,” (2023), arXiv:2311.15065 [astro-ph.CO].
- [71] Guillem Domènec, Shi Pi, Ao Wang, and Jianing Wang, “Induced gravitational wave interpretation of PTA data: a complete study for general equation of state,” *JCAP* **08**, 054 (2024), arXiv:2402.18965 [astro-ph.CO].
- [72] Zu-Cheng Chen and Lang Liu, “Can we distinguish the adiabatic fluctuations and isocurvature fluctuations with pulsar timing arrays?” (2024), arXiv:2402.16781 [astro-ph.CO].
- [73] Sayantan Choudhury, Siddhant Ganguly, Sudhakar Panda, Soumitra SenGupta, and Pranjal Tiwari, “Obviating PBH overproduction for SIGWs generated by pulsar timing arrays in loop corrected EFT of bounce,” *JCAP* **09**, 013 (2024), arXiv:2407.18976 [astro-ph.CO].
- [74] Sayantan Choudhury and M. Sami, “Large fluctuations and Primordial Black Holes,” *Phys. Rept.* **1103**, 1–276 (2025), arXiv:2407.17006 [gr-qc].
- [75] Zu-Cheng Chen, Jun Li, Lang Liu, and Zhu Yi, “Probing the speed of scalar-induced gravitational waves with pulsar timing arrays,” *Phys. Rev. D* **109**, L101302 (2024), arXiv:2401.09818 [gr-qc].
- [76] Chang Han et al., “Constraining inflation with nonminimal derivative coupling with the Parkes Pulsar Timing Array third data release,” (2024), arXiv:2412.09755 [gr-qc].
- [77] Sayantan Choudhury, Kritartha Dey, Siddhant Ganguly, Ahaskar Karde, Swapnil Kumar Singh, and Pranjal Tiwari, “Negative non-Gaussianity as a salvager for PBHs with PTAs in bounce,” (2024), arXiv:2409.18983 [astro-ph.CO].
- [78] Rong-gen Cai, Shi Pi, and Misao Sasaki, “Gravitational Waves Induced by non-Gaussian Scalar Perturbations,” *Phys. Rev. Lett.* **122**, 201101 (2019), arXiv:1810.11000 [astro-ph.CO].
- [79] Caner Unal, “Imprints of Primordial Non-Gaussianity on Gravitational Wave Spectrum,” *Phys. Rev. D* **99**, 041301 (2019), arXiv:1811.09151 [astro-ph.CO].
- [80] Chen Yuan and Qing-Guo Huang, “Gravitational waves induced by the local-type non-Gaussian curvature perturbations,” *Phys. Lett. B* **821**, 136606 (2021), arXiv:2007.10686 [astro-ph.CO].
- [81] Peter Adshead, Kaloian D. Lozanov, and Zachary J. Weiner, “Non-Gaussianity and the induced gravitational wave background,” *JCAP* **10**, 080 (2021), arXiv:2105.01659 [astro-ph.CO].
- [82] Chen Yuan, De-Shuang Meng, and Qing-Guo Huang, “Full analysis of the scalar-induced gravitational waves for the curvature perturbation with local-type non-Gaussianities,” *JCAP* **12**, 036 (2023), arXiv:2308.07155 [astro-ph.CO].
- [83] Xiang-Xi Zeng, Rong-Gen Cai, and Shao-Jiang Wang, “Multiple peaks in gravitational waves induced from primordial curvature perturbations with non-Gaussianity,” *JCAP* **10**, 045 (2024), arXiv:2406.05034 [astro-ph.CO].
- [84] Zhe Chang, Yu-Ting Kuang, Di Wu, Jing-Zhi Zhou, and Qing-Hua Zhu, “New constraints on primordial non-Gaussianity from missing two-loop contributions of scalar induced gravitational waves,” *Phys. Rev. D* **109**, L041303 (2024), arXiv:2311.05102 [astro-ph.CO].
- [85] Eiichiro Komatsu and David N. Spergel, “Acoustic signatures in the primary microwave background bispectrum,” *Phys. Rev. D* **63**, 063002 (2001), arXiv:astro-ph/0005036.
- [86] Daniel Babich, Paolo Creminelli, and Matias Zaldarriaga, “The Shape of non-Gaussianities,” *JCAP* **08**, 009 (2004), arXiv:astro-ph/0405356.
- [87] Nicola Bartolo, Valerie Domcke, Daniel G. Figueroa, Juan García-Bellido, Marco Peloso, Mauro Pieroni, Angelo Ricciardone, Mairi Sakellariadou, Lorenzo Sorbo, and Gianmassimo Tasinato, “Probing non-Gaussian Stochastic Gravitational Wave Backgrounds with LISA,” *JCAP* **11**, 034 (2018), arXiv:1806.02819 [astro-ph.CO].
- [88] Michele Maggiore, “Gravitational wave experiments and early universe cosmology,” *Phys. Rept.* **331**, 283–367 (2000), arXiv:gr-qc/9909001.
- [89] Michael S. Turner, Martin J. White, and James E. Lidsey, “Tensor perturbations in inflationary models as a probe of cosmology,” *Phys. Rev. D* **48**, 4613–4622 (1993), arXiv:astro-ph/9306029.
- [90] Latham A. Boyle and Paul J. Steinhardt, “Probing the early universe with inflationary gravitational waves,” *Phys. Rev. D* **77**, 063504 (2008), arXiv:astro-ph/0512014.
- [91] Wen Zhao and Yang Zhang, “Relic gravitational waves and their detection,” *Phys. Rev. D* **74**, 043503 (2006), arXiv:astro-ph/0604458.
- [92] Massimo Giovannini, “Primordial backgrounds of relic gravitons,” *Prog. Part. Nucl. Phys.* **112**, 103774 (2020), arXiv:1912.07065 [astro-ph.CO].
- [93] R. w. Hellings and G. s. Downs, “UPPER LIMITS ON THE ISOTROPIC GRAVITATIONAL RADIATION BACKGROUND FROM PULSAR TIMING ANALYSIS,” *Astrophys. J. Lett.* **265**, L39–L42 (1983).
- [94] G. Agazie et al. (International Pulsar Timing Array), “Comparing Recent Pulsar Timing Array Results on the Nanohertz Stochastic Gravitational-wave Background,” *Astrophys. J.* **966**, 105 (2024), arXiv:2309.00693 [astro-ph.HE].
- [95] N. Aghanim et al. (Planck), “Planck 2018 results. VI. Cosmological parameters,” *Astron. Astrophys.* **641**, A6 (2020), [Erratum: Astron. Astrophys. 652, C4 (2021)], arXiv:1807.06209 [astro-ph.CO].
- [96] Thomas J. Clarke, Edmund J. Copeland, and Adam Moss, “Constraints on primordial gravitational waves from the Cosmic Microwave Background,” *JCAP* **10**, 002 (2020), arXiv:2004.11396 [astro-ph.CO].
- [97] Joshua S. Speagle, “dynesty: a dynamic nested sampling package for estimating Bayesian posteriors and evidences,” *Mon. Not. Roy. Astron. Soc.* **493**, 3132–3158 (2020), arXiv:1904.02180 [astro-ph.IM].
- [98] Gregory Ashton et al., “BILBY: A user-friendly Bayesian inference library for gravitational-wave astronomy,” *Astrophys. J. Suppl.* **241**, 27 (2019), arXiv:1811.02042 [astro-ph.IM].
- [99] I. M. Romero-Shaw et al., “Bayesian inference for compact binary coalescences with bilby: validation and application to the first LIGO–Virgo gravitational-wave transient catalogue,” *Mon. Not. Roy. Astron. Soc.* **499**, 3295–3319 (2020), arXiv:2006.00714 [astro-ph.IM].
- [100] Christian T. Byrnes, Kazuya Koyama, Misao Sasaki,

- and David Wands, “Diagrammatic approach to non-Gaussianity from inflation,” *JCAP* **11**, 027 (2007), [arXiv:0705.4096 \[hep-th\]](#).
- [101] De-Shuang Meng, Chen Yuan, and Qing-guo Huang, “One-loop correction to the enhanced curvature perturbation with local-type non-Gaussianity for the formation of primordial black holes,” *Phys. Rev. D* **106**, 063508 (2022), [arXiv:2207.07668 \[astro-ph.CO\]](#).
- [102] Jessica L. Cook and Lorenzo Sorbo, “An inflationary model with small scalar and large tensor nongaussianities,” *JCAP* **11**, 047 (2013), [arXiv:1307.7077 \[astro-ph.CO\]](#).
- [103] Maresuke Shiraishi, Angelo Ricciardone, and Shohei Saga, “Parity violation in the CMB bispectrum by a rolling pseudoscalar,” *JCAP* **11**, 051 (2013), [arXiv:1308.6769 \[astro-ph.CO\]](#).
- [104] Peter Adshead and Eugene A. Lim, “3-pt Statistics of Cosmological Stochastic Gravitational Waves,” *Phys. Rev. D* **82**, 024023 (2010), [arXiv:0912.1615 \[astro-ph.CO\]](#).
- [105] N. Bartolo, V. De Luca, G. Franciolini, A. Lewis, M. Peloso, and A. Riotto, “Primordial Black Hole Dark Matter: LISA Serendipity,” *Phys. Rev. Lett.* **122**, 211301 (2019), [arXiv:1810.12218 \[astro-ph.CO\]](#).
- [106] N. Bartolo, V. De Luca, G. Franciolini, M. Peloso, D. Racco, and A. Riotto, “Testing primordial black holes as dark matter with LISA,” *Phys. Rev. D* **99**, 103521 (2019), [arXiv:1810.12224 \[astro-ph.CO\]](#).
- [107] N. Bartolo, D. Bertacca, S. Matarrese, M. Peloso, A. Ricciardone, A. Riotto, and G. Tasinato, “Anisotropies and non-Gaussianity of the Cosmological Gravitational Wave Background,” *Phys. Rev. D* **100**, 121501 (2019), [arXiv:1908.00527 \[astro-ph.CO\]](#).
- [108] Cari Powell and Gianmassimo Tasinato, “Probing a stationary non-Gaussian background of stochastic gravitational waves with pulsar timing arrays,” *JCAP* **01**, 017 (2020), [arXiv:1910.04758 \[gr-qc\]](#).
- [109] Emanuela Dimastrogiovanni, Matteo Fasiello, and Gianmassimo Tasinato, “Searching for Fossil Fields in the Gravity Sector,” *Phys. Rev. Lett.* **124**, 061302 (2020), [arXiv:1906.07204 \[astro-ph.CO\]](#).
- [110] Rong-Gen Cai, Zong-Kuan Guo, Jing Liu, Lang Liu, and Xing-Yu Yang, “Primordial black holes and gravitational waves from parametric amplification of curvature perturbations,” *JCAP* **06**, 013 (2020), [arXiv:1912.10437 \[astro-ph.CO\]](#).
- [111] Yi-Fu Cai, Chao Chen, Xi Tong, Dong-Gang Wang, and Sheng-Feng Yan, “When Primordial Black Holes from Sound Speed Resonance Meet a Stochastic Background of Gravitational Waves,” *Phys. Rev. D* **100**, 043518 (2019), [arXiv:1902.08187 \[astro-ph.CO\]](#).
- [112] Zhi-Zhang Peng, Cheng-Jun Fang, and Zong-Kuan Guo, “Tensor bispectrum mediated by an excited scalar field during inflation,” *Phys. Rev. D* **110**, 043538 (2024), [arXiv:2403.04617 \[astro-ph.CO\]](#).