A Fixed-Point Approach to History-Dependent Sweeping Processes

Matías Godoy, Manuel Torres-Valdebenito and Emilio Vilches

Received: date / Accepted: date

Abstract In this paper, we study the well-posedness of state-dependent and state-independent sweeping processes driven by prox-regular sets and perturbed by a history-dependent operator. Our approach, based on an enhanced version of Gronwall's lemma and fixed-point arguments, provides an efficient framework for analyzing sweeping processes. In particular, our findings recover all existing results for the class of Volterra sweeping processes and provide new insights into history-dependent sweeping processes. Finally, we apply our theoretical results to establish the well-posedness of a viscoelastic model with long memory.

Keywords sweeping processes \cdot fixed point theory \cdot history-dependent operator \cdot prox-regular sets

Mathematics Subject Classification (2000) 34A60 · 49J52 · 49J53

1 Introduction

The sweeping process is a differential inclusion involving normal cones to a family of moving sets. Since its introduction by J.-J. Moreau in a series of papers [20–22], it has become a natural mathematical modeling tool for problems with constraints, as well as for modeling various phenomena in contact

Manuel Torres-Valdebenito Departamento de Ingeniería Matemática, Universidad de Chile, Santiago, Chile manuel.torres@ug.uchile.cl

Emilio Vilches

Instituto de Ciencias de la Ingeniería, Universidad de O'Higgins, Rancagua, Chile emilio.vilches@uoh.cl

Matías Godoy

Facultad de Ingeniería, Arquitectura y Diseño, Universidad San Sebastián, Santiago, Chile matias.godoy@uss.cl

mechanics, electrical circuits, crowd motion problems, among others. We refer to [1, 2, 24] for further details.

The sweeping process, originally studied by Moreau for convex sets, was later extended to the context of non-convex sets by various authors. Notably, the seminal works of Lionel Thibault [29, 30] developed the well-posedness theory for a family of uniformly prox-regular sets, which constitute a natural and broad framework for sweeping processes. In parallel, the perturbed and state-dependent cases were developed, generalizing classical results in differential equations and paving the way for further advancements in mathematical modeling. See, for example, [6, 7, 14, 16, 25, 27, 32].

Recently, a new variant of the sweeping process was proposed in [10]. This variant, now known as the Volterra Sweeping Process, incorporates an integral term into the classical perturbed sweeping process. This integral term facilitates the generalization of Volterra differential equations and enables the modeling of constrained processes where the velocity depends on the trajectory at previous times. For well-posedness results, we refer to [4, 5, 33].

In this paper, we generalize all the aforementioned developments by considering a history-dependent operator, which accounts for constrained problems where the velocity depends on its history but not necessarily as an integral form. Our approach, which is based on an enhanced version of Gronwall's lemma proved in [33] and fixed-point arguments, provides an efficient framework for analyzing state-dependent and state-independent Volterra sweeping processes.

It is worth emphasizing that history-dependent operators have been used to model viscoelastic materials [19, 23, 24, 28], as in these materials, the displacement field depends on the trajectory. Many materials can be considered viscoelastic due to their properties, including soft and hard tissues (e.g., skin, cartilage, bone), synthetic polymers, elastomers, and others. For more details, we refer to [8,18]. We apply our theoretical results to prove the well-posedness of a problem in contact mechanics with long memory.

The paper is organized as follows. After presenting some mathematical preliminaries, in Section 3, we summarize the main assumptions used throughout the paper. In Section 4, we establish the existence and uniqueness of solutions for state-independent sweeping processes driven by prox-regular sets and perturbed by history-dependent operators. Subsequently, in Section 5, we demonstrate the existence of solutions for state-dependent sweeping processes driven by prox-regular sets and perturbed by history-dependent operators. Then, in Section 6, as a consequence of our results, we establish the existence of solutions for Volterra sweeping processes. The paper concludes with an application to viscoelastic models with long memory.

2 Preliminaries

Let $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ be a real Hilbert space. As usual, the norm of \mathcal{H} is defined as $||x|| := \sqrt{\langle x, x \rangle}$ and the closed unit ball is denoted as \mathbb{B} . The real numbers

will be denoted by \mathbb{R} and for T > 0, we set I := [0, T]. Let $S \subset \mathcal{H}$ be a nonempty closed set. The distance from a set S to a point $x \in \mathcal{H}$ is defined as $d_S(x) := \inf_{y \in S} ||x - y||$. The set of points where the distance from S to $x \in \mathcal{H}$ is attained is denoted by $\operatorname{Proj}_S(x)$. This set is possibly empty and, when it consists of a single point, is denoted by $\operatorname{Proj}_S(x)$.

Given two closed sets $A, B \subset \mathcal{H}$, the excess of A over B is defined as

$$\operatorname{exc}(A;B) := \sup_{x \in A} d_B(x)$$

Moreover, it is well-known that $exc(A; B) = \inf\{\varepsilon > 0 : A \subset B + \varepsilon \mathbb{B}\}$. Additionally, the Hausdorff distance between A and B is defined as

$$\operatorname{Haus}(A, B) := \max\{\operatorname{exc}(A; B), \operatorname{exc}(B; A)\},\$$

which can be characterized through the formula (see, e.g., [3, Lemma 3.74]):

$$\operatorname{Haus}(A, B) = \sup_{x \in \mathcal{H}} |d_A(x) - d_B(x)|.$$

We refer to [3] for more details.

2.1 Convex and Variational Analysis Tools

Given a function $f: \mathcal{H} \to \mathbb{R} \cup \{+\infty\}$, we say that $f \in \Gamma_0(\mathcal{H})$ if f is proper, convex, and lower semicontinuous (l.s.c.). Given $f \in \Gamma_0(\mathcal{H})$, we say that x^* is an element of the Fénchel subdifferential of f at $x \in \mathcal{H}$, denoted by $\partial f(x)$, if

$$f(y) \ge f(x) + \langle x^*, y - x \rangle$$
 for all $y \in \mathcal{H}$.

The Legendre-Fenchel conjugate $f^* \colon \mathcal{H} \to \mathbb{R} \cup \{+\infty\}$ of a function $f \in \Gamma_0(\mathcal{H})$ is defined by

$$f^*(x^*) = \sup_{x \in \mathcal{H}} \{ \langle x^*, x \rangle - f(x) \}$$
 for all $x^* \in \mathcal{H}$.

Moreover, it is well-known that the following equivalences hold:

$$x^* \in \partial f(x) \iff x \in \partial f^*(x^*) \iff \langle x^*, x \rangle = f(x) + f^*(x^*).$$

Besides, given a non-empty, closed, and convex set $C \subset \mathcal{H}$, the indicator function of C is defined as $\iota_C : \mathcal{H} \to \mathbb{R} \cup \{+\infty\}$, given by

$$\iota_C(x) = \begin{cases} 0 & \text{if } x \in C; \\ +\infty & \text{if } x \notin C, \end{cases}$$

and the support function of C is defined as $\sigma_C \colon \mathcal{H} \to \mathbb{R} \cup \{+\infty\}$, given by

$$\sigma_C(x^*) = \sup_{x \in C} \langle x^*, x \rangle \quad \text{for all } x^* \in \mathcal{H}.$$

Moreover, the following relations hold $\sigma_C = (\iota_C)^*$ and $\iota_C = (\sigma_C)^*$.

Given a closed and nonempty set $S \subset \mathcal{H}$ and $x \in S$. We say that v belongs to the proximal normal cone $N^P(S; x)$ if there exists $\sigma \geq 0$ such that

$$\langle v, y - x \rangle \le \sigma \|y - x\|^2$$
 for all $y \in S$.

Whenever $x \notin S$, we set $N^P(S; x) = \emptyset$. We refer to [9] for more details about proximal calculus. Now, we recall the concept of a uniformly prox-regular set. Introduced by Federer in the finite-dimensional setting (see [15]) and later developed by Rockafellar, Poliquin, and Thibault in [26], prox-regularity generalizes and unifies the classes of convex sets and nonconvex bodies with C^2 boundary.

Definition 2.1 Let S be a closed subset of \mathcal{H} and $\rho \in]0, +\infty]$. We say that S is ρ -uniformly prox-regular provided that, for all $x \in S$ and all $v \in N^P(S; x) \cap \mathbb{B}$ one has

$$x \in \operatorname{Proj}_{S}(x+tv)$$
 for any $t \leq \rho$.

It is important to emphasize that convex sets are ρ -uniformly prox-regular for any $\rho > 0$. The following result summarizes the main characterization of uniform prox-regularity. We refer to [11,31] for further results.

Proposition 2.1 Let S be a nonempty closed subset of \mathcal{H} and $\rho \in]0, +\infty]$. The following assertions are equivalent:

- (i) The set S is ρ -uniformly prox-regular.
- (ii) For all $x, x' \in S$ and $\zeta \in N^P(S; x)$ one has

$$\langle \zeta, x' - x \rangle \le \frac{1}{2\rho} \|\zeta\| \|x' - x\|^2.$$

(iii) For any $x_i \in S$, $\zeta_i \in N^P(S; x_i) \cap \mathbb{B}$ with i = 1, 2 one has

$$\langle \zeta_1 - \zeta_2, x_1 - x_2 \rangle \ge -\frac{1}{\rho} \|x_1 - x_2\|^2.$$

that is, the set-valued mapping $x \mapsto N^P(S; x) \cap \mathbb{B}$ is $\frac{1}{\rho}$ -hypomonotone.

(iv) For any positive $\gamma < 1$ the projection mapping $\operatorname{proj}_S^{\cdot}$ is well-defined and Lipschitz continuous on $U_{\rho}^{\gamma}(S) := \{z \in \mathcal{H} : d_S(x) < \gamma \rho\}$ with $1/(1-\gamma)$ as a Lipschitz constant, i.e.,

$$\|\operatorname{proj}_{S}(x_{1}) - \operatorname{proj}_{S}(x_{2})\| \leq \frac{1}{1-\gamma} \|x_{1} - x_{2}\| \quad \text{for all } x_{1}, x_{2} \in U_{\rho}^{\gamma}(S).$$

2.2 Nonlinear Analysis Tools

Let A be a bounded subset of \mathcal{H} . We define the Kuratokwki measure of noncompactness of A, $\alpha(A)$, as

 $\alpha(A) := \inf \left\{ d > 0 : A \text{ admits a finite cover by sets of diameter} \le d \right\},\$

and the Hausdorff measure of non-compactness of A, $\beta(A)$, as

 $\beta(A) := \inf \{ r > 0 : A \text{ can be covered by finitely many ball of radius } r \}.$

In *Hilbert spaces* the relation between these two concepts is given by the inequality for $A \subset \mathcal{H}$ bounded:

$$\sqrt{2}\beta(A) \le \alpha(A) \le 2\beta(A).$$

The following proposition gathers the main properties of *Kuratowski and Hausdorff measures of non-compactness* (see [12, Proposition 7.2], [13, Section 9.2, Proposition 9.1]).

Proposition 2.2 Let \mathcal{H} be an infinite dimensional Hilbert space and B, B_1 , B_2 be bounded subsets of \mathcal{H} . Let γ be either the Kuratowski or the Hausdorff measures of non-compactness. Then:

(a) $\gamma(B) = 0$ if and only if \overline{B} is compact. (a) $\gamma(\lambda B) = |\lambda|\gamma(B)$ for every $\lambda \in \mathbb{R}$. (b) $\gamma(B_1 + B_2) \leq \gamma(B_1) + \gamma(B_2)$. (c) $B_1 \subset B_2$ implies $\gamma(B_1) \leq \gamma(B_2)$. (d) $\gamma(\operatorname{conv} B) = \gamma(B)$. (e) $\gamma(\overline{B}) = \gamma(B)$.

We will denote by $C(I; \mathcal{H})$ the set of all *continuous functions* from I to \mathcal{H} . The norm of the *uniform convergence* on $C(I; \mathcal{H})$ will be denoted by $\|\cdot\|_{\infty}$. We denote by $L^1([0,T]; \mathcal{H})$ the space of \mathcal{H} -valued Lebesgue integrable functions defined over the interval [0,T]. We say that $x \in AC([0,T]; \mathcal{H})$ if there exists $f \in L^1([0,T]; \mathcal{H})$ and $x_0 \in \mathcal{H}$ such that

$$x(t) = x_0 + \int_0^t f(s) ds \ \forall t \in [0, T].$$

The following result, proved in [33], is an enhanced version of the classical Gronwall inequality.

Lemma 2.1 (Enhanced Gronwall Inequality) Let I := [0,T], and let $\rho: I \to \mathbb{R}$ be a nonnegative absolutely continuous function. Let $K_1, K_2, \varepsilon: I \to \mathbb{R}_+$, and $K_3: I \times I \to \mathbb{R}_+$ be nonnegative measurable functions such that

$$t \mapsto K_1(t) \text{ and } t \mapsto K_2(t) \int_0^t K_3(t,s) ds \text{ are integrable}$$

Suppose that

$$\dot{\rho}(t) \le \varepsilon(t) + K_1(t)\rho(t) + K_2(t) \int_0^t K_3(t,s)\rho(s)ds \quad \text{for a.e. } t \in I.$$

Then, one has

$$\rho(t) \le \rho(0) \exp\left(\int_0^t \gamma(s) ds\right) + \int_0^t \varepsilon(s) \exp\left(\int_s^t \gamma(\tau) d\tau\right) ds \quad \text{for all } t \in I,$$

where $\gamma(t) := K_1(t) + K_2(t) \int_0^t K_3(t,s) ds.$

The following result will be used in the proof of Theorem 4.1.

Lemma 2.2 Let $\Theta: I \to \mathbb{R}$ be a nonnegative continuous function. Assume that α and β are two nonnegative integrable functions such that

$$\Theta^{2}(t) \leq \int_{0}^{t} \alpha(s)\Theta(s)ds + \int_{0}^{t} \beta(s)\Theta^{2}(s)ds \quad \text{ for all } t \in I.$$

Then, for all $t \in I$, one has

$$\Theta(t) \leq \int_0^t \exp\left(\int_s^t \beta(\tau) d\tau\right) \alpha(s) ds.$$

Proof We will prove that

$$\Theta(t) \le \int_0^t \alpha(\tau) d\tau + \int_0^t \beta(\tau) \Theta(\tau) d\tau \text{ for all } t \in I,$$
(1)

which, by virtue of the classical Gronwall's inequality, will imply the result. If $\Theta \equiv 0$ the result is obvious. Otherwise, assume that $\Theta \not\equiv 0$ and define $t^* := \inf\{s \in I : \Theta(s) > 0\}$. On the one hand, we observe that $\Theta(t) = 0$ for any $t \in [0, t^*]$, hence, (1) holds for any $t \in [0, t^*]$. On the other hand, from hypothesis, it is clear that for all $t \in]t^*, T]$, one has

$$\sup_{s \in [0,t]} \Theta(s)^2 \le \sup_{s \in [0,t]} \Theta(s) \left(\int_0^t \alpha(\tau) d\tau + \int_0^t \beta(\tau) \Theta(\tau) d\tau \right),$$

which, being $\sup_{s \in [0,t]} \Theta(s) > 0$ for all $t \in [t^*, T]$, implies that for $t \in [t^*, T]$

$$\Theta(t) \le \int_0^t \alpha(\tau) d\tau + \int_0^t \beta(\tau) \Theta(\tau) d\tau,$$

which finishes the proof.

An operator $\mathcal{R}: C(I; \mathcal{H}) \to C(I; \mathcal{H})$ is called an *History-Dependent Operator* if there exists $\kappa > 0$, such that for all $x, y \in C(I; \mathcal{H})$ and $t \in I$, one has

$$\|\mathcal{R}(x)(t) - \mathcal{R}(y)(t)\| \le \kappa \int_0^t \|x(s) - y(s)\| ds.$$

For more details, we refer to [28, Chapter 2]. The following result provides a fixed point result for history-dependent operators (see, e.g., [28, Chapter 2]).

Proposition 2.3 Let $S: C(I; E) \to C(I; F)$ be a history dependent operator, where E, F are two Banach spaces. Then, the operator S has a unique fixed point.

3 Technical assumptions

For ease of presentation, in this section, we gather the hypotheses used along the paper.

- (\mathcal{H}^f) The function $f: I \times \mathcal{H} \to \mathcal{H}$ satisfies
 - (i) For each $x \in \mathcal{H}$, the map $t \mapsto f(t, x)$ is measurable.
 - (ii) There exists two nonnegative integrable functions α and β such that

$$||f(t,x)|| \le \alpha(t)||x|| + \beta(t) \quad \text{for all } (t,x) \in I \times \mathcal{H}.$$

(iii) For each r > 0, there exists a nonnegative constant κ_f^r such that

 $||f(t,x) - f(t,y)|| \le \kappa_f^r ||x - y|| \quad \text{for all } x, y \in r\mathbb{B} \text{ and } t \in I.$

- (\mathcal{H}^g) The function $g: I \times I \times \mathcal{H} \to \mathcal{H}$ satisfies
 - (a) For each $x \in \mathcal{H}$, the map $(t, s) \mapsto g(t, s, x)$ is measurable.
 - (b) For all r > 0, there exists an integrable function $\mu_r \colon I \to \mathbb{R}_+$ such that for all $(t, s) \in D$

$$\|g(t,s,x) - g(t,s,y)\| \le \mu_r(t) \|x - y\| \text{ for all } x, y \in r\mathbb{B}$$

Here $D := \{(t, s) \in I \times I : s \leq t\}.$

(c) There exists a nonnegative integrable function $\sigma\colon D\to\mathbb{R}$ such that

$$||g(t,s,x)|| \le \sigma(t,s)(1+||x||)$$
 for all $(t,s) \in D$ and $x \in \mathcal{H}$.

 $(\mathcal{H}^{\mathcal{R}})$ The operator $\mathcal{R}: C(I; \mathcal{H}) \to C(I; \mathcal{H})$ is history-dependent of constant $\kappa_{\mathcal{R}}$, that is, there exists $\kappa_{\mathcal{R}} \ge 0$ such that for all $x, y \in C(I; \mathcal{H})$ one has

$$\|\mathcal{R}(x)(t) - \mathcal{R}(y)(t)\| \le \kappa_{\mathcal{R}} \int_0^t \|x(s) - y(s)\| ds \text{ for all } t \in I.$$

 (\mathcal{H}^C) The map $C: [0,T] \Rightarrow \mathcal{H}$ has nonempty, closed and ρ -uniformly proxregular values, for some $\rho > 0$. Moreover, there exists an absolutely continuous function $v: [0,T] \to \mathbb{R}$ such that

$$\operatorname{Haus}(C(t), C(s)) \le |v(t) - v(s)| \quad \text{for all } t, s \in I.$$

- (\mathcal{H}_x^C) The map $C \colon [0,T] \times \mathcal{H} \rightrightarrows \mathcal{H}$ has nonempty, closed and ρ -uniformly proxregular values, for some $\rho > 0$. Moreover, the following conditions hold:
 - (i) There exists an absolutely continuous function $v \colon [0,T] \to \mathbb{R}$ and $L \in [0,1[$, such that

 $\operatorname{Haus}(C(t,x),C(s,y)) \le |v(t) - v(s)| + L ||x - y|| \text{ for all } t, s \in I, x, y \in \mathcal{H}.$

(ii) For any $A, B \subset \mathcal{H}$ bounded and r > 0, the set $C(t, A) \cap B \cap r\mathbb{B}$ is relatively compact.

4 History-Dependent Sweeping Processes

In this section, we provide an existence and uniqueness result for the solutions of the history-dependent sweeping process:

$$\begin{cases} \dot{x}(t) \in -N_{C(t)}(x(t)) + f(t, x(t)) + \mathcal{R}(x)(t) & \text{for a.e. } t \in I, \\ x(0) = x_0 \in C(0), \end{cases}$$
(2)

where C, f and \mathcal{R} satisfy (\mathcal{H}^C) , (\mathcal{H}^f) and $(\mathcal{H}^{\mathcal{R}})$, respectively. Before providing the main result of this section, we provide a basic existence result for the sweeping process with integrable perturbation. We refer to [14, Proposition 1] for the proof.

Proposition 4.1 Let \mathcal{H} be a real Hilbert space, and suppose that $C(\cdot)$ satisfies (\mathcal{H}^C) . Let $h: I \to \mathcal{H}$ be a single-valued integrable mapping. Then, for any $x_0 \in C(0)$ there exists a unique absolutely continuous solution $x(\cdot)$ for the following differential inclusion:

$$\begin{cases} \dot{x}(t) \in -N_{C(t)}(x(t)) + h(t) & a.e. \ t \in I, \\ x(0) = x_0. \end{cases}$$

Moreover, $x(\cdot)$ satisfies the following inequality:

$$\|\dot{x}(t) - h(t)\| \le \|h(t)\| + |\dot{v}(t)|$$
 a.e. $t \in I$.

Now, we present the main result of this section. Based on a fixed-point result for history-dependent operators (see Proposition 2.3), we prove the wellposedness for the problem (2).

Theorem 4.1 Assume that (\mathcal{H}^C) , (\mathcal{H}^f) and $(\mathcal{H}^{\mathcal{R}})$ hold. Then, for any initial condition $x_0 \in C(0)$ there exists a unique absolutely continuous solution $x(\cdot)$ for the problem (2). Moreover, the following bound holds:

$$||x(t)|| \leq r(t)$$
 for all $t \in I$ and $||\dot{x}(t)|| \leq q(t)$ for a.e. $t \in I$,

where, $r(\cdot)$ and $q(\cdot)$ are the functions defined by

$$r(t) := \|x_0\| \exp\left(2\int_0^t (\alpha(s) + \kappa_{\mathcal{R}})ds\right) + \int_0^t \varepsilon(s) \exp\left(2\int_s^t (\alpha(\tau) + \kappa_{\mathcal{R}})d\tau\right)ds,$$

$$q(t) := |\dot{v}(t)| + 2\alpha(t)r(t) + 2\beta(t) + 2\kappa_{\mathcal{R}}\int_0^t r(s)ds + 2\|\mathcal{R}(0)(t)\|,$$

(3)

and $\varepsilon(t) := |\dot{v}(t)| + 2 \|\mathcal{R}(0)(t)\| + 2\beta(t).$

Proof Given $y \in C(I; \mathcal{H})$, let us consider the function

$$h_y(t) := f(t, \tilde{y}(t)) + \mathcal{R}(\tilde{y})(t),$$

where $\tilde{y}(t) := \operatorname{proj}_{r(t)\mathbb{B}}(y(t))$. By virtue of (\mathcal{H}^f) and $(\mathcal{H}^{\mathcal{R}})$, for $t \in I$, one has

$$\begin{aligned} \|h_y(t)\| &\leq \alpha(t)\|\tilde{y}(t)\| + \beta(t) + \|\mathcal{R}(\tilde{y})(t) - \mathcal{R}(0)(t)\| + \|\mathcal{R}(0)(t)\| \\ &\leq \alpha(t)r(t) + \beta(t) + \kappa_{\mathcal{R}} \int_0^t r(s)ds + \|\mathcal{R}(0)(t)\|, \end{aligned}$$

which shows that the map $t \mapsto h_y(t)$ is integrable. Hence, according to Proposition 4.1, the differential inclusion:

$$\begin{cases} \dot{x}(t) \in -N_{C(t)}(x(t)) + h_y(t) & \text{ for a.e. } t \in I, \\ x(0) = x_0 \in C(0), \end{cases}$$
(4)

admits a unique solution $x(\cdot)$, which is absolutely continuous and

$$\|\dot{x}(t) - h_y(t)\| \le |\dot{v}(t)| + \|h_y(t)\|$$
 a.e. $t \in I$,

where $r(\cdot)$ is defined in formula (3). Therefore, for a.e. $t \in I$, one has

$$\|\dot{x}(t) - h_y(t)\| \le \psi(t) := |\dot{v}(t)| + \alpha(t)r(t) + \beta(t) + \kappa_{\mathcal{R}} \int_0^t r(s)ds + \|\mathcal{R}(0)(t)\|.$$

Hence, the operator $\mathcal{F}: C(I; \mathcal{H}) \to C(I; \mathcal{H})$ defined as $\mathcal{F}(y) := x_y$ is a welldefined single-valued function. We proceed to prove that the operator \mathcal{F} is history-dependent. Indeed, let $y_1, y_2 \in C(I; \mathcal{H})$ and set $x_i = \mathcal{F}(y_i)$ for i = 1, 2. Hence, according to (4), for a.e. $t \in I$,

$$\frac{\dot{x}_i(t) - h_{y_i}(t)}{\psi(t)} \in -N_{C(t)}(x_i(t)) \cap \mathbb{B} \quad \text{for } i = 1, 2,$$

which, by the ρ -uniformly prox-regularity of the sets C(t) and assertion (*iii*) of Proposition 2.1, implies that for a.e. $t \in I$, one has

$$\langle \dot{x}_1(t) - \dot{x}_2(t) - [h_{y_1}(t) - h_{y_2}(t)], x_1(t) - x_2(t) \rangle \leq \frac{\psi(t)}{\rho} ||x_1(t) - x_2(t)||^2.$$

Therefore, for a.e. $t \in I$,

$$\frac{1}{2}\frac{d}{dt}\|x_1(t) - x_2(t)\|^2 \le \|h_{y_1}(t) - h_{y_2}(t)\| \cdot \|x_1(t) - x_2(t)\| + \frac{\psi(t)}{\rho}\|x_1(t) - x_2(t)\|^2.$$

Then, by integrating the above inequality and using Lemma 2.2, we obtain that for all $t \in I$, one has

$$\|x_1(t) - x_2(t)\| \le 2\int_0^t \exp\left(2\int_s^t \frac{\psi(\tau)}{\rho} d\tau\right) \|h_{y_1}(s) - h_{y_2}(s)\|ds.$$
(5)

Now, we proceed to estimate the term $||h_{y_1}(t) - h_{y_2}(t)||$. Indeed, by virtue of (\mathcal{H}^f) and $(\mathcal{H}^{\mathcal{R}})$, we obtain that for a.e. $t \in I$, one has

$$\begin{aligned} \|h_{y_1}(t) - h_{y_2}(t)\| &\leq \|f(t, \tilde{y}_1(t)) - f(t, \tilde{y}_1(t))\| + \|\mathcal{R}(\tilde{y}_1)(t) - \mathcal{R}(\tilde{y}_2)(t)\| \\ &\leq \kappa_f^{r(T)} \|\tilde{y}_1(t) - \tilde{y}_2(t)\| + \kappa_{\mathcal{R}} \int_0^t \|\tilde{y}_1(t) - \tilde{y}_2(t)\| ds \\ &\leq \kappa_f^{r(T)} \|y_1(t) - y_2(t)\| + \kappa_{\mathcal{R}} \int_0^t \|y_1(s) - y_2(s)\| ds, \end{aligned}$$

where we have used that the map $x \mapsto \operatorname{proj}_{r(T)\mathbb{B}}(x)$ is Lipschitz of constant 1. Hence, from the previous calculation and inequality (5), we deduce that for all $t \in I$, one has

$$\|x_{1}(t) - x_{2}(t)\| \leq 2\kappa_{f}^{r(T)} \int_{0}^{t} \exp\left(2\int_{s}^{t} \frac{\psi(\tau)}{\rho} d\tau\right) \|y_{1}(s) - y_{2}(s)\| ds + 2\kappa_{\mathcal{R}} \int_{0}^{t} \exp\left(2\int_{s}^{t} \frac{\psi(\tau)}{\rho} d\tau\right) \int_{0}^{s} \|y_{1}(\tau) - y_{2}(\tau)\| d\tau ds,$$

from which we deduce that \mathcal{F} is a state-dependent operator. Therefore, by virtue of Proposition 2.3, the history-dependent operator \mathcal{F} has a unique fixed point $x(\cdot)$, which solves the differential inclusion:

$$\begin{cases} \dot{x}(t) \in -N_{C(t)}(x(t)) + h_x(t) & \text{for a.e. } t \in I, \\ x(0) = x_0 \in C(0), \end{cases}$$
(6)

where $h_x(t) := f(t, \tilde{x}(t)) + \mathcal{R}(\tilde{x}(t))$ and $\tilde{x}(t) := \text{proj}_{r(t)\mathbb{B}}(x(t))$. To finish the proof, it remains to show that

$$||x(t)|| \le r(t)$$
 for all $t \in I$ and $||\dot{x}(t)|| \le q(t)$ for a.e. $t \in I$.

Indeed, by virtue of (\mathcal{H}^f) and $(\mathcal{H}^{\mathcal{R}})$, for all $t \in I$, one has

$$\begin{split} \|x(t)\| &\leq \|x_0\| + \int_0^t \|\dot{x}(s)\| ds \\ &\leq \|x_0\| + \int_0^t (|\dot{v}(s)| + \|h_x(s)\|) ds + \int_0^t \|h_x(s)\| ds \\ &\leq \|x_0\| + \int_0^t |\dot{v}(s)| ds + 2\int_0^t \|f(s,\tilde{x}(s))\| ds + 2\int_0^t \|\mathcal{R}(\tilde{x})(s)\| ds \\ &\leq \|x_0\| + \int_0^t |\dot{v}(s)| ds + 2\int_0^t \|\mathcal{R}(0)(s)\| ds + 2\int_0^t \|f(s,\tilde{x}(s))\| ds \\ &+ 2\int_0^t \|\mathcal{R}(\tilde{x})(s) - \mathcal{R}(0)(s)\| ds \\ &\leq \|x_0\| + \int_0^t |\dot{v}(s)| ds + 2\int_0^t \|\mathcal{R}(0)(s)\| ds + 2\int_0^t \beta(s) ds \\ &+ 2\int_0^t \alpha(s)\|\ddot{x}(s)\| ds + 2\kappa_{\mathcal{R}}\int_0^t \int_0^s \|\ddot{x}(\tau)\| d\tau ds \\ &\leq \|x_0\| + \int_0^t |\dot{v}(s)| ds + 2\int_0^t \|\mathcal{R}(0)(s)\| ds + 2\int_0^t \beta(s) ds \\ &+ 2\int_0^t \alpha(s)\|x(s)\| ds + 2\kappa_{\mathcal{R}}\int_0^t \int_0^s \|x(\tau)\| d\tau ds, \end{split}$$

where we have used that the map $x \mapsto \text{proj}_{r(t)\mathbb{B}}(x)$ is Lipschitz of constant 1. Therefore, for all $t \in I$, one has

$$\|x(t)\| \le \|x_0\| + \int_0^t \varepsilon(s)ds + 2\int_0^t \alpha(s)\|x(s)\|ds + 2\kappa_{\mathcal{R}} \int_0^t \int_0^s \|x(\tau)\|d\tau ds,$$

where $\varepsilon(t) := |\dot{v}(t)| + 2 \|\mathcal{R}(0)(t)\| + 2\beta(t)$. Hence, by virtue of Lemma 2.1, we obtain that, for all $t \in I$, one has

$$\|x(t)\| \le r(t) := \|x_0\| \exp\left(2\int_0^t (\alpha(s) + \kappa_{\mathcal{R}})ds\right) + \int_0^t \varepsilon(s) \exp\left(2\int_s^t (\alpha(\tau) + \kappa_{\mathcal{R}})d\tau\right)ds,$$

which implies that $x(\cdot)$ solves the history-dependent sweeping process (2). Finally, similarly to the above calculations, for a.e. $t \in I$, one has

$$\begin{aligned} \|\dot{x}(t)\| &\leq \|\dot{x}(t) - h_{x}(t)\| + \|h_{x}(t)\| \\ &\leq |\dot{v}(t)| + 2\|h_{x}(t)\| \\ &\leq |\dot{v}(t)| + 2\|\mathcal{R}(0)(s)\| + 2\alpha(t)\|x(t)\| + 2\beta(t) + 2\kappa_{\mathcal{R}} \int_{0}^{t} \|x(s)\| ds \\ &\leq q(t) := |\dot{v}(t)| + 2\|\mathcal{R}(0)(s)\| + 2\alpha(t)r(t) + 2\beta(t) + 2\kappa_{\mathcal{R}} \int_{0}^{t} r(s) ds \end{aligned}$$

which ends the proof.

5 State-Dependent Sweeping Process

In this section, we consider the following state-dependent history-dependent sweeping process:

$$\begin{cases} \dot{x}(t) \in -N_{C(t,x(t))}(x(t)) + f(t,x(t)) + \mathcal{R}(x)(t) & \text{ for a.e. } t \in [0,T], \\ x(0) = x_0 \in C(0,x_0), \end{cases}$$
(7)

where C, f and \mathcal{R} satisfy (\mathcal{H}_x^C) , (\mathcal{H}^f) and $(\mathcal{H}^{\mathcal{R}})$, respectively. By means of the Schauder Fixed Point Theorem, we prove the existence of solutions for the above differential inclusion.

Theorem 5.1 Assume that (\mathcal{H}_x^C) , (\mathcal{H}^f) and $(\mathcal{H}^{\mathcal{R}})$ hold. Then, for any $x_0 \in C(0, x_0)$, there exists at least one absolutely continuous solution $x(\cdot)$ for the problem (7). Moreover, the following bound holds:

$$\begin{aligned} \|x(t)\| &\leq r(t) + L \int_0^t \psi(s) \exp\left(2\int_s^t (\alpha(\tau) + \kappa_{\mathcal{R}}) d\tau\right) ds \quad \text{for all } t \in I, \\ \|\dot{x}(t)\| &\leq \psi(t) \quad \text{for a.e. } t \in I, \end{aligned}$$

where r(t) is defined in formula (3) and

$$\psi(t) := \frac{q(t)}{1-L} + \frac{2L}{1-L}(\alpha(t) + \kappa_{\mathcal{R}}) \int_0^t \exp\left(2\int_t^s (\alpha(\tau) + \kappa_{\mathcal{R}})d\tau\right) ds$$

Proof We say that $y \in \mathcal{K}$ if there exists $f \in L^1(I; \mathcal{H})$ such that $||f(t)|| \le \psi(t)$ for a.e. $t \in I$ and

$$y(t) := x_0 + \int_0^t f(s)ds \text{ for all } t \in I.$$

It is clear that the set \mathcal{K} seen as a subset of $C(I; \mathcal{H})$ is nonempty, closed and convex. We observe that for a given $y \in \mathcal{K}$ the set-valued map $t \mapsto C(t, y(t))$ satisfies assumption (\mathcal{H}^C) . Indeed, according to (\mathcal{H}^C_x) , for any $t, s \in I$ with $s \leq t$, one has

$$\begin{aligned} \operatorname{Haus}(C(t, y(t)), C(s, y(s))) &\leq |v(t) - v(s)| + L \|y(t) - y(s)\| \\ &\leq |v(t) - v(s)| + L \int_{s}^{t} \psi(\tau) d\tau. \end{aligned}$$

Therefore, by virtue of Theorem 4.1, for any $y \in \mathcal{K}$, there exists a unique absolutely continuous solution $x(\cdot)$ to the following differential inclusion:

$$\begin{cases} \dot{x}(t) \in -N_{C(t,y(t))}(x(t)) + f(t,x(t)) + \mathcal{R}(x)(t) & \text{ for a.e. } t \in I, \\ x(0) = x_0. \end{cases}$$
(8)

Moreover, the following bounds hold:

$$\begin{split} \|x(t)\| &\leq r_0(t) := r(t) + L \int_0^t \psi(s) \exp\left(2\int_s^t (\alpha(\tau) + \kappa_{\mathcal{R}})d\tau\right) ds & \text{for all } t \in I, \\ \|\dot{x}(t)\| &\leq q_0(t) := q(t) + L\psi(t) + 2\alpha(t)L \exp\left(2\int_0^t (\alpha(\tau) + \kappa_{\mathcal{R}})d\tau\right)\nu(t) \\ &+ 2\kappa_{\mathcal{R}}L \int_0^t \exp\left(2\int_0^t (\alpha(\tau) + \kappa_{\mathcal{R}})d\tau\right)\nu(s)ds & \text{a.e. } t \in I, \end{split}$$

where

$$\nu(t) := \int_0^t \psi(s) \exp\left(2\int_0^s (\alpha(\tau) + \kappa_{\mathcal{R}}) d\tau\right) ds.$$

Hence, we can consider the operator $\mathcal{F} \colon \mathcal{K} \to C(I; \mathcal{H})$ which assigns to each $y \in \mathcal{K}$ the unique solution $x(\cdot) := \mathcal{F}(y)$ of the problem (8). Moreover, for a.e. $t \in I$, one has

$$\begin{aligned} \|\dot{x}(t)\| &\leq q(t) + L\psi(t) + 2\alpha(t)L\exp\left(2\int_0^t (\alpha(\tau) + \kappa_{\mathcal{R}})d\tau\right)\nu(t) \\ &+ 2\kappa_{\mathcal{R}}L\int_0^t \exp\left(2\int_0^t (\alpha(\tau) + \kappa_{\mathcal{R}})d\tau\right)\nu(s)ds = \psi(t), \end{aligned}$$

which proves that the operator \mathcal{F} takes values in \mathcal{K} .

Claim 1: The operator $\mathcal{F} \colon \mathcal{K} \to \mathcal{K}$ is continuous.

Proof of Claim 1: Let us consider a sequence $(y_n) \subset \mathcal{K}$ converging to $y \in \mathcal{K}$. Set $x_n := \mathcal{F}(y_n)$ and $x := \mathcal{F}(y)$. By using (8), we obtain that

$$\begin{aligned} \dot{x}_n(t) &\in -N_{C(t,y_n(t))}(x_n(t)) + h_n(t) & \text{ a.e. } t \in I, \\ \dot{x}(t) &\in -N_{C(t,y(t))}(x(t)) + h(t) & \text{ a.e. } t \in I, \end{aligned}$$

where $h_n(t) := f(t, x_n(t)) + \mathcal{R}(x_n)(t)$ and $h(t) := f(t, x(t)) + \mathcal{R}(x)(t)$. Hence, by virtue of Proposition 4.1 and assumptions (\mathcal{H}^f) and $(\mathcal{H}^{\mathcal{R}})$, we obtain that, for a.e. $t \in I$, one has

$$\begin{aligned} \|\dot{x}_{n}(t) - h_{n}(t)\| &\leq |\dot{v}(t)| + L\psi(t) + \|h_{n}(t)\| \\ &\leq m(t) := |\dot{v}(t)| + L\psi(t) + \alpha(t)r_{0}(t) + \beta(t) \\ &+ \kappa_{\mathcal{R}} \int_{0}^{t} r_{0}(s)ds + \|\mathcal{R}(0)(t)\|, \end{aligned}$$

which implies that for all $n \in \mathbb{N}$, one has

$$\frac{\dot{x}_n(t) - h_n(t)}{m(t)} \in -N_{C(t,y_n(t))}(x_n(t)) \cap \mathbb{B} \quad \text{a.e. } t \in I.$$

Similarly,

$$\frac{\dot{x}(t) - h(t)}{m(t)} \in -N_{C(t,y(t))}(x(t)) \cap \mathbb{B} \quad \text{a.e. } t \in I.$$

Hence, By virtue of Proposition 2.1, we get that for all $c \in C(t, y_n(t))$

$$\langle -\dot{x}_n(t) + h_n(t), c - x_n(t) \rangle \le \frac{m(t)}{2\rho} \|c - x_n(t)\|^2$$
 (9)

Similarly, for all $c \in C(t, y(t))$

$$\langle -\dot{x}(t) + h(t), c - x(t) \rangle \le \frac{m(t)}{2\rho} \|c - x(t)\|^2.$$
 (10)

Next, according to (\mathcal{H}_x^C) , for all $t \in I$, we have

$$C(t, y_n(t)) \subset C(t, y(t)) + L ||y(t) - y_n(t)|| \mathbb{B},$$

$$C(t, y(t)) \subset C(t, y_n(t)) + L ||y(t) - y_n(t)|| \mathbb{B},$$

which implies the existence of $d_n(t), \tilde{d}_n(t) \in \mathbb{B}$ such that

$$x(t) - L \|y_n(t) - y(t)\| d_n(t) \in C(t, y_n(t)),$$

$$x_n(t) - L \|y_n(t) - y(t)\| d_n(t) \in C(t, y(t)).$$

Therefore, by using the above quantities in (9) and (10), respectively, we obtain that for a.e. $t \in I$

$$\begin{aligned} \langle -\dot{x}_n(t) + h_n(t), x(t) - x_n(t) \rangle &\leq Lm(t) \|y_n(t) - y(t)\| + \frac{m(t)}{\rho} \|x(t) - x_n(t)\|^2 \\ &+ \frac{m(t)L^2}{\rho} \|y(t) - y_n(t)\|^2 \\ \langle -\dot{x}(t) + h(t), x_n(t) - x(t) \rangle &\leq Lm(t) \|y_n(t) - y(t)\| + \frac{m(t)}{\rho} \|x_n(t) - x(t)\|^2 \\ &+ \frac{m(t)L^2}{\rho} \|y(t) - y_n(t)\|^2. \end{aligned}$$

Therefore, for a.e. $t \in I$,

$$\frac{1}{2} \frac{d}{dt} \|x_n(t) - x(t)\|^2 = \langle \dot{x}_n(t) - \dot{x}(t), x_n(t) - x(t) \rangle$$

$$\leq 2Lm(t) \|y_n(t) - y(t)\| + 2\frac{m(t)L^2}{\rho} \|y_n(t) - y(t)\|^2$$

$$+ 2\frac{m(t)}{\rho} \|x_n(t) - x(t)\|^2 + \langle h_n(t) - h(t), x_n(t) - x(t) \rangle.$$

Moreover, by using (\mathcal{H}^f) and $(\mathcal{H}^{\mathcal{R}})$, we obtain that

$$\begin{aligned} \langle h_n(t) - h(t), x_n(t) - x(t) \rangle &\leq \|h_n(t) - h(t)\| \cdot \|x_n(t) - x(t)\| \\ &\leq k_f^R \|x_n(t) - x(t)\|^2 + \kappa_{\mathcal{R}} \|x_n(t) - x(t)\| \int_0^t \|x_n(s) - x(s)\| ds, \end{aligned}$$

where $R := \sup_{t \in I} r_0(t)$ and k_f^R is the constant given by (\mathcal{H}^f) . Therefore, for all $t \in I$, one has

$$||x_n(t) - x(t)||^2 \le 4L \int_0^t m(s) ||y_n(s) - y(s)|| ds + \frac{4L^2}{\rho} \int_0^t m(s) ||y_n(s) - y(s)||^2 ds + \int_0^t \left(4\frac{m(s)}{\rho} + 2\kappa_f^R\right) ||x_n(s) - x(s)||^2 ds + \kappa_R \left(\int_0^t ||x_n(s) - x(s)|| ds\right)^2$$

Set $\sigma(t) := \limsup \|x_n(t) - x(t)\|$. By taking limit in the above inequality, we obtain that for all $t \in I$

$$\sigma^{2}(t) \leq \int_{0}^{t} \left(4\frac{m(s)}{\rho} + 2\kappa_{f}^{R}\right) \sigma^{2}(s)ds + \kappa_{\mathcal{R}} \left(\int_{0}^{t} \sigma(s)ds\right)^{2} \\ \leq \int_{0}^{t} \left(4\frac{m(s)}{\rho} + 2\kappa_{f}^{R}\right) \sigma^{2}(s)ds + \kappa_{\mathcal{R}}t \int_{0}^{t} \sigma^{2}(s)ds,$$

where we have used Holder's inequality. Finally, from the classical Gronwall's inequality, we conclude that $\sigma \equiv 0$, which proves the continuity of \mathcal{F} . \Box To apply Schauder's Fixed Point Theorem, it remains to show that $\mathcal{F}(\mathcal{K})$ is relatively compact in $C(I; \mathcal{H})$.

Claim 2: The set $\mathcal{F}(\mathcal{K})$ relatively compact in $C(I; \mathcal{H})$.

Proof of Claim 2: On the one hand, since \mathcal{K} is bounded in $W^{1,1}(I; \mathcal{H})$ by definition. It is clear that $\mathcal{F}(\mathcal{K})$ is equicontinuous. On the other hand, fix $t \in I$ and define

$$\mathcal{K}(t) := \{ y(t) \colon y \in \mathcal{K} \} \text{ and } \mathcal{F}(\mathcal{K})(t) := \{ \mathcal{F}(y)(t) \colon y \in \mathcal{K} \}.$$

We can observe that $\mathcal{F}(\mathcal{K})(t) \subset C(t, \mathcal{K}(t)) \cap r_0(t)\mathbb{B}$. Hence, by using the notion measure of non-compactness, we obtain that

$$\gamma(\mathcal{F}(\mathcal{K})(t)) \leq \gamma\left(C(t,\mathcal{K}(t)) \cap r_0(t)\mathbb{B}\right) = 0,$$

where we have used assumption (\mathcal{H}_x^C) and the monotony of γ . Therefore, the set $\mathcal{F}(\mathcal{K})(t)$ is relatively compact in \mathcal{H} . Finally, by virtue of the Arzelà-Ascoli Theorem, we conclude that the set $\mathcal{F}(\mathcal{K})$ is relatively compact. \Box Finally, by Schauder's Fixed Point Theorem, the operator \mathcal{F} has a fixed point, which is a solution of (7).

6 Volterra Sweeping Processes

In this section, we show that our main results (Theorems 4.1 and 5.1) enables us to obtain the existence of solutions for the Volterra Sweeping Process:

$$\begin{cases} \dot{x}(t) \in -N_{C(t)}(x(t)) + f(t, x(t)) + \int_0^t g(t, s, x(s)) ds \text{ for a.e. } t \in I, \\ x(0) = x_0 \in C(0), \end{cases}$$
(11)

and the State-Dependent Volterra Sweeping Process:

$$\begin{cases} \dot{x}(t) \in -N_{C(t,x(t))}(x(t)) + f(t,x(t)) + \int_0^t g(t,s,x(s))ds \text{ for a.e. } t \in I, \\ x(0) = x_0 \in C(0,x_0). \end{cases}$$
(12)

Here we assume that f and g satisfy (\mathcal{H}^f) and (\mathcal{H}^g) , respectively. Hence, to address the above dynamical systems, we can consider the operator

$$\mathcal{R}(x)(t) := \int_0^t f(t, s, x(s)) ds$$

Unfortunately, this operator does not satisfy Assumption $(\mathcal{H}^{\mathcal{R}})$; hence, we cannot directly apply Theorems 4.1 and 5.1, respectively. Nevertheless, by using the reparametrization technique developed in [33], we can, without loss of generality, assume that \mathcal{R} satisfies Assumption $(\mathcal{H}^{\mathcal{R}})$. Hence, on the one hand, the existence result for the Volterra Sweeping Process is the following:

Theorem 6.1 Assume that (\mathcal{H}^C) , (\mathcal{H}^f) and (\mathcal{H}^g) hold. Then, for any initial condition $x_0 \in C(0)$ there exists a unique absolutely continuous solution $x(\cdot)$ for the problem (11).

On the other hand, the existence result for the State-Dependent Volterra Sweeping Process is the following:

Theorem 6.2 Assume that (\mathcal{H}_x^C) , (\mathcal{H}^f) and (\mathcal{H}^g) hold. Then, for any $x_0 \in C(0, x_0)$, there exists at least one absolutely continuous solution $x(\cdot)$ for the problem (12).

Remark 6.1 On the one hand, we note that Theorem 6.1 is not new; in fact, it was previously proven using different methods in [5,33]. Our contribution lies in demonstrating that it can be derived through a fixed-point argument. On the other hand, to the best of our knowledge, the result presented in 6.1 is novel.

7 An Application to Viscoelastic Models with Long Memory

In this section, we illustrate one of our theoretical results (Theorem 4.1) through the modeling of a contact mechanical problem with long memory.

Let $\Omega \subset \mathbb{R}^d$, with $d \in \{2, 3\}$, be an open, bounded and connected set with a Lipschitz boundary $\Gamma := \partial \Omega$. We assume that Γ can be decomposed into mutually disjoint measurable sets Γ_D , Γ_N and Γ_C , with Γ_D having a positive Hausdorff measure. We are interested in the situation where Ω describes a viscoelastic material with long memory, which is clamped on Γ_D , subjected to a normal traction on Γ_N , and experiences dry friction on Γ_C (see Fig. 1).

To describe the mathematical model, let us consider \mathbb{S}^d as the space of symmetric second-order tensors over \mathbb{R}^d . On \mathbb{R}^d and \mathbb{S}^d , we consider the inner



Fig. 1 A viscoelastic material with long memory subjected to a external body force f_0 (left), normal traction f_2 and dry friction f_3 (right).

products

$$\langle u, v \rangle_{\mathbb{R}^d} := \sum_{i=1}^d u_i v_i \text{ and } \langle \sigma, \tau \rangle_{\mathbb{S}^d} := \sum_{i,j=1}^d \sigma_{ij} \tau_{ij} \text{ for all } u, v \in \mathbb{R}^d, \sigma, \tau \in \mathbb{S}^d.$$

These inner products induce the norms $||u||_{\mathbb{R}^d} := \sqrt{\langle u, u \rangle_{\mathbb{R}^d}}$ and $||\sigma||_{\mathbb{S}^d} := \sqrt{\langle \sigma, \sigma \rangle_{\mathbb{S}^d}}$ on \mathbb{R}^d and \mathbb{S}^d , respectively. For simplicity, we will denote these norms by $|| \cdot ||$ whenever no ambiguity arises.

Given $u \in (H^1(\Omega))^d$ we denote by $\varepsilon(u)$ the strain operator defined by $\varepsilon(u) := (\varepsilon_{ij}(u))_{i,j=1,\dots,d} \in \mathbb{S}^d$, where

$$\varepsilon_{ij}(u) := \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right)$$

To model the situation depicted in Fig. 1, we consider the vector spaces:

$$V := \{ u \in (H^1(\Omega))^d : u = 0 \text{ on } \Gamma_D \} \text{ and } W := \{ \sigma \in \mathbb{S}^d : \sigma_{ij} = \sigma_{ji} \in L^2(\Omega) \},\$$

where the equality u = 0 on Γ_D is understood in the sense of traces. The above spaces are endowed with the inner products:

$$\langle u,v\rangle_V:=\int_{\varOmega}\varepsilon(u(x))\cdot\varepsilon(v(x))dx\quad \text{ and }\quad \langle\sigma,\tau\rangle_W:=\int_{\varOmega}\sigma(x)\cdot\tau(x)dx.$$

We denote by ν the unit outward vector to Γ . The normal and tangential components of the vector $u \in \mathbb{R}^d$ are $u_{\nu} := u \cdot \nu$ and $u_{\tau} := u - u_{\nu}\nu$, respectively. Similarly, $\sigma_{\nu} := (\sigma\nu) \cdot \nu$ and $\sigma_{\tau} := \sigma\nu - \sigma_{\nu}\nu$ are the normal and tangential components of the tensor $\sigma \in \mathbb{S}^d$, respectively.

Description of the mechanical model: The considered constitutive law has the structure of a general viscoelastic law with long memory, including a non-linear integral part (see [19]), given by

$$\sigma(t,x) := \mathcal{A}(x,\varepsilon(u(t,x))) + \int_0^t \mathcal{B}(t-s,x,\varepsilon(u(t,x))) \, ds,$$

for all $(t, x) \in I \times \Omega$, where $\sigma(t, x)$ denotes the stress tensor, u(t, x) is the displacement field and \mathcal{A} and \mathcal{B} denote the *elasticity* and the *relaxation* operators, respectively. The above constitutive law describes the stress response of a material that exhibits both immediate elastic behavior and time-dependent memory effects, which are present, for example, in polymers, biological tissues, and certain metals. It provides a framework for predicting phenomena such as creep and stress relaxation. For further details on applications and notation, we refer to [2, 19, 24, 28].

The problem can be formulated as follows.

Problem 7.1 (Contact problem) Find a displacement vector field $u: I \times \Omega \to \mathbb{R}^d$ such that

$$\operatorname{div}(\sigma(t,x)) + f_0(t,x) = 0 \qquad \text{for all } (t,x) \in I \times \Omega, \tag{13}$$

$$u(t,x) = 0$$
 for all $(t,x) \in I \times \Gamma_D$, (14)

$$\sigma(t, x)\nu(x) = f_2(t, x) \quad \text{for all } (t, x) \in I \times \Gamma_N, \tag{15}$$

$$u_{\nu}(t,x) = 0$$
 for all $(t,x) \in I \times \Gamma_C$. (16)

Moreover, dry friction is acting on Γ_C , i.e., for all $(t, x) \in I \times \Gamma_C$

$$\|\sigma_{\tau}(t,x)\| \le f_3(t,x) \text{ and } -\sigma_{\tau}(t,x) = f_3(t,x) \frac{\dot{u}_{\tau}(t,x)}{\|\dot{u}_{\tau}(t,x)\|} \text{ if } \dot{u}_{\tau}(t,x) \ne 0.$$
 (17)

Equation (13) in Problem 7.1 represents the equilibrium balance between the stress tensor and an external force field $f_0(t, x)$, while (14) indicates that the set Ω is clamped on Γ_D . Moreover, (15) states that a normal traction force $f_2(t, x)$ is applied on Γ_N . Equation (16) specifies that no normal displacements are allowed in the contact region Γ_C . Finally, (17) represents the dry friction, defined by a friction bound $f_3(t, x)$.

Assumptions on the data: In order to provide the well-posedness for the Problem 7.1, we consider the following hypotheses: $(2t^{\Gamma})$: Assumptions on f, f, and f:

 (\mathcal{H}^{Γ}) : Assumptions on f_0 , f_2 and f_3 :

(i) There exists an absolutely continuous function ϑ_0 such that

$$\|f_0(t,\cdot) - f_0(s,\cdot)\|_{(L^2(\Omega))^d} \le |\vartheta_0(t) - \vartheta_0(s)| \text{ for all } t, s \in I.$$

(ii) There exists an absolutely continuous function ϑ_2 such that

$$\|f_2(t,\cdot) - f_2(s,\cdot)\|_{(L^2(\Omega))^d} \le |\vartheta_2(t) - \vartheta_2(s)| \text{ for all } t, s \in I.$$

- (iii) For all $t \in I$, the map $x \to f_3(t, x)$ is measurable and $f_3(0, x) = 0$ for a.e. $x \in \Gamma_C$.
- (\mathcal{H}^{σ}) : Assumptions on \mathcal{A} and \mathcal{B} :

- (i) The elasticity operator $\mathcal{A} = (a_{ijkl})$ defined from $\Omega \times \mathbb{S}^d$ into \mathbb{S}^d , is a linear application respect his second variable, such that satisfies:
 - (a) The components are symmetric, i.e., $a_{ijkl} = a_{jikl} = a_{klij}$, and a_{ijkl} is essentially bounded, i.e., $a_{ijkl} \in L^{\infty}(\Omega \times \mathbb{S}^d)$
 - (b) There exists $m_{\mathcal{A}} > 0$ such that

$$\langle \mathcal{A}(x,\varepsilon),\varepsilon\rangle_{\mathbb{S}^d} \ge m_{\mathcal{A}} \|\varepsilon\|_{\mathbb{S}^d}^2$$
 for all $\varepsilon \in \mathbb{S}^d$.

- (ii) The relaxation operator $\mathcal{B} = (b_{ijkl})$ defined from $I \times \Omega \times \mathbb{S}^d$ into \mathbb{S}^d satisfies the following conditions:
 - (a) The components are symmetric, i.e., $b_{ijkl} = b_{jikl} = b_{klij}$.
 - (b) For all $\varepsilon \in \mathbb{S}^d$ the map $t \mapsto b_{ijkl}(t, \cdot, \varepsilon)$ belongs to $W^{1,2}(I; L^{\infty}(\Omega))$.
 - (c) There exist nonnegative constants $\kappa_{\mathcal{B}}$ and $\kappa_{\mathcal{B}'}$ such that
 - For all $(t, x) \in I \times \Omega$ and any $\varepsilon_1, \varepsilon_2 \in \mathbb{S}^d$ one has

$$\|\mathcal{B}(t,x,\varepsilon_1) - \mathcal{B}(t,x,\varepsilon_2)\|_{\mathbb{S}^d} \le \kappa_{\mathcal{B}} \|\varepsilon_1 - \varepsilon_2\|_{\mathbb{S}^d}.$$

- For a.e. $t \in I$, for all $x \in \Omega$ and any $\varepsilon_1, \varepsilon_2 \in \mathbb{S}^d$ one has

$$\|\mathcal{B}'(t,x,\varepsilon_1) - \mathcal{B}'(t,x,\varepsilon_2)\|_{\mathbb{S}^d} \le \kappa_{\mathcal{B}'} \|\varepsilon_1 - \varepsilon_2\|_{\mathbb{S}^d}$$

(d) For any $(t, x) \in I \times \Omega$ one has

$$\langle \mathcal{B}(t, x, \varepsilon_1) - \mathcal{B}(t, x, \varepsilon_2), \varepsilon_1 - \varepsilon_2 \rangle_{\mathbb{S}^d} \ge 0 \text{ for all } \varepsilon_1, \varepsilon_2 \in \mathbb{S}^d$$

Now, based on standard calculations, the weak formulation of 7.1 is given by:

Problem 7.2 (Variational inequality) Find a displacement field u(t, x) such that the following inequality holds: for all $v \in V$,

$$\begin{split} \int_{\Omega} \sigma(t,x) &: (\varepsilon(v(x)) - \varepsilon(\dot{u}(t,x)))dx + \int_{\Gamma_{C}} f_{3}(t,x)(\|v_{\tau}(x)\| - \|\dot{u}_{\tau}(t,x)\|)da(x) \\ &\geq \int_{\Omega} f_{0}(t,x)(v(x) - \dot{u}(t,x))dx + \int_{\Gamma_{N}} f_{2}(t,x)(v(x) - \dot{u}(t,x))da(x). \end{split}$$

To apply our theoretical result (Theorem 4.1), we introduce the following auxiliary functions.

Definition 7.1 (Auxiliary functions) Let $f_0, f_2, f_3, \mathcal{A}$ and \mathcal{B} satisfying the assumptions (\mathcal{H}^{Γ}) and (\mathcal{H}^{σ}) . We consider the following functions:

1. Let $j: V \to \mathbb{R}$ defined by

$$j(u) = \int_{\Gamma_C} f_3(t, x) \| u_\tau(x) \| da(x) \quad \text{ for all } t \in I$$

2. Let $f: I \to V$ defined by

$$\langle f(t), u \rangle_V := \int_{\Omega} f_0(t, x) \cdot u(x) dx + \int_{\Gamma_N} f_2(t, x) \cdot u(x) da(x) \quad \text{ for all } t \in I.$$

3. Let $A: V \to V$ and $B: C(I; V) \to C(I; V)$ be such that:

$$\langle A(u), v \rangle_{V} = \int_{\Omega} \mathcal{A}(x, \varepsilon(u(x))) \cdot \varepsilon(v(x)) dx$$

$$\langle B(u)(t), v \rangle_{V} = \int_{\Omega} \left(\int_{0}^{t} \mathcal{B}(t - s, x, \varepsilon(u(s, x))) ds \right) \cdot \varepsilon(v(x)) dx.$$

Furthermore, we set $P: V \to V$ be such that $A = P^*P$, where P^* denotes the adjoint operator of P and $Q := (P^*)^{-1}$.

4. Let $C: I \rightrightarrows V$ be a set-valued map defined by $C(t) := Q(f(t) - \partial j(0))$. 5. Let $R: C(I; V) \rightarrow C(I; V)$ be defined by

$$\langle R(u)(t), v \rangle_{V} = \int_{\Omega} \mathcal{B}(0, x, \varepsilon(u(t, x))) \cdot \varepsilon(v(x)) dx + \int_{\Omega} \left(\int_{0}^{t} \mathcal{B}'(t - s, x, \varepsilon(u(s, x))) ds \right) \cdot \varepsilon(v(x)) dx.$$

We now present the main result of this section, which establishes the existence and uniqueness of a weak solution to Problem 7.2. Furthermore, we demonstrate that this solution satisfies a related history-dependent sweeping process.

Theorem 7.1 Assume that (\mathcal{H}^{Γ}) and (\mathcal{H}^{σ}) hold. Then, the Problem (7.2) admits a unique absolutely continuous solution $u: I \to V$ to the differential inclusion:

$$\begin{cases} \dot{u}(t) \in -N_{C(t)}(u(t)) + QR(u)(t) & a.e. \ t \in I, \\ u(0) = Pu_0, \end{cases}$$

where $Q = (P^*)^{-1}$ and $A = P^*P$.

Proof According to Definition 7.1, the Problem 7.2 is equivalent to the following variational inequality: find $u: I \to V$ such that

$$j(\dot{u}(t)) \ge j(v) + \langle f(t) - A(u(t)) - B(u)(t), \dot{u}(t) - v \rangle_V \text{ for all } (t,v) \in I \times V.$$

Since the function $j: V \to \mathbb{R}$ is convex, the above inequality is equivalent to:

$$f(t) - A(u(t)) - B(u)(t) \in \partial j(\dot{u}(t)) \quad \text{for all } t \in I.$$

Moreover, we observe that j is positively homogeneous, which implies that

$$\partial j(\dot{u}(t)) = \partial \sigma_{\partial j(0)}(\dot{u}(t))$$
 for a.e. $t \in I$.

Therefore, for a.e. $t \in I$, one has

$$\begin{split} \dot{u}(t) &\in \partial \sigma^*_{\partial j(0)}(f(t) - A(u(t)) - B(u)(t)) \\ &= N_{\partial j(0)}(f(t) - A(u(t)) - B(u)(t)) \\ &= N_{\partial j(0) - f(t)}(-P^*P(u(t)) - B(u)(t)) \\ &= -N_{f(t) - \partial j(0)}(P^*P(u(t)) + B(u)(t)). \end{split}$$

Following the ideas of [17], we consider the operator $\mathcal{G} \colon V \to V$ defined by

$$\mathcal{G}(u)(t) := Pu(t) + QB(u)(t).$$

It is clear that \mathcal{G} is invertible with Lipschitz inverse (see, e.g., [34, Proposition 31.4]). Therefore, setting $w(t) := \mathcal{G}(u)(t)$, we obtain that which gives

$$\dot{w}(t) \in -N_{C(t)}(w(t)) + QR(\mathcal{G}^{-1}(w))(t)$$
 a.e. $t \in I$.

It can be noted that $QR\mathcal{G}^{-1}$, and C satisfy the hypotheses (\mathcal{H}^R) and (\mathcal{H}^C) , respectively. Therefore, by virtue of Theorem 4.1, there exists a unique solution $u \in AC(I; V)$ for the history-dependent sweeping process.

Acknowledgements The third author was supported by ANID Chile under grants Fondecyt Regular N° 1240120, Fondecyt Regular N° 1220886, Proyecto de Exploración 13220097, CMM BASAL funds for Center of Excellence FB210005, Project ECOS230027, MATH-AMSUD 23-MATH-17).

References

- S. Adly. A variational approach to nonsmooth dynamics. SpringerBriefs Math. Springer, Cham, 2017.
- S. Adly and M. Sofonea. Time-dependent inclusions and sweeping processes in contact mechanics. Z. Angew. Math. Phys., 70(2):Paper No. 39, 19 pages, 2019.
- 3. C. D. Aliprantis and K. C. Border. *Infinite dimensional analysis*. Springer, Berlin, third edition, 2006.
- A. Bouach, T. Haddad, and B. S. Mordukhovich. Optimal control of nonconvex integrodifferential sweeping processes. J. Differential Equations, 329:255–317, 2022.
- 5. A. Bouach, T. Haddad, and L. Thibault. Nonconvex integro-differential sweeping process with applications. SIAM J. Control Optim., 60(5):2971–2995, 2022.
- M. Bounkhel and L. Thibault. Nonconvex sweeping process and prox-regularity in Hilbert space. J. Nonlinear Convex Anal., 6(2):359–374, 2005.
- N. Chemetov and M. D. P. Monteiro Marques. Non-convex quasi-variational differential inclusions. Set-Valued Anal., 15(3):209-221, 2007.
- 8. R. Christensen. Theory of Viscoelasticity: An Introduction. Academic Press, 2012.
- F. H. Clarke, Yu. S. Ledyaev, R. J. Stern, and P. R. Wolenski. Nonsmooth analysis and control theory, volume 178 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1998.
- 10. G. Colombo and C. Kozaily. Existence and uniqueness of solutions for an integral perturbation of Moreau's sweeping process. J. Convex Anal., 27(1):229–238, 2020.
- G. Colombo and L. Thibault. Prox-regular sets and applications. In *Handbook of nonconvex analysis and applications*, pages 99–182. Int. Press, Somerville, MA, 2010.
 K. Drimling, Nucleur for the standard set of the Society of Society and Societ
- 12. K. Deimling. Nonlinear functional analysis. Springer-Verlag, Berlin, 1985.
- K. Deimling. Multivalued differential equations, volume 1 of De Gruyter Series in Nonlinear Analysis and Applications. Walter de Gruyter & Co., Berlin, 1992.
- J. F. Edmond and L. Thibault. Relaxation of an optimal control problem involving a perturbed sweeping process. *Math. Program.*, 104(2-3):347–373, 2005.
- 15. H. Federer. Curvature measures. Trans. Amer. Math. Soc., 93:418-491, 1959.
- A. Jourani and E. Vilches. Moreau-Yosida regularization of state-dependent sweeping processes with nonregular sets. J. Optim. Theory Appl., 173(1):91–116, 2017.
- A. Jourani and E. Vilches. A differential equation approach to implicit sweeping processes. J. Differential Equations, 266(8):5168–5184, 2019.
- 18. R.S. Lakes. Viscoelastic Materials. Cambridge University Press, 2009.
- S. Migórski, A. Ochal, and M. Sofonea. Nonlinear inclusions and hemivariational inequalities, volume 26 of Adv. Mech. Math. Springer, New York, 2013.

- 20. J.-J. Moreau. Rafle par un convexe variable. I. In Travaux du Séminaire d'Analyse Convexe, Vol. I, volume No. 118 of Secrétariat des Mathématiques, Publication, pages Exp. No. 15, 43. Univ. Sci. Tech. Languedoc, Montpellier, 1971.
- J.-J. Moreau. Rafle par un convexe variable. II. In Travaux du Séminaire d'Analyse Convexe, Vol. II, volume No. 122 of Secrétariat des Mathématiques, Publication, pages Exp. No. 3, 36. Univ. Sci. Tech. Languedoc, Montpellier, 1972.
- J.-J. Moreau. Evolution problem associated with a moving convex set in a hilbert space. J. Differential Equations, 26(3):347–374, 1977.
- F. Nacry and M. Sofonea. History-dependent operators and prox-regular sweeping processes. Fixed Point Theory Algorithms Sci. Eng., pages Paper No. 5, 23, 2022.
- F. Nacry and M. Sofonea. History-dependent sweeping processes in contact mechanics. J. Convex Anal., 29(1):77–100, 2022.
- D. Narváez and E. Vilches. Moreau-Yosida regularization of degenerate state-dependent sweeping processes. J. Optim. Theory Appl., 193(1-3):910–930, 2022.
- R. A. Poliquin, R. T. Rockafellar, and L. Thibault. Local differentiability of distance functions. *Trans. Amer. Math. Soc.*, 352(11):5231–5249, 2000.
- M. Sene and L. Thibault. Regularization of dynamical systems associated with proxregular moving sets. J. Nonlinear Convex Anal., 15(4):647–663, 2014.
- M. Sofonea and S. Migórski. Variational-hemivariational inequalities with applications. Monogr. Res. Notes Math. CRC Press, Boca Raton, FL, 2018.
- 29. L. Thibault. Sweeping process with regular and nonregular sets. J. Differential Equations, 193(1):1–26, 2003.
- L. Thibault. Regularization of nonconvex sweeping process in Hilbert space. Set-Valued Anal., 16(2-3):319–333, 2008.
- L. Thibault. Unilateral Variational Analysis in Banach spaces. Part II: Special Classes of Functions and Sets. World Scientific Publishing Co. Pte. Ltd., 2023.
- E. Vilches. Regularization of perturbed state-dependent sweeping processes with nonregular sets. J. Nonlinear Convex Anal., 19(4):633–651, 2018.
- E. Vilches. Well-posedness for integro-differential sweeping processes of Volterra type. J. Convex Anal., 31(4):1273–1288, 2024.
- 34. E. Zeidler. Nonlinear functional analysis and its applications. II/B. Springer-Verlag, New York, 1990. Nonlinear monotone operators, Translated from the German by the author and Leo F. Boron.