

VIOLATION OF PARITY AND FLAVOR SYMMETRIES IN A NAMBU–JONA-LASINIO MODEL

YUKIMI GOTO¹ AND TOHRU KOMA²

Abstract: We study a lattice Nambu–Jona-Lasinio model with certain continuous chiral and two-flavor symmetries. For the Hamiltonian of the model, we construct a ground state which breaks the parity and flavor symmetries. In our argument, the chiral symmetry plays a crucial role for proving the violation of the parity and flavor symmetries, although the model does not contain the so-called Wilson term.

CONTENTS

1. Introduction	1
2. Hamiltonian of the Nambu–Jona-Lasinio model	4
3. Continuous chiral symmetry of the Hamiltonian	7
4. Spontaneous breaking of both flavor and parity symmetries	8
5. Reflection positivity: Preliminary	10
6. Reflection with respect to the plane $x^{(1)} = 1/2$	12
6.1. Two unitary transformations	14
6.2. Kinetic Hamiltonian	15
6.3. Interaction Hamiltonian $\tilde{H}_{\text{int,R}}^{(\Lambda)}(h)$	19
6.4. Interaction Hamiltonian $\tilde{H}_{\text{int,I}}^{(\Lambda)}$	21
7. Reflection with respect to the plane $x^{(2)} = 1/2$	22
8. Gaussian domination and infrared bound	23
9. Long-range order	25
Appendix A. Chiral rotational symmetry	28
Appendix B. A lower bound for $\mathcal{E}^{(\Lambda)}$	29
Acknowledgements	30
References	30

1. INTRODUCTION

In the present paper, we deal with a lattice Nambu–Jona-Lasinio model with certain continuous chiral and two-flavor symmetries. A model of this type in the lattice

¹ Gakushuin University, Department of Mathematics, Mejiro, Toshima-ku, Tokyo 171-8588, Japan.
Email: yukimi.goto@gakushuin.ac.jp.

² Gakushuin University (retired), Department of Physics, Mejiro, Toshima-ku, Tokyo 171-8588, Japan.

regularization with Wilson fermion is expected to exhibit the violation of the chiral symmetry and flavor symmetry in the ground state [5]. In lattice quantum chromodynamics (QCD) literature, such a phase structure is referred to as Aoki phase, going back to the pioneering works by Aoki [1, 2]. For Wilson fermions, it was predicted in [1, 2] that, in the broken phase for the parity and flavor symmetries, the charged pions can be interpreted as Nambu–Goldstone (massless) bosons, despite the absence of chiral symmetry. This expectation was strengthened by computer simulations of QCD [4, 6].

For the present Nambu–Jona-Lasinio model, we construct a ground state which breaks the parity and flavor symmetries. More precisely, we introduce certain two observables to detect the violation of the parity and flavor symmetries, and for these observables, we prove the existence of a non-vanishing spontaneous magnetization in the ground state. Technically, the method of fermion reflection positivity [17] is crucial for the proof, as in our previous works [13, 14]. However, it should be noted that if we introduce a symmetry-breaking field, the existence of spontaneous magnetization is always equivalent to the non-vanishing of the expectation value of $\bar{\Psi}\Psi$ due to the symmetries of the present model. Thus, the usual procedure to show spontaneous symmetry breaking does not work. To overcome this difficulty, we use the trial state introduced by Horsch and von der Linden [16]. In particular, we show the simultaneous non-vanishing of spontaneous magnetizations in two directions using the two order parameters. This approach is one of the principal novel ingredients in our paper. Then, the above conjecture is proved for our model without any approximation. However, we have not been able to prove that there exists an infinite-volume ground state such that it is pure and breaks the parity and flavor symmetries. Namely, the ground state which we constructed may consist of two or more pure states, and each pure state may not break the two symmetries simultaneously. Unfortunately, we cannot deny the possibility, but we do not expect such a situation to occur, since two observables can be simultaneously diagonalized in finite volume systems by $[\gamma_0\tau_1, i\gamma_0\gamma_5\tau_2] = 0$ in the Hilbert space. For further details, see Remark 4.2.

In this section, we roughly describe the present model and our idea to prove the violation of the parity and flavor symmetries. The precise mathematical definitions will be given later. For this purpose, we start from our motivation. Namely, the numerical simulations of QCD showed the violation of the parity and flavor symmetries [6]. The Lagrangian of QCD is given by

$$\mathcal{L}_0 = \sum_{i=1}^{N_f} \bar{\Psi}_i \gamma_\mu (\partial^\mu + ig_0 A_\mu) \Psi_i + \frac{1}{4} F_{\mu\nu} F^{\mu\nu},$$

where Ψ_i are the quark fields with N_f flavors, A_μ is the gauge field, $F_{\mu\nu}$ is its field strength, and g_0 is the coupling constant in the standard notation. Clearly, this Lagrangian \mathcal{L}_0 has the symmetry of the $SU(N_f)$ rotation about the flavors. In addition,

it is invariant under the usual chiral transformation,

$$\Psi_j \rightarrow e^{i\theta_j \gamma_5} \Psi_j \quad \text{for } j = 1, 2, \dots, N_f, \quad (1.1)$$

with the N_f angles, $\theta_j \in [0, 2\pi)$, $j = 1, 2, \dots, N_f$.

Next, we go back to the original idea of Nambu [24] for getting the effective Lagrangian. In the following, we will treat only the case with two flavors, i.e., $N_f = 2$, because the generalization is straightforward. We also write

$$\Psi = \begin{pmatrix} \Psi_1 \\ \Psi_2 \end{pmatrix} = \begin{pmatrix} \Psi_u \\ \Psi_d \end{pmatrix}.$$

By formally integrating out the gauge fields A_μ , the resulting Lagrangian \mathcal{L} can be written in terms of only the quark fields Ψ_i , and the Lagrangian becomes the effective one for the self-interacting fermion system. In the following, we will not treat the color degrees of freedom. As an approximation for the dominant contributions in the interaction terms, the four-fermion interactions have been widely used so far. The explicit form of the effective Lagrangian is given by

$$\mathcal{L} = \bar{\Psi} \gamma_\mu \partial^\mu \Psi + \tilde{g} \left\{ (\bar{\Psi} \Psi)^2 + (\bar{\Psi} i \gamma_5 \Psi)^2 + \sum_{j=1}^3 [(\bar{\Psi} \tau_j \Psi)^2 + (\bar{\Psi} i \gamma_5 \tau_j \Psi)^2] \right\} \quad (1.2)$$

with the effective coupling constant \tilde{g} , where τ_j are the Pauli matrices which act on only the flavor indices. From a physical viewpoint, we assume that the effective interactions between two quarks, which are mediated by the gauge fields A_μ , are attractive. Therefore, we choose $\tilde{g} > 0$ for the effective coupling constant. The above \mathcal{L} is nothing but the Lagrangian of a generalized Nambu–Jona-Lasinio model [15, 25, 26]. Clearly, it is invariant under the SU(2) flavor rotation. Further, this Lagrangian \mathcal{L} is also invariant under the above chiral transformation (1.1). The latter invariance will be checked in Sec. 3 below for the corresponding Hamiltonian on a lattice. Since the chiral symmetry plays a crucial role for proving the violation of the parity and flavor symmetries, we cannot include the so-called Wilson term into the Hamiltonian. Therefore, the well-known doubling problem arises in our model. In the original works of Aoki phase [1–4], it was also argued that the expectation value of the flavor singlet pseudoscalar operator, $i\bar{\Psi} \gamma_5 \Psi$, must vanish as a result of the Vafa–Witten theorem [31]. This means that the η meson remains massive in the broken phase for the parity and flavor symmetries, thereby solving the U(1) problem on the lattice. However, since our model presumably lacks a desirable continuum limit that reproduces the chiral anomaly, due to the presence of doublers [9, 23, 27, 28], it may have a different physical interpretation compared to the standard picture of Aoki phase.

Since the interactions of the Lagrangian \mathcal{L} are attractive, the corresponding interaction Hamiltonian, whose explicit form is given by (2.6) below on a lattice, looks very similar to the ferromagnetic Heisenberg spin Hamiltonian, although their internal degrees of freedom are totally different from each other. As is well known, in the ground

states of the Heisenberg ferromagnet, the spins align in the same direction, and the rotational symmetry of the spins is spontaneously broken.

In the model given by the above Lagrangian \mathcal{L} of (1.2), the observables which correspond to the spin components of the Heisenberg model, contain two observables [6], $\bar{\Psi}\tau_1\Psi$ and $\bar{\Psi}i\gamma_5\tau_2\Psi$. In the present paper, we construct a ground state such that the ground-state expectation values for these two observables are simultaneously non-vanishing. This result implies the violation of the parity and flavor symmetries. In our proof, we need to treat the above two observables because of the symmetries of the present system.

The present paper is organized as follows: We give the precise definition of the Hamiltonian of the Nambu–Jona-Lasinio model in Sec. 2, and the chiral symmetry of the Hamiltonian is checked in Sec. 3. Our main result is given in Sec. 4. The reflection positivity of the model is proved in the three Sections 5–7. The Gaussian domination and the infrared bound are obtained in Sec. 8. The existence of the long-range order is proved in Sec. 9. Appendices A and B are respectively devoted to technical relation and estimate.

2. HAMILTONIAN OF THE NAMBU–JONA-LASINIO MODEL

In order to derive the Hamiltonian of the Nambu–Jona-Lasinio model (1.2) on a lattice, consider first the case of the single flavor. The corresponding Lagrangian of the Nambu–Jona-Lasinio model is given by [15, 25, 26]

$$\mathcal{L} = \bar{\Psi}\gamma_\mu\partial^\mu\Psi + \tilde{g}[(\bar{\Psi}\Psi)^2 - (\bar{\Psi}\gamma_5\Psi)^2]$$

with the effective coupling constant $\tilde{g} \geq 0$. The four-fermion interaction for the Dirac fermion fields Ψ is formally written

$$\begin{aligned} & (\bar{\Psi}\Psi)^2 - (\bar{\Psi}\gamma_5\Psi)^2 \\ &= \frac{1}{2}(\bar{\Psi}\Psi + \bar{\Psi}\gamma_5\Psi)(\bar{\Psi}\Psi - \bar{\Psi}\gamma_5\Psi) + \frac{1}{2}(\bar{\Psi}\Psi - \bar{\Psi}\gamma_5\Psi)(\bar{\Psi}\Psi + \bar{\Psi}\gamma_5\Psi) \\ &= 2\left[\bar{\Psi}\frac{1+\gamma_5}{2}\Psi \cdot \bar{\Psi}\frac{1-\gamma_5}{2}\Psi + \bar{\Psi}\frac{1-\gamma_5}{2}\Psi \cdot \bar{\Psi}\frac{1+\gamma_5}{2}\Psi\right] \\ &= 2\left[\Psi^\dagger\gamma_0\frac{1+\gamma_5}{2}\Psi \cdot \Psi^\dagger\gamma_0\frac{1-\gamma_5}{2}\Psi + \Psi^\dagger\gamma_0\frac{1-\gamma_5}{2}\Psi \cdot \Psi^\dagger\gamma_0\frac{1+\gamma_5}{2}\Psi\right]. \end{aligned}$$

We want to define the corresponding interaction on the three-dimensional cubic lattice $\Lambda \subset \mathbb{Z}^3$ in the Hamiltonian formalism. We simply introduce

$$\Gamma^{(+)}(x) := \Psi^\dagger(x)\gamma_0 \cdot \frac{1}{2}(1 + \gamma_5)\Psi(x)$$

and

$$\Gamma^{(-)}(x) := \Psi^\dagger(x)\gamma_0 \cdot \frac{1}{2}(1 - \gamma_5)\Psi(x),$$

where $\Psi(x)$ is the four-component Dirac-field operator at the site $x \in \Lambda$. Then, from the above observations, one can consider the nearest-neighbor interaction given by

$$\Gamma^{(+)}(x)\Gamma^{(-)}(y) + \Gamma^{(-)}(x)\Gamma^{(+)}(y)$$

for the nearest-neighbor sites, $x, y \in \Lambda$. We also introduce

$$\Gamma^{(1)}(x) := \Gamma^{(+)}(x) + \Gamma^{(-)}(x)$$

and

$$\Gamma^{(2)}(x) := i[\Gamma^{(+)}(x) - \Gamma^{(-)}(x)].$$

Clearly, one has

$$\Gamma^{(1)}(x) = \Psi^\dagger(x)\gamma_0\Psi(x) \quad \text{and} \quad \Gamma^{(2)}(x) = \Psi^\dagger(x)i\gamma_0\gamma_5\Psi(x). \quad (2.1)$$

In addition, one notices

$$\Gamma^{(1)}(x)\Gamma^{(1)}(y) + \Gamma^{(2)}(x)\Gamma^{(2)}(y) = 2[\Gamma^{(+)}(x)\Gamma^{(-)}(y) + \Gamma^{(-)}(x)\Gamma^{(+)}(y)].$$

From these observations, one can consider the interaction Hamiltonian,

$$H_{\text{int}} = -g \sum_{x,y:|x-y|=1} [\Gamma^{(1)}(x)\Gamma^{(1)}(y) + \Gamma^{(2)}(x)\Gamma^{(2)}(y)],$$

with the coupling constant $g > 0$, i.e., we assume that the interaction is attractive, following the idea of [25]. Here, the sum is over all the nearest-neighbor pairs, $x, y \in \Lambda$. The precise definition of the finite lattice which we consider is given by

$$\Lambda := \{x = (x^{(1)}, x^{(2)}, x^{(3)}) \in \mathbb{Z}^3: -L + 1 \leq x^{(i)} \leq L, i = 1, 2, 3\}$$

with a positive integer L , and the periodic boundary condition. The lattice Λ can be considered as the three-dimensional torus. We write e_μ for the unit vector whose μ -th component is 1.

Next, consider the two flavor Dirac fermions. The four-component Dirac field $\Psi_f(x)$ with the flavor $f = u, d$ at the site $x \in \Lambda$ is given by

$$\Psi_f(x) := \begin{pmatrix} \psi_f^{(1)}(x) \\ \psi_f^{(2)}(x) \\ \psi_f^{(3)}(x) \\ \psi_f^{(4)}(x) \end{pmatrix}, \quad \Psi_f^\dagger(x) := \left(\psi_f^{(1)}(x)^\dagger, \psi_f^{(2)}(x)^\dagger, \psi_f^{(3)}(x)^\dagger, \psi_f^{(4)}(x)^\dagger \right)$$

where $\psi_f^{(i)}(x)$ is the fermion operator for the component $i \in \{1, 2, 3, 4\}$ at the site $x \in \Lambda$ with the flavor $f \in \{u, d\}$. These satisfy the anti-commutation relations,

$$\{\psi_f^{(i)}(x), [\psi_{f'}^{(j)}(y)]^\dagger\} = \delta_{x,y}\delta_{i,j}\delta_{f,f'}, \quad \{\psi_f^{(i)}(x), \psi_{f'}^{(j)}(y)\} = 0,$$

for the sites $x, y \in \Lambda$, $i, j \in \{1, 2, 3, 4\}$ and $f, f' \in \{u, d\}$. We write

$$\Psi(x) := \begin{pmatrix} \Psi_u(x) \\ \Psi_d(x) \end{pmatrix}, \quad \Psi^\dagger(x) := \left(\Psi_u^\dagger(x), \Psi_d^\dagger(x) \right)$$

for short. In order to simplify the notation, we will use $\mathcal{M}\Psi(x)$ to mean

$$\mathcal{M}\Psi(x) = \begin{pmatrix} \mathcal{M}\Psi_u(x) \\ \mathcal{M}\Psi_d(x) \end{pmatrix},$$

for any 4×4 matrix \mathcal{M} . Our Hamiltonian of the Nambu–Jona-Lasinio model [15] is given by

$$H^{(\Lambda)} := H_K^{(\Lambda)} + H_{\text{int}}^{(\Lambda)}, \quad (2.2)$$

where the first term is the usual kinetic Hamiltonian given by

$$H_K^{(\Lambda)} := i\kappa \sum_{x \in \Lambda \subset \mathbb{Z}^3} \left\{ [\Psi^\dagger(x)\alpha_1\Psi(x+e_1) - \Psi^\dagger(x+e_1)\alpha_1\Psi(x)] \right. \quad (2.3)$$

$$\left. + [\Psi^\dagger(x)\alpha_2\Psi(x+e_2) - \Psi^\dagger(x+e_2)\alpha_2\Psi(x)] \right. \quad (2.4)$$

$$\left. + [\Psi^\dagger(x)\alpha_3\Psi(x+e_3) - \Psi^\dagger(x+e_3)\alpha_3\Psi(x)] \right\}, \quad (2.5)$$

with the hopping parameter $\kappa \in \mathbb{R}$, where

$$\alpha_i = \begin{pmatrix} 0 & \sigma_i \\ \sigma_i & 0 \end{pmatrix} \quad \text{for } i = 1, 2, 3$$

with the Pauli matrices,

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \text{and} \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix};$$

The second term is the interaction Hamiltonian given by

$$\begin{aligned} H_{\text{int}}^{(\Lambda)} := & -g \sum_{x \in \Lambda} \sum_{\mu=1}^3 [\Gamma^{(1)}(x)\Gamma^{(1)}(x+e_\mu) + \Gamma^{(2)}(x)\Gamma^{(2)}(x+e_\mu)] \\ & - g \sum_{x \in \Lambda} \sum_{\mu=1}^3 \sum_{j=1}^3 [S^{(j)}(x)S^{(j)}(x+e_\mu) + S_5^{(j)}(x)S_5^{(j)}(x+e_\mu)] \end{aligned} \quad (2.6)$$

with the coupling constant $g > 0$, where

$$\Gamma^{(1)}(x) := \Psi^\dagger(x)\gamma_0\Psi(x), \quad \Gamma^{(2)}(x) := \Psi^\dagger(x)i\gamma_0\gamma_5\Psi(x),$$

$$S^{(j)}(x) := \Psi^\dagger(x)\gamma_0\tau_j\Psi(x) \quad \text{and} \quad S_5^{(\ell)}(x) := \Psi^\dagger(x)i\gamma_0\gamma_5\tau_\ell\Psi(x)$$

for $j, \ell = 1, 2, 3$, where τ_j are the Pauli matrices which act on only the flavor indices.

The explicit forms are given by

$$\tau_1 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \tau_2 := \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \text{and} \quad \tau_3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

In the following, we will use the expressions,

$$\gamma_0 = \begin{pmatrix} 1_2 & 0 \\ 0 & -1_2 \end{pmatrix} \quad \text{and} \quad \gamma_5 = \begin{pmatrix} 0 & 1_2 \\ 1_2 & 0 \end{pmatrix}, \quad (2.7)$$

with the 2-by-2 unit matrix 1_2 . We also impose the anti-periodic boundary condition [13] for the boundary of the kinetic Hamiltonian (2.3), in order to realize the fermion reflection positivity [17].

Originally, reflection positivity was introduced in quantum field theory by Osterwalder and Schrader [30], and it is closely related to the existence of a positive self-adjoint transfer matrix [29]. Subsequently, this concept was applied to the study of phase transitions in statistical mechanical models [10–12]. In the present paper, reflection positivity is the essential tool to establish the existence of long-range order.

3. CONTINUOUS CHIRAL SYMMETRY OF THE HAMILTONIAN

As is well-known, the form (2.6) of the interaction Hamiltonian $H_{\text{int}}^{(\Lambda)}$ is invariant under the usual chiral transformation,

$$\Psi_{\text{u}}(x) \rightarrow e^{i\theta\gamma_5}\Psi_{\text{u}}(x), \quad \Psi_{\text{d}}(x) \rightarrow e^{i\theta\gamma_5}\Psi_{\text{d}}(x), \quad (3.1)$$

with a real θ for $x \in \Lambda$. As we will see below, the present Hamiltonian $H^{(\Lambda)}$ of (2.2) is also invariant under another chiral transformation,

$$\Psi_{\text{u}}(x) \rightarrow e^{i\theta\gamma_5}\Psi_{\text{u}}(x), \quad \Psi_{\text{d}}(x) \rightarrow e^{-i\theta\gamma_5}\Psi_{\text{d}}(x), \quad (3.2)$$

where the two flavors are rotated by the opposite angles. Since $\gamma_0 e^{i\theta\gamma_5} \gamma_0 = e^{-i\theta\gamma_5}$, one notices that the quantity $\Psi^\dagger(x)\gamma_0\tau_1\Psi(y)$ is invariant under this transformation. More generally, the operators, $S^{(j)}(x)$ and $S_5^{(j)}(x)$ for $j = 1, 2$, are all invariant under this transformation.

The rest of the terms are given by the operators,

$$\Gamma^{(1)}(x)\Gamma^{(1)}(x + e_\mu), \Gamma^{(2)}(x)\Gamma^{(2)}(x + e_\mu), S^{(3)}(x)S^{(3)}(x + e_\mu) \text{ and } S_5^{(3)}(x)S_5^{(3)}(x + e_\mu).$$

Note that

$$\Gamma^{(1)}(x) = \Psi_{\text{u}}^\dagger(x)\gamma_0\Psi_{\text{u}}(x) + \Psi_{\text{d}}^\dagger(x)\gamma_0\Psi_{\text{d}}(x)$$

and

$$S_5^{(3)}(x) = \Psi_{\text{u}}^\dagger(x)i\gamma_0\gamma_5\Psi_{\text{u}}(x) - \Psi_{\text{d}}^\dagger(x)i\gamma_0\gamma_5\Psi_{\text{d}}(x).$$

Under the present chiral transformation (3.2), one has

$$\Gamma^{(1)}(x) \rightarrow \Gamma^{(1)}(x) \cos 2\theta + S_5^{(3)}(x) \sin 2\theta,$$

$$S_5^{(3)}(x) \rightarrow -\Gamma^{(1)}(x) \sin 2\theta + S_5^{(3)}(x) \cos 2\theta.$$

Therefore, the following quantity is invariant under the chiral transformation (3.2):

$$\Gamma^{(1)}(x)\Gamma^{(1)}(y) + S_5^{(3)}(x)S_5^{(3)}(y)$$

for $x, y \in \Lambda$. Similarly, the following quantity is also invariant:

$$\Gamma^{(2)}(x)\Gamma^{(2)}(y) + S^{(3)}(x)S^{(3)}(y).$$

These observations imply that the interaction Hamiltonian $H_{\text{int}}^{(\Lambda)}$ of (2.6) is invariant under the chiral transformation (3.2). Since the kinetic Hamiltonian $H_{\text{K}}^{(\Lambda)}$ is invariant,

the present Hamiltonian $H^{(\Lambda)}$ of (2.2) is invariant under the chiral transformation (3.2).

4. SPONTANEOUS BREAKING OF BOTH FLAVOR AND PARITY SYMMETRIES

The aim of the present paper is to prove the spontaneous breaking of both flavor and parity symmetries.

We write

$$\omega_0^{(\Lambda)}(\dots) := \lim_{\beta \nearrow \infty} \frac{\text{Tr}(\dots) e^{-\beta H^{(\Lambda)}}}{\text{Tr} e^{-\beta H^{(\Lambda)}}}$$

for the expectation value with respect to the ground state without the symmetry breaking field. In order to consider a phase transition, we introduce two order parameters

$$O_\Lambda^{(1)} = \frac{1}{|\Lambda|} \sum_{x \in \Lambda} S^{(1)}(x) = \frac{1}{|\Lambda|} \sum_{x \in \Lambda} \Psi^\dagger(x) \gamma_0 \tau_1 \Psi(x)$$

and

$$O_{5,\Lambda}^{(2)} = \frac{1}{|\Lambda|} \sum_{x \in \Lambda} S_5^{(2)}(x) = \frac{1}{|\Lambda|} \sum_{x \in \Lambda} \Psi^\dagger(x) i \gamma_0 \gamma_5 \tau_2 \Psi(x).$$

By a standard argument (see also Appendix A), it is easy to see that for any $\theta \in \mathbb{R}$,

$$e^{i\theta \gamma_5} \gamma_0 e^{-i\theta \gamma_5} = \gamma_0 \cos(2\theta) - i \gamma_0 \gamma_5 \sin(2\theta).$$

This relation implies that under the chiral transformation (3.1), $S^{(1)}(x)$ is transformed into $\Psi^\dagger(x) i \gamma_0 \gamma_5 \tau_1 \Psi(x)$. Then, the chiral and flavor symmetries of the Hamiltonian $H^{(\Lambda)}$ lead to the following equalities

$$\omega_0^{(\Lambda)}(O_\Lambda^{(1)}) = \omega_0^{(\Lambda)}(O_{5,\Lambda}^{(2)}) = 0,$$

and

$$\omega_0^{(\Lambda)}([O_\Lambda^{(1)}]^2) = \omega_0^{(\Lambda)}([O_{5,\Lambda}^{(2)}]^2).$$

As we will prove below, there exists a long-range order in the sense that

$$\omega_0^{(\Lambda)}([O_\Lambda^{(1)}]^2) = \omega_0^{(\Lambda)}([O_{5,\Lambda}^{(2)}]^2) > c_0, \quad (4.1)$$

where c_0 is a strictly positive constant and independent of the volume $|\Lambda|$. We consider a state [16, 18, 21, 22],

$$\omega^{(\Lambda)}(\dots) := \mathcal{N} \omega_0^{(\Lambda)}((1 + O_\Lambda)(\dots)(1 + O_\Lambda)),$$

where \mathcal{N} is the normalization constant, and we have written

$$O_\Lambda := O_\Lambda^{(1)} + O_{5,\Lambda}^{(2)}.$$

This is also a ground-state expectation. To see this, we compute the expectation value of $H^{(\Lambda)} - E_0^{(\Lambda)}$, where $E_0^{(\Lambda)}$ is the energy eigenvalue of the ground state. We have

$$\begin{aligned} \omega^{(\Lambda)}([H^{(\Lambda)} - E_0^{(\Lambda)}]) &= \mathcal{N}\omega_0^{(\Lambda)}((1 + O_\Lambda)[H^{(\Lambda)} - E_0^{(\Lambda)}](1 + O_\Lambda)) \\ &= \mathcal{N}\omega_0^{(\Lambda)}(O_\Lambda[H^{(\Lambda)} - E_0^{(\Lambda)}]O_\Lambda) \\ &= \frac{\mathcal{N}}{2}\{\omega_0^{(\Lambda)}(O_\Lambda[H^{(\Lambda)}, O_\Lambda]) - \omega_0^{(\Lambda)}([H^{(\Lambda)}, O_\Lambda]O_\Lambda)\} \\ &= \frac{\mathcal{N}}{2}\omega_0^{(\Lambda)}([O_\Lambda, [H^{(\Lambda)}, O_\Lambda]]) \leq \frac{c_1}{|\Lambda|}, \end{aligned}$$

where c_1 is a positive constant. Thus, the energy is degenerate into the energy of the ground state in the infinite-volume limit $|\Lambda| \nearrow \infty$.

By using the flavor rotational symmetry of the Hamiltonian $H^{(\Lambda)}$ of (2.2) and the existence of the long-range order (4.1), we obtain

$$\omega^{(\Lambda)}(O_\Lambda^{(1)}) \geq 2\mathcal{N}c_0 \quad \text{and} \quad \omega^{(\Lambda)}(O_{5,\Lambda}^{(2)}) \geq 2\mathcal{N}c_0.$$

Thus, the two order parameters show the non-vanishing spontaneous magnetizations in the infinite-volume ground state $\omega(\cdots) := \text{weak}^* - \lim_{\Lambda \nearrow \mathbb{Z}^3} \omega^{(\Lambda)}(\cdots)$.

These observations imply that the chiral, flavor and parity symmetries are all spontaneously broken in the infinite-volume ground state $\omega(\cdots)$.

Our main result is summarized as follows:

Theorem 4.1. *There exists a positive number α_0 small enough such that*

$$\lim_{\Lambda \nearrow \mathbb{Z}^\nu} \omega^{(\Lambda)}(O_\Lambda^{(1)}) > 0 \quad \text{and} \quad \lim_{\Lambda \nearrow \mathbb{Z}^\nu} \omega^{(\Lambda)}(O_{5,\Lambda}^{(2)}) > 0$$

for all the hopping parameter κ and the coupling constant g of the interaction which satisfy $|\kappa|/g \leq \alpha_0$.

Remark 4.2. In passing, we should remark the following:

- (i) In order to prove the violation of the parity, we need the two observables because of the symmetries of the present system. Actually, as shown in Appendix A, a chiral rotation yields the transformation $\Psi^\dagger(x)i\gamma_0\gamma_5\tau_2\Psi(x) \rightarrow \Psi^\dagger(x)\gamma_0\tau_2\Psi(x)$.
- (ii) We have not been able to prove that the state $\omega(\cdots)$ is pure¹. Therefore, when we consider the pure state decomposition for the state $\omega(\cdots)$, it is not obvious that there exists a pure state which shows a non-vanishing spontaneous magnetization for both of the order parameters. Physically, since each of the two flavors can take any direction for their chiral rotations, one can expect the existence of a pure state having the desired symmetry breaking character.

In other words, the present internal degrees of freedom allow to yield the two

¹In general, it is quite difficult to prove whether a given state is pure or not. Here, the term ‘‘pure state’’ refers to the concept in the context of infinite systems [7, 8], rather than its usage in usual quantum mechanics of finite degrees of freedom.

non-vanishing spontaneous magnetizations. In fact, these magnetizations can be expected from the commutativity $[\gamma_0\tau_1, i\gamma_0\gamma_5\tau_2] = 0$. Namely, the first spontaneous magnetization does not seem to prevent the second observable from magnetizing. Thus, the assumption of the two flavors is essential for our proof of the violation of the parity.

5. REFLECTION POSITIVITY: PRELIMINARY

Let $\Lambda' \subset \Lambda$ be a subset and $\mathcal{A}(\Lambda')$ be the algebra generated by $\psi_f^{(i)}(x)$ and $[\psi_{f'}^{(j)}(y)]^\dagger$ for $x, y \in \Lambda'$, $f, f' \in \{u, d\}$ and $i, j \in \{1, 2, 3, 4\}$. Since our Λ is symmetric with respect to a plane with the periodic boundary condition, there are a natural decomposition $\Lambda = \Lambda_- \cup \Lambda_+$ with $\Lambda_- \cap \Lambda_+ = \emptyset$ and a reflection map $r : \Lambda_\pm \rightarrow \Lambda_\mp$ satisfying $r(\Lambda_\pm) = \Lambda_\mp$. We write $\mathcal{A} = \mathcal{A}(\Lambda)$ and $\mathcal{A}_\pm = \mathcal{A}(\Lambda_\pm)$. The reflection has an anti-linear representation $\vartheta : \mathcal{A}_\pm \rightarrow \mathcal{A}_\mp$ requiring [17]

$$\begin{aligned} \vartheta(\psi_f^{(i)}(x)) &= \psi_f^{(i)}(\vartheta(x)), & \vartheta([\psi_f^{(i)}(x)]^\dagger) &= [\psi_f^{(i)}(\vartheta(x))]^\dagger, \\ \vartheta(AB) &= \vartheta(A)\vartheta(B), & \vartheta(A)^\dagger &= \vartheta(A^\dagger) \quad \text{for } A, B \in \mathcal{A}. \end{aligned}$$

For $x \in \Lambda$, we introduce Majorana fermion operators $\xi_f^{(i)}(x), \eta_f^{(i)}(x)$ by

$$\xi_f^{(i)}(x) := [\psi_f^{(i)}(x)]^\dagger + \psi_f^{(i)}(x), \quad \eta_f^{(i)}(x) := i\{[\psi_f^{(i)}(x)]^\dagger - \psi_f^{(i)}(x)\}, \quad (5.1)$$

or equivalently,

$$\psi_f^{(i)}(x) = \frac{1}{2}[\xi_f^{(i)}(x) + i\eta_f^{(i)}(x)], \quad [\psi_f^{(i)}(x)]^\dagger = \frac{1}{2}[\xi_f^{(i)}(x) - i\eta_f^{(i)}(x)]. \quad (5.2)$$

These satisfy $[\xi_f^{(i)}(x)]^\dagger = \xi_f^{(i)}(x)$, $[\eta_f^{(i)}(x)]^\dagger = \eta_f^{(i)}(x)$, and the anti-commutation relations

$$\begin{aligned} \{\xi_f^{(i)}(x), \xi_{f'}^{(j)}(y)\} &= 2\delta_{x,y}\delta_{i,j}\delta_{f,f'}, & \{\eta_f^{(i)}(x), \eta_{f'}^{(j)}(y)\} &= 2\delta_{x,y}\delta_{i,j}\delta_{f,f'}, \\ \{\xi_f^{(i)}(x), \eta_{f'}^{(j)}(y)\} &= 0. \end{aligned}$$

Next, following the idea of [12], we will introduce certain functions, h_μ , on the lattice Λ , and rewrite the interaction Hamiltonian. For this purpose, we first decompose the interaction Hamiltonian into two parts,

$$H_{\text{int}}^{(\Lambda)} = H_{\text{int,R}}^{(\Lambda)} + H_{\text{int,I}}^{(\Lambda)}$$

where

$$\begin{aligned} H_{\text{int,R}}^{(\Lambda)} &:= -g \sum_{x \in \Lambda} \sum_{\mu=1}^3 [\Gamma^{(1)}(x)\Gamma^{(1)}(x + e_\mu) + S^{(1)}(x)S^{(1)}(x + e_\mu) \\ &\quad + S^{(3)}(x)S^{(3)}(x + e_\mu) + S_5^{(2)}(x)S_5^{(2)}(x + e_\mu)] \end{aligned} \quad (5.3)$$

and

$$\begin{aligned}
 H_{\text{int,I}}^{(\Lambda)} := & -g \sum_{x \in \Lambda} \sum_{\mu=1}^3 [\Gamma^{(2)}(x) \Gamma^{(2)}(x + e_\mu) + S^{(2)}(x) S^{(2)}(x + e_\mu) \\
 & + S_5^{(1)}(x) S_5^{(1)}(x + e_\mu) + S_5^{(3)}(x) S_5^{(3)}(x + e_\mu)].
 \end{aligned} \tag{5.4}$$

Note that

$$\sum_{x \in \Lambda} \sum_{\mu=1}^3 \Gamma^{(1)}(x) \Gamma^{(1)}(x + e_\mu) = -\frac{1}{2} \sum_{x \in \Lambda} \sum_{\mu=1}^3 [\Gamma^{(1)}(x) - \Gamma^{(1)}(x + e_\mu)]^2 + 3 \sum_{x \in \Lambda} [\Gamma^{(1)}(x)]^2.$$

Therefore, one has

$$\begin{aligned}
 H_{\text{int,R}}^{(\Lambda)} = & \frac{g}{2} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \{ [\Gamma^{(1)}(x) - \Gamma^{(1)}(x + e_\mu)]^2 + [S^{(1)}(x) - S^{(1)}(x + e_\mu)]^2 \\
 & + [S^{(3)}(x) - S^{(3)}(x + e_\mu)]^2 + [S_5^{(2)}(x) - S_5^{(2)}(x + e_\mu)]^2 \} \\
 & - 3g \sum_{x \in \Lambda} \{ [\Gamma^{(1)}(x)]^2 + [S^{(1)}(x)]^2 + [S^{(3)}(x)]^2 + [S_5^{(2)}(x)]^2 \}.
 \end{aligned} \tag{5.5}$$

Similarly, one has

$$\sum_{x \in \Lambda} \sum_{\mu=1}^3 \Gamma^{(2)}(x) \Gamma^{(2)}(x + e_\mu) = \frac{1}{2} \sum_{x \in \Lambda} \sum_{\mu=1}^3 [\Gamma^{(2)}(x) + \Gamma^{(2)}(x + e_\mu)]^2 - 3 \sum_{x \in \Lambda} [\Gamma^{(2)}(x)]^2.$$

Therefore,

$$\begin{aligned}
 H_{\text{int,I}}^{(\Lambda)} = & -\frac{g}{2} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \{ [\Gamma^{(2)}(x) + \Gamma^{(2)}(x + e_\mu)]^2 + [S^{(2)}(x) + S^{(2)}(x + e_\mu)]^2 \\
 & + [S_5^{(1)}(x) + S_5^{(1)}(x + e_\mu)]^2 + [S_5^{(3)}(x) + S_5^{(3)}(x + e_\mu)]^2 \} \\
 & + 3g \sum_{x \in \Lambda} \{ [\Gamma^{(2)}(x)]^2 + [S^{(2)}(x)]^2 + [S_5^{(1)}(x)]^2 + [S_5^{(3)}(x)]^2 \}
 \end{aligned} \tag{5.6}$$

Let $h_\mu(x)$ be a real-valued function on the lattice Λ for $\mu = 1, 2, 3$, and we introduce [10, 12]

$$\begin{aligned}
 H_{\text{int,R}}^{(\Lambda)}(h) := & \frac{g}{2} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \{ [\Gamma^{(1)}(x) - \Gamma^{(1)}(x + e_\mu) + h_\mu(x)]^2 + [S^{(1)}(x) - S^{(1)}(x + e_\mu)]^2 \\
 & + [S^{(3)}(x) - S^{(3)}(x + e_\mu)]^2 + [S_5^{(2)}(x) - S_5^{(2)}(x + e_\mu)]^2 \} \\
 & - 3g \sum_{x \in \Lambda} \{ [\Gamma^{(1)}(x)]^2 + [S^{(1)}(x)]^2 + [S^{(3)}(x)]^2 + [S_5^{(2)}(x)]^2 \}.
 \end{aligned} \tag{5.7}$$

We write

$$H_{\text{int}}^{(\Lambda)}(h) := H_{\text{int,R}}^{(\Lambda)}(h) + H_{\text{int,I}}^{(\Lambda)}. \tag{5.8}$$

Clearly, when $h_\mu = 0$ for all $\mu = 1, 2, 3$, this equals the interaction Hamiltonian $H_{\text{int}}^{(\Lambda)}$. We also write

$$H^{(\Lambda)}(h) := H_{\text{K}}^{(\Lambda)} + H_{\text{int}}^{(\Lambda)}(h) \quad (5.9)$$

for the whole Hamiltonian.

6. REFLECTION WITH RESPECT TO THE PLANE $x^{(1)} = 1/2$

Let us consider first the reflection with respect to the $x^{(1)} = 1/2$ plane. By this plane, we divide our finite lattice Λ into two parts,

$$\Lambda_- := \{x \in \Lambda : -L + 1 \leq x^{(1)} \leq 0\} \quad \text{and} \quad \Lambda_+ := \{x \in \Lambda : 1 \leq x^{(1)} \leq L\}.$$

In order to show that the present Hamiltonian has a reflection positivity with respect to this plane, we need some preparations.

We first introduce a transformation,

$$\Psi_{\text{u}}(x) \rightarrow \gamma_0 \Psi_{\text{u}}(x), \quad \Psi_{\text{d}}(x) \rightarrow \Psi_{\text{d}}(x),$$

and write $U(\gamma_0)$ for the unitary transformation on the fermion Fock space. We also introduce a transformation,

$$\Psi(x) \rightarrow e^{i\pi x^{(2)}/2} \Psi(x),$$

and write U_2 for the corresponding unitary transformation on the fermion Fock space.

Let us consider first the kinetic Hamiltonian $H_{\text{K}}^{(\Lambda)}$ of (2.3) about these two transformations. We have written $H_{\text{K},f}^{(\Lambda)}$ for the kinetic Hamiltonian $H_{\text{K}}^{(\Lambda)}$ (2.3) for the flavor $f \in \{\text{u}, \text{d}\}$. Clearly, the first transformation $U(\gamma_0)$ changes the sign of the kinetic Hamiltonian for the flavor u, i.e.,

$$U^\dagger(\gamma_0) H_{\text{K},\text{u}}^{(\Lambda)} U(\gamma_0) = -H_{\text{K},\text{u}}^{(\Lambda)}.$$

Therefore, by these two transformations, $U(\gamma_0)$ and U_2 , the kinetic Hamiltonians $H_{\text{K},f}^{(\Lambda)}$ for the flavors $f = \text{u}, \text{d}$ are transformed into the following forms:

$$\begin{aligned} \tilde{H}_{\text{K},\text{u}}^{(\Lambda)} &:= [U(\gamma_0) U_2]^\dagger H_{\text{K},\text{u}}^{(\Lambda)} U(\gamma_0) U_2 \\ &= -i\kappa \sum_{x \in \Lambda \subset \mathbb{Z}^3} \left\{ [\Psi_{\text{u}}^\dagger(x) \alpha_1 \Psi_{\text{u}}(x + e_1) - \Psi_{\text{u}}^\dagger(x + e_1) \alpha_1 \Psi_{\text{u}}(x)] \right. \\ &\quad + i[\Psi_{\text{u}}^\dagger(x) \alpha_2 \Psi_{\text{u}}(x + e_2) + \Psi_{\text{u}}^\dagger(x + e_2) \alpha_2 \Psi_{\text{u}}(x)] \\ &\quad \left. + [\Psi_{\text{u}}^\dagger(x) \alpha_3 \Psi_{\text{u}}(x + e_3) - \Psi_{\text{u}}^\dagger(x + e_3) \alpha_3 \Psi_{\text{u}}(x)] \right\} \end{aligned} \quad (6.1)$$

and

$$\begin{aligned}
 \tilde{H}_{\text{K,d}}^{(\Lambda)} &:= [U(\gamma_0)U_2]^\dagger H_{\text{K,d}}^{(\Lambda)} U(\gamma_0)U_2 \\
 &= i\kappa \sum_{x \in \Lambda \subset \mathbb{Z}^3} \left\{ [\Psi_{\text{d}}^\dagger(x)\alpha_1\Psi_{\text{d}}(x+e_1) - \Psi_{\text{d}}^\dagger(x+e_1)\alpha_1\Psi_{\text{d}}(x)] \right. \\
 &\quad + i[\Psi_{\text{d}}^\dagger(x)\alpha_2\Psi_{\text{d}}(x+e_2) + \Psi_{\text{d}}^\dagger(x+e_2)\alpha_2\Psi_{\text{d}}(x)] \\
 &\quad \left. + [\Psi_{\text{d}}^\dagger(x)\alpha_3\Psi_{\text{d}}(x+e_3) - \Psi_{\text{d}}^\dagger(x+e_3)\alpha_3\Psi_{\text{d}}(x)] \right\}. \tag{6.2}
 \end{aligned}$$

When we use the real representation for the Dirac fermion field $\Psi(x)$, these right-hand sides become pure imaginary hermitian by the expressions of the matrices, $\alpha_1, \alpha_2, \alpha_3$. This property is crucial for the reflection positivity [17].

In order to deal with the interaction Hamiltonian, we note that

$$\begin{aligned}
 \tilde{\Gamma}^{(1)}(x) &:= [U(\gamma_0)U_2]^\dagger \Gamma^{(1)}(x) U(\gamma_0)U_2 = \Psi^\dagger(x)\gamma_0\Psi(x) = \Gamma^{(1)}(x), \\
 \tilde{\Gamma}^{(2)}(x) &:= [U(\gamma_0)U_2]^\dagger \Gamma^{(2)}(x) U(\gamma_0)U_2 = -\Psi^\dagger(x)i\gamma_0\gamma_5\tau_3\Psi(x) = -S_5^{(3)}(x), \tag{6.3} \\
 \tilde{S}^{(1)}(x) &:= [U(\gamma_0)U_2]^\dagger S^{(1)}(x) U(\gamma_0)U_2 = \Psi^\dagger(x)\tau_1\Psi(x), \\
 \tilde{S}^{(2)}(x) &:= [U(\gamma_0)U_2]^\dagger S^{(2)}(x) U(\gamma_0)U_2 = \Psi^\dagger(x)\tau_2\Psi(x), \\
 \tilde{S}^{(3)}(x) &:= [U(\gamma_0)U_2]^\dagger S^{(3)}(x) U(\gamma_0)U_2 = \Psi^\dagger(x)\gamma_0\tau_3\Psi(x) = S^{(3)}(x),
 \end{aligned}$$

and

$$\begin{aligned}
 \tilde{S}_5^{(1)}(x) &:= [U(\gamma_0)U_2]^\dagger S_5^{(1)}(x) U(\gamma_0)U_2 = -\Psi^\dagger(x)\gamma_5\tau_2\Psi(x), \\
 \tilde{S}_5^{(2)}(x) &:= [U(\gamma_0)U_2]^\dagger S_5^{(2)}(x) U(\gamma_0)U_2 = \Psi^\dagger(x)\gamma_5\tau_1\Psi(x)
 \end{aligned}$$

and

$$\tilde{S}_5^{(3)}(x) := [U(\gamma_0)U_2]^\dagger S_5^{(3)}(x) U(\gamma_0)U_2 = -\Psi^\dagger(x)i\gamma_0\gamma_5\Psi(x) = -\Gamma^{(2)}(x). \tag{6.4}$$

Then

$$\tilde{H}_{\text{int}}^{(\Lambda)}(h) := [U(\gamma_0)U_2]^\dagger H_{\text{int}}^{(\Lambda)}(h) U(\gamma_0)U_2 = \tilde{H}_{\text{int,R}}^{(\Lambda)}(h) + \tilde{H}_{\text{int,I}}^{(\Lambda)},$$

where we have written

$$\begin{aligned}
 \tilde{H}_{\text{int,R}}^{(\Lambda)}(h) &:= [U(\gamma_0)U_2]^\dagger H_{\text{int,R}}^{(\Lambda)}(h) U(\gamma_0)U_2 \\
 &= \frac{g}{2} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \left\{ [\tilde{\Gamma}^{(1)}(x) - \tilde{\Gamma}^{(1)}(x+e_\mu) + h_\mu(x)]^2 + [\tilde{S}^{(1)}(x) - \tilde{S}^{(1)}(x+e_\mu)]^2 \right. \\
 &\quad + [\tilde{S}^{(3)}(x) - \tilde{S}^{(3)}(x+e_\mu)]^2 + [\tilde{S}_5^{(2)}(x) - \tilde{S}_5^{(2)}(x+e_\mu)]^2 \left. \right\} \\
 &\quad - 3g \sum_{x \in \Lambda} \left\{ [\tilde{\Gamma}^{(1)}(x)]^2 + [\tilde{S}^{(1)}(x)]^2 + [\tilde{S}^{(3)}(x)]^2 + [\tilde{S}_5^{(2)}(x)]^2 \right\} \tag{6.5}
 \end{aligned}$$

and

$$\begin{aligned}
\tilde{H}_{\text{int,I}}^{(\Lambda)} &:= [U(\gamma_0)U_2]^\dagger H_{\text{int,I}}^{(\Lambda)} U(\gamma_0)U_2 \\
&= -\frac{g}{2} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \{ [\tilde{\Gamma}^{(2)}(x) + \tilde{\Gamma}^{(2)}(x + e_\mu)]^2 + [\tilde{S}^{(2)}(x) + \tilde{S}^{(2)}(x + e_\mu)]^2 \\
&\quad + [\tilde{S}_5^{(1)}(x) + \tilde{S}_5^{(1)}(x + e_\mu)]^2 + [\tilde{S}_5^{(3)}(x) + \tilde{S}_5^{(3)}(x + e_\mu)]^2 \} \\
&\quad + 3g \sum_{x \in \Lambda} \{ [\tilde{\Gamma}^{(2)}(x)]^2 + [\tilde{S}^{(2)}(x)]^2 + [\tilde{S}_5^{(1)}(x)]^2 + [\tilde{S}_5^{(3)}(x)]^2 \}.
\end{aligned} \tag{6.6}$$

The whole Hamiltonian $H^{(\Lambda)}(h)$ of (5.9) is transformed into

$$\begin{aligned}
\tilde{H}^{(\Lambda)}(h) &:= [U(\gamma_0)U_2]^\dagger H^{(\Lambda)}(h) U(\gamma_0)U_2 \\
&= \tilde{H}_{\text{K,u}}^{(\Lambda)} + \tilde{H}_{\text{K,d}}^{(\Lambda)} + \tilde{H}_{\text{int,R}}^{(\Lambda)}(h) + \tilde{H}_{\text{int,I}}^{(\Lambda)}.
\end{aligned} \tag{6.7}$$

6.1. Two unitary transformations. Further, we will use two unitary transformations below. First we define $U(\alpha_1)$ as follows:

$$[U(\alpha_1)]^\dagger \Psi_{\text{u}}(x) U(\alpha_1) = \begin{cases} -\alpha_1 \Psi_{\text{u}}(x) & \text{for } x \in \Lambda_+; \\ \Psi_{\text{u}}(x) & \text{for } x \in \Lambda_-, \end{cases} \tag{6.8}$$

and

$$[U(\alpha_1)]^\dagger \Psi_{\text{d}}(x) U(\alpha_1) = \begin{cases} \Psi_{\text{d}}(x) & \text{for } x \in \Lambda_+; \\ \alpha_1 \Psi_{\text{d}}(x) & \text{for } x \in \Lambda_-. \end{cases} \tag{6.9}$$

We also introduce [11, 13]

$$u_{\text{PH,f}}^{(i)}(x) := \left[\prod_{\substack{y \in \Lambda, j \in \{1,2,3,4\}, f' \in \{\text{u,d}\}: \\ (y,j,f') \neq (x,i,f)}} (-1)^{n_{f'}^{(j)}(y)} \right] \{ [\psi_f^{(i)}(x)]^\dagger + \psi_f^{(i)}(x) \}, \tag{6.10}$$

where we have written

$$n_{f'}^{(j)}(y) := [\psi_{f'}^{(j)}(y)]^\dagger \psi_{f'}^{(j)}(y)$$

for $y \in \Lambda$ and $f' \in \{\text{u,d}\}$. Then, one has

$$[u_{\text{PH,f}}^{(i)}(x)]^\dagger \psi_{f'}^{(j)}(y) u_{\text{PH,f}}^{(i)}(x) = \begin{cases} [\psi_f^{(i)}(x)]^\dagger, & \text{for } (y, j, f') = (x, i, f); \\ \psi_{f'}^{(j)}(y), & \text{otherwise.} \end{cases}$$

By using these operators, we define a particle-hole transformation on a sublattice by [11, 13]

$$U_{\text{odd,u}} := \prod_{x \in \Lambda_{\text{odd}}} \prod_{j \in \{1,2,3,4\}} u_{\text{PH,u}}^{(j)}(x),$$

where we have written

$$\Lambda_{\text{odd}} := \{x \in \Lambda \mid x^{(1)} + x^{(2)} + x^{(3)} = \text{odd}\}.$$

For all $i \in \{1, 2, 3, 4\}$, one has

$$(U_{\text{odd,u}})^\dagger \psi_{\text{u}}^{(i)}(x) U_{\text{odd,u}} = \begin{cases} [\psi_{\text{u}}^{(i)}(x)]^\dagger & \text{for } x \in \Lambda_{\text{odd}}, \\ \psi_{\text{u}}^{(i)}(x) & \text{for } x \in \Lambda \setminus \Lambda_{\text{odd}}. \end{cases} \quad (6.11)$$

Similarly, we define

$$U_{\text{even,d}} := \prod_{x \in \Lambda_{\text{even}}} \prod_{j \in \{1,2,3,4\}} u_{\text{PH,d}}^{(j)}(x).$$

for the flavor d and the sublattice $\Lambda_{\text{even}} := \Lambda \setminus \Lambda_{\text{odd}}$. Further, we write

$$U_{\text{stagg}} := U_{\text{odd,u}} U_{\text{even,d}}, \\ \tilde{U}_1 := U(\alpha_1) U_{\text{stagg}},$$

and

$$\hat{H}^{(\Lambda)}(h) := (\tilde{U}_1)^\dagger \tilde{H}^{(\Lambda)}(h) \tilde{U}_1$$

for the transformed Hamiltonian from (6.7). We want to decompose this Hamiltonian into some parts [13] in the following way.

6.2. Kinetic Hamiltonian. Let us consider the kinetic Hamiltonians (6.1) and (6.2). We first deal with $\tilde{H}_{\text{K,u}}^{(\Lambda)}$ of (6.1). It can be written

$$\tilde{H}_{\text{K,u}}^{(\Lambda)} = \sum_{\mu=1}^3 \tilde{H}_{\text{K,u},\mu}^{(\Lambda)}$$

with

$$\tilde{H}_{\text{K,u},\mu}^{(\Lambda)} := -i\kappa \sum_{x \in \Lambda} [\Psi_{\text{u}}^\dagger(x) \alpha_\mu \Psi_{\text{u}}(x + e_\mu) - \Psi_{\text{u}}^\dagger(x + e_\mu) \alpha_\mu \Psi_{\text{u}}(x)] \quad \text{for } \mu = 1, 3,$$

and

$$\tilde{H}_{\text{K,u},2}^{(\Lambda)} := \kappa \sum_{x \in \Lambda} [\Psi_{\text{u}}^\dagger(x) \alpha_2 \Psi_{\text{u}}(x + e_2) + \Psi_{\text{u}}^\dagger(x + e_2) \alpha_2 \Psi_{\text{u}}(x)].$$

Clearly, the hopping Hamiltonians in the second and third directions can be decomposed into two parts as follows:

$$\tilde{H}_{\text{K,u},2}^{(\Lambda)} = \tilde{H}_{\text{K,u},2}^+ + \tilde{H}_{\text{K,u},2}^-$$

with

$$\tilde{H}_{\text{K,u},2}^\pm := \kappa \sum_{x \in \Lambda_\pm} [\Psi_{\text{u}}^\dagger(x) \alpha_2 \Psi_{\text{u}}(x + e_2) + \Psi_{\text{u}}^\dagger(x + e_2) \alpha_2 \Psi_{\text{u}}(x)] \quad (6.12)$$

and

$$\tilde{H}_{\text{K,u},3}^{(\Lambda)} = \tilde{H}_{\text{K,u},3}^+ + \tilde{H}_{\text{K,u},3}^-$$

with

$$\tilde{H}_{\text{K,u},3}^\pm := -i\kappa \sum_{x \in \Lambda_\pm} [\Psi_{\text{u}}^\dagger(x) \alpha_3 \Psi_{\text{u}}(x + e_3) - \Psi_{\text{u}}^\dagger(x + e_3) \alpha_3 \Psi_{\text{u}}(x)]. \quad (6.13)$$

The kinetic term in the first direction is decomposed into three parts as follows:

$$\tilde{H}_{\text{K,u},1} = \tilde{H}_{\text{K,u},1}^+ + \tilde{H}_{\text{K,u},1}^- + \tilde{H}_{\text{K,u},1}^0,$$

where

$$\tilde{H}_{K,u,1}^+ := -i\kappa \sum_{x \in \Lambda_+ : x^{(1)} \neq L} [\Psi_u^\dagger(x) \alpha_1 \Psi_u(x + e_1) - \Psi_u^\dagger(x + e_1) \alpha_1 \Psi_u(x)], \quad (6.14)$$

$$\tilde{H}_{K,u,1}^- := -i\kappa \sum_{x \in \Lambda_- : x^{(1)} \neq 0} [\Psi_u^\dagger(x) \alpha_1 \Psi_u(x + e_1) - \Psi_u^\dagger(x + e_1) \alpha_1 \Psi_u(x)], \quad (6.15)$$

and

$$\begin{aligned} \tilde{H}_{K,u,1}^0 &:= -i\kappa \sum_{x \in \Lambda : x^{(1)} = 0} [\Psi_u^\dagger(x) \alpha_1 \Psi_u(x + e_1) - \Psi_u^\dagger(x + e_1) \alpha_1 \Psi_u(x)] \\ &\quad - i\kappa \sum_{x \in \Lambda : x^{(1)} = L} [\Psi_u^\dagger(x_L^-) \alpha_1 \Psi_u(x) - \Psi_u^\dagger(x) \alpha_1 \Psi_u(x_L^-)], \end{aligned} \quad (6.16)$$

where $x_L^- := (-L+1, x^{(2)}, x^{(3)})$, and we have used the anti-periodic boundary condition for the fermions.

By using the unitary transformation $U(\alpha_1)$ of (6.8), we have

$$\begin{aligned} [U(\alpha_1)]^\dagger \tilde{H}_{K,u,1}^0 U(\alpha_1) &= i\kappa \sum_{x \in \Lambda : x^{(1)} = 0} [\Psi_u^\dagger(x) \Psi_u(x + e_1) - \Psi_u^\dagger(x + e_1) \Psi_u(x)] \\ &\quad + i\kappa \sum_{x \in \Lambda : x^{(1)} = L} [\Psi_u^\dagger(x_L^-) \Psi_u(x) - \Psi_u^\dagger(x) \Psi_u(x_L^-)], \end{aligned}$$

where we have used α_1 is self-adjoint, and $(\alpha_1)^2 = 1$. In passing, we note the following: The negative sign in front of the right-hand side of (6.1) is canceled out with the negative sign in the right-hand side of (6.8). On the other hand, in the case of the flavor d of (6.2), both of the corresponding two signs are plus. Therefore, we obtain the same result in the case of the flavor d. Namely, for the flavor d, one has

$$\begin{aligned} [U(\alpha_1)]^\dagger \tilde{H}_{K,d,1}^0 U(\alpha_1) &= i\kappa \sum_{x \in \Lambda : x^{(1)} = 0} [\Psi_d^\dagger(x) \Psi_d(x + e_1) - \Psi_d^\dagger(x + e_1) \Psi_d(x)] \\ &\quad + i\kappa \sum_{x \in \Lambda : x^{(1)} = L} [\Psi_d^\dagger(x_L^-) \Psi_d(x) - \Psi_d^\dagger(x) \Psi_d(x_L^-)] \end{aligned}$$

from (6.2) and (6.9).

By using the Majorana fermions of (5.1), one has

$$[\psi_u^{(i)}(x)]^\dagger \psi_u^{(i)}(y) - [\psi_u^{(i)}(y)]^\dagger \psi_u^{(i)}(x) = \frac{1}{2} [\xi_u^{(i)}(x) \xi_u^{(i)}(y) + \eta_u^{(i)}(x) \eta_u^{(i)}(y)] \quad \text{for } x \neq y.$$

Therefore, we have

$$\begin{aligned} [U(\alpha_1)]^\dagger \tilde{H}_{K,u,1}^0 U(\alpha_1) &= \frac{i\kappa}{2} \sum_{x \in \Lambda : x^{(1)} = 0} \sum_{i=1}^4 [\xi_u^{(i)}(x) \xi_u^{(i)}(x + e_1) + \eta_u^{(i)}(x) \eta_u^{(i)}(x + e_1)] \\ &\quad + \frac{i\kappa}{2} \sum_{x \in \Lambda : x^{(1)} = L} \sum_{i=1}^4 [\xi_u^{(i)}(x_L^-) \xi_u^{(i)}(x) + \eta_u^{(i)}(x_L^-) \eta_u^{(i)}(x)]. \end{aligned} \quad (6.17)$$

From the representations (5.1) of the Majorana fermions and (6.11), one has

$$(U_{\text{odd,u}})^\dagger \xi_{\text{u}}^{(i)}(x) U_{\text{odd,u}} = \xi_{\text{u}}^{(i)}(x) \quad \text{for all } x \in \Lambda \text{ and } i = 1, 2, 3, 4,$$

and

$$(U_{\text{odd,u}})^\dagger \eta_{\text{u}}^{(i)}(x) U_{\text{odd,u}} = \begin{cases} -\eta_{\text{u}}^{(i)}(x) & \text{for } x \in \Lambda_{\text{odd}}; \\ \eta_{\text{u}}^{(i)}(x) & \text{for } x \in \Lambda \setminus \Lambda_{\text{odd}} \end{cases}$$

for all $i = 1, 2, 3, 4$. By combining this with $\tilde{U}_1 = U(\alpha_1)U_{\text{stag}}$, the definition of the reflection map ϑ and (6.17), we obtain the desired expression,

$$\begin{aligned} \hat{H}_{\text{K,u,1}}^0 &:= (\tilde{U}_1)^\dagger \tilde{H}_{\text{K,u,1}}^0 \tilde{U}_1 = \frac{i\kappa}{2} \sum_{x \in \Lambda : x^{(1)}=0} \sum_{i=1}^4 [\xi_{\text{u}}^{(i)}(x) \xi_{\text{u}}^{(i)}(x + e_1) - \eta_{\text{u}}^{(i)}(x) \eta_{\text{u}}^{(i)}(x + e_1)] \\ &\quad + \frac{i\kappa}{2} \sum_{x \in \Lambda : x^{(1)}=L} \sum_{i=1}^4 [\xi_{\text{u}}^{(i)}(x_L^-) \xi_{\text{u}}^{(i)}(x) - \eta_{\text{u}}^{(i)}(x_L^-) \eta_{\text{u}}^{(i)}(x)] \\ &= \frac{i\kappa}{2} \sum_{x \in \Lambda : x^{(1)}=0} \sum_{i=1}^4 [\xi_{\text{u}}^{(i)}(x) \vartheta(\xi_{\text{u}}^{(i)}(x)) + \eta_{\text{u}}^{(i)}(x) \vartheta(\eta_{\text{u}}^{(i)}(x))] \\ &\quad + \frac{i\kappa}{2} \sum_{x \in \Lambda : x^{(1)}=-L+1} \sum_{i=1}^4 [\xi_{\text{u}}^{(i)}(x) \vartheta(\xi_{\text{u}}^{(i)}(x)) + \eta_{\text{u}}^{(i)}(x) \vartheta(\eta_{\text{u}}^{(i)}(x))]. \end{aligned} \tag{6.18}$$

This is nothing but the desired form for the reflection positivity [13, 17].

As to $\tilde{H}_{\text{K,u,1}}^\pm$ of (6.14) and (6.15), one notices that both of the two Hamiltonians do not change under the $U(\alpha_1)$ transformation of (6.8). Further, since the matrix α_1 is symmetric, i.e., its transpose equals itself, one has

$$\begin{aligned} &(U_{\text{odd,u}})^\dagger [\Psi_{\text{u}}^\dagger(x) \alpha_1 \Psi_{\text{u}}(x + e_1) - \Psi_{\text{u}}^\dagger(x + e_1) \alpha_1 \Psi_{\text{u}}(x)] U_{\text{odd,u}} \\ &= \Psi_{\text{u}}^\dagger(x) \alpha_1 {}^t \Psi_{\text{u}}^\dagger(x + e_1) - {}^t \Psi_{\text{u}}(x + e_1) \alpha_1 \Psi_{\text{u}}(x) \end{aligned}$$

for any $x \in \Lambda$, where the superscript ‘t’ denotes the transpose, namely

$${}^t \Psi_{\text{f}}^\dagger(x) = \begin{pmatrix} \psi_{\text{f}}^{(1)}(x)^\dagger \\ \psi_{\text{f}}^{(2)}(x)^\dagger \\ \psi_{\text{f}}^{(3)}(x)^\dagger \\ \psi_{\text{f}}^{(4)}(x)^\dagger \end{pmatrix}, \quad {}^t \Psi_{\text{f}}(x) = \left(\psi_{\text{f}}^{(1)}(x), \psi_{\text{f}}^{(2)}(x), \psi_{\text{f}}^{(3)}(x), \psi_{\text{f}}^{(4)}(x) \right).$$

Therefore, we have

$$\hat{H}_{\text{K,u,1}}^+ := (\tilde{U}_1)^\dagger \tilde{H}_{\text{K,u,1}}^+ \tilde{U}_1 = -i\kappa \sum_{x \in \Lambda_+ : x^{(1)} \neq L} [\Psi_{\text{u}}^\dagger(x) \alpha_1 {}^t \Psi_{\text{u}}^\dagger(x + e_1) - {}^t \Psi_{\text{u}}(x + e_1) \alpha_1 \Psi_{\text{u}}(x)]$$

and

$$\hat{H}_{\text{K,u,1}}^- := (\tilde{U}_1)^\dagger \tilde{H}_{\text{K,u,1}}^- \tilde{U}_1 = -i\kappa \sum_{x \in \Lambda_- : x^{(1)} \neq 0} [\Psi_{\text{u}}^\dagger(x) \alpha_1 {}^t \Psi_{\text{u}}^\dagger(x + e_1) - {}^t \Psi_{\text{u}}(x + e_1) \alpha_1 \Psi_{\text{u}}(x)].$$

In addition, since the reflection map ϑ changes the hopping direction, we obtain

$$\hat{H}_{K,u,1}^+ = \vartheta(\hat{H}_{K,u,1}^-),$$

where we have used that the matrix α_1 is real hermitian. In passing, since this argument is independent of the sign of the hopping amplitude, the relation holds in the case of the flavor d.

Next, consider the kinetic Hamiltonian $\tilde{H}_{K,u,3}^\pm$ of (6.13) in the third direction. From the properties of the matrices α_i , one notices that

$$(U(\alpha_1))^\dagger \tilde{H}_{K,u,3}^+ U(\alpha_1) = -\tilde{H}_{K,u,3}^+ \quad \text{and} \quad (U(\alpha_1))^\dagger \tilde{H}_{K,u,3}^- U(\alpha_1) = \tilde{H}_{K,u,3}^-.$$

Further, one has

$$\begin{aligned} & (U_{\text{odd},u})^\dagger [\Psi_u^\dagger(x) \alpha_3 \Psi_u(x + e_3) - \Psi_u^\dagger(x + e_3) \alpha_3 \Psi_u(x)] U_{\text{odd},u} \\ &= \Psi_u^\dagger(x) \alpha_3 {}^t \Psi_u^\dagger(x + e_3) - {}^t \Psi_u(x + e_3) \alpha_3 \Psi_u(x) \end{aligned}$$

because the matrix α_3 is symmetric. From these observations, we have

$$\hat{H}_{K,u,3}^\pm := (\tilde{U}_1)^\dagger \tilde{H}_{K,u,3}^\pm \tilde{U}_1 = \pm i\kappa \sum_{x \in \Lambda_\pm} [\Psi_u^\dagger(x) \alpha_3 {}^t \Psi_u^\dagger(x + e_3) - {}^t \Psi_u(x + e_3) \alpha_3 \Psi_u(x)]. \quad (6.19)$$

This implies

$$\hat{H}_{K,u,3}^+ = \vartheta(\hat{H}_{K,u,3}^-)$$

because the matrix α_3 is real hermitian.

Finally, let us consider $\tilde{H}_{K,u,2}^\pm$ of (6.12). In the same way, one has

$$[U(\alpha_1)]^\dagger \tilde{H}_{K,u,2}^\pm U(\alpha_1) = \mp \kappa \sum_{x \in \Lambda_\pm} [\Psi_u^\dagger(x) \alpha_2 \Psi_u(x + e_2) + \Psi_u^\dagger(x + e_2) \alpha_2 \Psi_u(x)].$$

Note that

$$\begin{aligned} & (U_{\text{odd},u})^\dagger [\Psi_u^\dagger(x) \alpha_2 \Psi_u(x + e_2) + \Psi_u^\dagger(x + e_2) \alpha_2 \Psi_u(x)] U_{\text{odd},u} \\ &= \Psi_u^\dagger(x) \alpha_2 {}^t \Psi_u^\dagger(x + e_2) + {}^t \Psi_u(x + e_2) \alpha_2 \Psi_u(x) \end{aligned}$$

for any $x \in \Lambda$, where we have used that the matrix α_2 is anti-symmetric. By combining these two, we obtain

$$\hat{H}_{K,u,2}^\pm := (\tilde{U}_1)^\dagger \tilde{H}_{K,u,2}^\pm \tilde{U}_1 = \mp \kappa \sum_{x \in \Lambda_\pm} [\Psi_u^\dagger(x) \alpha_2 {}^t \Psi_u^\dagger(x + e_2) + {}^t \Psi_u(x + e_2) \alpha_2 \Psi_u(x)].$$

Since the matrix α_2 is pure imaginary hermitian, we have

$$\hat{H}_{K,u,2}^+ = \vartheta(\hat{H}_{K,u,2}^-).$$

As mentioned in some comments above, these arguments hold for the case of the flavor d.

6.3. Interaction Hamiltonian $\tilde{H}_{\text{int,R}}^{(\Lambda)}(h)$. Next, consider the interaction Hamiltonian $\tilde{H}_{\text{int,R}}^{(\Lambda)}(h)$ of (6.5). For the part about the operator $\tilde{\Gamma}^{(1)}(x)$, we write

$$\tilde{H}_{\tilde{\Gamma}^{(1)}}^{(\Lambda)}(h) = \sum_{\mu=1}^3 \tilde{H}_{\tilde{\Gamma}^{(1)},\mu}^{(\Lambda)}(h) - 3g \sum_{x \in \Lambda} [\tilde{\Gamma}^{(1)}(x)]^2, \quad (6.20)$$

where

$$\tilde{H}_{\tilde{\Gamma}^{(1)},\mu}^{(\Lambda)}(h) := \frac{g}{2} \sum_{x \in \Lambda} [\tilde{\Gamma}^{(1)}(x) - \tilde{\Gamma}^{(1)}(x + e_\mu) + h_\mu(x)]^2.$$

The Hamiltonian $\tilde{H}_{\tilde{\Gamma}^{(1)},1}^{(\Lambda)}(h)$ can be decomposed into three parts as follows:

$$\tilde{H}_{\tilde{\Gamma}^{(1)},1}^{(\Lambda)}(h) = \tilde{H}_{\tilde{\Gamma}^{(1)},1}^+(h) + \tilde{H}_{\tilde{\Gamma}^{(1)},1}^-(h) + \tilde{H}_{\tilde{\Gamma}^{(1)},1}^0(h),$$

where

$$\tilde{H}_{\tilde{\Gamma}^{(1)},1}^+(h) := \frac{g}{2} \sum_{\substack{x \in \Lambda_+ \\ x^{(1)} \neq L}} [\tilde{\Gamma}^{(1)}(x) - \tilde{\Gamma}^{(1)}(x + e_1) + h_1(x)]^2,$$

$$\tilde{H}_{\tilde{\Gamma}^{(1)},1}^-(h) := \frac{g}{2} \sum_{\substack{x \in \Lambda_- \\ x^{(1)} \neq 0}} [\tilde{\Gamma}^{(1)}(x) - \tilde{\Gamma}^{(1)}(x + e_1) + h_1(x)]^2,$$

and

$$\tilde{H}_{\tilde{\Gamma}^{(1)},1}^0(h) := \frac{g}{2} \sum_{\substack{x \in \Lambda \\ x^{(1)} = 0, L}} [\tilde{\Gamma}^{(1)}(x) - \tilde{\Gamma}^{(1)}(x + e_1) + h_1(x)]^2. \quad (6.21)$$

Similarly, we have

$$\tilde{H}_{\tilde{\Gamma}^{(1)},\mu}^{(\Lambda)}(h) = \tilde{H}_{\tilde{\Gamma}^{(1)},\mu}^+(h) + \tilde{H}_{\tilde{\Gamma}^{(1)},\mu}^-(h) \quad \text{for } \mu = 2, 3,$$

where

$$\tilde{H}_{\tilde{\Gamma}^{(1)},\mu}^\pm(h) := \frac{g}{2} \sum_{x \in \Lambda_\pm} [\tilde{\Gamma}^{(1)}(x) - \tilde{\Gamma}^{(1)}(x + e_\mu) + h_\mu(x)]^2 \quad \text{for } \mu = 2, 3.$$

From the definitions of $\tilde{\Gamma}^{(1)}(x)$ and $U(\alpha_1)$, one has

$$[U(\alpha_1)]^\dagger \tilde{\Gamma}^{(1)}(x) U(\alpha_1) = \begin{cases} -\Psi_u^\dagger(x) \gamma_0 \Psi_u(x) + \Psi_d^\dagger(x) \gamma_0 \Psi_d(x) & \text{if } x \in \Lambda_+; \\ \Psi_u^\dagger(x) \gamma_0 \Psi_u(x) - \Psi_d^\dagger(x) \gamma_0 \Psi_d(x) & \text{if } x \in \Lambda_-. \end{cases}$$

Further, by using $\tilde{U}_1 := U(\alpha_1) U_{\text{stagg}}$, we have

$$(\tilde{U}_1)^\dagger \tilde{\Gamma}^{(1)}(x) \tilde{U}_1 = \mp (-1)^{x^{(1)} + x^{(2)} + x^{(3)}} \tilde{\Gamma}^{(1)}(x) \quad \text{for } x \in \Lambda_\pm, \quad (6.22)$$

where we have used the expression of the matrix γ_0 in (2.7). For the Hamiltonian $\tilde{H}_{\tilde{\Gamma}^{(1)},1}^0(h)$ of (6.21), we have

$$\begin{aligned} \tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(1)},1}^0(h) \tilde{U}_1 &= \frac{g}{2} \sum_{\substack{x \in \Lambda: \\ x^{(1)}=0}} [\tilde{\Gamma}^{(1)}(x) - \tilde{\Gamma}^{(1)}(x + e_1) + (-1)^{x^{(1)}+x^{(2)}+x^{(3)}} h_1(x)]^2 \\ &\quad + \frac{g}{2} \sum_{\substack{x \in \Lambda: \\ x^{(1)}=L}} [\tilde{\Gamma}^{(1)}(x) - \tilde{\Gamma}^{(1)}(x + e_1) - (-1)^{x^{(1)}+x^{(2)}+x^{(3)}} h_1(x)]^2. \end{aligned}$$

This is the desired form [10, 13] for getting a Gaussian domination. Similarly, one has

$$\tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(1)},1}^+(h) \tilde{U}_1 = \frac{g}{2} \sum_{\substack{x \in \Lambda_+: \\ x^{(1)} \neq L}} [\tilde{\Gamma}^{(1)}(x) + \tilde{\Gamma}^{(1)}(x + e_1) - \tilde{h}_1(x)]^2$$

with $\tilde{h}_1(x) := (-1)^{x^{(1)}+x^{(2)}+x^{(3)}} h_1(x)$, and

$$\tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(1)},1}^-(h) \tilde{U}_1 = \frac{g}{2} \sum_{\substack{x \in \Lambda_-: \\ x^{(1)} \neq 0}} [\tilde{\Gamma}^{(1)}(x) + \tilde{\Gamma}^{(1)}(x + e_1) + \tilde{h}_1(x)]^2.$$

Moreover for $\mu = 2, 3$, we have

$$\tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(1)},\mu}^+(h) \tilde{U}_1 = \frac{g}{2} \sum_{x \in \Lambda_+} [\tilde{\Gamma}^{(1)}(x) + \tilde{\Gamma}^{(1)}(x + e_\mu) - \tilde{h}_\mu(x)]^2$$

and

$$\tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(1)},\mu}^-(h) \tilde{U}_1 = \frac{g}{2} \sum_{x \in \Lambda_-} [\tilde{\Gamma}^{(1)}(x) + \tilde{\Gamma}^{(1)}(x + e_\mu) + \tilde{h}_\mu(x)]^2$$

with $\tilde{h}_\mu(x) := (-1)^{x^{(1)}+x^{(2)}+x^{(3)}} h_\mu(x)$. In particular, when $h = 0$, these imply

$$\vartheta(\hat{H}_{\tilde{\Gamma}^{(1)},\mu}^-(0)) = \hat{H}_{\tilde{\Gamma}^{(1)},\mu}^+(0) \quad \text{for } \mu = 1, 2, 3,$$

where we have written

$$\hat{H}_{\tilde{\Gamma}^{(1)},\mu}^\pm(h) := \tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(1)},\mu}^\pm(h) \tilde{U}_1$$

for $\mu = 1, 2, 3$.

Clearly, the second sum in the right hand side of (6.20) can be treated in the same way.

In the above argument, the relation (6.22) is crucial for the reflection positivity. Therefore, as to the four operators, $\tilde{S}^{(1)}(x)$, $\tilde{S}^{(3)}(x)$ and $\tilde{S}_5^{(2)}(x)$, in the Hamiltonian $\tilde{H}_{\text{int,R}}^{(\Lambda)}(h)$, it is enough to check the corresponding relations. Actually, in the same way, we have

$$\tilde{U}_1^\dagger \tilde{S}^{(1)}(x) \tilde{U}_1 = \pm (-1)^{x^{(1)}+x^{(2)}+x^{(3)}} [\text{t}\Psi_u(x) \alpha_1 \Psi_d(x) + \Psi_d^\dagger(x) \alpha_1 \text{t}\Psi_u^\dagger(x)] \quad \text{for } x \in \Lambda_\pm,$$

$$\tilde{U}_1^\dagger \tilde{S}^{(3)}(x) \tilde{U}_1 = \pm (-1)^{x^{(1)}+x^{(2)}+x^{(3)}} [-\Psi_u^\dagger(x) \gamma_0 \Psi_u(x) + \Psi_d^\dagger(x) \gamma_0 \Psi_d(x)] \quad \text{for } x \in \Lambda_\pm$$

and

$$\tilde{U}_1^\dagger \tilde{S}_5^{(2)}(x) \tilde{U}_1 = \mp (-1)^{x^{(1)}+x^{(2)}+x^{(3)}} [\Psi_u^\dagger(x) \gamma_5 \alpha_1 {}^t\Psi_d^\dagger(x) + {}^t\Psi_d(x) \gamma_5 \alpha_1 \Psi_u(x)] \quad \text{for } x \in \Lambda_\pm.$$

6.4. Interaction Hamiltonian $\tilde{H}_{\text{int,I}}^{(\Lambda)}$. Let us consider the interaction Hamiltonian $\tilde{H}_{\text{int,I}}^{(\Lambda)}$ of (6.6). The part about the operator $\tilde{\Gamma}^{(2)}(x)$ of (6.3) is given by

$$\tilde{H}_{\tilde{\Gamma}^{(2)}}^{(\Lambda)} := \sum_{\mu=1}^3 \tilde{H}_{\tilde{\Gamma}^{(2)},\mu}^{(\Lambda)} + 3g \sum_{x \in \Lambda} [\tilde{\Gamma}^{(2)}(x)]^2, \quad (6.23)$$

where

$$\tilde{H}_{\tilde{\Gamma}^{(2)},\mu}^{(\Lambda)} := -\frac{g}{2} \sum_{x \in \Lambda} [\tilde{\Gamma}^{(2)}(x) + \tilde{\Gamma}^{(2)}(x + e_\mu)]^2.$$

The Hamiltonian $\tilde{H}_{\tilde{\Gamma}^{(2)},1}^{(\Lambda)}$ can be decomposed into three parts as follows:

$$\tilde{H}_{\tilde{\Gamma}^{(2)},1}^{(\Lambda)} = \tilde{H}_{\tilde{\Gamma}^{(2)},1}^+ + \tilde{H}_{\tilde{\Gamma}^{(2)},1}^- + \tilde{H}_{\tilde{\Gamma}^{(2)},1}^0,$$

where

$$\tilde{H}_{\tilde{\Gamma}^{(2)},1}^+ := -\frac{g}{2} \sum_{\substack{x \in \Lambda_+ \\ x^{(1)} \neq L}} [\tilde{\Gamma}^{(2)}(x) + \tilde{\Gamma}^{(2)}(x + e_1)]^2,$$

$$\tilde{H}_{\tilde{\Gamma}^{(2)},1}^- := -\frac{g}{2} \sum_{\substack{x \in \Lambda_- \\ x^{(1)} \neq 0}} [\tilde{\Gamma}^{(2)}(x) + \tilde{\Gamma}^{(2)}(x + e_1)]^2,$$

and

$$\tilde{H}_{\tilde{\Gamma}^{(2)},1}^0 := -\frac{g}{2} \sum_{\substack{x \in \Lambda \\ x^{(1)}=0,L}} [\tilde{\Gamma}^{(2)}(x) + \tilde{\Gamma}^{(2)}(x + e_1)]^2. \quad (6.24)$$

Similarly, we have

$$\tilde{H}_{\tilde{\Gamma}^{(2)},\mu}^{(\Lambda)} = \tilde{H}_{\tilde{\Gamma}^{(2)},\mu}^+ + \tilde{H}_{\tilde{\Gamma}^{(2)},\mu}^- \quad \text{for } \mu = 2, 3,$$

where

$$\tilde{H}_{\tilde{\Gamma}^{(2)},\mu}^\pm := -\frac{g}{2} \sum_{x \in \Lambda_\pm} [\tilde{\Gamma}^{(2)}(x) + \tilde{\Gamma}^{(2)}(x + e_\mu)]^2 \quad \text{for } \mu = 2, 3.$$

One has

$$[U(\alpha_1)]^\dagger \tilde{\Gamma}^{(2)}(x) U(\alpha_1) = \begin{cases} \Psi_u^\dagger(x) i \gamma_0 \gamma_5 \Psi_u(x) + \Psi_d^\dagger(x) i \gamma_0 \gamma_5 \Psi_d(x) & \text{if } x \in \Lambda_+; \\ -\Psi_u^\dagger(x) i \gamma_0 \gamma_5 \Psi_u(x) - \Psi_d^\dagger(x) i \gamma_0 \gamma_5 \Psi_d(x) & \text{if } x \in \Lambda_-. \end{cases}$$

Therefore, we obtain

$$(\tilde{U}_1)^\dagger \tilde{\Gamma}^{(2)}(x) \tilde{U}_1 = \pm \Gamma^{(2)}(x) \quad \text{for } x \in \Lambda_\pm, \quad (6.25)$$

where we have used that the matrix $\gamma_0 \gamma_5$ is anti-symmetric. For the Hamiltonian $\tilde{H}_{\tilde{\Gamma}^{(2)},1}^0$ of (6.24), we have

$$\tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(2)},1}^0 \tilde{U}_1 = -\frac{g}{2} \sum_{\substack{x \in \Lambda \\ x^{(1)}=0,L}} [\Gamma^{(2)}(x) - \Gamma^{(2)}(x + e_1)]^2.$$

This is the desired form [10, 13] for getting a Gaussian domination. Similarly, one has

$$\tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(2)},1}^+ \tilde{U}_1 = -\frac{g}{2} \sum_{\substack{x \in \Lambda_+ : \\ x^{(1)} \neq L}} [\Gamma^{(2)}(x) + \Gamma^{(2)}(x + e_1)]^2$$

and

$$\tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(2)},1}^- \tilde{U}_1 = -\frac{g}{2} \sum_{\substack{x \in \Lambda_- : \\ x^{(1)} \neq 0}} [\Gamma^{(2)}(x) + \Gamma^{(2)}(x + e_1)]^2.$$

Further, we have

$$\tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(2)},\mu}^+ \tilde{U}_1 = -\frac{g}{2} \sum_{x \in \Lambda_+} [\Gamma^{(2)}(x) + \Gamma^{(2)}(x + e_\mu)]^2,$$

and

$$\tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(2)},\mu}^- \tilde{U}_1 = -\frac{g}{2} \sum_{x \in \Lambda_-} [\Gamma^{(2)}(x) + \Gamma^{(2)}(x + e_\mu)]^2$$

for $\mu = 2, 3$. From these observations, we have

$$\vartheta \left(\hat{H}_{\tilde{\Gamma}^{(2)},\mu}^- \right) = \hat{H}_{\tilde{\Gamma}^{(2)},\mu}^+ \quad \text{for } \mu = 1, 2, 3,$$

where we have written

$$\hat{H}_{\tilde{\Gamma}^{(2)},\mu}^\pm := \tilde{U}_1^\dagger \tilde{H}_{\tilde{\Gamma}^{(2)},\mu}^\pm \tilde{U}_1 \quad \text{for } \mu = 1, 2, 3,$$

and we have used that the matrix $i\gamma_0\gamma_5$ in $\Gamma^{(2)}(x)$ is pure imaginary hermitian.

Clearly, the second sum in the right-hand side of (6.23) can be treated in the same way.

In the above argument, the relation (6.25) is crucial again. Therefore, as to the rest of the operators in the interaction Hamiltonian $\tilde{H}_{\text{int},I}^{(\Lambda)}$ of (6.6), it is enough to check the corresponding relations. One can check that

$$\tilde{U}_1^\dagger \tilde{S}^{(2)}(x) \tilde{U}_1 = \pm i [\Psi_u^\dagger(x) \alpha_1 {}^t \Psi_d^\dagger(x) - {}^t \Psi_d(x) \alpha_1 \Psi_u(x)] \quad \text{for } x \in \Lambda_\pm,$$

$$\tilde{U}_1^\dagger \tilde{S}_5^{(1)}(x) \tilde{U}_1 = \mp i [\Psi_u^\dagger(x) \alpha_1 \gamma_5 {}^t \Psi_d^\dagger(x) - {}^t \Psi_d(x) \alpha_1 \gamma_5 \Psi_u(x)] \quad \text{for } x \in \Lambda_\pm$$

and

$$\tilde{U}_1^\dagger \tilde{S}_5^{(3)}(x) \tilde{U}_1 = \pm S_5^{(3)}(x) \quad \text{for } x \in \Lambda_\pm.$$

7. REFLECTION WITH RESPECT TO THE PLANE $x^{(2)} = 1/2$

As to the reflection with respect to the $x^{(1)}$ - $x^{(2)}$ plane, the argument is the same as in the above case of the $x^{(1)} = 1/2$ plane. Therefore, it is enough to deal with the case of the reflection with respect to the $x^{(2)} = 1/2$ plane.

We write

$$u_3(\theta) := \exp \left[i \frac{\theta}{2} \sigma_3 \right]$$

for the spin rotation by the angle $\theta \in [0, 2\pi)$ about the third axis, and

$$\mathcal{U}_3(\theta) := \begin{pmatrix} 0 & u_3(\theta) \\ u_3(\theta) & 0 \end{pmatrix}$$

for the four-component Dirac spinor. We also write $U_3(\theta)$ for the corresponding unitary operator on the fermion Fock space, i.e.,

$$[U_3(\theta)]^\dagger \Psi(x) U_3(\theta) = \mathcal{U}_3(\theta) \Psi(x) \quad \text{for } x \in \Lambda.$$

Note that

$$[\mathcal{U}_3(\theta)]^\dagger \alpha_i \mathcal{U}_3(\theta) = \begin{pmatrix} 0 & [u_3(\theta)]^\dagger \sigma_i u_3(\theta) \\ [u_3(\theta)]^\dagger \sigma_i u_3(\theta) & 0 \end{pmatrix} \quad \text{for } i = 1, 2, 3,$$

$$[u_3(\theta)]^\dagger \sigma_1 u_3(\theta) = \sigma_1 \cos \theta + \sigma_2 \sin \theta,$$

$$[u_3(\theta)]^\dagger \sigma_2 u_3(\theta) = \sigma_2 \cos \theta - \sigma_1 \sin \theta,$$

and

$$[u_3(\theta)]^\dagger \sigma_3 u_3(\theta) = \sigma_3.$$

Therefore, one has

$$[\mathcal{U}_3(-\pi/2)]^\dagger \alpha_1 \mathcal{U}_3(-\pi/2) = -\alpha_2, \quad (7.1)$$

$$[\mathcal{U}_3(-\pi/2)]^\dagger \alpha_2 \mathcal{U}_3(-\pi/2) = \alpha_1, \quad (7.2)$$

and

$$[\mathcal{U}_3(-\pi/2)]^\dagger \alpha_3 \mathcal{U}_3(-\pi/2) = \alpha_3. \quad (7.3)$$

Moreover,

$$[\mathcal{U}_3(-\pi/2)]^\dagger \gamma_0 \mathcal{U}_3(-\pi/2) = -\gamma_0 \quad (7.4)$$

and

$$[\mathcal{U}_3(-\pi/2)]^\dagger \gamma_5 \mathcal{U}_3(-\pi/2) = \gamma_5. \quad (7.5)$$

Clearly, these change only the matrices, α_1 and α_2 , of the hopping terms in the $x^{(1)}$ and $x^{(2)}$ directions in the kinetic Hamiltonian. In addition, although the coefficient κ in front of the matrix α_2 changes its sign, it does not affect the above argument about the reflection positivity in the case of the $x^{(1)} = 1/2$ plane.

8. GAUSSIAN DOMINATION AND INFRARED BOUND

By reflection positivity, we obtain the Gaussian domination bound

$$\text{Tr exp}[-\beta H^{(\Lambda)}(h)] \leq \text{Tr exp}[-\beta H^{(\Lambda)}(0)],$$

for the Hamiltonian $H^{(\Lambda)}(h)$ of (5.9). The proof proceeds in a similar manner to that in [13,14], and is omitted here to avoid repetition. In order to expand this with respect to the function h , we note that

$$\begin{aligned} & \frac{g}{2} \sum_{x \in \Lambda} \sum_{\mu=1}^3 [\Gamma^{(1)}(x) - \Gamma^{(1)}(x + e_\mu) + h_\mu(x)]^2 \\ &= \frac{g}{2} \sum_{x \in \Lambda} \sum_{\mu=1}^3 [\Gamma^{(1)}(x) - \Gamma^{(1)}(x + e_\mu)]^2 \\ & \quad + g \sum_{x \in \Lambda} \sum_{\mu=1}^3 [\Gamma^{(1)}(x) - \Gamma^{(1)}(x + e_\mu)] h_\mu(x) + \frac{g}{2} \sum_{x \in \Lambda} \sum_{\mu=1}^3 [h_\mu(x)]^2. \end{aligned}$$

The second sum in the right-hand side is written

$$g \sum_{x \in \Lambda} \sum_{\mu=1}^3 [\Gamma^{(1)}(x) - \Gamma^{(1)}(x + e_\mu)] h_\mu(x) = g \sum_{x \in \Lambda} \sum_{\mu=1}^3 \Gamma^{(1)}(x) \partial_\mu h_\mu(x),$$

where we have written

$$\partial_\mu h_\mu(x) := h_\mu(x) - h_\mu(x - e_\mu).$$

Therefore, the second order of the expansion yields

$$(\beta g)^2 \left(\sum_{x \in \Lambda} \sum_{\mu=1}^3 \Gamma^{(1)}(x) \partial_\mu h_\mu(x), \sum_{y \in \Lambda} \sum_{\mu'=1}^3 \Gamma^{(1)}(y) \partial_{\mu'} h_{\mu'}(y) \right)_\beta^{(\Lambda)} - \frac{\beta g}{2} \sum_{x \in \Lambda} \sum_{\mu=1}^3 [h_\mu(x)]^2 \leq 0,$$

where $(\dots, \dots)_\beta^{(\Lambda)}$ denotes the Duhamel two-point function [10]. Since this inequality can be extended to [10] complex-valued functions $h_\mu(x)$, we have

$$\left(\sum_{x \in \Lambda} \sum_{\mu=1}^3 \Gamma^{(1)}(x) \overline{\partial_\mu h_\mu(x)}, \sum_{y \in \Lambda} \sum_{\mu'=1}^3 \Gamma^{(1)}(y) \partial_{\mu'} h_{\mu'}(y) \right)_\beta^{(\Lambda)} \leq \frac{1}{2\beta g} \sum_{x \in \Lambda} \sum_{\mu=1}^3 |h_\mu(x)|^2, \quad (8.1)$$

where $\overline{\dots}$ denotes its complex conjugate. We choose

$$h_\mu(x) = \frac{1}{\sqrt{|\Lambda|}} [e^{ip(x+e_\mu)} - e^{ipx}]$$

with the wavenumber $p = (p^{(1)}, p^{(2)}, p^{(3)})$, where we have written $px = p^{(1)}x^{(1)} + p^{(2)}x^{(2)} + p^{(3)}x^{(3)}$. Then, one has

$$\partial_\mu h_\mu(x) = -\frac{2}{\sqrt{|\Lambda|}} e^{ipx} (1 - \cos p^{(\mu)})$$

and

$$\sum_{x \in \Lambda} \sum_{\mu=1}^3 |h_\mu(x)|^2 = 2E_p,$$

where we have written

$$E_p := \sum_{\mu=1}^3 (1 - \cos p^{(\mu)}).$$

From these, we have

$$\sum_{x \in \Lambda} \sum_{\mu=1}^3 \Gamma^{(1)}(x) \partial_\mu h_\mu(x) = -2\hat{\Gamma}_p^{(1)} E_p,$$

where we have written

$$\hat{\Gamma}_p^{(1)} := \frac{1}{\sqrt{|\Lambda|}} \sum_{x \in \Lambda} \Gamma^{(1)}(x) e^{ipx}. \quad (8.2)$$

By substituting these into the above bound (8.1), we obtain

$$\mathfrak{b}_p := (\hat{\Gamma}_{-p}^{(1)}, \hat{\Gamma}_p^{(1)})_\beta^{(\Lambda)} \leq \frac{1}{4\beta g E_p}. \quad (8.3)$$

9. LONG-RANGE ORDER

Now, let us prove the existence of the long-range order for $\Gamma^{(1)}(x)$. We write

$$\mathfrak{g}_p := \frac{1}{2} [\langle \hat{\Gamma}_p^{(1)} \hat{\Gamma}_{-p}^{(1)} \rangle_\beta^{(\Lambda)} + \langle \hat{\Gamma}_{-p}^{(1)} \hat{\Gamma}_p^{(1)} \rangle_\beta^{(\Lambda)}]$$

and

$$\mathfrak{c}_p := \langle [\hat{\Gamma}_{-p}^{(1)}, [H^{(\Lambda)}(0), \hat{\Gamma}_p^{(1)}]] \rangle_\beta^{(\Lambda)} \quad (9.1)$$

for their thermal expectation values $\langle \dots \rangle_\beta^{(\Lambda)}$. We use the inequality, [10]

$$\mathfrak{g}_p \leq \frac{1}{2} \left[\mathfrak{b}_p + \sqrt{\mathfrak{b}_p^2 + \beta \mathfrak{b}_p \mathfrak{c}_p} \right] \leq \mathfrak{b}_p + \frac{1}{2} \sqrt{\beta \mathfrak{b}_p \mathfrak{c}_p}.$$

By using the bound (8.3) for \mathfrak{b}_p , we have

$$\frac{1}{2} [\langle \hat{\Gamma}_p^{(1)} \hat{\Gamma}_{-p}^{(1)} \rangle_\beta^{(\Lambda)} + \langle \hat{\Gamma}_{-p}^{(1)} \hat{\Gamma}_p^{(1)} \rangle_\beta^{(\Lambda)}] \leq \frac{1}{4\beta g E_p} + \frac{1}{4} \sqrt{\frac{\mathfrak{c}_p}{g E_p}}.$$

Further, following [19, 20], we have

$$\frac{1}{3|\Lambda|} \sum_p \langle \hat{\Gamma}_p^{(1)} \hat{\Gamma}_{-p}^{(1)} \rangle_\beta^{(\Lambda)} \left[\sum_{\mu=1}^3 \cos p^{(\mu)} \right] \leq \delta(\beta) + \frac{1}{12|\Lambda|} \sum_{p \neq 0} \sqrt{\frac{\mathfrak{c}_p}{g E_p}} \left\{ \sum_{\mu=1}^3 \cos p^{(\mu)} \right\}_+ + \Sigma_{\text{LRO}}^{(\Lambda)}, \quad (9.2)$$

where $\{a\}_+ = \max\{a, 0\}$, and we have written

$$\delta(\beta) := \frac{1}{12|\Lambda|} \sum_{p \neq 0} \frac{1}{\beta g E_p} \left\{ \sum_{\mu=1}^3 \cos p^{(\mu)} \right\}_+$$

and

$$\Sigma_{\text{LRO}}^{(\Lambda)} := \frac{1}{|\Lambda|} \langle \hat{\Gamma}_0^{(1)} \hat{\Gamma}_0^{(1)} \rangle_\beta^{(\Lambda)} = \frac{1}{|\Lambda|^2} \sum_{x, y \in \Lambda} \langle \Gamma^{(1)}(x) \Gamma^{(1)}(y) \rangle_\beta^{(\Lambda)}.$$

Clearly, the quantity $\delta(\beta)$ becomes small for a large β , and $\Sigma_{\text{LRO}}^{(\Lambda)}$ is nothing but the long-range order. Our goal is to show that the long-range order $\Sigma_{\text{LRO}}^{(\Lambda)}$ is strictly positive in the infinite-volume limit $\Lambda \nearrow \mathbb{Z}^3$.

For this purpose, let us first estimate the second term in the right-hand side of (9.2). By using the Schwarz inequality, one has

$$\frac{1}{|\Lambda|} \sum_{p \neq 0} \frac{1}{\sqrt{E_p}} \left\{ \sum_{\mu=1}^3 \cos p^{(\mu)} \right\}_+ \sqrt{\mathbf{c}_p} \leq \sqrt{3} I_3 \sqrt{\frac{1}{|\Lambda|} \sum_p \mathbf{c}_p}, \quad (9.3)$$

where we have written

$$I_3 := \sqrt{\frac{1}{3|\Lambda|} \sum_{p \neq 0} \frac{1}{E_p} \left(\left\{ \sum_{\mu=1}^3 \cos p^{(\mu)} \right\}_+ \right)^2}.$$

Let us estimate the right-hand side of (9.3). From the definition (9.1) of \mathbf{c}_p , one has

$$\begin{aligned} \mathbf{c}_p &= \langle [\hat{\Gamma}_{-p}^{(1)}, [H^{(\Lambda)}(0), \hat{\Gamma}_p^{(1)}]] \rangle_{\beta}^{(\Lambda)} \\ &= \langle [\hat{\Gamma}_{-p}^{(1)}, [H_K^{(\Lambda)}, \hat{\Gamma}_p^{(1)}]] \rangle_{\beta}^{(\Lambda)} + \langle [\hat{\Gamma}_{-p}^{(1)}, [H_{\text{int}}^{(\Lambda)}, \hat{\Gamma}_p^{(1)}]] \rangle_{\beta}^{(\Lambda)}. \end{aligned}$$

For the contribution of the kinetic Hamiltonian $H_K^{(\Lambda)}$ in the right-hand side, we have

$$\begin{aligned} \frac{1}{|\Lambda|} \sum_p \langle [\hat{\Gamma}_{-p}^{(1)}, [H_K^{(\Lambda)}, \hat{\Gamma}_p^{(1)}]] \rangle_{\beta}^{(\Lambda)} &= \frac{1}{|\Lambda|^2} \sum_p \sum_{x, y \in \Lambda} \langle [\Gamma^{(1)}(x), [H_K^{(\Lambda)}, \Gamma^{(1)}(y)]] \rangle_{\beta}^{(\Lambda)} e^{-ip(x-y)} \\ &= \frac{1}{|\Lambda|} \sum_{x \in \Lambda} \langle [\Gamma^{(1)}(x), [H_K^{(\Lambda)}, \Gamma^{(1)}(x)]] \rangle_{\beta}^{(\Lambda)}. \end{aligned}$$

Note that

$$\begin{aligned} &[H_K^{(\Lambda)}, \Gamma^{(1)}(x)] \\ &= i\kappa \sum_{\mu=1,3} \sum_{y=x, x-e_{\mu}} \{ [\Psi^{\dagger}(y) \alpha_{\mu} \Psi(y+e_{\mu}), \Gamma^{(1)}(x)] - [\Psi^{\dagger}(y+e_{\mu}) \alpha_{\mu} \Psi(y), \Gamma^{(1)}(x)] \} \\ &\quad - \kappa \sum_{y=x, x-e_2} \{ [\Psi^{\dagger}(y) \alpha_{\mu} \Psi(y+e_2), \Gamma^{(1)}(x)] + [\Psi^{\dagger}(y+e_2) \alpha_{\mu} \Psi(y), \Gamma^{(1)}(x)] \}. \end{aligned}$$

Clearly, this right-hand side is of order of 1. Therefore, we have

$$\left| \frac{1}{|\Lambda|} \sum_p \langle [\hat{\Gamma}_{-p}^{(1)}, [H_K^{(\Lambda)}, \hat{\Gamma}_p^{(1)}]] \rangle_{\beta}^{(\Lambda)} \right| \leq \mathcal{C}_0 |\kappa|,$$

where \mathcal{C}_0 is a positive constant.

The contribution for the interaction Hamiltonian $H_{\text{int}}^{(\Lambda)}$ is written

$$\begin{aligned}
 & \frac{1}{|\Lambda|} \sum_p \langle [\hat{\Gamma}_{-p}^{(1)}, [H_{\text{int}}^{(\Lambda)}, \hat{\Gamma}_p^{(1)}]] \rangle_\beta^{(\Lambda)} \\
 &= \frac{1}{|\Lambda|^2} \sum_p \sum_{x,y \in \Lambda} \langle [\Gamma^{(1)}(x), [H_{\text{int}}^{(\Lambda)}, \Gamma^{(1)}(y)]] \rangle_\beta^{(\Lambda)} e^{-ip(x-y)} \\
 &= \frac{1}{|\Lambda|} \sum_{x \in \Lambda} \langle [\Gamma^{(1)}(x), [H_{\text{int}}^{(\Lambda)}, \Gamma^{(1)}(x)]] \rangle_\beta^{(\Lambda)} \\
 &= -\frac{g}{|\Lambda|} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \{ \langle [\Gamma^{(1)}(x), [\Gamma^{(2)}(x), \Gamma^{(1)}(x)]] \Gamma^{(2)}(x + e_\mu) \rangle_\beta^{(\Lambda)} \\
 & \quad + \langle [\Gamma^{(1)}(x), [\Gamma^{(2)}(x), \Gamma^{(1)}(x)]] \Gamma^{(2)}(x - e_\mu) \rangle_\beta^{(\Lambda)} \} \\
 & \quad - \frac{g}{|\Lambda|} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \sum_{\ell=1}^3 \{ \langle [\Gamma^{(1)}(x), [S_5^{(\ell)}(x), \Gamma^{(1)}(x)]] S_5^{(\ell)}(x + e_\mu) \rangle_\beta^{(\Lambda)} \\
 & \quad + \langle [\Gamma^{(1)}(x), [S_5^{(\ell)}(x), \Gamma^{(1)}(x)]] S_5^{(\ell)}(x - e_\mu) \rangle_\beta^{(\Lambda)} \},
 \end{aligned}$$

where we have used the commutation relations, (A.4) in Appendix A for getting the third equality. Further, by using the commutation relations (A.3), (A.5), (A.6), and the chiral and the flavor rotational symmetries of the present Hamiltonian, we obtain

$$\begin{aligned}
 \frac{1}{|\Lambda|} \sum_p \langle [\hat{\Gamma}_{-p}^{(1)}, [H_{\text{int}}^{(\Lambda)}, \hat{\Gamma}_p^{(1)}]] \rangle_\beta^{(\Lambda)} &= \frac{8g}{|\Lambda|} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \langle \Gamma^{(2)}(x) \Gamma^{(2)}(x + e_\mu) \rangle_\beta^{(\Lambda)} \\
 & \quad + \frac{8g}{|\Lambda|} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \sum_{\ell=1}^3 \langle S_5^{(\ell)}(x) S_5^{(\ell)}(x + e_\mu) \rangle_\beta^{(\Lambda)} \\
 &= \frac{32g}{|\Lambda|} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \langle \Gamma^{(1)}(x) \Gamma^{(1)}(x + e_\mu) \rangle_\beta^{(\Lambda)}.
 \end{aligned}$$

Consequently, we obtain

$$\frac{1}{|\Lambda|} \sum_p \mathfrak{c}_p \leq \mathcal{C}_0 |\kappa| + 32g \mathcal{E}^{(\Lambda)}, \quad (9.4)$$

where we have written

$$\mathcal{E}^{(\Lambda)} := \frac{1}{|\Lambda|} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \langle \Gamma^{(1)}(x) \Gamma^{(1)}(x + e_\mu) \rangle_\beta^{(\Lambda)}. \quad (9.5)$$

As shown in Appendix B, this quantity $\mathcal{E}^{(\Lambda)}$ is positive at sufficiently low temperatures in the strong coupling regime $g/|\kappa| \gg 1$.

The left-hand side of (9.2) is written

$$\frac{1}{3|\Lambda|} \sum_p \langle \hat{\Gamma}_p^{(1)} \hat{\Gamma}_{-p}^{(1)} \rangle_\beta^{(\Lambda)} \left[\sum_{\mu=1}^3 \cos p^{(\mu)} \right] = \frac{1}{3|\Lambda|} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \langle \Gamma^{(1)}(x) \Gamma^{(1)}(x + e_\mu) \rangle_\beta^{(\Lambda)} = \frac{1}{3} \mathcal{E}^{(\Lambda)}$$

from (8.2). By combining this, (9.2), (9.3) and (9.4), we obtain

$$\begin{aligned} \frac{1}{3} \mathcal{E}^{(\Lambda)} &\leq \delta(\beta) + \frac{\sqrt{3}I_3}{12\sqrt{g}} \sqrt{\mathcal{C}_0|\kappa| + 32g\mathcal{E}^{(\Lambda)}} + \Sigma_{\text{LRO}}^{(\Lambda)} \\ &\leq \delta(\beta) + \frac{\sqrt{3}I_3}{12} \sqrt{\frac{\mathcal{C}_0|\kappa|}{g}} + \frac{\sqrt{6}}{3} I_3 \sqrt{\mathcal{E}^{(\Lambda)}} + \Sigma_{\text{LRO}}^{(\Lambda)}. \end{aligned}$$

Therefore, the lower bound for the long-range order $\Sigma_{\text{LRO}}^{(\Lambda)}$ is given by

$$\frac{1}{3} \sqrt{\mathcal{E}^{(\Lambda)}} \left[\sqrt{\mathcal{E}^{(\Lambda)}} - \sqrt{6}I_3 \right] - \delta(\beta) - \frac{\sqrt{3}I_3}{12} \sqrt{\frac{\mathcal{C}_0|\kappa|}{g}} \leq \Sigma_{\text{LRO}}^{(\Lambda)}.$$

The lower bound for $\mathcal{E}^{(\Lambda)}$ is given by

$$\mathcal{E}^{(\Lambda)} \geq 6 - \frac{3|\kappa|}{g} - \frac{1}{\beta g} \log 2$$

as shown in Appendix B. Since $I_3 = 0.68 \dots$, these imply the existence of the long-range order at sufficiently low temperatures in the strong coupling regime $|\kappa|/g \ll 1$.

APPENDIX A. CHIRAL ROTATIONAL SYMMETRY

One can check that, for any two 4×4 matrices, \mathcal{M} and \mathcal{M}' , the following relation holds:

$$[\Psi^\dagger(x) \mathcal{M} \Psi(x), \Psi^\dagger(x) \mathcal{M}' \Psi(x)] = \Psi^\dagger(x) [\mathcal{M}, \mathcal{M}'] \Psi(x). \quad (\text{A.1})$$

We write

$$\Gamma^{(5)}(x) := \Psi^\dagger(x) \gamma_5 \Psi(x). \quad (\text{A.2})$$

Then, by using the above formula, one has the commutation relations,

$$\begin{aligned} [\Gamma^{(1)}(x), \Gamma^{(2)}(x)] &= 2i\Gamma^{(5)}(x), \quad [\Gamma^{(2)}(x), \Gamma^{(5)}(x)] = 2i\Gamma^{(1)}(x), \\ \text{and } [\Gamma^{(5)}(x), \Gamma^{(1)}(x)] &= 2i\Gamma^{(2)}(x). \end{aligned} \quad (\text{A.3})$$

In addition, from the definitions of $S^{(j)}(x)$ for $j = 1, 2, 3$, we have

$$[\Gamma^{(1)}(x), S^{(j)}(x)] = 0 \quad \text{for } j = 1, 2, 3. \quad (\text{A.4})$$

Moreover, one has

$$[\Gamma^{(1)}(x), S_5^{(\ell)}(x)] = 2i\Psi^\dagger(x) \gamma_5 \tau_\ell \Psi(x) \quad (\text{A.5})$$

and

$$[\Gamma^{(1)}(x), \Psi^\dagger(x) \gamma_5 \tau_\ell \Psi(x)] = -2iS_5^{(\ell)}(x). \quad (\text{A.6})$$

for $\ell = 1, 2, 3$.

Next consider the rotation by the generator $\Gamma^{(5)}(x)$. Let $\theta \in \mathbb{R}$, and note that

$$\begin{aligned} \frac{d}{d\theta} e^{i\theta\Gamma^{(5)}(x)}\Gamma^{(1)}(x)e^{-i\theta\Gamma^{(5)}(x)} &= ie^{i\theta\Gamma^{(5)}(x)}[\Gamma^{(5)}(x), \Gamma^{(1)}(x)]e^{-i\theta\Gamma^{(5)}(x)} \\ &= -2e^{i\theta\Gamma^{(5)}(x)}\Gamma^{(2)}(x)e^{-i\theta\Gamma^{(5)}(x)}, \end{aligned}$$

where we have used the above commutation relation $[\Gamma^{(5)}(x), \Gamma^{(1)}(x)] = 2i\Gamma^{(2)}(x)$. Further,

$$\begin{aligned} \frac{d^2}{d\theta^2} e^{i\theta\Gamma^{(5)}(x)}\Gamma^{(1)}(x)e^{-i\theta\Gamma^{(5)}(x)} &= -2ie^{i\theta\Gamma^{(5)}(x)}[\Gamma^{(5)}(x), \Gamma^{(2)}(x)]e^{-i\theta\Gamma^{(5)}(x)} \\ &= -4e^{i\theta\Gamma^{(5)}(x)}\Gamma^{(1)}(x)e^{-i\theta\Gamma^{(5)}(x)}. \end{aligned}$$

These imply

$$e^{i\theta\Gamma^{(5)}(x)}\Gamma^{(1)}(x)e^{-i\theta\Gamma^{(5)}(x)} = \Gamma^{(1)}(x) \cos 2\theta - \Gamma^{(2)}(x) \sin 2\theta.$$

In particular, for $\theta = -\pi/4$, the following holds:

$$e^{-i(\pi/4)\Gamma^{(5)}(x)}\Gamma^{(1)}(x)e^{i(\pi/4)\Gamma^{(5)}(x)} = \Gamma^{(2)}(x). \quad (\text{A.7})$$

APPENDIX B. A LOWER BOUND FOR $\mathcal{E}^{(\Lambda)}$

In this Appendix, we obtain a lower bound for $\mathcal{E}^{(\Lambda)}$ of (9.5).

In the same way as in [14], we first use Peierls's inequality. Namely, we have

$$Z_\beta^{(\Lambda)} := \text{Tr} \exp[-\beta H^{(\Lambda)}(0)] \geq \exp[-\beta \langle \Omega_{\text{var}}^{(\Lambda)}, H^{(\Lambda)}(0) \Omega_{\text{var}}^{(\Lambda)} \rangle] \quad (\text{B.1})$$

for any variational wavefunction $\Omega_{\text{var}}^{(\Lambda)}$. We recall the expression of the operator $\Gamma^{(1)}(x)$, i.e.,

$$\Gamma^{(1)}(x) = \Psi_{\text{u}}^\dagger(x)\gamma_0\Psi_{\text{u}}(x) + \Psi_{\text{d}}^\dagger(x)\gamma_0\Psi_{\text{d}}(x).$$

Since the matrix γ_0 in $\Gamma^{(1)}(x)$ has the four eigenvalues, $+1, +1, -1, -1$, we choose the variational state $\Omega_{\text{var}}^{(\Lambda)}$ so that it is the eigenstate of $\Gamma^{(1)}(x)$ for both of the two flavors, u and d, and all the sites $x \in \Lambda$ with the eigenvalue $+4$, i.e., $\Gamma^{(1)}(x)\Omega_{\text{var}}^{(\Lambda)} = 4\Omega_{\text{var}}^{(\Lambda)}$. In other words, each site x is occupied by the two eigenstates of γ_0 with the eigenvalue $+1$ for both of the two flavors, u and d. Then, clearly, one has

$$\langle \Omega_{\text{var}}^{(\Lambda)}, H_{\text{K}}^{(\Lambda)} \Omega_{\text{var}}^{(\Lambda)} \rangle = 0$$

for the kinetic Hamiltonian $H_{\text{K}}^{(\Lambda)}$. Further, we have

$$\langle \Omega_{\text{var}}^{(\Lambda)}, \Gamma^{(2)}(x) \Omega_{\text{var}}^{(\Lambda)} \rangle = 0$$

for all $x \in \Lambda$ because the matrix $\gamma_0\gamma_5$ in $\Gamma^{(2)}(x)$ satisfies the commutation relation $\gamma_0\gamma_0\gamma_5 + \gamma_0\gamma_5\gamma_0 = 0$, which yields $\langle v, \gamma_0\gamma_5 v \rangle + \langle v, \gamma_0\gamma_5 v \rangle = 2\langle v, \gamma_0\gamma_5 v \rangle = 0$ for any eigenvector v of γ_0 . Therefore, except for the term having the operator $\Gamma^{(1)}(x)$, all the expectation values of the terms in the Hamiltonian are vanishing. From these observations, one has

$$\langle \Omega_{\text{var}}^{(\Lambda)}, H^{(\Lambda)}(0) \Omega_{\text{var}}^{(\Lambda)} \rangle = -16 \cdot 3g|\Lambda|.$$

By substituting this into the right-hand side of (B.1), we have

$$\log Z_\beta^{(\Lambda)} \geq 16 \cdot 3\beta g |\Lambda|. \quad (\text{B.2})$$

On the other hand, the following holds:

$$\log Z_\beta^{(\Lambda)} = -\beta \langle H^{(\Lambda)}(0) \rangle_\beta^{(\Lambda)} + S_\beta^{(\Lambda)} \leq -\beta \langle H^{(\Lambda)}(0) \rangle_\beta^{(\Lambda)} + \log 2^{8|\Lambda|},$$

where $S_\beta^{(\Lambda)}$ is the entropy, and we have used its upper bound. By combining this with the above inequality (B.2), one has

$$-\frac{1}{|\Lambda|} \langle H^{(\Lambda)}(0) \rangle_\beta^{(\Lambda)} \geq 16 \cdot 3g - \frac{8}{\beta} \log 2.$$

The right-hand side is written

$$\begin{aligned} -\frac{1}{|\Lambda|} \langle H^{(\Lambda)}(0) \rangle_\beta^{(\Lambda)} &= -\frac{1}{|\Lambda|} \langle H_K^{(\Lambda)} \rangle_\beta^{(\Lambda)} + \frac{8g}{|\Lambda|} \sum_{x \in \Lambda} \sum_{\mu=1}^3 \langle \Gamma^{(1)}(x) \Gamma^{(1)}(x + e_\mu) \rangle_\beta^{(\Lambda)} \\ &\leq 8 \cdot 3|\kappa| + 8g\mathcal{E}^{(\Lambda)}, \end{aligned}$$

where we have used the rotational symmetries of the Hamiltonian $H^{(\Lambda)}(0)$. Therefore, we obtain the desired lower bound,

$$\mathcal{E}^{(\Lambda)} \geq 6 - \frac{3|\kappa|}{g} - \frac{1}{\beta g} \log 2. \quad (\text{B.3})$$

Acknowledgements. We would like to thank Sinya Aoki for valuable discussions. The present paper is an answer to his question whether the methods developed in [13, 14] for the Nambu–Jona-Lasinio models are applicable to the mathematical proof of the parity violation which is conjectured in the papers [5, 6]. We are also grateful to Etsuko Itou for her helpful comments on the presentation of the paper. Y. G. is partially supported by JSPS Kakenhi Grant Number 23K12989.

REFERENCES

- [1] S. Aoki, New phase structure for lattice QCD with Wilson fermions, *Phys. Rev. D* **30**, 2653 (1984).
- [2] S. Aoki, Solution to the U(1) problem on a lattice, *Phys. Rev. Lett.* **57**, 3136 (1986).
- [3] S. Aoki, U(1) problem and lattice QCD, *Nucl. Phys. B* **314**, 79–111 (1989).
- [4] S. Aoki, On the Phase Structure of QCD with Wilson Fermions *Prog. Theor. Phys.* **122**, 179 (1996).
- [5] S. Aoki, S. Boettcher and A. Gocksch, Spontaneous breaking of flavor symmetry and parity in the Nambu–Jona-Lasinio model with Wilson fermions, *Phys. Lett. B* **331** 157–164 (1994).
- [6] S. Aoki and A. Gocksch, Spontaneous breaking of flavor symmetry and parity in lattice QCD with Wilson fermions, *Phys. Rev. D* **45**, 3845–3853 (1992).
- [7] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 1*, Springer-Verlag, New York, Berlin Heidelberg, 2nd ed. (1987).
- [8] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 2*, Springer-Verlag, New York, Berlin Heidelberg, 2nd ed. (1997).

- [9] H. Cornean, H. Garde, A. Jensen, Discrete approximations to Dirac operators and norm resolvent convergence, *J. Spectr. Theory* **12**, 1589–1622 (2022).
- [10] F. J. Dyson, E. H. Lieb and B. Simon, Phase Transitions in Quantum Spin Systems with Isotropic and Nonisotropic Interactions, *J. Stat. Phys.* **13**, 335–383 (1978).
- [11] J. Fröhlich, R. B. Israel, E. H. Lieb and B. Simon, Phase Transitions and Reflection Positivity. II. Lattice Systems with Short-Range and Coulomb Interactions, *J. Stat. Phys.* **22**, 297–347 (1980).
- [12] J. Fröhlich, B. Simon, and T. Spencer, Infrared bounds, phase transitions and continuous symmetry breaking, *Commun. Math. Phys.* **50** 79–95 (1976).
- [13] Y. Goto and T. Koma, Spontaneous Mass Generation and Chiral Symmetry Breaking in a Lattice Nambu–Jona-Lasinio Model, *Commun. Math. Phys.* **404(3)**, 1463–1493 (2023), arXiv:2209.06031.
- [14] Y. Goto and T. Koma, Nambu-Goldstone Modes in a Lattice Nambu-Jona-Lasinio Model with Multi Flavor Symmetries, *J. Math. Phys.* **65**, 102301 (2024), arXiv:2310.15922.
- [15] T. Hatsuda, T. Kunihiro, QCD Phenomenology based on a Chiral Effective Lagrangian, *Phys. Rept.* **247**, 221–367 (1994), arXiv:hep-ph/9401310.
- [16] P. Horsch and W. von der Linden, Spin-Correlations and Low Lying Excited States of the Spin-1/2 Heisenberg Antiferromagnet on a Square Lattice, *Z. Phys. B* **72**, 181–193 (1988).
- [17] A. Jaffe and F. L. Pedrocchi, Reflection Positivity for Majoranas, *Ann. Henri Poincaré*, **16(1)**, 189–203 (2015).
- [18] T. A. Kaplan, P. Horsch, and W. von der Linden, Order parameter in quantum antiferromagnets, *J. Phys. Soc. Jpn.* **58**, 3894–3898 (1989).
- [19] T. Kennedy, E. H. Lieb, and B. S. Shastry, Existence of Néel order in some spin-1/2 Heisenberg antiferromagnets, *J. Stat. Phys.* **53**, **1019** (1988).
- [20] T. Kennedy, E. H. Lieb, and B. S. Shastry, The XY Model Has Long-Range Order for All Spins and All Dimensions Greater than One, *Phys. Rev. Lett.* **61**, 2582 (1988).
- [21] T. Koma, H. Tasaki, Symmetry breaking in Heisenberg antiferromagnets, *Commun. Math. Phys.* **158**, 191–214 (1993).
- [22] T. Koma, H. Tasaki, Symmetry Breaking and Finite Size Effects in Quantum Many-Body Systems, *J. Stat. Phys.* **76**, 745–803 (1994).
- [23] S. Nakamura, Remarks on discrete Dirac operators and their continuum limits, *J. Spectr. Theory* **14** 255–269 (2024).
- [24] Y. Nambu, Axial Vector Current Conservation in Weak Interactions, *Phys. Rev. Lett.* **4** 380–382 (1960).
- [25] Y. Nambu, and G. Jona-Lasinio, Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. I, *Phys. Rev.* **122** 345–358 (1961).
- [26] Y. Nambu, and G. Jona-Lasinio, Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. II, *Phys. Rev.* **124**, 246–254 (1961).
- [27] H.B. Nielsen, M. Ninomiya, Absence of neutrinos on a lattice: (I). Proof by homotopy theory, *Nucl. Phys. B*, **185**, 20–40 (1981); Erratum: *B* **195** 541–542 (1982).
- [28] H.B. Nielsen, M. Ninomiya, Absence of neutrinos on a lattice: (II). Intuitive topological proof, *Nucl. Phys. B*, **193**, 173–194 (1981).
- [29] K. Osterwalder, E. Seiler, Gauge Field Theories on the Lattice, *Ann. Phys.* , **110** 440–471 (1978).
- [30] K. Osterwalder, R. Schrader, Axioms for Euclidean Green’s functions, *Commun. Math. Phys.* **31** 83–112 (1973).
- [31] C. Vafa and E. Witten, Restrictions on symmetry breaking in vector-like gauge theories, *Nucl. Phys. B* **234**, 173–188 (1984).