# Phase transitions in low-dimensional long-range random field Ising models

Jian Ding<sup>1</sup>, Fenglin Huang<sup>1</sup>, João Maia<sup>2</sup>

<sup>1</sup> School of Mathematical Sciences, Peking University, China <sup>2</sup> Beijing International Center for Mathematical Research, Peking University, China

#### **Abstract**

We consider the long-range random field Ising model in dimension  $d = 1, 2$ , whereas the long-range interaction is of the form  $J_{xy} = |x - y|^{-\alpha}$  with  $1 < \alpha < 3/2$  for  $d = 1$  and with  $2 < \alpha \leq 3$  for  $d = 2$ . Our main results establish phase transitions in these regimes. In one dimension, we employ a Peierls argument with some novel modification, suitable for dealing with the randomness coming from the external field; in two dimensions, our proof follows that of Affonso, Bissacot, and Maia (2023) with some adaptations, but new ideas are required in the critical case of  $\alpha = 3$ .

## **1 Introduction**

The Ising model is one of the most studied models in statistical mechanics. In this paper, we work with the combination of two variations of the Ising model, the random field and the long-range models. The long-range variation, introduced first by Kac and Thompson [\[23\]](#page-39-0), considers a model on the lattice  $\mathbb{Z}^d$  with the same configuration space  $\Omega := \{-1, +1\}^{\mathbb{Z}^d}$  (as in the nearest-neighbor case), but with a long-range interaction  ${J_{xy}}_{x,y \in \mathbb{Z}^d}$  given by

$$
J_{xy} = \begin{cases} \frac{1}{|x-y|^{\alpha}} & \text{if } x \neq y, \\ 0 & \text{otherwise,} \end{cases}
$$
 (1.1)

with  $\alpha > d$  ensuring that the model is regular. A second variation, the random field Ising model (RFIM), is the usual Ising model with the addition of a random external field. Our model of interest is the long-range random field Ising model, which has a *local Hamiltonian* given by

$$
H_{\Lambda,\varepsilon h}^{\eta}(\sigma) := -\sum_{x,y \in \Lambda} J_{xy} \sigma_x \sigma_y - \sum_{x \in \Lambda, y \in \Lambda^c} J_{xy} \sigma_x \eta_y - \sum_{x \in \Lambda} \varepsilon h_x \sigma_x, \tag{1.2}
$$

where  $\Lambda \subset \mathbb{Z}^d$  is finite,  $\eta \in \Omega$  is the *boundary condition*,  $\{h_x\}_{x \in \mathbb{Z}^d}$  is a collection of i.i.d. Gaussian random variables in a probability space denoted as  $(\tilde{\Omega}, \mathcal{A}, \mathbb{P})$  and the parameter  $\varepsilon > 0$  controls the variance of the external field. As in the usual Ising model, the local Gibbs measure is a probability measure given by

$$
\mu_{\Lambda;\beta,\varepsilon h}^\eta(\sigma) \coloneqq \mathbbm{1}_{\Omega_\Lambda^\eta}(\sigma) \frac{e^{-\beta H_{\Lambda,\varepsilon h}^\eta(\sigma)}}{Z_{\Lambda;\beta,\varepsilon}^\eta(h)},
$$

where  $Z_{\Lambda;\beta,\varepsilon}^{\eta}(h)$  is a normalizing constant and  $\Omega_{\Lambda}^{\eta} := \{ \omega \in \Omega : \omega_x = \eta_x, \forall x \in \Lambda^c \}$  is the collection of the configurations matching *η* outside Λ. For the extremal boundary conditions  $\eta \equiv +1$  ( $\eta_x$ )  $+1, \forall x \in \mathbb{Z}^d$ , and analogously,  $\eta \equiv -1$ , the weak<sup>\*</sup>-limits

$$
\mu_{\beta,\varepsilon h}^{\pm}[\sigma] \coloneqq \lim_{N \to \infty} \mu_{[-N,N]^d; \beta, \varepsilon h}^{\pm}[\sigma],
$$

are well-defined (see [\[10,](#page-38-0) Theorem 7.2.2]) and are our main objects of study. When the limiting measures are distinct, i.e.,  $\mu_{\beta, \epsilon h}^+ \neq \mu_{\beta, \epsilon h}^-$  P-a.s. for some  $\beta > 0$ , we say there is *phase transition*, also called *long-range order*.

Both the long-range model and the RFIM have a rich history in statistical physics. Determining for which dimensions *d* the RFIM presents a phase transition was a topic of heated debate in the physics community until it was settled in the seminal works of Aizenmann and Wehr [\[7\]](#page-38-1) and Bricmont and Kupiainen [\[11\]](#page-38-2). Aizenmann and Wehr showed that the model has only one Gibbs measure in dimensions  $d = 1, 2$ , while Bricmont and Kupiainen proved phase transition for  $d \geq 3$ . Several interesting phenomena occur in the RFIM, and interesting results were obtained in recent years: we mention for example  $[5, 6, 13, 14, 16, 17]$  $[5, 6, 13, 14, 16, 17]$  $[5, 6, 13, 14, 16, 17]$  $[5, 6, 13, 14, 16, 17]$  $[5, 6, 13, 14, 16, 17]$  $[5, 6, 13, 14, 16, 17]$  for results in two dimensions, and  $[15, 18]$  $[15, 18]$  for three and higher dimensions.

For the long-range model in one dimension, Dobrushin and Ruelle proved that there is no phase transition if  $\alpha > 2$  [\[19,](#page-39-1) [24\]](#page-39-2). On the other hand, Kac and Thompson [\[23\]](#page-39-0) conjectured that the longrange interaction should induce a phase transition, different from the Ising model. This was proved by Dyson [\[20\]](#page-39-3) for 1 *< α <* 2; for this reason, the model is also known as the *Dyson model*. For the critical exponent  $\alpha = 2$ , phase transition was proved in the seminal work of Fröhlich and Spencer [\[22\]](#page-39-4), where the authors introduced a notion of long-range one-dimensional contours. This contour system was extended to the multidimensional setting by Affonso, Bissacot, Endo and Handa [\[2\]](#page-38-11), where the authors proved phase transition at low temperatures, even in the presence of a decaying field.

For the long-range random field Ising model, very little was known until very recently. In low dimensions, the same work of Aizenman and Wehr [\[7\]](#page-38-1) proved uniqueness for  $d = 1, 2$  and  $3d/2 < \alpha$ , and in one dimension they also proved the uniqueness for the critical case of  $\alpha = 3/2$ . Phase transition for the nearest-neighbor Ising model with no external field implies phase transition for the long-range model, again with no external field, by standard correlation inequalities. However, a different argument is required for models with a random field, since some correlation inequalities are lost in this setting. A recent work of Affonso, Bissacot, and Maia [\[3\]](#page-38-12) proved phase transition for dimensions  $d \geq 3$  in the entire region  $\alpha > d$ . In one dimension, the only known result prior to this work was proved by Cassandro, Orlandi, and Picco [\[12\]](#page-38-13), with the extra restrictions that  $3 - \frac{\ln 3}{\ln 2} < \alpha < 3/2$  and  $J(1)$  large enough, where  $J(1)$  is the nearest-neighbor interaction.

A natural follow-up question is if the restriction of  $J(1)$  being large can be lifted since this is not required in the long-range model with no field. Another question of interest, as pointed out by [\[4\]](#page-38-14), is if  $\alpha = 3$  is indeed the critical exponent for phase transition in two dimensions in the presence of a random field. In this paper we answer both these questions, completing the picture of phase transition.

For the long-range random field Ising model, we prove phase transition in one dimension for  $1 < \alpha < 3/2$ , without the assumption of  $J(1)$  being large. We also show that there is a phase transition in two dimensions for all  $2 < \alpha \leq 3$ , including the critical exponent  $\alpha = 3$ . Notice that this is distinct from the one-dimensional critical case  $\alpha = 3/2$ , where there is uniqueness at all temperatures. Our results can be summarized in the next two theorems.

<span id="page-1-0"></span>**Theorem 1.1.** Fix  $d = 1$  and  $1 < \alpha < \frac{3}{2}$ . For any constant  $c_1 > 0$ , there exist constants  $\varepsilon_1 := \varepsilon_1(\alpha, c_1) > 0$  and  $\beta_1 := \beta_1(\alpha, c_1) > 0$  such that for any  $\varepsilon < \varepsilon_1$  and any  $\beta > \beta_1$ , the following *holds with*  $\mathbb{P}$ *-probability at least*  $1 - c_1$ *:* 

$$
\mu_{\beta,\varepsilon h}^+(\sigma_0=-1)
$$

<span id="page-2-0"></span>**Theorem 1.2.** Fix  $d = 2$  and  $2 < \alpha \leq 3$ . For any constant  $c_2 > 0$ , there exist constants  $\varepsilon_2 := \varepsilon_2(\alpha, c_2) > 0$  and  $\beta_2 := \beta_2(\alpha, c_2) > 0$  such that for any  $\varepsilon < \varepsilon_2$  and any  $\beta > \beta_2$ , the following *holds with*  $\mathbb{P}$ *-probability at least*  $1 - c_2$ *:* 

$$
\mu_{\beta,\varepsilon h}^+(\sigma_0=-1)
$$

Both previous works on the long-range RFIM employed a notion of contour suitable for the long-range Ising model with no field and then did the necessary adaptations to control the external field. Despite the notions of contours in [\[12\]](#page-38-13) and [\[3\]](#page-38-12) being different, both of them were inspired by the contours of Fröhlich and Spencer [\[22\]](#page-39-4).

In one dimension, it does not seem possible to use the same contours as in [\[12\]](#page-38-13) and remove the condition on  $J(1)$ . Removing the condition on  $J(1)$  was achieved in [\[9\]](#page-38-15) for the model with no field, but an ever stronger restriction on  $\alpha$  where  $3/2 < \alpha^* < \alpha < 2$  is required. On the other hand, one could try to use the same notion of contour as in [\[22\]](#page-39-4), since they were recently extended to the entire region  $1 < \alpha \leq 2$  in [\[1\]](#page-37-0), again for the model with no random field. This however, would require extending the entropy arguments of [\[3\]](#page-38-12) to one dimension, which seems possible but not trivial, and then applying our improved coarse-graining procedure as presented in Section [3.](#page-26-0)

We opt to use a different approach, introducing a new Peierls-type argument, tailored for longrange interactions with random perturbations. Our construction of the Peierls map resembles that of [\[22\]](#page-39-4), and it is also done in a multiscale fashion. A noticeable difference is that we do not rely on spin flips, only on the local densities of the plus and minus spins in the configuration. Using our construction, the entropy arguments are simpler, and a coarse-graining argument similar to that in [\[21\]](#page-39-5) is enough to control the field. In addition, we can also show phase transition for  $\alpha \in (1,2)$ when there is no external field, which gives an alternative proof to [\[1\]](#page-37-0) (see Remark [2.22\)](#page-15-0).

#### **1.1 Proof strategy in one dimension**

Our argument in one dimension follows the general principle of a Peierls argument: denoting by  $\Lambda \in \mathbb{Z}$  a finite subset of  $\mathbb{Z}$ , for each configuration  $\sigma \in \Omega^+ := \cup_{\Lambda \in \mathbb{Z}} \Omega^+_{\Lambda}$  with  $\sigma_0 = -1$ , we map  $\sigma$  to a configuration  $\omega$  with  $\omega_0 = 1$  and compare their energies. Our first intuition for the Peierls map is to flip all the minuses in a suitable interval containing the origin. However, two difficulties arise:

- 1. Spins now interact with all the others, so it is hard to define what is a suitable interval.
- 2. If we flip all the minuses in an interval, the minus spins surrounded by many pluses result in a huge entropy that we can not control.

To overcome the difficulties above, especially the second one, in Section [2.1](#page-4-0) we define a balancing procedure. The intuition is that minus spins surrounded by a vast majority of plus spins are, in a sense, 'isolated', so they should be treated as plus spins and vice-versa. We will continue this procedure until no such spin appears. In the end, we will flip all the minus spins left in a suitable interval to gain energy for which we can control the entropy.

In Lemma [2.29,](#page-17-0) we establish the key energy bound, which essentially states that for a configuration without 'isolated' spins, there will be a significant number of plus spins as long as the configuration contains at least one plus spin. Lemma [2.28](#page-16-0) plays a crucial role in the proof of Lemma [2.29,](#page-17-0) and its validity heavily depends on the one-dimensional structure.

Then in Proposition [2.20](#page-13-0) we provide the necessary entropy estimations to do a Peirls argument. To control the random field we use the approach of [\[18\]](#page-38-10), so phase transition follows after we control the probability of some bad events (see [\(2.15\)](#page-13-1)). This is done in Proposition [2.18,](#page-13-2) using the coarsegraining argument of [\[21\]](#page-39-5) together with the previously established entropy bounds.

### **1.2 Proof strategy in two dimensions**

The argument of [\[3\]](#page-38-12) applies the strategy of [\[18\]](#page-38-10) to the long-range model, and the main complication is to extend the coarse-graining argument of [\[21\]](#page-39-5) to long-range contours, which are in general not connected. Their approach is robust, which allows us to consider the same contour system in two dimensions. For  $2 < \alpha < 3$ , a simple modification of their argument suffices to show phase transition. Such adaption, however, does not suffice to prove the critical case  $\alpha = 3$ . A rough intuition underlying the existence of phase transition is the following Imry-Ma argument: the contribution of the external field from any  $A \subset \mathbb{Z}^2$  should be of order  $\sqrt{|A|}$ , so there is phase transition as long as the energy contribution of flipping the spins in larger than  $\sqrt{|A|}$ . For the box  $A = [-N, N]^2$ , this contribution is at least  $N^{4-\alpha}$ , which beats  $\sqrt{|A|} = N$  only when  $\alpha < 3$ . However, when  $\alpha = 3$ the interaction between spins inside and outside A is actually of order  $N \log(N)$ , see [\[8\]](#page-38-16). So, for the critical case, we must introduce some improved energy estimations, which capture the correct order of the interaction, see Lemma [3.13.](#page-34-0)

Even with this improvement, the same coarse-graining argument does not hold. In order to compare the entropy bound with the energy bound we need to introduce a finer coarse-graining procedure. Instead of directly using the entropy bound of [\[3\]](#page-38-12), we give our entropy bound according to its coarse-graining level. This improvement allows us to apply a union bound over all levels and control the enumeration coming from each level.

The structure of the paper is as follows. Section [2](#page-3-0) contains the proof of Theorem [1.1.](#page-1-0) In Section [2.1](#page-4-0) we define our Peierls map in one dimension. In Section [2.2,](#page-6-0) we analyze the Peierls map and in Section [2.2.1](#page-12-0) we present the outline of our argument and prove Theorem [1.1](#page-1-0) assuming some required estimations on the energy and entropy. Sections [2.3](#page-15-1) and [2.4](#page-24-0) are devoted to proving such estimations. Section [3](#page-26-0) contains the proof of Theorem [1.2,](#page-2-0) and it is rather independent from Section [2.](#page-3-0) In Section [3.1,](#page-29-0) we prove the case that  $2 < \alpha < 3$  and in Section [3.2](#page-31-0) we prove the critical case  $\alpha = 3$ .

## <span id="page-3-0"></span>**2 Phase Transition in 1d**

In this section, we focus on the case  $d = 1$  and prove Theorem [1.1.](#page-1-0) In Section [2.1,](#page-4-0) we construct the Peierls map, whereas a key ingredient is a balancing procedure; in Section [2.2,](#page-6-0) we analyze the Peierls map and prove some properties of the balancing procedure, and in Section [2.2.1](#page-12-0) we prove Theorem [1.1](#page-1-0) assuming some suitable bounds on the energy and the entropy; in Sections [2.3](#page-15-1) and [2.4,](#page-24-0) we provide the aforementioned bounds on the energy and the entropy respectively.

Throughout this section, we fix  $d = 1, 1 < \alpha < \frac{3}{2}$  and  $\Lambda \in \mathbb{Z}$  containing the origin. We also fix the following constants:  $c_1 \geq 10$  an arbitrary constant,  $\delta := \min\{0.001, \frac{1.5 - \alpha}{20}\} > 0$ , and  $M_0 :=$  $M_0(\alpha, c_1) > 2$  large enough, whose value will be chosen later. We reserve notations  $C_1, C_2, \ldots$ for arbitrary constants, possibly taking different values in different proofs, but depending only on *M*<sub>0</sub>, δ, or *α*. In addition, we use  $c_1$ ,  $c_2$ , ... and  $\overline{c}_1$ ,  $\overline{c}_2$ , ... for constants whose values are fixed (throughout the paper) upon their first occurrences.

### <span id="page-4-0"></span>**2.1 The construction of the Peierls map**

Given a configuration  $\sigma \in \Omega_{\Lambda}^+$  such that  $\sigma_0 = -1$ , we aim to determine a set of sites  $A \subset \mathbb{Z}$ such that our Peierls map will obtain  $\tau_A(\sigma)$  from  $\sigma$  by flipping spins in *A*. Such a set has to be carefully chosen so that on the one hand we can compare the energy change from  $\sigma$  to  $\tau_A(\sigma)$ , and on the other hand, the multiplicity of the Peierls map (i.e., the entropy loss) can be controlled. To this end, we will employ a balancing procedure (see Definition [2.4](#page-5-0) below).

As a notation convention, when writing an interval  $[a, b)$  we always mean its restriction to  $\mathbb{Z}$ , [ $a, b$ ) ∩ Z. At each scale  $\ell \geq 0$ , we consider intervals  $I_{\ell}(x) = [2^{\ell-4}x - 2^{\ell-1}, 2^{\ell-4}x + 2^{\ell-1}),$  with  $x \in \mathbb{Z}$  and side length  $2^{\ell}$ . Each such interval will be called an *ℓ*-interval, and we write I<sub>ℓ</sub> (i.e., omitting the center *x*) to denote an arbitrary  $\ell$ -interval. The family of all such intervals is denoted as  $\mathcal{I} := \{I_{\ell}(x)\}_{x \in \mathbb{Z}, \ell \geq 0}$ , and the restriction to  $\ell$ -intervals is  $\mathcal{I}_{\ell} := \{I_{\ell}(x)\}_{x \in \mathbb{Z}}$ . We can further split each  $\mathcal{I}_{\ell}$  into 16 sub-collections

<span id="page-4-2"></span>
$$
\mathcal{I}_{\ell}^{i} \coloneqq \{ [2^{\ell-4}(16x+i), 2^{\ell-4}(16x+16+i)) \}_{x \in \mathbb{Z}}, \ i = 0, 1, \dots 15,
$$
\n
$$
(2.1)
$$

whereas each sub-collection forms a partition of Z.

For any configuration  $\sigma \in \Omega$  and any interval I, we split I into its positive and negative sites  $I^{\pm}(\sigma) := \{x \in I : \sigma_x = \pm 1\}.$  We say that  $I_{\ell}$  is *minus dense* with respect to a configuration  $\sigma$ when  $|I_{\ell}^{-}(\sigma)| > 2^{\ell} \left(1 - \frac{1}{M_{\ell}}\right)$ ), where  $M_{\ell} := M_0 2^{\delta \ell}$ . When the minus density satisfies  $|\mathcal{I}_{\ell}(\sigma)| \leq \frac{2^{\ell}}{M_{\ell}}$  $\frac{2^{\infty}}{M_{\ell}}$ there are very few minuses in the interval, so we say I*<sup>ℓ</sup>* is *minus vacant*. If the density satisfies  $|I_{\ell}^-(\sigma)| > \frac{2^{\ell}}{M_{\ell}}$  $\frac{2^k}{M_\ell}$ , we say  $I_\ell$  is *minus occupied*. We remark that a minus dense interval is also a minus occupied interval. In a completely analogous fashion, we define the *plus dense*, *plus vacant*, and *plus occupied* intervals. Notice that the notions of occupied are not exclusionary, that is, an interval  $I_{\ell}$  can be both minus occupied and plus occupied since  $\frac{1}{M_{\ell}} \leq \frac{1}{2}$ . However, the dense and vacant properties are complementary: a plus vacant interval has minus density larger than  $(1 - \frac{1}{M_{\ell}})$ , and it is therefore minus dense.

In the Peierls argument, if  $I_\ell \ni 0$  is a minus dense interval, the natural attempt is to compare  $\sigma$ with a configuration after we flip all minus spins in I*ℓ*. The long-range interaction, however, makes this very costly since there can be many other minus dense intervals near I*ℓ*. We therefore need a notion to identify when I*<sup>ℓ</sup>* is isolated.

<span id="page-4-1"></span>**Definition 2.1.** An interval  $I_{\ell}(x)$  is *plus favored* with respect to  $\sigma$  if it satisfies the following two conditions:

- **(I)**  $\sigma_y = 1$  whenever  $1 \le d(y, I_\ell(x)) < 2^{\ell-1}$  (this will be used in e.g. Lemma [2.34](#page-21-0) later).
- **(II)** All the  $2M_{\ell}$  closest neighboring  $\ell$ -intervals are plus dense, that is, for all  $1 \leq k \leq M_{\ell}$ ,  $I_{\ell}(x+16k)$  and  $I_{\ell}(x-16k)$  are plus dense.

We may also consider the following weaker notion of dense and favored: I*<sup>ℓ</sup>* is *weakly* plus dense (with respect to  $\sigma$ ) when  $|I_{\ell}^{+}(\sigma)| > 2^{\ell} \left(1 - \frac{c_1}{M_{\ell}}\right)$ ), and  $I_{\ell}$  is *weakly plus favored* (with respect to  $\sigma$ ) if it satisfies **(I)** and **(II)** after we replace  $M_{\ell}$  by  $M'_{\ell} := 2 \left| \frac{M_{\ell}}{2c_1} \right|$  $\Big|$  (so  $M'_{\ell}$  is even, which is only for notational convenience later when divided by 2) and replace plus dense by weakly plus dense, where  $c_1 \geq 10$  $c_1 \geq 10$  $c_1 \geq 10$  is a constant fixed at the beginning of this section. See Figure 1 for an illustration of plus favored interval. Similarly, we can define the *minus favored*, *weakly minus dense*, and *weakly minus favored* intervals.

<span id="page-5-1"></span>

Figure 1: The figure depicts a configuration  $\sigma$  where the red cubes denote plus spins and blue cubes denote minus spins. Taking  $M_0 = 1$  and  $\delta = \frac{2}{3}$ , the interval  $I_3(x)$  is plus favored, since its  $2^{3-1} = 4$ closest neighbors in both directions are positive, and the number of minuses in each interval of the form I<sub>3</sub>( $x \pm 16$ ) and I<sub>3</sub>( $x \pm 32$ ) is smaller than  $\frac{2^3}{2^3}$  $\frac{2^3}{2^{3}\cdot\frac{1}{3}}$  = 4. Moreover, as there are minuses inside I<sub>3</sub>(*x*) and it is plus favored, it is also isolated.

**Definition 2.2.** Given any  $r \geq 1$  and any interval  $I \subset \mathbb{Z}$ , we define  $\rho_r(I) := \{x \in \mathbb{Z} : d(x,I) \leq I\}$  $(r-1)|I|\},\$  the expansion of I by a  $2r-1$  factor.

<span id="page-5-2"></span>**Definition 2.3.** An interval  $I_{\ell}$  is *isolated* with respect to  $\sigma$  when it is plus favored and  $I_{\ell}^-(\sigma) \neq \emptyset$ , or minus favored and  $I_{\ell}^{+}(\sigma) \neq \emptyset$ . Similarly, we define the *weakly isolated* intervals. We say  $\sigma$  is *balanced* in I if for every  $\ell \geq 0$ , every  $\ell$ -interval  $I' \in I_{\ell}$  with  $\rho_{M_{\ell}}(I') \subset I$  is not isolated with respect to *σ*. A plus favored isolated interval will be called a *plus isolated* interval. Similarly, we define the *minus isolated* interval.

We denote by  $\mathscr{C}_{\ell}^{\pm}(\sigma)$  the plus/minus isolated intervals in the  $\ell$ -th scale. We also denote  $\mathscr{C}^{\pm}(\sigma) \coloneqq$  $\cup_{\ell \geq 0} \mathscr{C}_{\ell}^{\pm}(\sigma)$  and  $\mathscr{C}(\sigma) := \mathscr{C}^{+}(\sigma) \cup \mathscr{C}^{-}(\sigma)$ . In addition, let  $\mathscr{C}^{+,0}(\sigma) := \{I \in \mathscr{C}^{+}(\sigma) : 0 \in I\}$  be the set of all plus isolated intervals in  $\sigma$  containing the origin.

To determine the spins of  $\sigma$  to be flipped, we transform it into a balanced configuration by erasing all isolated intervals, one by one. Next, we precisely describe this erasing process (as we will see below, we erase a plus/minus isolated interval by flipping all minuses/pluses therein).

For any  $\sigma \in \Omega$  and  $A \subset \mathbb{Z}$ , let us denote by  $\tau_A(\sigma)$  the configuration  $\sigma$  after we flip all spins in *A*, so  $\tau_A(\sigma) := ((-1)^{1_A(x)} \sigma_x)_{x \in \mathbb{Z}}$ . Given any  $\sigma \in \Omega^+$ , at each step  $s \geq 0$ , starting with the configuration  $\sigma^0 := \sigma$ , we employ the following *balancing procedure*.

- 1. Take  $\ell \geq 0$  the smallest integer such that there exists an isolated  $\ell$ -interval I<sub>l</sub> with I<sub>l</sub>  $\notin$  $\mathscr{C}^{+,0}(\sigma^s)$  (that is, if I<sub>ℓ</sub> is a plus isolated interval then it cannot contain the origin; this is because we do not want to flip the origin). If there is more than one isolated *ℓ*-interval, pick the leftmost one and denote it by  $B_s$ ; if there is no such isolated interval, we then stop our procedure.
- 2. Flip all minuses from B*<sup>s</sup>* if it is a plus isolated interval, and flip all pluses if it is minus isolated. That is, we take  $\sigma^{s+1} := \tau_{B^+_s(\sigma^s)}(\sigma^s) 1\!\!1_{\{B_s \in \mathscr{C}^- \}} + \tau_{B^-_s(\sigma^s)}(\sigma^s) 1\!\!1_{\{B_s \in \mathscr{C}^+ \}}$  (recall that  $B^{\pm}(\sigma) = \{v \in B : \sigma_v = \pm 1\}$  for a set B).

At each step, the balancing procedure selects one isolated interval and erases it. See Figure [2](#page-6-1) for an illustration. Lemma [2.6](#page-7-0) bellow assures that our procedure will stop in a finite time.

<span id="page-5-0"></span>**Definition 2.4.** We let  $S := S(\sigma)$  be the last step of the balancing procedure and let  $I_{\sigma}$  be the smallest isolated interval with respect to  $\sigma^S$  containing the origin (if there is more than one, pick the leftmost one; we let  $I_{\sigma} = \emptyset$  if such an interval does not exist). The spins to be flipped in the Peierls argument are  $A_{\sigma} := I_{\sigma}^{-}(\sigma^{S}).$ 

<span id="page-6-1"></span>

Figure 2: Here is an illustration of the balancing procedure. As in Figure [1,](#page-5-1) red regions denote plus spins, while blue regions denote minus spins. We also consider  $M_0 = 1$  and  $\delta = \frac{2}{3}$ . Starting from configuration  $\sigma^0$ , the first isolated interval selected is I<sub>3</sub>. As it is plus favored, it is flipped to plus. Next,  $I'_3$  is selected and flipped to minus for an analogous reason. Then,  $I_4$  is selected and flipped to minus. Notice that I<sub>4</sub> is isolated with respect to  $\sigma^2$  but not in  $\sigma^1$ , since  $M_4 = 2^{\frac{8}{3}} \approx 6.34$  and all its first 6 nearest-neighboring intervals are minus dense with respect to  $\sigma^2$  but not  $\sigma^1$ . Notice that, in  $\sigma^3$ , I<sub>5</sub> is not plus favored, since  $M_5 = 2^{\frac{10}{3}} \approx 10.07$ , but not all of its 10 closest neighbors are plus dense. The procedure stops at  $\sigma^3$ , since the only isolated interval  $(I_9)$  is plus favored but contains 0.

## <span id="page-6-0"></span>**2.2 The analysis of the Peierls map**

In this subsection, we analyze the Peierls map constructed in Section [2.1.](#page-4-0) To this end, we first show that the balancing procedure eventually stops, for which the following lemma will be useful. For any interval B selected in the balancing procedure, let  $s_B$  denote the step in which it was selected.

<span id="page-6-2"></span>**Lemma 2.5.** Let I be an interval selected at step  $s<sub>I</sub>$  (in the balancing procedure). Then for any  $k \geq 1$ ,  $\sigma^{s_1+k}|_{\rho_{\frac{3}{2}}(I)}$  *is constant (here*  $\sigma|$ *. means the restriction of*  $\sigma$ *). Furthermore, for all intervals* 2 B *selected in step*  $s_B > s_I$ *, we have*  $B \nsubseteq I$ *.* 

*Proof.* By symmetry, we assume that I is flipped to plus at step  $s_I$ . Note that  $\sigma^{s_I+1}|_{\rho_{\frac{3}{2}}(I)} = +1$  for the following reasons: for  $x \in \rho_{\frac{3}{2}}(I) \setminus I$  we have  $\sigma^{s_1} = 1$  by Definition [2.1](#page-4-1) **(I)** (this obviously holds at step  $s_1 + 1$ ); for  $x \in I$ , we have  $\sigma_x^{s_1+1} = 1$  by the definition of the balancing procedure. Suppose now there is a step  $s > s_1$  such that  $\sigma^s|_{\rho_{\frac{3}{2}}(I)} = u$  is constant and  $\sigma^{s+1}|_{\rho_{\frac{3}{2}}(I)}$  is not constant. Let B be the interval selected at step *s* in the balancing procedure. Since we flipped B to  $-u$ , the neighboring sites of B must have sign  $-u$  in  $\sigma^s$ , by the definition of our balancing procedure. As  $\sigma^s|_{\rho_{\frac{3}{2}}(I)} = u$ , these neighboring sites of B cannot be contained in  $\rho_{\frac{3}{2}}(I)$ . Hence we have either  $\rho_{\frac{3}{2}}(I) \cap B = \emptyset$  or  $\rho_{\frac{3}{2}}(I) \subset B$ . This contradicts the fact that  $\sigma^{s}|_{\rho_{\frac{3}{2}}(I)}$  is constant and  $\sigma^{s+1}|_{\rho_{\frac{3}{2}}(I)}$  is not constant.

In addition, for any interval B, if in some step we flipped B to  $u \in \{-1, 1\}$ , then B must have two neighboring *u* sites by the definition of our balancing procedure. Since  $\sigma^{s_1+k}|_{\rho_{\frac{3}{2}}(I)}$  is always constant for  $k \geq 1$ , we get that  $B \nsubseteq I$ . Altogether, this completes the proof of the lemma.  $\Box$  <span id="page-7-0"></span>**Lemma 2.6.** For any finite interval  $\Lambda \subset \mathbb{Z}$  containing the origin and any  $\sigma \in \Omega_{\Lambda}^{+}$ , the balancing *procedure starting from*  $\sigma$  *will end in finite steps. Furthermore, the final configuration*  $\sigma$ <sup>S</sup> *is balanced in*  $\rho_{\frac{3}{2}}(I_{\sigma})$ *. In addition, if*  $\sigma_0 = -1$ *, then*  $I_{\sigma} \neq \emptyset$  *and*  $A_{\sigma} \subset \Lambda$ *.* 

*Proof.* We first prove that the balancing procedure will end in a finite step. By Lemma [2.5,](#page-6-2) any interval  $B \in \mathcal{I}$  will be selected in the procedure at most once. Thus, it suffices to show that the number of intervals that may be selected is finite. We first claim that spins outside  $\Lambda$  are never flipped to minus. Assuming otherwise, we can take B as the first minus isolated interval selected in the balancing procedure with  $B \cap \Lambda^c \neq \emptyset$ . Supposing that B was selected in step *s*<sub>B</sub>, since B is minus isolated with respect to  $\sigma^{s_B}$ , we get that the nearest neighbor of B outside  $\Lambda$  must take value  $-1$  in  $\sigma$ <sup>sB</sup>. This contradicts our assumption that B was the first minus isolated interval intersecting Λ *c* selected in the procedure.

From the above claim, it follows in particular that

<span id="page-7-1"></span>
$$
\sigma^S|_{\Lambda^c} = 1. \tag{2.2}
$$

We also get that any selected interval B with  $B \cap \Lambda^c \neq \emptyset$  in the balancing procedure can only be flipped to plus. By the definition of plus isolated interval, we get that B contains at least one minus spin and thus  $B \cap \Lambda \neq \emptyset$ . As a result, we get that no interval B with  $B \cap \Lambda = \emptyset$  can be selected during the balancing procedure. Moreover, the number of intervals  $B \subset \Lambda$  is clearly finite, so it suffices to show that the number of intervals B selected in the balancing procedure satisfying  $B \cap \Lambda \neq \emptyset$  and  $B \cap \Lambda^c \neq \emptyset$  is finite.

To this end, we note that every such interval B must be plus favored, as argued in the first paragraph of this proof. Recall that  $B \in \mathcal{I}$ , so if B has length  $2^n$ , we must have  $\min_{x \in B} |x| \geq 2^{n-4}$ . This is because each *n*-interval in I can be written as  $[2^{n-4}x, 2^{n-4}(x+16))$  for some  $x \in \mathbb{Z}$ , and in addition we have  $x > 0$  or  $x \le -16$  since  $0 \notin B$ . Combining with the fact  $B \cap \Lambda \ne \emptyset$ , we get that the length of B is smaller than  $32|\Lambda|$  and therefore the number of intervals in this case is finite.

Next we prove that  $\sigma^S$  is balanced in  $\rho_{\frac{3}{2}}(I_{\sigma})$ . Otherwise there would exist an  $\ell \geq 0$  and an  $\ell$ interval  $B \subset \rho_{\frac{3}{2}}(I_{\sigma})$  isolated with respect to  $\sigma^S$ . By Definition [2.3,](#page-5-2) we get that  $\rho_{M_{\ell}}(B) \subset \rho_{\frac{3}{2}}(I_{\sigma})$  and thus B is smaller than  $I_{\sigma}$ . Moreover, since  $\sigma^S$  is the final configuration of the balancing procedure, B must be plus favored and contain the origin. This, however, contradicts (the 'smallest' in) the definition of I*σ*.

Finally, we prove that  $I_{\sigma} \neq \emptyset$  and  $A_{\sigma} \subset \Lambda$  whenever  $\sigma_0 = -1$ . By the definition of the balancing procedure, a plus favored interval containing 0 has never been selected through the balancing procedure. Thus, if  $\sigma_0 = -1$ , the spin at the origin has never been flipped to +1 and hence we have  $\sigma_0^S = -1$ . From this, it follows that  $0 \in B^-(\sigma)$  for every plus favored interval  $B \in \mathcal{I}$  containing the origin.

From [\(2.2\)](#page-7-1), we get that there exists at least one plus favored interval containing the origin (any interval  $B \in \mathcal{I}$  with  $\Lambda \subset B$  is plus isolated), and thus  $I_{\sigma}$  exists. Moreover,  $A_{\sigma} = I_{\sigma}^{-}(\sigma^{S}) \subset \Lambda$ , since by [\(2.2\)](#page-7-1) all spins outside must be plus.

One desirable feature of our balancing procedure, as shown in Lemma [2.9,](#page-8-0) is that the final configuration inherits structural properties of the original configuration: under some suitable conditions, dense regions of pluses/minuses in the original configuration remain to be weakly dense regions of pluses/minuses in the final configuration. We introduce some auxiliary definitions and lemmas before proving Lemma [2.9.](#page-8-0)

<span id="page-8-3"></span>**Definition 2.7.** An interval I is *tame* with respect to  $\sigma$  up to step *T* if  $|I \cap B_t^c| \geq |I|/16$  for all steps  $t < T$ , where  $B_t$  is the interval selected at step t in the balancing procedure of  $\sigma$ .

<span id="page-8-4"></span>**Lemma 2.8.** *Suppose that*  $B = [b, b + 2^{\ell})$  *is a plus favored interval with respect to*  $\sigma$ *. Then for any integer*  $0 < p \leq M_{\ell} 2^{\ell}$ *, we have* 

<span id="page-8-1"></span>
$$
|\{x \in [b + 2^{\ell}, b + 2^{\ell} + p) : \sigma_x = 1\}| \ge (1 - \frac{2}{M_{\ell}})p,
$$
  

$$
|\{x \in [b - p, b) : \sigma_x = 1\}| \ge (1 - \frac{2}{M_{\ell}})p.
$$
 (2.3)

*Proof.* By symmetry, it suffices to prove [\(2.3\)](#page-8-1). We consider three cases according to the value of p.

- 1. If  $p \leq 2^{\ell-1}$ , then all spins in  $[b+2^{\ell}, b+2^{\ell}+p)$  must be plus, and thus [\(2.3\)](#page-8-1) holds immediately.
- 2. If  $2^{\ell-1} < p \le 2^{\ell}$ , as the number of minuses in  $[b + 2^{\ell}, b + 2^{\ell} + p)$  is at most  $\frac{2^{\ell}}{M}$  $\frac{2^s}{M_{\ell}}$ , we get  $(2.3)$ immediately since  $\frac{2^{\ell}}{M}$  $\frac{2^{\ell}}{M_{\ell}} \leq \frac{2p}{M_{\ell}}.$
- 3. If  $p > 2^{\ell}$ , let  $k = \lfloor \frac{p}{2^{\ell}} \rfloor$ , so  $p \ge k \cdot 2^{\ell}$ . Since  $p \le M_{\ell} 2^{\ell}$ , the  $\ell$ -interval  $[b+i \cdot 2^{\ell}, b+(i+1) \cdot 2^{\ell})$  is  $\sum_{i=1}^{n} p_i > 2$ ,  $\sum_{i=1}^{n} p_i = \sum_{i=1}^{n} p_i > 2$ . Since  $p \leq m_i/2$ , the value of minus vacant for any integer  $0 < i \leq k$ , implying that the number of minuses in  $[b+2^{\ell}, b+2^{\ell}+p)$ is at most  $\frac{(k+1)2^{\ell}}{M_{\ell}}$ . Again, the trivial bound  $\frac{(k+1)2^{\ell}}{M_{\ell}} \leq \frac{2k \cdot 2^{\ell}}{M_{\ell}}$  $\frac{k \cdot 2^{\ell}}{M_{\ell}} \leq \frac{2p}{M_{\ell}}$  yields the desired [\(2.3\)](#page-8-1).

<span id="page-8-0"></span>**Lemma 2.9.** Let I be an interval with length  $2^n \leq |I| < 2^{n+1}$ . Let  $\sigma \in \Omega_{\Lambda}^+$  and let  $T \geq 1$  be a step such that I is tame with respect to  $\sigma$  up to step T. If  $|I^+(\sigma)| \leq \frac{|I|}{\sqrt{M_n}}$ , then  $|\bigcup_{t=0}^T I^+(\sigma^t)| \leq c_1 |I^+(\sigma)|$ .

*Proof.* Our proof of Lemma [2.9](#page-8-0) is by induction on *n*. If  $n = 1$  then  $|I| = 1$ . Thus, no interval intersecting I is allowed to be selected, implying the claim immediately. Now we assume the lemma holds for any  $n \leq N-1$ , any starting configuration  $\sigma$  and any  $T \geq 1$ . We next prove by induction the case when  $n = N$ . To this end, let  $T' + 1$  be the first step before T such that an  $\ell$ -interval B with  $M_{\ell}2^{\ell} > 2^n$  and  $B \cap I \neq \emptyset$  was selected and flipped to plus (if no such interval exists before step *T*, then we take  $T' = T$ ). We claim that it suffices to prove that

<span id="page-8-2"></span>
$$
|\bigcup_{t=0}^{T'} \mathbf{I}^+(\sigma^t)| \le c_1 |\mathbf{I}^+(\sigma)|,\tag{2.4}
$$

i.e., the desired bound holds at step  $T'$ . To this end, we next prove that in fact, we must have  $T' = T$ assuming  $(2.4)$  holds. Otherwise, if  $T' < T$ , as B was selected at step  $T' + 1$  and I was assumed to be tame up to step *T* (recall Definition [2.7\)](#page-8-3), we must have  $|I \cap B^c| \ge \frac{|I|}{16}$ . Since  $2^n < M_{\ell} 2^{\ell} \le M_n 2^n$ , by Lemma [2.8](#page-8-4) the minus density in  $I \ B$  is at most  $\frac{2}{M_{\ell}}$ . Hence we have  $|I^+(\sigma^{T'})| \geq (1 - \frac{2}{M_{\ell}})|I \ B| \geq \frac{|I|}{32}$ , which contradicts [\(2.4\)](#page-8-2). This then proves that  $T = T'$  and the desired result follows. The rest of the proof is dedicated to proving [\(2.4\)](#page-8-2).

From now on, we fix a starting configuration  $\sigma$ . In order to keep track of the change during the balancing procedure, for any  $x \in I$  we record the first selected interval  $D^x \ni x$  which results in flipping *x* to plus (*x* may or may not be flipped again, but that is not relevant to us). If *x* was never flipped to plus we just take  $D^x = \emptyset$ . Let  $\mathcal{F}_{\ell} := \{D^x \subset \mathbb{Z} : |D^x| = 2^{\ell}, x \in I\}$ be the collection of  $\ell$ -intervals we have recorded. For any  $B \in \mathcal{F}_{\ell}$ , let  $s_B$  be the step in which B was flipped and let  $P_{\ell}(\mathbf{B}) = |\bigcap_{t=0}^{s_{\mathbf{B}}} \mathbf{B}^{-}(\sigma^{t})|$  be the number of new pluses created at the flip. We say  $B \in \mathcal{F}_{\ell}$  is *negligible* when  $P_{\ell}(B) \leq \frac{2^{\ell}}{M_{\ell}}$  $\frac{2^{\circ}}{M_{\ell}}$ , i.e., not many new pluses were created when flipping minuses in *B*; otherwise we say B is *non-negligible*. We further split F*<sup>ℓ</sup>* into the union of  $\mathcal{F}_{\ell}^{j} := \mathcal{F}_{\ell} \cap \{[2^{\ell}(x + \frac{j}{16}), 2^{\ell}(x + \frac{j}{16} + 1))\}_{x \in \mathbb{Z}}$  for  $j = 0, \ldots, 15$ .

<span id="page-9-0"></span>**Claim 2.10.** For negligible intervals, if  $\ell < n$  we have

<span id="page-9-2"></span>
$$
\sum_{\substack{\mathbf{B}\in\mathcal{F}_{\ell}^j\\ \mathbf{B} \text{ negligible}}} P_{\ell}(\mathbf{B}) \le \frac{8|\mathbf{I}^+(\sigma)|}{\sqrt{M_{\ell}}}.\tag{2.5}
$$

<span id="page-9-1"></span>**Claim 2.11.** For non-negligible intervals, if  $\ell < n$  we have

<span id="page-9-3"></span>
$$
\sum_{\substack{\mathbf{B}\in\mathcal{F}_{\ell}^j\\ \mathbf{B} \text{ non-}negligible}} P_{\ell}(\mathbf{B}) \le \frac{2|\mathbf{I}^+(\sigma)|}{M_{\ell}^{1/3}}.
$$
\n(2.6)

For smooth presentation, we will first complete the proof of the lemma (i.e., proving  $(2.4)$ ) assuming Claims [2.10](#page-9-0) and [2.11,](#page-9-1) and provide their proofs afterward. We avoid stating these claims as independent lemmas since we are under the induction hypothesis of the present lemma.

Note that  $\bigcup_{t=0}^{T'} I^+(\sigma^t) \subset I^+(\sigma) \cup \left($  $\cup_{t=1}^{T'} I^+(\sigma^t) ∩ I^-(\sigma)$ . Recall the definition of  $\mathcal{F}_{\ell}$  and  $P_{\ell}(B)$ . Every  $x \in \bigcup_{t=0}^{T'} I^+(\sigma^t) \cap I^-(\sigma)$  must be in an interval that was selected in the balancing procedure and flipped to plus (i.e., this selected interval was an isolated plus interval). Then we have

$$
\left|\bigcup_{t=0}^{T'} \mathrm{I}^+(\sigma^t) \cap \mathrm{I}^-(\sigma)\right| \leq \sum_{\ell=0}^{\ell_n} \sum_{\mathrm{B} \in \mathcal{F}_{\ell}} P_{\ell}(\mathrm{B}),
$$

where  $\ell_n := \lfloor \frac{n - \log_2(M_0)}{1 + \delta} \rfloor$  is the maximal integer  $\ell$  such that  $M_{\ell} 2^{\ell} \leq 2^n$ .

Combining  $(2.5)$  and  $(2.6)$ , as the upper bounds do not depend on  $j$ , we get

$$
|\bigcup_{t=1}^{T'} \mathcal{I}^+(\sigma^t) \cap \mathcal{I}^-(\sigma)| \le \sum_{\ell=0}^{\ell_n} \sum_{\mathbf{B} \in \mathcal{F}_{\ell}} |\mathbf{B}^-(\sigma^{s_{\mathbf{B}}})|
$$
  

$$
\le \sum_{\ell=0}^{\ell_n} 16 \times \left( \frac{2|\mathcal{I}^+(\sigma)|}{M_{\ell}^{1/3}} + \frac{8|\mathcal{I}^+(\sigma)|}{\sqrt{M_{\ell}}} \right) \le (c_1 - 1)|\mathcal{I}^+(\sigma)|,
$$

where the last inequality comes from choosing  $M_0 > 0$  large enough. Thus we complete the proof of [\(2.4\)](#page-8-2).  $\Box$ 

*Proof of Claim [2.10](#page-9-0).* Recall that for any negligible interval  $B \in \mathcal{F}_{\ell}^j$ ,  $s_B$  is the step that B was flipped. Let  $C = \rho_{\frac{3}{2}}(B) \cap I$ . We will prove that C is tame with respect to  $\sigma$  up to step  $s_B$  (and then we may apply our induction hypothesis to C, i.e., apply Lemma [2.9](#page-8-0) to C; note that in the context of induction we have  $|C| \leq 2^n$ ). If an interval I' with  $|(I')^c \cap C| \leq |C|/16$  was selected in the balancing procedure before step  $s_B$ , the length of I' must be at least  $\frac{15}{16} \cdot |C|$  and thus  $C \subset \rho_{\frac{3}{2}}(I')$ . By Lemma [2.5,](#page-6-2)  $\rho_{\frac{3}{2}}(I')$  would be either all plus or all minus after that flip. In addition, since B is plus favored at step  $s_B$ , the only possibility is that  $\rho_{\frac{3}{2}}(I')$  is all plus at step  $s_B$  and this contradicts the fact that at least one vertex in B∩I was flipped to plus at step *s*<sub>B</sub>. Therefore, only intervals I' with  $|(I')^c \cap C| > |C|/16$  can be selected in the balancing procedure before step  $s_B$ . This implies that C satisfies the claimed tame condition.

Next, we want to control the number of negligible  $\ell$ -intervals in  $\mathcal{F}_{\ell}^j$ . For any negligible interval  $B \in \mathcal{F}_{\ell}^j$ ,  $\rho_{\frac{3}{2}}(B) \setminus B$  is the union of two intervals, and one of them must be contained in I since  $\ell < n$ . In

<span id="page-10-0"></span>

Figure 3: The red region denotes the interval  $B_i$ ; the black line denotes the interval  $C_i$ , and is divided into 16 parts. The black box delimits the interval I' that covers more than a  $\frac{15}{16}$  fraction of  $C_i$ , and the gray box represents  $\rho_{\frac{3}{2}}(I')$ . The dotted line splits I' in half.

addition, since B is plus favored at step  $s_B$  we have  $\sigma^{s_B}|_{\rho_{\frac{2}{3}}(B)\setminus B} = 1$ , and thus  $|\bigcup_{t=0}^{s_B} C^+(\sigma^t)| \geq 2^{\ell-1}$ . Applying the induction hypothesis combined with the fact that  $|C| \leq 2^{\ell+1}$  and  $\frac{1}{c_1}(2^{\ell-1}) > \frac{2^{\ell-1}}{\sqrt{M_{\ell+1}}}$ , we get  $|C^+(\sigma)| \ge \frac{2^{\ell-1}}{\sqrt{M_{\ell+1}}}$ .

Note that for any non-neighboring intervals  $B, B' \in \mathcal{F}_{\ell}^j$  we have  $\rho_{\frac{3}{2}}(B) \cap \rho_{\frac{3}{2}}(B') = \emptyset$ , and thus  $C \cap C' = \emptyset$ , where  $C' = \rho_{\frac{3}{2}}(B') \cap I$ . This implies that, for any  $x \in \mathbb{Z}$ , there are at most two  $B \in \mathcal{F}_{\ell}^j$ with  $x \in \rho_{\frac{3}{2}}(B)$ . As there are  $|I^+(\sigma)|$  pluses in I at the start of the balancing procedure, we conclude that the number of negligible intervals in  $\mathcal{F}_{\ell}^{j}$  is at most  $2|I^{+}(\sigma)|$  $\sqrt{M_{\ell+1}}$  $\frac{2^{\ell-1}}{2^{\ell-1}}$ . Combining with the fact that  $P_{\ell}(\mathbf{B}) \leq \frac{2^{\ell}}{M_{\ell}}$  $\frac{2^{\ell}}{M_{\ell}}$  for any negligible interval  $B \in \mathcal{F}_{\ell}^{j}$ , we get that

$$
\sum_{\substack{\mathbf{B}\in\mathcal{F}_{\ell}^j\\ \mathbf{B}_i \text{ negligible}}} P_{\ell}(\mathbf{B}) \leq 2|\mathbf{I}^+(\sigma)|\frac{\sqrt{M_{\ell+1}}}{2^{\ell-1}} \cdot \frac{2^{\ell}}{M_{\ell}} \leq \frac{8|\mathbf{I}^+(\sigma)|}{\sqrt{M_{\ell}}},
$$

 $\Box$ 

concluding the proof of Claim [2.10.](#page-9-0)

*Proof of Claim [2.11](#page-9-1).* We list the non-negligible intervals in  $\mathcal{F}_{\ell}^{j}$  as  $\{B_1, \ldots, B_K\}$ . For  $1 \leq i \leq K$ , we write  $s_i := s_{\text{B}_i}$ . We want to find an interval  $\text{C}_i$  neighboring  $\text{B}_i$  such that  $\text{C}_i$  has a large density of pluses. The set  $\{x \in \mathbb{Z} : 1 \leq d(x, B_i) \leq \frac{M_{\ell}-1}{2} 2^{\ell}\}\)$  is the union of two intervals, and one of them (which we denote as  $C_i$ ) must be contained in I, since  $M_{\ell}2^{\ell} \leq 2^n$ . In addition, we have  $|C_i| < 2^n$ . Similar to the proof of Claim [2.10,](#page-9-0) we want to apply the induction hypothesis to get a lower bound on the number of pluses in C*<sup>i</sup>* at the start of the balancing procedure, and then give an upper bound on the number of non-negligible *ℓ*-intervals.

As in the proof of Claim [2.10,](#page-9-0) we now show that  $C_i$  is tame with respect to  $\sigma$  up to step  $s_i$ . Suppose this is not the case. So, there is an interval I' with  $|(I')^c \cap C_i| \leq |C_i|/16$  selected in the balancing procedure at some step  $s'_i < s_i$ . Then by Lemma [2.5,](#page-6-2) we get that  $\rho_{3/2}(\mathbf{I}')$  is constant in  $\sigma^{s'_i+k}$  for any  $k \geq 1$ . Note that  $|I'| \geq \frac{15|C_i|}{16}$  and thus  $\frac{|I'|}{2} \geq \frac{15|C_i|}{32} > \frac{|C_i|}{8} + |B_i|$  (see Figure [3](#page-10-0) for an illustration), so we have  $B_i \subset \rho_{3/2}(I')$ . Since  $B_i$  is plus isolated with respect to  $\sigma^{s_i}$ , this implies that  $\sigma^{s_i}|_{B_i}$  contains at least a minus spin. Thus  $\sigma^{s_i}|_{\rho_{\frac{3}{2}}(I')} = -1$ . However, this contradicts the fact that  $B_i$  is plus favored with respect to  $\sigma^{s_i}$ . This then verifies the desired tame condition for  $C_i$ .

Next, we want to control the number of non-negligible  $\ell$ -intervals in  $\mathcal{F}_{\ell}^j$ . Since  $B_i$  is plus favored *ℓ* at step  $s_i$ , we get that  $|\mathcal{C}_i^+(\sigma^{s_i})| \geq \frac{M_{\ell}-1}{2}$  $\left(1-\frac{1}{M_{\ell}}\right)$  $\left(2^{\ell} \cdot \text{Take } k < n \text{ satisfying } 2^{k} \le \frac{M_{\ell}-1}{2} 2^{\ell} \le 2^{k+1}. \right)$ As in the proof of Claim [2.10,](#page-9-0) we apply the induction hypothesis and combine it with the fact that  $\frac{1}{c_1}(\frac{M_{\ell}-1}{2})$  $\left(1-\frac{1}{M_{\ell}}\right)$  $\left(2^{\ell}\right)>\frac{1}{\sqrt{\Lambda}}$  $\frac{1}{M_k} \frac{M_\ell - 1}{2} 2^\ell$  to get

<span id="page-11-0"></span>
$$
|C_i^+(\sigma)| \ge \frac{1}{\sqrt{M_k}} \frac{M_\ell - 1}{2} 2^\ell \ge \frac{M_\ell^{1/3} 2^\ell}{2},\tag{2.7}
$$

where the last inequality holds since  $M_0 > 0$  is big enough and  $\delta < \frac{1}{20}$ .

To count the number of  $C_i$ 's, we first show that they are disjoint by proving that for any  $B_i, B_j$ , we have

<span id="page-11-3"></span>
$$
d(B_i, B_j) > (M_{\ell} - 1)2^{\ell}.
$$
\n(2.8)

To this end, we can assume without loss of generality that  $s_i < s_j$ , i.e.,  $B_i$  was flipped first. If  $d(B_i, B_j) \leq (M_{\ell} - 1)2^{\ell}$ , since  $B_i$  is plus isolated with respect to  $\sigma^{s_i}$ ,  $B_j$  must be minus vacant with respect to  $\sigma^{s_i}$ . Thus,  $|B_j^-(\sigma^{s_i})| \leq \frac{2^{\ell}}{M_i}$  $\frac{2^{\circ}}{M_{\ell}}$ , which contradicts to the fact that B<sub>j</sub> is non-negligible.

Note that the choice of  $C_i$  ensures that  $C_i \subset I$ . Combining with the fact that  $C_i$ 's do not intersect, we get from [\(2.7\)](#page-11-0) that the maximum number of non-negligible intervals is at most  $|I^+(\sigma)|\frac{2}{M_{\ell}^{1/3}2^{\ell}},$  and therefore

$$
\sum_{i=1}^{K} P_{\ell}(\mathbf{B}_{i}) \leq |\mathbf{I}^{+}(\sigma)| \frac{2}{M_{\ell}^{1/3} 2^{\ell}} \cdot 2^{\ell} \leq \frac{2|\mathbf{I}^{+}(\sigma)|}{M_{\ell}^{1/3}}.
$$

Thus we complete the proof of Claim [2.11.](#page-9-1)

Suppose at some step of the balancing procedure we selected a plus favored interval I and flipped all minus spins in I. In the following steps of the balancing procedure, we may have created minuses close to I, so that it is no longer plus favored with respect to the final configuration. We can however show that I will still be weakly plus favored with respect to the final configuration, as incorporated in the next proposition.

<span id="page-11-2"></span>**Proposition 2.12.** Let  $\sigma^T$  be the configuration obtained via the balancing procedure starting from *σ after T steps.* Given  $x \in \mathbb{Z}$  *with*  $\sigma_x^s \neq \sigma_x^T$  *for some*  $0 \leq s < T$ , *let* I *be the last interval selected in the balancing procedure before step T that flips x. Then* I *is weakly plus (resp. minus) favored with respect to*  $\sigma^T$  *if*  $\sigma_x^T = 1$  (*resp.*  $\sigma_x^T = -1$ ).

*Proof.* Let us assume that  $\sigma_x^s = -\sigma_x^T = -1$ . Let  $s_I$  be the step in the balancing procedure in which I was flipped. First notice that all sites in  $\rho_{\frac{3}{2}}(I)$  are positive in  $\sigma^{s_I+1}$ . Since I is the last interval selected before step *T* that flips *x*, we see that after this no interval containing I can be selected after step  $s_I$  and flipped to minus. Combined with Lemma [2.5,](#page-6-2) it implies that all spins of  $\sigma^T$  in  $\rho_{\frac{3}{2}}(I)$  are still plus.

Now, let  $\ell \geq 0$  and  $x \in \mathbb{Z}$  be such that  $I = I_{\ell}(x)$ . For any  $1 \leq |k| \leq M_{\ell}/c_1$  and any interval  $B_k = I_{\ell}(x + 16k)$ , we have  $|B_k(\sigma^{s_1})| \leq \frac{2^{\ell}}{M_k}$  $\frac{2^{\ell}}{M_{\ell}}$  (since I is plus favored with respect to  $\sigma^{s_1}$ ). Without loss of generality, we only consider  $k \geq 0$ . We want to apply induction on  $k$  to show that

<span id="page-11-1"></span>
$$
\left|\bigcup_{t=s_1}^T \mathbf{B}_k^-(\sigma^t)\right| \le \frac{c_1 2^\ell}{M_\ell}.\tag{2.9}
$$

 $\Box$ 

The case  $k = 0$  holds since I remains all plus after step  $s<sub>1</sub>$ . Now we suppose [\(2.9\)](#page-11-1) holds for  $0 \leq k \leq m-1$ , and wish to prove (2.9) for  $k = m$ . To this end, we claim that there does not exist an interval C flipped to minus after step  $s_1$  such that  $|B_m \cap C^c| \leq \frac{|B_m|}{16}$ . If such a C exists, then  $|C| \geq 2^{\ell}$  and hence  $|\rho_{\frac{3}{2}}(C) \cap B_{m-1}| \geq \frac{2^{\ell}}{2} - \frac{2^{\ell}}{16} \geq 7 \cdot 2^{\ell-4}$ . By the definition of minus favored, we get that  $\rho_{\frac{3}{2}}(C)$  is all minus in  $\sigma^{s_c+1}$ . Thus  $B_{m-1}^-(\sigma^{s_c+1}) \geq 7 \cdot 2^{\ell-4}$  which contradicts to the induction hypothesis [\(2.9\)](#page-11-1) for  $k = m - 1$ , completing the verification of the claim.

We now prove  $(2.9)$  using the above claim and Lemma [2.9.](#page-8-0) Let  $T_m + 1$  be the first step that an  $\text{interval } C \text{ with } |B_m \cap C^c| \leq \frac{|B_m|}{16}$  was selected in the balancing procedure (if no such interval exists, then take  $T_m = T$ ). Then  $B_m$  is tame with respect to  $\sigma^{s_1}$  up to step  $T_m$ . Thus, by Lemma [2.9,](#page-8-0)

<span id="page-12-1"></span>
$$
\left|\cup_{t=s_1}^{T_m} \mathcal{B}_m^{-}(\sigma^t)\right| \le \frac{c_1 2^{\ell}}{M_{\ell}}.\tag{2.10}
$$

Furthermore, letting C be the interval selected at step  $T_m + 1$ , we see from our aforementioned claim that C is flipped to plus. Thus, we get by Lemma [2.5](#page-6-2) that  $\sigma^{T_m+t}|_{\rho_{\frac{3}{2}}(C)}$  is constant for all  $t \geq 1$ . Moreover, since no interval D with  $|\mathcal{B}_m \cap \mathcal{D}^c| \leq \frac{|\mathcal{B}_m|}{16}$  (in particular, no interval containing C) was selected and flipped to  $-1$ , we must have  $\sigma^{T_m+t}|_{\rho_{\frac{3}{2}}(C)} = +1$  for all  $t \ge 1$ . Combined with  $\Box$ the fact that  $B_m \subset \rho_{\frac{3}{2}}(C)$  and [\(2.10\)](#page-12-1), this completes the proof of [\(2.9\)](#page-11-1) for  $k = m$ .

#### <span id="page-12-0"></span>**2.2.1 Proof of Theorem [1.1](#page-1-0)**

In this subsection, we prove Theorem [1.1](#page-1-0) assuming Propositions [2.13,](#page-12-2) [2.14](#page-12-3) and Propositions [2.18,](#page-13-2) [2.20,](#page-13-0) whose proofs are postponed until Sections [2.3](#page-15-1) and [2.4](#page-24-0) respectively. Fix

<span id="page-12-5"></span>
$$
\theta := \min\{2 - \alpha - 10\delta, \log_{3.9}(2)\} > \frac{1}{2}.\tag{2.11}
$$

<span id="page-12-2"></span>Moreover, for all  $A, B \subset \mathbb{Z}$  we define  $J(A, B) = \sum_{x \in A, y \in B} J_{xy}$ .

**Proposition 2.13.** *Recall Definition [2.4.](#page-5-0) There is a constant*  $c_2 > 0$  *such that for any configuration*  $\sigma \in \Omega_{\Lambda}^{+},$ 

$$
J(A_{\sigma}, A_{\sigma}^c) \ge c_2 |\mathbf{I}_{\sigma}|^{\theta}.
$$

<span id="page-12-3"></span>**Proposition 2.14.** For any configuration  $\sigma \in \Omega_{\Lambda}^{+}$ ,

<span id="page-12-6"></span>
$$
H_{\Lambda,0}^{+}(\sigma) - H_{\Lambda,0}^{+}(\tau_{A_{\sigma}}(\sigma)) \ge J(A_{\sigma}, A_{\sigma}^{c}).
$$
\n(2.12)

Proposition [2.14](#page-12-3) shows that flipping spins in  $A_{\sigma}$  will gain energy in the interaction term. In order to control the term from the external field in the Hamiltonian, we want to flip the external field on  $A_{\sigma}$  as in [\[18\]](#page-38-10). Similar to  $\sigma$ , for any set  $A \subset \mathbb{Z}$  we define  $\tau_A(h) := ((-1)^{1_A(x)} h_x)_{x \in \mathbb{Z}}$ . We define the error function

<span id="page-12-4"></span>
$$
\Delta_A(h) := -\frac{1}{\beta} \log \frac{Z_{\Lambda;\beta,\varepsilon}^+(h)}{Z_{\Lambda;\beta,\varepsilon}^+(r_A(h))},\tag{2.13}
$$

where  $Z_{\Lambda;\beta,\varepsilon}^+(h)$  is the *partition function* (note that  $H_{\Lambda,\varepsilon}^{\eta}(\cdot)$  can be naturally regarded as a function of  $\sigma \in \{-1, 1\}^{\Lambda}$ )

$$
Z^{\eta}_{\Lambda;\beta,\varepsilon}(h)\coloneqq \sum_{\sigma\in\{-1,1\}^\Lambda}e^{-\beta H^n_{\Lambda,\varepsilon h}(\sigma)}.
$$

The key property required for the error functions [\(2.13\)](#page-12-4) is their subgaussian behavior, proved in [\[18\]](#page-38-10).

<span id="page-13-8"></span>**Lemma 2.15.** *For any*  $A, A' \in \mathbb{Z}$  *and*  $\lambda > 0$ *, we have* 

$$
\mathbb{P}(|\Delta_A(h) - \Delta_{A'}(h)| > \lambda \mid h_{A \cup A'^c}) \le 2e^{-\frac{\lambda^2}{8\varepsilon^2 |A \Delta A'|}}, \tag{2.14}
$$

<span id="page-13-7"></span>*where A*∆*A*′ *is the symmetric difference.*

**Definition 2.16.** Let  $\mathcal{A}(I)$  denote the collection of subsets  $A \subset I$  such that there exists a balanced configuration  $\sigma$  on I satisfying  $A = \mathbb{I}^-(\sigma)$ . In addition, for any constant  $Q > 0$  we define  $\mathcal{A}_Q(I)$  to be the collection of subsets  $A \in \mathcal{A}(\mathbf{I})$  such that  $J(A, A^c) \in [Q, 2Q)$ .

**Definition 2.17.** Suppose that  $I_n$  is an *n*-interval containing the origin. Let  $\hat{\mathcal{A}}(I_n)$  be the collection of subsets  $A \in \mathcal{A}(\mathbf{I}_n)$  such that  $J(A, A^c) \ge c_2 \cdot 2^{\theta n}$ . We define the good event to be

<span id="page-13-1"></span>
$$
\mathcal{E}(\mathbf{I}_n) := \left\{ \Delta_A(h) \le \frac{J(A, A^c)}{10}, \text{ for all } A \in \hat{\mathcal{A}}(\mathbf{I}_n) \right\}.
$$
 (2.15)

The next proposition implies that the probability of the bad event  $\mathcal{E}(\mathbf{I}_n)^c$  is small.

<span id="page-13-2"></span>**Proposition 2.18.** Suppose that  $I_n$  is an *n*-interval containing the origin. Then there exists  $c_3$  :=  $c_3(\alpha)$  *such that* 

$$
\mathbb{P}(\mathcal{E}(\mathbf{I}_n)^c) \le c_3^{-1} \exp(-c_3 \frac{2^{(2\theta - 1 - 2\delta)n}}{\varepsilon^2}).
$$

Now, it suffices to control the entropy, as incorporated in Proposition [2.20](#page-13-0) below. Recall that we have a partition of  $\mathbb{Z}$  into  $\ell$ -intervals as  $\mathcal{I}_{\ell}^0 = \{ [x \cdot 2^{\ell}, (x+1) \cdot 2^{\ell} \} \}_{x \in \mathbb{Z}}$ . We start with some definitions.

<span id="page-13-3"></span>**Definition 2.19.** Suppose that  $I_n$  is an *n*-interval. For any  $A \in \mathcal{A}(I_n)$  and  $0 \leq \ell \leq n$ , we define  $\Psi_{\ell}(A): \mathcal{I}_{\ell}^{0} \to \{-1, 0, 1\}$  as:

<span id="page-13-6"></span>
$$
\Psi_{\ell}(A, I_{\ell}(x)) = \begin{cases}\n-1 & \text{if } I_{\ell}(x) \subset A; \\
1 & \text{if } I_{\ell}(x) \cap A = \emptyset; \\
0 & \text{otherwise.} \n\end{cases}
$$
\n(2.16)

See Figure [4](#page-14-0) for an illustration of  $\Psi(A)$ .

<span id="page-13-0"></span>**Proposition 2.20.** Suppose that  $I_n$  is an *n*-interval containing the origin. Let  $E(\ell, Q, I_n) :=$  $|\{\Psi_{\ell}(A): A \in \mathcal{A}_{Q}(I_n)\}|$  denote the number of images in  $\Psi_{\ell}(\mathcal{A}_{Q}(I_n))$  (here  $\Psi_{\ell}(A)$  is treated as a *map*), and recall our definition of  $\theta$  from [\(2.11\)](#page-12-5). Then there exists a constant  $c_4 > 0$  such that the *following inequality holds for any*  $M \geq 1$  *and*  $\ell \leq n-4$ *:* 

<span id="page-13-4"></span>
$$
E(\ell, Q, I_n) \le E(\ell + 1, Q, I_n) \times c_4^{-1} \exp\left(\frac{c_4 Q}{2^{\ell \theta}}\right).
$$
 (2.17)

<span id="page-13-5"></span>**Corollary 2.21.** *Suppose that* I*<sup>n</sup> is an n-interval containing the origin. There exists a constant*  $c_5 > 0$  *such that the following inequality holds for any*  $M \geq 1$  *and*  $\ell \leq n-3$ *:* 

<span id="page-13-9"></span>
$$
E(\ell, Q, I_n) \le c_5^{-1} \exp\left(\frac{c_5 Q}{2^{\ell \theta}}\right). \tag{2.18}
$$

<span id="page-14-0"></span>

Figure 4: Considering a configuration  $\sigma$  in an interval I, and taking  $A = \Gamma^-(\sigma)$ , the picture depicts, from bottom to top,  $\Psi_0(A), \Psi_1(A), \Psi_2(A)$  and  $\Psi_3(A)$ . For every  $\ell = 0, 1, 2, 3$ , intervals  $I_\ell$  are painted red if  $\Psi_{\ell}(A, I_{\ell}) = 1$ , painted blue if  $\Psi_{\ell}(A, I_{\ell}) = -1$  and painted white otherwise. We write  $\sigma = \Psi_0(A)$  since both functions attribute the same value to each site.

*Proof.* Since  $I_n$  is an *n*-interval containing the origin, we get that  $A \subset [-2^n, 2^n)$  for any  $A \in \mathcal{A}(I_n)$ . Recall Definition [2.19.](#page-13-3) Thus we get that  $E(n-3, Q, I_n) \leq 3^{16}$ . Hence applying [\(2.17\)](#page-13-4) iteratively, we get that for  $\ell \leq n-4$ 

$$
E(\ell, Q, I_n) \le 3^{16} \cdot \prod_{k=\ell}^{n-4} c_4^{-1} \exp\left(\frac{c_4 Q}{2^{k\theta}}\right) \le c_5^{-1} \exp\left(\frac{c_5 Q}{2^{\ell\theta}}\right)
$$

with some  $c_5$  depending on  $c_4$  and  $\theta$ .

Now we are ready to prove Theorem [1.1.](#page-1-0)

*Proof of Theorem [1.1](#page-1-0).* Let  $\mathcal{J}_n$  denote the collection of *n*-intervals I such that  $0 \in I$ , and take  $\mathcal{H}_n \coloneqq \bigcap_{\mathbf{I} \in \mathcal{J}_n} \mathcal{E}(\mathbf{I}).$  Since  $|\mathcal{J}_n| \leq 16$ , we get from Proposition [2.18](#page-13-2) and a union bound that

<span id="page-14-1"></span>
$$
\mathbb{P}(\mathcal{H}_n) \ge 1 - 16c_3^{-1} \exp(-c_3 \frac{2^{(2\theta - 1 - 2\delta)n}}{\varepsilon^2}).
$$
\n(2.19)

Letting  $\mathcal{H} = \bigcap_{n=0}^{\infty} \mathcal{H}_n$ , we get from [\(2.19\)](#page-14-1) that

<span id="page-14-3"></span>
$$
\mathbb{P}(\mathcal{H}) \ge 1 - 16 \sum_{n=0}^{\infty} c_3^{-1} \exp(-c_3 \frac{2^{(2\theta - 1 - 2\delta)n}}{\varepsilon^2}) \ge 1 - C_1 \exp(\frac{1}{C_1 \varepsilon^2}).
$$
 (2.20)

Recall Definition [2.4.](#page-5-0) For any integer  $n \geq 0$  and constant  $Q > 0$ , let  $\Sigma_{Q,n}$  denote the collection of configurations such that

- 1.  $J(A_{\sigma}, A_{\sigma}^c) \in [Q, 2Q)$ .
- 2.  $I_{\sigma}$  is an *n*-interval.

By [\(2.12\)](#page-12-6), [\(2.13\)](#page-12-4)and [\(2.15\)](#page-13-1) (as well as the definition of  $\mathcal{H}$ ), we get that for any  $\sigma \in \Sigma_{Q,n}$  and  $h \in \mathcal{H}$ 

<span id="page-14-2"></span>
$$
\mu_{\Lambda;\beta,\varepsilon h}^+(\sigma) \le \mu_{\Lambda;\beta,\varepsilon \tau_{A_\sigma}(h)}^+(\tau_{A_\sigma}(\sigma)) \times \exp(-\beta \cdot 0.9Q). \tag{2.21}
$$

By Definition [2.19,](#page-13-3)  $\Psi_0(A, I_0(x)) = -\mathbb{1}_A(x)$ , so  $A \to \Psi_0(A)$  is clearly a bijection. In addition, since  $A_{\sigma} = I_{\sigma}^{-}(\sigma^{S})$  and  $\sigma^{S}$  is balanced on  $I_{\sigma}$  (by Lemma [2.6\)](#page-7-0), we get that  $A_{\sigma} \in \mathcal{A}(I_{\sigma})$ . Hence the number of choices for  $A_{\sigma}$  with  $\sigma \in \Sigma_{Q,n}$  is at most  $\sum_{I \in \mathcal{J}_n} E(0, Q, I)$ . By Corollary [2.21,](#page-13-5) we have

 $\Box$ 

 $E(0, Q, I) \leq c_5^{-1} \exp(c_5 Q)$  for any  $I \in \mathcal{J}_n$ . Combining with the fact that  $|\mathcal{J}_n| \leq 16$ , we get that the multiplicity of the map  $(\sigma, h) \to (\tau_{A_{\sigma}}(\sigma), \tau_{A_{\sigma}}(h))$  is at most 16  $c_5^{-1}$  exp( $c_5Q$ ) to 1. Hence, summing over  $\sigma \in \Sigma_{Q,n}$  in [\(2.21\)](#page-14-2) we get that, for  $h \in \mathcal{H}$ ,

$$
\sum_{\epsilon \Sigma_{Q,n}} \mu_{\Lambda;\beta,\varepsilon h}^+(\sigma) \le c_5^{-1} \exp(c_5 Q) \times \exp(-\beta \cdot 0.9Q)
$$

$$
= c_5^{-1} \exp(-(0.9\beta - c_5)Q). \tag{2.22}
$$

By Proposition [2.13,](#page-12-2) we have  $\Sigma_{Q,n} = \emptyset$  whenever  $Q < c_2 2^{n\theta}$ . Letting  $Q_{n,\ell} \coloneqq c_2 2^{n\theta+\ell}$ , we have  $\{\sigma \in \Omega_{\Lambda}^+ : |I_{\sigma}| = 2^n\} = \bigcup_{\ell=0}^{\infty} \Sigma_{Q_{n,\ell},n}$ . Therefore, from [\(2.22\)](#page-15-2) we get that, for  $h \in \mathcal{H}$ ,

$$
\sum_{\sigma: |I_{\sigma}|=2^n} \mu_{\Lambda;\beta,\varepsilon h}^+(\sigma) \le \sum_{\ell=0}^\infty c_5^{-1} \exp(-(0.9\beta - c_5)Q_{n,\ell})
$$
  

$$
\le C_1^{-1} \exp(-C_1(0.9\beta - c_5)2^{n\theta}),
$$

where the second inequality holds under the assumption that  $0.9\beta - c_5 > 0$ . Recalling Lemma [2.6](#page-7-0) and summing over *n* we obtain

$$
\mu_{\Lambda;\beta,\varepsilon h}^+(\sigma_0 = -1) \le \sum_{n=0}^{\infty} C_1^{-1} \exp(-C_1(0.9\beta - c_5)2^{n\theta})
$$
  

$$
\le C_2^{-1} \exp(-C_2(0.9\beta - c_5)).
$$

<span id="page-15-0"></span>Combining this with  $(2.20)$ , we get the desired result by letting  $\beta > 0$  large enough.

<span id="page-15-2"></span> $\Box$ 

**Remark 2.22.** *Notice that Propositions* [2.13,](#page-12-2) [2.14](#page-12-3) *and* [2.20](#page-13-0) *hold for*  $1 < \alpha < 2$  *with*  $\delta = \frac{2-\alpha}{20}$  $and \ \theta = \min\{2 - \alpha - 10\delta, \log_{3.9}(2)\} > 0$ . Thus the long-range order without disorder in this regime *follows from a standard Peierls argument.*

**Remark 2.23.** *Notice that our argument does not rely on the state space being* {−1*,* 1}*, so our results can be extended to the q-Potts model with*  $q \geq 3$ *. One can define the notion of p-favored for*  $p = 1, \dots, q$ *, repeat the balancing procedure, and get the long-range order for the q-Potts model at a low enough temperature.*

### <span id="page-15-1"></span>**2.3 Energy bounds**

*σ*∈Σ*Q,n*

This subsection is dedicated to proving the energy estimation, namely, Propositions [2.13](#page-12-2) and [2.14.](#page-12-3) We state some auxiliary lemmas before proving Proposition [2.13.](#page-12-2)

<span id="page-15-3"></span>**Lemma 2.24.** *There exists a constant*  $\overline{c}_1 := \overline{c}_1(\alpha)$  *such that, for any interval*  $I \subset \mathbb{Z}$  *and any*  $\text{configuration } \sigma \in \Omega, \text{ if } \min\{|I^-(\sigma)|, |I^+(\sigma)|\} = m, \text{ then}$ 

$$
\sum_{x,y\in I} \mathbb{1}_{\{\sigma_x \neq \sigma_y\}} J_{xy} \geq \overline{c}_1 m^{2-\alpha}.
$$

*Proof.* We can assume without loss of generality that  $|I^-(\sigma)| \leq |I^+(\sigma)|$ . It is enough to show that, for all  $x \in I^-(\sigma)$ 

$$
\sum_{y \in I^+(\sigma)} J_{xy} \ge \overline{c}_1 m^{1-\alpha},\tag{2.23}
$$

since we can sum both sides over  $x \in I^-(\sigma)$  to get the desired bound. As we assumed  $m = |I^-(\sigma)| \le$  $|I^+(\sigma)|$ , it is obvious that  $\{|x-y| : y \in I^+(\sigma)\}$  is dominated by the set  $\{m+1,\ldots,2m\}$  (here the domination is in the sense of the domination between the empirical measures of these two sets), implying that

$$
\sum_{y \in \mathcal{I}^+(\sigma)} J_{xy} \ge \sum_{k=m+1}^{2m} k^{-\alpha} \ge \overline{c}_1 m^{1-\alpha}.
$$

<span id="page-16-4"></span>**Corollary 2.25.** *For any*  $A \subset \mathbb{Z}$ *,* 

$$
J(A, A^c) \ge \overline{c}_1 |A|^{2-\alpha}.
$$

*Proof.* Let I be the smallest interval with  $A \subset I$  and  $|I| \ge 2|A|$ . Applying Lemma [2.24](#page-15-3) for  $\sigma$  with  $\sigma_x = \mathbb{1}_A(x) - \mathbb{1}_{A^c}(x)$  vields the desired result (note that  $|I^{\pm}(\sigma)| > |A|$ ).  $\sigma_x = \mathbb{1}_A(x) - \mathbb{1}_{A^c}(x)$  yields the desired result (note that  $|\mathcal{I}^{\pm}(\sigma)| \geq |A|$ ).

When there are very few minuses or very few pluses in an interval, the lower bound on Lemma [2.24](#page-15-3) is not very useful. To get a better bound, we need to take advantage of the balanced property. To this end, it will be useful to approximate arbitrary intervals by *ℓ*-intervals, as carried out in the next lemma.

<span id="page-16-2"></span>**Lemma 2.26.** *Let* I *be any interval. There exists*  $\ell \geq 0$  *and an*  $\ell$ -interval  $I_{\ell} = [a, b)$  *such that*  $I \subset I_\ell$  *and*  $\max \{d(I, a), d(I, b)\} \leq 0.7|I|$  *(see Figure [5](#page-16-1) for an illustration).* 

*Proof.* Take  $\ell \geq 0$  satisfying  $\frac{15}{8} 2^{\ell-2} \leq |I| \leq \frac{15}{8} 2^{\ell-1}$ . As  $|I| \leq 2^{\ell} - \frac{2^{\ell}}{16}$ , we can find an  $\ell$ -interval  $I_{\ell}(x)$ covering I. Supposing *x* is the smallest integer with  $I \subset I_{\ell}(x)$ , we can always find a translation  $I_{\ell}(x+n) = [a, b),$  with  $0 \leq n \leq 15$ , such that  $|d(a, 1) - d(b, 1)| \leq \frac{2^{\ell}}{16}$ . Thus, for this interval  $I_{\ell}(x+n)$ , we have  $d(a, I) \leq \frac{1}{2} \left(2^{\ell} + 2^{\ell-4} - \frac{15}{8} 2^{\ell-2}\right) \leq \frac{7}{10} |I|$ . A similar estimation holds for  $d(b, I)$ , concluding the proof.

<span id="page-16-1"></span>

Figure 5: The gray region delimits the interval I. We take  $\ell$  satisfying  $\frac{15}{8}2^{\ell-2} \leq |I| \leq \frac{15}{8}2^{\ell-1}$ . The black line represents  $\mathbb Z$  divided into intervals of length  $2^{\ell}/16$ . The  $\ell$ -interval  $I_{\ell}$  with endpoints in the red regions satisfies the desired inequality.

In Lemma [2.29,](#page-17-0) we will show that there is a restriction on the distance between intervals covering the negative spins. To describe that precisely we use the next definition and lemma.

<span id="page-16-3"></span><span id="page-16-0"></span>**Definition 2.27.** Given  $\lambda > 0$ , we say that a sequence  $\mathscr{P} = (p_i)_{i=1}^N \subset \{0,1\}^N$  is  $\lambda$ -good, if for any interval  $I \subsetneq [1, N]$  with  $\max_{i \in I} p_i = 1$ , there exists  $x \in [1, N] \setminus I$  with  $p_x = 1$  and  $d(x, I) \leq \lambda |I| + 1$ . **Lemma 2.28.** Let  $N \geq 1$  and  $\mathscr{P} = (p_i)_{i=1}^N$  be an  $\lambda$ -good sequence with  $p_1 = p_N = 1$ , Then,

$$
|\{1 \le i \le N : p_i = 1\}| \ge N^{\log_{\lambda+2}(2)}
$$
.

*Proof.* Our proof is by induction. If  $N = 1$  then  $p_1 = 1$  and the desired result holds. Then we assume the lemma holds for any  $N \leq M - 1$  (for some  $M \geq 2$ ) and we next prove it for  $N = M$ . Let  $I = [n, m]$  be the largest interval such that  $\max_{i \in I} p_i = 0$ . We now prove that  $\mathscr{P}_{[1,n)} \coloneqq (p_i)_{i=1}^n$ and  $\mathscr{P}_{(m,M)} := (p_i)_{i=m}^M$  are *λ*-good. Take any interval  $B \subsetneq [1,n)$  with  $\max_{i \in B} p_i = 1$ , and take  $x \in [1, M] \setminus B$  with  $p_x = 1$  such that  $d(x, B) = d(B, \{y \in [1, M] \setminus B : p_y = 1\}).$  When there are two possible choices for x, we take the one on the left. As  $\mathscr P$  is  $\lambda$ -good, we already know that  $d(x, B) \leq \lambda |B| + 1$ , by the definition of  $\lambda$ -good, so we just need to show that  $x \in [1, n) \setminus B$ . Otherwise, we have  $x = m + 1$ . Since  $p_1 = p_{n-1} = 1$  ( $p_{n-1} = 1$  follows from the maximality of I), by the definition of *x* and the assumption that  $x = m + 1$ , we can assume that  $B = [b, n)$  for some *b* > 1. Moreover, as  $d(x, B) = |I| + 1$ , we must have  $\max_{i \in [b-|I|-2, b)} p_i = 0$  (by  $p_1 = 1$ ), contradicting the fact that  $|I|$  is the largest interval containing only 0's. The same argument shows that  $\mathscr{P}_{(m,M]}$  is  $\lambda$ -good, and therefore, by the induction hypothesis

$$
\{x \in [1, M] : p_x = 1\} = |\{x \in [1, n) : p_x = 1\}| + \{x \in (m, M] : p_x = 1\}
$$
  
\n
$$
\ge (n - 1)^{\log_{\lambda + 2}(2)} + (M - m - 1)^{\log_{\lambda + 2}(2)}.
$$
\n(2.24)

Denoting  $N_1 = n - 1$  and  $N_2 = M - m - 1$ , we can assume without loss of generality that  $N_1 \leq N_2$ . As  $\mathscr P$  is  $\lambda$ -good and  $[1,n) \subsetneq [1,N]$  satisfies  $\max_{i\in[1,n]} p_i = 1$ , we must have  $|I| =$  $d([1, n), (m, M]) - 1 \leq \lambda N_1$ . Hence  $M \leq N_1(1 + \lambda) + N_2$  and  $(2 + \lambda)N_2 \geq M$ . Plugging this into [\(2.24\)](#page-17-1) we get

$$
\{x \in [1, N] : p_x = 1\} \ge N_2^{\log_{\lambda+2}(2)} + (\frac{M - N_2}{\lambda + 1})^{\log_{\lambda+2}(2)} \ge 2\left(\frac{M}{\lambda + 2}\right)^{\log_{\lambda+2}(2)} = M^{\log_{\lambda+2}(2)},
$$

where in the last inequality we used the fact that  $N_2^{\log_{\lambda+2}(2)} + (\frac{M-N_2}{\lambda+1})^{\log_{\lambda+2}(2)}$  increases in  $N_2$  when  $N_2 \geq \frac{M}{2+\lambda}$ .  $\Box$ 

<span id="page-17-0"></span>**Lemma 2.29.** *Recall our definition of*  $\theta$  *in* [\(2.11\)](#page-12-5)*. Let* I *be an interval, and let*  $\sigma \in \Omega$  *be a balanced configuration in*  $\rho_{\frac{3}{2}}(I)$ *. If*  $\sigma$  *is not constant on* I, then there *is a constant*  $\overline{c}_2 := \overline{c}_2(\alpha)$  *such that* 

<span id="page-17-1"></span>
$$
\sum_{x,y \in \rho_{\frac{3}{2}}(I)} \mathbb{1}_{\{\sigma_x \neq \sigma_y\}} J_{xy} \geq \overline{c}_2 |I|^{\theta}.
$$

*Proof.* To simplify the notation let us write  $\overline{I} := \rho_{\frac{3}{2}}(I)$ . Taking *N* such that  $2^N \leq |I| < 2^{N+1}$ , we want to show that

<span id="page-17-2"></span>
$$
J(\overline{1}^-(\sigma), \overline{1}^+(\sigma)) \ge \overline{c}_2 2^{N\theta + \theta}.
$$
\n(2.25)

Our proof is again by induction on *N*. When  $N \leq 1000 \log_2(M_0) + 1$ , [\(2.25\)](#page-17-2) holds trivially by taking  $\overline{c}_2$  sufficiently small depending on  $M_0$ . Now, given  $n > 1000 \log_2(M_0) + 1$ , we suppose that  $(2.25)$  holds for all intervals I with  $2^N \leq |I| < 2^{N+1}$  for all  $N < n$ . We will show that for I with  $2^n \leq |I| < 2^{n+1}$  the inequality [\(2.25\)](#page-17-2) also holds with  $N = n$ . We can assume that  $|I| = 2^n$ , since otherwise we take I' the subinterval of I with length  $2^n$  such that  $\sigma$  is not constant on I' and prove [\(2.25\)](#page-17-2) for I'. Furthermore, we can assume without loss of generality that  $I = [0, 2^n)$  and  $|\overline{\mathbf{I}}^-(\sigma)| \leq |\overline{\mathbf{I}}^+(\sigma)|.$ 

Considering the scale  $m = \lfloor n(1 - C_1\delta) - C_1 \log_2(M_0) \rfloor$  with  $C_1 = 500$ , we have  $2^{n-m} \ge M_n^{C_1}$ . By the assumption  $n > 1000 \log_2(M_0) + 1$ , we get that  $m > 0$ . For scales  $\ell \geq m$ , if there is a minus occupied  $\ell$ -interval  $I_{\ell} \subset I$ , we can apply Lemma [2.24](#page-15-3) to I and get [\(2.25\)](#page-17-2).

We can thus assume all  $\ell$ -intervals  $I_{\ell} \subset I$  with  $\ell \geq m$  are minus vacant. Under this assumption, if  $B \subset I$  is an  $\ell$ -interval containing a minus with  $\ell \geq m$  and  $\rho_{M_{\ell}}(I_{\ell}) \subset I$ , then there exists  $x \in \overline{I}^{-}(\sigma) \cap B^{c}$ such that  $d(x, B) \leq 2^{\ell-1}$ ; this is because B cannot be plus favored due to the balanced assumption for I.

Let  $\{B_1, \ldots, B_k\} \subset \mathcal{I}_m^0$  be a covering of  $\overline{\mathcal{I}}^-(\sigma)$  by disjoint *m*-intervals, such that each  $B_i$  (for  $1 \leq i \leq k$ ) contains a minus spin. We want to use Lemma [2.28](#page-16-0) to give a lower bound on k. Let  $\mathcal{B} = \bigcup_{i=1}^{k} B_i$  and let I<sub>B</sub> be the smallest interval that contains B. We claim that  $|I_B| \ge \frac{2^n}{5M}$  $\frac{2^n}{5M_n}$ . Otherwise, by Lemma [2.26,](#page-16-2) there exists an interval  $D = [a, b) \in \mathcal{I}$  containing  $I_B$  such that  $|D| \leq$  $d(I_B, a) + d(I_B, b) + |I_B| \leq 2.4|I_B| \leq 2.4 \cdot \frac{2^n}{5M}$  $\frac{2^n}{5M_n}$ . Moreover,  $\rho_{M_n}(D) \subset \overline{I}$  (since  $M_n|D| < 2^{n-1}$ ) and  $\sigma|_{\overline{I}\setminus I_B} = 1$ , by the definition of *B*. Thus D is a plus isolated interval in  $\sigma$  which contradicts the balanced assumption for I, yielding that  $|I_{\mathcal{B}}| \ge \frac{2^n}{5M_n}$ .

Taking  $K := |I_{\mathcal{B}}|/2^m$ , we partition  $I_{\mathcal{B}} = \cup_{i=1}^K C_i$  with  $\{C_1, \ldots, C_K\} \subset \mathcal{I}_{m}^0$  being a collection of disjoint *m*-intervals. Notice that  $\{B_1, \ldots, B_k\} \subset \{C_1, \ldots, C_K\}$ . We then define  $\mathscr{P}_{\mathcal{B}} = (p_i)_{i=1}^K$  by taking  $p_i = 1$  when  $C_i \in \{B_1, \ldots, B_k\}$  and  $p_i = 0$  otherwise. We want to show that this sequence is 1.9-good (recall Definition [2.27\)](#page-16-3). Take any interval  $[k_1, k_2] \subset [1, K]$  with  $\max_{k_1 \leq i \leq k_2} p_i = 1$ . Let  $C = \bigcup_{i=k_1}^{k_2} C_i$  be the union of *m*-intervals associated to  $(p_i)_{i=k_1}^{k_2}$ . Let  $L \geq m$  be such that  $2^L \leq |C| \leq 2^{L+1}.$ 

By Lemma [2.26,](#page-16-2) there exists  $D = [a, b)$ , either an  $(L+1)$ -interval or an  $(L+2)$ -interval, such that  $\max \{d(a, C), d(b, C)\} \leq 0.7|C|$  and  $C \subset D$ . Since C contains a minus spin, so does D. In addition, since all *ℓ*-intervals in I with  $\ell \geq m$  are assumed to be minus vacant, there must be a minus spin  $x \in$  $\mathbb{Z}\setminus D$  with  $d(x,D) \leq \frac{|D|}{2}$ , as otherwise D would be isolated (so in particular,  $x \in \mathcal{B}\setminus C$ ). Thus, by the triangle inequality we have  $d(C, \mathcal{B} \setminus C) \le \max\{d(a, C), d(b, C)\} + d(D, \mathcal{B} \setminus C) \le 0.7|C| + \frac{|D|}{2} \le 1.9|C|.$ This shows in particular that there exists  $y \in [1, K]$  with  $p_y = 1$  and  $d(y, [k_1, k_2]) \leq 1.9(k_1 - k_2)$ , implying that  $\mathscr{P}_\mathcal{B}$  is a 1.9-good sequence. Therefore, Lemma [2.28](#page-16-0) gives us

$$
k = |\{1 \le i \le K : p_i = 1\}| \ge \left(\frac{|I_B|}{2^m}\right)^{\log_{3.9}(2)} \ge \left(\frac{2^{n-m}}{5M_n}\right)^{\log_{3.9}(2)},
$$

where in the last inequality we used the lower bound  $|I_B| \geq \frac{2^n}{5M}$  $\frac{2^n}{5M_n}$ . Let  $B_i = \rho_{\frac{3}{2}}(B_i)$ . As the lemma holds for each B<sub>i</sub> by the induction hypothesis, we use the fact that  $\overline{B}_i \cap \overline{B}_{i+2} = \emptyset$  to conclude

$$
J(\overline{1}^{-}(\sigma), \overline{1}^{+}(\sigma)) \geq \sum_{i=1}^{\lceil k/2 \rceil} J(\overline{B}_{2i-1}^{-}(\sigma), \overline{B}_{2i-1}^{+}(\sigma))
$$
  

$$
\geq \frac{k}{2} \cdot \overline{c}_{2} 2^{m\theta} \geq \frac{\overline{c}_{2}}{2 \cdot 5^{\log_{3.9}(2)}} \cdot 2^{n\theta} \cdot 2^{(n-m)(\log_{3.9}(2)-\theta)} \cdot M_{n}^{-\log_{3.9}(2)}
$$
  

$$
= \frac{\overline{c}_{2}}{2 \cdot 5^{\log_{3.9}(2)}} \cdot 2^{n\theta} \cdot M_{n}^{C_{1}(\log_{3.9}(2)-\theta) - \log_{3.9}(2)} \geq \overline{c}_{2} 2^{n\theta + \theta},
$$

where in the last inequality we used that  $C_1 (\log_{3.9}(2) - \theta) = 500(\log_{3.9}(2) - 0.5) \ge 1$  and chose  $M_0 > 0$  large enough depending on  $\theta$ .

We are ready to prove Proposition [2.13.](#page-12-2)

*Proof of Proposition [2.13](#page-12-2).* Since  $I_{\sigma}$  is plus favored with respect to  $\sigma^S$ , we see that  $\sigma^S$  is all plus in  $\rho_{3/2}(I_{\sigma}) \setminus I_{\sigma}$ . If  $A_{\sigma} = I_{\sigma}$  (i.e.,  $\sigma^{S}$  is all minus in  $I_{\sigma}$ ), then we must have  $|\{x \in \rho_{\frac{3}{2}}(A_{\sigma}) : \sigma_{x}^{S} =$  $|+1\rangle = |\{x \in \rho_{\frac{3}{2}}(A_{\sigma}) : \sigma_x^S = -1\}| = |A_{\sigma}|.$  We can therefore bound

$$
J(A_\sigma,A_\sigma^c)\geq \sum_{x,y\in \rho_{\frac32}(A_\sigma)}\mathbbm{1}_{\{\sigma_x^S\neq \sigma_y^S\}}J_{xy}\geq c_1'|{\rm I}_\sigma|^{2-\alpha},
$$

where the second inequality is due to Lemma [2.24.](#page-15-3)

If  $A_{\sigma} \neq I_{\sigma}$ , then  $\sigma^{S}$  is not a constant on  $I_{\sigma}$ . Since in addition  $I_{\sigma}$  is balanced with respect to  $\sigma^{S}$ , we can apply Lemma [2.29](#page-17-0) to get the desired bound. П

<span id="page-19-2"></span>We now want to prove Proposition [2.14.](#page-12-3) Again, we first introduce some auxiliary lemmas.

**Lemma 2.30.** For any configuration  $\sigma \in \Omega^+$ ,

$$
\sum_{\substack{x \in A_{\sigma} \\ y \in I_{\sigma}^c}} 1_{\{\sigma_x = \sigma_y^S = -1\}} J_{xy} \leq 0.1 J(A_{\sigma}, A_{\sigma}^c).
$$

*Proof.* To ease the notation we drop the subscript in  $A_{\sigma}$  and  $I_{\sigma}$  throughout the proof. Let  $n \geq 0$ and  $x \in \mathbb{Z}$  be such that  $I = I_n(x)$ . For every  $i \in \mathbb{Z}$ , take  $E_i := I_n(x + 16i)$ . Recalling that  $I_n(x) =$  $[2^{n-4}x - 2^{n-1}, 2^{n-4}x - 2^{n-1})$ , we see that  $I_n(x)$  and  $I_n(x \pm 16)$  are disjoint and thus  $\{E_i\}_{i \in \mathbb{Z}}$  is a partition of Z. For  $1 \le i \le M_n$ , as  $\sigma^S$  is plus favored with respect to I, we must have  $|E_i^-(\sigma^S)| \le \frac{2^n}{M_n}$ . Note that  $E_i$  is to the right-hand side of I. Letting  $E_i = [x_i, x_{i+1})$ , since the interaction is decreasing with respect to the distance, one can get that  $\min_{x \in \mathbb{E}_i^-} d(x, A) \ge d(x_i, A) \ge \max_{x \in \mathbb{E}_{i-1}^+} d(x, A)$  and

thus  $J(\mathbf{E}_i^-(\sigma^S), A) \leq \frac{2^n}{M_r}$  $\frac{2^n}{M_n} J(x_i, A)$  and  $J(\mathbf{E}_{i-1}^+(\sigma^S), A) \geq \left(2^n - \frac{2^n}{M_n}\right)$ *M<sup>n</sup>*  $J(x_i, A)$ . Thus we get for all  $1 < i \leq M_n$ 

<span id="page-19-1"></span>
$$
J(\mathbf{E}_{i}^{-}(\sigma^{S}), A) \leq \frac{1}{M_{n}-1} J(\mathbf{E}_{i-1}^{+}(\sigma^{S}), A).
$$

To get a similar bound for  $E_1(\sigma^S)$ , we notice that the first half of  $E_1$  must be all plus in  $\sigma^S$ , that is,  $E_1^+(\sigma^S) \supset [x_1, \frac{x_1+x_2}{2}]$ . This is, again, a consequence of I being plus favored. As the interaction decreases with respect to the distance, we get that  $J(E_1^-(\sigma^S), A) \leq \frac{2^n}{M_r}$  $\frac{2^n}{M_n} J(\frac{x_1+x_2}{2}, A)$  and  $J(E_1^+(\sigma^S), A) \geq 2^{n-1} J(\frac{x_1+x_2}{2}, A)$ , and therefore  $J(E_1^-(\sigma^S), A) \leq \frac{2}{M_n} J(E_1^+(\sigma^S), A)$ . We can thus bound

$$
\sum_{i=1}^{M_n} J(\mathbf{E}_i^-(\sigma^S), A) \le \frac{2}{M_n} J(\mathbf{E}_1^+(\sigma^S), A) + \sum_{i=1}^{M_n - 1} \frac{1}{M_n - 1} J(\mathbf{E}_i^+(\sigma^S), A)
$$
  

$$
\le \frac{3}{M_n - 1} J(\bigcup_{i=1}^{M_n} \mathbf{E}_i^+(\sigma^S), A). \tag{2.26}
$$

For the distant intervals  $E_i$  with  $i > M_n$ , as  $d(A, E_i) > M_0 2^{n(1+\delta)}$ , we can simply bound

<span id="page-19-0"></span>
$$
J(A, \bigcup_{i>M_n} E_i) \le |A| \sum_{R \ge M_0 2^{n(1+\delta)}} R^{-\alpha} \le CM_0^{1-\alpha} 2^{n(1+\delta)(1-\alpha)} |A|
$$
  

$$
\le CM_n^{1-\alpha} |A|^{2-\alpha}, \qquad (2.27)
$$

where the last inequality comes from  $|A| \leq 2^n$ . By Corollary [2.25,](#page-16-4) we have  $\overline{c}_1|A|^{2-\alpha} \leq J(A, A^c)$ . Combined with [\(2.27\)](#page-19-0), this yields

<span id="page-20-0"></span>
$$
J(A, \bigcup_{i>M_n} \mathcal{E}_i) \le \frac{C}{\overline{c}_1} M_n^{1-\alpha} J(A, A^c). \tag{2.28}
$$

By symmetry, we have the same estimations  $(2.26)$  and  $(2.28)$  for  $i < 0$ , and therefore

$$
\sum_{\substack{x \in A \\ y \in \Gamma^c}} 1_{\{\sigma_x = \sigma_y^S = -1\}} J_{xy} \le \frac{3}{M_n - 1} J(A, \bigcup_{1 \le |i| \le M_n} E_i^+(\sigma^S)) + \frac{2C}{\overline{c}_1} M_n^{1-\alpha} J(A, A^c).
$$

The desired bound then comes from choosing  $M_0$  large enough.

$$
\Box
$$

Next, we control the interaction between  $A^c_\sigma$  and the sites in  $A_\sigma$  that changed signs in the balancing procedure. Let us first introduce some definitions.

<span id="page-20-2"></span>**Definition 2.31.** Let  $F(-, \sigma) := \{x \in A_{\sigma} : \sigma_x = 1\}$  be the collection of spins flipped to minus in the balancing procedure. For each  $x \in F(-, \sigma)$ , let I<sup>x</sup> be the last interval containing x which flipped *x* to minus in the balancing procedure of  $\sigma$ . Let  $\mathscr{F}_{\ell}(-,\sigma) = {\{\mathbf{I}^x : x \in F(-,\sigma), |\mathbf{I}^x| = 2^{\ell}\}}.$ Given  $I \in \mathscr{F}_{\ell}(-,\sigma)$ , recall that  $s_I$  denotes the step in which I was selected. Therefore, we have  $F(-, \sigma) \subset \bigcup_{\ell \geq 0} F_{\ell}(-, \sigma)$ , where  $F_{\ell}(-, \sigma) := \bigcup_{I \in \mathscr{F}_{\ell}(-, \sigma)} I^{+}(\sigma^{s_{I}})$ . Similarly, we define  $F(+, \sigma) := \{x \in I\}$  $A^c_\sigma: \sigma_x = -1, \sigma_x^S = 1$  to be the collection of spins flipped to plus in the balancing procedure and define  $\mathscr{F}_{\ell}(+, \sigma) = \{I^x : x \in F(+, \sigma), |I^x| = 2^{\ell}\}.$ 

Next, we prove a few lemmas and the last two will be used in the proof of Proposition [2.14.](#page-12-3) The lemmas are stated for  $\mathscr{F}_{\ell}(-,\sigma)$ , but we stress that both Lemmas [2.33](#page-21-1) and [2.34](#page-21-0) hold for  $\mathscr{F}_{\ell}(+,\sigma)$ .

<span id="page-20-1"></span>**Lemma 2.32.** *Recall Definition* [2.4](#page-5-0) *and recall that*  $M'_{\ell} = 2 \left| \frac{M_{\ell}}{2c_1} \right|$  $\int$ *. Given σ* ∈ Ω<sup>+</sup>, let *n* ≥ 0 *be such that*  $|I_{\sigma}| = 2^n$ . Then, for all  $\ell < n$  and  $I_{\ell} \in \mathscr{F}_{\ell}(-, \sigma)$ , we have  $\rho_{\frac{M'_{\ell}}{2}}(I_{\ell}) \subset I_{\sigma}$ .

*Proof.* Let *x* be the center of  $I_{\ell}$ , and thus we have  $I_{\ell} = I_{\ell}(x)$ . Let  $B_k := I_{\ell}(x + 16k)$  for all  $1 \leq |k| \leq M'_{\ell}$ . Supposing there exists  $1 \leq |k| \leq \frac{M'_{\ell}}{2}$  such that  $B_k \cap I_{\sigma}^c \neq \emptyset$ , we next derive a contradiction. Without loss of generality, we can assume that *k* has the smallest absolute value. Our derivation of the contradiction is divided into the following two cases.

If  $\ell < n-1$ , since  $B_k \cap I_{\sigma}^c \neq \emptyset$  we must have  $B_{k'} \subset \rho_{\frac{3}{2}}(I_{\sigma}) \setminus I_{\sigma}$ , where  $k' = \text{sign}(k) \cdot (|k| + 1)$ and thus  $B_{k'}$  is the neighboring interval of  $B_k$  which is further away from  $I_{\sigma}$ . As  $I_{\sigma}$  is plus favored with respect to  $\sigma^S$ , all spins of  $\sigma^S$  in  $\rho_{\frac{3}{2}}(I_{\sigma}) \setminus I_{\sigma}$  are plus, and so are all spins in  $B_{k'}$ . However, by Proposition [2.12,](#page-11-2) I<sub>ℓ</sub> is weakly minus favored, and thus  $B_{k'}$  must be weakly minus dense, arriving at a contradiction.

If  $\ell = n - 1$ , then we have  $\rho_{\frac{3}{2}}(I_{\sigma}) \setminus I_{\sigma} \subset B_{-2} \cup B_{-1} \cup B_1 \cup B_2$ . Recall that all spins of  $\sigma^{S}$ in  $\rho_{\frac{3}{2}}(I_{\sigma}) \setminus I_{\sigma}$  are plus. Recalling also that  $I_{\ell}$  is weakly minus favored with respect to  $\sigma^{S}$  (see Proposition [2.12\)](#page-11-2), we see that  $B_i$  must be weakly minus dense for any  $i = -2, -1, 1, 2$ . Hence the number of pluses of  $\sigma^S$  in B<sub>2</sub> ∪ B<sub>1</sub> ∪ B<sub>−1</sub> ∪ B<sub>−2</sub> is at most  $4 \cdot \frac{2^{\ell}}{M_{\ell}^{\ell}}$ . We have again arrived at a contradiction since  $4 \cdot \frac{2^{\ell}}{M_{\ell}} < 2^{n}$  (and that  $\rho_{3/2}(I_{\sigma}) \setminus I_{\sigma}$  is all plus).  $\Box$ 

Provided with Lemma [2.32,](#page-20-1) we can prove the following lemma.

<span id="page-21-1"></span>**Lemma 2.33.** *Given*  $\sigma \in \Omega^+$  *and*  $I \in \mathscr{F}_\ell(-,\sigma)$ *, let*  $\tilde{I} := \rho_{\frac{M'_\ell}{2}}(I)$ *. For any*  $B \subset \tilde{I}^c$  *we have* 

$$
J(\mathcal{I}^+(\sigma^{s_{\mathcal{I}}}),A^c_\sigma \cap B) \leq \frac{4}{M'_\ell} J(A_\sigma \cap \widetilde{\mathcal{I}},A^c_\sigma \cap B) \frac{|\mathcal{I}^+(\sigma^{s_{\mathcal{I}}})|}{2^\ell}.
$$

*An analogous result holds for*  $I \in \mathcal{F}_{\ell}(+, \sigma)$  *where in the above*  $I^+$  *is replaced with*  $I^-$ *.* 

*Proof.* To ease the notation, we drop the subscript in  $A_{\sigma}$  in the proof. For every  $y \in A^c \cap B$ , we wish to lower-bound  $J(A \cap I, y)$ . Take a, b such that  $I = [a, b)$ . We assume  $y \ge b$ , and the argument for  $y < a$  follows by symmetry. Let  $x \in \mathbb{Z}$  be such that  $I = I_{\ell}(x)$ . By Proposition [2.12,](#page-11-2) I is weakly minus favored with respect to  $\sigma^S$ . So, for all  $1 \leq k \leq M'_{\ell}$ , we have  $I_{\ell}(x + 16k)$  is weakly minus dense. Moreover, by Lemma [2.32](#page-20-1) and Definition [2.4,](#page-5-0) we get that  $\{w \in I_\ell(x + 16k) : \sigma_w^S = -1\} \subset A$ for every  $1 \leq |k| \leq \frac{M'_{\ell}}{2}$ . Therefore we have

$$
|\{z \in A \cap \tilde{1} : z \ge b\}| \ge \sum_{1 \le k \le \frac{M'_\ell}{2}} |\{w \in I_\ell(x + 16k) : \sigma_w^S = -1\}|
$$
  

$$
\ge \left(1 - \frac{1}{M'_\ell}\right) \frac{M'_\ell}{2} \cdot 2^\ell \ge \frac{M'_\ell 2^\ell}{4}.
$$

In addition, since the interaction decreases with respect to the distance, we have  $J(A \cap \tilde{I}, y) \geq$  $(y - b + 1)^{-\alpha} \frac{M_{\ell}' 2^{\ell}}{4}$  $\frac{d^2}{4}$ . Similarly we can bound  $J(y, I^+(\sigma^{s_1})) \le (y - b + 1)^{-\alpha} |I^+(\sigma^{s_1})|$  and therefore

$$
J(A \cap \widetilde{I}, y) \ge \frac{J(y, I^+(\sigma^{s_1}))}{|I^+(\sigma^{s_1})|} \frac{M_{\ell}' 2^{\ell}}{4}
$$

Summing over  $y \in A^c \cap B$  we get the desired bound.

For the interaction within  $\rho_{\frac{M'_{\ell}}{2}}(I)$ , we need a tighter control. Given  $I = I_{\ell}(x) \in \mathscr{F}_{\ell}(-,\sigma)$ , for each  $1 \leq |k| \leq \frac{M'_{\ell}}{2}$ , let us denote  $I^k := I_{\ell}(x + 16k)$ .

<span id="page-21-0"></span>**Lemma 2.34.** *There exists a constant*  $\overline{c}_3 := \overline{c}_3(\alpha) > 0$  *such that, for any*  $\sigma$ *,* I  $\in \mathscr{F}_{\ell}(-, \sigma)$  *and*  $1 \leq |k| \leq \frac{M'_{\ell}}{2},$ 

$$
J(\mathrm{I}^+(\sigma^{s_1}), \mathrm{I}^k \cap A^c_\sigma) \leq \overline{c}_3(M'_\ell)^{1-\alpha} J(\mathrm{I}^k \cap A_\sigma, \mathrm{I}^k \cap A^c_\sigma) \frac{|\mathrm{I}^+(\sigma^{s_1})|}{2^\ell}.
$$

*An analogous result holds for*  $I \in \mathcal{F}_{\ell}(+, \sigma)$ *, where in the above*  $I^+$  *is replaced with*  $I^-$ *.* 

*Proof.* For notation clarity, in this proof we drop the subscript in  $A_{\sigma}$ . As I is weakly minus favored with respect to  $\sigma^S$  (by Proposition [2.12\)](#page-11-2), all its first  $2^{\ell-1}$  neighbors are minus, hence  $d(\mathbf{I}, \mathbf{I}^k \cap A^c_{\sigma}) \geq 2^{\ell-1}$ . Thus, on the one hand, we have

<span id="page-21-2"></span>
$$
J(\mathbf{I}^+(\sigma^{s_1}), \mathbf{I}^k \cap A^c) \le (2^{\ell-1})^{-\alpha} |\mathbf{I}^+(\sigma^{s_1})| |\mathbf{I}^k \cap A^c|.
$$
 (2.29)

*.*

On the other hand, as  $I^k$  is weakly plus vacant with respect to  $\sigma^S$  (recalling that I is weakly minus favored), we have  $|I_k \cap A^c| = |\{x \in I^k : \sigma_x^S = 1\}| \leq \frac{1}{M'_\ell} 2^{\ell}$ . As this is smaller than  $|I^k|/2$ , we can apply Lemma [2.24](#page-15-3) to get  $J(I_k \cap A^c, I_k \cap A) \ge \overline{c}_1 | I_k \cap A^c |^{2-\alpha}$ . Combined with [\(2.29\)](#page-21-2), it yields that

$$
J(\mathbf{I}^+(\sigma^{s_1}), \mathbf{I}^k \cap A^c) \le (2^{\ell-1})^{-\alpha} |\mathbf{I}^+(\sigma^{s_1})| |\mathbf{I}^k \cap A^c_\sigma| \frac{J(\mathbf{I}_k \cap A^c, \mathbf{I}_k \cap A)}{\overline{c}_1 |\mathbf{I}_k \cap A^c|^{2-\alpha}}
$$

$$
\Box
$$

$$
\leq \frac{2^{\alpha}}{\overline{c}_1} J(I_k \cap A^c, I_k \cap A) \frac{|I^+(\sigma^{s_1})|}{2^{\ell}} \left( \frac{|I_k \cap A^c|}{2^{\ell}} \right)^{\alpha - 1}
$$
  

$$
\leq \frac{2^{\alpha}}{\overline{c}_1} J(I_k \cap A^c, I_k \cap A) \frac{|I^+(\sigma^{s_1})|}{2^{\ell}} (M'_{\ell})^{1 - \alpha}.
$$

We are ready to prove Proposition [2.14.](#page-12-3)

*Proof of Proposition [2.14](#page-12-3).* Using the identity  $\sigma_x \sigma_y = 21_{\{\sigma_x = \sigma_y\}} - 1$  for all  $x, y \in \mathbb{Z}$ , we can write

<span id="page-22-1"></span>
$$
H_{\Lambda}^{+}(\sigma) - H_{\Lambda}^{+}(\tau_{A_{\sigma}}(\sigma)) = - \sum_{x \in A_{\sigma}, y \in A_{\sigma}^{c}} 2J_{xy}\sigma_{x}\sigma_{y} = 2J(A_{\sigma}, A_{\sigma}^{c}) - 4 \sum_{x \in A_{\sigma}, y \in A_{\sigma}^{c}} J_{xy} \mathbb{1}_{\{\sigma_{x} = \sigma_{y}\}}.
$$
 (2.30)

We then split  $\mathbb{1}_{\{\sigma_x=\sigma_y\}}$  according to the signs of the original configuration and the final configuration  $\sigma^S$ , getting that

$$
\mathbb{1}_{\{\sigma_x = \sigma_y\}} = \mathbb{1}_{\{\sigma_x = \sigma_y = \sigma_y^S = -1\}} + \mathbb{1}_{\{\sigma_x = \sigma_y = -\sigma_y^S = -1\}} + \mathbb{1}_{\{\sigma_x = \sigma_y = 1\}} \leq \mathbb{1}_{\{\sigma_x = \sigma_y = \sigma_y^S = -1\}} + \mathbb{1}_{\{\sigma_x = \sigma_y = -\sigma_y^S = -1\}} + \mathbb{1}_{\{\sigma_x = 1\}}.
$$
\n(2.31)

Recall that  $I_{\sigma}$  is the *n*-interval such that  $A_{\sigma} = I_{\sigma}^{-}(\sigma^{S})$ . Recalling also Definition [2.31,](#page-20-2) we have  ${x \in A_{\sigma} : \sigma_x \neq \sigma_x^S} = F(-, \sigma)$  and  ${y \in A_{\sigma}^c : -\sigma_y = \sigma_y^S = 1} = F(+, \sigma)$ . With this, we can plug  $(2.31)$  into  $(2.30)$  and get that

$$
H_{\Lambda}^{+}(\sigma) - H_{\Lambda}^{+}(\tau_{A_{\sigma}}(\sigma)) \ge 2J(A_{\sigma}, A_{\sigma}^{c}) - 4\sum_{\substack{x \in A_{\sigma} \\ y \in \Gamma_{\sigma}^{c}}} \mathbb{1}_{\{\sigma_{x} = \sigma_{y} = \sigma_{y}^{S} = -1\}} J_{xy}
$$

$$
- 4J(F(-, \sigma), A_{\sigma}^{c}) - 4J(A_{\sigma} \setminus F(-, \sigma), F(+, \sigma)). \quad (2.32)
$$

The sum on the right-hand side of [\(2.32\)](#page-22-2) is only over  $y \in I_{\sigma}^c$  since all spins in  $I_{\sigma} \setminus A_{\sigma}$  are plus. We claim that it suffices to prove the following:

<span id="page-22-3"></span><span id="page-22-2"></span><span id="page-22-0"></span>
$$
J(F(-, \sigma), A^c_{\sigma}) \le \frac{C}{M_0} J(A_{\sigma}, A^c_{\sigma}),
$$
\n(2.33)

<span id="page-22-4"></span>
$$
J(F(+,\sigma), A_{\sigma}) \le \frac{C}{M_0} J(A_{\sigma}, A_{\sigma}^c),
$$
\n(2.34)

for an appropriate constant  $C > 0$  depending only on  $\alpha$  and  $\delta$ . Indeed, combining [\(2.33\)](#page-22-3) and [\(2.34\)](#page-22-4) with Lemma  $2.30$  (which bounds the second term on the right-hand side of  $(2.32)$ ), we get that

$$
H_{\Lambda}^{+}(\sigma) - H_{\Lambda}^{+}(\tau_{A_{\sigma}}(\sigma)) \geq (2 - 4/10 - \frac{2C}{M_0})J(A_{\sigma}, A_{\sigma}^{c}),
$$

and the desired result follows by taking  $M_0$  large enough.

We now prove [\(2.33\)](#page-22-3). Recall Definition [2.31.](#page-20-2) Moreover, we can restrict each  $\mathscr{F}_{\ell}(-,\sigma)$  to disjoint intervals by taking  $\mathscr{F}_{\ell,i}(-,\sigma) := \mathscr{F}_{\ell}(-,\sigma) \cap \mathcal{I}_{\ell}^{i}$ , for  $i = 0, \ldots, 15$ , where  $\mathcal{I}_{\ell}^{i}$  was defined in [\(2.1\)](#page-4-2). We

can also write  $F_{\ell}(-, \sigma) \subset \bigcup_{i=1}^{15} F_{\ell,i}(-, \sigma)$  where  $F_{\ell,i}(-, \sigma) := \bigcup_{I \in \mathscr{F}_{\ell,i}(-, \sigma)} I^+( \sigma^{s_I}),$  for  $i = 0, \ldots, 15$ . Then,

<span id="page-23-4"></span>
$$
J(F(-, \sigma), A^c_{\sigma}) \le \sum_{\ell=0}^n \sum_{i=0}^{15} J(F_{\ell, i}(-, \sigma), A^c_{\sigma}) \le \sum_{\ell=0}^n \sum_{i=0}^{15} \sum_{B \in \mathcal{I}^i_{\ell}} J(F_{\ell, i}(-, \sigma), A^c_{\sigma} \cap B). \tag{2.35}
$$

Fixing  $B \in \mathcal{I}_{\ell}^i$ , we now upper-bound  $J(F(-, \sigma), A_{\sigma}^c \cap B)$ . Recall that for any  $I \in \mathscr{F}_{\ell}(-, \sigma)$ , we denote  $\widetilde{I} = \rho_{M'_\ell}(I)$ . As  $\mathcal{I}^i_\ell$  forms a disjoint partition of  $\mathbb{Z}$ , for all  $I_\ell \in \mathscr{F}_{\ell,i}(-,\sigma)$  either  $B \subset \widetilde{I}_\ell$  or  $B \cap \widetilde{I}_{\ell} = \emptyset.$ 

We first consider  $I_{\ell} \in \mathscr{F}_{\ell}(-,\sigma)$  with  $B \cap I_{\ell} = \emptyset$ . As in the proof of Lemma [2.9,](#page-8-0) we say  $I_{\ell} \in \mathscr{F}_{\ell}(-,\sigma)$  is *negligible* when  $|I_{\ell}^{+}(\sigma^{s_{I_{\ell}}})| \leq \frac{2^{\ell}}{M'_{\ell}}$ , i.e., not many spins were flipped to minus. Otherwise, we say I*<sup>ℓ</sup>* is *non-negligible*. Similarly to [\(2.8\)](#page-11-3), we want to lower-bound the distance between two non-negligible intervals. We cannot simply use [\(2.8\)](#page-11-3) since it was proved for intervals in  $\mathcal{F}_{\ell}$ , not in  $\mathcal{F}_{\ell}(-,\sigma)$  as we need. Next we show that two non-negligible intervals  $I_{\ell}, I'_{\ell}$  must satisfy

<span id="page-23-0"></span>
$$
d(\mathbf{I}_{\ell}, \mathbf{I}_{\ell}') \ge (M_{\ell}' - 1)2^{\ell}.
$$
\n
$$
(2.36)
$$

Indeed, for two non-negligible intervals  $I_{\ell}, I'_{\ell}$ , let *s*, *s'* be the step in which they were flipped respectively. We can assume without loss of generality that  $s < s'$ , i.e., I<sub>ℓ</sub> was flipped first. We get from Proposition [2.12](#page-11-2) that  $I_{\ell}$  is weakly minus favored with respect to  $\sigma^{s'}$ . If  $d(I_{\ell}, I'_{\ell}) \leq (M'_{\ell} - 1)2^{\ell}$ , then  $|(I'_{\ell})^{-}(\sigma^{s'})| \geq 2^{\ell} - \frac{2^{\ell}}{M_{\ell}}$  $\frac{2^{\infty}}{M'_{\ell}}$ , which contradicts to the fact that B<sub>j</sub> is non-negligible. This implies, in particular, that  $\widetilde{I}_{\ell} \cap \widetilde{I}'_{\ell} = \emptyset$ . By Lemma [2.33,](#page-21-1) when  $B \cap \widetilde{I}_{\ell} = \emptyset$  we have  $J(I_{\ell}^{+}(\sigma^{s_{I_{\ell}}}), A_{\sigma}^{c} \cap B) \leq$  $\frac{4}{M'_\ell} J(\tilde{\mathfrak{l}}_\ell \cap A_\sigma, A_\sigma^c \cap \mathcal{B})$ . Summing over non-negligible intervals we get

<span id="page-23-1"></span>
$$
\sum_{\substack{\mathbf{I}_{\ell} \in \mathscr{F}_{\ell,i}(-,\sigma), \ \mathbf{B} \cap \mathbf{I}_{\ell} = \emptyset}} J(\mathbf{I}_{\ell}^{+}(\sigma^{s_{\mathbf{I}_{\ell}}}), A_{\sigma}^{c} \cap \mathbf{B}) \le \frac{4}{M'_{\ell}} J(A_{\sigma}, A_{\sigma}^{c} \cap \mathbf{B}).
$$
\n(2.37)

For negligible  $I_{\ell} \in \mathscr{F}_{\ell,i}^{-}$ , by Lemma [2.33](#page-21-1) we have

$$
J(\mathcal{I}^+_\ell(\sigma^{s_{\mathcal{I}_\ell}}),A^c_\sigma \cap \mathcal{B}) \leq \frac{4}{M'_\ell}J(\widetilde{\mathcal{I}}_\ell \cap A_\sigma,A^c_\sigma \cap \mathcal{B})\frac{|\mathcal{I}^+_\ell(\sigma^{s_{\mathcal{I}}_\ell})|}{2^\ell} \leq \frac{4}{M'_\ell}J(\widetilde{\mathcal{I}}_\ell \cap A_\sigma,A^c_\sigma \cap \mathcal{B})\frac{1}{M'_\ell}.
$$

In this case, not all  $I_\ell$ 's are disjoint, but every  $x \in I_\ell$  can be contained in at most  $M'_\ell$  other intervals  $\widetilde{\mathbf{I}}'_{\ell}$  with  $\mathbf{I}'_{\ell} \in \mathscr{F}_{\ell}(-, \sigma)$ . Therefore,

<span id="page-23-2"></span>
$$
\sum_{\substack{\mathbf{I}_{\ell} \in \mathscr{F}_{\ell,i}(-,\sigma), \ \mathbf{B} \cap \mathbf{I}_{\ell} = \emptyset}} J(\mathbf{I}_{\ell}^{+}(\sigma^{s_{\mathbf{I}_{\ell}}}), A_{\sigma}^{c} \cap \mathbf{B}) \leq \frac{4}{M'_{\ell}} J(A_{\sigma}, A_{\sigma}^{c} \cap \mathbf{B}).
$$
\n(2.38)

We now consider the interval  $I_\ell \in \mathscr{F}_{\ell,i}(-,\sigma)$  with  $B \subset \tilde{I}_\ell$ . If  $I_\ell$  is negligible, Lemma [2.34](#page-21-0) yields  $J(I_{\ell}^+(\sigma^{s_1}_{\ell}), A_{\sigma}^c \cap B) \leq \overline{c}_3(M_{\ell}')^{-\alpha} J(B \cap A_{\sigma}, B \cap A_{\sigma}^c)$ . Summing over all  $M_{\ell}'$  possible such intervals we get

<span id="page-23-3"></span>
$$
\sum_{\substack{\mathbf{I}_{\ell} \in \mathscr{F}_{\ell,i}(-,\sigma), \ \mathbf{B} \subset \mathbf{I}_{\ell} \\ \mathbf{I}_{\ell} \ \text{negligible}}} J(\mathbf{I}_{\ell}^{+}(\sigma^{s_{\mathbf{I}_{\ell}}}), A_{\sigma}^{c} \cap \mathbf{B}) \leq \overline{c}_{3}(M_{\ell}')^{1-\alpha} J(\mathbf{B} \cap A_{\sigma}, \mathbf{B} \cap A_{\sigma}^{c}).
$$
\n(2.39)

We note that at most one interval  $I_\ell \in \mathscr{F}_{\ell,i}(-,\sigma)$  with  $B \subset I_\ell$  can be non-negligible since all non-negligible intervals are at least  $(M'_{\ell}-1)2^{\ell}$  distance away from each other (see [\(2.36\)](#page-23-0)). For this interval Lemma [2.34](#page-21-0) again yields  $J(I_{\ell}^+(\sigma^{s_1}_{\ell}), A_{\sigma}^c \cap B) \leq \overline{c}_3(M_{\ell}')^{1-\alpha} J(B \cap A_{\sigma}, B \cap A_{\sigma}^c)$ . Together with [\(2.37\)](#page-23-1), [\(2.38\)](#page-23-2) and [\(2.39\)](#page-23-3), this yields that

$$
\sum_{\mathbf{B}\in\mathcal{J}_{\ell}^{i}} J(F_{\ell,i}(-,\sigma), A_{\sigma}^{c}\cap \mathbf{B}) \leq (64 + 2\overline{c}_{3})(M_{\ell}')^{1-\alpha}J(A_{\sigma}, A_{\sigma}^{c}).
$$

Plugging the preceding inequality into [\(2.35\)](#page-23-4) we get [\(2.33\)](#page-22-3). To prove [\(2.34\)](#page-22-4) we proceed similarly. We first get an expression like [\(2.35\)](#page-23-4), with  $F(-, \sigma)$  replaced by  $F(+, \sigma)$  and with  $\mathcal{F}_{\ell,i}(-, \sigma)$  replaced by  $\mathcal{F}_{\ell,i}(+,\sigma)$ . To upper-bound  $J(F(-,\sigma), A^c_{\sigma} \cap B)$ , for  $B \in \mathcal{I}_{\ell}^i$ , we now proceed in the same manner as that for  $(2.33)$ ; this is valid since to prove inequalities  $(2.37)$ ,  $(2.38)$  and  $(2.39)$  with  $J(I_{\ell}^+(\sigma^{s_1}_{\ell}), A_{\sigma}^c \cap B)$  replaced by  $J(I_{\ell}^-(\sigma^{s_1}_{\ell}), A_{\sigma} \cap B)$  we would only use Lemmas [2.34](#page-21-0) and [2.33.](#page-21-1)

### <span id="page-24-0"></span>**2.4 Entropy bound**

In this subsection, we first prove Proposition [2.20](#page-13-0) and then use it to show Proposition [2.18.](#page-13-2)

*Proof of Proposition [2.20](#page-13-0).* Take any  $A \in \mathcal{A}_Q(I_n)$ . Recalling Definition [2.19,](#page-13-3) we can view  $\Psi_{\ell}(A)$ as a map with input given by the intervals in  $\mathcal{I}_{\ell}^0$ , see Figure [4.](#page-14-0) In order to prove [\(2.17\)](#page-13-4), we first upper-bound  $|\Psi_{\ell}(A') : A' \in \mathcal{A}_{Q}(\mathbf{I}_n), \Psi_{\ell+1}(A') = \Psi_{\ell+1}(A)|$ , the number of possible maps  $\Psi_{\ell}(A')$ that satisfy  $\Psi_{\ell}(A', I_{\ell+1}) = \Psi_{\ell+1}(A, I_{\ell+1})$  for every interval  $I_{\ell+1} \in \mathcal{I}_{\ell+1}^0$ . For any interval  $B \in \mathcal{I}_{\ell}^0$ , let  $\hat{B}$  be the  $(\ell + 1)$ -interval in  $\mathcal{I}_{\ell+1}^0$  such that  $B \subset \hat{B}$ . Then if  $\Psi_{\ell+1}(A, \hat{B}) \neq 0$ , we get from  $(2.16)$ that

<span id="page-24-1"></span>
$$
\Psi_{\ell}(A, B) = \Psi_{\ell+1}(A, \hat{B}). \tag{2.40}
$$

Letting

<span id="page-24-5"></span>
$$
\mathcal{I}(A) := \{ \mathcal{I}_{\ell+1} \in \mathcal{I}_{\ell+1}^0 : \Psi_{\ell+1}(A, \mathcal{I}_{\ell+1}) = 0 \},\tag{2.41}
$$

by [\(2.40\)](#page-24-1), we only need to choose the values for the *ℓ*-intervals B such that  $\hat{B} \in \mathcal{I}(A)$ . Since there are  $2|\mathcal{I}(A)|$  such intervals and each has three possible values, the number of possible choices is upper-bounded by  $3^{2|\mathscr{I}(A)|}$ , i.e.,

<span id="page-24-4"></span>
$$
|\Psi_{\ell}(A') : A' \in \mathcal{A}_Q(\mathcal{I}_n), \Psi_{\ell+1}(A') = \Psi_{\ell+1}(A)| \le 3^{2|\mathcal{I}(A)|}.
$$
 (2.42)

By Lemma [2.6,](#page-7-0)  $\sigma^S$  is balanced in  $\rho_{\frac{3}{2}}(I_{\sigma})$ . As every  $\hat{B} \in \mathscr{I}(A)$  is contained in  $\rho_{\frac{3}{2}}(I_{\sigma})$ , we get that  $\sigma^S$  is balanced in B. Lemma [2.29](#page-17-0) then yields that  $J(A \cap \rho_{\frac{3}{2}}(\hat{B}), A^c \cap \rho_{\frac{3}{2}}(\hat{B})) \geq \overline{c}_2 2^{(\ell+1)\theta}$ . Summing over  $\hat{B} \in \mathscr{I}(A)$  we get that

<span id="page-24-2"></span>
$$
|\mathcal{I}(A)|\overline{c}_2 \cdot 2^{(\ell+1)\theta} \le \sum_{\hat{\mathbf{B}} \in \mathcal{I}(A)} J(A \cap \rho_{\frac{3}{2}}(\hat{\mathbf{B}}), A^c \cap \rho_{\frac{3}{2}}(\hat{\mathbf{B}})) \le 3J(A, A^c),\tag{2.43}
$$

where the last inequality comes from the fact that each  $J(x, y)$  with  $x \in A, y \in A^c$  is accounted for at most 3 times in total when summing  $J(A \cap \rho_{\frac{3}{2}}(\hat{B}), A^c \cap \rho_{\frac{3}{2}}(\hat{B}))$  over  $\hat{B} \in \mathscr{I}(A)$ . From [\(2.43\)](#page-24-2) we get that

<span id="page-24-3"></span>
$$
|\mathcal{I}(A)| \le \frac{3J(A, A^c)}{\overline{c_2 2^{(\ell+1)\theta}}} \le \frac{6Q}{\overline{c_2 2^{(\ell+1)\theta}}}.
$$
\n(2.44)

Plugging [\(2.44\)](#page-24-3) into [\(2.42\)](#page-24-4) we get  $|\Psi_{\ell}(A') : A' \in \mathcal{A}_Q(I_n), \Psi_{\ell+1}(A') = \Psi_{\ell+1}(A)| \le \exp\left\{ \frac{12 \ln(3) Q}{\bar{c}_2 2^{(\ell+1)\theta}} \right\}$  $\frac{12 \ln(3) Q}{c_2 2^{(\ell+1)\theta}}$ . As this upper bound does not depend on  $\Psi_{\ell+1}(A)$ , we can sum over all possible choices of  $\Psi_{\ell+1}(A)$ in  $\{\Psi_{\ell+1}(A) : A \in \mathcal{A}_Q(I_n)\}\$  and get

$$
E(\ell, Q, I_n) \le E(\ell + 1, Q, I_n) \times \frac{12 \ln(3) Q}{\overline{c}_2 2^{(\ell+1)\theta}},
$$

and the desired result follows from taking  $c_4 > 0$  small enough.

*Proof of Proposition [2.18](#page-13-2).* Using Proposition [2.20,](#page-13-0) our proof follows [\[21\]](#page-39-5) closely (see also [\[10\]](#page-38-0) for a more detailed version). Recall Definition [2.16.](#page-13-7) Given  $A \in \mathcal{A}(\mathcal{I}_n)$ , for each  $1 \leq \ell \leq n-3$ , let *A*<sup> $\ell$ </sup> to be the union of  $\ell$ -intervals  $I_{\ell} \in \mathcal{I}_{\ell}^0$  such that  $A \cap I_{\ell}(x) \neq \emptyset$ , that is,  $A_{\ell} \coloneqq$  $\bigcup$  $I_{\ell} \in \mathcal{I}_{\ell}^0 : A \cap I_{\ell}(x) \neq \emptyset$  $I_{\ell}(x)$ .

By the definition of  $\Psi(A)$ , we can also write  $A_{\ell} =$  $\bigcup$  $I_{\ell} \in \mathcal{I}_{\ell}^0 : \Psi(A, I_{\ell}) \neq 1$  $I_{\ell}$ . Recalling  $(2.12)$ , we write

$$
\Delta_A(h) = \sum_{\ell=0}^{n-3} \left( \Delta_{A_{\ell}}(h) - \Delta_{A_{\ell+1}}(h) \right), \tag{2.45}
$$

with the convention that  $A_0 = A$  and  $h_{A_{n-2}} = h_{\emptyset} = 0$ . By Lemma [2.15,](#page-13-8) we have for any  $z_{\ell} > 0$ 

$$
\mathbb{P}\left(|\Delta_{A_{\ell}}(h)-\Delta_{A_{\ell}+1}(h)|\geq z_{\ell}\right)\leq C_1 \exp\left(-\frac{z_{\ell}^2}{2\varepsilon^2|A_{\ell}\Delta A_{\ell+1}|}\right).
$$

Recall our definition of  $\mathscr{I}(A)$  in [\(2.41\)](#page-24-5), and recall that for any  $B \in \mathcal{I}_{\ell}^0$ ,  $\hat{B} \in \mathcal{I}_{\ell+1}^0$  is the  $(\ell+1)$ -interval such that  $B \subset \hat{B}$ . To upper-bound the symmetric difference, first notice that  $A_{\ell} \subset A_{\ell+1}$ , by the definition of the  $\Psi_{\ell}(A)$  map as in [\(2.16\)](#page-13-6). Moreover, for any  $x \in A_{\ell+1} \setminus A_{\ell}$ , there exists an  $\ell$ -interval  $B \in \mathcal{I}_{\ell}^0$  such that  $x \in B$ ,  $\Psi_{\ell}(A, B) = 1$  and  $\Psi_{\ell+1}(A, \hat{B}) = 0$ . Thus we get that  $|A_{\ell} \Delta A_{\ell+1}| \leq 2^{\ell} |\{B \in \mathcal{I}_{\ell}^0 | \}$  $\mathcal{I}_{\ell}^{0}: \Psi_{\ell+1}(A, \hat{B}) = 0$ }| =  $2^{\ell}|\mathscr{I}(A)|$ . Using [\(2.44\)](#page-24-3), we get  $|A_{\ell}\Delta A_{\ell+1}| \leq \frac{3J(A, A^{c})}{\bar{c}_{2}2^{(\ell+1)\theta}}$  $\frac{3J(A,A^c)}{c_2 2^{(\ell+1)\theta}} \cdot 2^{\ell}$  for  $0 \le \ell \le n-3$ , and therefore

<span id="page-25-0"></span>
$$
\mathbb{P}(|\Delta_{A_{\ell}}(h) - \Delta_{A_{\ell+1}}(h)| \ge z_{\ell}) \le C_1 \exp\left\{-\frac{z_{\ell}^2 \cdot \overline{c}_2 2^{(\ell+1)\theta}}{24\varepsilon^2 J(A, A^c) 2^{\ell}}\right\}.
$$
\n(2.46)

In order to give a union bound, we define  $\hat{\mathcal{A}}_Q(I_n) = \hat{\mathcal{A}}(I_n) \cap \mathcal{A}_Q(I_n)$ . As the interval  $I_n$  is fixed, we will omit it from the notation of  $\hat{\mathcal{A}}_Q$  for simplicity. Recalling the definition  $E(\ell, Q, I_n)$  from Proposition [2.20,](#page-13-0) since each  $\Psi_{\ell}(A)$  (seen as a map), uniquely determines  $A_{\ell}$ , we have  $|\{A_{\ell}: A \in A\}|$  $|\mathcal{A}_Q\rangle| \leq |\{\Psi_\ell(A) : A \in \mathcal{A}_Q\}| = E(\ell, Q, I_n)$ . Thus, from [\(2.46\)](#page-25-0) we get that for  $z_{\ell,Q} \geq 0$ ,

<span id="page-25-1"></span>
$$
\mathbb{P}(\sup_{A \in \hat{\mathcal{A}}_Q} |\Delta_{A_{\ell}(h)} - \Delta_{A_{\ell+1}}(h)| \ge z_{\ell,Q})
$$
\n
$$
\le E(\ell, Q, I_n) \cdot E(\ell+1, Q, I_n) \cdot C_1 \exp(-\frac{z_{\ell,Q}^2 \cdot \overline{c}_2 2^{(\ell+1)\theta}}{48Q}). \tag{2.47}
$$

Applying [\(2.18\)](#page-13-9) to [\(2.47\)](#page-25-1) and letting  $z_{\ell,Q} = \frac{C_2 Q}{2^{(\ell+1)\delta}}$ , we get that

$$
\mathbb{P}(\sup_{A\in \hat{\mathcal{A}}_Q} |\Delta_{A_\ell}(h) - \Delta_{A_{\ell+1}}(h)| \geq z_{\ell,Q}) \leq \frac{C_1}{C_3} \exp(\frac{C_3Q}{2^{\ell\theta}} - \frac{C_4Q}{\varepsilon^2 \cdot 2^{\ell(1-\theta+2\delta)}}).
$$



By the definition of  $\hat{\mathcal{A}}(I_n)$ , we only need to consider  $Q \ge c_2 \cdot 2^{\theta n}$ , and by our choice of  $\delta$  (at the beginning of this section) we have  $\theta > 1 - \theta + 2\delta$ . Thus we get that

<span id="page-26-2"></span><span id="page-26-1"></span>
$$
\mathbb{P}(\sup_{A\in\hat{\mathcal{A}}_Q} |\Delta_{A_\ell}(h) - \Delta_{A_{\ell+1}}(h)| \ge \frac{C_2Q}{2^{\ell\delta}}) \le C_5 \exp(-\frac{Q}{C_5\varepsilon^2 \cdot 2^{\ell(1-\theta+2\delta)}}). \tag{2.48}
$$

Let  $C_2 > 0$  be small enough depending on  $\delta$ , we get that  $\sum_{\ell=0}^{n-3} \frac{C_2 Q}{\varepsilon_2 \ell \delta} \leq \frac{0.1 Q}{\varepsilon}$ . Thus we get from [\(2.48\)](#page-26-1) and a union bound that

$$
\mathbb{P}(\sup_{A \in \hat{\mathcal{A}}_Q} |\Delta_A(h)| \ge 0.1J(A, A^c)) \le \sum_{\ell=0}^{n-3} \mathbb{P}(\sup_{A \in \hat{\mathcal{A}}_Q} |\Delta_{A_{\ell}}(h) - \Delta_{A_{\ell+1}}(h)| \ge \frac{C_2 Q}{2^{\ell\delta}})
$$
  

$$
\le \sum_{\ell=0}^{n-3} C_5 \exp(-\frac{Q}{C_5 \varepsilon^2 \cdot 2^{\ell(1-\theta+2\delta)}}) \le C_6 \exp(-\frac{Q}{C_6 \varepsilon^2 \cdot 2^{(n-2)(1-\theta+2\delta)}}). \tag{2.49}
$$

Summing over  $Q \ge c_2 \cdot 2^{\theta n}$  in [\(2.49\)](#page-26-2), taking a union bound and choosing  $\varepsilon > 0$  small enough gives the desired result (recall  $\theta > 1 - \theta + 2\delta$ ). □

## <span id="page-26-0"></span>**3 Phase Transition in 2d**

Throughout this section, we will always consider the long-range Ising model in *d* = 2 with  $2 < \alpha \leq 3$ .

Our proof of phase transition in two dimensions builds upon the ideas of [\[3\]](#page-38-12). Their argument uses an improved version of the long-range contour system of [\[2\]](#page-38-11), together with the approach of [\[18\]](#page-38-10) and the coarse-graining technique of [\[21\]](#page-39-5). For  $2 < \alpha < 3$ , phase transition follows after we make a simple modification: we replace the coarse-graining estimations depending on the boundary of a contour with estimations depending on the interactions between the flipped region and the outside. This, however, is not enough to reach the critical value  $\alpha = 3$ , and we must also improve the counting argument in [\[21\]](#page-39-5). For the convenience of comparing notes with [\[3\]](#page-38-12), we use the same notations of [\[3\]](#page-38-12) whenever possible.

Let us first introduce the long-range contour system. Given  $\sigma \in \Omega^+$ , the *incorrect* points of *σ* are defined to be  $\partial \sigma := \{x \in \mathbb{Z}^2 : \sigma_x = -\sigma_y \text{ for some } y \in \mathbb{Z}^2, |x - y| = 1\}.$  For a finite  $A \in \mathbb{Z}^2$ , we denote by  $V(A)$  the points that are separated by *A* from infinite (when viewing  $\mathbb{Z}^2$  as a nearest-neighbor graph), including *A* itself. We use the following definition to partition *∂σ*.

**Definition 3.1.** Let  $M, a > 0$ . For each  $A \subseteq \mathbb{Z}^2$ , a set  $\Gamma := {\overline{\gamma} : \overline{\gamma} \subset A}$  is called an  $(M, a)$ -*partition* when the following two conditions are satisfied.

**(A)** They form a partition of *A*, i.e.,  $\bigcup_{\overline{\gamma} \in \Gamma} \overline{\gamma} = A$  and  $\overline{\gamma} \cap \overline{\gamma}' = \emptyset$  for distinct elements of  $\Gamma$ .

**(B)** For all  $\overline{\gamma}, \overline{\gamma}' \in \Gamma$ ,

<span id="page-26-3"></span>
$$
d(\overline{\gamma}, \overline{\gamma}') > M \min\left\{ |V(\overline{\gamma})|, |V(\overline{\gamma}')|\right\}^{\frac{a}{2}}.
$$
\n(3.1)

We choose the constants *M* and *a*, similar to [\[3\]](#page-38-12), that is,  $a := a(\alpha) = \frac{6}{\alpha - 2}$  and  $M > 0$  is a large constant to be determined later. Moreover, for any  $A \subset \mathbb{Z}^2$ , we let  $\Gamma(A)$  be the finest  $(M, a)$ partition of *A*, in the following sense: given any other  $(M, a)$ -partition  $\Gamma'$ , for all  $\overline{\gamma} \in \Gamma$  there exists

<span id="page-27-0"></span>

Figure 6: The gray regions on the left, both light and dark gray, represent the set of incorrect points *∂σ* of a configuration *σ*. In this case,  $\Gamma$ (*σ*) = {*γ*, *γ*<sub>1</sub>, *γ*<sub>2</sub>}, and the dark gray region depicts sp(*γ*), while the light gray regions delimits the supports  $sp(\gamma_1)$  and  $sp(\gamma_2)$ . For each connected component of  $sp(\gamma)^c$ , a red boundary indicates that the label function attributes  $+1$  to this region, while a blue boundary indicates the complementary case. On the right, we depict the minus interior Int<sub>−</sub>( $\gamma$ ) of *γ*.

 $\overline{\gamma}' \in \Gamma'(A)$  with  $\overline{\gamma} \subset \overline{\gamma}'$ . There is an explicit construction for the finest  $(M, a)$ -partition: given any two  $(M, a)$ -partitions Γ and Γ', the intersection of Γ and Γ', defined as  $\Gamma \cap \Gamma' := {\overline{\gamma} \cap \overline{\gamma}' : \overline{\gamma} \in \overline{\gamma}}$  $\Gamma, \ \overline{\gamma} \in \Gamma', \ \overline{\gamma} \cap \overline{\gamma'} \neq \emptyset$ , is an  $(M, a)$ -partition (both conditions **(A)** and **(B)** are easy to check). As the number of  $(M, a)$ -partitions of a finite set is finite, the intersection of all  $(M, a)$ -partitions of *A* is a well-defined  $(M, a)$ -partition and it is also the finest. We write  $\Gamma(\sigma) = \Gamma(\partial \sigma)$  to simplify the notation. The contours are then pairs  $\gamma = (\overline{\gamma}, \text{lab}_{\overline{\gamma}})$ , where  $\overline{\gamma} \in \Gamma(\sigma)$  for some configuration *σ*, and lab<sub>*γ*</sub> is a function that attributes a sign to each connected component of  $\mathbb{Z}^2 \setminus \overline{\gamma}$  (for every connected component  $C \in \mathbb{Z}^2 \setminus \overline{\gamma}$ , lab $\overline{\gamma}(C)$  is the sign of  $\sigma$  on the inner boundary of *C*. The sign on the boundary is constant by [\[2,](#page-38-11) Lemma 3.10], see Figure [6](#page-27-0) for an illustration). The *support* of *γ* is sp(*γ*) :=  $\overline{\gamma}$ , and its *interior* is Int(*γ*) :=  $V(\gamma) \setminus sp(\Gamma)$ , where  $V(\gamma) := V(sp(\gamma))$ . We also denote  $|\gamma| := |\text{sp}(\gamma)|$ . Denoting by  $\text{Int}^{(1)}(\gamma), \text{Int}^{(2)}(\gamma), \dots, \text{Int}^{(k)}(\gamma)$  the connected components of  $\text{Int}(\gamma)$ , we can split the interior into the *plus interior* Int<sub>+</sub>( $\gamma$ ) and the *minus interior* Int<sub>−</sub>( $\gamma$ ), defined as

$$
\text{Int}_{\pm}(\gamma) = \bigcup_{\substack{1 \le j \le k, \\ \text{lab}_{\overline{\gamma}}(\text{Int}^{(j)}(\gamma)) = \pm 1}} \text{Int}^{(j)}(\gamma).
$$

A contour  $\gamma \in \Gamma(\sigma)$  is *external* when  $V(\gamma) \not\subset V(\gamma')$  for all  $\gamma' \in \Gamma(\sigma) \setminus {\gamma}$ . See Figure [6](#page-27-0) for an illustration of contours and minus interior. The operation that erases a contour is  $\tau_{\gamma}$ , defined as

$$
\tau_{\gamma}(\sigma)_x = \begin{cases}\n\sigma_x & \text{if } x \in \text{Int}_{+}(\gamma) \cup V(\gamma)^c, \\
-\sigma_x & \text{if } x \in \text{Int}_{-}(\gamma), \\
+1 & \text{if } x \in \text{sp}(\gamma).\n\end{cases}
$$

With these contours, in [\[3\]](#page-38-12) they can control the energy cost of erasing a contour. Although their paper focuses only on the case  $d \geq 3$ , their proofs on the energy and entropy bounds apply to  $d = 2$ directly.

**Proposition 3.2.** [\[3,](#page-38-12) Proposition 2.10] For *M* large enough, there exists a constant  $b_1 := b_1(\alpha) > 0$ , *such that for any*  $\Lambda \in \mathbb{Z}^2$ ,  $\sigma \in \Omega_{\Lambda}^+$  *and*  $\gamma \in \Gamma(\sigma)$  *external, it holds that* 

$$
H^+_{\Lambda;0}(\sigma)-H^+_{\Lambda;0}(\tau_\gamma(\sigma))\geq b_1\left(|\gamma|+J(\text{Int}_-(\gamma))\right),
$$

*with*  $J(A) \coloneqq J(A, A^c)$  *for every*  $A \subset \mathbb{Z}^2$ .

Defining  $\Omega^+ := \cup_{\Lambda \in \mathbb{Z}^2} \Omega^+_{\Lambda}$ , let

$$
\mathcal{C}_0 \coloneqq \{ \gamma \subset \mathbb{Z}^2 : 0 \in V(\gamma), \gamma \in \Gamma(\sigma) \text{ for some } \sigma \in \Omega^+, \text{ and } \gamma \text{ external} \}
$$

be the collection of external contours surrounding the origin. These long-range contours are in general not connected, but the control of the mutual distances between connected components of a contour (ensured by [\(3.1\)](#page-26-3) and the finest condition) is enough to yield an upper bound on the enumeration of these contours.

**Proposition 3.3.** [\[3,](#page-38-12) Corollary 3.28] Let  $\Lambda \in \mathbb{Z}^2$ . There exists a constant  $b_2 := b_2(\alpha) > 0$  such *that, for all*  $n \geq 1$ ,  $|\mathcal{C}_0(n)| \leq e^{b_2 n}$ .

As their approach relies on the ideas of [\[18\]](#page-38-10), we define the good event

$$
\mathcal{E} \coloneqq \left\{ \Delta_{\mathrm{Int}_-(\gamma)}(h) {\leq} \frac{b_1 |\gamma| + b_1 J(\mathrm{Int}_-(\gamma))}{4}, \ \forall \gamma \in \mathcal{C}_0 \right\},
$$

where  $\Delta_A$  is the error function defined as in [\(2.13\)](#page-12-4), but in two dimensions. We remark that Lemma [2.15](#page-13-8) also holds when  $d = 2$ . Then, the proof of phase transition in two dimensions follows the same steps of [\[3,](#page-38-12) Theorem 4.1] once we prove the following result.

<span id="page-28-0"></span>**Proposition 3.4.** For any  $2 < \alpha \leq 3$ , there exists a constant  $C := C(\alpha)$  such that  $\mathbb{P}(\mathcal{E}^c) \leq e^{-\frac{C}{\varepsilon^2}}$ *for all*  $\epsilon \leq C$ *.* 

As in [\[3\]](#page-38-12), we use a coarse-graining argument to control the external field. The argument in [\[3\]](#page-38-12) uses Dudley's entropy bound, but we opt to follow the original work of [\[21\]](#page-39-5) for two main reasons: it is the same argument used in the one-dimensional case (when proving Proposition [2.18\)](#page-13-2), and it would be clearer to state the adaptations needed to reach the critical exponent  $\alpha = 3$ .

The idea of the coarse-graining argument is to, at each scale  $\ell \geq 0$ , approximate each interior Int<sub>−</sub>( $\gamma$ ) by a region  $B_{\ell}(\text{Int}_{-}(\gamma))$  close to the region  $B_{\ell-1}(\text{Int}_{-}(\gamma))$ , in the sense that  $|B_{\ell}(\text{Int}_{-}(\gamma))\Delta$  $B_{\ell-1}(\text{Int}_-(\gamma))$  is small, and we should also control the number of possible approximations at each step. We take the same approximations as in [\[21\]](#page-39-5) and [\[3\]](#page-38-12), described next.

Fixing a parameter  $r > 4$ , at each scale  $\ell \geq 0$  we partition  $\mathbb{Z}^2$  into cubes

$$
C_{\ell}(x) := \left(\prod_{i=1}^{2} \left[2^{r\ell}x_i, \ 2^{r\ell}(x_i+1)\right)\right) \cap \mathbb{Z}^2,
$$

with  $x \in \mathbb{Z}^2$ . This gives a collection of disjoint cubes partitioning  $\mathbb{Z}^2$ . As in the one-dimensional case, each such cube will be called an *ℓ*-cube, and C*<sup>ℓ</sup>* will denote an arbitrary *ℓ*-cube. Given any  $A \in \mathbb{Z}^2$ , let  $\mathscr{C}_{\ell}(A)$  be the collection of  $\ell$ -cubes which intersect *A*. We write simply  $\mathscr{C}_{\ell}$  to denote an arbitrary collection of  $\ell$ -cubes. For any  $A \subset \mathbb{Z}^2$ , we define the collection of *admissible*  $\ell$ -cubes to be

<span id="page-28-1"></span>
$$
\mathfrak{C}_{\ell}(A) := \left\{ \mathcal{C}_{\ell} : |\mathcal{C}_{\ell} \cap A| \ge \frac{1}{2} |\mathcal{C}_{\ell}| \right\}.
$$
\n(3.2)

Then, at each scale  $\ell$ , we approximate  $\text{Int}_{-}(\gamma)$  by  $B_{\ell}(\text{Int}_{-}(\gamma)) := \bigcup_{\alpha \in \mathcal{A}}$ <sup>C</sup>∈C*ℓ*(Int−(*γ*)) C, the union of admissible

 $l$ -cubes. For any  $n, Q > 0$ , let

<span id="page-29-5"></span>
$$
\mathfrak{Int}(n,Q) \coloneqq \{ \text{Int}_{-}(\gamma) : \gamma \in \mathcal{C}_0, \ n \le |\gamma| < 2n \text{ and } Q \le J(\text{Int}_{-}(\gamma)) < 2Q \}. \tag{3.3}
$$

We first give an upper bound on  $|B_\ell(\text{Int}(n, Q))| := |\{B_\ell(\text{Int}): \text{Int} \in \text{Int}(n, Q)\}|$ . To do so, we estimate the *edge boundary* of admissible cubes, where the edge boundary between two collections of  $\ell$ -cubes  $\mathscr{C}_{\ell}$  and  $\mathscr{C}'_{\ell}$  is

<span id="page-29-6"></span>
$$
\partial(\mathscr{C}_{\ell}, \mathscr{C}'_{\ell}) \coloneqq \{ \{ C_{\ell}, C'_{\ell} \} : C_{\ell} \in \mathscr{C}_{\ell}, C'_{\ell} \in \mathscr{C}'_{\ell} \text{ and } C'_{\ell} \text{ shares an edge with } C_{\ell} \}. \tag{3.4}
$$

To ease the notation, we will write  $\partial \mathscr{C}_{\ell} = \partial(\mathscr{C}_{\ell}, \mathscr{C}_{\ell}^{c})$ , where  $\mathscr{C}_{\ell}^{c} = \mathscr{C}_{\ell}(\mathbb{Z}^{2}) \setminus \mathscr{C}_{\ell}$ . We will also write  $(C, C') \in \partial \mathfrak{C}_{\ell}(\text{Int})$  to denote a pair in  $\partial \mathfrak{C}_{\ell}(\text{Int})$  with  $C \in \mathfrak{C}_{\ell}(\text{Int})$  and  $C' \notin \mathfrak{C}_{\ell}(\text{Int})$ , for any Int = Int<sub>−</sub>( $\gamma$ ) for some  $\gamma \in C_0$ . It is clear that  $B_\ell(\text{Int}_-(\gamma))$  is uniquely determined by  $\partial \mathfrak{C}_\ell(\text{Int}_-(\gamma))$ .

Adapting verbatim the arguments in [\[3,](#page-38-12) Proposition 3.30] we can prove the next proposition.

<span id="page-29-1"></span>**Proposition 3.5.** [\[3,](#page-38-12) Proposition 3.30] Given  $n, Q, \ell \geq 0$  and  $M_{n,Q,\ell} > 0$ , let  $\tilde{\mathfrak{Im}}(n,Q)$  be the *collection of* Int  $\in$   $\mathfrak{Int}(n, Q)$  *such that* 

$$
|\partial \mathfrak{C}_{\ell}(\text{Int})| \leq M_{n,Q,\ell}.\tag{3.5}
$$

*Then, there exist constants*  $b_3 := b_3(\alpha)$  *and*  $\kappa = \kappa(\alpha)$  *such that,* 

$$
|B_{\ell}\left(\widehat{\mathfrak{Int}}(n,Q)\right)| \leq \exp\left\{b_3 \frac{\ell^{\kappa+1}n}{2^{2r\ell}} + b_3 M_{n,Q,\ell}\right\}.
$$
 (3.6)

In light of Proposition [3.5,](#page-29-1) the key remaining task is to bound  $|\partial \mathfrak{C}_{\ell}(\text{Int})|$  for Int ∈  $\mathfrak{Int}(n, Q)$ . For  $2 < \alpha < 3$ , a uniform bound suffices, but the same does not hold in the critical case of  $\alpha = 3$ , for which a more delicate estimate is required. The next subsections are dedicated to proving Proposition [3.4,](#page-28-0) first for  $2 < \alpha < 3$ , then for the critical  $\alpha = 3$ .

## <span id="page-29-0"></span>**3.1 Phase transition for** 2 *< α <* 3

In previous works, the upper bound on  $\partial \mathfrak{C}_{\ell}(\text{Int}_{-}(\gamma))$  is given in terms of  $|\gamma|$ , see [\[21,](#page-39-5) Proposition 2] and [\[3,](#page-38-12) Proposition 3.17]. Our idea is to give an upper bound for  $\partial \mathfrak{C}_{\ell}(\text{Int}_{-}(\gamma))$  in terms of  $J(\text{Int}_-(\gamma))$ , taking advantage of the large interaction between pairs in edge boundary  $\partial \mathfrak{C}_\ell(\text{Int}_-(\gamma))$ .

<span id="page-29-2"></span>**Proposition 3.6.** *There exist constants*  $b_4$ ,  $b_5$  *depending only on r such that, for every* Int ∈  $\mathfrak{Int}(n,Q)$ 

<span id="page-29-3"></span>
$$
|\partial \mathfrak{C}_{\ell}(\text{Int})| \le b_4 \frac{Q}{2^{r\ell(4-\alpha)}},\tag{3.7}
$$

*and*

<span id="page-29-4"></span>
$$
|B_{\ell}(\text{Int})\Delta B_{\ell+1}(\text{Int})| \le b_5 2^{r\ell(\alpha-2)}Q,\tag{3.8}
$$

*for every*  $\ell \geq 0$ *.* 

The proof of this proposition is very similar to that of [\[21,](#page-39-5) Proposition 2] and [\[3,](#page-38-12) Proposition 3.17]. Moreover, in the next subsection, we will prove Proposition [3.9,](#page-32-0) which immediately implies Proposition [3.6](#page-29-2) (see Remark [3.10\)](#page-32-1). As a result, we omit the proof of Proposition [3.6.](#page-29-2)

Using Propositions [3.5](#page-29-1) and [3.6](#page-29-2) one can prove Proposition [3.4](#page-28-0) following the arguments of [\[3\]](#page-38-12), with some minor modifications. We present a short alternative proof here so an interested reader can see all the required adaptations to treat the case  $\alpha = 3$ .

*Proof of Proposition* [3.4](#page-28-0) *for*  $2 < \alpha < 3$ . If  $J(\text{Int}_-(\sigma)) < 2$ , then we get that  $\text{Int}_-(\sigma) = \emptyset$  and thus  $\Delta_{\text{Int}-(\sigma)}=0$ . Hence in order to upper-bound  $\mathbb{P}(\mathcal{E}^c)$ , it suffices to consider Int ∈ Jnt(*n*, *Q*) for  $n \geq 1$  and  $Q \geq 2$ . A simple union bound yields

$$
\mathbb{P}\left(\mathcal{E}^c\right) = \mathbb{P}\left(\sup_{\gamma \in \mathcal{C}_0} \frac{\Delta_{\text{Int}_{-}(\gamma)}(h)}{|\gamma| + J(\text{Int}_{-}(\gamma))} > \frac{b_1}{4}\right) \le \sum_{n \ge 1, Q \ge 2} \mathbb{P}\left(\sup_{\text{Int}_{-} \in \mathfrak{Int}(n, Q)} \Delta_{\text{Int}}(h) > \frac{b_1}{4}(n + Q)\right). (3.9)
$$

For any  $L \geq 0$  we can write

<span id="page-30-3"></span><span id="page-30-1"></span><span id="page-30-0"></span>
$$
\Delta_{\text{Int}}(h) = \sum_{\ell=0}^{L} (\Delta_{B_{\ell}(\text{Int})}(h) - \Delta_{B_{\ell+1}(\text{Int})}(h)) + \Delta_{B_{L+1}(\text{Int})}(h), \tag{3.10}
$$

with the convention that  $\Delta_{\emptyset}(h) = 0$ . Write  $\Delta(\ell, \text{Int}, h) = \Delta_{B_{\ell}(\text{Int})}(h) - \Delta_{B_{\ell+1}(\text{Int})}(h)$  for notational convenience. By [\(3.7\)](#page-29-3), there is no admissible cube when  $2^{rL(4-\alpha)} > 2b_4Q$ . Thus, taking  $L := L_Q =$  $\lceil \frac{\log_2(2b_4Q)}{r(4-\alpha)} \rceil$  $\frac{g_2(2b_4Q)}{r(4-\alpha)}$  we have  $B_{L+1}(\text{Int}(n, Q)) = \emptyset$ , so the last term on the right hand side of [\(3.10\)](#page-30-0) vanishes. Let  $g_{\ell} = \min\{\ell + 1, L_Q - \ell + 1\}$ . As  $\frac{b_1}{4}(n+Q) \ge \frac{b_1}{40} \sum_{\ell=0}^{L} \left(\frac{n+Q}{g_{\ell}^2}\right)$  $\frac{+Q}{g_{\ell}^2}$ ), we can bound

$$
\mathbb{P}\left(\sup_{\text{Int}\in\mathfrak{Int}(n,Q)}\Delta_{\text{Int}}(h) > \frac{b_1}{4}(n+Q)\right) \le \sum_{\ell=0}^{L} \mathbb{P}\left(\sup_{\text{Int}\in\mathfrak{Int}(n,Q)}|\Delta(\ell,\text{Int},h)| > \frac{b_1(n+Q)}{40g_{\ell}^2}\right). \quad (3.11)
$$

Using Lemma [2.15,](#page-13-8) [\(3.8\)](#page-29-4) and a union bound, we have

$$
\mathbb{P}\left(\sup_{\text{Int}\in\mathfrak{Int}(n,Q)}|\Delta(\ell,\text{Int},h)| > \frac{b_1}{40}\left(\frac{n+Q}{g_\ell^2}\right)\right)
$$
\n
$$
\leq |B_\ell(\mathfrak{Int}(n,Q))| \cdot |B_{\ell+1}(\mathfrak{Int}(n,Q))| \cdot \exp\left\{-\frac{C_1[(n+Q)g_\ell^{-2}]^2}{\varepsilon^2 2^{r\ell(\alpha-2)}Q}\right\}
$$
\n
$$
\leq |B_\ell(\mathfrak{Int}(n,Q))| \cdot |B_{\ell+1}(\mathfrak{Int}(n,Q))| \cdot \exp\left\{-\frac{C_1}{\varepsilon^2 2^{r\ell(\alpha-2)}}\left(\frac{n+Q}{g_\ell^4}\right)\right\}.
$$

Combined with Proposition [3.5](#page-29-1) applied to  $M_{n,Q,\ell} = 2b_4 \frac{Q}{2r^{\ell}(4)}$  $\frac{Q}{2^{r\ell(4-\alpha)}}$  and  $M_{n,Q,\ell+1} = 2b_4 \frac{Q}{2^{r(\ell+1)}}$  $\frac{Q}{2^{r(\ell+1)(4-\alpha)}}$ , it yields that

<span id="page-30-2"></span>
$$
\mathbb{P}\left(\sup_{\text{Int}\in\mathfrak{In}_{t}(n,Q)}|\Delta(\ell,\text{Int},h)| > \frac{b_1}{40}\left(\frac{n+Q}{g_{\ell}^{2}}\right)\right)
$$
\n
$$
\leq \exp\left\{C_2\frac{(\ell+1)^{\kappa+1}n}{2^{2r\ell}} + C_2\frac{Q}{2^{r\ell(4-\alpha)}}\right\} \cdot \exp\left\{-\frac{C_1}{\varepsilon^2 2^{r\ell(\alpha-2)}}\left(\frac{n+Q}{g_{\ell}^{4}}\right)\right\}
$$
\n
$$
\leq \exp\left\{-\frac{C_3n}{\varepsilon^2 2^{r\ell(\alpha-2)}g_{\ell}^{4}} - \frac{C_3Q}{\varepsilon^2 2^{r\ell(\alpha-2)}g_{\ell}^{4}}\right\}.
$$
\n(3.12)

The last inequality holds since  $\alpha < 3$  and thus  $2^{r\ell(4-\alpha)} \geq C2^{r\ell(\alpha-2)} g_\ell^4$  for some constant  $C > 0$  and  $\varepsilon > 0$  small enough.

The result follows now from some straightforward computations. Since  $\alpha > 2$ , we get that  $2^{r(\alpha-2)(L_Q-\ell+1)} > Cg_{\ell}^4$  for some  $C > 0$  depending only on  $\alpha$ . Thus, by [\(3.11\)](#page-30-1) and [\(3.12\)](#page-30-2), we get that

$$
\mathbb{P}\left(\mathcal{E}^c\right) \leq \sum_{Q\geq 2} \sum_{n\geq 1} \sum_{\ell=0}^{L_Q} \exp\left\{-\frac{C_3 n}{\varepsilon^2 2^{rL_Q(\alpha-2)} g_\ell^4} - C_3 \frac{Q}{\varepsilon^2 2^{r\ell(\alpha-2)} g_\ell^4}\right\}
$$
\n
$$
\leq \sum_{Q\geq 2} \sum_{n\geq 1} \exp\left\{-\frac{C_3 n}{\varepsilon^2 (L_Q+1)^4 2^{rL_Q(\alpha-2)}}\right\} \sum_{\ell=0}^{L_Q} \exp\left\{-C_4 \frac{Q}{\varepsilon^2 2^{rL_Q(\alpha-2)}}\right\}.
$$
\n(3.13)

Recalling that  $L_Q = \left[\frac{\log_2(2b_4Q)}{r(4-\alpha)}\right]$  $r(4-$ , we get that  $2^{rL_Q(4-\alpha)} \leq 2^{r(4-\alpha)} \cdot 2b_4Q$ . Thus, we have

$$
\sum_{\ell=0}^{L_Q} \exp\left\{-C_4 \frac{Q}{\varepsilon^2 2^{rL_Q(\alpha-2)}}\right\} \le (L_Q + 1) \exp\left\{-\frac{C_5}{\varepsilon^2} Q^{1 - \frac{\alpha - 2}{4 - \alpha}}\right\}
$$

$$
\le 2 \left[\frac{\log_2(2b_4Q)}{r(4 - \alpha)}\right] \exp\left\{-\frac{C_5}{\varepsilon^2} Q^{\frac{6 - 2\alpha}{4 - \alpha}}\right\} \le C_6^{-1} \exp\left\{-\frac{C_6}{\varepsilon^2} Q^{\frac{6 - 2\alpha}{4 - \alpha}}\right\}.
$$
(3.14)

In addition, since  $2 < \alpha < 3$  and using again that  $2^{rL_Q} \leq 2^r (2b_4)^{\frac{1}{4-\alpha}} Q^{\frac{1}{4-\alpha}}$ , we can upper-bound  $(L_Q+1)^4 2^{r(\alpha-2)L_Q} \leq C_7 Q$  for some constant  $C_7 > 0$  depending only on *r* and  $\alpha$ . Thus we get that

<span id="page-31-2"></span>
$$
\sum_{n\geq 1} \exp\left\{-\frac{C_3 n}{\varepsilon^2 (L_Q + 1)^4 2^{rL_Q(\alpha - 2)}}\right\} \leq C_8 (L_Q + 1)^4 2^{rL_Q(\alpha - 2)} \leq C_9 Q. \tag{3.15}
$$

Plugging  $(3.14)$  and  $(3.15)$  into  $(3.13)$ , we get that

$$
\mathbb{P}\left(\mathcal{E}^c\right) \leq \sum_{Q \geq 2} C_9 Q \cdot C_6^{-1} \exp \left\{-\frac{C_6 Q^{\frac{6-2\alpha}{4-\alpha}}}{\varepsilon^2}\right\}.
$$

The result follows by taking  $\varepsilon^2$  sufficiently small.

### <span id="page-31-0"></span>**3.2 Phase Transition for**  $\alpha = 3$

To reach the critical  $\alpha = 3$  we want to take advantage of interactions in each level. We start by considering interactions restricted to smaller cubes inside C and C'. Given any  $C_{\ell}(x)$ , let

<span id="page-31-4"></span>
$$
\widehat{C}_{\ell}(x) := \left(\prod_{i=1}^{2} \left[2^{r\ell}x_i + 2^{r(\ell-1)}, \ 2^{r\ell}(x_i+1) - 2^{r(\ell-1)}\right)\right) \cap \mathbb{Z}^2 \tag{3.16}
$$

be the cube  $C_{\ell}$  after we remove an outer layer with thickness  $2^{r(\ell-1)}$ , see Figure [7.](#page-33-0) For an arbitrary  $\ell$ -cube  $C_\ell$ , we denote  $\widehat{C}_\ell$  the cube of the form [\(3.16\)](#page-31-4) sharing the same center as  $C_\ell$ .

<span id="page-31-5"></span>**Lemma 3.7.** *There exists a constant*  $b_6 := b_6(\alpha)$  *such that, for any*  $A \in \mathbb{Z}^2$  *and*  $(C, C') \in \partial \mathfrak{C}_{\ell}(A)$ *with*  $\ell \geq 1$ *,* 

$$
J(A \cap \widehat{C}, A^c \cap \widehat{C}') \ge b_6 2^{r\ell(4-\alpha)}.
$$

<span id="page-31-3"></span><span id="page-31-1"></span> $\Box$ 

*Proof.* For any  $x \in \widehat{C}$  and  $y \in \widehat{C}'$ , we have  $J_{xy} \geq (3 \cdot 2^{r\ell})^{-\alpha}$ . Thus we can bound

<span id="page-32-2"></span>
$$
J(A \cap \widehat{C}, A^c \cap \widehat{C}') \ge \frac{1}{3^{\alpha}} |A \cap \widehat{C}| |A^c \cap \widehat{C}'| 2^{-r\ell\alpha}.
$$
 (3.17)

Moreover, we have  $|A \cap \hat{C}| \geq |A \cap C| - |C \setminus \hat{C}| \geq 2^{2r\ell}(\frac{1}{2} - \frac{4}{2^r})$ , where the second inequality follows from the admissibility condition as in [\(3.2\)](#page-28-1). Recalling that  $r > 4$ , we have  $\frac{1}{2} - \frac{4}{2r} \geq \frac{1}{4}$  and thus  $|A \cap \hat{C}|$  ≥ 2<sup>2*rℓ*−2</sup>. An analogous argument shows that  $|A^c \cap \hat{C}'|$  ≥ 2<sup>2*rℓ*−2</sup>, and therefore [\(3.17\)](#page-32-2) yields

$$
J(A \cap \widehat{C}, A^c \cap \widehat{C}') \ge \frac{1}{16 \cdot 3^{\alpha}} 2^{r\ell(4-\alpha)},
$$

proving the lemma for  $b_6 = \frac{1}{16 \cdot 3^{\alpha}}$ .

The key advantage of considering smaller boxes  $\hat{C}$  is that their interactions do not overlap in the coarse-graining approximations. This is demonstrated in the next lemma.

<span id="page-32-6"></span>**Lemma 3.8.** *For every*  $\ell \geq 0$  *and*  $A \in \mathbb{Z}^2$ *, defining* 

<span id="page-32-4"></span>
$$
Q_{\ell}(A) := \sum_{(C,C') \in \partial \mathfrak{C}_{\ell}(A)} J(A \cap \widehat{C}, A^c \cap \widehat{C}'),
$$
\n(3.18)

 $\Box$ 

*we have*

$$
\sum_{\ell \ge 0} Q_{\ell}(A) \le J(A).
$$

*Proof.* Given any  $(C_{\ell}, C'_{\ell}) \in \partial \mathfrak{C}_{\ell}(A)$ , it is enough to show that no interaction between  $\widehat{C}_{\ell}$  and  $\widehat{C}'_{\ell}$ appear in  $Q_k(A)$  for all  $k > \ell$ . An interaction  $J_{xy}$  with  $x \in A \cap \widehat{C}_{\ell}$  and  $y \in A^c \cap \widehat{C}_{\ell}^{\prime}$  appears in  $Q_k(A)$  only if there exists  $(C_k, C'_k) \in \partial \mathfrak{C}_k(A)$  with  $C_\ell \subset C_k$  and  $C'_\ell \subset C'_k$ . As both pairs  $\{C_\ell, C'_\ell\}$ and  $\{C_k, C'_k\}$  share an edge, one of the edges of  $C_\ell$  must be contained in an edge of  $C_k$ , and the same holds for  $C'_{\ell}$  and  $C'_{k}$ , see Figure [7](#page-33-0) for an illustration with  $k = \ell + 1$ . As  $2^{r(k-1)} \geq 2^{r\ell}$ , we must have  $C_k \cap C_\ell = \emptyset$  and similarly  $C'_k \cap C'_\ell = \emptyset$ . П

The next proposition is a modification of [\[21,](#page-39-5) Proposition 1] and [\[3,](#page-38-12) Proposition 3.16], which is also a generalization of Proposition [3.6.](#page-29-2)

<span id="page-32-0"></span>**Proposition 3.9.** *There exists a constant*  $b_7$  *depending only on*  $r$  *such that, for any*  $A \in \mathbb{Z}^2$  *and*  $\ell \geq 0$ *, we have* 

<span id="page-32-3"></span>
$$
|\partial \mathfrak{C}_{\ell}(A)| \le b_6 \frac{Q_{\ell}(A)}{2^{r\ell(4-\alpha)}},\tag{3.19}
$$

*and*

<span id="page-32-5"></span>
$$
|B_{\ell}(A)\Delta B_{\ell+1}(A)| \le b_7 2^{r\ell(\alpha-2)} Q_{\ell}(A). \tag{3.20}
$$

<span id="page-32-1"></span>**Remark 3.10.** By the definition of  $Q_{\ell}(A)$  we can directly bound  $Q_{\ell}(A) \leq J(A)$ , so Proposition *[3.6](#page-29-2) follows immediately from Proposition [3.9.](#page-32-0)*

*Proof of Proposition [3.9](#page-32-0).* For [\(3.19\)](#page-32-3), combining Lemma [3.7](#page-31-5) with [\(3.18\)](#page-32-4) yields that

$$
Q_{\ell}(A) \ge b_6 2^{4-\alpha} |\partial \mathfrak{C}_{\ell}(A)|.
$$

<span id="page-33-0"></span>

Figure 7: In this picture, we take  $r = 3$ . The dashed lines delimit two  $(\ell+1)$ -cubes,  $C_{\ell+1}$  and  $C'_{\ell+1}$ , while the dark gray areas are  $C_{\ell+1}$  and  $C'_{\ell+1}$ . The black cube represents one of the  $\ell$ -cubes inside  $C_{\ell+1}$ , and the light gray cubes are two neighboring  $\ell$ -cubes  $C_{\ell} \subset C_{\ell+1}$  and  $C'_{\ell} \subset C'_{\ell+1}$ .

We now turn to the proof of [\(3.20\)](#page-32-5). Given  $C \in \mathscr{C}_{\ell+1}(B_{\ell+1}(A) \setminus B_{\ell}(A))$ , there exists a pair  ${C_{\ell}, C'_{\ell}} \in \partial \mathfrak{C}_{\ell}(A)$  with  $C_{\ell} \cup C'_{\ell} \subset C$ . By Lemma [\(3.7\)](#page-31-5) we have

$$
| (B_{\ell+1}(A) \setminus B_{\ell}(A)) \cap C | \leq |C| = 2^{2r(\ell+1)} = 2^{2r} 2^{r\ell(\alpha-2)} 2^{r\ell(4-\alpha)}
$$
  

$$
\leq b_6^{-1} 2^{2r} 2^{r\ell(\alpha-2)} J(A \cap \widehat{C}_{\ell}, A^c \cap \widehat{C}'_{\ell}).
$$

Therefore, as all cubes  $C \in \mathscr{C}_{\ell+1}(B_{\ell+1}(A) \setminus B_{\ell}(A))$  are disjoint,

$$
|B_{\ell+1}(A) \setminus B_{\ell}(A)| = \sum_{\substack{C \in \mathscr{C}_{\ell+1}(B_{\ell+1}(A) \setminus B_{\ell}(A)) \\ \leq \sum_{\substack{C \in \mathscr{C}_{\ell+1}(B_{\ell+1}(A) \setminus B_{\ell}(A)) \\ \leq b_6^{-1} 2^{2r} 2^{r\ell(\alpha-2)}}} \frac{|(B_{\ell+1}(A) \setminus B_{\ell}(A)) \cap C|}{b_6^{-1} 2^{2r} 2^{r\ell(\alpha-2)} 3^{r\ell(\alpha-2)}} J(A \cap \widehat{C}_{\ell}, A^c \cap \widehat{C}_{\ell}')
$$

To get the same bound for  $|B_{\ell}(A) \setminus B_{\ell+1}(A)|$  we use a similar argument, covering  $B_{\ell}(A) \setminus B_{\ell+1}(A)$ with  $(\ell + 1)$ -cubes and considering the edge boundaries in those  $(\ell + 1)$ -cubes. This concludes the proof taking  $b_7 = b_6^{-1} 2^{2r+1}$ . □

**Remark 3.11.** *Notice that the proof of* [\(3.20\)](#page-32-5) *also works when*  $|B_{\ell+1}(A)\Delta B_{\ell}(A)| = |B_{\ell}(A)|$ *, that is, when ℓ is the last scale with non-empty admissible cubes. In this case,* [\(3.20\)](#page-32-5) *actually bounds*  $|B_{\ell}(A)|$ *.* 

For  $\alpha = 3$ , we can prove a stronger version of Lemma [2.24](#page-15-3) in two dimensions. Let us recall the following definition from the previous sections. Given  $A \subset \mathbb{Z}^2$  and a configuration  $\sigma \in \Omega$ ,  $A^+(\sigma) := \{x \in A : \sigma_x = 1\}$  is the collection of plus spins in *A*, and similarly  $A^-(\sigma)$  is the collection of minus spins in *A*. We start with an auxiliary lemma.

<span id="page-33-1"></span>**Lemma 3.12.** For any  $0 \leq \ell < L$ , let  $\mathscr{C}, \mathscr{C}'$  be two disjoint collections of  $\ell$ -cubes such that  $\cup_{C\in\mathscr{C}\cup\mathscr{C}'} C$  *is an L*-cube. If  $|\mathscr{C}| \ge c2^{2r(L-\ell)}$  *and*  $|\mathscr{C}'| \ge c2^{2r(L-\ell)}$  *for some constant*  $0 < c \le 0.5$ *, then*

$$
|\partial(\mathscr{C}, \mathscr{C}')| \geq \sqrt{c}2^{r(L-\ell)}.
$$

*Proof.* By the isoperimetric inequality,  $|\partial \mathscr{C}| \geq 4\sqrt{|\mathscr{C}|}$  and  $|\partial \mathscr{C}'| \geq 4\sqrt{|\mathscr{C}'|}$ . Notice that  $|\partial \mathscr{C}|$  +  $|\partial \mathscr{C}'| = 2|\partial(\mathscr{C}, \mathscr{C}')| + |\partial(\mathscr{C} \cup \mathscr{C}')|$ . In addition, since the union of the *ℓ*-cubes in  $\mathscr{C} \cup \mathscr{C}'$  is an *L*-cube, thus we get that  $|\partial(\mathscr{C} \cup \mathscr{C}')| = 4 \cdot 2^{r(L-\ell)}$ . Therefore

$$
|\partial(\mathscr{C}, \mathscr{C}')| = 2\sqrt{|\mathscr{C}|} + 2\sqrt{|\mathscr{C}'|} - 2 \cdot 2^{r(L-\ell)} = 2 \cdot \frac{(\sqrt{|\mathscr{C}|} + \sqrt{|\mathscr{C}'|} - 2^{r(L-\ell)}) \cdot (\sqrt{|\mathscr{C}|} + \sqrt{|\mathscr{C}'|} + 2^{r(L-\ell)})}{\sqrt{|\mathscr{C}|} + \sqrt{|\mathscr{C}'|} + 2^{r(L-\ell)}}.
$$
(3.21)

Note that  $|\mathscr{C}| + |\mathscr{C}'| = 2^{2r(L-\ell)}$ , we get that

<span id="page-34-1"></span>
$$
\left(\sqrt{|\mathscr{C}|} + \sqrt{|\mathscr{C}'|} - 2^{r(L-\ell)}\right) \cdot \left(\sqrt{|\mathscr{C}|} + \sqrt{|\mathscr{C}'|} + 2^{r(L-\ell)}\right) = 2\sqrt{|\mathscr{C}| \cdot |\mathscr{C}'|} \ge \sqrt{2c} \cdot 2^{2r(L-\ell)}.\tag{3.22}
$$

In addition, we have

<span id="page-34-2"></span>
$$
\sqrt{|\mathscr{C}|} + \sqrt{|\mathscr{C}'|} + 2^{r(L-\ell)} \le (\sqrt{2}+1) \cdot 2^{r(L-\ell)}.
$$
\n(3.23)

<span id="page-34-0"></span>Plugging [\(3.22\)](#page-34-1) and [\(3.23\)](#page-34-2) into [\(3.21\)](#page-34-3), we complete the proof of Lemma [3.12.](#page-33-1)

**Lemma 3.13.** *There exists a constant*  $b_8 > 0$  *depending only on r such that for any L*-*cube* C *and any configuration*  $\sigma$ *, if* min{ $|C^+(\sigma)|$ ,  $|C^-(\sigma)|$ } = *m, then we have* 

<span id="page-34-4"></span>
$$
J(\mathcal{C}^+(\sigma), \mathcal{C}^-(\sigma)) \ge b_8 \sqrt{m} \ln(m). \tag{3.24}
$$

<span id="page-34-3"></span> $\Box$ 

*Proof.* Our proof is based on an induction on *L*. If  $L = 1$ , the desired result follows by taking  $b_8 > 0$ small enough. Now we assume [\(3.24\)](#page-34-4) holds for any  $\ell \leq L-1$  and then prove the case for  $\ell = L$ . Without loss of generality one can assume  $|C^+(\sigma)| \geq |C^-(\sigma)|$ . Recall the definition of  $\mathfrak{C}_{\ell}(C^-(\sigma))$ from [\(3.2\)](#page-28-1). Let  $a_{\ell} := |\mathfrak{C}_{\ell}(\mathrm{C}^{-}(\sigma))|$ . We consider four different cases.

*Case (i):*  $m \leq 2^{2r(L-1)-1}$ . We first write C as the disjoint union of  $(L-1)$ -cubes  $C_1, C_2, \cdots, C_{2^{2r}}$ . Let  $x_i = |C_i^-(\sigma)|$ , for  $1 \leq i \leq 2^{2r}$ , be the number of minus spins in  $C_i$ . Since  $\sum_{i=1}^k x_i = m \leq$  $2^{2r(L-1)-1}$ , we must have min $\{|C_i^+(\sigma)|, |C_i^-(\sigma)|\} = x_i$  for all  $1 \leq i \leq 2^{2r}$ . Thus, we can apply the induction hypothesis to get

$$
J(C^+(\sigma), C^-(\sigma)) \ge \sum_{i=1}^{2^{2r}} J(C_i^+(\sigma), C_i^-(\sigma)) \ge \sum_{i=1}^{2^{2r}} b_8 \sqrt{x_i} \ln(x_i).
$$
 (3.25)

The desired result follows once we notice that  $f(x) = \sqrt{x} \ln(x)$  is a concave function, and thus we get from Karamata's inequality that  $\sum_{i=1}^{2^{2r}} \sqrt{x_i} \ln(x_i) \geq \sqrt{m} \ln(m)$ .

Case (ii):  $m > 2^{2r(L-1)-1}$  and  $a_{\ell} < \frac{m}{2^{2r\ell+1}}$  for some  $\ell \leq \frac{L}{2}$ . In this case, there are at least  $\frac{m}{2}$  minuses contained in non-admissible  $\ell$ -cubes with respect to C (recall [\(3.2\)](#page-28-1)). Let  $B_1, \dots, B_k$  be *ℓ*-cubes in C that are not in  $\mathfrak{C}_{\ell}(\mathcal{C}^{-}(\sigma))$ , and as in Case (i), let  $x_i = |B_i^-(\sigma)|$ , for  $1 \leq i \leq k$ , be the number of minuses in those cubes. Then we get from the induction hypothesis that

<span id="page-34-5"></span>
$$
J(\mathcal{C}^+(\sigma), \mathcal{C}^-(\sigma)) \ge \sum_{i=1}^k J(\mathcal{B}_i^+(\sigma), \mathcal{B}_i^-(\sigma)) \ge \sum_{i=1}^k b_8 \sqrt{x_i} \ln(x_i). \tag{3.26}
$$

Again we can use that  $f(x) = \sqrt{x} \ln(x)$  is concave, and thus by Karamata's inequality we get that

$$
\sum_{i=1}^{k} \sqrt{x_i} \ln(x_i), \ 0 \le x_i \le 2^{2r\ell - 1}
$$

reaches its minimum if at most one  $x_i$  takes value in  $(0, 2^{2r\ell-1})$  and others have values either 0 or  $2^{2r\ell-1}$ . Thus we get that

<span id="page-35-0"></span>
$$
\sum_{i=1}^{k} \sqrt{x_i} \ln(x_i) \ge \left\lfloor \frac{m}{2^{2r\ell}} \right\rfloor \cdot \sqrt{2^{2r\ell - 1}} \ln(2^{2r\ell - 1}) \ge m \cdot 2^{-r\ell - 0.5}.
$$
 (3.27)

Combining [\(3.26\)](#page-34-5), [\(3.27\)](#page-35-0) and the fact that  $2^{2r(L-1)-1} \le m \le 2^{2rL-1}$ , we get that

$$
J(\mathcal{C}^+(\sigma), \mathcal{C}^-(\sigma)) \ge m \cdot 2^{-r\ell - 0.5} \ge \sqrt{m} \cdot 2^{rL - r\ell - 1} \ge C_2 \sqrt{m} \cdot \ln(m),
$$

for some small absolute constant  $C_2$ , where the last transition used the assumption that  $\ell \leq L/2$ . This then completes the verification of  $(3.24)$  (we may choose  $b_8$  smaller than the absolute constant  $C_2$  here).

Case (iii):  $m > 2^{2r(L-1)-1}$ ,  $a_j \geq \frac{m}{2^{2r_j+1}}$  for all  $j \leq \frac{L}{2}$  and  $a_\ell > 3 \cdot 2^{2r(L-\ell)-2}$  for some  $\ell \leq \frac{L}{2}$ . Let  $D_1, \dots, D_k$  be the  $\ell$ -cubes in  $\mathfrak{C}_{\ell}(\mathbf{C}^{-}(\sigma))$  with  $x_i := |D_i(\sigma)| \leq 3 \cdot 2^{2r\ell-2}$ . Then we get that  $m \geq k \cdot 2^{2r\ell-1} + (a_{\ell} - k) \cdot 3 \cdot 2^{2r\ell-2}$ . Combining with the fact that  $m \leq 2^{2rL-1}$  and  $a_{\ell} > 3 \cdot 2^{2r(L-\ell)-2}$ , we get that  $k \geq 3a_{\ell} - \frac{m}{2^{2r(\ell-2)}} \geq 2^{2r(L-\ell)-2}$ . Applying induction hypothesis to  $D_i$  we conclude that

$$
J(\mathrm{C}^+(\sigma),\mathrm{C}^-(\sigma)) \ge \sum_{i=1}^k J(\mathrm{D}_i^+(\sigma),\mathrm{D}_i^-(\sigma)) \ge 2^{2r(L-\ell)-2} \cdot b_8 \sqrt{2^{2r\ell-2}} \ln(2^{2r\ell-2}).
$$

Since  $\ell \leq L/2$  and  $m \leq \frac{2^{2rL}}{2}$  $\frac{n}{2}$ , the desired lower-bound holds.

*Case (iv):*  $m > 2^{2r(L-1)-1}$  *and*  $\frac{m}{2^{2r(\ell+1)}} \leq a_{\ell} \leq 3 \cdot 2^{2r(L-\ell)-2}$  *for all*  $\ell \leq \frac{L}{2}$ . In this case, we recall [\(3.18\)](#page-32-4) and claim that for any  $\ell \leq \frac{L}{2}$ , we have  $Q_{\ell}(C^{-}(\sigma)) \geq C_3 \sqrt{m}$  for some absolute constant  $C_3 > 0$ . Having assumed  $m > 2^{2r(L-1)-1}$  and  $\frac{m}{2^{2r\ell+1}} \le a_{\ell}$ , we can apply Lemma [3.12](#page-33-1) (since  $\min\{a_{\ell}, 2^{2r(L-\ell)}-a_{\ell}\}\geq \frac{m}{2^{2r(\ell+1)}}$ ) to get that  $|\partial(\mathfrak{C}_{\ell}(C^-(\sigma)), \mathfrak{C}_{\ell}(C^+(\sigma)))|\geq \sqrt{\frac{m}{2^{2r(\ell+1)}}}$ . Then, recalling [\(3.18\)](#page-32-4), we get from Lemma [3.7](#page-31-5) that

$$
Q_{\ell}(\mathcal{C}^{-}(\sigma)) \geq |\partial \left( \mathfrak{C}_{\ell} \left( \mathcal{C}^{-}(\sigma) \right), \mathfrak{C}_{\ell} \left( \mathcal{C}^{+}(\sigma) \right) \right)| \cdot b_2 2^{r\ell} \geq \sqrt{\frac{m}{2^{2r\ell+1}}} \cdot b_2 2^{r\ell} \geq C_3 \sqrt{m}.
$$

Summing over  $0 \leq \ell \leq \frac{L}{2}$  and combining with Lemma [3.8](#page-32-6) we conclude that

$$
J(\mathrm{C}^+(\sigma), \mathrm{C}^-(\sigma)) \ge \frac{L}{2} \cdot C_3 \sqrt{m}.
$$

 $\Box$ 

The desired result follows by noticing that  $\ln(m) \leq 2rL$  (and we may choose  $b_8 \leq \frac{C_3}{4r}$ ).

We are ready to prove the main result of this section.

*Proof of Proposition* [3.4](#page-28-0) *for*  $\alpha = 3$ . Recall [\(3.9\)](#page-30-3) and [\(3.10\)](#page-30-0). Recall the definition of  $\mathfrak{Int}(n, Q)$ from [\(3.3\)](#page-29-5). Consider Int  $\in \mathfrak{Int}(n, Q)$  for  $n \geq 1$  and  $Q \geq 2$ . Given any pair  $(C, C') \in \partial \mathfrak{C}_{\ell}(\text{Int})$  (recall the definition of edge boundary from [\(3.4\)](#page-29-6)), there exists a cube  $C_1 \subset C \cup C'$  with side length  $2^{r\ell}$ and satisfying min  $\{|C_1 \cap Int|, |C_1 \cap Int^c|\} \geq 2^{2r\ell-2}$ . Hence, by Lemma [3.13](#page-34-0) we have

<span id="page-36-0"></span>
$$
J(\text{Int} \cap \text{C}, \text{Int}^c \cap \text{C}') \ge \overline{b}_9 \ell 2^{r\ell} \tag{3.28}
$$

for a suitable constant  $\overline{b}_9$ . Fixing  $L := L_Q$ , where  $L_Q$  is the largest integer  $\ell$  satisfying  $\overline{b}_9 \ell 2^{r\ell} \leq Q$ , by [\(3.28\)](#page-36-0) we have  $B_{L+1}(\text{Int}) = \emptyset$  for every  $\text{Int} \in \mathfrak{Int}(n, Q)$ . Let  $\mathcal{H}_{\ell} = \mathcal{H}_{\ell,n,Q}$  be the event such that for all Int ∈  $\mathfrak{Int}(n, Q)$ ,  $|\Delta(\ell, \text{Int}, h)|$  ≤  $C_1(Q_{\ell}(\text{Int}) + \frac{n}{(\ell+1)^2} + \frac{Q}{(L-\ell+1)^2})$ with  $C_1 < 0.1b_1(2\pi)^{-2}$ . Lemma [3.8](#page-32-6) guarantees that  $C_1 \sum_{\ell=0}^{L} (Q_{\ell}(\text{Int}) + \frac{n}{(\ell+1)^2} + \frac{Q}{(L-\ell+1)^2}) \leq \frac{b_1}{4}(n+Q)$ , so by [\(3.10\)](#page-30-0) we have

<span id="page-36-2"></span><span id="page-36-1"></span>
$$
\mathcal{E}^c \subset \cup_{\ell=0}^L \mathcal{H}_\ell^c. \tag{3.29}
$$

For  $0 \leq \ell \leq L$ , we wish to split Int  $\in \mathfrak{Int}(n, Q)$  according to  $Q_{\ell}(\text{Int})$  (recall [\(3.18\)](#page-32-4)). To this end, we note that Lemma [3.7](#page-31-5) assures that  $Q_{\ell}(\text{Int}) \geq b_6 2^{r\ell}$  as long as  $\mathfrak{C}_{\ell}(\text{Int}) \neq \emptyset$ . Thus, defining  $\mathfrak{Int}_{\ell,k}(n,Q) := \{\text{Int} \in \mathfrak{Int}(n,Q) : Q_{\ell}(\text{Int}) \in [b_6 2^{r_{\ell}} 2^k, b_6 2^{r_{\ell}} 2^{k+1})\},\}$  we can bound

$$
\mathbb{P}(\mathcal{H}_{\ell}^{c}) \leq \sum_{k=0}^{K_{\ell}} \mathbb{P}\Bigg(\sup_{\text{Int}\in\mathfrak{Int}_{\ell,k}(n,Q)} |\Delta(\ell,\text{Int},h)| \geq z_{\ell,k,n,Q}\Bigg). \tag{3.30}
$$

where  $z_{\ell,k,n,Q} = C_1 \left( b_6 2^{r\ell} 2^k + \frac{n}{(\ell+1)^2} + \frac{Q}{(L-\ell+1)^2} \right)$ ), and  $K_{\ell}$  is the smallest integer  $K$  satisfying  $\overline{b}_9 L 2^{rL} < b_6 2^{\ell} 2^{K+1}$ . By our choice of  $L = L_Q$ , we have  $L 2^{rL} > \overline{b}_9^{-1} Q$ , and thus  $\mathfrak{Int}_{\ell,k}(n, Q) = \{\emptyset\}$ for  $k > K_{\ell}$ .

By [\(3.19\)](#page-32-3), we can apply Proposition [3.5](#page-29-1) with  $M_{n,Q,\ell} = b_6^2 2^{k+1}$ , getting that  $|B_{\ell}(\text{Int}_{\ell,k}(n,Q))| \le$  $\exp\left\{C_2 \frac{\ell^{k+1}n}{2^{2r\ell}} + C_2 2^k\right\}$ . Using Lemma [2.15,](#page-13-8) [\(3.20\)](#page-32-5) and a union bound, we get that

$$
\mathbb{P}\Bigg(\sup_{\text{Int}\in\mathfrak{Int}_{\ell,k}(n,Q)}|\Delta(\ell,\text{Int},h)| \geq z_{\ell,k,n,Q}\Bigg)
$$
\n
$$
\leq |B_{\ell}(\mathfrak{Int}_{\ell,k}(n,Q))| \cdot |B_{\ell+1}(\mathfrak{Int}_{\ell,k}(n,Q))| \cdot \exp\Bigg\{-\frac{C_3[2^{r\ell}2^k + n(\ell+1)^{-2} + Q(L-\ell+1)^{-2}]^2}{\varepsilon^2 2^{2r\ell} 2^k}\Bigg\}
$$
\n
$$
\leq |B_{\ell}(\mathfrak{Int}_{\ell,k}(n,Q))| \cdot |B_{\ell+1}(\mathfrak{Int}_{\ell,k}(n,Q))| \cdot \exp\Bigg\{-\frac{C_4}{\varepsilon^2}\Bigg[\frac{n}{(\ell+1)^2 2^{r\ell}} + \frac{Q}{2^{r\ell}(L-\ell+1)^2} + 2^k\Bigg]\Bigg\}.
$$

Combined with [\(3.20\)](#page-32-5), it yields

$$
\sum_{k=0}^{K_{\ell}} \mathbb{P}\left(\sup_{\text{Int}\in\mathfrak{In}_{\ell,k}(n,Q)} |\Delta(\ell,\text{Int},h)| \geq z_{\ell,k,n,Q}\right)
$$
\n
$$
\leq \sum_{k=0}^{K_{\ell}} \exp\left\{C_2 \frac{(\ell+1)^{\kappa+1}n}{2^{2r\ell}} + C_2 2^k\right\} \exp\left\{-\frac{C_4}{\varepsilon^2} \left[\frac{n}{(\ell+1)^2 2^{r\ell}} + \frac{Q}{2^{r\ell}(L-\ell+1)^2} + 2^k\right]\right\}
$$
\n
$$
\leq \exp\left\{-\frac{C_5 n}{\varepsilon^2 (\ell+1)^2 2^{r\ell}} - C_5 \frac{Q}{\varepsilon^2 2^{r\ell}(L-\ell+1)^2}\right\} \sum_{k\geq 0} \exp\left\{-\frac{C_5 2^k}{\varepsilon^2}\right\}
$$

$$
\leq \exp\left\{-\frac{C_6 n}{\varepsilon^2 (\ell+1)^2 2^{r\ell}} - C_6 \frac{Q}{\varepsilon^2 2^{r\ell} (L-\ell+1)^2}\right\},\tag{3.31}
$$

where the second and the last inequalities hold for  $\varepsilon$  small enough. Combining[\(3.9\)](#page-30-3), [\(3.29\)](#page-36-1), [\(3.30\)](#page-36-2) and [\(3.31\)](#page-37-1), we have

<span id="page-37-1"></span>
$$
\mathbb{P}\left(\mathcal{E}^c\right) \leq \sum_{Q\geq 2} \sum_{n\geq 1} \sum_{\ell=0}^L \exp\left\{-\frac{C_6 n}{\varepsilon^2 (\ell+1)^2 2^{r\ell}} - C_6 \frac{Q}{\varepsilon^2 2^{r\ell} (L-\ell+1)^2}\right\}
$$

$$
\leq \sum_{Q\geq 2} \sum_{n\geq 1} \exp\left\{-\frac{C_6 n}{\varepsilon^2 (L+1)^2 2^{rL}}\right\} \sum_{\ell=0}^L \exp\left\{-C_6 \frac{Q}{\varepsilon^2 2^{r\ell} (L-\ell+1)^2}\right\}.
$$

Again by our choice of  $L = L_Q$  we have  $ln(Q)2^{rL} \leq C_7Q$ , for  $C_7$  large enough. Thus,

<span id="page-37-2"></span>
$$
\sum_{\ell=0}^{L} \exp\left\{-C_6 \frac{Q}{\varepsilon^2 2^{r\ell} (L-\ell+1)^2}\right\} \le \sum_{\ell=0}^{L} \exp\left\{-C_6 C_7 \frac{\ln(Q) \cdot 2^{r(L-\ell)}}{\varepsilon^2 (L-\ell+1)^2}\right\}.
$$
(3.32)

Recall that  $r > 4$ , thus we have  $\frac{2^{r(L-\ell)}}{(L-\ell+1)^2} \ge (L-\ell+1)$ . Combined with [\(3.32\)](#page-37-2), it yields that

$$
\sum_{\ell=0}^{L} \exp \left\{ -C_6 \frac{Q}{\varepsilon^2 2^{r\ell} (L - \ell + 1)^2} \right\} \le \sum_{\ell=0}^{L} \exp \left\{ -C_6 C_7 \frac{\ln(Q) \cdot (L - \ell + 1)}{\varepsilon^2} \right\}
$$

$$
= \sum_{\ell=0}^{L} Q^{-\frac{C_6 C_7 (L - \ell + 1)}{\varepsilon^2}} \le C_8 Q^{-\frac{C_8}{\varepsilon^2}}.
$$

Moreover, one can easily bound

$$
\sum_{n\geq 1} \exp\left\{-\frac{C_6 n}{\varepsilon^2 (L+1)^2 2^{rL}}\right\} \leq \sum_{n\geq 1} \exp\left\{-\frac{C_9 n}{\varepsilon^2 \ln(Q)^2 Q}\right\} \leq C_{10} Q (\ln Q)^2,
$$

and we conclude that  $\mathbb{P}(\mathcal{E}^c) \leq \sum_{Q \geq 2} C_8 C_{10} Q^{1-C_8/\varepsilon^2} (\ln Q)^2$ , and the desired result follows by taking *ε* large enough.  $\Box$ 

## **Acknowledgements**

J. Ding is supported by NSFC Tianyuan Key Program Project No.12226001, NSFC Key Program Project No.12231002, and by New Cornerstone Science Foundation through the XPLORER PRIZE. F. Huang is supported by the Beijing Natural Science Foundation's Undergraduate Initiating Research Program (Grant No. QY23006). We thank Lucas Affonso for discussions during his visit to Peking University.

## <span id="page-37-0"></span>**References**

[1] L. Affonso, R. Bissacot, H. Corsini, and K. Welsch. Phase Transitions on 1d Long-Range Ising Models with Decaying Fields: A Direct Proof via Contours, 2024, [ArXiv:2412.07098.](http://arxiv.org/abs/2412.07098)

- <span id="page-38-12"></span><span id="page-38-11"></span>[2] L. Affonso, R. Bissacot, E. O. Endo, and S. Handa. Long-range Ising models: Contours, phase transitions and decaying fields. *Journal of the European Mathematical Society*, 2024.
- <span id="page-38-14"></span>[3] L. Affonso, R. Bissacot, and J. Maia. Phase Transitions in Multidimensional Long-Range Random Field Ising Models, 2024, [ArXiv:2307.14150.](http://arxiv.org/abs/2307.14150)
- [4] M. Aizenman, R. L. Greenblatt, and J. L. Lebowitz. Proof of rounding by quenched disorder of first order transitions in low-dimensional quantum systems. *Journal of Mathematical Physics*, 53(2), 2012.
- <span id="page-38-3"></span>[5] M. Aizenman, M. Harel, and R. Peled. Exponential Decay of Correlations in the 2D Random Field Ising Model. *Journal of Statistical Physics*, 180(1-6):304–331, 2020.
- <span id="page-38-4"></span>[6] M. Aizenman and R. Peled. A power-law upper bound on the correlations in the 2*D* random field Ising model. *Communications in Mathematical Physics*, 372(3):865–892, 2019.
- <span id="page-38-1"></span>[7] M. Aizenman and J. Wehr. Rounding effects of quenched randomness on first-order phase transitions. *Communications in Mathematical Physics*, 130:489–528, 1990.
- <span id="page-38-16"></span>[8] M. Biskup, L. Chayes, and S. A. Kivelson. On the absence of ferromagnetism in typical 2D ferromagnets. *Communications in Mathematical Physics*, 274:217–231, 2007.
- <span id="page-38-15"></span>[9] R. Bissacot, E. O. Endo, A. C. van Enter, B. Kimura, and W. M. Ruszel. Contour methods for long-range Ising models: weakening nearest-neighbor interactions and adding decaying fields. *Annales Henri Poincaré*, 19:2557–2574, 2018.
- <span id="page-38-0"></span>[10] A. Bovier. *Statistical Mechanics of Disordered Systems: A Mathematical Perspective*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2006.
- <span id="page-38-2"></span>[11] J. Bricmont and A. Kupiainen. Phase transition in the 3d random field Ising model. *Communications in Mathematical Physics*, 116:539–572, 1988.
- <span id="page-38-13"></span>[12] M. Cassandro, E. Orlandi, and P. Picco. Phase Transition in the 1d Random Field Ising Model with long range interaction. *Communications in Mathematical Physics*, 288:731–744, 2009.
- <span id="page-38-5"></span>[13] S. Chatterjee. On the decay of correlations in the random field Ising model. *Communications in Mathematical Physics*, 362(1):253–267, 2018.
- <span id="page-38-6"></span>[14] J. Ding, F. Huang, and A. Xia. A phase transition and critical phenomenon for the twodimensional random field Ising model, 2024, [ArXiv:2310.12141.](http://arxiv.org/abs/2310.12141)
- <span id="page-38-9"></span>[15] J. Ding, Y. Liu, and A. Xia. Long range order for three-dimensional random field Ising model throughout the entire low temperature regime. *Inventiones mathematicae*, 2024.
- <span id="page-38-7"></span>[16] J. Ding and M. Wirth. Correlation length of the two-dimensional random field Ising model via greedy lattice animal. *Duke Mathematical Journal*, 172(9):1781–1811, 2023.
- <span id="page-38-8"></span>[17] J. Ding and J. Xia. Exponential decay of correlations in the two-dimensional random field Ising model. *Inventiones mathematicae*, 224:1–47, 2021.
- <span id="page-38-10"></span>[18] J. Ding and Z. Zhuang. Long range order for random field Ising and Potts models. *Communications on Pure and Applied Mathematics*, 77(1):37–51, 2024.
- <span id="page-39-3"></span><span id="page-39-1"></span>[19] R. L. Dobrushin. Gibbsian random fields. General case. *Funkcional. Anal. i Priložen.*, 3(1):27– 35, 1969.
- <span id="page-39-5"></span>[20] F. J. Dyson. Existence of a phase-transition in a one-dimensional Ising ferromagnet. *Communications in Mathematical Physics*, 12:91–107, 1969.
- [21] D. S. Fisher, J. Fröhlich, and T. Spencer. The Ising model in a random magnetic field. *Journal of Statistical Physics*, 34(5-6):863–870, 1984.
- <span id="page-39-4"></span>[22] J. Fröhlich and T. Spencer. The Phase Transition in the one-dimensional Ising model with 1*/r*<sup>2</sup> interaction energy. *Communications in Mathematical Physics*, 84:87–101, 1982.
- <span id="page-39-2"></span><span id="page-39-0"></span>[23] M. Kac and C. J. Thompson. Critical Behaviour of Several Lattice Models with Long-Range Interaction. *Journal of Mathematical Physics*, 10:1373–1386, 1969.
- [24] D. Ruelle. Statistical mechanics of a one-dimensional lattice gas. *Communications in Mathematical Physics*, 9:267–278, 1968.