SCHRÖDINGER OPERATORS WITH NON-INTEGER POWER-LAW POTENTIALS AND LIE-RINEHART ALGEBRAS

IVAN BESCHASTNYI, CATARINA CARVALHO, VICTOR NISTOR, AND YU QIAO

ABSTRACT. We study Schrödinger operators $H := -\Delta + V$ with potentials V that have power-law growth (not necessarily polynomial) at 0 and at ∞ using methods of Lie theory (Lie-Rinehart algebras) and microlocal analysis. More precisely, we show that H is "generated" in a certain sense by an explicit Lie-Rinehart algebra. This allows then to construct a suitable (microlocal) calculus of pseudodifferential operators that provides further properties of H. Classically, this microlocal analysis method was used to study H when the power-laws describing the potential V have integer exponents. Thus, the main point of this paper is that this integrality condition on the exponents is not really necessary for the microlocal analysis method to work. While we consider potentials following (possibly non-integer) power-laws both at the origin and at infinity, our results extend right away to potentials having power-law singularities at several points. The extension of the classical microlocal analysis results to potentials with non-integer power-laws is achieved by considering the setting of Lie-Rinehart algebras and of the continuous family groupoids integrating them. (The classical case relies instead on Lie algebroids and Lie groupoids.)

CONTENTS

1. Introduction	2
Contents of the paper	3
Acknowledgements	3
2. Continuous family groupoids and the associated Lie-Rinehart algebras	4
2.1. Locally compact groupoids	4
2.2. Continuous family groupoids	4
2.3. The Lie-Rinehart algebra associated to a CF-groupoid	9
2.4. Universal enveloping algebras, differential, and pseudodifferential operators	11
2.5. CF-groupoids and associated Lie-Rinehart algebras over intervals	12
2.6. Properties of the algebras $(\mathcal{C}_{\phi}^{(\infty)}, \mathcal{V}_{\phi}^{(\infty)})$	13
3. The basic fibered boundary case	18
3.1. Formulation of the problem and main integration results	18
3.2. The case $\psi(\alpha) > 0$	20
3.3. The case $\psi(\alpha) = 0$	22
3.4. Localization and gluing of CF-groupoids	25
3.5. The groupoid associated to Schrödinger operators	26
4. An application to Schrödinger operators	27
References	29

I.B. was supported by the French government through the France 2030 investment plan managed by the National Research Agency (ANR), as part of the Initiative of Excellence Université Côte d'Azur under reference number ANR-15-IDEX-01, C.C. was supported by FCT/Portugal through project UIDB/04459/2020 with DOI identifier 10-54499/UIDP/04459/2020. V.N. was supported by ANR grant OpART ANR-23-CE40-0016, Y.Q. was supported by NSFC (11971282, 12271323).

1. INTRODUCTION

We study the Schrödinger operator with power-law potential:

(1)
$$\begin{cases} H := -\Delta + V, \\ V(x) \sim |x|^{-a}, & \text{as } x \to 0, \text{ and} \\ V(x) \sim |x|^{a'}, & \text{as } x \to \infty, a, a' \in \mathbb{R} \end{cases}$$

using Lie-Rinehart algebras and microlocal analysis. (See Section 4 for precise definitions). This type of potentials arises often in Physics, where *a* is referred to as the "hardness" of an effective potential. In classical physics, this type of potentials models non-hard collisions of particles in real systems, such as colloids [52]. There exist many classical mathematical works dealing with Schrödinger operators with power-law potentials *if a and a' are integers*. The main goal of this paper is to show how the framework of *Lie-Rinehart algebras* and of *continuous family groupoids* can be used to extend some of these classical methods (i.e. for *a* and *a'* integers) to the general case $a, a' \in \mathbb{R}$ not necessarily integers.

Here is a concrete description of our results for the Schrödinger operator $H := -\Delta + V$ defined in Equation (1). Let $M = S^{n-1} \times [0, \infty]$ (where $S^k \subset \mathbb{R}^{k+1}$ is the unit sphere) and let $\phi : M \to [0, \infty]$ be a smooth function such that: $\phi(x) = |x|^a$ if |x| is small, $\phi(x) = |x|^{-a'}$ if |x| is large, and $\phi > 0$ on the rest of M.

(i) A first result is to construct an algebra $\mathcal{A}_{\mathcal{S}}$ of functions

(2)
$$\mathcal{C}_c^{\infty}(M) \subset \mathcal{A}_{\mathcal{S}} \subset \mathcal{C}(M)$$

and a Lie algebra of vector fields $\mathcal{L}_S \subset \mathcal{C}^{\infty}(M;TM)$ acting by derivations on \mathcal{A}_S and such that

(3)
$$\phi(-\Delta+V) \in \operatorname{Diff}(\mathcal{A}_{\mathcal{S}}, \mathcal{L}_{\mathcal{S}})$$

(that is, $\phi(-\Delta + V)$ belongs to the algebra of differential operators on M generated by $\mathcal{A}_{\mathcal{S}}$ and $\mathcal{L}_{\mathcal{S}}$) see Proposition 4.1. This type of results are known to be relevant for the study of the essential spectrums of $\phi(-\Delta + V)$ and $-\Delta + V$ via general results for operators in Diff $(\mathcal{A}_{\mathcal{S}}, \mathcal{L}_{\mathcal{S}})$.

- (ii) A second result is to construct a suitable pseudodifferential calculus Ψ(S) that "quantizes" the algebra Diff(A_S, L_S) and which *contains its parametrices*. Some canonical completions of Ψ(S) will also contain the resolvents of the differential operators in Diff(A_S, L_S). While -Δ + V is not in Ψ(S) (it is φ(-Δ + V) who belongs to Ψ(S)), with some additional work, we can describe also the parametrices and resolvents of -Δ + V (Theorem 4.2).
- (iii) Classically, the construction of pseudodifferential calculi is often achieved by "integrating" $(\mathcal{A}_{\mathcal{S}}, \mathcal{L}_{\mathcal{S}})$ to a Lie groupoid. This is not possible in our case because the functions in $\mathcal{A}_{\mathcal{S}}$ and the vector fields in $\mathcal{L}_{\mathcal{S}}$ are not smooth enough to come from a Lie algebroid (Lie algebroids are the infinitesimal forms of Lie groupoids). Nevertheless, we prove that the pair $(\mathcal{A}_{\mathcal{S}}, \mathcal{L}_{\mathcal{S}})$ is a Lie-Rinehart algebra (Theorem 3.3), (a *Lie-Rinehar* algebra is an algebraic generalization of a Lie algebroid).
- (iv) The natural object integrating a Lie-Rinehart algebra (when this is possible) is a CF-groupoid (the short hand for "continuous family groupoid" [39, 40]). The general problem of integrating a Lie-Rinehart algebra (formulated here) has not yet been solved. It will be studied in more detail in our forthcoming paper [6]. In this paper, we content to integrate the Lie-Rinehart algebra (*A*_S, *L*_S) to a CF-groupoid *S* = *S*_{*γ*,*γ*-1,*γ*'+2,*γ*'+1} for some explicit parameters *γ* and *γ*' given in terms of *a* and *a*' (and hence, in terms of the potentials). Our pseudodifferential calculus is the

pseudodifferential calculus associated to the CF-groupoid S [11, 27, 39]. As mentioned above, it can be shown to contain the parametrices of differential operators in Diff (A_S, \mathcal{L}_S) .

Technically, we first deal with the points (iii) and (iv) above and then we use them to fully establish (i) and (ii). Of course, we establish many other intermediate results, some of which are of independent interest. For instance, a lot of the technical work goes into the construction of the CF-groupoid $S = S_{\tilde{\gamma},\tilde{\gamma}-1,\tilde{\gamma}'+2,\tilde{\gamma}'+1}$ integrating our Lie-Rinehart algebra $(\mathcal{A}_{\mathcal{S}}, \mathcal{L}_{\mathcal{S}})$. The extensive background material that we include is needed for this construction. Nevertheless, we try to go as quickly as possible to the applications related to Schrödinger operators, choosing to include further theoretical developments in the next paper in this series [6].

In addition to allowing us to treat non-integer power-law singularities, our approach (based on Lie-Rinehart algebras and CF-groupoids), allows us to treat differential operators whose coefficients have lower regularity near the boundary. This is critical in dealing with the Schrödinger operator (1).

Another example similar to the power-law Schrödinger operators of Equation (1) involves Riemannian manifolds with boundary and metrics that, in a tubular neighborhood of the boundary, are of the form

$$f = \frac{dx^2 + g_x}{x^a},$$

where x is a boundary-defining function on our manifold with boundary (i.e. $x \ge 0$, the boundary is given by x = 0, and dx does not vanish on the boundary) and g_x is a family of Riemannian metrics on the level sets of the defining function x. As it is known, the case a = 2 corresponds to asymptotically hyperbolic spaces. However, the Laplace-Beltrami operator on the same spaces with fractional values of the parameter a has been proposed recently for studying acoustic modes of gas giants [15]. After a change of variables, these metrics can be put in the form

$$f = dx^2 + x^{-\beta}g_x$$

Manifolds with metrics of this type are called *Grushin manifolds*. They have been studied in the cases of non-integer real parameter β in a number of works [5, 8, 9, 18, 42].

Contents of the paper. In Section 2, we review the notions of Lie-Rinehart algebras and CF-groupoids. We also briefly discuss locally compact and Lie groupoids. In particular, we introduce the Lie-Rinehart algebra, the differential operators, and the pseudodifferential calculus associated to a CF-groupoid [27, 39, 40]. We also recall the concept of the universal enveloping algebra of a Lie-Rinehart algebra and discuss its relation to the algebra of differential operators associated to a CF-groupoid. At the end of this section, we examine one-dimensional examples in detail. In Section 3 we construct the CF-groupoids used to study Schrödinger operators. Our construction relies on the one dimensional examples of the previous section. It is an instance of "integrating" a suitable Lie-Rinehart algebra. In the last section, we apply the results of the previous sections to study Schrödinger operators with (possibly non-integer) power-law singularities at 0 and at ∞ .

Acknowledgements. We thank Camille Laurent-Gengoux and Robert Yuncken for useful discussions. Some results of this paper were announced in [7].

2. CONTINUOUS FAMILY GROUPOIDS AND THE ASSOCIATED LIE-RINEHART ALGEBRAS

In this section we review continuous family groupoids and their associated Lie-Rinehart algebras [27, 39, 40]. We also briefly discuss locally compact and Lie groupoids.

2.1. Locally compact groupoids. Let us begin by recalling the definition of groupoids in general [11, 14, 27, 35, 39, 43]. A *groupoid* is a small category with all morphisms invertible. More precisely, we have the following concrete description of groupoids.

Remark 2.1. A groupoid is determined by the following data:

- (1) a set \mathcal{G} of *morphisms*;
- (2) a set of *composable pairs of morphisms*, denoted by $\mathcal{G}^{(2)} \subset \mathcal{G} \times \mathcal{G}$, together with a *multiplication map* $\mu : \mathcal{G}^{(2)} \to \mathcal{G}$, also written $\mu(g, h) = gh$; and
- (3) an *inverse map* $\iota : \mathcal{G} \to \mathcal{G}$, also written $\iota(q) = q^{-1}$.

This data is required to satisfy the following conditions:

(1) if $(g_1, g_2) \in \mathcal{G}^{(2)}$ and $(g_2, g_2) \in \mathcal{G}^{(2)}$, then

$$(g_1g_2)g_3 = g_1(g_2g_3),$$

(in particular, $(g_1g_2, g_3), (g_1, g_2g_3) \in \mathcal{G}^{(2)}$, i.e. these pairs are composable);

(2) for all $g \in \mathcal{G}$, we have $(g, g^{-1}) \in \mathcal{G}^{(2)}$ and if $(g, h) \in \mathcal{G}^{(2)}$, then

$$g^{-1}(gh) = h$$
 and $(gh)h^{-1} = g$.

We then define the *domain* and *range* maps $d: \mathcal{G} \to \mathcal{G}^{(0)}, r: \mathcal{G} \to \mathcal{G}^{(0)}$ by

$$d(g) := g^{-1}g$$
 and $r(g) := gg^{-1}$,

where the *unit space* $\mathcal{G}^{(0)}$ is defined to be $d(\mathcal{G}) = r(\mathcal{G})$.

Usually, we will write $(\mathcal{G}, d, r, i, \mu)$ or, simply, \mathcal{G} , to denote a groupoid.

A locally compact groupoid is a groupoid $(\mathcal{G}, d, r, i, \mu)$ that is second countable, locally compact space¹ with the space of units $\mathcal{G}^{(0)}$ a closed, Hausdorff subset such that all structural maps are continuous. In the sequel, for subsets $A, B \subset \mathcal{G}^{(0)}$, we denote by $\mathcal{G}_A := d^{-1}(A), \mathcal{G}^B := r^{-1}(B)$, and $\mathcal{G}^B_A := d^{-1}(A) \cap r^{-1}(B)$. In particular, if $x \in \mathcal{G}^{(0)}$, then \mathcal{G}_x and \mathcal{G}^x are the *d*-fiber and *r*-fiber at *x*, respectively. The set \mathcal{G}^A_A is also a groupoid, called the *reduction* of \mathcal{G} to *A*. Moreover, if $\mathcal{G}^A_A = \mathcal{G}_A = \mathcal{G}^A$, then *A* is called *invariant*. In this case, \mathcal{G}_A is also a groupoid. For any $x \in \mathcal{G}^{(0)}, \mathcal{G}^x_x$ is a group, called the *isotropy group* at *x*.

2.2. **Continuous family groupoids.** Our motivation for studying groupoids is that Lie groupoids have been used to study certain partial differential equations with *singularities*, see [1, 2, 3, 4, 5, 7, 11, 14, 25, 32, 45, 48] and the references therein. As explained in the introduction, certain differential operators (such as our Schrödinger operators with non-integer power-law singularities) cannot be studied using Lie groupoids. We prove, however, that they can be studied using continuous family groupoids. We will do that for Schrödinger operators in the last subsection of this paper.

Continuous family groupoids (*CF-groupoids*, for short) were introduced by Paterson [39], who has studied index problems in this setting [40]. Broadly speaking, a continuous family groupoid is such that all *d*-fibers $\mathcal{G}_x := d^{-1}(x)$ and all *r*-fibers $\mathcal{G}^x := r^{-1}(x)$

¹Our locally compact spaces are not required to be Hausdorff; some authors use the terminology "locally quasi-compact" spaces for these spaces.

are smooth manifolds for $x \in \mathcal{G}^{(0)}$, and these fibers vary continuously on x, forming a *continuous family of manifolds* over the units.

We next briefly review the necessary material on continuous families of manifolds and on continuous family groupoids from [39].

2.2.1. *Fiber spaces.* Let X and Y be topological spaces and $p: X \to Y$ be a continuous, open surjection. The pair (X, p) is then called a fiber space over Y with fibers $X^y := p^{-1}(y)$ for $y \in Y$. Let (X_1, p_1) and (X_2, p_2) be fiber spaces over Y_1 and Y_2 , respectively, and let $q: Y_1 \to Y_2$ be a continuous map. A continuous map $f: X_1 \to X_2$ will be called *fiber-preserving with respect to q* if $p_2 \circ f = q \circ p_1$. In case $Y_1 \subset Y_2$ and $q: Y_1 \to Y_2$ is the inclusion, a fiber-preserving map $f: X_1 \to X_2$ with respect to the inclusion q will be called simply *fiber-preserving*.

Let (X, p) be a fiber space over Y (so $p : X \to Y$). Let Z be a topological space and $\rho : Z \to Y$ be a continuous map. The *pull-back* $(\rho^{-1}(X), \pi_2)$ of (Y, p) over Z (along the map ρ) is defined to be

(4)
$$\rho^{-1}(X) := \{(x, z) \in X \times Z \mid \rho(z) = p(x) \in Y\} \subset X \times Z,$$

with fibration map $\pi_2(x, z) := z$ and the topology induced from the product $X \times Z$. It is a fiber space over Z. Let (W, q) be a fiber space over Z. Then a fiber-preserving map $f: W \to X$ over ρ is the same thing as a fiber preserving map $f_{\rho}: W \to \rho^{-1}(X)$ (as Z-fiber spaces), where $f_{\rho}(w) = (f(w), q(w))$. Indeed, f is fiber-preserving over ρ if, and only if, $\rho(q(w)) = p(f(w))$, which in turn is equivalent to $f_{\rho}(w) \in \rho^{-1}(X)$. Moreover, f is continuous if, and only if, f_{ρ} is continuous. This allows to reduce the study of "fiberpreserving maps over a map ρ " to simply "fiber-preserving maps" (i.e. "fiber-preserving maps over the identity").

2.2.2. Continuous families of manifolds and $C^{\infty,0}$ maps. The main difference between Lie groupoids and CF-groupoids is that, in the later, smooth manifolds are replaced by "continuous families of manifolds," a concept that we introduce in this section. To this end, we now look at a particular case of fiber-preserving maps.

Remark 2.2. Let Y be a topological space, A and B be open subsets of $\mathbb{R}^k \times Y$ for some $k \ge 1$, and π_2 be the canonical projection from $\mathbb{R}^k \times Y$ to Y. Then (A, π_2) and (B, π_2) are fiber spaces over subsets of Y and a continuous map $f : A \to B$ is fiber-preserving if, and only if, for every $y \in A$, we have $\pi_2(f(y)) = \pi_2(y)$. The definition implies that $\pi_2(A) \subset \pi_2(B)$. Let us consider a *fiber-preserving function* $f : A \to B$ and let $U_i \subset \mathbb{R}^k$ and $V_i \subset Y$, i = 1, 2, be open subsets such that

$$U_1 \times V_1 \subset A$$
, $U_2 \times V_2 \subset B$, and $f(U_1 \times V_1) \subset U_2 \times V_2$.

Then, for each $y \in V_1$, there exists a unique function $f_y : U_1 \to U_2$ such that

(5)
$$f(x,y) = (f_y(x), y) \in U_2 \times V_2, \text{ where } x \in U_1 \text{ and } y \in V_1.$$

We let $\mathcal{C}^{\infty}(U_1, U_2)$ denote the set of *smooth* functions $U_1 \to U_2$ endowed with the topology of uniform convergence on compacts for all partial derivatives.

We shall often use the notation and concepts introduced in the above remark, without further discussion.

Definition 2.3 (Connes [13], Paterson [39]). Let $A, B \subset \mathbb{R}^k \times Y$ with the induced fiber spaces structures over (subsets of) Y, as in Remark 2.2, whose notation we continue to use. A fiber-preserving function $f : A \to B$ is called a $C^{\infty,0}$ -function (or longitudinally smooth) if, whenever $U_i \subset \mathbb{R}^k$ and $V_i \subset Y$, i = 1, 2, are open subsets such that $U_1 \times V_1 \subset$

A, $U_2 \times V_2 \subset B$, and $f(U_1 \times V_1) \subset U_2 \times V_2$, the induced map $y \to f_y$ is continuous from V_1 to $\mathcal{C}^{\infty}(U_1, U_2)$ (see Equation 5 for the definition of f_y). We let $\mathcal{C}^{\infty,0}(A, B)$ denote the set of $\mathcal{C}^{\infty,0}$ -functions from A to B. A $\mathcal{C}^{\infty,0}$ fiber-preserving homeomorphism $f: A \to B$ such that f^{-1} exists and $f^{-1} \in \mathcal{C}^{\infty,0}(B, A)$ will be called a $\mathcal{C}^{\infty,0}$ -homeomorphism.

This definition allows us now to introduce continuous families of manifolds, as in [39].

Definition 2.4. A fiber space (X, p) over Y is called a continuous family of smooth manifolds over Y if it is paracompact and there exists a set of coordinate charts $\{(U_{\tau}, \varphi_{\tau})\}_{\tau \in I}$, where every U_{τ} is an open subset of X and $\bigcup_{\tau \in I} U_{\tau} = X$, such that

- (1) for every index τ , the map φ_{τ} is a fiber-preserving homeomorphism from U_{τ} to an open subset of $\mathbb{R}^k \times Y$ (recall that fiber preserving means $\pi_2 \circ \varphi_{\tau} = p|_{U_{\tau}}$);
- (2) for every pair of indices τ and β , the composition map

$$\varphi_{\beta} \circ \varphi_{\tau}^{-1} : \varphi_{\tau}(U_{\tau} \cap U_{\beta}) \to \varphi_{\beta}(U_{\tau} \cap U_{\beta})$$

is a $C^{\infty,0}$ -homeomorphism (see Definition 2.3).

(3) The fiber $p^{-1}(y) \subset X$ of p is Hausdorff for every $y \in Y$.

It follows right away from this definition that, if (X, p) is a continuous family of smooth manifolds over Y, then, for every $y \in Y$, the space $p^{-1}(y)$ has a natural smooth manifold structure in the usual sense. More generally, if $q : Z \to Y$ is a continuous map and (X, p)is a continuous family of smooth manifolds over Y, then the pull-back $(q^{-1}(X), \pi_2)$ is a continuous family of smooth manifolds over Z [39] (see Equation (4) for the definition of $(q^{-1}(X), \pi_2)$).

Recall also that our smooth manifolds *do not have boundaries or corners*. Thus a "manifold with corners" is not a "smooth manifold" in our sense. We shall need the following lemma.

Lemma 2.5. Let (X, p) be a continuous family of smooth manifolds over Y and let $f : X \to \mathbb{C}$ be a continuous function such that $f(x_1) = f(x_2)$ for all $x_1, x_2 \in X$ with $p(x_1) = p(x_2)$. Then there exists a continuous function $f_0 : Y \to \mathbb{C}$ such that $f = f_0 \circ p$. The converse is also true (and trivial). Similarly, let $E \to Y$ be a (locally trivial) vector bundle and ξ be a continuous section of $p^*(E)$. There exists a continuous section $\xi_0 : Y \to E$ such that $\xi = \xi_0 \circ p$ if, and only if, $\xi(x_1) = \xi(x_2)$ for all $x_1, x_2 \in X$ with $p(x_1) = p(x_2)$.

Proof. The existence of a function f_0 with these properties is immediate. We only need to prove that it is continuous. Indeed, by the definition of a continuous family of manifolds, it follows that f_0 is locally continuous. But a locally continuous function is continuous. The converse is trivial since p is continuous and hence $f := f_0 \circ p$ is also continuous and has the desired properties. To prove the extension of the statement to vector bundles, we first notice that it is a direct consequence of the first part if $E = Y \times \mathbb{R}^k$ (a trivial vector bundle). By localizing, we can then reduce the general case to that of a trivial vector bundle.

We now extend the definition of $\mathcal{C}^{\infty,0}$ -maps between subsets of $\mathbb{R}^k \times Y$, Definition 2.3 to arbitrary fiber spaces.

Definition 2.6. To define a $C^{\infty,0}$ -*function* between two continuous families of manifolds (X_1, p_1) and (X_2, p_2) , it is convenient to distinguish two cases.

(1) If two continuous families (X_1, p_1) and (X_2, p_2) of manifolds are over the same base Y, a $\mathcal{C}^{\infty,0}$ -function from (X_1, p_1) to (X_2, p_2) (covering the identity of Y) is simply a function $f: X_1 \to X_2$ such that, whenever (U, φ) and (U', φ') are local

SCHRÖDINGER OPERATORS

charts for X_1 and X_2 respectively with $p_1(U) \subset p_2(U')$ and $f(U) \subset U'$ (see Definition 2.4), then $\varphi' \circ f \circ \varphi^{-1} \in \mathcal{C}^{\infty,0}(\varphi(U), \varphi'(U'))$. If this is the case, we write $f \in \mathcal{C}^{\infty,0}(X_1, X_2)$.

(2) Let (X₁, p₁) and (X₂, p₂) be two continuous families of manifolds over Y₁ and Y₂ respectively. Let q : Y₁ → Y₂ be a continuous map and f : X₁ → X₂ be a continuous fiber-preserving map with respect to q, that is, p₂∘f = q∘p₁. Recall the pull-back q⁻¹(X₂), Equation (4). Then f defines a continuous, fiber preserving map f_q : X₁ → q⁻¹(X₂) ⊂ X₂ × Y₁ by the formula f_q(x) = (f(x), p₁(x)). Notice that both X₁ and q⁻¹(X₂) are families of smooth manifolds over the same base Y₁. We say that f is a C^{∞,0}-function (covering q : Y₁ → Y₂) if f_q is a C^{∞,0}-function (covering the identity map of Y₁).

It is not hard to show that the composite of two $C^{\infty,0}$ -maps of continuous families of manifolds is still a $C^{\infty,0}$ -map. A $C^{\infty,0}$ -map is, in particular, fiber preserving and smooth on each fiber.

Remark 2.7. As in the case of usual (single) smooth manifolds, to a continuous family of manifolds (X, p) over Y, we can associate in a functorial way its (vertical) tangent and cotangent bundles $T_{ver}X$ and T_{ver}^*X , see [39].

2.2.3. Fibered products and continuous family groupoids. Suppose that (X_1, p_1) and (X_2, p_2) are two continuous families of manifolds over Y. The fibered product of (X_1, p_1) and (X_2, p_2) is defined to be

(6)
$$(X_1, p_1) * (X_2, p_2) := \{ (x_1, x_2) \in X_1 \times X_2 | p_1(x_1) = p_2(x_2) \} \subset X_1 \times X_2$$

with fiber map $p: (X_1, p_1) * (X_2, p_2) \to Y$ given by $p(x_1, x_2) = p_1(x_1) = p_2(x_2)$. We shall usually write $X_1 * X_2$ instead of $(X_1, p_1) * (X_2, p_2)$ when the maps p_1 and p_2 are obvious. The fiber space $(X_1 * X_2, p)$ is a continuous family of smooth manifolds over Y with respect to the subspace topology. Moreover, the fibered product $X_1 * X_2$ can naturally be regarded as a continuous family of manifolds over X_1 and X_2 respectively. For instance, the pullback of (X_2, p_2) along the map $p_1: X_1 \to Y$ gives the continuous family $(p_1^{-1}(X_2), \pi_1) = (X_1 * X_2, \pi_1)$ of manifolds over X_1 , which is illustrated in the following commutative diagram:

$$X_{1} * X_{2} = p_{1}^{-1}(X_{2}) \xrightarrow{\pi_{2}} > X_{2}$$

$$x_{1} \xrightarrow{\gamma} p_{1} \qquad y = Y$$

$$X_{1} \xrightarrow{\gamma} Y$$

Here we switch the first and second components in the definition of the pullback (see Equation (4)) to match Equation (6). Likewise, we are able to obtain a continuous family $(X_1 * X_2, \pi_2)$ of manifolds over X_2 . We are now in position to recall the definition of continuous family groupoids from [39].

Definition 2.8. A groupoid $(\mathcal{G}, d, r, i, \mu)$ is called a *continuous family groupoid* (*CF*-*groupoid* for short) if the following conditions hold:

- (1) The set $\mathcal{G}^{(0)}$ is locally compact Hausdorff and both (\mathcal{G}, d) and (\mathcal{G}, r) are continuous families of manifolds over $\mathcal{G}^{(0)}$;
- (2) the inverse map $\iota : (\mathcal{G}, d) \to (\mathcal{G}, r)$, where $i(g) = g^{-1}$, is a $\mathcal{C}^{\infty,0}$ -homeomorphism of continuous families of manifolds;

(3) the multiplication map $\mu : ((\mathcal{G}, d) * (\mathcal{G}, r), \pi_1) \to (\mathcal{G}, r)$ is a $\mathcal{C}^{\infty, 0}$ -map of continuous families of manifolds with respect to the range map r:

$$\begin{array}{cccc} \mathcal{G} * \mathcal{G} & \stackrel{\mu}{\longrightarrow} & \mathcal{G} \\ & & & \\ & & & \\ & & & \\ \mathcal{G} & \stackrel{r}{\longrightarrow} & \mathcal{G}^{(0)} \end{array}$$

If \mathcal{G} and \mathcal{G}' are two CF-groupoids, a *CF-groupoids isomorhism* $\Phi : \mathcal{G} \to \mathcal{G}'$ is an isomorphism of groupoids that is also a $\mathcal{C}^{\infty,0}$ -diffeomorphism.

Roughly speaking, a continuous family groupoid $(\mathcal{G}, d, r, i, \mu)$ is such that all *d*-fibers $\mathcal{G}_x := d^{-1}(x)$ (respectively *r*-fibers $\mathcal{G}^x := r^{-1}(x)$) are smooth manifolds for $x \in \mathcal{G}^{(0)}$, and these fibers are required to vary continuously on *x*. Thus \mathcal{G}_x does not have boundary or corners. The structural maps are required to be smooth along the fibers and continuous along the units. An equivalent definition of continuous family groupoids using coordinate charts is given in [27].

Examples 2.9. The following examples will play an important role.

- (1) Let M_1 be a smooth manifold and $\mathcal{P} := M_1 \times M_1$ with d the second projection and r the first projection. Then \mathcal{P} is a Lie groupoid, and hence a CF-groupoid as well, called the *pair groupoid*. However, if M_1 is a manifold with a non-empty boundary, then the pair groupoid M_1^2 is not a CF-groupoid.
- (2) If, moreover, Y is a topological space, then Y × P will be a CF-groupoid. It will be a Lie groupoid precisely if Y is a manifold (possibly with boundary) with d(y, m₁, m₂) := (y, m₂) and r(y, m₁, m₂) := (y, m₁).
- (3) If G is a groupoid and G and G⁽⁰⁾ are manifolds (possibly with corners), all the structural maps d, r, μ, ι, u are smooth, and d is a submersion, then G is a *Lie groupoid*. (Recall that the condition that d be a submersion implies that the fibers G_x := d⁻¹(x) are smooth, boundaryless manifolds.) All Lie groupoids are, in particular, CF-groupoids.
- (4) Let G be a Lie group acting on a CF-groupoid H continuously via C^{∞,0}-isomorphisms. Then we form the *action* (or *semi-direct product*) groupoid H ⋊ G with the same units as H and with structural maps d(h,g) = d(h), r(h,g) = g(r(h)), and product (h,g)(h',g') = (g⁻¹(h')h,gg'). Then H ⋊ G is a CF-groupoid, [39, Proposition 5]
- (5) Let G be a continuous family groupoid with units G⁽⁰⁾ and (X, p) be a fiber space over G⁽⁰⁾. We consider the fiber space (G, d) and the fiber product G * X, see Equation (6). A *continuous action* of G on X is a continuous map ρ : G * X → X (ρ(q, x) = qx, for short) that satisfies the following conditions:
 - (a) $(g_1g_2)y = g_1(g_2y)$, if $(g_1, g_2) \in \mathcal{G}^{(2)}$ (the set of composable arrows),
 - (b) p(gy) = r(g), and
 - (c) $q^{-1}(qy) = y$,

whenever they make sense. Given this data, we can form the *semi-direct product* $X \rtimes \mathcal{G} := (\mathcal{G}, d) \ast (X, p)$. Then the semi-direct product $X \rtimes \mathcal{G} := \mathcal{G} \ast X$ is a CF-groupoid with units X [39, Proposition 5]. If \mathcal{G} is a Lie groupoid, X is a smooth manifold, and the action is smooth, then the semi-direct product groupoid $X \rtimes \mathcal{G} := \mathcal{G} \ast X$ is still a Lie groupoid [35, Page 125].

We shall not use Lie groupoids in this paper for purposes other than to provide examples, but the reader can find more information on Lie groupoids in [7, 14, 31, 31].

2.3. The Lie-Rinehart algebra associated to a CF-groupoid. Lie-Rinehart algebras constitute an algebraic generalisation of Lie algebroids. Recall that a Lie algebroid is given by a smooth vector bundle $A \rightarrow M$ with a Lie algebra structure on its space of smooth sections and a vector bundle map $\rho : A \rightarrow TM$ that satisfies a Leibniz rule. Before discussing Lie-Rinehart algebras associated to continuous family groupoids, let us recall the definition of Lie-Rinehart algebras [44] and give some examples.

Definition 2.10 ([44]). A (unital) *Lie-Rinehart algebra* is a pair $(\mathcal{A}, \mathcal{L})$ that consists of a unital commutative algebra \mathcal{A} and a Lie algebra \mathcal{L} with the following additional structures:

- (1) \mathcal{L} is a left \mathcal{A} -module;
- (2) \mathcal{L} acts by derivations on \mathcal{A} through an \mathcal{A} -module Lie algebra homomorphism $\rho: \mathcal{L} \to \text{Der}(\mathcal{A})$
- (3) The homomorphism ρ satisfies the Leibnitz rule:

$$[X, aY] = \rho(X)(a)Y + a[X, Y]$$

for all $X, Y \in \mathcal{L}, a, b \in \mathcal{A}$.

The map ρ is called the anchor morphism. If $(\mathcal{A}, \mathcal{L})$ and $(\mathcal{B}, \mathcal{M})$ are two Lie-Rinehart algebras, a *Lie-Rinehart isomorhism* $\Phi := (\Phi_{\mathcal{A}}, \Phi_{\mathcal{L}}) : (\mathcal{A}, \mathcal{L}) \to (\mathcal{B}, \mathcal{M})$ is a pair of an algebra and a Lie-algebra isomorphisms compatible with the respective actions.

Lie-Rinehart algebras were considered also in [20, 21, 22, 23, 26, 36, 38, 44]. From now on everywhere in the text we assume that \mathcal{A} is an algebra over \mathbb{K} , where \mathbb{K} is either \mathbb{C} or \mathbb{R} . The same is true for all function spaces, which we assume to be either real or complex valued. Generally, our vector bundles will be over \mathbb{R} , whereas all algebras (Lie or otherwise) will be over \mathbb{C} .

Let $(\mathcal{G}, d, r, i, \mu)$ be a CF-groupoid. We now introduce its associated Lie-Rinehart algebra by analogy with the corresponding construction of the Lie algebroid of a Lie groupoid. Let $\mathcal{G}_x := d^{-1}(x)$, x a unit of \mathcal{G} (that is, $x \in \mathcal{G}^{(0)}$). Then \mathcal{G}_x is endowed with the structure of *a Hausdorff, smooth manifold*, by the definition of a CF-groupoid (recall that our smooth manifolds do not have boundaries or corners). Let $g \in \mathcal{G}$, then, again by definition, the right multiplication by g induces a *diffeomorphism*

(7)
$$\mathcal{G}_{r(g)} \ni h \to hg \in \mathcal{G}_{d(g)}$$

such that R_g varies continuously with g [39, 40]. The same holds for left translations. Recall [39] that the tangent spaces to the fibers $d^{-1}(x)$ of d fit into a (continuous) vector bundle $T_{\text{ver}}\mathcal{G}$ determined by

(8)
$$T\mathcal{G}_x = T_{\mathrm{ver}}\mathcal{G}|_{\mathcal{G}_x}.$$

The vector bundle $T_{ver}\mathcal{G}$ is called the *d*-vertical tangent bundle to \mathcal{G} .

Definition 2.11. Let \mathcal{G} be a CF-groupoid. We let $\mathcal{A}(\mathcal{G})$ consist of all the continuous functions $f : \mathcal{G} \to \mathbb{C}$ with the following properties:

- (1) f is $\mathcal{C}^{\infty,0}$ (recall that this means that it is continuous and its restriction to each $\mathcal{G}_x := d^{-1}(x)$ is smooth).
- (2) f is right G-invariant (that is, for all $g, h \in G$, d(h) = r(g), we have f(hg) = f(h)).

Similarly, we let $\mathcal{L}(\mathcal{G})$ consist of all the continuous sections X of $T_{ver}\mathcal{G}$ with the following properties:

(1) X is $\mathcal{C}^{\infty,0}$ (or longitudinally smooth) in the following sense: it is a continuous section of $T_{\text{ver}}\mathcal{G}$ and its restriction to each $\mathcal{G}_x := d^{-1}(x)$ is a smooth vector field.

(2) X is right \mathcal{G} -invariant (that is, for all $g, h \in \mathcal{G}$, d(h) = r(g), we have $X(hg) = g_*X(h)$).

Then the pair $(\mathcal{A}(\mathcal{G}), \mathcal{L}(\mathcal{G}))$ is called *the Lie-Rinehart algebra associated to* \mathcal{G} .

This definition is justified by the following lemma.

Lemma 2.12. The pair $(\mathcal{A}(\mathcal{G}), \mathcal{L}(\mathcal{G}))$ in Definition 2.11 is a Lie-Rinehart algebra.

Proof. Both the longitudinal smoothness and the right invariance properties are stable under multiplication or under the Lie bracket, and hence $\mathcal{A}(\mathcal{G})$ is a commutative algebra for the pointwise multiplication and $\mathcal{L}(\mathcal{G})$ is a Lie algebra for the Lie bracket of vector fields. Let $f \in \mathcal{A}(\mathcal{G})$ and $X \in \mathcal{L}(\mathcal{G})$. Then $fX \in \mathcal{L}(\mathcal{G})$ and $X(f) \in \mathcal{A}(\mathcal{G})$ by the definitions of $\mathcal{A}(\mathcal{G})$, of $\mathcal{L}(\mathcal{G})$, and of a continuous family of smooth manifolds (in this case, applied to $d: \mathcal{G} \to \mathcal{G}^{(0)}$). Therefore $\mathcal{L}(\mathcal{G})$ is also a $\mathcal{A}(\mathcal{G})$ -module and acts on $\mathcal{A}(\mathcal{G})$, by derivations. The Leibnitz rule is satisfied since the Lie bracket of vector fields satisfies it. \Box

Definition 2.13. Let \mathcal{G} be a CF-groupoid and $T_{\text{ver}}\mathcal{G}$ be the vertical tangent bundle to \mathcal{G} (tangent to the fibers of $d: \mathcal{G} \to \mathcal{G}^{(0)}$). Then

$$\mathbf{A}(\mathcal{G}) := T_{\mathrm{ver}}\mathcal{G}|_{\mathcal{G}^{(0)}}$$

is called the *algebroid associated to* \mathcal{G} . We shall also say that \mathcal{G} *integrates* any Lie-Rinehart algebra isomorphic to $(\mathcal{A}(\mathcal{G}), \mathcal{L}(\mathcal{G}))$.

Clearly, the algebroids of two isomorphic CF-groupoids are also isomorphic. The same is true for their associated Lie-Rinehart algebras.

Remark 2.14. Let us assume that \mathcal{G} is a Lie groupoid with units $\mathcal{G}^{(0)}$. Recall that any Lie groupoid has an associated Lie algebroid $\mathbf{A}(\mathcal{G})$. In turn, to $\mathbf{A}(\mathcal{G})$, one can associate the Lie-Rinehart algebra $(\mathcal{C}^{\infty}(\mathcal{G}^{(0)}), \mathcal{C}^{\infty}(\mathcal{G}^{(0)}, \mathbf{A}(\mathcal{G})))$. This is *not*, however, the Lie-Rinehart algebra associated to \mathcal{G} when regarded as a CF-groupoid (although the algebroid $\mathbf{A}(\mathcal{G})$) will be the same). The reason is that, while $\mathcal{C}^{\infty}(\mathcal{G}^{(0)})$ is always contained in $\mathcal{A}(\mathcal{G})$, the two algebras are usually different. The same comment applies to $\mathcal{C}^{\infty}(\mathcal{G}^{(0)}, \mathbf{A}(\mathcal{G}))$ and $\mathcal{L}(\mathcal{G})$: the second space always contains the first one, but is usually larger. See Remark 2.15. below.

Recall that C(K) denotes the space of continuous functions on some locally compact space K and that C(K; E) denotes the space of continuous sections of some vector bundle $E \to K$.

Remark 2.15. We use the notation of Definition 2.13, then $\mathbf{A}(\mathcal{G})$ is a continuous vector bundle on $\mathcal{G}^{(0)}$ with global (continuous) sections denoted $\mathcal{C}(\mathcal{G}^{(0)}; \mathbf{A}(\mathcal{G}))$. The restriction maps

$$\begin{aligned} \mathcal{A}(\mathcal{G}) &\ni f \to f|_{\mathcal{G}^{(0)}} \in \mathcal{C}(\mathcal{G}^{(0)}) \quad \text{and} \\ \mathcal{L}(\mathcal{G}) &\ni X \to X|_{\mathcal{G}^{(0)}} \in \mathcal{C}(\mathcal{G}^{(0)}; \mathbf{A}(\mathcal{G})) \end{aligned}$$

are injective and we have a canonical isomorphism $T_{ver}\mathcal{G} \simeq r^* \mathbf{A}(\mathcal{G})$.

The algebroid $\mathbf{A}(\mathcal{G})$ associated to the CF-groupoid \mathcal{G} was introduced by Paterson [39]. The algebroid $\mathbf{A}(\mathcal{G})$ was used in [39] to construct a continuous Haar system on \mathcal{G} using the 1-density bundle of $\mathbf{A}(\mathcal{G})$. Let us record the following simple lemma for further use.

Lemma 2.16. Let $f : \mathcal{G} \to \mathbb{C}$ be a continuous, \mathcal{G} -right invariant function (that is, f(hg) = f(h), whenever $g, h \in \mathcal{G}$ are such that d(h) = r(g)). Then there exists a continuous function $f_0 : \mathcal{G}^{(0)} \to \mathbb{C}$ such that, for all $g \in \mathcal{G}$, $f(g) = f_0(r(g))$. The converse is trivially

10

true: any function of the form $f_0 \circ r$ is \mathcal{G} -right invariant. A similar statement holds for sections of $T_{ver}\mathcal{G}$: if X is a right invariant vertical vector field on \mathcal{G} , then there exists a section Y of $\mathbf{A}(\mathcal{G})$ such that $r_*(X) = Y$.

Proof. This is a consequence of the fact that $r : \mathcal{G} \to \mathcal{G}^{(0)}$ is a continuous family of smooth manifolds, of Lemma 2.5, and (for the last part) of the definition relation $T_{\text{ver}}\mathcal{G} \simeq r^*(\mathbf{A}(\mathcal{G}))$.

Remark 2.17. We let the spaces of restriction to $\mathcal{G}^{(0)}$ be given by

$$\mathcal{A}(\mathcal{G})_0 := \{ f_0 \in \mathcal{C}(\mathcal{G}^{(0)}) \mid f_0 \circ r \in \mathcal{A}(\mathcal{G}) \} \text{ and} \\ \mathcal{L}(\mathcal{G})_0 := \{ X \in \mathcal{C}(\mathcal{G}^{(0)}; \mathbf{A}(\mathcal{G})) \mid X \circ r \in \mathcal{L}(\mathcal{G}) \}.$$

The restriction maps define isomorphisms $\mathcal{A}(\mathcal{G}) \simeq \mathcal{A}(\mathcal{G})_0 \subset \mathcal{C}(\mathcal{G}^{(0)})$ and $\mathcal{L}(\mathcal{G}) \simeq \mathcal{L}(\mathcal{G})_0 \subset \mathcal{C}(\mathcal{G}^{(0)}, \mathbf{A}(\mathcal{G}))$. In particular, $(\mathcal{A}(\mathcal{G})_0, \mathcal{L}(\mathcal{G})_0)$ is a Lie-Rinehart algebra.

2.4. Universal enveloping algebras, differential, and pseudodifferential operators. From now on, $(\mathcal{A}, \mathcal{L})$ will denote a unital Lie-Rinehart algebra (Definition 2.10) over $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$. We shall need the concept of the *universal enveloping algebra* $\mathcal{U}(\mathcal{A}, \mathcal{L})$ of $(\mathcal{A}, \mathcal{L})$ [26, 36, 44], which we recall now. First, we endow the \mathbb{K} -vector space $\mathcal{A} \oplus \mathcal{L}$ with the Lie algebra structure with bracket

$$[(a, X), (b, Y)] := (X(b) - Y(a), [X, Y])$$

Denote by $\mathcal{U}(\mathcal{A} \oplus \mathcal{L})$ the *universal enveloping algebra* of the Lie algebra $\mathcal{A} \oplus \mathcal{L}$, which is isomorphic to the quotient of the tensor algebra

$$T(\mathcal{A} \oplus \mathcal{L}) := \mathbb{K} \oplus \left[\oplus_{n \ge 1} (\mathcal{A} \oplus \mathcal{L})^{\otimes n} \right] \supset (\mathcal{A} \oplus \mathcal{L})^{\otimes 1} := \mathcal{A} \oplus \mathcal{L}$$

by the ideal generated by the elements of the form $u \otimes v - v \otimes u - [u, v] \in T(\mathcal{A} \oplus \mathcal{L})$, $u, v \in \mathcal{A} \oplus \mathcal{L}$. Let $i : \mathcal{A} \oplus \mathcal{L} \to \mathcal{U}(\mathcal{A} \oplus \mathcal{L})$ be the canonical inclusion map above, and $\tilde{\mathcal{U}}(\mathcal{A} \oplus \mathcal{L})$ be the subalgebra of $\mathcal{U}(\mathcal{A} \oplus \mathcal{L})$ generated by the image of *i*. Then

(9)
$$\mathcal{U}(\mathcal{A} \oplus \mathcal{L}) = \mathbb{K} \oplus \tilde{\mathcal{U}}(\mathcal{A} \oplus \mathcal{L}).$$

Definition 2.18. Let $\tilde{\mathcal{U}}(\mathcal{A} \oplus \mathcal{L})$ be as in Equation (9). The *enveloping algebra of the Lie-Rinehart algebra* $(\mathcal{A}, \mathcal{L})$ is, by definition, the quotient

$$\mathcal{U}(\mathcal{A},\mathcal{L}) := \mathcal{U}(\mathcal{A} \oplus \mathcal{L})/I,$$

where *I* is the ideal generated by elements of the form i(b, 0)i(a, X) - i(ba, bX).

The enveloping algebra $\mathcal{U}(\mathcal{A}, \mathcal{L})$ can be characterized by the following universal property: if we are given an algebra B, an algebra homomorphism $\phi_{\mathcal{A}} : \mathcal{A} \to B$, and a Lie algebra homomorphism $\phi_{\mathcal{L}} : \mathcal{L} \to B$ (regarding B as a Lie algebra for the commutator) such that $\phi_{\mathcal{A}}(a)\phi_{\mathcal{L}}(X) = \phi_{\mathcal{L}}(aX)$ and $[\phi_{\mathcal{L}}(X), \phi_{\mathcal{A}}(a)] = \phi_{\mathcal{A}}(X(a))$, then there exists a unique homomorphism $\Phi : \mathcal{U}(\mathcal{A}, \mathcal{L}) \to B$ such that $\Phi \circ i_{\mathcal{A}} = \phi_{\mathcal{A}}$ and $\Phi \circ i_{\mathcal{L}} = \phi_{\mathcal{L}}$. This allows us to introduce the following important representation of $\mathcal{U}(\mathcal{A}, \mathcal{L})$.

Definition 2.19 ([36]). Using the universal property of $\mathcal{U}(\mathcal{A}, \mathcal{L})$, we let

$$\rho_{\mathcal{A},\mathcal{L}}: \mathcal{U}(\mathcal{A},\mathcal{L}) \to \operatorname{End}(\mathcal{A})$$

be the unique morphism such that $\rho_{\mathcal{A},\mathcal{L}}(i_{\mathcal{A}}(a))$ acts by left multiplication by $a \in \mathcal{A}$ and $\rho_{\mathcal{A},\mathcal{L}}(i_{\mathcal{L}}(\mathcal{G}))$ acts by derivations on \mathcal{A} . We also let $\text{Diff}(\mathcal{A},\mathcal{L}) := \rho_{\mathcal{A},\mathcal{L}}(\mathcal{U}(\mathcal{A},\mathcal{L}))$ be the *algebra of differential operators* associated to $(\mathcal{A},\mathcal{L})$.

In case $(\mathcal{A}, \mathcal{L}) = (\mathcal{A}(\mathcal{G}), \mathcal{L}(\mathcal{G}))$ is associated to the CF-groupoid \mathcal{G} , then we shall write $\text{Diff}(\mathcal{A}, \mathcal{L}) = \text{Diff}(\mathcal{G})$.

In the examples considered in this article, $\rho_{\mathcal{A},\mathcal{L}}$ is injective and hence

$$\mathrm{Diff}(\mathcal{A},\mathcal{L}) := \rho_{\mathcal{A},\mathcal{L}}(\mathcal{U}(\mathcal{A},\mathcal{L})) \simeq \mathcal{U}(\mathcal{A},\mathcal{L})$$

General injectivity results of this kind for $\rho_{\mathcal{A},\mathcal{L}}$ will be included in [6]. If $\rho_{\mathcal{A},\mathcal{L}}$ is injective, we may regard Diff $(\mathcal{A},\mathcal{L})$ as the subalgebra of End (\mathcal{A}) generated by \mathcal{A} , acting as multiplication operators, and by \mathcal{L} , acting as derivations. If, moreover, $(\mathcal{A},\mathcal{L})$ is associated to the CF-groupoid \mathcal{G} , then from Remark 2.15, we have $\mathcal{A} = \mathcal{A}(\mathcal{G})_0$ acting injectively on \mathcal{A} as multiplication operators, and $\mathcal{L} = \mathcal{L}(\mathcal{G})_0 \subset \mathcal{C}(\mathcal{G}^{(0)}; \mathbf{A}(\mathcal{G}))$ also acting injectively as derivations, so Diff (\mathcal{G}) is isomorphic to the subalgebra of End (\mathcal{A}) generated by functions in $\mathcal{A}(\mathcal{G})_0$ and vector fields in $\mathcal{L}(\mathcal{G})_0$, which can then be regarded as differential operators of order 1.

To obtain our final results on the Schrödinger operator, we will need pseudodifferential operators. One important feature is that $\text{Diff}(\mathcal{G})$ maps injectively to (it is even included in) the much larger class $\Psi(\mathcal{G})$ of pseudodifferential operators on \mathcal{G} . Recall the following definition [27, 39].

Definition 2.20. The set $\Psi^m(\mathcal{G})$ consists of right invariant, continuous families of pseudodifferential operators of order m on $\mathcal{G}_x := d^{-1}(x)$ with compactly supported convolution kernels k_P (which are distributions on \mathcal{G}).

Similarly to the representation $\rho_{\mathcal{A},\mathcal{L}}$, an operator $P \in \Psi^m(\mathcal{G})$ acts on $\mathcal{A}(\mathcal{G})$ via a map $\pi_0(P)$, and we have $\text{Diff}(\mathcal{G}) := \text{Diff}(\mathcal{A}(\mathcal{G}), \mathcal{L}(\mathcal{G})) \subset \pi_0(\Psi^\infty(\mathcal{G}))$. The morphism π_0 is not injective in general, but it will be in the examples considered in this paper. Note that we will not need this class of operators up to section 4.

2.5. **CF-groupoids and associated Lie-Rinehart algebras over intervals.** We now look at some of the simplest examples of Lie-Rinehart algebras. The Lie-Rinehart algebras present some analytic subtleties, as it will be apparent even from the one dimensional examples studied in this section. These examples will also be an important building block in the constructions of the next section. We agree that $\mathbb{N} = \{1, 2, ...\}$ and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. Let $\mathcal{C}(K)$ denote the space of continuous functions on some compact space K, as before.

Let us introduce now some basic spaces. To this end, let $N \in \mathbb{Z}_+ \cup \{\infty\}$ and $\alpha, \beta \in [-\infty, \infty] =: \overline{\mathbb{R}}, \alpha < \beta$. We consider

(10)
$$X(t) := \phi(t)\partial_t, \quad \phi : (\alpha, \beta) \to (0, \infty) \quad continuous,$$
$$\mathcal{C}_{\phi}^{(N)} := \{ u : [\alpha, \beta] \to \mathbb{R} \mid X^k u \in \mathcal{C}([\alpha, \beta]) \text{ for all } 0 \le k \le N \}, \quad \text{and}$$
$$\mathcal{V}_{\phi}^{(N)} := \{ uX \mid u \in \mathcal{C}_{\phi}^{(N)} \}.$$

We shall write $\mathcal{C}_{\phi}^{(N)} = \mathcal{C}_{\phi}^{(N)}([\alpha,\beta])$ and $\mathcal{V}_{\phi}^{(N)} = \mathcal{V}_{\phi}^{(N)}([\alpha,\beta])$ when we want to be more precise. The relation " $X^{k}u \in \mathcal{C}([\alpha,\beta])$ " is defined by induction to mean that " $X^{k-1}u$ is differentiable on (α,β) and $X^{k}u := \phi(X^{k-1}u)'$ extends to a continuous function on $[\alpha,\beta]$ ". Note that $\mathcal{C}_{\phi}^{(\infty)} = \bigcap_{n \in \mathbb{Z}_{+}} \mathcal{C}_{\phi}^{(n)}$ and $\mathcal{C}_{\phi}^{(0)} = \mathcal{C}([\alpha,\beta])$. We fix a function ϕ and a vector field X as in Equation (10) for the rest of this section.

We fix a function ϕ and a vector field X as in Equation (10) for the rest of this section. The main examples that will concern us will be $\phi(t) = (t - \alpha)^a (\beta - t)^b$, with a, b > 0(even ≥ 1 for the most part). Most of our example will be such that ϕ is differentiable on the open interval (α, β) , but this is by no means necessary for the development of the theory, so it will not be assumed. We have the following simple remark. **Remark 2.21.** Let us assume that ϕ is smooth on (α, β) . The assumption that $\phi(t) > 0$ gives then that every $u \in \mathcal{C}_{\phi}^{(\infty)}$ is smooth on (α, β) . In fact, we even have

$$\mathcal{C}^{\infty}_{c}((\alpha,\beta)) \subset \mathcal{C}^{(\infty)}_{\phi} \subset \mathcal{C}([\alpha,\beta]) \cap \mathcal{C}^{\infty}((\alpha,\beta)).$$

If ϕ is smooth on a *bounded* interval $[\alpha, \beta]$, then $\mathcal{C}^{\infty}([\alpha, \beta]) \subset \mathcal{C}^{(\infty)}_{\phi}$, but, in general, $\mathcal{C}^{\infty}([\alpha,\beta]) \neq \mathcal{C}^{(\infty)}_{\phi}$, and this is one of the point of considering Lie-Rinehart algebras: it allows for more general functions than the usual smooth functions.

We have the following simple, technical result.

Proposition 2.22. We shall use the notation of Equation (10). Let $n \in \mathbb{Z}_+ \cup \{\infty\}$. Then:

(1) $\mathcal{C}_{\phi}^{(n)}$ is an algebra. (2) $\mathcal{V}_{\phi}^{(n)}\mathcal{C}_{\phi}^{(n+1)} \subset \mathcal{C}_{\phi}^{(n)}.$ (3) $[\mathcal{V}_{\phi}^{(n+1)}, \mathcal{V}_{\phi}^{(n+1)}] \subset \mathcal{V}_{\phi}^{(n)}.$

In particular, $(\mathcal{C}^{(\infty)}_{\phi}, \mathcal{V}^{(\infty)}_{\phi})$ is a Lie-Rinehart algebra.

Proof. We proceed in order. It suffices to assume $n \in \mathbb{Z}^+$, as the results for $n = \infty$ will then follow. (1) Let us show that $\mathcal{C}_{\phi}^{(n)}$ is an algebra. This is clear if n = 0, so let us assume that n > 0. We have X(fg) = X(f)g + fX(g), as $X := \phi \partial_t$ is a derivation. We obtain, for $k \in \mathbb{Z}_+, k \leq n$,

(11)
$$X^{k}(uv) = \sum_{j=0}^{n} C_{k}^{j} X^{j}(u) X^{k-j}(v),$$

where $C_k^j = \frac{k!}{j!(k-j)!}$, $0 \le j \le n$ are the usual binomial coefficients. It follows that, if $X^{j}(u)$ and $X^{k-j}(v)$ are continuous on $[\alpha, \beta]$ for all $j = 0, \ldots, k$, then $X^{k}(uv)$ will also be continuous on $[\alpha, \beta]$. In particular, if $u, v \in C_{\phi}^{(n)}$ then $uv \in C_{\phi}^{(n)}$, so $C_{\phi}^{(n)}$ is an algebra. (2) follows immediately from (1). Indeed, we have $XC_{\phi}^{(n+1)} \subset C_{\phi}^{(n)}$ by definition

(Equation 10). Let $u \in \mathcal{C}_{\phi}^{(n)}$ and $v \in \mathcal{C}_{\phi}^{(n+1)}$. Then

$$(uX)(v) := u(X(v)) \in \mathcal{C}_{\phi}^{(n)}$$

since $u, X(v) \in C_{\phi}^{(n)}$, which was proved to be an algebra. (3) also follows immediately from (1) and (2) in view of the Leibnitz rule, which yields the identity [uX, vX] = [uX(v) - vX(u)]X.

In particular, $\mathcal{C}_{\phi}^{(\infty)}$ is a commutative algebra, $\mathcal{V}_{\phi}^{(\infty)}$ is a Lie algebra, by (3), which is also a left \mathcal{A} -module, and acts by derivations on $\mathcal{C}_{\phi}^{(\infty)}$, as X does. Moreover, if $a \in \mathcal{C}_{\phi}^{(\infty)}$ and $uX, vX \in \mathcal{V}_{\phi}^{(\infty)}$, then

$$[uX, avX] = (uX(av) - avX(u))X = (uX)(a)vX + a[uX, vX].$$

Hence, the Leibnitz rule is satisfied and $(\mathcal{C}_{\phi}^{(\infty)}, \mathcal{V}_{\phi}^{(\infty)})$ is a Lie-Rinehart algebra.

2.6. Properties of the algebras $(\mathcal{C}_{\phi}^{(\infty)}, \mathcal{V}_{\phi}^{(\infty)})$. We continue to fix $\alpha, \beta \in \mathbb{R}$ and a continuous function $\phi : (\alpha, \beta) \to (0, \infty)$, as in the definition of the Lie-Rinehart algebra $(\mathcal{C}^{(\infty)}_{\phi}, \mathcal{V}^{(\infty)}_{\phi})$ of Equation (10).

Lemma 2.23. Let $\psi : (\alpha, \beta) \to (0, \infty)$ be a continuous function and $\mathcal{C}_{\psi}^{(\infty)}$ be defined as in (10).

(1) If $\phi/\psi \in \mathcal{C}_{\psi}^{(\infty)}$, then $\mathcal{C}_{\psi}^{(\infty)} \subset \mathcal{C}_{\phi}^{(\infty)}$. (2) Consequently, if $\phi/\psi \in \mathcal{C}_{\psi}^{(\infty)}$ and $\psi/\phi \in \mathcal{C}_{\phi}^{(\infty)}$, then $\mathcal{C}_{\phi}^{(\infty)} = \mathcal{C}_{\psi}^{(\infty)}$.

Proof. Let $Y := \psi \partial_t$. Then $\mathcal{C}_{\psi}^{(\infty)} := \{u : [\alpha, \beta] \to \mathbb{C} \mid Y^k u \in \mathcal{C}_{\psi}^{(\infty)}, k \in \mathbb{Z}_+\}$, by definition, Equation (10). Let $X := \phi \partial_t = \frac{\phi}{\psi} Y$. The assumption that $\frac{\phi}{\psi} \in \mathcal{C}_{\psi}^{(\infty)}$ gives that $X \in \mathcal{V}_{\psi}^{(\infty)}$. Since $\mathcal{V}_{\psi}^{(\infty)}$ acts by derivations on $\mathcal{V}_{\psi}^{(\infty)}$, we have that $X^k u \in \mathcal{V}_{\psi}^{(\infty)} \subset \mathcal{C}$, for all $k \in \mathbb{Z}_+$, and hence $u \in \mathcal{V}_{\phi}^{(\infty)}$, by definition. The second statement follows from the first one, by symmetry.

We shall need the following lemma.

Lemma 2.24. Let $X := \phi \partial_t$, as before, and $n, j \in \mathbb{Z}_+$.

(1) Let us assume $\phi' \in C_{\phi}^{(n+j-2)}$ if $n \ge 2$ (we impose no condition if n = 0 or n = 1). Then there exist $a_k \in C_{\phi}^{(j)}$, k = 0, ..., n - 1, such that

$$X^{n}u = \sum_{k=0}^{n-1} a_{k}\phi^{k}\partial_{t}^{k}u + \phi^{n}\partial_{t}^{n}u.$$

(2) Let us assume $\phi' \in C_{\phi}^{(n-2)}$ if $n \geq 2$. We have $u \in C_{\phi}^{(n)}$ if, and only if $\phi^k \partial_t^k u \in C([\alpha, \beta])$, for all k = 0, 1, ..., n.

Proof. (1) Let $S_n := \{a_j \neq 0\}$ be the set of non-zero coefficients of $X^n = \sum_{k=0}^n a_k \phi^k \partial_t^k$. We shall prove the statement (i.e. $S_n \subset C_{\phi}^{(j)}$ if $\phi' \in C_{\phi}^{(n+j-2)}$) by induction on n (and for all j). We have $S_0 = S_1 = \{1\}$ by definition and $S_2 = \{1, \phi'\}$. Hence the statement is true for $n \leq 2$ (for n = 2 this is because we have assumed that $\phi' \in C_{\phi}^{(j)}$).

Let us prove the induction step. Let us assume $\phi' \in C_{\phi}^{(n+1+j-2)}$ for some $n \ge 2$. Then $S_n \subset C_{\phi}^{(j+1)}$, by the induction hypothesis. Next,

$$X^{n+1}u = X\left(\sum_{k=0}^{n} a_k \phi^k \partial_t^k u\right)$$
$$= \sum_{k=0}^{n} \left[X(a_k) + ka_k \phi' + a_{k-1}\right] \phi^k \partial_t^k u + a_n \phi^{n+1} \partial_t^{n+1} u,$$

(where we have set $a_{-1} := 0$). Hence $\mathcal{S}_{n+1} \subset \mathcal{C}_{\phi}^{(j)}$.

(2) Follows by induction from definitions using (1) for j = 0.

We obtain the following consequence.

Corollary 2.25. Let $n \in \mathbb{Z}_+ \cup \{\infty\}$. We have $\phi' \in \mathcal{C}_{\phi}^{(n)}$ if, and only if, $\phi^k \partial_t^{k+1} \phi \in \mathcal{C}([\alpha, \beta])$ for all $0 \leq k \leq n$.

Proof. The proof is again by induction on n. For n = 0 and n = 1 this is checked directly. Let us assume the statement to be true for some $n \ge 1$ and prove it for n + 1. Assume $\phi' \in C_{\phi}^{(n+1)}$. Then Lemma 2.24 (2) for $u = \phi'$ gives $\phi^k \partial_t^k \phi' \in C([\alpha, \beta])$ for all $0 \le k \le n + 1$. Conversely, if $\phi^k \partial_t^k \phi' \in C([\alpha, \beta])$ for all $0 \le k \le n + 1$, then the

induction hypothesis gives $\phi' \in C_{\phi}^{(n)}$ Hence we can use Lemma 2.24 (2) again for $u = \phi'$ to conclude that, in fact, $\phi' \in C_{\phi}^{(n+1)}$.

Similarly, we obtain the following consequence.

Corollary 2.26. Let us assume that $\phi^k \partial_t^{k+1} \phi \in \mathcal{C}([\alpha, \beta])$ for all $k \in \mathbb{Z}_+$ and let $\eta \in \mathcal{C}([\alpha, \beta])$ be such that $\phi^k \partial_t^k \eta \in \mathcal{C}([\alpha, \beta])$ for all $k \in \mathbb{Z}_+$. Then $\eta \in \mathcal{C}_{\phi}^{(\infty)}$.

Let $\psi: [\alpha, \beta] \to [0, \infty), \psi(t) > 0$ for $t \in (\alpha, \beta)$ be continuous. Also, let

(12)
$$C_{\psi,\phi} := \frac{\phi\psi'}{\psi} = X(\psi)/\psi,$$

which is defined initially for $t \in (\alpha, \beta)$, but which we extend to α and β by continuity, when possible. We shall need yet the following corollary.

Corollary 2.27. Let us assume that $\phi(t) = (t - \alpha)^a$ and $\psi(t) = (t - \alpha)^b$ for t close to α and that $\phi(t) = (\beta - t)^{a'}$ and $\psi(t) = (\beta - t)^{b'}$ for t close to β . Assume $a, a' \ge 1$ and $b, b' \ge 0$. Then $\psi \in C_{\phi}^{(\infty)}$ and also

$$\phi', C_{\psi,\phi} := \phi \psi' / \psi \in \mathcal{C}_{\phi}^{(\infty)}$$

Proof. The fact that $\psi \in C_{\phi}^{(\infty)}$ follows directly from the definition since, for instance, near α , $X^k \psi = b(b + a - 1) \dots (b + (k - 1)(a - 1))(x - \alpha)^{b+k(a-1)}$, which is continuous. Recall that $C_{\psi,\phi} := \frac{\phi \psi'}{\psi} = X(\psi)/\psi$. Let us check the conditions of Corollaries 2.25 and 2.26 near α if b > 0. Indeed, for $t \in [\alpha, \beta]$ close to α , $C_{\psi,\phi}(t) = b(t - \alpha)^{a-1}$ and

$$a^{-1}\phi^k\phi^{(k+1)}(t) = b^{-1}\phi^k\partial_t^k C_{\psi,\phi}(t) = (a-1)\dots(a-k)(t-\alpha)^{(k+1)(a-1)}$$

are both continuous near α (including α). If b = 0, everything is the same, except the last formula, in which we have $C_{\psi,\phi} = 0$, and hence the same conclusion. The calculations near β are similar.

We remark that, if ϕ and ψ are as in the last corollary, then

$$\begin{cases} C_{\psi,\phi}(\alpha) = b & \text{if } a = 1 \\ C_{\psi,\phi}(\alpha) = 0 & \text{if } a > 1 \end{cases}$$

A similar relation holds true at β :

$$\begin{cases} C_{\psi,\phi}(\beta) = b' & \text{if } a' = 1\\ C_{\psi,\phi}(\beta) = 0 & \text{if } a' > 1. \end{cases}$$

As we have seen $(\mathcal{C}_{\phi}^{(\infty)}, \mathcal{V}_{\phi}^{(\infty)})$ is a Lie-Rinehart algebra, it is natural to ask when it comes from a CF-groupoid? We shall see that this is the case if ϕ (or $X := \phi \partial_t$) is *complete* in the sense that

(13)
$$\int_{\alpha}^{\gamma} \frac{1}{\phi(t)} dt = \int_{\gamma}^{\beta} \frac{1}{\phi(t)} dt = \infty$$

for one (equivalently, for any) $\gamma \in (\alpha, \beta)$. We fix from now on $\gamma \in (\alpha, \beta)$. Let, for further reference,

(14)
$$F(x) := \int_{\gamma}^{x} \frac{dt}{\phi(t)}$$

This notation will be fixed throughout the rest of the paper.

Proposition 2.28. Let us assume that our continuous function $\phi : (\alpha, \beta) \to (0, \infty)$ satisfies the completeness conditions of Equation (13). Then $X := \phi \partial_t$ generates a one parameter group of homeomorphisms σ_s , $s \in \mathbb{R}$, of $[\alpha, \beta]$ that are C^1 on (α, β) .

What this proposition says is that under the completeness condition we have a well defined flow

(15)
$$\sigma_s : [\alpha, \beta] \to [\alpha, \beta], \quad \sigma_s \circ \sigma_t = \sigma_{s+t}, \ s, t \in \mathbb{R},$$

which as usually means

(16)
$$\frac{d}{ds}\sigma_s(x) = \phi(\sigma_s(x)), \text{ for } x \in (\alpha, \beta),$$

and which is equivalent to the usual relation

(17)
$$\frac{d}{ds}f(\sigma_s(x)) = [Xf](\sigma_s(x)).$$

Proof. Let F be as in Equation (14). Then, our assumptions on ϕ give that $F : (\alpha, \beta) \to (-\infty, \infty)$ is \mathcal{C}^1 -homeomorphism. Then the desired flow σ_s is given by

(18)
$$\sigma_s(x) = F^{-1}(F(x) + s).$$

Note that even though ϕ is not Lipschitz, the flow σ_s is well defined. Indeed, Any two solutions t(s) and $\tilde{t}(s)$ of the differential equation

$$y'(t) = \phi(y(t)),$$

with $y(0) = \tilde{y}(0) = \gamma$ must satisfy $F(y(s)) = F(\tilde{y}(s))$, for all $s \in \mathbb{R}$. By assumption we have $\phi(x) > 0$, for $x \in (\alpha, \beta)$. For this reason, F is a strictly monotone increasing function and therefore one-to-one. Hence the two solutions y and \tilde{y} must agree and we get uniqueness of the Cauchy problem and a well-defined flow on the open interval (α, β) .

Since $\lim_{x\to\alpha} F(x) = -\infty$ and $\lim_{y\to-\infty} F^{-1}(y) = \alpha$, we obtain that $\lim_{x\to\alpha} \sigma_s(x) = \alpha$. Similarly, $\lim_{x\to\beta} \sigma_s(x) = \beta$. We shall then set

(19)
$$\sigma_s(\alpha) := \alpha \text{ and } \sigma_s(\beta) := \beta.$$

In this way, the resulting maps σ_s define a one-parameter group of *homeomorphisms* of $[\alpha, \beta]$ that are C^1 -homeomorphisms in the interior (α, β) .

Let us give some concrete examples, i.e., start with the simplest example, namely, the one that underpins the so called "*b*-calculus" of Melrose [33] and Schulze [47]. See also [19, 29, 30, 46].

Example 2.29. Let $(\alpha, \beta) = (0, \infty)$ and $\phi(t) = t$. Then the function F of Equation (14) is given by $F(x) = \ln x$, which is a diffeomorphism $F : (0, \infty) \to \mathbb{R}$ with $F^{-1}(y) = \exp(y) = e^y$. Hence the flow $\sigma_s, s \in \mathbb{R}$, of Proposition 2.28 will be given by

$$\sigma_s(x) = F^{-1}(F(x) + s) = \exp(\ln x + s) = e^s x.$$

We note that, in this example, σ_s is smooth also at 0.

This example can be generalized as follows.

Example 2.30. Let $\alpha = 0$, $\beta > \gamma = 1$, and a > 1 (the case a = 1 was treated in the previous example and here *a* is *not* assumed to be an integer!). Let $\phi(t) = t^a$ for $t \le 1$ and

arbitrary for t > 1, as long as it satisfies the needed conditions, namely ϕ is continuous and > 0 on (α, β) (Equation 10) and Equation (13). Then

$$\begin{cases} F(x) = \frac{1-x^{1-a}}{a-1} & \text{for } x \le 1 \text{ and} \\ F^{-1}(y) = [1-(a-1)y]^{\frac{1}{1-a}} & \text{for } y \le 0 \,. \end{cases}$$

Therefore, if $x \leq 1$ and $F(x) + s \leq 0$, we obtain

$$\sigma_s(x) = F^{-1}(F(x) + s)$$

= $[1 - (a - 1)sx^{a-1}]^{\frac{1}{1-a}}x$

In particular, $\sigma_s: [0,\beta] \to [0,\beta]$ is \mathcal{C}^{∞} at 0 if, and only if, a is an integer.

Yet another generalization of Example 2.29 is to a *bounded* interval.

Example 2.31. Let $[\alpha, \beta] = [-1, 1]$ and $\phi(t) = (1 + t)(1 - t)$. Then $F(x) = \frac{1}{2} \ln \left| \frac{1+x}{1-x} \right|$ is a diffeomorphism $F : (-1, 1) \to \mathbb{R}$ with $F^{-1}(y) = \tanh(y)$. Therefore $\sigma_s(x) = \tanh(F(x) + s)$ and, in this case, σ_s extends to a \mathcal{C}^{∞} diffeomorphism $[-1, 1] \to [-1, 1]$, for all s.

The point is that if ϕ satisfies the completeness condition of Equation (13), we can use the flow σ_s to construct a CF-groupoid whose Lie-Rinehart algebra is exactly $(\mathcal{C}_{\phi}^{(\infty)}, \mathcal{V}_{\phi}^{(\infty)})$. More precisely, consider the group-action groupoid:

(20)
$$\mathcal{G}_{\phi} := [\alpha, \beta] \rtimes_{\sigma} \mathbb{R}$$

with the usual structural maps (see [13, 14, 43, 53])

(21)
$$\begin{cases} d(x,t) := x \\ r(x,t) := \sigma_t(x) \\ (y,s)(x,t) = (x,s+t) & \text{if } \sigma_t(x) = y \end{cases}$$

where σ_s is the flow given by Proposition 2.28. The particular case $\phi \equiv 1$ and $[\alpha, \beta] = [-\infty, +\infty] = \overline{\mathbb{R}}$ will play a special role; the associated groupoid (namely \mathcal{G}_{ϕ}) will also be denoted $\mathcal{G}_{\phi} = \mathcal{G}_1$, for simplicity. So $\mathcal{G}_1 := \overline{\mathbb{R}} \rtimes \mathbb{R}$, with the action of \mathbb{R} on $\overline{\mathbb{R}}$ by translations.

Theorem 2.32. Let us assume that $\phi : (\alpha, \beta) \to (0, \infty)$ is continuous and that it satisfies the completeness condition of Equation (13), as before. Then the group-action groupoid \mathcal{G}_{ϕ} of Equations (20) and (21) is a CF-groupoid such that $F : [\alpha, \beta] \to \mathbb{R}$ induces an isomorphism

$$\mathcal{G}_{\phi} \simeq \mathcal{G}_1$$
.

We hence have $\mathcal{A}(\mathcal{G}_{\phi})_0 = \mathcal{C}_{\phi}^{(\infty)}$ and $\mathcal{L}(\mathcal{G}_{\phi})_0 = \mathcal{V}_{\phi}^{(\infty)}$ (see Equation (10) and Remark (2.17) for notation). Let us assume that the interval $[\alpha, \beta]$ is bounded and it is endowed with the usual smooth structure, then \mathcal{G}_{ϕ} is a Lie groupoid if, and only if, the function ϕ is smooth on $[\alpha, \beta]$.

Proof. That \mathcal{G}_{ϕ} is a CF-groupoid follows directly from [39, Proposition 5] (see the Example 2.9(4) for the needed statement).

The relations $\mathcal{A}(\mathcal{G}_{\phi})_0 = \mathcal{C}_{\phi}^{(\infty)}$ and $\mathcal{L}(\mathcal{G}_{\phi})_0 = \mathcal{V}_{\phi}^{(\infty)}$ can be checked directly if $[\alpha, \beta] = \overline{\mathbb{R}}$ and $\phi \equiv 1$. They are hence true in general since the diffeomorphism $F : (\alpha, \beta) \to \mathbb{R}$ of Equation (14) maps $X := \phi \partial_t$ to ∂_t , by construction, hence $\mathcal{C}_{\phi}^{(\infty)} = \mathcal{C}_1^{(\infty)}, \mathcal{V}_{\phi}^{(\infty)} = \mathcal{V}_1^{(\infty)}$. Moreover, $\mathcal{G}_{\phi} \simeq \mathcal{G}_1$, so the associated Lie-Rinehart algebras are also isomorphic.

If the groupoid \mathcal{G}_{ϕ} is a Lie groupoid, then r must be smooth, and hence the action σ_t (with σ_t the flow of Proposition 2.28) will be smooth on $[\alpha, \beta]$. Consequently, ϕ will also be smooth on $[\alpha, \beta]$. Conversely, if ϕ is smooth on $[\alpha, \beta]$, then the standard results on the differentiability of solutions of ODEs gives that σ_t is smooth on $[\alpha, \beta]$.

Corollary 2.33. For example, assume in addition that $a, b \ge 1$ are such that $\phi(t) = (t - \alpha)^a$ for t close to α and $\phi(t) = (\beta - t)^b$ for t close to β . Then \mathcal{G}_{ϕ} is a Lie groupoid if, and only if, $a, b \in \mathbb{N}$.

Our results raise the following integrability question: "Which Lie-Rinehart algebras are associated to some CF-groupoids?" The gluing construction of [37] (integrating on each orbit and then gluing) seems to be still relevant here. In the following, we shall provide some constructions of CF-groupoids associated to some given Lie-Rinehart algebras particularly useful for studying Schrödinger operators with singular potentials.

3. The basic fibered boundary case

We now construct the CF-groupoids used to study Schrödinger operators. Our construction relies on the one dimensional examples of the previous section. It is an instance of "integrating" a suitable Lie-Rinehart algebra.

3.1. Formulation of the problem and main integration results. We now use the results proved so far to investigate more complicated Lie-Rinehart algebras and their associated CF-groupoids. Let M_1 be a smooth, compact manifold of dimension d (i.e. a closed manifold) and, for the rest of this section, we let

$$M := [\alpha, \beta] \times M_1$$
.

We continue to consider a continuous function $\phi : (\alpha, \beta) \to (0, \infty)$ that is subject to the same assumptions as before (namely, Equations 10 and 13). To fix ideas, we shall also assume that ϕ is *smooth on* (α, β) and extends by continuity to $[\alpha, \beta]$ such that $\phi(\alpha) = \phi(\beta) = 0$. We then lift ϕ to a function $[\alpha, \beta] \times M_1 \to [0, \infty)$, still denoted ϕ , by abuse of notation. In addition to the function ϕ , we consider a second function ψ with properties similar to those of ϕ (and those of the function ψ in the previous subsection), namely,

(22)
$$\begin{cases} \psi \in \mathcal{C}_{\phi}^{(\infty)} \\ \psi(t) > 0 \text{ on } (\alpha, \beta) \\ C_{\phi,\psi} := \phi(t)\psi'(t)/\psi(t) \text{ extends to a function in } \mathcal{C}_{\phi}^{(\infty)} \end{cases}$$

(Recall that $\mathcal{C}_{\phi}^{(\infty)}$ was introduced in Equation (10) and that $\mathcal{C}_{\phi}^{(\infty)} \subset \mathcal{C}([\alpha, \beta])$.) For simplicity, we shall also assume that $\alpha, \beta \in \mathbb{R}$ (that is, that $[\alpha, \beta]$ is bounded).

Let us begin with a heuristical formulation of our problem. Let us place ourselves in a local coordinate chart, so M_1 is an open subset of \mathbb{R}^d . We shall use the functions ϕ and ψ described in the previous paragraph. We consider vector fields of the form

$$X := Y_0 := \phi \partial_t, \text{ and } Y_k := \psi \partial_{x_k}, \quad 1 \le k \le d,$$

on $[\alpha, \beta] \times M_1 \ni (t, x)$ and we will construct a CF-groupoid whose Lie-Rinehart algebra is generated by these vector fields.

To formulate more precisely our problem, let us introduce the following notation. Let $\pi_0 : [\alpha, \beta] \times M_1 \to [\alpha, \beta]$ and $\pi_1 : [\alpha, \beta] \times M_1 \to M_1$ be the two projections. Then we have a canonical isomorphism of vector bundles

(23)
$$TM \simeq \pi_0^*(T[\alpha,\beta]) \oplus \pi_1^*(TM_1) \simeq \mathbb{R} \times \pi_1^*(TM_1),$$

where the direct sum \oplus is the direct sum of vector bundles on M and \times is the usual Cartesian product of sets. A section of $\pi_0^*(T[\alpha, \beta])$ will be called a *horizontal vector* field (on M) and a section of $\pi_1^*(TM_1)$ will be called a *vertical vector* field (on M). Accordingly, a vector field Z on M will decompose as

(24)
$$Z := (Z_0, Z_1) \in \pi_0^*(T[\alpha, \beta]) \oplus \pi_1^*(TM_1),$$

where Z_0 is a vector field on $M := [\alpha, \beta] \times M_1$ in the direction of $[\alpha, \beta]$ and Z_1 is a vector field on M in the direction of M_1 . Thus Z_0 is a horizontal vector field and Z_1 is a vertical vector field. The copy of \mathbb{R} on the right hand side is due to the fact that every horizontal vector field Z_0 is of the form $v_0\partial_t$, for some suitable function v_0 on M, so, at each point of M, it is determined by a single scalar (in \mathbb{R}). We shall often use in what follows the isomorphisms $\mathcal{A}(\mathcal{G}) \simeq \mathcal{A}(\mathcal{G})_0 \subset \mathcal{C}(\mathcal{G}^{(0)})$ and $\mathcal{L}(\mathcal{G}) \simeq \mathcal{L}(\mathcal{G})_0 \subset \mathcal{C}(\mathcal{G}^{(0)}, \mathbf{A}(\mathcal{G}))$ defined by restriction to units (as in Remark (2.17)).

Problem 3.1. To find a CF-groupoid $\mathcal{G}_{\psi,\phi}$ with units $\mathcal{G}_{\psi,\phi}^{(0)} = M := [\alpha,\beta] \times M_1$ whose associated Lie-Rinehart algebra of vector fields $\mathcal{L}(\mathcal{G}_{\psi,\phi})_0$ is given by restriction to units (see Remark (2.17)) by vector fields of the form

$$Z = \left(\phi Z_0, \psi Z_1\right),$$

with Z_0 horizontal and Z_1 vertical vector fields on M (see Equation (24) for notation; additional smoothness properties may be required of Z_0 and Z_1 , in particular, ϕ will be as in Equation (10) and ψ will be as in Equation (22)).

Of course, our Problem 3.1 is an "integration problem" for Lie-Rinehart algebras. More precisely, let $(\mathcal{A}(S), \mathcal{L}(S))$ be the Lie-Rinehart algebra generated by vector fields of the form $Z = (\phi Z_0, \psi Z_1)$, as above. Then we will construct a CF-groupoid $\mathcal{G}_{\psi,\phi}$ integrating $(\mathcal{A}(S), \mathcal{L}(S))$, that is, such that $(\mathcal{A}(S), \mathcal{L}(S)) = (\mathcal{A}(\mathcal{G}), \mathcal{L}(\mathcal{G}))$.

Our construction of the groupoid $\mathcal{G}_{\psi,\phi}$ depends on whether or not ψ vanishes at the end points (α and β). The result, however, is independent of this condition. More precisely, we will construct two CF-groupoids \mathcal{H}_{ψ} and $\mathcal{G}_{\psi,\phi}$ with units M satisfying, in turn, he following two results (recall the notation and isomorphisms of Remark (2.17)). The main properties of the groupoid \mathcal{H}_{ψ} are described by the following proposition.

Proposition 3.2. Let ψ be as in Equation (22). There exists an explicit CF-groupoid \mathcal{H}_{ψ} (given by Equation (26), if $\psi(\alpha) > 0$, and by Equation (33), if $\psi(\alpha) = 0$), which is minimal (and hence unique up to isomorphism) such that the Lie-Rinehart algebra of \mathcal{H}_{ψ} is given by

$$\mathcal{A}(\mathcal{H}_{\psi})_{0} = \{ u \mid \psi^{j} Y_{1} \dots Y_{j} u \in \mathcal{C}(M), \ j \in \mathbb{Z}_{+} \ and \ Y_{i} \in \mathcal{C}^{\infty}(M; \pi_{1}^{*}(TM_{1})) \}$$
$$\mathcal{L}(\mathcal{H}_{\psi})_{0} = \psi \mathcal{A}(\mathcal{H}_{\psi}) \mathcal{C}^{\infty}(M; \pi_{1}^{*}(TM_{1})) \subset \mathcal{C}(M; TM).$$

In turn, the main properties of the groupoid $\mathcal{G}_{\psi,\phi}$ are described by the following proposition. Before stating it, let us notice that the vector fields Y_i above are *vertical* vector fields on M. In particular, the CF-groupoid \mathcal{H}_{ψ} integrates the vertical vector fields on M that are multiples of ψ . In order to simplify the statement of the result satisfied by $\mathcal{G}_{\psi,\phi}$, it is useful to notice that, in the last proposition, we can replace $\mathcal{C}^{\infty}(M; \pi_1^*(TM_1))$ (the set of smooth, vertical vector fields on M) with $\mathcal{C}^{\infty}(M_1; TM_1) \subset \mathcal{C}^{\infty}(M; \pi_1^*(TM_1))$ (the smooth, vertical vector fields on $M = [\alpha, \beta] \times M_1$ that are independent of $t \in [\alpha, \beta]$).

Theorem 3.3. Let ϕ and ψ be as in Equations (10) and (22) and \mathcal{H}_{ϕ} be the CF-groupoid of Proposition 3.2. Then there exists an action of \mathbb{R} on \mathcal{H}_{ψ} whose infinitesimal generator

is $X = \phi \partial_t$ and the action (or semi-direct product) groupoid $\mathcal{G}_{\psi,\phi} := \mathcal{H}_{\psi} \rtimes \mathbb{R}$ is a CFgroupoid with Lie-Rinehart algebra given by

$$\mathcal{A}(\mathcal{G}_{\psi,\phi})_0 = \{ u \mid \psi^j X^k Y_1 \dots Y_j u \in \mathcal{C}(M), \ j,k \in \mathbb{Z}_+ \ and \ Y_i \in \mathcal{C}^\infty(M_1;TM_1) \}$$
$$\mathcal{L}(\mathcal{G}_{\psi,\phi})_0 = \phi \mathcal{A}(\mathcal{G}_{\psi,\phi})\partial_t + \psi \mathcal{A}(\mathcal{G}_{\psi,\phi})\mathcal{C}^\infty(M_1;TM_1) \subset \mathcal{C}(M;TM) .$$

If $\frac{\psi}{\psi_1}, \frac{\psi_1}{\psi} \in \mathcal{C}_{\phi}^{(\infty)}$, then we have canonical isomorphisms $\mathcal{H}_{\psi} \simeq \mathcal{H}_{\psi_1}$ and $\mathcal{G}_{\psi,\phi} \simeq \mathcal{G}_{\psi_1,\phi}$.

Let $\psi, \psi_1 : [\alpha, \beta] \to [0, \infty)$. In view of the above theorem, we shall write

(25)
$$\psi \sim_{\phi} \psi_1 \Leftrightarrow \frac{\psi}{\psi_1}, \frac{\psi_1}{\psi} \in \mathcal{C}_{\phi}^{(\infty)}$$

(More precisely, the quotients $\frac{\psi}{\psi_1}$ and $\frac{\psi_1}{\psi}$ extend by continuity to functions defined on $[\alpha, \beta]$ and that these extensions are in $C_{\phi}^{(\infty)}$.)

Remark 3.4. It follows that the Lie-Rinehart algebra $(\mathcal{A}(\mathcal{G}_{\psi,\phi}), \mathcal{L}(\mathcal{G}_{\psi,\phi}))$ associated to the CF-groupoid $\mathcal{G}_{\psi,\phi}$ is "generated" in an obvious sense by $X = \phi \partial_t$ and $\psi \mathcal{C}^{\infty}(M_1; TM_1)$.

The following two subsections are devoted to proving the above two results assuming $\psi(\beta) > 0$. The proof will be split in two according to the values of ψ at α . (The case $\psi(\beta) = 0$ is completely similar and can also be obtained from the case $\psi(\beta) > 0$ using a gluing procedure.)

3.2. The case $\psi(\alpha) > 0$. We thus consider first the case when our given continuous function $\psi : [\alpha, \beta] \to [0, \infty)$ is such that $\psi(\alpha)\psi(\beta) \neq 0$. Then $\psi > 0$ on $[\alpha, \beta]$. In this case, our construction will independent of ψ with these properties. Our first step is to construct a groupoid \mathcal{H}_{ψ} of Proposition 3.2. Explicitly,

(26)
$$\mathcal{H}_{\psi} := [\alpha, \beta] \times (M_1 \times M_1),$$

with structural maps:

(27)
$$\begin{aligned} d(s,x,y) &:= (s,y) \quad \text{and} \quad r(s,x,y) &:= (x,s), \quad \text{and} \\ (s,x,y)(s,y,z) &:= (s,x,z) \quad s \in [\alpha,\beta] \text{ and } x,y \in M_1 \end{aligned}$$

In the standard terminology (see, for instance, [11]), \mathcal{H}_{ψ} is a Lie groupoid, product of the space $[\alpha, \beta]$ (a groupoid that has only units) with the pair groupoid $M_1 \times M_1$. (The pair groupoid $M_1 \times M_1$ is obtained by removing the factor s in the above formulas, see Example 2.9(1)). The minimality of \mathcal{H}_{ψ} follows from the fact that the pair groupoid $M_1 \times M_1$ is the minimal groupoid with algebroid TM_1 . Recall that $M := [\alpha, \beta] \times M_1$ and a vector field on M is *vertical* if it is tangent to all the copies $\{c\} \times M_1$ of $M_1, c \in [\alpha, \beta]$. We see right away that \mathcal{H}_{ϕ} is independent on ψ .

We want to prove Proposition 3.2 in this case. Since ψ is non-vanishing and continuous the required identifications of Proposition 3.2 become

(28)
$$\mathcal{A}(\mathcal{H}_{\psi})_{0} = \{ u \mid \psi^{j} Y_{1} \dots Y_{j} u \in \mathcal{C}(M) \} = \mathcal{C}^{\infty,0}(M)$$

(for all $j \in \mathbb{Z}_+$ and $Y_i \in \mathcal{C}^{\infty}(M; \pi_1^*(TM_1))$) and

(29)
$$\mathcal{L}(\mathcal{H}_{\psi})_{0} = \psi \mathcal{A}(\mathcal{H}_{\psi}) \mathcal{C}^{\infty}(M; \pi_{1}^{*}(TM_{1})) = \psi \mathcal{C}^{\infty,0}(M; \pi_{1}^{*}(TM_{1})).$$

This amounts to the fact that $\mathcal{A}(\mathcal{H}_{\psi})_0$ consists of continuous functions on $[\alpha, \beta] \times M_1$ that are smooth in the direction of M_1 (the vertical direction) and $\mathcal{L}(\mathcal{H}_{\psi})_0$ consists of continuous vertical vector fields on M that are smooth in the direction of M_1 (again the vertical direction).

SCHRÖDINGER OPERATORS

Proof of Proposition 3.2 for $\psi(\alpha)\psi(\beta) \neq 0$. We notice that \mathcal{H}_{ψ} , $\mathcal{A}(\mathcal{H}_{\psi})_0$, $\mathcal{L}(\mathcal{H}_{\psi})_0$, and the other spaces of Equations (28) and (29) $(\mathcal{C}^{\infty,0}(M) \text{ and } \psi\mathcal{C}^{\infty,0}(M; \pi_1^*(TM_1)))$ are independent on $\psi > 0$ because $1/\psi \in \mathcal{C}_{\phi}^{(\infty)}$ (which is due, in turn, to the fact that $\psi \in \mathcal{C}_{\phi}^{(\infty)}$ and $\psi > 0$ on $[\alpha, \beta]$). It is enough then to assume that $\psi = 1$. By definition, $\mathcal{A}(\mathcal{H}_{\psi})_0$ consists of longitudinally smooth right-invariant functions. For this Lie groupoid we have d(s, x, y) = (s, y), where $s \in [\alpha, \beta]$ and $x, y \in M_1$. Hence, given a function f(s, x, y), it must be smooth in the *x*-variable and continuous in the other ones. Now if *f* is rightinvariant, then

$$f(s, x, y) = f((s, x, y)(s, y, z)) = f(s, x, z).$$

From here, we see that right-invariant functions are actually independent of the second variable, and thus we can identify them with functions f(s, x) on M, which are smooth along the variable $x \in M_1$. This is exactly the first set given in Proposition 3.2.

Let us now identify the second set in Proposition 3.2. First, we have that a vector field $X \in \mathcal{L}(\mathcal{H}_{\psi})_0$ must be a derivation along the *d*-fibers. In local coordinates, this means that it must be of the form:

$$X = \sum_{i=1}^{\dim M_1} X_i(s, x, y) \partial_{x^i} \,.$$

where X a priori is only smooth in the x variables. Now we need to check what conditions on $X_i(s, x, y)$ are imposed by the right invariance. For this, we need to compute the differential $(R_{(s,y,z)})_*$ of the right multiplication. Since, (s, x, y)(s, y, z) = (s, x, z), we have

$$(R_{(s,y,z)})_* = \begin{pmatrix} id & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 0 \end{pmatrix}.$$

Thus

$$(R_{(s,y,z)})_*X(s,x,y)=\sum_{i=1}^{\dim M_1}X_i(s,x,z)\partial_{x^i}.$$

If $(R_{(s,y,z)})_*X(s,x,y) = X(s,x,y)$, then we must have $X_i(s,x,y) = X_i(s,x,z)$, which again allows us to conclude that X_i is independent of the last variable, and hence that $X_i \in \mathcal{A}(\mathcal{H}_{\psi})_0$. This completes the proof.

Let σ be the action of \mathbb{R} on $[\alpha, \beta]$ associated to ϕ as in the previous section (See Proposition 2.28 and the discussion following it). We extend this action to an action of \mathbb{R} on $\mathcal{H}_{\psi} := [\alpha, \beta] \times M_1 \times M_1$ that is trivial on $M_1 \times M_1$. We then let

(30)
$$\mathcal{G}_{\psi,\phi} := \mathcal{H}_{\psi} \rtimes_{\tau} \mathbb{R} \simeq M_1 \times M_1 \times \mathcal{G}_{\phi} ,$$

(see Theorem 3.3). We see from the definitions that $\mathcal{G}_{\psi,\phi} = \mathcal{G}_{1,\phi}$ is independent of ψ , as long as ψ is smooth and it does not vanish at the end points.

Proof of Theorem 3.3 for $\psi(\alpha)\psi(\beta) \neq 0$. The groupoid $\mathcal{G}_{\psi,\phi} := \mathcal{G}_{\psi} \rtimes \mathbb{R}$ of Theorem 3.3 (see also Equation (30)) is the Cartesian product of the action groupoid \mathcal{G}_{ϕ} and the pair groupoid $M_1 \times M_1$. We shall identify it, as a set, with $M_1 \times M_1 \times [\alpha, \beta] \times \mathbb{R}$ via

(31)
$$\mathcal{G}_{\psi,\phi} := \mathcal{H}_{\psi} \rtimes_{\tau} \mathbb{R} \simeq M_1 \times M_1 \times \mathcal{G}_{\phi} \simeq M_1 \times M_1 \times [\alpha,\beta] \times \mathbb{R} \,.$$

that is, the product of the pair groupoid $M_1 \times M_1$ with the action groupoid \mathcal{G}_{ϕ} introduced in the previous section. See Equations (21), (21), and (27) for the products on these groupoids. Hence, we have

$$(x, y, s, t)(y, z, \sigma_{-\tau}(s), \tau) = (x, z, \sigma_{-\tau}(s), t + \tau).$$

In particular, if a function f is right-invariant, then

$$f(x, y, s, t) = f(x, z, \sigma_{-\tau}(s), t + \tau).$$

As a consequence if f(x, y, s, t) is right invariant, then,

- it is independent of y and,
- it is entirely determined by the restriction to $t + \tau = 0$.

Thus $f(x, y, s, t) = f_0(x, \sigma_t(s))$, for some function f_0 on the set of units $M_1 \times [\alpha, \beta] \simeq M$ (the diffeomorphism is given by switching $[\alpha, \beta]$ and M_1). Since $f \in \mathcal{A}(\mathcal{G}_{\psi,\phi})_0$ must be smooth along the *d*-fibres, it should be differentiable in the *x* and *t* variables. To establish the isomorphism from the statement, we notice that the reduced function $f_0(x, \sigma_t(s))$ depends only on r(x, y, s, t). Hence it can be seen as the pull-back r^*u of a function $u \in \mathcal{C}^{\infty}(M)$. Fix *t* and consider a new variable $t'(s) = \sigma_t(s)$ and extend it by continuity to $t'(-\infty) = \alpha$, $t'(-\infty) = \beta$. Then $f(x, \sigma_t(s)) = f(x, t')$ and the differentiability in *t* transfers to differentiability via the vector field $\phi(t')\partial_{t'}$. The differentiability properties now follow.

Next we prove the second isomorphism. The vector fields in $\mathcal{L}(\mathcal{G}_{\psi,\phi})_0$ must be tangent to the *d*-fibers and right invariant. Since $d(x, y, \tau, t) = (y, \tau)$, those vector fields must be derivations in *t* and *x*. Vector fields ∂_t and $V \in \mathcal{C}^{\infty}(M_1, TM_1) \subset \mathcal{C}^{\infty}(M, TM)$ can be seen tautologically as vector fields on the *d*-fibers. Note that all of them are already rightinvariant: ∂_t because the right multiplication is just a translation in the *t*-variable, and *V* by an argument similar to the proof of Proposition 3.2 for our case. Since those vector fields generate any other *d*-tangent vector field, all the invariant vector fields must be linear combinations of those ones with invariant functions as coefficients. So it only remains to take $t' = \sigma_t(s)$ as a new variable and we get exactly $\mathcal{A}(\mathcal{G}_{\psi,\phi})_0$. In the case $\psi > 0$, all the groupoids and associated spaces are independent on ψ . This gives

(32)
$$\mathcal{G}_{\psi_1,\phi} = \mathcal{G}_{1,\phi} = \mathcal{G}_{\psi,\phi}, \qquad \psi, \psi_1 > 0,$$

that is, the isomorphism of the last statement in Theorem 3.3.

3.3. The case $\psi(\alpha) = 0$. Recall that $\psi(\beta) > 0$. As before, we begin by defining the CF-groupoid \mathcal{H}_{ψ} integrating $\psi \mathcal{C}^{\infty}(M; \pi_1^*(TM_1))$. Let

(33)
$$\mathcal{H}_{\psi} := TM_1 \times \{\alpha\} \cup M_1 \times M_1 \times (\alpha, \beta)$$

as a set. The groupoid structure is the following. The domain, range, and multiplication maps on $M_1 \times M_1 \times (\alpha, \beta]$ are induced from the inclusion

$$M_1 \times M_1 \times (\alpha, \beta] \times \mathbb{R} \subset \mathcal{G}_{1,\phi} := M_1 \times M_1 \times [\alpha, \beta] \times \mathbb{R}$$

in the groupoid $\mathcal{G}_{1,\phi}$ of the previous subsection. Let $\pi_{TM_1} : TM_1 \to M_1$ be the canonical projection. It will be convenient to write $M = [\alpha, \beta] \times M_1$ (we have switched the two factors). On the other hand, if $s = \alpha$, these maps are given by

(34)
$$\begin{cases} d(v,\alpha) = r(v,\alpha) := (\pi_{TM_1}(v),\alpha) \in M_1 \times \{\alpha\} \subset M & \text{and} \\ (v,\alpha)(w,\alpha) := (v+w,\alpha), \end{cases}$$

where $v, w \in TM_1$ are vectors in the same fiber over M_1 , that is, $\pi_{TM_1}(v) = \pi_{TM_1}(w)$. Thus, \mathcal{H}_{ψ} is a "family" of groupoids indexed by $s \in [\alpha, \beta]$, such that, to $s \in (\alpha, \beta]$ we associate the pair groupoid $M_1 \times M_1$ and to $s = \alpha$ we associate the tangent bundle TM_1 with the associated groupoid structure of a family of Lie groups.

To define the topology and CF-groupoid structure on \mathcal{H}_{ψ} , we consider a variant of the "deformation to the normal cone" considered by Connes, Debord, Skandalis, and others [14, 17]. Thus, we first consider on $TM_1 \times \{\alpha\}$ and on $M_1 \times M_1 \times \{\alpha, \beta\}$ the natural smooth

(manifold) structures and the Lie groupoid structures. To glue these smooth structures, we use the exponential coordinates of [37] to define the smooth structure in the neighborhood of a point $(v, \alpha) \in TM_1 \times \{\alpha\}$. This is done as follows. Then, let us fix some arbitrary connection on M_1 (that is, on its tangent space TM_1) and let $\exp : U \subset TM_1 \to M_1$ be its associated exponential map [41] (here U is its maximal domain of definition, a subset of TM_1). Let $\pi_{TM_1} : TM_1 \to M_1$ be the projection map. It is known that

(35)
$$\Psi := (\pi_{TM_1}, \exp) : U \to M_1 \times M_1$$

is a diffeomorphism from an open neighborhood of the zero section of TM_1 onto an open neighborhood of the diagonal $\delta_{M_1} \subset M_1 \times M_1$. Let $v \in TM_1$. Let us define now a neighborhood of (v, α) in \mathcal{H}_{ψ} . Let first $1 \geq s_0 > 0$ be such that $sv \in U$ for all $|s| \leq s_0$. Then, let us choose a relatively compact neighborhood W of sv in TM_1 such that $[-1, 1]\overline{W} \subset U$. (So the exponential $\exp(sw)$ is defined for all $w \in W$ and $|s| \leq 1$.) Let $\epsilon > 0$ be such that $\psi(s) < s_0$ if $\alpha \leq s < \alpha + \epsilon < \beta$. We then let $\Phi : W \times [0, \epsilon) \to \mathcal{H}_{\psi,\phi}$,

(36)
$$\Phi(w,s) := \begin{cases} (w,\alpha) & \text{if } s = 0\\ (\Psi(\psi(s)w), \alpha + s) \in M_1 \times M_1 \times (\alpha,\beta) & \text{if } s > 0. \end{cases}$$

(We recall from Equation (35) that $(\Psi(\psi(s)w), s') = (\pi_{TM_1}(w), \exp(\psi(s)w), s')$.) We stress the appearance of the factor $\psi(s)$ in these formulas. Except this "rescaling term", our formulas are as in the aforementioned papers. Finally, the desired neighborhood of v consists of

$$(37) V := \Phi(W \times [0, \epsilon))$$

with the induced topological and continuous family of manifolds structure from Φ . (The continuous family of manifolds structure on V comes from the map d.) We are ready now to prove Proposition 3.2 for $\psi(\alpha) = 0$ and $\psi(\beta) > 0$.

Proof of Proposition 3.2 for $\psi(\alpha)\psi(\beta) = 0$. Similar to the proof of this Proposition in the case $\psi(\alpha) > 0$, the invariant functions for $s \in (\alpha, \beta)$ satisfy

$$u(x, y, s) = u_0(x, s)$$

and must be smooth in x for all $s \in (\alpha, \beta)$. The difference now is in the behavior as s approaches α or β . We shall concentrate on the case s close to α , the case s close to β being completely similar. So let us change the coordinates using the map Φ . In this coordinates the domain and the range maps are given by

$$d(w,t) = \pi_{TM_1}(w)$$

$$r(w,t) = \exp(\psi(t)w),$$

where $(w, t) \in W \times [0, \epsilon) \subset TM_1 \times [0, \epsilon)$ (see the definition of the map Φ , Equation (36)) So differentiating along *d*-fibres in these coordinates is differentiating along vertical fibres of $w \in TM_1$. Since $x = \exp(\psi(t)w)$ we have that any derivation in the *w* variables will be a derivation in *x* pushforwarded by the differential of *r*, that is, by

(38)
$$d(\exp(\psi(t)\cdot)) = \psi(t)d\exp .$$

Note that d here indicates the differential. Since the exponential map is a diffeomorphism for sufficiently small t, we have that the derivative along the d-fiber exists if, and only if, $\psi Y u$ is continuous for any derivation $Y \in C^{\infty}(M_1, TM_1)$. The claimed description of $\mathcal{A}(\mathcal{H}_{\psi})_0$ is then obtained by noticing that ψ and Y commute (i.e. $Y(\psi) = 0$).

Now we characterize $\mathcal{L}(\mathcal{H}_{\psi})_0$. Again as in the proof of this Proposition in the case $\psi(\alpha) > 0$, the right invariant vector fields for $t > \alpha$ agree with continuous families of

derivations in the x variables parametrized by the t variable. The only thing to understand is what happens as we approach $t = \alpha$. For this we change to the w-variables using the diffeomorphism Φ of Equation (36). If $X_t \in \mathcal{C}^{\infty}(M_1, TM_1)$ is a family of such derivations, then in the w variables we get via (38):

$$(\exp(\psi(t))\cdot)_*^{-1}X_t = \frac{1}{\psi(t)}(\exp)_*^{-1}X_t.$$

Hence, we must have that $X_t/\psi(t)$ is continuous in t. From here the characterization follows. \square

We want to define an action of \mathbb{R} on \mathcal{H}_{ψ} integrating $X = \phi \partial_t$ to define $\mathcal{G}_{\psi,\phi} := \mathcal{H}_{\psi} \rtimes \mathbb{R}$ Theorem 3.3. First, the automorphisms σ_s , $s \in \mathbb{R}$, extend (as for the case $\psi > 0$ in the previous subsection) to $M_1 \times M_1 \times (\alpha, \beta]$. Recall the spaces $\mathcal{C}^{(\infty)}_{\phi} \subset \mathcal{C}([\alpha, \beta])$ introduced in Equation (10).

Proposition 3.5. We continue to assume that $\psi(\alpha) = 0$ (as throughout this subsection). Then the assumption $C_{\psi,\phi} := \frac{X(\psi)}{\psi} := \frac{\phi\psi'}{\psi} \in C_{\phi}^{(\infty)}$, of Equation (22) (see also Equation (10)) allows us to conclude that σ_s extend to CF-groupoid automorphism of \mathcal{H}_{ψ} , for all $s \in \mathbb{R}$. This extension to the end point α is such that

$$\sigma_s(v,\alpha) = \left(e^{-C_{\psi,\phi}(\alpha)s}v,t\right).$$

Proof. Let $s \in \mathbb{R}$. Let us check the continuity and longitudinal smoothness of σ_s near α (near the other points $(\alpha, \beta]$, it is obvious; it follows, for instance from case already proved of Theorem 3.3). Let $\lambda := C_{\psi,\phi}(\alpha)$. Let Φ and Ψ be as in Equations (35) and (36). We have

(39)
$$\sigma_s(\Phi(w,t)) = \begin{cases} (e^{-\lambda s}w,\alpha) & \text{if } t = \alpha\\ (\Psi(\psi(t)w),\sigma_s(t)) \in M_1 \times M_1 \times (\alpha,\beta) & \text{if } t \in (\alpha,\alpha+\epsilon) \,. \end{cases}$$

Hence, the map σ_s maps the fibers of $d: \mathcal{H}_{\psi} \to M$ to themselves smoothly. In other words, σ_s is longitudinally smooth. To complete the proof, we need to show that it is continuous. We have

$$\Phi(\psi(t)w,\sigma_s(t)) = \left(\pi_{TM_1}(w), \exp\left(\frac{\psi(t)}{\psi(\sigma_s(t))}\psi(\sigma_s(t))w\right), \sigma_s(t)\right)$$

So it only remains to show that $\lim_{t\to\alpha} \frac{\psi(t)}{\psi(\sigma_s(t))} = e^{-\lambda s}$. Indeed, we can compute:

$$\ln \frac{\psi(\sigma_s(t))}{\psi(t)} = \int_0^s \partial_\tau \ln \psi(\sigma_\tau(t)) d\tau = \int_0^s \frac{\partial_\tau \psi(\sigma_\tau(t))}{\psi(\sigma_\tau(t))} d\tau$$
$$= \int_0^s \frac{(X\psi)(\sigma_\tau(t))}{\psi(\sigma_\tau(t))} d\tau = \int_0^s C_{\psi,\phi}(\sigma_\tau(t)) d\tau \to C_{\psi,\phi}(\alpha) s = \lambda s$$
$$\text{sume } \sigma_\tau(t) \to \alpha \text{ as } t \to \alpha \text{ uniformly in } \tau \in [0, s].$$

because $\sigma_{\tau}(t) \to \alpha$ as $t \to \alpha$ uniformly in $\tau \in [0, s]$.

Proof. Proof of Theorem 3.3 for $\psi(\alpha) = 0$ and $\psi(\beta) > 0$. The proof is very similar to that for the case $\psi(\alpha) > 0$. (All the difficulties were transferred to the definition of \mathcal{H}_{ψ} .) The action of \mathbb{R} on \mathcal{H}_{ψ} , for $t \in (\alpha, \beta]$, agrees exactly with the action of \mathbb{R} on the groupoid (31). From the proof of the case $\psi(\alpha)$ positive of Proposition 3.2 it follows, that the invariant functions for $t > \alpha$ can be identified with functions of the form $f(x, y, t, s) = f(x, \sigma_s(t))$ (notice that t and s have been switched compared to the proof of Proposition 3.2). As in the proof of Theorem 3.3 for $\psi(\alpha) = 0$ by compactifying the \mathbb{R} -action and introducing a new variable $t' = \sigma_s(t)$, the differentiability in s, implies that $\phi_{t'} \partial_{t'} f$ must be continuous. On the other hand, the differentiability in x is exactly as in Proposition 3.2. The same holds for higher derivatives and hence the characterization follows.

The characterization of $\mathcal{V}_{\psi,\phi}(\mathcal{G}_{\psi,\phi})_0$ is again very similar to the proofs of the above propositions. We have that elements of this space must be differentiations in (x, s). For derivations in x we repeat word-by-word the argument in the proof of Proposition 3.2 and find that $\psi \mathcal{C}^{\infty}(M_1, TM_1)$ belong to $\mathcal{V}_{\psi,\phi}(\mathcal{G}_{\psi,\phi})_0$. Next we consider a derivation in s. It is a differentiation of a smooth family of automorphisms of \mathcal{H}_{ψ} by Proposition. 3.5. Hence it is enough to describe this differentiation for $t \in (\alpha, \beta]$ and then extend it to whole of \mathcal{H}_{ψ} . The argument in the proof of Theorem 3.3 when $\psi(\alpha) \neq 0$ says that for $t \in (\alpha, \beta]$ the derivative ∂_s is pushforwarded to $\phi(t')\partial_{t'}$ after a change of variables. Any other invariant vector field must be a linear combination of mentioned ones with invariant functions as coefficients.

Finally, again, as a set $\mathcal{G}_{\psi,\phi}$ is independent of ψ and the identity map $\mathcal{G}_{\psi,\phi} \to \mathcal{G}_{\psi_1,\phi}$ is an isomorphism of (plain) groupoids. If $\psi \sim_{\phi} \psi_1$ (see Equation (25) for the notation), then the identity map $\mathcal{G}_{\psi,\phi} \to \mathcal{G}_{\psi_1,\phi}$ is also a $\mathcal{C}^{\infty,0}$ -map. This completes the proof. \Box

Remark 3.6. Theorem 2.32 can be cast in a more general framework as follows. Let \mathcal{H} be a CF-groupoid and let G be a Lie group acting by $\mathcal{C}^{\infty,0}$ -homeomorphisms on \mathcal{H} (thus they are continuous and smooth on each of the fibers of the maps d or r). Then $\mathcal{G} := \mathcal{H} \rtimes G$ is a CF groupoid (see also Example 2.9 (4) and (5)). We have that $\mathcal{A}(\mathcal{G})_0$ identifies with the smooth vectors in $\mathcal{A}(\mathcal{H})$ and $\mathcal{L}(\mathcal{G})_0$ identifies with the direct sum of the set of smooth vectors in $\mathcal{L}(\mathcal{H})_0$ with $\mathcal{A}(\mathcal{G})_0 Lie(G)$.

The case of a general interval $[\alpha, \beta] \subset \mathbb{R}$ is dealt with by using a homeomorphism of $[\alpha, \beta]$ to a bounded interval, provided that this homeomorphism is a diffeomorphism in the interior.

As already mentioned, the remaining case $\psi(\beta) = 0$ can be treated similarly. Better, yet, it can be treated by localization and by gluing groupoids, as in [10, 12]. We recall now the gluing construction, for the benefit of the reader.

3.4. Localization and gluing of CF-groupoids. In this section, the symbol \mathcal{G} (possibly decorated with various indices) will denote a CF-groupoid. The set of units of \mathcal{G}_i will be denoted $\mathcal{G}_i^{(0)}$, unless otherwise stated. The domain and range maps will be denoted by d and, respectively, r, as usual. Let \mathcal{G} be a CF-groupoid with units $\mathcal{G}^{(0)}$ and $U \subset \mathcal{G}^{(0)}$ be an open subset. Let

$$\mathcal{G}_U^U := d^{-1}(U) \cap r^{-1}(U),$$

Then \mathcal{G}_U^U is a CF-groupoid with units U. We can glue CF-groupoids in a similar way as it was done for Lie groupoids in [10, 12].

Theorem 3.7. Let M be a locally compact space and \mathcal{G}_i , i = 1, ..., N, be CF-groupoids with units open subsets $\mathcal{G}_i^{(0)} \subset M$ such that $\bigcup_{i=1}^N \mathcal{G}_i^{(0)} = M$. We assume that these groupoids satisfy the gluing conditions:

(1) (Isomorphism on the common part) There are CF-groupoid isomorphisms

$$f_{ji}: (\mathcal{G}_j)_{\mathcal{G}_i^{(0)} \cap \mathcal{G}_j^{(0)}}^{\mathcal{G}_i^{(0)} \cap \mathcal{G}_j^{(0)}} \to (\mathcal{G}_j)_{\mathcal{G}_i^{(0)} \cap \mathcal{G}_j^{(0)}}^{\mathcal{G}_i^{(0)} \cap \mathcal{G}_j^{(0)}}$$

such that $f_{ij} = f_{ji}^{-1}$ and $f_{ij}f_{jk} = f_{ik}$ on the common domains, for all $i, j, k \in \{1, 2, \dots, N\}$,.

(2) (The orbits are included completely in at least one groupoid) For each $x \in M$, there exist $i_x \in \{1, 2, \dots, N\}$, such that

$$\mathcal{G}_j \cdot x := \{ r(g) \, | \, d(g) = x \} \subset \mathcal{G}_{i_x}^{(0)}$$

for all $j \in \{1, 2, \cdots, N\}$.

Then their union $\mathcal{G} := \bigsqcup_{i=1}^{N} \mathcal{G}_i / \sim$, where \sim is the equivalent relation generated by $g \sim f_{ji}(g)$, for all $i, j \in \{1, 2, \dots, N\}$ and $g \in \mathcal{G}_i$, is a CF-groupoid. The Lie-Rinehart algebra of \mathcal{G} can be identified as $\mathcal{A}(\mathcal{G})_0 = \{u \in \mathcal{C}(M) \mid u|_{\mathcal{G}_i^{(0)}} \in \mathcal{A}(\mathcal{G}_i)_0\}$, and $\mathcal{L}(\mathcal{G})_0 = \{u \in \mathcal{C}(M, \mathbf{A}(\mathcal{G})) \mid u|_{\mathcal{G}_i^{(0)}} \in \mathcal{L}(\mathcal{G}_i)_0\}$.

The proof goes along the same lines as the one in [10]. Details will be included in [6]. To use the gluing theorem, we observe that the reduction of the groupoid $\mathcal{G}_{\psi,\phi}$ to any relatively compact subset of (α, β) is independent of ψ and ϕ . More precisely, if $I := [\alpha', \beta'] \subset (\alpha, \beta)$, then

$$(\mathcal{G}_{\psi,\phi})_I^I \simeq (I \times M_1)^2$$

(the pair groupoid, see 2.9(1)). Note that this relies on the minimality of \mathcal{H}_{ψ} .

3.5. The groupoid associated to Schrödinger operators. Let us consider now $M = S^{n-1} \times [0, \infty]$ (thus $[\alpha, \beta] = [0, \infty]$) and four parameters a, b, a', b', with $a, a' \ge 1$ and $b, b' \ge 0$. We then choose our functions ϕ and ψ to satisfy

(40)
$$\begin{cases} \phi(t) = t^{a} & \text{if } t \in [0, \epsilon/2] \\ \phi(t) = t^{-a'} & \text{if } t \in [2\epsilon^{-1}, \infty] \\ \psi(t) = t^{b} & \text{if } t \in [0, \epsilon/2] \\ \psi(t) = t^{-b'} & \text{if } t \in [2\epsilon^{-1}, \infty] , \end{cases}$$

for some $0 < \epsilon < 1$. (This is, of course, in addition to the usual assumptions: ϕ is strictly positive and continuous on (α, β) and ψ satisfies the conditions (22) on $[\alpha, \beta]$.) Moreover, for simplicity, we shall assume that ϕ is smooth on (α, β) . Up to an isomorphism, the resulting groupoid $\mathcal{G}_{\psi,\phi}$ is independent of ϵ and of the choices of ϕ and ψ satisfying Equation (40), by the last part of Theorem 3.3. Therefore, it makes sense to set

(41)
$$\mathcal{S} = \mathcal{S}_{a,a',b,b'} := \mathcal{G}_{\psi,\phi}, \quad \phi, \psi \text{ satisfying Equation (40)}.$$

We now look at some specific properties of the groupoid $S = S_{a,a',b,b'}$. Recall that, for a CF-groupoid G, the algebra of differential operators Diff(G) was defined as the image of the enveloping algebra of $\text{Diff}(\mathcal{A}(G)_0, \mathcal{L}(G)_0)$ in $\text{End}(\mathcal{A}(G)_0)$, see Subsection 2.4. Let

(42)
$$\begin{cases} \rho_0, \rho_\infty : S^{n-1} \times [0, \infty] \to [0, 1] \\ \rho_0(x, t) = t \text{ and } \rho_\infty(x, t) = 1 & \text{ for } t < \epsilon \\ \rho_\infty(x, t) = 1/t \text{ and } \rho_0(x, t) = 1 & \text{ for } t > 1/\epsilon \end{cases}$$

be smooth, depend only on the second variable, and be such that ρ_0 vanishes only when t = 0 and ρ_{∞} vanishes only when $t = \infty$. Thus these two functions are the defining functions of the two components of the boundary of $S^{n-1} \times [0, \infty]$. In view of the notation (25), we can thus take

(43)
$$\phi = \rho_0^a \rho_\infty^{a'} \quad \text{and} \quad \psi = \rho_0^b \rho_\infty^{b'}.$$

(The function ϕ used in the introduction is the same as the function ϕ used here.)

Proposition 3.8. We shall denote generically by $D_j = Y_1 Y_2 \dots Y_j$ a differential monomial of order j in smooth, vertical vector fields on M (so Y_i are tangent to all $S^{n-1} \times \{t\}$) and $S = S_{a,a',b,b'}$, with $\phi = \rho_0^a \rho_\infty^{a'}$ and $\psi = \rho_0^b \rho_\infty^{b'}$. Then

(44)
$$u \in \mathcal{A}(\mathcal{S})_0 \quad \Leftrightarrow \quad i, j \in \mathbb{Z}_+, \quad \phi^i \psi^j \, \partial_t^i D_j u \in \mathcal{C}(M) \,,$$

Similarly, a differential operator on the interior of M is in Diff(S) if, and only if, it is a linear combinations of terms of the form $c_{i,j}\phi^i\psi^j\partial_t^i D_j$, where $c_{i,j} \in \mathcal{A}(S)_0$, $i, j \in \mathbb{Z}_+$.

Proof. This follows from Theorem 3.3, more precisely, from the characterization of the Lie-Rinehart algebras in that theorem, and from Corollary 2.26. (See also Remark 3.4.)

Definition 3.9. Let $S = S_{a,a',b,b'}$ and let D be a differential operator on $M := S^{n-1} \times [0,\infty]$, as above. We shall say that D is in the $c_{a,b}$ -calculus (respectively, in the $c_{a',b'}$ -calculus) if it coincides with a differential operator in Diff(S) in a neighborhood of $M_1 \times \{0\}$ (respectively, in a neighborhood of $M_1 \times \{\infty\}$).

Related calculi to the $c_{a,b}$ calculi were used in see, for instance in [24, 50, 51], also in connection to Schrödinger operators. See also [16, 25, 32, 34].

Let d and r be the structural maps of the groupoid $S = S_{a,b,a',b'}$. We shall need the following property of this groupoid.

Proposition 3.10. Let ρ_0 and ρ_{∞} be the functions introduced in Equation (42), then

(45)
$$\zeta_0 := \frac{\rho_0 \circ d}{\rho_0 \circ r} \quad and \quad \zeta_\infty := \frac{\rho_\infty \circ d}{\rho_\infty \circ r}$$

extend to nowhere vanishing $C^{\infty,0}$ functions on $S_{a,b,a',b'}$.

Proof. The defining formula $\mathcal{G}_{\psi,\phi} := \mathcal{H}_{\psi} \rtimes_{\sigma} \mathbb{R}$ (Theorem 3.3), we see that the quotients ζ_0 and ζ_{∞} depend only on $[0,\infty] \rtimes \mathbb{R}$, via the projection $\mathcal{H}_{\psi} \to [0,\infty]$. The required smoothness then follows from the explicit formulas for $\sigma_s(x)/x$ of Examples 2.29 (for a = 1 or a' = 1) or 2.30 (for a > 1 or a' > 1).

Consider the class $\Psi(\mathcal{G})$ of pseudodifferential operators on \mathcal{G} , Definition 2.20. The previous lemma then yields the following result.

Proposition 3.11. Let ρ_0 and ρ_∞ be the functions introduced Equation (42), $S = S_{a,b,a',b'}$, $P \in \Psi^m(S)$. Let also ζ_0 and ζ_∞ be as in Proposition 3.10 and $t, t' \in \mathbb{R}$. Then the operator

$$\tilde{P} := \rho_{\infty}^{-t'} \rho_0^{-t} P \rho_0^t \rho_{\infty}^{t'}$$

has distribution kernel $k_{\tilde{P}} = \zeta_0^t \zeta_{\infty}^{t'} k_P$, and hence it belongs to $\Psi^m(\mathcal{S})$ as well.

Proof. The proof is a direct consequence of definitions. Indeed, the distribution kernel of fPg is $k_{fPg}(\gamma) = f(r(\gamma))k_P(\gamma)g(d(\gamma))$. This gives the claimed form of the convolution kernel of \tilde{P} . Since ζ_0 and ζ_∞ are $\mathcal{C}^{\infty,0}$ and > 0, by Proposition 3.10, $\zeta_0^t \zeta_\infty^{t'}$ is $\mathcal{C}^{\infty,0}$ on \mathcal{S} . This gives the result by the identification of the convolution kernels k_P in [27].

4. AN APPLICATION TO SCHRÖDINGER OPERATORS

We now apply the results of the previous sections to study Schrödinger operators with power-law singularities at 0 and at ∞ introduced in Equation (1). We continue to consider the functions ρ_0 and ρ_∞ of Equation (42) (introduced right before Proposition 3.10). Recall that the operator

$$H := -\Delta + V$$

of Equation (1) is such that the potential V is of the form $V = \rho_0^{-2\gamma} \rho_{\infty}^{2\gamma'} V_0$, with V_0 smooth on $S^{n-1} \times [0, \infty]$ in spherical coordinates (ρ, x') , $\rho \in [0, \infty]$, $x' \in S^{n-1}$. (We can relax the smoothness assumptions on V_0 , but the most general assumptions on V_0 are much more difficult to state.)

We begin by studying the operator $-\Delta + V$ in a neighbourhood of the origin, where $\rho_0 = \rho$. We may also assume $\rho_{\infty} = 1$ there. Let us assume first that $\gamma \leq 1$. Proposition 3.8 and the last part of Corollary 2.27 show that the operator

(46)
$$\rho^2(-\Delta + V) = (\rho\partial_{\rho})^2 + (n-2)\rho\partial_{\rho} + \Delta_{S^{n-1}} + \rho^{2-2\gamma}V_0$$

is in the $c_{1,0}$ -calculus of the previous section (see Definition 3.9); this is the case even if 2γ is not an integer due to Corollary 2.27. (That is, our coefficients *do not* have to be smooth up to the boundary. By contrast, $\rho^2(-\Delta + V)$ is in the *b*-calculus precisely if $2\gamma \in \mathbb{Z}$ and $2\gamma \leq 2$.) On the other hand, if $\gamma \geq 1$ (but keeping the assumption that V_0 is smooth in polar coordinates), we have that

(47)
$$\rho^{2\gamma}(-\Delta+V) = (\rho^{\gamma}\partial_{\rho})^2 + (n-1-\gamma)\rho^{\gamma-1}\rho^{\gamma}\partial_{\rho} + \rho^{2\gamma-2}\Delta_{S^{n-1}} + V_0$$

is in the $c_{\gamma,\gamma-1}$ calculus, since its coefficients, $\rho^{\gamma-1}$ and $\rho^{2\gamma-2}$, are in the algebra $\mathcal{C}_{\rho^{\gamma}}^{(\infty)} = \mathcal{C}_{\rho^{\gamma}}^{(\infty)}([0,\infty))$ of Equation (10) by Corollary 2.27.

We can carry a similar analysis at $+\infty$. Our assumption is that our potential V is of the form $V(\rho x') = \rho^{2\gamma'} V'_0(\rho^{-1}, x')$, with V'_0 smooth on $[0, \infty) \times S^{n-1}$. Let us set $r := \rho^{-1}$ and let us write the Schrödinger operator in the coordinates $(r, x') \in [0, \infty) \times S^{n-1}$ (that is, "near ∞ "). We have $\rho \partial_{\rho} = -r \partial_r$. This gives us

(48)
$$-\Delta + V = (r^2 \partial_r)^2 - (n-1)r(r^2 \partial_r) + r^2 \Delta_{S^{n-1}} + r^{-2\gamma'} V_0'.$$

Similar to the analysis at $\rho = 0$ we see that if $\gamma' \leq 0$, then at the infinity this operator is in the $c_{2,1}$ -calculus of Definition 3.9. If $\gamma' \geq 0$, then we factor out $r^{-2\gamma'}$. It is a straightforward computation to check that:

$$(r^{2+\gamma'}\partial_r)^2 = r^{2\gamma'}(r^2\partial_r)^2 + \gamma' r^{2\gamma+3}\partial_r \,.$$

Then

(49)
$$r^{2\gamma'}(-\Delta+V) = (r^{2+\gamma'}\partial_r)^2 - (n-1+\gamma')r^{1+\gamma'}(r^{2+\gamma'}\partial_r) + r^{2\gamma'+2}\Delta_{S^{n-1}} + V_0'$$

is in the $c_{2+\gamma',\gamma'+1}$ -calculus. We have thus proved the following proposition.

Proposition 4.1. Let ρ_0 and ρ_∞ be the functions introduced right before Proposition 3.10, in particular, $\rho_0(t) = t$ for t small and $\rho_\infty(t) = t^{-1}$ for t large, see Equation (42). Let $\gamma, \gamma' \in \mathbb{R}$ and V be such that $\rho_0^{-2\gamma} \rho_\infty^{-2\gamma'} V$ be smooth on $[0, \infty] \times S^{n-1}$. Let $\tilde{\gamma} := \max\{\gamma, 1\}$ and, similarly, $\tilde{\gamma}' := \max\{\gamma', 0\}$. Then the operator

$$P := \rho_0^{\tilde{\gamma}} \rho_{\infty}^{\tilde{\gamma}'}(-\Delta + V) \in \text{Diff}(\mathcal{S}), \quad \mathcal{S} := \mathcal{S}_{\tilde{\gamma}, \tilde{\gamma} - 1, \tilde{\gamma}' + 2, \tilde{\gamma}' + 1}$$

We let $S := S_{\tilde{\gamma},\tilde{\gamma}-1,\tilde{\gamma}'+2,\tilde{\gamma}'+1}$ from now on, with the parameters $\tilde{\gamma}$ and $\tilde{\gamma}'$ as in the statement of Proposition 4.1. We have $\text{Diff}(S_{a,b,a',b'}) \subset \Psi(S_{a,b,a',b'})$ (this is valid for any CF-groupoid) and, because the action of $\Psi(S_{a,b,a',b'})$ on $C_c^{\infty}(S^{n-1} \times [0,\infty])$ is injective, we see that the operator $P := \rho_0^{\tilde{\gamma}} \rho_{\infty}^{\tilde{\gamma}'}(-\Delta + V)$ corresponds to a unique operator in $\Psi^2(S_{\tilde{\gamma},\tilde{\gamma}-1,\tilde{\gamma}'+2,\tilde{\gamma}'+1})$, which is elliptic

by Equations (46-49). Hence it has a parametrix $Q_0 \in \Psi^{-2}(S_{\tilde{\gamma},\tilde{\gamma}-1,\tilde{\gamma}'+2,\tilde{\gamma}'+1})$. We obtain the following result on the structure of the parametrices of $-\Delta + V$.

Theorem 4.2. There exists $Q \in \Psi^{-2}(S_{\tilde{\gamma},\tilde{\gamma}-1,\tilde{\gamma}'+2,\tilde{\gamma}'+1})$ such that

$$(-\Delta+V)\left(Q\rho_0^{\tilde{\gamma}}\rho_{\infty}^{\tilde{\gamma}'}\right), \quad \left(Q\rho_0^{\tilde{\gamma}}\rho_{\infty}^{\tilde{\gamma}'}\right)(-\Delta+V) \in \Psi^{-\infty}(\mathcal{S}_{\tilde{\gamma},\tilde{\gamma}-1,\tilde{\gamma}'+2,\tilde{\gamma}'+1})$$

Let $\phi := \rho_0^{\tilde{\gamma}} \rho_{\infty}^{\tilde{\gamma}'}$ (it is the same ϕ as the one appearing in the Introduction.)

Proof. Let $Q := Q_0$ be the differential operators (parametrix) considered in the paragraph before the statement of the Theorem.

The definition of Q_0 is such that

$$\phi(-\Delta+V)Q$$
, $Q\phi(-\Delta+V) \in \Psi^{-\infty}(\mathcal{S}_{\tilde{\gamma},\tilde{\gamma}-1,\tilde{\gamma}'+2,\tilde{\gamma}'+1})$.

Then

$$(-\Delta+V)(Q\phi) = \phi^{-1} \left[\phi(-\Delta+V)Q \right] \phi \in \Psi^{-\infty}(\mathcal{S}_{\tilde{\gamma},\tilde{\gamma}-1,\tilde{\gamma}'+2,\tilde{\gamma}'+1}),$$

by Proposition 3.11.

Let us now sketch how one can treat the resolvents in our setting. Let $S := S_{\tilde{\gamma}, \tilde{\gamma}-1, \tilde{\gamma}'+2, \tilde{\gamma}'+1}$, $\phi := \rho_0^{\tilde{\gamma}} \rho_{\infty}^{\tilde{\gamma}'}$, as before, and let $\overline{\Psi}^{-\infty}(S)$, be the closure of $\Psi^{-\infty}(S)$ in the norms

(50)
$$T \mapsto \| (\phi \Delta)^{i} T(\phi \Delta)^{j} \|_{\mathcal{L}(L^{2}(\mathbb{R}^{n}))}, \qquad i, j \in \mathbb{Z}_{+},$$

where $\mathcal{L}(L^2(\mathbb{R}^n))$ is the space of bounded operators on $L^2(\mathbb{R}^n)$ with the usual norm. Let $\overline{\Psi}^m(\mathcal{S}) := \Psi^m(\mathcal{S}) + \overline{\Psi}^{-\infty}(\mathcal{S})$. Then

Theorem 4.3. Let $z \in \mathbb{C}$ and let us assume that $\phi(-\Delta + V) - z$ is invertible on $L^2(\mathbb{R}^n)$. Then $[\phi(-\Delta + V) - z]^{-1} \in \overline{\Psi}^{-2}(S)$.

Proof. This follows from Proposition 4.1 and the results of [28, 49] (see also [46]). Indeed, $\phi(-\Delta + V) - z \in \text{Diff}(S) \subset \Psi(S) \subset \overline{\Psi}(S)$. The latter is stable under inverses, by the results of [28, 49], hence $[\phi(-\Delta + V) - z]^{-1} \in \overline{\Psi}^{-2}(S)$, as claimed.

Note that, while our statements about the structure of $-\Delta + V$ require only Lie-Rinehart algebras, the statement about its parametrices does require a CF-groupoid integrating the relevant Lie-Rinehart algebras.

REFERENCES

- AMMANN, B., LAUTER, R., AND NISTOR, V. On the geometry of Riemannian manifolds with a Lie structure at infinity. Int. J. Math. Math. Sci., 1-4 (2004), 161–193.
- [2] ANDROULIDAKIS, I., MOHSEN, O., AND YUNCKEN, R. The convolution algebra of Schwartz kernels along a singular foliation. J. Oper. Theory 85, 2 (2021), 475–503.
- [3] ANDROULIDAKIS, I., MOHSEN, O., AND YUNCKEN, R. A pseudodifferential calculus for maximally hypoelliptic operators and the Helffer-Nourrigat conjecture, 2022.
- [4] ANDROULIDAKIS, I., AND SKANDALIS, G. The holonomy groupoid of a singular foliation. J. Reine Angew. Math. 626 (2009), 1–37.
- [5] BESCHASTNYI, I. Closure of the Laplace-Beltrami operator on 2D almost-Riemannian manifolds and semi-Fredholm properties of differential operators on Lie manifolds. *Results Math.* 78, 2 (2023), Paper No. 59, 56.
- [6] BESCHASTNYI, I., CARVALHO, C., NISTOR, V., AND QIAO, Y. The integration of Lie-Rinehart algebras and pseudodifferential operators. In final preparation.
- [7] BESCHASTNYI, I., CARVALHO, C., NISTOR, V., AND QIAO, Y. Analysis on noncompact manifolds and index theory: Fredholm conditions and pseudodifferential operators. In *Extended abstracts* 2021/2022. *Methusalem lectures, Ghent, Belgium.* 2024, pp. 3–14.
- [8] BESCHASTNYI, I., AND QUAN, H. Self-adjointness criteria and self-adjoint extensions of the laplacebeltrami operator on α-grushin manifolds, 2024.

- [9] BOSCAIN, U., AND PRANDI, D. Self-adjoint extensions and stochastic completeness of the Laplace-Beltrami operator on conic and anticonic surfaces. J. Differential Equations 260, 4 (2016), 3234–3269.
- [10] CARVALHO, C., CÔME, R., AND QIAO, Y. Gluing action groupoids: Fredholm conditions and layer potentials. *Rev. Roumaine Math. Pures Appl.* 64, 2-3 (2019), 113–156.
- [11] CARVALHO, C., NISTOR, V., AND QIAO, Y. Fredholm conditions on non-compact manifolds: theory and examples. In *Operator theory, operator algebras, and matrix theory*, vol. 267 of *Oper. Theory Adv. Appl.* Birkhäuser/Springer, Cham, 2018, pp. 79–122.
- [12] CÔME, R. The Fredholm property for groupoids is a local property. *Results Math.* 74, 4 (2019), Paper No. 160, 33.
- [13] CONNES, A. Sur la théorie non commutative de l'intégration. In Algèbres d'opérateurs (Sém., Les Planssur-Bex, 1978), vol. 725 of Lecture Notes in Math. Springer, Berlin, 1979, pp. 19–143.
- [14] CONNES, A. Non commutative differential geometry. Publ. Math. IHES 62 (1985), 41-144.
- [15] DE VERDIÈRE, Y. C., DIETZE, C., DE HOOP, M. V., AND TRÉLAT, E. Weyl formulae for some singular metrics with application to acoustic modes in gas giants, 2024.
- [16] DEBORD, C., LESCURE, J.-M., AND ROCHON, F. Pseudodifferential operators on manifolds with fibred corners. Ann. Inst. Fourier 65, 4 (2015), 1799–1880.
- [17] DEBORD, C., AND SKANDALIS, G. Adiabatic groupoid, crossed product by \mathbb{R}^*_+ and pseudodifferential calculus. *Adv. Math.* 257 (2014), 66–91.
- [18] GALLONE, M., MICHELANGELI, A., AND POZZOLI, E. Quantum geometric confinement and dynamical transmission in Grushin cylinder. *Rev. Math. Phys.* 34, 7 (2022), Paper No. 2250018, 91.
- [19] GRIESER, D. Basics of the b-calculus. In Approaches to singular analysis. Based on the workshop, Berlin, Germany, April 8–10, 1999. Basel: Birkhäuser, 2001, pp. 30–84.
- [20] HERZ, J.-C. Pseudo-algèbres de Lie I, II, C. R. Acad. Sci. Paris 236 (1953), 1935–1937, 2289–2291.
- [21] HUEBSCHMANN, J. Poisson cohomology and quantization. J. Reine Angew. Math. 408 (1990), 57–113.
- [22] HUEBSCHMANN, J. Duality for Lie-Rinehart algebras and the modular class. J. Reine Angew. Math. 510 (1999), 103–159.
- [23] HUEBSCHMANN, J. Lie-Rinehart algebras, descent, and quantization. In Papers from the workshop on categorical structures for descent and Galois theory, Hopf algebras, and semiabelian categories, Toronto, ON, Canada, September 23–28, 2002. (2004), G. a. Janelidze, Ed., vol. 43, Fields Institute Communications, pp. 295–316.
- [24] KOTTKE, C. Functorial compactification of linear spaces. Proc. Amer. Math. Soc. 147, 9 (2019), 4067–4081.
- [25] KOTTKE, C., AND ROCHON, F. L²-cohomology of quasi-fibered boundary metrics. *Invent. Math.* 236, 3 (2024), 1083–1131.
- [26] LAURENT-GENGOUX, C., LOUIS, R., AND RYVKIN, L. An invitation to singular foliations. in final preparation.
- [27] LAUTER, R., MONTHUBERT, B., AND NISTOR, V. Pseudodifferential analysis on continuous family groupoids. Doc. Math. 5 (2000), 625–655 (electronic).
- [28] LAUTER, R., MONTHUBERT, B., AND NISTOR, V. Spectral invariance for certain algebras of pseudodifferential operators. J. Inst. Math. Jussieu 4, 3 (2005), 405–442.
- [29] LAUTER, R., AND SEILER, J. Pseudodifferential analysis on manifolds with boundary a comparison of b-calculus and cone algebra. In Oper. Theory Adv. Appl., vol. 125 of Adv. Partial Differ. Equ. 2001, pp. 131– 166.
- [30] LESCH, M. Operators of Fuchs type, conical singularitites, and asymptotic methods, vol. 136 of Teubner-Texte Math. Stuttgart: B. G. Teubner, 1997.
- [31] MACKENZIE, K. Lie groupoids and Lie algebroids in differential geometry, vol. 124 of LMS Lect. Note Series. Cambridge U. Press, Cambridge, 1987.
- [32] MAZZEO, R. Elliptic theory of differential edge operators. I. Commun. Partial Differ. Equations 16, 10 (1991), 1615–1664.
- [33] MELROSE, R. Transformation of boundary problems. Acta Math. 147 (1981), 149-236.
- [34] MELROSE, R. B. Geometric scattering theory. Stanford Lectures. Cambridge University Press, Cambridge, 1995.
- [35] MOERDIJK, I., AND MRČUN, J. Introduction to foliations and Lie groupoids, vol. 91 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2003.
- [36] MOERDIJK, I., AND MRCUN, J. On the universal enveloping algebra of a Lie algebroid. Proc. Amer. Math. Soc. 138 (2010), 3135–3145.
- [37] NISTOR, V. Groupoids and the integration of Lie algebroids. J. Math. Soc. Japan 52, 4 (2000), 847-868.
- [38] PALAIS, R. S. The cohomology of Lie rings. Proc. Sympos. Pure Math., III (1961), 130-137.
- [39] PATERSON, A. L. T. Continuous family groupoids. Homology Homotopy Appl. 2 (2000), 89-104.

SCHRÖDINGER OPERATORS

- [40] PATERSON, A. L. T. The equivariant analytic index for proper groupoid actions. *K-Theory 32*, 3 (2004), 193–230.
- [41] PETERSEN, P. Riemannian geometry, 2nd ed. ed., vol. 171 of Grad. Texts Math. New York, NY: Springer, 2006.
- [42] POZZOLI, E. Quantum confinement in α-Grushin planes. In Mathematical challenges of zero-range physics—models, methods, rigorous results, open problems, vol. 42 of Springer INdAM Ser. Springer, Cham, [2021] ©2021, pp. 229–237.
- [43] RENAULT, J. A groupoid approach to C*-algebras, vol. 793 of LNM. Springer, 1980.
- [44] RINEHART, G. Differential forms on general commutative algebras. *Trans. Amer. Math. Soc. 108* (1963), 195–222.
- [45] SAVIN, A. Elliptic operators on manifolds with singularities and *K*-homology. *K-Theory 34*, 1 (2005), 71–98.
- [46] SCHROHE, E. Fréchet algebra techniques for boundary value problems on noncompact manifolds: Fredholm criteria and functional calculus via spectral invariance. *Math. Nachr. 199* (1999), 145–185.
- [47] SCHULZE, B.-W. Pseudo-differential operators on manifolds with singularities, vol. 24 of Studies in Mathematics and its Applications. North-Holland Publishing Co., 1991.
- [48] VAN ERP, E., AND YUNCKEN, R. A groupoid approach to pseudodifferential calculi. J. Reine Angew. Math. 756 (2019), 151–182.
- [49] VASSOUT, S. Unbounded pseudodifferential calculus on Lie groupoids. J. Funct. Anal. 236, 1 (2006), 161– 200.
- [50] VASY, A. Propagation of singularities in many-body scattering. Ann. Sci. École Norm. Sup. (4) 34, 3 (2001), 313–402.
- [51] VASY, A. Geometry and analysis in many-body scattering. In *Inside out: inverse problems and applications*, vol. 47 of *Math. Sci. Res. Inst. Publ.* Cambridge Univ. Press, Cambridge, 2003, pp. 333–379.
- [52] WHEATLEY, R. J. Inverse power potentials: Virial coefficients and a general equation of state. *The Journal of Physical Chemistry B 109*, 15 (2005), 7463–7467. PMID: 16851856.
- [53] WILLIAMS, D. Crossed products of C*-algebras, vol. 134 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2007.

UNIVERSITÉ CÔTE D'AZUR, CENTRE INRIA D'UNIVERSITÉ CÔTE D'AZUR, 2004, ROUTE DES LUCI-OLES, 06902, SOPHIA ANTIPOLIS, FRANCE

Email address: ivan.beschastnyi@inria.fr

CENTRO DE ANÁLISE MATEMÁTICA, GEOMETRIA E SISTEMAS DINÂMICOS, DEP. MATEMÁTICA, INSTI-TUTO SUPERIOR TÉCNICO, UNIVERSITY OF LISBON, AV. ROVISCO PAIS 1, 1049-001 LISBON, PORTUGAL *Email address*: catarinaccarvalho@tecnico.ulisboa.pt

UNIVERSITÉ DE LORRAINE, CNRS, IECL, F-57000 METZ, FRANCE *Email address*: victor.nistor@univ-lorraine.fr

SCHOOL OF MATHEMATICS AND STATISTICS, SHAANXI NORMAL UNIVERSITY, XI'AN, 710119, CHINA *Email address*: yqiao@snnu.edu.cn