RANDOM MATRIX STATISTICS AND ZEROES OF L-FUNCTIONS VIA PROBABILITY IN λ-RINGS

SEAN HOWE

ABSTRACT. We introduce a theory of probability in λ -rings designed to efficiently describe random variables valued in multisets of complex numbers, varieties over a field, or other similar enriched settings. A key role is played by the σ -moment generating function based on the plethystic exponential, which allows us to describe distributions and argue with independence in a way that is as simple as classical probability theory. As a first application, we use this theory to obtain a concise description of the asymptotic σ -moment generating functions describing distributions of eigenvalues of Haar random matrices in compact classical groups. Beyond the theory of probability in λ -rings, the proof uses only classical invariant theory. Using our description we reprove the results of Diaconis and Shahshahani on the joint distributions of traces of powers of matrices, and we also treat symmetric groups. Next, we use Poonen's sieve to establish equidistribution results for the zeroes of L-functions in some natural families: simple Dirichlet characters for $\mathbb{F}_q(t)$ and the vanishing cohomology of smooth hypersurface sections. We give concise descriptions of the asymptotic σ -moment generating functions in these families, then compare them to the associated random matrix distributions. These equidistribution results are sideways in that we fix q and take the degree d to infinity, as opposed to Deligne equidistribution for fixed d as $q \to \infty$, and the large d-limits are related to explicit descriptions of stable homology with twisted coefficients.

CONTENTS

2 SEAN HOWE

1. INTRODUCTION

In this work we give a concise description of the eigenvalue distributions of orthogonal, symplectic, unitary, and symmetric groups, then use this description to compare the distributions of zeroes of function field L-functions to random matrix statistics in two natural families. We note that our comparison is "sideways" compared to [\[19\]](#page-60-1), in that we first fix q and compute a limiting distribution as the degree d goes infinity, then afterwards study how the limit depends on q as $q \to \infty$ to compare to random matrix statistics. This is in contrast to the approach of [\[19\]](#page-60-1), where one first applies Deligne equidistribution for a fixed d as $q \to \infty$, and then takes $d \to \infty$ purely at the level of random matrices (see also Remark [1.3.6\)](#page-10-0).

Along the way we obtain a new proof of the results of Diaconis-Shahshahani [\[11\]](#page-60-2) on the joint moments of traces of powers of matrices in classical groups and a probabilistic interpretation of certain generating functions that have previously arisen in stable cohomology computations. The main new tool is the theory of

probability in λ -rings and the accompanying σ -moment generating function. To make precise statements we begin by describing our results on random matrices in essentially classical terms, which will lead us naturally to the theory of λ -probability.

1.1. **Random matrices.** Let $G \subseteq GL_n(\mathbb{C})$ be a compact subgroup equipped with Haar measure, and let X be the function which sends a matrix $M \in G$ to its multiset of eigenvalues. From this function, we can extract many complex-valued random variables by taking symmetric functions of the eigenvalues. Concretely, let $\Lambda_n = \mathbb{Z}[t_1,\ldots,t_n]^{S_n}$ be the ring of symmetric polynomials in *n*-variables. Then, for any $f \in \Lambda_n$, we can evaluate f on the eigenvalues of a matrix $M \in G$ to obtain a random variable X_f on G .

Example 1.1.1.

• For the power sum symmetric functions $p_i = \sum_{j\geq 1} t_j^i$

$$
X_{p_i}(M) = \text{Tr}(M^i).
$$

• For the complete symmetric functions h_i , obtained by summing all monomials of degree i ,

$$
\frac{1}{\det(\mathrm{Id} - xM)} = \sum_{i \ge 0} X_{h_i}(M) x^i.
$$

• For the elementary symmetric functions e_i , obtained by summing all monomials of degree i where each variable appears at most once,

$$
\det(\mathrm{Id} + xM) = \sum_{i \ge 0} X_{e_i}(M)x^i.
$$

Note that we have $p_1 = h_1 = e_1$. In the above and below we also adopt the convention that $p_0 = h_0 = e_0 = 1$.

In order to compare the distributions of such random variables across different n , it will be useful consider the ring of symmetric functions $\Lambda = \lim_{n} \Lambda_n$. Symmetric functions can be evaluated on a multiset of n complex numbers by projection to the nth component. Then, we can encode the joint moments of all the random variables $X_f, f \in \Lambda$ simultaneously by considering the Λ -distribution μ_X of X,

$$
\mu_X: \Lambda \to \mathbb{C}, f \mapsto \mathbb{E}[X_f].
$$

In classical probability, the distribution of a (real-valued^{[1](#page-2-1)}) random variable Y is, in many cases, determined by its moments. One can thus encode the distribution of such a random variable using the moment generating function:

$$
\mathbb{E}[\exp(Yt)] = \sum_{i=0}^{\infty} \mathbb{E}\left[\frac{Y^i}{i!}\right]t^i.
$$

This is often a very concise description: for example, a Gaussian distribution of mean 0 and variance 1 is determined by its moment generating function $\exp(t^2)$.

To define the analog in λ -probability, we will need to name some elements of Λ indexed by partitions. By a partition, we mean a non-increasing sequence of non-negative integers (τ_1, τ_2, \ldots) that is eventually zero. Given a partition τ , let

 1 For a complex-valued random variable one should instead take joint moments with the complex conjugate. Most, but not all, of the classical random variables we encounter below are actually real-valued; for the ones that are not we will use joint moments with the complex conjugate.

 $m_{\tau} \in \Lambda$ denote the monomial symmetric function obtained by formally summing the monomials that can be obtained from $\prod t_j^{\tau_j}$ by permutation. Let

$$
p_{\tau} = \prod p_{\tau_j}, h_{\tau} = \prod h_{\tau_j}, \text{ and } e_{\tau} = \prod e_{\tau_j}.
$$

where the p_i , h_i , and e_i are as in Example [1.1.1.](#page-2-2) The m_{τ} , h_{τ} , and e_{τ} each form Z-bases for Λ as τ varies over all partitions, and the p_{τ} form a Q-basis for $\Lambda_{\mathbb{Q}}$.

We define the σ -moment generating function for any random variable X valued in multisets of complex numbers as above by

(1.1.1.1)
$$
\mathbb{E}[\text{Exp}_{\sigma}(Xh_1)] = \sum_{\tau \text{ a partition}} \mathbb{E}[X_{h_{\tau}}]m_{\tau} \in \Lambda_{\mathbb{C}}^{\wedge}.
$$

Here $\Lambda_{\mathbb C}^{\wedge}$ is the ring of symmetric power series obtained by completing $\Lambda_{\mathbb C}$ for the filtration by polynomial degree. Since the h_{τ} and m_{τ} are each Z-bases for Λ , giving the σ -moment generating function of X is equivalent to giving its Λ -distribution.

Remark 1.1.2. Here Exp_{σ} is the plethystic exponential which, for R a λ -ring, can be defined on a symmetric power series $x \in \Lambda_R^{\wedge}$ with constant term zero by

$$
Exp_{\sigma}(x) = \sum_{i=0}^{\infty} h_i \circ x,
$$

where $h_i \circ x$ denotes a plethystic action of h_i on the element x. In the theory of $λ$ -rings, $h_i \circ x$ is often written $σ_i(x)$. It is frequently a decategorified *i*th symmetric power, and thus a natural analog of $\frac{x^i}{i!}$ $\frac{x^i}{i!}$ in the usual exponential $\exp(x) = \sum_{i=0}^{\infty} \frac{x^i}{i!}$ $\frac{x}{i!}$.

Remark 1.1.3. The equivalence between Λ -distributions and σ -moment generating functions is the isomorphism from $\Lambda_{\mathbb C}^{\wedge}$ to $\text{Hom}_{\mathbb Z}(\Lambda,{\mathbb C})$ given by the Hall inner product $\langle \cdot, \cdot \rangle$ (this follows because $\langle m_{\tau}, h_{\gamma} \rangle = \delta_{\tau \gamma}$).

For a partition τ , we write $||\tau||$ for the number of entries in τ , i.e. the number of distinct variables appearing in each monomial of m_{τ} , and $|\tau|$ for the sum of τ , i.e. the degree of each monomial in m_{τ} . For example, $||(2, 2, 1)|| = 3$ and $|(2, 2, 1)| = 5$.

Theorem A (see Theorem [4.2.1\)](#page-25-1). For $n \geq 1$ and G_n a compact subgroup of $GL_n(\mathbb{C})$, we view G_n as a probability space using unit volume Haar measure, and we write X_n for the random variable sending $M \in G_n$ to the multiset of eigenvalues of M. With this notation:

(1) Let $G_n = O(n)$, the subgroup of matrices which are both real and unitary. Modulo the m_{τ} with $||\tau|| > n$,

$$
\mathbb{E}[\operatorname{Exp}_{\sigma}(X_n h_1)] \equiv \operatorname{Exp}_{\sigma}(h_2) = \prod_{1 \le i \le j} \frac{1}{1 - t_i t_j}.
$$

(2) If n is even, let $G_n = \text{Sp}(n)$, the subgroup of matrices which are both symplectic and unitary. Modulo the m_{τ} with $||\tau|| > n$,

$$
\mathbb{E}[\operatorname{Exp}_{\sigma}(X_n h_1)] \equiv \operatorname{Exp}_{\sigma}(e_2) = \prod_{1 \le i < j} \frac{1}{1 - t_i t_j}.
$$

(3) Let $G_n = \Sigma_n$, the subgroup of permutation matrices. Modulo the m_τ with $|\tau| > n$,

$$
\mathbb{E}[\text{Exp}_{\sigma}(X_n h_1)] \equiv \text{Exp}_{\sigma}(\text{Exp}_{\sigma}(h_1) - 1) = \text{Exp}_{\sigma}(h_1 + h_2 + ...)
$$

$$
= \prod_{\substack{non-constant \\ monomials m \\ \text{in } h_1, h_2, ...}} \frac{1}{1 - m}.
$$

(4) Let $G_n = U(n)$, the subgroup of unitary matrices. Modulo the $m_{\tau} \overline{m}_{\tau}$, with $||\tau|| > n$ or $||\tau'|| > n$,

$$
\mathbb{E}[\exp_{\sigma}(X_n h_1 + \overline{X}_n \overline{h}_1)] \equiv \exp_{\sigma}(h_1 \overline{h}_1) = \prod_{1 \le i,j} \frac{1}{1 - t_i \overline{t}_j}.
$$

Here \overline{X}_n is the conjugate while \overline{t}_i , etc., is meant just as another variable with no relation to t_i . The left-hand side is the joint σ -moment generating function describing the joint Λ -distribution of X_n and \overline{X}_n ,

$$
\mathbb{E}[\operatorname{Exp}_{\sigma}(X_n h_1 + \overline{X}_n \overline{h}_1)] = \sum_{\tau_1, \tau_2} \mathbb{E}[X_{n, h_{\tau_1}} \overline{X}_{n, h_{\tau_2}}] m_{\tau_1} \overline{m}_{\tau_2}.
$$

In particular, in each of (1)-(3) (resp. (4)), as $n \to \infty$, the Λ -distributions (resp. joint Λ-distributions) converge to the unique Λ-distribution (resp. joint $Λ$ -distribution) with the given $σ$ -moments (resp. joint $σ$ -moments).

Remark 1.1.4. In case (1)-(3) of Theorem [A,](#page-3-0) the random variables $X_{n,f}$ for $f \in \Lambda$ are all real valued functions on G_n , so it is only in case (4) that we need to consider not just the Λ-distribution but the joint Λ-distribution with the complex conjugate to encode the distributions of the associated classical random variables.

Remark 1.1.5. By Lemma [2.8.1,](#page-19-2) the standard involution ω on the ring of symmetric function swaps the asymptotic generating functions in the orthogonal and symplectic cases of Theorem [A.](#page-3-0) We thank Michael Magee for suggesting this.

To prove Theorem [A,](#page-3-0) we interpret the relevant coefficients of the moment generating functions as invariants in symmetric powers. For example, in $(1)-(3)$, using the expansion Eq. [\(1.1.1.1\)](#page-3-1) and orthogonality of characters, one finds the coefficient of m_{τ} is the multiplicity of the trivial representation of G_n in $Sym^{\tau} \mathbb{C}^n = \bigotimes_i Sym^{\tau_i} \mathbb{C}^n$. When $||m_{\tau}|| \leq n$, for (1) and (2) the first fundamental theorem of invariant theory gives a description of these coefficients that matches the counting problem for the coefficient of m_{τ} in the power series we have written down, while for (3) the matching is a combinatorial exercise. The argument in (4) is similar to $(1)-(2)$.

In Corollary [4.4.1,](#page-29-1) we deduce from Theorem $A-(1),(2)$ $A-(1),(2)$, and (4) the results of Diaconis and Shahshahani [\[11,](#page-60-2) Theorems 2, 4, and 6] on the joint moments of traces of powers of matrices in classical groups. By Example [1.1.1,](#page-2-2) to accomplish this we need to extract the values of the Λ -distribution on power sum monomials from the σ -moment generating function. By Remark [1.1.3,](#page-3-2) this can be done by taking the Hall inner product of the σ -moment generating function with power sum monomials. This inner product is straightforward to compute after using the alternative expansion of the σ -exponential via power sum plethysms described in Lemma [2.5.4.](#page-17-1) We note that our proof makes no use of classical, orthogonal, or symplectic Schur polynomials; in particular, this eliminates the use of Ram's theory

6 SEAN HOWE

for symplectic and orthogonal Schur polynomials that is necessary in the original proof in [\[11\]](#page-60-2).

For the symmetric group, [\[11,](#page-60-2) Theorem 7] computes the joint moments of the cycle counting functions. These can also be derived from the computation of the moment generating function in Theorem $A-(3)$ $A-(3)$, as we explain in Section [4.5.](#page-32-0)

Remark 1.1.6. Since each of h_2 , e_2 , and $h_1\overline{h}_1$ is a degree 2 homogeneous polynomial, the Λ -distributions appearing in Theorem [A-](#page-3-0)(1), (2), and (4) should be thought of as different λ -incarnations of Gaussians (recalling that a Gaussian of mean zero and variance 1 has moment generating function e^{t^2}). This is related to the fact that, when considering distributions of traces in Corollary [4.4.1,](#page-29-1) the traces of powers of matrices behave asymptotically as independent Gaussians. Similarly, the Λ -distribution appearing in Theorem [A-](#page-3-0)(3) should be though of as a λ incarnation of a Poisson distribution (noting that a classical Poisson distribution of mean 1 has moment generating function e^{e^t-1} , and this is related to the fact that the cycle counting functions behave asymptotically as independent Poisson random variables (see Section [4.5\)](#page-32-0). It would be interesting to formulate a more general principle governing this relation in all of these cases.

Remark 1.1.7. It is perhaps interesting to note that we discovered the formulas in Theorem [A](#page-3-0) by computing the arithmetic limits related to zeroes of L-functions discussed in Section [1.3](#page-6-0) below. We knew in advance that these limits should match random matrix statistics, but it was the final form of these arithmetic limits that led us to the simple description of the random matrix statistics.

1.2. λ -probability. Theorem [A](#page-3-0) is most naturally interpreted using a theory of probability in λ -rings. Here a λ -ring is a ring R equipped with a plethystic action of Λ , denoted $f \circ r$ for $f \in \Lambda$ and $r \in R$, satisfying certain natural compatibilities. A key example is the monoid ring $\mathbb{Z}[\mathbb{C}]$, where $\mathbb C$ is a monoid under multiplication so that $[z_1][z_2] = [z_1z_2]$ and the plethystic action is determined by $h_i \circ [z] = p_i \circ [z] = z^i$. A function on a probability space valued in multisets of complex numbers can be viewed as a function valued in $\mathbb{Z}[\mathbb{C}]$ to obtain a random variable valued in a λ -ring.

One can formulate Theorem [A](#page-3-0) in this setting, and such a formulation is useful for making basic probabilistic arguments.

Example 1.2.1. In Theorem [A-](#page-3-0)(3), note that the permutation representation of Σ_n can be decomposed as the direct sum of the standard irreducible representation V_n of dimension $n-1$ and the trivial representation. Let Z_n be the random variable on Σ_n sending $g \in \Sigma_n$ to the eigenvalues of g acting on V_n . If we view these multisets as elements of $\mathbb{Z}[\mathbb{C}]$, we have $Z_n = X_n - [1]$. Since any random variable is independent to the constant random variable [1], our formalism gives

$$
\mathbb{E}[\text{Exp}_{\sigma}(Z_n h_1)] = \mathbb{E}[\text{Exp}_{\sigma}(X_n h_1)] \mathbb{E}[\text{Exp}_{\sigma}(-[1]h_1)]
$$

= $\mathbb{E}[\text{Exp}_{\sigma}(X_n h_1)] \text{Exp}_{\sigma}(-h_1).$

From Theorem [A-](#page-3-0)(3), we deduce that, modulo the m_{τ} with $||\tau|| > n$,

$$
\mathbb{E}[\text{Exp}_{\sigma}(Z_n h_1)] \equiv \text{Exp}_{\sigma}(h_2 + h_3 + \ldots) = \prod_{\substack{\text{monomials } m \text{ with } \\ \deg m \ge 2 \text{ in } t_1, t_2, \ldots}} \frac{1}{1 - m}
$$

.

In fact there is an even more natural formulation of the general theory and Theorem [A:](#page-3-0) instead of considering random variables valued in λ -rings, once can consider abstract λ -rings of random variables equipped with an expectation functional. In particular, one can take the Grothendieck ring of representations of a compact group as a λ -ring of random variables and equip it with the expectation functional given by the integral of the trace or, equivalently, the multiplicity of the trivial representation. This is how we will understand Theorem [A](#page-3-0) in the body of the text (see Theorem [4.2.1\)](#page-25-1). In particular, this perspective explains why the moment generating functions are all power series with coefficients in $\mathbb Z$ even though the random variables X_f discussed above were all complex-valued.

1.3. Zeroes of L-functions. There is a large body of work in number theory relating the distributions of zeroes of L-functions to the distributions of eigenvalues of random matrices. The theory of λ -probability is well-suited for this type of question because of the simple form of the σ -moment generating functions in Theorem [A.](#page-3-0)

We illustrate this by studying the average behavior of zeroes of L-functions for two natural families: simple idèle class characters for $\mathbb{F}_q(t)$, and the vanishing cohomology of smooth hypersurface sections of smooth projective varieties over \mathbb{F}_q . As in the theory of random matrices, there are two natural λ -probability spaces that one can use for this study, one a literal function space and one more abstract.

Given a variety S/\mathbb{F}_q , the natural abstract ring of random variables is the Grothendieck ring of constructible ℓ -adic sheaves on S. Indeed, the families of L-functions we consider all arise from ℓ -adic local systems on such a variety S. However, the methods we will use in the present work depend only on a much simpler notion: in fact, we could formulate and prove most of our results without even knowing the definition of étale cohomology. Instead, for $W(\mathbb{C})$ the big Witt ring of $\mathbb C$, which we can think of concretely as the ring $\prod_{i=1}^{\infty} \mathbb C$ with plethystic operations determined by $p_j \circ (a_1, a_2, \ldots) = (a_j, a_{2j,\ldots})$, we consider the space $C(S(\overline{\mathbb{F}}_q), W(\mathbb{C}))$ of $W(\mathbb{C})$ -valued function on $S(\overline{\mathbb{F}}_q)$ that are invariant under the $Gal(\overline{\mathbb{F}}_q/\overline{\mathbb{F}}_q)$ -action (equivalently, one could think of these as functions on the closed points of S). This ring has a natural pointwise λ -ring structure, and an expectation functional

$$
\mathbb{E}: C(S(\overline{\mathbb{F}}_q), W(\mathbb{C})) \to W(\mathbb{C})
$$

that, for the *i*th coordinate, averages over the \mathbb{F}_{q^i} -points $s \in S(\mathbb{F}_{q^i}) \subseteq S(\mathbb{F}_q)$ the values $f(s)_{i/|s|}$, where |s| is the size of the Galois-orbit of s. To understand the motivation for this expectation, we note that the additive of group $W(\mathbb{C})$ is also naturally isomorphic to $1 + t\mathbb{C}[[t]]$, and under this isomorphism the formula for the expectation functional comes from the Euler product expansion of zeta functions.

After choosing a prime ℓ coprime to q and an isomorphism $\mathbb{Q}_{\ell} \cong \mathbb{C}$, it follows from [\[21,](#page-60-3) Théoréme (1.1.2)] that the Grothendieck ring of constructible \mathbb{Q}_{ℓ} sheaves embeds into $C(S(\overline{\mathbb{F}}_q), W(\mathbb{C}))$ by taking the sequence of traces of powers of Frobenius at each point (or, viewing $W(\mathbb{C})$ as $1+t\mathbb{C}[[t]]$, by taking the characteristic power series of Frobenius). This matches the pointwise λ -ring structure on $C(S(\overline{\mathbb{F}}_q), W(\mathbb{C}))$ with the usual λ -ring structure on ℓ -adic sheaves where the plethystic operations $h_k \circ$ (resp. $e_k \circ$) decategorify symmetric (resp. exterior) powers of local systems.

1.3.1. Some characters. We fix a finite field \mathbb{F}_q of order q and a prime ℓ dividing q − 1. We write $\mu_{\ell} \subseteq \mathbb{F}_q^{\times}$ for the ℓ th roots of unity in \mathbb{F}_q , a cyclic group of order l, and fix a non-trivial character $\chi : \mu_{\ell} \to \mathbb{C}^{\times}$. For any lth power free monic polynomial $f \in \mathbb{F}_q[x]$, we obtain a μ_{ℓ} -extension of $\mathbb{F}_q(x)$ by Kummer theory and thus, by composition with χ , a character χ_f of Gal($\mathbb{F}_q(x)/\mathbb{F}_q(x)$). By class field

8 SEAN HOWE

theory, we can also view χ_f as an idéle class character (which can also be described explicitly in terms of f). The χ_f of this form are a natural family of characters: as we vary over ℓ th-power-free monic polynomials of positive degree, they hit every equivalence class of character of order ℓ up to globally unramified twist exactly once. Moreover, if we vary only over degrees d not divisible by ℓ , then we obtain exactly the characters which are ramified at ∞ . Note that, for κ/\mathbb{F}_q a finite extension and $f \in \kappa[x]$ an ℓ th-power-free monic polynomial, we may also form χ_f as a character of $Gal(\kappa(x)/\kappa(x))$, and this is compatible with restriction.

Example 1.3.2. If $\ell = 2$ and $f \in \mathbb{F}_q[x]$ is monic of odd degree > 2, then χ_f is ramified at ∞ and, as in [\[2,](#page-60-4) §11], the L-function associated to χ_f is given by the Dirichlet series

$$
L(s, \chi_f) = \sum_{m \in \mathbb{F}_q[x] \text{ monic}} \binom{f}{m} |m|^{-s}
$$

where $\binom{f}{m}$ denotes the usual Legendre symbol for $\mathbb{F}_q[x]$. The zeta function $Z_{C_f}(t)$ of the hyperelliptic curve C_f with affine equation $y^2 = f(x)$ is of the form

$$
Z_{C_f}(t) = \frac{\mathcal{L}(t, \chi_f)}{(1-t)(1-qt)}
$$

where $\mathcal{L}(q^{-s}, \chi_f) = L(s, \chi_f)$. In particular, by the Weil conjectures, the roots of $\mathcal{L}(t, \chi_f)$ have absolute value $q^{-1/2}$.

In general, for $f \in \overline{\mathbb{F}}_q[x]$ monic ℓ -power free, and χ_f the associated idéle class character of $\mathbb{F}_q(f)(x)$ (where $\mathbb{F}_q(f)$ is the extension of \mathbb{F}_q obtained by adjoining the coefficients of f), there is a polynomial $\mathcal{L}(t, \chi_f) \in \mathbb{C}[t]$ whose roots have absolute value $(\# \mathbb{F}_q(f))^{-\frac{1}{2}} = q^{-\frac{1}{2} [\mathbb{F}_q(f) : \mathbb{F}_q]}$ such that

$$
L(s,\chi_f) = \mathcal{L}(t,\chi_f)|_{t=q^{-s}}.
$$

In what follows, we note that the plethystic exponential Exp_{σ} and logarithm Log_{σ} can be used to define powers $f^r = \text{Exp}_{\sigma}(r\text{Log}_{\sigma}(f))$ whenever f is power series with constant term 1 and coefficients in a λ -ring R and $r \in R$ (see Sec-tion [2.6\)](#page-18-0). We will use this to give a succint expression of the limiting σ -moment generating function. We also note that there is a natural embedding of λ -rings $\mathbb{Z}[\mathbb{C}] \hookrightarrow W(\mathbb{C}), [z] \mapsto (z, z^2, z^3, \ldots).$ In particular, we may view a multiset of complex numbers as an element of $W(\mathbb{C})$. Finally, we note that we write $W(\mathbb{C})^{\text{bdd}}$ for the sub- λ -ring of $W(\mathbb{C}) = \prod_{i \geq 1} \mathbb{C}$ consisting of (a_1, a_2, \ldots) such that there exists an $M > 0$ with $|a_i| \leq M$ for all i. The following result will discuss congruences in $W(\mathbb{C})^{\text{bdd}}$ modulo $[q^{-1}]$; this is a convenient algebraic way to express that two quantities described over \mathbb{F}_{q^n} (obtained by projecting to the *n*th component of $W(\mathbb{C}))$ agree up to $O(q^{-n})$ — see Section [9](#page-54-0) for more details on this translation.

Theorem B (Theorem [7.3.2](#page-48-0) and Proposition [9.2.1;](#page-56-1) cf. also Theorem [7.2.2\)](#page-43-1). Let q be a prime power, let ℓ be a prime dividing $q-1$, and let $\chi : \mu_{\ell} \to \mathbb{C}^{\times}$ be a non-trivial character of the group μ_{ℓ} of ℓ th roots of unity in \mathbb{F}_q . Let $U_{d,\ell}(\overline{\mathbb{F}}_q)$ denote the set of degree d monic ℓ th-power-free polynomials over $\overline{\mathbb{F}}_q$. Let X_d be the $W(\mathbb{C})$ -valued random variable on $U_{d,\ell}(\overline{\mathbb{F}}_q)$ sending f to the set of zeroes of $\mathcal{L}(\chi_f,(\#\mathbb{F}_q(f))^{-\frac{1}{2}}t)$, where $\mathbb{F}_q(f)$ is the extension generated by the coefficients of f and χ_f is the idèle class character of $\mathbb{F}_q(f)(x)$ associated to f and χ by Kummer theory as above.

For $\ell = 2$, *i.e.* for χ quadratic,

$$
\lim_{\substack{d \to \infty \\ \ell \nmid d}} \mathbb{E}[\exp_{\sigma}(X_d h_1)] = \left(1 + \frac{[q]}{[q] + 1} \sum_{k \ge 1} [q^{-k}] e_{2k}\right)^{[q]}
$$
\n
$$
\equiv \exp_{\sigma}(e_2) \mod [q^{-1}] \Lambda_{W(\mathbb{C})^{\text{bdd}}}^{\wedge}.
$$

where we recall that, by Theorem [A-](#page-3-0)(2), $Exp_{\sigma}(e_2)$ encodes the eigenvalue distribution of a random compact symplectic matrix .

For $\ell > 2$, i.e. for χ non-quadratic,

$$
\lim_{\substack{d \to \infty \\ \ell \nmid d}} \mathbb{E}[\exp_{\sigma}(X_d h_1 + \overline{X}_d \overline{h}_1)] =
$$
\n
$$
\left(1 + \frac{[q^{\ell-1}]}{[q^{\ell-1}] + \ldots + [q] + 1} \sum_{\substack{k_1 \geq 1 \text{ or } k_2 \geq 1 \\ k_1 \equiv k_2 \mod \ell}} (-1)^{k_1 + k_2} [q^{-\frac{k_1 + k_2}{2}}] e_{k_1} \overline{e}_{k_2} \right)^{[q]}
$$
\n
$$
\equiv \exp_{\sigma}(h_1 \overline{h}_1) \mod [q^{-1}] \Lambda_{W(\mathbb{C})^{\text{bdd}}}^{\wedge}.
$$

where we recall that, by Theorem $A-(4)$ $A-(4)$, $Exp_{\sigma}(h_1h_2)$ encodes the eigenvalue distribution of a random unitary matrix.

Note that we equip $W(\mathbb{C})$ with the product topology, so that convergence can be treated after projection to the ith component. By the definition of the expectation functional, this reduces to a problem on the finite λ -probability space of $W(\mathbb{C})$ valued functions on $U_{d,\ell}(\mathbb{F}_{q^i})$ equipped with the C-valued expectation obtained by averaging the first component. The $W(\mathbb{C})$ -valued random variables on these finite probability spaces can be made explicit as infinite sums of random variables parameterized by closed points in $\mathbb{A}^1_{\mathbb{F}_{q^i}}$, and an application of Poonen's [\[23\]](#page-60-5) sieve shows that they are asymptotically independent and computes their asymptotic σ -moment generating functions; the asymptotic independence implies that the moment generating function we are interested in is a product of the moment generating functions of these variables. The ith component of the power appearing on the right-hand side can also be made very explicit, allowing us to match it with the result.

Remark 1.3.3. In Example [7.2.3](#page-46-0) we show that the case of Theorem [B](#page-7-0) where χ is quadratic (i.e. $\ell = 2$) implies [\[2,](#page-60-4) Theorem 11.2.3] by projecting to the first component and applying the standard involution ω on the ring of symmetric functions. In [\[2\]](#page-60-4) this result is proved in two ways: first, it is deduced from the Poincar´e series for stable cohomology of $[2,$ Theorem 1.6, the description of the Frobenius action in $[2,$ Theorem 1.10], and bounds on the size of unstable cohomology, by substituting in $[q^{-1}]$ for z and $[q^k]h_{2k}$ for h_{2k} as described in [\[2,](#page-60-4) Top of p.4]. Second, it is deduced from a more direct argument using generating functions. By contrast, our proof of Theorem [B](#page-7-0) is by an application of Poonen's sieve [\[23\]](#page-60-5). The relation between the second method of [\[2\]](#page-60-4) and our proof here seems to be similar to the relation between studying square-free polynomials as a configuration space via the methods of [\[17\]](#page-60-6) or as a special case of smooth hypersurface sections via the methods of [\[16\]](#page-60-7).

Example 1.3.4. As in [\[2\]](#page-60-4), one classical random variable we are interested in is the central value $L(\chi_f, 1/2)$. Theorem [B](#page-7-0) is insufficient to obtain the refined results of [\[2\]](#page-60-4), but it is sufficient to study natural truncations: indeed, for $f \in U_{d,\ell}(\overline{\mathbb{F}}_q)$,

$$
L(\chi_f, 1/2) = \sum_{i=0}^{\infty} (-1)^i (e_i \circ X_d)_1(f),
$$

where the subscript 1 denotes the first projection from $W(\mathbb{C})$ to \mathbb{C} , thus for any k we can take the truncation

$$
L^{[k]}(\chi_f, 1/2) = \sum_{i=0}^k (-1)^i (e_i \circ X_d)_1(f).
$$

Projecting onto the first component in Theorem [B,](#page-7-0) we find that the distribution of $L^{[k]}(\chi_f, 1/2)$ as we vary over $f \in U_{d,\ell}(\mathbb{F}_q)$ approximates, as $d \to \infty$, the distribution of the value at 1 of the k-truncation of the characteristic polynomial of a random compact symplectic matrix if χ is quadratic or a random unitary matrix if χ has order $\ell > 2$. Projecting onto the *n*th component gives a similar statement for $f \in U_{d,\ell}(\mathbb{F}_{q^n})$, and the approximation is up to an error term controlled by q in the sense that, if we vary the limit over \mathbb{F}_{q^n} for all n, the error term is $O(q^{-n})$.

1.3.5. Vanishing cohomology of smooth hypersurface sections. By a similar method, we compute the Λ-distributions of zeroes of L-functions of vanishing cohomology of smooth hypersurface sections of smooth projective varieties over \mathbb{F}_q . Recall (e.g. from [\[16,](#page-60-7) §4]) that the vanishing cohomology of a smooth hypersurface section H/\mathbb{F}_q of a smooth projective variety Y/\mathbb{F}_q is the part of the cohomology of H that is not visible already in the cohomology of Y (by weak Lefschetz and Poincaré duality, it lies in the middle degree cohomology of H). As in the case of characters considered above, we write $\mathcal{L}_H(t) \in \mathbb{C}[t]$ for the polynomial version of the L-function of the vanishing cohomology — its roots have absolute value $q^{-\frac{\dim H}{2}}$ and, for

$$
L_H(s) = \mathcal{L}_H(q^{-s}),
$$

 $L_H(s)^{(-1)^{\dim Y}}$ is a factor of $\zeta_H(s) = Z_H(q^{-s})$ (which is a rational function in q^{-s}). We write $[Y/\mathbb{F}_q] = (\#Y(\mathbb{F}_q), \#Y(\mathbb{F}_{q^2}), \ldots) \in W(\mathbb{C})$ and $[H^i(Y)] \in W(\mathbb{C})$ for the

sequence of traces of powers of Frobenius on the *i*th étale cohomology of $Y_{\overline{\mathbb{F}}_q}$.

Theorem C (Theorem [8.4.2](#page-53-2) and Proposition [9.2.2\)](#page-58-0). Let $Y \subseteq \mathbb{P}_{\mathbb{F}_q}^m$ be an $(n +$ 1)-dimensional smooth irreducible closed sub-variety, and let $U_d(\overline{\mathbb{F}}_q)$ be the set of nonzero homogenenous degree d polynomials F in $m + 1$ variables over $\overline{\mathbb{F}}_q$ whose vanishing locus $V(F)$ intersects Y transversely. Let X_d be the $W(\mathbb{C})$ -valued random variable on $U_d(\overline{\mathbb{F}}_q)$ sending F to the set of zeroes of $L_{V(F)\cap Y}(t(\#\mathbb{F}_q(F))^{-\frac{n}{2}})$, where $\mathbb{F}_q(F)$ is the extension generated by the coefficients of F and $V(F) \cap Y$ is a variety over $\mathbb{F}_q(F)$. Then,

$$
\lim_{d \to \infty} \mathbb{E}[\exp_{\sigma}(X_d h_1)] = (1 + p \sum_{i \ge 1} [q^{-in/2}] \epsilon^i f_i]^{[Y/\mathbb{F}_q]} \cdot \exp_{\sigma}(\epsilon \mu h_1)
$$

where $f_i = e_i$ and $\epsilon = -1$ if n is odd and $f_i = h_i$ and $\epsilon = 1$ if n is even,

$$
p = \frac{[q]^{n+1} - 1}{[q]^{n+2} - 1}, \text{ and } \mu = -\epsilon \left(\sum_{i=0}^{n-1} (-1)^i \left([q^{\frac{-n}{2}}] + [q^{\frac{n-2i}{2}}] \right) [H^i(Y)] \right) - [q^{-n/2}] [H^n(Y)].
$$

In particular, modulo $[q^{-\frac{1}{2}}] \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}}$ this agrees with:

(1) For $n > 0$ even, $Exp_{\sigma}(h_2)$ (as in Theorem [A-](#page-3-0)(1), the orthogonal case).

PROBABILITY IN λ -RINGS 11

- (2) For n odd, $Exp_{\sigma}(e_2)$ (as in Theorem [A-](#page-3-0)(2), the symplectic case).
- (3) For $n = 0$, $Exp_{\sigma}(h_2 + h_3 + ...)$ (the distribution associated to the standard irreducible representation of the symmetric group as in Example [1.2.1\)](#page-5-1).

Thus the asymptotic Λ -distribution of X_d approximates the asymptotic eigenvalue Λ -distribution of the associated random matrices modulo $[q^{-1/2}] \Lambda_{W(\mathbb{C})^{\text{bdd}}}$.

Remark 1.3.6. In each case $(1), (2), (3)$ enumerated at the end of Theorem [C,](#page-9-0) one can show that for a smooth hypersurface section one obtains a normalized Frobenius conjugacy class in the associated group whose eigenvalues are the zeroes of the normalized L-function (an orthogonal group in case (1), a symplectic group in case (2), and a symmetric group Σ_m acting through its irreducible $m-1$ dimensional representation in case (3)). This is essentially because in case (1) the cup product on vanishing cohomology is a non-degenerate symmetric pairing, in case (2) it is a non-degenerate antisymmetric pairing, and, in the zero-dimensional case (3), when we pass to cohomology taking the vanishing part corresponds exactly to splitting of the standard irreducible from the standard permutation representation. However, the fact that one has such a conjugacy class plays no role whatsoever in the proof of Theorem [C.](#page-9-0) It is interesting to contrast this to the situation when one takes first a limit as $q \to \infty$ and then afterwards as $d \to \infty$ as in [\[19\]](#page-60-1), where the computation of the monodromy group is crucial in order to apply Deligne equidistribution; we note further that with that order of limits, only the degree 0 cohomology of local systems matter, whereas our results detect cohomology in all degrees. In [\[8\]](#page-60-8) we will obtain an analog of Theorem [C](#page-9-0) using cohomological realizations in place of the point-counting realization that can be used, in particular, to determine the degree zero cohomology of local systems and thus also the monodromy groups.

Remark 1.3.7. Theorem [C](#page-9-0) and its proof are closely related to [\[16,](#page-60-7) Theorems B and C. The main improvement here is the use of the σ -moment generating function in place of what amounts, in hindsight, to a λ -variant of a falling moment generating function. In particular, this improvement is what makes possible the explicit comparison with random matrix statistics. Note also that, in [\[16\]](#page-60-7) we attached to each partition τ a sequence of local systems \mathcal{V}_{τ} over the moduli of smooth hypersurface sections by taking the irreducible representation of the symplectic, orthogonal, or symmetric group associated to τ and composing it with the monodromy representation on vanishing cohomology, and our main goal in [\[16\]](#page-60-7) was to show that homological stability held for these families of local systems at the level of enriched Euler characteristics. From Theorem [C](#page-9-0) we obtain better explicit formulas than those described in [\[16\]](#page-60-7): to extract the stable point count for a family of local system associated to a partition τ from Theorem [C,](#page-9-0) one needs only compute the Hall inner product between the moment generating function and the associated orthogonal, symplectic, or symmetric Schur polynomial. This is, both conceptually and practically, simpler than the method described in [\[16,](#page-60-7) Appendix A]. We note that there has also been recent progress towards establishing homological stability in this setting [\[1\]](#page-60-9), but since there is no known control over the unstable cohomology, the homological stability results do not imply point counting results.

1.3.8. Generalizations, independence, and motivic Euler products. The methods used to treat these families can be used in other cases: for example, they apply also to the L-functions of the vanishing cohomology of complete intersections by using the version of Poonen's sieve treated in [\[9\]](#page-60-10), and to other families using other generalizations of Poonen's sieve such as [\[12\]](#page-60-11) and [\[3\]](#page-60-12); we plan to give further examples in $[4]$. In fact, even with just the results of $[23]$, it should be possible to prove a much more general result from which both Theorem [B](#page-7-0) and Theorem [C](#page-9-0) can be deduced by a computation of local factors in a motivic Euler product. Indeed, a more robust way to prove Theorem [B](#page-7-0) and Theorem [C](#page-9-0) would be to show that the random variables appearing are infinite sums of asymptotically independent random variables parameterized by the $\overline{\mathbb{F}}_q$ -points of a variety (\mathbb{A}^1 in the case of Theorem [B](#page-7-0) and Y in the case of Theorem [C\)](#page-9-0), and that the σ -moment generating function of such a sum is given by a motivic Euler product of the σ -moment generating functions of the random variables in the family. However, formulating a good notion of independence for such a family is a bit delicate, so that the statement of a result in the most natural level of generality and its proof requires the development of considerably more machinery than we have introduced here. In particular, this will require a theory of independence for such families that will be developed in $[8]$ (along with analogs in the Grothendieck ring of varieties and Hodge structures), and a compatibility between motivic Euler products and plethystic exponentials that will be explained in $[6]$. The reason we are able to get away with less in the special cases of Theorem [A](#page-3-0) and Theorem [B](#page-7-0) is essentially because the moment generating functions are powers rather than more general motivic Euler products.

1.4. **Organization.** In Section [2](#page-12-0) we recall the theory of pre- λ and λ -rings. The main novelty in our treatment is that we emphasize monoid rings as the primitive example, and this facilitates many of the explicit computations we make with plethysm later in the paper. In Section [3,](#page-20-0) we introduce the basic notions of probability in λ -rings. In fact, we work in the generality of pre- λ rings — none of the foundational results depend on the added conditions that cut out λ -rings among pre- λ -rings, and there are interesting examples of pre- λ rings that may not be λ rings, such as Grothendieck rings of varieties, where one hopes to apply the theory. In addition to the σ -moment generating function, we also discuss more general moment generating functions and give examples of some important Λ-distributions. In Section [4](#page-24-1) we prove Theorem [A](#page-3-0) and deduce from it the main results of [\[11\]](#page-60-2).

In Section [5](#page-33-0) we setup an abstract point counting formalism and establish several basic results and lemmas that go into the proofs of Theorem [B](#page-7-0) and Theorem [C.](#page-9-0) In Section [6](#page-40-0) we apply Poonen's sieve to give a general computation of σ -moment generating function for certain $W(\mathbb{C})$ -valued random variables on finite probability spaces cut out by Taylor conditions (Proposition $6.2.1$); this is used in the proofs of Theorem [B](#page-7-0) and Theorem [C.](#page-9-0) In Section [7](#page-42-0) we compute the limits in Theorem [B](#page-7-0) and discuss complements, and in Section [8](#page-49-0) we compute the limits in Theorem [C](#page-9-0) and discuss complements. Finally, in Section [9](#page-54-0) we complete the proofs of Theorem [B](#page-7-0) and Theorem [C](#page-9-0) by comparing the limiting σ -moment generating functions computed therein with the limiting σ -moment generating functions for the associated random matrix distributions as computed in Theorem [A](#page-3-0) (and Example [1.2.1\)](#page-5-1).

1.5. Acknowledgements. This work owes an enormous debt to our collaborations with Margaret Bilu and Ronno Das, whom we thank both for these collaborations and for other helpful conversations. We thank Guillaume Dubach and Michael Magee for separate helpful conversations about random matrices and the Weingarten calculus (which we do not use anywhere, though perhaps we should!). We thank Dan Petersen for a helpful conversation about [\[2\]](#page-60-4). We thank Ravi Vakil for encouraging us several years ago to write up some of our ideas on this subject, and Helmut Höfer for inviting us to give a mathematical conversation at the Institute for Advanced Study, which provided the perfect opportunity to revisit that project.

During the preparation of this work, Sean Howe was supported by the National Science Foundation through grant DMS-2201112 and as a member at the Institute for Advanced Study during the academic year 2023-24 by the Friends of the Institute for Advanced Study Membership.

2. $(PRE-) \lambda$ -RINGS

In this section we recall the theory of pre- λ and λ -rings. One novelty in our treatment is that we emphasize monoid rings as the atomic examples of pre- λ rings. In particular, we use this to construct the pre- λ ring structure on the ring of symmetric functions Λ by embedding it in an infinite power series ring. This perspective is extremely helpful for making computations with plethysm, since the plethystic action on the power series ring (and other monoid rings) is very simple. In particular, we use this in Section [3](#page-20-0) to give explicit computations of σ -moment generating functions (including the crucial expansion Eq. $(1.1.1.1)$) of the σ -moment generating function). Some of the results on powers in Section [2.6](#page-18-0) are also of independent interest; for example, Example [2.6.5](#page-18-1) and the surrounding discussion explain a general construction of symmetric functions that, in particular, simplifies the treatment of configuration symmetric functions given in [\[17\]](#page-60-6).

2.1. Symmetric functions. We recall some standard definitions and results for symmetric functions (see, e.g., [\[22\]](#page-60-15)). Let $\Lambda_n := \mathbb{Z}[t_1,\ldots,t_n]^{S_n}$, the ring of symmetric polynomials in *n*-variables, and let $\Lambda := \lim_{n} \Lambda_n$, the ring of symmetric functions, where the transition map from Λ_{n+1} to Λ_n sends t_i to t_i for $1 \leq i \leq n$ and sends t_{n+1} to 0. We will use the following elements of Λ :

- The power sum symmetric functions $p_i = \sum_j t_j^i$. j
- The complete symmetic functions $h_i = \sum_{\sum a_j=i, a_j \in \mathbb{Z}_{\geq 0}} \prod_j t_j^{a_j}$
- • The elementary symmetric functions $e_i = \sum_{n=1}^{\infty} a_{i} = i, a_j \in \{0,1\} \prod_j t_j^{a_j}$.

These give polynomial bases:

(2.1.0.1)
$$
\Lambda = \mathbb{Z}[e_1, e_2, \ldots] = \mathbb{Z}[h_1, h_2, \ldots] \text{ and } \Lambda_{\mathbb{Q}} = \mathbb{Q}[p_1, p_2, \ldots].
$$

By a partition, we mean a non-increasing sequence of non-negative integers (τ_1, τ_2, \ldots) that is eventually zero. For $i \geq 0$, we write $\tau(i)$ for the multiplicity of i in τ , i.e. number of occurences of i in τ . Thus we can also view a partition as a function from $\mathbb{Z}_{\geq 1}$ to $\mathbb{Z}_{\geq 0}$ with finite support. For τ a partition, we write

$$
p_\tau := \prod_i p_i^{\tau(i)}, h_\tau := \prod_i h_i^{\tau(i)}, e_\tau := \prod_i e_i^{\tau(i)}.
$$

By Eq. [\(2.1.0.1\)](#page-12-2), $\{e_{\tau}\}_{\tau}$ and $\{h_{\tau}\}_{\tau}$ each give a Z-basis for Λ, while $\{p_{\tau}\}_{\tau}$ is a basis for $\Lambda_{\mathbb{Q}}$. There is also a Z-basis $\{m_{\tau}\}_{\tau}$ for Λ , where m_{τ} is the monomial symmetric function obtained by summing the monomials that can be obtained from permutations of $t^{\tau} = \prod_{i \geq 1} t_i^{\tau(i)}$, e.g.

$$
m_{(2,1,0,0,...)} = (t_1^2 t_2 + t_1^2 t_3 + \ldots) + (t_2^2 t_1 + t_2^2 t_3 + \ldots) + \ldots
$$

In particular, note that

$$
p_1 = p_{(1,0,0,\ldots)} = e_1 = e_{(1,0,0,\ldots)} = h_1 = h_{(1,0,0,\ldots)} = m_{(1,0,0,\ldots)} = \sum_i t_i.
$$

We recall some well-known relations between complete, elementary, and power sum symmetric functions: first, there are the obvious identities in $\Lambda[[z]]$

$$
\sum h_k z^k = \prod_{j \ge 1} \frac{1}{1 - t_j z} \text{ and } \sum e_k z^k = \prod_{j \ge 1} (1 + t_j z).
$$

It follows, in particular, by substituting $-z$ for z in the second identity that

(2.1.0.2)
$$
\left(\sum h_k z^k\right) \left(\sum (-1)^k e_k z^k\right) = 1.
$$

By taking the logarithm in $\Lambda_{\mathbb{Q}}[[z]]$ we also find

(2.1.0.3)
$$
\prod_i \exp\left(\frac{p_i z}{i}\right) = \prod_{j\geq 1} \frac{1}{1 - t_j z} = \sum h_k z^k.
$$

2.2. Pre- λ -rings. We now recall the theory of pre- λ rings. Some of this material is standard and can be found in, e.g., $[17, 14]$ $[17, 14]$ or elsewhere, but we are not aware of another source for the specific material on monoid rings developed below.

Definition 2.2.1. A pre- λ ring is a commutative ring R equipped with a *plethystic* action map

$$
R \to \mathrm{Hom}_{\mathrm{ring}}(\Lambda, R), r \mapsto (f \mapsto f \circ r)
$$

such that

(1)
$$
h_k \circ (r+s) = (h_k \circ r) + (h_{k-1} \circ r)(h_1 \circ s) + \ldots + (h_k \circ s).
$$

(2) $h_1 \circ r = r.$

This matches with, e.g., [\[17,](#page-60-6) Definition 2.2], by setting $h_k \circ = \sigma_k$. We note that part (2) is not always included in the definition of a pre- λ ring, but as in [\[17\]](#page-60-6) it is a natural condition for our purposes.

It can be deduced from Eq. $(2.1.0.2)$ and Eq. $(2.1.0.3)$ that, in any pre- λ ring R,

$$
e_k \circ (r+s) = (e_k \circ r) + (e_{k-1} \circ r)(e_1 \circ s) + \ldots + (e_k \circ s)
$$

$$
p_k \circ (r+s) = (p_k \circ r) + (p_k \circ s).
$$

Moreover, verifying the first of these is equivalent to verifying the analogous condition for $h_k \circ$ in Definition [2.2.1,](#page-13-3) and the second is equivalent if R is torsion-free as a Z-algebra.

In particular, the operations $p_k \circ$ are endomorphisms of the underlying abelian group $(R, +)$. One often writes $\psi_k = p_k \circ$ and refers to it as the kth Adams operation.

Example 2.2.2. Z admits a natural pre- λ ring structure such that the σ -operations $\sigma_k = h_k \circ$ decategorify symmetric powers of vector spaces and the λ -operations $e_k \circ$ decategorify exterior powers of vector spaces:

for
$$
n \ge 1
$$
, $h_k \circ n = \dim \text{Sym}^k \mathbb{C}^n = {k+n-1 \choose n-1}$, $e_k \circ n = \dim \bigwedge^k \mathbb{C}^n = {n \choose k}$.

One can show for this structure that $p_k \circ$ is the identity morphism for all k.

The ring of symmetric functions Λ itself has a natural pre- λ ring structure, and using this we will define the notion of a λ -ring. For making computations, it will be useful to understand this pre- λ ring structure on Λ as the restriction of a very simple pre- λ ring structure on a ring of power series. More generally, monoid rings over pre- λ rings have a natural pre- λ structure, as we explain now.

We use the following notation: given a set S and $k \geq 0$, we write $\text{Sym}^k S =$ S^k/Σ_k . A commutative monoid is a set M equipped with a commutative and associative multiplication law $M \times M \to M$, $(m_1, m_2) \mapsto m_1 m_2$ admitting an identity element 1_M . It is locally finite if the fibers of $M \times M \to M$ are finite.

Definition 2.2.3. Let M be a commutative monoid and let R be a ring.

(1) We write $R[M]$ for the associated monoid ring, i.e. the convolution algebra of finitely supported R -valued functions on M , addition is pointwise and

$$
fg(m) = \sum_{\substack{(m_1,m_2)\in M^2\\ \text{s.t. } m_1m_2=m}} f(m_1)f(m_2).
$$

For $m \in M$, we write $[m]$ for the function that send m to 1_R and all other elements to zero. Thus any element is a finite sum $\sum a_m[m]$, and the multiplication is determined by $[m_1][m_2] = [m_1m_2]$ for $m_1, m_2 \in M$. It is a commutative ring with unit $[1_M]$.

(2) If M is locally finite then we write $R[[M]]$ for convolution algebra of all R-valued functions on M. It is a commutative ring containing $R[M]$.

If R is a pre- λ ring, we equip $R[M]$ and, if M is locally finite, $R[[M]]$, with the plethystic action of Λ determined by

$$
(h_k \circ f)(m) = \sum_{\substack{\sum a_n n \in \text{Sym}^k M \\ \text{s.t. } \prod n^{a_n} = m}} \prod h_{a_n} \circ f(n)
$$

Note that here the a_n are non-negative integers and are zero for all but finitely many n , so that the sum and product are both finite.

When M is locally finite, we topologize $R[[M]]$ with the weak topology (where R is equipped with the discrete topology),. so that a net $(f_i)_{i\in I}$ in $R[[M]]$ converges to $f \in R[[M]]$ if and only if $f_i(m)$ is eventually constant equal to $f(m)$. Then $R[M]$ is dense in $R[[M]]$, and the ring structure and plethystic actions of symmetric functions $f \in \Lambda$ are evidently continuous for this topology.

Lemma 2.2.4. If R is a pre- λ ring and M is a commutative monoid, then $R[M]$ is a pre- λ ring for the plethystic action as in Definition [2.2.3.](#page-14-0) If M is furthermore locally finite, then $R[[M]]$ is a pre- λ ring for the plethystic action as in Definition [2.2.3.](#page-14-0) In particular, in these pre-λ-ring structures we have

$$
h_k \circ (a[m]) = (h_k \circ a)[m^k] \text{ and } e_k \circ (a[m]) = (e_k \circ a)[m^k].
$$

If R is torsion free, then we have moreover

$$
(p_k \circ f)(m) = \sum_{\substack{n \in M \\ s.t. \; n^k = m}} p_k \circ f(n) \text{ and, in particular } p_k \circ (a[m]) = (p_k \circ a)[m^k].
$$

Proof. We first show that the plethystic action defined in Definition [2.2.3](#page-14-0) is a pre- λ ring structure. The verification is essentially the same as the verification that convolution distributes over addition, but we spell it out for completeness.

$$
(h_k \circ (f+g))(m) = \sum_{\substack{\sum c_n n \in \text{Sym}^k M \\ \text{s.t. } \prod n^{c_n} = m}} \prod_{n^{c_n} n} h_{c_n} \circ (f(n) + g(n))
$$

\n
$$
= \sum_{\substack{\sum c_n n \in \text{Sym}^k M \\ \text{s.t. } \prod n^{c_n} = m}} \prod_{i+j = c_n} (h_i \circ f(n))(h_j \circ g(n))
$$

\n
$$
= \sum_{\substack{\sum a_n n \in \text{Sym}^{k_1} M, \sum b_n n \in \text{Sym}^{k_2} M \\ \text{s.t. } k_1 + k_2 = k, \prod n^{a_n + b_n} = m}} \prod_{n^{a_n + b_n} = m} (h_{a_n} \circ f(n))(h_{b_n} \circ g(n))
$$

\n
$$
= \sum_{\substack{m_1 m_2 = m \\ k_1 + k_2 = k}} \left(\sum_{\substack{\sum a_n n \in \text{Sym}^{k_1} M \\ \text{s.t. } \prod n^{a_n} = m_1}} h_{a_n} \circ f(n) \right) \left(\sum_{\substack{\sum b_n n \in \text{Sym}^{k_2} M \\ \text{s.t. } \prod n^{b_n} = m_2}} h_{b_n} \circ g(n) \right)
$$

\n
$$
= \sum_{k_1 + k_2 = k} ((h_{k_1} \circ f)(h_{k_2} \circ g))(m).
$$

The formula for $h_k \circ (a[m])$ is immediate from the definition. It follows that, in $R[M][[z]],$

$$
\sum_{k\geq 0} (h_k \circ (a[m])) z^k = \sum_{k\geq 0} (h_k \circ a)[m^k] z^k.
$$

Thus, applying Eq. $(2.1.0.2)$ to the element a in R and substituting $[m]z$ for z, we find the multiplicative inverse of this series is $\sum_{k\geq 0}(-1)^k(e_k\circ a)[m^k]z^k$. But since $R[M]$ is a pre- λ ring, by the same logic its multiplicative inverse is also $\sum_{k\geq 0} (-1)^k (e_k \circ (a[m])) z^k$. Thus we obtain the claimed formula for $e_k \circ (a[m])$.

Now assume R is torsion free. To obtain the description of the plethystic action of p_k , note that the given formula is evidently an additive homomorphism on $R[M]$, so to obtain it on $R[M]$ it suffices to verify the formula $p_k \circ a([m]) = (p_k \circ a)[m^k]$. This also gives the formula in the locally finite case by continuity, since $R[M]$ is dense in R[[M]]. To verify $p_k \circ a([m]) = (p_k \circ a)[m^k]$, we argue similarly to the above but invoking Eq. $(2.1.0.3)$ instead of Eq. $(2.1.0.2)$.

Example 2.2.5.

(1) Taking the monoid $M_n = \mathbb{Z}_{\geq 0}^n$ and and equipping $\mathbb Z$ with the pre- λ structure of Example [2.2.2,](#page-13-4) from Lemma [2.2.4](#page-14-1) we obtain natural pre- λ structures on $\mathbb{Z}[M_n] = \mathbb{Z}[t_1,\ldots,t_n]$ and $\mathbb{Z}[[M_n]] = \mathbb{Z}[[t_1,\ldots,t_n]]$. Note that, since $e_k \circ 1 = 0$ for all $k > 1$, for any monomial \underline{t}^j we find

$$
e_1 \circ \underline{t}^{\underline{j}} = \underline{t}^{\underline{j}}
$$
 and $e_k \circ \underline{t}^{\underline{j}} = 0 \forall k > 2$.

(2) The invariants Λ_n and Λ_n^{\wedge} for the action of Σ_n on $\mathbb{Z}[t_1,\ldots,t_n]$ and $\mathbb{Z}[[t_1,\ldots,t_n]]$ are preserved by the pre- λ structure, so have natural pre- λ structures.

We also have the following evident functorialities for the construction.

Lemma 2.2.6. Suppose $\varphi : R \to S$ is a map of pre- λ rings and $f : M \to N$ is a map of monoids. Then there is a unique map of pre- λ rings $R[M] \rightarrow S[N]$ such that $r[m] \mapsto \varphi(r)[f(m)]$. If M and N are both locally finite and f has finite fibers, then it extends uniquely to a continuous map $R[[M]] \to S[[N]]$.

2.3. Symmetric power series. Building on Example [2.2.5,](#page-15-0) we may view

$$
R[t_1, t_2, \ldots] = R[t_i, i \in \mathbb{N}]
$$

as $R[M]$ for $M_N = \bigoplus_N \mathbb{Z}_{\geq 0}$ the free commutative monoid on \mathbb{N} , and we write

 $R[[\underline{t}_{\mathbb{N}}]] := R[[M_{\mathbb{N}}]] \cong \lim_n R[[t_1,\dots,t_n]].$

We write Λ_R^{\wedge} for the subset of functions in $R[[t_{\mathbb{N}}]]$ that are invariant under all automorphisms induced by automorphism of \mathbb{N} , and Λ_R for the subset of Λ_R^{\wedge} consisting of those functions that are zero on $\mathbb{Z}_{\geq n}^{\mathbb{N}}$ for some $n \gg 0$. In particular, for $R = \mathbb{Z}$ we recover the symmetric functions $\overline{\Lambda} = \Lambda_{\mathbb{Z}}$ and its natural completion $\Lambda_{\mathbb{Z}}^{\wedge}$ for the filtration by degree. More generally, Λ_R is naturally identified with $\Lambda \otimes_{\mathbb{Z}} R$, and Λ_R^{\wedge} is identified with the completion of Λ_R for the filtration by degree.

If R is a pre- λ ring, then Λ_R and Λ_R^{\wedge} are evidently preserved by the pre- λ opera-tions of Definition [2.2.3.](#page-14-0) Taking $R = \mathbb{Z}$ with the pre- λ structure of Example [2.2.2,](#page-13-4) we obtain, in particular, a natural pre- λ ring structure on Λ itself.

Example 2.3.1. For $f \in \Lambda$, from Lemma [2.2.4](#page-14-1) we find that $p_i \circ f = f(t_1^i, t_2^i, \ldots)$. Thus, comparing with [\[22,](#page-60-15) Chapter 8, eq. (8.4) on p. 135], we have reconstructed the usual plethystic action of Λ on itself.

2.4. Witt vectors and λ -rings. We now recall some standard results about Witt vectors and their relation with pre- λ rings as in [\[15,](#page-60-17) §16]. There is a natural coalgebra structure on Λ, where the addititive counit $\Lambda \to \mathbb{Z}$ sends h_i to 0, $i \geq 0$, the multiplicative counit $\Lambda \to \mathbb{Z}$ sends h_i to 1, $i \geq 0$, the coaddition $\Lambda \to \Lambda \otimes_{\mathbb{Z}} \Lambda$ is uniquely determined by

$$
p_i \mapsto p_i \otimes 1 + 1 \otimes p_i,
$$

or, equivalently,

(2.4.0.1)
$$
h_k \mapsto \sum_{i+j=k} h_i \otimes h_k.
$$

The comultiplication $\Lambda \to \Lambda \otimes_{\mathbb{Z}} \Lambda$ is uniquely determined by

$$
p_i\mapsto p_i\otimes p_i.
$$

For R a ring, $W(R) := \text{Hom}_{\text{Ring}}(\Lambda, R)$ is thus a ring. It is moreover a pre- λ ring for the plethystic action of Λ by $(f \circ w)(g) = w(g \circ f)$. If R is a pre- λ ring, there is a canonical homomorphism of abelian groups

$$
R \mapsto W(R), r \mapsto (f \mapsto f \circ r).
$$

Conversely, to give a pre- λ ring structure on R it is equivalent to give such a homomomorphism (by comparing the defining property with Eq. $(2.4.0.1)$).

Definition 2.4.1. A pre- λ ring R is a λ -ring if the canonical map $R \to W(R)$ defining the pre $-\lambda$ structure is a morphism of pre- λ rings.

Example 2.4.2. For any ring R, $W(R)$ is a λ -ring. It is the cofree λ -ring on R, i.e., for any λ -ring S, Hom_{λ -Ring} (S, R) = Hom_{Ring} (S, R) via evaluation at h_1 .

It follows, in particular, that if R is a λ -ring then $f \circ (g \circ r) = (f \circ g) \circ r$ and the Adams operations $p_k \circ$ are λ -ring homomorphisms (note that a λ -ring homomoprhism is the same thing as a pre- λ ring homomorphism of λ -rings). Using the definition of the algebra structure and pletyhysm on $W(R)$, we have:

Proposition 2.4.3. A ($\mathbb{Z}-$)torsion free pre- λ ring R is a λ -ring if and only if

- (1) For all $k, j, p_k \circ (p_j \circ r) = p_{jk} \circ r$
- (2) For all k, $p_k \circ : R \to R$ is a ring homomorphism.

In particular, using this criterion we obtain

Lemma 2.4.4. If R is a torsion-free λ -ring and M is a commutative monoid, then $R[M]$ is a λ -ring. If M is locally fnite, then $R[[M]]$ is a λ -ring. In particular, if R is a λ -ring then Λ_R and Λ_R^{\wedge} are both λ -rings.

Proof. For $R[M]$ and $R[[M]]$, the criteria of Proposition [2.4.3](#page-16-3) follow immediately from Lemma [2.2.4.](#page-14-1) The result then follows for Λ_R and Λ_R^{\wedge} since any sub-pre- λ ring of a λ -ring is a λ -ring.

2.5. Plethystic exponential and logarithm. Let R be a pre- λ ring. For

$$
A = R[[\underline{t}_{\mathbb{N}}]]
$$
 or Λ_R^{\wedge} ,

let \mathfrak{m}_A be the ideal of elements with constant term 0. For $f \in \mathfrak{m}_A$, we define

$$
Exp_{\sigma}(f) = \sum_{k \ge 0} h_k \circ f,
$$

where the sum evidently converges. We have

Lemma 2.5.1. Exp_{σ} is a continuous group isomorphism from m_A , equipped with addition, to $1 + \mathfrak{m}_A$, equipped with multiplication.

Proof. We first treat the case $A = R[[t_{N}]]$. That it is a homomorphism follows from $h_k(x+y) = \sum_{i+j=k} h_i(x)h_j(y)$. We thus compute

$$
Exp_{\sigma}(r_{\underline{j}}\underline{t}^{\underline{j}}) = \prod_{\underline{j}} \sum_{k \ge 0} h_k(r_{\underline{j}}) \underline{t}^{k\underline{j}}
$$

Since $h_1(r_j) = r_j$, any element of $1 + \mathfrak{m}_A$ has a unique expansion of the form on the right, so we deduce that Exp_{σ} is a bijection. It commutes with the action of Aut(N), thus we also obtain the case of Λ_R^{\wedge} . \overline{R} .

Example 2.5.2. Suppose $X \in R$. Then we compute

$$
\operatorname{Exp}_{\sigma}(Xh_1) = \operatorname{Exp}_{\sigma}(Xt_1 + Xt_2 + \ldots) = \prod_{i \ge 1} \operatorname{Exp}_{\sigma}(Xt_i) = \prod_{i \ge 1} \left(\sum_{k \ge 0} h_k(X)t_i^k \right)
$$

where in the first equality we use Lemma [2.5.1](#page-17-2) and in the third equality we use Lemma [2.2.4](#page-14-1) to obtain $h_k(Xt_i) = h_k(X)t_i^k$. Expanding the product, we obtain

(2.5.2.1)
$$
\operatorname{Exp}_{\sigma}(Xh_1) = \sum_{\tau} (h_{\tau} \circ X) m_{\tau}.
$$

From Eq. $(2.1.0.2)$, we obtain

Lemma 2.5.3.

$$
Exp_{\sigma}(-x) = \sum_{i=0}^{\infty} (-1)^{i} e_i \circ x.
$$

From Eq. $(2.1.0.3)$, we obtain

Lemma 2.5.4. If R is torsion-free then

$$
Exp_{\sigma}(x) = \prod_{i \ge 1} exp\left(\frac{p_i \circ x}{i}\right) \text{ where } exp(t) := \sum_{i \ge 0} \frac{t^i}{i!}.
$$

Definition 2.5.5. We write $\text{Log}_{\sigma} : 1 + \mathfrak{m}_A \to \mathfrak{m}_A$ for the inverse map to Exp_{σ} , which exists by Lemma [2.5.1.](#page-17-2)

Lemma 2.5.6. If R is a λ -ring, then

$$
\operatorname{Log}_{\sigma}(x) = \sum \ell_i \circ (x - 1) \text{ for } \ell_i = \frac{-1}{i} \sum_{d \mid i} \mu(d) (-p_d)^{i/d} \in \Lambda
$$

Proof. This follows from a plethystic inversion formula of Stanley [\[24,](#page-60-18) Exercise 7.88e on p.479] (see also [\[13,](#page-60-19) Prop 8.6] and [\[2,](#page-60-4) Prop 2.3.7]).

Remark 2.5.7. Everything in the present section works for symmetric power series rings in arbitrary variables over pre- λ rings, but we only need the countable case. In fact, the plethystic exponential and logarithm can be treated much more generally for filtered pre- λ rings, extending the case of filtered λ -rings in [\[2\]](#page-60-4), or even for more general Hausdorff topological pre- λ rings as will be discussed in [\[6\]](#page-60-14).

2.6. Powers. Using the exponential we can also define powers:

Definition 2.6.1. For $f \in 1 + \mathfrak{m}_A$ and $N \in A$,

 $f^A := \text{Exp}_{\sigma}(N\text{Log}_{\sigma}(f)).$

Lemma 2.6.2. $f^{N_1+N_2} = f^N f^N$ and $(fg)^N = f^N g^N$.

Proof. Immediate from Lemma [2.5.1.](#page-17-2) □

This definition recovers the power structures of $[14]$ (cf. also $[17, §2.6]$) — indeed, the compatibility follows from continuity of Exp_{σ} and Log_{σ} and the computation

$$
\left(\frac{1}{1-\underline{t}^{\underline{j}}}\right)^N = \operatorname{Exp}_{\sigma}(N\underline{t}^{\underline{j}}) = \sum_{k\geq 0} (h_k \circ N) \underline{t}^{\underline{j}k},
$$

which is immediate from the formulas in Lemma [2.2.4.](#page-14-1)

Example 2.6.3.

$$
(1 + h_1 + h_2 + \ldots)^N = (\text{Exp}_{\sigma}(h_1))^N = \text{Exp}_{\sigma}(Nh_1).
$$

We note that, if $f \in \Lambda^{\wedge}$, then the assignment $N \mapsto f^N = \text{Exp}_{\sigma}(X \text{Log}_{\sigma}(f))$ is functorial in maps of pre- λ rings, in the sense that if $\varphi: R_1 \to R_2$ is a map of pre- λ rings, then $f^{\varphi(N)}$ is obtained from f^N by applying the continuous extension of $\varphi \otimes \text{Id}$ from $\Lambda_{R_1}^{\wedge}$ to $\Lambda_{R_2}^{\wedge}$. We claim that it follows that there are unique elements $a_{\tau,f} \in \Lambda$ such that, for any pre- λ ring R and $N \in R$

$$
f^N = \sum (a_{\tau,f} \circ N) m_{\tau}.
$$

Indeed, since Λ is the free pre- λ ring on one generator $(h_1 = e_1 = p_1)$, taking R to be Λ itself and writing its variables as s_1, s_2, \ldots , we can take the unique expansion in Λ^\wedge_Λ with respect to the m_τ

$$
f^{h_1(s_1,s_2,\ldots)} = \sum a_{\tau,f}(\underline{s})m_{\tau}.
$$

Since $a_{\tau,f}(\underline{s}) = a_{\tau,f} \circ h_1(\underline{s})$, the functoriality of the construction shows these are the desired elements.

Example 2.6.4. Combining Example [2.6.3](#page-18-2) and Example [2.5.2,](#page-17-3) we find

$$
a_{\tau,(1+h_1+h_2+\ldots)} = h_{\tau}.
$$

Example 2.6.5. We let $c_{\tau} := a_{\tau,(1+h_1)}$, so that

$$
(1+h_1)^N = \sum_{\tau} (c_{\tau} \circ N) m_{\tau}.
$$

This recovers the configuration symmetric functions denoted c_{τ} in [\[17,](#page-60-6) Lemma 3.5] an surrounding discussion]. We can express the h_{τ} in terms of the c_{τ} as follows: we consider $\Lambda[[\underline{v}_N]]$, but where we name our countable family of variables v_{ij} , $1 \le i, j$, and where we write s_i for the variables in Λ . Then, if we specialize^{[2](#page-19-3)}

$$
(1+\sum_{ij}v_{ij})^{h_1(\underline{s})}
$$

by substituting t_i^j for v_{ij} , then we obtain

$$
(1 + h_1 + h_2 + ...)^{h_1(\underline{s})} = \text{Exp}_{\sigma}(h_1(\underline{s})h_1(\underline{t})) = \sum h_{\tau}(\underline{s})m_{\tau}
$$

Expanding, one obtains expressions for h_{τ} as Z-linear combination of the c_{τ} . In particular, one finds $h_{\tau} = c_{\tau} + \dots$ where the other partitions τ' appearing with non-trivial coefficients have $|\tau'| < |\tau|$, thus the matrix can be inverted and we find that the c_{τ} are also a basis for Λ (as explained from an essentially equivalent geometric perspective in [\[17\]](#page-60-6)). We note that this computation is also closely related to constructions used in the theory of motivic Euler products [\[5\]](#page-60-20).

2.7. The Hall inner product. The Hall inner product on Λ is the inner product \langle , \rangle determined by

$$
(2.7.0.1) \t\t \langle m_{\tau}, h_{\mu} \rangle = \delta_{\tau\mu}.
$$

For s_{τ} the Schur symmetric functions, it satisfies

$$
\langle s_{\tau}, s_{\mu} \rangle = \delta_{\tau\mu}.
$$

On power sum symmmetric functions it satisfies

(2.7.0.2)
$$
\langle p_{\tau}, p_{\mu} \rangle = z_{\tau} \delta_{\tau \mu} \text{ where } z_{\tau} = \prod_{i} \tau(i)! i^{\tau(i)},
$$

recalling that $\tau(i)$ denotes the multiplicity of i in τ .

For R any ring, the Hall inner product induces an isomorphism of R -modules

(2.7.0.3)
$$
\Lambda_R^{\wedge} \xrightarrow{f \mapsto \langle f, \bullet \rangle} \text{Hom}_{\mathbb{Z}}(\Lambda, R).
$$

2.8. The involution ω . We write ω for the standard involution on Λ , so that

$$
e_i \stackrel{\omega}{\leftrightarrow} h_i, \; p_i \stackrel{\omega}{\leftrightarrow} (-1)^{i-1} p_i, \; s_{\tau} \stackrel{\omega}{\leftrightarrow} s_{\tau'}.
$$

For any ring R, it extends to an involution of Λ_R^{\wedge} .

We say $f \in \Lambda_R^{\wedge}$ is even if it is a formal sum of monomials of even degree. It is immediate that the even elements are a subring and, if R is a pre- λ ring, a pre- λ subring.

Lemma 2.8.1. Let R be a torsion-free pre- λ -ring. If $f \in \mathfrak{m}_{\Lambda_R^{\wedge}}$ is even, then

$$
\operatorname{Exp}_{\sigma}(\omega(f)) = \omega(\operatorname{Exp}_{\sigma}(f)) \text{ and } \operatorname{Log}_{\sigma}(\omega(1+f)) = \omega(\operatorname{Log}_{\sigma}(1+f)).
$$

²One can make this specialization using Lemma [2.2.6](#page-15-1) because it is induced by the map of monoids $\bigoplus_{i,j\geq 1}\mathbb{Z}_{\geq 0}\to\bigoplus_{i\geq 1}\mathbb{Z}_{\geq 0}$ sending e_{ij} to je_i , for e_{ij} (resp. e_i) the vector that is 1 in the ijth (resp. ith) component and zero in all other component which has finite fibers.

Proof. It suffices to treat Exp_{σ} , since Log_{σ} is its inverse.

We can write $f = \sum_{|\lambda| \text{ even}} a_{\lambda} p_{\lambda}$ where each $a_{\lambda} \in A_{\mathbb{Q}}$. Then

$$
\omega(\text{Exp}_{\sigma}(f)) = \prod \omega(\text{Exp}_{\sigma}(a_{\lambda}p_{\lambda})) \text{ and } \text{Exp}_{\sigma}(\omega(f)) = \prod \text{Exp}_{\sigma}(a_{\lambda}\omega(p_{\lambda}))
$$

so it suffices to treat the case $f = a_{\lambda} p_{\lambda}$.

By Lemma [2.5.4,](#page-17-1)

$$
\operatorname{Exp}(a_{\lambda}p_{\lambda}) = \prod_{i \geq 1} \sum_{k \geq 0} \frac{p_i(a)^k p_{i\lambda}^k}{i^k k!} \text{ and } \operatorname{Exp}(\omega(a_{\lambda}p_{\lambda})) = \prod_{i \geq 1} \sum_{k \geq 0} \frac{p_i(a)^k (p_i \circ \omega(p_{\lambda}))^k}{i^k k!}.
$$

Thus it suffices to show that, for each i ,

$$
\omega(p_{i\lambda})=p_i\circ\omega(p_\lambda).
$$

Expanding, we find this is equivalent to showing

$$
i|\lambda| - ||\lambda|| \equiv |\lambda| - ||\lambda|| \mod 2.
$$

But the difference is $(i - 1)|\lambda|$, which is even since $|\lambda|$ is even by assumption. \square

3. $(PRE-) \lambda$ -PROBABILITY

In this section we develop the basic theory of probability in λ -rings. In fact, we work everywhere with pre- λ rings: although all of the ring of random variables in the examples of interest in the present work are λ -rings, in [\[8\]](#page-60-8) we will want to work also with rings of random variables given by relative Grothendieck rings of varieties, and for these the natural pre- λ ring structure is not known to be a λ -ring structure. Working in the added generality of pre- λ rings costs us nothing, as all of the basic theory depends only on the properties of a pre- λ ring.

After giving basic defintions in Section [3.1,](#page-21-0) we arrive at the meat of the theory in our study of moment generating functions in Section [3.2.](#page-21-1) We establish there the basic properties of the σ -moment generating function, and also explain how to construct more general moment generating functions using the powers of Section [2.6,](#page-18-0) including a falling (or factorial) moment generating function that plays the same role as the classical falling moment generating function. We note that, for random variables of a combinatorial nature, the falling moment generating function often takes a simpler form (see Remark [3.2.5\)](#page-22-1).

In Section [3.3,](#page-22-0) we discuss binomial and Poission distributions and their σ moment and falling moment generating functions. It seems remarkable to us that there is a single definition in the λ -setting that matches so closely with the simple forms of the moment generating functions of binomial and Poission distributions in classical probability theory. We note that these λ -analogs of binomial distributions are ubiqitous in computations in arithmetic statistics because they arise from summing identically distributed independent Bernoulli random variables (just as in the case of classical binomial random variables, but see Remark [3.3.3\)](#page-23-0).

Finally, in Section [3.4,](#page-24-0) we make our notions of convergence of sequences of random variables precise for use in our main results.

Remark 3.0.1. A shadow of the theory described here was developed already in [\[17,](#page-60-6) §4]. The systematic use of the (pre-) λ ring structure and the σ -moment generating function makes the version developed here much more robust.

22 $\,$ $\,$ SEAN HOWE $\,$

3.1. Basic definitions.

Definition 3.1.1.

- (1) A (pre-) λ -probability space is a pair (R, \mathbb{E}) where R is a $(\text{pre}-\lambda)$ -ring (the random variables) and $\mathbb E$ is a $\mathbb Z$ -linear functional on R (the expectation functional) valued in another ring C and satisfying $\mathbb{E}[1_R] = 1_C$.
- (2) For (R, \mathbb{E}) a (pre-) λ -probability space and $X \in R$ a random variable, the $Λ$ -distribution of X is the C-valued functional μ_X on Λ sending $f \in Λ$ to $\mathbb{E}[f \circ X].$

Definition 3.1.2. For (R, \mathbb{E}) a (pre-) λ -probability space and $X_1, X_2, \ldots, X_n \in R$ random variables,

(1) the *joint distribution* of X_1, X_2, \ldots, X_n is the the element μ_{X_1,\ldots,X_n} of $\text{Hom}_{\mathbb{Z}}(\Lambda^{\otimes n}, C)$ determined by

$$
f_1 \otimes \ldots \otimes f_n \mapsto \mathbb{E}[(f_1 \circ X_1)(f_2 \circ X_2) \ldots (f_n \circ X_n)],
$$
 and

(2) we say $X_1, X_2, \ldots X_n$ are independent if μ_{X_1,\ldots,X_n} is equal to the element of Hom_Z $(\Lambda^{\otimes n}, C)$ determined by

$$
f_1 \otimes \ldots \otimes f_n \mapsto \mathbb{E}[f_1 \circ X_1] \cdot \mathbb{E}[f_2 \circ X_2] \cdot \ldots \cdot \mathbb{E}[f_n \circ X_n].
$$

More generally, if $(X_i)_{i\in I}$ is a possibly infinite family of random variables, we say $(X_i)_{i\in I}$ is independent if any finite subset is independent.

3.2. Moment generating functions. The main tool we will use for studying Λ-distributions is the σ-moment generating function.

Definition 3.2.1. For $(R, \mathbb{E}: R \to C)$ a (pre-) λ -probability space, the σ -moment generating function of $X \in R$ is

$$
\mathbb{E}[\mathrm{Exp}_{\sigma}(Xh_1)] \in \Lambda_C^{\wedge}.
$$

where here Xh_1 is viewed as an element of Λ_R^{\wedge} and $\mathbb E$ is applied coefficient-wise, i.e. it is the continuous extension of Id $\otimes \mathbb{E}$ on $\Lambda_R = \Lambda \otimes R$ to Λ_R^{\wedge} .

By Example [2.5.2,](#page-17-3) we find

(3.2.1.1)
$$
\mathbb{E}[\text{Exp}_{\sigma}(Xh_1)] = \sum_{\tau} \mathbb{E}[h_{\tau} \circ X]m_{\tau}.
$$

Because the m_{τ} and the h_{τ} are both Z-bases of Λ , it follows that the σ -moment generating function of X determines its Λ -distribution and vice-versa.

The Λ -distribution can be obtained from the σ -moment generating function by the Hall inner product via the isomorphism Eq. [\(2.7.0.3\)](#page-19-4):

Lemma 3.2.2. For any $f \in \Lambda$,

$$
\mathbb{E}[f \circ X] = \langle \mathbb{E}[\text{Exp}_{\sigma}(Xh_1)], f \rangle.
$$

Proof. It suffices to verify the two sides agree on h_{γ} for all partitions γ . We obtain this by invoking Eq. $(3.2.1.1)$ and Eq. $(2.7.0.1)$:

$$
\langle \mathbb{E}[\operatorname{Exp}_{\sigma}(Xh_1)], h_{\gamma} \rangle = \langle \sum_{\tau} \mathbb{E}[h_{\tau} \circ X]m_{\tau}, h_{\gamma} \rangle = \sum_{\tau} \langle \mathbb{E}[h_{\tau} \circ X]m_{\tau}, h_{\gamma} \rangle = \mathbb{E}[h_{\gamma} \circ X].
$$

We can also construct other moment generating functions: for $f \in 1+\mathfrak{m}_{\Lambda}$, with notation as in Section [2.6](#page-18-0) we obtain a moment generating function

(3.2.2.1)
$$
\mathbb{E}[f^X] = \sum_{\tau} (a_{\tau,f} \circ X) m_{\tau}.
$$

Example 3.2.3.

- (1) If $f = 1 + h_1 + h_2 + ... = \text{Exp}_{\sigma}(h_1)$, then, by Example [2.6.3,](#page-18-2) $f^X = \text{Exp}_{\sigma}(X)$ and we recover the σ -moment generating function.
- (2) If $f = (1 + h_1)$, then, with notation as in Example [2.6.5,](#page-18-1)

$$
\mathbb{E}[(1+h_1)^X] = \sum_{\tau} \mathbb{E}[c_{\tau} \circ X] m_{\tau}.
$$

We refer to this as the *falling moment generating function of* X . Since, as explained in Example [2.6.5,](#page-18-1) the c_{τ} are a Z-basis of Λ , we find that to give a Λ-distribution it is also equivalent to give its falling moment generating function.

Lemma 3.2.4. If random variables X_1 and X_2 in R are independent then, for any $f \in 1 + \mathfrak{m}_{\Lambda}$ ^{\wedge},

$$
\mathbb{E}\left[f^{(X_1+X_2)}\right] = \mathbb{E}[f^{X_1}]\mathbb{E}[f^{X_2}].
$$

In particular, taking $f = 1 + h_1 + h_2 + \ldots = \operatorname{Exp}_{\sigma}(h_1)$,

$$
[\operatorname{Exp}_{\sigma}((X_1 + X_2)h_1)] = \mathbb{E}[\operatorname{Exp}_{\sigma}(Xh_1)]\mathbb{E}[\operatorname{Exp}_{\sigma}(Xh_1)].
$$

Proof. By Lemma [2.6.2,](#page-18-3)

 $\mathbb E$

$$
\mathbb{E}\left[f^{(X_1+X_2)}\right] = \mathbb{E}[f^{X_1}f^{X_2}].
$$

Expanding as in Eq. [\(3.2.2.1\)](#page-22-2) and applying the definition of independence, we find that we can commute the expectation with the product. \square

Remark 3.2.5. In practice, it is simpler to work with the falling moment generating function for random variables of a combinatorial nature (e.g. valued in finite sets or varieties over a fixed variety), and with the σ -moment generating function for random variables of a linear nature (e.g. valued in multisets of complex numbers or sheaves on a fixed variety). We will default to working with the σ -moment generating function when we expect to mix the two types of random variable: in the combinatorial setting the σ -moment generating function is typically only mildy more complicated (see, e.g., the difference in Definition [3.3.2\)](#page-23-1), whereas falling moment generating functions tend to be completely intractable in the linear setting.

3.3. Some Λ-distributions. We recall that, in classical probability theory, a binomial random variable X with parameters p and N is the sum of N independent Bernoulli random variables, each of which takes values 1 and 0 with probability p and $(1 - p)$, and has falling moment and moment generating functions

$$
\mathbb{E}[(1+t)^{X}] = (1+pt)^{N} \text{ and } \mathbb{E}[e^{X}] = (1+p(e^{t}-1))^{N}.
$$

We also recall that, in classical probability theory, a Poisson random variable X of mean λ has falling moment and moment generating functions

$$
\mathbb{E}[(1+t)^X] = e^{\lambda t} \text{ and } \mathbb{E}[e^X] = e^{\lambda(e^t - 1)}.
$$

The following shows that there are perfect Λ -analogs of these distributions.

Lemma 3.3.1. Let R be a pre- λ ring.

(1) Let $p, N \in R$. The unique Λ -distribution with falling moment generating function

$$
\mathbb{E}[(1+h_1)^X] = (1+ph_1)^N
$$

has σ -moment generating function

$$
\mathbb{E}[\exp_{\sigma}(Xh_1)]^{\mathbb{N}} = (1 + p(\exp_{\sigma}(h_1) - 1))^N = (1 + p(h_1 + h_2 + ...))^N.
$$

(2) Let $\lambda \in R$. The unique Λ -distribution with falling moment generating function

$$
\mathbf{E}[(1+h_1)^X]^n = \operatorname{Exp}_{\sigma}(\lambda h_1)
$$

has σ -moment generating function

$$
\mathbb{E}[\operatorname{Exp}_{\sigma}(Xh_1)]^{\sigma} = \operatorname{Exp}_{\sigma}(\lambda(\operatorname{Exp}_{\sigma}(h_1) - 1)) = \operatorname{Exp}_{\sigma}(\lambda(h_1 + h_2 + \ldots)).
$$

Here we put quotes to indicate that we have not actually fixed a random variable X with the given Λ -distribution.

Proof. Both follow by making the same substitution as in Example [2.6.5.](#page-18-1) \Box

Definition 3.3.2. The following notions are well-defined by Lemma [3.3.1.](#page-22-3)

(1) For R a pre-λ-ring and $p, n \in R$, the (p, N) -binomial Λ -distribution is the unique Λ-distribution with falling moment generating function

$$
(1+ph_1)^N
$$

and σ -moment generating function

$$
1 + p \left(\text{Exp}_{\sigma}(h_1) - 1 \right))^N = (1 + p(h_1 + h_2 + \ldots))^{N}.
$$

(2) For R a pre-λ-ring and $\lambda \in R$, the λ-Poisson A-distribution is the unique Λ-distribution with falling moment generating function

$$
\operatorname{Exp}_{\sigma}(\lambda h_1)
$$

and σ -moment generating function

$$
Exp_{\sigma}(\lambda(Exp_{\sigma}(h_1)-1)) = Exp_{\sigma}(\lambda(h_1 + h_2 + ...)).
$$

Remark 3.3.3. In practice, the random variables we will encounter that have binomial Λ -distributions can be interpreted, at least morally, as sums of N independent Bernoulli random variables. When N is a positive integer, this makes sense already using Lemma [3.2.4.](#page-22-4) When R is the decategorification of a combinatorial category such as algebraic varieties or admissible Z-sets (see Section [5\)](#page-33-0), this can be made precise also when N is a more general effective class, as we will explain in [\[8\]](#page-60-8).

We note also that we can view our Poisson Λ -distributions with parameter λ as limits of binomial distributions with parameters p and N such that $pN \to \lambda$. This will also be explained precisely and treated in examples in [\[8\]](#page-60-8).

3.4. Convergence.

Definition 3.4.1. When $\mathbb E$ is valued in a Hausdorff topological ring, we say a sequence of tuples $(X_{1,d},...,X_{n,d})$ of n random variables indexed by natural numbers d converges in distribution to a joint Λ -distribution μ if, for each $f \in \Lambda^{\otimes n}$,

$$
\lim_{d \to \infty} \mu_{X_{1,d}, X_{2,d}, ..., X_{n,d}}(f) = \mu(f)
$$

It suffices to consider just f of the form $f = f_1 \otimes f_2 \otimes \ldots \otimes f_n$, for which this says

$$
\lim_{d \to \infty} \mathbb{E}[f_1 \circ X_{1,d} \cdot f_2 \circ X_{2,d} \dots f_n \circ X_{n,d}] = \mu(f).
$$

Similarly, it suffices to consider just the f_i as they range over any fixed basis of Λ .

Definition 3.4.2. When $\mathbb E$ is valued in a Hausdorff topological ring, we say a sequence indexed by natural numbers d of tuples $(X_{i,d},\ldots,X_{i,d})_{i \in [n]}$ of random variables is asymptotically independent if there exists for each $i \in I$ a Λ -distribution $\mu_{i,\infty}$ such that, for any finite collection of distinct indices $i_1, \ldots, i_n \in I, (X_{i_1,d}, \ldots, X_{i_n,d})$ converges in distribution to the joint Λ -distribution $\mu_{i_1} \otimes \mu_{i_2} \otimes \ldots \otimes \mu_{i_n}$.

Arguing as in Lemma [3.2.4,](#page-22-4) we obtain

Lemma 3.4.3. If $(X_{1,d},...,X_{n,d})$ is a sequence of asymptotically independent random variables, then

$$
\lim_{d \to \infty} \mathbb{E}[\exp_{\sigma}((X_{1,d} + \dots X_{n,d})h_1)] = \prod_{i=1}^n \lim_{d \to \infty} \mathbb{E}[\exp_{\sigma}(X_{i,d})h_1]
$$

where the limits in Λ_C^{\wedge} can be taken individually in C for each coefficient of m_{τ} .

4. Random matrices

In this section we prove our main result on random matrices, Theorem [A](#page-3-0) (see Theorem [4.2.1\)](#page-25-1), and give some complements. First in Section [4.1](#page-24-2) we situate the result more squarely within the theory of λ -probability that was developed in Section [3.](#page-20-0) Then in Section [4.2](#page-25-0) we prove the main result, admitting a computation of invariants described in Section [4.3.](#page-27-0) In Section [4.4](#page-29-0) we deduce as a corollary (Corollary [4.4.1\)](#page-29-1) the Diaconis-Shahshahani theorem on traces of powers of random matrices in classical groups. Finaly, in Section [4.5](#page-32-0) we explain the relation between the symmetric group case of Theorem A/T A/T heorem [4.2.1](#page-25-1) and cycle-counting.

4.1. The probability space. Let G be a compact Hausdorff topological group. The Grothendieck ring $K_0(\text{Rep}_{\mathbb{C}}G)$ of complex representations of G admits a natural structure as a λ -ring such that $h_k \circ [V] = [\text{Sym}^k V]$. For V a representation of G we write χ_V for its character $\chi_V(q) = \text{Tr}(q \text{ acting on V})$, and we write E for the expectation functional $K_0(\text{Rep} G) \to \mathbb{Z}$ which sends [V] to

$$
\int_G \chi_V dg
$$
 for dg the Haar measure on G .

These integrals are a priori complex numbers, but here they evaluate to integers since, by the usual orthogonality relations, the integral of χ_V is equal to the multiplicity of the trivial representation in V. In other words, viewing $K_0(\text{Rep}_{\mathbb{C}}G)$ as the free Z-module on the isomorphism classes of irreducible representations of G , $\mathbb E$ is the unique functional sending the trivial representation to 1 and any other irreducible representation to 0. We will consider the λ -probability space $(K_0(\text{Rep}_{\mathbb{C}}), \mathbb{E})$. 26 SEAN HOWE

Remark 4.1.1. The assignment $[V] \mapsto \chi_V$ extends to an embedding of $K_0(\text{Rep}G)$ into the ring of continuous class functions on G. This subring is thus equipped with a natural λ -ring structure, so we can view our ring of random variables as a subring of the ring of $\mathbb{C}\text{-}$ valued continuous functions on G. In particular, the classical $\mathbb{C}\text{-}$ valued expectation functional given by integration against Haar measure restricts to our Z-valued measure under this embedding.

Remark 4.1.2. Studying $K_0(\text{Rep}_{\mathbb{C}}G)$ with its Z-valued measure will suffice for the results we wish to prove here. To bring more general problems in random matrix theory into the realm of λ -probability, it is natural to study the the λ -ring of continuous $W(\mathbb{C}) \cong \prod_{i=1}^{\infty} \mathbb{C}$ -valued functions on G (where $W(\mathbb{C})$ is equipped with the product topology). The expectation on this ring is valued in $W(\mathbb{C})$ and is given by integration in each term, and the λ -ring structure is the pointwise structure $(f \circ F)(g) = f \circ (F(g))$. Builiding on Remark [4.1.1,](#page-24-3) we claim the ring of random variables $K_0(\text{Rep}G)$ embeds naturally into $Cont(G, W(\mathbb{C}))$ compatibly with this λ -ring structure and with the expectation functionals (under the unique map $\mathbb{Z} \to W(\mathbb{C})$. Indeed, we can first view [V] as the function sending $g \in G$ to its multiset of eigenvalues of g acting on V to obtain a function valued in $\mathbb{Z}[\mathbb{C}],$ and then compose with the usual embedding of $\mathbb{Z}[\mathbb{C}]$ into $W(\mathbb{C})$ (which sends [q] to (q, q^2, q^3, \ldots)). Thus [V] is mapped to the function which sends g to $(\chi_V(g^i))_i$. We will not use this perspective further in the current section, but rings of $W(\mathbb{C})$ -valued functions play an essential role in our study of L-functions starting in Section [5.](#page-33-0)

4.2. Random matrix moment generating functions. We now rephrase Theorem [A](#page-3-0) in the language of Section [4.1,](#page-24-2) then give a proof. In the following, we consider the multiplicity filtration on Λ^{\wedge}

$$
\Lambda_{\text{mult} > i}^{\wedge} = \{ \sum_{||\tau|| > i} a_{\tau} m_{\tau} \, | \, a_{\tau} \in \mathbb{Z} \}
$$

Similarly, we consider the disjoint multiplicity filtration on $(\Lambda \otimes \Lambda)^{\wedge}$,

$$
(\Lambda \otimes \Lambda)_{\text{mult}>i}^{\wedge} = \{ \sum_{||\tau||>i \text{ or } ||\overline{\tau}||>i} a_{\tau,\overline{\tau}} m_{\tau} m_{\overline{\tau}} \mid a_{\tau,\overline{\tau}} \in \mathbb{Z} \}
$$

where we recall $||\tau||$ is the number of distinct entries in the partition, so that, e.g, $||(3, 1)|| = 2$. We will also consider the degree filtration on Λ^{\wedge} , which can be written

$$
\Lambda_{\deg > i}^{\wedge} = \{ \sum_{|\tau| > i} a_{\tau} m_{\tau} \, | \, a_{\tau} \in \mathbb{Z} \}
$$

wher we recall $|\tau|$ is the sum of the partition so that, e.g., $|(3,1)|=4$.

Theorem 4.2.1.

(1) For $n \geq 1$, let V_n be the standard representation of $O(n)$ on \mathbb{C}^n . Then,

 $\mathbb{E}[\text{Exp}_{\sigma}([V_n]h_1)] \equiv \text{Exp}_{\sigma}(h_2) \mod \Lambda_{\text{mult}>n}^{\wedge}$

where we note

$$
Exp_{\sigma}(h_2) = \prod_{i \leq j} \frac{1}{1 - t_i t_j} \in \Lambda^{\wedge} \subseteq \mathbb{Z}[[\underline{t}_{\mathbb{N}}]].
$$

(2) For $n \geq 1$ even, let V_n be the standard representation of $\text{Sp}(n)$ on \mathbb{C}^n . Then,

$$
\mathbb{E}[\text{Exp}_{\sigma}([V_n]h_1)] = \text{Exp}_{\sigma}(e_2) \mod \Lambda_{\text{mult}>n}^{\wedge}
$$

where we note

$$
Exp_{\sigma}(h_2) = \prod_{i < j} \frac{1}{1 - t_i t_j} \in \Lambda^{\wedge} \subseteq \mathbb{Z}[[\underline{t}_{\mathbb{N}}]].
$$

(3) For $n \geq 1$, let V_n be the standard permutation representation of Σ_n on \mathbb{C}^n . Then,

$$
\mathbb{E}[\operatorname{Exp}_{\sigma}([V_n]h_1)] \equiv \operatorname{Exp}_{\sigma}(\operatorname{Exp}_{\sigma}(h_1) - 1) \mod \Lambda_{\deg > n}^{\wedge}
$$

where we note

$$
\operatorname{Exp}_{\sigma}(\operatorname{Exp}_{\sigma}(h_1) - 1) = \operatorname{Exp}_{\sigma}(h_1 + h_2 + \ldots)
$$

=
$$
\prod_{\substack{monomials \ m \text{ in } t_1, t_2, \ldots}} \frac{1}{1 - m} \in \Lambda^{\wedge} \subseteq \mathbb{Z}[[t_N]].
$$

(4) For $n \geq 1$, let V_n be the standard representation of $U(n)$ on \mathbb{C}^n . Then,

 $\mathbb{E}[\operatorname{Exp}_{\sigma}([V_n] \cdot h_1 + [V_n^*] \cdot \overline{h}_1)] \equiv \operatorname{Exp}_{\sigma}(h_1 \overline{h}_1) \mod \operatorname{Fil}^{n+1}(\Lambda \otimes_{\mathbb{Z}} \Lambda)_{\text{mult}>n}^{\wedge}$

where we note

$$
\operatorname{Exp}_{\sigma}(h_1\overline{h}_1)=\prod_{i,j}\frac{1}{1-t_i\overline{t}_j}\in (\Lambda\otimes_{\mathbb{Z}}\Lambda)^{\wedge}\subseteq\mathbb{Z}[[\underline{t}_{\mathbb{N}},\underline{\overline{t}}_{\mathbb{N}}]].
$$

Proof. We first discuss cases (1)-(3), writing G_n for the relevant group. For $\tau =$ $(\tau_1, \tau_2, \ldots, \tau_k)$, we write $\text{Sym}^{\tau}V_n = \text{Sym}^{\tau_1}V_n \otimes \ldots \otimes \text{Sym}^{\tau_k}V_n$. It follows from the definition of the λ -ring structure on $\text{Rep}_{\mathbb{C}}G$, where $h_k \circ [V] = [\text{Sym}^k V]$ and multiplication is given by tensor product, that

$$
h_{\tau} \circ [V_n] = (h_{\tau_1} \dots h_{\tau_k}) \circ [V_n] = (h_{\tau_1} \circ [V_n]) \dots (h_{\tau_k} \circ V_n) = [\text{Sym}^{\tau} V_n].
$$

Thus, using the expansion of Eq. [\(3.2.1.1\)](#page-21-2), we have

$$
\mathbb{E}[\text{Exp}_{\sigma}([V_n]h_1)] = \sum_{\tau} \mathbb{E}[h_{\tau} \circ [V_n]]m_{\tau}
$$

$$
= \sum_{\tau} \mathbb{E}[[\text{Sym}^{\tau}V_n]]m_{\tau}
$$

$$
= \sum_{\tau} \dim_{\mathbb{C}}(\text{Sym}^{\tau}V_n)^{G_n}m_{\tau}.
$$

where the final equality follows from the discussion surrounding the definition of E in Section [4.1.](#page-24-2) The result in these cases then follows from Lemma [4.3.3](#page-28-0) below after verifying that the given identities hold for the σ -exponentials in each case. This is clear by working in the non-symmetric power series ring $\mathbb{Z}[[t_{\mathrm{N}}]]$ containing Λ^{\wedge} of Section [2.3,](#page-16-0) where, e.g., in the orthogonal case we find

$$
Exp_{\sigma}(h_2) = Exp_{\sigma}(\sum_{i \leq j} t_i t_j) = \prod_{i \leq j} Exp_{\sigma}(t_i t_j) = \prod_{i \leq j} \frac{1}{1 - t_i t_j}.
$$

In case (4), writing $G_n = U(n)$ and expanding similarly we find

$$
\mathbb{E}[\operatorname{Exp}_{\sigma}([V_n] \cdot h_1 + [V_n^*] \cdot \overline{h}_1)] = \sum_{\tau, \overline{\tau}} \dim_{\mathbb{C}} (\operatorname{Sym}^{\tau} V_n \otimes \operatorname{Sym}^{\overline{\tau}} V_n^*)^{G_n} m_{\tau} \overline{m}_{\tau}
$$

and the result in this case then follows similarly from Lemma [4.3.3.](#page-28-0) \Box

 28 $\,$ SEAN HOWE $\,$

4.3. Computation of invariants. In the proof of Theorem [4.2.1](#page-25-1) we used a computation of the generating functions of invariants to be established as Lemma [4.3.3](#page-28-0) below. In the orthogonal, symplectic, and unitary cases, it will be a simple consequence of the first fundamental theorem of classical invariant theory, which we recall as Proposition [4.3.1](#page-27-1) below. In the symmetric case we make the relevant computation in Proposition [4.3.2.](#page-27-2)

Proposition 4.3.1 (First fundamental theorem of classical invariant theory).

(1) Let V be the standard representation of $O(n,\mathbb{C})$ on \mathbb{C}^n . Then, for any $m > 0$, the map of algebraic varieties

$$
V^{\oplus m}/O(n, \mathbb{C}) \to \prod_{1 \le i \le j \le m} \mathbb{A}^1
$$

$$
(v_1, \dots, v_m) \mapsto (\langle v_i, v_j \rangle)_{i \le j}
$$

is a closed immersion. Equivalently, the map

$$
\mathbb{C}[t_{ij}]_{1\leq i\leq j\leq m}\xrightarrow{t_{ij}\mapsto \langle v_i,v_j\rangle} \left(\text{Sym}^{\bullet}(V^{\oplus m})^*\right)^{O(n,\mathbb{C})}
$$

is surjective, where the action of $O(n, \mathbb{C})$ is diagonal.

(2) Let n be even and let V be the standard representation of $\text{Sp}(n,\mathbb{C})$ on \mathbb{C}^n . Then, for any $m \geq 0$, the map of algebraic varieties

$$
V^{\oplus m}/\mathrm{Sp}(n,\mathbb{C}) \to \prod_{1 \leq i < j \leq m} \mathbb{A}^1, (v_1, \dots, v_m) \mapsto \prod_{i < j} \langle v_i, v_j \rangle
$$

is a closed immersion. Equivalently, the map

$$
\mathbb{C}[t_{ij}]_{1\leq i
$$

is surjective, where the action of $O(n, \mathbb{C})$ is diagonal.

(3) Let V be the standard representation of $GL_n = U(n, \mathbb{C})$ on \mathbb{C}^n . Then, for any $m_1, m_2 \geq 0$, the map of algebraic varieties

$$
(V^{\oplus m_1} \oplus V^{*\oplus m_2})/\mathrm{GL}_n \to \prod_{\substack{1 \le i \le m_1 \\ 1 \le j \le m_2}} \mathbb{A}^1
$$

$$
(v_1, \dots, v_{m_1}, \delta_1, \dots, \delta_{m_2}) \mapsto (\langle v_i, \delta_j \rangle)_{\substack{1 \le i \le m_1 \\ 1 \le j \le m_2}}.
$$

is a closed immersion. Equivalently, the map

$$
\mathbb{C}[t_{ij}]_{\substack{1 \leq i \leq m_1 \\ 1 \leq j \leq m_2}} \xrightarrow{t_{ij} \mapsto \langle v_i, \delta_j \rangle} \left(\text{Sym}^{\bullet} \left(V^{\oplus m_1} \oplus V^{* \oplus m_2}\right)^{*}\right)^{\text{GL}_n}
$$

is surjective, where the action of GL_n is diagonal on $V^{\oplus m_1} \oplus V^{*\oplus m_2}$.

Proof. Part (1) follows from [\[25,](#page-60-21) Theorem 2.9.A] or, for a more modern reference, $[20, 10.2-(1)$ $[20, 10.2-(1)$ on p.114. Part (2) follows from $[25,$ Theorem 6.1.A or, for a more modern reference, [\[20,](#page-60-22) 10.3 on p.114]. Part (3) follows from [\[25,](#page-60-21) Theorem 2.6.A] or, for a more modern reference, [\[20,](#page-60-22) First Fundamental Theorem for $GL(V)$ in 2.1 on $p.16$].

Proposition 4.3.2. For $V = \mathbb{C}^n$ with the permutation representation of Σ_n ,

$$
\dim \left(\mathrm{Sym}^{k_1} V \otimes \mathrm{Sym}^{k_2} V \otimes \ldots \otimes \mathrm{Sym}^{k_\ell} V\right)^{\Sigma_n}
$$

is equal to the number of partitions of

$$
(k_1,\ldots,k_\ell)=\sum_{\vec{v}\in\mathbb{Z}_{\geq 0}^\ell}a_{\vec{v}}\vec{v} \text{ such that }\sum_{\vec{v}\in\mathbb{Z}_{\geq 0}^\ell}a_{\vec{v}}=n.
$$

Proof. Note that Sym^kV can be viewed the permutation representation associated to the Σ_n -set of labelings of $\{1,\ldots,n\}$ by elements of $\mathbb{Z}_{\geq 0}$ such that the total sum of the labels is k . The tensor product

$$
\mathrm{Sym}^{k_1}V \otimes \mathrm{Sym}^{k_2}V \otimes \ldots \otimes \mathrm{Sym}^{k_\ell}V
$$

is thus the permutation representation associated to the product of these Σ_n -sets, which can be viewed as the permutation representation associated to the Σ_n -set of labelings $\{1, \ldots, n\}$ by vectors $\vec{v} \in \mathbb{Z}_{\geq 0}$ whose sum is (k_1, \ldots, k_l) . The dimension of the Σ_n -invariants is thus equal to the number of orbits in the permutation action of Σ_n on this set of labelings. These orbits are determined uniquely by the multiplicities $a_{\vec{v}}$ with which each vector \vec{v} occurs as a label, and since there are n elements in the set we are labeling, the sum of the multiplicities $a_{\vec{v}}$ is equal to n. \Box

The following computation of generating functions that was used in the proof of Theorem [4.2.1](#page-25-1) is a simple consequence of the above results in invariant theory. We state the results in non-symmetric power series rings, where one has multiplicity and degree filtrations defined similarly (that restrict to the multiplicity filtrations on the symmetric power series rings).

Lemma 4.3.3.

(1) Let $G = O(n)$ and let V denote its standard representation on \mathbb{C}^n . Then

$$
\sum_{\tau} \dim_{\mathbb{C}}(\operatorname{Sym}^{\tau} V)^{G} m_{\tau} = \prod_{i \leq j} \frac{1}{1 - t_{i} t_{j}} \mod \mathbb{Z}[[\underline{t}_{\mathbb{N}}]]_{\mathrm{mult} > n}.
$$

(2) For n even, let $G = Sp(n)$ and let V denote its standard representation on \mathbb{C}^n . Then

$$
\sum_{\tau} \dim_{\mathbb{C}}(\operatorname{Sym}^{\tau}V)^{G} m_{\tau} = \prod_{i < j} \frac{1}{1 - t_{i}t_{j}} \mod \mathbb{Z}[[\underline{t}_{\mathbb{N}}]]_{\mathrm{mult} > n}.
$$

(3) Let $G = \sum_n$ and let V denote its standard permutation representation on \mathbb{C}^n . Then

$$
\sum_{\tau} \dim_{\mathbb{C}}(\operatorname{Sym}^{\tau} V)^{G} m_{\tau} = \prod_{\substack{n \text{ on-constant} \\ n \text{ in } t_1, t_2, \ldots \\ n \text{ in } t_1, t_2, \ldots}}^{n-term} \frac{1}{1 - m}
$$

$$
\equiv \prod_{\substack{n \text{ on-constant} \\ n \text{ non-constant} \\ n \text{ in } t_1, t_2, \ldots}} \frac{1}{1 - m} \mod \mathbb{Z}[[\underline{t}_{\mathbb{N}}]]_{\text{deg}>n}
$$

where the superscript "n-term" denotes that, when expanding the product, we only consider terms corresponding to at most n entries not equal to 1. (4) Let $G = U(n)$ and let V denote its standard representation on \mathbb{C}^n . Then

$$
\sum_{\tau,\overline{\tau}} \dim_{\mathbb{C}} \left(\operatorname{Sym}^{\tau} V_n \otimes \operatorname{Sym}^{\overline{\tau}} V_n^* \right)^{G_n} m_{\tau} m_{\overline{\tau}} \equiv \prod_{i,j} \frac{1}{1 - t_i \overline{t}_j} \mod \mathbb{Z}[[\underline{t}_{\mathbb{N}}, \underline{\overline{t}}_{\mathbb{N}}]]_{\mathrm{mult} > n}.
$$

Proof. The symmetric group case (3) is a direct rephrasing of Proposition [4.3.2](#page-27-2) using generating functions.

In cases (1) and (2) (resp. (4)), taking $m = n$ (resp. $m_1 = m_2 = n$) in Proposition [4.3.1,](#page-27-1) we claim the given closed immersion is actually an isomorphism. Since both sides are reduced, it suffices to see that the maps are dominant. But this is easily verified by considering the pairings of the standard basis vectors. For example, in the orthogonal case (1), to hit $(a_{ij})_{i\leq j}\in \prod_{i\leq j}\mathbb{C}$ with nonzero diagonal terms $a_{ii} \neq 0$ as the image of v_1, \ldots, v_n , we may take

$$
v_1 = \sqrt{a_{11}} \cdot e_1, v_2 = \frac{a_{12}}{\sqrt{a_{11}}} \cdot e_1 + \sqrt{a_{22}} \cdot e_2, v_3 = \frac{a_{13}}{\sqrt{a_{11}}} \cdot e_1 + \frac{a_{12} - \frac{a_{13}a_{12}}{a_{11}}}{\sqrt{a_{22}}} \cdot e_2 + \sqrt{a_{33}} \cdot e_3 \dots
$$

It follows that, in each of these cases, the map of rings in Proposition [4.3.1](#page-27-1) is an isomorphism. Then, to conclude cases (1) and (2), we expand

$$
\mathrm{Sym}^{\bullet}(V^{\oplus n})=\sum_{(k_1,\ldots,k_n)\in\mathbb{Z}_{\geq 0}^n}\mathrm{Sym}^{k_1}V\otimes \mathrm{Sym}^{k_2}V\otimes\ldots\otimes \mathrm{Sym}^{k_n}V,
$$

and the computation of the generating functions then follows immediately from the isomorphisms on the rings of invariants: in the notation of Proposition [4.3.1,](#page-27-1) each monomial $\prod t_{ij}^{b_{ij}}$ in the variables t_{ij} in the ring of invariants contributes a one dimensional space in Sym^{$\vec{k}V$} where $\vec{k} = \sum b_{ij} \cdot (e_i + e_j)$. The verification of case (4) is similar starting from the analogous decomposition of $\text{Sym}^{\bullet}(V^{\oplus n} \oplus V^{* \oplus n})$. \Box

4.4. Traces of random matrices. We now deduce the results of Diaconis and Shahshahani [\[11\]](#page-60-2) on traces of Haar random matrices in classical groups from Theorem $A/$ Theorem [4.2.1-](#page-25-1) (1) , (2) , and (4) .

Corollary 4.4.1 (Diaconis and Shahshahani^{[3](#page-29-2)} [\[11\]](#page-60-2), Theorems 2, 4, and 6).

Let
$$
g_j(a) = \begin{cases} 0 & \text{if } j \text{ and } a \text{ are both odd.} \\ j^{a/2}(a-1)(a-3)\cdots 1 & \text{if } j \text{ is odd and } a \text{ is even.} \\ \sum_k {a \choose 2k} j^k (2k-1)(2k-3)\cdots 1 & \text{if } j \text{ is even.} \end{cases}
$$

(1) For $\sum ia_i \leq n/2$ and M a Haar-random matrix in $O(n)$,

(4.4.1.1)
$$
\mathbb{E}[\prod \text{Tr}(M^i)^{a_i}] = \prod_i g_i(a_i)
$$

(2) For n even, $\sum ia_i \leq n/2$, and M a Haar-random matrix in $\text{Sp}(n)$,

(4.4.1.2)
$$
\mathbb{E}[\prod \text{Tr}(M^{i})^{a_i}] = \prod_{i} (-1)^{a_i - 1} g_i(a_i)
$$

(3) For $\sum ia_i \leq n$, $\sum_i ib_i \leq n$, and M a Haar-random matrix in $U(n)$,

(4.4.1.3)
$$
\mathbb{E}[\prod \text{Tr}(M^{i})^{a_i} \text{Tr}(\overline{M}^{i})^{b_i}] = \begin{cases} 0 & \text{if } (a_i) \neq (b_i) \\ \prod_i i^{a_i} a_i! & \text{if } (a_i) = (b_i) \end{cases}
$$

 3 The lower bounds on n in these results as stated in [\[11\]](#page-60-2) are not correct and do not follow from the proofs given there. The correct bounds are stated, e.g., in [\[18,](#page-60-23) Propositions 2.5 and 3.6].

Proof of Corollary [4.4.1.](#page-29-1) We first treat the orthogonal case. For V the standard representation of $O(n)$ and with notation as in Theorem [4.2.1,](#page-25-1) we have

$$
\mathbb{E}[\prod \text{Tr}(M^i)^{a_i}] = \mathbb{E}[\prod_i p_i^{a_i} \circ [V]] = \langle \mathbb{E}[\text{Exp}_{\sigma}([V]h_1)], \prod_i p_i^{a_i} \rangle,
$$

where the last equality follows from Lemma [3.2.2.](#page-21-3) We claim that, under the condition $\sum ia_i \leq n$, Theorem [4.2.1](#page-25-1) implies this final inner product is equal to

$$
\langle \text{Exp}_{\sigma}(h_2), \prod_i p_i^{a_i} \rangle.
$$

Indeed, $\prod_i p_i^{a_i}$ lives in degree $\sum i a_i \leq n$, and the degree $\leq n$ part of the moment generating function and $Exp_{\sigma}(h_2)$ agree by Theorem [4.2.1](#page-25-1) (since $|\tau| \leq ||\tau||$).

We now compute this inner product by expanding $Exp_{\sigma}(h_2)$ in terms of power sum monomials. Using Lemma [2.5.4,](#page-17-1) we have

$$
\begin{split} \operatorname{Exp}_{\sigma}(h_{2}) &= \operatorname{Exp}_{\sigma}\left(\frac{p_{1}^{2} + p_{2}}{2}\right) \\ &= \prod_{i=1}^{\infty} \exp\left(\frac{p_{i}}{i} \circ \frac{p_{1}^{2} + p_{2}}{2}\right) \\ &= \prod_{i=1}^{\infty} \exp\left(\frac{p_{i}^{2} + p_{2i}}{2i}\right) \\ &= \prod_{i \text{ odd}} \exp\left(\frac{p_{i}^{2}}{2i}\right) \prod_{i \text{ even}} \exp\left(\frac{p_{i}^{2}}{2i}\right) \exp\left(\frac{p_{i}}{i}\right) \\ &= \prod_{i \text{ odd}} \left(\sum_{k\geq 0} \frac{p_{i}^{2k}}{(2i)^{k}k!}\right) \prod_{i \text{ even}} \left(\sum_{k\geq 0} \frac{p_{i}^{2k}}{(2i)^{k}k!}\right) \left(\sum_{k\geq 0} \frac{p_{i}^{k}}{k!i^{k}}\right) \\ &= \prod_{i \text{ odd}} \left(\sum_{k\geq 0} \frac{p_{i}^{2k}}{(2k)!1i^{k}}\right) \prod_{i \text{ even}} \left(\sum_{k\geq 0} \frac{p_{i}^{2k}}{(2k)!1i^{k}}\right) \left(\sum_{k\geq 0} \frac{p_{i}^{k}}{k!i^{k}}\right) \end{split}
$$

where $(2k)!! = (2k)(2k-2)(2k-4) \ldots (2)$. Expanding, the coefficient of $\prod p_i^{a_i}$ is

$$
\prod_{i \text{ odd}} \left(\left\{ \frac{\frac{1}{a_i!!i^{a_i/2}}}{0} \text{ if } a_i \text{ odd} \right\} \prod_{i \text{ even}} \left(\sum_{2k \leq a_i} \frac{1}{(2k)!!(a_i - 2k)!i^{a_i - k}} \right) = \prod_i \frac{g_i(a_i)}{a_i!i^{a_i}}.
$$

We then obtain the formula on the right-hand side of Eq. [\(4.4.1.1\)](#page-29-3) by taking the Hall inner product with $\prod_i p_i^{a_i}$ and using Eq. [\(2.7.0.2\)](#page-19-6) (which shows that distinct power sum monomials are orthogonal and that $\langle \prod_i p_i^{a_i}, \prod_i p_i^{a_i} \rangle = \prod_i a_i! i^{a_i}$.

The argument in the symplectic case is the same except for the inner product computation, which differs only in a sign (alternatively, one can apply Lemma [2.8.1](#page-19-2)

to the computation above since $\omega(h_2) = e_2$:

$$
\begin{split} \operatorname{Exp}_{\sigma}(e_{2}) &= \operatorname{Exp}_{\sigma}\left(\frac{p_{1}^{2}-p_{2}}{2}\right) \\ &= \prod_{i=1}^{\infty} \exp\left(\frac{p_{i}}{i} \circ \frac{p_{1}^{2}-p_{2}}{2}\right) \\ &= \prod_{i=1}^{\infty} \exp\left(\frac{p_{i}^{2}-p_{2i}}{2i}\right) \\ &= \prod_{i \text{ odd}} \exp\left(\frac{p_{i}^{2}}{2i}\right) \prod_{i \text{ even}} \exp\left(\frac{p_{i}^{2}}{2i}\right) \exp\left(\frac{-p_{i}}{i}\right) \\ &= \prod_{i \text{ odd}} \left(\sum_{k\geq 0} \frac{p_{i}^{2k}}{(2i)^{k}k!}\right) \prod_{i \text{ even}} \left(\sum_{k\geq 0} \frac{p_{i}^{2k}}{(2i)^{k}k!}\right) \left(\sum_{k\geq 0} \frac{(-p_{i})^{k}}{k!i^{k}}\right) \\ &= \prod_{i \text{ odd}} \left(\sum_{k\geq 0} \frac{p_{i}^{2k}}{(2k)!!i^{k}}\right) \prod_{i \text{ even}} \left(\sum_{k\geq 0} \frac{p_{i}^{2k}}{(2k)!!i^{k}}\right) \left(\sum_{k\geq 0} \frac{(-p_{i})^{k}}{k!i^{k}}\right) \end{split}
$$

so that the coefficient of $\prod p_i^{a_i}$ is

$$
\prod_{i \text{ odd}} \left(\left\{ \frac{\frac{1}{a_i!!i^{a_i/2}}}{0} \text{ if } a_i \text{ odd} \right\} \prod_{i \text{ even}} (-1)^{a_i} \left(\sum_{2k \le a_i} \frac{1}{(2k)!!(a_i - 2k)!i^{a_i - k}} \right) \right)
$$
\n
$$
= \prod_i \frac{(-1)^{a_i} g_i(a_i)}{a_i!i^{a_i}}.
$$

and we obtain the desired inner product giving Eq. [\(4.4.1.2\)](#page-29-4) exactly as in the orthogonal case treated above.

Finally we treat the unitary case, following the same strategy. For V the standard representation of $U(n)$ and with notation as in Theorem [4.2.1,](#page-25-1) we have

$$
\mathbb{E}[\prod \text{Tr}(M^{i})^{a_i} \text{Tr}(\overline{M}^{i})^{b_i}] = \mathbb{E}\left[\left(\prod p_i^{a_i} \circ [V]\right) \left(\prod p_i^{b_i} \circ [V^*]\right)\right]
$$

$$
= \langle \mathbb{E}[\text{Exp}_{\sigma}([V]h_1 + [V^*]\overline{h}_i)], \prod_i p_i^{a_i} \prod_i \overline{p}_i^{b_i}.\rangle
$$

where on $(\Lambda \otimes \Lambda)$ ^{\wedge} we have taken the obvious tensor extension of the Hall inner product and are using the analog of Lemma [3.2.2](#page-21-3) in this setting.

We claim that, under the condition $\sum ia_i \leq n$ and $\sum ib_i \leq n$, Theorem [4.2.1](#page-25-1) implies this final inner product is equal to

$$
\langle \operatorname{Exp}_{\sigma}(h_1 \overline{h}_1), \prod p_i^{a_i} \prod_i \overline{p}_i^{b_i} \rangle.
$$

Indeed, $\prod_i p_i^{a_i}$ lives in degree $\sum i a_i \leq n$ while $\prod_i \overline{p}_i^{b_i}$ lives in degree $\sum i b_i \leq n$, and the part of the moment generating function where each degree is $\leq n$ and $\text{Exp}_{\sigma}(h_1h_1)$ agree by Theorem [4.2.1](#page-25-1) (again since $||\tau_i|| \leq |\tau_i|$). We now compute this inner product by expanding $Exp_{\sigma}(h_1h_1)$ in terms of power sum monomials.

$$
\begin{aligned} \operatorname{Exp}_{\sigma}(h_1 \overline{h}_1) &= \operatorname{Exp}_{\sigma}(p_1 \overline{p}_1) \\ &= \prod_{i=1}^{\infty} \operatorname{exp}\left(\frac{p_i}{i} \circ p_1 \overline{p}_1\right) \\ &= \prod_{i=1}^{\infty} \operatorname{exp}\left(\frac{p_i \overline{p}_i}{i}\right) \\ &= \prod_i \left(\sum_{k \ge 0} \frac{p_i^k \overline{p}_i^k}{i^k k!}\right). \end{aligned}
$$

The coefficient of $\prod p_i^{a_i} \prod \overline{p}_i^{b_i}$ is thus zero if $(a_i) \neq (b_i)$, and $\prod_i \frac{1}{i^{a_i}k!}$ otherwise. Thus, by Eq. [\(2.7.0.2\)](#page-19-6), taking the inner product with $\prod p_i^{a_i} \prod \overline{p}_i^{b_i}$ gives zero if $(a_i) \neq (b_i)$ and $\prod_i \frac{(i^{a_i}k!)^2}{i^{a_i}k!}$ $\frac{a_i^{a_i}k!}{i^{a_i}k!} = \prod_i i^{a_k}k!$ if $(a_i) = (b_i)$, recovering the right hand side of Eq. $(4.4.1.3)$.

Remark 4.4.2. In [\[11\]](#page-60-2), Corollary [4.4.1](#page-29-1) is interpreted as an asymptotic description of traces of powers as independent Gaussian random variables. The control on the moments in Corollary [4.4.1](#page-29-1) is enough to give a bound on the rate of convergence, and Johansson [\[18\]](#page-60-23) has established a much stronger estimate. Our method of proving Theorem [A](#page-3-0) shows that each coefficient in the moment generating function is nonnegative and bounded above by the limiting value. Using the second fundamental theorem of invariant theory, one can in principle control the discrepancy exactly. It would be interesting to understand what bounds this method gives on the rate of convergence, and to compare to the method of [\[18\]](#page-60-23) and related work.

4.5. Cycle counting in the symmetric group. We now explain how to recover [\[11,](#page-60-2) Theorem 7] on the joint moments of cycle counting functions on symmetric groups from Theorem $A/$ Theorem [4.2.1-](#page-25-1)(3).

We adopt the notation of Theorem [4.2.1-](#page-25-1)(3). First, we note that, by Definition [3.3.2/](#page-23-1)Lemma [3.3.1,](#page-22-3) Theorem [4.2.1](#page-25-1) implies that

$$
\mathbb{E}\left[(1+h_1)^{[V_n]} \right] \equiv \operatorname{Exp}_{\sigma}(h_1) \mod \Lambda_{\deg > n}^{\wedge}.
$$

Indeed, the degree for the congruence does not change when switching between moment generating functions because deg $h_{\tau} = \deg c_{\tau}$ (note, however, that the c_{τ} are not homogeneous).

Remark 4.5.1. One can also make a combinatorial argument to directly compute the falling moments to obtain

$$
\mathbb{E}\left[(1+h_1)^{[V_n]} \right] \equiv \operatorname{Exp}_{\sigma}(h_1) \mod \Lambda_{\deg > n}^{\wedge}.
$$

Moreover, in the spirit of Remark [3.2.5,](#page-22-1) because $[V_n]$ is secretly a combinatorial random variable, the argument is much simpler in the falling moment case: here to obtain the coefficient of m_{τ} , one is counting orbits in the configuration space $C^{\tau}(\{1,\ldots,n\})$, and there is exactly one orbit if it is non-empty.

Letting C_k denote the kth cycle counting function, we have

$$
(1 + h_1)^{[V_n]} = \prod_{i=1}^{\infty} \exp(C_i \log(1 + p_i))
$$

34 SEAN HOWE

where the exp and log on the right are computed as the classical exponential and logarithm series. This identity can be verified by hand, or viewed as a special case of [\[17,](#page-60-6) Lemma 3.5]. Then, since $\text{Exp}_{\sigma}(h_1) = \prod_{i=1}^{\infty} \text{exp}(\frac{p_i}{i})$ by Lemma [2.5.4,](#page-17-1) we deduce that

$$
\mathbb{E}\left[\prod_{i\geq 1} \exp(C_i \log(1+p_i))\right] \equiv \prod_{i\geq 1} \exp\left(\frac{p_i}{i}\right) \mod \Lambda_{\deg>n}^{\wedge}
$$

Note that $\exp(C_i \log(1 + p_i)) = (1 + p_i)^{C_i}$, where the power is interpreted in the classical sense. Since the p_i are a polynomial basis for Λ , this congruence says that, up to terms of degree $> n$, the joint falling moments of the cycle counting functions C_i are the joint falling moments of independent Poission distributions of mean $\frac{1}{i}$, which is precisely the statement of [\[11,](#page-60-2) Theorem 7].

5. Abstract point-counting

We are interested in studying the L-functions of ℓ -adic local systems and constructible sheaves on varieties over a finite field \mathbb{F}_q . Although a priori defined as Euler products over closed points, these are rational functions by the results of [\[10\]](#page-60-24), and we want to study the distribution of their zeroes, viewed as multisets of complex numbers by a fixed isomorphism $\mathbb{Q}_{\ell} \cong \mathbb{C}$, and discuss convergence properties for the archimedean topology on \mathbb{C} . As in [\[6\]](#page-60-14), it is thus natural to pass from $\mathbb{Z}[\mathbb{C}]$ to its completion $W(\mathbb{C}) \cong 1 + t\mathbb{C}[[t]]$ (for the point-counting topology of loc. cit.), in which case we do not actually need to know that the L-function was rational, since already the Euler product gave us the element of $1 + t\mathbb{C}[[t]]$. Moreover, the notion of an Euler product does not depend on the actual variety, just on its set of closed points, or equivalently the \mathbb{F}_q -points equipped with the action of Z where 1 acts by the (geometric) Frobenius, i.e. the action sending coordinates to their q-powers. Thus we find that the structure needed to study this problem via λ -probability is very simple and can be abstracted completely away from algebraic geometry: we need to look at \mathbb{Z} -invariant $W(\mathbb{C})$ -valued functions on (admissible) \mathbb{Z} -sets, and take an expectation modeled after an Euler product. This is the theory developed here.

In Section [5.1](#page-34-0) we recall some coordinates on $W(\mathbb{C})$. In Section [5.2](#page-34-1) we define the category of admissible Z-sets and explain the relation with zeta functions and other basic properties. In particular, in Proposition [5.2.7](#page-35-0) we give a description of certain powers (in the sense of Section [2.6\)](#page-18-0) that will be used in the proofs of Theorem [B](#page-7-0) and Theorem [C.](#page-9-0) In Section [5.3](#page-36-0) we introduce the relevant rings of $W(\mathbb{C})$ -valued functions and study their functoriality properties; in particular, we explain how to integrate these over an admissible \mathbb{Z} -set. In Section [5.4](#page-38-0) (see also Lemma [5.5.1\)](#page-39-2) we explain a key relation between integration over admissible Z-sets and integration over finite sets that we will use to reduce the proofs of Theorem [B](#page-7-0) and Theorem [C](#page-9-0) to naïve point-counting problems that can be accessed by Poonen's sieve. In Section [5.5,](#page-39-0) we define the relevant λ -probability space for an admissible \mathbb{Z} -set (generalizing the simple case of $W(\mathbb{C})$ -valued functions on a finite set with the uniform measure).

Finally, in Section [5.6](#page-39-1) we briefly explain the compatibility with the theory of constructible ℓ -adic sheaves and their L -functions; let us emphasize though that most of the results and proofs of this paper can be phrased so that they are completely independent of this interpretation!

5.1. The λ -ring $W(\mathbb{C})$ and zeta functions. Recall from Section [2.4](#page-16-1) that $W(\mathbb{C})$ is the ring $\text{Hom}_{\text{Ring}}(\Lambda, \mathbb{C})$. We recall here some more explicit interpretations in this case (see, e.g., [\[7,](#page-60-25) §3] for more details and several related constructions).

We identify $W(\mathbb{C})$ as an additive group with $1 + t\mathbb{C}[[t]]$ by

$$
\alpha \in \mathrm{Hom}_{\mathrm{Ring}}(\Lambda,\mathbb{C}) \mapsto \sum_{k\geq 0} \alpha(h_k)t^k \in 1 + t\mathbb{C}[[t]].
$$

We identify $W(\mathbb{C})$ as a ring with $\prod_{i\geq 1} \mathbb{C}$ by

$$
\alpha \mapsto \sum_{k \ge 1} \alpha(p_k) t^k,
$$

i.e. by passing to the ghost coordinates (which is always an isomorphism for any $\mathbb Q$ -algebra). To pass from the first identification to the second, one applies td log to an element of $1 + t\mathbb{C}[[t]]$ then takes the coefficients, i.e. for $\alpha \in W(\mathbb{C}),$

$$
td\log(\sum_{k\geq 0} \alpha(h_k)t^k) = \sum_{k\geq 1} \alpha(p_k)t^k.
$$

 $\prod_{i\geq 1} \mathbb{C}$, the plethystic action of Λ on $W(\mathbb{C})$ is determined by This identity follows from Eq. [\(2.1.0.3\)](#page-13-2). Note that, in the identification with

(5.1.0.1)
$$
p_j \circ (a_i)_i = (a_{ij})_i.
$$

Given $f \in \mathbb{C}[[t]]$ and k a positive integer, we write $f(t^k)$ for the series obtained by substituting t^k for t . Note that, for $f \in 1 + t\mathbb{C}[[t]],$

(5.1.0.2)
$$
td\log(f(t^k)) = k(d\log(f(t)))(t^k).
$$

Given $\alpha \in W(\mathbb{C})$, by $\alpha(t^k)$ we will always mean the element obtained by viewing α as an element of $1 + t\mathbb{C}[[t]]$ and then substituting t^k . On $\prod_{i\geq 1}\mathbb{C}$, Eq. [\(5.1.0.2\)](#page-34-2) shows this acts as

(5.1.0.3)
$$
(a_i)_i \mapsto (ka_{i/k})_i, \text{ where } a_{i/k} := 0 \text{ if } i/k \notin \mathbb{Z}_{\geq 1}.
$$

It follows that, for [**k**] the element of $W(\mathbb{C})$ corresponding to $\frac{1}{1-t^k} \in 1 + t\mathbb{C}[[t]]$ and k k

thus to ($\overline{0,\ldots,0,k},$ $\overline{0,\ldots,0,k},\ldots\rangle\in\prod_{i\geq 1}\mathbb{C},$ (5.1.0.4) $(p_k \circ \alpha)(t^k) = [\mathbf{k}] \cdot \alpha \text{ and } p_k \circ (\alpha(t^k)) = k\alpha$

5.2. Admissible Z-sets.

Definition 5.2.1. A \mathbb{Z} -set is a set with an action of \mathbb{Z} .

- (1) An Z-set S is admissible^{[4](#page-34-3)} if each point $s \in S$ is stabilized by a finite index subgroup $k\mathbb{Z}$, and $S^{k\mathbb{Z}}$ is finite for each $k \in \mathbb{Z}_{>0}$.
- (2) For k a positive integer, we write **k** for the admissible Z-set $\mathbb{Z}/k\mathbb{Z}$.
- (3) For S an admissible Z-set, and $s \in S$, we write |s| for the orbit of s and $deg(s) = deg(|s|)$ for the size of the orbit. We write $S(\mathbf{k}) := Hom_{\mathbb{Z}-set}(\mathbf{k}, S)$.

Example 5.2.2.

(1) Any finite set is an admissible Z-set when equipped with the trivial action. This identifies finite sets with a full subcategory of admissible Z-sets.

 4 An admissible Z-set is, in particular, a $\hat{\mathbb{Z}}$ -set, and the terminology is by analogy with the theory of admissible representations of profinite groups.

36 SEAN HOWE

(2) Let \mathbb{F}_q be a finite field and let X/\mathbb{F}_q be a variety. Then $X(\overline{\mathbb{F}}_q)$ is an admissible Z-set, where $1 \in \mathbb{Z}$ acts by the geometric Frobenius (which, if X is affine, acts by raising coordinates to the qth power). We have $|X(\overline{\mathbb{F}}_q)| = |X|$, the set of closed points of X, and $(X(\overline{\mathbb{F}}_q))(\mathbf{k}) = X(\mathbb{F}_{q^k})$. For $x \in X(\overline{\mathbb{F}}_q)$, $\deg x = [\mathbb{F}_q(x) : \mathbb{F}_q]$.

For S an admissible \mathbb{Z} -set, its zeta function is defined by

$$
Z_S(t) := \prod_{s \in |S|} \frac{1}{1 - t^{\deg(s)}} \in 1 + t\mathbb{C}[[t]] = W(\mathbb{C}).
$$

We also write $[S] = Z_S(t)$. We note that, with this definition $[\mathbf{k}] = \frac{1}{1-t^k}$, so that this notation does not conflict with the notation introduced above in Section [5.1.](#page-34-0)

Lemma 5.2.3. Let S be an admissible \mathbb{Z} -set. Then

$$
td\log(Z_s(t))=\sum_{k\geq 1}\#S([\mathbf{k}])t^k
$$

.

In particular, under our identification $W(\mathbb{C}) = \prod_{i \geq 1} \mathbb{C}$,

$$
[S] = (\#S([1]), \#S([2]), \ldots)
$$

and we have

$$
Z_{S_1 \sqcup S_2}(t) = Z_{S_1}(t) + _W Z_{S_2}(t) = Z_{S_1}(t)Z_{S_2}(t) \text{ and } Z_{S_1 \times S_2}(t) = Z_{S_1}(t) \cdot _W Z_{S_2}(t),
$$

were the subscript W emphasizes addition and multiplication in $W(\mathbb{C})$ vs $\mathbb{C}[[t]]$.

Proof. The computation of td log is the usual elementary expansion as for zeta functions of varieties over finite fields. The remaining equalities follow immediately since the coefficients of $td \log$ give the identification $W(\mathbb{C}) = \prod_{i \geq 1} \mathbb{C}$.

Remark 5.2.4. The assignment $S \mapsto [S] = Z_s(t)$ extends to an identification of $K_0(\text{Admissible } \mathbb{Z}\text{-sets}) = W(\mathbb{Z}) \subseteq W(\mathbb{C}),$ which explains the notation. The λ -ring structure is determined on the Grothendieck ring by $h_k \circ [S] = [\text{Sym}^k S].$

Definition 5.2.5. Let $\varphi: V \to S$ be a morphism of admissible Z-sets. For $s \in S$, we define the fiber of φ over s, V_s to be the admissible Z-set $\varphi^{-1}(s)$ with Z-action obtained from the Z-action on V by precomposition with multiplication by deg(s).

Example 5.2.6. If S is an admissible Z-set, then the fiber $(S \times [k])_1$ over $1 \in [k]$ is naturally identified with the set S but where the $\mathbb Z$ action is precomposed with multiplication by k. In particular, let \mathbb{F}_q be a finite field and let X/\mathbb{F}_q be a variety, and let $X' = X_{\mathbb{F}_{q^k}}$. Then $X(\overline{\mathbb{F}}_q) = X'(\overline{\mathbb{F}}_q)$ as sets, but as admissible Z-sets we have $X'(\overline{\mathbb{F}}_q) = (X(\overline{\mathbb{F}}_q) \times [k])_1.$

We will need the following result computing certain powers in this setting.

Proposition 5.2.7. Let V be an admissible \mathbb{Z} -set, and let $F \in W(\mathbb{C})[[\underline{t}_{\mathbb{N}}]]$ with constant coefficent 1. Then, using subscripts to denote projection to coordinates in $W(\mathbb{C}) = \prod_{i \geq 1} \mathbb{C},$

$$
(F^{[V]})_i = \prod_{|v| \in |(V \times \mathbf{i})_1|} F_{i \deg(|v|)}(\underline{t}_I^{\deg(|v|)}).
$$

In particular, for V/\mathbb{F}_q a variety, using Example [5.2.6,](#page-35-1)

$$
(F^{[V(\overline{\mathbb{F}}_q)]})_i = \prod_{|v| \in |V_{\overline{\mathbb{F}}_{q^i}}|} F_{i \deg(|v|)}(\underline{t}_I^{\deg(v)})
$$

where, $|V_{\mathbb{F}_{q^i}}|$ denotes the set of closed points and in computing the degree of a closed point $|v| \in |V_{\mathbb{F}_{q^i}}|$, $V_{\mathbb{F}_{q^i}}$ is treated as a variety over \mathbb{F}_{q^i} (rather than over \mathbb{F}_q).

Proof. We first note that

$$
F^{[V]} = F^{\sum_{|v| \in |V|} |v|} = \prod_{|v| \in |V|} F^{|v|}.
$$

Thus we are reduced to treating the case of a single orbit, i.e. $V = d$, so that

$$
[V] = [\mathbf{d}] = \frac{1}{1 - t^d} = \underbrace{(0, \dots, 0, d, 0, \dots, 0, d, \dots)}_{(0, \dots, 0, d, 0, \dots, 0, d, \dots)}
$$

Now, let $\mu = \frac{\text{lcm}(i,d)}{i} = \frac{d}{\text{gcd}(i,d)}$. In the following we will write $p_i * F$ for the plethystic action of power sum function on F acting only on the coefficients and not the variables; we note that this commutes with the usual plethystic action $p_i \circ$. Expanding via Lemma [2.5.4,](#page-17-1) we find

$$
(\text{Exp}_{\sigma}([V]\text{Log}_{\sigma}(F)))_i = \prod_{j\geq 1} \exp\left(\frac{p_j \circ [V]\text{Log}_{\sigma}(F)}{j}\right)_i
$$

\n
$$
= \prod_{j\geq 1} \exp\left(\frac{(p_j \circ [V])\text{Log}_{\sigma}(p_j \circ F)}{j}\right)_i
$$

\n
$$
= \prod_{j\geq 1} \exp\left(\frac{(p_{ij} \circ [V])\text{Log}_{\sigma}(p_j \circ p_i * F))}{j}\right)_1
$$

\n
$$
= \prod_{k\geq 1} \exp\left(\frac{\text{Log}_{\sigma}(p_{k\mu} \circ p_i * F)}{k\mu/d}\right)_1
$$

\n
$$
= \prod_{k\geq 1} \exp\left(\frac{d}{\mu} \cdot \frac{p_k \circ \text{Log}_{\sigma}(p_{\mu} \circ p_i * F)}{k}\right)_1
$$

\n
$$
= (\text{Exp}_{\sigma}(\text{Log}_{\sigma}(p_{\mu} \circ p_i * F))^{d/\mu})_1
$$

\n
$$
= (p_{\mu} \circ p_i * F)^{d/\mu}_1.
$$

Since $([\mathbf{d}] \times \mathbf{i})_1$ has $d/\mu = \gcd(i, d)$ orbits, each consisting of points of degree μ or \mathbb{F}_{a^i} , we conclude. over \mathbb{F}_{q^i} , we conclude.

5.3. $W(\mathbb{C})$ -valued functions.

Definition 5.3.1. For any admissible Z-set S, we write $C(S, W(\mathbb{C}))$ for the λ ring of Z-equivariant functions from S to $W(\mathbb{C})$, where the action of Z on $W(\mathbb{C})$ is trivial and the λ -ring structure is pointwise.

Note that, by the Z-equivariance, any $f \in C(S, W(\mathbb{C}))$ is constant on its orbit, so this ring is naturally identified with $C(|S|, W(\mathbb{C}))$.

Definition 5.3.2. Given an admissible Z-set V and $f \in C(V, W(\mathbb{C}))$, we define

$$
\int_V f = \sum_{v \in V} f(t^{\deg(v)})
$$

where we emphasizes that this is addition in $W(\mathbb{C})$ and that the infinite sum makes sense because, viewing it as an infinite product of power series in $1 + t\mathbb{C}[[t]]$, only finitely many terms contribute to the coefficient of each power of t (alternatively, viewing the sum as addition in $\prod_{\geq 1} \mathbb{C}$, only finitely many terms contribute to each component by Eq. $(5.1.0.3)$.

Definition 5.3.3. Given a map $\varphi: V \to S$ of admissible Z-sets, we define

(1) pull back of functions $\varphi^* : C(S, W(\mathbb{C})) \to C(V, W(\mathbb{C}))$ by

$$
\varphi^* f(z) := p_{\frac{\deg(|z|)}{\deg(|f(z)|)}} \circ f(\varphi(s)).
$$

(2) integration over fibers $\varphi_! : C(V, W(\mathbb{C})) \to C(S, W(\mathbb{C}))$ by

$$
(\varphi_!f)(s) := \int_{V_s} f|_{V_s},
$$

where the fiber V_s is as defined in Definition [5.2.5.](#page-35-2) Equivalently, viewing f and $\varphi_! f$ as functions on orbits,

$$
(\varphi_!f)(|s|)=\sum_{|v|\subseteq\varphi^{-1}(|s|)}(f(|v|))\left(t^{\frac{\deg(|v|)}{\deg(|s|)}}\right).
$$

Example 5.3.4. For the constant function 1 in $C(V, W(\mathbb{C}))$,

$$
\int_{V} 1 = \int_{V} \frac{1}{1 - t} = Z_{V}(t) = [V].
$$

More generally, if $\varphi: V \to S$ is a map of admissible Z-sets. Then we define $[V/S]$ to be the function sending s to $[V_s]$ as in Definition [5.2.5,](#page-35-2) so that $[V/S] = \varphi_1 1$.

Note that, since the power sum plethysms are λ -ring endomorphism on any λ -ring (see Section [2.4\)](#page-16-1), φ^* is a λ -ring morphism.

Example 5.3.5. Let $\varphi: V \to \{*\}$ be the final map. Then, $\varphi^* : W(\mathbb{C}) \to \varphi^* : V(\mathbb{C})$ $C(V, W(\mathbb{C}))$ sends $w \in W(\mathbb{C})$ to the function f_w defined by

(5.3.5.1)
$$
f_w(v) = p_{\deg v} \circ w \text{ for } v \in V.
$$

In particular, the W(\mathbb{C})-algebra structure given by φ^* agrees with the $W(\mathbb{C})$ -algebra structure coming from literal constant $W(\mathbb{C})$ -valued functions only when V is a finite set viewed as an admissible \mathbb{Z} -set under the embedding of Example [5.2.2-](#page-34-5)(1) (i.e. when V contains only points of degree 1). Note however that, for elements of $\mathbb C$ embedded diagonally in $W(\mathbb{C}) = \prod_{i \geq 1} \mathbb{C}$, the pullback structure and the constant structure always agree (using Eq. [\(5.1.0.1\)](#page-34-6)).

Warning: We always equip $C(V, W(\mathbb{C}))$ with the $W(\mathbb{C})$ - λ -algebra structure obtained by pullback from the final map $V \rightarrow \{*\}$ as in Example [5.3.5,](#page-37-0) and never the algebra structure from literal constant functions. In particular, if $w \in W(\mathbb{C})$, if we write $\int_V w$ this means to treat w as pulled back from the point, i.e. to integrate f_w from Example [5.3.5.](#page-37-0)

For $\varphi: V \to S$ a map of admissible Z-sets, the integration over fibers map $\varphi_!$ is additive but does not typically preserve products or plethysms. However, it does preserve the $W(\mathbb{C})$ -module structure, and even more:

Lemma 5.3.6 (Projection formula). Given a map $\varphi: V \to S$ of admissible Z-sets, we view $C(V, W(\mathbb{C}))$ as a $C(S, W(\mathbb{C}))$ -module by φ^* . Then, $\varphi_!$ is a morphism of $C(S, W(\mathbb{C}))$ -modules. In particular, φ_1 is always a morphism of $W(\mathbb{C})$ -modules.

Proof. The last "in particular" follows from the first claim after observing that, for any composable morphisms φ_1 and φ_2 , $(\varphi_2 \circ \varphi_1)^* f = \varphi_1^* \varphi_2^* f$.

For the first claim, by taking fibers one immediately reduces to the case $S = \{ * \}$ so that φ ! = \int_V . We for $w \in W(\mathbb{C})$, we write f_w for its pullback to V as in Exam-ple [5.3.5.](#page-37-0) To verify \int_V is $W(\mathbb{C})$ -linear, we note that for $w = (a_1, a_2, \ldots) \in W(\mathbb{C})$ and $v \in |V|$ of degree d, $f_w(v)$ is identified, via td log, with (a_d, a_{2d}, \ldots) . It follows that, for v of degree d and $g \in C(V,W(\mathbb{C}))$, $f_w g(v) = (a_d g_1(v), a_{2d} g_2(v), \ldots)$, so that, by Eq. [\(5.1.0.3\)](#page-34-4), $((f_w g(v))(t^d))_{di} = a_{di} dg_i(v)$ while the entries of index not divisible by d are all zero. In particular,

$$
\left(\int_{V} f_{w}g\right)_{k} = \sum_{v \text{ s.t. } \deg(v)|k} (\deg(v)g_{\frac{k}{\deg(v)}}(v)a_{k})
$$

$$
= a_{k} \sum_{v \text{ s.t. } \deg(v)|k} (\deg(v)g_{\frac{k}{\deg(v)}}(v))
$$

$$
= a_{k} \left(\int_{V} g\right)_{k}.
$$

Since this holds for all k, the map is $W(\mathbb{C})$ -linear.

5.4. A relation between integration over finite sets and integration over admissible Z-sets. For S an admissible Z-set and $k \geq 1$, there is a natural map of admissible Z-sets

$$
c_k: S(\mathbf{k}) \times \mathbf{k} \to S, (s, i) \mapsto (s(i))
$$

where here the finite set $S(\mathbf{k})$ is viewed as an admissible Z-set with the trivial action as in Example [5.2.2-](#page-34-5)(1). By pullback we obtain a map of $W(\mathbb{C})$ - λ -algebras.

$$
c_k^*: C(S, W(\mathbb{C})) \to C(S(\mathbf{k}) \times \mathbf{k}, W(\mathbb{C}))
$$

We can identify the right-hand side with $W(\mathbb{C})$ -valued functions on $|S(\mathbf{k}) \times \mathbf{k}| =$ $S(\mathbf{k})$ in the usual way, or equivalently by restricting to $S(\mathbf{k}) \times 1$. We thus obtain a map of $W(\mathbb{C})$ - λ -algebras

$$
res_k : C(S, W(\mathbb{C})) \to C(S(\mathbf{k}), W(\mathbb{C}))^{(k)}
$$

where the superscript on the right indicates that we take the $W(\mathbb{C})$ -algebra structure via $p_k \circ : W(\mathbb{C}) \to W(\mathbb{C})$. Explicity,

$$
\operatorname{res}_k(f)(s: \mathbf{k} \to S) = p_{\frac{k}{\deg(s(|\mathbf{k}|))}} \circ f(s(|\mathbf{k}|)).
$$

Proposition 5.4.1. For S an admissible Z-set, $k \geq 1$, and $f \in C(S, W(\mathbb{C}))$,

$$
\left(\int_{S} f\right)_k = \left(\int_{S(\mathbf{k})} \text{res}_k(f)\right)_1.
$$

where the subscripts denote components under $W(\mathbb{C}) = \prod_{i \geq 1} \mathbb{C}$.

Proof. For $f = (f_1, f_2, \ldots)$ in $C(S, W(\mathbb{C}))$, combining Definition [5.3.2](#page-36-1) with Eq. [\(5.1.0.3\)](#page-34-4), we obtain

$$
\left(\int_{S} f\right)_{k} = \sum_{\substack{|s| \in |S| \text{ s.t.} \\ \deg(|s|)|k}} \deg(|s|) f_{\frac{k}{\deg(|s|)}}(|s|)
$$

$$
= \sum_{\substack{|s| \in |S| \text{ s.t.} \\ \deg(|s|)|k}} \sum_{\substack{s \in S(\mathbf{k}) \\ \deg(|s|)|k}} f_{\frac{k}{\deg(|s|)}}(|s|)
$$

$$
= \sum_{s \in S(\mathbf{k})} f_{\frac{k}{\deg(s(\mathbf{k}))}}(s(\mathbf{k}))
$$

$$
= \sum_{s \in S(\mathbf{k})} (p_{\frac{k}{\deg(s(\mathbf{k}))}} \circ f(s(\mathbf{k})))_1
$$

$$
= \sum_{s \in S(\mathbf{k})} (\text{res}_k(f)(s))_1.
$$

5.5. Probability spaces. Let S be an admissible $\mathbb Z$ set. Then, it follows from Lemma [5.2.3](#page-35-3) that

$$
[S] = Z_S(t) \in 1 + t\mathbb{C}[[t]] = W(\mathbb{C}) = \prod_{\geq 1} \mathbb{C}
$$

is invertible if and only if $S(1) \neq \emptyset$ (since this implies $S(k) \neq \emptyset$ for all $k \geq 1$). In this case, we can define

$$
\mathbb{E}: C(S, W(\mathbb{C})) \to W(\mathbb{C}), E[f] = \frac{\int_S f}{[S]}.
$$

By Example [5.3.4,](#page-37-1) this is an expectation (i.e. it sends 1 to 1) and is $W(\mathbb{C})$ -linear. Note that, for $k \geq 1$, the projection $\mathbb{E}_k = \frac{(\int_S f)_k}{\#S(k)}$, so that we can define \mathbb{E}_k as long as $S(\mathbf{k}) \neq \emptyset$.

Using the restriction maps of Proposition [5.4.1,](#page-38-1) we can relate the projections of the expectation to probability on a the finite set of order k points of S :

Lemma 5.5.1. Let S be an admissible Z-set and $k \geq 1$. If $S(\mathbf{k}) \neq \emptyset$, then the map res_k of Proposition [5.4.1](#page-38-1) induces a map of λ -probability spaces

$$
(C(S, W(\mathbb{C})), \mathbb{E}_k) \to (C(S(\mathbf{k}), W(\mathbb{C}))^{(k)}, \mathbb{E}_1).
$$

Proof. Immediate by Proposition [5.4.1](#page-38-1) □

5.6. Varieties, ℓ -adic sheaves, and L-functions. Let \mathbb{F}_q be a finite field of order q, fix an algebraic closure $\overline{\mathbb{F}}_q/\mathbb{F}_q$, and let S/\mathbb{F}_q be a variety. For ℓ a prime coprime to q, we write $K_0(\text{Cons}(S, \mathbb{Q}_\ell))$ for the Grothendieck ring of constructible ℓ -adic sheaves on S. Any element can be written as a formal sum of the extensions by zero of local systems on locally closed subsets, and this sum is unique up to passing to a common constructible refinement. There is a unique λ -ring structure on $K_0(\text{Cons}(S,\mathbb{Q}_\ell))$ such that, for $\mathcal V$ a \mathbb{Q}_ℓ local system on a locally closed $Z \subseteq S$, $h_k \circ [j_! \mathcal{V}] = [j_! \text{Sym}^k \mathcal{V}]$ for $j : Z \hookrightarrow S$ the inclusion. For $f : V \to S$ a morphism of varieties over \mathbb{F}_q , there is a natural map

$$
f^*: K_0(\text{Cons}(S,\overline{\mathbb{Q}}_{\ell})) \to K_0(\text{Cons}(V,\overline{\mathbb{Q}}_{\ell}))
$$

 \Box

induced by pullback of constructible sheaves and an integration over fibers map

$$
f_! : K_0(\text{Cons}(V, \overline{\mathbb{Q}}_\ell)) \to K_0(\text{Cons}(S, \overline{\mathbb{Q}}_\ell))
$$

induced by (derived) proper pushforward of complexes. The map f^* is a map of λ -rings, and by the projection formula, $f_!$ is $K_0(\text{Cons}(V,\mathbb{Q}_\ell))$ -linear.

We fix an isomorphism $\overline{\mathbb{Q}}_{\ell} \to \mathbb{C}$. Then, we obtain a map

$$
\iota_S: K_0(\mathrm{Cons}(S, \overline{\mathbb{Q}}_\ell))) \to C(S(\overline{\mathbb{F}}_q), W(\mathbb{C}))
$$

by sending $[V]$ for V a constructible sheaf to

$$
f_{\mathcal{V}}: s \in S(\overline{\mathbb{F}}_q) \mapsto L_{\mathcal{V}_s}(t) \in 1 + t\mathbb{C}[[t]] = W(\mathbb{C})
$$

where $L_{\mathcal{V}_s}(t)$ is the characteristic polynomial of the (geometric) Frobenius element for $\mathbb{F}_q(s)$ acting on the finite dimensional $\overline{\mathbb{Q}}_\ell$ vector space \mathcal{V}_s , viewed as a C-vector space by the fixed isomorphism. Viewed as a sequence in $\prod_{i\geq 1} \mathbb{C}$, $f_{\mathcal{V}}(s)$ is given by the sequences of traces of powers of the $q^{\deg(s)}$ -power (geometric) Frobenius.

Proposition 5.6.1. The map ι_S is an injective map of λ -rings, and, for each map of varieties $f: V \to S$, $\iota_V \circ f^* = (f|_{V(\overline{\mathbb{F}}_q)})^* \iota_S$ and $\iota_S \circ f_! = (f|_{V(\overline{\mathbb{F}}_q)})_! \circ \iota_V$.

Proof. To see that it is a map of λ -rings, we can note that any $f_{\mathcal{V}}$ as above takes values in $\mathbb{Z}[\mathbb{C}] \subseteq W(\mathbb{C})$ and compare the symmetric power plethysms. That it is injective follows from [\[21,](#page-60-3) Théoréme $(1.1.2)$]. The commutativity for f^* is trivial, and the commutativity for $f_!$ follows, in the usual way, from the Grothendieck-Lefschetz fixed point formula.

Example 5.6.2. Let $f: V \to S$ be a map of varieties. Then, for $[V(\overline{\mathbb{F}}_q)/S(\overline{\mathbb{F}}_q)]$ as in Example [5.3.4,](#page-37-1)

$$
[V(\overline{\mathbb{F}}_q)/S(\overline{\mathbb{F}}_q)] = \iota_S \left(\sum (-1)^i [R^i f_! \overline{\mathbb{Q}}_\ell] \right).
$$

6. An application of Poonen's sieve

In this section, we apply [\[23,](#page-60-5) Theorem 1.3] to compute the λ -distributions of some natural $W(\mathbb{C})$ -valued probability problems on finite sets of homogeneous polynomials over a finite field \mathbb{F}_q . The main result is Proposition [6.2.1.](#page-41-1) This "naïve" point-counting result will be used in conjunction with Lemma [5.5.1](#page-39-2) in the next sections to prove our motivic point-counting results, Theorem [B](#page-7-0) and Theorem [C.](#page-9-0)

6.1. **Setup.** We fix an integer n and let $S_d(\mathbb{F}_q)$ denote the set of degree d homogeneous polynomials in $n+1$ variables x_1, \ldots, x_{n+1} with coefficients in \mathbb{F}_q . For every closed point $P \in |\mathbb{P}_{\mathbb{F}_q}^n|$, we fix a nonnegative integer M_P and a nonvanishing coordinate x_{j_P} . Given $F \in S_d(\mathbb{F}_q)$, we write F_P for the image of $F/x_{j_P}^d$ in $\mathcal{O}_{\mathbb{P}_{\mathbb{F}_q}^n,x}/\mathfrak{m}_x^{M_P+1}$. Given $A \subseteq \mathcal{O}_{\mathbb{P}^n_{\mathbb{F}_q},P}}/\mathfrak{m}_P^{M_P+1}$, we write

$$
\mu_P(A) := \frac{\#A}{\#\left(\mathcal{O}_{\mathbb{P}^n_{\mathbb{F}_q},x}/\mathfrak{m}_x^{M_P+1}\right)}.
$$

We fix, for each P, a non-empty subset $A_P \subseteq \mathcal{O}_{\mathbb{P}_{\mathbb{F}_q}^n,x}/\mathfrak{m}_x^{M_P+1}$ of "allowable" Taylor expansions. We suppose furthermore that these have been chosen so that there exist smooth quasi-projective subvarieties $X_1, \ldots, X_u \subseteq \mathbb{P}^n$ of dimensions $m_i = \dim X_i$, such that, for all but finitely many P, A_P contains F_p for any $F \in S_d(\mathbb{F}_q)$ such that $V(F) \cap X_i$ is smooth of dimension $m_i - 1$ at P for all i.

We let $U_d(\mathbb{F}_q)$ denote the subset of $S_d(\mathbb{F}_q)$ consisting of all $F \in S_d(\mathbb{F}_q)$ such that $F \in A_P$ for all $P \in | \mathbb{P}_{\mathbb{F}_q}^n |$. Note that, by [\[23,](#page-60-5) Theorem 1.3],

$$
\lim_{d \to \infty} \frac{\#U_d(\mathbb{F}_q)}{\#S_d} = \prod_{P \in \mathbb{P}_{\mathbb{F}_q}^n} \mu(A_P).
$$

In particular, for $d \gg 0$, $U_d(\mathbb{F}_q) \neq 0$, and we can consider the λ -probability space

$$
(C(U_d(\mathbb{F}_q), W(\mathbb{C})), \mathbb{E}_1) \text{ for } \mathbb{E}_1(X) = \frac{1}{\#U_d(\mathbb{F}_q)} \sum_{u \in U_d(\mathbb{F}_q)} X(u)_1,
$$

where the subscript 1 denotes projection to the first coordinate in $W(\mathbb{C}) = \prod_{i \geq 1} \mathbb{C}$.

Remark 6.1.1. We could take the $W(\mathbb{C})$ -valued expectation \mathbb{E} everywhere below instead of its first component, but it simplifies the proof to use \mathbb{E}_1 and this will suffice for computing the motivic λ -distributions via Proposition [5.4.1.](#page-38-1)

6.2. **Random variables.** Continuing with the above notation, for each P , we fix a function $X_P : A_P \to W(\mathbb{C}) = 1 + t\mathbb{C}[[t]]$. We may view X_P as a $W(\mathbb{C})$ -valued random variable on the finite set A_P , equipped with the uniform measure. It induces a random variable $X_{P,d}$ on $U_d(\mathbb{F}_q)$ sending F to $X_P(F_P)$.

Proposition 6.2.1. With notation as above, suppose that, for any $M > 0$, for all but finitely many points $P \in \mathbb{P}^n$, X_P factors through $1 + t^M \mathbb{C}[[t]]$. Then, for any $d, \sum_{P \in \mathbb{P}_{\mathbb{F}_q}^n} |X_{P,d} \text{ converges to a function } X_d : U_d(\mathbb{F}_q) \to W(\mathbb{C}), \text{ where we note that }$ the sum is point-wise in $W(\mathbb{C})$, i.e. product of power series in $1 + t\mathbb{C}[[t]]$, and

$$
\lim_{d \to \infty} \mathbb{E}_1[\mathrm{Exp}_{\sigma}(X_d h_1)] = \prod_{P \in |\mathbb{P}_{\mathbb{F}_q}^n|} \mathbb{E}_1[\mathrm{Exp}_{\sigma}(X_P h_1)].
$$

More generally, suppose for each point P given $X_{i,P}$, $i = 1, \ldots m$ satisfying this property, and define $X_{i,P,d}$, $X_{i,d}$ similarly. Then

$$
\lim_{d \to \infty} \mathbb{E}_1[\exp_{\sigma}(X_{1,d}h_1(t_{1,1}, t_{1,2} + \dots) + X_{2,d}h_1(t_{2,1}, t_{2,2} + \dots) + \dots)] =
$$
\n
$$
\prod_{P \in \mathbb{P}_{\mathbb{F}_q}^n} \mathbb{E}_1[\exp_{\sigma}(X_{1,P}h_1(t_{1,1}, t_{1,2} + \dots) + X_{2,P}h_1(t_{2,1}, t_{2,2} + \dots) + \dots)].
$$

Proof. We treat only the case of the σ -moment generating function when given a single random variable at each point; the argument for the joint σ -moment generating function given m random variables is essentially the same.

Let S_m be the finite collection of points such that X_P factors through $1 +$ $t^{m+1}\mathbb{C}[[t]]$ for all $P \notin S_m$. We claim that, for all $f \in \Lambda$ with $\deg f \leq m$ and any $d > 0$, we have an identity of C-valued functions on $U_d(\mathbb{F}_q)$

$$
(f \circ X_d)_1 = (f \circ \sum_{P \in S_m} X_{d,P})_1.
$$

To establish this claim, it suffices to show that, for any power sum symmetric function p_i with $i \leq m$ and any $P \notin S_m$, $(p_i \circ X_P)_1 = 0$ (since this implies that, for all d, $(p_i \circ X_{P,d})_1 = 0$. If we write $X_P = (X_{P,1}, X_{P,2}, \ldots)$, then for $P \notin S_m$, $X_{P,k} = 0$ for all $k \leq m$ since $td \log(1 + t^{m+1} \mathbb{C}[[t]]) \subseteq t^{m+1} \mathbb{C}[[t]]$. Then, as $(p_i \circ X_P)_1 = X_{P,i}$, we find that $(p_i \circ X_P)_1 = 0$ for $P \notin S_m$.

With this claim established, we obtain from Eq. [\(3.2.1.1\)](#page-21-2) that

$$
\mathbb{E}_1[\mathrm{Exp}_{\sigma}(X_d h_1)] \equiv \mathbb{E}_1\left[\mathrm{Exp}_{\sigma}\left(\left(\sum_{P\in S_m} X_{d,P}\right)h_1\right)\right] \mod \Lambda_{\mathbb{C},\deg>m}^{\wedge}.
$$

Similarly, we find that

$$
\prod_{P \in |\mathbb{P}_{\mathbb{F}_q}^n|} \mathbb{E}_1[\mathrm{Exp}_{\sigma}(X_P h_1)] \equiv \prod_{P \in S_m} \mathbb{E}_1[\mathrm{Exp}_{\sigma}(X_P h_1)] \mod \Lambda_{\mathbb{C},\deg>m}^{\wedge}.
$$

Thus, by Lemma [3.4.3,](#page-24-4) it suffices to show that for any P, for the expectation \mathbb{E}_1 , the Λ-distribution of $X_{P,d}$ limits as $d \to \infty$ to the Λ-distribution of X_P , and, for any m, the sequence of collections of random variables $(X_{P,d})_{P \in S_m}$ are asymptotically independent as $d \to \infty$.

To that end, we note that, by [\[23,](#page-60-5) Theorem 1.3], for any choice of subsets $B_P \subseteq$ A_P for each $P \in S_m$,

$$
\lim_{d\to\infty} \mathbb{P}(\substack{F_P \in B_P \text{ for all } P \in S_m \text{ and } P}}{F_P \in A_P \text{ for all } P \notin S_m} \mu_P(B_P) \prod_{P \in S_m} \mu_P(A_P).
$$

It follows that for any collection C-valued functions $g_P, P \in S_m$, the classical random variables $F \mapsto g_P(F_P)$ on $U_d(\mathbb{F}_q)$ with the uniform measure are asymptotically independent, and the distribution of each approaches the distribution of g_P treated as a classical random variable on A_P with the uniform measure. Applying this to g_P of the form $(f_P \circ X_P)_1$ for $f_P \in \Lambda$, we obtain the result.

7. L-functions of characters

In this section we prove Theorem [7.3.2,](#page-48-0) which computes the limiting distributions in Theorem [B,](#page-7-0) and discuss some complements — in particular, in Example [7.2.3,](#page-46-0) we compare the case of quadratic characters to some results in [\[2\]](#page-60-4) (see also Remark [1.3.3\)](#page-8-0). The proof of Theorem [B](#page-7-0) will be completed in Section [9](#page-54-0) by comparing the limit σ -moment generating functions to those in the associated random matrix statistics (Proposition [9.2.1\)](#page-56-1).

To prove Theorem [7.3.2,](#page-48-0) we use Lemma [5.5.1](#page-39-2) to reduce to a problem over a finite probability space, which we analyze using Proposition [6.2.1.](#page-41-1)

7.1. Setup and a note about a normalization. We fix a finite field \mathbb{F}_q , a prime $\ell | q - 1$, and a non-trivial character $\chi : \mu(\mathbb{F}_q) \to \mathbb{C}^\times$. For any ℓ -power free monic $f \in \mathbb{F}_q[x]$, we write χ_f for the character $Gal(\overline{\mathbb{F}_q(x)}/\mathbb{F}_q) \to \mathbb{C}^\times$ obtained by composing χ with the map $Gal(\mathbb{F}_q(x)/\mathbb{F}_q) \to \mu_\ell(\mathbb{F}_q) = \mu_\ell(\mathbb{F}_q(x))$ determined from f by Kummer theory. In other words, $\chi_f(\sigma) = \chi \left(\frac{\sigma(f^{1/\ell})}{f^{1/\ell}} \right)$.

We write $L^{\infty}(\chi_f, s)$ for the associated L-function with the Euler factor at ∞ removed. If $\ell \nmid \deg f$, then χ_f is ramified at ∞ and $L^{\infty}(\chi_f, s) = L(\chi_f, s)$. In general, if we set $\chi(0) = 0$, then we have an Euler product

$$
L^{\infty}(\chi_f, s) = \mathcal{L}(\chi_f, t)|_{t=q^{-s}} \text{ for } \mathcal{L}(\chi_f, t) = \prod_{|z| \in |\mathbb{A}_{\mathbb{F}_q}^1|} \frac{1}{1 - t^{\deg(z)} \chi\left(f(|z|)^{\frac{q^{\deg(|z|)-1}}{\ell}}\right)}.
$$

The function $\mathcal{L}(\chi_f, t)$ is a polynomial with constant term 1 whose degree depends only on the degree of f and, when $\ell \nmid \deg f$, its roots have absolute value $q^{-\frac{1}{2}}$ so that the roots of $\mathcal{L}(\chi_f, tq^{-\frac{1}{2}})$ all have absolute value 1 (this can be deduced from

44 SEAN HOWE

the Weil conjectures for the curve with affine model $y^{\ell} = f(x)$, but we do not actually need this fact for anything below except motivation).

We would like to study the distribution of symmetric functions of these roots using λ-probability. To that end, let us higlight a small issue regarding normalizations: if we write the multiset of roots of $\mathcal{L}(\chi_f, tq^{-\frac{1}{2}})$ as ζ_1, \ldots, ζ_m , then the class $[\zeta_1] + \dots [\zeta_m]$ in $\mathbb{Z}[\mathbb{C}]$ is identified, by our embedding into $W(\mathbb{C}) = 1 + t\mathbb{C}[[t]]$, with $\mathcal{L}(\chi_f, tq^{-\frac{1}{2}})^{-1}$. Thus this is the element of $W(\mathbb{C})$ that will appear below.

7.2. The point-counting result. Continuing with the notation of the previous subsection, we let $U_{d,\ell}(\mathbb{F}_q)$ be the set of degree d monic ℓ -power free polynomials in $\mathbb{F}_q[x]$. From the discussion above, it follows that the random variable on $U_{d,\ell}(\mathbb{F}_q)$ sending f to the multiset of zeroes of $\mathcal{L}(\chi_f, tq^{-\frac{1}{2}})$, viewed as an element of $Z[\mathbb{C}]$ is identified, after embedding $Z[\mathbb{C}]$ in to $W(\mathbb{C})$, with the random variable $X_{\chi,q,d}$ on $U_{d,\ell}(\mathbb{F}_q)$ defined by

$$
X_{\chi,q,d}(f) = \mathcal{L}(\chi_f, tq^{-\frac{1}{2}})^{-1} = \prod_{|z| \in |\mathbb{A}_{\mathbb{F}_q}^1|} \left(1 - (tq^{-\frac{1}{2}})^{\deg(|z|)} \chi\left(f(|z|)^{\frac{q^{\deg(|z|)}-1}{\ell}}\right)\right)
$$

$$
\in 1 + t\mathbb{C}[[t]] = W(\mathbb{C}),
$$

where, for $|z|$ a closed point of $\mathbb{A}^1_{\mathbb{F}_q}$, $f(|z|)$ denotes the image of the f in the residue field $\mathbb{F}_q(|z|)$. In particular, for any $g \in \Lambda$, $\mathbb{E}_1[g \circ X_{\chi,q,d}]$ is the average value on $U_{d,\ell}(\mathbb{F}_q)$ of the symmetric function g evaluated on the roots of $\mathcal{L}(\chi_f, tq^{-\frac{1}{2}})$.

Remark 7.2.1. Let us emphasize this random variable can be defined without knowing that $\mathcal{L}(\chi_f, t)$ is a polynomial or the absolute value of its roots, since the Euler product makes sense immediately as an element of $W(\mathbb{C})!$

Since the infinite product of power series is a sum in $W(\mathbb{C}),$

$$
X_{\chi,q,d} = \sum_{|z| \in |\mathbb{A}_{\mathbb{F}_q}^1|} X_{\chi,q,d,|z|} \text{ where } X_{\chi,q,d,|z|}(f) = 1 - (tq^{-\frac{1}{2}})^{\deg(|z|)} \chi \left(f(|z|)^{\frac{q^{\deg(|z|)}-1}{\ell}}\right)
$$

.

We will compute its asymptotic σ -moment generating function for the expectation \mathbb{E}_1 on $C(U_{d,\ell}(\mathbb{F}_q),W(\mathbb{C}))$ using Proposition [6.2.1.](#page-41-1)

Theorem 7.2.2. Let

$$
c_{\ell} = \frac{[q^{\ell-1}]}{[q]^{\ell-1} + [q]^{\ell-2} + \ldots + 1} \in W(\mathbb{C}) \text{ so that } (c_{\ell})_i = \frac{q^{i(\ell-1)}}{q^{i(\ell-1)} + q^{i(\ell-2)} + \ldots + 1} \in \mathbb{C}.
$$

For χ of order $\ell = 2$, the asymptotic σ -moment generating function of $X_{\chi,q,d}$ is

$$
\lim_{d \to \infty} \mathbb{E}_1[\operatorname{Exp}_{\sigma}(X_{\chi,q,d}h_1)] = \left(\left(1 + c_2 \sum_{k \ge 1} [q^{-k}] e_{2k} \right)^{[q]} \right)_1
$$
\n
$$
= \prod_{P \in |\mathbb{A}_{\mathbb{F}_q}^1} \left(1 + (c_2)_{\deg(P)} \sum_{k \ge 1} q^{-k \deg(P)} (p_{\deg(P)} \circ e_{2k}) \right).
$$

where the subscript denotes projection to the first component in $W(\mathbb{C})$ (note that the power on the right-hand side lies, by definition, in $\Lambda_{W(\mathbb{C})}^{\wedge})$).

Similarly, for χ of order $\ell > 2$ prime, the asymptotic joint σ -moment generating function of $X_{\chi,q,d}$ and its complex conjugate $\overline{X}_{\chi,q,d}$ is

$$
\lim_{d \to \infty} \mathbb{E}_{1}[\text{Exp}_{\sigma}(X_{\chi,q,d}h_{1} + \overline{X}_{\chi,q,d}\overline{h}_{1})] =
$$
\n
$$
\left(\left(1 + c_{\ell} \sum_{\substack{k_{1} \geq 1 \text{ or } k_{2} \geq 1 \\ k_{1} \equiv k_{2} \mod \ell}} (-1)^{\ell k} [q^{-\frac{k_{1} + k_{2}}{2}}] e_{k_{1}} \overline{e}_{k_{2}} \right)^{[q]} \right)_{1} =
$$
\n
$$
\prod_{P \in |\mathbb{A}_{\mathbb{F}_{q}}^{1}} \left(1 + (c_{\ell})_{\deg(P)} \sum_{\substack{k_{1} \geq 1 \text{ or } k_{2} \geq 1 \\ k_{1} \equiv k_{2} \mod \ell}} (-1)^{k_{1} + k_{2}} q^{-\frac{k_{1} + k_{2}}{2}} \deg(P) (p_{\deg(P)} \circ (e_{k_{1}} \overline{e}_{k_{2}})) \right).
$$

Proof. We first note that the first components of the powers appearing in the statement can be expanded as Euler products as claimed by invoking the $i = 1$ case of Proposition [5.2.7](#page-35-0) (to see this, note that, for any $z \in \mathbb{C}$, $[z] = (z, z^2, z^3, \ldots)$ as an element of $W(\mathbb{C}) = \prod_{i \geq \mathbb{C}} \mathbb{C}$, and that $p_{\text{deg }P} \circ f = f(t_1^{\text{deg }P}, t_2^{\text{deg }P}, \ldots)).$

We now explain how to put ourselves into the setup of Proposition $6.2.1$. In the notation of Section [6,](#page-40-0) we work with $n = 1$, i.e. on \mathbb{P}^1 with variables $[x_0 : x_1]$. For $\infty = [0 : 1],$ we set $M_{\infty} = 0$, $j_{\infty} = 0$, and $A_{\infty} = 1 \in \mathbb{F}_q = \mathcal{O}_{\mathbb{P}^1_{\mathbb{F}_q}, \infty}/\mathfrak{m}_{\infty}$. For $P \in |\mathbb{A}_{\mathbb{F}_q}^1| = |\mathbb{P}_{\mathbb{F}_q}^1| \setminus \infty$, we set $M_P = \ell - 1$, $j_{\infty} = 1$, and A_P to be the complement of $0 \in \mathcal{O}_{\mathbb{P}^1_{\mathbb{F}_q},P}/\mathfrak{m}_P^{\tilde{\ell}}$.

The condition at ∞ says that, for $F \in S_d(\overline{\mathbb{F}}_q)$ and $x := \frac{x_0}{x_1}, F_{\infty} \in A_{\infty}$ if and only if F/x_1^d is monic of degree d in $\mathbb{F}_q[x]$. Thus, the set $U_d(\mathbb{F}_q)$ of Section [6](#page-40-0) can be viewed, by $F \mapsto F/x_1^d$, as a subset of the degree d monic polynomials. Then, the conditions at the other primes says exactly that $U_d(\mathbb{F}_q)$ consists of those monic polynomials f that are ℓ th-power free, i.e. $U_d(\mathbb{F}_q) = U_{d,\ell}(\mathbb{F}_q)$.

Now, we define our random variable X_{∞} on A_{∞} to be the constant random variable with value $1 \in 1 + t\mathbb{C}[[t]] = W(\mathbb{C})$ (i.e. the zero element of $W(\mathbb{C})$), and, for any $P \in |A_{\mathbb{F}_q}^1|$, we define a random variable X_P on A_P as follows: first, to simplify notation, we write $q_P = q^{\deg(P)}$ and, for g the germ of a fuction at P, we write $\chi_P(g) := \chi\left(g(P)^{\frac{qp-1}{\ell}}\right)$. Then, we set

$$
X_P(g) = 1 - (q_P^{-\frac{1}{2}} \chi_P(g)) t^{\deg(P)} \in 1 + t\mathbb{C}[[t]] = W(\mathbb{C}).
$$

The random variables $X_{\chi,q,d,P}$ we have defined above are the associated random variables $X_{P,d}$ of Section [6,](#page-40-0) thus by Proposition [6.2.1,](#page-41-1)

$$
\text{for } \ell = 2: \lim_{d \to \infty} \mathbb{E}_1[\exp_{\sigma}(X_{\chi,q,d}h_1)] = \prod_{P \in |\mathbb{P}^1_{\mathbb{F}_q}|} \mathbb{E}_1[\exp_{\sigma}(X_Ph_1)] \text{ and,}
$$
\n
$$
\text{for } \ell > 2: \lim_{d \to \infty} \mathbb{E}_1[\exp_{\sigma}(X_{\chi,q,d}h_1 + \overline{X}_{\chi,q,d}\overline{h}_1)] = \prod_{P \in |\mathbb{P}^1_{\mathbb{F}_q}|} \mathbb{E}_1[\exp_{\sigma}(X_Ph_1 + \overline{X}_P\overline{h}_1)]
$$

The factors at $P = \infty$ on the right are 1. We must compute them for $P \in |\mathbb{A}_{\mathbb{F}_q}^1|$. We first observe that

(7.2.2.1)
$$
(\text{Exp}_{\sigma}(X_{P}h_{1}))_{1} = 1 + \sum_{k \geq 1} (-q_{P}^{-\frac{1}{2}} \chi(g))^{k} p_{\text{deg}(P)} \circ e_{k}
$$

Indeed, when we write an element $\alpha \in W(\mathbb{C}) = \text{Hom}(\Lambda, \mathbb{C})$ as a power series in $1 + t\mathbb{C}[[t]]$, then the coefficient of t^k is $\alpha(h_k)$. On the other hand, when we pass to the first component, we are evaluating on p_1 , and thus

$$
(h_k \circ \alpha)_1 = (h_k \circ \alpha)(p_1) = \alpha(p_1 \circ h_k) = \alpha(h_k).
$$

Since in our power series only the coefficient of $t^{\deg(P)}$ is non-zero, in the expansion of Eq. [\(2.5.2.1\)](#page-17-4)

$$
\operatorname{Exp}_{\sigma}(X_{P}h_1) = \sum_{\tau} (h_{\tau} \circ X_{P}) m_{\tau}
$$

only the terms corresponding to h_{τ} of the form $h_{\deg(P)}^k$ are non-zero. These are the terms corresponding to the monomial symmetric functions

$$
m_{\deg(P)^k} = p_{\deg(P)} \circ m_{1^k} = p_{\deg(P)} \circ e_k
$$

$$
k
$$

where i^k denotes the partition $\overline{(i, \ldots, i)}$. Moreover, the coefficient of such a term is exactly the coefficient of $t^{\deg(P)}$ to the kth power, giving Eq. [\(7.2.2.1\)](#page-44-0).

We now specialize to the case $\ell = 2$. Then,

$$
\mathbb{E}_1[\text{Exp}_{\sigma}(X_{P}h_1)] = \mathbb{E}[(\text{Exp}_{\sigma}(X_{P}h_1))_1]
$$

where the right-hand side is classical expectation (coefficient-wise in the powerseries) for the uniform probability measure on the finite set A_P . Now, for each $\zeta = \pm 1$, $\chi_P(g)$ is equal to ζ with probability

$$
\frac{1}{2}\frac{(q_P - 1)q_P}{q_P^2 - 1} = \frac{1}{2}(c_2)_{\deg(P)},
$$

and the rest of the time it is equal to 0. The expectation of the coefficient $(-q_P^{-\frac{1}{2}} \chi_P(g))^k$ of $p_{\text{deg}(P)} \circ e_k$, $k \geq 1$ in Eq. [\(7.2.2.1\)](#page-44-0) is thus 0 if k is odd and $q_P^{-\frac{k}{2}}$ if k is even. So,

$$
\mathbb{E}_1[\operatorname{Exp}_{\sigma}(X_P h_1)] = 1 + (c_2)_{\deg(P)} \sum_{k \ge 1} q_P^{-k} p_{\deg(P)} \circ e_{2k}.
$$

Plugging this back into the product and recalling $q_P = q^{\deg(P)}$, we find

$$
\lim_{d \to \infty} \mathbb{E}_1[\mathrm{Exp}_{\sigma}(X_{d,q}h_1)] = \prod_{P \in |\mathbb{A}_{\mathbb{F}_q}^1|} \left(1 + (c_2)_{\deg(P)} \sum_{k \ge 1} q^{-k \deg(P)} p_{\deg(P)} \circ e_{2k}\right).
$$

This concludes the proof for $\ell = 2$.

We now treat the case $\ell > 2$, arguing similarly. Then

$$
\operatorname{Exp}_{\sigma}(X_{P}h_1 + \overline{X}_{P}\overline{h}_1) = \operatorname{Exp}_{\sigma}(X_{P}h_1)\operatorname{Exp}_{\sigma}(\overline{X}_{P}\overline{h}_1)
$$

= 1 +
$$
\sum_{k_1 \geq 1 \text{ or } k_2 \geq 1} (-q_P^{-1/2})^{k_1 + k_2} \chi_{P}(g)^{k_1 - k_2} (p_{\deg(P)} \circ e_{k_1} \overline{e}_{k_2}).
$$

Note that, for each ℓ th root of unity ζ , $\chi_P(g)$ is equal to ζ with probability

$$
\frac{1}{\ell} \frac{(q_P - 1)q_P^{(\ell-1)}}{q_P^{\ell} - 1} = \frac{1}{2} (c_{\ell})_{\deg(P)},
$$

and the rest of the time it is equal to 0. The expectation of the coefficient $(-q_P^{-1/2})$ $(P_P^{-1/2})^{k_1+k_2}\chi_P(g)^{k_1-k_2}$ of $(p_{\deg(P)}\circ e_{k_1}\overline{e}_{k_2})$ is thus 0 if $k_1 \not\equiv k_2 \mod l$ (since

the sum of all ℓ th roots of unity is zero), and $(c_{\ell})_{\deg(P)}(-q_P^{-1/2})$ $\binom{-1/2}{P}^{k_1+k_2}$ if $k_1 \equiv k_2$ mod ℓ . So,

$$
\mathbb{E}_{1}[\text{Exp}_{\sigma}(X_{P}h_{1})] = 1 + (c_{\ell})_{\deg(P)} \sum_{\substack{k_{1} \geq 1 \ \text{or} \ k_{2} \geq 1 \\ k_{1} \equiv k_{2} \mod \ell}} (-1)^{k_{1}+k_{2}} q_{P}^{-\frac{k_{1}+k_{2}}{2}} (p_{\deg(P)} \circ e_{k_{1}} \overline{e}_{k_{2}}).
$$

Plugging this back into the Euler product, we conclude. \Box

Example 7.2.3. Consider, in Theorem [7.2.2,](#page-43-1) the case $\ell = 2$ of the quadratic character χ . Then, for d odd, the random variable $[q^{1/2}]X_{\chi,d,q}$ for $X_{\chi,d,q}$ as in Theorem [7.2.2](#page-43-1) is the random variable associated to V_d , the local system of first cohomology of the universal hyperelliptic curve $y^2 = f(x)$ over C_d , the space of monic square free polynomials. We note that, because the Schur polynomials s_{τ} are an orthonormal basis for the Hall inner product, Lemma [3.2.2](#page-21-3) implies that, for any random variable X , its σ -moment generating function can be written as

$$
\mathbb{E}_1[\operatorname{Exp}_{\sigma}(Xh_1)]] = \sum_{\tau} \mathbb{E}_1[s_{\tau} \circ X]s_{\tau}.
$$

We note that $s_{\tau} \circ [\mathcal{V}_d] = [S^{\tau}(\mathcal{V}_d)]$ for S^{τ} the Schur functor associated to τ . Applying this expression of the moment generating function to $[\mathcal{V}_d]$ and using Theorem [7.2.2,](#page-43-1) we find

$$
\sum_{\tau} \lim_{d \to \infty} \left(\mathbb{E}_{1}[S^{\tau}(\mathcal{V}_{d})] \right) s_{\tau} = \left(1 + \frac{[q]}{[q] + 1} \sum_{k \ge 1} e_{2k} \right)^{[q]}
$$

where $S^{\tau}(\bullet)$ is the Schur functor associated to τ . By the Grothendieck-Lefschetz formula, the left-hand side expands $as⁵$ $as⁵$ $as⁵$

$$
\sum_{\tau} \lim_{d \to \infty} \left(\mathbb{E}_{1}[S^{\tau}(\mathcal{V}_{d})] \right) s_{\tau} = \sum_{\tau} \lim_{d \to \infty} \left(\frac{1}{[C_{d}]} \sum_{i=0}^{\infty} (-1)^{i} [H_{c}^{i}(C_{d}, S^{\tau}(\mathcal{V}_{d}))] \right) s_{\tau}.
$$

Using that $\lim_{d\to\infty}$ [C_d]/[q^d] = 1 – [q⁻¹], and noting that, by Poincaré duality,

$$
[H_i(C_d, S^{\tau}(\mathcal{V}_d))] = \frac{[H_c^i(C_d, S^{\tau}(\mathcal{V}_d))]}{[q^d]},
$$

we find that (7.2.3.1)

$$
\sum_{\tau} \lim_{d \to \infty} \left(\sum_{i=0}^{\infty} (-1)^i [H_i(C_d, S^{\tau}(V_d))] \right) s_{\tau} = (1 - [q^{-1}]) \left(1 + \frac{[q]}{[q] + 1} \sum_{k \ge 1} e_{2k} \right)^{[q]}.
$$

We claim this formula is equivalent to a computation of the stable traces of Frobenius for Schur functors applied to V_d given in [\[2,](#page-60-4) Theorem 11.2.3]^{[6](#page-46-2)}: (7.2.3.2)

$$
\sum_{\tau} \lim_{d \to \infty} \left(\sum_{i=0}^{\infty} (-1)^i [H_i(C_d, S^{\tau}(\mathcal{V}_d))] \right) s_{\tau'} = \operatorname{Exp}_{\sigma} \left([q] \operatorname{Log}_{\sigma} \left([q^{-1}] + \sum_{k \geq 0} h_{2k} \right) - 1 \right).
$$

⁵We are projecting everywhere from $W(\mathbb{C})$ to its first component in $\mathbb C$ below, but we drop the subscripts 1 since the formulas are already complicated. Using our motivic results in the next section, one finds the computation hold already in $W(\mathbb{C})!$

⁶Note that to form the exponential in the formula in loc. cit., q^{-1} and q should be viewed as elements of $\Lambda((q^{-1}))$ with $h_k(q^{\pm 1}) = q^{\pm k}$.

48 SEAN HOWE

Here τ' denotes the partition conjugate to τ . Indeed, Eq. [\(7.2.3.2\)](#page-46-3) can be ob-tained from Eq. [\(7.2.3.1\)](#page-46-4) by applying the standard involution ω on symmetric func-tions of Section [2.8](#page-19-1) (which extends naturally to $\Lambda_{W(\mathbb{C})}^{\wedge}$). This is immediate from examining the left-hand side since $\omega(s_{\tau}) = s_{\tau'}$. However, to help understand the formulas, let us also explain how to make this comparison directly on the righthand side. We begin by rewriting the right-hand side of Eq. $(7.2.3.1)$ by substituting $(1 - [q^{-1}]) = \text{Exp}_{\sigma}(-[q^{-1}])$ and expressing the [q]th-power in terms of the σ -exponential:

$$
(1 - [q^{-1}]) \left(1 + \frac{[q]}{[q] + 1} \sum_{k \ge 1} e_{2k} \right)^{[q]} = \operatorname{Exp}_{\sigma} \left([q] \operatorname{Log}_{\sigma} \left(1 + \frac{[q]}{[q] + 1} \sum_{k \ge 1} e_{2k} \right) - [q^{-1}] \right).
$$

We then factor out $\frac{[q]}{[q]+1}$ inside the logarithm we obtain

$$
\operatorname{Exp}_{\sigma}\left([q] \operatorname{Log}_{\sigma}\left(\frac{[q]+1}{[q]} + \sum_{k \ge 1} e_{2k} \right) + [q] \operatorname{Log}_{\sigma}\left(\frac{[q]}{[q]+1} \right) - [q^{-1}] \right)
$$

We now simplify the fraction in the first logarithm term, and make the substitution $\text{Log}_{\sigma}\left(\frac{[q]}{[q]+1]}\right) = -[q^{-1}] + [q^{-2}],$ which follows from the computation

$$
Exp_{\sigma}(-[q^{-1}] + [q^{-2}]) = \frac{Exp_{\sigma}(-[q^{-1}])}{Exp_{\sigma}(-[q^{-2}])} = \frac{1 - [q^{-1}]}{1 - [q^{-2}]} = \frac{1}{1 + [q^{-1}]} = \frac{[q]}{[q] + 1},
$$

to obtain

$$
\mathrm{Exp}_{\sigma}\left([q]\mathrm{Log}_{\sigma}\left([q^{-1}]+\sum_{k\geq 0}e_{2k}\right)-1\right).
$$

Now, applying ω , we obtain the right-hand side of Eq. [\(7.2.3.2\)](#page-46-3): because all of the symmetric functions appearing are of *even* degree, we can commute ω with the σ -exponential and σ -logarithm (see Lemma [2.8.1\)](#page-19-2), and the comparison then follows because $\omega(e_i) = h_i$.

7.3. The motivic result. We now upgrade Theorem [7.2.2](#page-43-1) to a motivic version that encodes the result simultaneously for all finite extensions of \mathbb{F}_q and has the advantage that it is a statement at the level of $W(\mathbb{C})$ rather than a statement after projection to the first component.

We fix a finite field \mathbb{F}_q , an algebraic closure $\overline{\mathbb{F}}_q$, a prime $\ell|q-1$, and a non-trivial character $\chi : \mu_{\ell}(\mathbb{F}_q) \to \mathbb{C}^{\times}$. For $d > 0$, let $U_{d,\ell}(\overline{\mathbb{F}}_q)$ be the admissible Z-set of degree d monic ℓ th-power free polynomials in $\overline{\mathbb{F}}_q[x]$. It admits a degree 1 point, so we have the λ -probability space of Section [5.5,](#page-39-0)

$$
(C(U_{d,\ell}(\overline{\mathbb{F}}_q),W(\mathbb{C})),\mathbb{E})
$$

For $z \in \overline{\mathbb{F}}_q$, we write

$$
\mathcal{X}_{\chi,d} \in C(U_{d,\ell}(\overline{\mathbb{F}}_q) \times \mathbb{A}^1(\overline{\mathbb{F}}_q), W(\mathbb{C}))
$$

for the function sending (f, z) to

$$
\left(1 - (tq^{-\frac{1}{2}})^{[\mathbb{F}_q(z,f) : \mathbb{F}_q(f)]}\chi_f\left(f(z)^{\frac{q^{[\mathbb{F}_q(z,f) : \mathbb{F}_q]-1}}{\ell}}\right)\right) \in 1 + t\mathbb{C}[[t]] = W(\mathbb{C})
$$

We view $\mathcal{X}_{\chi,d}$ as a family of random variables on $U_{d,\ell}(\overline{\mathbb{F}}_q)$ parameterized by $\mathbb{A}^1(\overline{\mathbb{F}}_q)$. The random variable we are interested is then obtained by integrating this family over $\mathbb{A}^1(\overline{\mathbb{F}}_q)$:

$$
X_{\chi,d} := \pi_{1!} \mathcal{X}_{\chi,d} \in C(U_{d,\ell}(\overline{\mathbb{F}}_q), W(\mathbb{C})).
$$

It follows from the definition of π_{1} and the discussion of Section [7.1](#page-42-1) that $X_{\chi,d}$ sends $f \in U_{d,\ell}(\overline{\mathbb{F}}_q)$ to $\mathcal{L}(\chi_f, tq^{-\lfloor \mathbb{F}_q(f) \cdot \mathbb{F}_q \rfloor/2})^{-1} \in 1 + t\mathbb{C}[[t]] = W(\mathbb{C}),$ where χ_f is a representation of $Gal(\overline{\mathbb{F}_q(f)(x)}/\mathbb{F}_q(f)(x))$ for $\mathbb{F}_q(f)$ the field generated by the coefficients of f. As in Section [7.1,](#page-42-1) under the embedding $\mathbb{Z}[\mathbb{C}] \hookrightarrow W(\mathbb{C})$, this is identified with the multiset of zeroes of $\mathcal{L}(\chi_f, tq^{-[\mathbb{F}_q(f):\mathbb{F}_q]/2})$, so that this is precisely the random variable appearing in Theorem [B.](#page-7-0)

Recall the restriction functors res_i of Section [5.4.](#page-38-0)

Lemma 7.3.1. For each $i \geq 1$, $res_i(X_{\chi,d})$ is the random variable X_{χ,d,q^i} on $U_{d,\ell}(\mathbb{F}_{q^i})$ of Theorem [7.2.2](#page-43-1) (where in Theorem 7.2.2 we work over \mathbb{F}_{q^i}).

Proof. This follows from the definition of res_i , and the observation that, if we restrict χ_f to a character of $\mathbb{F}_{q^i}(x)$ rather than $\mathbb{F}_q(f)(x)$, then the L-function changes by an application of $p_{\frac{i}{\deg(x)}} \circ -$ this is a very general statement, but if one prefers to deduce it directly from the Euler product, it suffices to apply Eq. [\(5.1.0.4\)](#page-34-7) to see that the local factors at the closed points of $\mathbb{A}^1_{\mathbb{F}_{q^i}}$ above a closed point $|z|$ of $\mathbb{A}^1_{\mathbb{F}_q(f)}$ give a contribution equal to $p_{\frac{i}{\deg(x)}}$ applied to the local factor at $|z|$.

Combining this with Theorem [7.2.2,](#page-43-1) we obtain

Theorem 7.3.2. Let

$$
c_{\ell} = \frac{[q^{\ell-1}]}{[q]^{\ell-1} + [q]^{\ell-2} + \ldots + 1} \in W(\mathbb{C}).
$$

For χ of order $\ell = 2$, the asymptotic σ -moment generating function of $X_{\chi,d}$ is

$$
\lim_{d \to \infty} \mathbb{E}[\exp_{\sigma}(X_{\chi,d}h_1)] = \left(1 + c_2 \sum_{k \ge 1} [q^{-k}]e_{2k}\right)^{[q]}.
$$

For χ of order $\ell > 2$ prime, the asymptotic joint σ -moment generating function of $X_{\ell,d}$ and its complex conjugate $\overline{X}_{\ell,d}$ is

$$
\lim_{d \to \infty} \mathbb{E}[\exp_{\sigma}(X_{\ell,d}h_1 + \overline{X}_{\ell,d}\overline{h}_1)] =
$$
\n
$$
\left(1 + c_{\ell} \sum_{\substack{k_1 \geq 1 \ \text{or} \ k_2 \geq 1 \\ k_1 \equiv k_2 \mod \ell}} (-1)^{k_1 + k_2} [q^{-\frac{k_1 + k_2}{2}}] e_{k_1} \overline{e}_{k_2} \right)^{[q]}.
$$

Proof. For the case $\ell = 2$, combining Lemma [5.5.1](#page-39-2) and Lemma [7.3.1,](#page-48-1) we find the ith component of the left-hand side is

$$
\lim_{d \to \infty} \mathbb{E}_1[\exp_{\sigma}(X_{2,d,q} : h_1)].
$$

By Theorem [7.2.2,](#page-43-1) this is

$$
\prod_{P \in |\mathbb{A}_{\mathbb{F}_{q^i}}^1|} \left(1 + (c_2)_{i \deg(P)} \sum_{k \ge 1} (q^i)^{-k \deg(P)} (p_{\deg(P)} \circ e_{2k}) \right).
$$

By Proposition [5.2.7,](#page-35-0) this is the ith component of

$$
\left(1 + c_2 \sum_{k \ge 1} [q^{-k}] e_{2k} \right)^{[q]}
$$

This concludes the case $\ell = 2$. For $\ell > 2$, the argument is almost identical.

Remark 7.3.3. In the statement Theorem [7.2.2,](#page-43-1) the moment generating function was also expanded as an Euler product over the closed points of $\mathbb{A}^1_{\mathbb{F}_q}$. As will be explained in [\[6\]](#page-60-14), the powers appearing in Theorem [7.3.2](#page-48-0) can also be expanded as motivic Euler products over $\mathbb{A}^1_{\mathbb{F}_q}$.

8. The L-functions for vanishing cohomology of smooth hypersurface sections

In this section we compute the limiting Λ -distribution in Theorem [C](#page-9-0) (here as Theorem [8.4.2\)](#page-53-2). The remainder of Theorem [C,](#page-9-0) which gives a comparison between this limiting Λ-distribution and the random matrix statistics of Theorem [A](#page-3-0) / Theorem [4.2.1,](#page-25-1) is treated in Section [9](#page-54-0) as Proposition [9.2.2.](#page-58-0)

To compute the limit in question, it would be possible to argue directly with the random variable sending a smooth hypersurface section to the zeroes of the Lfunction of vanishing cohomology as in Section [7.](#page-42-0) However, the argument is much clearer if we instead first prove a result about the geometric random variable which counts points on smooth hypersurface sections, then subtract off the constant part of cohomology to obtain the random variable we are interested in. We prove the point counting version for this geometric random variable in Theorem [8.2.1](#page-51-1) and then the motivic version for this geometric random variable in Theorem [8.3.1.](#page-53-3) Note that part of the point-counting version can also be deduced from [\[16,](#page-60-7) Theorem C] (see Remark [8.2.2\)](#page-52-0), and that Theorem [C/](#page-9-0)Theorem [8.4.2](#page-53-2) implies, by projection to the first component, a much more precise version of [\[16,](#page-60-7) Theorem B] (see Remark [1.3.7\)](#page-10-1).

8.1. **Setup.** We fix a finite field \mathbb{F}_q and an algebraic closure $\overline{\mathbb{F}}_q/\mathbb{F}_q$.

We fix a prime ℓ coprime to q in order to use ℓ -adic étale cohomology, and for any variety V/\mathbb{F}_q , we write $H^i(V)$ as an abbreviation for $H^i_{\text{\'et}}(V_{\overline{\mathbb{F}}_q},\mathbb{Q}_\ell)$, which is equipped with an action of $\mathbb Z$ where the generator 1 acts by the (geometric) q-power Frobenius. We write $[H^i(V)]$ for the associated element of $W(\mathbb{C})$, which, viewed as an element of $1 + t\mathbb{C}[[t]]$ is the characteristic power series of Frobenius, or, viewed as an element of $\prod_{i\geq 1} \mathbb{C}$, is the sequence of traces of powers of Frobenius. If V is smooth and projective of dimension n, $H^i(V)$ is zero for $i < 0$ or $i > 2n$, and, fixing an embedding $\mathbb{Q}_{\ell} \hookrightarrow \mathbb{C}$, by the Weil conjectures the eigenvalues of Frobenius on $H^{i}(V)$ have absolute value $q^{i/2}$. By the Grothendieck-Lefschetz formula, we have

(8.1.0.1)
$$
[V] = \sum (-1)^{i} [H^{i}(V)]
$$

where we recall that we write $[V] = [V(\overline{\mathbb{F}}_q)]$ for the element of $W(\mathbb{C})$ that, viewed $\prod_{i\geq 1} \mathbb{C}$, is the sequence $(\#V(\mathbb{F}_q), \#V(\mathbb{F}_{q^2}), \ldots).$ as an element of $1 + t\mathbb{C}[[t]]$, is the zeta function $Z_V(t)$, and, viewed as element of

Let $Y \subseteq \mathbb{P}^m_{\mathbb{F}_q}$ be an $n+1$ -dimensional smooth closed subvariety. For F a degree d homogeneous polynomial in $m + 1$ variables with coefficients in \mathbb{F}_q , we write $V(F)$ for the vanishing locus of F, a closed subvariety of $\mathbb{P}^m_{\mathbb{F}_q}$. If $V(F)$ intersects Y transversely (in the scheme-theoretic sense; equivalently, the first order Taylor expansion of F at any closed point of Y is non-zero), then $Z = V(F) \cap Y \subseteq Y$ is a smooth projective subvariety of dimension n. By the weak Lefschetz theorem $[10,$ 4.1.6], we have $[H^{i}(Z)]=[H^{i}(Y)]$ for $i < n$ and $H^{n}(Y) \hookrightarrow H^{n}(Z)$. The vanishing cohomology $\mathcal{V}_Z \subseteq H^n(Y)$ is a complement to this inclusion (see [\[16,](#page-60-7) §4] for more details and further references); in particular, we have $[Hⁿ(Z)] = [Hⁿ(Y)] + [V_Z]$ (note that for our purposes below we could also simply define the class $[\mathcal{V}_Z]$ to be $[Hⁿ(Z)] - [Hⁿ(Y)]$ without knowing anything about vanishing cohomology). The remaining degrees of cohomology of Z are also determined by Poincaré duality, or, even more simply, the hard Lefschetz theorem $[10,$ Théorème 4.1.1]:

for
$$
i < n
$$
, $[H^{2n-i}(Z)] = [H^i(Z)][q^{n-i}].$

Combining these computations with Eq. $(8.1.0.1)$, we find that

$$
(8.1.0.2) \qquad [Z] = (-1)^n [\mathcal{V}_Z] + (-1)^n [H^n(Y)] + \sum_{i=0}^{n-1} (-1)^i (1 + [q^{n-i}])[H^i(Y)].
$$

The zeroes of the normalized L-function for vanishing cohomology $\mathcal{L}_Z(tq^{-n/2})$ as appearing in Theorem [C,](#page-9-0) viewed as a multiset of complex numbers and then, via $\mathbb{Z}[\mathbb{C}] \hookrightarrow W(\mathbb{C})$, an element of $W(\mathbb{C})$, is given by $[q^{-n/2}][\mathcal{V}_Z]$. We can solve for this quantity in Eq. $(8.1.0.2)$ to obtain (8.1.0.3)

$$
[q^{-n/2}][\mathcal{V}_Z] = [q^{-n/2}] \left((-1)^n [Z] - [H^n(Y)] - (-1)^n \sum_{i=0}^{n-1} (-1)^i (1 + [q^{n-i}])[H^i(Y)] \right).
$$

In particular, if we vary Z and treat $[\mathcal{V}_Z]$ as a random variable, then the only term on the right that is not constant is $[Z]$. Since any random variable is independent to a constant random variable, we can compute the σ moment generating function of $[q^{-n/2}]V_Z$ as soon as we can compute the σ -moment generating function of $(-1)^n[q^{-n/2}][Z]$, or, since scaling by a power of [q] has a simple effect, just $(-1)^n [Z]$. The difference in parity here accounts for the differences between the odd and even cases in Theorem $C/$ Theorem [8.4.2.](#page-53-2) The strategy is thus to first compute the distribution of $(-1)^n[Z]$ for all d, then apply Eq. [\(8.1.0.3\)](#page-50-1) to deduce the distribution of $(-1)^n[q^{-n/2}][\mathcal{V}_Z]$. As in Section [7,](#page-42-0) to compute the motivic distribution of $(-1)^n[Z]$ we first make a computation on finite probability spaces using Proposition [6.2.1](#page-41-1) and then apply Lemma [5.5.1](#page-39-2) and Proposition [5.2.7](#page-35-0) to obtain the motivic distribution.

Remark 8.1.1. For a random variable X , we do not have a nice way to extract the σ -moment generating function for $-X$ from the σ -moment generating function for X. Indeed, by Lemma [2.5.3,](#page-17-5) to obtain the σ -moment generating function for $-X$, one needs to know the value of the Λ -distribution on the elementary symmetric functions e_{τ} , however, by Lemma [3.2.2,](#page-21-3) this amounts to computing the inner products with the e_{τ} , and it is not clear to us how to do this in a useful way in general. Even for a binomial random variable, we do not see how to do it without using a stronger notion of independence as discussed in Remark [3.3.3](#page-23-0) that allows to make the interpretation as a sum of independent Bernoulli random variables parameterized by an admissible Z-set literal. Thus below we make the computations for both $[Z]$ and $-[Z]$ at the same time; fortunately neither is more difficult than the other (and moreover both are even simpler than the analogous computations in the proof of Theorem [7.2.2!](#page-43-1)).

8.2. The geometric random variable: point-counting. We fix an integer m and a smooth closed subvariety $Y \subseteq \mathbb{P}^m_{\mathbb{F}_q}$. We let $U_d(\mathbb{F}_q)$ be the set degree d homogeneous equations F in $m + 1$ variables such that $V(F)$ intersects Y transversely. By [\[23,](#page-60-5) Theorem 1.1], $U_d(\mathbb{F}_q) \neq \emptyset$ for $d \gg 0$, so we can consider the λ -probability space

$$
(C(U_d(\mathbb{F}_q),W(\mathbb{C})),\mathbb{E}_1).
$$

We write $Z_{q,d}$ for the random variable on $U_d(\mathbb{F}_q)$ that sends F to $[V(F) \cap Y] \in W(\mathbb{C}),$ i.e. to $Z_{V(F)\cap Y}(t) \in 1 + t\mathbb{C}[[t]] = W(\mathbb{C}).$

Theorem 8.2.1. With notation as above and denoting with a subscript 1 the projection from $W(\mathbb{C}) = \prod_{i \geq 1} \mathbb{C}$ to the first component,

$$
\lim_{d \to \infty} \mathbb{E}_1[\exp_{\sigma}(Z_{q,d}h_1)] = \left(\left(1 + \frac{[q]^{n+1} - 1}{[q]^{n+2} - 1} (h_1 + h_2 + h_3 + ...) \right)^{[Y]} \right)_1
$$
\n
$$
= \prod_{P \in |Y|} \left(1 + \frac{q^{(n+1)\deg(P)} - 1}{q^{(n+2)\deg(P)} - 1} p_{\deg(P)} \circ (h_1 + h_2 + h_3 + ...) \right)
$$

and

$$
\lim_{d \to \infty} \mathbb{E}_1[\exp_{\sigma}(-Z_{q,d}h_1)] = \left(\left(1 + \frac{[q]^{n+1} - 1}{[q]^{n+2} - 1}(-e_1 + e_2 - e_3 + ...) \right)^{[Y]} \right)_1
$$
\n
$$
= \prod_{P \in |Y|} \left(1 + \frac{q^{(n+1) \deg(P)} - 1}{q^{(n+2) \deg(P)} - 1} p_{\deg(P)} \circ (-e_1 + e_2 - e_3 + ...) \right)
$$

Proof. The identity of the first components of the powers and the claimed Euler products follows from Proposition [5.2.7.](#page-35-0) To compute the expectations, we will use Proposition [6.2.1.](#page-41-1)

We explain how to put ourselves into the situation of Proposition [6.2.1.](#page-41-1) In the notation of Section [6,](#page-40-0) for each $P \in | \mathbb{P}_{\mathbb{F}_q}^m |$, we take $M_P = 1$ so that we are looking at first-order Taylor expansions; the choice of jp to fix a trivializing coordinate x_{jp} does not effect the definition of anything below so we ignore it. For $P \notin |Y|$ we take $A_P = \mathcal{O}_{\mathbb{P}_{\mathbb{F}_q}^m,P}/\mathfrak{m}_P^2$ to be all first order expansions and for $P \in |Y|$ we take A_P to be the pre-image in $\mathcal{O}_{\mathbb{P}_{\mathbb{F}_q}^m,P}/\mathfrak{m}_P^2$ of the complement of 0 in $\mathcal{O}_{Y,P}/\mathfrak{m}_P^2$. For $P \notin |Y|$ we take X_P the be the constant random variable with value $0 \in W(\mathbb{C})$ (i.e. $1 \in 1 + t\mathbb{C}[[t]])$ and for $P \in |Y|$ we define,

$$
X_P(g) = \begin{cases} \frac{1}{1 - t^{\deg(P)}} \in 1 + t\mathbb{C}[[t]] = W(\mathbb{C}) & \text{if } g(p) = 0, \text{ i.e. } g \in \mathfrak{m}_P/\mathfrak{m}_P^2\\ 1 \in 1 + t\mathbb{C}[[t]] = W(\mathbb{C}) & \text{otherwise.} \end{cases}
$$

Then, for $X_{P,d}$ the associated random variables as in Section [6,](#page-40-0) we have

$$
Z_{q,d} = \sum_{P \in |\mathbb{P}_{\mathbb{F}_q^n}^n|} X_{P,d} = \sum_{P \in |Y|} X_{P,d}.
$$

Indeed: the sum is in $W(\mathbb{C})$ so is multiplication of power series in $1+t\mathbb{C}[[t]]$ and thus, when evaluating on F this is just the usual Euler product expansion of $Z_{q,d}(F)$ = $Z_{V(F)\cap Y}(t)$. Then, Proposition [6.2.1](#page-41-1) implies

$$
\lim_{d\to\infty} \mathbb{E}[Z_{q,d}] = \prod_{P \in \mathbb{P}_{\mathbb{F}_q}^n} \mathbb{E}[\exp_{\sigma}(X_P h_1)] = \prod_{P \in |Y|} \mathbb{E}[\exp_{\sigma}(X_P h_1)].
$$

Arguing as in the proof of Theorem [7.2.2,](#page-43-1) we have

$$
(\text{Exp}_{\sigma}(X_{P}h_{1}))_{1} = \begin{cases} 1 + p_{\text{deg}(P)} \circ (h_{1} + h_{2} + h_{3} + \dots) & \text{if } g(p)=0\\ 1 & \text{otherwise.} \end{cases}
$$

Indeed, the argument given in the proof of Theorem [7.2.2](#page-43-1) shows that $(h_k \circ \frac{1}{1-t^{\deg(P)}})_1$ is 1 if deg(P)|k and 0 otherwise, so that in Eq. [\(2.5.2.1\)](#page-17-4), only the terms m_{τ} with τ a multiple of deg(P) survive, and each with coefficient 1; this gives exactly the series written. Since $g(p) = 0$ with probability $\frac{q^{(n+1)\deg(P)}-1}{q^{(n+2)\deg(P)}-1}$ $\frac{q^{(n+2)\deg(Y)-1}}{q^{(n+2)\deg(P)}-1}$, taking expectation and passing to the product we obtain

$$
\lim_{d \to \infty} \mathbb{E}[Z_{q,d}] = \prod_{P \in |Y|} \left(1 + \frac{q^{(n+1)\deg(P)} - 1}{q^{(n+2)\deg(P)} - 1} p_{\deg(P)} \circ (h_1 + h_2 + h_3 + \ldots) \right).
$$

This concludes the computation for $Z_{q,d}$.

To treat $-Z_{q,d}$, we argue in exactly the same way using the sum

$$
-Z_{q,d} = \sum_{P \in |Y|} -X_{P,d}.
$$

Then, since, for $P \in |Y|$,

$$
-X_P(g) = \begin{cases} 1 - t^{\deg(P)} & \text{if } g(p) = 0, \text{ i.e. } g \in \mathfrak{m}_P/\mathfrak{m}_P^2 \\ 1 & \text{otherwise.} \end{cases}
$$

arguing as in the positive case above (cf. also the proof of Theorem [7.2.2](#page-43-1) where a similar minus sign appears), we find

$$
(\text{Exp}_{\sigma}(-X_{P}h_{1}))_{1} = \begin{cases} 1 + p_{\text{deg}(P)} \circ (-e_{1} + e_{2} - e_{3} + \dots) & \text{if } g(p)=0\\ 1 & \text{otherwise.} \end{cases}
$$

Passing to expectation and taking the product, we conclude exactly as in the positive case. \Box

Remark 8.2.2. We can also deduce the computation of $\lim_{d\to\infty} \mathbb{E}_1[\text{Exp}_{\sigma}(Z_{q,d}h_1)]$ in Theorem [8.2.1](#page-51-1) from [\[16,](#page-60-7) Theorem C] by an application of Lemma [3.3.1](#page-22-3) — indeed, the statement of the result in loc. cit. can be shown to be equivalent to the computation of the falling moment generating function $\mathbb{E}[(1+t)^{Z_{q,d}}]$ (see Section [3.2\)](#page-21-1). However, the method we use to prove Theorem [8.2.1](#page-51-1) is already closely related to the proof [\[16,](#page-60-7) Theorem C], just made much better organized and more robust via the use of λ-probability. Similarly the method of proof in Theorem [8.2.1](#page-51-1) could also be applied to directly compute the falling moment generating function and, since $Z_{q,d}$ is a "combinatorial random variable," the computation of the local factors at

54 SEAN HOWE

closed points P would be even simpler for the falling moments (see Remark [3.2.5](#page-22-1) for more on this along with an explanation of why we didn't just do this).

8.3. The geometric random variable motivically.

Theorem 8.3.1. Let $Y \subseteq \mathbb{P}_{\mathbb{F}_q}^m$ be an $(n+1)$ -dimensional smooth irreducible closed sub-variety, and let U_d/\mathbb{F}_q be the variety of nonzero homogenenous degree d polynomials F such that $V(F)$ intersects Y transversely. Let Z_d be the $W(\mathbb{C})$ -valued random variable on $U_d(\mathbb{F}_q)$ sending $F \in U_d(\mathbb{F}_q)$ to $Z_{V(F) \cap Y}(t)$, where $V(F) \cap Y$ is a variety over $\mathbb{F}_q(F)$, the extension of \mathbb{F}_q generated by the coefficients of F. Then the sequence of random variables Z_d converge in distribution to a binomial distribution with parameters $N = [Y/\mathbb{F}_q]$ and $p = \frac{[q]^{n+1}-1}{[q]^{n+2}-1}$ $\frac{[q]}{[q]^{n+2}-1}$ as in Definition [3.3.2,](#page-23-1) i.e.

$$
\lim_{d \to \infty} \mathbb{E}[\exp_{\sigma}(Z_d)] = \left(1 + \frac{[q]^{n+1} - 1}{[q]^{n+2} - 1}(h_1 + h_2 + h_3 + ...) \right)^{[Y]}
$$

Moreover, we also have

$$
\lim_{d \to \infty} \mathbb{E}[\exp_{\sigma}(-Z_d)] = \left(1 + \frac{[q]^{n+1} - 1}{[q]^{n+2} - 1}(-e_1 + e_2 - e_3 + ...) \right)^{[Y]}.
$$

Proof. As in the proof of Theorem [7.3.2,](#page-48-0) one computes immediately that $res_i Z_d =$ $Z_{q^i,d}$, for the latter the random variable on $U_d(\mathbb{F}_q^i)$ of Theorem [8.2.1](#page-51-1) for \mathbb{F}_{q^i} . Then, applying Lemma [5.5.1,](#page-39-2) we find

$$
\lim_{d \to \infty} \mathbb{E}_i[\exp_{\sigma}(Z_d h_1)] = \lim_{d \to \infty} \mathbb{E}_1[\exp_{\sigma}(Z_{q^i,d} h_1)].
$$

We then find that this is the *i*th component of the power by comparing the de-scription of the limit as an Euler product in Theorem [8.2.1](#page-51-1) (keeping in mind we are applying it over \mathbb{F}_{q^i} to the description of the *i*th component of the power in Proposition [5.2.7.](#page-35-0) This concludes for Z_d ; for $-Z_d$ we argue the same way. \Box

8.4. The cohomological random variable. For Z_d the $W(\mathbb{C})$ -valued random variable on $U_d(\overline{\mathbb{F}}_q)$ of Theorem [8.3.1,](#page-53-3) we let X_d be the $W(\mathbb{C})$ -valued random variable on $U_d(\overline{\mathbb{F}}_q)$ defined by $(8.4.0.1)$

$$
X_d = [q^{-n/2}] \left((-1)^n Z_d - [H^n(Y)] - (-1)^n \sum_{i=0}^{n-1} (-1)^i (1 + [q^{n-i}])[H^i(Y)] \right)
$$

By Eq. [\(8.1.0.3\)](#page-50-1) and the surrounding discussion this is precisely the random variable sending F to the zeroes of $L_{V(F)\cap Y}(t(\# \mathbb{F}_q(F))^{-n/2})$, where $V(F)\cap Y$ is viewed here as a variety over $\mathbb{F}_q(F)$, the extension generated by the coefficients of F.

Remark 8.4.1. Note that this identification depends crucially on the fact that when we view an element of $W(\mathbb{C})$ such as $[H^{n}(Y)]$ as an element of $C(U_{d}(\overline{\mathbb{F}}_{q}), W(\mathbb{C}))$ it is always via pullback from the point, which includes an application of $p_{\text{deg}(z)} \circ$ at any point z, instead of as the literal constant function! (See Example [5.3.5](#page-37-0) and the warning immediately afterwards).

In particular, X_d is the random variable appearing in the statement of Theorem [C.](#page-9-0) In the present notation, the computation of the limit in Theorem [C](#page-9-0) becomes: Theorem 8.4.2. With notation as above,

$$
\lim_{d \to \infty} \mathbb{E}[\exp_{\sigma}(X_d h_1)] = (1 + p \sum_{i \ge 1} [q^{-in/2}] \epsilon^i f_i]^{[Y/\mathbb{F}_q]} \cdot \exp_{\sigma}(\mu h_1)
$$

where $f_i = e_i$ and $\epsilon = -1$ if n is odd and $f_i = h_i$ and $\epsilon = 1$ if n is even,

$$
p = \frac{[q]^{n+1} - 1}{[q]^{n+2} - 1}, \text{ and } \mu = -\epsilon \left(\sum_{i=0}^{n-1} (-1)^i \left([q^{\frac{-n}{2}}] + [q^{\frac{n-2i}{2}}] \right) [H^i(Y)] \right) - [q^{-n/2}] [H^n(Y)].
$$

Proof. We simplify the expression Eq. [\(8.4.0.1\)](#page-53-4) to

$$
X_d = \epsilon[q^{-n/2}]Z_d + \mu
$$

Since μ is a constant and constant random variables are independent to any other random variable, by Lemma [3.2.4](#page-22-4) we have

$$
\mathbb{E}[\exp_{\sigma}(X_d h_1)] = \mathbb{E}\left[\exp_{\sigma}([q^{-n/2}]\epsilon Z_d)\right] \exp_{\sigma}(\mu h_1).
$$

The computation of the limit then follows from Theorem [8.3.1](#page-53-3) (since scaling a random variable by [z] just changes the degree d part of its σ -moment generating function by $[z]^d$).

9. Comparison of L-function statistics and random matrix statistics

In this section we complete the proofs of Theorem [B](#page-7-0) and Theorem [C](#page-9-0) by comparing the limiting expressions for the σ -moment generating functions obtained in Theorem [7.3.2](#page-48-0) and Theorem [8.4.2](#page-53-2) to the random matrix statistics of Theorem [A](#page-3-0) / Theorem [4.2.1](#page-25-1) (and Example [1.2.1\)](#page-5-1).

As in Remark [1.3.6,](#page-10-0) we note that, in the usual approach to comparing random matrix distributions to L -functions as in [\[19\]](#page-60-1), one would first apply Deligne equidistribution to show that, for a fixed degree d and varying over \mathbb{F}_{q^n} as $n \to \infty$, one obtains an equidistribution of conjugacy classes in the monodromy group, and then afterwards take the degree d to ∞ . In particular, the large q equidistribution turns the second limit in d into a problem purely in the study of random matrices. In our results the order of the limits is exchanged: in Theorem [7.3.2](#page-48-0) and Theorem [8.4.2](#page-53-2) we have taken the limit as $d \to \infty$ for a fixed q, and we would like to know what happens if afterwards we take the limit over q^n as $n \to \infty$.

Now, because we have expressed our results using $W(\mathbb{C})$, the limiting formulas actually include the relevant information over \mathbb{F}_{q^n} for any n by projection to the nth coordinate. A convenient way to express the rate of convergence to random matrix statistics as $q \to \infty$ is thus to work in the sub- λ -ring $W(\mathbb{C})^{\text{bdd}}$ consisting of elements whose components are uniformly bounded. In Section [9.1](#page-55-0) we introduce this ring, establish some basic lemmas that will be useful for establishing congruences of moment generating functions, and explain the relation to big O-notation.

In the case of families of idèle class characters as in Theorem [B,](#page-7-0) we show in Proposition [9.2.1](#page-56-1) that our large d limits agree with the random matrix distributions modulo $[q^{-1}]\Lambda_{W(\mathbb{C})^{\mathrm{bdd}}}^{\wedge}$. This is equivalent to saying that, for any symmetric function $f \in \Lambda$, if over \mathbb{F}_{q^n} we take the large d-limit of the average value of f evaluated on the normalized roots of the L-functions in the family, then the result agrees with the relevant random matrix distribution up to $O(q^{-n})$. For vanishing cohomology as in Theorem [C](#page-9-0) we obtain in Proposition [9.2.2](#page-58-0) a congruence modulo $[q^{-\frac{1}{2}}] \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}},$ which is equivalent to agreement with the random matrix distribution up to $O(q^{-\frac{1}{2}})$.

9.1. The λ -ring $W(\mathbb{C})^{\text{bdd}}$. Let

$$
W(\mathbb{C})^{\text{bdd}} := \{ (a_1, a_2, \ldots) \in W(\mathbb{C}) = \prod_{i \ge 1} \mathbb{C} \text{ s.t. } \exists M > 0 \text{ s.t. } |a_i| \le M \,\forall i. \}
$$

Then $W(\mathbb{C})^{\text{bdd}}$ is evidently a sub-ring of $W(\mathbb{C})$. It is also a sub- λ ring: because it is a \mathbb{O} -algebra, it suffices to show that it is preserved by power sum plethysms $p_i \circ$. Since $p_i \circ (a_1, a_2, \ldots) = (a_i, a_{2i}, \ldots)$, the boundedness property is preserved.

The following example shows that congruences can encode big O asymptotics.

Example 9.1.1. Suppose $z \in \mathbb{C}$ with $0 < |z| < 1$. Then $[z] = (z, z^2, z^3, \ldots)$ is contained in $W(\mathbb{C})^{\text{bdd}}$, but $[z^{-1}]$ is evidently not! Thus $[z]W(\mathbb{C})^{\text{bdd}} \subseteq W(\mathbb{C})^{\text{bdd}}$ is a proper ideal; we can also view it as a subgroup of $W(\mathbb{C})$ under addition. In particular, if (a_1, a_2, \ldots) and (b_1, b_2, \ldots) are elements of $W(\mathbb{C})$, then

$$
(a_1, a_2, \ldots) \equiv (b_1, b_2, \ldots) \mod [z]W(\mathbb{C})^{\text{bdd}}
$$

if and only if $a_i = b_i + O(z^i)$, i.e. if and only if there exists a constant $M > 0$ such that $|a_i - b_i| \le M |z^i|$ for all i (or, equivalently by changing M, for $i \gg 0$).

The point of stating our big O asymptotics in this way is that it is relatively straightforward to understand how such congruences interact with the plethystic exponential and logarithm. In particular, we have the following analogs of the usual first order approximations for the classical exponential and logarithm:

Proposition 9.1.2. Let $z \in \mathbb{C}$ with $0 < |z| < 1$.

(1) Suppose $f, g \in W(\mathbb{C})^{\text{bdd}}[[\underline{t}_{\mathbb{N}}]]$ have constant term 0. If

$$
f \equiv g \mod [z] \cdot W(\mathbb{C})^{\text{bdd}}[[\underline{t}_{\mathbb{N}}]],
$$

then

$$
\operatorname{Exp}_{\sigma}(f) \equiv \operatorname{Exp}_{\sigma}(g) \mod [z] \cdot W(\mathbb{C})^{\text{bdd}}[[\underline{t}_{\mathbb{N}}]] \ and
$$

$$
\operatorname{Log}_{\sigma}(1+f) \equiv \operatorname{Log}_{\sigma}(1+g) \mod [z] \cdot W(\mathbb{C})^{\text{bdd}}[[\underline{t}_{\mathbb{N}}]].
$$

(2) Suppose $a \in W(\mathbb{C})^{\text{bdd}}[[\underline{t}_{\mathbb{N}}]]$ has constant term zero, and $a \in [z] \cdot W(\mathbb{C})^{\text{bdd}}[[\underline{t}_{\mathbb{N}}]].$ Then

$$
\operatorname{Exp}_{\sigma}(a) \equiv 1 + a \mod [z^2] \cdot W(\mathbb{C})^{\text{bdd}}[[\underline{t}_{\mathbb{N}}]] \text{ and}
$$

$$
\operatorname{Log}_{\sigma}(1 + a) \equiv a \mod [z^2] \cdot W(\mathbb{C})^{\text{bdd}}[[\underline{t}_{\mathbb{N}}]].
$$

In particular, the same statements hold also in $\Lambda_{W(\mathbb{C})^{\text{bdd}}}^{\wedge}$ or $(\Lambda \otimes \Lambda)_{W(\mathbb{C})}^{\wedge}$.

Proof. The "in particular" at the end holds given the first statements because, for any λ -ring R, we can view Λ_R^{\wedge} or $(\Lambda \otimes \Lambda)_R^{\wedge}$ as a sub- λ ring of $R[[t_{\mathbb{N}}]]$ as in Section [2.](#page-12-0) To verify the exponential case of (1), note that

$$
Exp_{\sigma}(g) = Exp_{\sigma}(f)Exp_{\sigma}(g - f).
$$

Thus, taking $a = g - f$, it suffices to verify that if $a \in [z] \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}}$ has constant term zero then $Exp_{\sigma}(a) \in 1 + [z] \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}}$. This follows from the exponential case

of (2), which we verify now: if we write $a = [z]b$, then using Lemma [2.5.4](#page-17-1) we find

$$
\begin{aligned} \operatorname{Exp}_{\sigma}(a) &= \operatorname{Exp}_{\sigma}([z]b) \\ &= \prod_{i\geq 1} \operatorname{exp}\left(\frac{p_i \circ ([z]b)}{i}\right) \\ &= (1+a+[z]^2 \frac{b^2}{2} + \ldots) \prod_{i\geq 2} \left(1+[z]^i \left(\frac{p_i \circ b}{i}\right) + [z]^{2i} \frac{\left(\frac{p_i \circ b}{i}\right)^2}{2} + \ldots\right) \\ &\equiv (1+a) \mod [z]^2 \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}} .\end{aligned}
$$

To verify the logarithmic case of (1), note that

$$
Log_{\sigma}(1 + f) - Log_{\sigma}(1 + g) = Log_{\sigma}((1 + f)/(1 + g)).
$$

Since $(1 + f)/(1 + g) \in 1 + \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}},$ by taking $a = (1 + f)/(1 + g) - 1$, it suffices to verify that if $a \in [z] \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}}$ then $\log_{\sigma}(1+a) \in [z] \Lambda_{W(\mathbb{C})^{\text{bdd}}}$. This follows from the logarithmic case of (2), which we verify now: if we write $a = [z]b$, then using Lemma [2.5.6:](#page-18-4)

$$
Log_{\sigma}(1 + a) = Log_{\sigma}([z]b) = \sum_{i \ge 1} \ell_i \circ ([z]b)
$$

= $a + \sum_{i \ge 2} \frac{-1}{i} \sum_{d|i} \mu(d)[z]^i (-p_d \circ b)^{i/d}$
\equiv a mod $[z]^2 \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}}$.

 \Box

9.2. Computations.

Proposition 9.2.1. For ℓ a prime, let

$$
c_{\ell} = \frac{[q^{\ell-1}]}{[q]^{\ell-1} + [q]^{\ell-2} + \ldots + 1} \in W(\mathbb{C})^{\text{bdd}}.
$$

Then,

• Taking
$$
\ell = 2
$$
,

$$
\left(1 + c_2 \sum_{k \ge 1} [q^{-k}] e_{2k} \right)^{[q]} \equiv \text{Exp}_{\sigma}(e_2) \mod [q^{-1}] \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}}
$$

where the left-hand side is as in the quadratic limit of σ -moment generating functions in Theorem $B /$ $B /$ Theorem [7.3.2](#page-48-0) and the right-hand side is as in the symplectic case (part (2)) of Theorem [A](#page-3-0) / Theorem [4.2.1.](#page-25-1)

• For $\ell > 2$,

$$
\left(1 + c_{\ell} \sum_{\substack{k_1 \geq 1 \text{ or } k_2 \geq 1 \\ k_1 \equiv k_2 \mod \ell}} (-1)^{k_1 + k_2} [q^{-\frac{k_1 + k_2}{2}}] e_{k_1} \overline{e}_{k_2} \right)^{[q]}
$$

$$
\equiv \text{Exp}_{\sigma}(h_1 \overline{h}_1) \mod [q^{-1}] \Lambda_{W(\mathbb{C})^{\text{bdd}}}^{\wedge}
$$

where we note the left-hand side is as in the non-quadratic limit of joint σ moment generating functions in Theorem $B /$ $B /$ Theorem [7.3.2](#page-48-0) and the righthand side is as in the unitary case (part (4)) of Theorem [A](#page-3-0) / Theorem [4.2.1.](#page-25-1)

Proof. We first treat the $\ell = 2$ case. Then, by the definition of powers (Section [2.6\)](#page-18-0),

$$
\left(1+c_2\sum_{k\geq 1}[q^{-k}]e_{2k}\right)^{[q]}=\mathrm{Exp}_{\sigma}\left([q]\mathrm{Log}_{\sigma}\left(1+c_2\sum_{k\geq 1}[q^{-k}]e_{2k}\right)\right).
$$

Note that $c_2 \equiv 1 \mod [q^{-1}]$. Thus $c_2 \sum_{k \geq 1} [q^{-k}]e_{2k} \equiv [q^{-1}]e_2 \mod [q^{-2}]$. Using both parts of Proposition [9.1.2,](#page-55-1) we find

$$
\operatorname{Log}_{\sigma} \left(1 + c_2 \sum_{k \ge 1} [q^{-k}] e_{2k} \right) \equiv \operatorname{Log}_{\sigma} (1 + [q^{-1}] e_2) \equiv [q^{-1}] e_2 \mod [q^{-2}].
$$

Thus $[q] \text{Log}_{\sigma} \left(1 + c_2 \sum_{k \geq 1} [q^{-k}] e_{2k} \right)$ is in $W(\mathbb{C})^{\text{bdd}}$ and is $\equiv e_2 \mod [q^{-1}]$. Exponentiating, and applying Proposition [9.1.2-](#page-55-1)(1), we find

$$
\left(1 + c_2 \sum_{k \ge 1} [q^{-k}] e_{2k} \right)^{[q]} \equiv \operatorname{Exp}_{\sigma}(e_2) \mod [q^{-1}].
$$

We now treat the $\ell > 2$ case. By the definition of powers (Section [2.6\)](#page-18-0),

$$
\left(1 + c_{\ell} \sum_{\substack{k_1 \geq 1 \text{ or } k_2 \geq 1 \\ k_1 \equiv k_2 \mod \ell}} (-1)^{k_1 + k_2} [q^{-\frac{k_1 + k_2}{2}}] e_{k_1} \overline{e}_{k_2} \right)^{[q]} =
$$

$$
\operatorname{Exp}_{\sigma} \left([q] \operatorname{Log}_{\sigma} \left(1 + c_{\ell} \sum_{\substack{k_1 \geq 1 \text{ or } k_2 \geq 1 \\ k_1 \equiv k_2 \mod \ell}} (-1)^{k_1 + k_2} [q^{-\frac{k_1 + k_2}{2}}] e_{k_1} \overline{e}_{k_2} \right) \right).
$$

Note that $c_{\ell} \equiv 1 \mod [q^{-1}]$. Thus,

$$
c_{\ell} \sum_{\substack{k_1 \geq 1 \text{ or } k_2 \geq 1 \\ k_1 \equiv k_2 \mod \ell}} (-1)^{k_1 + k_2} [q^{-\frac{k_1 + k_2}{2}}] e_{k_1} \overline{e}_{k_2} \equiv [q^{-1}] e_1 \overline{e}_1 \equiv [q^{-1}] h_1 \overline{h}_1 \mod [q^{-2}].
$$

Using both parts of Proposition [9.1.2,](#page-55-1) we find

$$
\operatorname{Log}_{\sigma} \left(1 + c_{\ell} \sum_{\substack{k_1 \geq 1 \text{ or } k_2 \geq 1 \\ k_1 \equiv k_2 \mod \ell}} (-1)^{k_1 + k_2} [q^{-\frac{k_1 + k_2}{2}}] e_{k_1} \overline{e}_{k_2} \right) \equiv \operatorname{Log}_{\sigma} (1 + [q^{-1}] h_1 \overline{h}_1) \mod [q^{-2}]
$$

$$
\equiv [q^{-1}] h_1 \overline{h}_1 \mod [q^{-2}].
$$

Multiplying by [q] we get a congruence mod $[q^{-1}]$ of elements in $\Lambda_{W(\mathbb{C})^{\text{bdd}}}^{\wedge}$, and exponentiating and applying Proposition [9.1.2-](#page-55-1)(1), we find

$$
\left(1 + c_{\ell} \sum_{\substack{k_1, k_2 \ge 1 \\ k_1 \equiv k_2 \mod \ell}} (-1)^{k_1 + k_2} [q^{-\frac{k_1 + k_2}{2}}] e_{k_1} \overline{e}_{k_2} \right)^{[q]} \equiv \operatorname{Exp}_{\sigma}(h_1 \overline{h}_1) \mod [q^{-1}].
$$

Proposition 9.2.2. Let Y be a smooth projective variety of dimension $n + 1$. [C](#page-9-0)onsider the quantity appearing as the limit in Theorem $C /$ Theorem [8.4.2,](#page-53-2)

$$
Q:=(1+p\sum_{i\geq 1}[q^{-in/2}]\epsilon^if_i)^{[Y/\mathbb{F}_q]} \cdot \mathrm{Exp}_\sigma(\mu h_1)\in \Lambda_{W(\mathbb{C})}^\wedge
$$

where $f_i = e_i$ and $\epsilon = -1$ if n is odd and $f_i = h_i$ and $\epsilon = 1$ if n is even,

$$
p = \frac{[q]^{n+1} - 1}{[q]^{n+2} - 1}, \text{ and } \mu = -\epsilon \left(\sum_{i=0}^{n-1} (-1)^i \left([q^{\frac{-n}{2}}] + [q^{\frac{n-2i}{2}}] \right) [H^i(Y)] \right) - [q^{-n/2}] [H^n(Y)].
$$

Then $Q \in \Lambda_{W(\mathbb{C})^{\text{bdd}}}^{\wedge}$, and,

- (1) For $n > 0$ even, $Q \equiv \text{Exp}_{\sigma}(h_2) \mod [q^{-\frac{1}{2}}] \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}},$ where we note the right-hand side is as in the orthogonal case (part (1)) of Theorem [A](#page-3-0) / Theorem [4.2.1.](#page-25-1)
- (2) For n odd, $Q \equiv \text{Exp}_{\sigma}(e_2) \mod [q^{-\frac{1}{2}}] \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}},$ where we note the righthand side is as in the symplectic case (part (2)) of Theorem [A](#page-3-0) / Theorem [4.2.1.](#page-25-1)
- (3) For $n = 0$, $Q \equiv \text{Exp}_{\sigma}(h_2 + h_3 + ...)$ mod $[q^{-\frac{1}{2}}] \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}},$ where we note the right-hand side is as in the action of the symmetric group on the its standard irreducible representation as in Example [1.2.1.](#page-5-1)

Proof. We have

$$
Log_{\sigma}(Q) = [Y]Log_{\sigma}(1 + p \sum_{i \ge 1} [q^{-in/2}] \epsilon^{i} f_i) + \mu.
$$

We will analyze this quantity to apply Proposition [9.1.2.](#page-55-1) First, let us note

$$
p = \frac{[q]^{n+1} - 1}{[q]^{n+2} - 1} = \frac{[q^{-1}] - [q^{-(n+2)}]}{1 - [q^{-(n+2)}]} = [q^{-1}] \mod [q^{-(n+2)}].
$$

We first treat the case $n \geq 1$. Then,

$$
p\sum_{i\geq 1} [q^{-in/2}] \epsilon^i f_i \equiv [q^{-1-n/2}] \epsilon f_1 + [q^{-1-n}] f_2 \mod [q^{-(n+3/2)}].
$$

Applying both parts of Proposition [9.1.2,](#page-55-1) we thus find

$$
\begin{aligned} \text{Log}_{\sigma}(1+p\sum_{i\geq 1} [q^{-in/2}] \epsilon^i f_i) &= \text{Log}_{\sigma}(1+[q^{-1-n/2}] \epsilon f_1 + [q^{-1-n}] f_2) \\ &= [q^{-1-n/2}] \epsilon f_1 + [q^{-1-n}] f_2 \mod [q^{-(n+3/2)}]. \end{aligned}
$$

Note that, by the Weil conjectures, $[q^{-i/2}][H^i(Y)] \in W(\mathbb{C})^{\text{bdd}}$. Furthermore, we have $[H^{2(n+1)}(Y)] = [q^{n+1}]$. Writing

$$
[Y] = \sum_i (-1)^i [H^i(Y)]
$$

and combining with the above computations we find that

$$
[Y]\text{Log}_{\sigma}(1+p\sum_{i\geq 1}[q^{-in/2}]\epsilon^i f_i) \equiv f_2 + \left(\epsilon[q^{-1-n/2}]\sum_{i=n+2}^{2n+2}[H^i(Y)]\right)f_1 \mod [q^{-1/2}]\Lambda^{\wedge}_{(W(\mathbb{C})^{\text{bdd}})}.
$$

Rewriting $[H^{n+1+k}(Y)]=[q^k][H^{n+1-k}(Y)]$, we can rewrite this as

$$
[Y]\text{Log}_{\sigma}(1+p\sum_{i\geq 1}[q^{-in/2}]\epsilon^i f_i) \equiv f_2 + \left(\epsilon \sum_{i=0}^n (-1)^i [q^{n/2-i}][H^i(Y)]\right) f_1 \mod [q^{-1/2}]\Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}}.
$$

On the other hand, by the definition of μ , we have

$$
\mu \equiv -\epsilon \left(\sum_{i=0}^{n-1} (-1)^i [q^{n/2-i}] [H^i(Y)] \right) - [q^{-n/2}] [H^n(Y)] \mod [q^{-1/2}] \Lambda_{W(\mathbb{C})^{\text{bdd}}}.
$$

Combining, we find

$$
Log_{\sigma}(Q) = [Y]Log_{\sigma}(1 + p \sum_{i \ge 1} [q^{-in/2}] \epsilon^i f_i) + \mu h_1 \equiv f_2 \mod [q^{-1/2}] \Lambda_{(W(\mathbb{C})^{\text{bdd}})}.
$$

In particular, $[Y]Log_{\sigma}(Q) + \mu h_1 \in W(\mathbb{C})^{bdd}$, and thus by Proposition [9.1.2-](#page-55-1)(1) we can exponentiate to obtain

$$
Q \equiv \text{Exp}_{\sigma}(f_2) \mod [q^{-1/2}]\Lambda_{(W(\mathbb{C})^{\text{bdd}}}.
$$

It remains to treat the case $n = 0$. Note that, in this case, $\mu = -[H^0(Y)] = -1$, $\epsilon = 1$, and $f_i = h_i$. In particular

$$
Log_{\sigma}(Q) = [Y]Log_{\sigma}(1 + p \sum_{i \ge 1} h_i) - h_1.
$$

Since $p \equiv [q^{-1}] \mod [q^{-2}]$, applying both parts of Proposition [9.1.2](#page-55-1) we have

$$
Log_{\sigma}(1 + p \sum_{i \ge 1} h_i) \equiv Log_{\sigma}(1 + [q^{-1}] \sum_{i \ge 1} h_i) \equiv [q^{-1}] \sum_{i \ge 1} h_i \mod [q^{-2}].
$$

Then, expanding $[Y] = [H^0(Y)] - [H^1(Y)] + [H^2(Y)] = [1] - [H^1(Y)] + [q]$ and using the bounds on on $[H^1(Y)]$ as in the case above, we find

$$
Log_{\sigma}(Q) = [Y]Log_{\sigma}(1 + p \sum_{i \ge 1} h_i) - h_1 \equiv \sum_{i \ge 2} h_i \mod [q^{-1/2}] \Lambda^{\wedge}_{(W(\mathbb{C})^{\text{bdd}}}.
$$

In particular, $Log_{\sigma}(Q)$ is in $W(\mathbb{C})^{bdd}$, and exponentiating and apply Proposition $9.1.2-(1)$ $9.1.2-(1)$, we obtain

$$
Q \equiv \operatorname{Exp}_{\sigma} \left(\sum_{i \ge 2} h_i \right) \mod [q^{-1/2}] \Lambda^{\wedge}_{W(\mathbb{C})^{\text{bdd}}}.
$$

PROBABILITY IN λ -RINGS 61

REFERENCES

- [1] Alexis Aumonier and Ronno Das. Homological stability for the space of hypersurfaces with marked points. *arXiv:2312.03355*, 2023.
- [2] Jonas Bergström, Adrian Diaconu, Dan Petersen, and Craig Westerland. Hyperelliptic curves, the scanning map, and moments of families of quadratic l-functions. *arXiv:2302.07664*, 2024.
- Matthew Bertucci. Taylor conditions over finite fields. $arXiv:2412.08744$, 2024.
- [4] Matthew Bertucci and Sean Howe. More λ-distributions of point counting random variables. *In preparation*.
- [5] Margaret Bilu. Motivic Euler products and motivic height zeta functions. *Mem. Amer. Math. Soc.*, 282(1396):v+185, 2023.
- [6] Margaret Bilu, Ronno Das, and Sean Howe. Special values of motivic euler products. *In preparation*.
- [7] Margaret Bilu, Ronno Das, and Sean Howe. Zeta statistics and Hadamard functions. *Adv. Math.*, 407:Paper No. 108556, 68, 2022.
- [8] Margaret Bilu and Sean Howe. Motivic random variables. *In preparation*.
- [9] Alina Bucur and Kiran S. Kedlaya. The probability that a complete intersection is smooth. *J. Th´eor. Nombres Bordeaux*, 24(3):541–556, 2012.
- [10] Pierre Deligne. La conjecture de Weil. II. *Inst. Hautes Etudes Sci. Publ. Math. ´* , (52):137–252, 1980.
- [11] Persi Diaconis and Mehrdad Shahshahani. On the eigenvalues of random matrices. volume 31A, pages 49–62. 1994. Studies in applied probability.
- [12] Daniel Erman and Melanie Matchett Wood. Semiample Bertini theorems over finite fields. *Duke Math. J.*, 164(1):1–38, 2015.
- [13] E. Getzler and M. M. Kapranov. Modular operads. *Compositio Math.*, 110(1):65–126, 1998.
- [14] S. M. Guseĭn-Zade, I. Luengo, and A. Melle-Ernandez. On the power structure over the Grothendieck ring of varieties and its applications. *Tr. Mat. Inst. Steklova*, 258(Anal. i Osob. Ch. 1):58–69, 2007.
- [15] Michiel Hazewinkel. Witt vectors. I. In *Handbook of algebra. Vol. 6*, volume 6 of *Handb. Algebr.*, pages 319–472. Elsevier/North-Holland, Amsterdam, 2009.
- [16] Sean Howe. Motivic random variables and representation stability II: Hypersurface sections. *Adv. Math.*, 350:1267–1313, 2019.
- [17] Sean Howe. Motivic random variables and representation stability I: Configuration spaces. *Algebr. Geom. Topol.*, 20(6):3013–3045, 2020.
- [18] Kurt Johansson. On random matrices from the compact classical groups. *Ann. of Math. (2)*, 145(3):519–545, 1997.
- [19] Nicholas M. Katz and Peter Sarnak. *Random matrices, Frobenius eigenvalues, and monodromy*, volume 45 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1999.
- [20] Hanspeter Kraft and Claudio Procesi. *Classical invariant theory – A primer*. Course notes available at https://dmi.unibas.ch/fileadmin/user_upload/dmi/Personen/Kraft_Hanspeter/Classical_Invariant_Theory.p 1996.
- [21] G. Laumon. Transformation de Fourier, constantes d'équations fonctionnelles et conjecture de Weil. *Inst. Hautes Etudes Sci. Publ. Math. ´* , (65):131–210, 1987.
- [22] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Classic Texts in the Physical Sciences. The Clarendon Press, Oxford University Press, New York, second edition, 2015. With contribution by A. V. Zelevinsky and a foreword by Richard Stanley.
- [23] Bjorn Poonen. Bertini theorems over finite fields. *Ann. of Math. (2)*, 160(3):1099–1127, 2004.
- [24] Richard P. Stanley. *Enumerative combinatorics. Vol. 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999. With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin.
- [25] H. Weyl. *The Classical Groups: Their Invariants and Representations*. Number no. 1, pt. 1 in Princeton Landmarks in Mathematics and Physics. Princeton University Press, 1946.