Mem. S.A.It. Vol. 75, 282 © SAIt 2023

Memorie della



GRB multi-TeV detection: Beyond standard physics?

G. Galanti¹, L. Nava^{2,3}, M. Roncadelli^{4,2}, F. Tavecchio², and G. Bonnoli²

¹ INAF, Istituto di Astrofisica Spaziale e Fisica Cosmica di Milano, Via Alfonso Corti 12, I – 20133 Milano, Italy, e-mail: gam.galanti@gmail.com

² INAF, Osservatorio Astronomico di Brera, Via Emilio Bianchi 46, I – 23807 Merate, Italy

³ INFN, Sezione di Trieste, Via Alfonso Valerio 2, I – 34127 Trieste, Italy

⁴ INFN, Sezione di Pavia, Via Agostino Bassi 6, I – 27100 Pavia, Italy

Received: Day Month Year; Accepted: Day Month Year

Abstract. The recent detection by LHAASO up to 18 TeV of the gamma ray burst GRB 221009A at redshift z = 0.151 challenges standard physics because of the strong absorption due to the extragalactic background light (EBL) for photons with energies above 10 TeV. Emission models partially avoiding EBL absorption proposed to explain such an event are unsatisfactory since they require peculiar and contrived assumptions. By introducing in magnetized media the interaction of photons with axion-like particles (ALPs) – which are a generic prediction of most theories extending the standard model of particle physics towards a more satisfying theory – the detection of GRB 221009A can be naturally explained, thereby providing a strong hint at ALP existence.

Key words. Particle astrophysics - Gamma ray bursts - Particle interactions - Axions

1. Introduction

The detection of the Gamma Ray Burst GRB 221009A at very-high-energy (VHE, $\mathcal{E} > 100 \text{ GeV}$) by the LHAASO Collaboration (2022) with the highest photon energy reaching 18 TeV and less reliably by the Carpet-2 Collaboration (2022) up to 251 TeV challenges standard physics expectations: at the redshift z = 0.151 of GRB 221009A measured by the Stargate Collaboration (2022), photons with energy $\mathcal{E} > 10$ TeV are extremely difficult to detect because of their interaction with the extragalactic background light (EBL) photons (Dwek & Krennrich (2013)).

In particular, the attenuation of photons with $\mathcal{E} = 18 \text{ TeV}$ is at least $O(10^6 - 10^8)$, as

shown by the several existing EBL models (see e.g., Domínguez et al. (2011); Gilmore et al. (2012); Franceschini & Rodighiero (2017); Saldana-Lopez et al. (2021)). The luminosity required to justify the observed VHE photons from GRB 221009A turns out to be excessively high within all conventional GRB models.

In order to find a solution to such a problem, in Galanti et al. (2022a) it is proposed that photons oscillate into axion-like particles (ALPs) in the presence of external magnetic fields permeating the media crossed by the beam (see also Galanti et al. (2022b)). As a result, the effective photon optical depth gets reduced, since when the beam photons behave as ALPs they are *not* EBL absorbed. This scenario has been deeply investigated by us in the last years (Galanti & Roncadelli (2022)).

In Galanti et al. (2022a) we also analyze the effects on GRB 221009A produced by the Lorentz invariance violation (LIV).

2. Standard view

VHE photons originated at cosmological distances suffer absorption caused by their interaction with the EBL, namely photons emitted by stars and possibly reprocessed by dust during the whole Universe history.

All existing EBL models (see e.g., Domínguez et al. (2011); Gilmore et al. (2012); Franceschini & Rodighiero (2017); Saldana-Lopez et al. (2021)) are quite consistent among each other in the optical-ultraviolet range, but they differ in the infrared band which is responsible for photon absorption above ~ 5 TeV.

As based on observations from space, thereby avoiding the zodiacal light problem, the most robust EBL model appears to be that by Saldana-Lopez et al. (2021). Therefore, we hereafter employ the Saldana-Lopez et al. (2021) EBL model which predicts - within conventional physics - a photon survival probability amounting to $P_{\rm CP} \simeq 1 \times 10^{-8}$ at \mathcal{E} = 18 TeV and $P_{\rm CP} \simeq 3 \times 10^{-6}$ at $\mathcal{E} = 15$ TeV. The latter energy value is chosen in order to take into account an uncertainty of (15 - 20)%concerning the detection by the LHAASO Collaboration (2022). Regarding the possible detection by the Carpet-2 Collaboration (2022), the model by Saldana-Lopez et al. (2021) predicts $P_{CP} \sim 0$ both at 251 TeV and at 100 TeV, with the latter energy value chosen to account for an uncertainty of ~ 50%.

Because of the very low values assumed by P_{CP} , photons with $\mathcal{E} > 10 \text{ TeV}$ are strongly absorbed and their detection requires a very high TeV luminosity, which is very difficult to explain within conventional emission models.

3. Axion-like particles

In order to solve the tension between the observation at VHE of GRB 221009A and conventional emission models, we invoke the possibility that photons oscillate into axion-like particles (ALPs).

ALPs are predicted by superstring and superbrane theories (see e.g., Jaeckel & Ringwald (2010); Ringwald (2012)) and turn out to be extremely light neutral pseudo-scalar bosons with mass m_a interacting with photons with coupling $g_{a\gamma\gamma}$ through the Lagrangian

$$\mathcal{L}_{a\gamma\gamma} = -\frac{1}{4} g_{a\gamma\gamma} F_{\mu\nu} \tilde{F}^{\mu\nu} a = g_{a\gamma\gamma} \mathbf{E} \cdot \mathbf{B} a , \quad (1)$$

where *a* represents the ALP field, while $F_{\mu\nu}$ (with dual $\tilde{F}^{\mu\nu}$) is the electromagnetic tensor whose electric and magnetic components are **E** and **B**, respectively. The most solid bound on the ALP parameter space $(m_a, g_{a\gamma\gamma})$ is represented by $g_{a\gamma\gamma} < 0.66 \times 10^{-10} \text{ GeV}^{-1}$ for $m_a < 0.02 \text{ eV}$ at 2σ level by the CAST Collaboration (2017), while the most stringent ones read: (i) $g_{a\gamma\gamma} < 6.3 \times 10^{-13} \text{ GeV}^{-1}$ for $m_a < 10^{-12} \text{ eV}$ by Sisk-Reynés et al. (2022) and (ii) $g_{a\gamma\gamma} < 5.4 \times 10^{-12} \text{ GeV}^{-1}$ for $m_a < 3 \times 10^{-7} \text{ eV}$ by Dessert et al. (2022). The QED vacuum polarization (Heisenberg & Euler (1936); Weisskopf (1936); Schwinger (1951)), and the photon dispersion on the CMB (Dobrynina et al. (2015)) effects must also be considered in this context.

In the presence of an external magnetic field **B**, photons represented by **E** in Eq. (1)can convert into ALPs producing: (i) photon-ALP oscillations (Maiani et al. (1986); Raffelt & Stodolsky (1988)) and (ii) change of the polarization state of photons (Maiani et al. (1986); Raffelt & Stodolsky (1988)). As a result, photon-ALP interaction produces many consequences both on observed astrophysical spectra (see e.g., De Angelis et al. (2007, 2011); Simet et al. (2008); Sánchez-Conde et al. (2009); Tavecchio et al. (2012, 2015); Wouters & Brun (2012); Kohri & Kodama (2017); Galanti et al. (2019, 2020b)) and on the detected photon polarization (see e.g., Jain et al. (2002); Bassan et al. (2010); Agarwal et al. (2011); Payez et al. (2011); Perna et al. (2012); Day & Krippendorf (2018); Galanti (2022, 2023); Galanti et al. (2023a,b)).

In GRB 221009A VHE photons are produced in the GRB jet and they can then oscillate into ALPs inside all crossed magnetized media (the GRB jet, the galaxy hosting the GRB, the extragalactic space, and the Milky Way). As demonstrated in Galanti et al. (2022a), the photon-ALP conversion inside the GRB jet is negligible since the propagation length of the photon-ALP beam inside the source is too small to produce an efficient photon-ALP conversion. Following the results presented in Levan et al. (2023), the galaxy hosting GRB 221009A is a disclike one observed edge-on with the GRB located close to the nuclear region. Therefore, we use previous information to compute the photon-ALP conversion in the host, and we consider the two most likely possibilities concerning the nature of the host galaxy: (i) a typical spiral (see e.g. Beck (2016)) and (ii) a starburst with intermediate features similar to M82 (Thompson et al. (2006); Lopez-Rodriguez et al. (2021)). Photon-ALP conversion inside the host is efficient in the case of both a spiral and a starburst galaxy (see Galanti et al. (2022a)). Concerning photon-ALP interaction inside the extragalactic space, we follow the results reported in Galanti & Roncadelli (2018a,b) by considering both an efficient photon-ALP conversion with an extragalactic magnetic field strength $B_{\text{ext}} = 1 \text{ nG}$ (more likely, see e.g., Rees & Setti (1968); Hoyle (1969); Kronberg et al. (1999); Furlanetto & Loeb (2001)) and a negligible photon-ALP conversion with $B_{\text{ext}} < 10^{-15} \,\text{G}$ (extremely conservative, see e.g., Neronov & Vovk (2010); Durrer & Neronov (2013); Pshirkov et al. (2016)). Finally, we use the magnetic field map by Jansson & Farrar (2012a,b) and the electron number density model by Yao et al. (2017) in order to evaluate the photon-ALP conversion in the Milky Way. Once the transfer matrices of the photon-ALP beam in all the previous regions are evaluated and combined, we obtain the photon survival probability in the presence of photon-ALP oscillations P_{ALP} , as discussed in Galanti et al. (2022a).

In Fig. 1 we plot P_{ALP} at $\mathcal{E} = 15 \text{ TeV}$ as a function of m_a in the range $10^{-12} \text{ eV} \leq m_a \leq 10^{-6} \text{ eV}$ and of $g_{a\gamma\gamma}$ in the interval $10^{-13} \text{ GeV}^{-1} \leq g_{a\gamma\gamma} \leq 10^{-10} \text{ GeV}^{-1}$ by assuming the EBL model of Saldana-Lopez et al.




Fig. 1. Behavior of P_{ALP} at $\mathcal{E} = 15$ TeV as a function of m_a and $g_{a\gamma\gamma}$ by assuming the EBL model of Saldana-Lopez et al. (2021), $B_{\text{ext}} = 1$ nG and a starburst hosting galaxy. ALP bounds are also plotted (CAST Collaboration (2017); Sisk-Reynés et al. (2022); Dessert et al. (2022)). (Credit: Galanti et al. (2022a)).




Fig. 2. Behavior of P_{CP} and of P_{ALP} (for $B_{ext} = 1 \text{ nG}$ and $B_{ext} < 10^{-15} \text{ G}$) versus \mathcal{E} for the case of a starburst hosting galaxy and employing the EBL model of Saldana-Lopez et al. (2021). We assume $m_a = 10^{-10} \text{ eV}$ and $g_{a\gamma\gamma} = 4 \times 10^{-12} \text{ GeV}^{-1}$. (Credit: Galanti et al. (2022a)).

(2021), $B_{\text{ext}} = 1 \text{ nG}$ and a starburst hosting galaxy.

In Fig. 2 we report both P_{CP} and P_{ALP} (for $B_{ext} = 1 \text{ nG}$ and $B_{ext} < 10^{-15} \text{ G}$) versus \mathcal{E} for the case of a starburst hosting galaxy and employing the EBL model of Saldana-Lopez et al. (2021) by assuming $m_a = 10^{-10} \text{ eV}$ and $g_{a\gamma\gamma} = 4 \times 10^{-12} \text{ GeV}^{-1}$, which represent the values of

the ALP parameter space $(m_a, g_{a\gamma\gamma})$ maximizing P_{ALP} at $\mathcal{E} = 15$ TeV, as shown in Fig. 1, within the most stringent ALP bounds (Sisk-Reynés et al. (2022); Dessert et al. (2022)).

Still, even lower $g_{a\gamma\gamma}$ values and different situations – i.e. both a spiral and a starburst hosting galaxy and both the cases $B_{\text{ext}} =$ 1 nG and $B_{\text{ext}} < 10^{-15}$ G – give rise to a P_{ALP} large enough to allow the observability of GRB 221009A above 10 TeV (Galanti et al. (2022a,b)).

4. Lorentz invariance violation

A competitive scenario – with respect to photon-ALP oscillations – to explain the TeV detection of GRB 221009A can be represented by the possibility of Lorentz invariance violation (LIV), predicted by quantum gravity theories, above a threshold energy \mathcal{E}_{LIV} (for a review, see Addazi et al. (2022)). A consequence of LIV is the modification of the photon dispersion relation, reading

$$\mathcal{E}^2 - p^2 = -\frac{\mathcal{E}^{n+2}}{\mathcal{E}^n_{\text{LIV}}},$$
(2)

with \mathcal{E} and p the photon energy and momentum, respectively, and with a resulting modification of the effective photon optical depth (see e.g., Tavecchio & Bonnoli (2016); Galanti et al. (2020a)), which is relevant for the TeV detection of GRB 221009A (Galanti et al. (2022a)).

In Fig. 3 we plot both P_{CP} and the photon survival probability in the presence of LIV effects P_{LIV} by taking $\mathcal{E}_{LIV,n=1} = 3 \times 10^{29}$ eV in the case n = 1 and $\mathcal{E}_{LIV,n=2} = 5 \times 10^{21}$ eV in the case n = 2, which are values within the current LIV bounds (Lang et al. (2019)).

From Fig. 3 we infer that LIV is unable to explain the detection by the LHAASO Collaboration (2022), while it is quite effective to justify that by the Carpet-2 Collaboration (2022), but the latter is less reliable.

5. Discussion and Conclusions

Emission models not invoking new physics – as synchrotron self Compton (SSC) radiation




Fig. 3. Behavior of P_{CP} and of P_{LIV} versus \mathcal{E} . (Credit: adapted from Galanti et al. (2022a)).

and secondary emission from ultra-high energy protons (Mirabal (2023); Gonzalez et al. (2023); Das & Razzaque (2023); Zhao et al. (2023); Sahu et al. (2023)) - can hardly explain the detection of $\mathcal{E} > 10 \,\text{TeV}$ photons from GRB 221009A even assuming ad hoc and often contrived choices of the parameters because of the EBL absorption. So, a tension arises with the LHAASO Collaboration (2022) detection, when realistic EBL models are taken into account. Moreover, realistic EBL absorption models could become even more severe, as indicated by recent results of the JWST, which suggest a higher infrared background level than currently expected (Labbé et al. (2022)).

Instead, the photon-ALP oscillation scenario proposed in Galanti et al. (2022a) (see also Galanti et al. (2022b)) turns out to reduce the EBL absorption largely in the energy region of interest, and is extremely effective in justifying both the LHAASO observation and the possible Carpet 2 detection of GRB 221009A within the most stringent ALP bounds (Sisk-Reynés et al. (2022); Dessert et al. (2022)). The LIV model proposed in Galanti et al. (2022a) explains the less reliable Carpet 2 detection of GRB 221009A, but is ineffective at the lower energies observed by the LHAASO collaboration.

In conclusion, although a firm assessment will be possible only when the spectral data

of the LHAASO detection are known, we have at least a strong indication of the existence of ALPs coming from GRB 221009A (Galanti et al. (2022a)) with parameters compatible with two previous ALP hints (Tavecchio et al. (2012); Galanti et al. (2020b)) and with values which make ALPs good candidates for cold dark matter (Arias et al. (2012)).

Acknowledgements. We thank Felix Aharonian, Patrizia Caraveo, Elisabete de Gouveia Dal Pino, Giovanni Pareschi, Isabella Prandoni and Franco Vazza for discussions and useful information. The work of G. G. is supported by a contribution from the grant ASI-INAF 2015-023-R.1. The work of M. R. is supported by an INFN grant. This work was made possible also by the funding by the INAF Mini Grant 'High-energy astrophysics and axion-like particles', PI: Giorgio Galanti. The work by L. N. is partially supported by the INAF Mini Grant 'Shock acceleration in Gamma Ray Bursts', PI: Lara Nava.

References

- Addazi, A., et al. 2022, Progr. in Part. and Nucl. Phys., 125, 103948
- Agarwal, N., Kamal, A. & Jain, P. 2011, Phys. Rev. D, 83, 065014
- Arias, P., et al. 2012, JCAP, 06, 013
- Bassan, N., Mirizzi, A. & Roncadelli, M. 2010, JCAP, 05, 010
- Beck, R. 2016, A&A Rev., 24, 4
- Carpet-2 Collaboration 2022, ATel #15669, https://astronomerstelegram.org/ ?read=15669
- CAST Collaboration 2017, Nature Physics, 13, 584
- Das, S. & Razzaque, S. 2023, A&A, 670, 12
- Day, F. & Krippendorf, S. 2018, Galaxies, 6, 45
- De Angelis, A., Galanti, G. & Roncadelli, M. 2011, Phys. Rev. D, 84, 105030
- De Angelis, A., Roncadelli, M. & Mansutti, O. 2007, Phys. Rev. D, 76, 121301
- Dessert, C., Dunsky, D. & Safdi, B. R. 2022, Phys. Rev. D, 105, 103034
- Dobrynina, A., Kartavtsev, A. & Raffelt, G. 2015, Phys. Rev. D, 91, 083003
- Domínguez, A., et al. 2011, MNRAS, 410, 2556
- Durrer, R. & Neronov, A. 2013 A&A Rev., 21, 62

- Dwek, E. & Krennrich, F. 2013, Astropart. Phys., 43, 112
- Franceschini, A. & Rodighiero, G. 2017, A&A, 603, 34
- Furlanetto, S. & Loeb, A. 2001, ApJ, 556, 619
- Galanti, G. 2022, Phys. Rev. D,105, 083022
- Galanti, G., 2023, Phys. Rev. D, 107, 043006
- Galanti, G., Nava, L., Roncadelli, M., Tavecchio, F. & Bonnoli, G., 2022a, arXiv:2210.05659
- Galanti, G. & Roncadelli, M. 2018a, Phys. Rev. D, 98, 043018
- Galanti, G. & Roncadelli, M. 2018b, J. High Energy Astrophys., 20, 1
- Galanti, G. & Roncadelli, M. 2022, Universe, 8, 253
- Galanti, G., Roncadelli, M., De Angelis, A. & Bignami, G. F. 2020b, MNRAS, 493, 1553
- Galanti, G., Roncadelli, M. & Tavecchio, F. 2022b, arXiv:2211.06935.
- Galanti, G., Roncadelli, M. & Tavecchio, F. 2023b, arXiv:2301.08204
- Galanti, G., Roncadelli, M., Tavecchio, F. & Costa, E. 2023a, Phys. Rev. D, 107, 103007
- Galanti, G., Tavecchio, F. & Landoni, M. 2020a, MNRAS, 491, 5268
- Galanti, G., Tavecchio, F., Roncadelli, M. & Evoli, C. 2019, MNRAS, 487, 123
- Gilmore, R. C., Somerville, R. S., Primack, J. R. & Domínguez, A. 2012, MNRAS, 422, 3189
- Gonzalez, M. M., et al. 2023, ApJ, 944, 178
- Heisenberg, W. & Euler, H. 1936, Z. Phys., 98, 714
- Hoyle, F. 1969, Nature, 223, 936
- Jaeckel, J. & Ringwald, A. 2010, Ann. Rev. Nucl. Part. Sci., 60, 405
- Jain, P., Panda, S. & Sarala, S. 2002, Phys. Rev. D, 66, 085007
- Jansson, R. & Farrar, G. R. 2012a, ApJ, 757, 14
- Jansson, R. & Farrar, G. R. 2012b, ApJ, 761, L11
- Kohri, K. & Kodama, H. 2017, Phys. Rev. D, 96, 051701
- Kronberg, P. P., Lesch, H. & Hopp, U. 1999, ApJ, 511, 56
- Labbé, I., et al. 2022, arXiv:2207.12446
- Lang, R. G., Martínez-Huerta, H. & de Souza, V. 2019, Phys. Rev. D, 99, 043015

Levan, A. J., et al. 2023, arXiv:2302.07761

- LHAASO Collaboration 2022, GCN Circular n. 32677, https://gcn.gsfc.nasa.gov/ gcn3/32677.gcn3
- Lopez-Rodriguez, E., Guerra, J. A., Asgari-Targhi, M. & Schmelz, J. T. 2021, ApJ, 914, 24
- Maiani, L., Petronzio, R. & Zavattini, E., 1986, Phys. Lett. B, 175, 359
- Mirabal, N. 2023, MNRAS, 519, 85
- Neronov, A. & Vovk, I. 2010, Science, 328, 73
- Payez, A., Cudell, J. R. & Hutsemékers, D. 2011, Phys. Rev. D, 84, 085029
- Perna, R., et al. 2012, ApJ, 748, 116
- Pshirkov, M. S., Tinyakov, P. G. & Urban, F. R. 2016, Phys. Rev. Lett., 116, 191302
- Raffelt, G. & Stodolsky, L. 1988, Phys. Rev. D, 37, 1237
- Rees, M. J. & Setti, G. 1968, Nature, 219, 127
- Ringwald, A. 2012, Phys. Dark Univ., 1, 116
- Sahu, S., Medina-Carrillo, B., Sánchez-Colón, G. & Rajpoot, S. 2023, ApJ, 942, 30
- Saldana-Lopez, A., et al. 2021, MNRAS, 507, 5144
- Sánchez-Conde, M. A., et al. 2009,

Phys. Rev. D, 79, 123511

- Schwinger, J. 1951, Phys. Rev., 82, 664
- Simet, M., Hooper, D. & Serpico, P. D. 2008, Phys. Rev. D, 77, 063001
- Sisk-Reynés, J., et al. 2022, MNRAS, 510, 1264
- Stargate Collaboration 2022, GCN Circular n. 32648, https://gcn.gsfc.nasa.gov/ gcn3/32648.gcn3
- Tavecchio, F. & Bonnoli, G. 2016, A&A, 585, A25
- Tavecchio, F., Roncadelli, M. & Galanti, G. 2015, Phys. Lett. B, 744, 375
- Tavecchio, F., Roncadelli, M., Galanti, G. & Bonnoli, G. 2012, Phys. Rev. D, 86, 085036
- Thompson, T. A., et al. 2006, ApJ, 645, 186 Weisskopf, V. S. 1936, K. Dan. Vidensk. Selsk. Mat. Fys. Medd., 14, 6
- Wouters, D. & Brun, P. 2012, Phys. Rev. D, 86, 043005
- Yao, J. M., Manchester, R. N. & Wang, N. 2017, ApJ, 835, 29
- Zhao, Z.-C., Zhou, Y. & Wang, S. 2023, EPJC, 83, 92