

# Double Exponents in $SL(2, \mathbb{Z})$ Cosmology

---

**Renata Kallosh and Andrei Linde**

*Stanford Institute for Theoretical Physics and Department of Physics,  
Stanford University, Stanford, CA 94305, USA*

*E-mail:* [kallosh@stanford.edu](mailto:kallosh@stanford.edu), [alinde@stanford.edu](mailto:alinde@stanford.edu)

ABSTRACT:  $SL(2, \mathbb{Z})$  invariant  $\alpha$ -attractor models proposed in [1] have plateau potentials with respect to the inflaton and axion fields. The slope of the potential in the inflaton direction is exponentially suppressed at large values of the inflaton field, but the slope of the potential in the axion direction is *double-exponentially suppressed*. Therefore, the axion field remains nearly massless and practically does not change during inflation. The inflationary trajectory in such models is stable with respect to quantum fluctuations of the axion field. We show that isocurvature perturbations do not feed into the curvature perturbations during inflation, and argue that such transfer may remain inefficient at the post-inflationary stage.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Inflationary potentials and power of double exponents</b>	<b>3</b>
<b>3</b>	<b>Inflation in <math>SL(2, \mathbb{Z})</math> invariant models</b>	<b>7</b>
<b>4</b>	<b>Background invariant action for linear perturbations</b>	<b>9</b>
<b>5</b>	<b><math>SL(2, \mathbb{Z})</math> models with hyperbolic geometry and ultra-light axion</b>	<b>11</b>
<b>6</b>	<b>Adiabatic perturbations from the isocurvature ones?</b>	<b>14</b>
<b>7</b>	<b>Discussion</b>	<b>16</b>

---

## 1 Introduction

In this paper, we continue investigation of the cosmological  $\alpha$ -attractor models with  $SL(2, \mathbb{Z})$  symmetry proposed in [1–3], where the potentials depend on  $SL(2, \mathbb{Z})$  invariants.

We focus on a class of potentials introduced in [1] depending on Klein’s  $j$ -function

$$\left(\frac{\mathcal{L}(\tau, \bar{\tau})}{\sqrt{-g}}\right)^{SL(2, \mathbb{Z})} = \frac{R}{2} + \frac{3\alpha}{4} \frac{\partial\tau\partial\bar{\tau}}{(\text{Im}\tau)^2} - V(j(\tau), \bar{j}(\tau)). \quad (1.1)$$

All other  $SL(2, \mathbb{Z})$  invariant potentials in [1, 2] have analogous features with regard to the axion  $\theta$  and the inflaton  $\varphi$ , where

$$\tau = x + iy = \theta + ie\sqrt{\frac{2}{3\alpha}}\varphi. \quad (1.2)$$

Three features of our cosmological models play an important role in the analysis.

1. *The geodesics in hyperbolic geometry in half-plane coordinates (1.2) are straight vertical lines*

$$\theta = \text{const} \quad (1.3)$$

as well as certain semi-circles, see Fig. 1 and discussion in Appendix in [4]. The reason for the straight line geodesics in curved space (line of the shortest distance) is the property of the hyperbolic geometry that  $\Gamma_{\varphi\varphi}^{\theta} = 0$  and the geodesic equation

$$\theta'' - 2\sqrt{\frac{2}{3\alpha}}\varphi'\theta' = 0. \quad (1.4)$$

In general geometry, this equation would have a term  $\Gamma_{\varphi\varphi}^\theta \varphi' \varphi'$ , which would prevent a straight line solution for the geodesics  $\theta' = 0$ .

2. *The potential at large positive values of the inflaton field  $\varphi$  has the  $\alpha$ -attractor behavior,*

$$V(\varphi, \theta) \propto 1 - e^{-\sqrt{\frac{2}{3\alpha}}\varphi} + \dots \quad (1.5)$$

3. *The derivatives of the potential with respect to the axion field are double-exponentially suppressed during inflation*

$$\frac{\partial V}{\partial \theta} \propto e^{-2\pi e\sqrt{\frac{2}{3\alpha}}\varphi} \sin(2\pi\theta). \quad (1.6)$$

This feature makes the geodesic  $\theta = \text{const}$  a correct solution even with an account of the potential since the contribution from the potential in the  $\theta$  equation during inflation can be neglected.

The number of e-foldings of inflation beginning at some value of the inflaton field  $\varphi$  in  $\alpha$ -attractor models is given by

$$N_e = \frac{3\alpha}{8} e^{\sqrt{\frac{2}{3\alpha}}\varphi}. \quad (1.7)$$

Therefore the double-exponential factor in (1.6) can also be represented as

$$e^{-2\pi e\sqrt{\frac{2}{3\alpha}}\varphi} = e^{-\frac{16\pi N_e}{3\alpha}} \sim 10^{-1200} = \text{googol}^{-12} \quad (1.8)$$

for  $\alpha = 1/3$  and  $N_e \sim 55$ . Thus the double exponential behavior of the axion potential during inflation brings in the googol scale into cosmology. One googol<sup>1</sup> equals  $10^{100}$  and  $\text{googol}^{-12} = 10^{-1200}$ . These are not the numbers that we often encounter in physics. Amazingly, these numbers appear in the description of derivatives of the  $SL(2, \mathbb{Z})$  invariant potentials. The origin of the double-exponential suppression (1.6) will be discussed in section 2.

It is instructive to compare inflation in the theory of  $SL(2, \mathbb{Z})$  invariant  $\alpha$ -attractors [1] with inflation in closely related models in [5, 6], where the universality of a broad class of two-field  $\alpha$ -attractor predictions was established. It was found there that once inflation takes place near the boundary at  $y \rightarrow 0$ , the general inflationary predictions of T and E two-field models coincide with general predictions of single-field  $\alpha$ -attractors for a large number of e-foldings  $N_e$  [7–10]:

$$A_s = \frac{V_0 N_e^2}{18\pi^2 \alpha}, \quad n_s = 1 - \frac{2}{N_e}, \quad r = \frac{12\alpha}{N_e^2}. \quad (1.9)$$

Here  $A_s$  describes the magnitude of perturbations,  $n_s$  is the slope of their spectrum, and  $r$  is the tensor-to-scalar ratio.

---

<sup>1</sup>Google, the name of the search engine and the verb that refers to searching the Internet using the Google search engine are plays on the mathematical expression  $\text{googol} = 10^{100}$ .

It would be nice to use these results for investigation of the two-field evolution in the models with the  $SL(2, \mathbb{Z})$  invariant plateau potentials [1]. However, there is a substantial difference between these models and the models explored in [5, 6]. The main difference is in the choice of the variable  $y$  as a function of the canonically normalized inflaton field  $\varphi$ .

There are two such choices at large  $\varphi$

$$y = e^{-\sqrt{\frac{2}{3\alpha}}\varphi} , \quad (1.10)$$

$$y = e^{\sqrt{\frac{2}{3\alpha}}\varphi} . \quad (1.11)$$

In the first case, implemented in the models of Refs. [5, 6],  $y$  becomes exponentially small at large  $\varphi$ ; in the second case, studied in [1],  $y$  it becomes exponentially large. Thus, large  $\varphi$  corresponds to inflation near the  $y \rightarrow 0$  boundary as in eq. (1.10).

Meanwhile inflation in the  $SL(2, \mathbb{Z})$  invariant  $\alpha$ -attractors occurs in the limit  $y \rightarrow \infty$  as in eq. (1.11). Therefore the results of [5, 6] do not automatically apply to the new models.

As we will show in this paper, the axion field also remains frozen during inflation in the theory of  $SL(2, \mathbb{Z})$  invariant  $\alpha$ -attractors, but for a different reason. At large values of the inflaton field  $\varphi$ , the slope of the potential in the axion direction practically vanishes. As a result, even if the axion field was moving initially, it eventually stops, and after that, the axion does not move at all until the field  $\varphi$  becomes small and inflation ends.

Thus, inflation happens just like in the single-field  $\alpha$ -attractors. However, one may wonder whether quantum fluctuations of the nearly massless axion field may destabilize the inflationary trajectory and/or affect the standard  $\alpha$ -attractor predictions. We are going to investigate these issues in the present paper.

## 2 Inflationary potentials and power of double exponents

Cosmological  $\alpha$ -attractor models with  $SL(2, \mathbb{Z})$  symmetry were developed only recently [1–3]. For definiteness, we focus here on potentials depending on Klein’s Absolute Invariant  $J(\tau)$  or  $j(\tau)$ -function where  $j(\tau) = 12^3 J(\tau)$  [1]. These modular functions are defined in the hyperbolic geometry of the upper half plane  $\tau = x + iy$  where  $y > 0$ :

$$j(\tau) = q^{-1} + \sum_{n=0} c_n q^n , \quad c_0 = 744 , \quad (2.1)$$

where

$$q = e^{2\pi i\tau} = e^{-2\pi y + 2\pi i x} \quad q^{-1} = e^{-2\pi i\tau} = e^{2\pi y - 2\pi i x} . \quad (2.2)$$

Properties of the hyperbolic geometry in half-plane coordinates can be illustrated by Escher’s picture of Heaven and Hell, Fig. 1.



**Figure 1:** Escher's picture of Heaven and Hell in half-plane variables. The boundary of the half-plane  $y \rightarrow 0$  is the absolute, which cannot be reached. The angels and devils look smaller and smaller near  $y \rightarrow 0$ . Inflation in  $SL(2, \mathbb{Z})$  invariant potentials in [1] takes place in the upper part of the half-plane close to  $y \rightarrow +\infty$ . We presented and discussed this picture and the relevant cosmological  $\alpha$ -attractor models, mostly in disk variables in [11].



**Figure 2:** Fredrik Johansson's complex plot of the  $j$ -function, see fredrikj.net in My blog: Modular forms in Arb. Here Arb is for a C library for arbitrary-precision ball arithmetic.

We have presented the complex plot of the  $j$ -function in Fig. 2. At  $y \gg 1$ , the function  $j(\tau)$  can be well represented by its first two terms:

$$j(\tau)|_{y \gg 1} = \left( e^{2\pi y - 2\pi i x} + c_0 + \sum_{n=1} c_n e^{-2n\pi y + 2n\pi i x} \right) |_{y \gg 1} \rightarrow e^{2\pi y - 2\pi i x} + 744 . \quad (2.3)$$

It is known from [12, 13] that at  $n \geq 1$

$$c_n \leq \frac{e^{4\pi\sqrt{n}}}{\sqrt{2} n^{3/4}} , \quad (2.4)$$

and therefore the terms of order  $n$  as a function of  $y$  are constrained to be

$$c_n q^n \leq \frac{e^{4\pi\sqrt{n}}}{\sqrt{2} n^{3/4}} e^{-2\pi n y} . \quad (2.5)$$

Therefore, these series rapidly converge even for the values of  $y$  close to the end of inflation. For example, already for  $y = 2$  the series converges very fast,


$$c_n q^n |_{y=2} \leq \frac{e^{-4\pi(n-\sqrt{n})}}{\sqrt{2} n^{3/4}} . \quad (2.6)$$

Meanwhile, the  $j(\tau)$ -function near the boundary  $y \rightarrow 0$  is given by the series in eq. (2.1) where all terms have to be taken into account. Nevertheless, one can investigate its global properties by using  $SL(2, \mathbb{Z})$  invariance [4].

As an example, consider the potential

$$V = V_0 \left( 1 - \frac{\ln j(i)^2}{\ln (|j(\tau)|^2 + j(i)^2)} \right). \quad (2.7)$$

Note that  $j[\tau = e^{\frac{2\pi i}{3}}] = 0$  and the potential vanishes,  $V(\tau = e^{\frac{2\pi i}{3}}) = 0$  and it has a minimum there. At  $\tau = i$  the potential is positive and has a de Sitter saddle point. We show the plot of this potential in Fig. 3. As one can see, at large  $\varphi$  the potential is very flat in the  $\theta$  direction.



**Figure 3:** Potential (2.7), for  $\tau = \theta + ie^{\sqrt{\frac{2}{3\alpha}}\varphi}$  as a function of  $\varphi$  and  $\theta$  in case  $3\alpha = 1$ .

We may also consider a slightly more general potential

$$V = V_0 \left( 1 - \frac{\ln \beta^2}{\ln (|j(\tau)|^2 + \beta^2)} \right). \quad (2.8)$$

The potential in (2.7) is a special choice of the one in (2.8) when  $\beta = j(i) = 12^3$ . The so-called Klein invariant  $J(\tau) = 12^{-3}j(\tau)$  at this point is  $J(i) = 1$ .

To understand the origin of the remarkable flatness of this potential in  $\theta$  at large  $\varphi$ , we will evaluate the derivative  $V_\theta$  at large  $\varphi$ . We can use the expansion in eq. (2.1)  $j(\tau) = q^{-1} + \sum_{n=0} c_n q^n$ , where  $q^{-1} = e^{2\pi(y-i\theta)}$  and  $q = e^{-2\pi(y-i\theta)}$ . At large  $y$ , as we argued in eqs. (2.3), (2.6),

$$j(\tau)|_{y \rightarrow \infty} \rightarrow e^{2\pi y} e^{-i2\pi\theta} + 744, \quad (2.9)$$

where we have omitted all terms of the order  $e^{-2\pi n y}$  with  $n \geq 1$ , so that

$$|j(\tau)|^2 = |e^{2\pi e^{\sqrt{\frac{2}{3\alpha}}\varphi}} + 744 e^{i2\pi\theta}|^2. \quad (2.10)$$

We can now calculate the derivative  $V_\theta$  of the potential (2.8) during inflation.

As we already mentioned in the Introduction, during inflation in  $\alpha$ -attractor models, we have  $e^{\sqrt{\frac{2}{3\alpha}}\varphi} = \frac{8N_e}{3\alpha}$ , where  $N_e$  is the number of e-folds of inflation beginning at  $\varphi = \varphi_{N_e}$ . Therefore the exponential coefficient  $2\pi e^{\sqrt{\frac{2}{3\alpha}}\varphi}$  in this equation can be represented as

$$e^{2\pi e^{\sqrt{\frac{2}{3\alpha}}\varphi}} = e^{2\pi e^{\frac{8N_e}{3\alpha}}}. \quad (2.11)$$

This term remains extraordinarily large all the way to the end of inflation with  $N_e = 0$ . Therefore one can keep only two leading terms in (2.10):

$$|j(\tau)|^2 = e^{4\pi e^{\sqrt{\frac{2}{3\alpha}}\varphi}} \left( 1 + 1488 e^{-2\pi e^{\sqrt{\frac{2}{3\alpha}}\varphi}} \cos(2\pi\theta) \right). \quad (2.12)$$

As one can see, the  $\theta$ -dependent term is double-exponentially suppressed as compared to the first term.

The double-exponential growth of  $|j(\tau)|^2$  disappears when we calculate the potential (2.8), but the double-exponential suppression of its  $\theta$ -dependence remains. Using (2.12), we find that the  $\theta$ -derivative of the potential (2.8) during inflation is

$$\frac{\partial V}{\partial \theta} = -372 V_0 \sin(2\pi\theta) \ln(\beta^2) y^{-2} e^{-2\pi y}. \quad (2.13)$$

Note that  $V_\theta$  is multiplied by the exponentially large factor  $\frac{2}{3\alpha} e^{2\sqrt{\frac{2}{3\alpha}}\varphi} = \frac{2}{3\alpha} y^2$  in the equation for  $\theta$  (3.5), so the relevant quantity is

$$\frac{2}{3\alpha} y^2 \frac{\partial V}{\partial \theta} = -\frac{248 V_0}{\alpha} \sin(2\pi\theta) \ln(\beta^2) e^{-2\pi y}. \quad (2.14)$$

The most important factor in this expression is the double-exponential factor  $e^{-2\pi y} = e^{-2\pi e^{\sqrt{\frac{2}{3\alpha}}\varphi}}$  which makes the last term in equation (3.5) vanishingly small during inflation despite of the factor  $y^2 = \frac{2}{3\alpha} e^{2\sqrt{\frac{2}{3\alpha}}\varphi}$ . Replacing  $y$  by  $e^{\sqrt{\frac{2}{3\alpha}}\varphi}$  in eq. (2.14) we find

$$\frac{2}{3\alpha} y^2 \frac{\partial V}{\partial \theta} = -\frac{248 V_0}{\alpha} \sin(2\pi\theta) \ln(\beta^2) e^{-2\pi e^{\sqrt{\frac{2}{3\alpha}}\varphi}}. \quad (2.15)$$

The second derivative of the potential over  $\theta$ ,  $V_{\theta\theta}$  is also suppressed by the double exponential factor.

To understand the strength of this effect, we once again use the fact that during inflation in  $\alpha$ -attractor models,  $y = e^{\sqrt{\frac{2}{3\alpha}}\varphi} = \frac{8N_e}{3\alpha}$ . This means that

$$\frac{2}{3\alpha} e^{2\sqrt{\frac{2}{3\alpha}}\varphi} \frac{\partial V}{\partial \theta} = -\frac{496 V_0}{\alpha^2} \sin(2\pi\theta) \ln(\beta^2) e^{-\frac{16\pi N_e}{3\alpha}}. \quad (2.16)$$

As shown in the Introduction, the last coefficient in (2.16) is extremely small. For example, for  $N_e = 55$  and  $\alpha = 1/3$ , one finds that the derivative of the potential with respect to  $\theta$ , even multiplied by  $e^{2\sqrt{\frac{2}{3\alpha}}\varphi}$ , as we need for eq. (3.7), is suppressed by the factor  $10^{-1200}$ , see (1.8).

This explains why the potential is incredibly flat in the axion direction during inflation. Note that the approach to the plateau in the inflaton direction is linear in  $1/y \sim e^{-\sqrt{\frac{2}{3\alpha}}\varphi}$ . The flatness is exponential for the inflaton but double exponential for the axion.

Now we will find the value of  $N_e$  for which the maximal value of the right-hand side of (2.16) becomes of the same order as the derivative of the potential with respect to  $\varphi$ , which is given by

$$V_\varphi = V_0 \sqrt{\frac{2}{3\alpha}} e^{-\sqrt{\frac{2}{3\alpha}}\varphi} = V_0 \frac{8N_e}{3\alpha} \sqrt{\frac{2}{3\alpha}} . \quad (2.17)$$

One can easily check that, for  $\alpha \lesssim 1$ , the term  $V_\varphi$  in the equation (3.6) remains much greater than the term  $\frac{2}{3\alpha} e^{2\sqrt{\frac{2}{3\alpha}}\varphi} \frac{\partial V}{\partial \theta}$  in (3.7) all the way to the end of inflation at  $N_e < 1$ . Thus, during inflation one can safely ignore the term  $\frac{2}{3\alpha} e^{2\sqrt{\frac{2}{3\alpha}}\varphi} \frac{\partial V}{\partial \theta}$  in the equations (3.5) and (3.7) for  $\theta$ . After the end of inflation, the coefficient  $\frac{2}{3\alpha} e^{2\sqrt{\frac{2}{3\alpha}}\varphi}$  becomes  $O(1)$ , so the evolution of the fields  $\varphi$  and  $\theta$  shortly after the end of inflation can be well understood ignoring hyperbolicity of space-time.

### 3 Inflation in $SL(2, \mathbb{Z})$ invariant models

Our cosmological models belong to a class of models where the field space has a curved geometry

$$\mathcal{S} = \int d^4x \sqrt{-g} \left[ -\frac{1}{2} \mathcal{G}_{IJ}(\phi^K) \partial_\mu \phi^I \partial^\mu \phi^J - V(\phi^K) \right] , \quad (3.1)$$

where  $\mathcal{G}_{IJ}(\phi^K)$  is the metric on the field space. Equations of motions are

$$\mathcal{D}_t \dot{\phi}^I + 3H \dot{\phi}^I + V^{,I} = 0 , \quad (3.2)$$

where  $\mathcal{D}_t \phi^I = \dot{\phi}^I + \Gamma_{JK}^I \dot{\phi}^J \phi^K$ .

In hyperbolic geometry with  $\tau = \theta + ie\sqrt{\frac{2}{3\alpha}}\varphi$  and  $\phi^1 = \varphi$  and  $\phi^2 = \theta$ ,  $G_{11} = 1$ ,  $G_{22} = \frac{3\alpha}{2} e^{-2\sqrt{\frac{2}{3\alpha}}\varphi}$ . The action of the scalar fields in hyperbolic geometry is given by<sup>2</sup>

$$\int d^4x \sqrt{-g} \left( -\frac{1}{2} \partial^\mu \varphi \partial_\mu \varphi - \frac{3\alpha}{4} e^{-2\sqrt{\frac{2}{3\alpha}}\varphi} \partial_\mu \theta \partial^\mu \theta - V(\varphi, \theta) \right) . \quad (3.3)$$

Equations of motion (3.2) for the homogeneous fields  $\varphi$  and  $\theta$  are

$$\ddot{\varphi} + 3H\dot{\varphi} + \sqrt{\frac{3\alpha}{2}} e^{-2\sqrt{\frac{2}{3\alpha}}\varphi} \dot{\theta}^2 + V_\varphi = 0 , \quad (3.4)$$

---

<sup>2</sup>Inflation and quantum fluctuations in the context of very similar models were studied in detail in [5, 6]. In these models the metric of the  $\theta$  field at large  $\varphi$  is of the form  $\sim e^{2\sqrt{\frac{2}{3\alpha}}\varphi}$



$$\ddot{\theta} + 3H\dot{\theta} - 2\sqrt{\frac{2}{3\alpha}}\dot{\varphi}\dot{\theta} + \frac{2}{3\alpha}e^{2\sqrt{\frac{2}{3\alpha}}\varphi}V_{\theta} = 0 . \quad (3.5)$$

These two equations at  $\varphi \gg \sqrt{\alpha}$  in the slow-roll approximation are

$$3H\dot{\varphi} = -V_{\varphi} , \quad (3.6)$$

$$3H\dot{\theta} = -\frac{2}{3\alpha}e^{2\sqrt{\frac{2}{3\alpha}}\varphi}V_{\theta} . \quad (3.7)$$

Note that although  $V_{\theta}$  is small, it is multiplied by  $e^{2\sqrt{\frac{2}{3\alpha}}\varphi}$ , which is very large during inflation. This could destabilize the trajectory with  $\dot{\theta} = 0$ .

However, in all  $SL(2, \mathbb{Z})$  invariant potentials presented in [1], the plateau of the potentials at  $\sqrt{\frac{2}{3\alpha}}\varphi \gg 1$  is very flat not only with respect to the inflaton field  $\varphi$ , which is a general property of all  $\alpha$ -attractors, but also with respect to the axion field  $\theta$ . Moreover, the smallness of  $V_{\theta}$  is quite extraordinary, which can be explained by the fact that the  $SL(2, \mathbb{Z})$  invariants at large  $\varphi$  behave as double exponents in  $\varphi$ , see eq. (2.12).

Let us return to equations (3.4) and (3.5) with this new understanding of the properties of the axion potential. The results obtained above imply that we can simply ignore the last term in (3.5) during inflation, and this equation becomes

$$\ddot{\theta} + 3H\dot{\theta} \left(1 - \frac{2}{3}\sqrt{\frac{2}{3\alpha}}\frac{\dot{\varphi}}{H}\right) = 0 . \quad (3.8)$$

In the slow-roll regime with  $\dot{\phi} < 0$ , this equation can be written as

$$\ddot{\theta} + 3H\dot{\theta} \left(1 + \frac{4}{3}\sqrt{\frac{\epsilon}{3\alpha}}\right) = 0 , \quad (3.9)$$

where  $\epsilon = \frac{1}{2}\frac{V_{\varphi}^2}{V^2} \ll 1$ . In  $SL(2, \mathbb{Z})$  invariant models, the values of  $3\alpha$  are discrete, 1, 2, 3, ..., 7, see [1]. In such models, the last term in (3.9) can be neglected, the Hubble constant at the beginning of inflation is approximately constant, and the solution of this equation is given by

$$\theta = \theta_0(1 - e^{-3Ht}). \quad (3.10)$$

This implies that during the early stages of inflation with  $H \approx const$ , the speed of the field  $\theta$  decreases exponentially,  $\dot{\theta} \propto e^{-3Ht}$ , and within several e-foldings the field  $\theta$  freezes,

$$\theta = \theta_0 . \quad (3.11)$$

Equation (3.4) with  $\dot{\theta} = 0$  becomes the standard equation for the single-field  $\alpha$ -attractors, decoupled from the field  $\theta$ .

$$\ddot{\varphi} + 3H\dot{\varphi} + V_{\varphi} = 0 . \quad (3.12)$$

Therefore, one may expect that the general inflationary predictions of the new class of models in the models with  $\dot{\theta} = 0$  coincide with general predictions of single-field  $\alpha$ -attractors (1.9),

see [7, 8, 10]. However, one must check whether the perturbations of the ultra-light axion field may lead to significant isocurvature perturbations, which could destabilize the inflationary trajectory and alter the amplitude of the curvature perturbations.

Similar issues were recently discussed in [14–16] in the context of Fibre inflation derived in perturbative string theory. The potential of the inflaton field  $\varphi$  in Fibre inflation during the last 50-60 e-foldings nearly coincides with the  $\alpha$ -attractor potentials with  $\alpha = 2$  [17]. The Fibre model of perturbative string theory was extended with non-perturbative terms of the type  $e^{-aT}$ , which resulted in double-exponentially flat axion directions of the extended potential, similar to what we found in our  $SL(2, \mathbb{Z})$  invariant models. The investigation performed in [15, 16] shows that the isocurvature perturbations in their models do not affect their cosmological predictions. Since the models studied in [15, 16] are somewhat similar to our models with  $SL(2, \mathbb{Z})$  invariant potentials, one may expect that their conclusions regarding adiabatic and isocurvature perturbations apply to our models as well.

However, the results obtained in [15, 16] were based in part on the numerical investigation of the model with substantially different parameters, and the model itself was recently modified [18]. In the following sections, we will perform an independent investigation of these issues in the context of our models with  $SL(2, \mathbb{Z})$  invariant potentials.

## 4 Background invariant action for linear perturbations

Following [19], we will investigate here the cosmological models (3.1). The equation of motion for the Fourier modes of scalar field perturbations *on flat hypersurfaces*  $Q^I = \delta\phi^I$  was derived in [20]

$$\mathcal{D}_t \mathcal{D}_t Q^I + 3H \mathcal{D}_t Q^I + \frac{k^2}{a^2} Q^I + \mathcal{M}^I{}_J Q^J = 0, \quad (4.1)$$

where

$$\mathcal{M}^I{}_J \equiv V_{;J}^I - \mathcal{R}^I{}_{KLJ} \dot{\phi}^K \dot{\phi}^L - \frac{1}{a^3} \mathcal{D}_t \left( \frac{a^3}{H} \dot{\phi}^I \dot{\phi}_J \right). \quad (4.2)$$

Here the covariant in field-space geometry “long” derivatives involve Christoffel symbols due to the non-trivial geometry of the field-space

$$\mathcal{D}_t A^I = \dot{A}^I + \Gamma_{JK}^I \dot{\phi}^J A^K, \quad (4.3)$$

$$V_{;IJ} \equiv V_{,IJ} - \Gamma_{IJ}^K V_{,K}. \quad (4.4)$$

The second term in eq. (4.2) depends on Riemann tensor  $\mathcal{R}^I{}_{KLJ}$ . For a two-dimensional field space,  $I = 1, 2$  the only independent component of the Riemann tensor is  $\mathcal{R}_{IJKL} = \frac{1}{2} \mathcal{R}_{\text{fs}} (\mathcal{G}_{IK} \mathcal{G}_{JL} - \mathcal{G}_{IL} \mathcal{G}_{JK})$ , where  $\mathcal{R}_{\text{fs}}$  is the intrinsic scalar curvature of the field space. In our case of hyperbolic geometry in eq. (1.1)  $\mathcal{R}_{\text{fs}} = -\frac{4}{3\alpha}$  (note that the supergravity Kähler curvature is  $\mathcal{R}_K = -\frac{2}{3\alpha}$ ). The third term in eq. (4.2) presents the gravitational backreaction due to spacetime metric perturbations induced by the field fluctuations at first order.

If one prefers to work with gauge-invariant perturbations, one has to replace  $Q^I = \delta\phi^I$  as follows

$$Q^I = \delta\phi^I - \mathcal{R} \frac{\dot{\phi}^I}{H} . \quad (4.5)$$

Here  $\mathcal{R}$  is the intrinsic curvature perturbation of the constant time hypersurfaces, defined in [20]. The variables (4.5) represent a multi-field version of the Mukhanov-Sasaki variables [21–23]. They are described, for example, in [24].

To study cosmology with ultra-light axions in FRW background, it is important to take into account an identity derived in [25] for the second order action for perturbations  $Q^I$  in multi-field inflation models with kinetic terms described by a non-trivial metric  $G_{IJ}(\phi^K)$  in field space<sup>3</sup>.

In the modes which we study, without higher derivatives  $P(X, \phi^I) = X - V(\phi^K)$ , the identity proven in [25] takes a simple form, which we will present below and apply to our cosmological models with hyperbolic geometry.

The identity in our case states that the manifestly covariant second-order action  $S_{(2)}$ , where all derivatives are “long” derivatives as in eqs. (4.3), (4.4), is equal (up to total derivatives) to the action  $S_{(2)}^{non-cov}$  where derivatives are “short”. This action  $S_{(2)}^{non-cov}$  is not manifestly invariant, but it is equal to a manifestly invariant action  $S_{(2)}$ . The identity is

$$S_{(2)}^{non-cov} = S_{(2)} . \quad (4.6)$$

Here the manifestly invariant action is

$$S_{(2)} = \frac{1}{2} \int dt d^3x a^3 [G_{IJ}(\mathcal{D}_t Q^I \mathcal{D}_t Q^J - \frac{1}{a^2} \partial_i Q^I \partial^i Q^J) - \mathcal{M}_{IJ} Q^I Q^J] , \quad (4.7)$$

where  $\mathcal{M}_{IJ}$  is related to the mass term in eq. (4.2) as follows

$$\mathcal{M}_{IJ} = G_{IK} \mathcal{M}^K{}_J \equiv V_{;IJ} - \mathcal{R}_{IKLJ} \dot{\phi}^K \dot{\phi}^L - \frac{1}{a^3} \mathcal{D}_t \left( \frac{a^3}{H} \dot{\phi}_I \dot{\phi}_J \right) . \quad (4.8)$$

Note that the space derivatives  $\partial_i$  are background covariant in the FRW metric. Thus, each term in this action is background invariant.

The action (4.7) can be brought to the form  $S_{(2)}^{non-cov}$  in which it is not invariant term by term; only the sum of all terms is invariant due to the Langlois-Renaux-Petel identity (4.6).

$$S_{(2)}^{non-cov} = \frac{1}{2} \int dt d^3x a^3 [G_{IJ}(\dot{Q}^I \dot{Q}^J - \frac{1}{a^2} \partial_i Q^I \partial^i Q^J) + 2G_{IJ,K} \dot{\phi}^I Q^K \dot{Q}^J - (\mathbb{M}^2)_{IJ} Q^I Q^J] . \quad (4.9)$$

Here

$$(\mathbb{M}^2)_{IJ} = V_{;IJ} - \frac{1}{2} G_{KL,IJ} \dot{\phi}^K \dot{\phi}^L + (3 - \epsilon) \dot{\phi}_I \dot{\phi}_J + \frac{1}{H} (V_{,I} \dot{\phi}_J + \dot{\phi}_I V_{,J}) . \quad (4.10)$$

---

<sup>3</sup>In [25] the action depends on a general function  $P(X, \phi^I)$  where  $X = -\frac{1}{2} \mathcal{G}_{IJ}(\phi^K) \partial_\mu \phi^I \partial^\mu \phi^J$ . In that way, it covers cosmological models with higher derivatives, like the Dirac-Born-Infeld inflationary model.

Here we used the fact that

$$-\frac{1}{a^3}\mathcal{D}_t\left(\frac{a^3}{H}\dot{\phi}_I\dot{\phi}_J\right) = (3-\epsilon)\dot{\phi}_I\dot{\phi}_J + \frac{1}{H}(V_{,I}\dot{\phi}_J + \dot{\phi}_IV_{,J}) . \quad (4.11)$$

To prove (4.11) we have used the background field equations

$$\mathcal{D}_t\dot{\phi}^I + 3H\dot{\phi}^I + V^{,I} = 0, \quad a^{-3}\mathcal{D}_t(a^3\dot{\phi}^I) + V^{,I} = 0 . \quad (4.12)$$

We may try to compare the action (4.9) for linear scalar perturbations in the FRW background with the quadratic part of the classical action for scalar fields in the FRW background. The action (4.9) can be given as

$$S_{(2)}^Q = \frac{1}{2} \int dt d^3x a^3 \left( -G_{IJ}\partial_\mu Q^I \partial^\mu Q^J - V_{,IJ}(\phi)Q^I Q^J + \quad (4.13)$$

$$2G_{IJ,K}\dot{\phi}^I Q^K \dot{Q}^J V_{,IJ} - \left[ \frac{1}{2}G_{KL,IJ}\dot{\phi}^K \dot{\phi}^L - (3-\epsilon)\dot{\phi}_I\dot{\phi}_J - \frac{1}{H}(V_{,I}\dot{\phi}_J + \dot{\phi}_IV_{,J}) \right] Q^I Q^J \right) \quad (4.14)$$

This is to be compared with the quadratic part of the classical action in eq. (3.1)

$$\mathcal{S}_{(2)}^{\text{cl}} = \frac{1}{2} \int d^4x \sqrt{-g} \left[ -\mathcal{G}_{IJ}\partial_\mu \phi^I \partial^\mu \phi^J - V_{IJ}\phi^I \phi^J \right] . \quad (4.15)$$

In general metric  $G_{IJ}(\phi)$  the actions for perturbations  $S_{(2)}^Q$  and classical action  $\mathcal{S}_{(2)}^{\text{cl}}$  differ significantly, e.g. all terms in the second line in eq. (4.14) are absent in (4.15).

We will now show that in  $SL(2, \mathbb{Z})$  models with hyperbolic geometry and ultra-light axions, the action for the perturbations  $Q^\theta \sim \delta\theta$  at small momentum  $k \rightarrow 0$  is the same as the action for the homogeneous classical field  $\theta$ .

## 5 $SL(2, \mathbb{Z})$ models with hyperbolic geometry and ultra-light axion

Consider the gauge-invariant perturbations in eq. (4.5). The term correcting  $\delta\phi^I$  to gauge-covariant  $Q^I$  depends on the intrinsic curvature perturbation of the constant time hypersurfaces  $\mathcal{R}$ , but it is also proportional to velocity  $\dot{\phi}^I$ . If a consistent background solution in some direction  $\hat{I}$  with  $\dot{\phi}^{\hat{I}} = 0$  exists we find that during inflation

$$Q^{\hat{I}}|_{\dot{\phi}^{\hat{I}}=0} = \delta\phi^{\hat{I}} . \quad (5.1)$$

It means that even in a gauge where  $\mathcal{R} \neq 0$ , in some direction  $\hat{I}$  where  $\dot{\phi}^{\hat{I}} = 0$  the gauge-invariant perturbation is just a scalar field perturbation.

This is precisely the case of  $SL(2, \mathbb{Z})$  models with hyperbolic geometry and ultra-light axion  $\theta$ , where during inflation we have an ultra-light background field  $\theta$  with vanishing background velocity and vanishing derivatives of the potential.

$$\dot{\theta} = 0, \quad V_{,\theta} = 0, \quad V_{,\theta\varphi} = 0, \quad V_{,\theta\theta} = 0 . \quad (5.2)$$

We will now apply systematically the study of perturbations above for general metric  $G_{IJ}(\phi)$  and arbitrary potentials to the case of  $SL(2, \mathbb{Z})$  cosmological models in FWR background. Our fields are  $\phi^1 = \varphi$ ,  $\phi^2 = \theta$ , and  $G_{11} = 1, G_{12} = 0, G_{22} = \sqrt{3\alpha/2}e^{-2\sqrt{2/3\alpha}\varphi}$ , see eq. (3.3). In such case, the action for inflationary perturbations takes the form

$$S_{(2)} = \frac{1}{2} \int d^4 x a^3 [G_{IJ}(\dot{Q}^I \dot{Q}^J - \frac{1}{a^2} \partial_i Q^I \partial^i Q^J) + 2G_{\theta\theta, \varphi} \dot{\theta} Q^\varphi \dot{Q}^\theta - (\mathbb{M}^2)_{IJ} Q^I Q^J]. \quad (5.3)$$

Here

$$(\mathbb{M}^2)_{IJ} = V_{,IJ} - \frac{1}{2} G_{\theta\theta, IJ} \dot{\theta} \dot{\theta} + (3 - \epsilon) \dot{\phi}_I \dot{\phi}_J + \frac{1}{H} (V_{,I} \dot{\phi}_J + \dot{\phi}_I V_{,J}). \quad (5.4)$$

During inflation, the expression for the mass matrix simplifies since  $\dot{\theta} = \frac{\partial V}{\partial \theta} = \frac{\partial^2 V}{\partial \theta \partial \varphi} = \frac{\partial^2 V}{\partial \theta \partial \theta} = 0$ . Therefore

$$(\mathbb{M}^2)_{IJ}^{\text{infl}} = \begin{pmatrix} (\mathbb{M}^2)_{\varphi\varphi} & 0 \\ 0 & 0 \end{pmatrix}. \quad (5.5)$$

This result confirms the related result obtained in [15] by a different method.

Thus, perturbations in the inflationary background have the following action

$$S_{(2)} = \frac{1}{2} \int d^4 x a^3 [(\dot{Q}^\varphi \dot{Q}^\varphi - \frac{1}{a^2} \partial_i Q^\varphi \partial^i Q^\varphi) + G_{\theta\theta}(\dot{Q}^\theta \dot{Q}^\theta - \frac{1}{a^2} \partial_i Q^\theta \partial^i Q^\theta) + (\mathbb{M}^2)_{\varphi\varphi} Q^\varphi Q^\varphi]. \quad (5.6)$$

The action for  $Q^\theta$  perturbations is decoupled from the action for  $Q^\varphi$  perturbations and, for inflationary perturbations with exponentially large wavelengths, has the same form as the action for the background field  $\theta$

$$\frac{1}{2} \int d^4 x a^3 G_{\theta\theta} \dot{Q}^\theta \dot{Q}^\theta \quad \Leftrightarrow \quad \frac{1}{2} \int d^4 x a^3 G_{\theta\theta} \dot{\theta} \dot{\theta} \quad (5.7)$$

The action for  $Q^\varphi$  perturbations even for  $k \rightarrow 0$  is different from the action for a background scalar  $\varphi$  since the last term in the expression

$$(\mathbb{M}^2)_{\varphi\varphi} = V_{,\varphi\varphi} - \frac{1}{a^3} \mathcal{D}_t \left( \frac{a^3}{H} \dot{\varphi} \dot{\varphi} \right). \quad (5.8)$$

is due to spacetime metric perturbations. This term is not present in the equation for the background field  $\varphi$ .

It is important that the action (5.6) does not have any terms depending simultaneously on  $Q^\varphi$  and  $Q^\theta$ . Therefore we have two independent equations for  $Q^\varphi$  and  $Q^\theta$  in the background  $\dot{\theta} = 0$ :

$$\ddot{Q}^\varphi + 3H \dot{Q}^\varphi + \left( \frac{k^2}{a^2} + \mathcal{M}^\varphi_\varphi \right) Q^\varphi = 0, \quad (5.9)$$

$$\ddot{Q}^\theta + (3H + \Gamma_{\theta\varphi}^\theta \dot{\varphi}) \dot{Q}^\theta + \left( \frac{k^2}{a^2} \right) Q^\theta = 0. \quad (5.10)$$

The first of these two equations is the standard equation for adiabatic perturbations in the single-field inflation models; its solutions are well-known. Therefore, all curvature perturbations produced at that stage with  $\dot{\theta} = 0$  coincide with the perturbations in the single-field  $\alpha$ -attractors.

The second equation can be further simplified. First of all, during inflation with  $\dot{\theta} = 0$ , the second term in the expression  $Q^\theta = \delta\theta - \mathcal{R}\frac{\dot{\theta}^i}{H}$  vanishes, so the amplitude of isocurvature perturbations  $Q^\theta$  is equal to the amplitude of the fluctuations  $\delta\theta$ .

Also,  $\Gamma_{\theta\varphi}^\theta = -\sqrt{3\alpha/2}$ , which means that  $3H + \Gamma_{\theta\varphi}^\theta\dot{\varphi} = 3H\left(1 - \sqrt{\frac{\alpha}{6}}\frac{\dot{\varphi}}{H}\right)$ . The full equation for perturbations  $Q^\theta = \delta\theta$  is

$$\ddot{\delta\theta} + 3H\dot{\delta\theta}\left(1 - \frac{2}{3}\sqrt{\frac{2}{3\alpha}}\frac{\dot{\varphi}}{H}\right) + \left(\frac{k^2}{a^2}\right)\delta\theta = 0. \quad (5.11)$$

In the limit  $k/a \ll H$ , i.e., for the modes with wavelengths greater than the horizon  $\sim H^{-1}$ , one can neglect the last term, and the equation for the perturbations  $\delta\theta$  becomes the same as the equation for the homogeneous field  $\theta$  (3.8).

In the slow-roll regime with  $\alpha = O(1)$  one has  $-\frac{2}{3}\sqrt{\frac{2}{3\alpha}}\frac{\dot{\varphi}}{H} = \frac{4}{3}\sqrt{\frac{c}{3\alpha}} \ll 1$ , and the equation (5.11) becomes

$$\ddot{\delta\theta} + 3H\dot{\delta\theta} + \left(\frac{k^2}{a^2}\right)\delta\theta = 0. \quad (5.12)$$

This equation coincides with equation (7.3.5) from the book [26], so we can use its solution presented there in equation (7.3.8):

$$\delta\theta_k(t) = \frac{iH}{k\sqrt{2k}} \left(1 + \frac{k}{iH}e^{-Ht}\right) \exp\left(\frac{ik}{H}e^{-Ht}\right). \quad (5.13)$$

At large  $t$ , when  $ke^{-Ht}$  becomes smaller than  $H$ , which corresponds to the horizon crossing, the solution ceases to oscillate and becomes equal to

$$\delta\theta_k(t)|_{ke^{-Ht} < H} \rightarrow \frac{iH}{k\sqrt{2k}}. \quad (5.14)$$

Eq. (7.3.12) in [26] explains that this leads to the well-known expression for the amplitude of the perturbations generated during a single e-folding:

$$\delta\theta = \frac{H}{2\pi}. \quad (5.15)$$

Thus we find that during each e-folding of inflation with  $\dot{\theta} = 0$ , the isocurvature perturbations

$$Q^\theta = \delta\theta - \mathcal{R}\frac{\dot{\theta}}{H} = \delta\theta \quad (5.16)$$

freeze at a constant value  $\frac{H}{2\pi}$  when the wavelengths of these perturbations become greater than  $H^{-1}$ .

One may wonder whether these perturbations may “unfreeze” and begin to grow at the very end of inflation when the field  $\varphi$  continues rolling down and the slow-roll approximation is no longer valid. This could happen if the friction coefficient  $3H\left(1 - \sqrt{\frac{\alpha}{6}} \frac{\dot{\varphi}}{H}\right)$  in equation (5.10) changes its sign. But  $\dot{\varphi} < 0$  when the field  $\phi$  rolls down. Therefore the friction coefficient is even greater than  $3H$  for  $\dot{\varphi} < 0$ . This leads to a strong stabilization of the inflationary trajectory.

These considerations address the issue of stability of the inflationary trajectory  $\dot{\theta} = 0$ : during inflation and shortly after its ends, the classical trajectory  $\dot{\theta} = 0$  of the ultra-light axion field is stable, as long as one can ignore the double-exponentially suppressed derivatives  $V_{\theta}$  and  $V_{\theta,\theta}$ .

However, there are small inflationary perturbations (5.15) near each classical trajectory  $\theta = \text{const}$ . At the end of inflation, the trajectory may turn towards one of the minima of the potential. That is the time when isocurvature perturbations could feed into the adiabatic ones. We will discuss this possibility in the next section.

## 6 Adiabatic perturbations from the isocurvature ones?

As we already mentioned, the field  $\theta$  remains constant during inflation in our models. In this sense, such models are similar to the single-field inflationary models, where the amplitude of curvature perturbations on comoving hypersurfaces  $\mathcal{R}_c$  has a simple interpretation in the  $\delta N$  approach [20, 27]:

$$\mathcal{R}_c = \delta N_{\varphi} = H\delta t = H \frac{\delta\varphi}{\dot{\varphi}} = \frac{H^2}{2\pi\dot{\varphi}} . \quad (6.1)$$

Here  $\delta N_{\varphi}$  is the change of the number of e-foldings due to the fluctuations with the average amplitude  $|\delta\varphi| = \frac{H}{2\pi}$ , which may either increase or decrease the number of e-folds until the end of inflation, depending on the sign of  $\delta\varphi$ . All quantities here should be evaluated at the moment  $N_e$  e-foldings before the end of inflation. To compare the results with the CMB data, one should consider  $N_e \sim 50$ .

The amplitude  $A_s$  usually presented in CMB data is given by the square of this quantity,

$$A_s = \delta N_{\varphi}^2 = \left( \frac{H^2}{4\pi^2\dot{\varphi}} \right)^2 . \quad (6.2)$$

For  $\alpha$ -attractors it is given by [7–10]

$$A_s = \frac{V_0 N_e^2}{18\pi^2\alpha} . \quad (6.3)$$

A comparison of (6.2) and (6.3) shows that the amplitude of perturbations  $\delta N_{\varphi}$  in these models is directly proportional to  $N_e$ . Since the Hubble constant does not change much until the end of inflation in these models, these equations imply that  $\dot{\varphi}$  is inversely proportional

to  $N_e$ . This means, for example, that the velocity of the field 50 e-foldings prior to the end of inflation is approximately 10 times smaller than the velocity of the field at  $N_e = 5$ , and  $O(50)$  times smaller than the velocity of the field at  $N_e = 1$ . This is a rough estimate, but the result is qualitatively correct, especially for  $\alpha < 1$ : The value of  $\dot{\varphi}$  at the end of inflation is significantly greater than the value of  $\dot{\varphi}$  at the moment corresponding to  $N_e \sim 50$ .

As long as the inflationary trajectory remains straight, the perturbations of the field  $\theta$  do not induce any perturbations  $\delta N_\theta$  because they are orthogonal to the inflationary trajectory. If, for example, the field  $\theta$  at the beginning was close to 0.5, then this field trajectory never changes its direction even long after inflation ends because  $V_\theta(\pm 0.5) = 0$  and  $V_{\theta,\theta}(\pm 0.5) > 0$  for all  $\theta$ . This trajectory ends in one of the minima at  $\theta_0 = 0.5$ . In that case (no turns of the field trajectory), the perturbations  $\delta\theta$  do not affect the spectrum of curvature perturbations (6.1) and the cosmological predictions of this class of models coincide with the standard predictions of  $\alpha$ -attractors.

However, for  $\theta_0 \neq \pm 0.5$ , each field trajectory eventually turns towards one of the minima of the potential at  $\varphi < 0$ . At that time, some of the perturbations  $\delta\theta$  generated at early stages of inflation may feed into the curvature perturbations.

For the sake of argument, suppose that after the turn, the trajectory turns to the direction  $\theta$ , which brings the field to one of the minima. At the beginning of the turn, the field was not exactly at  $\theta = \theta_0$ , but at  $\theta = \theta_0 \pm \delta\theta$ . The parts of the universe that were closer to the minimum of the potential by  $\delta\theta$  will reach it earlier by  $\delta t = \delta\theta/\dot{\theta}$ . The perturbations  $\delta\theta$  formed at  $N_e = 50$  have an initial magnitude  $|\delta\theta| = \frac{H}{2\pi}$ . As shown in the previous section, this amplitude does not change until the trajectory begins to turn.

If the turn occurs during inflation, such perturbations may lead to a change in the number of e-foldings by

$$\delta N_\theta \sim H\delta t = H \frac{\delta\theta}{\dot{\theta}} \approx \frac{H^2}{2\pi\dot{\theta}} . \quad (6.4)$$

These perturbations are adiabatic, and they are added to the previously generated adiabatic perturbations (6.1) with the same large wavelength  $\sim e^{N_e}$ .

In the previous section, we have shown that the inflationary trajectory does not bend,  $\theta = \text{const}$ , all the way to the end of inflation. This is a rather general result valid for a broad class of the  $SL(2, \mathbb{Z})$  invariant  $\alpha$ -attractor models. Moreover, the trajectory typically remains straight and stable all the way to  $\varphi \approx 0$ , after which the trajectory may turn towards one of the minima of the potential at  $\varphi < 0$  [28].

Thus, in the simplest models that we study, the inflationary trajectory *does not* turn during inflation, and its turning after the end of inflation should produce a much smaller effect, if any. Nevertheless, we will use (6.4) to estimate, perhaps too generously, a possible impact of the perturbations  $\delta\theta$  on the curvature perturbations.

To make this estimate, we would need to know the value of  $\dot{\theta}$  shortly after the fields turn.



The turn typically happens soon after the field  $\varphi$  approaches  $\varphi = 0$ . If the trajectory turns quickly, energy conservation implies that  $\dot{\theta}$  should not be much smaller than  $\dot{\varphi}$  at that time. An exact value of  $\dot{\varphi}$  at that time is model-dependent, but an investigation of the simplest T-model  $\alpha$ -attractor with the potential  $V \sim \tanh^2 \frac{\varphi}{\sqrt{6\alpha}}$  can be quite instructive. For  $\alpha = 1/3$ , one finds that the value of  $\dot{\varphi}$  at the  $\varphi = 0$  is approximately 50 times greater than the value of  $\dot{\varphi}$  in equation (6.1) at the moment corresponding to  $N_e \sim 50$ . This result matches the general estimates made at the beginning of this section. The value of  $\dot{\theta}$  after the turn in our models is expected to be just slightly smaller. The results of a numerical investigation of inflation in the simplest  $SL(2, \mathbb{Z})$  invariant  $\alpha$ -attractor models discussed in this paper confirm these expectations [28].

Suppose for definiteness that  $\dot{\theta}$  shortly after the turn is 30 times smaller than the value of  $\dot{\varphi}$  50 e-foldings from the end of inflation. This would mean that  $\delta N_\theta$  (6.4) is approximately 30 times smaller than  $\delta N_\varphi$  (6.1). And the contribution of  $\delta N_\theta$  to  $A_s$  is even much smaller.

Indeed, perturbations  $\delta\varphi$  and  $\delta\theta$  (and their sign) are not correlated. Therefore, one may estimate the resulting value of  $A_s$  as

$$A_s \approx \delta N_\varphi^2 + \delta N_\theta^2 . \quad (6.5)$$

Thus if  $\delta N_\theta$  is 30 times smaller than  $\delta N_\varphi$ , its relative contribution to  $A_s$  is 900 times smaller than the standard single-field contribution  $\delta N_\varphi^2$ . Even if  $\delta N_\theta$  is only 10 times smaller than  $\delta N_\varphi$ , its contribution to  $A_s$  is 100 times smaller than the standard single-field contribution  $\delta N_\varphi^2$ .

Finally, we should remember that the field trajectory in our models begins to turn only after inflation ends. In that case, quantum fluctuations  $\delta\theta$  do not affect the number of e-foldings of inflation. They may only slightly affect the time of the end of reheating, which has a much smaller impact on the equation of state than the end of inflation. This suggests that the conversion of isocurvature perturbations to adiabatic perturbations is even less efficient than what our simple estimates indicate.

## 7 Discussion

In this paper, we explored the structure of the inflationary plateau of the  $SL(2, \mathbb{Z})$  invariant  $\alpha$ -attractor potentials introduced in [1, 2]. We have found that the derivatives of these potentials in the axion direction  $\theta$  are double-exponentially suppressed during inflation at large values of the inflaton field  $\varphi$ . For example, during inflation the potential in eq. (2.7) is

$$V \rightarrow V_0 \left( 1 - c_1 e^{-\sqrt{\frac{2}{3\alpha}}\varphi} + c_2 e^{-2\sqrt{\frac{2}{3\alpha}}\varphi} e^{-2\pi e\sqrt{\frac{2}{3\alpha}}\varphi} \cos(2\pi\theta) \right) . \quad (7.1)$$

where the coefficients  $c_1$  and  $c_2$  depend on the choice of the modular potentials in [1, 2].

Because of the extraordinary flatness of the axion potential, the axion field freezes and does not evolve during inflation. Therefore in typical models of this type, the only evolving field is the inflaton. We developed a theory of curvature and isocurvature perturbations in these models and demonstrated that the isocurvature perturbations do not destabilize the inflationary trajectory. Thus, inflation in this theory is effectively single-field.

Our results are consistent with the results obtained in [15, 16] in Fibre inflation models, which have potentials with very similar properties. Effectively they can be described as an  $\alpha = 2$  case of supergravity  $\alpha$ -attractor models, see [18] and references therein.

However, a more detailed investigation is required to obtain a complete and reliable expression of the amplitude of adiabatic perturbations if the conversion of isocurvature perturbations to adiabatic may happen after the end of inflation, where the field trajectory turns towards one of the many minima of the potential. The corresponding results may be somewhat model-dependent because the post-inflationary dynamics in the models with  $SL(2, \mathbb{Z})$  invariant potentials depend on the structure of potentials close to their minima.

A preliminary investigation of this issue contained in the last section suggests that for the simplest models considered in this paper, the contribution to the amplitude  $A_s$  of the curvature perturbations generated from the isocurvature perturbations is strongly suppressed by a factor inversely proportional to  $N_e^2$ . In such case, the inflationary predictions of this class of models should nearly coincide with the predictions of the single-field  $\alpha$  attractors (1.9).

However, our investigation is based on the simplest  $SL(2, \mathbb{Z})$  invariant  $\alpha$ -attractor potentials, such as the potential (2.7) shown in Fig. 3. Modifications of the model parameters may lead to changes in the post-inflationary dynamics or even introduce a secondary stage of inflation. In particular, as mentioned in [1], the potential (2.8) has a complicated structure in the limit  $\beta \rightarrow 1$ , see Fig. 6 in [1]. In such models, inflation may consist of two separate stages. At the first stage, the field  $\theta$  does not change, as in the models considered above, but then the field trajectory turns in the  $\theta$ -direction, and a new stage of inflation begins. In that case, the spectrum of the perturbations may change considerably, which may even lead to the formation of primordial black holes, as in the models of hybrid attractors [29–31].

We plan to return to the investigation of  $SL(2, \mathbb{Z})$  invariant  $\alpha$ -attractors in the subsequent publication [28].

## Acknowledgement

We are grateful to M. Braglia, D. Wands, T. Wrase, and Y. Yamada for helpful discussions and especially to J.J. Carrasco and D. Roest for many fruitful discussions of  $SL(2, \mathbb{Z})$  cosmology and an ongoing collaboration on inflation in these models. This work is supported by SITP and by the US National Science Foundation Grant PHY-2310429.

## References

- [1] R. Kallosh and A. Linde, *SL(2, Z) Cosmological Attractors*, [2408.05203](#).
- [2] G.F. Casas and L.E. Ibáñez, *Modular Invariant Starobinsky Inflation and the Species Scale*, [2407.12081](#).
- [3] R. Schimrigk, *Large and small field inflation from hyperbolic sigma models*, *Phys. Rev. D* **105** (2022) 063541 [[2108.05400](#)].
- [4] R. Kallosh and A. Linde, *Landscape of Modular Cosmology*, [2411.07552](#).
- [5] A. Achúcarro, R. Kallosh, A. Linde, D.-G. Wang and Y. Welling, *Universality of multi-field  $\alpha$ -attractors*, *JCAP* **1804** (2018) 028 [[1711.09478](#)].
- [6] A. Achúcarro, A. Linde, D.-G. Wang, Y. Welling and Y. Yamada, *Hypernatural inflation*, *JCAP* **1807** (2018) 035 [[1803.09911](#)].
- [7] R. Kallosh, A. Linde and D. Roest, *Superconformal Inflationary  $\alpha$ -Attractors*, *JHEP* **11** (2013) 198 [[1311.0472](#)].
- [8] M. Galante, R. Kallosh, A. Linde and D. Roest, *Unity of Cosmological Inflation Attractors*, *Phys. Rev. Lett.* **114** (2015) 141302 [[1412.3797](#)].
- [9] J.J.M. Carrasco, R. Kallosh, A. Linde and D. Roest, *Hyperbolic geometry of cosmological attractors*, *Phys. Rev.* **D92** (2015) 041301 [[1504.05557](#)].
- [10] R. Kallosh and A. Linde, *BICEP/Keck and cosmological attractors*, *JCAP* **12** (2021) 008 [[2110.10902](#)].
- [11] R. Kallosh and A. Linde, *Escher in the Sky*, *Comptes Rendus Physique* **16** (2015) 914 [[1503.06785](#)].
- [12] H. Petersson, *Über die Entwicklungskoeffizienten der automorphen Formen*, *Acta Math.* **58** (1932) 169.
- [13] N. Brisebarre and G. Philibert, *Effective lower and upper bounds for the Fourier coefficients of powers of the modular invariant  $j$* , *Journal of the Ramanujan Mathematical Society* **20** (2005) 255.
- [14] M. Cicoli, C.P. Burgess and F. Quevedo, *Fibre Inflation: Observable Gravity Waves from IIB String Compactifications*, *JCAP* **0903** (2009) 013 [[0808.0691](#)].
- [15] M. Cicoli, V. Guidetti, F. Muia, F.G. Pedro and G.P. Vacca, *On the choice of entropy variables in multifield inflation*, *Class. Quant. Grav.* **40** (2023) 025008 [[2107.03391](#)].
- [16] M. Cicoli, V. Guidetti, F. Muia, F.G. Pedro and G.P. Vacca, *A fake instability in string inflation*, *Class. Quant. Grav.* **39** (2022) 195012 [[2107.12392](#)].
- [17] R. Kallosh, A. Linde, D. Roest, A. Westphal and Y. Yamada, *Fibre Inflation and  $\alpha$ -attractors*, *JHEP* **02** (2018) 117 [[1707.05830](#)].
- [18] M. Cicoli, A. Grassi, O. Lacombe and F.G. Pedro, *Chiral global embedding of Fibre Inflation with  $\overline{\text{D3}}$  uplift*, [2412.08723](#).
- [19] L. Iacconi, H. Assadullahi, M. Fasiello and D. Wands, *Revisiting small-scale fluctuations in  $\alpha$ -attractor models of inflation*, *JCAP* **06** (2022) 007 [[2112.05092](#)].

- [20] M. Sasaki and E.D. Stewart, *A General analytic formula for the spectral index of the density perturbations produced during inflation*, *Prog. Theor. Phys.* **95** (1996) 71 [[astro-ph/9507001](#)].
- [21] V.F. Mukhanov, *Gravitational Instability of the Universe Filled with a Scalar Field*, *Formen, Acta Math.* **58** (1932) 169.
- [22] M. Sasaki, *Large Scale Quantum Fluctuations in the Inflationary Universe*, *Prog. Theor. Phys.* **76** (1986) 1036.
- [23] V.F. Mukhanov, *Quantum Theory of Gauge Invariant Cosmological Perturbations*, *Sov. Phys. JETP* **67** (1988) 1297.
- [24] C.M. Peterson and M. Tegmark, *Testing Two-Field Inflation*, *Phys. Rev. D* **83** (2011) 023522 [[1005.4056](#)].
- [25] D. Langlois and S. Renaux-Petel, *Perturbations in generalized multi-field inflation*, *JCAP* **04** (2008) 017 [[0801.1085](#)].
- [26] A.D. Linde, *Particle physics and inflationary cosmology*, Harwood Academic Publishers (1990), [[hep-th/0503203](#)].
- [27] M. Sasaki and T. Tanaka, *Superhorizon scale dynamics of multiscalar inflation*, *Prog. Theor. Phys.* **99** (1998) 763 [[gr-qc/9801017](#)].
- [28] J.J. Carrasco, R. Kallosh, A. Linde and D. Roest, *work in progress*, .
- [29] R. Kallosh and A. Linde, *Hybrid cosmological attractors*, *Phys. Rev. D* **106** (2022) 023522 [[2204.02425](#)].
- [30] M. Braglia, A. Linde, R. Kallosh and F. Finelli, *Hybrid  $\alpha$ -attractors, primordial black holes and gravitational wave backgrounds*, *JCAP* **04** (2023) 033 [[2211.14262](#)].
- [31] L. Iacconi, M. Bacchi, L.F. Guimarães and F.T. Falciano, *Testing inflation on all scales: a case study with  $\alpha$ -attractors*, [2412.02544](#).