

The nucleus of the Johnson graph $J(N, D)$

Kazumasa Nomura and Paul Terwilliger

December 30, 2024

Abstract. This paper is about the nucleus of the Johnson graph $\Gamma = J(N, D)$ with $N > 2D$. The nucleus is described as follows. Let X denote the vertex set of Γ . Let $A \in \text{Mat}_X(\mathbb{C})$ denote the adjacency matrix of Γ . Let $\{E_i\}_{i=0}^D$ denote the Q -polynomial ordering of the primitive idempotents of A . Fix $x \in X$. The corresponding dual adjacency matrix A^* is the diagonal matrix in $\text{Mat}_X(\mathbb{C})$ such that for $y \in X$ the (y, y) -entry of A^* is equal to the (x, y) -entry of $|X|E_1$. For $0 \leq i \leq D$ the diagonal matrix $E_i^* \in \text{Mat}_X(\mathbb{C})$ is the projection onto the i th subconstituent of Γ with respect to x . The matrices $\{E_i^*\}_{i=0}^D$ are the primitive idempotents of A^* . The subalgebra T of $\text{Mat}_X(\mathbb{C})$ generated by A, A^* is called the subconstituent algebra of Γ with respect to x . Let $V = \mathbb{C}^X$ denote the standard module of Γ . For $0 \leq i \leq D$ define

$$\mathcal{N}_i = (E_0^*V + E_1^*V + \cdots + E_i^*V) \cap (E_0V + E_1V + \cdots + E_{D-i}V).$$

It is known that the sum $\mathcal{N} = \sum_{i=0}^D \mathcal{N}_i$ is direct, and \mathcal{N} is a T -module. This T -module \mathcal{N} is called the nucleus of Γ with respect to x . The sum $\mathcal{N} = \sum_{i=0}^D E_i^*\mathcal{N}$ is direct. In this paper we consider the following vectors in V . For a subset $\alpha \subseteq x$ let $\alpha^\vee \in V$ denote the characteristic vector of the set of vertices in X that contain α . Let $\alpha^\mathcal{N} \in V$ denote the characteristic vector of the set of vertices in X whose intersection with x is equal to α . We show that (i) the vectors $\{\alpha^\vee\}_{\alpha \subseteq x}$ form a basis of \mathcal{N} ; (ii) the vectors $\{\alpha^\mathcal{N}\}_{\alpha \subseteq x}$ form a basis of \mathcal{N} ; (iii) for $0 \leq i \leq D$, the vectors $\{\alpha^\mathcal{N} \mid \alpha \subseteq x, |\alpha| = D - i\}$ form a basis of $E_i^*\mathcal{N}$; (iv) for $0 \leq i \leq D$, the vectors $\{\alpha^\vee \mid \alpha \subseteq x, |\alpha| = D - i\}$ form a basis of \mathcal{N}_i . We give the transition matrices between the bases $\{\alpha^\vee\}_{\alpha \subseteq x}$ and $\{\alpha^\mathcal{N}\}_{\alpha \subseteq x}$. We give the actions of A, A^* on the bases $\{\alpha^\vee\}_{\alpha \subseteq x}$ and $\{\alpha^\mathcal{N}\}_{\alpha \subseteq x}$. For $0 \leq i \leq D$ we characterize the basis vectors for $E_i^*\mathcal{N}$ in terms of the connected components of a certain graph obtained by adjusting the edges of the subgraph of Γ induced on the i th subconstituent of Γ with respect to x .

1 Introduction

In [21] the second author introduced the nucleus of a Q -polynomial distance-regular graph. In this paper we describe the nucleus of the Johnson graph $J(N, D)$.

Before we summarize our results, we describe the context and history. Let Γ denote a distance-regular graph with vertex set X and diameter D , see [4, Section 4.1.A]. Let ∂

denote the path-length distance function. Let $A \in \text{Mat}_X(\mathbb{C})$ denote the adjacency matrix of Γ , see [1, p. 2]. Let $\{E_i\}_{i=0}^D$ denote a Q -polynomial ordering of the primitive idempotents of A , see [19, Section 2], [1, Definition 2.80]. Fix a vertex $x \in X$. For $0 \leq i \leq D$, the set $\Gamma_i(x) = \{y \in X \mid \partial(x, y) = i\}$ is called the i th subconstituent of Γ with respect to x . The corresponding dual primitive idempotent is the diagonal matrix $E_i^* \in \text{Mat}_X(\mathbb{C})$ such that for $y \in X$ the (y, y) -entry of E_i^* is 1 if $y \in \Gamma_i(x)$ and 0 if $y \notin \Gamma_i(x)$. Define a diagonal matrix $A^* \in \text{Mat}_X(\mathbb{C})$ such that for $y \in X$ the (y, y) -entry of A^* is equal to the (x, y) -entry of $|X|E_1$. The matrices $\{E_i^*\}_{i=0}^D$ are the primitive idempotents of A^* , see [15, Section 3]. The subalgebra T of $\text{Mat}_X(\mathbb{C})$ generated by A, A^* is called the subconstituent algebra of Γ with respect to x , see [15, Section 3]. Let $V = \mathbb{C}^X$ denote the vector space over \mathbb{C} consisting of the column vectors with coordinates indexed by X and all entries in \mathbb{C} . For $y \in X$ the vector $\hat{y} \in V$ has y -coordinate 1 and all other coordinates 0. The vectors $\{\hat{y}\}_{y \in X}$ form a basis of V . For $0 \leq i \leq D$ define a subspace \mathcal{N}_i of V by

$$\mathcal{N}_i = (E_0^*V + E_1^*V + \cdots + E_i^*V) \cap (E_0V + E_1V + \cdots + E_{D-i}V).$$

By [18, Corollary 5.8] the sum $\mathcal{N} = \sum_{i=0}^D \mathcal{N}_i$ is direct. By [18, Theorem 7.1] \mathcal{N} is a T -module. This T -module \mathcal{N} is the nucleus of Γ with respect to x . In [21] the nucleus is described for the hypercubes, the Odd graphs, the Hamming graphs, and the nonbipartite dual polar graphs.

In this paper we consider the Johnson graph $J(N, D)$. This graph is defined as follows. Fix nonnegative integers N, D such that $2D < N$. Let Ω denote a set of size N . Let the set X consist of the subsets of Ω that have size D . The graph $J(N, D)$ has vertex set X . Vertices $x, y \in X$ are adjacent whenever $|x \cap y| = D - 1$. By [4, Theorem 9.1.2] $J(N, D)$ is distance-regular with diameter D . The adjacency matrix A has eigenvalues

$$\theta_i = (D - i)(N - D - i) - i \quad (0 \leq i \leq D).$$

For $0 \leq i \leq D$ let E_i denote the primitive idempotent of A associated with θ_i . The ordering $\{E_i\}_{i=0}^D$ is Q -polynomial, see [2, Corollary to Theorem 2.6]. Recall the fixed vertex $x \in X$. We consider the following vectors in V . For a subset $\alpha \subseteq x$, define

$$\alpha^\vee = \sum_{\substack{y \in X \\ \alpha \subseteq y}} \hat{y}, \quad \alpha^\mathcal{N} = \sum_{\substack{y \in X \\ y \cap x = \alpha}} \hat{y}.$$

The vector α^\vee is the characteristic vector of the set of vertices in X that contain α , and $\alpha^\mathcal{N}$ is the characteristic vector of the set of vertices in X whose intersection with x is equal to α . We show the following:

- (i) the vectors $\{\alpha^\vee\}_{\alpha \subseteq x}$ form a basis of \mathcal{N} ;
- (ii) the vectors $\{\alpha^\mathcal{N}\}_{\alpha \subseteq x}$ form a basis of \mathcal{N} ;
- (iii) for $0 \leq i \leq D$, the vectors $\{\alpha^\vee \mid \alpha \subseteq x, |\alpha| = D - i\}$ form a basis of \mathcal{N}_i ;
- (iv) for $0 \leq i \leq D$, the vectors $\{\alpha^\mathcal{N} \mid \alpha \subseteq x, |\alpha| = D - i\}$ form a basis of $E_i^*\mathcal{N}$.

We give the transition matrices between the bases $\{\alpha^\vee\}_{\alpha \subseteq x}$ and $\{\alpha^N\}_{\alpha \subseteq x}$. We give the actions of A, A^* on the bases $\{\alpha^\vee\}_{\alpha \subseteq x}$ and $\{\alpha^N\}_{\alpha \subseteq x}$. We characterize the vectors $\{\alpha^N\}_{\alpha \subseteq x}$ as follows. For $0 \leq i \leq D$ we show that the subgraph $\Gamma_i(x)$ is isomorphic to the Cartesian product $J(D, i) \times J(N - D, i)$. So $\Gamma_i(x)$ has two types of edges, some come from $J(D, i)$ and some come from $J(N - D, i)$. Consider a new graph obtained from $\Gamma_i(x)$ by removing the edges that come from $J(D, i)$. The vectors $\{\alpha^N \mid \alpha \subseteq x, |\alpha| = D - i\}$ are the characteristic vectors of the connected components of this new graph.

This paper is organized as follows. Section 2 contains some preliminaries. In Section 3 we review Q -polynomial distance-regular graphs and the subconstituent algebra T . In Section 4 we review T -modules. In Section 5 we recall the notion of the nucleus. In Sections 6–11 we consider the Johnson graph. In Section 6 we review the definition and basic properties of the Johnson graph $J(N, D)$. In Section 7 we investigate the structure of the subgraph $\Gamma_i(x)$. In Section 8 we introduce the vectors α^N and α^\vee , and give relations between these vectors. In Section 9 we describe the action of A, A^* on the vectors α^N and α^\vee . In Section 10 we prove some results concerning α^\vee . In Section 11 we use the vectors α^N and α^\vee to obtain two bases for the nucleus.

2 Preliminaries

We now begin our formal argument. The following notation and concepts will be used throughout the paper. Let \mathbb{C} denote the field of complex numbers. Let X denote a finite nonempty set. Let $\text{Mat}_X(\mathbb{C})$ denote the \mathbb{C} -algebra consisting of the matrices with rows and columns indexed by X and all entries in \mathbb{C} . The identity matrix in $\text{Mat}_X(\mathbb{C})$ is denoted by I , and let $J \in \text{Mat}_X(\mathbb{C})$ have all entries 1. Let \mathbb{C}^X denote the vector space over \mathbb{C} consisting of the column vectors with coordinates indexed by X and all entries in \mathbb{C} . The algebra $\text{Mat}_X(\mathbb{C})$ acts on \mathbb{C}^X by left multiplication. We call \mathbb{C}^X the *standard module*. For a subset $Y \subseteq X$, the vector $\hat{Y} = \sum_{y \in Y} \hat{y}$ is called the *characteristic vector of Y* .

All the graphs we discuss are assumed to be finite, undirected, and have no loops and no multiple edges. Let Γ denote a graph with vertex set X . Then Γ is said to be *regular with valency k* whenever each vertex of Γ is adjacent to exactly k vertices. Define $A \in \text{Mat}_X(\mathbb{C})$ that has (x, y) -entry

$$A_{x,y} = \begin{cases} 1 & \text{if } x, y \text{ are adjacent,} \\ 0 & \text{if } x, y \text{ are not adjacent} \end{cases} \quad (x, y \in X).$$

We call A the *adjacency matrix of Γ* . We recall the Cartesian product of graphs. Given graphs Γ and Γ' with vertex set X and X' , the Cartesian product $\Gamma \times \Gamma'$ is the graph with vertex set $X \times X'$ and adjacency defined as follows. Two vertices $(x, x'), (y, y')$ are adjacent in $\Gamma \times \Gamma'$ whenever either (i) x, y are adjacent in Γ and $x' = y'$ or (ii) $x = y$ and x', y' are adjacent in Γ' . In case (i) (resp. (ii)) we say that the edge $(x, x'), (y, y')$ in $\Gamma \times \Gamma'$ *comes from Γ* (resp. *comes from Γ'*). We make an observation.

Lemma 2.1 *Let Γ and Γ' denote connected graphs with vertex set X and X' , respectively. Consider a new graph obtained from $\Gamma \times \Gamma'$ by removing the edges that come from Γ . Then the connected components of the new graph are*

$$\{x\} \times X', \quad x \in X.$$

In particular, every connected component of the new graph is isomorphic to Γ' , and the number of connected components is $|X|$.

Definition 2.2 For an integer $n \geq 0$ and a set Y of size at least n , let the set $\binom{Y}{n}$ consist of the subsets of Y that have size n . Thus

$$\binom{Y}{n} = \{y \mid y \subseteq Y, |y| = n\}.$$

Definition 2.3 For sets Y, Z , let the set $Y \setminus Z$ consist of the elements in Y that are not in Z . Thus

$$Y \setminus Z = \{y \in Y \mid y \notin Z\}.$$

3 Q -polynomial distance-regular graphs

In this section we recall Q -polynomial distance-regular graphs and the subconstituent algebra. Let Γ denote a connected graph with vertex set X . For $x, y \in X$, let $\partial(x, y)$ denote the path-length distance between x, y . The integer

$$D = \max\{\partial(x, y) \mid x, y \in X\}$$

is called the *diameter* of Γ . For $x \in X$ and $0 \leq i \leq D$ define

$$\Gamma_i(x) = \{y \in X \mid \partial(x, y) = i\}.$$

The graph Γ is said to be *distance-regular* whenever for $0 \leq h, i, j \leq D$ and $x, y \in X$ at distance $\partial(x, y) = h$, the number

$$p_{i,j}^h = |\Gamma_i(x) \cap \Gamma_j(y)|$$

is a constant that is independent of x, y . The parameters $p_{i,j}^h$ are called the *intersection numbers* of Γ .

For the rest of this paper, we assume that Γ is distance-regular. Define

$$k_i = p_{i,i}^0 \quad (0 \leq i \leq D).$$

For $x \in X$ we have $|\Gamma_i(x)| = k_i$ for $0 \leq i \leq D$. The graph Γ is regular, with valency $k = k_1$ if $D \geq 1$ and $k = 0$ if $D = 0$. By the triangle inequality, the following hold for $0 \leq h, i, j \leq D$:

- (i) $p_{i,j}^h = 0$ if one of h, i, j is greater than the sum of the other two;

(ii) $p_{i,j}^h \neq 0$ if one of h, i, j is equal to the sum of the other two.

By [4, Lamma 4.1.7] all the intersection numbers are determined by the following intersection numbers:

$$c_i = p_{1,i-1}^i \quad (1 \leq i \leq D), \quad a_i = p_{1,i}^i \quad (0 \leq i \leq D), \quad b_i = p_{1,i+1}^i \quad (0 \leq i \leq D-1).$$

We have

$$c_i + a_i + b_i = k \quad (0 \leq i \leq D),$$

where $c_0 = 0$ and $b_D = 0$. We recall the Bose-Mesner algebra of Γ . For $0 \leq i \leq D$, define $A_i \in \text{Mat}_X(\mathbb{C})$ that has (x, y) -entry

$$(A_i)_{x,y} = \begin{cases} 1 & \text{if } \partial(x, y) = i, \\ 0 & \text{if } \partial(x, y) \neq i \end{cases} \quad (x, y \in X).$$

We have (i) $A_0 = I$; (ii) $J = \sum_{i=0}^D A_i$; (iii) $A_i^t = A_i$ ($0 \leq i \leq D$); (iv) $\overline{A_i} = A_i$ ($0 \leq i \leq D$); (v) $A_i A_j = A_j A_i = \sum_{h=0}^D p_{i,j}^h A_h$ ($0 \leq i, j \leq D$). Therefore, the matrices $\{A_i\}_{i=0}^D$ form a basis for a commutative subalgebra M of $\text{Mat}_X(\mathbb{C})$, called the *Bose-Mesner algebra* of Γ . The adjacency matrix A of Γ satisfies $A = A_1$ if $D \geq 1$ and $A = 0$ if $D = 0$. By [20, Corollary 3.4] A generates M . Since A is symmetric with real entries, A is diagonalizable over \mathbb{R} . Consequently M has a second basis $\{E_i\}_{i=0}^D$ such that (i) $E_0 = |X|^{-1} J$; (ii) $I = \sum_{i=0}^D E_i$; (iii) $E_i^t = E_i$ ($0 \leq i \leq D$); (iv) $\overline{E_i} = E_i$ ($0 \leq i \leq D$); (v) $E_i E_j = \delta_{i,j} E_i$ ($0 \leq i, j \leq D$). We call $\{E_i\}_{i=0}^D$ the *primitive idempotents* of Γ . We abbreviate $V = \mathbb{C}^X$. By construction,

$$V = \sum_{i=0}^D E_i V \quad (\text{direct sum}). \quad (1)$$

The summands in (1) are the eigenspaces of A . For $0 \leq i \leq D$ let θ_i denote the eigenvalue of A for $E_i V$. We have $A E_i = \theta_i E_i = E_i A$ ($0 \leq i \leq D$) and $A = \sum_{i=0}^D \theta_i E_i$. By [4, p 128] we have $\theta_0 = k$. We define some polynomials in an indeterminate λ . For $0 \leq i \leq D$ define

$$\begin{aligned} \tau_i(\lambda) &= (\lambda - \theta_0)(\lambda - \theta_1) \cdots (\lambda - \theta_{i-1}), \\ \eta_i(\lambda) &= (\lambda - \theta_D)(\lambda - \theta_{D-1}) \cdots (\lambda - \theta_{D-i+1}). \end{aligned}$$

For $0 \leq r \leq D$ we have

$$E_r = \prod_{\substack{0 \leq i \leq D \\ i \neq r}} \frac{A - \theta_i I}{\theta_r - \theta_i} = \frac{\tau_r(A) \eta_{D-r}(A)}{\tau_r(\theta_r) \eta_{D-r}(\theta_r)}. \quad (2)$$

Next we recall the Krein parameters of Γ . We have $A_i \circ A_j = \delta_{i,j} A_i$ ($0 \leq i, j \leq D$), where \circ denotes the entry-wise multiplication. Therefore, M is closed under \circ . Consequently, there exist scalars $q_{i,j}^h \in \mathbb{C}$ ($0 \leq h, i, j \leq D$) such that

$$E_i \circ E_j = |X|^{-1} \sum_{h=0}^D q_{i,j}^h E_h \quad (0 \leq i, j \leq D).$$

The scalars $q_{i,j}^h$ are called the *Krein parameters* of Γ . By [2, p. 69] the Krein parameters are non-negative real numbers.

Next we recall the Q -polynomial property. The graph Γ is said to be Q -polynomial (with respect to the ordering $\{E_i\}_{i=0}^D$) whenever the following hold for $0 \leq h, i, j \leq D$:

- (i) $q_{i,j}^h = 0$ if one of h, i, j is greater than the sum of the other two;
- (ii) $q_{i,j}^h \neq 0$ if one of h, i, j is equal the sum of the other two.

For the rest of this paper, we assume that Γ is Q -polynomial with respect to the ordering $\{E_i\}_{i=0}^D$.

Next we recall the dual Bose-Mesner algebra of Γ . Fix a vertex $x \in X$. For $0 \leq i \leq D$, define a diagonal matrix $E_i^* = E_i^*(x) \in \text{Mat}_X(\mathbb{C})$ that has (y, y) -entry

$$(E_i^*)_{y,y} = \begin{cases} 1 & \text{if } \partial(x, y) = i, \\ 0 & \text{if } \partial(x, y) \neq i \end{cases} \quad (y \in X).$$

We have (i) $I = \sum_{i=0}^D E_i^*$; (ii) $E_i^* E_j^* = \delta_{i,j} E_i^*$ ($0 \leq i, j \leq D$). Therefore the matrices $\{E_i^*\}_{i=0}^D$ form a basis for a commutative subalgebra $M^* = M^*(x)$ of $\text{Mat}_X(\mathbb{C})$. We call M^* the *dual Bose-Mesner algebra of Γ with respect to x* . We call $\{E_i^*\}_{i=0}^D$ the *dual primitive idempotents of Γ with respect to x* . We have

$$V = \sum_{i=0}^D E_i^* V \quad (\text{direct sum}). \quad (3)$$

Define a matrix $A^* = A^*(x) \in \text{Mat}_X(\mathbb{C})$ as follows. If $D = 0$ then $A^* = 0$. If $D \geq 1$ then A^* is diagonal with (y, y) -entry

$$(A^*)_{y,y} = |X|(E_1)_{x,y} \quad (y \in X).$$

The matrix A^* is called the *dual adjacency matrix of Γ with respect to x* . By [15, p. 279], there exist scalars $\theta_i^* \in \mathbb{C}$ ($0 \leq i \leq D$) such that

$$A^* = \sum_{i=0}^D \theta_i^* E_i^*.$$

By [15, Lemma 3.11], $\{\theta_i^*\}_{i=0}^D$ are mutually distinct. For $0 \leq i \leq D$, θ_i^* is the eigenvalue of A^* for the eigenspace $E_i^* V$, and $E_i^* A^* = A^* E_i^* = \theta_i^* E_i^*$. The scalars $\{\theta_i^*\}_{i=0}^D$ are called the *dual eigenvalues* of Γ . By [20, Corollary 11.6], A^* generates M^* . Let $T = T(x)$ denote the subalgebra of $\text{Mat}_X(\mathbb{C})$ generated by A and A^* . The algebra T is called the *subconstituent algebra of Γ with respect to x* .

4 T -modules

We continue to discuss the Q -polynomial distance-regular graph Γ with vertex set X . Fix $x \in X$ and write $T = T(x)$. Recall the standard module $V = \mathbb{C}^X$. In this section we discuss the T -modules.

By a T -module we mean a subspace W of V such that $TW \subseteq W$. We recall the notion of isomorphism for T -modules. Let W, W' denote T -modules. By a T -module isomorphism from W to W' we mean a \mathbb{C} -linear bijection $\sigma : W \rightarrow W'$ such that $\sigma B = B\sigma$ for all $B \in T$. The T -modules W and W' are said to be *isomorphic* whenever there exists a T -module isomorphism from W to W' .

A T -module W is said to be *irreducible* whenever $W \neq 0$ and W does not contain a T -submodule besides 0 and W . Let W denote an irreducible T -module. By the *endpoint of W* we mean

$$\min\{i \mid 0 \leq i \leq D, E_i^*W \neq 0\}.$$

By the *dual endpoint of W* we mean

$$\min\{i \mid 0 \leq i \leq D, E_iW \neq 0\}.$$

By [21, Lemma 5.6],

$$|\{i \mid 0 \leq i \leq D, E_iW \neq 0\}| = |\{i \mid 0 \leq i \leq D, E_i^*W \neq 0\}|. \quad (4)$$

By the *diameter of W* we mean the common value in (4) minus 1.

Lemma 4.1 (See [21, Lemma 5.4].) *Let W denote an irreducible T -module, with endpoint r , dual endpoint t , and diameter d . Then for $0 \leq i \leq D$ the following hold:*

- (i) $E_iW \neq 0$ if and only if $t \leq i \leq t + d$;
- (ii) $E_i^*W \neq 0$ if and only if $r \leq i \leq r + d$.

Lemma 4.2 (See [21, Lemma 5.7].) *Let W denote an irreducible T -module, with endpoint r , dual endpoint t , and diameter d . Then for $0 \leq i \leq d$ the following subspaces have the same dimension:*

$$E_{t+i}W, \quad E_{t+d-i}W, \quad E_{r+i}^*W, \quad E_{r+d-i}^*W. \quad (5)$$

Definition 4.3 We refer to the irreducible T -module W from Lemma 4.2. For $0 \leq i \leq d$ let ρ_i denote the common dimension of the four subspaces (5). By construction $\rho_i \neq 0$ and $\rho_i = \rho_{d-i}$. We call the sequence $\{\rho_i\}_{i=0}^d$ the *shape of W* .

Definition 4.4 Let W denote an irreducible T -module, with diameter d and shape $\{\rho_i\}_{i=0}^d$. Then W is said to be *thin* whenever $\rho_i = 1$ for $0 \leq i \leq d$.

Lemma 4.5 (See [15, Lemma 3.4].) *Every T -module is a direct sum of irreducible T -modules.*

By Lemma 4.5, there exist irreducible T -modules $\{W_h\}_{h=1}^n$ such that

$$V = \sum_{h=1}^n W_h \quad (\text{direct sum}). \quad (6)$$

Let W denote an irreducible T -module. By [5, Lemma 3.3], the number of summands in (6) that are isomorphic to W is independent of the choice of $\{W_h\}_{h=1}^n$. This number is denoted by $\text{mult}(W)$.

5 The nucleus

We continue to discuss the Q -polynomial distance-regular graph Γ with vertex set X . Fix $x \in X$ and write $T = T(x)$. In this section we recall the nucleus of Γ with respect to x .

Lemma 5.1 (See [21, Proposition 6.3].) *Let W denote an irreducible T -module, with endpoint r , dual endpoint t , and diameter d . Then*

$$r + t - D + d \geq 0. \quad (7)$$

Moreover, equality holds in (7) if and only if both $t = r$ and $d = D - 2r$. In this case W is thin.

Definition 5.2 (See [21, Definition 6.6].) Let W denote an irreducible T -module, with endpoint r , dual endpoint t , and diameter d . By the *displacement* of W we mean the integer

$$r + t - D + d.$$

Definition 5.3 (See [21, Definition 6.8].) By the *nucleus* of Γ with respect to x we mean the span of the irreducible T -modules that have displacement 0.

Lemma 5.4 (See [21, Proposition 6.10].) *Let W denote an irreducible T -submodule of the nucleus, with endpoint r , dual endpoint t , and diameter d . Then W is thin, and*

$$0 \leq r \leq D/2, \quad t = r, \quad d = D - 2r.$$

Definition 5.5 (See [21, Definition 7.2].) For $0 \leq i \leq d$, define

$$\mathcal{N}_i = (E_0^*V + E_1^*V + \cdots + E_i^*V) \cap (E_0V + E_1V + \cdots + E_{D-i}V).$$

Note 5.6 We have $\mathcal{N}_0 = E_0^*V$ and $\mathcal{N}_D = E_0V$.

Lemma 5.7 (See [21, Lemma 7.1].) *The following sum is direct:*

$$\mathcal{N}_0 + \mathcal{N}_1 + \cdots + \mathcal{N}_D.$$

Definition 5.8 (See [21, Definition 7.4].) Define a subspace \mathcal{N} by

$$\mathcal{N} = \mathcal{N}_0 + \mathcal{N}_1 + \cdots + \mathcal{N}_D.$$

Lemma 5.9 (See [21, Theorem 7.5].) *The following are the same:*

- (i) *the subspace $\mathcal{N} = \mathcal{N}(x)$ from Definition 5.8;*
- (ii) *The nucleus of Γ with respect to x .*

6 The Johnson graph $J(N, D)$

In this section we recall the Johnson graph. Let N, D denote integers such that $0 \leq D \leq N$. Let Ω denote a set of size N . The graph $J(N, D)$ has vertex set $\binom{\Omega}{D}$, and two vertices x, y are adjacent whenever $|x \cap y| = D - 1$. This graph $J(N, D)$ is called a *Johnson graph*. The graph $J(N, D)$ is isomorphic to $J(N, N - D)$. For this reason and to avoid degenerate situations, we assume that

$$N > 2D.$$

By [1, Section 2.10.3], the graph $J(N, D)$ is distance-regular with diameter D . For vertices x, y of $J(N, D)$ we have

$$\partial(x, y) = D - |x \cap y|.$$

By [4, Theorem 9.1.2], the intersection numbers of $J(N, D)$ satisfy

$$\begin{aligned} c_i &= i^2 & (1 \leq i \leq D), \\ b_i &= (D - i)(N - D - i) & (0 \leq i \leq D - 1). \end{aligned}$$

By [1, Proposition 2.87],

$$k_i = \binom{D}{i} \binom{N - D}{i} \quad (0 \leq i \leq D). \quad (8)$$

Consider the adjacency matrix A of $J(N, D)$. By [1, Theorem 2.95], the eigenvalues of A are

$$\theta_i = (D - i)(N - D - i) - i \quad (0 \leq i \leq D). \quad (9)$$

For $0 \leq i \leq D$, let E_i denote the primitive idempotent of A associated with θ_i . By [4, Corollary 8.4.2], $J(N, D)$ is Q -polynomial with respect to the ordering $\{E_i\}_{i=0}^D$. Fix a vertex x of $J(N, D)$, and abbreviate $T = T(x)$. By [1, Theorem 2.95], the eigenvalues of $A^* = A^*(x)$ are as follows. If $D = 0$ then $\theta_0^* = 0$. If $D \geq 1$ then

$$\theta_i^* = N - 1 - \frac{iN(N - 1)}{D(N - D)} \quad (0 \leq i \leq D). \quad (10)$$

7 The subgraph $\Gamma_i(x)$

We continue to discuss the Johnson graph $J(N, D)$ with the fixed vertex x . In this section we investigate the structure of the subgraph $\Gamma_i(x)$ for $0 \leq i \leq D$. We construct a graph isomorphism

$$\Gamma_i(x) \rightarrow J(D, i) \times J(N - D, i). \quad (11)$$

This isomorphism shows that the subgraph $\Gamma_i(x)$ has two types of edges, some come from $J(D, i)$ and some come from $J(N - D, i)$. We consider a new graph obtained from $\Gamma_i(x)$ by removing the edges that come from $J(D, i)$. We describe the connected components of the new graph.

The fixed vertex x is a subset of Ω with $|x| = D$. For the rest of this section, fix an integer i with $0 \leq i \leq D$. Consider the Johnson graph $J(D, i)$ with vertex set $\binom{x}{i}$, and the Johnson graph $J(N - D, i)$ with vertex set $\binom{\Omega \setminus x}{i}$. The Cartesian product $J(D, i) \times J(N - D, i)$ has vertex set

$$\binom{x}{i} \times \binom{\Omega \setminus x}{i}.$$

For $y \in \Gamma_i(x)$ we have

$$x \setminus y \in \binom{x}{i}, \quad y \setminus x \in \binom{\Omega \setminus x}{i}.$$

We consider a map

$$\Gamma_i(x) \rightarrow J(D, i) \times J(N - D, i) \quad (12)$$

that sends


$$y \mapsto (x \setminus y, y \setminus x), \quad y \in \Gamma_i(x).$$

Our next goal is to show that the map (12) is a graph isomorphism.

Lemma 7.1 *Let y, z denote vertices in $\Gamma_i(x)$. Then y, z are adjacent in the subgraph $\Gamma_i(x)$ if and only if one of the following holds:*

- (i) $x \setminus y = x \setminus z$, and $y \setminus x, z \setminus x$ are adjacent in $J(N - D, i)$;
- (ii) $y \setminus x = z \setminus x$, and $x \setminus y, x \setminus z$ are adjacent in $J(D, i)$.

Proof. Consider the Venn diagram for x, y, z :



Since $|x| = |y| = |z| = D$,

$$|a| + |e| + |f| + |g| = D, \quad (13)$$

$$|b| + |d| + |e| + |g| = D, \quad (14)$$

$$|c| + |d| + |f| + |g| = D. \quad (15)$$

Since $y, z \in \Gamma_i(x)$, we have $|x \cap y| = D - i$ and $|x \cap z| = D - i$. So

$$|e| + |g| = D - i, \quad |f| + |g| = D - i. \quad (16)$$

The vertices y, z are adjacent if and only if

$$|d| + |g| = D - 1. \quad (17)$$

Condition (i) holds if and only if

$$|e| = 0, \quad |f| = 0, \quad |d| = i - 1. \quad (18)$$

Condition (ii) holds if and only if

$$|b| = 0, \quad |c| = 0, \quad |a| = i - 1. \quad (19)$$

Using (13)–(16) we routinely find that (17) holds if and only if one of (18), (19) holds. \square

Lemma 7.2 *The map (12) is a graph isomorphism.*

Proof. We first show that the map (12) is a bijection. Clearly, the map (12) is injective. By (8) the graphs $\Gamma_i(x)$ and $J(D, i) \times J(N - D, i)$ have the same number of vertices. By these comments, the map (12) is a bijection. The map (12) respects adjacency by Lemma 7.1. The result follows. \square

Referring to the isomorphism (12), consider a new graph that is obtained from the subgraph $\Gamma_i(x)$ by removing the edges that come from $J(D, i)$. Our next goal is to describe

the connected components of the new graph. By Lemma 2.1, these connected components are the preimages under (12) of the sets

$$\{\alpha\} \times \binom{\Omega \setminus x}{i}, \quad \alpha \in \binom{x}{i}.$$

Lemma 7.3 *Referring to the isomorphism (12), for $\alpha \in \binom{x}{i}$, the preimage of $\{\alpha\} \times \binom{\Omega \setminus x}{i}$ is the set*

$$\{y \in X \mid x \setminus y = \alpha\}. \quad (20)$$

Proof. Clear by construction. \square

Lemma 7.4 *Consider a new graph obtained from the subgraph $\Gamma_i(x)$ by removing the edges that come from $J(D, i)$. For the new graph the connected components are*

$$\{y \in X \mid y \cap x = \alpha\}, \quad \alpha \subseteq x, \quad |\alpha| = D - i. \quad (21)$$

Proof. By the claim above Lemma 7.3 and Lemmas 7.2 and 7.3. \square

Corollary 7.5 *Referring to Lemma 7.4, each connected component of the new graph is isomorphic to $J(N - D, i)$. For the new graph the number of the connected components is $\binom{D}{i}$.*

Proof. By Lemmas 7.3 and 7.4. \square

8 The vectors α^\vee and $\alpha^\mathcal{N}$

We continue to discuss the Johnson graph $J(N, D)$ with the fixed vertex x . Recall the standard module $V = \mathbb{C}^X$. We introduce some vectors in V that will form a basis of the nucleus with respect to x .

Definition 8.1 For a subset $\alpha \subseteq x$, define vectors in V :

$$\alpha^\vee = \sum_{\substack{y \in X \\ \alpha \subseteq y}} \hat{y}, \quad \alpha^\mathcal{N} = \sum_{\substack{y \in X \\ y \cap x = \alpha}} \hat{y}.$$

The vector α^\vee is the characteristic vector of the set of vertices in X that contain α , and $\alpha^\mathcal{N}$ is the characteristic vector of the set of vertices in X whose intersection with x is equal to α .

The vectors α^\vee and $\alpha^\mathcal{N}$ are related as follows.

Lemma 8.2 For $\alpha \subseteq x$ the following hold:

$$(i) \alpha^\vee = \sum_{\alpha \subseteq \beta \subseteq x} \beta^\mathcal{N}.$$

$$(ii) \alpha^\mathcal{N} = \sum_{\alpha \subseteq \beta \subseteq x} (-1)^{|\beta| - |\alpha|} \beta^\vee.$$

Proof. (i) We argue

$$\alpha^\vee = \sum_{\substack{y \in X \\ \alpha \subseteq y}} \hat{y} = \sum_{\alpha \subseteq \beta \subseteq x} \sum_{\substack{y \in X \\ y \cap x = \beta}} \hat{y} = \sum_{\alpha \subseteq \beta \subseteq x} \beta^\mathcal{N}.$$

(ii) Using (i) we obtain

$$\begin{aligned} \sum_{\alpha \subseteq \beta \subseteq x} (-1)^{|\beta| - |\alpha|} \beta^\vee &= \sum_{\alpha \subseteq \beta \subseteq x} (-1)^{|\beta| - |\alpha|} \sum_{\beta \subseteq \gamma \subseteq x} \gamma^\mathcal{N} \\ &= \sum_{\alpha \subseteq \gamma \subseteq x} \gamma^\mathcal{N} \sum_{\alpha \subseteq \beta \subseteq \gamma} (-1)^{|\beta| - |\alpha|}. \end{aligned}$$

For $\alpha \subseteq \gamma \subseteq x$ we have

$$\begin{aligned} \sum_{\alpha \subseteq \beta \subseteq \gamma} (-1)^{|\beta| - |\alpha|} &= \sum_{j=0}^{|\gamma| - |\alpha|} \binom{|\gamma| - |\alpha|}{j} (-1)^j \\ &= (1 - 1)^{|\gamma| - |\alpha|} \\ &= \begin{cases} 1 & \text{if } \gamma = \alpha, \\ 0 & \text{if } \gamma \neq \alpha. \end{cases} \end{aligned}$$

The result follows. \square

Lemma 8.3 For $\alpha \subseteq x$,

$$\alpha^\mathcal{N} = E_i^* \alpha^\vee,$$

where $i = D - |\alpha|$.

Proof. For $y \in X$ and $0 \leq j \leq D$, we have $j = D - |x \cap y|$ if and only if $y \in \Gamma_j(x)$. The result follows from this and Definition 8.1. \square

By Lemma 7.4,

$$\{\alpha^\mathcal{N} \mid \alpha \subseteq x\} \quad \text{are linearly independent.} \quad (22)$$

By Lemma 8.3 and (22),

$$\{\alpha^\vee \mid \alpha \subseteq x\} \quad \text{are linearly independent.} \quad (23)$$

9 The action of A, A^* on the vectors $\alpha^\vee, \alpha^\mathcal{N}$

We continue to discuss the Johnson graph $J(N, D)$ with the fixed vertex x . Recall the vectors $\alpha^\vee, \alpha^\mathcal{N}$ from Definition 8.1. In this section we display the action of A, A^* on these vectors.

Theorem 9.1 For $\alpha \subseteq x$,

$$A\alpha^\vee = \theta_{|\alpha|}\alpha^\vee + (D - |\alpha| + 1) \sum_{\substack{\beta \subseteq \alpha \\ |\beta| = |\alpha| - 1}} \beta^\vee. \quad (24)$$

Proof. Let the set S_0 consist of the vertices in X that include α . The characteristic vector of S_0 is equal to α^\vee . Let the set S_1 consist of the vertices in $X \setminus S_0$ that are adjacent to at least one vertex in S_0 . The characteristic vector of S_1 is equal to

$$\sum_{\substack{\beta \subseteq \alpha \\ |\beta| = |\alpha| - 1}} (\beta^\vee - \alpha^\vee).$$

By combinatorial counting, we find that each vertex in S_0 is adjacent to exactly $(D - |\alpha|)(N - D)$ vertices in S_0 . Also, each vertex in S_1 is adjacent to exactly $D - |\alpha| + 1$ vertices in S_0 . By these comments,

$$A\alpha^\vee = (D - |\alpha|)(N - D)\alpha^\vee + (D - |\alpha| + 1) \sum_{\substack{\beta \subseteq \alpha \\ |\beta| = |\alpha| - 1}} (\beta^\vee - \alpha^\vee).$$

By this and (9) we obtain (24). \square

Theorem 9.2 For $\alpha \subseteq x$,

$$\begin{aligned} A\alpha^\mathcal{N} &= (D - |\alpha|)(N - 2D + |\alpha|)\alpha^\mathcal{N} + (N - 2D + |\alpha| + 1) \sum_{\substack{\alpha \subseteq \beta \subseteq x \\ |\beta| = |\alpha| + 1}} \beta^\mathcal{N} \\ &+ \sum_{\substack{\beta \subseteq x \\ |\beta| = |\alpha| \\ |\beta \cap \alpha| = |\alpha| - 1}} \beta^\mathcal{N} + (D - |\alpha| + 1) \sum_{\substack{\beta \subseteq \alpha \\ |\beta| = |\alpha| - 1}} \beta^\mathcal{N}. \end{aligned}$$

Proof. The vector $\alpha^\mathcal{N}$ is the characteristic vector of the set

$$\{y \in X \mid y \cap x = \alpha\}. \quad (25)$$

So we have

$$A\alpha^\mathcal{N} = \sum_{z \in X} e_z \widehat{z},$$

where e_z is the number of edges from z into the set (25). We now compute e_z for $z \in X$. Let z be given and define $\beta = z \cap x$. By combinatorial counting, we find that $e_z = 0$ or e_z is given in the table below.

Case	e_z
$\beta = \alpha$	$(D - \alpha)(N - 2D + \alpha)$
$\alpha \subseteq \beta$ and $ \beta = \alpha + 1$	$N - 2D + \alpha + 1$
$ \beta = \alpha $ and $ \beta \cap \alpha = \alpha - 1$	1
$\beta \subseteq \alpha$ and $ \beta = \alpha - 1$	$D - \alpha + 1$

Using the data in the table, we routinely obtain the result. \square

Theorem 9.3 For $\alpha \subseteq x$,

$$A^* \alpha^N = \theta_{D-|\alpha|}^* \alpha^N.$$

Proof. Clear since $\alpha^N \in E_i^* V$, where $i = D - |\alpha|$. \square

Theorem 9.4 For $\alpha \subseteq x$,

$$A^* \alpha^\vee = \theta_{D-|\alpha|}^* \alpha^\vee + \frac{N(N-1)}{D(N-D)} \sum_{\substack{\alpha \subseteq \gamma \subseteq x \\ |\gamma| = |\alpha| + 1}} \gamma^\vee.$$

Proof. Using Lemma 8.2(i), Theorem 9.3, and Lemma 8.2(ii) in order,

$$\begin{aligned} A^* \alpha^\vee &= \sum_{\alpha \subseteq \beta \subseteq x} A^* \beta^N. \\ &= \sum_{\alpha \subseteq \beta \subseteq x} \theta_{D-|\beta|}^* \beta^N \\ &= \sum_{\alpha \subseteq \beta \subseteq x} \theta_{D-|\beta|}^* \sum_{\beta \subseteq \gamma \subseteq x} (-1)^{|\gamma|-|\beta|} \gamma^\vee. \end{aligned}$$

Thus

$$A^* \alpha^\vee = \sum_{\alpha \subseteq \gamma \subseteq x} \gamma^\vee \sum_{\alpha \subseteq \beta \subseteq \gamma} (-1)^{|\gamma|-|\beta|} \theta_{D-|\beta|}^*. \quad (26)$$

For $\alpha \subseteq \beta \subseteq \gamma$ we have

$$\begin{aligned} \sum_{\alpha \subseteq \beta \subseteq \gamma} (-1)^{|\gamma|-|\beta|} \theta_{D-|\beta|}^* &= \sum_{i=|\alpha|}^{|\gamma|} (-1)^{|\gamma|-i} \binom{|\gamma|-\alpha}{i-|\alpha|} \theta_{D-i}^* \\ &= \sum_{j=0}^{|\gamma|-|\alpha|} (-1)^{|\gamma|-|\alpha|-j} \binom{|\gamma|-|\alpha|}{j} \theta_{D-|\alpha|-j}^*. \end{aligned}$$

Evaluating this using (10) we obtain

$$\sum_{\alpha \subseteq \beta \subseteq \gamma} (-1)^{|\gamma|-|\beta|} \theta_{D-|\beta|}^* = \begin{cases} 0 & \text{if } |\gamma| - |\alpha| \geq 2, \\ \frac{N(N-1)}{D(N-D)} & \text{if } |\gamma| - |\alpha| = 1, \\ \theta_{D-|\alpha|}^* & \text{if } \gamma = \alpha \end{cases}$$

By this and (26) we get the result. \square

10 Some results concerning the vectors α^\vee and $\alpha^\mathcal{N}$

We continue to discuss the Johnson graph $J(N, D)$ with the fixed vertex x . Recall the vector α^\vee and $\alpha^\mathcal{N}$ from Definition 8.1. In this section we show that the vector α^\vee and $\alpha^\mathcal{N}$ are contained in the nucleus of $J(N, D)$ with respect to x .

The following result is known [14]. We give a short direct proof for the sake of completeness.

Lemma 10.1 (See [14, Example (7.1)].) *For $\alpha \subseteq x$,*

$$\alpha^\vee \in E_0V + E_1V + \cdots + E_{|\alpha|}V.$$

Proof. We may assume that $\alpha \neq x$; otherwise the result is trivial. We have $|\alpha| < D$. By Theorem 9.1, for $\gamma \subseteq x$ we have

$$(A - \theta_{|\gamma|}I)\gamma^\vee \in \text{Span}\{\beta^\vee \mid \beta \subseteq \gamma, |\beta| = |\gamma| - 1\}.$$

Using this and induction on $|\alpha|$, we obtain

$$\tau_{|\alpha|+1}(A)\alpha^\vee = 0.$$

By this and (2),

$$E_r\alpha^\vee = 0 \quad (|\alpha| < r \leq D).$$

The result follows. \square

Lemma 10.2 *For $\alpha \subseteq x$,*

$$\alpha^\vee \in E_0^*V + E_1^*V + \cdots + E_{D-|\alpha|}^*V.$$

Proof. For $y \in X$ we have $\hat{y} \in E_i^*V$, where $i = D - |y \cap x|$. By Definition 8.1,

$$\alpha^\vee = \sum_{\substack{y \in X \\ \alpha \subseteq y}} \hat{y}.$$

By these comments we get the result. \square

Lemma 10.3 For $\alpha \subseteq x$,

$$\alpha^\vee \in \mathcal{N}_{D-|\alpha|}. \quad (27)$$

Proof. By Definition 5.5 and Lemmas 10.1, 10.2. \square

Lemma 10.4 For $\alpha \subseteq x$,

$$\alpha^N \in E_{D-|\alpha|}^* \mathcal{N}.$$

Proof. By Lemmas 8.3 and 10.3. \square

11 Some bases for the nucleus

We continue to discuss the Johnson graph $J(N, D)$ with the fixed vertex x . In this section we obtain some bases for the nucleus of with respect to x . Let W denote an irreducible T -module, with endpoint r , dual endpoint s , and diameter d . By [11, Section 2.3], the isomorphism class of W is determined by r, t, d . Assume that W has displacement 0. Then by Lemma 5.1, $r = t$ and $0 \leq r \leq D/2$ and $d = D - 2r$. In this case the isomorphism class of W is determined by r . Abbreviate

$$\text{mult}_r = \text{mult}(W).$$

Lemma 11.1 (See [7, Theorem 10.2], [11, Remark 2].) *We have $\text{mult}_0 = 1$ and*

$$\text{mult}_r = \binom{D}{r} - \binom{D}{r-1} \quad (1 \leq r \leq D/2).$$

Lemma 11.2 For $0 \leq i \leq D$,

$$\dim E_i^* \mathcal{N} = \binom{D}{i}. \quad (28)$$

Proof. By Lemmas 4.5 and 5.4, there exist irreducible T -modules $\{W_h\}_{h=1}^m$ with displacement 0 such that

$$\mathcal{N} = \sum_{h=1}^m W_h \quad (\text{direct sum}).$$

So

$$\dim E_i^* \mathcal{N} = \sum_{h=1}^m \dim E_i^* W_h.$$

For $1 \leq h \leq m$ we have $\dim E_i^* W_h = \dim E_{D-i}^* W_h$ by Lemmas 4.2 and 5.4. So $\dim E_i^* \mathcal{N} = \dim E_{D-i}^* \mathcal{N}$. By this we may assume $0 \leq i \leq D/2$. For $1 \leq h \leq m$ let r_h denote the endpoint of W_h . By Lemma 5.1 and the definition of the endpoint, we have

$$\dim E_i^* W_h = \begin{cases} 0 & \text{if } i < r_h, \\ 1 & \text{if } r_h \leq i \end{cases} \quad (1 \leq h \leq m).$$

For $0 \leq r \leq i$, the number of h ($1 \leq h \leq m$) such that $r_h = r$ is equal to mult_r . By these comments,

$$\dim E_i^* \mathcal{N} = \sum_{r=0}^i \text{mult}_r.$$

Evaluating this sum using Lemma 11.1, we routinely obtain (28). \square

Lemma 11.3 *We have*

- (i) $\dim \mathcal{N} = 2^D$;
- (ii) $\dim \mathcal{N}_i = \binom{D}{i}$ for $0 \leq i \leq D$.

Proof. (i) We have $\mathcal{N} = \sum_{i=0}^D E_i^* \mathcal{N}$ (direct sum). By this and Lemma 11.2, we get the result.

(ii) By Lemma 5.7, $\mathcal{N} = \sum_{i=0}^D \mathcal{N}_i$ (direct sum). By this and (i),

$$2^D = \dim \mathcal{N} = \sum_{i=0}^D \dim \mathcal{N}_i.$$

By (23) and (27),

$$\dim \mathcal{N}_i \geq \binom{D}{i} \quad (0 \leq i \leq D).$$

By these comments we get the result. \square

Theorem 11.4 *The following hold:*

- (i) *The vectors $\{\alpha^{\mathcal{N}}\}_{\alpha \subseteq x}$ form a basis of \mathcal{N} .*
- (ii) *The vectors $\{\alpha^{\vee}\}_{\alpha \subseteq x}$ form a basis of \mathcal{N} .*
- (iii) *For $0 \leq i \leq D$, the vectors $\{\alpha^{\mathcal{N}} \mid \alpha \subseteq x, |\alpha| = D - i\}$ form a basis of $E_i^* \mathcal{N}$.*
- (iv) *For $0 \leq i \leq D$, the vectors $\{\alpha^{\vee} \mid \alpha \subseteq x, |\alpha| = D - i\}$ form a basis of \mathcal{N}_i .*

Proof. (i) By Lemma 10.4, $\alpha^{\mathcal{N}} \in \mathcal{N}$. Now the result follows from (22) and Lemma 11.3(i).

(ii) By Lemma 10.3, $\alpha^{\vee} \in \mathcal{N}$. Now the result follows from (23) and Lemma 11.3(i).

(iii) By Lemma 10.4, $\alpha^{\mathcal{N}} \in E_i^* \mathcal{N}$ if $|\alpha| = D - i$. Now the result follows from Lemma 11.2 and (22).

(iv) By Lemma 10.3, $\alpha^{\vee} \in \mathcal{N}_i$ if $|\alpha| = D - i$. Now the result follows from (23) and Lemma 11.3(ii). \square

Recall by Lemma 7.2 that for $0 \leq i \leq D$ the subgraph $\Gamma_i(x)$ is isomorphic to $J(D, i) \times J(N - D, i)$.

Corollary 11.5 *For $0 \leq i \leq D$, consider a new graph obtained from the subgraph $\Gamma_i(x)$ by removing the edges that come from $J(D, i)$. Then the vectors in Theorem 11.4(iii) are the characteristic vectors of the connected components of the new graph.*

Proof. By Lemma 7.4 the sets

$$\{y \in X \mid y \cap x = \alpha\}, \quad \alpha \subseteq x, |\alpha| = D - i$$

are the connected components of the new graph. Recall that α^N is the characteristic vector of $\{y \in X \mid y \cap x = \alpha\}$. By these comments, we get the result. \square

References

- [1] E. Bannai, Et. Bannai, T. Ito, R. Tanaka, Algebraic Combinatorics, De Gruyter Series in Discrete Mathematics and Applications 5, De Gruyter, 2021.
- [2] E. Bannai, T. Ito, Algebraic Combinatorics, I. Association Schemes, Benjamin/Cummings, Menlo Park, CA, 1984.
- [3] N. L. Biggs, Algebraic Graph Theory, Cambridge Mathematical Library 67, Cambridge University Press, London 1974.
- [4] A. E. Brouwer, A. M. Cohen, A. Neumaier, Distance-Regular Graphs, Springer-Verlag, Berlin, 1989.
- [5] B. Curtin, Bipartite distance-regular graphs, part I, Graphs Combin. 15 (1999) 143–158.
- [6] C. Curtis, I. Reiner, Representation Theory of Finite Groups and Associative Algebras, AMS Chelsea Publishing, 1962.
- [7] J. Go, The Terwilliger algebra of the hypercube, European J. Combin. 23 (2002) 399–429.
- [8] H. Huang, The Clebsch-Gordan coefficients of $U(\mathfrak{sl}_2)$ and the Terwilliger algebras of Johnson graphs, J. Combin. Theory A 203 (2024), 105833.
- [9] T. Ito, K. Tanabe, P. Terwilliger, Some algebra related to P - and Q -polynomial association schemes, Codes and Association Schemes (Piscataway NJ, 1999) 167–192.
- [10] X. Liang, Y. Tan, T. Ito, An observation on Leonard system parameters for the Terwilliger algebra of the Johnson scheme $J(N, D)$, Graphs Combin. 33 (2017) 149–156.
- [11] Y. Tan, Y. Z. Fan, T. Ito, X. Liang, The Terwilliger algebra of the Johnson scheme $J(N, D)$ revisited from the viewpoint of group representations, European J. Combin. 80 (2019) 157–171.

- [12] A. Pascasio, On the multiplicities of the primitive idempotents of a Q -polynomial distance-regular graph, *European J. Combin.* 23 (2002) 1073–1078.
- [13] J. Rotman, *Advanced Modern Algebra*, 2nd edition, AMS, Providence, Rhode Island, 2010.
- [14] H. Tanaka, Vertex subsets with minimal width and dual width in Q -polynomial distance-regular graphs, *Electron. J. Combin.* 18 (2011) P167.
- [15] P. Terwilliger, The subconstituent algebra of an association scheme I, *J. Algebraic Combin.* 1 (1992) 363–388.
- [16] P. Terwilliger, The subconstituent algebra of an association scheme II, *J. Algebraic Combin.* 2 (1993) 73–103.
- [17] P. Terwilliger, The subconstituent algebra of an association scheme III, *J. Algebraic Combin.* 2 (1993) 177–210.
- [18] P. Terwilliger, The displacement and split decompositions for a Q -polynomial distance-regular graph, *Graphs Combin.* 21 (2005) 263–276.
- [19] P. Terwilliger, Notes on the Leonard system classification, *Graphs Combin.* 37 (2021) 1687–1748.
- [20] P. Terwilliger, Distance-regular graphs, the subconstituent algebra, and the Q -polynomial property, *London Math. Soc. Lecture Note Ser.*, 487, Cambridge University Press, London, 2024, 430–491.
- [21] P. Terwilliger, The nucleus of a Q -polynomial distance-regular graph, preprint; arXiv:2408.11282.

Kazumasa Nomura
 Institute of Science Tokyo
 Kohnodai, Ichikawa, 272-0827 Japan
 email: knomura@pop11.odn.ne.jp

Paul Terwilliger
 Department of Mathematics
 University of Wisconsin
 480 Lincoln Drive
 Madison, Wisconsin, 53706 USA
 email: terwilli@math.wisc.edu

Keywords. Johnson graph, distance-regular graph, Q -polynomial. nucleus.
2020 Mathematics Subject Classification. 05E30, 15B10.