

CLOSURE OF KNITTED SURFACES AND SURFACE-LINKS

INASA NAKAMURA AND JUMPEI YASUDA

ABSTRACT. A knitted surface is a surface with boundary properly embedded in $D^2 \times B^2$, which is a generalization of a braided surface. A knitted surface is called a 2-dimensional knit if its boundary is the closure of a trivial braid. From a 2-dimensional knit S , we obtain a surface-link in \mathbb{R}^4 by taking the closure of S . We show that any surface-link is ambiently isotopic to the closure of some 2-dimensional knit. Further, we consider another type of the closure of a knitted surface, called the plat closure. It is known that any trivial surface-knot is ambiently isotopic to the plat closure of a knitted surface of degree 2. We show that the plat closure of any knitted surface of degree 2 is a trivial surface-link, and any trivial surface-link is ambiently isotopic to the plat closure of a knitted surface of degree 2. We also show the same result for the closure of 2-dimensional knits of degree 2.

1. INTRODUCTION

Let n be a positive integer. An n -knit [1, 13] is a tangle obtained from an n -braid in a cylinder $D^2 \times I$ by splicing at some crossings. Splicing a crossing results in a ‘‘hook pair’’, and we describe the information of each hook pair by a segment called a pairing [14]. We give an equivalence relation to the set of n -knits so that two n -knits are equivalent if they are related by an isotopy of n -knits. The set of equivalence classes of n -knits forms a monoid D_n by the multiplication induced from the composition of knits, which is called the n -knit monoid. The n -knit monoid D_n is generated by $3(n-1)$ elements, $\sigma_1, \dots, \sigma_{n-1}, \sigma_1^{-1}, \dots, \sigma_{n-1}^{-1}, \tau_1, \dots, \tau_{n-1}$, where σ_i and σ_i^{-1} are the standard generator and its inverse of the n -braid group B_n , and τ_i is the hook pair between the i th and $(i+1)$ th strings as in Figure 1 ($i = 1, \dots, n-1$). Let e be the unit element of D_n , which is represented by the trivial braid. The n -knit monoid D_n has the monoid presentation

$$\left\langle \begin{array}{l} \sigma_1, \dots, \sigma_{n-1}, \\ \sigma_1^{-1}, \dots, \sigma_{n-1}^{-1}, \\ \tau_1, \dots, \tau_{n-1} \end{array} \middle| \begin{array}{l} \sigma_i \sigma_i^{-1} = \sigma_i^{-1} \sigma_i = e, \quad \sigma_i \tau_i = \tau_i \sigma_i = \tau_i, \\ \sigma_i \sigma_k = \sigma_k \sigma_i, \quad \sigma_i \tau_k = \tau_k \sigma_i, \quad \tau_i \tau_k = \tau_k \tau_i \quad (|i-k| > 1), \\ \sigma_i \sigma_j \sigma_i = \sigma_j \sigma_i \sigma_j, \quad \sigma_i \sigma_j \tau_i = \tau_j \sigma_i \sigma_j \quad (|i-j| = 1) \end{array} \right\rangle_{\text{monoid}}.$$

A knitted surface [14] in a bidisk $D^2 \times B^2$ is defined as an analogue to the notion of an n -knit in $D^2 \times I$. A knitted surface is an extended notion of a braided surface [15], which is a surface in $D^2 \times B^2$ in the form of a branched covering over B^2 . A knitted surface is constructed as the

2020 *Mathematics Subject Classification*. Primary: 57K45, Secondary: 57Q35 57K10.

Key words and phrases. surfaces in 4-space; surface-links; surface braids; tangles; knits; closure; plat closure.




FIGURE 1. Generators σ_i , σ_i^{-1} and τ_i of the monoid D_n of n -knits.

trace of deformations of knits, as follows. Let $\beta_0 \rightarrow \beta_1 \rightarrow \beta_2 \rightarrow \cdots \rightarrow \beta_m$ be a sequence of elements of the free monoid consisting of σ_i, σ_i^{-1} and τ_i ($i = 1, \dots, n-1$) such that β_{j-1} and β_j are related by one of the following for each j ($j = 1, \dots, m$):

- (1) One of the relations of the n -knot monoid D_n .
- (2) Replacement of e and σ_i^ϵ , or replacement of e and τ_i , for some i and $\epsilon \in \{+1, -1\}$.
- (3) Replacement of τ_i and $\tau_i \tau_i$ for some i .

We decompose $I = [0, 1]$ into m subintervals: $I = I_1 \cup \cdots \cup I_m$, $I_j = [(j-1)/m, j/m]$ ($j = 1, \dots, m$). A knitted surface S in $D^2 \times I \times I$ is constructed for each $S \cap (D^2 \times I \times I_j)$ as follows. We denote by the same notation β_j a geometric n -knot given by the presentation β_j . For case (1), let $\{S_{[t]}\}_{t \in I_j}$ be an isotopy of n -knits in $D^2 \times I$ relating β_{j-1} and β_j which describes the knot monoid relation. Then we construct $S \cap (D^2 \times I \times I_j)$ by $\cup_{t \in I_j} S_{[t]} \times \{t\}$. This construction is called the *trace* of $\{S_{[t]}\}_{t \in I_j}$. The construction is determined by giving $S_{[t]}$, which is called a *slice* of S at t .

For case (2), we construct $S \cap (D^2 \times I \times I_j)$ by giving slices as follows: $S_{[t]} = \beta_{j-1}$ ($t \in [(j-1)/m, (2j-1)/2m)$), $S_{[t]} = \beta_j$ ($t \in (2j-1)/2m, j/m]$) and $S_{[t]}$ ($t = (2j-1)/2m$) is the union of β_{j-1} and a twisted band (respectively, an untwisted band) corresponding to the σ_i^ϵ (respectively τ_i). See Figure 2. The n -knot β_j is the result of band surgery along a band attaching to β_{j-1} , and the band corresponds to a branch point of the knitted surface.

For case (3), let $S_{[t]} = \beta_{j-1}$ ($t \in [(j-1)/m, (2j-1)/2m)$), $S_{[t]} = \beta_j$ ($t \in ((2j-1)/2m, j/m]$) and let $S_{[t]}$ ($t = (2j-1)/2m$) be the union of β_{j-1} and a disk whose boundary is the simple closed curve formed by the $\tau_i \tau_i$. See Figure 3. The knitted surface is the trace of deformations of knits where a simple closed curve is born or dies.

A *surface-link* is a closed surface smoothly embedded in \mathbb{R}^4 . In particular, a surface-link with one component is called a *surface-knot*. A braided/knitted surface is called a *2-dimensional braid/knit* if its boundary is the closure of the trivial braid. From a 2-dimensional braid S , we obtain an orientable surface-link in \mathbb{R}^4 by taking the closure of S , and Kamada [7] showed that every orientable surface-link is ambiently isotopic to the closure of some 2-dimensional braid. For a 2-dimensional knot S , we can also define the closure of S by a similar way, which provides a surface-link. In this paper, we show Claim 9.1 in [14] as follows.

Theorem 1.1. *Every surface-link in \mathbb{R}^4 is ambiently isotopic to the closure of some 2-dimensional knit.*

Further, we consider another type of the closure of a knitted surface, called the plat closure. In [14, Theorem 8.1], we showed that any trivial surface-knot is ambiently isotopic to the plat closure of some knitted surface of degree 2. In this paper, we show the following.

Theorem 1.2. *The plat closure of any knitted surface of degree 2 is a trivial surface-link, and any trivial surface-link is ambiently isotopic to the plat closure of a knitted surface of degree 2.*

Theorem 1.2 follows from Corollary 5.7 and Theorem 5.13, where Theorem 5.13 is [14, Claim 8.3]. We show this by using a chart description of a knitted surface, which is a generalization of a chart description of a simple braided surface [6, 9]. As a corollary of Theorem 1.2, we have the following.

Corollary 1.3. *The closure of any 2-dimensional knit degree 2 is a trivial surface-link, and any trivial surface-link is ambiently isotopic to the closure of some 2-dimensional knit of degree 2.*

The paper is organized as follows. In Section 2, we review knitted surfaces. In Section 3, we review the closure of knitted surfaces, and we show Theorem 1.1. In Section 4, we review a chart description of knitted surfaces of degree 2. In Section 5, we show Theorems 5.13 and 1.2 and Corollary 1.3. We give references [3, 5, 10] for the basics of 1-dimensional and 2-dimensional knot theory. In this paper, we assume that surfaces and tangles are smooth.

2. KNITTED SURFACES

In this section, we review knitted surfaces [14].

2.1. Motion picture method. In order to describe surfaces in 4-space, we use motion pictures, regarding surfaces as the trace of slices of the surfaces by 3-spaces along the 4th dimensional axis. In this section, we review motion picture method.

Let S be a surface properly embedded in $D^2 \times B^2$, and we denote $B^2 = I_3 \times I_4$, where $I_3 = I_4 = [0, 1]$. We modify the saddle points and minimal/maximal points of S into saddle bands and minimal/maximal disks, respectively; see [5, Section 8.4]. Let $p : D^2 \times B^2 \rightarrow D^2 \times I_3$ be the projection. For each $t \in I_4$, the *slice* of S at t , denoted by $S_{[t]}$, is defined as the image by p of the intersection $S \cap (D^2 \times I_3 \times \{t\})$: that is, that is, $S_{[t]} = p(S \cap (D^2 \times B^2))$.

When each slice $S_{[t]}$ is either (1) a tangle (or a link) or (2) a union of a tangle and several disks (and bands), we describe S by a finite sequence of diagrams of $S_{[t_0]}, \dots, S_{[t_m]}$ consisting unions of (2) and tangles before and after the time of (2). This sequence of diagrams is called a *motion picture* of S .

2.2. Band surgery of a tangle along a band set. Let β be a tangle in $D^2 \times I$. A *band* attaching to β is a 2-disk B in $D^2 \times I$ such that $B \cap \beta$ consists of a pair of intervals in ∂B . The result of *band surgery along* B ,

band attaching to e so that the result of the band surgery along B is s_i . Then we define S by

$$S_{[t]} = \begin{cases} e & (0 \leq t < \frac{1}{2}) \\ e \cup B & (t = \frac{1}{2}) \\ s_i & (\frac{1}{2} < t \leq 1), \end{cases}$$

and for $s_i \rightarrow e$, the presented surface S' is given by the mirror image of S with respect to $D^2 \times \{1/2\}$, that is, the surface S' given by the slice $S'_{[t]} = S_{[1-t]}$ for each $t \in [0, 1]$.

(Case 3) A transformation $\tau_i \leftrightarrow \tau_i\tau_i$. In this case, we define S by the motion picture as given in Figure 3; we define S by slices as follow: we define S for $\tau_i \rightarrow \tau_i\tau_i$; the other case is given by the same method as in (Case 2). We denote by D a disk whose boundary is the simple closed curve in $\tau_i\tau_i$. Further, we assume that $\tau_i = \tau_i\tau_i \setminus \partial D$. Then we define S by

$$S_{[t]} = \begin{cases} \tau_i & (0 \leq t < \frac{1}{2}) \\ \tau_i \cup D & (t = \frac{1}{2}) \\ \tau_i\tau_i & (\frac{1}{2} < t \leq 1). \end{cases}$$




FIGURE 3. Motion picture of the knitted surfaces presented by $\tau_i \leftrightarrow \tau_i\tau_i$, where we omit the j th strings for $j \neq i, i + 1$. This figure is similar to [14, Figure 3].

For a transformation $\beta \rightarrow \beta'$ such that β' is obtained from β by a finite number of mutually disjoint transformations as given in (Case 1)–(Case 3), the knitted surface presented by $\beta \rightarrow \beta'$ is defined as the surface determined from those of (Case 1)–(Case 3). We remark that the construction is called the “vertical product” [5, Section 15.1].

2.4. Knitted surfaces. Knitted surfaces are defined by using “pairings” and “knit structure” [14]. Two knitted surfaces are said to be equivalent if they are related by an “isotopy of knitted surfaces”. Every knitted surface is equivalent to a knitted surface in a normal form. In this paper, we give a knitted surface as a knitted surface derived from a normal form, by using deformations of knits.

Definition 2.1. Let S be a surface properly embedded in $D^2 \times B^2$. Then, S is called a *knitted surface* of degree n if there exists a partition $0 = t_0 < t_1 < \dots < t_m = 1$ for some m satisfying the following conditions, where $\beta_j = S_{[t_j]}$ ($j = 0, \dots, m$).

- (1) Each slice β_j is an n -knit ($j = 0, \dots, m$).

(2) The surface $S \cap (D^2 \times I_3 \times [t_{j-1}, t_j])$ is the knitted surface presented by $\beta_{j-1} \rightarrow \beta_j$ for $j \in \{0, \dots, m\}$ such that the knit β_j is obtained from β_{j-1} by a finite number of mutually disjoint transformations associated with the relations of the n -knit monoid D_n as follows ($j = 1, \dots, m$), where $\beta \leftrightarrow \beta'$ denotes the sequences $\beta \rightarrow \beta'$ and $\beta' \rightarrow \beta$:

- $e \leftrightarrow \sigma_i^\epsilon$,
- $e \leftrightarrow \tau_i$,
- $\tau_i \tau_i \leftrightarrow \tau_i$,
- $\tau_i \leftrightarrow \sigma_i^\epsilon \tau_i$, $\tau_i \leftrightarrow \tau_i \sigma_i^\epsilon$,
- $\sigma_i^\epsilon \sigma_i^{-\epsilon} \leftrightarrow e$,
- $\sigma_i^\epsilon \sigma_j^\delta \leftrightarrow \sigma_j^\delta \sigma_i^\epsilon$ ($|i - j| > 1$),
- $\sigma_i^\epsilon \tau_j \leftrightarrow \tau_j \sigma_i^\epsilon$ ($|i - j| > 1$),
- $\tau_i \tau_j \leftrightarrow \tau_j \tau_i$ ($|i - j| > 1$),
- $\sigma_i^\epsilon \sigma_j^\epsilon \sigma_i^\epsilon \leftrightarrow \sigma_j^\epsilon \sigma_i^\epsilon \sigma_j^\epsilon$, $\sigma_i^\epsilon \sigma_j^\epsilon \sigma_i^{-\epsilon} \leftrightarrow \sigma_j^{-\epsilon} \sigma_i^\epsilon \sigma_j^\epsilon$ ($|i - j| = 1$),
- $\sigma_i^\epsilon \sigma_j^\epsilon \tau_i \leftrightarrow \tau_j \sigma_i^\epsilon \sigma_j^\epsilon$ ($|i - j| = 1$),

where $\epsilon \in \{+1, -1\}$, $i, j \in \{1, \dots, n-1\}$.

We denote by Q_n the starting point set of n -knits, that is a fixed set of n interior points of D^2 such that $\beta \cap (D^2 \times \partial I_3) = Q_n \times \partial I_3$ for any n -knit β . We call the knitted surface $Q_n \times B^2$ the *trivial* knitted surface.

A knitted surface of degree n is called a *braided surface* of degree n if the transformations $\beta_0 \rightarrow \dots \rightarrow \beta_m$ are restricted to the relations of the n -braid group B_n ; that is, the relations do not contain τ_i ($i = 1, \dots, n-1$).

A knitted surface S of degree n is called a *2-dimensional knit* if $\partial S = Q_n \times \partial B^2$. A 2-dimensional knit is called a *2-dimensional braid* if it is a braided surface.

3. THE CLOSURE OF 2-DIMENSIONAL KNITS

In this section, we show Theorem 1.1.

3.1. Closure of knits and 2-dimensional knits. Let $m = 1, 2$. Let B_i^m be copies of m -disks B^m . Let S be an n -knit in $D^2 \times B_1^2$ or a 2-dimensional knit in $D^2 \times B_2^2$, and let $S_0^m = Q_n \times B_2^m \subset D^2 \times B_2^m$ be the trivial knits/2-dimensional knits. Then, $\text{cl}(S) := S \cup S_0$ is a closed 1-manifold or a closed surface in $D^2 \times S^m$, where $S^m = B_1^m \cup_\partial B_2^m$.

Let F_0^m be a circle (respectively a 2-sphere) standardly embedded in \mathbb{R}^{m+2} when $m = 1$ (respectively $m = 2$), and let N_m be a regular neighborhood of F_0^m . We identify N_m with $D^2 \times S^m$ ($m = 1, 2$). The *closure* of S is a link or a surface-link in \mathbb{R}^{m+2} which is given by $\text{cl}(S) \subset N_m \subset \mathbb{R}^{m+2}$.

As an analogue of Alexander's theorem for oriented links, it is known that:

Theorem 3.1 ([8]). *Every orientable surface-link is ambiently isotopic to the closure of some 2-dimensional braid.*

3.2. The plat closure of braids. For a set $w = \{w^1, \dots, w^l\}$, we denote by $\cup w$ the union $w^1 \cup \dots \cup w^l$. A *wicket* is a semi-circle properly embedded in $D^2 \times I$ which meets $D^2 \times \{0\}$ orthogonally. A *configuration of n wickets* is

a set of mutually disjoint n wickets. \mathcal{W}_n denotes the space consisting of configurations of n wickets. Let Q_{2n} be a set of $2n$ interior points $\{q_1, \dots, q_{2n}\}$ of D^2 such that q_1, \dots, q_{2n} lie in $\{1/2\} \times I \subset I \times I = D^2$ in this order. Let w_0 denotes the configuration of n wickets such that $\partial w_0 = Q_{2n}$ and q_i and q_{i+1} are boundaries of the same component of w_0 for each $i \in \{1, \dots, n\}$. We denote by w_0^* the mirror image of w_0 with respect to $D^2 \times \{1/2\}$; remark that $\cup w_0$ and $\cup w_0^*$ are disjoint. The union $(\cup w_0) \cup (\cup w_0^*)$ represents the $2n$ -knit $\tau := \tau_1 \tau_3 \dots \tau_{2n-1}$.

For a $2n$ -braid β , the *plat closure* of β , denoted by $\tilde{\beta}$, is a link in $D^2 \times [-1, 2] \subset \mathbb{R}^3$ obtained by taking the union of $w_0 \subset D^2 \times [-1, 0]$, $\beta \subset D^2 \times [0, 1]$, and $w_0^* \subset D^2 \times [1, 2]$. See Figure 4. The plat closure $\tilde{\beta}$ can be regarded as the closure of the knit $\tau\beta$, that is, $\tilde{\beta} = \text{cl}(\tau\beta)$.




FIGURE 4. The plat closure of a braid.

3.3. The plat closure of braided surfaces. We recall the definition of the plat closure of a braided surface [16]. To take the plat closure, a braided surface is required to be “adequate”.

For a loop $f : (I, \partial I) \rightarrow (\mathcal{W}_n, w_0)$, we define the $2n$ -braid β_f by $\beta_f = \bigcup_{t \in I} \partial f(t) \times \{t\} \subset D^2 \times \{0\} \times I$. A braid β is called *adequate* if there exists a loop $f : (I, \partial I) \rightarrow (\mathcal{W}_n, w_0)$ such that $\beta = \beta_f$. The set of adequate $2n$ -braids forms a group called the *Hilden subgroup* of the $2n$ -braid group B_{2n} , denoted by K_{2n} , and K_{2n} is generated by σ_{2i-1} , $\sigma_{2j}\sigma_{2j-1}\sigma_{2j+1}\sigma_{2j}$, and $\sigma_{2j}\sigma_{2j-1}\sigma_{2j+1}^{-1}\sigma_{2j}^{-1}$ for $i \in \{1, \dots, m\}$ and $j \in \{1, \dots, m-1\}$ [2] (see Figure 5 for the case $n = 2$).




FIGURE 5. Adequate braids which generate the Hilden subgroup K_4 : from the left, σ_1 , σ_3 , $\sigma_2\sigma_1\sigma_3\sigma_2$, and $\sigma_2\sigma_1\sigma_3^{-1}\sigma_2^{-1}$. This figure is [14, Figure 16].

Let β be an adequate $2n$ -braid in $D^2 \times I$. There exists a unique loop $f : (I, \partial I) \rightarrow (\mathcal{W}_n, w_0)$ satisfying $\beta_f = \beta$. Let $S^1 = [0, 1]/\sim$ with $0 \sim 1$. Let $N = I \times S^1$, an annulus. We define the surface A_β in $D^2 \times N$ by

$$A_\beta = \bigcup_{t \in [0, 1]} ((\cup f(t)) \times \{t\}) \subset (D^2 \times I) \times S^1 = D^2 \times N. \quad (3.1)$$

A braided surface S is called *adequate* if ∂S is the closure of an adequate braid. We remark that an adequate braided surface is of degree $2n$ for some n , and every 2-dimensional braid of degree $2n$ is adequate.

For an adequate braided surface S in $D^2 \times B^2$, we denote by β the adequate braid whose closure is ∂S . We take the surface A_β given by (3.1). Recall that $B^2 = [0, 1] \times [0, 1]$. We identify N with $[-1, 2] \times [-1, 2] \setminus B^2$, and we identify ∂B^2 with $\{0\} \times S^1 \subset I \times S^1 = N$, and we embed $D^2 \times N \subset \mathbb{R}^4$ by $D^2 \subset \mathbb{R}^2$ and $N \subset \mathbb{R}^2$. We remark that $\partial S = \partial A_\beta = S \cap A_\beta$. Then the *plat closure* of S is a surface-link in \mathbb{R}^4 defined by the union of S and A_β , denoted by \tilde{S} .

Theorem 3.2 ([16]). *Every surface-link is ambiently isotopic to the plat closure of some adequate braided surface.*

3.4. Proof of Theorem 1.1. Let F be a surface-link. By Theorem 3.2, we take an adequate braided surface S of degree $2n$ such that the plat closure \tilde{S} is ambiently isotopic to F .

(Case 1) First, we consider the case that S is a 2-dimensional braid, that is, \tilde{S} has the motion picture as in Figure 6. Put $\tau = \tau_1 \tau_3 \cdots \tau_{2n-1}$. We take a




FIGURE 6. Motion picture of the plat closure of S .

surface-link F_1 given by

$$(F_1)_{[t]} = \begin{cases} \mathcal{D} & (t = -1) \\ \text{cl}(\tau) & (-1 < t < 0) \\ \text{cl}(\tau S_{[t]}) & (0 \leq t \leq 1) \\ \text{cl}(\tau) & (1 < t < 2) \\ \mathcal{D} & (t = 2), \end{cases}$$

where \mathcal{D} is a union of n copies of disks whose boundaries are $\text{cl}(\tau)$; see Figure 7. The surface F_1 is ambiently isotopic to the plat closure \tilde{S} .

Let Δ^- be the surface properly embedded in $\mathbb{R}^3 \times (-\infty, 0]$ given by

$$(\Delta^-)_{[t]} = \begin{cases} \mathcal{D}' & (t = -2) \\ \text{cl}(e) & (-2 < t < -1) \\ \text{cl}(e) \cup (\cup \mathcal{B}) & (t = -1) \\ \text{cl}(\tau) & (-1 < t \leq 0), \end{cases}$$

where \mathcal{D}' is a union of $2n$ disks whose boundaries are $\text{cl}(e)$ and \mathcal{B} is a band set attaching to e such that $h(e; \mathcal{B}) = \tau$, and $\cup \mathcal{B}$ is the union of bands of \mathcal{B} ; see Figure 8. The surface Δ^- consists of n copies of disks. We denote by F_1^- the surface $F_1 \cap (\mathbb{R}^3 \times (-\infty, 0])$. Then both Δ^- and F_1^- consist of n copies of disks




FIGURE 7. Motion picture of F_1 .

with $\Delta^- \cap (\mathbb{R}^3 \times (-\infty, 0]) = \partial\Delta^- = \partial F_1^- = F_1^- \cap (\mathbb{R}^3 \times (-\infty, 0]) \subset \mathbb{R}^3 \times \{0\}$. These set of disks are called *trivial disk systems*, and it is known [11] that they are ambiently isotopic rel $\mathbb{R}^3 \times [0, \infty)$.




FIGURE 8. Motion picture of Δ^- .

Similarly, we take the surface Δ^+ properly embedded in $\mathbb{R}^3 \times [1, \infty)$ given by the mirror image of Δ^- with respect to $\mathbb{R}^3 \times \{1/2\}$, and we denote by F_1^+ the surface $F_1 \cap (\mathbb{R}^3 \times [1, \infty))$. Then Δ^+ and F_1^+ are ambiently isotopic rel $\mathbb{R}^3 \times (-\infty, 1]$. Let F_2 be a surface obtained from F_1 by replacing F_1^+ and F_1^- with Δ^+ and Δ^- , respectively; that is,

$$F_2 = \Delta^- \cup (F_1 \cap (\mathbb{R}^3 \times [0, 1])) \cup \Delta^+.$$

By construction, F_2 is ambiently isotopic to F_1 .

Let $D^2 \times I$ be the cylinder associated with the knits in slices. Put $S_1 = F_2 \cap (D^2 \times I \times [-3/2, 5/2])$. Then S_1 is a 2-dimensional knit given by

$$(S_1)_{[t]} = \begin{cases} e & (-3/2 \leq t \leq -1) \\ e \cup (\cup \mathcal{B}) & (t = -1) \\ \tau & (-1 < t < 0) \\ \tau \cdot S_{[t]} & (0 \leq t \leq 1) \\ \tau & (1 < t < 2) \\ \tau \cup (\cup \mathcal{B}) & (t = 2) \\ e & (2 < t \leq 5/2). \end{cases}$$

Further, we see that the closure of S_1 is F_2 . Thus, the closure of the 2-dimensional knit S_1 is ambiently isotopic to the given surface-link F .

(Case 2) Next, we consider the case when F is ambiently isotopic to the plat closure \tilde{S} of an adequate braided surface $S \subset D^2 \times B^2$ of degree $2n$. By considering a self-homeomorphism f of B^2 and the induced self-homeomorphism of $D^2 \times B^2$ given by $\text{id} \times f$, we assume that $\partial S \cap (D^2 \times I \times \{1\})$ has a presentation of an adequate braid whose closure is ∂S . We denote the adequate braid $\partial S \cap (D^2 \times I \times \{1\})$ by γ . By the same isotopy used to obtain F_1 from \tilde{S} in (Case 1), we have a surface-link F_1 . Let S' be a knitted surface in $D^2 \times B^2$ associated with the adequate braid γ given in [14, Section 6.4.2]. Then, the surface S' satisfies

$$S'_{[t]} = \begin{cases} \tau \cdot \gamma & (t = 0) \\ \tau & (t = 1) \end{cases}.$$

Thus, let S_1 be the knitted surface given by

$$(S_1)_{[t]} = \begin{cases} \tau \cdot S_{[2t]} & (0 \leq t \leq 1/2) \\ S'_{[2t-1]} & (1/2 \leq t \leq 1). \end{cases}$$

The Let F_2 be the surface given by

$$(F_2)_{[t]} = \begin{cases} \mathcal{D} & (t = -1) \\ \text{cl}(\tau) & (-1 < t < 0) \\ \text{cl}((S_1)_{[t]}) & (0 \leq t \leq 1) \\ \text{cl}(\tau) & (1 < t < 2) \\ \mathcal{D} & (t = 2), \end{cases}$$

where \mathcal{D} is a union of n copies of disks whose boundaries are $\text{cl}(\tau)$.

By construction of S' , $F_2 \cap (\mathbb{R}^3 \times [1/2, \infty))$ is a trivial disk system (see [14, Proposition 6.6]) and F_2 is ambiently isotopic to the surface-link F_1 .

Then, by the same process with Case 1, we obtain a new surface-link F_3 from F_2 by replacing $F_2^+ = F_2 \cap \mathbb{R}^3 \times [1, \infty)$ and $F_2^- = F_2 \cap \mathbb{R}^3 \times (-\infty, 0]$ with Δ^+ and Δ^- , respectively. The surface-link F_3 is ambiently isotopic to F_2 . Then, $S_2 := F_3 \cap (D^2 \times I \times [-3/2, 5/2])$ is a 2-dimensional knit and the closure of S_2 is the surface-link F_3 . Hence, the closure of the 2-dimensional knit S_2 is ambiently isotopic to the given surface-link F . \square

4. CHART DESCRIPTION

A knitted surface of degree n is presented by a finite graph on B^2 called a “chart” of degree n [14], which is a generalization of a chart for braided surfaces [6, 7, 9]. Here, we give the definition of charts of degree 2. See [14] for the definition of degree n .

Definition 4.1. Let Γ be a finite graph in a 2-disk B^2 . Then, Γ is a *knit chart* of degree 2, or simply a *2-chart*, if it satisfies the following conditions.

- (1) The intersection $\Gamma \cap \partial B^2$ consists of a finite number of endpoints of edges of Γ meeting ∂B^2 orthogonally. Though they are vertices of degree one, we call elements of $\Gamma \cap \partial B^2$ *boundary points*, and we call only a vertex in $\text{Int}(B^2)$ a *vertex* of Γ .
- (2) Each edge is equipped with the label 1 and moreover, each edge is either oriented or unoriented. We call an oriented/unoriented edge a σ -/ τ -edge, respectively.
- (3) Each vertex is of degree 1 or 3 as depicted in Figure 9.

A vertex of degree 1 connected with a σ -edge (respectively τ -edge) is depicted by a small black (respectively red) disk, called a *black σ -vertex* (respectively *black τ -vertex*). A vertex of degree 3 is called a *mixed trivalent vertex* (respectively a *trivalent τ -vertex*) if it is connected with a σ -edge and a pair of τ -edges (respectively three τ -edges).




FIGURE 9. Vertices of a 2-chart.

A knitted surface has a chart description. From a knitted surface of degree n , we obtain a chart of degree n , and from a given chart Γ , we can construct a knitted surface whose chart is Γ ([14]).

In the rest of this paper, we focus on knitted surfaces of degree 2 and their 2-charts. Thus we only review the knitted surface $S(\Gamma)$ of degree 2 constructed from a 2-chart Γ as follows. We decompose B^2 into a union of a finite number of 2-disks such that for each disk E , by modification, $\Gamma \cap E$ is either an empty graph (called an *empty chart*) or one of the 2-charts as in Figures 9 and 10, where the dotted square in Figure 10 denotes ∂E . We identify E with $I_3 \times I_4$, where the direction of I_3 is the vertical direction from up to down and the direction of I_4 is the horizontal direction from left to right in the figures. When $\Gamma \cap E$ is an empty chart, we define $S(\Gamma \cap E)$ by the trivial knitted surface $Q_2 \times E$. When $\Gamma \cap E$ is one of the 2-charts as in Figure 10, we define $S(\Gamma \cap E)$ by the knitted surface $\sigma_1 \times I_4$

for Figure 10(1), $\sigma_1^{-1} \times I_4$ for Figure 10(2), and $\tau_1 \times I_4$ for Figure 10(3). When $\Gamma \cap E$ is one of the 2-charts as in Figure 9, we define $S(\Gamma \cap E)$ by the knitted surface presented by $s \rightarrow e$ for the 2-chart containing one black σ/τ -vertex for appropriate $s \in \{\sigma_1, \sigma_1^{-1}, \tau_1\}$, and the knitted surface presented by $\tau_1 \rightarrow \sigma_1\tau_1$, $\tau_1 \rightarrow \sigma_1^{-1}\tau_1$, or $\tau_1\tau_1 \rightarrow \tau_1$ for the 2-chart containing a trivalent vertex. We construct the knitted surface presented by Γ as the union of the knitted surfaces presented by $\Gamma \cap E$.




FIGURE 10. The 2-chart $\Gamma \cap E$ consisting of one edge.

4.1. Chart moves. In this section, we give the definition of chart moves for 2-charts, that are local moves of charts. This includes C-moves of charts of braided surfaces [5, 6, 9].

Definition 4.2. Let Γ and Γ' be 2-charts in B^2 . Then, a *knit chart move*, or a *C-move*, is a local move in a 2-disk $E \subset B^2$ such that Γ (respectively Γ') is changed to Γ' (respectively Γ), satisfying the following.

- (1) The boundary ∂E does not contain any vertices of Γ and Γ' except for boundary points, and $\Gamma \cap \partial E$ (respectively $\Gamma' \cap \partial E$) consists of transverse intersection points of edges of Γ (respectively Γ') and ∂E .
- (2) The charts are identical in the complement of E : $\Gamma \cap (B^2 \setminus E) = \Gamma' \cap (B^2 \setminus E)$.
- (3) The charts in E , $\Gamma \cap E$ and $\Gamma' \cap E$, are as in Figure 11 and the orientation-reversed figures of Figure 11.

We say that Γ and Γ' are *C-move equivalent* if they are related by a finite sequence of C-moves and ambient isotopies of B^2 .

Let Γ and Γ' be 2-charts and let S and S' be the knitted surfaces presented by Γ and Γ' , respectively. Theorem 7.5 in [14] implies that S and S' are ambiently isotopic if Γ and Γ' are C-move equivalent. Thus, the plat closures of S and S' are ambiently isotopic if Γ and Γ' are C-move equivalent.

5. PLAT CLOSURE OF KNITTED SURFACES OF DEGREE 2

In this section, we consider knitted surfaces of degree 2. We give a normal form of a 2-chart (Definition 5.1 and Proposition 5.2). We show that any trivial surface-link is ambiently isotopic to the plat closure of knitted surfaces of degree 2 (Corollary 5.7). We show that the plat closure of any knitted surface of degree 2 is a trivial surface-link (Theorem 5.13), and we show Theorem 1.2 and Corollary 1.3.




FIGURE 11. C-moves for 2-charts, where we omit labels, which are all 1. C-moves include the orientation-reversed version.

5.1. Knitted surfaces of degree 2. A σ -/ τ -edge of a chart is called a *free* σ -/ τ -edge if its endpoints are black σ -/ τ -vertices. A σ -edge of a chart is called a *half* σ -edge if its endpoints consists of a black σ -vertex and either a boundary point or a mixed trivalent vertex. A τ -edge of a chart is called a *half* τ -edge if its endpoints consists of a black τ -vertex and either a boundary point or a trivalent τ -vertex.

For a chart Γ , the τ -chart of Γ , denoted by Γ_τ , is defined as the subgraph of Γ consisting of τ -edges and the connecting vertices. The τ -chart Γ_τ may contain 2-valent vertices which are obtained from mixed trivalent vertices. Then, we merge the two edges connected by each 2-valent vertex after removing the 2-valent vertex. Thus, Γ_τ consists of intervals, circles, and graphs consisting of 1-valent vertices and 3-valent vertices. We call a connected component of Γ_τ a τ -component. When Γ_τ is contained in B^2 , a *region* of Γ_τ is the closure of a connected component of $B^2 \setminus \Gamma_\tau$.

Definition 5.1. Let Γ be a 2-chart, let R_1, \dots, R_m be regions of the τ -chart Γ_τ , and put $\Gamma^j := \Gamma \cap R_j$, the subchart of Γ ($j = 1, \dots, m$). Then, we say that Γ is in a *normal form* if for each $j = 1, \dots, m$, there exists a τ -component G_j of Γ^j such that every σ -edge of Γ_j is either a free σ -edge or connected with G_j , and half σ -edges connected with G_j have coherent orientations. We call G_j the *initial* τ -component of Γ_j .

See Figure 12 for an example.

Proposition 5.2. *Any 2-chart is C-move equivalent to a 2-chart in a normal form given in Definition 5.1.*

Proof. Let Γ be a 2-chart. The labels of edges are all one. We denote by the same notation Γ the charts obtained from Γ by C-moves. From now on, we




FIGURE 12. An example of a 2-chart in a normal form, where we omit labels, which are all 1. The τ -edge in the form of the circle in the center is the initial τ -component associated with the region whose closure contains the larger part of ∂B^2 .

consider each subchart $\Gamma^j = \Gamma \cap R_j$ of Γ as given in Definition 5.1. We fix a τ -component G_j of Γ^j . By the definition of 2-charts, σ -edges are classified by combinations of their endpoints; a σ -edge is either a simple closed curve (called a circle component), a free σ -edge, a half σ -edge, a σ -edge connecting with $\partial B^2/\tau$ -components. Let \mathbf{e} be a σ -edge of Γ^j disjoint with G_j . Since R_j is connected, we take a simple path from a point of \mathbf{e} to G_j . Applying (A1), (A2) and (D1)-moves, each circle component of σ -edges is deformed into a σ -edge connecting τ -components. By applying (A2) and (D1)-moves, we deform half σ -edges and σ -edges connecting τ -components into σ -edges connected with G_j , see Figure 13; the detail is left to the reader as exercise. Further, by (A2)-moves, we deform half σ -edges connected with G_j to a set of free σ -edges and half σ -edges with coherent orientations. Thus we arrange Γ into a normal form. \square




FIGURE 13. Deforming σ -edges in Γ^j to σ -edges connecting with the initial τ -component G_j . We omit the labels of the 2-charts.

Remark 5.3. We can deform a 2-chart Γ to a more regulated form, also denoted by Γ , satisfying that (1) every black τ -vertex is an endpoint of a free τ -edge, (2) free σ -/ τ -edges are in a 2-disk which contains no other parts of Γ , (3) σ -edges connecting a τ -component and the initial τ -component have coherent orientations, and (4) there are no σ -edges both of whose endpoints are connected with the initial τ -component.

5.2. The plat closure of knitted surfaces. Let S be a knitted surface of degree 2. We define the *plat closure* of S in such a way as follows. If S is a braided surface, then S is adequate; thus we define the plat closure \tilde{S} of S by that of a braided surface, given by $\tilde{S} = S \cup A_\beta \subset \mathbb{R}^4$, where A_β is the surface of (3.1) in Section 3.3. Further, if ∂S is the closure of a 2-braid, then we define the plat closure of S by the same way, given by $\tilde{S} = S \cup A_\beta$. We consider the case that ∂S is the closure of a 2-knit having hook pairs. Then, ∂S consists of m simple closed curves, where m is the number of hook pairs in ∂S . We define the plat closure of S by the closed surface obtained from S by attaching m copies of 2-disks trivially along ∂S .

For knitted surfaces of degree n , we do not give the exact definition in terms of wickets in this paper, but we briefly give the idea of the plat closure, as follows. We call an $2n$ -knit *adequate* if it is an element of the submonoid of the $2n$ -knit monoid D_{2n} generated by adequate $2n$ -braids and τ_{2k-1} ($k = 1, \dots, n$). We call a knitted surface S *adequate* if ∂S is the closure of an adequate knit. For an adequate knitted surface S , we construct the *plat closure* of S as follows. Let β be the adequate knit whose closure is ∂S . We denote by β' the braid obtained from the adequate knit β by changing each letter in the form of τ_i for some i to σ_i . We denote by s_1, \dots, s_m the letters of β' coming from the letters of β in the form of τ_i ($i = 1, \dots, n-1$). The $2n$ -braid β' is adequate. Let $A_{\beta'}$ be the surface associated with β' given in (3.1) in Section 3.3. The part of $A_{\beta'}$ corresponding to s_j , that is $A_{\beta'} \cap (D^2 \times I) \times [t_j, t'_j]$ for some $0 \leq t_j < t'_j \leq 1$ with $\partial S \cap (D^2 \times [t_j, t'_j]) = s_j$, is a band with one twist. We delete the band from $A_{\beta'}$ and take the closure for each s_j ($j = 1, \dots, m$). Then, slightly modifying $A_{\beta'}$, we obtain a surface A' in $D^2 \times N$ with $\partial A' = \partial S = S \cap A'$. We define the *plat closure* of S by the surface-link $S \cup A'$ in \mathbb{R}^4 .

Remark 5.4. The plat closure of a knitted surface S of degree 2 is a special case of the closure of a 2-dimensional knit of degree 2 by the following meaning. Let Γ be a 2-chart of S , $y \in \partial B^2$ a point disjoint with Γ , and let N be a small neighborhood of y disjoint from Γ . Let \tilde{B}^2 be a 2-disk such that B^2 is contained in the interior of \tilde{B}^2 . Let Γ' be the 2-chart in \tilde{B}^2 obtained from Γ by adding to Γ a union of τ -edges in the form $\partial B^2 \setminus (N \cup \Gamma)$. Since Γ' does not have boundary points, it represents a 2-dimensional knit of degree 2, denoted by S' . By applying the argument in the proof of Theorem 1.1, the plat closure of S and the closure of S' is ambiently isotopic as surface-links. In particular, when Γ has no boundary points, then Γ' is obtained from Γ by an addition of a single free τ -edge.

We can calculate the Euler characteristics of knitted surfaces of degree 2 by using the information derived from charts. The proof is left to the reader; see [5, Exercise 9.15]. For simplicity, we consider a knitted surface $S \subset D^2 \times S^2$, presented by a 2-chart in a 2-sphere S^2 .

Proposition 5.5. *Let $F \subset D^2 \times S^2$ be a knitted surface of degree 2, and let $\Gamma \subset S^2$ be its 2-chart. Let R_1, \dots, R_m be regions of the τ -chart Γ_τ , and put $\Gamma^j := \Gamma \cap R_j$, the subchart of Γ ($j = 1, \dots, m$). Then F is a split union of F_1, \dots, F_m , where F_j is the plat closure of a knitted surface $S(\Gamma^j)$,*

$j = 1, \dots, m$, and the Euler characteristic $\chi(F_j)$ is calculated by

$$\chi(F_j) = 2\chi(R'_j) - b(\Gamma^j),$$

where $b(\Gamma^j)$ is the number of black σ -vertices of Γ^j , and $\chi(R'_j)$ is the Euler characteristic of the closure of $R_j \cap (B^2 \setminus N(\Gamma_\tau))$, where $N(\Gamma_\tau)$ is a regular neighborhood of Γ_τ ($j = 1, \dots, m$).

5.3. Plat closure of knitted surfaces of degree 2 presenting trivial surface-links. A standard 2-sphere (respectively a standard torus) in \mathbb{R}^4 is the boundary of a 3-ball (respectively an unknotted solid torus) in $\mathbb{R}^3 \times \{0\}$, and a standard positive projective plane (respectively a standard negative projective plane) is the surface given by Figure 14(1) (respectively (2)). These surfaces are called *standard surfaces*. A surface-knot in \mathbb{R}^4 is called *trivial* if it is a connected sum of a finite number of standard surfaces in \mathbb{R}^4 , and a surface-link is *trivial* if it is a split union of a finite number of trivial surface-knots.




FIGURE 14. (1) A standard positive projective plane P_+ and (2) a standard negative projective plane P_- .

When a half σ -edge has an orientation toward (respectively from) the black σ -vertex, we call it *positive* (respectively *negative*). We denote by T a standard torus, and by P_+ (respectively P_-) a standard positive (respectively negative) projective plane.

Theorem 5.6 ([14]). *Let F be a trivial surface-knot, which is a connected sum of g copies of T and m copies of P_+ and n copies of P_- . Then F is the plat closure of the knitted surface presented by a 2-chart Γ such that Γ consists of $\min\{m, n\}$ copies of free σ -edges, g copies of free τ -edges, and $|m - n|$ copies of positive (respectively negative) half σ -edges if $m \geq n$ (respectively $m < n$).*

Corollary 5.7. *Any trivial surface-link is ambiently isotopic to the plat closure of some knitted surface of degree 2.*

Proof. Let F be a trivial surface-link, and let F_1, \dots, F_m denote connected components of F . Let l_j be the segment $I \times \{j/m\}$ in $B^2 = I \times I$ ($j = 1, \dots, m - 1$). Then, l_1, \dots, l_{m-1} divide B^2 into m 2-disks $E_j = I \times [(j - 1)/m, j/m]$, $j = 1, \dots, m$. Applying Theorem 5.6, we obtain a 2-chart Γ_j in E_j such that the plat closure of a knitted surface $S(\Gamma_j)$ is ambiently isotopic to F_j . Let Γ be the union of Γ_j ($j = 1, \dots, m$) and τ -edges in the form of $\cup_{j=1}^{m-1} l_j$. Then, the plat closure of $S(\Gamma)$ is a split union of the plat closures of $S(\Gamma_j)$, $j = 1, \dots, m$, which is ambiently isotopic to F . \square

5.4. Knit plat index and the knit index. The *plat index* [16] of a surface-link F , denoted by $\text{Plat}(F)$, is the half of the minimum of the degrees of braided surfaces whose plat closures are ambiently isotopic to F . We define the *knit plat index*, denoted by $\text{knit.Plat}(F)$, as the half of the minimum of the degrees of knitted surfaces whose plat closure is ambiently isotopic to F . By definition, for a surface-link F ,

$$\text{knit.Plat}(F) \leq \text{Plat}(F).$$

It is known [16] that the plat index of surface-link is one if and only if it is a trivial 2-sphere or nonorientable trivial surface-knot, and the plat index is two if the surface-link is a trivial orientable surface-knot with positive genus. Corollary 5.7 implies the following:

Corollary 5.8. *Let F be any trivial surface-link. Then, the knit plat index of F is one:*

$$\text{knit.Plat}(F) = 1.$$

The *braid index* [5] of an oriented surface-link F , denoted by $\text{Braid}(F)$, is the minimum of the degrees of 2-dimensional braids whose closures are ambiently isotopic to F . We define the *knit index* of a surface-link F , denoted by $\text{Knit}(F)$, by the minimum of the degrees of 2-dimensional knits whose closures are ambiently isotopic to F . By definition, for an oriented surface-link F ,

$$\text{Knit}(F) \leq \text{Braid}(F).$$

By the argument in the proof of Theorem 1.1, we have the following.

Corollary 5.9. *Let F be any surface-link. Then, the knit index of F is bounded from above by twice the plat index of F :*

$$\text{Knit}(F) \leq 2\text{Plat}(F).$$

By Corollary 1.3, we have the following.

Corollary 5.10. *Let F be a trivial surface-link. Then, if F is not a trivial 2-sphere, then,*

$$\text{Knit}(F) = 2.$$

5.5. Remark on 2-charts presenting trivial surface-knots. The trivial surface-knot $P_+\#P_-$ is presented by a 2-chart other than that given in Theorem 5.6.

Proposition 5.11. *The trivial surface-knot $P_+\#P_-$ is ambiently isotopic to the plat closure of the knitted surfaces of degree 2 presented by the 2-charts as in Figure 15.*

Proof. The trivial surface-knot $P_+\#P_-$ has the motion picture as in Figure 16(1). The surface is ambiently isotopic to the surface as in Figure 16(2). This surface is ambiently isotopic to the surface as in Figure 16(3), which is the surface given by the gray regions of the 2-charts in Figure 17; so it is the plat closure of the knitted surface presented by the 2-chart in the left figure of Figure 15. By Theorem 5.6, $P_+\#P_-$ is ambiently isotopic to the plat closure of the knitted surface presented by the 2-chart in the right figure of Figure 15. Thus the 2-charts as in Figure 15 both present $P_+\#P_-$. \square



FIGURE 15. A pair of 2-charts whose plat closures are ambiently isotopic as surface-knots to $P_+ \# P_-$, where we omit labels. The left chart is the chart given in Theorem 5.6.




FIGURE 16. Motion pictures of $P_+ \# P_-$.




FIGURE 17. The gray regions of the 2-charts present the same surface of Figure 16(3).

Let F be a trivial surface-knot, which is a connected sum of g copies of T and m copies of P_+ and n copies of P_- . If $m \neq n$, then F is a connected sum of $\min\{m, n\} + g$ copies of T and $|m - n|$ copies of P_+ (respectively P_-) if $m > n$ (respectively $m < n$). We can show this by using 2-charts, applying Theorem 5.6, Proposition 5.11 and C-moves. The proof is left to the reader as exercise. In the following, we give the result in terms of 2-charts.

Corollary 5.12. *Let Γ be a 2-chart consisting of μ copies of free σ -edges, ν copies of free τ -edges and λ copies of positive (respectively negative) half σ -edges. If $\lambda \neq 0$, then the plat closure of the knitted surface presented by Γ is ambiently isotopic to the plat closure of the knitted surface presented by the 2-chart Γ' consisting of $\mu + \nu$ copies of free τ -edges and λ copies of positive (respectively negative) half σ -edges.*

5.6. Proof of Theorem 1.2 and Corollary 1.3. Let F be a surface-link in \mathbb{R}^4 . A *1-handle attaching to F* is a 3-ball $b = D^2 \times I$ embedded in \mathbb{R}^4 such that $b \cap F = (D^2 \times \partial I) \cap F$. The arc $\{0\} \times I \subset D^2 \times I$ is called the *core* of the 1-handle. The *1-handle surgery* is the operation which changes F to a new surface-link F' given by $F' = \text{Cl}(F \setminus (D^2 \times \partial I) \cup (\partial D^2 \times I))$. A 1-handle b attaching to F is called *trivial* if we can move the 1-handle b by ambient isotopy so that b is contained in a 4-ball U in \mathbb{R}^4 such that the pair $(U, F \cap U)$ is homeomorphic to the standard $(4, 2)$ -ball pair (B^4, B^2) .

Theorem 5.13. *Let F be the plat closure of an arbitrary knitted surface of degree 2. Then, F is a trivial surface-link.*

Proof. Let Γ be an arbitrary 2-chart. It suffices to show the result for the subchart of each region divided by the τ -chart Γ_τ of Γ . So we assume that $\Gamma \subset B^2$ is in a normal form such that $B^2 \setminus \Gamma_\tau$ is connected. We call a σ -edge which is not a half σ -edge a *non-half σ -edge*. Let l_1, \dots, l_s be segments in B^2 such that each l_j coincides with each non-half σ -edge. Since we take the plat closure, we regard ∂B^2 as the initial τ -component.

Let Γ' be the 2-chart obtained from Γ by changing each non-half σ -edge into a τ -edge. Let F (respectively F') be the plat closure of the knitted surface presented by Γ (respectively Γ'). We see that F' is a split union of the plat closures of knitted surfaces of 2-charts of regions divided by the τ -chart of Γ' , and F is obtained from F' by 1-handle surgery along 1-handles b_1, \dots, b_s such that the projection to B^2 of the core of b_j transversely intersects with the segment l_j ($j = 1, \dots, s$). We remark that since Γ is in a normal form, each region is a 2-disk. We consider the case $s = 1$, when the number of non-half σ -edges is one. We put $b := b_1$, the 1-handle along which F is obtained from F' by 1-handle surgery. The plat closure F is a surface-knot, and F' is either a surface-knot or a surface-link with two components. By Theorem 5.6, using (T2)-moves (see Figure 11) if necessary, we see that F and F' are trivial surface-knots/links. If F' is a surface-link with two components, denoted by F'_1 and F'_2 , then, F is a connected sum of F'_1 and F'_2 . If F' is a surface-knot, then, since F' is trivial, the 1-handle b is trivial [4]. Hence, the result of 1-handle surgery, F , is a connected sum of F' and either a standard torus T or a connected sum $P_+ \# P_-$ (see Figures 16(3) and 17). For the general case, since F' is a trivial surface-link by Theorem 5.6, by induction on the number s of non-half σ -edges, we see that F is obtained from components of F' by taking connected sum with several copies of T and $P_+ \# P_-$. Hence we see that F is a trivial surface-knot. Thus we have the required result. \square

Proof of Theorem 1.2. By Corollary 5.7 and Theorem 5.13, we have the required result. \square

Proof of Corollary 1.3. By Remark 5.4, Corollary 5.7 implies that any trivial surface-link is ambiently isotopic to the closure of some 2-dimensional knit. Let Γ be a 2-chart in B^2 . Theorem 1.1 implies that if Γ contains at least one τ -edge, then the closure of the 2-dimensional knit presented by Γ is a trivial surface-link. If Γ does not contain τ -edges, then it Γ presents a 2-dimensional braid, the closure of which is an orientable trivial surface-link [5]. \square

ACKNOWLEDGEMENTS

The first author was partially supported by JST FOREST Program, Grant Number JPMJFR202U. The second author was partially supported by JSPS KAKENHI Grant Number 22J20494.

REFERENCES

- [1] J. S. Birman, H. Wenzl, *Braids, link polynomial and a new algebra*, Trans. Amer. Math. Soc. **313** (1989) 249–273.
- [2] T. Brendle, A. Hatcher, *Configuration spaces of rings and wickets*, Comment. Math. Helv., **88** (2008) 131–162.
- [3] J. S. Carter, S. Kamada, M. Saito, *Surfaces in 4-space*, Encyclopaedia of Mathematical Sciences 142, Low-Dimensional Topology III, Berlin, Springer-Verlag, 2004.
- [4] S. Kamada, *Cords and 1-handles attached to surface-knots*, Bol. Soc. Mat. Mex. **20** (2014) 595–609.
- [5] S. Kamada, *Braid and Knot Theory in Dimension Four*, Math. Surveys and Monographs 95, Amer. Math. Soc., 2002.
- [6] S. Kamada, *An observation of surface braids via chart description*, J. Knot Theory Ramifications **5** (1996), no. 4, 517–529.
- [7] S. Kamada, *A characterization of groups of closed orientable surfaces in 4-space*, Topology **33** (1994) 113–122.
- [8] S. Kamada, *Alexander’s and Markov’s theorems in dimension four*, Bull. Amer. Math. Soc. (N.S.), **31** (1994), no. 1, 64–67.
- [9] S. Kamada, *Surfaces in \mathbb{R}^4 of braid index three are ribbon*, J. Knot Theory Ramifications **1** (1992), no. 2, 137–160.
- [10] A. Kawachi, *A Survey of Knot Theory*, Birkhäuser Verlag, Basel, 1996.
- [11] A. Kawachi, T. Shibuya, S. Suzuki, *Descriptions on surfaces in four-space. I. Normal forms*, Math. Sem. Notes Kobe Univ., **10** (1982), no. 1, 75–125.
- [12] H. M. Hilden, *Generators for two groups related to the braid group*, Pacific J. Math., **59** (1975), no. 2, 475–486.
- [13] J. Murakami, *The Kauffman polynomial of links and representation theory*, Osaka J. Math. **24** (1987) 745–758.
- [14] I. Nakamura and J. Yasuda, *Surfaces in the 4-ball constructed using generators of knits and their graphical description*, arXiv: 2306.10479v2.
- [15] L. Rudolph, *Braided surfaces and Seifert ribbons for closed braids*, Comment. Math. Helv. **58** (1983) 1–37.
- [16] J. Yasuda, *A plat form presentation for surface-links*, arXiv:2105.08634, to appear in Osaka J. Math.

DEPARTMENT OF MATHEMATICS, INFORMATION SCIENCE AND ENGINEERING, SAGA UNIVERSITY, 1 HONJOMACHI, SAGA, 840-8502, JAPAN. TEL: +81-952-28-8113
Email address: `inasa@cc.saga-u.ac.jp`

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, 1-1, MACHIKANAYAMA, TOYONAKA, OSAKA 560-0043, JAPAN
Email address: `u444951d@ecs.osaka-u.ac.jp`