THE COZERO PART OF THE POINTFREE VERSION OF $C_c(X)$

ALI AKBAR ESTAJI^{1*} AND MARYAM TAHA

ABSTRACT. Let $C_c(L) := \{\alpha \in \mathcal{R}(L) \mid R_\alpha \text{ is a countable subset of } \mathbb{R}\}$, where $R_\alpha := \{r \in \mathbb{R} \mid \cos(\alpha - r) \neq \top\}$ for every $\alpha \in \mathcal{R}(L)$. By using idempotent elements, it is going to prove that $\operatorname{Coz}_c[L] := \{\operatorname{coz}(\alpha) \mid \alpha \in C_c(L)\}$ is a σ -frame for every completely regular frame L, and from this, we conclude that it is regular, paracompact, perfectly normal and an Alexandroff algebra frame such that each cover of it is shrinkable. Also, we show that L is a zero-dimensional frame if and only if L is a c-completely regular frame.

Keywords: σ -frame, Idempotent element, Cozero element, Perfectly normal frame, Alexandroff algebra frame.

MSC(2010): Primary: 06D22; Secondary: 54C05; 54C30; 17C27.

1. INTRODUCTION

Some authors, such as Ball and Hager [2], studied the ring $\mathcal{R}(L)$ of real-valued continuous functions on a frame, namely the pointfree version of the ring C(X), prior to 1996. Banaschewski [5] did a systematic study of the ring of real-valued continuous functions in pointfree topology in 1996. Also, to read more about $\mathcal{R}(L)$, you can see [3, 4, 8, 17, 19].

The largest subring of C(X), whose elements are countable images, is $C_c(X)$, which leads to research on the pointfree version of its, that is, $\mathcal{R}_c(L)$, by Estaji et al. [12, 18, 20, 10]. Estaji at al. [13] defined $R_\alpha := \{r \in \mathbb{R} \mid \operatorname{coz}(\alpha - r) \neq \top\}$ for every $\alpha \in \mathcal{R}(L)$. The frame of open sets of a space X is called O(X). Aliabadi and Mahmodi [1] studied $\operatorname{pim}(\alpha) := \bigcap \{\omega \in O(\mathbb{R}) \mid \alpha(\omega) = \top\}$ for every $\alpha \in \mathcal{R}(L)$, and proved that $\operatorname{pim}(\alpha) = R_\alpha$.

By using the definition of R_{α} , authors of [21] introduced another pointfree version of $C_c(X)$ via the range of functions and denoted it by $C_c(L)$, which is more compatible with the functions of countable range. Also, in [22], it was proved that every radical ideal in it is an absolutely convex ideal. Characterization of the clean elements of $\mathcal{R}(L)$ has been done and proved when $C_c(L)$ is clean (see [14]).

The cozero map plays a key role in a way not usually encountered in ordinary topology; in fact, its main essence is that it provides a pointfree way of dealing with important point-dependent classical arguments. The collection of all cozero elements of *L* is called the *cozero part* and is usually denoted by Coz[L]. General properties of cozero elements and Coz[L] can be found in [6, 8]. Cozero elements play a particular role in $\mathcal{R}(L)$, $\mathcal{R}_c(L)$, and $C_c(L)$. The cozero part of a frame *L* is seen to be a sublattice of *L*, and also, Coz[L] is a σ -frame.

Date: Received: , Accepted: .

^{*}Corresponding author.

The main purpose of this article is to show that $\text{Coz}_c[L]$ is a σ -frame for every completely regular frame *L* and that *L* is a zero-dimensional frame if and only if *L* is a *c*-completely regular frame.

2. Preliminaries

A complete lattice *L* is said to be a *frame* if $a \land \bigvee B = \bigvee_{b \in B}(a \land b)$, for any $a \in L$ and $B \subseteq L$. We denote the *top* element and the *bottom* element of a frame *L* by \top and \bot , respectively. For every element *a* of a frame *L*, set $\uparrow a := \{b \in L \mid a \leq b\}$ and $a \rightarrow b := \{x \in L \mid a \land x \leq b\}$ for every $a, b \in L$. The *pseudocomplement* of $a \in L$ is the element $a^* := a \rightarrow \bot = \bigvee \{x \in L \mid x \land a = \bot\}$. If $a \lor a^* = \top$, then *a* is said to be *complemented* and the complement of *a* denoted by *a'*.

Recall from [5] that the *frame of reals* is the frame $\mathcal{L}(\mathbb{R})$ generated by all ordered pairs (p, q) with $p, q \in \mathbb{Q}$, subject to the following relations:

(R1) $(p,q) \land (r,s) = (p \lor r, q \land s),$ (R2) $(p,q) \lor (r,s) = (p,s),$ whenever $p \le r < q \le s,$ (R3) $(p,q) = \bigvee \{(r,s) \mid p < r < s < q\},$ (R4) $\top = \bigvee \{(p,q) \mid p, q \in \mathbb{Q}\}.$

A *continuous real function* on a frame is a frame homomorphism $\mathcal{L}(\mathbb{R}) \to L$. The set of all continuous real functions on a frame *L* is denoted by $\mathcal{R}(L)$. We recall from [5] that the set $\mathcal{R}(L)$ with the operator $\diamond \in \{+, \cdot, \land, \lor\}$ defined by

$$(\alpha \diamond \beta)(p,q) = \bigvee \{ \alpha(r,s) \land \beta(u,v) \mid < r, s > \diamond < u, v > \subseteq < p,q > \} (\forall \alpha, \beta \in \mathcal{R}(L))$$

is an *f*-ring. Here, $\langle p, q \rangle := \{r \in \mathbb{R} \mid p < r < q\}$ for every $p, q \in \mathbb{Q}$, and

 $< r, s > \diamond < u, v >= \{ y \diamond z \mid y \in < r, s >, z \in < u, v > \}.$

To facilitate the work, for every $p, q \in \mathbb{Q}$, we put

$$(p,-) := \bigvee_{\substack{r,s>p,\\r,s\in\mathbb{Q}}} (r,s) \text{ and } (-,q) := \bigvee_{\substack{r,s< q,\\r,s\in\mathbb{Q}}} (r,s).$$

For $\alpha \in \mathcal{R}(L)$, $\alpha(-, 0) \lor \alpha(0, -)$ is called *cozero element* of *L* and denoted by $coz(\alpha)$. In [5, 19], a frame is completely regular if and only if it is generated by its cozero elements, that is, for every $x \in L$, there exists $\{\alpha_i\}_{i \in I} \subseteq \mathcal{R}(L)$ such that

$$x = \bigvee_{\operatorname{coz}(\alpha_i) \le x} \operatorname{coz}(\alpha_i) \, .$$

Apart from frames, we shall also consider σ -frames, defined as below. A σ -frame is a distributive lattice L in which any countable subset has a join such that

$$a \land \bigvee S = \bigvee_{s \in S} (a \land s),$$

for all $a \in L$ and any countable subset $S \subseteq L$. For example,

$$\operatorname{Coz}[L] := \{\operatorname{coz}(\alpha) \mid \alpha \in \mathcal{R}(L)\}$$

is a sub- σ -frame of *L* (see [8, Corollary 2]).

We recall from [13] that for each $\alpha \in \mathcal{R}(L)$, $R_{\alpha} := \{r \in \mathbb{R} \mid \operatorname{coz}(\alpha - r) \neq \top\}$ and also,

$$C_c(L) := \{ \alpha \in \mathcal{R}(L) \mid R_\alpha \text{ is a countable subset of } \mathbb{R} \}$$

as the pointfree topology version of the ring $C_c(X)$ (for more details see [21, 22, 14]) and we set

$$\operatorname{Coz}_{c}[L] := {\operatorname{coz}(\alpha) \mid \alpha \in C_{c}(L)}.$$

3. Algebraic operations of the idempotent elements of $\mathcal{R}(L)$

It is written B(A) for the set of all the idempotent elements of a given ring A. If $e \in B(\mathcal{R}(L))$, then $\operatorname{coz}(e)$ is complemented. For $e_1, e_2 \in B(\mathcal{R}(L))$ in this section, we check $(e_1 \diamond e_2)(-, x)$ and $(e_1 \diamond e_2)(x, -)$ for every $x \in \mathbb{Q}$, where $\diamond \in \{+, -, \cdot, \lor, \land\}$.

Remark 3.1. Let *e* be an idempotent element of $\mathcal{R}(L)$. Then the following statements are true:

(1) For every $x \in \mathbb{Q}$,

$$e(x,-) = \begin{cases} \top, & x < 0, \\ \cos(e), & 0 \le x < 1, \text{ and } e(-,x) = \begin{cases} \bot, & x \le 0, \\ (\cos(e))', & 0 < x \le 1, \\ \top, & x \ge 1, \end{cases}$$

(2) If $x, k \in \mathbb{Q}$ with k > 0, then

$$ke(x,-) = \begin{cases} \top, & x < 0, \\ \cos(e), & 0 \le x < k, \text{ and } ke(-,x) = \begin{cases} \bot, & x \le 0, \\ (\cos(e))', & 0 < x \le k, \end{cases} \\ \top, & x \ge k, \end{cases}$$

(3) If $x, k \in \mathbb{Q}$ with k < 0, then

$$ke(x,-) = \begin{cases} \bot, & x \ge 0, \\ (\cot(e))', & k \le x < 0, \text{ and } ke(-,x) = \\ \top, & x < k, \end{cases} \quad \forall x > 0, \\ \cot(e), & k < x \le 0, \\ \bot, & x \le k. \end{cases}$$

Lemma 3.2. If $e_1, e_2 \in B(\mathcal{R}(L))$, then for every $x \in \mathbb{Q}$, we have

$$(e_1 + e_2)(x, -) = \begin{cases} \top & x < 0, \\ \cos(e_1 + e_2), & 0 \le x < 1, \\ \cos(e_1) \land \cos(e_2), & 1 \le x < 2, \\ \bot, & x \ge 2, \end{cases}$$

and

$$(e_1 + e_2)(-, x) = \begin{cases} \bot, & x \le 0, \\ (\cos(e_1))' \land (\cos(e_2))', & 0 < x \le 1, \\ (\cos(e_1))' \lor (\cos(e_2))', & 1 < x \le 2, \\ \bot, & x > 2. \end{cases}$$

Proof. For every $x \in \mathbb{Q}$, if x < 0, then $(e_1 + e_2)(x, -) = \top$, since $e_1 + e_2 \ge \mathbf{0}$. It is evident that $\operatorname{coz}(e_1 + e_2) = (e_1 + e_2)(0, -)$. For every $x, r \in \mathbb{Q}$, we define $H_{(x,r)} = e_1(r, -) \land e_2(x - r, -)$. Let $x \in \mathbb{Q}$ with 0 < x < 1 be given. Then there exist $r, s \in \mathbb{Q}$ such that 0 < r < x < s < 1, which implies that $H_{(x,x-s)} = \operatorname{coz}(e_2)$, $H_{(x,s)} = \operatorname{coz}(e_1)$, and $H_{(x,r)} = \operatorname{coz}(e_1) \land \operatorname{coz}(e_2)$. Therefore,

$$(e_1 + e_2)(x, -) = \operatorname{coz}(e_2) \lor \operatorname{coz}(e_1) \lor (\operatorname{coz}(e_1) \land \operatorname{coz}(e_2))$$
$$= \operatorname{coz}(e_1) \lor \operatorname{coz}(e_2)$$
$$= \operatorname{coz}(e_1 + e_2)$$

for every $x \in \mathbb{Q}$ with $0 \le x < 1$. Let $x \in \mathbb{Q}$ with $1 \le x < 2$ be given. Then for every $r \in \mathbb{Q}$,

 $\begin{aligned} x - r < 0 &\Rightarrow 1 \le x < r \Rightarrow H_{(x,r)} = \bot, \\ 0 \le x - r < 1 \Rightarrow (0 \le r < 1 \text{ and } 0 \le x - r < 1) \Rightarrow H_{(x,r)} = \operatorname{coz}(e_1) \land \operatorname{coz}(e_2), \text{ and} \\ x - r \ge 1 \Rightarrow H_{(x,r)} = \bot. \end{aligned}$

Hence,

$$(e_1 + e_2)(x, -) = \operatorname{coz}(e_1) \wedge \operatorname{coz}(e_2)$$

for every $x \in \mathbb{Q}$ with $1 \le x < 2$. Let $x \in \mathbb{Q}$ with $x \ge 2$ be given. Then for every $r \in \mathbb{Q}$, if $0 \le x - r < 1$ or x - r < 0, then r > 1 or $2 \le x < r$, which implies that $H_{(x,r)} = \bot$. Therefore, for every $x \in \mathbb{Q}$ with $2 \le x$, we have $(e_1 + e_2)(x, -) = \bot$. \Box

Lemma 3.3. If $e_1, e_2 \in B(\mathcal{R}(L))$, then for every $x \in \mathbb{Q}$, it holds that

$$(e_1e_2)(x,-) = \begin{cases} \top, & x < 0, \\ \cos(e_1) \wedge \cos(e_2), & 0 \le x < 1, \\ \bot, & x \ge 1, \end{cases}$$

and

$$(e_1e_2)(-, x) = \begin{cases} \bot, & x \le 0, \\ (\operatorname{coz}(e_1))' \lor (\operatorname{coz}(e_2))', & 0 < x \le 1, \\ \top, & x > 1. \end{cases}$$

Proof. For every $x \in \mathbb{Q}$, if x < 0, then $(e_1e_2)(x, -) = \top$, since $e_1e_2 \ge \mathbf{0}$. For every $x, s \in \mathbb{Q}$, we define $H_{(x,s)} = e_1(s, -) \land e_2(\frac{x}{s}, -)$. Let $0 < s \in \mathbb{Q}$. Then for x = 0, we have

 $0 < s < 1 \Rightarrow H_{(x,s)} = \operatorname{coz}(e_1) \wedge \operatorname{coz}(e_2),$ $s \ge 1 \Rightarrow H_{(x,s)} = \bot.$

Therefore, $(e_1e_2)(0, -) = (\operatorname{coz}(e_1) \wedge \operatorname{coz}(e_2)) \lor \bot = \operatorname{coz}(e_1) \wedge \operatorname{coz}(e_2).$

Let 0 < x < 1. Then

$$x < s < 1 \Rightarrow H_{(x,s)} = \operatorname{coz}(e_1) \land \operatorname{coz}(e_2),$$

 $s < x \text{ and } s > 1 \Rightarrow H_{(x,s)} = \bot.$

Hence, $(e_1e_2)(x, -) = (\operatorname{coz}(e_1) \wedge \operatorname{coz}(e_2)) \lor \bot = \operatorname{coz}(e_1) \wedge \operatorname{coz}(e_2)$. Let $x \ge 1$. Then

$$x \ge s \Rightarrow H_{(x,s)} = \bot,$$

$$s > x \Rightarrow H_{(x,s)} = \bot.$$

Thus, $(e_1e_2)(x, -) = \bot$.

For every $x \in \mathbb{Q}$, if x < 0, then $(e_1e_2)(x, -) = \bot$. For every $x, s \in \mathbb{Q}$, we define $H_{(s,x)} = e_1(-, s) \land e_2(-, \frac{x}{s})$. Let $0 < s \in \mathbb{Q}$. For x = 0, we have

$$0 < s \le 1 \Rightarrow H_{(x,s)} = (\operatorname{coz}(e_1))' \land \bot = \bot,$$

$$s > 1 \Rightarrow H_{(x,s)} = \top \land \bot = \bot.$$

Therefore, $(e_1e_2)(-, x) = \bot$.

Let $0 < x \le 1$. Then

$$0 < s \le x \Rightarrow H_{(s,x)} = (\operatorname{coz}(e_1))' \land \top = (\operatorname{coz}(e_1))',$$

$$x < s \le 1 \Rightarrow H_{(s,x)} = (\operatorname{coz}(e_1))' \land (\operatorname{coz}(e_2))',$$

$$s > 1 \Rightarrow H_{(s,x)} = \top \land (\operatorname{coz}(e_2))' = (\operatorname{coz}(e_2))'.$$

Thus,

$$(e_1e_2)(-, x) = (\operatorname{coz}(e_1))' \lor ((\operatorname{coz}(e_1))' \land (\operatorname{coz}(e_2))') \lor (\operatorname{coz}(e_2))' = (\operatorname{coz}(e_1))' \lor (\operatorname{coz}(e_2))'.$$

Let x > 1. Then

$$s \le 1 \Rightarrow H_{(s,x)} = (\operatorname{coz}(e_1))' \land \top = (\operatorname{coz}(e_1))',$$

$$x < s \Rightarrow H_{(s,x)} = \top \land (\operatorname{coz}(e_2))' = (\operatorname{coz}(e_2))',$$

$$1 < s < x \Rightarrow H_{(s,x)} = \top.$$

Therefore, $(e_1e_2)(-, x) = (\operatorname{coz}(e_1))' \lor (\operatorname{coz}(e_2))' \lor \top = \top$.

Similar to the process of proving the above lemma, the next lemma was proved. Lemma 3.4. If $e_1, e_2 \in B(\mathcal{R}(L))$, then for every $x \in \mathbb{Q}$, we have

$$(e_1 \lor e_2)(x, -) = \begin{cases} \top, & x < 0, \\ \cos(e_1 + e_2), & 0 \le x < 1, \\ \bot, & x \ge 1, \end{cases}$$
$$(e_1 \lor e_2)(-, x) = \begin{cases} \bot, & x \le 0, \\ (\cos(e_1))' \land (\cos(e_2))', & 0 < x \le 1, \\ \top, & x > 1, \end{cases}$$
$$(e_1 \land e_2)(x, -) = \begin{cases} \top, & x < 0, \\ \cos(e_1 \land e_2), & 0 \le x < 1, \\ \bot, & x \ge 1, \end{cases}$$

and

$$(e_1 \wedge e_2)(-, x) = \begin{cases} \bot, & x \le 0, \\ (\operatorname{coz}(e_1))' \lor (\operatorname{coz}(e_2))', & 0 < x \le 1, \\ \top, & x > 1. \end{cases}$$

4. Dedekind cuts for ${\mathbb R}$ by idempotent elements and prime ideals

In article [1], $A_p(\alpha)$ and $B_p(\alpha)$ were introduced. We found that it is better to use $P_u(\alpha)$ and $P_l(\alpha)$ symbols instead of them, which would be easier for the reader to understand the concepts. They are given below. For every $\alpha \in \mathcal{R}(L)$ and every prime ideal *P* in *L*, let

$$P_u(\alpha) := \{x \in \mathbb{Q} \mid \alpha(x, -) \in P\}$$
 and $P_l(\alpha) := \{x \in \mathbb{Q} \mid \alpha(-, x) \in P\}.$

To express our arguments, we need the following lemma and result from [1].

Lemma 4.1. [1] For every $\alpha \in \mathcal{R}(L)$ and every prime ideal P in L, the following statements are true:

- (1) $P_u(\alpha) \cup P_l(\alpha) = \mathbb{Q}$.
- (2) Any element of $P_u(\alpha)$ is an upper bound of $P_l(\alpha)$ and any element of $P_l(\alpha)$ is a lower bound of $P_u(\alpha)$.
- (3) $\uparrow P_u(\alpha) = P_u(\alpha)$ and $\downarrow P_l(\alpha) = P_l(\alpha)$.

Corollary 4.2. [1] Let P be a prime ideal of L and let $\alpha \in \mathcal{R}(L)$ be given. Then the following statements are equivalent:

- (1) inf $P_u(\alpha) \in \mathbb{R}$.
- (2) $P_l(\alpha) \neq \emptyset \neq P_u(\alpha)$.
- (3) sup $P_l(\alpha) \in \mathbb{R}$.
- (4) inf $P_u(\alpha) = \sup P_l(\alpha) \in \mathbb{R}$.

Corollary 4.3. Let P be a prime ideal of L and let $\alpha \in \mathcal{R}(L)$ be given. Then inf $P_u(\alpha) \in \mathbb{Q}$ if and only if inf $P_u(\alpha) \in P_u(\alpha)$ if and only if $|P_u(\alpha) \cap P_l(\alpha)| = 1$.

Proof. By Corollary 4.2, it is evident.

Lemma 4.4. Let P be a prime ideal of L. If e be an idempotent element of $\mathcal{R}(L)$, then the following statements are true:

(1) inf $P_u(e) = \sup P_l(e) \in P_u(e) \cap P_l(e)$ and

$$\inf P_u(e) = \begin{cases} 0, & \operatorname{coz}(e) \in P, \\ 1, & \operatorname{coz}(e) \notin P. \end{cases}$$

(2) For every $k \in \mathbb{Q}$, $\inf P_u(ke) = k \inf P_u(e)$.

Proof. (1). It is evident that $coz(e) \in P$ or $(coz(e))' \in P$. If $coz(e) \in P$, then

$$P_u(e) = \{x \in \mathbb{Q} \mid x \ge 0\}$$
 and $P_l(e) = \{x \in \mathbb{Q} \mid x \le 0\}$.

Therefore,

$$\inf P_u(e) = \sup P_l(e) = 0 \in P_u(e) \cap P_l(e)$$

If $(\operatorname{coz}(e))' \in P$, then

$$P_u(e) = \{x \in \mathbb{Q} \mid x \ge 1\}$$
 and $P_l(e) = \{x \in \mathbb{Q} \mid x \le 1\}.$

Hence,

$$\inf P_u(e) = \sup P_l(e) = 1 \in P_u(e) \cap P_l(e).$$

(2). Let $k \in \mathbb{Q}$ with k > 0 be given. Then

$$x \in P_u(ke) \Leftrightarrow ke(x, -) \in P \Leftrightarrow e(\frac{x}{k}, -) \in P \Leftrightarrow x \in kP_u(e),$$

which implies that $\inf P_u(ke) = k \inf P_u(e)$.

Let $k \in \mathbb{Q}$ with k < 0 be given. Then

$$x \in P_u(ke) \Leftrightarrow ke(x, -) \in P \Leftrightarrow e(-, \frac{x}{k}) \in P \Leftrightarrow x \in kP_l(e),$$

which implies that

$$\inf P_u(ke) = \inf kP_l(e) = k \sup P_l(e) = k \inf P_u(e).$$

In [1, Theorem 2.23], the prime ideal *P* is considered as countably \lor -complete, which here, since our discussion is about idempotent elements, we are able to remove this condition and express it as follows.

Proposition 4.5. Let *P* be a prime ideal of *L*. If $e_1, e_2 \in B(\mathcal{R}(L))$, then $\inf P_u(e_1 \diamond e_2) = \sup P_l(e_1 \diamond e_2) \in P_u(e_1 \diamond e_2) \cap P_l(e_1 \diamond e_2)$ and $\inf P_u(e_1 \diamond e_2) = \inf P_u(e_1) \diamond \inf P_u(e_2)$ for every $\diamond \in \{+, \lor, \land, \cdot\}$.

Proof. We only prove it for $\diamond = +$; the rest will be proved similarly. It is evident that

 $P_u(e_1 + e_2) \cap (-\infty, 0) = \emptyset$ and $P_l(e_1 + e_2) \cap (0, +\infty) = \emptyset$.

Case (1): If $coz(e_1) \in P$ and $coz(e_2) \in P$, then $coz(e_1 + e_2) = coz(e_1) \lor coz(e_2) \in P$, which implies that $P_u(e_1+e_2) = \{x \in \mathbb{Q} \mid x \ge 0\}$ and $P_l(e_1+e_2) = \{x \in \mathbb{Q} \mid x \le 0\}$. Therefore,

$$\inf P_u(e_1 + e_2) = \sup P_l(e_1 + e_2) = 0 \in P_u(e_1 + e_2) \cap P_l(e_1 + e_2)$$

and $\inf P_u(e_1 + e_2) = \inf P_u(e_1) + \inf P_u(e_2)$.

Case (2): If $coz(e_1) \in P$ and $(coz(e_2))' \in P$, then $coz(e_1 + e_2) \lor (coz(e_2))' = \top$, which implies that $coz(e_1 + e_2) \notin P$. Hence, $P_u(e_1 + e_2) \cap [0, 1) \cap \mathbb{Q} = \emptyset$. Since $coz(e_1) \land coz(e_2) \le coz(e_1) \in P$, we conclude that $[1, 2) \cap \mathbb{Q} \subseteq P_u(e_1 + e_2)$. It is evident that $[2, +\infty) \subseteq P_u(e_1 + e_2)$ and implies that $P_u(e_1 + e_2) = [1, +\infty) \cap \mathbb{Q}$. A similar proof shows that $P_l(e_1 + e_2) = (-\infty, 1] \cap \mathbb{Q}$. Therefore,

$$\inf P_u(e_1 + e_2) = \sup P_l(e_1 + e_2) = 1 \in P_u(e_1 + e_2) \cap P_l(e_1 + e_2)$$

and $\inf P_u(e_1 + e_2) = \inf P_u(e_1) + \inf P_u(e_2)$.

Case (3): Let $(coz(e_1))' \in P$ and let $coz(e_2) \in P$. Similar to the proof of Case (2), we have

$$\inf P_u(e_1 + e_2) = \sup P_l(e_1 + e_2) = 1 \in P_u(e_1 + e_2) \cap P_l(e_1 + e_2)$$

and $\inf P_u(e_1 + e_2) = \inf P_u(e_1) + \inf P_u(e_2)$.

Case (4): Let $(coz(e_1))' \in P$ and let $(coz(e_2))' \in P$. We have

$$(\operatorname{coz}(e_1 + e_2))' = (\operatorname{coz}(e_1))' \land (\operatorname{coz}(e_2))' \in P \Rightarrow \operatorname{coz}(e_1 + e_2) \notin P$$
$$\Rightarrow P_u(e_1 + e_2) \cap [0, 1) \cap \mathbb{Q} = \emptyset.$$

If $coz(e_1) \wedge coz(e_2) \in P$, then $coz(e_1) \in P$ or $coz(e_2) \in P$, which implies that $\top = coz(e_1) \vee (coz(e_1))' \in P$ or $\top = coz(e_2) \vee (coz(e_2))' \in P$, a contradiction. Hence, $P_u(e_1 + e_2) \cap [1, 2) \cap \mathbb{Q} = \emptyset$, which follows that $P_u(e_1 + e_2) = [2, +\infty) \cap \mathbb{Q}$. A similar proof shows that $P_l(e_1 + e_2) = (-\infty, 2] \cap \mathbb{Q}$. Therefore,

$$\inf P_u(e_1 + e_2) = \sup P_l(e_1 + e_2) = 2 \in P_u(e_1 + e_2) \cap P_l(e_1 + e_2)$$

and $\inf P_u(e_1 + e_2) = \inf P_u(e_1) + \inf P_u(e_2)$.

Corollary 4.6. Let P be a prime ideal of L. Suppose that $e_1, e_2 \in B(\mathcal{R}(L))$. If $x = \inf P_u(e_1 \diamond e_2)$, then $x \in R_{e_1 \diamond e_2}$ for every $\diamond \in \{\cdot, \lor, \land\}$.

Proof. By Proposition 4.5, it is obvious.

We recall that for every $p \in \mathbb{R}$,

$$\mathbf{p}(-,r) = \begin{cases} \bot, & r \le p, \\ \top, & r > p, \end{cases} \text{ and } \mathbf{p}(r,-) = \begin{cases} \top, & r < p, \\ \bot, & r \ge p. \end{cases}$$

A function f(x) is said to be bounded on a given interval if there exist two numbers M and N such that: $M \leq f(x) \leq N$. Also, $\mathcal{R}^*(L)$ is the set of bounded elements of $\mathcal{R}(L)$.

Lemma 4.7. Let P be a prime ideal of L. If $\alpha \in \mathbb{R}^*(L)$, then $\inf P_u(\alpha) = \sup P_l(\alpha)$.

Proof. There exists $n \in \mathbb{N}$ such that $-\mathbf{n} \leq \alpha \leq \mathbf{n}$, which implies that

$$T = \mathbf{n}(-, x) \le \alpha(-, x) \text{ for every } x > n \text{ and}$$

$$T = -\mathbf{n}(y, -) \le \alpha(y, -) \text{ for every } y < -n.$$

Hence, $(n, +\infty) \cap \mathbb{Q} \subseteq P_u(\alpha)$ and $(-\infty, -n) \cap \mathbb{Q} \subseteq P_l(\alpha)$. It follows from Corollary 4.2 that $\inf P_u(\alpha) = \sup P_l(\alpha)$.

Proposition 4.8. Let P be a prime ideal of L and let $n \in \mathbb{N}$. If $e_1, \ldots, e_n \in B(\mathcal{R}(L))$, then

$$\inf P_u(\mathbf{r}_1 e_1 + \dots + \mathbf{r}_n e_n) = \sup P_l(\mathbf{r}_1 e_1 + \dots + \mathbf{r}_n e_n)$$

belongs to

$$P_u(\mathbf{r_1}e_1 + \cdots + \mathbf{r_n}e_n) \cap P_l(\mathbf{r_1}e_1 + \cdots + \mathbf{r_n}e_n)$$

and

$$\inf P_u(\mathbf{r}_1 e_1 + \dots + \mathbf{r}_n e_n) = r_1 \inf P_u(e_1) + \dots + r_n \inf P_u(e_n)$$

for every $r_1, \ldots, r_n \in \mathbb{Q}$.

Proof. We prove it by induction on $n \in \mathbb{N}$. For n = 1, the result is trivial. Let $r_1, \ldots, r_n \in \mathbb{Q}$ be given and let $n \ge 2$. We put $\alpha = \mathbf{r_1}e_1 + \cdots + \mathbf{r_{n-1}}e_{n-1}$. Let $x \in P_u(\alpha + \mathbf{r_n}e_n)$ be given. Then

$$\bigvee_{a \in \mathbb{Q}} \alpha(a, -) \wedge \mathbf{r}_{\mathbf{n}} e_n(x - a, -) = (\alpha + \mathbf{r}_{\mathbf{n}} e_n)(x, -) \in P,$$

which implies that $\alpha(a, -) \wedge \mathbf{r_n} e_n(x - a, -) \in P$ for every $a \in \mathbb{Q}$, and so $\alpha(a, -) \in P$ or $\mathbf{r_n} e_n(x - a, -) \in P$ for every $a \in \mathbb{Q}$. If $a < \inf P_u(\alpha)$, then $\alpha(a, -) \notin P$, which implies from Remark 3.1 that

$$\perp = \mathbf{r_n} e_n(x-a,-) \in P \text{ or } \operatorname{coz}(e_n)' = \mathbf{r_n} e_n(x-a,-) \in P.$$

From

$$v := \mathbf{r}_{\mathbf{n}} e_n(x - \inf P_u(\alpha), -) = \bigvee_{\substack{a \in \mathbb{Q} \\ a < \inf P_u(\alpha)}} r_n e_n(x - a, -),$$

we conclude that $v = \bot \in P$ or $v = coz(e_n)' \in P$, which implies that inf $P_u(\mathbf{r}_n e_n) \le x - \inf P_u(\alpha)$ and hence,

$$\inf P_u(\alpha) + \inf P_u(\mathbf{r_n}e_n) \le \inf P_u(\alpha + \mathbf{r_n}e_n).$$

Let $x \in P_l(\alpha + \mathbf{r_n}e_n)$ be given. We set

i

$$w := \mathbf{r_n} e_n(-, x - \sup P_l(\alpha)) = \bigvee_{\substack{s \in \mathbb{Q} \\ s > \sup P_l(\alpha)}} r_n e_n(-, x - s).$$

Therefore, similar to the above, it conclude that for every $s > \sup P_l(\alpha)$, which we conclude from Remark 3.1 that $w = \bot \in P$ or $w = \operatorname{coz}(e_n) \in P$. Thus we have $x - \sup P_l(\alpha) \le \sup P_l(\mathbf{r}_n e_n)$ and hence,

$$\sup P_l(\alpha + \mathbf{r_n}e_n) \le \sup P_l(\alpha) + \sup P_l(\mathbf{r_n}e_n),$$

and finally, by Lemma 4.7,

$$\inf P_u(\alpha) + \inf P_u(\mathbf{r_n}e_n) = \inf P_u(\alpha + \mathbf{r_n}e_n).$$

The rest follows with Lemma 4.4.

5.
$$\operatorname{Coz}(B(C_c(L)))$$
 as a base for $\operatorname{Coz}_c[L]$

In [22], for every $r \in \mathbb{R}$, it was defined that $u_r := \bigvee_{n \in \mathbb{N}} (-, a_n)$ and $l_r := \bigvee_{n \in \mathbb{N}} (b_n, -)$, where $a_n, b_n \subseteq \mathbb{Q}$ are sequences such that $a_n, b_n \to r$ and for every $n \in \mathbb{N}$,

$$a_n < a_{n+1} < r < b_{n+1} < b_n$$
.

It was also shown that $\alpha(u_r)$ and $\alpha(l_r)$ are complemented in *L* and that the complement of $\alpha(l_r)$ is $\alpha(u_r)$. Also, see Lemma 4.1 in [13].

The following results are counterpart of [13, Proposition 5.1].

Proposition 5.1. If $\alpha \in C_c(L)$, then $coz(\alpha)$ is a countable join of complemented elements of L.

Proof. For any $n \in \mathbb{N}$, by [22, Remark 7.2], there exist $r_n, s_n \in \mathbb{R} \setminus R_\alpha$ satisfying all of the following conditions:

(1) $\frac{-1}{n} < r_n < 0 < s_n < \frac{1}{n}$, (2) $u_{r_n}, u_{s_n}, l_{r_n}$ and l_{s_n} are complemented, (3) $\alpha(\frac{1}{n}, -) \leq \alpha(l_{s_n}) \leq \operatorname{coz}(\alpha)$ and $\alpha(-, \frac{-1}{n}) \leq \alpha(u_{r_n}) \leq \operatorname{coz}(\alpha)$, (4) $\alpha(u_{r_n} \lor l_{s_n})$ is complemented.

Therefore,

$$\operatorname{coz}(\alpha) = \bigvee_{n \in \mathbb{N}} \alpha(-, \frac{-1}{n}) \lor \alpha(\frac{1}{n}, -) \leqslant \alpha(u_{r_n}) \lor \alpha(l_{s_n}) \leqslant \operatorname{coz}(\alpha),$$

which shows that $coz(\alpha) = \bigvee_{n \in \mathbb{N}} \alpha(u_{r_n} \vee l_{s_n})$.

Remark 5.2. It is easy to see that if *a* is complemented in *L* and $\alpha \in \mathcal{R}(L)$, then $\beta : \mathcal{L}(\mathbb{R}) \longrightarrow L$ given by

$$\beta(p,q) = \begin{cases} \alpha(p,q) \lor a' & \text{if } 0 \in \tau(p,q), \\ \alpha(p,q) \land a & \text{if } 0 \notin \tau(p,q), \end{cases}$$

is a continuous real-valued function. Since β is a continuous real-valued function, and by [11, Proposition 4.1], $e_a^2 = e_a$ and $\cos(e_\alpha) = a$. It is easily seen that *e* is an idempotent element of $\mathcal{R}(L)$ is and only if $e = e_{coz(\alpha)}$.

The following results are counterpart of [13, Theorem 5.2].

Proposition 5.3. For every $\alpha \in C_c(L)$, there exists a sequence $\{\alpha_n\}_{n \in \mathbb{N}} \subseteq B(C_c(L))$ with $\operatorname{coz}(\alpha) = \bigvee_{n \in \mathbb{N}} \operatorname{coz}(\alpha_n)$.

Proof. By Proposition 5.1 and Remark 5.2, it is obvious.

Definition 5.4. A frame *L* is a *c*-completely regular frame if and only if $\text{Coz}_c[L]$ is a base for *L*.

The following results are counterpart of [15, Proposition 4.4].

Proposition 5.5. *Let L be a frame. Then L is a zero-dimensional frame if and only if L is a c-completely regular frame.*

Proof. Necessity. Let $a \in L$. Then there exists a sequence $\{a_{\lambda}\}_{\lambda \in \Lambda}$ of complemented elements of *L* such that $a = \bigvee_{\lambda \in \Lambda} a_{\lambda}$. Also, we know $\operatorname{coz}(e_{a_{\lambda}}) = a_{\lambda}$, which implies that $a = \bigvee_{\lambda \in \Lambda} \operatorname{coz}(e_{a_{\lambda}})$. Since for every $\lambda \in \Lambda$, $e_{a_{\lambda}}$ is an idempotent element and all the idempotent elements belong to $C_c(L)$, the proof is complete.

Sufficiency. By Definition 5.4, $\text{Coz}_c[L]$ is a base for *L*, and by Proposition 5.1, for every $\alpha \in C_c(L)$, $\text{coz}(\alpha)$ is a countable join of complemented elements of *L*. Then *L* is a zero-dimensional frame.

6.
$$\operatorname{Coz}_{c}[L]$$
 as a σ -frame

In this section, the main purpose of writing this article is going to be proved, that is, $\text{Coz}_c[L]$ is a σ -frame for every completely regular frame L. To achieve this purpose, we need some lemmas and propositions, which we will express and prove them at the beginning.

Lemma 6.1. Let α be an element of $\mathcal{R}(L)$ and let $x \in \mathbb{R}$. Then the following statements are true:

- (1) If $x \in R_{\alpha}$, then there exists a prime ideal P in L such that $x = \inf P_u(\alpha)$.
- (2) If there exists a countably \lor -complete prime ideal P in L such that $x = \inf P_u(\alpha)$, then $x \in R_{\alpha}$.
- (3) Let P be a prime ideal of L and let $n \in \mathbb{N}$ be given. Suppose that $e_1, \ldots, e_n \in B(\mathcal{R}(L))$. If $r_1, \ldots, r_n \in \mathbb{Q}$ and $x = \inf P_u(\mathbf{r}_1 e_1 + \cdots + \mathbf{r}_n e_n)$, then $x \in R_{\mathbf{r}_1 e_1 + \cdots + \mathbf{r}_n e_n}$.

Proof. (1). By the hypothesis we have $coz(\alpha - \mathbf{x}) \neq \top$. Then there exists a prime ideal *P* in *L* such that $coz(\alpha - \mathbf{x}) \in P$, which implies that

$$\bigvee_{\substack{r \in \mathbb{Q} \\ r < x}} \alpha(-, r) \lor \bigvee_{\substack{r \in \mathbb{Q} \\ r > x}} \alpha(r, -) \in P$$

Therefore,

$$\forall r \in \mathbb{Q} \ (r < x \Rightarrow \alpha(-, r) \in P \Rightarrow r \in P_l(\alpha)) \Rightarrow x \le \sup P_l(\alpha),$$

and

$$\forall r \in \mathbb{Q} \ (r > x \Rightarrow \alpha(-, r) \in P \Rightarrow r \in P_u(\alpha)) \Rightarrow x \ge \inf P_u(\alpha)$$

Since, by Corollary 4.2, $\sup P_l(\alpha) = \inf P_u(\alpha)$, we conclude that $x = \inf P_u(\alpha)$.

(2). Let P be a prime ideal of L such that $x = \inf P_u(\alpha)$. Then, by Lemma 4.1,

$$\forall r \in \mathbb{Q} \ (r < x \Rightarrow r \in P_l(\alpha) \Rightarrow \alpha(-, r) \in P)$$

and

$$\forall r \in \mathbb{Q} \ \left(r > x \Rightarrow r \in P_u(\alpha) \Rightarrow \alpha(r, -) \in P \right)$$

By the hypothesis we have

$$\operatorname{coz}(\alpha - \mathbf{x}) = \bigvee_{\substack{r \in \mathbb{Q} \\ r < x}} \alpha(-, r) \lor \bigvee_{\substack{r \in \mathbb{Q} \\ r > x}} \alpha(r, -) \in P,$$

and so $x \in R_{\alpha}$.

(3). By Lemma 6.1,

$$x \in P_u(\mathbf{r}_1 e_1 + \cdots + \mathbf{r}_n e_n) \cap P_l(\mathbf{r}_1 e_1 + \cdots + \mathbf{r}_n e_n).$$

Then

 $\operatorname{coz}(\mathbf{r}_{1}e_{1} + \dots + \mathbf{r}_{n}e_{n} - x) = (\mathbf{r}_{1}e_{1} + \dots + \mathbf{r}_{n}e_{n})(-, x) \lor (\mathbf{r}_{1}e_{1} + \dots + \mathbf{r}_{n}e_{n})(x, -) \in P,$ which implies that $\operatorname{coz}(\mathbf{r}_{1}e_{1} + \dots + \mathbf{r}_{n}e_{n} - \mathbf{x}) \neq \top$. Therefore, $x \in R_{\mathbf{r}_{1}e_{1} + \dots + \mathbf{r}_{n}e_{n}}$. \Box

For every $A, B \subseteq \mathbb{R}$, we put $A \diamond B := \{a \diamond b \mid a \in A \text{ and } b \in B\}$ for every $\diamond \in \{+, \cdot, \wedge, \vee\}$.

Proposition 6.2. Let *n* be an element of \mathbb{N} . Suppose that $e_1, \ldots, e_n \in B(\mathcal{R}(L))$. If $r_1, \ldots, r_n \in \mathbb{Q}$, then

$$R_{\mathbf{r}_1e_1+\cdots+\mathbf{r}_ne_n} \subseteq R_{\mathbf{r}_1e_1}+\cdots+R_{\mathbf{r}_ne_n}.$$

Proof. Suppose that $x \in R_{\mathbf{r}_1 e_1 + \dots + \mathbf{r}_n e_n}$. Thus, by Lemma 6.1(1), there exists a prime ideal *P* in *L* such that $x = \inf P_u(\mathbf{r}_1 e_1 + \dots + \mathbf{r}_n e_n)$. Hence, by Proposition 4.8, $x = x_1 + \dots + x_n$, where $x_i = \inf P_u(\mathbf{r}_i e_i)$ for every $1 \le i \le n$. Then, by Lemma 6.1(3), $x_i \in R_{\mathbf{r}_i e_i}$. Therefore,

$$R_{\mathbf{r}_1 e_1 + \dots + \mathbf{r}_n e_n} \subseteq R_{\mathbf{r}_1 e_1} + \dots + R_{\mathbf{r}_n e_n}.$$

Proposition 6.3. Suppose that $e_1, e_2 \in B(\mathcal{R}(L))$. Then, $R_{e_1 \diamond e_2} \subseteq R_{e_1} \diamond R_{e_2}$ for every $\diamond \in \{\cdot, \lor, \land\}$.

Proof. By Proposition 4.5, Corollary 4.6, and Lemma 6.1, this proposition will be easily proved.

Remark 6.4. If α and β are elements of $\mathcal{R}(L)$ with $\alpha \leq \beta$, then $P_u(\beta) \subseteq P_u(\alpha)$ and $P_l(\alpha) \subseteq P_l(\beta)$.

Let $(c_n)_{n=0}^{\infty} \in \text{Coz}[L]$ and $(\alpha_n)_{n=0}^{\infty} \in \text{Coz}[L]$ with $\mathbf{0} \leq \alpha_n \leq \mathbf{1}$ and $\text{coz}(\alpha_n) = c_n$ for every *n*, be given. Banaschewski [6] showed that $\alpha := \sum_{i=0}^{\infty} \frac{\alpha_i}{2^i} \in \mathcal{R}(L)$, $\text{coz}(\alpha) = \bigvee_{i=0}^{\infty} c_n$ and $\alpha \leq \sum_{i=0}^{n} \frac{\alpha_i}{2^i} + \frac{1}{2^n}$ for every *n*.

 $\operatorname{coz}(\alpha) = \bigvee_{i=0}^{\infty} c_n \text{ and } \alpha \leq \sum_{i=0}^{n} \frac{\alpha_i}{2^i} + \frac{1}{2^n} \text{ for every } n.$ Given any countable family $(e_n)_{n=0}^{\infty}$ in $B(\mathcal{R}(L))$, consider $\alpha = \sum_{n=0}^{\infty} \frac{e_n}{2^n}$. Then for every $n \in \mathbb{N}$, we have

$$\alpha \leq \sum_{i=0}^{n} \frac{e_i}{2^i} + \frac{1}{2^n} \Rightarrow P_u(\sum_{i=0}^{n} \frac{e_i}{2^i} + \frac{1}{2^n}) \subseteq P_u(\alpha) \Rightarrow \inf P_u(\alpha) \leq \sum_{i=0}^{n} \inf P_u(\frac{e_i}{2^i}) + \frac{1}{2^n},$$

which implies that $\inf P_u(\alpha) \leq \sum_{i=0}^{\infty} \inf P_u(\frac{e_i}{2^i})$. On the other hand, for every $n \in \mathbb{N}$, it follows that

$$\sum_{i=0}^{n} \frac{e_i}{2^i} \le \alpha \Rightarrow P_u(\alpha) \subseteq P_u(\sum_{i=0}^{n} \frac{e_i}{2^i}) \Rightarrow \sum_{i=0}^{n} \inf P_u(\frac{e_i}{2^i}) \le \inf P_u(\alpha),$$

which implies that $\sum_{i=0}^{\infty} \inf P_u(\frac{e_i}{2^i}) \leq \inf P_u(\alpha)$. Hence, $\inf P_u(\alpha) = \sum_{i=0}^{\infty} \inf P_u(\frac{e_i}{2^i})$. **Proposition 6.5.** Let $(e_n)_{n=0}^{\infty}$ be a countable family of $B(\mathcal{R}(L))$. If $\alpha = \sum_{n=0}^{\infty} \frac{e_n}{2^n}$,

then $R_{\alpha} \subseteq \sum_{n=0}^{\infty} R_{\frac{e_n}{2^n}}$.

Proof. Suppose that $x \in R_{\alpha}$. Thus, by Lemma 6.1(1), there exists a prime ideal *P* in *L* such that $x = \inf P_u(\alpha)$. Hence, it follows from Remark 6.4 that $x = \sum_{n=0}^{\infty} x_i$, where

 $x_i = \inf P_u(\frac{e_i}{2^i})$ for every $i \in \mathbb{N} \cup \{0\}$. Then, by Lemma 6.1(3), we have $x_i \in R_{\frac{e_i}{2^i}}$.

Theorem 6.6. For every completely regular frame L, $Coz_c[L]$ is a σ -frame.

Proof. By Proposition 5.3, it suffices to show that if $(e_n)_{n=0}^{\infty}$ is a countable family of $B(\mathcal{R}(L))$, then $\bigvee_{n=0}^{\infty} \operatorname{coz}(e_n) \in \operatorname{Coz}_c[L]$. We put $\alpha = \sum_{n=0}^{\infty} \frac{e_n}{2^n}$. It is well known that $\operatorname{coz}(\alpha) = \bigvee_{n=0}^{\infty} \operatorname{coz}(e_n)$. Since $\sum_{n=0}^{\infty} R_{\frac{e_n}{2^n}}$ is a countable set, we conclude from Proposition 6.5 that R_{α} is too, which implies that $\alpha \in C_c(L)$. Therefore, $\bigvee_{n=0}^{\infty} \operatorname{coz}(e_n) \in \operatorname{Coz}_c[L]$.

An element *a* of frame *L* is said to *rather below* an element *b* or *a* is *well inside b*, written as a < b, if there is an element *x* such that $a \land x = \bot$ and $x \lor b = \top$. In [9], a σ -frame *L* is *regular* if for every $a \in L$, there exist $(a_n)_{n=1}^{\infty}$ such that $a_n < a$ and $a = \bigvee_{n=1}^{\infty} a_n$ for every *n*.

Corollary 6.7. For every completely regular frame L, $Coz_c[L]$ is regular.

A bounded lattice *L* is *normal* if given $a \lor b = \top$ that $a, b \in L$, there exist $u, v \in L$ such that $a \lor u = \top = v \lor b$ and $u \land v = \bot$. Also, a normal σ -frame *L* is *perfectly normal* if for each $a \in L$ there exist $(a_n)_{n=1}^{\infty}$ such that $a \land b = a \land c$ if and only $a_n \lor b = a_n \lor c$ for every $a, b \in L$ (see [9]).

By [16, Proposition 1-1], it is obvious that $\text{Coz}_c[L]$ is normal and perfectly normal for every completely regular frame *L*.

An Alexandroff algebra is a normal σ -lattice L that for every $a \in L$, there exist $(a_n, b_n)_{n=1}^{\infty}$ such that $b_n \lor a = \top$, $b_n \land a_n = \bot$ and $\bigvee_{n=1}^{\infty} a_n = a$.

Corollary 6.8. For every completely regular frame L, $Coz_c[L]$ is an Alexandroff algebra frame.

Proof. Let $a \in L$ be given. By Proposition 5.3, there exists $(e_n)_{n=1}^{\infty} \subseteq B(\mathcal{R}(L))$ such that $a = \bigvee_{\lambda \in \Lambda} \operatorname{coz}(e_{a_{\lambda}})$. We put $(a_n, b_n)_{n=1}^{\infty} = (\operatorname{coz}(e_n), \operatorname{coz}(e_n)')_{n=1}^{\infty}$, then we have

- (1) $\bigvee_{n=1}^{\infty} a_n = a$,
- (2) $b_n \wedge a_n = \bot$ for every $n \in \mathbb{N}$, and
- (3) $b_m \lor a = \operatorname{coz}(e_m)' \lor \bigvee_{n=1}^{\infty} \operatorname{coz}(e_n) = \top$ for every $m \in \mathbb{N}$.

Therefore, the proof is complete.

A cover of σ -frame *L* is a family $(a_n)_{n=1}^{\infty} \subseteq L$ such that $\bigvee_{n=1}^{\infty} a_n = \top$. A cover $(b_n)_{n=1}^{\infty}$ of σ -frame *L* is *locally finite* if there is a cover $(c_n)_{n=1}^{\infty}$ of *L* and, for each *m*, a finite subset p(m) of \mathbb{N} such that $c_m \wedge b_n = \bot$ if $n \notin p(m)$. A σ -frame is *paracompact* if for each countable cover $\mathcal{A} = (a_n)_{n=1}^{\infty}$ of *L* there is a locally finite cover $\mathcal{B} = (b_m)_{m=1}^{\infty}$ of *L* such that for each *m* there is some *n* with $b_m \leq a_n$. A cover $(a_n)_{n=1}^{\infty}$ of σ -frame *L* is *shrinkable* if there is a cover $(d_n)_{n=1}^{\infty}$ of *L* with $d_n \leq a_n$ for every *n*.

Proposition 6.9. For every completely regular frame L, the following statements are true:

- (1) $\operatorname{Coz}_{c}[L]$ is paracompact.
- (2) Each cover of $Coz_c[L]$ is shrinkable.

Proof. By [7, Proposition 4] and Corollary 6.7, it is obvious.

References

- [1] A. R. Aliabad and M. Mahmoudi, *Pre-image of function in* $C_c(L)$, Categ. Gen. Algebr. Struct. Appl. **15** (2021), no. 1, 35–58. https://doi.org/10.1007/s10472--007--9064--3.1, 6, 7
- [2] R.N. Ball and A.W. Hager, On the localic Yosida representation of an Archimedean lattice ordered group with weak order unit, J. Pure Appl. Algebra **70** (1991), 17–43. https://doi.org/10.1016/0022--4049(91)90004--L.1
- [3] R.N. Ball and J. Walters-Wayland, C- and C^*- quotients in pointfree topology, Dissertationes Math. (Rozprawy Mat.), Warszawa **412** (2002), 1–62. https://doi.org/10.4064/dm412--0--1.1
- [4] B. Banaschewski, *Pointfree topology and the spectra of f-rings*, Ordered algebraic structures(Curaçoa, 1995), Kluwer Acad. Publ., Dordrecht (1997), 123–148. https://doi.org/10.1007/978--94--011--5640--0_5.1
- [5] _____, *The real numbers in pointfree topology*, Textos de Mathematica (Series B), Univ. Coimbra, Dept. Math. Coimbra **12** (1997), 1–96. 1, 2

^[6] _____, *A new aspect of the cozero lattice in pointfree topology*, Topology Appl. **156** (2009), 2028–2038. https://doi.org/10.1016/j.topol.2009.03.041. 1, 12

^[7] B. Banaschewski and C.R.A. Gilmour, Stone-Čech compactification and dimension theory for regular σ -frames, J. London Math. Soc. **39** (1989), 1–8. https://doi.org/10.1112/jlms/s2-39.1.1.13

- [8] _____, *Pseudocompactness and the cozero part of a frame*, Comment. Math. Univ. Carolinae **37** (1996), no. 3, 577–587. http://eudml.org/doc/247922. 1, 2
- [9] M. G. Charalambous, *Dimension theory for* σ -*frames*, J. London Math. Soc. **2** (1974), no. 8, 149–160. https://doi.org/10.1112/JLMS. 12
- [10] M. Elyasi, A.A. Estaji, and M. Robat Sarpoushi, *Locally functionally countable subalgebra of* $\mathcal{R}(L)$, Arch. Math. (Brno) Tomus. **56** (2020), 127–140. https://doi.org/10.5817/AM2020--3--127.1
- [11] A.A. Estaji and M. Abedi, On injectivity of the ring of real valued continuous function on a frame, Bull. Belg. Math. Soc. Simon Stevin 25 (2018), no. 14, 467–480 https://doi.org/10.36045/bbms/1536631239.10
- [12] A.A. Estaji, A. Karimi Feizabadi, and M. Robat Sarpoushi, z_c -ideals and prime ideals in ring $\mathcal{R}_c(L)$, Faculty of Sci. Math., Univ. Nis, Serbia. **32** (2018), 6741–6752. https://doi.org/10.2298/FIL1819741E.1
- [13] A.A. Estaji, M. Robat Sarpoushi, and M. Elyasi, *Further thoughts on the ring* $\mathcal{R}_c(L)$ *in frames*, Algebra Universalis **80** (2019), no. 43, 14 pp. https://doi.org/10.1007/s00012-019-0619-z. 1, 3, 9, 10
- [14] A.A. Estaji and M. Taha, *The clean elements of the ring* $\mathcal{R}(L)$, Accepted in Czechoslovak Math. J. (2023). 1, 3
- [15] M. Ghadermazi, O.A.S. Karamzadeh, and M. Namdari, On the functionally countable subalgebra of C(X), Rend. Sem. Mat. Univ. Padova **129** (2013), 47–69. https://doi.org/10.4171/RSMUP/129--4.10
- [16] R. A. Gilmour, *Realcompact spaces and regular* σ *-frames*, Rend. Sem. Mat. Univ. Padova **96** (1984), 73–79. https://doi.org/10.1017/S0305004100061946 13
- [17] P.T. Johnstone, *Stone spaces*, Cambridge University Press, Cambridge, 1982. 1
- [18] A. Karimi-Feizabadi, A.A. Estaji, and M. Robat Sarpoushi, *Pointfree version of image of real-valued continuous functions*, Categ. Gen. Algebr. Struct. Appl. **9** (2018), no. 1, 59 –75. 1
- [19] J. Picado and A. Pultr, *Frames and Locales: Topology without Points*, Front. Math., Springer Basel, 2012. https://doi.org/10.1007/978-3-0348-0154-6.1, 2
- [20] M. Robat Sarpoushi, *Pointfree topology version of continuous functions with countable image*, Ph.D. Thesis, Univ. Hakim Sabzevari, Sabzevar, Iran, 2017. 1
- [21] M. Taha, A.A. Estaji, and M. Robat Sarpoushi, *On the regularity of* $C_c(L)$, 53^{nd} Annual Iranian Mathematics Conference University of Science, September 5-8, Technology of Mazandaran (2022). 1, 3
- [22] _____, *The pointfree version of* $C_c(X)$ *via the range of functions*, Accepted in Math. Slovaca (2023). 1, 3, 9

(Ali Akbar Estaji) Faculty of Mathematics and Computer Sciences, Hakim Sabzevari University, Sabzevar, Iran.

Email address: aaestaji@hsu.ac.ir

(Maryam Taha) Faculty of Mathematics and Computer Sciences, Hakim Sabzevari University, Sabzevar, Iran.

Email address: tahamaryam872@gmail.com