Significant circular Unruh effect at small acceleration

Yuebing Zhou,^{1,2} Jiawei Hu,^{2,*} and Hongwei Yu^{2,†}

¹Department of Physics, Huaihua University, Huaihua, Hunan 418008, China

²Department of Physics, Synergetic Innovation Center for Quantum Effects and Applications,

and Institute of Interdisciplinary Studies,

Hunan Normal University, Changsha, Hunan 410081, China

We study the transition rates of an atom rotating in a circular orbit, which is coupled with fluctuating electromagnetic fields in vacuum. We find that when the rotational angular velocity exceeds the transition frequency of the atom, the excitation rate can reach the same order of magnitude as the emission rate, even with an extremely low centripetal acceleration resulting from a very small orbital radius. For experimentally accessible centripetal accelerations, the excitation rate of centripetally accelerated atoms can be up to ten to the power of two hundred thousand times that of linearly accelerated atoms with the same acceleration. Our result suggests that the circular version of the Unruh effect can be significant even at very small centripetal accelerations, contrary to the common belief that a large Unruh effect requires large acceleration. This finding sheds new light on the experimental detection of the circular Unruh effect.

Introduction.—It is well known that, in vacuum, an atom in an excited state can spontaneously transition to its ground state and emit a photon, whereas an atom in the ground state can never transition to an excited state. Remarkably, however, spontaneous excitation can occur for uniformly accelerated ground-state atoms, and the spontaneous excitation rate Γ_{-} is related to the spontaneous emission rate Γ_{+} as

$$\Gamma_{-} = e^{-2\pi\omega_0/a} \Gamma_{+} = e^{-\omega_0/T_U} \Gamma_{+}, \qquad (1)$$

where ω_0 and a are the transition frequency and proper acceleration of the atom respectively. This means that a uniformly accelerated atom would excite as if it were immersed in a bath of thermal radiation at a temperature $T_U = a/2\pi$. This striking phenomenon is known as the Unruh effect [1-3] (see Refs. [4-6], for reviews). The Unruh effect is not only intriguing on its own but also closely related to other quantum effects in curved spacetimes, such as the Hawking radiation [7], from the point of view of the equivalence principle regarding acceleration and gravity. Eq. (1) reveals that the spontaneous excitation rate of a uniformly accelerated atom will be much smaller than the spontaneous emission rate when the acceleration is small compared with the transition frequency of the atom. This indicates that a very large acceleration compared with the atomic transition frequency is required in experimental detection of the effect, which is extremely challenging.

One may wonder what happens if an atom experiences centripetal acceleration, which is another common type of acceleration besides uniform acceleration. In this context, it has been found that a centripetally accelerated observer, similar to a uniformly accelerated one, would also perceive radiation in a vacuum, although the spectrum perceived is non-thermal [8–13]. Correspondingly, atoms undergoing centripetal acceleration can also get excited spontaneously [14–17], although the relation between the excitation rate and the emission rate is now not as simple as Eq. (1). This phenomenon is dubbed the circular Unruh effect. Intuitively, one may expect that a large centripetal acceleration is also necessary to obtain a relatively large spontaneous excitation rate. In fact, extensive studies on the circular Unruh effect over recent decades align with this expectation [18–30]. Many of these studies were conducted in the ultrarelativistic limit, i.e., in the limit where the linear velocity approaches the speed of light while the centripetal acceleration remains finite [11, 17-22]. In this limit, the relation between the upward and downward transition rates is similar to that in the linear acceleration case described by Eq. (1), and an effective temperature can thus be defined using the ratio between these two rates, which is approximately proportional to the centripetal acceleration. There are also investigations in which the ultrarelativistic limit is not taken [23-30]. However, to the best of our knowledge, in all known works, the spontaneous excitation rate of centripetally accelerated atoms approaches zero in the limit of vanishingly small centripetal acceleration. This is consistent with what we have learned from the linear Unruh effect and seems to imply that an extremely large centripetal acceleration is indeed necessary to detect the circular Unruh effect with spontaneous excitation.

Contrary to the common wisdom that a large Unruh effect demands a large acceleration, we discover in this Letter that an atom rotating in a circular orbit can exhibit a large spontaneous excitation rate even at a vanishingly small centripetal acceleration. Specifically, we investigate the transition rates of a centripetally accelerated atom coupled with fluctuating electromagnetic fields in vacuum. We show in detail that when the angular velocity exceeds the transition frequency of the atom, the spontaneous excitation rate, even in the limit of a vanishingly small orbital radius and consequently a vanishingly small centripetal acceleration, is not only nonvanishing but also, unexpectedly, of the same order of magnitude as the emission rate. Moreover, the effective temperature defined by the ratio of the excitation and emission rates is approximately proportional to the rotational angular velocity when the angular velocity is much larger than the transition frequency of the atom. This is remarkably different from the linear acceleration case, where the temperature is proportional to the acceleration. This suggests that we can observe a significant circular Unruh effect at an almost vanishing centripetal acceleration. Units with $\hbar = c = \epsilon_0 = k_B = 1$ are used in this Letter, where \hbar is the reduced Planck constant, c the speed of light, ϵ_0 the vacuum permittivity, and k_B the Boltzmann constant.

The transition rates for a rotating atom.—We consider a circularly moving two-level atom in interaction with the fluctuating electromagnetic fields in vacuum. The atom is assumed to be rotating at an angular velocity Ω in a circular orbit with a radius R, and so its trajectory can be described in the cylindrical coordinates as

$$r(t) = R, \qquad \theta(t) = \Omega t, \qquad z(t) = 0, \qquad (2)$$

where t is the coordinate time. The atom is assumed to be polarizable but does not possess a permanent dipole moment. However, it can be instantaneously polarized by fluctuating electromagnetic fields in vacuum. In the proper frame of the atom, the Hamiltonian describing the atom-field interaction can be written in a manifestly invariant form as [18] $H_I^{(0)} = -D^{\mu}F_{\mu\nu}u^{\nu}$, where $F_{\mu\nu}$ is the electromagnetic tensor, D^{μ} is the four-electric dipole moment, which can be expressed as $(0, D_r, D_{\theta}, D_z)$ in the proper frame, and u^{ν} is the four-velocity. In the following, we work in the laboratory frame, in which $D^{\mu} =$ $(\gamma R\Omega D_{\theta}, D_r, \gamma D_{\theta}/R, D_z)$, and $u^{\nu} = (\gamma, 0, \gamma \Omega, 0)$. Thus, the interaction Hamiltonian in the laboratory frame can be written as

$$H_I = -\sum_{i=r,\theta,z} D_i \mathcal{E}_i(t, \mathbf{x}), \tag{3}$$

where

$$\begin{aligned} \mathcal{E}_r(t, \mathbf{x}) &= E_r(t, \mathbf{x}) + R\Omega B_z(t, \mathbf{x}), \\ \mathcal{E}_\theta(t, \mathbf{x}) &= \gamma^{-1} E_\theta(t, \mathbf{x}), \\ \mathcal{E}_z(t, \mathbf{x}) &= E_z(t, \mathbf{x}) - R\Omega B_r(t, \mathbf{x}). \end{aligned}$$
(4)

Here $E_i(t, \mathbf{x})$ and $B_i(t, \mathbf{x})$ are the electric field and magnetic induction strength, respectively, and (t, \mathbf{x}) is the abbreviation for (t, r, θ, z) .

In the framework of open quantum systems, the emission rate Γ_+ and the excitation rate Γ_- can be derived as,

$$\Gamma_{\pm}(\omega) = \sum_{i,j=r,\theta,z} d_i d_j^* \int_{-\infty}^{\infty} G_{ij}(\Delta t) \ e^{\pm i\omega\Delta t} d\Delta t, \quad (5)$$

where $d_i = \langle 1|D_i|0\rangle$ is the dipole transition matrix element, ω is the energy level spacing of the atom in the laboratory frame, which is related to that in the proper frame of the atom ω_0 via $\omega = \omega_0/\gamma > 0$, and

$$G_{ij}(\Delta t) = \left\langle \mathcal{E}_i(t, \mathbf{x}(t)) \mathcal{E}_j(t', \mathbf{x}(t')) \right\rangle \tag{6}$$

are the electromagnetic field correlation functions, with $\langle \cdots \rangle$ denoting the expectation value with respect to the vacuum state of the electromagnetic field. Note that the field correlation functions here are invariant under temporal translations, i.e., they are functions of $\Delta t = t - t'$. For simplicity, we assume that the transition matrix elements d_i are real, then the cross terms in Eq. (5) will be vanishing. By substituting the trajectory of the atom Eq. (2) into the general form of the field correlation functions shown in Eq. (6), and with the help of the quantized vector potential of the electromagnetic field in the cylindrical coordinates [31, 32], one obtains the explicit forms of the field correlation functions in the laboratory frame as

$$G_{rr}(\Delta t) = \frac{1}{4\pi^2} \sum_{m=-\infty}^{+\infty} \sum_{\zeta,\alpha,\beta=-1}^{1} \int_{0}^{+\infty} dk_{\perp} \int_{-\infty}^{+\infty} dk_{z} \frac{k_{\perp}}{4\omega_{k}} J_{|m+\zeta|+\alpha}(k_{\perp}R) J_{|m+\zeta|+\beta}(k_{\perp}R) e^{i(m\Omega-\omega_{k})\Delta t} \\ \times \left\{ \frac{|\zeta| \left(R^{2}\Omega^{2}+2\right)-2}{4} k_{\perp}^{2}\alpha\beta + (-1)^{\frac{|\alpha|+|\beta|+\alpha+\beta}{2}} \frac{|\zeta|(1-|\alpha\beta|)(R\Omega k_{\perp})^{|\alpha|+|\beta|}}{2[\Omega+\zeta(m\Omega-\omega_{k})]^{|\alpha|+|\beta|-2}} \right\},$$
(7)

$$G_{\theta\theta}(\Delta t) = \frac{1}{4\pi^2} \sum_{m=-\infty}^{+\infty} \sum_{\zeta=-1}^{1} \int_{0}^{+\infty} dk_{\perp} \int_{-\infty}^{+\infty} dk_{z} \frac{k_{\perp}}{4\omega_{k}} J_{|m+\zeta|}^{2}(k_{\perp}R) e^{i(m\Omega-\omega_{k})\Delta t} \gamma^{-2} \left[\frac{2m^{2}}{R^{2}} (|\zeta|-1) + \omega_{k}^{2} |\zeta| \right], \quad (8)$$

$$G_{zz}(\Delta t) = \frac{1}{4\pi^2} \sum_{m=-\infty}^{+\infty} \sum_{\zeta=-1}^{1} \int_{0}^{+\infty} dk_{\perp} \int_{-\infty}^{+\infty} dk_{z} \frac{k_{\perp}}{4\omega_{k}} J_{|m+\zeta|}^{2}(k_{\perp}R) e^{i(m\Omega-\omega_{k})\Delta t} \\ \times \left\{ 2(m\Omega-\omega_{k})^{2}(1-|\zeta|) + k_{z}^{2} \left[(3-\gamma^{-2})|\zeta|-2 \right] \right\},$$
(9)

where $J_{\nu}(z)$ is the Bessel function of the first kind. Plugging the explicit forms of the field correlation functions

Eqs. (7)-(9) into Eq. (5), the transition rates Γ_{\pm} can be found to be

$$\Gamma_{\pm}(\omega) = \sum_{m=-\infty}^{+\infty} \frac{(m\Omega \pm \omega)^3}{3\pi} \Theta(m\Omega \pm \omega) P_m^{\pm}(\Omega/\omega, R\omega),$$
(10)

where $\Theta(x)$ the Heaviside theta function which equals 0 for x < 0 and 1 for x > 0, and

$$P_m^{\pm}(\Omega/\omega, R\omega) = \sum_{\zeta,\alpha,\beta} \sum_{k=0}^{+\infty} \frac{3\left(d_r^2 \mathcal{M}_m^{\pm} + d_{\theta}^2 \mathcal{N}_m^{\pm} + d_z^2 \mathcal{Z}_m^{\pm}\right) (-1)^{k+h} \left[(k+p+q)!\right]^2 (2k+2p)! \left[R\omega(m\Omega/\omega\pm 1)\right]^{2(k+p+q)}}{4\left(\left||m+\zeta|+\alpha\right|+k\right)! \left(\left||m+\zeta|+\beta\right|+k\right)! (2k+2p+2q+1)! (k+2p)! k!} \right]} .$$
(11)

Here,
$$\zeta, \alpha, \beta \in \{-1, 0, 1\}, p = \frac{\left||m+\zeta|+\alpha\right|+\left||m+\zeta|+\beta\right|}{2}, q = \frac{|\alpha|+|\beta|}{2}, g = \frac{\alpha+\beta}{2}, h = p - g - |m+\zeta|$$
, and

$$\mathcal{M}_{m}^{\pm} = \frac{(-1)^{q+g} |\zeta| (1 - |\alpha\beta|) (\Omega/\omega)^{2q}}{(\Omega/\omega \mp \zeta)^{2q-2} (m\Omega/\omega \pm 1)^{2}} + \frac{|\zeta| \left[(R\omega)^{2} (\Omega/\omega)^{2} + 2 \right] - 2}{(R\omega)^{2} (m\Omega/\omega \pm 1)^{2}} \alpha\beta,$$
(12)

$$\mathcal{N}_{m}^{\pm} = \left[|\zeta| - \frac{2m^{2}(1 - |\zeta|)}{(R\omega)^{2}(m\Omega/\omega \pm 1)^{2}} \right] \frac{(1 - |\alpha|)(1 - |\beta|)}{\gamma^{2}}, \tag{13}$$

$$\mathcal{Z}_{m}^{\pm} = \left[\frac{2(1-|\zeta|)}{(m\Omega/\omega\pm1)^{2}} + \frac{(3-\gamma^{-2})|\zeta|-2}{2|m+\zeta|+2k+3}\right](1-|\alpha|)(1-|\beta|).$$
(14)

In the following, we will separately calculate the contributions to the transition rates of the polarization of the atom along the axis of rotation and that perpendicular to it.

(I) Atomic polarization along the axis of rotation.

First, we consider the transition rate contributed by the atomic polarization along the direction of the rotation axis (i.e., the z-axis). When the orbital radius R is small such that $v = R\Omega \ll 1$ and $R\omega_0 \ll 1$, the emission and excitation rates of the atom Eq. (10) can be approximated as

$$\Gamma_{+} = \frac{d_{z}^{2}\omega_{0}^{3}}{3\pi} + O\left[(R\omega_{0})^{2}\right],$$

$$\Gamma_{-} = O\left[(R\omega_{0})^{2\lceil\omega_{0}/\Omega\rceil}\right],$$
(15)

where $O[x^n]$ denotes that infinitesimals of the nth and higher orders of x are omitted, and $\lceil x \rceil$ gives the smallest integer greater than or equal to x. From Eq. (15), it is obvious that: 1) The excitation rate of the atom is always a higher-order small quantity compared with the emission rate. 2) When the orbital radius R tends to zero, the emission and excitation rates of the atom tend to those of inertial atoms in vacuum. This aligns with one's intuition since the centripetal acceleration $a = \gamma^2 R \Omega^2$ vanishes when $R \to 0$.

(II) Atomic polarization perpendicular to the axis of rotation.

We now consider the transition rate contributed by the atomic polarization perpendicular to the axis of rotation (i.e., the radial and tangential direction of the orbit). In this case, the approximated expressions of the emission and excitation rates in the limit of a small orbital radius R ($v = R\Omega \ll 1$ and $R\omega_0 \ll 1$) are obtained from Eq.

(10) as follows

$$\Gamma_{+} = \left(d_{r}^{2} + d_{\theta}^{2}\right) \frac{\omega_{0}^{3} + 3\Omega^{2}\omega_{0}}{3\pi} + O[(R\omega_{0})^{2}],$$

$$\Gamma_{-} = O\left[(R\omega_{0})^{2\lceil\omega_{0}/\Omega\rceil - 2}\right],$$
(16)

for $\Omega < \omega_0$, and

$$\Gamma_{\pm} = \left(d_r^2 + d_{\theta}^2\right) \frac{(\Omega \pm \omega_0)^3}{6\pi} + O\left[(R\omega_0)^2\right], \quad (17)$$

for $\Omega \geq \omega_0$. From Eqs. (16) and (17), we can draw the following conclusions.

1) When the rotational angular velocity does not reach the transition frequency of the atom, i.e, $\Omega < \omega_0$, the excitation rate is a higher-order small quantity compared with the emission rate. As the orbital radius R tends to zero, the excitation rate approaches that of an inertial atom in vacuum, meaning it will vanish. In contrast, the emission rate does not vanish because the rotation introduces an extra term $\left(d_r^2 + d_\theta^2\right) \frac{\Omega^2 \omega_0}{\pi}$ to the emission rate.

2) When the rotational angular velocity exceeds the transition frequency of the atom, i.e, $\Omega > \omega_0$, the excitation rate will be of the same order of magnitude as the emission rate, even in the limit of a small orbital radius R such that the centripetal acceleration $a = \gamma^2 R \Omega^2$ is vanishing.

3) Comparing Eqs. (16) and (17) with Eq. (15), one can find that, when the orbital radius is small, the excitation rate contributed by the atomic polarization along

the direction of the axis of rotation is always an infinitesimal of higher order than that contributed by the atomic polarization perpendicular to the axis of rotation, regardless of whether the rotational angular velocity exceeds the transition frequency of the atom.

The effective temperature.—Usually, atoms are isotropically polarizable. Since the atomic polarizability is related to the dipole transition matrix elements as $\alpha_i =$ $2|d_i|^2/\omega_0$ for a two-level atom [33], and we have assumed that d_i are real, then isotropic polarizability implies that $d_r^2 = d_{\theta}^2 = d_z^2 \equiv d^2$. Then, according to Eqs. (15), (16) and (17), one obtains

$$\Gamma_{+} = d^{2} \frac{\omega_{0}^{3} + 2\Omega^{2}\omega_{0}}{\pi} + O\left[(R\omega_{0})^{2}\right],$$

$$\Gamma_{-} = O\left[(R\omega_{0})^{2\lceil\omega_{0}/\Omega\rceil - 2}\right],$$
(18)

for $\Omega < \omega_0$, and

$$\Gamma_{+} = d^{2} \frac{(\Omega + \omega_{0})^{3} + \omega_{0}^{3}}{3\pi} + O\left[(R\omega_{0})^{2}\right],$$

$$\Gamma_{-} = d^{2} \frac{(\Omega - \omega_{0})^{3}}{3\pi} + O\left[(R\omega_{0})^{2}\right],$$
(19)

for $\Omega \geq \omega_0$.

As the spontaneous transition persists for a sufficiently long time, the ratio of the probabilities for the atom in the ground and excited states reaches a steady value, and the steady state of the atom $\rho(t)$ becomes a thermal state

$$\lim_{t \to \infty} \rho(t) = \frac{e^{-H_A/T_{\text{eff}}}}{\text{Tr}[e^{-H_A/T_{\text{eff}}}]},$$
(20)

where H_A is the Hamiltonian of the two-level atom, and

$$T_{\rm eff} = \frac{\omega_0}{\ln(\Gamma_+/\Gamma_-)},\tag{21}$$

is the effective temperature. For an ensemble of rotating two-level atoms, the effective temperature characterizes the ratio of the population of atoms in the ground and excited states in the steady state. See, e.g., Refs. [8, 12]. According to the definition of the effective temperature (21) and Eqs. (18)-(19), when the rotational angular velocity is less than the transition frequency of the atom, i.e, $\Omega < \omega_0$, the effective temperature is

$$T_{\rm eff} \sim \omega_0 / \ln \left[(R\omega_0)^{2-2\lceil \omega_0/\Omega \rceil} \right],$$
 (22)

which depends on the orbital radius R, and vanishes as Rand thus the centripetal acceleration tend to zero. However, when the rotational angular velocity exceeds the transition frequency of the atom, i.e., $\Omega > \omega_0$, the effective temperature becomes

$$T_{\rm eff} = \frac{\omega_0}{\ln \{ [(\Omega + \omega_0)^3 + \omega_0^3] / (\Omega - \omega_0)^3 \}},$$
 (23)

which is independent of the orbital radius and the centripetal acceleration. In principle, the orbital radius and

thus the centripetal acceleration can be arbitrarily small while keeping the rotational angular velocity larger than the transition frequency of the atom. Therefore, surprisingly, a large effective temperature in the centripetal acceleration case can be obtained at an almost vanishing centripetal acceleration. Furthermore, when the rotational angular velocity is much larger than the transition frequency, i.e., when $\Omega \gg \omega_0$, the effective temperature can further be approximated as

$$T_{\rm eff} \approx \frac{\Omega}{6}.$$
 (24)

This shows that the effective temperature is approximately proportional to the angular velocity when the rotational angular velocity is much larger than the energy level spacing, which is remarkably different from the linear acceleration case, where the temperature is proportional to the acceleration.

Discussion.—A few comments are now in order.

First, the reason that the transition rate for the rotating atom does not reduce to that in the inertial case in the limit of a vanishingly small orbital radius and thus a vanishingly small centripetal acceleration can be traced back to the interaction Hamiltonian (3). In the limit of $R \to 0$ and thus $v \to 0$ and $\gamma \to 1$, but the angular velocity Ω is finite, the interaction Hamiltonian reduces to $H_I = -\sum_{i=r,\theta,z} D_i E_i(t, \mathbf{x})$. Although this seems to be the usual dipole interaction Hamiltonian, one should bear in mind that D_i are components of the dipole operator in the proper frame of the rotating atom, which are related to those in the laboratory frame by a rotation transformation. For atoms polarizable along the direction of the axis of rotation, the dipole is invariant under the rotation transformation. However, this is not the case for atoms polarizable perpendicular to the direction of the axis of rotation. This explains why the transition rate for a rotating atom with a finite angular velocity does not reduce to that in the inertial case in the limit of a vanishingly small centripetal acceleration. Note that this phenomenon does not exist in the toy model in previous studies where the atom is assumed to be in the monopole interaction with a scalar field [11, 14–18, 21–23, 25–29].

Second, why does the excitation rate contributed by the atomic polarization perpendicular to the axis of rotation change dramatically when the rotational angular velocity Ω crosses the energy level spacing ω_0 in the limit of a small orbital radius R? To answer this question, let us note that when the orbital radius R is small such that $v = R\Omega \ll 1$, the field correlation functions (7)-(9) can be approximated as,

$$G_{rr}(\Delta t) = G_0(\Delta t) \frac{e^{i\Omega\Delta t} + e^{-i\Omega\Delta t}}{2} + O[(R\Omega)^2], \quad (25)$$

$$G_{\theta\theta}(\Delta t) = G_0(\Delta t) \frac{e^{\kappa \Delta t} + e^{-\kappa \Delta t}}{2} + O[(R\Omega)^2], \quad (26)$$
$$G_{zz}(\Delta t) = G_0(\Delta t) + O[(R\Omega)^2], \quad (27)$$

$$G_{zz}(\Delta t) = G_0(\Delta t) + O[(R\Omega)^2], \qquad (27)$$

where $G_0(\Delta t) = \frac{1}{\pi^2(\Delta t - i\epsilon)^4}$ is the field correlation function obtained by taking the trajectory of an inertial atom into the general expression (6), with ϵ being a positive infinitesimal. Now, one can easily obtain, in the leading order, from Eqs. (25)-(27) that, for the case of the atomic polarization along the radial or tangential direction of the orbit,

$$\Gamma_{\pm}(\omega_0) = \frac{1}{2} \left[\Gamma^0_{\pm}(\omega_0 + \Omega) + \Gamma^0_{\pm}(\omega_0 - \Omega) \right].$$
 (28)

Here, $\Gamma^{0}_{+}(\omega_{0}) = d^{2}\omega_{0}^{3}/(3\pi)$ and $\Gamma^{0}_{-}(\omega_{0}) = 0$ are the spontaneous emission rate and excitation rate for inertial atoms respectively. Moreover, from Eq. (5), we have that

$$\Gamma_{\pm}(-\omega) = \Gamma_{\mp}(\omega). \tag{29}$$

Therefore, for the case of the atomic polarization along the direction perpendicular to the axis of rotation, when the orbital radius is sufficiently small and the rotational angular velocity Ω is smaller than the energy level spacing ω_0 , i.e., $\Omega < \omega_0$, the excitation rate is approximately half of the sum of the excitation rates of two inertial atoms in vacuum with the energy level spacing being $\omega_0 + \Omega$ and $\omega_0 - \Omega$ respectively, which is vanishing. However, when $\Omega > \omega_0$, i.e., $\omega_0 - \Omega < 0$, the excitation rate is approximately half of the sum of the excitation rate of an inertial atom with an energy level spacing $\omega_0 + \Omega$ (which is vanishing), and the *emission* rate of an inertial atom with an energy level spacing $\Omega - \omega_0$ (which is finite). This explains why the excitation rate undergoes a sudden change when the rotational angular velocity Ω crosses the energy level spacing ω_0 .

Third, the rotating atoms are treated as point-like particles in this Letter. However, actually, the size of an atom must be nonvanishing. In order to avoid the consideration of the finite-size effect, the orbital radius Rshould be much larger than the radius of the atom characterized by the Bohr radius $R_B \sim 10^{-10}$ m, although it has been assumed to be small such that $R\omega_0 \ll 1$ and $R\Omega \ll 1$. Note that the typical energy level spacing of an atom is of the order of $\omega_0 \sim 1$ eV, which corresponds to a transition wavelength $\omega_0^{-1} \sim 10^{-6}$ m. Therefore, if we assume that the rotational angular velocity Ω is of the same order of ω_0 , there exists plenty parameter space for the orbital radius R such that it is much larger than the radius of the atom and at the same time much smaller than the length characterized by the rotational angular velocity Ω and the energy level spacing ω_0 .

Fourth, let us note that the transition rates shown in Eq. (5) is derived under the Markov approximation. By adopting the method presented in Ref. [34], it can be shown that the Markov approximation is justified regardless of whether the rotational angular velocity Ω exceeds the transition frequency ω_0 .

Finally, let us discuss the possibility of verifying this effect in experiment. Recently, there have been efforts

to achieve hyperfast rotation in an optically levitated nanoparticle system [35-37], and the highest rotation angular frequency obtained so far is $\Omega \approx 6$ GHz for nanoparticles with an average radius R = 95 nm [37]. Then, a natural idea is that we can manage to attach the atom to a hyperfast rotating nanoparticle, as proposed in Ref. [30]. Based on the data in Ref. [37], we assume that $\omega_0 \sim 1$ GHz, $\Omega = 6\omega_0$, $R \sim 10^{-7}$ m, corresponding to a centripetal acceleration $a \sim 10^{11} \text{ m/s}^2$. This is much smaller than the transition frequency of the atom since $a/\omega_0 \sim 10^{-6} \ll 1$. For a uniformly accelerated atom at such an acceleration, the excitation rate is completely negligible compared with the emission rate, since $\Gamma_{-}^{\text{linear}} \sim 10^{-272876} \Gamma_{+}^{\text{linear}}$. For a rotating atom, in contrast, the excitation rate is of the same order of the emission rate, and at the same time it is extraordinarily larger than that in the linear acceleration case, as now $\Gamma_{-}^{\text{circular}} \sim \Gamma_{+}^{\text{circular}} \sim 10^{272878} \Gamma_{-}^{\text{linear}}$. This shows that, compared with the linear Unruh effect, the circular Unruh effect can be significant even at a centripetal acceleration extremely small compared with the transition frequency of the atom. In fact, the excitation rate will be even larger if a larger rotational angular velocity can be reached. For an optically levitated nanoparticle system, the rotational angular velocity increases as the size of the particle decreases [36]. Moreover, from Eq. (19), we can see that the leading term of the transition rate is independent of the orbital radius R when the angular velocity exceeds the energy level spacing. This indicates that, one can achieve a higher rotational angular velocity by reducing the size of the nanoparticle to which the atom is attached, thereby obtaining a more significant circular Unruh effect. Therefore, the phenomenon discovered in this Letter mcircularay potentially be observed with the help of an optically levitated nanoparticle system.

Summary.—In this Letter, we have investigated the transition rate of an atom rotating in a circular orbit, which is coupled with fluctuating electromagnetic fields in vacuum. We have shown that when the rotational angular velocity exceeds the transition frequency of the atom, the spontaneous excitation rate, even in the limit of a vanishingly small orbital radius and thus a vanishingly small centripetal acceleration, is not only nonvanishing but also unexpectedly of the same order of magnitude as that of the emission rate. Moreover, the effective temperature of the circular Unruh effect, defined by the ratio of the excitation and emission rates, is approximately proportional to the rotational angular velocity when the orbital radius is vanishingly small and the rotational angular velocity is much larger than the energy level spacing. This is remarkably different from the linear acceleration case, where the temperature is proportional to the acceleration. Our result, therefore, suggests that a significant circular Unruh effect can be observed at a vanishingly small centripetal acceleration. This phenomenon may potentially be observed with the help of an optically

levitated nanoparticle system.

We would like to thank Wenting Zhou for helpful discussions. We also extend our gratitude to the anonymous referees for their insightful comments and helpful suggestions. This work was supported in part by the NSFC under Grant No. 12075084, No. 12375047, the innovative research group of Hunan Province under Grant No. 2024JJ1006, the Hunan Provincial Natural Science Foundation of China under Grant No. 2023JJ40515, and the Scientific Research Program of Education Department of Hunan Province of China under Grant No. 22B0762.

* Corresponding author: jwhu@hunnu.edu.cn † Corresponding author: hwyu@hunnu.edu.cn

- S. A. Fulling, Nonuniqueness of canonical field quantization in Riemannian space-time, Phys. Rev. D 7, 2850 (1973).
- [2] P. C. W. Davies, Scalar production in Schwarzschild and Rindler metrics, J. Phys. A 8, 609 (1975).
- [3] W. G. Unruh, Notes on black-hole evaporation, Phys. Rev. D 14, 870 (1976).
- [4] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, United Kingdom, 1982).
- [5] R. M. Wald, Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics (University of Chicago Press, Chicago, 1994).
- [6] L. C. B. Crispino, A. Higuchi, and G. E. A. Matsas, The Unruh effect and its applications, Rev. Mod. Phys. 80, 787 (2008).
- [7] S. W. Hawking, Black hole explosions? Nature (London) 248, 30 (1974).
- [8] J. S. Bell and J. M. Leinaas, Electrons as accelerated thermometers, Nucl. Phys. B212, 131 (1983).
- [9] S. Hacyan, A. Sarmiento, Vacuum energy of the electromagnetic field in a rotating system, Phys. Lett. B 179, 287 (1986).
- [10] J. S. Bell and J. M. Leinaas, The Unruh effect and quantum fluctuations of electrons in storage rings, Nucl. Phys. B284, 488 (1987).
- [11] S. K. Kim, K. S. Soh, and J. H. Yee, Zero-point field in a circular-motion frame, Phys. Rev. D 35, 557 (1987).
- [12] W. G. Unruh, Acceleration radiation for orbiting electrons, Phys. Rep. 307, 163 (1998).
- [13] H. C. Rosu, Quantum vacuum radiation and detection proposals, Int. J. Theor. Phys. 44, 493 (2005).
- [14] J. R. Letaw, Stationary world lines and the vacuum excitation of noninertial detectors, Phys. Rev. D 23, 1709 (1981).
- [15] J. I. Korsbakken and J. M. Leinaas, Fulling-Unruh effect in general stationary accelerated frames, Phys. Rev. D 70, 084016 (2004).
- [16] M. Good, B. A. Juárez-Aubry, D. Moustos, and M. Temirkhan, Unruh-like effects: Effective temperatures along stationary worldlines, J. High Energy Phys. 06 059 (2020).
- [17] S. Takagi, On the response of a particle detector in a circular motion, Prog. Theor. Phys. 72, 1270 (1984)
- [18] S. Takagi, Vacuum Noise and Stress Induced by Uniform

Acceleration: Hawking-Unruh Effect in Rindler Manifold of Arbitrary Dimension, Prog. Theor. Phys. Suppl. 88, 1 (1986).

- [19] Y. Jin, J. Hu, and H. Yu, Spontaneous excitation of a circularly accelerated atom coupled to electromagnetic vacuum fluctuations, Ann. Phys. (Amsterdam) 344, 97 (2014).
- [20] Y. Jin, J. Hu, and H. Yu, Dynamical behavior and geometric phase for a circularly accelerated two-level atom, Phys. Rev. A 89, 064101 (2014).
- [21] S. Biermann, S. Erne, C. Gooding, J. Louko, J. Schmiedmayer, W. G. Unruh, and S. Weinfurtner, Unruh and analogue Unruh temperatures for circular motion in 3+1and 2+1 dimensions, Phys. Rev. D 102, 085006 (2020).
- [22] C. R. D. Bunney and J. Louko, Circular motion analogue Unruh effect in a thermal bath: robbing from the rich and giving to the poor, Class. Quant. Grav. 40, 155001 (2023).
- [23] J. R. Letaw and J. D. Pfautsch, Quantized scalar field in rotating coordinates, Phys. Rev. D 22, 1345 (1980).
- [24] J. Rogers, Detector for the temperaturelike effect of acceleration, Phys. Rev. Lett. 61, 2113 (1988).
- [25] O. Levin, Y. Peleg and A. Peres, Unruh effect for circular motion in a cavity, J. Phys. A: Math. Gen. 26, 3001 (1993)
- [26] P. C. W. Davies, T. Dray, and C. A. Manogue, Detecting the rotating quantum vacuum, Phys. Rev. D 53, 4382 (1996).
- [27] V. A. De Lorenci, R. D. M. De Paola and N. F. Svaiter, The rotating detector and vacuum fluctuations, Class. Quant. Grav. 17, 4241 (2000).
- [28] Y. Gim, H. Um, and W. Kim, Unruh temperatures in circular and drifted Rindler motions, arXiv.1806.11439.
- [29] C. R. D. Bunney, L. Parry, T. R. Perche, and J. Louko, Ambient temperature versus ambient acceleration in the circular motion Unruh effect, arXiv:2310.05700.
- [30] K. Lochan, H. Ulbricht, A. Vinante, and S. K. Goyal, Detecting acceleration-enhanced vacuum fluctuations with atoms inside a cavity, Phys. Rev. Lett. **125**, 241301 (2020).
- [31] A. N. Aliev and D. V. Gal'tsov, Gravitational Aharonov-Bohm radiation in string-generated conical space-time, Ann. Phys. (N.Y.) 193, 142 (1989).
- [32] H. Cai, H. Yu, and W. Zhou, Spontaneous excitation of a static atom in a thermal bath in cosmic string spacetime, Phys. Rev. D 92, 084062 (2015).
- [33] H. Friedrich, *Theoretical Atomic Physics* (Springer-Verlag, New York, 1990).
- [34] D. Moustos and C. Anastopoulos, Non-Markovian time evolution of an accelerated qubit, Phys. Rev. D 95, 025020 (2017).
- [35] R. Reimann, M. Doderer, E. Hebestreit, R. Diehl, M. Frimmer, D. Windey, F. Tebbenjohanns, and L. Novotny, GHz Rotation of an optically trapped nanoparticle in vacuum, Phys. Rev. Lett. **121**, 033602 (2018).
- [36] J. Ahn, Z. Xu, J. Bang, Y.-H. Deng, T. M. Hoang, Q. Han, R.-M. Ma, and T. Li, Optically levitated nanodumbbell torsion balance and GHz nanomechanical rotor, Phys. Rev. Lett. **121**, 033603 (2018).
- [37] Y. Jin, J. Yan, S. J. Rahman, J. Li, X. Yu, and J. Zhang, 6 GHz hyperfast rotation of an optically levitated nanoparticle in vacuum, Photon. Res. 9, 1344 (2021).