

REVISITING 3-SASAKIAN AND G_2 -STRUCTURES

SIMON SALAMON AND RAGINI SINGHAL

ABSTRACT. The algebra of exterior differential forms on a regular 3-Sasakian 7-manifold is investigated, with special reference to nearly-parallel G_2 3-forms. This is applied to the study of 3-forms invariant under cohomogeneity-one actions by $SO(4)$ on the 7-sphere and on Berger's space $SO(5)/SO(3)$.

CONTENTS

Introduction	1
1. Notation	2
2. $Spin(7)$ and G_2 -structures	4
3. The cohomogeneity-one action of $SO(4)$ on S^7	8
4. The cohomogeneity-one action of $SO(4)$ on B^7	12
5. Hypersurface calculations	18
Acknowledgements	20
References	20

INTRODUCTION

Let M be an oriented self-dual Einstein 4-manifold M with positive scalar curvature. The bundle of 2-forms

$$(0.1) \quad \Lambda^2 T^*M = \Lambda^+ \oplus \Lambda^-$$

splits into self-dual (SD) and anti-self-dual (ASD) subbundles, in line with the double covering

$$SO(4) \rightarrow SO(3)^+ \times SO(3)^-$$

of Lie groups. Our hypothesis implies that Λ^- has constant curvature in the sense of (1.5) below. The reason for focussing on Λ^- is that (with its standard orientation and Fubini-Study metric) the projective plane $\mathbb{C}\mathbb{P}^2$ satisfies the hypothesis. Much of the theory extends to orbifolds, and sources of examples include quaternion-Kähler quotients [19, 14], and the $SO(3)$ -invariant orbifolds discovered by Hitchin [24]. The latter are parametrized by an integer $k \geq 3$ and the first two examples are S^4 and $\mathbb{C}\mathbb{P}^2/\mathbb{Z}_2$.

The (total space of the) $SO(3)^-$ principal bundle P^- associated to Λ^- is 3-Sasakian [28], and therefore admits an Einstein metric and a family of nearly-parallel G_2 (for short, NP_2) structures. An equivalent way of expressing the 3-Sasakian condition is to say that the Riemannian cone $P^- \times \mathbb{R}^+$ is hyperkähler, meaning that it has a metric with holonomy $Sp(2)$ [7]. However, it is well known that a squashing of the fibres of P^- produces a second Einstein metric and canonical NP_2 structure whose cone has holonomy *equal* to $Spin(7)$.

Now suppose (restricting to an open set if necessary) that M is spin, so that its structure group lifts to $Spin(4) \cong SU(2)^2$. The negative spin bundle Σ^- can be regarded as a quaternionic line bundle over M , and admits a family of metrics with $Spin(7)$ holonomy, all asymptotic to the cone over (a $2 : 1$ cover of) the squashed NP_2 manifold [12]. An initial purpose of this article is to relate well-known constructions of the relevant differential forms on these spaces, one via a $Spin(7)$ 4-form on a vector bundle, the other via 3-Sasakian 1-forms on a principal bundle. This is motivated by the situation in which M itself admits an action by $SO(3)$ with 3-dimensional orbits. In this case, P^- or its double cover \tilde{P}^- will admit an action by $SO(4)$ or $SU(2)^2$ of cohomogeneity one, preserving the G_2 structures. This happens when M is S^4 or $\mathbb{C}P^2$, and more generally for the Hitchin orbifolds.

We shall make this explicit for the 7-sphere $S^7 \cong \tilde{P}^-S^4$, in order to express its squashed NP_2 structure in canonical form. The result is consistent with a known expression for the round metric on S^7 subject to $SO(4)$ and \mathbb{Z}_3 symmetry [35]. The same technique is applied to the isotropy irreducible space $B^7 = SO(5)/SO(3)$, which Berger showed has positive sectional curvature [9]. This space also has a homogeneous NP_2 structure [32], a cohomogeneity-one action by $SO(4)$ [31], and is diffeomorphic to \tilde{P}^+ for a non-trivial Hitchin orbifold [23]. Although not our main concern, exhaustive classifications of associative submanifolds in B^7 and in squashed 3-Sasakian manifolds are undertaken in [5, 6] using twistor correspondences, pursuing work in [30, 28] for S^7 .

Paying debts to this previous mountain of work, our survey highlights the distinction between the homogeneous NP_2 structures on S^7 and B^7 , when expressed in terms of the local differential geometry on an interval times $SO(4)$. The choice of geodesics orthogonal to the orbits allows one to construct bases (X_i) and (Y_i) of invariant 1-forms that enable the respective 3-forms to be built up using upper triangular 2×2 matrices; see Corollaries 3.4 and 4.4. These bases ultimately arise from different conjugacy classes of 3-dimensional subgroups of $SO(4)$, and different group diagrams for the cohomogeneity-one actions.

Just as a hypersurface of a G_2 holonomy manifold acquires a so-called half-flat $SU(3)$ structure, so a hypersurface of a NP_2 manifold has an analogous nearly-half-flat (NHF) property. There is a sense in which a nearly-Kähler manifold has both properties. We conclude with a brief investigation of induced structures on hypersurfaces of S^7 and B^7 , and we show that constants defining the homogeneous NP_2 structures are to some extent determined by the NHF condition. The results of this section form part of a broader project to describe exceptional structures in dimensions 6 and 7.

1. NOTATION

In the first part of this paper, we shall adopt the technique of [12], which relies on quaternion-valued differential forms on the total spaces of Λ^- and Σ^- , though only the latter will concern us.

The Riemannian metric of M^4 will be denoted by g_4 and its positive scalar curvature by 12κ , where κ is another constant. The sphere $S^n(r)$ of radius r has scalar curvature $n(n-1)/r^2$, so $\kappa = 1$ for $S^4(1)$. **From now on, we shall scale g_4 so that $\kappa = 1$.**

The induced connection on Σ^- is defined by a vector-valued 1-form

$$\phi = i\phi_1 + j\phi_2 + k\phi_3$$

on the total space of Σ^- . We shall refer to a differential form with values in $\text{Im } \mathbb{H}$ as ‘vector valued’, as opposed to ‘quaternion-valued’ (for \mathbb{H}). Thus, $\bar{\phi} = -\phi$, where a bar denotes quaternionic conjugate. The curvature of Σ^- is

$$(1.2) \quad \begin{aligned} \Phi &= d\phi + \phi \wedge \phi \\ &= i(d\phi_1 + 2\phi_{23}) + i(d\phi_2 + 2\phi_{31}) + i(d\phi_3 + 2\phi_{12}) \end{aligned}$$

where $\phi_{23} = \phi_2 \wedge \phi_3$. See [12, p 841].

The choice of a unit section of Σ^- determines a local orthonormal basis of ASD 2-forms via the isomorphism $S^2\Sigma^- \cong (\Lambda_-^2)_c$ of complex vector spaces or bundles. These 2-forms are the components of the tautological vector-valued 2-form

$$\frac{1}{2}(\bar{\omega} \wedge \omega) = i\Omega_1 + i\Omega_2 + k\Omega_3 = \Omega.$$

We use a Greek letter without an index to stand for a quantity with values in \mathbb{H} that encodes all the individual components. Scalings are fixed by the existence pointwise of an orthonormal basis $\{\omega_0, \omega_1, \omega_2, \omega_3\}$ of 1-forms for g_4 such that

$$(1.3) \quad \Omega_k = \omega_{0k} - \omega_{ij}$$

for $(ijk) = (123)$.

It follows from (1.2) that

$$(1.4) \quad d\Phi = -2\phi \wedge \Phi$$

where \wedge combines multiplication of imaginary quaternions (cross product) with wedging. With the notation above, the Einstein ASD condition becomes

$$(1.5) \quad \Phi = \frac{1}{2}\kappa\Omega$$

as stated in [12, p 842], although we are assuming throughout that $\kappa = 1$.

Now let

$$(1.6) \quad a = a_0 + a_1i + a_2j + a_3k$$

be a quaternionic coordinate on the fibre of Σ^- . Note that $\bar{a}a = a\bar{a} = R$ is the radius squared (denoted r in [12, p 846]). Set

$$(1.7) \quad \alpha = da - a\phi = \alpha_0 + i\alpha_1 + j\alpha_2 + k\alpha_3.$$

At each point, the annihilator of α is the horizontal subspace determined by the connection. We can then determine the effect of the action $a \rightsquigarrow aq$ on differential forms. It yields the transformations $\phi \rightsquigarrow \bar{q}\phi q$ and $\Omega \rightsquigarrow \bar{q}\Omega q$.

Next, consider the quaternion-valued 1-form

$$(1.8) \quad \beta = \bar{a}\alpha = \beta_0 + i\beta_1 + j\beta_2 + k\beta_3$$

as defined in [12, p 846]. In particular,

$$\beta_1 = a_0\alpha_1 - a_1\alpha_0 - a_2\alpha_3 + a_3\alpha_2$$

is the result of applying a fibrewise ASD almost complex structure to $\beta_0 = \frac{1}{2}dR$. However, from now on, we work on \tilde{P}^- , so that $R = 1$, and β becomes a vector-valued 1-form. In this setting,

Lemma 1.1. $d\beta + \beta \wedge \beta = -2 \operatorname{Im}(\phi \wedge \beta) - \Phi$.

Proof. Using (1.2), (1.5), and (1.7),

$$(1.9) \quad d\alpha = -(\alpha + a\phi) \wedge \phi - ad\phi = -\alpha \wedge \phi - a\Phi.$$

Therefore

$$\begin{aligned} d\beta &= d(\bar{a}\alpha) = d\bar{a} \wedge \alpha + \bar{a} d\alpha \\ &= d\bar{a} \wedge \alpha + \bar{a}(-\alpha \wedge \phi - a\Phi) \\ &= \bar{a} \wedge \alpha - (\phi \wedge \beta + \beta \wedge \phi) - \Phi. \end{aligned}$$

Now restrict to $R = 1$, so that β (and, as always, ϕ) have values in $\operatorname{Im} \mathbb{H}$. Because of the wedging, the real parts of $\phi \wedge \beta$ and $\beta \wedge \phi$ cancel out. Moreover,

$$(1.10) \quad \bar{a} \wedge \alpha = \bar{a}a \wedge \bar{a}\alpha = \bar{\beta} \wedge \beta = -\beta \wedge \beta$$

and the result follows. \square

The first component of β is

$$(1.11) \quad d\beta_1 + 2\beta_{23} = -2(\phi_2 \wedge \beta_3 - \phi_3 \wedge \beta_2) - \Phi_1.$$

Note that β_{23} is totally vertical whereas $\Phi_1 = \frac{1}{2}\kappa\Omega_1$ ‘lives’ on M , so their exterior derivatives cannot be proportional, which forces the appearance of the mixed terms on the right-hand side. The fibrewise ASD property of $d\beta$ is indicated by the presence of $\bar{a} \wedge \alpha$ rather than the self-dual $\alpha \wedge \bar{a}$.

2. SPIN(7) AND G_2 -STRUCTURES

One of the main results in [12, §4], by now well known, is that the total space of Σ^- admits a family of metrics with holonomy equal to $Spin(7)$ and asymptotically conical. The induced Einstein metric h_7 on the link of this conical limit therefore underlies a so-called nearly-parallel G_2 structure (our abbreviation is NP_2). The latter comprises a 3-form φ with stabilizer G_2 for which

$$(2.12) \quad d\varphi = \mu * \varphi$$

where $21\mu^2/8$ is known to equal the scalar curvature of h_7 [9]. The metric h_7 is also said to have weak holonomy G_2 [22]. We proceed to write down φ explicitly.

In order to extract useful real-valued differential forms from those defined in Section 1, we need to exploit the obvious invariants for the action of $SO(3)$ on $\operatorname{Im} \mathbb{H} \cong \mathbb{R}^3$, namely inner product and determinant. In this way, we can define the real 3-forms

$$\begin{aligned} \Upsilon &= \beta_{123} &= -\frac{1}{6} \beta \wedge \beta \wedge \beta \\ \Theta &= \Phi_1 \wedge \beta_1 + \Phi_2 \wedge \beta_2 + \Phi_3 \wedge \beta_3 &= -\operatorname{Re}[\Phi \wedge \beta] \end{aligned}$$

and the real 4-forms

$$\begin{aligned} -d\Upsilon &= \Phi_1 \wedge \beta_{23} + \Phi_2 \wedge \beta_{31} + \Phi_3 \wedge \beta_{12} = -\frac{1}{2}\text{Re}[\Phi \wedge \beta \wedge \beta] \\ \chi &= \Phi_1 \wedge \Phi_1 + \Phi_2 \wedge \Phi_2 + \Phi_3 \wedge \Phi_3 = -\text{Re}[\Phi \wedge \Phi]. \end{aligned}$$

Using these, we can build G_2 structures. In particular:

Proposition 2.1. *Let $\lambda = 2/\sqrt{5}$. The 3-form*

$$(2.13) \quad \varphi = 2\lambda\Theta + \lambda^3\Upsilon$$

defines an NP_2 structure on \tilde{P}^- with $\mu = 6/\sqrt{5}$. The associated Einstein metric is

$$(2.14) \quad h_7 = \frac{4}{5} \sum_{i=1}^3 \beta_i \otimes \beta_i + \pi^*g_4.$$

Proof. The calculation of $d\Theta$ is a little delicate:

$$\begin{aligned} d\Theta &= -\text{Re}[d\Phi \wedge \beta + \Phi \wedge d\beta] \\ &= -\text{Re}[-2\phi \wedge \Phi \wedge \beta] - \text{Re}[\Phi \wedge (-\beta \wedge \beta - 2\phi \wedge \beta - \Phi)] \\ &= 2d\Upsilon - \chi. \end{aligned}$$

Terms in $\phi \wedge \Phi$ and $\Phi \wedge \phi$ cancel out because wedging is commutative but imaginary quaternion multiplication is not.

Recall from (1.3) that $\Phi_i = \frac{1}{2}(\omega_{0i} - \omega_{jk})$ is a horizontal ASD 2-form, so $\chi = -\frac{3}{2}\omega_{0123}$, the volume form of (M, g_4) being ω_{0123} . It follows that (2.13) defines a G_2 structure whose associated metric has an orthonormal basis

$$(2.15) \quad \omega_0, \omega_1, \omega_2, \omega_3, \lambda\beta_1, \lambda\beta_2, \lambda\beta_3$$

so

$$*\varphi = 2\lambda^2 d\Upsilon - \frac{2}{3}\chi.$$

On the other hand,

$$d\varphi = \lambda(4 + \lambda^2)d\Upsilon - 2\lambda\chi$$

so the NP_2 condition is satisfied when

$$\mu = \frac{1}{2}\lambda^{-1}(4 + \lambda^2) = 3\lambda$$

and $\lambda^2 = 4/5$. The sign of λ and μ is merely a matter of orientation, so we choose them to be positive for this proposition. The form of h_7 follows from (2.15). \square

The complete metric g_Φ with holonomy $Spin(7)$ appears on [12, p848]. If we take the asymptotic limit and set $r = 1$, we obtain $\frac{1}{5}h_7$. It is well known that (2.14) is the so-called squashed Einstein metric that exists on the S^3 (or \mathbb{RP}^3) bundle over M^4 . The more standard Einstein metric (the round metric in the case of $\pi: S^7 \rightarrow S^4$) has the same form, but without the factor $1/5$.

The proof above is modelled on [20, Prop. 2.5] from the theory of 3-Sasakian 7-manifolds. Indeed, we shall explain next that the 3-form (2.13) coincides with one that can be defined using globally-defined 1-forms η_1, η_2, η_3 dual to the Lie algebra of $SU(2)$ acting on

the fibre of a regular 3-Sasakian 7-manifold. This is accomplished by means of a gauge transformation that converts the geometry of a vector bundle to that of a principal bundle.

Modify (1.8) by conjugation and (partly for convention) a sign change:

$$(2.16) \quad \eta = -a\beta\bar{a} = -\alpha\bar{a} = a\bar{\alpha}$$

where a is a unit quaternion. Retaining the condition $R = 1$, we can write

$$\eta = i\eta_1 + j\eta_2 + k\eta_3.$$

In particular,

$$\eta_1 = -a_0\alpha_1 + a_1\alpha_0 - a_2\alpha_3 + a_3\alpha_2.$$

Observe that, as a direct consequence of the conjugation, η is *invariant* under the action $a \rightsquigarrow aq$ of $SU(2)^-$. Moreover, η , like β , annihilates the horizontal distribution on \tilde{P}^- . It follows that each real component η_i is globally defined on \tilde{P}^- . Moreover,

$$\alpha \wedge \bar{\alpha} = \alpha\bar{a} \wedge a\bar{\alpha} = -\eta \wedge \eta$$

in analogy to (1.10), but with a fibrewise self-dual property. Converting from β to η furnishes a more succinct version of Lemma 1.1 on \tilde{P}^- , namely

Lemma 2.2. $d\eta + \eta \wedge \eta = a\Phi\bar{a}$.

Proof. Applying (1.7) and (1.9),

$$\begin{aligned} d\eta &= -d(\alpha\bar{a}) = -d\alpha\bar{a} + \alpha \wedge d\bar{a} \\ &= (\alpha \wedge \phi + a\Phi)\bar{a} + \alpha \wedge (\bar{a} - \phi\bar{a}) \\ &= \alpha \wedge \bar{a} + a\Phi\bar{a}. \end{aligned}$$

As has to be, the individual terms in the last line are unchanged when a, α, Φ are replaced by $aq, \alpha q, \bar{q}\Phi q$, respectively. \square

It is instructive to compare Lemmas 1.1 and 2.2. In analogy to (1.2), set

$$(2.17) \quad H = a\Phi\bar{a} = i(d\eta_1 + 2\eta_{23}) + i(d\eta_2 + 2\eta_{31}) + i(d\eta_3 + 2\eta_{12}).$$

Then H is ‘3-horizontal’, i.e. its individual real components are pointwise the pullbacks of forms on the 4-manifold M . However, they are not invariant by $SU(2)^-$, only their quaternionic combination enjoys this property, the price to pay for avoiding analogues of the terms $\phi \wedge \beta$ in Lemma 1.1. If we replace the symbol H_i by $\bar{\phi}^i$ (the overline is just notation), this coincides with [20, p 50].

Remark 2.3. Conjugation of an imaginary quaternion by the unit quaternion (1.6) describes the double covering $SU(2)^- \rightarrow SO(3)^-$ by mapping (1.6) to the orthogonal matrix

$$(2.18) \quad (P_{i,j}) = \begin{pmatrix} a_0^2 + a_1^2 - a_2^2 - a_3^2 & 2(-a_0a_3 + a_1a_2) & 2(a_0a_2 + a_3a_1) \\ 2(a_0a_3 + a_1a_2) & a_0^2 - a_1^2 + a_2^2 - a_3^2 & 2(-a_0a_1 + a_2a_3) \\ 2(-a_0a_2 + a_3a_1) & 2(a_0a_1 + a_2a_3) & a_0^2 - a_1^2 - a_2^2 + a_3^2 \end{pmatrix}$$

that features in many articles on the quaternions, such as [1, Prop. 23.12]. The nine entries of (2.18) are tracefree quadratics, elements of $S_0^2(\mathbb{R}^4) \cong \Lambda_+^2 \otimes \Lambda_-^2$. Indeed, $P_{i,j}$ equals minus the contraction of standard basis elements of Λ^+ and Λ^- . The triple $(P_{1,i})$ is the hyperkähler moment map $\mathbb{R}^4 \rightarrow \mathfrak{so}(3)$ for the action of $U(1)$ on \mathbb{R}^4 tangent to I , and is familiar from the Gibbons-Hawking construction. In components,

$$(2.19) \quad d\eta_1 + 2\eta_{23} = 2 \sum_p P_{1,p} \Omega_p.$$

It is easy to compute dH using (2.17) twice. In particular,

$$dH_1 = 2(d\eta_2 \wedge \eta_3 - \eta_2 \wedge d\eta_3) = -2(\eta_2 \wedge H_3 - \eta_3 \wedge H_2)$$

which (as it has to) matches (1.4):

$$d\Phi_1 = -2(\phi_2 \wedge \Phi_3 - \phi_3 \wedge \Phi_2)$$

All the factors of the right-hand side of the last line are 3-horizontal.

Proposition 2.4. *Let $\lambda = 2/\sqrt{5}$. Define*

$$\varphi' = 2\lambda \sum_{i=1}^3 H_i \wedge \eta_i + \lambda^3 \eta_{123}.$$

Then φ' coincides with minus the 3-form (2.13), and therefore gives an equivalent description of the squashed NP_2 structure on \tilde{P}^- .

Proof. The definitions of η and H , and some telescoping, imply that

$$\varphi' = -a\varphi\bar{a} = -\varphi$$

since both φ and φ' are real differential forms. \square

The definition of φ' in Proposition 2.4 is modelled on that of φ , but is perhaps more familiar. However, there is a significant difference between Lemmas 1.1 and 2.2, either of which can be used to compute $d\varphi' = -d\varphi$. The necessity to change the sign of λ arises from a minus sign that we inserted in the definition (2.16) of η to preserve the combination $d\eta + \eta^2$ to match $d\beta + \beta^2$.

As explained in [20, p 51], in the 3-Sasakian context, one can define a hyperkähler triple of exact 2-forms

$$(2.20) \quad \varpi_i = d(2R\eta_i) = 4\eta_{0i} - 4R\eta_{jk} + 2RH_i$$

on the cone over \tilde{P}^- . Pointwise, each $2H_i = \omega_{0i} - \omega_{jk}$ is a horizontal ASD 2-form that matches the other terms, see (2.15) with $\lambda = 2$. The existence of such a hyperkähler structure is well known in the setting of twistor space and quaternionic geometry [34], though we are here more interested in induced G_2 structures on the 3-sphere bundle \tilde{P}^- .

The metric associated to (2.20) is the cone over X , and this is one way of defining a 3-Sasakian structure [7, 10]. Since the holonomy of the cone lies in $Sp(2)$, it also lies in $Spin(7)$ and \tilde{P}^- must have an NP_2 structure. However, such a subgroup $Spin(7)$ is

conjugate to the stabilizer of a 4-form $-\varpi_1^2 + \varpi_2^2 + \varpi_3^2$, so one needs to break the $SO(3)$ symmetry of the hyperkähler triple, whereupon there emerges a family of NP_2 structures.

We can replace each β_i by η_i in (2.14) without affecting the sum. More general families of metrics of the form

$$dt^2 + \sum_{k=1}^3 A_k(t)^2 \eta_k \otimes \eta_k + B(t)^2 g_4,$$

with holonomy $Spin(7)$ were constructed by Bazaikin by deforming 3-Sasakian structures [8]. This has guided our own presentation of (or rather, hope to generalize) the homogeneous NP_2 structures, in which trigonometric functions of t appear on subspaces $\langle \beta_k, \omega_k \rangle$ for $k = 1, 2, 3$, see Corollaries 3.4 and 4.4.

3. THE COHOMOGENEITY-ONE ACTION OF $SO(4)$ ON S^7

The group $Sp(n)$ acts transitively on the sphere S^{4n-1} in quaternionic space \mathbb{H}^n and, in particular,

$$(3.21) \quad S^7 \cong \frac{Sp(2)}{Sp(1)} \cong \frac{SO(5)}{SU(2)}.$$

This is the homogeneous model of 3-Sasakian geometry in dimension 7, with its triple of invariant 1-forms. From the point of view of Section 2, it is more appropriate to start from the description

$$(3.22) \quad S^7 \cong \frac{Sp(2) \times Sp(1)}{Sp(1) \times Sp(1)} = \frac{Sp(2)Sp(1)}{Sp(1)Sp(1)},$$

which features in the classification [18] of homogeneous NP_2 spaces. The second factor of the isotropy group $Sp(1)Sp(1) \cong SO(4)$ acts diagonally, and juxtaposition indicates a quotient by ± 1 across both factors. In quaternion-Kähler jargon, S^7 is here regarded as the unit sphere in the $Sp(2)Sp(1)$ module \mathfrak{m} , with complexification $\mathfrak{m}_\mathbb{C} = E \otimes H$, where $Sp(2)$ acts on $E \cong \mathbb{C}^4$ and $Sp(1)$ acts on $H \cong \mathbb{C}^2$.

Let $\Sigma^+ \oplus \Sigma^-$ denote the spin representation of $Spin(4) \cong Sp(1) \times Sp(1)$, so that $Sp(1)^+$ acts on $\Sigma^+ \cong \mathbb{C}^2$ and (as before) $Sp(1)^-$ acts on $\Sigma^- \cong H$. The isotopy subgroup in (3.22) then decomposes $E = \Sigma^+ \oplus \Sigma^-$, so that

$$\mathfrak{m}_\mathbb{C} \cong (\Sigma^+ \oplus \Sigma^-) \otimes \Sigma^-.$$

Since $S^2\Sigma^-$ is the complexification of the space Λ^- of ASD 2-forms on \mathbb{R}^4 , the isotropy subgroup $SO(4)$ acts on the real tangent space to S^7 as $\mathbb{R}^4 \oplus \Lambda^-$. It is well known that this action factors through G_2 . Moreover there are independent $SO(4)$ invariant 3-forms φ_1, φ_2 on S^7 , a linear combination φ of which must satisfy (2.12) with $\mu = 6/\sqrt{5}$, in accordance with Proposition 2.1.

To obtain an action by $SO(4) \cong Sp(1)^+Sp(1)^-$ of cohomogeneity one on S^7 , we move from isotropy subgroups to isometry groups and set $E = S^3\Sigma^+$, so that

$$\mathfrak{m}_\mathbb{C} \cong S^3\Sigma^+ \otimes \Sigma^-$$

is the isotropy representation of the Wolf space $G_2/SO(4)$, as in [35]. Then

$$(3.23) \quad \begin{aligned} S^2\mathfrak{m}_c &\cong (S^2E \otimes S^2H) \oplus (\Lambda^2E \otimes \Lambda^2H) \\ &\cong (S^6\Sigma^+ \otimes S^2\Sigma^-) \oplus (S^2\Sigma^+ \otimes S^2\Sigma^-) \oplus S^4\Sigma^+ \oplus \mathbb{C}. \end{aligned}$$

The penultimate summand is the complexification of the space $S_0^2(\mathbb{R}^3) \cong \mathbb{R}^5$ of 3×3 tracefree symmetric matrices. Projection to this module of $v \otimes v$ (where v is a non-zero vector in $\mathfrak{m} = \mathbb{R}^8$) defines the $Sp(2)Sp(1)$ -equivariant fibration $S^7 \rightarrow S^4$.

Remark 3.1. The 9-dimensional summand of (3.23) is isomorphic to $S_0^2(\mathbb{R}^4)$ (cf. Remark 2.3), and is the model for a geometrical structures on 9-manifolds investigated in [16]. In the present context, it gives rise to an embedding of $\mathbb{R}\mathbb{P}^7$ as a hypersurface of S^8 .

Let $A \in S^4 \subset S_0^2(\mathbb{R}^3)$ be a 3×3 matrix with distinct eigenvalues. Then the orbit of $SO(3) = SU(2)^+/\mathbb{Z}_2$ containing A has dimension 3 and principal isotropy group \mathbb{Z}_2^2 . There are two singular orbits, each copies of $\mathbb{R}\mathbb{P}^2$, corresponding to repeated eigenvalues, which can be negative or positive. To study the round metric g_4 on S^4 relative to the action by $SO(3)$ of cohomogeneity one, set $\vartheta_k = 2\pi(k-1)/3$, and

$$(3.24) \quad \begin{aligned} c_k &= c_k(t) = \cos(t + \vartheta_k) \\ s_k &= s_k(t) = \sin(t + \vartheta_k) \end{aligned}$$

for $k = 1, 2, 3$. These trigonometric quantities satisfy

$$(3.25) \quad c_1 + c_2 + c_3 = 0 = s_1 + s_2 + s_3$$

as well as the identity

$$(3.26) \quad c_j c_k + s_j s_k = -\frac{1}{2},$$

for $(ijk) = (123)$.

Now choose left-invariant 1-forms on $SO(3)$ satisfying

$$(3.27) \quad de_1 = -2e_{23}, \quad de_2 = -2e_{31}, \quad de_3 = -2e_{12}.$$

This convention is chosen to match expressions such as (1.11). Combined with the constraint on κ , it leads us to adopt an orthonormal basis of 1-forms on S^4 formed from dt and $4s_k e^k$ ($k = 1, 2, 3$), so that

$$(3.28) \quad g_4 = dt^2 + 16 \sum_{k=1}^3 s_k(t)^2 e_k \otimes e_k$$

away from the singular orbits. The associated basis of ASD 2-forms consists of

$$\begin{aligned} \omega_1 &= 4s_1 dt \wedge e_1 - 16s_2 s_3 e_{23} \\ \omega_2 &= 4s_2 dt \wedge e_2 - 16s_3 s_1 e_{31} \\ \omega_3 &= 4s_3 dt \wedge e_3 - 16s_1 s_2 e_{12}. \end{aligned}$$

Lemma 3.2. *With this choice, the connection 1-form ϕ for $\Lambda_-^2 T^*S^4$ is given by*

$$(3.29) \quad -\phi = i(2c_2 + 1)e_1 + j(2c_2 + 1)e_2 + k(2c_3 + 1)e_3,$$

and $\kappa = 1$.

Proof. It suffices to verify (1.4) and (1.5), since these equations have a unique solution by the theories of Cartan or Levi-Civita.

First observe that

$$d\omega_1 = 8(s_1 - 2(c_2s_3 + s_2c_3))dt \wedge e_{23},$$

whereas

$$\begin{aligned} -2(\phi_2 \wedge \omega_3 - \phi_3 \wedge \omega_2) &= 8(ac_2 - 1)s_3dt \wedge e_{23} + 8(ac_3 - 1)s_2dt \wedge e_{23} \\ &= 8(s_1 - 2(c_2s_3 + c_3s_2))dt \wedge e_{23}. \end{aligned}$$

This verifies (1.4) after cyclic permutations.

As regards (1.5), we have

$$\begin{aligned} d\phi_1 + 2\phi_{23} &= 2s_1dt \wedge e_1 + 2(2c_1 + 1)e_{23} + 2(2c_2 + 1)(2c_3 + 1)e_{23} \\ &= 2s_1dt \wedge e_1 + (4 + 8c_2c_3)e_{23} \\ &= 2s_1dt \wedge e_1 - 8s_2s_3e_{23} \\ &= \frac{1}{2}\omega_1, \end{aligned}$$

which is consistent with (1.5) and $\kappa = 1$. □

Next, consider the locally-defined 1-form

$$(3.30) \quad f = \bar{a} da = \bar{a}(\alpha + a\phi) = \beta + \phi$$

on Σ^- over a self-dual Einstein 4-manifold. It is vector valued on the 3-sphere bundle, and transforms as $f \rightsquigarrow \bar{q}fq$. It arises from the Maurer-Cartan form of the fibre, and

$$df = d(\bar{a} da) = d\bar{a} \wedge da = (d\bar{a} a) \wedge (\bar{a} da) = -f \wedge f$$

so in components,

$$(3.31) \quad df_1 = -2f_{23}, \quad df_2 = -2f_{31}, \quad f_3 = -2f_{12}.$$

Regard $S^7 = \tilde{P}^-S^4$ as the S^3 bundle over S^4 . Consider the 1-forms (e_k) and (f_k) satisfying (3.31) and (3.27). Lemma 3.2 yields

Theorem 3.3. *For $k = 1, 2, 3$, we have $\beta_k = (2c_k + 1)e_k + f_k$.*

These equations relate the 1-forms from Lemma 1.1 to the action of $SO(4)$, which is a significant step in understanding the cohomogeneity-one setup. The trigonometric terms arise from the action of the Weyl group in the cohomogeneity-one theory [23], and is suggestive of a discrete Fourier transform. One cannot usefully substitute (β_k) by the 3-Sasakian 1-forms (η_k) in the theorem, since this would yield $a\phi\bar{a}$ in (3.30).

An orthonormal basis of 1-forms for the round metric g_7 on S^7 is given by

$$dt, 2\beta_1, 2\beta_2, 2\beta_3, 4s_1e_1, 4s_2e_2, 4s_3e_3.$$

Set $\tau = t/2$. Then

$$\begin{aligned} g_7 &= dt^2 + 4 \sum_{k=1}^3 (\beta_k)^2 + \pi^* g_4 \\ &= dt^2 + 4 \sum_{k=1}^3 (\beta_k)^2 + 16 \sum_{k=1}^3 (s_k e_k)^2 \\ &= dt^2 + 4 \sum_{k=1}^3 [f_k^2 + \gamma_k (e_k f_k + f_k e_k) + \delta_k e_k^2], \end{aligned}$$

where

$$\gamma_k = 2c_k + 1 = -1 + 4 \cos^2 \tau, \quad \delta_k = (2c_k + 1)^2 + (2s_k)^2 = 1 + 8 \cos^2 \tau.$$

It follows that the round metric can be expressed as dt^2 plus a 6×6 matrix with 2×2 symmetric blocks, of which the first is

$$(3.32) \quad 4 \begin{pmatrix} 1 & -1 + 4 \cos^2 \tau \\ -1 + 4 \cos^2 \tau & 1 + 8 \cos^2 \tau \end{pmatrix}.$$

This can be reconciled with [35, p7] up to scale, in which the period of t in (3.24) has been halved to $\pi/6$. A change of off-diagonal sign results from our ASD convention.

We now turn to the squashed metric (2.14) on S^7 and its NP_2 structure. Consider the orthonormal basis 1-forms (X_i) defined by $X_{2k-1} = \lambda\beta_k$, $X_{2k} = 4s_k e_k$, and $X_7 = -dt$. Theorem 3.3 implies

Corollary 3.4. *Let $\kappa = 1$ and $\lambda = 2/\sqrt{5}$. Then*

$$\begin{pmatrix} X_{2k-1} \\ X_{2k} \end{pmatrix} = \begin{pmatrix} \lambda & \lambda(2c_k + 1) \\ 0 & 4s_k \end{pmatrix} \begin{pmatrix} f_k \\ e_k \end{pmatrix}$$

for $k = 1, 2, 3$.

The behaviour of both g_7 and h_7 at the two singular orbits of $SO(4)$ can be investigated by setting $t = 0$ and $t = \pi/3$, so that $X_2 = 0$ and $X_4 = 0$, respectively.

Referring to (2.13), we now see that $\lambda^3 \Upsilon = X_{135}$ and

$$2\lambda\Theta = (X_{72} - X_{46}) \wedge X_1 + (X_{74} - X_{62}) \wedge X_3 + (X_{76} - X_{24}) \wedge X_5.$$

Proposition 2.1 therefore yields

Theorem 3.5. *The natural NP_2 structure on S^7 has*

$$\varphi = dt \wedge (X_{12} + X_{34} + X_{56}) + X_{135} - X_{146} - X_{236} - X_{245}.$$

The theorem asserts that in the stated basis, φ assumes a standard form

$$\begin{aligned}\varphi &= dt \wedge \xi + \operatorname{Re} \Xi \\ * \varphi &= -dt \wedge \operatorname{Im} \Xi + \frac{1}{2} \xi^2\end{aligned}$$

relative to the $SO(4)$ orbits, where

$$(3.33) \quad \begin{aligned}\xi &= X_{12} + X_{34} + X_{56} \\ \Xi &= (X_1 + iX_2) \wedge (X_3 + iX_4) \wedge (X_5 + iX_6).\end{aligned}$$

To sum up: we have described the ‘squashed’ 3-form φ from Propositions 2.1 and 2.4 in terms of $SO(4)$ invariant 1-forms, via Corollary 3.4. This provides a model for the construction of NP_2 structures in the cohomogeneity-one situation of an interval times (a finite quotient of) $SO(4)$. In the next section, we shall discover a strikingly similar description on the principal part of the isotropy irreducible space $SO(5)/SO(3)$.

4. THE COHOMOGENEITY-ONE ACTION OF $SO(4)$ ON B^7

Berger discovered two homogeneous spaces of positive sectional curvature, of respective dimensions 7 and 13, namely

$$B^7 = \frac{SO(5)}{SO(3)}, \quad B^{13} = \frac{SU(5)}{Sp(2)U(1)},$$

which he studied in [9]. The former is the subject of this section; it is the coset space that parametrizes principal subalgebras $\mathfrak{so}(3)$ of $\mathfrak{so}(5)$, equivalently Veronese surfaces in S^4 [6].

The subgroup $SO(3)$ of $SO(5)$ acts irreducibly on \mathbb{R}^5 , which can be identified with the space $S_0^2(\mathbb{R}^3)$, equivalently the space of traceless symmetric 3×3 matrices. The isotropy subgroup of B^7 is a principal 3-dimensional subgroup that acts irreducibly on \mathbb{R}^7 . Indeed,

$$(4.34) \quad \mathfrak{so}(5) = \mathfrak{so}(3) \oplus \mathfrak{m},$$

where \mathfrak{m} can be identified with the space $S_0^3(\mathbb{R}^3)$ of primitive cubic forms on \mathbb{R}^3 . After complexification,

$$(4.35) \quad \mathfrak{so}(3)_c \cong S^2(\mathbb{C}^2), \quad (\mathbb{R}^5)_c \cong S^4(\mathbb{C}^2), \quad \mathfrak{m}_c \cong S^6(\mathbb{C}^2)$$

where \mathbb{C}^2 denotes the faithful representation of the double-cover $SU(2)$ of $SO(3)$.

It is known that the action of $SO(3)$ on \mathfrak{m} factors through G_2 . Indeed, there is an $SU(2)$ equivariant isomorphism

$$(4.36) \quad \Lambda^3(S^6(\mathbb{C}^2)) \cong S^{12} \oplus S^8 \oplus S^6 \oplus S^4 \oplus S^0$$

(with obvious abbreviation), and the trivial 1-dimensional submodule $S^0 \cong \mathbb{C}$ is spanned by a G_2 -admissible 3-form φ . The latter also arises from the projection to \mathfrak{m} of the Lie bracket $[\mathfrak{m}, \mathfrak{m}]$ within (4.34). The fact that $[\mathfrak{m}, \mathfrak{m}]$ does not lie in $\mathfrak{so}(3)$ implies that B^7 is not a symmetric space, and its holonomy group equals $SO(7)$.

Now, φ determines an $SO(5)$ -invariant 3-form on B^7 (which we denote with the same symbol), and an invariant 4-form again unique up to scaling (which we can denote $*\varphi$).

Since there is no non-zero invariant 2-form, $d\varphi$ must be a nonzero multiple of $*\varphi$, which is the NP_2 condition (2.12). Hence, the well-known

Proposition 4.1. B^7 has an $SO(5)$ -invariant NP_2 structure.

It follows that the Riemannian cone $B^7 \times \mathbb{R}^+$ has holonomy $Spin(7)$. It is not known whether this singularity can be resolved in the manner of [8], but it motivates the study of metrics with holonomy $Spin(7)$ admitting a codimension-two action by $SO(4)$.

Remark 4.2. The space B^{13} is not isotropy irreducible. There is a fibration $B^{13} \rightarrow \mathbb{C}P^4$, with fibre $SU(4)/Sp(2) \cong S^5$. Its complexified isotropy representation is $(V \oplus \bar{V}) \oplus \Lambda_0^2 E$, where $V = E \otimes L \cong \mathbb{C}^4$ is a standard representation of $Sp(2) \times U(1)$. The space B^{13} has a 3-dimensional space of invariant 4-forms, which are relevant to the study of ‘strongly positive curvature’ [2].

The 3-form φ on B^7 was computed explicitly in [21] and in [6], but in different ways. The calculations below are equivalent, but our approach emphasizes the underlying 3-fold symmetry.

Denote by S_{ij} the symmetric 3×3 matrix with (i, j) and (j, i) entry equal to 1 and 0 elsewhere. We then adopt the following basis of traceless symmetric matrices:

$$(4.37) \quad \begin{aligned} \bar{S}_1 &= \frac{1}{\sqrt{6}} \text{diag}(1, 1, -2), & \bar{S}_2 &= \frac{1}{\sqrt{2}} \text{diag}(1, -1, 0), \\ \bar{S}_3 &= S_{12}, & \bar{S}_4 &= S_{13}, & \bar{S}_5 &= S_{23}. \end{aligned}$$

We represent elements of the Lie algebra (4.35) by skew-symmetric 5×5 matrices. Let E_{ij} denote the matrix with 1 at (i, j) , -1 at (j, i) , and 0 elsewhere. For the subalgebra $\mathfrak{so}(3)$, we may choose a basis $\{\varepsilon_1, \varepsilon_2, \varepsilon_3\}$, where

$$\begin{aligned} \varepsilon_1 &= \frac{1}{\sqrt{5}}(2E_{23} + E_{45}) \\ \varepsilon_2 &= \frac{1}{\sqrt{5}}(E_{24} + E_{35} + \sqrt{3}E_{14}) \\ \varepsilon_3 &= \frac{1}{\sqrt{5}}(-E_{25} + E_{34} + \sqrt{3}E_{15}). \end{aligned}$$

One can check that $\text{tr}(\varepsilon_i \varepsilon_j) = -2\delta_{ij}$, and that $[\varepsilon_i \varepsilon_j] = -\lambda \varepsilon_k$, where $(ijk) = (123)$.

Now set

$$R(t) = \begin{pmatrix} \cos t & \sin t & 0 & 0 & 0 \\ -\sin t & \cos t & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix}.$$

We have chosen the lower block to be a 3×3 permutation matrix (rather than the identity) because $\rho = R(\frac{2\pi}{3})$ has special significance; it is derived from the effect on (4.37) of permuting the basis of \mathbb{R}^3 . It therefore lies in the intersection of the principal $SO(3)$ with the subgroup $SO(2) \times SO(3)$ of $SO(5)$. Conjugation by this element of order 3 cycles the ε_i , so that $\rho^{-1}\varepsilon_1\rho = \varepsilon_2$, etc.

Let $a = \cos(2\pi/3)$ and $b = \sin(2\pi/3)$. A basis for \mathfrak{m} is given by $\{\gamma_1, \dots, \gamma_7\}$ where

$$(4.38) \quad \begin{aligned} \gamma_1 &= \frac{1}{\sqrt{5}}(E_{23} - 2E_{45}) \\ \gamma_2 &= -E_{13} \\ \gamma_3 &= \frac{1}{\sqrt{5}}(bE_{15} + aE_{25} - 2E_{34}) \\ \gamma_4 &= -aE_{15} + bE_{25} \\ \gamma_5 &= \frac{1}{\sqrt{5}}(-bE_{14} + aE_{24} + 2E_{35}) \\ \gamma_6 &= -aE_{14} - bE_{24} \\ \gamma_7 &= E_{12}. \end{aligned}$$

One can check that $\text{tr}(\gamma_i \gamma_j) = -2\delta_{ij}$, and that $\text{tr}(\gamma_i \varepsilon_j) = 0$ for all i, j . Observe that ρ cycles the pairs $(\gamma_{2k-1}, \gamma_{2k})$ for $k = 1, 2, 3$, in the sense that $\rho^{-1}(\gamma_1, \gamma_2)\rho = (\gamma_3, \gamma_4)$ etc. In fact, it was only necessary to choose the orthogonal basis γ_1, γ_2 orthogonal to $\mathfrak{so}(3)$; the rest were generated by ρ . On the other hand, γ_7 commutes with ρ .

Having made these choices, the $SO(5)$ -invariant 3-form φ must be proportional to

$$\frac{1}{10} \sum_{i < j < k} \text{tr}([\gamma_i, \gamma_j] \gamma_k) \gamma_i^* \wedge \gamma_j^* \wedge \gamma_k^*,$$

where (γ_i^*) is the dual basis of invariant 1-forms. With the given normalization, we discover

Proposition 4.3. $\varphi = \gamma_{127}^* + \gamma_{135}^* - \gamma_{146}^* - \gamma_{236}^* - \gamma_{245}^* + \gamma_{347}^* + \gamma_{567}^*$.

Thus is in canonical form for a G_2 structure. Since B^7 has a unique $SO(5)$ -invariant Riemannian metric g_B (up to an overall constant), it is a corollary of the proposition that (γ_i^*) is an orthonormal basis for g_B .

There is a cohomogeneity-one action of $SO(4)$ on B^7 [31]. The principal orbits are $SO(4)/\mathbb{Z}_2^2$, and the singular orbits are both isomorphic to $SO(4)/O(2)$. To analyse this, we start with bases $(e_k), (f_k)$, whose exterior derivatives are normalized as in (3.27), (3.31). We then define positive and negative combinations $p_k = \frac{1}{2}(f_k + e_k)$ and $n_k = \frac{1}{2}(-f_k + e_k)$, so that

$$(4.39) \quad \begin{pmatrix} p_k \\ n_k \end{pmatrix} = L \begin{pmatrix} f_k \\ e_k \end{pmatrix}, \quad L = \frac{1}{2} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$$

for $k = 1, 2, 3$. This switch of bases will play a significant role in simplifying subsequent expressions, as we are about to see. A coefficient of $1/2$ is chosen in the light of the differential relations (4.43) below.

The Maurer-Cartan form of the subgroup $SO(4)$ in $SO(5)$ can be expressed as

$$(4.40) \quad A = 2 \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & n_3 & n_2 & n_1 \\ 0 & -n_3 & 0 & -p_1 & p_2 \\ 0 & -n_2 & p_1 & 0 & -p_3 \\ 0 & -n_1 & -p_2 & p_3 & 0 \end{pmatrix}$$

so that e_1 corresponds to the combination $-E_{25} - E_{34}$, and f_1 to $E_{25} - E_{34}$, etc. These choices match the conventions (3.27),(3.31) and (the subsequent) (4.43).

The pullback of A on $[0, \frac{\pi}{3}] \times SO(4)$ is given by

$$R(t)^{-1}AR(t) + \gamma_7^* \wedge dt = \begin{pmatrix} 0 & dt & -sn_1 & -sn_3 & -sn_2 \\ -dt & 0 & cn_1 & cn_3 & cn_2 \\ sn_1 & -cn_1 & 0 & -p_2 & p_3 \\ sn_2 & -cn_3 & p_2 & 0 & -p_1 \\ sn_3 & -cn_2 & -p_3 & p_1 & 0 \end{pmatrix}$$

where $c = \cos t$ and $s = \sin t$ as temporary abbreviations.

Let Y_i denote the pullback of $2\gamma_i^*$. These 1-forms satisfy

$$\begin{aligned} Y_1 &= \lambda(cn_1 + 2p_1) \\ Y_2 &= 2sn_1 \\ Y_3 &= \lambda((ca - sb)n_2 + 2p_2) \\ Y_4 &= 2(sa + cb)n_2 \\ Y_5 &= \lambda((ca + sb)n_3 + 2p_3) \\ Y_6 &= 2(sa - cb)n_3 \\ Y_7 &= 2dt \end{aligned}$$

where $\lambda = 2/\sqrt{5}$ and (as before) $a = \cos(2\pi/3)$ and $b = \sin(2\pi/3)$.

These equations manifest the \mathbb{Z}_3 symmetry arising from the action of the Weyl group for the action of $SO(4)$ [23], and can be restated as

Corollary 4.4. *Let $\kappa = 1$ and $\lambda = 2/\sqrt{5}$. Then*

$$(4.41) \quad \begin{pmatrix} Y_{2k-1} \\ Y_{2k} \end{pmatrix} = \begin{pmatrix} 2\lambda & \lambda c_k \\ 0 & 2s_k \end{pmatrix} \begin{pmatrix} p_k \\ n_k \end{pmatrix}$$

for $k = 1, 2, 3$.

The 3-form from Proposition 4.3 defining the NP_2 structure on B^7 can be written

$$\varphi = \xi \wedge Y_7 + \text{Re } \Xi,$$

where Ξ is defined by (3.33) with Y_i in place of X_i . If we denote the 2×2 matrix in Corollary 4.4 by $M_k = M_k(t)$, the analogue of (3.32) for the term of g_B involving the plane $\langle f_1, e_1 \rangle$ equals

$$(4.42) \quad L^\top M_1^\top M_1 L = \frac{1}{5} \begin{pmatrix} 5 + 4 \sin^2 t + 4 \cos t & 1 - 4 \cos^2 t \\ 1 - 4 \cos^2 t & 5 + 4 \sin^2 t - 4 \cos t \end{pmatrix}.$$

The change of basis (4.39) reveals a striking similarity between Corollary 3.4 and Corollary 4.4, which we explore next.

The 1-forms in (4.39) satisfy

$$(4.43) \quad \begin{cases} dp_k &= -2(p_{ij} + n_{ij}) \\ dn_k &= -2(p_i n_j - p_j n_i) \end{cases}$$

for $(ijk) = (123)$. If ∇ is a connection on $T^*SO(4)$ that satisfies

$$\nabla n_k = p_i \otimes n_j - p_j \otimes n_i$$

then we may regard the p_i as connection 1-forms on \mathcal{D} , whose curvature 2-forms are

$$dp_k + 2p_{ij} = -2n_{ij}.$$

The second equation of (4.43) implies that $\mathcal{D} = \langle Y_2, Y_4, Y_6 \rangle$ is a differential ideal in $SO(4)$. Regarding \mathcal{D} as a subspace of $\mathfrak{so}(4)^*$, its annihilator is the diagonal subalgebra

$$\mathfrak{so}(3)_d \subset \mathfrak{so}(3)_+ + \mathfrak{so}(3)_- = \mathfrak{so}(4)$$

represented by the bottom 3×3 block in (4.40). This subalgebra corresponds to the choice of isometry $\sigma: \langle e_1, e_2, e_3 \rangle \rightarrow \langle f_1, f_2, f_3 \rangle$ defined by our indexing, or (up to sign) the vector in \mathbb{R}^4 that is fixed by our choice of geodesic.

Inside each generic $SO(4)$ orbit, the integral submanifolds of \mathcal{D} consist of \mathbb{Z}_2^2 quotients of conjugacy classes of a diagonal $SO(3)_d$, and are parametrized by $SO(4)/SO(3)_d \cong S^3$. Moreover,

$$\text{Im } \Xi = -Y_{246} + Y_{235} + Y_{145} + Y_{136}$$

vanishes on these submanifolds, and they are associative (see (3.33)). They play a role analogous to that of the orbits of $SU(2)^-$ acting on S^7 , insofar as the homogeneous NP_2 metrics are concerned. Note that the complete theory of associative submanifolds of B^7 is very rich because of the many different orbit types of a principal $SO(3)$ subgroup of G_2 acting on the space $G_2/SO(4)$ of associative subspaces of \mathbb{R}^7 [6].

Remark 4.5. In the wider homogeneous context, conjugacy classes of 3-dimensional subalgebras of $\mathfrak{so}(5)$ (namely $\mathfrak{su}(2)^-$, $\mathfrak{so}(3)_d$ and principal $\mathfrak{so}(3)$) correspond to critical submanifolds (namely S^4 , $\mathbb{G}r_3(\mathfrak{so}(5))$ and B^7) for the natural Morse-Bott functional on the 21-dimensional Grassmannian $\mathbb{G}r_3(\mathfrak{so}(5))$. Two associated unstable manifolds of dimensions 8 and 12 then admit incomplete quaternion-Kähler metrics by the theory in [27], and it is tempting to tie this in with the results of this article.

The behaviour of the 1-forms Y_i at $t = 0$ and $\pi/3$ gives some information about the singular locus and $SO(4)$ action. For example, Y_2 vanishes at $t = 0$, so the stabilizer $O(2)$ is tangent to the annihilator of $\langle Y_1, Y_3, Y_4, Y_5, Y_6 \rangle$. At our distinguished point, $SO(2)$ rotates $\langle E_4, E_5 \rangle$ by θ and $\langle E_2, E_3 \rangle$ by 2θ . Its induced action on (0.1) fixes $E_{23} \pm E_{45}$, but rotates $\langle E_{24} + E_{53}, E_{25} + E_{34} \rangle$ by 3θ , whereas $\langle E_{24} - E_{53}, E_{25} - E_{34} \rangle$ by $-\theta$. The singular orbit $SO(4)/O(2)$ is therefore covered by the circle bundle in $\mathcal{O}(3, 1) \rightarrow \mathbb{C}P^1 \times \mathbb{C}P^1$.

The subgroup $SU(2)^+$ of $SO(4)$ (defined by our choice of oriented basis) has stabilizer \mathbb{Z}_3 at $t = 0$, but $SU(2)^-$ acts freely. The opposite is true at $t = \pi/3$, giving rise to a group diagram with weights $(3, 1)$ at $t = 0$, and $(1, 3)$ at $t = \pi/3$, relative to $SO(4) = SO(3)^+SO(3)^-$. It follows that $B^7/SU(2)^+$ is smooth for $t \in (0, \pi/3]$, but not at $t = 0$.

Since $SO(4)/SU(2)^+ \cong SO(3)$, the commutator $SU(2)^-$ acts on the quotient with generic stabilizer $V = \mathbb{Z}_2^2$. This observation leads to the $SO(4)$ -equivariant diffeomorphism $B^7 \cong \tilde{P}^+\mathcal{O}_5$ established by Grove, Wilking, and Ziller [23], where \mathcal{O}_k is Hitchin's orbifold with a \mathbb{Z}_{k-2} singularity transverse to $\mathbb{R}\mathbb{P}^2$ (we use $k \geq 3$ as in [24]). We outline this setup next because of its implications for NP_2 structures with an $SO(4)$ symmetry, working backwards from \mathcal{O}_k . Recall our convention that a self-dual conformal structure has vanishing Weyl component W_- (as for $\mathbb{C}\mathbb{P}^2$ whose Kähler 2-form ω lies in Λ^+).

The irreducible action of $SO(3) = G$ on S^4 has generic orbit G/V , and singular orbits $\mathcal{L} \cong G/L$ and $\mathcal{R} \cong G/R$ where L and R are copies of $O(2)$ in different blocks of $SO(3)$. Topologically, \mathcal{O}_k is constructed by gluing disk bundles over \mathcal{L} and \mathcal{R} , each a copy of $\mathbb{R}\mathbb{P}^2$. If we choose \mathcal{L} to be the singular locus, then

$$\mathcal{O}_k \approx (G \times_{L/\ell} (D^2/\ell) \bigcup_{G/V} (G \times_R D^2),$$

and this space is still homeomorphic to S^4 . The subgroups L, R act on the tangent and normal spaces T, T^\perp as $(e^{i\ell\theta}, e^{-2i\theta})$ at $x \in \mathcal{L}$, and as $(e^{i\theta}, e^{2i\theta})$ at $y \in \mathcal{R}$. The factors of 2 are needed to match the generic stabilizer V . The actions on $\Lambda^+(T \oplus T^\perp)$ are therefore $e^{(\ell-2)i\theta}$ at x and $e^{3i\theta}$ at y . The inclusion

$$(4.44) \quad U(1) \subset L \subset SO(3) \times SU(2)^+ SU(2)^-$$

is given by $(e^{i\ell\theta}, e^{(\ell-2)i\theta}, e^{(\ell+2)i\theta})$ at x , and the differing weights $\ell, \ell+2$ at x show that the $SO(3)^+$ principal bundle $P^+\mathcal{O}_k$ is smooth. The same is true of $P^-\mathcal{O}_k$, though the SD/ASD distinction can be verified by considering $\mathbb{C}\mathbb{P}^2$ for which $\Lambda^+ = \langle \omega \rangle \oplus [\Lambda^{2,0}]$ extends the canonical line bundle. The singular locus \mathcal{L} of $\mathcal{O}_4 = \mathbb{C}\mathbb{P}^2/\mathbb{Z}_2$ is totally real, reflecting the weight $\ell-2=0$. When $\ell=3$, the group diagram for $\tilde{P}^+\mathcal{O}_5$ with weights $(3, 1)$ from (4.44) and (at \mathcal{R}) $(1, 3)$ coincides with that of B^7 , if we ignore the third factor in (4.44), and let $SO(3)$ play the role of $SU(2)^-$ on B^7 . Whence the equivalence $\tilde{P}^+\mathcal{O}_5 \cong B^7$.

Remark 4.6. Although B^7 cannot be the total space of a smooth principal S^3 bundle, it is known to be diffeomorphic to a smooth S^3 bundle over S^4 with Euler class ± 10 and (in particular) $p_1 = \pm 16$ [21].

The orbifolds \mathcal{O}_k arise from those on the moduli spaces of monopoles of charge 2 and mass 1 on hyperbolic space $\mathcal{H}^3(-16/(k-2)^2)$ for $k \geq 3$. Their metrics can (in theory, all) be constructed from solutions of Painlevé VI, and approach the hyperkähler metric on the Atiyah-Hitchin space M_0^2 [24, 4] as $k \rightarrow \infty$. Explicit algebraic expressions and fascinating graphs for $\ell = 3, 4, 6$ can be found in [35]. The Konishi bundle $P^-\mathcal{O}_k$ inherits an $SO(4)$ -invariant 3-Sasakian metric. Therefore both $P^-\mathcal{O}_5$ and $P^+\mathcal{O}_5$ admit (albeit, very different) NP_2 structures, and it is natural to ask whether the same could be true for $k > 5$. There is an analogous question for positive sectional curvature, since modification of the 3-Sasakian metric on $P^-\mathcal{O}_5$ has this property for $k = 5$ [23, 13].

5. HYPERSURFACE CALCULATIONS

In the setup of (3.33), we resort to the description of an NP_2 manifold by means of the induced $SU(3)$ structures on the hypersurfaces for which $t \in (0, \pi/3)$ is constant, so the generic $SO(4)$ orbits. The condition (2.12) is equivalent to the next two equations:

$$(5.45) \quad d\xi - \frac{\partial}{\partial t} \text{Re } \Xi = \mu \text{Im } \Xi$$

$$(5.46) \quad d(\text{Re } \Xi) = \mu \frac{1}{2} \xi^2.$$

The second characterizes the so-called nearly-half-flat (NHF) condition introduced by [15], and studied in the present context by the second author [33].

Recall the NP_2 structure on S^7 determined by the the basis (X_i) and the 2×2 matrices that feature in Corollary 3.4. Let us just consider 6-dimensional orbits of $SO(4)$ with slightly more general $SU(3)$ structures. The next result shows that it is to some extent possible to reverse engineer the constants occurring in Corollary 3.4 by merely applying the NHF condition for all values of t :

Proposition 5.1. *Define ξ and Ξ by (3.33) with*

$$\begin{pmatrix} X_{2k-1} \\ X_{2k} \end{pmatrix} = \begin{pmatrix} \lambda & \lambda(ac_k + b) \\ 0 & 4s_k \end{pmatrix} \begin{pmatrix} f_k \\ e_k \end{pmatrix},$$

where λ, a, b are (unconstrained) constants. Then (5.46) holds for all t iff

- (i) $(a, b) = (0, -1)$ or $(0, 0)$, in which case $\mu = -2/\lambda$, or
- (ii) $(a, b) = (\pm 2, 1)$, in which case $\mu = -\frac{1}{2}(\lambda^2 + 4)/\lambda$.

In this formulation, the coefficient ‘4’ is retained as a normalization.

Proof. Equations (3.33) yield $\xi = -4\lambda(s_1e_1f_1 + s_2e_2f_2 + s_3e_3f_3)$ so

$$\frac{1}{2}\xi^2 = -16\lambda^2(s_2s_3e_{23}f_{23} + s_3s_1e_{31}f_{31} + s_1s_2e_{12}f_{12}).$$

Moreover

$$\begin{aligned} d\Xi &= (-2\lambda f_{23} - 2\lambda(ac_1 + b)e_{23} - 8is_1e_{23}) \wedge \\ &\quad (\lambda f_2 + \lambda(ac_2 + b)e_2 + 4is_2e_2) \wedge (\lambda f_3 + \lambda(ac_3 + b)e_3 + 4is_3e_3) + \dots \end{aligned}$$

Therefore the $e_{23}f_{23}$ component, call it Γ_1 , of $d(\text{Re } \Xi)$ equals

$$\begin{aligned} &-2\lambda[\lambda^2(ac_2 + b)(ac_3 + b) - 16s_2s_3] - 2\lambda^3(ac_1 + b) \\ &= -2\lambda^3[a^2c_2c_3 - abc_1 + b^2] + 32\lambda s_2s_3 - 2\lambda^3(ac_1 + b) \\ &= -2\lambda^3(b + b^2) - 2\lambda^3a^2c_2c_3 + 32\lambda s_2s_3 + 2\lambda^3(ab - a)c_1 \\ &= -2\lambda^3(b + b^2) - 16\lambda - 2\lambda(16 + a^2\lambda^2)c_2c_3 + 2\lambda^3(ab - a)c_1, \end{aligned}$$

the last line by (3.26). This must equal μ times $-16\lambda^2s_2s_3 = 8\lambda^2(1 + 2c_2c_3)$ for all t . To eliminate the coefficient of c_1 above, we need $a = 0$ or $b = 1$.

If $a = 0$, we obtain the system

$$\begin{cases} -32\lambda = 16\lambda^2\mu \\ -2\lambda^3b(1+b) - 16\lambda = 8\lambda^2\mu, \end{cases}$$

which implies that $\lambda\mu = -2$ and $b = 0$ or -1 . If $b = 1$, we obtain

$$-4\lambda^3 - 2\lambda(8 + a^2\lambda^2)c_2c_3 - 8\lambda = \mu(8a^2\lambda^2 + 16c_2c_3),$$

which provides the system

$$\begin{cases} \lambda^2 + 2\mu\lambda + 4 = 0 \\ a^2\lambda^2 + 2\mu\lambda + 4 = 0, \end{cases}$$

and $a = \pm 2$. The unique quadratic then gives the constraint between λ and μ . \square

As a corollary, any positive definite linear combination of g_7 and h_7 on S^7 is associated to an NHF structure. Note however that $\lambda = 2/\sqrt{5}$ implies that $\mu = -6/\sqrt{5}$, as in Proposition 2.1 modulo orientation.

The analysis of (5.45) requires more extensive calculation. Turning to B^7 , we have established that the solution in Corollary 4.4 is essentially unique within a 3-parameter family. This is part (i) of

Theorem 5.2. *Define ξ and Ξ by (3.33) with Y_i in place of X_i , where now*

$$\begin{pmatrix} Y_{2k-1} \\ Y_{2k} \end{pmatrix} = \begin{pmatrix} 2\lambda & 2\lambda ac_k \\ b & 2s_k \end{pmatrix} \begin{pmatrix} p_k \\ n_k \end{pmatrix},$$

and λ, a, b are (unconstrained) constants. Then

- (i) (5.45) and (5.46) both hold iff $(\lambda, a, b) = (\pm \frac{2}{\sqrt{5}}, \frac{1}{2}, 0)$;
- (ii) (5.46) holds if either $\lambda^2(4 - a^2) = 3$ and $b = 0$, or $(\lambda, a, b) = (\frac{1}{2}, 2, \pm\sqrt{3})$.

In all cases, μ is determined by λ .

We omit the proof of (i), since analysis of (5.45) is somewhat fruitless. The proof of (ii) is analogous to that of Proposition 5.1, but more involved since it is based upon (4.43). Both parts were checked by hand and using *Maple*, but in different orders.

The significance of the condition $b = 0$ is that it guarantees smooth extension to the singular orbits. This justifies our presentation of the S^7 and B^7 solutions using upper triangular 2×2 matrices, and contrasting geometry at the singular orbits. The special NHF solutions in this section will be analysed elsewhere.

There is an obvious generalization of the hypotheses of Theorem 5.2 that will also capture the NP_2 solution on the principal part $SO(4) \times (0, \pi/3)$ of B^7 . Namely one can saturate the first row of the 2×2 matrix by linear combinations 1 and c_k , and its second row by terms proportional to s_k . For that matter, one could insert less trivial Fourier series in each slot. The prospect of adapting the methods to [17] to find new compact NP_2 manifolds of cohomogeneity one appears remote. Nonetheless, the results of this section motivate a computational search for other NP_2 solutions of the above type.

ACKNOWLEDGEMENTS

Both authors were supported by the Simons Foundation (#488635 Simon Salamon). Some of the results of this paper were presented at the conference ‘Special Holonomy and Geometric Structures on Complex Manifolds’ at IMPA in March 2024, and we thank the organizers.

REFERENCES

- [1] E. Abbena, S. Salamon, A. Gray: *Modern Differential Geometry of Curves and Surfaces with Mathematica*, 3rd edition, Chapman & Hall/CRC, 2006. [7](#)
- [2] I. Agricola, G. Dileo, L. Stecker: Curvature properties of 3- (α, δ) -Sasaki manifolds. arXiv:2206.05150. [13](#)
- [3] I. Agricola, T. Friedrich: 3-Sasakian manifolds in dimension seven, their spinors and G_2 -structures. *J. Geom. Physics* 60 (2010) 326–332.
- [4] M.F. Atiyah, N.J. Hitchin: *The Geometry and Dynamics of Magnetic Monopoles*. Porter lectures, Princeton University Press, 1988. [17](#)
- [5] G. Ball, J. Madnick: Associative submanifolds of squashed 3-Sasakian manifolds. arXiv:2208.10622 [2](#)
- [6] G. Ball, J. Madnick: Associative Submanifolds of the Berger Space. *Commun. Analysis Geometry*, to appear. [2](#), [12](#), [13](#), [16](#)
- [7] C. Bär: Real Killing spinors and holonomy. *Commun. Math. Phys.* 154 (1993) 509–521. [1](#), [7](#)
- [8] Ya.V. Bazaikin: Noncompact Riemannian spaces with the holonomy group $Spin(7)$ and 3-Sasakian manifolds. *Proc. Steklov Inst. Math.* 263(2008) 2–12. [8](#), [13](#)
- [9] M. Berger: Les variétés riemanniennes homogènes normales simplement connexes à courbure strictement positive. *Ann. Scuola Norm. Sup. Pisa* 15 (1961) 179–246. [2](#), [4](#), [12](#)
- [10] C. Boyer, K. Galicki: *Sasakian Geometry*. Oxford Mathematical Monographs, Oxford University Press, 2008. [7](#)
- [11] R.L. Bryant: Some remarks on G_2 structures. *Proc. Gokova Geometry-Topology Conference 2005*, eds. S. Akbulut, T. Onder, R.J. Stern, International Press (2006) 75–109.
- [12] R.L. Bryant, S.M. Salamon: On the construction of complete metrics with exceptional holonomy. *Duke Math. J.* 58 (1989) 829–850. [2](#), [3](#), [4](#), [5](#)
- [13] O. Dearnicott: A 7-manifold with positive curvature. *Duke Math. J.* 158 (2011) 307–346. [17](#)
- [14] O. Dearnicott: Positively curved self-dual Einstein metrics on weighted projective planes. *Ann. Global Anal. Geom.* 27 (2005) 79–86. [1](#)
- [15] M. Fernández, S. Ivanov, V. Muñoz, L. Ugarte, Nearly hypo structures and compact nearly Kähler 6-manifolds with conical singularities. *J. Lond. Math. Soc.* 78 (2008) 580–604. [18](#)
- [16] A. Fino, P. Nuruowski: Analog of selfduality in dimension nine. *J. Reine Angew. Math.* 699 (2015) 67–110. [9](#)
- [17] L. Foscolo, M. Haskins: New G_2 holonomy cones and exotic nearly Kähler structures on S^6 and $S^3 \times S^3$. *Annals of Math.* 185 (2017) 59–130. [19](#)
- [18] Th. Friedrich, I. Kath, A. Moroianu, U. Semmelmann: On nearly parallel G_2 -structures. *J. Geom. Physics* 23 (1997) 259–286. [8](#)
- [19] K. Galicki, H.B. Lawson Jr.: Quaternionic reduction and quaternionic orbifolds. *Math. Annalen* 282 (1988) 1–22. [1](#)
- [20] K. Galicki, S. Salamon: Betti numbers of 3-Sasakian manifolds. *Geom. Dedicata* 63 (1996) 45–68. [5](#), [6](#), [7](#)
- [21] S. Goette, N. Kitchloo, K. Shankar: Diffeomorphism type of the Berger space $SO(5)/SO(3)$. *Amer. J. Math.* 126 (2004) 395–416. [13](#), [17](#)
- [22] A. Gray: The structure of nearly Kähler manifolds. *Math. Annalen* 223 (1976) 233–248 [4](#)

- [23] K. Grove, B. Wilking, W. Ziller: Positively curved cohomogeneity one manifolds and 3-Sasakian geometry. *J. Differ. Geom.* 78 (2008) 33–111. [2](#), [10](#), [15](#), [17](#)
- [24] N.J. Hitchin: A new family of Einstein metrics. In: *Manifolds and Geometry* (Pisa, 1993). *Sympos. Math.* XXXVI (1996) 190–222. [1](#), [17](#)
- [25] K. Kawai: Some associative submanifolds of the squashed 7-sphere. *Quart. J. Math.* 66 (2015) 861–893. [1](#), [2](#)
- [26] A. Kennon, J. Lotay: Geometric flows of G_2 -structures on 3-Sasakian 7-manifolds. *J. Geom. Physics* 187 (2023) 104793.
- [27] P.Z. Kobak, A. Swann: Classical nilpotent orbits as hyperkähler quotients. *Intern. J. Math.* 7 (1996) 193–210. [16](#)
- [28] M. Konishi: On manifolds with Sasakian 3-structure over quaternion Kaehler manifolds. *Kodai Math. Sem. Rep.* 26(1975) 194–200. [1](#), [2](#)
- [29] P.B. Kronheimer: Instantons and the geometry of the nilpotent variety. *J. Differ. Geom.* 32 (1990) 473–490.
- [30] J.D. Lotay: Associative submanifolds of the 7-sphere. *Proc. London Math. Soc.* 105 (2012) 1183–1214. [2](#)
- [31] F. Podestà, L. Verdiani: Positively curved 7-dimensional manifolds. *Quart. J. Math.* 50 (1999) 497–504. [2](#), [14](#)
- [32] S. Salamon: *Riemannian Geometry and Holonomy Groups*, Longman Scientific and Technical, Harlow, Essex, 1989. [2](#)
- [33] R. Singhal: Nearly half-flat $SU(3)$ -structures on $S^3 \times S^3$. *Differ. Geom. Appl.* 97 (2024) 102187. [18](#)
- [34] A. Swann: HyperKähler and quaternionic Kähler geometry. *Math. Ann.* 289 (1991) 421–450. [7](#)
- [35] W. Ziller: On the geometry of cohomogeneity one manifolds with positive curvature. In: *Riemannian Topology and Geometric Structures on Manifolds*, *Progress Math.* 271 (2009) 233–262. [2](#), [8](#), [11](#), [17](#)

University of Münster, Einsteinstrasse 62, 48149 Münster, Germany.
`rsinghal@uni-muenster.de`

King's College London, Strand, London, WC2R 2LS, UK.
`simon.salamon@kcl.ac.uk`