Composite nature of the T_{cc} state

Xian-Wei Kang and Wen-Shuo Ding

Key Laboratory of Beam Technology of Ministry of Education,

School of Physics and Astronomy, Beijing Normal University, Beijing 100875, China and Institute of Radiation Technology,

Beijing Academy of Science and Technology, Beijing 100875, China

In 2021, LHCb collaboration reported a very narrow state in the $D^0D^0\pi^+$ mass spectrum just below the $D^{*+}D^0$ mass threshold. We consider the influence of the Castillejo-Dalitz-Dyson (CDD) pole in the scattering amplitude to derive a general treatment for the two-body final state interaction near its threshold. The line shape (or the energy dependent event distribution) are then obtained, where the parameters can be fixed by fitting to the experimental data on the $D^0D^0\pi^+$ mass spectrum. Within our method the data are quite well reproduced. The pole structure in the complex energy plane indicates the bound state structure of the T_{cc} state. The compositeness as a measure of molecule component in its hadron wave function is predicted to be $0.23^{+0.40}_{-0.09}$. The non-molecular component, e.g., the compact tetraquark also takes a non-negligible portion.

I. INTRODUCTION

In 2021, LHCb reported a narrow state named by T_{cc} in $D^0D^0\pi^+$ mass spectrum just below the $D^{*+}D^0$ mass threshold. The pole in the second Reimann sheet of the $D^{*+}D^0$ scattering amplitude with respect to the $D^{*+}D^0$ threshold is found to be [1, 2]

$$\delta m_{\text{pole}} = -360 \pm 40^{+4}_{-0} \text{ keV},$$

$$\Gamma_{\text{pole}} = 48 \pm 2^{+0}_{-14} \text{ keV}.$$
(1)

Since the state is so close to the $D^{*+}D^0$ threshold, Breit-Wigner parameterization is not appropriate and we should focus this pole parameter as the mass and width values.

This observation arose great interest concerning its feature in the hadron world. It was understood as the molecular state generated by DD^* scattering [3–6], or the compact tetraquark [7–10], or the virtual state by a refined data analysis utilizing K-matrix with the Chew-Mandelstam formalism [11], or kinematic singularity [12]. The

lattice analysis assign the observed T_{cc} as the virtual state [13]. For a review one may refer to Ref. [14]. In fact, a genuine state may involve several configurations. There are recent developments for constructing the compositeness to quantify the molecular component [15–18].

The Castillejo-Dalitz-Dyson (CDD) pole is proposed in the context of Low equation in 1956 [19]. The pointed out that infinite number of parameters (pole location and residue corresponding to CDD pole) can appear in the Low equation, and also those appearance of CDD pole signifies the amount of the molecular component. If the CDD pole sits very close to the threshold of the two scattering hadrons, elementary degree of freedom is dominant for the generated resonances by the scattering, and otherwise, the two-body molecular component is dominant.

Combining the concenpt of the compositeness and the scattering amplitude with inclusion of the CDD pole has been an effective method to analyze the inner structure of hadron state. The application along this line can be found in Refs. [20, 21] for X(3872); in Refs. [22, 23] for $Z_b(10610)$ and $Z_b(10650)$; in Ref. [24] for $f_0(980)$; in Ref. [25] for $\Lambda_c^+(2595)$ and in Ref. [26] for other near-threshold heavy flavor resonances.

We next in Sec. II derive the expression for the partial-wave amplitude with the impact of the CDD pole, from which we derive the event distribution. The parameters are obtained by fit to the mass spectrum data. In Sec. III we show our theoretical results with the analysis of the compositeness value. In Sec. IV we render the summary and outlook.

II. SCATTERING AMPLITUDE AND EVENT DISTRIBUTION

Due to the extreme proximity between the state T_{cc} and $D^{*+}D^0$ mass threshold, within one MeV, it is enough that to consider the $D^{*+}D^0$ scattering in non-relativistic form. The spin-parity quantum number of T_{cc} is determined to be 1⁺, so the $D^{*+}D^0$ scattering mainly occurs in S wave. For such case, $D^{*+}D^0$ scattering amplitude in terms of the energy of DD^* can be written as [21, 23, 25]

$$t(E) = \left[\frac{\lambda}{E - M_{\rm CDD}} + \beta - ik\right]^{-1},\tag{2}$$

where M_{CDD} and λ are the position and residue, respectively, for the CDD pole; $m_{\text{th}} = m_{D^*} + m_{D^0}$ is the mass threshold; and $k = \sqrt{2\mu_{D^*D}(E - m_{\text{th}})}$ is the magnitude of

three-momentum with $\mu_{D^*D} = m_{D^0}m_{D^{*+}}/(m_{D^0} + m_{D^{*+}})$ denoting the reduced mass of D^{*+} and D^0 . The unitarity condition is $\operatorname{Im} t^{-1}(E) = -k$. In the second Riemann Sheet (RS), indicated by a superscript II, it becomes

$$t^{\rm II}(E) = \left[\frac{\lambda}{E - M_{\rm CDD}} + \beta + ik\right]^{-1}.$$
 (3)

Note that there is a change of sign in front of k comparing with t(E) for the first (physical) RS in Eq. (2). In this convention k is always calculated such that Im k > 0. Comparing to the effective range expansion $t(E) = \left(-\frac{1}{a} + \frac{1}{2}rk^2 - ik\right)^{-1}$, one can obtain the scattering length a and effective range r as

$$\frac{1}{a} = \frac{\lambda}{M_{\rm CDD} - m_{\rm th}} - \beta,$$

$$r = -\frac{\lambda}{\mu_{D^*D}(M_{\rm CDD} - m_{\rm th})^2}.$$
(4)

Clearly, when $M_{\text{CDD}} \approx m_{\text{th}}$, i.e., the CDD pole position is extremely close to the twomeson threshold, r goes to infinity. In this circumstance, the effective range expansion approach fails due to the too limited convergence radius. We do encounter this situation of $r \to \infty$ in Refs. [21] and [23]. Or more precisely, the data quality is not capable of pinning down the location of CDD pole.

As a matter of the fact, CDD pole is zero of the scattering amplitude t(E), which strongly distorts the energy dependence of t(E) near $E \approx M_{\text{CDD}}$. The production process is mediated by the following d(E) (but not t(E)) by removing the extra $E - M_{\text{CDD}}$ factor in t(E) [25]:

$$d(E) = \left(1 + \frac{E - M_{\text{CDD}}}{\lambda}(\beta - ik)\right)^{-1}$$
(5)

We use the function d(E) to treat the final state interaction. The factor $|d(E)|^2$ constituents the parametrization for the signal. Near the resonance region, one has the form

$$d(E) \simeq \frac{\gamma_E}{E - E_P},\tag{6}$$

with $E_P = M_P - i \Gamma_P/2$ and γ_E denoting the pole and its residue in the complex *E*-plane. The residue can be calculated by an integration along a closed contour around the pole:

$$\gamma_E = \frac{1}{2\pi i} \oint d(E) \mathrm{d}E. \tag{7}$$

Then the mass distribution can be written as

$$\frac{\mathrm{d}N}{\mathrm{d}E} = \frac{\Gamma_P |d(E)|^2}{2\pi |\gamma_E|^2},\tag{8}$$

with Γ_P denoting the pole width. We take the normalization condition such that for a narrow resonance case, the following integration

$$\mathcal{N} = \int_{-\infty}^{+\infty} \mathrm{d}E \frac{\mathrm{d}N}{\mathrm{d}E} \tag{9}$$

is close to 1. In such scenario, the following Y_{const} can be understood as the yield for this decay process. Similar choice is also taken in Refs. [21, 27]. When E_P corresponds to a virtual state or other situations for which the final-state interaction function d(E)has a shape that strongly departs from a non-relativistic Breit-Wigner one, \mathcal{N} could be far away from 1. Our description Eq. (5) can display various line shapes beyond one shown in Eq. (6) and provides a more general treatment for the final state interaction. So we claim Eq. (5) is applicable in a wider range.

We now consider the experimental mass resolution and the corresponding background. Following the experiment [2], the energy resolution function is described by a sum of two Gaussian functions

$$R(E', E) = \frac{0.778}{\sqrt{2\pi\sigma_1}} \exp\left[\frac{(E'-E)^2}{-2\sigma_1^2}\right] + \frac{0.222}{\sqrt{2\pi\sigma_2}} \exp\left[\frac{(E'-E)^2}{-2\sigma_2^2}\right],$$
(10)

with $\sigma_1 = 276.15$ keV and $\sigma_2 = 666.35$ keV. The background contribution is parameterized by

$$B(E) = P_2 \times \Phi_{D^{*+}D^0},$$
(11)

with $P_2 = aE^2 + bE + c$ denoting a positive second-order polynomial, and

$$\Phi_{D^{*+}D^0} = \frac{\sqrt{(E^2 - (m_{D^0} + m_{D^{*+}})^2)(E^2 - (m_{D^0} - m_{D^{*+}})^2)}}{2E}$$
(12)

the two-body phase space factor. Finally, we obtain the energy-dependent event number distribution in an energy bin of width $\Delta = 500$ keV centered at E_i :

$$N(E_i) = \int_{E_i - \Delta/2}^{E_i + \Delta/2} dE' \int_{-\infty}^{+\infty} dE \left[Y_{\text{const}} \left| d(E) \right|^2 + B(E) \right] R(E', E).$$
(13)

We have 7 parameters in total: λ , M_{CDD} , β in d(E) function, a, b, c in the background, and Y_{const} as the overall normalization constant. These parameters could be fixed by fit to the data. In the fit, we should include the finite D^{*+} with, $\Gamma_{D^{*+}} = 83.4$ keV [28], and as a result,

$$k = \sqrt{2\mu_{D^*D}(E - m_{\rm th} + i\Gamma_{D^{*+}}/2)}.$$
(14)

The T_{cc} state is so close to threshold (recalling the value of δm_{pole} in Eq. (1)) such that the inclusion of 83.4 keV is necessary, as has also been verified by the numerical calculation.

We use the optimization algorithm of MINUIT package to perform the minimization of χ^2 [29]. The parameter values are shown in Table I. In the calculation, all the mass, energy and momentum are expressed in units of MeV. Then the units for the parameters are also given in the table. The fit result is shown in Figure 1, where "Energy" corresponds to the invariant mass of $D^0 D^0 \pi^+$. In fact, near the pole (signal) region, the invariant mass $M_{D^{*+}D^0} \approx M_P \approx M_{D^0D^0\pi^+}$. From the following discussions, we claim the T_{cc} state can be understood as the $D^{*+}D^0$ bound state. For a bound state case, the aforementioned $\mathcal{N} \approx 1$ is clear since Lorentzian distribution reduces to a delta function. Y_{const} can then be understood as the yield corresponding to the product of production cross section of T_{cc} and branching fractions of $T_{cc} \to D^{*+}D^0$ and $D^{*+} \to D^0 \pi^+$ (note that cross section has dimension of [MeV]⁻²).

$Y_{\rm const} = (10.8 \pm 8.0) \ {\rm MeV}^{-2}$		
$\lambda = (83.6 \pm 63.8)~\mathrm{MeV^2}$	$\beta = (70.5 \pm 37.6) \text{ MeV}$	$M_{\rm CDD} - m_{\rm th} = (0.47 \pm 0.38) {\rm MeV}$
$a = -(81.4 \pm 2.8) \text{ MeV}^{-2}$	$b = -(99.2 \pm 10.8) \ {\rm MeV^{-3}}$	$c = (1653.5 \pm 41.9) \text{ MeV}^{-4}$

Table I: Our best values of the 7 fitted parameters, with χ^2 per degree of freedom being 0.93.

Figure 1: Mass spectrum for the $D^0 D^0 \pi^+$ decay channel. The data are from the LHCb collaboration [2]. The line represents our fit result.

In Ref. [23], we have imposed a pole in the second sheet with pole parameters given by the experimental determination. In this way, λ and β are expressed as functions of M_{CDD} , and then two parameters will be reduced. We have tried the similar exercise in our present study. The fit result is acceptable, but not as good as the one above. One of the reason could be that the width of T_{cc} is much narrower or the peak is more prominent than the ones for Z_b states. We will not show its parameter value and figure. However, the conclusion that both poles in the first and second sheet are found does not change. In the second sheet, the pole is at $-0.36 \text{ MeV} + m_{\text{th}} - i\,0.024 \text{ MeV}$ (cf. Eq. (1)), as required. In this case we can not provide its compositeness value since it locates below the threshold. For the bound state case, a small compositeness number is found, which implies the compact tetraquark component in its wave function can not be overlooked.

III. SCATTERING PARAMETERS AND COMPOSITENESS

Using the central values of the fitted parameters, one obtains the scattering length a = 1.8 fm and the effective range r = -77.2 fm. The value of r is so large that the effective range expansion method is not an appropriate tool in this case. Within uncertainties we have another set of parameter value, $\lambda = 19.8$, $\beta = 108.0$, $M_{\rm CDD} = 0.47 + m_{\rm th}$, which leads to a = -3.0 fm, r = -18.3 fm. The values of r are not the typical magnitude of a few fermi. So we will concentrate on the outcome of our CDD analysis.

The definition of compositeness for a bound state is well defined by the Weinberg formula [30, 31]. However, for resonance case, the compositeness calculated in that way will become a complex-valued number. As mentioned before, there are several developments for defining the compositeness. We will follow the one developed in Ref. [16] for a resonance:

$$X = |\gamma_s|^2 \left| \frac{dG(s)}{ds} \right|_{s=E_P^2},\tag{15}$$

for the case of $\sqrt{\text{Re}E_P^2}$ larger than the lightest threshold for a coupled-channel scattering. $-\gamma_s^2$ is the residue of t(s) at the resonance pole position, E_P^2 , of the complex *s* plane in the second RS:

$$t^{\rm II}(s) \to \frac{-\gamma_s^2}{s - E_P^2}.\tag{16}$$

 γ_s^2 is related to γ_E by [21, 23]

$$\gamma_s^2 = -16\pi\gamma_E E_P^2 \times \frac{E_P - M_{\rm CDD}}{\lambda}.$$
(17)

G(s) is the two-point Green function, and the infinity can be removed by cutoff regularization or the dimensional regularization [32]. Its form in the dimensional regularization scheme at the regularization scale μ can be written as [25]

$$G(s) = \alpha(\mu^2) + \frac{1}{(4\pi)^2} \left(\log \frac{m_2^2}{\mu^2} - \varkappa_+ \log \frac{\varkappa_+ - 1}{\varkappa_+} - \varkappa_- \log \frac{\varkappa_- - 1}{\varkappa_-} \right),$$
(18)

with

$$\varkappa_{\pm} = \frac{s + m_1^2 - m_2^2}{2s} \pm \frac{p}{\sqrt{s}},$$

$$p = \frac{\sqrt{(s - (m_1 - m_2)^2)(s - (m_1 + m_2)^2)}}{2\sqrt{s}},$$
(19)

It is equavalent to the one used in Ref. [33]. The G(s) defined in Eq. (18) corresponds to the physical value on the real axis, or $G^{I}(s + i\epsilon)$ in the physical sheet.

Resonance appears as a pole in the second RS. We thus need to make an analytical extrapolation of G(s) to the second RS. The result is given by

$$G^{\rm II}(s+i\epsilon) = G^{\rm I}(s+i\epsilon) + \frac{i}{4\pi\sqrt{s}} \frac{\sqrt{(s-(m_1-m_2)^2)(s-(m_1+m_2)^2)}}{2\sqrt{s}}.$$
 (20)

The left and right side of Eq. (20) implies the condition of Im s > 0. For Im s < 0 case, using the Schwarz reflection principle $G(s^*) = [G(s)]^*$, one has

$$G^{\rm II}(s-i\epsilon) = G^{\rm I}(s-i\epsilon) - \frac{i}{4\pi\sqrt{s}} \frac{\sqrt{(s-(m_1-m_2)^2)(s-(m_1+m_2)^2)}}{2\sqrt{s}}.$$
 (21)

It is G^{II} that should be used in Eq. (15) for a resonance. For a bound state (E_B) case, one simply has

$$X = -\gamma_s^2 \left. \frac{dG^{\mathrm{I}}(s)}{ds} \right|_{s=E_B^2},\tag{22}$$

and of course γ_s is calculated in the physical sheet. The situation of X = 1 corresponds to a pure bound state or molecular. The elementariness Z = 1 - X measures the weight of all other components in the hadron wave function.

We search for the pole in the complex energy plane. We first consider the welldefined elastic scattering case with $\Gamma_{D^{*+}} \rightarrow 0$. As a result, for the central value of the fitted parameter we find a pole 3874.72 MeV in the physical sheet, as a bound state; and a pole 3874.48 – i 1.74 MeV in the unphysical Riemann sheet, as a resonance. By slowly varying e.g., the values of $M_{\rm CDD}$ both poles move gradually. For the bound state pole, we have the residue $\gamma_s^2 = 5.04 \text{ GeV}^2$, and the corresponding compositeness X = 0.23. Considering the uncertainties in Table. I within one sigma region, we have $X = 0.23^{+0.40}_{-0.09}$. The finite but small D^{*+} width moves the aforementioned resonance pole to 3874.5 - i1.73 MeV and the bound state pole to 3874.72 - i0.0098 MeV. It is interesting to note that our such finding of a bound state pole agrees well with Ref. [34], where they find a bound state with width of 80 keV by using the complex scaling method, since the parameter values $\lambda = 19.8$, $\beta = 108.0$, $M_{\rm CDD} = 0.47 + m_{\rm th}$ leads to a bound state pole at 3875.4 - i0.039 with the width of 78 keV for the unstable bound state. On the other hand, the resonance pole width found is too large compared to the corresponding experimental value of the T_{cc} . From this viewpoint, it is not appropriate to be assigned as T_{cc} . However, the pole parameter in Eq. (1) derives from the analysis by an unitary Breit-Wigner model. The precise determination of the pole parameter still needs more refined study. For some of the resonance poles that are found, we can not provide its compositeness value since $\sqrt{\operatorname{Re} E_P^2} < m_{\text{th}}$. If the condition $\sqrt{\operatorname{Re} E_P^2} \ge m_{\text{th}}$ is fulfilled, the compositeness formula, Eq. (15), could be applied and the compositeness X can range from 0.07 to 0.57 for the resonance interpretation. As a short summary, we favor the bound state interpretation; and in any case, the twobody hadron component plays an important role in the T_{cc} state; however, the other configuration, e.g., the compact tetraquark component takes a comparable portion.

IV. CONCLUSION AND OUTLOOK

The mystery of T_{cc} state reported by LHCb has not been unveiled, where various interpretations are proposed. We utilize an amplitude including the CDD pole to incorporate the nontrivial and non-perturbative final state interaction of $D^{*+}D^0$. This method provides much more information than the effective range expansion method. The appearing parameters are λ , β and M_{CDD} , which will be fixed by fitting to the data. In the fit the experimental energy bin width and mass resolution function are considered. With the known parameters, we search for the pole in the complex energy plane. Both poles are found in the physical and unphysical Riemann sheet. Considering both the chi-squared value and the pole parameter, we favour the bound state interpretation. That is, the reported T_{cc} pole is understood as the $D^{*+}D^0$ bound state, with compositeness value of $X = 0.23^{+0.40}_{-0.09}$, while 1 - X provides a measure of other components, e.g., the compact tetraquark. The coupled-channel study following Ref. [21] may render more useful information and this work is ongoing.

Acknowledgments

We are indebted to J. A. Oller for fruitful discussion. This work is supported by the National Natural Science Foundation of China under Project No. 12275023.

- [1] R. Aaij et al. [LHCb], Nature Phys. 18, no.7, 751-754 (2022) [arXiv:2109.01038 [hep-ex]].
- [2] R. Aaij *et al.* [LHCb], Nature Commun. 13, no.1, 3351 (2022) [arXiv:2109.01056 [hep-ex]].
- [3] A. Feijoo, W. H. Liang and E. Oset, Phys. Rev. D 104, no.11, 114015 (2021)
 [arXiv:2108.02730 [hep-ph]].
- [4] M. Albaladejo, Phys. Lett. B 829, 137052 (2022) [arXiv:2110.02944 [hep-ph]].
- [5] M. Abolnikov, V. Baru, E. Epelbaum, A. A. Filin, C. Hanhart and L. Meng, [arXiv:2407.04649 [hep-ph]].
- [6] M. L. Du, V. Baru, X. K. Dong, A. Filin, F. K. Guo, C. Hanhart, A. Nefediev, J. Nieves and Q. Wang, Phys. Rev. D 105, no.1, 014024 (2022) [arXiv:2110.13765 [hep-ph]].
- [7] X. Z. Weng, W. Z. Deng and S. L. Zhu, Chin. Phys. C 46, no.1, 013102 (2022)
 [arXiv:2108.07242 [hep-ph]].
- [8] Y. Kim, M. Oka and K. Suzuki, Phys. Rev. D 105, no.7, 074021 (2022) [arXiv:2202.06520
 [hep-ph]].
- [9] M. Karliner and J. L. Rosner, Phys. Rev. D 105, no.3, 034020 (2022) [arXiv:2110.12054
 [hep-ph]].
- [10] W. C. Dong and Z. G. Wang, [arXiv:2407.19383 [hep-ph]].
- [11] L. Y. Dai, X. Sun, X. W. Kang, A. P. Szczepaniak and J. S. Yu, Phys. Rev. D 105, no.5, L051507 (2022) [arXiv:2108.06002 [hep-ph]].
- [12] N. N. Achasov and G. N. Shestakov, Phys. Rev. D 105, no.9, 096038 (2022)
 [arXiv:2203.17100 [hep-ph]].
- [13] T. Whyte, D. J. Wilson and C. E. Thomas, [arXiv:2405.15741 [hep-lat]].
- [14] H. X. Chen, W. Chen, X. Liu, Y. R. Liu and S. L. Zhu, Rept. Prog. Phys. 86, no.2, 026201 (2023) [arXiv:2204.02649 [hep-ph]].
- [15] T. Kinugawa and T. Hyodo, Phys. Rev. C 109, no.4, 045205 (2024) [arXiv:2303.07038
 [hep-ph]].

- [16] Z. H. Guo and J. A. Oller, Phys. Rev. D 93, no.9, 096001 (2016) [arXiv:1508.06400 [hep-ph]].
- [17] V. Baru, J. Haidenbauer, C. Hanhart, Y. Kalashnikova and A. E. Kudryavtsev, Phys. Lett. B 586, 53-61 (2004) [arXiv:hep-ph/0308129 [hep-ph]].
- [18] T. Sekihara, T. Hyodo and D. Jido, PTEP 2015, 063D04 (2015) [arXiv:1411.2308 [hep-ph]].
- [19] L. Castillejo, R. H. Dalitz and F. J. Dyson, Phys. Rev. 101, 453-458 (1956)
- [20] X. W. Kang, J. Z. Zhang and X. H. Guo, [arXiv:2410.14521 [hep-ph]].
- [21] X. W. Kang and J. A. Oller, Eur. Phys. J. C 77, no.6, 399 (2017) [arXiv:1612.08420 [hep-ph]].
- [22] X. W. Kang, Z. H. Guo and J. A. Oller, Phys. Rev. D 94, no.1, 014012 (2016) [arXiv:1603.05546 [hep-ph]].
- [23] L. Zhang, X. W. Kang and X. H. Guo, Eur. Phys. J. C 82, no.4, 375 (2022) [arXiv:2203.02301 [hep-ph]].
- [24] Z. Q. Wang, X. W. Kang, J. A. Oller and L. Zhang, Phys. Rev. D 105, no.7, 074016
 (2022) [arXiv:2201.00492 [hep-ph]].
 20
- [25] Z. H. Guo and J. A. Oller, Phys. Rev. D 93, no.5, 054014 (2016) [arXiv:1601.00862 [hep-ph]].
- [26] R. Gao, Z. H. Guo, X. W. Kang and J. A. Oller, Adv. High Energy Phys. 2019, 4651908
 (2019) [arXiv:1812.07323 [hep-ph]].
- [27] E. Braaten and J. Stapleton, Phys. Rev. D 81, 014019 (2010) [arXiv:0907.3167 [hep-ph]].
- [28] S. Navas et al. [Particle Data Group], Phys. Rev. D 110, no.3, 030001 (2024)
- [29] F. James, "MINUIT C Function Minimization and Error Analysis", CERN Program Library Long Writeup D506, Version 94.1.
- [30] S. Weinberg, Phys. Rev. **130**, 776-783 (1963).
- [31] S. Weinberg, Phys. Rev. **137**, B672-B678 (1965).
- [32] X. W. Kang and J. A. Oller, Phys. Rev. D 94, no.5, 054010 (2016) [arXiv:1606.06665 [hep-ph]].
- [33] J. A. Oller, Eur. Phys. J. A 28, 63-82 (2006) [arXiv:hep-ph/0603134 [hep-ph]].
- [34] Z. Y. Lin, J. B. Cheng and S. L. Zhu, Phys. Rev. D 110, no.5, 5 (2024) [arXiv:2205.14628

 $[\mathrm{hep}\text{-}\mathrm{ph}]].$