Stringy scaling of multi-tensor hard string scattering amplitudes and the K-identities

Sheng-Hong Lai,^{1,*} Jen-Chi Lee,^{1,2,[†](#page-0-1)} and Yi Yang^{1,2,3,[‡](#page-0-2)}

¹Department of Electrophysics, National Yang Ming Chiao Tung University, Hsinchu, Taiwan, R.O.C. ²Center for Theoretical and Computational Physics (CTCP),

National Yang Ming Chiao Tung University, Hsinchu, Taiwan, R.O.C.

³National Center for Theoretical Physics, R.O.C.

(Dated: December 30, 2024)

Abstract

We calculate n-point hard string scattering amplitudes $(HSSA)$ with $n-2$ tachyons and 2 tensor states at arbitrary mass levels. We discover the stringy scaling behavior of these HSSA. It is found that for HSSA with more than 2 transverse directions, the degree of stringy scaling $\dim M_2$ decreases comparing to the degree of stringy scaling $\dim M_1$ of the $n-1$ tachyons and 1 tensor HSSA calculated previously. Moreover, we propose a set of K-identities which is the key to demonstrate the stringy scaling behavior of HSSA. We explicitly prove both the diagonal and off-diagonal K-identities for the 4-point $HSSA$ and give numerical proofs of these K-identities for some higher point HSSA.

[∗]Electronic address: xgcj944137@gmail.com

[†]Electronic address: jcclee@cc.nctu.edu.tw

[‡]Electronic address: yiyang@mail.nctu.edu.tw

I. INTRODUCTION

Recently, it was discovered [\[1](#page-26-0)[–3](#page-26-1)] that the degree of freedom or number of kinematic variables reduced for the n-point with $n-1$ tachyons and 1 tensor hard string scattering amplitudes (HSSA) and the Regge string scattering amplitudes (RSSA). Historically, the first stringy scaling behavior was conjectured by Gross [\[4](#page-26-2)[–7\]](#page-26-3) for the 4-point HSSA and later explicitly proved in $[8-11]$. Indeed, all functional forms of 4-point $HSSA$ at each fixed mass level are found to be proportional to each other with constant ratios (independent of the scattering angle ϕ , or the deficit of the kinematics variable dim $\mathcal{M}_1 = 1 - 0 = 1$). See the recent reviews [\[12](#page-26-6), [13\]](#page-26-7).

This stringy scaling behavior, similar to the QCD Bjorken scaling [\[14](#page-26-8), [15](#page-27-0)] and its $GLAP$ correction [\[16](#page-27-1), [17\]](#page-27-2), was recently generalized to arbitrary n-point $(n-1)$ tachyons and 1 tensor) $HSSA$ with $n \geq 5$ [\[1](#page-26-0), [2](#page-26-9)]. As an example, the number of degree of freedom of 6-point HSSA (5 tachyons and 1 tensor) with 3 transverse directions reduced from 8 to 2, and dim $M_1 = 8 - 2 = 6$.

Since massive higher spin tensor string states receive more degree of freedom of spin polarizations, it is tempted to speculate that the degree of stringy scaling or dim \mathcal{M}_k ($k \leq n$) for the *n*-point *k*-tensor $HSSA$ with $k \geq 2$ may decrease and is smaller than $\dim \mathcal{M}_1$. In this paper, we calculate n-point ($n \geq 4$) HSSA with $n-2$ tachyons and 2 high energy tensor states at arbitrary mass levels. It is found that the degree of stringy scaling $\dim {\mathcal M}_2$ for HSSA with more than 2 transverse directions or $r \geq 3$ (see Eq.[\(4.10\)](#page-18-0)) indeed decreases comparing to that of the 1 tensor $HSSA$ dim \mathcal{M}_1 calculated previously.

In the saddle-point calculation of the $HSSA$, the saddle-point of the integration can be exactly solved only for some lower point HSSA calculation. For the cases of general n-point saddle-point $HSSA$ calculation, we propose a set of the so-called K-identities. Without directly solving the saddle-point equation, these K -identities can be used to simplify the HSSA calculation, and turn out to be the key step of the calculation to demonstrate the stringy scaling behavior.

Some of these K-identities were originally proposed for the calculation of n -point 1 tensor HSSA [\[1](#page-26-0), [2](#page-26-9)]. They are the direct results of the identification of two different calculation of $HSSA$, namely, the saddle-point calculation and the decoupling of zero-norm state (ZNS) calculation [\[12](#page-26-6), [13\]](#page-26-7). However, only K-identities for the 4-point $HSSA$ with 1 and 2 tensors will be analytically proved in this paper, other K-identities of higher point ($n \geq 5$) HSSA calculation will be tested numerically. Presumably, these K-identities are results of conservation of momenta in the hard scattering limit at the saddle-point.

For the saddle-point calculation of *n*-point 2 tensor $HSSA$, in addition to the diagonal K-identities one obtained during the calculation of n-point 1 tensor $HSSA$, it turns out that one needs to introduce1 more off-diagonal K -identity. This off-diagonal K -identity will contribute to the degree of stringy scaling $\dim M_2$ for $HSSA$ out of the scattering plane. It is conjectured that for general higher k tensor $HSSA$ calculation ($k \leq n$), these off-diaganal K-identities are crucial to show the consistent result that $\dim \mathcal{M}_k \geq 0$.

This paper is organized as following. In section II, we first calculate an example of 4 point 2 tensor HSSA for mass level $M^2 = 2$. We then extend the calculation to arbitrary mass levels and obtain the ratios consistent with calculation from the method of decoupling of ZNS . In section III, we calculate general *n*-point 2 tensor $HSSA$. One key step is to propose both the diagonal and off-diagonal K-identities to simplify the calculation and obtain the consistent ratios calculated from the method of ZNS. In section IV, we discuss in details the K -identities and apply them to the calculation of the degree of stringy scaling $\dim M_2$. We also explicitly verify all 4-point 1 and 2 tensors K-identities and discuss the numerical check of the higher point K -identities. A brief conclusion is given in section V.

II. THE FOUR-POINT TWO TENSOR HSSA

In this section, we will use the saddle-point method to calculate the 4-point $HSSA$ with 2 tachyons and 2 tensors. We begin with a brief review of the calculation of $HSSA$ with 3 tachyons and 1 tensor state [\[8](#page-26-4)[–11\]](#page-26-5). For the 4-point HSSA, we first note that in the hard scattering limit, components of polarization orthogonal to the scattering plane are subleading order in energy. Defining $e^P = \frac{1}{M}$ $\frac{1}{M_2}(E_2, \mathbf{k}_2, \vec{0}) = \frac{k_2}{M_2}$ the momentum polarization, $e^L = \frac{1}{M}$ $\frac{1}{M_2}$ (k₂, E_2 , $\vec{0}$) the longitudinal polarization and the transverse polarization $e^T = (0, 0, 1)$ on the scattering plane, it can then be shown that at each fixed mass level $M^2 = 2(N - 1)$ only states of the following form [\[8](#page-26-4)[–11](#page-26-5)] (in the hard scattering limit $e^P \simeq e^L$)

$$
|N, 2m, q\rangle = (\alpha_{-1}^{T})^{N-2m-2q} (\alpha_{-1}^{L})^{2m} (\alpha_{-2}^{L})^{q} |0; k\rangle
$$
 (2.1)

are leading order in energy. Note in particular that states with $(\alpha_{-1}^L)^{2m+1}$ are of subleading order in energy. To achieve BRST invariance, it can be shown that states in the physical state basis (PSB) can be written as a linear combination of states in the oscillator state basis (OSB) in Eq.[\(2.1\)](#page-6-0) [\[12](#page-26-6), [13](#page-26-7)].

There are two different methods to calculate the stringy scaling behavior of HSSA, the method of decoupling of ZNS and the saddle-point method. Both methods lead to constant ratios among 4-point HSSA at each fixed mass level [\[8](#page-26-4)[–11\]](#page-26-5)

$$
\frac{A^{(N,2m,q)}}{A^{(N,0,0)}} = \frac{(2m)!}{m!} \left(\frac{-1}{2M}\right)^{2m+q}.
$$
 (independent of ϕ !). (2.2)

In Eq.[\(2.2\)](#page-6-1), $A^{(N,m,q)}$ is the 4-point HSSA of any string vertex V_j with $j = 1, 3, 4$ and V_2 is the high energy state in Eq.[\(2.1\)](#page-6-0). $A^{(N,0,0)}$ is the 4-point HSSA of any string vertex V_j with $j = 1, 3, 4$, and V_2 is the leading Regge trajectory string state at mass level N. Note that we have omitted the tensor indice of V_j with $j = 1, 3, 4$ and keep only those of V_2 in $\mathcal{T}^{(N,2m,q)}$.

For the j tensor cases with $j = 2, 3$ and 4, it is easy to argue from the method of the decoupling of ZNS that the ratios are j products of ratio in Eq.[\(2.2\)](#page-6-1). However, it is highly nontrivial to calculate *n*-point multi-tensor $HSSA$ at arbitrary mass levels. Moreover, as will will see in section IV that one important step of this saddle-point calculation is to introduce the K vectors and K-identities, and give information to calculate the degree of stringy scaling dimM.

For later reference, we begin with a brief description of the 4-point saddle-point calculation with 3 tachyons and 1 tensor state here. The $t - u$ channel of $HSSA$ with 3 tachyons and 1 high energy state in Eq.[\(2.1\)](#page-6-0) can be calculated to be (after $SL(2, R)$ fixing)

$$
A^{(N,2m,q)} = \int_1^{\infty} dz z^{(1,2)} (1-z)^{(2,3)} \left[\frac{e^T \cdot k_1}{z} - \frac{e^T \cdot k_3}{1-z} \right]^{N-2m-2q}
$$

$$
\cdot \left[\frac{e^P \cdot k_1}{z} - \frac{e^P \cdot k_3}{1-z} \right]^{2m} \left[-\frac{e^P \cdot k_1}{z^2} - \frac{e^P \cdot k_3}{(1-z)^2} \right]^q
$$
(2.3)

where $(1, 2) = k_1 \cdot k_2$ etc. We can rewrite the amplitude above into the following form

$$
A^{(N,2m,q)}(K) = \int_{1}^{\infty} dz \ u(z) e^{-Kf(z)}
$$
\n(2.4)

where

$$
\Lambda \equiv -(1,2) \rightarrow \frac{s}{2} \rightarrow 2E^2,\tag{2.5}
$$

$$
\tau \equiv -\frac{(2,3)}{(1,2)} \to -\frac{t}{s} \to \sin^2 \frac{\phi}{2},\tag{2.6}
$$

$$
f(z) \equiv \ln z - \tau \ln(1 - z),\tag{2.7}
$$

$$
u(z) \equiv \left[\frac{(1,2)}{M}\right]^{2m+q} (1-z)^{-N+2m+2q} \underbrace{(f')^{2m}}_{*} (f'')^{q} (-e^T \cdot k_3)^{N-2m-2q}.
$$
 (2.8)

The saddle-point $z = \tilde{z}$, is defined by

$$
f'(\tilde{z}) = 0,\t\t(2.9)
$$

which can be exactly solved to be

$$
\tilde{z} = \frac{1}{1 - \tau} = \sec^2 \frac{\phi}{2}.
$$
\n(2.10)

It is now easy to see that

$$
u(\tilde{z}) = u'(\tilde{z}) = \dots = u^{(2m-1)}(\tilde{z}) = 0,
$$
\n(2.11)

and

$$
u^{(2m)}(\tilde{z}) = \left[\frac{(1,2)}{M}\right]^{2m+q} (1-\tilde{z})^{-N+2m+2q} (2m)! (f_0'')^{2m+q} (-e^T \cdot k_3)^{N-2m-2q}.
$$
 (2.12)

It is important to note that in calculating $HSSA$ containing high energy states in Eq.[\(2.1\)](#page-6-0) with $(\alpha_{-1}^L)^{2m}$, the *naive* energy order of the amplitudes based on the naive mass counting vanishes [\[8](#page-26-4)[–11](#page-26-5)], and the real leading energy order HSSA will in general drop by an even number of energy power as can be seen from Eq.[\(2.11\)](#page-8-0). On the other hand, $HSSA$ corresponding to states with $(\alpha_{-1}^L)^{2m+1}$ turn out to be of subleading order in energy.

With these inputs, one can evaluate the Gaussian integral in Eq.[\(2.4\)](#page-6-2) and show explicitly that the angular dependence of $HSSA$ depends only on the level N but not m and q

$$
A^{(N,2m,q)} = \sqrt{\frac{2\pi}{Kf_0''}} e^{-\Lambda f_0} \left[(-1)^{N-q} \frac{2^{N-2m-q} (2m)!}{m! M^{2m+q}} \tau^{-\frac{N}{2}} (1-\tau)^{\frac{3N}{2}} E^N + O(E^{N-2}) \right], \quad (2.13)
$$

and we have $[8-11]$

$$
\lim_{E \to \infty} \frac{A^{(N,2m,q)}}{A^{(N,0,0)}} = \frac{(-1)^q (2m)!}{m!(2M)^{2m+q}} = \left(-\frac{2m-1}{M}\right) \dots \left(-\frac{3}{M}\right) \left(-\frac{1}{M}\right) \left(-\frac{1}{2M}\right)^{m+q}
$$
\n
$$
= \left(\frac{1}{2}\right)^{m+q} \left(-\frac{1}{M}\right)^{q+2m} (2m-1)!!,\tag{2.14}
$$

which is remarkably consistent with calculation of decoupling of high energy ZNS obtained in Eq. (2.2) .

We now turn to calculate the 4-point HSSA with 2 tachyons and 2 tensor states. We choose to put the 2 tensors on z_2 and z_3 . The kinematics of 4-points $HSSA$ can be written as

$$
k_1 = \left(E - E \ 0\right),
$$

\n
$$
k_2 = \left(E E \ 0\right),
$$

\n
$$
k_3 = \left(-E - E \cos \phi - E \sin \phi\right),
$$

\n
$$
k_4 = \left(-E E \cos \phi E \sin \phi\right),
$$

\n(2.15)

which imply

$$
k_{12} = -2E^2,
$$

\n
$$
k_{13} = E^2 (1 - \cos \phi),
$$

\n
$$
k_{23} = E^2 (1 + \cos \phi).
$$
\n(2.16)

For the 2 tensor cases, we will use the following short notations for polarizations

$$
T_2 = \begin{pmatrix} 0 & 0 & 1 \end{pmatrix},
$$

\n
$$
T_3 = \begin{pmatrix} 0 & -\sin\phi & \cos\phi \end{pmatrix},
$$

\n
$$
L_2 = \frac{1}{M_2} \begin{pmatrix} E & E & 0 \end{pmatrix},
$$

\n
$$
L_3 = \frac{1}{M_3} \begin{pmatrix} -E & -E\cos\phi & -E\sin\phi \end{pmatrix}
$$
\n(2.17)

where T_2 means e^{T_2} etc. One can easily calculate the following projections

$$
k_1^{T_2} = 0, \ k_3^{T_2} = -E\sin\phi, \ k_1^{T_3} = E\sin\phi, \ k_2^{T_3} = -E\sin\phi; \tag{2.18}
$$

$$
k_1^{L_2} = \frac{-2E^2}{M_2}, \ k_3^{L_2} = \frac{2E^2}{M_2}\tau, \ k_1^{L_3} = \frac{2E^2}{M_3}(1-\tau), \ k_2^{L_3} = \frac{2E^2}{M_3}\tau; \tag{2.19}
$$

and after $SL(2, R)$ fixing

$$
f_0'' = -\frac{1}{\tilde{z}^2} + \frac{\tau}{(1-\tilde{z})^2} = \frac{(1-\tau)^3}{\tau}.
$$
\n(2.20)

A. An example of mass level $M^2 = 2$

As a simple example of 4-point HSSA with 2 tachyons and 2 tensor states, we first calculate HSSA for tensor states at mass level $N = 2$. By Eq.[\(2.1\)](#page-6-0), there are 3 high energy states in the OSB

$$
\left(\alpha_{-1}^{T}\right)^{2}\left|0;k\right\rangle, \quad \left(\alpha_{-2}^{L}\right)\left|0;k\right\rangle, \quad \left(\alpha_{-1}^{L}\right)^{2}\left|0;k\right\rangle. \tag{2.1}
$$

By Eq.[\(2.2\)](#page-6-1), the ratios of $HSSA$ with 3 tachyons and 1 tensor in Eq.[\(2.1\)](#page-6-0) are

$$
A^{T_2 T_2} : A^{L_2} : A^{L_2 L_2} = 4 : \sqrt{2} : 1.
$$
\n(2.2)

For the case of HSSA with 2 tachyons and 2 tensors, the calculation of the decoupling of zero-norm state (ZNS) gives

$$
(T_2T_2 \oplus L_2 \oplus L_2L_2) \otimes (T_3T_3 \oplus L_3 \oplus L_3L_3)
$$

=
$$
(4:\sqrt{2}:1) \times (4:\sqrt{2}:1).
$$
 (2.3)

We will recalculate the ratios by using the saddle-point method. In calculating the $HSSA$, one needs to pay attention to the state $(\alpha_{-1}^L)^2 |0; k\rangle$ with $m = 1$ in Eq.[\(2.1\)](#page-6-0), as has been mentioned in the paragraph after Eq.[\(2.12\)](#page-8-1). The calculation for $HSSA$ with $m = 0$ are straightforward and we obtain

$$
A^{T_2 T_2; T_3 T_3} = \int d^4 z e^{-\Lambda f} \left(\frac{k_1^{T_2}}{z_1 - z_2} + \frac{k_3^{T_2}}{z_3 - z_2} + \frac{k_4^{T_2}}{z_4 - z_2} \right)^2 \left(\frac{k_1^{T_3}}{z_1 - z_3} + \frac{k_2^{T_3}}{z_2 - z_3} + \frac{k_4^{T_3}}{z_4 - z_3} \right)^2
$$

= $e^{-\Lambda f_0} \left(4E^2 \frac{(1 - \tau)^3}{\tau} \right) \left(4E^2 \frac{1 - \tau}{\tau} \right) \sqrt{\frac{2\pi}{f_0''}},$ (2.4)

$$
A^{T_2 T_2; L_3} = \int d^4 z e^{-\Lambda f} \left(\frac{k_1^{T_2}}{z_1 - z_2} + \frac{k_3^{T_2}}{z_3 - z_2} + \frac{k_4^{T_2}}{z_4 - z_2} \right)^2 \left(\frac{k_1^{L_3}}{(z_1 - z_3)^2} + \frac{k_2^{L_3}}{(z_2 - z_3)^2} + \frac{k_4^{L_3}}{(z_4 - z_3)^2} \right)
$$

= $e^{-\Lambda f_0} \left(4E^2 \frac{(1 - \tau)^3}{\tau} \right) \left(\frac{2E^2 (1 - \tau)}{M_3 \tau} \right) \sqrt{\frac{2\pi}{f_0''}}.$ (2.5)

With $M_3 = \sqrt{2}$, Eq.[\(2.4\)](#page-6-2) and Eq.[\(2.5\)](#page-6-3) give the consistent ratios in Eq.[\(2.2\)](#page-6-1).

We now turn to calculate $HSSA$ containing state $(\alpha_{-1}^L)^2 |0; k\rangle$. The saddle-point calculation gives

$$
A^{T_2 T_2; L_3 L_3} = \int d^4 z e^{-\Lambda f} \left(\frac{k_1^{T_2}}{z_1 - z_2} + \frac{k_3^{T_2}}{z_3 - z_2} + \frac{k_4^{T_2}}{z_4 - z_2} \right)^2 \left(\frac{k_1^{L_3}}{z_1 - z_3} + \frac{k_2^{L_3}}{z_2 - z_3} + \frac{k_4^{L_3}}{z_4 - z_3} \right)^2
$$

=
$$
\int dz_2 e^{-\Lambda f} \left(-E \sin \phi \frac{\tau - 1}{\tau} \right)^2 \left(\frac{\frac{2E^2}{M_3} \cos^2 \frac{\phi}{2}}{-1} + \frac{\frac{2E^2}{M_3} \sin^2 \frac{\phi}{2}}{z_2 - 1} \right)^2.
$$
 (2.6)

We note that the above integrand vanish if we evaluate z_2 at the saddle point $z_2 = \tilde{z}_2$. To obtain the real energy order HSSA, we need to do Tayor expansion about the saddle point \tilde{z}_2 before performing the integration. By doing so, we obtain

$$
A^{T_2 T_2; L_3 L_3} = \int dz_2 e^{-\Lambda f} \left(-2E\sqrt{\tau (1-\tau)} \frac{\tau - 1}{\tau} + \cdots \right)^2
$$

$$
\cdot \left(-\frac{2E^2}{M_3} (1-\tau) + \frac{2E^2}{M_3} \tau \frac{1-\tau}{\tau} - \frac{\frac{2E^2}{M_3} \sin^2 \frac{\phi}{2}}{(z_{20}-1)^2} (z_2 - \tilde{z}_2) + \cdots \right)^2
$$

$$
= \int dz_2 e^{-\Lambda f} \left(-2E\sqrt{\tau (1-\tau)} \frac{\tau - 1}{\tau} + \cdots \right)^2 \left(0 - \frac{\frac{2E^2}{M_3} \sin^2 \frac{\phi}{2}}{(z_{20}-1)^2} (z_2 - \tilde{z}_2) + \cdots \right)^2
$$

$$
= e^{-\Lambda f_0} \left(4E^2 \frac{(1-\tau)^3}{\tau} \right) \left(\frac{2E^2}{M_3^2} \frac{(1-\tau)}{\tau} \right) \sqrt{\frac{2\pi}{f_0''}}, \tag{2.7}
$$

which gives the ratio consistent with Eq.[\(2.2\)](#page-6-1). Similar calculation can be performed for other $HSSA$ in Eq.[\(2.2\)](#page-6-1) and all consistent ratios can be obtained except the $A^{L_2L_2;L_3L_3}$ one which needs to be treated carefully.

We first note that in calculating HSSA in Eq.[\(2.3\)](#page-6-4), contraction terms from $\partial_2 X_2 \cdot \cdot$ and $\partial_3 X_3 \cdot \cdot$ are considered to be of subleading order in energy and can be ignored. However, for the special case of $A^{L_2L_2;L_3L_3}$ amplitude, one encounters the naive energy order problem, and it turns out that one has to include contraction terms from $\partial_2 X_2 \cdots$ and $\partial_3 X_3 \cdots$ in order to achieve the consistent ratios calculated by the method of decoupling of ZNS in Eq.[\(2.3\)](#page-6-4). We thus follow the standard KLT [\[18\]](#page-27-3) method to calculate the amplitude

$$
A^{L_2L_2;L_3L_3} = \int d^4 z e^{ik_1 X_1} e^{ik_2 X_2 + i\xi_2 \partial_2 X_2 + i\xi_2' \partial_2 X_2} e^{ik_3 X_3 + i\xi_3 \partial_3 X_3 + i\xi_3' \partial_3 X_3} e^{ik_4 X_4}
$$
\n
$$
= \int d^4 z e^{-\Lambda f} \exp \left\{ \begin{array}{c} \left[\frac{k_1 \cdot \xi_2}{z_1 - z_2} + \frac{k_3 \cdot \xi_2}{z_3 - z_2} \right] + \left[\frac{k_1 \cdot \xi_2'}{z_1 - z_3} + \frac{k_2 \cdot \xi_3'}{z_2 - z_3} \right] \\ + \left[\frac{k_1 \cdot \xi_3}{z_1 - z_3} + \frac{k_2 \cdot \xi_3'}{z_2 - z_3} \right] + \left[\frac{k_1 \cdot \xi_3'}{z_1 - z_2} + \frac{k_2 \cdot \xi_3'}{z_2 - z_3} \right] \end{array} \right\} \text{ multilinear}
$$
\n
$$
= \int d^4 z e^{-\Lambda f} \left\{ \begin{array}{c} \left[\frac{k_1 \cdot \xi_2}{z_1 - z_2} + \frac{k_3 \cdot \xi_2}{z_2 - z_3} \right] + \left[\frac{k_1 \cdot \xi_2'}{z_1 - z_2} + \frac{\xi_2' \cdot \xi_3}{z_3 - z_2} \right] + \frac{k_2 \cdot \xi_3'}{(z_3 - z_2)^2} + \frac{\xi_2' \cdot \xi_3'}{(z_3 - z_2)^2} \right] \frac{k_1 \cdot \xi_3'}{z_1 - z_3} + \frac{k_2 \cdot \xi_3'}{z_2 - z_3} \end{array} \right\}
$$
\n
$$
= \int d^4 z e^{-\Lambda f} \left\{ \begin{array}{c} \left[\frac{k_1 \cdot \xi_2}{z_1 - z_2} + \frac{k_3 \cdot \xi_2}{z_3 - z_2} \right] \left[\frac{k_1 \cdot \xi_2'}{z_1 - z_2} + \frac{k_2 \cdot \xi_2'}{z_3 - z_2} \right] \left[\frac{k_1 \cdot \xi_3'}{z_1 - z_3} + \frac{k_2 \cdot \xi
$$

The result in Eq.[\(2.10\)](#page-8-2) includes contraction terms from $\partial_2 X_2 \cdots$ and $\partial_3 X_3 \cdots$. By setting $\xi_2 = \xi'_2 = L_2$ and $\xi_3 = \xi'_3 = L_3$ in Eq.[\(2.10\)](#page-8-2), we obtain

$$
A^{L_2 L_2; L_3 L_3} = \int d^4 z e^{-\Lambda f} \left\{ \begin{array}{l} \left[\frac{k_1^{L_2}}{z_1 - z_2} + \frac{k_2^{L_2}}{z_3 - z_2} \right]^2 \left[\frac{k_1^{L_3}}{z_1 - z_3} + \frac{k_2^{L_3}}{z_2 - z_3} \right]^2 \right. \\ \left. + \frac{4L_2 \cdot L_3}{(z_3 - z_2)^2} \left[\frac{k_1^{L_2}}{z_1 - z_2} + \frac{k_3^{L_2}}{z_3 - z_2} \right] \left[\frac{k_1^{L_3}}{z_1 - z_3} + \frac{k_2^{L_3}}{z_2 - z_3} \right] \right\} \\ \left. + \frac{2(L_2 \cdot L_3)^2}{(z_3 - z_2)^4} \right\} \\ \simeq \int dz_2 e^{-\Lambda f} \left\{ \begin{array}{l} \left(0 + \frac{2E^2}{M_2} \left(\frac{(1-\tau)^3}{\tau} \right) (z_2 - z_{20}) + \cdots \right)^2 \left(0 - \frac{2E^2}{M_3} \frac{(1-\tau)^2}{\tau} (z_2 - z_{20}) + \cdots \right)^2 \right. \\ \left. + \frac{4\left(\frac{2E^2 \tau}{M_2 M_3} \right)}{(1 - \frac{1}{1 - \tau})^2} \left(0 + \frac{2E^2}{M_2} \left(\frac{(1-\tau)^3}{\tau} \right) (z_2 - z_{20}) + \cdots \right) \right. \\ \left. + \left(0 - \frac{2E^2}{M_3} \frac{(1-\tau)^2}{\tau} (z_2 - z_{20}) + \cdots \right) \right. \\ \left. + \frac{2\left(\frac{2E^2 \tau}{M_2 M_3} \right)^2}{(1 - \frac{1}{1 - \tau})^4} \right\} \end{array} \right\} \tag{2.12}
$$

.

In Eq.[\(2.12\)](#page-8-1) we have done Taylor expansion about the saddle point z_{20} for the first two terms and have put 0 for each naive energy order amplitude [\[8](#page-26-4)[–11\]](#page-26-5). Most importantly, the last two terms of $Eq. (2.12)$ $Eq. (2.12)$ involve crossing contraction terms and will be seen soon that they are indeed of the same energy order as the first term in $Eq.(2.12)$ $Eq.(2.12)$ which we have done Taylor expansion about the saddle point \tilde{z}_2 due to the naive energy order problem.

Finally, we perform the integration in $Eq.(2.12)$ $Eq.(2.12)$ to obtain

$$
A^{L_2 L_2; L_3 L_3} \simeq \int dz_2 e^{-\Lambda f} \begin{Bmatrix} \left(\frac{2E^2}{M_2}\right)^2 \left(\frac{2E^2}{M_3}\right)^2 \frac{(1-\tau)^{10}}{\tau^4} (z_2 - \tilde{z}_2)^4\\ -4 \left(\frac{2E^2}{M_2}\right)^2 \frac{2E^2}{M_3^2} \frac{(1-\tau)^7}{\tau^3} (z_2 - \tilde{z}_2)^2\\ +2 \frac{2E^2}{M_2^2} \frac{2E^2}{M_3^2} \frac{(1-\tau)^4}{\tau^2} \end{Bmatrix}
$$
(2.13)

$$
= e^{-\Lambda f_0} \begin{cases} \left(\frac{2E^2}{M_2}\right)^2 \left(\frac{2E^2}{M_3}\right)^2 \frac{(1-\tau)^{10}}{\tau^4} \left(\frac{-2}{1}\right)^2 \left(\frac{-1}{2}\right) \left(\frac{-3}{2}\right) \left(\frac{1}{f_0''}\right)^2 \sqrt{\frac{2\pi}{f_0''}}\\ -4 \left(\frac{2E^2}{M_2}\right)^2 \frac{2E^2}{M_3^2} \frac{(1-\tau)^7}{\tau^3} \left(\frac{-2}{1}\right) \left(\frac{-1}{2}\right) \frac{1}{f_0''} \sqrt{\frac{2\pi}{f_0''}}\\ +2 \frac{2E^2}{M_2^2} \frac{2E^2}{M_3^2} \frac{(1-\tau)^4}{\tau^2} \sqrt{\frac{2\pi}{f_0''}} \end{cases}
$$
(2.14)

$$
= e^{-\Lambda f_0} \frac{2E^2}{M_2^2} \frac{2E^2}{M_3^2} \frac{(1-\tau)^4}{\tau^2} \sqrt{\frac{2\pi}{f_0''}}.
$$
\n(2.15)

which gives the ratio consistent with Eq. (2.3) . This completes the saddle-point calculation of HSSA in Eq.[\(2.3\)](#page-6-4). We conclude that for the 4-point 2 tachyons 2 tensor HSSA, the degree of stringy scaling $\dim M_2 = 1$ is the same as that of 3 tachyons 1 tensor $HSSA$ $\dim \mathcal{M}_1 = 1$ calculated previously. However, as we will see in the next section, for the higher point $HSSA$, dim \mathcal{M}_2 will be in general smaller than dim \mathcal{M}_1 .

B. General mass levels

In this subsection, we extend our previous result to two general mass levels $M_i^2 = 2(N_i -$ 1), $i = 2, 3$. We first define the K-vectors

$$
K_i = \sum_{j \neq i} \frac{-k_j}{z_j - z_i} = (K_i^{L_i}, K_i^{T_i}),
$$
\n(2.16)

which will be important in our following discussion. By setting $z_1 = 0, z_4 = \infty$ in

$$
k_{12}f = \sum_{i>j} k_{ij} \ln (z_i - z_j), \qquad (2.17)
$$

we obtain

$$
k_{12}f = k_{12}\ln z_2 + k_{13}\ln z_3 + k_{23}\ln (z_3 - z_2). \tag{2.18}
$$

At the saddle point

$$
k_{12}\frac{\partial f}{\partial z_2} = \frac{k_{12}}{z_2} - \frac{k_{23}}{z_3 - z_2} = M_2 K_2^{L_2} = 0,
$$
\n(2.19)

$$
k_{12}\frac{\partial f}{\partial z_3} = \frac{k_{13}}{z_3} + \frac{k_{23}}{z_3 - z_2} = M_3 K_3^{L_3} = 0.
$$
 (2.20)

After setting $z_3 = 1$, we get

$$
K_2^{L_2} = -\frac{2E^2}{M_2} \frac{(1-\tau)^3}{\tau} (z_2 - \tilde{z}_2), \qquad (2.21)
$$

$$
K_3^{L_3} = \frac{2E^2}{M_3} \frac{(1-\tau)^2}{\tau} (z_2 - \tilde{z}_2).
$$
 (2.22)

One important property of the K_j vector is that its longitudinal component $K_j^{L_j}$ j^{L_j} is proportional to $\frac{\partial f}{\partial z_j}$ and thus vanishes at the saddle point

$$
\tilde{K}_j^{L_j} = 0,\t\t(2.23)
$$

The K -identities $[1-3]$ can then be written as

$$
\tilde{K}_j^2 + 2M_j \partial_j \tilde{K}_j^{L_j} = 0. \tag{2.24}
$$

We will discuss these K-identities in section IV.

The 4-point $HSSA$ with 2 tachyons and 2 tensors of Eq.[\(2.1\)](#page-6-0),

$$
|p_i, 2m_i, q_i\rangle = \left(\alpha_{-1}^{T_i}\right)^{p_i} \left(\alpha_{-1}^{L_i}\right)^{2m_i} \left(\alpha_{-2}^{L_i}\right)^{q_i} |0, k\rangle, i = 2, 3 \tag{2.25}
$$

with the mass levels

$$
N_i = p_i + 2m_i + 2q_i \text{ and } N = N_2 + N_3,
$$
\n(2.26)

can be calculated to be

$$
A^{(p_i, 2m_i, q_i)} = \int d^4 z_2 e^{-\Lambda f} \left(K_2^{T_2}\right)^{p_2} \left(K_3^{T_3}\right)^{p_3} \left(-\partial_2 K_2^{L_2}\right)^{q_2} \left(-\partial_3 K_3^{L_3}\right)^{q_3}
$$

\n
$$
\cdot \exp\left\{\sum_{j_2=1}^{2m_2} K_2 \cdot \xi_{2, j_2} + \sum_{j_3=1}^{2m_3} K_3 \cdot \xi_{3, j_3} + \sum_{j_2=1}^{2m_2} \sum_{j_3=1}^{2m_3} \left[\frac{\xi_{2, j_2} \cdot \xi_{3, j_3}}{(z_3 - z_2)^2}\right]\right\}_{multiplicar}
$$

\n
$$
= \int d^4 z e^{-\Lambda f} \left(K_2^{T_2}\right)^{p_2} \left(K_3^{T_3}\right)^{p_3} \left(-\partial_2 K_2^{L_2}\right)^{q_2} \left(-\partial_3 K_3^{L_3}\right)^{q_3}
$$

\n
$$
\sum_{j=0}^{2m} j! C_j^{2m_2} C_j^{2m_3} \left[\frac{L_2 \cdot L_3}{(z_3 - z_2)^2}\right]^j \left(K_2^{L_2}\right)^{2m_2 - j} \left(K_3^{L_3}\right)^{2m_3 - j} \tag{2.27}
$$

where in the last line we have set $\xi_{2,j} = L_2, \xi_{3,j} = L_3$ and $m = \min(m_2, m_3)$. Plugging in the kenematics, we have

$$
A^{(p_i,2m_i,q_i)} = \left(\frac{4E^2\left(1-\tau\right)^3}{\tau}\right)^{N_2/2+m_2+m_3} \left(\frac{4E^2\left(1-\tau\right)}{\tau}\right)^{N_3/2} \sum_{j=0}^{2m} j! C_j^{2m_2} C_j^{2m_3} \left[\frac{-\tau}{2E^2\left(1-\tau\right)^3}\right]^j
$$

$$
\cdot \left(\frac{-1}{2M_2}\right)^{2m_2+q_2} \left(\frac{-1}{2M_3}\right)^{2m_3+q_3} \int dz_2 e^{-\Lambda f} \left(z_2 - \tilde{z}_2\right)^{2m_2+2m_3-2j} . \tag{2.28}
$$

We can now perform the integration to get

$$
A^{(p_i,2m_i,q_i)} = \left(\frac{4E^2\left(1-\tau\right)^3}{\tau}\right)^{N_2/2+m_2+m_3} \left(\frac{4E^2\left(1-\tau\right)}{\tau}\right)^{N_3/2} \sum_{j=0}^{2m} j! C_j^{2m_2} C_j^{2m_3} \left[\frac{-\tau}{2E^2\left(1-\tau\right)^3}\right]^j
$$

$$
\cdot \left(\frac{-1}{2M_2}\right)^{2m_2+q_2} \left(\frac{-1}{2M_3}\right)^{2m_3+q_3} \left(\frac{-2d}{df_0''}\right)^{m_2+m_3-j} \sqrt{\frac{2\pi}{f_0''}}.
$$
(2.29)

After differentiation, we obtain

$$
A^{(p_i,2m_i,q_i)} = \left(\frac{4E^2(1-\tau)^3}{\tau}\right)^{N_2/2+m_2+m_3} \left(\frac{4E^2(1-\tau)}{\tau}\right)^{N_3/2} \sum_{j=0}^{2m} j! C_j^{2m_2} C_j^{2m_3} \left[\frac{-\tau}{2E^2(1-\tau)^3}\right]^j
$$

$$
\cdot \left(\frac{-1}{2M_2}\right)^{2m_2+q_2} \left(\frac{-1}{2M_3}\right)^{2m_3+q_3} \sqrt{\frac{2\pi}{f_0''}} \frac{[2(m_2+m_3-j)]!}{2^{m_2+m_3-j} (m_2+m_3-j)!} (f_0'')^{-(m_2+m_3-j)}
$$

$$
= \sqrt{\frac{2\pi}{f_0''}} \left(\frac{4E^2(1-\tau)^3}{\tau}\right)^{N_2/2} \left(\frac{4E^2(1-\tau)}{\tau}\right)^{N_3/2} \left(\frac{-1}{2M_2}\right)^{2m_2+q_2} \left(\frac{-1}{2M_3}\right)^{2m_3+q_3}
$$

$$
\cdot \sum_{j=0}^{2m} j! C_j^{2m_2} C_j^{2m_3} (-2)^j \frac{[2(m_2+m_3-j)]!}{(m_2+m_3-j)!}.
$$
(2.30)

Surprisingly, the summation in the last equation of Eq.[\(2.30\)](#page-11-0) can be calculated to be

$$
\sum_{j=0}^{2m} j! C_j^{2m_2} C_j^{2m_3} (-2)^j \frac{[2(m_2 + m_3 - j)]!}{(m_2 + m_3 - j)!} = \Gamma\left(m_2 + \frac{1}{2}\right) \Gamma\left(m_3 + \frac{1}{2}\right),\tag{2.31}
$$

and finally the 4-point 2 tensor HSSA is

$$
A^{(p_i,2m_i,q_i)} = \sqrt{\frac{2\pi}{f_0''}} \left(\frac{4E^2 (1-\tau)^3}{\tau} \right)^{N_2/2} \left(\frac{4E^2 (1-\tau)}{\tau} \right)^{N_3/2} \n\cdot \left(\frac{-1}{2M_2} \right)^{2m_2+q_2} \left(\frac{-1}{2M_3} \right)^{2m_3+q_3} \Gamma \left(m_2 + \frac{1}{2} \right) \Gamma \left(m_3 + \frac{1}{2} \right) \n= \sqrt{\frac{2\pi}{f_0''}} \left(\frac{4E^2 (1-\tau)^3}{\tau} \right)^{N_2/2} \left(\frac{4E^2 (1-\tau)}{\tau} \right)^{N_3/2} \n\cdot \left[\left(\frac{1}{2} \right)^{m_2+q_2} \left(\frac{-1}{M_2} \right)^{2m_2+q_2} (2m_2 - 1)!! \right] \left[\left(\frac{1}{2} \right)^{m_3+q_3} \left(\frac{-1}{M_3} \right)^{2m_3+q_3} (2m_3 - 1)!! \right].
$$
\n(2.32)

One can now easily calculate the ratio

$$
\frac{A^{(\{p_2,p_3\},2m_2,2m_3,q_2,q_3)}}{A^{(\{N_1,N_2\},0,0)}} = \left[\left(\frac{1}{2} \right)^{m_2+q_2} \left(-\frac{1}{M_1} \right)^{q_2+2m_2} (2m_2-1)!! \right] \cdot \left[\left(\frac{1}{2} \right)^{m_3+q_3} \left(-\frac{1}{M_2} \right)^{q_3+2m_3} (2m_3-1)!! \right],
$$
\n(2.33)

which is consistent with the calculation of ZNS method in Eq.[\(2.2\)](#page-6-1).

III. THE n-POINT TWO TENSOR HSSA

In this section, we calculate the general n-point $HSSA$ with $n-2$ tachyons and 2 tensors of Eq.[\(2.1\)](#page-6-0). For the cases of $n \geq 5$, one needs to consider scattering processes out of the scattering plane with r -dimensional transverse space. We begin with the case of $HSSA$ with $n-1$ tachyons and 1 tensor. The general high energy states at each fixed mass level $M^2 = 2(N - 1)$ can be written as

$$
|\{p_i\}, 2m, q\rangle = \left(\alpha_{-1}^{T_1}\right)^{N+p_1} \left(\alpha_{-1}^{T_2}\right)^{p_2} \cdots \left(\alpha_{-1}^{T_r}\right)^{p_r} \left(\alpha_{-1}^L\right)^{2m} \left(\alpha_{-2}^L\right)^{q} |0; k\rangle \tag{3.1}
$$

where $\sum_{i=1}^{r} p_i = -2(m+q)$ with $r \le 24$.

One generalizes the transverse polarization $e^T = (0, 0, 1)$ to $e^{\hat{T}} = (0, 0, \vec{\omega})$ where

$$
\omega_j = \cos \theta_j \prod_{\sigma=1}^{j-1} \sin \theta_\sigma, \text{ with } j = 1, \cdots, r, \ \theta_r = 0 \tag{3.2}
$$

are the solid angles in the transverse space spanned by r transverse directions e^{T_i} . For this case, the method of decoupling of ZNS leads to the ratios of *n*-point $HSSA$ [\[1](#page-26-0)[–3](#page-26-1)]

$$
\frac{A^{(\{p_i\},2m,q)}}{A^{(\{0_i\},0,0)}} = \frac{(2m)!}{m!} \left(\frac{-1}{2M}\right)^{2m+q} \prod_{j=1}^r \omega_j^{p_j} \ (\# \text{ of kinematic variables dependence reduced } !).
$$
\n(3.3)

These ratios are valid to all string loop orders, and for the string-tree level case, Eq.[\(3.3\)](#page-12-0) can be rederived from the saddle-point calculation [\[1](#page-26-0)[–3](#page-26-1)].

We are now ready to discuss the 2 tensor case. We first define

$$
L_i = \frac{k_i}{M_i}, i = 1, 2.
$$
\n(3.4)

For simplicity of the amplitude calculation, we choose to put 2 tensors at z_1 and z_2 . We consider 2 high energy states of Eq. (2.1) at vertex i

$$
|2m_i\rangle = \prod_{j=1}^r \left(\alpha_{-1}^{T_{ij}}\right)^{p_{ij}} \left(\alpha_{-1}^{L_i}\right)^{2m_i} \left(\alpha_{-2}^{L_i}\right)^{q_i} |0, k\rangle \tag{3.5}
$$

with the total mass level

$$
N_i = \sum_{j=1}^{r} p_{ij} + 2m_i + 2q_i
$$
\n(3.6)

where $i = 1, 2$ and $j = 1 \cdots r$.

The K_i vectors can be similarly defined as in the 4-point case in Eq.[\(2.16\)](#page-9-0)

$$
K_1 = \sum_{j \neq 1} \frac{-k_j}{z_j - z_1} = (K_1^{L_1}, K_1^{T_{11}}, K_1^{T_{12}}, \cdots, K_1^{T_{1r}}),
$$
\n(3.7)

$$
K_2 = \sum_{j \neq 2} \frac{-k_j}{z_j - z_2} = (K_2^{L_2}, K_2^{T_{21}}, K_2^{T_{22}}, \cdots, K_2^{T_{2r}}). \tag{3.8}
$$

Since the longitudinal component $K_i^{L_i}$ vanishes at the saddle point as in Eq.[\(2.23\)](#page-10-0), we can formally use two set of Eq.[\(3.2\)](#page-12-1), ω_1^j j_1 and ω_2^j $_2^j$ to parametrize the K vectors

$$
\tilde{K}_i^{T_{ij}} = \left| \tilde{K}_i \right| \omega_i^j, \ i = 1, 2 \text{ and } j = 1 \cdots r. \tag{3.9}
$$

We stress that not all kinematic variables of ω_1^j j_1 and ω_2^j $2₂$ are independent due to the various K-identities which put constraints on ω_i^j i_i . We will discuss this issue in section IV.

For the kinematics of *n*-point amplitudes in the hard scattering limit, $p = E \rightarrow \infty$, we define the 26-dimensional momenta in the CM frame to be

$$
k_1 = (E, -E, 0^r),
$$

\n
$$
k_2 = (E, +E, 0^r),
$$

\n
$$
k_j = (-q_j, -q_j \Omega_1^j, \dots - q_j \Omega_{25}^j)
$$
\n(3.10)

where $j = 3, 4, \cdots, n$, and

$$
\Omega_i^j = \cos \phi_i^j \prod_{\sigma=1}^{i-1} \sin \phi_\sigma^j \text{ with } \phi_{j-1}^j = 0, \ \phi_{i>r}^j = 0 \text{ and } r \le \min \{n-3, 24\} \tag{3.11}
$$

are the solid angles in the $(j-2)$ -dimensional spherical space with $\sum_{i=1}^{j-2}$ (Ω_i^j) $\binom{j}{i}^2 = 1$. In Eq.[\(3.10\)](#page-13-0), 0^r denotes the r-dimensional null vector. The condition $\phi_{j-1}^j = 0$ in Eq.[\(3.11\)](#page-13-1) was chosen to fix the frame by using the rotational symmetry. The independent kinematics variables can be chosen to be some φ_i^j i and some fixed ratios of infinite q_j . See section IV for the example of $n = 5$.

The HSSA of n-point string scattering with 2 general tensors at z_2 and z_1 can be expressed as

$$
A^{(\{p_{ij}\},2m_i,q_i)} = \int_0^1 dz_{n-2} \cdots \int_0^{z_4} dz_3 \int_0^{z_3} dz_2 \, u e^{-\Lambda f} \tag{3.12}
$$

where

$$
u = \left[\prod_{j=1}^{r} \left(K_2^{T_{2j}} \right)^{p_{2j}} \left(K_1^{T_{1j}} \right)^{p_{1j}} \right] \left(-\partial_2 K_2^{L_2} \right)^{q_2} \left(-\partial_1 K_1^{L_1} \right)^{q_1}
$$

$$
\cdot \exp \left\{ \sum_{j_2=1}^{2m_2} \left(K_2 \right) \cdot \xi_{2,j_2} + \sum_{j_1=1}^{2m_1} \left(K_1 \right) \cdot \xi_{1,j_1} + \sum_{j_2=1}^{2m_2} \sum_{j_1=1}^{2m_1} \left[\frac{\xi_{2,j_2} \cdot \xi_{1,j_1}}{\left(z_1 - z_2 \right)^2} \right] \right\}_{multiplace} . \tag{3.13}
$$

In the second line of the above equation, we can set $\xi_{2,k} = L_2, \xi_{1,k} = L_1$ and $m =$ $\min\left(m_2, m_1\right)$ to obtain

$$
u = (-1)^{q_1+q_2} \sum_{k=0}^{2m} \left[\prod_{j=1}^r \left(K_2^{T_{2j}} \right)^{p_{2j}} \left(K_1^{T_{1j}} \right)^{p_{1j}} \right] \left(\partial_2 K_2^{L_2} \right)^{q_2} \left(\partial_1 K_1^{L_1} \right)^{q_1} \n\cdot \left(K_2^{L_2} \right)^{2m_2-k} \left(K_1^{L_1} \right)^{2m_1-k} k! C_k^{2m_2} C_k^{2m_1} \left[\frac{k_{21}}{M_2 M_1 (z_1 - z_2)^2} \right]^k \n= (-1)^{q_1+q_2} \sum_{k=0}^{2m} u_k.
$$
\n(3.14)

Since $\tilde{K}_i^{L_i} = 0$, $i = 1, 2$ at the saddle-point, the leading order term in each term of the summation is

$$
\partial_2^{2m_2+2m_1-2k}\tilde{u}_k = C_{2m_1-k}^{2m_2+2m_1-2k} \left[\prod_{j=1}^r \left(\tilde{K}_2^{T_{2j}} \right)^{p_{2j}} \left(\tilde{K}_1^{T_{1j}} \right)^{p_{1j}} \right] \left(\partial_2 \tilde{K}_2^{L_2} \right)^{q_2+2m_2-k}
$$

$$
\left(\partial_1 \tilde{K}_1^{L_1} \right)^{q_1} \left(\partial_2 \tilde{K}_1^{L_1} \right)^{2m_1-k} (2m_2-k)!(2m_1-k)!k!C_k^{2m_2}C_k^{2m_1} \left[\frac{k_{21}}{M_2M_1 \left(\tilde{z}_1 - \tilde{z}_2 \right)^2} \right]^k \tag{3.15}
$$

where we have used

$$
\partial^{n}(fg) = \sum_{k=0}^{n} C_{k}^{n} f^{k} g^{n-k}.
$$
\n(3.16)

The next important step is to apply the diagonal and the off-diagonal K -identities

$$
\tilde{K}_i \tilde{K}_j + 2M_i \partial_j \tilde{K}_i^{L_i} = 0 \Rightarrow \begin{cases} \tilde{K}_i^2 + 2M_i \partial_i \tilde{K}_i^{L_i} = 0, \ i = j \\ \tilde{K}_i \tilde{K}_j + \frac{2k_{ij}}{(\tilde{z}_i - \tilde{z}_j)^2} = 0 \ i \neq j \end{cases}
$$
\n(3.17)

to obtain

$$
\partial_2^{2m_2+2m_1-2k}\tilde{u}_k = \left[\prod_{j=1}^r \left(\left|\tilde{K}_2\right|\omega_2^j\right)^{p_{2j}} \left(\left|\tilde{K}_1\right|\omega_1^j\right)^{p_{1j}}\right] \left(-\frac{\tilde{K}_2^2}{2M_2}\right)^{q_2+2m_2-k} \left(-\frac{\tilde{K}_1^2}{2M_1}\right)^{q_1} \left(-\frac{\tilde{K}_2\tilde{K}_1}{2M_1}\right)^{2m_1-k} (2m_2+2m_1-2k)!k!C_k^{2m_2}C_k^{2m_1} \left(-\frac{\tilde{K}_2\tilde{K}_1}{2M_2M_1}\right)^k.
$$
\n(3.18)

One can now use

$$
\tilde{K}_i^{T_{ij}} = \left| \tilde{K}_i \right| \omega_i^j,\tag{3.19}
$$

$$
N_i = \sum_{j=1}^{r} p_{ij} + 2m_i + 2q_i \text{ with } i = 1, 2 \text{ and } j = 1 \cdots r
$$
 (3.20)

to get

$$
\partial_{2}^{2m_{2}+2m_{1}-2k}\tilde{u}_{k} = \left[\prod_{j=1}^{r} \left(\left|\tilde{K}_{2}\right| \omega_{2}^{j}\right)^{p_{2j}} \left(\left|\tilde{K}_{1}\right| \omega_{1}^{j}\right)^{p_{1j}}\right] \left(-\frac{\tilde{K}_{2}^{2}}{2M_{2}}\right)^{q_{2}+2m_{2}} \cdot \left(-\frac{\tilde{K}_{1}^{2}}{2M_{1}}\right)^{q_{1}} \left(-\frac{\tilde{K}_{2}\tilde{K}_{1}}{2M_{1}}\right)^{2m_{1}} (2m_{2}+2m_{1}-2k)!k!C_{k}^{2m_{2}}C_{k}^{2m_{1}} \left(-\frac{\tilde{K}_{2}^{2}}{2}\right)^{-k} \left(3.21\right) \n= \left[\prod_{j=1}^{r} \left(\omega_{2}^{j}\right)^{p_{2j}} \left(\omega_{1}^{j}\right)^{p_{1j}}\right] \tilde{K}_{2}^{N_{2}+2m_{2}+2m_{1}} \tilde{K}_{1}^{N_{1}} \left(-\frac{1}{2M_{2}}\right)^{q_{2}+2m_{2}} \left(-\frac{1}{2M_{1}}\right)^{q_{1}+2m_{1}} \cdot (2m_{2}+2m_{1}-2k)!k!C_{k}^{2m_{2}}C_{k}^{2m_{1}} \left(-\frac{\tilde{K}_{2}^{2}}{2}\right)^{-k}.
$$
\n(3.22)

Finally, we can use the saddle-point method to evaluate the integral to obtain

$$
A^{(\{p_{ij}\},2m_i,q_i)} \simeq \int dz^{n-3} \sum_{k=0}^{2m} \frac{(-1)^{q_1+q_2} \partial_2^{2m_2+2m_1-2k} \tilde{u}_k}{(2m_2+2m_1-2k)!} (z_2 - \tilde{z}_2)^{2m_2+2m_1-2k} e^{-\Lambda \tilde{f}}
$$

\n
$$
\simeq e^{-\Lambda \tilde{f}} \sum_{k=0}^{2m} \frac{(-1)^{q_1+q_2} \partial_2^{2m_2+2m_1-2k} \tilde{u}_k}{(2m_2+2m_1-2k)!} \int_0^1 dz_2 (z_2 - \tilde{z}_2)^{2m_2+2m_1-2k} e^{-\frac{1}{4}\tilde{K}_2^2 (z_2-\tilde{z}_2)^2}
$$

\n
$$
= e^{-\Lambda f_0} \sum_{k=0}^{2m} \frac{(-1)^{q_1+q_2} \partial_2^{2m_2+2m_1-2k} \tilde{u}_k}{(2m_2+2m_1-2k)!} \left(-4\frac{d}{d\tilde{K}_2^2}\right)^{m_2+m_1-k} \sqrt{\frac{4\pi}{\tilde{K}_2^2}}.
$$
 (3.23)

After doing differentiation and putting in Eq.[\(3.22\)](#page-15-0), we obtain

$$
A^{(\{p_{ij}\},2m_i,q_i)} = e^{-\Lambda f_0} \sum_{k=0}^{2m} \frac{(-1)^{q_1+q_2} \partial_2^{2m_2+2m_1-2k} \tilde{u}_k}{(2m_2+2m_1-2k)!} \sqrt{\frac{4\pi}{\tilde{K}_2^2}} \frac{(2m_2+2m_1-2k)!}{(m_2+m_1-k)!} \left(\tilde{K}_2^2\right)^{-(m_2+m_1-k)}
$$

$$
= (-1)^{q_1+q_2} e^{-\Lambda f_0} \sqrt{\frac{4\pi}{\tilde{K}_2^2}} \tilde{K}_2^{N_2} \tilde{K}_1^{N_1} \left[\prod_{j=1}^r \left(\omega_2^j\right)^{p_{2j}} \left(\omega_1^j\right)^{p_{1j}}\right]
$$

$$
\cdot \left(-\frac{1}{2M_2}\right)^{q_2+2m_2} \left(-\frac{1}{2M_1}\right)^{q_1+2m_1} \sum_{k=0}^{2m} k! C_k^{2m_2} C_k^{2m_1} (-2)^k \frac{[2(m_2+m_1-k)]!}{(m_2+m_1-k)!}.
$$
(3.24)

We can now calculate the summation to get

$$
A^{(\{p_{ij}\},2m_i,q_i)} = e^{-\Lambda f_0} \sqrt{\frac{4}{\pi \tilde{K}_2^2}} \tilde{K}_2^{N_2} \tilde{K}_1^{N_1} \left[\prod_{j=1}^r \left(\omega_2^j \right)^{p_{2j}} \left(\omega_1^j \right)^{p_{1j}} \right]
$$

$$
\cdot \left(\frac{1}{2} \right)^{q_1+q_2} \left(-\frac{1}{M_2} \right)^{q_2+2m_2} \left(-\frac{1}{M_1} \right)^{q_1+2m_1} \Gamma \left(m_2 + \frac{1}{2} \right) \Gamma \left(m_1 + \frac{1}{2} \right)
$$

$$
= e^{-\Lambda f_0} \sqrt{\frac{4\pi}{\tilde{K}_2^2}} \tilde{K}_2^{N_2} \tilde{K}_1^{N_1} \left[\left(\frac{1}{2} \right)^{m_1+q_1} \left(-\frac{1}{M_1} \right)^{q_1+2m_1} (2m_1 - 1)!! \prod_{j=1}^r \left(\omega_1^j \right)^{p_{1j}} \right]
$$

$$
\cdot \left[\left(\frac{1}{2} \right)^{m_2+q_2} \left(-\frac{1}{M_2} \right)^{q_2+2m_2} (2m_2 - 1)!! \prod_{j=1}^r \left(\omega_2^j \right)^{p_{2j}} \right].
$$
 (3.25)

At last, the ratio can be easily calculated to be

$$
\frac{A^{(\{p_{1j}, p_{2j}\}, 2m_1, 2m_2, q_1, q_2)}}{A^{(\{N_1, N_2\}, 0, 0)}} = \left[\left(\frac{1}{2} \right)^{m_1 + q_1} \left(-\frac{1}{M_1} \right)^{q_1 + 2m_1} (2m_1 - 1)!! \prod_{j=1}^r \left(\omega_1^j \right)^{p_{1j}} \right]
$$

$$
\cdot \left[\left(\frac{1}{2} \right)^{m_2 + q_2} \left(-\frac{1}{M_2} \right)^{q_2 + 2m_2} (2m_2 - 1)!! \prod_{j=1}^r \left(\omega_2^j \right)^{p_{2j}} \right], \qquad (3.26)
$$

which is consistent with the calculation of ZNS method in Eq.[\(2.2\)](#page-6-1).

IV. THE K-IDENTITIES AND THE DEGREE OF STRINGY SCALING

In the general *n*-point multi-tensor $HSSA$ calculation, one key step is to introduce the diagonal and off-diagonal K-identities propsed in Eq. (3.17) . The K-identities were originally proposed only for the diagonal ones for the calculation of n -point 1 tensor $HSSA$ calculation [\[1](#page-26-0)[–3\]](#page-26-1). They are the direct results of the identification of two different calculation of $HSSA$, namely, the saddle-point calculation and the decoupling of zero-norm state (ZNS) calculation [\[1](#page-26-0)[–3](#page-26-1)].

We will show in this section that both diagonal and off-diagonal K -identities for the 4-point HSSA with 1 and 2 tensor can be analytically proved. For the 5-point HSSA calculation, although the saddle-point equation is quadratic and can be exactly solved, the explicit proof of the K-identities is too lengthy, and we will only prove them numerically. We have also used maple to numerically prove the K-identities for the 6-point $HSSA$. Presumably, these K-identities are direct results of conservation of momenta in the hard scattering limit at the saddle-point.

It is important to point out that the diagonal K-identities put constraints on the number of independent ω_i^j i_i , and the off-diagonal K-identities put further constraints on the number of independent ω_i^j i . So studying K-identities is crucial to calculate the degree of stringy scaling dim \mathcal{M}_k .

A. The K-identities

We begin our discussion of the K-identity with one example of 6-point 1 tensor $HSSA$ calculation. For $n = 6$ and $r = 3$, the ratio of HSSA can be calculated by using saddle-point method to be [\[1](#page-26-0)[–3\]](#page-26-1)

$$
\frac{A^{(p_1,p_2,p_3,2m,q)}}{A^{(N,0,0,0)}} = \frac{(2m)!}{m!} \left(\frac{-1}{2M}\right)^{2m+q} \frac{\left(\frac{\sum_{i\neq 2,n=6} \frac{k_i^{T_2}}{\tilde{z}_i - \tilde{z}_2}}{\tilde{z}_i - \tilde{z}_2}\right)^{p_2} \left(\frac{\sum_{i\neq 2,n=6} \frac{k_i^{T_3}}{\tilde{z}_i - \tilde{z}_2}}{\sum_{i\neq 2,n=6} \frac{k_i^{T_1}}{\tilde{z}_i - \tilde{z}_2}}\right)^{p_3}}{\left(\frac{\sum_{i\neq 2,n} \frac{k_i^{T_1}}{\tilde{z}_i - \tilde{z}_2}}{\sqrt{2K\tilde{p}_{2}}}\right)^{2m+2q}}.
$$
(4.1)

On the other hand, the decoupling of ZNS method gives [\[1](#page-26-0)[–3](#page-26-1)]

$$
\frac{A^{(p_1,p_2,2m,q)}}{A^{(N,0,0,0)}} = \frac{(2m)!}{m!} \left(\frac{-1}{2M}\right)^{2m+q} \omega_1^{p_1} \omega_2^{p_2} \omega_3^{p_3} = \frac{(2m)!}{m!} \left(\frac{-1}{2M}\right)^{2m+q} \frac{(\tan \theta_1 \cos \theta_2)^{p_2} (\tan \theta_1 \sin \theta_2)^{p_3}}{(\cos \theta_1)^{2m+2q}}.
$$
\n(4.2)

Eq.(??) and Eq.[\(4.2\)](#page-17-0) can be identified for any p_2 , p_3 , m and q if

$$
\sum_{i \neq 2, n=6} \frac{k_i^{T_1}}{\tilde{z}_i - \tilde{z}_2} = \sqrt{2\Lambda \tilde{f}_{22}} \cos \theta_1,\tag{4.3}
$$

$$
\frac{\sum_{i \neq 2, n=6} \frac{k_i^{T_2}}{\tilde{z}_i - \tilde{z}_2}}{\sum_{i \neq 2, n=6} \frac{k_i^{T_1}}{\tilde{z}_i - \tilde{z}_2}} = \sqrt{2\Lambda \tilde{f}_{22}} \tan \theta_1 \cos \theta_2, \tag{4.4}
$$

$$
\frac{\sum_{i \neq 2, n=6} \frac{k_i^{T_3}}{\tilde{z}_i - \tilde{z}_2}}{\sum_{i \neq 2, n=6} \frac{k_i^{T_1}}{\tilde{z}_i - \tilde{z}_2}} = \sqrt{2\Lambda \tilde{f}_{22}} \tan \theta_1 \sin \theta_2.
$$
\n(4.5)

which allow us to formally express θ_1 and θ_2 defined in Eq.[\(3.2\)](#page-12-1) in terms of kinematics variables defined in Eq.[\(3.10\)](#page-13-0) as

$$
\theta_1 = \arctan \frac{\sqrt{\left(\tilde{k}_{\perp}^{T_2}\right)^2 + \left(\tilde{k}_{\perp}^{T_3}\right)^2}}{\tilde{k}_{\perp}^{T_1}}, \ \theta_2 = \arctan \frac{\tilde{k}_{\perp}^{T_3}}{\tilde{k}_{\perp}^{T_2}}.
$$
\n
$$
(4.6)
$$

It is important to note from Eq.(??) that the ratios of 6-point $HSSA$ with $r = 3$ depends only on 2 kinematics variables θ_1 and θ_2 instead of 8. We see that Eq.[\(4.3\)](#page-17-1), Eq.[\(4.4\)](#page-17-2) and Eq.[\(4.5\)](#page-17-3) gives the K-identity

$$
\tilde{K}_2^2 + 2M_2 \partial_2 \tilde{K}_2^{L_2} = 0 \tag{4.7}
$$

in Eq.[\(2.24\)](#page-10-1) for the case of $n = 6$ and $r = 3$. We note that the K-identity puts 1 constraint on ω_1 , ω_2 and ω_3 . Thus there are only $r-1=3-1=2$ angles θ_1 and θ_2 , and $\dim \mathcal{M}=8-2=6$.

For the second example of the K-identities, we consider n -point 2 tensor with general r HSSA calculated in section III. For this multi-tensor case, in addition to the diagonal K -identities, we proposed the off-diagonal K -identity

$$
\tilde{K}_i \tilde{K}_j + 2M_i \partial_j \tilde{K}_i^{L_i} = 0 \Rightarrow \begin{cases} \tilde{K}_i^2 + 2M_i \partial_i \tilde{K}_i^{L_i} = 0, \ i = 1, 2, \\ \tilde{K}_1 \tilde{K}_2 + \frac{2k_{12}}{(\tilde{z}_1 - \tilde{z}_2)^2} = 0 \end{cases}
$$
\n(4.8)

in Eq.[\(3.17\)](#page-14-0). The diagonal and off-diagonal K-identities proposed for the calculation of 2 tensor HSSA are generalization of the diagonal K-identity proposed for the calculation of 1 tensor HSSA. They are the key step to obtain the consistent multi-tensor HSSA in the saddle-point calculation.

B. The degree of stringy scaling

For the general *n*-point 1 tensor $HSSA$, it was shown that [\[1](#page-26-0)[–3](#page-26-1)]

$$
\dim \mathcal{M}_1 = \frac{(r+1)(2n-r-6)}{2}.
$$
\n(4.9)

.

For a given (n, r) , we have $[1-3]$

For the general *n*-point 2 tensor $HSSA$, the off-diagonal K-identity in Eq.[\(3.17\)](#page-14-0) puts 1 further constraint except for the cases of $r = 1$ which contain only trivial $\omega_1^1 = \omega_2^1 = 1$ according to Eq. (3.2) . We thus have

$$
\dim M_2 = \frac{(r+1)(2n-r-6)}{2} - (r-1) + (1 - \delta_{r1})
$$

$$
= \dim M_1 - (r-1) + (1 - \delta_{r1}) \tag{4.10}
$$

where $\max r = \min (n-3, 24)$. The $(r-1)$ factor in Eq.[\(4.10\)](#page-18-0) is the number of angles of ω_2^j Z_2 , $j = 1, 2, \dots, r$, and $(1 - \delta_{r1})$ factor is the effect of the off-diagonal K-identity.

For a given (n, r) , we have

We note that for $r = 1, 2$, $\dim \mathcal{M}_2 = \dim \mathcal{M}_1$. For $r \geq 3$, $\dim \mathcal{M}_2 = \dim \mathcal{M}_1 - r + 2$. It is interesting to see that the off-diagonal K -identity contributes to the degree of scaling $\dim M_2$ for HSSA out of the scattering plane with $r \geq 2$. We see that for HSSA with more than 2 transverse directions or $r \geq 3$, the degree of scaling dim \mathcal{M}_2 decreases as one would have expected.

On the other hand, for a consistent check, for the case of $r = 24$ and $n \geq 27$, one can show that dim $\mathcal{M}_2 \geq 0$. For the case of $r = n - 3$ and $n \leq 27$, we also have the consistent result

$$
\dim \mathcal{M}_2 = \frac{(n-2)(n-3)}{2} - (n-3) + (2 - \delta_{r1})
$$

$$
= \left(\frac{n}{2} - 2\right)(n-3) + (2 - \delta_{r1}) \ge 0. \tag{4.11}
$$

.

For the general *n*-point k tensor $HSSA$, it will be a challenge to calculate $\dim \mathcal{M}_k$. It is conjectured that for general higher k tensor $HSSA$ calculation ($k \leq n$), the off-diaganal K-identities similar to Eq.[\(3.17\)](#page-14-0) are crucial to show the consistent result that $\dim \mathcal{M}_k \geq 0$.

C. Checking the K-identities

In this subsection, we will explicitly prove the K-identities of $HSSA$ for the 4-point 1 and 2 tensor cases. We will also discuss K-identities of some higher point HSSA. We begin with the 4-point kinematics in the hard scattering limit

$$
k_1 = (E, -E, 0),\nk_2 = (E, +E, 0),\nk_3 = (-E, -E\cos\phi, -E\sin\phi),\nk_4 = (-E, E\cos\phi, E\sin\phi),
$$
\n(4.12)

which can be used to calculate

$$
k_1 \cdot k_2 = -2E^2, \quad k_2 \cdot k_3 = 2E^2 \tau, \quad k_1 \cdot k_3 = 2E^2 (1 - \tau). \tag{4.13}
$$

For the 4-point case, the saddle-point equation is linear, and the saddle-point can be easily solved to be

$$
\tau = \sin^2 \frac{\phi}{2}, \, \tilde{x}_2 = \frac{1}{1 - \tau}.\tag{4.14}
$$

For the 4-point 1 tensor case, we want to show that

$$
\tilde{K}_2^2 + 2M_2 \partial_2 \tilde{K}_2^{L_2} = 0 \tag{4.15}
$$

where

$$
K_2(z_2) = \frac{k_1}{z_2} - \frac{k_3}{1 - z_2}.
$$
\n(4.16)

In the hard scattering limit

$$
\tilde{K}_2 = (1 - \tau) \left(k_1 + \frac{k_3}{\tau} \right),
$$
\n
$$
\tilde{K}_2^2 = (1 - \tau)^2 \left(k_1^2 + \frac{k_3^2}{\tau^2} + 2 \frac{k_1 \cdot k_3}{\tau} \right) = (1 - \tau)^2 \left(0 + \frac{0}{\tau^2} + \frac{4E^2 (1 - \tau)}{\tau} \right)
$$
\n
$$
= 4E^2 \frac{(1 - \tau)^3}{\tau}.
$$
\n(4.18)

On the other hand

$$
\partial_2 K_2(z_2) = -\frac{k_1}{z_2^2} - \frac{k_3}{(1-z_2)^2},\tag{4.19}
$$

so

$$
\partial_2 \tilde{K}_2 = - (1 - \tau)^2 \left(k_1 + \frac{k_3}{\tau^2} \right) \tag{4.20}
$$

and

$$
\partial_2 \tilde{K}_2^{L_2} = -(1-\tau)^2 \left(k_1 + \frac{k_3}{\tau^2}\right) \cdot \frac{k_2}{M_2}
$$

= $-2E^2 \frac{(1-\tau)^3}{\tau} \frac{1}{M_2}.$ (4.21)

The K-identity in Eq. (4.15) is thus proved.

For the 4-point 2 tensor cases, we define

$$
K_1(z_2) = -\frac{k_2}{z_2 - z_1} - \frac{k_3}{z_3 - z_1}, \quad K_2(z_2) = -\frac{k_1}{z_1 - z_2} - \frac{k_3}{z_3 - z_2}.
$$
 (4.22)

It is straightforward to calculate

and

One can now check the diagonal K -identities

$$
\left| \tilde{K}_1 \right| \left| \tilde{K}_1 \right| + 2M_1 \partial_1 \tilde{K}_1^{L_1} = \left| \tilde{K}_1 \right|^2 + 2M_1 \partial_1 \tilde{K}_1^{L_1}
$$

= $(-k_2 (1 - \tau) - k_3)^2 + 2M_1 (-k_2 (1 - \tau)^2 - k_3) \cdot \frac{k_1}{M_1}$

$$
k_2 \cdot k_3 (1 - \tau) - 2 (k_1 \cdot k_2 (1 - \tau)^2 + k_1 \cdot k_3)
$$

= $4E^2 \tau (1 - \tau) - 2 (-2E^2 (1 - \tau)^2 + 2E^2 (1 - \tau)) = 0$ (4.25)

and

$$
\left| \tilde{K}_2 \right| \left| \tilde{K}_2 \right| + 2M_2 \partial_2 \tilde{K}_2^{L_2} = \tilde{K}_2^2 + 2M_2 \partial_2 \tilde{K}_2^{L_2}
$$
\n
$$
= \left(k_1 + \frac{k_3}{\tau} \right)^2 (1 - \tau)^2 - 2M_2 \left(k_1 + \frac{k_3}{\tau^2} \right) \cdot \frac{k_2}{M_2} (1 - \tau)^2
$$
\n
$$
k_2 \left(1 - \tau \right)^2
$$
\n
$$
k_2 \left(1 - \tau \right)^2
$$
\n
$$
k_2 \left(1 - \tau \right)^2
$$
\n
$$
k_2 \left(1 - \tau \right)^2 - 2k_2 \cdot k_3 \frac{(1 - \tau)^2}{\tau^2}
$$
\n
$$
= (1 - \tau)^2 \left[\frac{4E^2 (1 - \tau)}{\tau} + 4E^2 - \frac{4E^2 \tau}{\tau^2} \right] = 0. \tag{4.26}
$$

Similarly, the off-diagonal K -identity is

$$
\left| \tilde{K}_1 \right| \left| \tilde{K}_2 \right| + 2M_1 \partial_2 \tilde{K}_1^{L_1} = \left| \tilde{K}_2 \right| \left| \tilde{K}_1 \right| + 2M_2 \partial_1 \tilde{K}_2^{L_2}
$$
\n
$$
= \sqrt{(-k_2 (1 - \tau) - k_3)^2} \sqrt{\left(\left(k_1 + \frac{k_3}{\tau} \right) (1 - \tau) \right)^2 + 2M_1 (1 - \tau)^2 \frac{k_2 \cdot k_1}{M_1}
$$
\n
$$
\left(\sqrt{2k_2 \cdot k_3 (1 - \tau)} \sqrt{2k_1 \cdot k_3 \frac{(1 - \tau)^2}{\tau}} + 2 (1 - \tau)^2 k_1 \cdot k_2 \right)
$$
\n
$$
= \sqrt{4E^2 \tau (1 - \tau)} \sqrt{4E^2 (1 - \tau) \frac{(1 - \tau)^2}{\tau}} - 4E^2 (1 - \tau)^2 = 0. \tag{4.27}
$$

We now turn to discuss the 5-point K -identities. The 5-point kinematics can be written as

$$
k_1 = \left(\sqrt{p^2 + M_1^2}, -p, 0, 0\right),
$$

\n
$$
k_2 = \left(\sqrt{p^2 + M_2^2}, p, 0, 0\right),
$$

\n
$$
k_3 = \left(-\sqrt{q_3^2 + M_3^2}, -q_3 \cos \phi_{32}, -q_3 \sin \phi_{32}, 0\right),
$$

\n
$$
k_4 = \left(-\sqrt{q_4^2 + M_4^2}, -q_4 \cos \phi_{42}, -q_4 \sin \phi_{42} \cos \phi_{43}, -q_4 \sin \phi_{42} \sin \phi_{43}\right),
$$

\n
$$
k_5 = \left(-\sqrt{q_5^2 + M_5^2}, -q_5 \cos \phi_{52}, -q_5 \sin \phi_{52} \cos \phi_{53}, -q_5 \sin \phi_{52} \sin \phi_{53}\right).
$$
\n(4.28)

The energy-momentum conservation laws imply

$$
\sqrt{p^2 + M_1^2} + \sqrt{p^2 + M_2^2} = \sqrt{q_3^2 + M_3^2} + \sqrt{q_4^2 + M_4^2} + \sqrt{q_5^2 + M_5^2},
$$

\n
$$
0 = q_3 \cos \phi_{32} + q_4 \cos \phi_{42} + q_5 \cos \phi_{52},
$$

\n
$$
0 = q_3 \sin \phi_{32} + q_4 \sin \phi_{42} \cos \phi_{43} + q_5 \sin \phi_{52} \cos \phi_{53},
$$

\n
$$
0 = 0 + q_4 \sin \phi_{42} \sin \phi_{43} + q_5 \sin \phi_{52} \sin \phi_{53}.
$$
 (4.29)

The energy conservation gives

$$
2p + \left(\frac{M_1^2}{2p} + \frac{M_2^2}{2p}\right) + \dots = q_3 + q_4 + q_5 + \left(\frac{M_3^2}{2q_3} + \frac{M_4^2}{2q_4} + \frac{M_5^2}{2q_5}\right) + \dots,\tag{4.30}
$$

which in the hard scattering limit means

$$
2p = q_3 + q_4 + q_5. \tag{4.31}
$$

We assume

$$
q_3: q_4: q_5 = 1: r_1: r_2 \tag{4.32}
$$

to get

$$
q_3 = \frac{2E}{1 + r_1 + r_2}, q_4 = \frac{2Er_1}{1 + r_1 + r_2}, q_5 = \frac{2Er_2}{1 + r_1 + r_2}.
$$
\n(4.33)

On the other hand, the momentum conservation laws become

$$
0 = \cos \phi_{32} + r_1 \cos \phi_{42} + r_2 \cos \phi_{52},
$$

\n
$$
0 = \sin \phi_{32} + r_1 \sin \phi_{42} \cos \phi_{43} + r_2 \sin \phi_{52} \cos \phi_{53},
$$

\n
$$
0 = r_1 \sin \phi_{42} \sin \phi_{43} + r_2 \sin \phi_{52} \sin \phi_{53},
$$
\n(4.34)

which can be used to solve r_2 , ϕ_{53} and ϕ_{52} . They are

$$
r_2(\phi_{32}, \phi_{42}, \phi_{43}, r_1)
$$

= $\sqrt{(\cos \phi_{32} + r_1 \cos \phi_{42})^2 + (\sin \phi_{32} + r_1 \sin \phi_{42} \cos \phi_{43})^2 + (r_1 \sin \phi_{42} \sin \phi_{43})^2},$
= $\sqrt{1 + r_1^2 + 2r_1 (\cos \phi_{32} \cos \phi_{42} + \sin \phi_{32} \sin \phi_{42} \cos \phi_{43})},$ (4.35)

$$
\phi_{53} (\phi_{32}, \phi_{42}, \phi_{43}, r_1)
$$

= arctan $\left(\frac{r_1 \sin \phi_{42} \sin \phi_{43}}{\sin \phi_{32} + r_1 \sin \phi_{42} \cos \phi_{43}} \right)$ (4.36)

and

$$
\phi_{52}(\phi_{32}, \phi_{42}, \phi_{43}, r_1) \tag{4.37}
$$

$$
= \arctan\left(\pm\sqrt{\frac{(r_1\sin\phi_{42}\sin\phi_{43})^2 + (\sin\phi_{32} + r_1\sin\phi_{42}\cos\phi_{43})^2}{(\cos\phi_{32} + r_1\cos\phi_{42})^2}}\right).
$$
 (4.38)

Now, all parameters can be express in terms of the following 5 ones

$$
E, \phi_{32}, \phi_{42}, \phi_{43}, r_1. \tag{4.39}
$$

The 5-point kinematics in the hard scattering limit can thus be written as

$$
k_1 = (E, -E, 0, 0),
$$

\n
$$
k_2 = (E, E, 0, 0),
$$

\n
$$
k_3 = (-q_3, -q_3 \cos \phi_{32}, -q_3 \sin \phi_{32}, 0),
$$

\n
$$
k_4 = (-q_4, -q_4 \cos \phi_{42}, -q_4 \sin \phi_{42} \cos \phi_{43}, -q_4 \sin \phi_{42} \sin \phi_{43}),
$$

\n
$$
k_5 = (-q_5, -q_5 \cos \phi_{52}, -q_5 \sin \phi_{52} \cos \phi_{53}, -q_5 \sin \phi_{52} \sin \phi_{53}),
$$
\n(4.40)

which can be used to calculate in the hard scattering limit

$$
k_{12} = -2E^2,
$$

\n
$$
k_{13} = \frac{2E^2}{1 + r_1 + r_2} (1 + \cos \phi_{32}),
$$

\n
$$
k_{23} = \frac{2E^2}{1 + r_1 + r_2} (1 - \cos \phi_{32}),
$$

\n
$$
k_{24} = \frac{2E^2 r_1}{1 + r_1 + r_2} (1 - \cos \phi_{42}),
$$

\n
$$
k_{34} = \frac{4E^2 r_1 r_2}{(1 + r_1 + r_2)^2} (-1 + \cos \phi_{32} \cos \phi_{42} + \sin \phi_{32} \sin \phi_{42} \cos \phi_{43}).
$$
\n(4.41)

We are now ready to discuss the 5-point K -identities. We first discuss 5-point 1 tensor K-identity. The saddle-point equation are

$$
\frac{\partial f(z_2, z_3)}{\partial z_2} = 0,\t\t(4.42)
$$

$$
\frac{\partial f(z_2, z_3)}{\partial z_3} = 0,\t\t(4.43)
$$

which can be expanded as

$$
\frac{1}{z_2} + \frac{k_{23}}{k_{12}} \frac{1}{z_2 - z_3} + \frac{k_{24}}{k_{12}} \frac{1}{z_2 - 1} = 0,
$$
\n(4.44)

$$
\frac{k_{13}}{k_{12}}\frac{1}{z_3} + \frac{k_{23}}{k_{12}}\frac{1}{z_3 - z_2} + \frac{k_{34}}{k_{12}}\frac{1}{z_3 - 1} = 0.
$$
 (4.45)

After some lengthy calculation, we end up with two quadratic equations for z_2 and z_3

$$
(k_{12} + k_{24} + k_{23}) (k_{12} + k_{13} + k_{23} + k_{24} + k_{34}) z_2^2
$$

- [(k₁₂ + k₁₃ + k₂₄ + k₂₃ + k₃₄)k₁₂ + (k₁₂ + k₂₄ + k₂₃)(k₁₂ + k₂₃ + k₁₃) + k₂₃k₃₄] z₂
+ (k₁₂ + k₂₃ + k₁₃)k₁₂ = 0,
(4.46)

$$
(k_{13} + k_{34} + k_{23}) (k_{12} + k_{13} + k_{23} + k_{24} + k_{34}) z_3^2
$$

- [(k₁₂ + k₁₃ + k₂₄ + k₂₃ + k₃₄) k₁₃ + (k₁₃ + k₃₄ + k₂₃) (k₁₃ + k₂₃ + k₁₂) + k₂₃k₂₄] z₃
+ (k₁₂ + k₂₃ + k₁₃) k₁₃ = 0. (4.47)

Although the saddle-point equation is exactly solvable, it turns out to be too lengthy to explicitly calculate it. We have instead successfully used maple to check the 5-point 1 and 2

tensor diagonal K-identities. The 5-point 2 tensor of diagonal K-identity can be similarly checked within the error range. In addition, we have also numerically checked the validity of 6-point 1 tensor diagonal K-identity.

V. CONCLUSION

In this paper, we use the saddle-point method to calculate n-point $HSSA$ with $n-2$ tachyons and 2 tensor states at arbitrary mass levels. During the multi-tensor saddle-point calculation, one encounters the naive energy order problem, and it turns out that one has to include contraction terms from $\partial_2 X_2 \cdots$ and $\partial_3 X_3 \cdots$ in order to achieve the consistent ratios calculated by the method of decoupling of ZNS in Eq.[\(2.3\)](#page-6-4).

More importantly, we discover the stringy scaling [\[19](#page-27-4)] behavior of these HSSA. For HSSA with $r = 1,2$, we found that $\dim \mathcal{M}_2 = \dim \mathcal{M}_1$. However, for HSSA with more than 2 transverse directions or $r \geq 3$, we found that the degree of stringy scaling dim \mathcal{M}_2 decreases comparing to the degree of scaling dim \mathcal{M}_1 of the $n-1$ tachyons and 1 tensor HSSA calculated previously.

In the saddle-point calculation of the multi-tensor $HSSA$, we propose a set of diagonal and off-diagonal K -identities which is the key to demonstrate the stringy scaling behavior of HSSA. We explicitly prove both the diagonal and off-diagonal K-identities for the 4-point HSSA and give a numerical proof of these K-identities for the higher point HSSA.

For the general n-point k tensor $HSSA$, we expect that there exist more types of Kidentities, and it will be a challenge to calculate the degree of stringy scaling $\dim M_k$. It is conjectured that for general higher k tensor $HSSA$ calculation $(k \leq n)$, the off-diaganal K-identities similar to Eq.[\(3.17\)](#page-14-0) are crucial to show the consistent result that $\dim \mathcal{M}_k \geq 0$.

Acknowledgments

We thank T. Yoneya and Y. Okawa for pointing out the importance of K-identities in the early stage of this work. This work is supported in part by National Science and Technology Council (NSTC) of Taiwan.

- [1] Sheng-Hong Lai, Jen-Chi Lee and Yi Yang, "Stringy scaling of n-point Regge string scattering amplitudes", JHEP 09 (2023), hep-th 2303.17909.
- [2] Sheng-Hong Lai, Jen-Chi Lee and Yi Yang, "Stringy scaling of n-point hard string scattering amplitudes", hep-th 2207.09236.
- [3] Sheng-Hong Lai, Jen-Chi Lee and Yi Yang, "Stringy scaling", hep-th 2408.12612, talk given by Jen-Chi Lee at Komaba, Tokyo university.
- [4] David J Gross and Paul F Mende. The high-energy behavior of string scattering amplitudes. *Phys. Lett. B*, 197(1):129–134, 1987.
- [5] David J Gross and Paul F Mende. String theory beyond the Planck scale. *Nucl. Phys. B*, 303(3):407–454, 1988.
- [6] David J. Gross. High-Energy Symmetries of String Theory. *Phys. Rev. Lett.*, 60:1229, 1988.
- [7] David J Gross and JL Manes. The high energy behavior of open string scattering. *Nucl. Phys. B*, 326(1):73–107, 1989.
- [8] Chuan-Tsung Chan and Jen-Chi Lee. Stringy symmetries and their high-energy limits. *Phys. Lett. B*, 611(1):193–198, 2005.
- [9] Chuan-Tsung Chan and Jen-Chi Lee. Zero-norm states and high-energy symmetries of string theory. *Nucl. Phys. B*, 690(1):3–20, 2004.
- [10] Chuan-Tsung Chan, Pei-Ming Ho, Jen-Chi Lee, Shunsuke Teraguchi, and Yi Yang. Highenergy zero-norm states and symmetries of string theory. *Phys. Rev. Lett.*, 96(17):171601, 2006.
- [11] Chuan-Tsung Chan, Pei-Ming Ho, Jen-Chi Lee, Shunsuke Teraguchi, and Yi Yang. Solving all 4-point correlation functions for bosonic open string theory in the high-energy limit. *Nucl. Phys. B*, 725(1):352–382, 2005.
- [12] Jen-Chi Lee and Yi Yang. Review on high energy string scattering amplitudes and symmetries of string theory. *arXiv preprint [arXiv:1510.03297](http://arxiv.org/abs/1510.03297)*, 2015.
- [13] Jen-Chi Lee and Yi Yang. Overview of high energy string scattering amplitudes and symmetries of string theory. *Symmetry*, 11(8):1045, 2019.
- [14] J. D. Bjorken, Phys. Rev. 179 (1969) 1547.
- [15] C. G. Callan and D. Gross, Phys. Rev. Lett. 22 (1969) 156.
- [16] V.N. Gribov and L. Lipatov, Sov. J. Nucl. Phys.15, 438 (1972).
- [17] G. Altarelli and G. Parisi, Nucl. Phys. B126, 298 (1977).
- [18] Hideyuki Kawai, David C Lewellen, and S-HH Tye. A relation between tree amplitudes of closed and open strings. *Nucl. Phys. B*, 269(1):1–23, 1986.
- [19] Oleg Andreev, Scaling laws in hadronic processes and string theory", [hep-th/0209256](http://arxiv.org/abs/hep-th/0209256) (2002).