Quantum Cluster State Model with Haagerup Fusion Category Symmetry

Zhian Jia^{1,2,*}

¹Centre for Quantum Technologies, National University of Singapore, SG 117543, Singapore

²Department of Physics, National University of Singapore, SG 117543, Singapore

(Dated: December 30, 2024)

We propose a (1+1)D lattice model, inspired by a weak Hopf algebra generalization of the cluster state model, which realizes Haagerup fusion category \mathcal{H}_3 symmetry and features a tensor product Hilbert space. The construction begins with a reconstruction of the Haagerup weak Hopf algebra \mathcal{H}_3 from the Haagerup fusion category, ensuring that the representation category of \mathcal{H}_3 is equivalent to \mathcal{H}_3 . Utilizing the framework of symmetry topological field theory (SymTFT), we develop an ultra-thin weak Hopf quantum double model, characterized by a smooth topological boundary condition. We show that this model supports Haagerup fusion category symmetry. Finally, we solve the ground state of the model in terms of a weak Hopf matrix product state, which serves as a natural generalization of the cluster state, embodying Haagerup fusion category symmetry.

Introduction. — Symmetry is one of the central topics in physics and is traditionally characterized by a group. From a modern perspective, symmetries are characterized by the algebraic structure of topological defects [1–16]. Symmetries associated with non-invertible topological defects are known as non-invertible symmetries. For an (n+1)-dimensional system, the symmetries are described by fusion *n*-categories [3, 17]. For (1+1)-dimensional systems, these symmetries are described by fusion 1-categories \mathbb{C} , i.e., the usual fusion categories; hence, they are also referred to as fusion category symmetries [15, 16, 18].

The topological defect lines (TDLs) for (1+1)D system are labeled by the simple objects in C, and these TDLs can be fused according to the fusion rule:

$$a \otimes b = \sum_{c \in \operatorname{Irr}(\mathfrak{C})} N_{ab}^c c, \tag{1}$$

where N_{ab}^c are non-negative integers known as fusion multiplicities. The associativity of the fusion is characterized by *F*-symbols. Group symmetry is a special case where the fusion category is chosen as Vect_{*G*}, the category of *G*-graded vector spaces [19].

Non-invertible symmetries also play a crucial role in conformal field theory (CFT), particularly in the investigation of its properties and classification. In the case of rational CFTs (RCFTs), which form the foundational building blocks of general CFTs, the underlying mathematical structure is identified as modular tensor categories (MTCs) [20]. An important open question is whether a corresponding CFT can be constructed for any given MTC. It has been proposed that subfactors can be used to construct CFTs, with a conjectured correspondence between subfactors and CFTs [21-25]. While there is significant evidence supporting this conjecture, several potential counterexamples exist, one of the most notable being the Haagerup subfactor [26–28]. There are three fusion categories arising from Haagerup subfactors, namely $\mathcal{H}_1, \mathcal{H}_2$, and \mathcal{H}_3 , each of which is Morita equivalent to the others. Recently, there has been considerable interest in constructing lattice models that exhibit Haagerup \mathcal{H}_3 symmetry [29–34]. The

Haagerup fusion category \mathcal{H}_3 [35, 36] has a Drinfeld center $\mathcal{Z}(\mathcal{H}_3)$ that is a modular tensor category (MTC).

Given that any unitary (multi)fusion category corresponds to the representation category Rep(H) of some weak Hopf algebra H, it is a natural progression to examine non-invertible symmetries within the framework of Hopf and weak Hopf algebras [37–44]. These symmetries, which are associated with structures such as Hopf, quasi-Hopf, and weak Hopf algebras, along with their module and comodule algebras, represent a substantial class of non-invertible symmetries [4, 45–50].

In this work, we introduce a (1+1)D lattice model, inspired by a weak Hopf algebra generalization of the cluster state model. This model realizes the Haagerup fusion category \mathcal{H}_3 symmetry and is constructed with a tensor product Hilbert



FIG. 1. Illustration of SymTFT sandwich and cluster ladder model. (a) The symmetry TFT consists of a symmetry boundary \mathcal{B}_{sym} which encodes the fusion category symmetry; a physical boundary, which may be gapped or gapless that encodes the dynamics of the theory; the bulk is a topological field theory $\mathcal{Z}(\mathcal{B}_{svm})$. (b) Depiction of the cluster ladder model, which is an ultra-thin quantum double model with two boundaries (the qudit is put on edges), one boundary is chosen as a smooth boundary that encodes the symmetry information. If the physical boundary is chosen as a rough boundary, then the model becomes a cluster state model. (c) The chessboard representation of the cluster ladder model, where the qudits are placed on vertices, and each vertex corresponds to an edge of the quantum double model. For the cluster state model, the vertices on the physical boundary (cyan vertices) are removed, leaving only two types of local stabilizers: one for the symmetry boundary vertex operator and the other for the face operator.

^{*} giannjia@foxmail.com

space, distinguishing it significantly from models constructed using anyonic chains [30–33].

The one-dimensional cluster state model is a quintessential example of an symmetry-protected topological (SPT) phase, exhibiting $\mathbb{Z}_2 \times \mathbb{Z}_2$ SPT order [51]. More recently, it has been proposed that the cluster state also hosts a noninvertible global symmetry, described by the fusion category $\operatorname{Rep}(D_8)$ [52]. Generalizations to finite groups are discussed in Refs. [53, 54]. The Hopf algebraic generalization is presented in Ref. [55], where it is shown that the cluster state model is, in fact, an ultra-thin quantum double model with one smooth boundary and one rough boundary. In a recent work [56], we develop a general theory for the weak Hopf cluster state model, which can realize arbitrary weak Hopf symmetries (with fusion category symmetry as a special case). Building on these insights, we introduce a lattice model that exhibits Haagerup fusion category symmetry (and, more generally, Haagerup weak Hopf symmetry). The main result can be summarized as follows:

Theorem 1. By applying the Tannaka-Krein reconstruction or the boundary tube algebra approach, we can recover a C^* weak Hopf algebra H_3 , whose representation category $\text{Rep}(H_3)$ is equivalent to the Haagerup fusion category \mathcal{H}_3 as fusion categories. When H_3 is used as input data for a weak Hopf cluster state model (or, more generally, a cluster ladder model) as described in Ref. [56], we obtain a cluster model that exhibits Haagerup \mathcal{H}_3 symmetry (see Eq. (23)) and, more generally, weak Hopf symmetry \hat{H}_3 (the dual weak Hopf algebra of H_3).

The Haagerup fusion category symmetric cluster state model $\mathbb{H}_{\text{cluster}} = -\sum_{v_s} \mathbb{A}_{v_s} - \sum_f \mathbb{B}_f$ possesses the following weak Hopf symmetry:

- On a closed manifold: $Sym = Cocom(\hat{H}_3) \times Cocom(H_3)$, where Cocom(H) is the set of all cocommutative elements in weak Hopf algebra H.
- On an open manifold: Sym = $\hat{H}_3 \times H_3$.

Note that $\mathcal{H}_3 \simeq \operatorname{Rep}(H_3) \subset \operatorname{Cocom}(\hat{H}_3) \subset \hat{H}_3$ (here we focus on the fusion algebra of the category), implying that both cases exhibit \mathcal{H}_3 symmetry.

Haagerup Fusion Category \mathcal{H}_3 Symmetry. — The Haagerup fusion category is one of the simplest examples that does not originate from finite groups or affine Lie algebras.

The category \mathcal{H}_3 contains six simple objects: **1**, α , α^2 , ρ , $_{\alpha}\rho$, $_{\alpha^2}\rho$. The fusion rules are summarized in Table I, with the nontrivial rules given by:

$$\alpha^{3} = \mathbf{1}, \quad {}_{\alpha}\rho = \alpha \otimes \rho, \quad {}_{\alpha^{2}}\rho = \alpha^{2} \otimes \rho,$$

$$\rho \otimes \alpha = {}_{\alpha^{2}}\rho, \quad \rho \otimes \alpha^{2} = {}_{\alpha}\rho,$$

$$\rho \otimes \rho = \mathbf{1} \oplus \rho \oplus {}_{\alpha}\rho \oplus {}_{\alpha^{2}}\rho.$$
(2)

All other fusion rules can be derived from Eq. (2). The quantum dimensions of the simple objects are $d_1 = d_{\alpha} = d_{\alpha^2} = 1$ and $d_{\rho} = d_{\alpha\rho} = d_{\alpha^2}\rho = \frac{3+\sqrt{13}}{2}$. The *F*-symbols of \mathcal{H}_3 have been computed in Refs. [57, 58].

TABLE I. Fusion rule for the Haagerup fusion category \mathcal{H}_3 . In this table, the elements of the first column are fused with the elements of the first row ("column labels" \otimes "row labels").

\otimes	1	α	α^2	ρ	αρ	$\alpha^2 \rho$
1	1	α	α^2	ρ	$_{\alpha}\rho$	$\alpha^2 \rho$
α	α	α^2	1	$\alpha \rho$	$\alpha^2 \rho$	ρ
α^2	α^2	1	α	$\alpha^2 \rho$	ρ	$\alpha \rho$
ρ	ρ	$_{\alpha^2} ho$	$_{\alpha}\rho$	$1 \oplus ho \oplus {}_{lpha} ho \oplus {}_{lpha^2} ho$	$\alpha^2 \oplus \rho \oplus {}_{\alpha}\rho \oplus {}_{\alpha^2} ho$	$\alpha \oplus ho \oplus {}_{lpha} ho \oplus {}_{lpha^2} ho$
$_{\alpha}\rho$	$_{\alpha}\rho$	ρ	$\alpha^2 \rho$	$\alpha \oplus \rho \oplus {}_{\alpha}\rho \oplus {}_{\alpha^2} ho$	$1 \oplus ho \oplus {}_{lpha} ho \oplus {}_{lpha^2} ho$	$\alpha^2 \oplus \rho \oplus {}_{\alpha}\rho \oplus {}_{\alpha^2} ho$
$\alpha^2 \rho$	$\alpha^2 \rho$	$_{\alpha}\rho$	ρ	$\alpha^2 \oplus \rho \oplus {}_{\alpha}\rho \oplus {}_{\alpha^2} ho$	$\alpha \oplus \rho \oplus {}_{\alpha}\rho \oplus {}_{\alpha^2} ho$	$1 \oplus ho \oplus {}_{lpha} ho \oplus {}_{lpha^2} ho$

The Drinfeld center $\mathcal{Z}(\mathcal{H}_3)$ is a MTC containing 12 simple objects (topological charges). These simple objects are written as $(X, \beta_{X,\bullet})$, where X is a simple object in \mathcal{H}_3 , and $\beta_{X,\bullet}$ denotes the half-braiding. Following Ref. [59], we denote these topological charges as $\mathbf{1}_{\mathcal{Z}(\mathcal{H}_3)}, \pi_{1,2}, \sigma_{1,2,3}, \mu_{1,2,\cdots,6}$. The quantum dimensions and topological spins of these topological charges are listed in Table II. The quantum dimension of $\mathcal{Z}(\mathcal{H}_3)$ is

$$\dim(\mathcal{Z}(\mathcal{H}_3)) = \sum_{a \in \operatorname{Irr}(\mathcal{Z}(\mathcal{H}_3))} d_a^2 = 99d_\rho^2 + 54d_\rho + 18.$$
(3)

The S-matrix is given in Ref. [59, Section 2.2].

TABLE II. Simple objects of $\mathcal{Z}(\mathcal{H}_3)$, specified by $(X, \beta_{X, \bullet})$ with $X \in \mathcal{H}_3$ and $\beta_{X, \bullet}$ as the half-braiding, along with their quantum dimensions and topological spins.

$\mathcal{Z}(\mathcal{H}_3)$	$X \in \mathcal{H}_3$	$\beta_{X,\bullet}$	quantum dimension	topological spin
$1_{\mathcal{Z}(\mathcal{H}_3)}$	1	id	1	1
π_1	$\pi_1 = 1 \oplus \rho \oplus_{\alpha} \rho \oplus_{\alpha^2} \rho$	$\beta_{\pi_1,\bullet}$	$3d_{\rho} + 1$	1
π_2	$\pi_2 = 1 \oplus 1 \oplus \rho \oplus_{\alpha} \rho \oplus_{\alpha^2} \rho$	$\beta_{\pi_2,\bullet}$	$3d_{\rho}+2$	1
$\sigma_{1,2,3}$	$\sigma = \alpha \oplus \alpha^2 \oplus \rho \oplus {}_{\alpha}\rho \oplus {}_{\alpha^2}\rho$	$\beta_{\sigma,\bullet}^{1,2,3}$	$3d_{\rho} + 2$	$1, e^{\pm 2\pi i/3}$
$\mu_{1,2,\cdots,6}$	$\mu = \rho \oplus_{\alpha} \rho \oplus_{\alpha^2} \rho$	$\beta^{1,2,\cdots,6}_{\mu,\bullet}$	3 <i>d</i> _p	$e^{\pm 4\pi i/13},$ $e^{\pm 10\pi i/13},$ $e^{\pm 12\pi i/13}$

Reconstructing weak Hopf H₃ symmetry from Haagerup fusion category H₃ symmetry. — Our model is based on the weak Hopf algebra H₃ for which H₃ is equivalent to Rep(H₃). By definition, a complex weak Hopf algebra [60] is a complex vector space H equipped with algebra (H, μ, η) and coalgebra (H, Δ, ε) structures, along with an antipode map $S: H \to H$, satisfying specific compatibility conditions. We use Sweedler's notation for comultiplication, denoting $\Delta(h) = \sum_{(h)} h^{(1)} \otimes h^{(2)} := \sum_i h_i^{(1)} \otimes h_i^{(2)}$. The comultiplication satisfies $\Delta(xy) = \Delta(x) \cdot \Delta(y)$, explicitly expressed as $\sum_{(xy)} (xy)^{(1)} \otimes (xy)^{(2)} = \sum_{(x),(y)} x^{(1)} \cdot y^{(1)} \otimes x^{(2)} \cdot y^{(2)}$. The unit satisfies weak comultiplicativity, $(\Delta \otimes id) \circ \Delta(1_H) = (\Delta(1_H) \otimes 1_H) \cdot (1_H \otimes \Delta(1_H))$, where $\Delta(1_H) = \sum_{(1_H)} 1_H^{(1)} \otimes 1_H^{(2)}$. The counit satisfies weak multiplicativity, meaning that for all $x, y, z \in H$, $\varepsilon(xyz) = \sum_{(y)} \varepsilon(xy^{(1)})\varepsilon(y^{(2)}z)$. The antipode S satisfies three

key properties. For the left counit, $\sum_{(x)} x^{(1)} S(x^{(2)}) = \varepsilon_L(x) := \sum_{(1_H)} \varepsilon(1_H^{(1)} x) 1_H^{(2)}$. For the right counit, $\sum_{(x)} S(x^{(1)}) x^{(2)} = \varepsilon_R(x) := \sum_{(1_H)} 1_H^{(1)} \varepsilon(x 1_H^{(2)})$. Additionally, the antipode satisfies the decomposition property, $\sum_{(x)} S(x^{(1)}) x^{(2)} S(x^{(3)}) = S(x)$.

The images of the left and right counit maps, $H_L = \varepsilon_L(H)$ and $H_R = \varepsilon_R(H)$, serve as the tensor unit in the representation category Rep(H), which is typically a multifusion category. If H is both connected [61] (also known as pure in [60]) and coconnected [62], then Rep(H) is a fusion category. For more information on weak Hopf algebras and their applications to lattice models, refer to Refs. [43, 56].

The Tannaka-Krein reconstruction offers a structured approach to obtaining a weak Hopf algebra from a fusion category. For any fusion category \mathcal{C} , there exists a \mathbb{C} -algebra A such that \mathcal{C} can be faithfully and exactly embedded into ${}_{A}\operatorname{Mod}_{A}$ via a monoidal functor $F : \mathcal{C} \to {}_{A}\operatorname{Mod}_{A}$. The algebra $H = \operatorname{End}(F)$, defined by the natural transformations of F, forms a weak Hopf algebra. Furthermore, the representation category [63, 64]. When applied to \mathcal{H}_3 , this process yields a weak Hopf algebra, denoted H_3 , such that $\mathcal{H}_3 \simeq \operatorname{Rep}(H_3)$. The weak Hopf algebras whose representation categories are equivalent to \mathcal{H}_3 are not unique; they are Morita equivalent to one another. We will call H_3 Haagerup weak Hopf algebra.

An intuitive and explicit approach involves the concept of the boundary tube algebra [44, 65–67], as applied to the string-net model, where the basis elements and Haar integral can be explicitly written. For a given bulk fusion category C, we consider a gapped boundary described by a C-module category ${}_{C}\mathcal{M}$ and construct the corresponding boundary tube algebra **Tube**(${}_{C}\mathcal{M}$), which is a C^* weak Hopf algebra. Setting $\mathcal{M} = C$, i.e., considering the smooth boundary, we obtain the boundary tube algebra spanned by the following basis:

$$\begin{cases} g \\ f \\ e \\ e \\ c \end{cases} & \mu \in \operatorname{Hom}(a \otimes f, g), \\ \psi \\ e \\ c \\ \mu \in \operatorname{Hom}(c, a \otimes e) \\ c \\ \end{cases} \right\}.$$
(4)

The unit is give by



which means we set a = 1 and c = e, f = g and $\mu = id$, $\nu = id$

in Eq. (4). The multiplication is defined as



After evaluating the diagram on the right-hand side using F-moves, parallel moves, and loop moves [44], we obtain a linear combination of the basis elements in Eq. (4). The counit is given by



After performing the topological evaluation of the right-hand side, the result will be a real number. The comultiplication is defined as



The antipode map is defined as



It can be proved that this is a C^* weak Hopf algebra [44]. The Haar integral is of the form:

$$\lambda = \frac{1}{\operatorname{rank} \mathcal{C}} \frac{1}{\operatorname{FPdim}(\mathcal{C})} \sum_{a,x,y,\mu} \sqrt{\frac{d_a}{d_x^3 d_y}} a \left(\begin{array}{c} y \\ x \\ x \\ y \end{array} \right). \quad (10)$$

A detailed proof will be provided in Ref. [67]. Since the dual weak Hopf algebra can be regarded as a **Tube**($\mathcal{C}_{\mathcal{C}}$), the Haar measure Λ (Haar integral of the dual weak Hopf algebra) can be constructed similarly (see also Ref. [66] for a slightly different construction, where the structure constants of the boundary tube algebra are chosen differently, although the two algebras remain Morita equivalent).

The fusion category $\mathcal{C} = \operatorname{Fun}_{\mathbb{C}}(\mathcal{C}, \mathcal{C})$ (the category of module category functors), interpreted as the boundary phase, can be embedded into the representation category of the tube algebra $\operatorname{Rep}(\operatorname{Tube}(_{\mathbb{C}}\mathcal{C}))$. It is believed that $\mathcal{C} \simeq \operatorname{Rep}(\operatorname{Tube}(_{\mathbb{C}}\mathcal{C}))$ [65, 66], and we will argue in Ref. [67] that this is, in fact, an embedding $\mathcal{C} \hookrightarrow \operatorname{Rep}(\operatorname{Tube}(_{\mathbb{C}}\mathcal{C}))$ for general case. Consequently, the fusion category $\mathcal{C}_{\mathcal{M}}^{\vee} := \operatorname{Fun}_{\mathbb{C}}(\mathcal{M}, \mathcal{M})$ can be realized through the weak Hopf algebra $\operatorname{Tube}(_{\mathbb{C}}\mathcal{C})$. For further details, we refer the reader to Refs. [44, 67]. When applied to \mathcal{H}_3 , we denote the corresponding tube algebra as $H_3 = \operatorname{Tube}(_{\mathcal{H}_3}\mathcal{H}_3)$, with a slight abuse of notation.

Symmetry Topological Field Theory (SymTFT) with Haagerup \mathcal{H}_3 symmetry. — SymTFT provides a general framework for studying non-invertible symmetry-protected topological (SPT) phases in both gapped and gapless cases (see, e.g., [3–13]). The SymTFT can be applied to weak Hopf lattice gauge theory and establish the general algebraic framework for (1+1)D non-invertible SPT phases [55, 56]. Based on this, we will construct a lattice model.

The SymTFT has a sandwich structure, as illustrated in Fig. 1 (a). The sandwich manifold is defined as $\mathbb{M}^{1,1} \times [0,1]$, where $\mathbb{M}^{1,1}$ represents the (1 + 1)-dimensional manifold on which our system resides. This manifold has two boundaries: the basic idea is to impose non-invertible symmetry on one boundary and place the physical system on the other. After compactifying over the interval [0,1], we obtain a (1 + 1)-dimensional system with non-invertible symmetry. To construct the SymTFT, the symmetry boundary must be chosen as a gapped topological boundary condition. There are many topological orders that do not have any gapped boundary, but non-chiral topological orders are guaranteed to have at least one gapped boundary. Thus, we put a non-chiral topological order on the 2d bulk.

For Haagerup symmetry $\mathcal{H}_3 = \operatorname{Rep}(H_3)$, we place $\mathcal{Z}(\mathcal{H}_3)$ on the (2+1)D bulk of SymTFT, which means the bulk is a weak Hopf quantum double model with input data given by H_3 . The bulk topological excitation is characterized by $\mathcal{Z}(\mathcal{H}_3) \simeq_{\otimes, \mathrm{br}} \operatorname{Rep}(D(H_3))$, where the equivalence is both monoidal and braided, and $D(H_3)$ denotes the quantum double of H_3 (see [43] for details on the weak Hopf quantum double model). The symmetry boundary is chosen as $\mathcal{B}_{\mathrm{sym}} = \mathcal{H}_3 =$ $\operatorname{Rep}(H_3)$, meaning that the symmetry boundary is a smooth boundary for the quantum double model. The boundary input data is H_3 (regarded as a comodule algebra over H_3). For the physical boundary, we can choose either a gapped or gapless boundary condition, and the resulting lattice model will possess Haagerup fusion category symmetry.

Haagerup Fusion Symmetric Cluster Ladder Model. — To construct the lattice model with Haagerup fusion category symmetry, we first introduce the generalized Pauli operators. The regular action of a Haagerup weak Hopf algebra H_3 on itself can be viewed as a generalization of Pauli X-type operators. For the left action $H \curvearrowright H$, we define:

$$\overrightarrow{X}_{g}|h\rangle = |gh\rangle, \quad \overleftarrow{X}_{g}|h\rangle = |hS^{-1}(g)\rangle, \quad \forall g, h \in H_{3}.$$
 (11)

The dual space $\hat{H}_3 := \text{Hom}(H_3, \mathbb{C})$ is also a weak Hopf algebra. There are also canonical actions of the dual weak Hopf algebra \hat{H}_3 on the Hopf qudit H_3 , defined using Sweedler's notation as follows:

$$\boldsymbol{\varphi} \rightharpoonup \boldsymbol{x} := \sum_{(x)} \boldsymbol{x}^{(1)} \langle \boldsymbol{\varphi}, \boldsymbol{x}^{(2)} \rangle, \quad \boldsymbol{x} \leftarrow \boldsymbol{\varphi} := \sum_{(x)} \langle \boldsymbol{\varphi}, \boldsymbol{x}^{(1)} \rangle \boldsymbol{x}^{(2)}, \quad (12)$$

for all $\varphi \in \hat{H}_3$ and $x \in H_3$. For the left action $\hat{H}_3 \frown H_3$, we define:

$$\overrightarrow{Z}_{\psi}|h\rangle = |\psi \rightharpoonup h\rangle = \sum_{(h)} \psi(h^{(2)})|h^{(1)}\rangle, \qquad (13)$$

$$\stackrel{\leftarrow}{Z}_{\psi}|h\rangle = |h \leftarrow \hat{S}(\psi)\rangle = \sum_{(h)} \psi(S(h^{(1)}))|h^{(2)}\rangle.$$
(14)

These operators can be regarded as generalized Pauli Z-type operators.

Our construction of local operators is based on the comodule algebra over H_3 . By definition, an algebra K is referred to as a left H_3 -comodule algebra [68] if there exists a map $\beta: K \to H_3 \otimes K$ that satisfies the following properties:

$$\beta(xy) = \beta(x)\beta(y), \tag{15}$$

and

$$(1_{H_3} \otimes x)\beta(1_K) = (\varepsilon_R \otimes \mathrm{id}_K) \circ \beta(x), \tag{16}$$

for all $x, y \in K$. Using Sweedler's notation, the coaction $\beta(x)$ is expressed as $\sum_{[x]} x^{[-1]} \otimes x^{[0]}$, with higher coactions represented as $\beta_2(x) = \sum_{[x]} x^{[-2]} \otimes x^{[-1]} \otimes x^{[0]}$, and so forth. Each topological boundary condition corresponds to an indecomposable comodule algebra *K* over *H*₃. This comodule algebra also corresponds to an indecomposable module category \mathcal{M}_K over \mathcal{H}_3 , as well as a Lagrangian algebra \mathcal{A}_K in the bulk quantum double phase $\mathcal{Z}(\mathcal{H}_3)$.

The model is put on an ultra-thin sandwich lattice (regarded as a weak Hopf quantum double model [43]):



Periodic or open boundary conditions can be applied in the horizontal direction. The bulk edges are drawn in black, with the edge Hilbert space chosen as the Haagerup weak Hopf qudit H_3 . The edges of the symmetry boundary are drawn in blue, with the edge space given by the left comodule algebra $K = H_3$. The edges of the physical boundary are drawn in red, where the edge space can be chosen as an arbitrary right H_3 -comodule algebra J. This means that the total Hilbert space has well-defined tensor-product structure.

Observe that, in the described lattice structure, the bulk resides to the left of the symmetry boundary and to the right of the physical boundary when moving along the positive boundary direction. Therefore, the associated comodule algebras Kand J should be selected as a left H_3 -comodule algebra and a right H_3 -comodule algebra, respectively. Modifying the configuration of edge orientations in the lattice leads to different models, but does not alter the underlying physics of the system.

The smooth boundary corresponds to $K = H_3$. In this case, the boundary vertex operator is given by:

where λ is the Haar integral of H_3 (Eq. (10)). The cocommutativity of λ ensures that the operator $\mathbb{A}_{v_s}^K$ is independent of the specific starting link $s = (v_s, f)$ chosen at the vertex v_s . Consequently, $\mathbb{A}_{v_s}^K$ depends only on the vertex v_s itself. The symmetry boundary Hamiltonian is defined as

$$\mathbb{H}_{\text{sym}} = -\sum_{\nu_s} \mathbb{A}_{\nu_s}^K,\tag{19}$$

where the sum runs over all vertices in the symmetry boundary.

The face operator is constructed from generalized Pauli *Z* operators as follows:

$$g \underbrace{\textcircled{1}}_{\mathcal{Y}}^{X} h \qquad \mathbb{B}_{f}^{\psi} = \overrightarrow{Z}_{\psi^{(1)}} \otimes \overleftarrow{Z}_{\psi^{(2)}}^{K} \otimes \overleftarrow{Z}_{\psi^{(3)}}^{J} \otimes \overrightarrow{Z}_{\psi^{(4)}}^{J}.$$

Written explicitly, we have

$$\mathbb{B}_{f}^{\Psi}|g,y,h,x\rangle = \sum \Psi\left(g^{(2)}S(y^{[-1]})S(h^{(1)})x^{[1]}\right)|g^{(1)},y^{[0]},h^{(2)},x^{[0]}\rangle.$$

By setting ψ to be the Haar measure $\Lambda \in \hat{H}_3$, we obtain the face stabilizer operator $\mathbb{B}_f = \mathbb{B}_f^{\Lambda}$, and the corresponding bulk Hamiltonian is

$$\mathbb{H}_{bk} = -\sum_{f} \mathbb{B}_{f}^{H}, \qquad (21)$$

(20)

where the sum runs over all faces of the lattice.

The physical boundary can be chosen to be either gapped or gapless. In the gapped case, we assign a right H_3 -comodule algebra to each edge of the physical boundary. The vertex operator can be constructed in a similar way as in Eq. (18) (see Ref. [56], where we need to use the symmetric separability idempotent of J). The physical boundary Hamiltonian is given by

$$\mathbb{H}_{\text{phys}} = -\sum_{\nu_p} \mathbb{A}^J_{\nu_p},\tag{22}$$

where the sum runs over all vertices v_p of the physical boundary.

The cluster ladder model that has Haagerup fusion category symmetry is thus given by

ŀ

$$\mathbb{H}_{H_3}^{K,J} = \mathbb{H}_{bk} + \mathbb{H}_{sym} + \mathbb{H}_{phys}.$$
 (23)

It can be proven that all local operators commute with each other. Fixing symmetry boundary as smooth boundary, viz., $K = H_3$, different choices of comodule algebra *J* over the Haagerup weak Hopf algebra H_3 give rise to different \mathcal{H}_3 -symmetric phases.

In fact, the model has a larger symmetry. Since $\mathcal{H}_3 \simeq \text{Rep}(\mathcal{H}_3)$, the Grothendieck group $\text{Gr}(\mathcal{H}_3)$ of the category is generated by the irreducible characters χ_{Γ} of \mathcal{H}_3 , and

$$\chi_{\Gamma} \cdot \chi_{\Phi} = \chi_{\Gamma \otimes \Phi} = \sum_{\Psi} N_{\Gamma, \Phi}^{\Psi} \chi_{\Psi}, \qquad (24)$$

where $N_{\Gamma,\Phi}^{\Psi} \in \mathbb{Z}_{\geq 0}$ are the fusion multiplicities. The character algebra is given by $R(\mathcal{H}_3) = \operatorname{Gr}(\mathcal{H}_3) \otimes_{\mathbb{Z}} \mathbb{C}$, which is a subalgebra of the dual weak Hopf algebra \hat{H}_3 . This means that the Haagerup fusion category symmetry is a sub-symmetry of the dual Haagerup weak Hopf symmetry \hat{H}_3 .

A crucial special case is the cluster state model, where J is chosen as the trivial comodule algebra. In this case, all physical boundary operators become trivial and can be removed from the Hamiltonian. This makes the model equivalent to a weak Hopf quantum double model that has one smooth boundary and one rough boundary.

To find the symmetry of the Haagerup cluster state model, we deform the ladder (with periodic boundary conditions) into a cone. We introduce a vertex v_{rough} on the rough boundary (the apex of the cone) and a face f_{smooth} on the smooth boundary (the base of the cone):



In this way, we can regard the model as a quantum double model on a sphere. The symmetry operator at the smooth boundary is the face operator $W_{\psi} = \mathbb{B}_{f_{smooth}}^{\psi}$. We have the relation $W_{\psi}W_{\phi} = W_{\psi\cdot\phi}$ with $\psi, \phi \in \hat{H}_3$. When $\psi \in Cocom(\hat{H}_3)$, the operator $\mathbb{B}_{f_{smooth}}^{\psi}$ will commute with the Hamiltonian [43, 69]. This implies that the model has a $Cocom(\hat{H}_3)$ symmetry. For $\mathcal{H}_3 = \text{Rep}(H_3)$, its character algebra is contained in $Cocom(\hat{H}_3)$, and thus the model also has Haagerup fusion category symmetry. The rough boundary also gives symmetry operators $W_h = \mathbb{A}_{v_{rough}}^h$, with the relation $W_h W_g = W_{hg}$. When $h \in Cocom(H_3)$, W_h commutes with the Hamiltonian. This implies that the model has a $Cocom(H_3)$ symmetry. For open manifolds, there is no need for the symmetry elements to be cocommutative, and thus the symmetry becomes $\hat{H}_3 \times H_3$. The result is summarized in Theorem 1.

For the general cluster ladder model, due to the fact that J is not trivial, the symmetry H_3 from the rough boundary is not

broken. Using a similar discussion as above, we find that the dual symmetry \hat{H}_3 still exists. Thus, for periodic boundary conditions, the model has $\text{Cocom}(\hat{H}_3)$ symmetry, while for open boundary conditions, the model has \hat{H}_3 symmetry. In both cases, the symmetry contains \mathcal{H}_3 as a sub-symmetry.

Matrix-Product State Representation of Ground State. — The cluster state model can be solved using the weak Hopf matrix product state developed in Refs. [42, 43, 55, 56]. Here, we will only provide the ground state for the cluster state model. For a more general cluster ladder model, the solution can be obtained based on the discussion in Ref. [56].

For edges on the smooth boundary, the local tensor is chosen as the comultiplication of the Haar integral $\lambda \in H_3$:

$$\Delta(\lambda) = \begin{bmatrix} \lambda^{(2)} \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \lambda^{(1)} \end{bmatrix}}.$$
 (25)

The local tensor for a bulk edge corresponds to $\Delta_2(\lambda)$:

$$\Delta(\lambda) = \frac{\lambda^{(3)}}{\lambda} \frac{\lambda}{\lambda^{(1)}} \frac{\lambda^{(2)}}{\lambda^{(1)}}.$$
 (26)

The local tensor for a face corresponds to the Haar measure $\Lambda \in \hat{H}_3$, which is introduced to glue the edge tensors together:

 $(\mathrm{id} \otimes \mathrm{id} \otimes \hat{S}) \circ \hat{\Delta}(\Lambda) = \bigwedge_{\Lambda^{(1)}}^{\hat{S}(\Lambda^{(3)})} \longrightarrow_{\Lambda^{(2)}}^{\Lambda^{(2)}} .$ (27)

The Haagerup cluster state is defined as the following ten-

sor network state:



The red free leg represents the physical degrees of freedom. This tensor network forms the ground state of the Haagerup cluster state model.

Conclusion and Discussion.- In this work, we propose a cluster state model that possesses Haagerup fusion category symmetry. The ground state and symmetry properties of the model are discussed in detail. Despite the progress made, there are still several directions to explore further: (i) The conformal field theory (CFT) perspective of the model needs to be studied. We have not addressed the central charge, entanglement properties, and other related aspects; however, we believe these can be investigated in a manner similar to that in Refs. [30-33], likely through numerical approaches. (ii) The gapless model is also an interesting direction to explore. We mainly discussed the gapped lattice model, but if we introduce a gapless physical boundary, the model becomes gapless. For anyonic chain-based models, this has been discussed in Ref. [32], and similar results should apply to our model. (iii) The application of the model in measurement-based quantum computation is another interesting avenue. For the finite group case, this has been partially discussed in Ref. [54], but for general non-Abelian groups, Hopf algebras, and weak Hopf algebras, the problem remains largely open. All these problems will be left for future work.

Acknowledgments. – I would like to thank Dagomir Kaszlikowski for his support. This work is supported by the National Research Foundation in Singapore and A*STAR under its CQT Bridging Grant and CQT- Return of PIs EOM YR1-10 Funding.

- D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, "Generalized global symmetries," Journal of High Energy Physics 2015, 1 (2015), arXiv:1412.5148 [hep-th].
- [2] K. Inamura, "Topological field theories and symmetry protected topological phases with fusion category symmetries," Journal of High Energy Physics 2021, 1 (2021), arXiv:2103.15588 [condmat.str-el].
- [3] L. Kong, T. Lan, X.-G. Wen, Z.-H. Zhang, and H. Zheng, "Algebraic higher symmetry and categorical symmetry: A holographic and entanglement view of symmetry," Phys. Rev. Res. 2, 043086 (2020), arXiv:2005.14178 [cond-mat.str-el].
- [4] S. Schäfer-Nameki, "ICTP lecture on (non-)invertible generalized symmetries," Physics Reports 1063, 1 (2024), arXiv:2305.18296 [hep-th].
- [5] S.-J. Huang and M. Cheng, "Topological holography, quantum criticality, and boundary states," (2023), arXiv:2310.16878 [cond-mat.str-el].
- [6] L. Bhardwaj, L. E. Bottini, S. Schafer-Nameki, and A. Tiwari,

"Lattice models for phases and transitions with non-invertible symmetries," (2024), arXiv:2405.05964 [cond-mat.str-el].

- [7] D. S. Freed, G. W. Moore, and C. Teleman, "Topological symmetry in quantum field theory," (2024), arXiv:2209.07471 [hep-th].
- [8] D. Gaiotto and J. Kulp, "Orbifold groupoids," Journal of High Energy Physics 2021, 1 (2021), arXiv:2008.05960 [hep-th].
- [9] L. Bhardwaj and S. Schafer-Nameki, "Generalized charges, part ii: Non-invertible symmetries and the symmetry TFT," (2023), arXiv:2305.17159 [hep-th].
- [10] F. Apruzzi, F. Bonetti, I. García Etxebarria, S. S. Hosseini, and S. Schäfer-Nameki, "Symmetry tfts from string theory," Communications in mathematical physics 402, 895 (2023), arXiv:2112.02092 [hep-th].
- [11] L. Bhardwaj, L. E. Bottini, D. Pajer, and S. Schafer-Nameki, "Gapped phases with non-invertible symmetries: (1+1)d," (2024), arXiv:2310.03784 [hep-th].
- [12] C. Zhang and C. Córdova, "Anomalies of (1+1)-dimensional

categorical symmetries," Phys. Rev. B 110, 035155 (2024).

- [13] W. Ji and X.-G. Wen, "Categorical symmetry and noninvertible anomaly in symmetry-breaking and topological phase transitions," Phys. Rev. Res. 2, 033417 (2020).
- [14] A. Chatterjee, W. Ji, and X.-G. Wen, "Emergent generalized symmetry and maximal symmetry-topological-order," (2024), arXiv:2212.14432 [cond-mat.str-el].
- [15] C.-M. Chang, Y.-H. Lin, S.-H. Shao, Y. Wang, and X. Yin, "Topological defect lines and renormalization group flows in two dimensions," Journal of High Energy Physics 2019, 1 (2019), arXiv:1802.04445 [hep-th].
- [16] L. Bhardwaj and Y. Tachikawa, "On finite symmetries and their gauging in two dimensions," Journal of High Energy Physics 2018, 1 (2018), arXiv:1704.02330 [hep-th].
- [17] L. Bhardwaj, L. E. Bottini, S. Schäfer-Nameki, and A. Tiwari, "Non-invertible higher-categorical symmetries," SciPost Phys. 14, 007 (2023), arXiv:2204.06564 [hep-th].
- [18] R. Thorngren and Y. Wang, "Fusion category symmetry. part i. anomaly in-flow and gapped phases," Journal of High Energy Physics 2024, 1 (2024), arXiv:1912.02817 [hep-th].
- [19] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, *Tensor categories*, Vol. 205 (American Mathematical Soc., 2016) pp. xvi+343.
- [20] G. Moore and N. Seiberg, "Classical and quantum conformal field theory," Communications in Mathematical Physics 123, 177 (1989).
- [21] V. F. Jones, "Von neumann algebras in mathematics and physics," International Congress of Mathematicians, Kyoto, Japan, August 1990.
- [22] M. Bischoff, "The relation between subfactors arising from conformal nets and the realization of quantum doubles," (2015), arXiv:1511.08931 [math-ph].
- [23] M. Bischoff, "A remark on cft realization of quantum doubles of subfactors: Case index \$\${ ; 4}\$\$," Letters in Mathematical Physics 106, 341 (2016), arXiv:1506.02606 [math-ph].
- [24] F. Xu, "Examples of subfactors from conformal field theory," Communications in Mathematical Physics 357, 61 (2018), arXiv:1608.04119 [math.OA].
- [25] F. Calegari, S. Morrison, and N. Snyder, "Cyclotomic integers, fusion categories, and subfactors," Communications in Mathematical Physics 303, 845 (2011), arXiv:1004.0665 [math.NT].
- [26] U. Haagerup, "Principal graphs of subfactors in the index range $4 < [m:N] < 3 + \sqrt{2}$," Subfactors (Kyuzeso, 1993) **1** (1994).
- [27] M. Asaeda and U. Haagerup, "Exotic subfactors of finite depth with jones indices and," Communications in Mathematical Physics 202, 1 (1999).
- [28] M. Asaeda and S. Yasuda, "On haagerup's list of potential principal graphs of subfactors," Communications in Mathematical Physics 286, 1141 (2009), arXiv:0711.4144 [math.OA].
- [29] R. Vanhove, L. Lootens, M. Van Damme, R. Wolf, T. J. Osborne, J. Haegeman, and F. Verstraete, "Critical lattice model for a haagerup conformal field theory," Phys. Rev. Lett. 128, 231602 (2022), arXiv:2110.03532 [cond-mat.stat-mech].
- [30] T.-C. Huang, Y.-H. Lin, K. Ohmori, Y. Tachikawa, and M. Tezuka, "Numerical evidence for a haagerup conformal field theory," Phys. Rev. Lett. **128**, 231603 (2022), arXiv:2110.03008 [cond-mat.stat-mech].
- [31] L. Corcoran and M. de Leeuw, "Integrable and critical haagerup spin chains," (2024), arXiv:2410.16356 [cond-mat.stat-mech].
- [32] L. E. Bottini and S. Schafer-Nameki, "A gapless phase with haagerup symmetry," (2024), arXiv:2410.19040 [hep-th].
- [33] Y. Liu, Y. Zou, and S. Ryu, "Operator fusion from wavefunction overlap: Universal finite-size corrections and application to the haagerup model," Phys. Rev. B 107, 155124 (2023),

arXiv:2203.14992 [cond-mat.stat-mech].

- [34] T.-C. Huang and Y.-H. Lin, "Topological field theory with haagerup symmetry," Journal of Mathematical Physics 63, 042306 (2022), arXiv:https://pubs.aip.org/aip/jmp/articlepdf/doi/10.1063/5.0079062/16561548/042306_1_online.pdf [hep-th].
- [35] P. Grossman and N. Snyder, "Quantum subgroups of the haagerup fusion categories," Communications in Mathematical Physics 311, 617 (2012), arXiv:1102.2631 [math.OA].
- [36] D. E. Evans and T. Gannon, "The exoticness and realisability of twisted haagerup-izumi modular data," Communications in Mathematical Physics **307**, 463 (2011), arXiv:1006.1326 [math.OA].
- [37] A. Kitaev, "Fault-tolerant quantum computation by anyons," Annals of Physics 303, 2 (2003), arXiv:quant-ph/9707021 [quant-ph].
- [38] A. F. Bais, B. J. Schroers, and J. K. Slingerland, "Hopf symmetry breaking and confinement in (2+1)-dimensional gauge theory," Journal of High Energy Physics 2003, 068 (2003), arXiv:hep-th/0205114 [hep-th].
- [39] C. Meusburger, "Kitaev lattice models as a Hopf algebra gauge theory," Communications in Mathematical Physics 353, 413 (2017), arXiv:1607.01144 [math.QA].
- [40] O. Buerschaper, J. M. Mombelli, M. Christandl, and M. Aguado, "A hierarchy of topological tensor network states," Journal of Mathematical Physics 54, 012201 (2013), arXiv:1007.5283 [cond-mat.str-el].
- [41] F. Girelli, P. K. Osei, and A. Osumanu, "Semidual Kitaev lattice model and tensor network representation," Journal of High Energy Physics 2021, 1 (2021), arXiv:1709.00522 [math.QA].
- [42] Z. Jia, D. Kaszlikowski, and S. Tan, "Boundary and domain wall theories of 2d generalized quantum double model," Journal of High Energy Physics 2023, 1 (2023), arXiv:2207.03970 [quant-ph].
- [43] Z. Jia, S. Tan, D. Kaszlikowski, and L. Chang, "On weak Hopf symmetry and weak Hopf quantum double model," Communications in Mathematical Physics 402, 3045 (2023), arXiv:2302.08131 [hep-th].
- [44] Z. Jia, S. Tan, and D. Kaszlikowski, "Weak Hopf symmetry and tube algebra of the generalized multifusion stringnet model," Journal of High Enegy Physics 07, 207 (2024), arXiv:2403.04446 [hep-th].
- [45] C. Cordova, T. T. Dumitrescu, K. Intriligator, and S.-H. Shao, "Snowmass white paper: Generalized symmetries in quantum field theory and beyond," (2022), arXiv:2205.09545 [hep-th].
- [46] T. D. Brennan and S. Hong, "Introduction to generalized global symmetries in QFT and particle physics," (2023), arXiv:2306.00912 [hep-ph].
- [47] J. McGreevy, "Generalized symmetries in condensed matter," Annual Review of Condensed Matter Physics 14, 57 (2023), arXiv:2204.03045 [cond-mat.str-el].
- [48] R. Luo, Q.-R. Wang, and Y.-N. Wang, "Lecture notes on generalized symmetries and applications," Physics Reports 1065, 1 (2024), arXiv:2307.09215 [hep-th].
- [49] S.-H. Shao, "What's done cannot be undone: Tasi lectures on non-invertible symmetries," (2024), arXiv:2308.00747 [hepth].
- [50] L. Bhardwaj, L. E. Bottini, L. Fraser-Taliente, L. Gladden, D. S. Gould, A. Platschorre, and H. Tillim, "Lectures on generalized symmetries," Physics Reports 1051, 1 (2024), arXiv:2307.07547 [hep-th].
- [51] W. Son, L. Amico, and V. Vedral, "Topological order in 1d cluster state protected by symmetry," Quantum Information Processing 11, 1961 (2012), arXiv:1111.7173 [quant-ph].

- [52] S. Seifnashri and S.-H. Shao, "Cluster state as a noninvertible symmetry-protected topological phase," Phys. Rev. Lett. 133, 116601 (2024), arXiv:2404.01369 [cond-mat.str-el].
- [53] C. G. Brell, "Generalized cluster states based on finite groups," New Journal of Physics 17, 023029 (2015), arXiv:1408.6237 [quant-ph].
- [54] C. Fechisin, N. Tantivasadakarn, and V. V. Albert, "Noninvertible symmetry-protected topological order in a groupbased cluster state," (2023), arXiv:2312.09272 [cond-mat.strel].
- [55] Z. Jia, "Generalized cluster states from hopf algebras: noninvertible symmetry and hopf tensor network representation," Journal of High Energy Physics 2024, 147 (2024), arXiv:2405.09277 [quant-ph].
- [56] Z. Jia, "Weak hopf non-invertible symmetry-protected topological spin liquid and lattice realization of (1+1)d symmetry topological field theory," (2024), arXiv:2412.15336 [hep-th].
- [57] T. J. Osborne, D. E. Stiegemann, and R. Wolf, "The fsymbols for the h3 fusion category," (2019), arXiv:1906.01322 [math.CT].
- [58] T.-C. Huang and Y.-H. Lin, "The *F*-symbols for transparent haagerup-izumi categories with $g = \mathbb{Z}_{2n+1}$," (2021), arXiv:2007.00670 [math.CT].
- [59] S.-M. Hong, E. Rowell, and Z. Wang, "On exotic modular tensor categories," Communications in Contemporary Mathematics 10, 1049 (2008), arXiv:0710.5761 [math.GT].
- [60] G. Böhm, F. Nill, and K. Szlachányi, "Weak Hopf algebras: I. Integral theory and C*-structure," Journal of Algebra 221, 385 (1999), arXiv:math/9805116 [math.QA].

- [61] P. Etingof, D. Nikshych, and V. Ostrik, "On fusion categories," Annals of Mathematics , 581 (2005), arXiv:math/0203060 [math.QA].
- [62] D. Nikshych, "Semisimple weak hopf algebras," Journal of Algebra 275, 639 (2004), arXiv:math/0304098 [math.QA].
- [63] K. Szlachányi, "Finite quantum groupoids and inclusions of finite type," in *Mathematical physics in mathematics and physics: quantum and operator algebraic aspects*, Fields Institute Communications, Vol. 30 (American Mathematical Soc., 2001) p. 393–407, arXiv:math/0011036 [math.QA].
- [64] V. Ostrik, "Module categories, weak Hopf algebras and modular invariants," Transformation Groups 8, 177 (2003), arXiv:math/0111139 [math.QA].
- [65] A. Kitaev and L. Kong, "Models for gapped boundaries and domain walls," Communications in Mathematical Physics 313, 351 (2012), arXiv:1104.5047 [cond-mat.str-el].
- [66] J. C. Bridgeman, L. Lootens, and F. Verstraete, "Invertible bimodule categories and generalized Schur orthogonality," Communications in Mathematical Physics 402, 2691 (2023), arXiv:2211.01947 [math.QA].
- [67] Z. Jia and S. Tan, "Weak Hopf tube algebra for domain walls between 2d gapped phases of Turaev-Viro TQFTs," in preparation.
- [68] G. Böhm, "Doi-hopf modules over weak Hopf algebras," Communications in Algebra 28, 4687 (2000), arXiv:math/9905027 [math.QA].
- [69] L. Chang, "Kitaev models based on unitary quantum groupoids," Journal of Mathematical Physics 55, 041703 (2014), arXiv:1309.4181 [math.QA].