Minimal dark matter in SU(5) grand unification

Takashi Toma^{1,2‡}

¹Institute for Theoretical Physics, Kanazawa University, Kanazawa 920-1192, Japan ²Institute of Liberal Arts and Science, Kanazawa University, Kanazawa 920-1192, Japan

Abstract

Minimal dark matter is an attractive candidate for dark matter because it is stabilized without the need to impose additional symmetries. It is known that the mass of the $SU(2)_L$ quintuplet fermion dark matter is predicted to be around 14 TeV, based on the thermal production mechanism. In this work, we embed the quintuplet dark matter within non-supersymmetric SU(5) grand unified theories. We find that two pairs of colored sextet fermions are required at the $\mathcal{O}(1-10)$ TeV scale to achieve gauge coupling unification, with the unification scale near the reduced Planck scale. These colored sextet fermions become metastable because their interactions are suppressed by the unification scale. Our model can be tested through comprehensive searches for colored sextet fermions in collider experiments, as well as through indirect and direct detection methods for minimal dark matter.

[‡]toma@staff.kanazawa-u.ac.jp

1 Introduction

Dark matter is known to exist in the universe, but its fundamental properties, such as mass, interactions, and production mechanism, remain unknown. One of the most promising candidates for dark matter is thermal dark matter, which is produced through the freeze-out mechanism. Despite extensive searches via direct detection, indirect detection, and collider experiments, no signals have been detected so far. Consequently, experimental constraints on interactions with Standard Model (SM) particles have become increasingly stringent. In particular, direct detection experiments place severe limits on the spin-independent cross section, with the current strongest bound provided by the LZ experiment [1, 2]. Even under these constraints, thermal dark matter scenarios remain appealing due to their strong predictiveness and the successful description of the universe's thermal history below MeV scales. For example, candidates such as minimal dark matter [3], including the wino in supersymmetric models in a broader sense, pseudo-Nambu-Goldstone dark matter [4], and pseudo-scalar interacting fermionic dark matter [5,6], continue to align with current experimental and observational bounds.

Minimal dark matter, in particular, can be naturally incorporated through simple extensions of the SM, such as introducing an $SU(2)_L$ quintuplet fermion or a septuplet scalar [3]. The interactions of these particles with the SM are highly constrained by gauge symmetries, leading to an accidental \mathbb{Z}_2 symmetry that stabilizes the lightest neutral state in the quintuplet or septuplet at the renormalizable level.

Separately, models that unify the SM field content and gauge interactions known as grand unified theories (GUTs) have been studied for decades. The SU(5) symmetry is well-established as the minimal GUT framework, embedding all SM matter particles within the $\overline{5}$ and 10 representations [7]. However, it is also well-known that gauge coupling unification is not realized in the minimal SU(5) GUT model when considering only SM particles.

In this work, we propose a simple embedding of the quintuplet minimal dark matter and all other matter particles into the SU(5) gauge symmetry.¹ We assume that the scale of new light particles in the multiplets is fixed by the dark matter mass, approximately 14 TeV, to reproduce the thermal relic abundance via the freeze-out mechanism, including the effects of Sommerfeld enhancement [10–12] and bound state formation [13]. To achieve gauge coupling unification, we will find that two additional $SU(3)_c$ sextet fermions, which can be embedded in a **15** representation, are required. These exotic sextet fermions are metastable due to interactions suppressed by the unification scale and can form bound states with quarks and gluons, which may be detectable in collider experiments. Together with gamma-ray observations and direct detection experiments for the quintuplet fermion

¹Alternative attempts to embed minimal dark matter in SO(10) GUTs have been explored [8], and a comprehensive study of dark matter stability in SO(10) unification has also been conducted [9].

dark matter, these collider searches provide a comprehensive test of our unified model.

2 Minimal dark matter in SU(5)

The Lagrangian for the fermionic $SU(2)_L$ quintuplet χ is given by

$$\mathcal{L} = \frac{1}{2} \overline{\chi} \left(i D - M_5 \right) \chi, \qquad (2.1)$$

where M_5 is the tree-level mass [3]. The other interactions are forbidden at the renormalizable level by SM gauge symmetry. An accidental \mathbb{Z}_2 symmetry ensures the stability of the lightest component in the quintuplet. At tree level, all components in the quintuplet are degenerate in mass. However, small mass splitting between the charged and neutral components ($\approx 166 \text{ MeV}$) arises at the loop level [3]. Consequently, the neutral component χ^0 becomes the lightest state and is identified as the dark matter.

The mass M_5 is the sole parameter of minimal dark matter, as all interactions are determined by the gauge symmetry. The dark matter mass is constrained by the requirement to reproduce the observed relic abundance through the thermal freeze-out mechanism. The dominant annihilation channels are $\chi^0 \chi^0 \to W^+ W^-$, $ZZ, Z\gamma$ and $\gamma\gamma$, mediated by gauge bosons. Accounting for the effects of Sommerfeld enhancement and bound state formation, the mass is determined to be $M_5 = 14$ TeV [13].

To further investigate, we propose embedding the minimal dark matter within the SU(5) gauge symmetry. In the Georgi-Glashow model [7], all SM fermions in a single generation are embedded in the $\overline{\mathbf{5}}_F$ and $\mathbf{10}_F$ representations, while the SM Higgs doublet H resides in the $\mathbf{5}_H$ representation. Additionally, the $\mathbf{24}_H$ adjoint Higgs field is introduced to achieve the symmetry-breaking pattern $SU(5) \rightarrow SU(3)_c \times SU(2)_L \times U(1)_Y$ via its vacuum expectation value (VEV). Beyond the Georgi-Glashow model, we introduce a **200** fermion representation, which is the smallest representation containing an $SU(2)_L$ quintuplet. The **200** representation can be decomposed under the $SU(3)_c \times SU(2)_L \times U(2)_L \times U(1)_Y$ subgroup as follows [14–16]:

$$\begin{aligned} \mathbf{200} &= \quad (\mathbf{6}, \mathbf{3})_{-5/3} + (\mathbf{15}, \mathbf{2})_{-5/6} + (\mathbf{3}, \mathbf{4})_{-5/6} + (\mathbf{3}, \mathbf{2})_{-5/6} \\ &+ (\mathbf{27}, \mathbf{1})_0 + (\mathbf{8}, \mathbf{3})_0 + (\mathbf{8}, \mathbf{1})_0 + (\mathbf{1}, \mathbf{5})_0 + (\mathbf{1}, \mathbf{3})_0 + (\mathbf{1}, \mathbf{1})_0 \\ &+ (\overline{\mathbf{6}}, \mathbf{3})_{5/3} + (\overline{\mathbf{15}}, \mathbf{2})_{5/6} + (\overline{\mathbf{3}}, \mathbf{4})_{5/6} + (\overline{\mathbf{3}}, \mathbf{2})_{5/6}, \end{aligned}$$
(2.2)

The introduction of the **200** representation ensures the incorporation of the quintuplet minimal dark matter into the SU(5) framework, providing a natural extension of the Georgi-Glashow model. Further exploration of this embedding will allow for a comprehensive analysis of its phenomenological implications, including gauge coupling unification and the search for additional states predicted by the model.

3 Gauge coupling unification

To investigate gauge coupling unification, we compute the β -functions for the gauge couplings g_i (i = 1, 2, 3) at the two-loop level. The gauge couplings are normalized as $g_1 = \sqrt{5/3}g_Y$. The renormalization group equation (RGE) for the gauge couplings is expressed as [17–21]

$$\frac{dg_i}{dt} = \frac{b_i g_i^3}{(4\pi)^2} + \sum_{j=1}^3 \frac{b_{ij} g_i^3 g_j^2}{(4\pi)^4} + \frac{c_i g_i^3 y_t^2}{(4\pi)^4},\tag{3.1}$$

where $t = \log(\mu/m_Z)$, y_t is the top Yukawa coupling, and the coefficients b_i , b_{ij} , and c_i are given by

$$b_{i} = \begin{pmatrix} 41/10 \\ -19/6 \\ -7 \end{pmatrix}, \quad b_{ij} = \begin{pmatrix} 199/50 & 27/10 & 44/5 \\ 9/10 & 35/6 & 12 \\ 11/10 & 9/2 & -26 \end{pmatrix}, \quad c_{i} = \begin{pmatrix} -17/10 \\ -3/2 \\ -2 \end{pmatrix}, \quad (3.2)$$

The contributions of other Yukawa couplings are neglected due to their small magnitudes compared to y_t .

The RGE for the top Yukawa coupling y_t is given by

$$\frac{dy_t}{dt} = \frac{\beta_{y_t}^{(1)}}{(4\pi)^2} + \frac{\beta_{y_t}^{(2)}}{(4\pi)^4},\tag{3.3}$$

where the one-loop $(\beta_{y_t}^{(1)})$ and two-loop $(\beta_{y_t}^{(2)})$ contributions are

$$\beta_{y_t}^{(1)} = \frac{9}{2} y_t^3 - y_t \left(\frac{17}{20} g_1^2 + \frac{9}{4} g_2^2 + 8g_3^2 \right), \tag{3.4}$$

$$\beta_{y_t}^{(2)} = y_t \left(-12y_t^4 - 6y_t^2 \lambda + \frac{3}{2}\lambda^2 \right) + y_t^3 \left(\frac{393}{80}g_1^2 + \frac{225}{16}g_2^2 + 36g_3^2 \right) + y_t \left(\frac{1187}{600}g_1^4 - \frac{9}{20}g_1^2g_2^2 + \frac{19}{15}g_1^2g_3^2 - \frac{23}{4}g_2^4 + 9g_2^2g_3^2 - 108g_3^4 \right).$$
(3.5)

Here, λ is the Higgs self-coupling normalized as $\mathcal{L} \supset -\frac{\lambda}{2}|H|^4$.

The RGE for the Higgs self-coupling λ is given by

$$\frac{d\lambda}{dt} = \frac{\beta_{\lambda}^{(1)}}{(4\pi)^2} + \frac{\beta_{\lambda}^{(2)}}{(4\pi)^4},$$
(3.6)

where the one-loop $(\beta_{\lambda}^{(1)})$ and two-loop $(\beta_{\lambda}^{(2)})$ contributions are

$$\beta_{\lambda}^{(1)} = 12\left(\lambda^{2} + y_{t}^{2}\lambda - y_{t}^{4}\right) - \lambda\left(\frac{9}{5}g_{1}^{2} + 9g_{2}^{2}\right) + \frac{9}{4}\left(\frac{3}{25}g_{1}^{4} + \frac{2}{5}g_{1}^{2}g_{2}^{2} + g_{2}^{4}\right), \qquad (3.7)$$
$$\beta_{\lambda}^{(2)} = -78\lambda^{3} - 72\lambda^{2}y_{t}^{2} + 18\lambda^{2}\left(\frac{3}{5}g_{1}^{2} + 3g_{2}^{2}\right) - 3\lambda y_{t}^{4} + 10\lambda y_{t}^{2}\left(\frac{17}{20}g_{1}^{2} + \frac{9}{4}g_{2}^{2} + 8g_{3}^{2}\right)$$

$$-\lambda \left(\frac{73}{8}g_2^4 - \frac{117}{20}g_1^2g_2^2 - \frac{1887}{200}g_1^4\right) - \frac{3411}{1000}g_1^6 - \frac{289}{40}g_1^2g_2^4 - \frac{1677}{200}g_1^4g_2^2 + \frac{305}{8}g_2^6 + 60y_t^6 - \frac{16}{5}y_t^4\left(g_1^2 + 20g_3^2\right) - \frac{9}{50}y_t^2\left(19g_1^4 - 70g_1^2g_2^2 + 25g_2^4\right).$$
(3.8)

The quintuplet introduces additional contributions to the β -functions in the energy scale above its mass M_5 . These corrections are given by

$$\Delta b_2 = \frac{20}{3}, \quad \Delta b_{22} = \frac{560}{3}, \quad \Delta \beta_{y_t}^{(2)} = 5g_2^4, \quad \Delta \beta_{\lambda}^{(2)} = -2g_2^4 \left(4g_1^2 + 20g_2^2 - 25\lambda\right). \tag{3.9}$$

We used SARAH to calculate the β -functions at the two loop level [22,23]. These contributions indicate that the gauge couplings do not unify at any scale. Furthermore, following these renormalization group equations, it is observed that a Landau pole for the g_2 coupling appears at a scale slightly above the Planck scale. This finding imposes a strong constraint: no additional particles charged under the $SU(2)_L$ gauge symmetry can be introduced without exacerbating this issue.

To achieve gauge coupling unification, two pairs of $SU(3)_c$ sextet fermions are introduced: $(\mathbf{6}, \mathbf{1})_{-2/3}$ and $(\mathbf{\overline{6}}, \mathbf{1})_{2/3}$. These fermions can originate from the $\mathbf{15}_F$ and $\mathbf{\overline{15}}_F$ representations of SU(5), whose branching rules under $SU(3)_c \times SU(2)_L \times U(1)_Y$ are given by [14]

$$\mathbf{15} = (\mathbf{6}, \mathbf{1})_{-2/3} + (\mathbf{3}, \mathbf{2})_{1/6} + (\mathbf{1}, \mathbf{3})_1, \tag{3.10}$$

$$\overline{\mathbf{15}} = (\overline{\mathbf{6}}, \mathbf{1})_{2/3} + (\overline{\mathbf{3}}, \mathbf{2})_{-1/6} + (\mathbf{1}, \mathbf{3})_{-1}.$$
(3.11)

A pair of sextet fermions contributes to the β -functions as follows:

$$\Delta b_1 = \frac{32}{15}, \quad \Delta b_{11} = \frac{128}{75}, \quad \Delta b_{13} = \frac{64}{3}, \tag{3.12}$$

$$\Delta b_3 = \frac{10}{3}, \quad \Delta b_{31} = \frac{8}{3}, \qquad \Delta b_{33} = \frac{250}{3}, \tag{3.13}$$

$$\Delta\beta_{y_t}^{(2)} = \frac{232}{225}g_1^4 + \frac{200}{9}g_3^4, \qquad \Delta\beta_{\lambda}^{(2)} = -\frac{16}{125}g_1^4 \left(12g_1^2 + 20g_2^2 - 25\lambda\right). \tag{3.14}$$

The masses of these sextet fermions are denoted as M_6 and M_6 . The contributions to the β -functions are included at the energy scales $\mu = M_6$ and $\mu = \tilde{M}_6$, respectively.

To investigate gauge coupling unification, we numerically solve the RGEs at the twoloop level. The integration is performed over the energy range $m_Z \leq \mu \leq M_U$, where M_U is the unification scale. The unified gauge coupling α_U^{-1} at the unification scale M_U is determined by minimizing the quantity [24]

$$\Delta \alpha \equiv \sqrt{\Delta \alpha_{12}^{-2}(M_U) + \Delta \alpha_{23}^{-2}(M_U)}, \qquad (3.15)$$

where $\Delta \alpha_{ij}^{-1} = \alpha_i^{-1} - \alpha_j^{-1}$, and $\alpha_i = g_i^2/(4\pi)$ are the gauge couplings. The gauge couplings at the reference scale $\mu = m_Z$ are determined using physical constants from experimental



Figure 1: (Left): Contours of $\Delta \alpha$ in M_6 - \tilde{M}_6 plane. The red line indicates the region where $\Delta \alpha$ is minimized, representing the best gauge coupling unification, and the gray region is excluded based on current experimental bounds from colored fermion searches at the LHC [26]. (Right): Example of the gauge coupling unification where the sextet fermion masses are fixed to be $M_6 = 2$ TeV and $\tilde{M}_6 = 9.7$ TeV. The gauge couplings unify at a scale $M_U \approx 2.1 \times 10^{18}$ GeV, close to the Planck scale, and the unified gauge coupling at $\mu = M_U$ is predicted as $\alpha_U^{-1} \approx 9.0$.

measurements [25]. Specifically: $\alpha_1 = 58.81$, $\alpha_2 = 29.48$ and $\alpha_3 = 8.475$. These values are derived using the following physical constants:

$$m_W = 80.377 \text{ GeV}, \quad G_F = 1.1663788 \text{ GeV}^{-2},$$

 $\sin^2 \theta_W = 0.23121, \qquad \alpha_s = 0.1180.$

The contours of $\Delta \alpha$ are shown in the left panel of Fig. 1 where the quintuplet fermion mass is fixed to be $M_5 = 14$ TeV. The optimal unification ($\Delta \alpha \approx 0.03$) is achieved when the sextet fermion masses satisfy the relation

$$\sqrt{M_6 \tilde{M}_6} \approx 4.3 \text{ TeV.}$$
 (3.16)

Under this condition, the unification scale and the unified gauge coupling are determined to be $M_U = 2.1 \times 10^{18}$ GeV and $\alpha_U^{-1} = 9.0.^2$ Notably, the unification scale is very close to the reduced Planck scale $M_P = 2.4 \times 10^{18}$ GeV.³ At such high energy scales, quantum gravity effects are expected to become significant and could influence the coupling unification. Indeed, the energy scale E_{new} at which tree-level unitarity breaks

²In fact, there is very small preference for the unification at $M_6 = \tilde{M}_6 = 4.3$ TeV. However the difference is invisible.

 $^{^{3}}$ A possibility of gauge coupling unification at the Planck scale has been studied [27]. Additionally, a mechanism for neutrino mass generation from the Planck scale has also been explored [28, 29].

down can be estimated as $E_{\text{new}}^2 = 20(G_N N)^{-1}$ where G_N is the Newton constant and $N = \frac{2}{3}N_s + N_f + 4N_V$ accounts for the number of complex scalars, Weyl fermions and real vector bosons [30]. In our model, we can obtain $E_{\text{new}} \approx 2.7 \times 10^{18}$ GeV, which is slightly exceeding the unification scale. In the purple region ($\Delta \alpha \leq 0.50$), the gauge coupling unification could still be acceptable. An example running of the gauge couplings is shown in Fig. 1 where the sextet fermion masses are fixed to be $M_6 = 2$ TeV and $\tilde{M}_6 = 9.7$ TeV, respectively. We have checked that the unification scale and the unified gauge coupling are almost constant along with the red line, and they change within the range of 1.7×10^{18} GeV $\lesssim M_U \lesssim 2.8 \times 10^{18}$ GeV and $8.4 \lesssim \alpha_U^{-1} \lesssim 9.6$ in the purple region. It should be noted that threshold corrections from heavy particles could influence the running of the gauge couplings. However, incorporating these effects is beyond the scope of this work.

using the unification scale M_U and unified gauge coupling strength α_U^{-1} , the proton lifetime can be roughly estimated as $\tau_p = \alpha_U^{-2} M_U^4 / m_p^5 \approx 3.9 \times 10^{39}$ years where m_p is the proton mass. The predicted lifetime is significantly longer than the current experimental upper bound $\tau_p^{\exp} = 2.4 \times 10^{34}$ years for the $p \to e^+ \pi^0$ decay channel, as reported by Super-Kamiokande [31]. Furthermore, it exceeds the expected future sensitivity of Hyper-Kamiokande, which is on the order of $\mathcal{O}(10^{35})$ years [32].

4 Experimental tests

Our model predicts the presence of sextet fermions at the energy scale of $\mathcal{O}(1 - 10)$ TeV. Such exotic colored fermions can be pair-produced through gluon fusion and quark-antiquark annihilations and searched for at the LHC [26] (see also [33,34]). Since the interactions that induce their decays are suppressed by the unification scale, these sextet fermions are metastable. They form *R*-hadrons, which are bound states with quarks and gluons. At the LHC, these metastable particles can be detected via their large ionization energy loss, akin to the long-lived gluinos in supersymmetric models. Although no significant deviations from the SM have been observed, upper bounds on the production cross section of long-lived gluinos have been derived [35, 36]. These bounds translate to a conservative mass limit of 1.58 TeV for the sextet fermions in our model. Discrimination between sextet fermions and gluinos (octets) could be achieved through detailed calculations of their production cross sections.

Meanwhile, the quintuplet fermion dark matter can be probed via high-energy gammaray observations, such as those by MAGIC [37] and CTA [38]. The annihilation cross section of the dark matter is expected to be significantly enhanced due to Sommerfeld enhancement and bound-state formation effects [13]. For instance, at the dark matter mass of $M_5 = 14$ TeV and a relative velocity of $v = 2 \times 10^{-3}$ (assuming a Burkert density profile), the annihilation cross sections for $\chi^0 \chi^0 \to W^+ W^-$ and $\gamma \gamma$ are predicted to be approximately 10^{-24} cm³/s and 2×10^{-26} cm³/s, respectively. These values are within the detection capabilities of CTA [38].

The quintuplet dark matter can also be probed through direct detection experiments. The strongest current constraint on the spin-independent cross section comes from LZ [2], with the bound of approximately 4.2×10^{-46} cm² extrapolated to a dark matter mass of 14 TeV. This value is close to the theoretical prediction of 1.0×10^{-46} cm² [39]. Therefore, our model can be thoroughly tested via a combination of searches for exotic colored fermions and minimal dark matter in the $\mathcal{O}(10)$ TeV range.

5 Additional comments

We must consider the possible new interactions between the SM particles and the exotic sextet fermions. In particular, one can write the following gauge-invariant Yukawa couplings:

$$\mathcal{L} \supset Y \mathbf{24}_H \mathbf{10}_F \mathbf{15}_F + \tilde{Y} \mathbf{5}_H \mathbf{5}_F \mathbf{15}_F.$$
 (5.1)

We have checked that all additional couplings between the sextet fermions and quarks are suppressed by the unification scale.

It is well-known that the Higgs potential in the Standard Model becomes unstable at high energy scales. This issue persists in our model, as we have introduced only fermions and not scalars. However, this could be mitigated by adding additional gauge-singlet scalars and mixed quartic couplings. These gauge-singlet scalars would not affect the gauge coupling unification.

Even if we assume that the triplet $(1, 3)_0$ in 200 is light instead of the quintuplet, it can also serve as stable dark matter, similar to the Wino dark matter in supersymmetric models. In this case, gauge coupling unification can be achieved by adding a $(8, 1)_0$ fermion and a $(1, 3)_0$ scalar at the $\mathcal{O}(1)$ TeV scale [40–42]. These extra fields originate from the existing representations of 200_F and the 24_H adjoint Higgs in our model, respectively. The triplet fermion dark matter mass is predicted to be $M_3 = 2.7$ TeV in order to reproduce the observed relic abundance via the thermal freeze-out mechanism [12,13]. The octet fermion shares the same quantum numbers as the gluino in supersymmetric models, so we can apply the gluino mass bound to the triplet model. The current lower bound on the gluino mass is 2.3 TeV [43–45].

For the $(\mathbf{1}, \mathbf{3})_0$ fermion, the lower mass bound is obtained at the LHC and is 790 GeV [46]. For the triplet scalar, the relevant mass bound can be inferred from the charged Higgs boson decaying into $\Delta^{\pm} \rightarrow t\bar{b}$. This bound is approximately 1 TeV [47]. The charged scalar in the triplet can mix with the Goldstone boson absorbed by the W boson in the SM via the cubic coupling $\mathbf{24}_H \mathbf{5}_H \mathbf{\overline{5}}_H$ after electroweak symmetry breaking.

6 Summary

The quintuplet fermion dark matter is embedded in a **200** representation of the nonsupersymmetric SU(5) model, with a mass fixed at 14 TeV to reproduce the thermal relic abundance. Gauge coupling unification is achieved without introducing new intermediate scales by adding two pairs of colored sextet fermions, which are embedded in the **15** and $\overline{15}$ representations. It was found that the unification scale is very close to the reduced Planck scale, ensuring the stability of protons. This model can be explored through comprehensive searches for the metastable exotic sextet fermions and quintuplet fermion dark matter at the O(1 - 10) TeV scale. The colored sextet fermions can be probed at collider experiments if their mass is less than a few TeV. In principle, they can be distinguished from other exotic colored particles, such as gluinos in supersymmetric models or leptoquarks, due to differences in the magnitudes of their production cross sections. Additionally, the minimal dark matter can be searched for and experimentally identified by combining results from indirect and direct detection experiments, such as those from CTA and LZ, at the 14 TeV scale.

Acknowledgments

This work was supported by a JSPS Grant-in-Aid for Scientific Research KAKENHI Grant No. 23H04004.

References

- LZ Collaboration, J. Aalbers *et al.*, "First Dark Matter Search Results from the LUX-ZEPLIN (LZ) Experiment," *Phys. Rev. Lett.* 131 no. 4, (2023) 041002, arXiv:2207.03764 [hep-ex].
- [2] S. Haselschwardt, "New Dark Matter Search Results from the LUX-ZEPLIN (LZ) Experiment," https://indico.uchicago.edu/event/427/contributions/1325/ attachments/359/548/lz_results_tevpa.pdf.
- [3] M. Cirelli, N. Fornengo, and A. Strumia, "Minimal dark matter," *Nucl. Phys. B* 753 (2006) 178–194, arXiv:hep-ph/0512090.
- [4] C. Gross, O. Lebedev, and T. Toma, "Cancellation Mechanism for Dark-Matter-Nucleon Interaction," *Phys. Rev. Lett.* **119** no. 19, (2017) 191801, arXiv:1708.02253 [hep-ph].
- [5] M. Freytsis and Z. Ligeti, "On dark matter models with uniquely spin-dependent detection possibilities," *Phys. Rev. D* 83 (2011) 115009, arXiv:1012.5317 [hep-ph].

- [6] T. Abe, M. Fujiwara, and J. Hisano, "Loop corrections to dark matter direct detection in a pseudoscalar mediator dark matter model," *JHEP* 02 (2019) 028, arXiv:1810.01039 [hep-ph].
- [7] H. Georgi and S. L. Glashow, "Unity of All Elementary Particle Forces," *Phys. Rev. Lett.* 32 (1974) 438–441.
- [8] G.-C. Cho, K. Hayami, and N. Okada, "SO(10) grand unification with minimal dark matter and color octet scalars," *Phys. Rev. D* 105 no. 1, (2022) 015027, arXiv:2110.03884 [hep-ph].
- [9] S. Ferrari, T. Hambye, J. Heeck, and M. H. G. Tytgat, "SO(10) paths to dark matter," *Phys. Rev. D* 99 no. 5, (2019) 055032, arXiv:1811.07910 [hep-ph].
- [10] J. Hisano, S. Matsumoto, and M. M. Nojiri, "Explosive dark matter annihilation," *Phys. Rev. Lett.* **92** (2004) 031303, arXiv:hep-ph/0307216.
- [11] J. Hisano, S. Matsumoto, M. M. Nojiri, and O. Saito, "Non-perturbative effect on dark matter annihilation and gamma ray signature from galactic center," *Phys. Rev. D* 71 (2005) 063528, arXiv:hep-ph/0412403.
- [12] J. Hisano, S. Matsumoto, M. Nagai, O. Saito, and M. Senami, "Non-perturbative effect on thermal relic abundance of dark matter," *Phys. Lett. B* 646 (2007) 34–38, arXiv:hep-ph/0610249.
- [13] A. Mitridate, M. Redi, J. Smirnov, and A. Strumia, "Cosmological Implications of Dark Matter Bound States," JCAP 05 (2017) 006, arXiv:1702.01141 [hep-ph].
- [14] N. Yamatsu, "Finite-Dimensional Lie Algebras and Their Representations for Unified Model Building," arXiv:1511.08771 [hep-ph].
- [15] R. Feger, T. W. Kephart, and R. J. Saskowski, "LieART 2.0 A Mathematica application for Lie Algebras and Representation Theory," *Comput. Phys. Commun.* 257 (2020) 107490, arXiv:1912.10969 [hep-th].
- [16] R. M. Fonseca, "GroupMath: A Mathematica package for group theory calculations," *Comput. Phys. Commun.* 267 (2021) 108085, arXiv:2011.01764 [hep-th].
- [17] M. E. Machacek and M. T. Vaughn, "Two Loop Renormalization Group Equations in a General Quantum Field Theory. 1. Wave Function Renormalization," *Nucl. Phys. B* 222 (1983) 83–103.
- [18] M. E. Machacek and M. T. Vaughn, "Two Loop Renormalization Group Equations in a General Quantum Field Theory. 2. Yukawa Couplings," *Nucl. Phys. B* 236 (1984) 221–232.
- [19] M. E. Machacek and M. T. Vaughn, "Two Loop Renormalization Group Equations in a General Quantum Field Theory. 3. Scalar Quartic Couplings," *Nucl. Phys. B* 249 (1985) 70–92.

- [20] H. Arason, D. J. Castano, B. Keszthelyi, S. Mikaelian, E. J. Piard, P. Ramond, and B. D. Wright, "Renormalization group study of the standard model and its extensions. 1. The Standard model," *Phys. Rev. D* 46 (1992) 3945–3965.
- [21] M.-x. Luo and Y. Xiao, "Two loop renormalization group equations in the standard model," *Phys. Rev. Lett.* **90** (2003) 011601, arXiv:hep-ph/0207271.
- [22] F. Staub, "SARAH," arXiv:0806.0538 [hep-ph].
- [23] F. Staub, "SARAH 4 : A tool for (not only SUSY) model builders," Comput. Phys. Commun. 185 (2014) 1773-1790, arXiv:1309.7223 [hep-ph].
- [24] J. Schwichtenberg, "Gauge Coupling Unification without Supersymmetry," Eur. Phys. J. C 79 no. 4, (2019) 351, arXiv:1808.10329 [hep-ph].
- [25] Particle Data Group Collaboration, R. L. Workman *et al.*, "Review of Particle Physics," *PTEP* 2022 (2022) 083C01.
- [26] L. M. Carpenter, T. Murphy, and T. M. P. Tait, "Phenomenological cornucopia of SU(3) exotica," *Phys. Rev. D* 105 no. 3, (2022) 035014, arXiv:2110.11359
 [hep-ph].
- [27] R. Howl and S. F. King, "Planck scale unification in a supersymmetric Standard Model," *Phys. Lett. B* 652 (2007) 331–337, arXiv:0705.0301 [hep-ph].
- [28] A. Ibarra, P. Strobl, and T. Toma, "Neutrino masses from Planck-scale lepton number breaking," *Phys. Rev. Lett.* **122** no. 8, (2019) 081803, arXiv:1802.09997 [hep-ph].
- [29] A. Ibarra, P. Strobl, and T. Toma, "Two-loop renormalization group equations for right-handed neutrino masses and phenomenological implications," *Phys. Rev. D* 102 no. 5, (2020) 055011, arXiv:2006.13584 [hep-ph].
- [30] T. Han and S. Willenbrock, "Scale of quantum gravity," *Phys. Lett. B* 616 (2005) 215-220, arXiv:hep-ph/0404182.
- [31] Super-Kamiokande Collaboration, A. Takenaka *et al.*, "Search for proton decay via p → e⁺π⁰ and p → μ⁺π⁰ with an enlarged fiducial volume in Super-Kamiokande I-IV," *Phys. Rev. D* 102 no. 11, (2020) 112011, arXiv:2010.16098 [hep-ex].
- [32] Hyper-Kamiokande Collaboration, Y. Itow, "Construction status and prospects of the Hyper-Kamiokande project," *PoS* ICRC2021 (2021) 1192.
- [33] T. Han, I. Lewis, and Z. Liu, "Colored Resonant Signals at the LHC: Largest Rate and Simplest Topology," JHEP 12 (2010) 085, arXiv:1010.4309 [hep-ph].
- [34] P. Richardson and D. Winn, "Simulation of Sextet Diquark Production," Eur. Phys. J. C 72 (2012) 1862, arXiv:1108.6154 [hep-ph].

- [35] **ATLAS** Collaboration, M. Aaboud *et al.*, "Search for metastable heavy charged particles with large ionization energy loss in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS experiment," *Phys. Rev. D* **93** no. 11, (2016) 112015, arXiv:1604.04520 [hep-ex].
- [36] **ATLAS** Collaboration, M. Aaboud *et al.*, "Search for heavy long-lived charged *R*-hadrons with the ATLAS detector in 3.2 fb⁻¹ of proton-proton collision data at $\sqrt{s} = 13$ TeV," *Phys. Lett. B* **760** (2016) 647–665, arXiv:1606.05129 [hep-ex].
- [37] MAGIC Collaboration, H. Abe *et al.*, "Search for Gamma-Ray Spectral Lines from Dark Matter Annihilation up to 100 TeV toward the Galactic Center with MAGIC," *Phys. Rev. Lett.* **130** no. 6, (2023) 061002, arXiv:2212.10527 [astro-ph.HE].
- [38] A. Montanari, E. Moulin, and N. L. Rodd, "Toward the ultimate reach of current imaging atmospheric Cherenkov telescopes and their sensitivity to TeV dark matter," *Phys. Rev. D* 107 no. 4, (2023) 043028, arXiv:2210.03140 [astro-ph.HE].
- [39] M. Farina, D. Pappadopulo, and A. Strumia, "A modified naturalness principle and its experimental tests," JHEP 08 (2013) 022, arXiv:1303.7244 [hep-ph].
- [40] E. Ma, "Efficacious additions to the standard model," *Phys. Lett. B* 625 (2005) 76-78, arXiv:hep-ph/0508030.
- [41] P. Cox, A. Kusenko, O. Sumensari, and T. T. Yanagida, "SU(5) Unification with TeV-scale Leptoquarks," JHEP 03 (2017) 035, arXiv:1612.03923 [hep-ph].
- [42] E. Ma, " $SO(10) \rightarrow SU(5) \times U(1)_{\chi}$ as the Origin of Dark Matter," *Phys. Rev. D* 98 no. 9, (2018) 091701, arXiv:1809.03974 [hep-ph].
- [43] **ATLAS** Collaboration, G. Aad *et al.*, "Search for squarks and gluinos in final states with jets and missing transverse momentum using 139 fb⁻¹ of $\sqrt{s} = 13$ TeV *pp* collision data with the ATLAS detector," *JHEP* **02** (2021) 143, arXiv:2010.14293 [hep-ex].
- [44] **ATLAS** Collaboration, G. Aad *et al.*, "Search for squarks and gluinos in final states with one isolated lepton, jets, and missing transverse momentum at $\sqrt{s} = 13$ with the ATLAS detector," *Eur. Phys. J. C* **81** no. 7, (2021) 600, arXiv:2101.01629 [hep-ex]. [Erratum: Eur.Phys.J.C 81, 956 (2021)].
- [45] **ATLAS** Collaboration, G. Aad *et al.*, "Searches for new phenomena in events with two leptons, jets, and missing transverse momentum in 139 fb⁻¹ of $\sqrt{s} = 13$ TeV *pp* collisions with the ATLAS detector," *Eur. Phys. J. C* 83 no. 6, (2023) 515, arXiv:2204.13072 [hep-ex].

- [46] **ATLAS** Collaboration, G. Aad *et al.*, "Search for type-III seesaw heavy leptons in dilepton final states in *pp* collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector," *Eur. Phys. J. C* **81** no. 3, (2021) 218, arXiv:2008.07949 [hep-ex].
- [47] **ATLAS** Collaboration, G. Aad *et al.*, "Search for charged Higgs bosons decaying into a top quark and a bottom quark at $\sqrt{s} = 13$ TeV with the ATLAS detector," *JHEP* **06** (2021) 145, arXiv:2102.10076 [hep-ex].