

CASIMIR RADIAL PARTS VIA MATSUKI DECOMPOSITION

PHILIP SCHLÖSSER AND MIKHAIL ISACHENKOV

ABSTRACT. We use Matsuki’s decomposition for symmetric pairs (G, H) of (not necessarily compact) reductive Lie groups to construct the radial parts for invariant differential operators acting on matrix-spherical functions. As an application, we employ this machinery to formulate an alternative, mathematically rigorous approach to obtaining radial parts of Casimir operators that appear in the theory of conformal blocks, which avoids poorly defined analytical continuations from the compact quotient cases. To exemplify how this works, after reviewing the presentation of conformal 4-point correlation functions via matrix-spherical functions for the corresponding symmetric pair, we for the first time provide a complete analysis of the Casimir radial part decomposition in the case of Lorentzian signature. As another example, we revisit the Casimir reduction in the case of conformal blocks for two scalar defects of equal dimension. We argue that Matsuki’s decomposition thus provides a proper mathematical framework for analysing the correspondence between Casimir equations and the Calogero–Sutherland-type models, first discovered by one of the authors and Schomerus.

2020 *Mathematics Subject Classification.* 33C55, 33C67, 33C80, 43A90, 81T40.

Key words and phrases. matrix-spherical function, symmetric pair, Cartan decomposition, conformal blocks, Calogero–Sutherland model, degenerate double-affine Hecke algebra, Cherednik–Dunkl operator, Heckman–Opdam hypergeometric function.

CONTENTS

1. Introduction	2
2. Matrix-Spherical Functions (MSFs)	4
3. Matsuki's Double Coset Decomposition	7
4. Radial Parts	9
4.1. Root Space Decomposition	9
4.2. Decomposition	10
4.3. Radial Part of the Quadratic Casimir Element $\Omega_{\mathfrak{g}}$	17
5. Mathematical Setup for Conformal Blocks	21
5.1. Conformal Compactification	21
5.2. Conformal Group G and its Structure	22
5.3. Point Configurations	28
5.4. 4-Point Functions as MSF	32
6. Obtaining the Casimir Equation	34
6.1. Cartan Subsets $(C_i)_{i \in I}$ of G	34
6.2. Coordinates	41
6.3. Root Spaces in the Euclidean Setting	52
6.4. Root Spaces in the Lorentzian Setting	53
6.5. Matrix Case	61
6.6. Scalar Case	63
6.7. Euclidean Spinor Case	65
7. Blocks of Conformal Defects	67
8. Conclusion and Outlook	70
9. Acknowledgements	72
References	72

1. INTRODUCTION

A matrix-spherical function for two (topological, real Lie, complex Lie, algebraic) groups (G, H) with $H \leq G$ finite-dimensional H -representations V, W is a (continuous, smooth, holomorphic, regular) function $f: G \rightarrow \text{Hom}(V, W)$ satisfying

$$\forall g \in G, h, h' \in H: f(hgh') = \pi_W(h)f(g)\pi_V(h').$$

The theory of matrix-spherical functions, and the matrix-valued orthogonal polynomials they give rise to, has been well studied, with references including [1, 2, 3]. Considerable attention has been given to matrix-spherical functions for symmetric pairs (G, K) , i.e. when there is an involutive isomorphism θ of G and $K = G^\theta$ is compact.

However, the literature becomes more sparse when we allow K to be non-compact. The reason for this is that the well-studied cases correspond to the situation (and slight variations thereof) that the reductive group (G, K, θ, B) (see e.g. [4, Section VII.2]) is quotiented by its maximal compact subgroup K , and thus there is an Abelian subgroup A such that $G = KAK$ [4, Section VII.3]. This allows to uniquely reconstruct a (G, K) -matrix-spherical functions from their restrictions to A . Leveraging this fact, invariant differential operators can be decomposed into matrix-valued differential operators on A , their *radial parts*, as is done in [2], following the earlier works of Harish-Chandra.

The proof that G decomposes as KAK quite crucially relies on the fact that K is compact, and such a statement at face value is usually wrong otherwise. Actually,

as is shown in [5], the best we can hope for with a general symmetric pair (G, H) is

$$G = \bigcup_i \overline{HC_iH}$$

for finitely many ‘‘affine tori’’ C_i , i.e. cosets with respect to an Abelian subgroup. Fortunately, it turns out that in this case matrix-spherical functions are still determined by their restrictions to all of the C_i . In the present work we will take this story one step further and explore how invariant differential operators, in particular the quadratic Casimir element, decompose in such setting.

This has a direct application to conformal field theory (CFT), more specifically the theory of conformal blocks. As is explained in [6, Section 3], conformal blocks for (Euclidean) 4-point correlators can be described as matrix-spherical functions for $(SO(d+1, 1)_0, SO(1, 1)_0 \times SO(d))$.¹ In the scalar case (i.e. $SO(d)$ acting trivially, and $SO(1, 1)_0 \cong \mathbb{R}_{>0}$ acting by characters), the action of the quadratic Casimir operator was first determined in [9]. The authors of [10] then showed that a change of coordinates transforms the resulting differential operator into the Hamiltonian of the Calogero–Sutherland model for BC_2 , with parameters that refer not only to multiplicities of a relevant root system, but also to the characters by which $SO(1, 1)_0$ acts from the left and from the right.

The Calogero–Sutherland model is a super-integrable model with quite some history. In particular, it has been extensively studied by Heckman and Opdam (see e.g. [11] and references therein) and has been known to capture the Casimir action on zonal spherical functions for the more conventionally studied symmetric pairs (G, K) , where K has to be compact. This connection, however, is not enough to explain the result from [10], as $SO(1, 1)_0 \times SO(d)$ is non-compact, and as the conformal blocks in question aren’t zonal spherical functions. What we will show in present paper is that, with Matsuki’s decomposition in hand, the steps from [11] can be reproduced very closely to yield parallel results in the vastly generalised setting, including for example the case of Lorentzian kinematics, where the configuration space of points on the group has richer structure compared to the Euclidean situation. As a matter of fact, we will develop most of the theory for the general symmetric pair context, then specialize to a general indefinite orthogonal group $G = SO(p+1, q+1)$, and only further down the road specialize to Euclidean and Lorentzian signatures ($q = 0, 1$), once the real rank of G will start to affect the amount of needed calculations in an essential way.

Lastly, conformal blocks for two defects of the same dimension can also be described as spherical functions, namely for $(SO(d+1, 1)_0, SO(d-p) \times SO(p+1, 1)_0)$. We will see that this setup also falls within the domain of validity for Matsuki’s decomposition, so that we will be able to similarly use it in the radial part decomposition of the quadratic Casimir element, clarifying the appearance of the Calogero–Sutherland model in this context, which was earlier argued in [12].

Let us stress that the issue of having a properly defined Cartan decomposition in non-compact case is also important from the physics perspective, perhaps contrary to what might seem at the first glance. Being sometimes perceived more as a mathematical nuisance, it is usually glossed over in the physics literature by declaring it to be a more or less straightforward ‘analytical continuation’ from the compact case. However, as is frequently the case with going from compact spaces to non-compact ones, the ‘boundary questions’ actually cause trouble and turn out to be rather non-trivial mathematically, especially in treating lower-dimensional

¹Notice, however, that a rigorous treatment of correlation functions in a CFT also needs to take into account their distributional nature. In the present paper we choose not to work in this generality. See e.g. [7, 8] for the recent relevant work in this direction and a review of earlier physics literature.

group orbits. Notice that, despite the possible lower dimensionality of such strata, good analytical control over them is of relevance, since various such regions might carry information about singularities of the physical correlation functions.

As we have just reviewed above, bits and pieces of this radial part analysis, mostly but not exclusively in Euclidean setting, have appeared in physics literature before, see, in particular, [13, 6, 14, 15, 16]. Spherical functions relevant for Lorentzian analysis in the scalar case were also recently discussed in [17]. However, the mathematically rigorous description of the Casimir reduction in the setting pertinent to the four-point (spinning) conformal blocks, including the case of Lorentzian signature, was, as far as we are aware, up to now missing from the literature. In particular, a careful analysis of the subtleties of the KAK -type matrix decomposition in non-compact case that we provide in this paper using Matsuki's theory is, to our knowledge, new.² This, besides the wish to streamline and clean up mathematical details, might be seen as the main motivation behind this work on the physics side. It puts the KAK -type decompositions used in the Calogero–Sutherland approach to conformal blocks on a firm mathematical ground.

The plan of the paper is as follows. In Section 2 we review some generalities on matrix-spherical functions, especially with respect to group actions on them. We then go on, in Section 3 to review Matsuki's ([5]) theory of decompositions in the special case where the two involutions $\sigma = \tau$ and the two subgroups $H = L$ are equal. In Section 4 we develop this into a theory of radial part decompositions and compute the radial part of the quadratic Casimir element. In Section 5 we then pivot to applications in the theory of conformal blocks and explain how conformal blocks for 4-point correlation functions can be viewed as matrix-spherical functions in the sense defined earlier. In particular, in Section 6 we do a thorough analysis of Matsuki's theory for the group $SO(p+1, q+1)_0$, derive the expression of the quadratic Casimir element and match it with the scalar (Heckman–Opdam) expression from [10] and the spinor expression from [15]. In Section 7 we then provide a brief discussion of defect blocks and matching our results with [12]. We close this paper with a short summary and discussion of possible future directions.

A word about notation: most of the notation is chosen in such a way that it is compatible with [5]. The notation of Section 5 that pertains to G, \mathfrak{g} and various subgroups and subalgebras is more in line with the standard treatment of the structure theory of simple Lie groups (as presented in [4]) and will, due to collisions with [5]'s notation, not be used in the subsequent sections. In Subsection 6.6 (and examples beforehand), we introduce the scalar parameters α, β , so then roots will not be called α, β , but γ .

2. MATRIX-SPHERICAL FUNCTIONS (MSFs)

Definition 2.1. *Let (G, H) be a pair of groups, i.e. G is a group and $H \leq G$ is a subgroup. Let $(V, \pi_V), (W, \pi_W)$ be representations of H . A function $f: G \rightarrow \text{Hom}(V, W)$ is said to be a matrix-spherical function (MSF) if*

$$\forall g \in G, h, h' \in H: \quad f(hgh') = \pi_W(h)f(g)\pi_V(h').$$

In the case where G, H are real Lie groups (complex Lie groups, algebraic groups), we usually also require f to be smooth (holomorphic, regular). Write $E^{V,W}(G, H)$ for the set of matrix-spherical functions for (G, H) with the representations V, W .

²For example, in [16] the existence of such a decomposition is attributed to the Gel'fand pair property of (G, H) . However, we are not aware of any general statements yielding such a group factorisation for a given non-compact Gel'fand pair.

In the more general case, where W is an H -bimodule, we can make the same definition: a function $f: G \rightarrow W$ is a MSF if

$$\forall g \in G, h, h' \in H: f(hgh') = h \cdot f(g) \cdot h'.$$

However, as we cannot interpret this function as being matrix-valued anymore, the name MSF is a misnomer. Write $E^W(G, H)$ for the functions thus described.

We're now going to consider the case where (G, H) are Lie groups and W an H -bimodule with two smooth actions.

Lemma 2.2. $E^W(G, H)$ has actions by $U(\mathfrak{g})^H$ in terms of left-invariant and right-invariant differential operators. Furthermore, these two actions are pointwise related: for $f \in E^W(G, H)$ (or $C^\infty(G, W)$ more generally), $g \in G, p \in U(\mathfrak{g})^H$ with $\text{Ad}(g)(p) \in U(\mathfrak{g})^H$ we have

$$(p \cdot f)(g) = (f \cdot \text{Ad}(g)(p))(g).$$

Proof. We first show that $C^\infty(G, W)$ is a $(U(\mathfrak{g}), H)$ -bimodule (which is derived from the left and right regular representations).

For a function $f \in C^\infty(G, W)$, and elements $g \in G, h, h' \in H, X \in \mathfrak{g}$ we define

$$\begin{aligned} (h \cdot f \cdot h')(g) &:= (h')^{-1} \cdot f(h'gh) \cdot h^{-1} \\ (X \cdot f)(g) &:= \left. \frac{d}{dt} f(g \exp(tX)) \right|_{t=0} \\ (f \cdot X)(g) &:= \left. \frac{d}{dt} f(\exp(tX)g) \right|_{t=0}. \end{aligned}$$

Since G is associative, the left and right actions commute. Furthermore, for $h \in H, p \in U(\mathfrak{g})$ we have

$$\begin{aligned} h \cdot p \cdot h^{-1} \cdot f &= \text{Ad}(h)(p) \cdot f \\ f \cdot h \cdot p \cdot h^{-1} &= f \cdot \text{Ad}(h)(p), \end{aligned}$$

so that the left actions (resp. the right actions) are compatible with each other, which establishes that $(U(\mathfrak{g}), H)$ -bimodule structure. Therefore, the invariants under the left action of H still have a left action of $U(\mathfrak{g})^H$, analogously for the right actions. In particular, the elements that are left and right invariant under H (i.e. elements of $E^W(G, H)$) have a left and a right action of $U(\mathfrak{g})^H$. As can be seen from the definition, the left action of $U(\mathfrak{g})$ involves the right regular representation on $C^\infty(G)$ and therefore commutes with the left regular representation, so that it acts by left-invariant differential operators. Similarly, the right action of $U(\mathfrak{g})$ is an action by means of right-invariant differential operators.

Lastly, we show the pointwise relation between the left and right actions without the H -invariance assumption, i.e. for $C^\infty(G, W)$ and $U(\mathfrak{g})$. Let $f \in C^\infty(G, W)$ and $X \in \mathfrak{g}$ then

$$(X \cdot f)(g) = \left. \frac{d}{dt} f(g \exp(tX)) \right|_{t=0} = \left. \frac{d}{dt} f(\exp(t \text{Ad}(g)(X))g) \right|_{t=0} = (f \cdot \text{Ad}(g)(X))(g).$$

Now assume, the claim already holds for $p \in U(\mathfrak{g})$, then we have

$$\begin{aligned} (Xp \cdot f)(g) &= (X \cdot (p \cdot f))(g) = ((p \cdot f) \cdot \text{Ad}(g)(X))(g) \\ &= (p \cdot (f \cdot \text{Ad}(g)(X)))(g) = ((f \cdot \text{Ad}(g)(X)) \cdot \text{Ad}(g)(p))(g) \\ &= (f \cdot \text{Ad}(g)(Xp))(g). \end{aligned} \quad \square$$

Corollary 2.3. The actions from Lemma 2.2 can be restricted to $U(\mathfrak{g})^G$. If G is connected, this yields two actions of $Z(U(\mathfrak{g}))$ that coincide.

Proof. If G is connected, G -invariance is the same as \mathfrak{g} -invariance, so $U(\mathfrak{g})^G = Z(U(\mathfrak{g}))$. Let $z \in Z(U(\mathfrak{g}))$, $g \in G$ and $f \in E^W(G, H)$. By Lemma 2.2, we have

$$(z \cdot f)(g) = (f \cdot \text{Ad}(g)(z))(g) = (f \cdot z)(g). \quad \square$$

Corollary 2.4. *For $p \in U(\mathfrak{h})$ and $q \in U(\mathfrak{g})$ and $f \in E^W(G, H)$ we have*

$$(pq \cdot f)(g) = (p \cdot f)(g) \cdot q, \quad (f \cdot qp)(g) = q \cdot (f \cdot p)(g).$$

Proof. We show the claim for q being a monomial and then use linearity. Let $X_1 \cdots X_r \in \mathfrak{h}$ and $p = X_1 \cdots X_r$. Let $g \in G$, then

$$\begin{aligned} (q \cdot f)(g) &= \frac{d}{dt_1} \cdots \frac{d}{dt_r} f(g \exp(tX_1) \cdots \exp(tX_r)) \Big|_{t=0} \\ &= \frac{d}{dt_1} \cdots \frac{d}{dt_r} f(g) \cdot \exp(t_1 X_1) \cdots \exp(t_r X_r) \Big|_{t_1=\dots=t_r=0} \\ &= f(g) \cdot q, \end{aligned}$$

whence $(pq \cdot f)(g) = (p \cdot (q \cdot f))(g) = (p \cdot f)(g) \cdot q$ as claimed. The result for the right action can be obtained analogously. \square

Suppose now that we can decompose a dense subset G' of G as follows:

$$(2.1) \quad G' = \bigsqcup_{i \in I} HC_i H$$

where $C_i \subseteq G$ are weakly embedded submanifolds (in our story these will turn out to be tori or right cosets of tori), i.e. the identity map $C_i \hookrightarrow G$ is an immersion. Parallel to the usual treatment, a spherical function is then determined by its restrictions to these tori C_i .

Lemma 2.5. *The map $|_C: E^W(G, H) \rightarrow \prod_{i \in I} C^\infty(C_i, W)$ given by*

$$f \mapsto (f|_{C_i})_{i \in I}$$

is injective.

Proof. Let f, f' with $f|_C = f'|_C$, i.e. $f|_{C_i} = f'|_{C_i}$ for all $i \in I$. Since f, f' are matrix-spherical, they agree on the corresponding $H \times H$ -orbits as well, i.e. on $HC_i H$ for all $i \in I$, so in particular on

$$\bigcup_{i \in I} HC_i H = G'.$$

Since both are continuous and G' is dense in G , we have $f = f'$. \square

Definition 2.6. *For $i \in I$ define*

$$\begin{aligned} N_{C_i} &:= \{(h, h') \in H^2 \mid hC_i h'^{-1} = C_i\} \\ Z_{C_i} &:= \{(h, h') \in H^2 \mid \forall x \in C_i: h x h'^{-1} = x\} \end{aligned}$$

and $J_{C_i} := N_{C_i}/Z_{C_i}$. They are all groups.

Lemma 2.7. *Let $f \in E^W(G, H)$ and $i \in I$, then*

$$f|_{C_i} \in C^\infty(C_i, W^{Z_{C_i}})^{J_{C_i}}.$$

Proof. Let $g \in C_i$ and $(h, h') \in N_{C_i}$, then

$$f(hgh'^{-1}) = h \cdot f(g) \cdot h'^{-1} = (h, h') \cdot f(g).$$

For the case where $(h, h') \in Z_{C_i}$ we have $hgh'^{-1} = g$ and hence $f(g) \in W^{Z_{C_i}}$. Thus we see that $f|_{C_i}$ intertwines the actions of N_{C_i} , which pass to the quotient J_{C_i} . \square

3. MATSUKI'S DOUBLE COSET DECOMPOSITION

We are now going to quote some results from [5] that will be relevant later. Note that Matsuki considers two (generally) different involutions σ, τ on G , whereas in this paper we shall always assume $\sigma = \tau$.

Let (G, H) be a symmetric pair of Lie groups, i.e. $H \leq G$ a Lie subgroup and $\sigma \in \text{Aut}(G)$ an involution such that $(G^\sigma)_0 \leq H \leq G^\sigma$, where we assume that $HG_0H = G$. We shall also assume that (G, K, θ, B) is a reductive Lie group (see e.g. [4, Section VII.2]), where θ and σ ('s derivative) commute.

We shall also assume that B is invariant under σ , which is no restriction by the following observation:

Lemma 3.1. *Let (G, K, θ, B) be a reductive Lie group and let $S \leq \text{Aut}(G)$ be a finite subgroup (the derivative of) whose action commutes with θ . Then B can without loss of generality be chosen to be S -invariant.*

Proof. If we choose

$$B'(X, Y) := \sum_{\sigma \in S} B(\sigma(X), \sigma(Y))$$

instead of B , we obtain a real inner product on \mathfrak{g} that is invariant under $\text{Ad}(G), \theta, S$, which is still positive(negative)-definite on \mathfrak{p} (\mathfrak{k}), and under which \mathfrak{p} and \mathfrak{k} are still orthogonal. In other words: (G, K, θ, B') is still a reductive Lie group. \square

Definition 3.2. *Let \mathfrak{t} be a maximal commutative subalgebra of $\mathfrak{k}^{-\sigma}$ and extend \mathfrak{t} to a maximal commutative subalgebra $\mathfrak{c} = \mathfrak{t} \oplus \mathfrak{a}$ of $\mathfrak{g}^{-\sigma}$. The subgroup $C := \exp(\mathfrak{c}) = : \exp(\mathfrak{a})T$ is called a fundamental Cartan subset.*

Definition 3.3. *Given a fundamental Cartan subset C as above, a subset $C' = \exp(\mathfrak{c}')t$ of G is called a standard Cartan subset if*

- (i) $t \in T$ and $\mathfrak{c}' \leq \mathfrak{g}^{-\sigma} \cap \text{Ad}(t)(\mathfrak{g}^{-\sigma})$ is a commutative subalgebra;
- (ii) $\mathfrak{c}' = \mathfrak{t}' \oplus \mathfrak{a}'$ where $\mathfrak{t}' \leq \mathfrak{t}$ and $\mathfrak{a} \leq \mathfrak{a}' \subseteq \mathfrak{p}$;
- (iii) $\dim(\mathfrak{c}') = \dim(\mathfrak{c})$.

Two standard Cartan subsets $\exp(\mathfrak{a}'_1)T'_1$ and $\exp(\mathfrak{a}'_2)T'_2$ are said to be conjugate if there are $h, h' \in K \cap H$ such that $hT'_1h'^{-1} = T'_2$. In particular, any two Cartan subsets with the same T' are conjugate.

Example 3.4. *For the case of $\sigma = \theta$ and $H = K$, i.e. the case of non-compact symmetric spaces, we pick $\mathfrak{t} := 0$ and \mathfrak{a} as a maximal commutative subalgebra of \mathfrak{p} . Then the fundamental Cartan subset C is the analytic subgroup A of \mathfrak{a} , i.e. the torus one finds in the usual treatment for this case, e.g. [4, Theorem 6.46]. There are also no other standard Cartan subsets (with respect to C) as \mathfrak{a} is already a maximal commutative subalgebra of $\mathfrak{g}^{-\sigma}$.*

From now on, we shall fix a fundamental Cartan subset $C \leq G$.

Lemma 3.5. *Let $C' = \exp(\mathfrak{c}')t$ be a standard Cartan subset, then $\text{Ad}(t)$ is an involution of $\mathfrak{g}^{-\sigma} \cap \text{Ad}(t)(\mathfrak{g}^{-\sigma})$. In particular, C' and $tC't^{-1}$ are conjugate standard Cartan subsets.*

Proof. Let $X \in \mathfrak{g}^{-\sigma} \cap \text{Ad}(t)(\mathfrak{g}^{-\sigma})$, so that $X, \text{Ad}(t^{-1})(X) \in \mathfrak{g}^{-\sigma}$. Note that $t \in \exp(\mathfrak{k}^{-\sigma})$, whence $\sigma(t^{-1}) = t$. Consequently, we have

$$-\text{Ad}(t^{-1})(X) = \sigma(\text{Ad}(t^{-1})(X)) = \text{Ad}(t)(\sigma(X)) = -\text{Ad}(t)(X),$$

which implies $\text{Ad}(t^{-2})(X) = X$. Therefore we have $\text{Ad}(t)(X), X \in \mathfrak{g}^{-\sigma}$, so that

$$\text{Ad}(t)(X) \in \mathfrak{g}^{-\sigma} \cap \text{Ad}(t)(\mathfrak{g}^{-\sigma}).$$

Now, note that $tC't^{-1} = t \exp(\mathfrak{c}') = \exp(\text{Ad}(t)(\mathfrak{c}'))t$ and that $\text{Ad}(t)(\mathfrak{c}')$ decomposes as

$$\mathfrak{t}' \oplus \text{Ad}(t)(\mathfrak{a}')$$

since $t \in \exp(\mathfrak{t})$ commutes with $\mathfrak{t}' \leq \mathfrak{t}$. Therefore, $tC't^{-1}$ is also a standard Cartan subset that shares its “compact part” $\exp(\mathfrak{t}')t$ with C' and is therefore conjugate to C' . \square

Definition 3.6. *The set G_{ss} of all semisimple elements of G is the set of all $g \in G$ such that $\sigma \circ \text{Ad}(g) \circ \sigma \circ \text{Ad}(g)^{-1}$ is a semisimple Lie algebra automorphism of \mathfrak{g} . Let*

$$G_{\text{rs}} := \{g \in G_{\text{ss}} \mid g^{-\sigma} \cap \text{Ad}(g)(\mathfrak{g}^{-\sigma}) \text{ is commutative}\}$$

be the set of all regular semisimple elements. Both are dense in G .

Lemma 3.7. *Let $C' = \exp(\mathfrak{c}')t$ be a standard Cartan subset and let $x \in C'$. Then*

$$\mathfrak{g}^{-\sigma} \cap \text{Ad}(x)(\mathfrak{g}^{-\sigma}) = \mathfrak{c}'$$

if and only if $x \in G_{\text{rs}}$.

Proof. “ \Leftarrow ”: “ \supset ”: Write $x = \exp(X)t$ and let $Y \in \mathfrak{c}'$. Then

$$\text{Ad}(x^{-1})(Y) = \text{Ad}(t^{-1}) \exp(-\text{ad}(X))(Y) = \text{Ad}(t^{-1})(Y) \in \mathfrak{g}^{-\sigma}$$

since $\mathfrak{c}' \leq \mathfrak{g}^{-\sigma} \cap \text{Ad}(t)(\mathfrak{g}^{-\sigma})$ by definition.

“ \subseteq ”: Since $x = \exp(X)t$ is regular, $\mathfrak{s} := \mathfrak{g}^{-\sigma} \cap \text{Ad}(x)(\mathfrak{g}^{-\sigma})$ is commutative and contains \mathfrak{c}' by the observation above. For $Y \in \mathfrak{s}$ we thus have $\text{ad}(X)(Y) = 0$, so that

$$\text{Ad}(t^{-1})(Y) = \text{Ad}(x^{-1}) \exp(-\text{ad}(X))(Y) = \text{Ad}(x^{-1})(Y) \in \mathfrak{g}^{-\sigma}.$$

Consequently,

$$\mathfrak{c}' \leq \mathfrak{s} \subseteq \mathfrak{g}^{-\sigma} \cap \text{Ad}(t)(\mathfrak{g}^{-\sigma}),$$

where \mathfrak{s} is commutative and \mathfrak{c}' is maximally commutative. Consequently, $\mathfrak{s} \leq \mathfrak{c}'$.

“ \Rightarrow ”: If x is not regular, then $\mathfrak{g}^{-\sigma} \cap \text{Ad}(x)(\mathfrak{g}^{-\sigma})$ is not commutative. Therefore, it cannot be \mathfrak{c}' . \square

Theorem 3.8 ([5, Theorem 3(i–iii)]). *Let G, H, K, σ, θ as before. Fix a fundamental Cartan subset and let $(C_i)_{i \in I}$ be representatives of all the conjugacy classes of standard Cartan subsets. Then*

$$G_{\text{ss}} = \bigcup_{i \in I} HC_iH$$

$$G_{\text{rs}} = \bigsqcup_{i \in I} H(C_i \cap G_{\text{rs}})H.$$

This establishes a decomposition as in (2.1). Every element of the group J_{C_i} we defined for such a case can be represented by $(h, h')N_{C_i}$ where $h, h' \in K \cap H$.

As a consequence, we see that any matrix-spherical function for (G, H) is entirely determined by its restrictions to some standard Cartan subsets (Theorem 3.8 and Lemma 2.5). This defines for us an appropriate setting in which we can now start computing radial parts of the differential operators encountered in Lemma 2.2, and especially of the quadratic Casimir element.

4. RADIAL PARTS

4.1. Root Space Decomposition.

Proposition 4.1. *Let $C' = \exp(\mathfrak{c}')t$ be a standard Cartan subset of G . For $\alpha \in (\mathfrak{c}'_{\mathbb{C}})^*$ define*

$$\mathfrak{g}_{\alpha} := \{X \in \mathfrak{g}_{\mathbb{C}} \mid \forall Z \in \mathfrak{c}'_{\mathbb{C}}: \text{ad}(Z)(X) = \alpha(Z)X\}.$$

Then

- (i) $\mathfrak{g}_{\mathbb{C}} = \bigoplus_{\alpha} \mathfrak{g}_{\alpha}$;
- (ii) $[\mathfrak{g}_{\alpha}, \mathfrak{g}_{\beta}] \subseteq \mathfrak{g}_{\alpha+\beta}$;
- (iii) $\sigma(\mathfrak{g}_{\alpha}) \subseteq \mathfrak{g}_{-\alpha}$
- (iv) $B(\mathfrak{g}_{\alpha}, \mathfrak{g}_{\beta}) = 0$ unless $\alpha + \beta = 0$.

Write $\Sigma(\mathfrak{g}; \mathfrak{c}')$ for the set of α with $\dim(\mathfrak{g}_{\alpha}) > 0$.

Proof. (i) Since \mathfrak{c}' is commutative, so is $\text{ad}(\mathfrak{c}')$. For simultaneous diagonalisability and hence the existence of the claimed root space decomposition, it suffices therefore to show semisimplicity. We show that there is a (sesquilinear, positive-definite) inner product on $\mathfrak{g}_{\mathbb{C}}$ with respect to which $\text{ad}(\mathfrak{c}')$ acts normally.

Recall that since (G, K, θ, B) is a reductive Lie group, the bilinear form B_{θ} is a positive-definite inner product on \mathfrak{g} . If we extend it sesquilinearly, we therefore obtain an inner product on $\mathfrak{g}_{\mathbb{C}}$. For any $X, Y \in \mathfrak{g}_{\mathbb{C}}$ and $Z \in \mathfrak{c}'$ we have

$$\begin{aligned} B_{\theta}(\text{ad}(Z)(X), Y) &= -B(\text{ad}(Z)(X), \theta(Y)) = B(X, \text{ad}(Z)(\theta(Y))) \\ &= B(X, \theta(\text{ad}(\theta(Z))(Y))) = -B_{\theta}(X, \text{ad}(\theta(Z))(Y)). \end{aligned}$$

Consequently, the adjoint of $\text{ad}(Z)$ (with respect to B_{θ}) is $-\text{ad}(\theta(Z))$. Since by definition, \mathfrak{c}' is θ -invariant, the element $-\text{ad}(\theta(Z))$ lies in the commutative algebra $\text{ad}(\mathfrak{c}')$ and therefore commutes with $\text{ad}(Z)$.

- (ii) Let $X \in \mathfrak{g}_{\alpha}, Y \in \mathfrak{g}_{\beta}$ and $Z \in \mathfrak{c}'_{\mathbb{C}}$, then

$$\begin{aligned} \text{ad}(Z)([X, Y]) &= [\text{ad}(Z)(X), Y] + [X, \text{ad}(Z)(Y)] \\ &= [\alpha(Z)X, Y] + [X, \beta(Z)Y] \\ &= (\alpha + \beta)(Z)[X, Y]. \end{aligned}$$

- (iii) Let $X \in \mathfrak{g}_{\alpha}, Z \in \mathfrak{c}'_{\mathbb{C}}$. By definition of \mathfrak{c}' , we have $\sigma(Z) = -Z$, hence

$$\text{ad}(Z)(\sigma(X)) = \sigma(\text{ad}(\sigma(Z))(X)) = -\sigma(\text{ad}(Z)(X)) = -\alpha(Z)\sigma(X).$$

- (iv) Let $X \in \mathfrak{g}_{\alpha}, Y \in \mathfrak{g}_{\beta}$ and $Z \in \mathfrak{c}'_{\mathbb{C}}$, then

$$\begin{aligned} 0 &= B(\text{ad}(Z)(X), Y) + B(X, \text{ad}(Z)(Y)) \\ &= \alpha(Z)B(X, Y) + \beta(Z)B(X, Y) \\ &= (\alpha + \beta)(Z)B(X, Y) \end{aligned}$$

due to B 's ad-invariance. If $B(X, Y) \neq 0$, the above implies that $(\alpha + \beta)(Z) = 0$ for all Z , and hence $\alpha + \beta = 0$. \square

Lemma 4.2. *Let $\exp(\mathfrak{c}_1)t_1, \exp(\mathfrak{c}_2)t_2$ be two Cartan subsets and suppose there exists $\phi \in \text{Aut}(\mathfrak{g}_{\mathbb{C}})$ with $\phi(\mathfrak{c}_1) = \mathfrak{c}_2$. Then the reduced root systems $\Sigma(\mathfrak{g}; \mathfrak{c}_1) = \phi^*(\Sigma(\mathfrak{g}; \mathfrak{c}_2))$ and*

$$\phi(\mathfrak{g}_{\phi^*(\alpha)}) = \mathfrak{g}_{\alpha}.$$

for all $\alpha \in (\mathfrak{c}_2, \mathbb{C})^*$, meaning that the root systems and root multiplicities are the same.

Proof. Let $\alpha \in (\mathfrak{c}_{2,\mathbb{C}})^*$, then $\phi^*(\alpha) = \alpha \circ \phi \in (\mathfrak{c}_{1,\mathbb{C}})^*$. Let $X \in \mathfrak{g}_{\phi^*(\alpha)}$ and $Z \in \mathfrak{c}_1$, then

$$\text{ad}(Z)(\phi(X)) = \phi(\text{ad}(\phi^{-1}(Z))(X)) = \phi(\phi^*(\alpha)(\phi^{-1}(Z))X) = \alpha(Z)\phi(X),$$

which shows that $\phi(X) \in \mathfrak{g}_\alpha$. Consequently, ϕ restricts to isomorphisms $\mathfrak{g}_{\phi^*(\alpha)} \cong \mathfrak{g}_\alpha$ for all $\alpha \in (\mathfrak{c}_{2,\mathbb{C}})^*$, which in turn also shows that the root systems are isomorphic via ϕ^* . \square

4.2. Decomposition. From now on we shall fix a standard Cartan subset $C' = \exp(\mathfrak{c}')t$. Assume that $\text{Ad}(t)$ leaves \mathfrak{c}' invariant and can be decomposed as follows: there is an involution $\phi \in O(\mathfrak{g}_{\mathbb{C}}, B)$ that commutes with σ , and $\epsilon: \Sigma(\mathfrak{g}; \mathfrak{c}') \rightarrow \mathbb{C}^\times$ such that

$$\forall \alpha \in \Sigma(\mathfrak{g}; \mathfrak{c}'), X \in \mathfrak{g}_\alpha: \quad \text{Ad}(t)(X) = \epsilon_\alpha \phi(X).$$

By Lemma 3.5, $\text{Ad}(t)$ is an involution of $\mathfrak{c}'_{\mathbb{C}}$ and hence also of $\Sigma(\mathfrak{g}; \mathfrak{c}')$. Unless this leads to ambiguity, we shall denote this involution by t .

This definition implies two facts about the function ϵ :

Lemma 4.3. *For $\alpha \in \Sigma(\mathfrak{g}; \mathfrak{c}')$ we have*

- (i) $\epsilon_\alpha \epsilon_{-\alpha} = 1$ and
- (ii) $\epsilon_\alpha = \epsilon_{t\alpha}$.

In particular, $\epsilon_0 = \pm 1$, which can be absorbed into ϕ .

Proof. (i) Let $0 \neq X \in \mathfrak{g}_\alpha$, then due to the non-degeneracy of B there exists $Y \in \mathfrak{g}_{-\alpha}$ such that $B(X, Y) \neq 0$. Then due to B 's invariance under $\text{Ad}(t)$ and ϕ we have

$$\begin{aligned} B(X, Y) &= B(\text{Ad}(t)X, \text{Ad}(t)Y) = B(\epsilon_\alpha \phi(X), \epsilon_{-\alpha} \phi(Y)) \\ &= \epsilon_\alpha \epsilon_{-\alpha} B(\phi(X), \phi(Y)) = \epsilon_\alpha \epsilon_{-\alpha} B(X, Y), \end{aligned}$$

hence $1 = \epsilon_\alpha \epsilon_{-\alpha}$ since $B(X, Y) \neq 0$.

- (ii) Let $0 \neq X \in \mathfrak{g}_\alpha$, then we leverage that $\sigma(t) = t^{-1}$ and that $\sigma(\text{Ad}(t)(X)) \in \mathfrak{g}_{-t\alpha}$:

$$\begin{aligned} \sigma(X) &= \text{Ad}(t) \text{Ad}(t)^{-1} \sigma(X) = \text{Ad}(t) \sigma \text{Ad}(t)(X) \\ &= \epsilon_{-t\alpha} \phi \sigma \text{Ad}(t)(X) \\ &= \epsilon_\alpha \epsilon_{-t\alpha} \phi \sigma \phi(X) = \epsilon_\alpha \epsilon_{-t\alpha} \sigma(X), \end{aligned}$$

which shows with (i) that $\epsilon_\alpha = \epsilon_{t\alpha}$. \square

We now investigate what $Z_{C'}$ and $N_{C'}$ look and act like.

Proposition 4.4. (i) $(h, h') \in Z_{C'}$ if and only if $h' = t^{-1}ht$ and $\text{Ad}(h)$ acts trivially on \mathfrak{c}' .

- (ii) $(h, h') \in N_{C'}$ if and only if $\text{Ad}(h)\mathfrak{c}' \subseteq \mathfrak{c}'$ and $hth'^{-1}t^{-1} \in \exp(\mathfrak{c}')$.

- (iii) If $(h, h') \in N_{C'}$, say $hth'^{-1}t^{-1} = \exp(Y)$, then (h, h') acts on C' as follows:

$$(h, h') \cdot \exp(X)t = \exp(\text{Ad}(h)(X) + Y)t$$

Proof. (i) “ \Rightarrow ”: We know $t \in C'$, then $hth'^{-1} = t$, whence $h' = t^{-1}ht$. Let $X \in \mathfrak{c}'$, then

$$\begin{aligned} X &= \left. \frac{d}{ds} \exp(tX)tt^{-1} \right|_{s=0} = \left. \frac{d}{ds} h \exp(sX)th'^{-1}t^{-1} \right|_{s=0} \\ &= \left. \frac{d}{ds} \exp(s \text{Ad}(h)(X)) \right|_{s=0} = \text{Ad}(h)(X). \end{aligned}$$

“ \Leftarrow ”: Let $(h, h') \in H \times H$ satisfy the conditions above. Every element of C' can be written as $\exp(X)t$, so that

$$(h, h') \cdot \exp(X)t = h \exp(X)th'^{-1} = \exp(\text{Ad}(h)(X))hth'^{-1} = \exp(X)t,$$

so that $(h, h') \in Z_{C'}$.

(ii) “ \Rightarrow ”: We apply the definition first to t , which yields

$$hth'^{-1} = \exp(Y)t$$

for some $Y \in \mathfrak{c}'$. Let now $X \in \mathfrak{c}'$ and $s \in \mathbb{R}$, then

$$h \exp(sX)th'^{-1} = \exp(\text{Ad}(h)(sX) + Y)t.$$

Multiplying with t^{-1} and taking the s -derivative at $s = 0$, shows that $\text{Ad}(h)(X) \in \mathfrak{c}'$. The reverse implication is clear.

(iii) Follows from (ii). □

Lemma 4.5. *Let $\mathfrak{m}' := Z_{\mathfrak{h}}(\mathfrak{c}')$. Then*

$$\mathfrak{g}_0 = \mathfrak{c}'_{\mathbb{C}} \oplus \mathfrak{m}'_{\mathbb{C}}.$$

This direct sum is orthogonal with respect to B and B_{σ} .

Proof. The sum is evidently direct as \mathfrak{c}' and \mathfrak{m}' lie in different eigenspaces of σ . The inclusion “ \supseteq ” is clear. For “ \subseteq ”, note that σ leaves \mathfrak{g}_0 invariant by Proposition 4.1(iii), hence we can decompose

$$\mathfrak{g}_0 = \mathfrak{g}_0^+ \oplus \mathfrak{g}_0^-,$$

where σ acts as ± 1 on \mathfrak{g}_0^{\pm} . Since σ is orthogonal with respect to both B and B_{σ} , these two eigenspaces are orthogonal with respect to both bilinear forms. It just remains to show that $\mathfrak{g}_0^+ \subset \mathfrak{m}'_{\mathbb{C}}$ and $\mathfrak{g}_0^- \subset \mathfrak{c}'_{\mathbb{C}}$ (then these inclusions are equalities and we get the claimed orthogonality).

For the first inclusion, let $X + iY \in \mathfrak{g}_0^+$, then both X and Y commute with \mathfrak{c}' and satisfy $\sigma(X) = X, \sigma(Y) = Y$, so that $X, Y \in \mathfrak{h}$, and hence $X, Y \in \mathfrak{m}'$.

For the second inclusion, let $X + iY \in \mathfrak{g}_0^-$, then $X, Y \in \mathfrak{g}^{-\sigma}$. Furthermore, we have

$$\begin{aligned} \sigma(\text{Ad}(t^{-1})(X)) &= \sigma(\epsilon_0^{-1}\phi(X)) = \epsilon_0^{-1}\phi\sigma(X) = -\epsilon_0^{-1}\phi(X) \\ &= -\text{Ad}(t^{-1})(X), \end{aligned}$$

so $X \in \text{Ad}(t)(\mathfrak{g}^{-\sigma})$ as well. Consequently, $X \in \mathfrak{g}^{-\sigma} \cap \text{Ad}(t)(\mathfrak{g}^{-\sigma})$ and commutes with \mathfrak{c}' . Since \mathfrak{c}' is maximal commutative, we have $X \in \mathfrak{c}'$. Similarly, we also see $Y \in \mathfrak{c}'$. □

Definition 4.6. *Let $x \in \exp(X)t \in C'$ and $\alpha \in \Sigma(\mathfrak{g}:\mathfrak{c}')$. Define*

$$x^{\alpha} := \epsilon_{\alpha} \exp(\alpha(X)) \in \mathbb{C}^{\times}.$$

Proposition 4.7. (i) *For $x \in C'$ and $Y \in \mathfrak{g}_{\alpha}$ we have*

$$\text{Ad}(x)(Y) = x^{t\alpha}\phi(Y) \quad \text{Ad}(x^{-1})(Y) = x^{-\alpha}\phi(Y),$$

so in particular $x \mapsto x^{\alpha}$ is well-defined.

(ii) *For all $x \in C', \alpha \in \Sigma(\mathfrak{g}:\mathfrak{c}')$ we have $(x^{\alpha})^{-1} = x^{-\alpha}$.*

(iii) *With respect to the group homomorphism*

$$\exp(\mathfrak{c}') \rightarrow \mathbb{C}^{\times}, \quad \exp(X') \mapsto \exp(\alpha(X')),$$

the map $x \mapsto x^{\alpha}$ intertwines the (left) group actions of $\exp(\mathfrak{c}')$ on C' and of \mathbb{C}^{\times} on \mathbb{C}^{\times} (is in particular a homomorphism of torsors).

(iv) Let $(h, h') \in N_{C'}$ and let $hth'^{-1} = x = \exp(Y)t$. Then

$$\left(\epsilon_\alpha x^{-\text{Ad}^*(h)(\alpha)}\right)^2 = 1.$$

Proof. (i) We have

$$\text{Ad}(x)(Y) = \exp(\text{ad}(X)) \text{Ad}(t)(Y) = \epsilon_\alpha \exp(\text{ad}(X))\phi(Y) = \epsilon_\alpha \exp((t\alpha)(X))\phi(Y),$$

which by Lemma 4.3(ii) equals $x^{t\alpha}\phi(Y)$ as claimed. Furthermore, we have

$$\text{Ad}(x^{-1})(Y) = \text{Ad}(t^{-1}) \exp(\text{ad}(-X))(Y) = \epsilon_{t\alpha}^{-1} \exp(-\alpha(X))\phi(Y) = x^{-\alpha}\phi(Y)$$

by Lemma 4.3(i,ii).

(ii) Follows from Lemma 4.3(i).

(iii) Follows from the definition and the fact that \mathfrak{c}' is commutative.

(iv) Rearranging the definition of x we obtain $h = xh't^{-1}$. Applying σ , this equals $tx^{-1}t^{-1}h't$. Writing $x = \exp(Y)t$, this equation becomes

$$\exp(Y)th't^{-1} = \exp(-Y)t^{-1}h't.$$

Let $X \in \mathfrak{g}_\alpha$, then $\text{Ad}(th't^{-1})(X)$ and $\text{Ad}(t^{-1}h't)(X)$ only differ by a constant:

$$\begin{aligned} \text{Ad}(th't^{-1})(X) &= \frac{\epsilon_{\text{Ad}(th't)(\alpha)}}{\epsilon_\alpha} \phi(\text{Ad}(h')(\phi(X))) \\ \text{Ad}(t^{-1}h't)(X) &= \frac{\epsilon_\alpha}{\text{Ad}(th't)(\alpha)} \phi(\text{Ad}(h')(\phi(X))). \end{aligned}$$

Consequently, we have

$$\exp(\text{Ad}^*(th't)(\alpha)(Y)) \frac{\epsilon_{\text{Ad}(th't)(\alpha)}}{\epsilon_\alpha} = \exp(-\text{Ad}^*(th't)(\alpha)(Y)) \frac{\epsilon_\alpha}{\epsilon_{\text{Ad}(th't)(\alpha)}},$$

which implies the claim using the fact that $\text{Ad}^*(th't)$ and $\text{Ad}^*(h)$ coincide on $(\mathfrak{c}'_{\mathbb{C}})^*$. □

Lemma 4.8. *We can characterise $C' \cap G_{rs}$ as follows:*

$$C' \cap G_{rs} = \{x \in C' \mid \forall \alpha \in \Sigma: x^\alpha \neq x^{-\alpha}\}.$$

Proof. Let $x \in C'$. By Lemma 3.7 we have $x \in C' \cap G_{rs}$ iff $\mathfrak{c}' = \mathfrak{g}^{-\sigma} \cap \text{Ad}(x)(\mathfrak{g}^{-\sigma})$. So we are going to show that this condition holds iff $x^\alpha \neq x^{-\alpha}$ for all roots $\alpha \in \Sigma$.

“ \Rightarrow ”: Assume that $\mathfrak{g}^{-\sigma} \cap \text{Ad}(x)(\mathfrak{g}^{-\sigma}) = \mathfrak{c}'$. Let $\alpha \in \Sigma$ and $0 \neq E \in \mathfrak{g}_\alpha$. Then $E - \sigma(E) \in \mathfrak{g}^{-\sigma}$ is a nontrivial linear combination of \mathfrak{c}' -root vectors for different root spaces and cannot be a root vector itself, in particular not an element of \mathfrak{c}' . This implies that $E - \sigma(E) \notin \text{Ad}(x)(\mathfrak{g}^{-\sigma})$. We have

$$\text{Ad}(x^{-1})(E - \sigma(E)) = x^{-\alpha}\phi(E) - x^\alpha\phi(\sigma(E)) = x^{-\alpha}\phi(E) - x^\alpha\sigma(\phi(E)).$$

That this element is not σ -antiinvariant, shows that $x^\alpha \neq x^{-\alpha}$.

“ \Leftarrow ”: Assume $x^\alpha \neq x^{-\alpha}$ for all roots $\alpha \in \Sigma$ and let $Y \in \mathfrak{g}^{-\sigma} \cap \text{Ad}(x)(\mathfrak{g}^{-\sigma})$. We can decompose Y according to

$$\mathfrak{g}_{\mathbb{C}} = \mathfrak{m}'_{\mathbb{C}} \oplus \mathfrak{c}'_{\mathbb{C}} \oplus \bigoplus_{\alpha \in \Sigma} \mathfrak{g}_\alpha$$

as

$$Y = Y_{0,+} + Y_{0,-} + \sum_{\alpha \in \Sigma} Y_\alpha.$$

That Y is σ -antiinvariant shows that $Y_{0,+} = 0$ and $Y_{-\alpha} = -\sigma(Y_\alpha)$ for all $\alpha \in \Sigma$. Furthermore,

$$\begin{aligned} \text{Ad}(x^{-1})(Y) &= \epsilon_0 \phi(Y_{0,-}) + \sum_{\alpha \in \Sigma^+} (x^{-\alpha} \phi(Y_\alpha) - x^\alpha \phi(\sigma(Y_\alpha))) \\ &= \epsilon_0 \phi(Y_{0,-}) + \sum_{\alpha \in \Sigma^+} (x^{-\alpha} \phi(Y_\alpha) - x^\alpha \sigma(\phi(Y_\alpha))). \end{aligned}$$

Since this is σ -antiinvariant as well, we obtain $x^{-\alpha} \phi(Y_\alpha) = x^\alpha \phi(Y_\alpha)$. Since $x^\alpha \neq x^{-\alpha}$, this reads $\phi(Y_\alpha) = 0$ or equivalently $Y_\alpha = 0$. Consequently, we have $Y = Y_{0,-} \in \mathfrak{c}'$. \square

Proposition 4.9. *Let $x \in C' \cap G_{rs}$ and $0 \neq E \in \mathfrak{g}_\alpha$. Let $H := E + \sigma(E)$, then*

$$\begin{aligned} E &= \frac{\text{Ad}(x)(\phi(H)) - x^{-\alpha} H}{x^\alpha - x^{-\alpha}} \\ \sigma(E) &= \frac{x^\alpha H - \text{Ad}(x)(\phi(H))}{x^\alpha - x^{-\alpha}} \\ \phi(E) &= \frac{x^\alpha \phi(H) - \text{Ad}(x^{-1})(H)}{x^\alpha - x^{-\alpha}} \\ \sigma(\phi(E)) &= \frac{\text{Ad}(x^{-1})(H) - x^{-\alpha} \phi(H)}{x^\alpha - x^{-\alpha}} \end{aligned}$$

Proof. From Proposition 4.7 we get

$$\begin{aligned} \text{Ad}(x)(\phi(H)) &= x^\alpha E + x^{-\alpha} \sigma(E) \\ \text{Ad}(x^{-1})(H) &= x^{-\alpha} \phi(E) + x^\alpha \sigma(\phi(E)), \end{aligned}$$

which leads to the following linear systems of equations.

$$\begin{aligned} \begin{pmatrix} H \\ \text{Ad}(x)(\phi(H)) \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ x^\alpha & x^{-\alpha} \end{pmatrix} \begin{pmatrix} E \\ \sigma(E) \end{pmatrix} \\ \begin{pmatrix} \phi(H) \\ \text{Ad}(x^{-1})(H) \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ x^{-\alpha} & x^\alpha \end{pmatrix} \begin{pmatrix} \phi(E) \\ \sigma(\phi(E)) \end{pmatrix}. \end{aligned}$$

Solving these systems yields the claimed expressions. Note that by Lemma 4.8, we always have $x^\alpha \neq x^{-\alpha}$, whence the determinant is nonzero. \square

Lemma 4.10. *For each $x \in C' \cap G_{rs}$ we can decompose \mathfrak{g} as follows:*

$$\begin{aligned} \mathfrak{g}_\mathbb{C} &= \mathfrak{c}'_\mathbb{C} \oplus (\mathfrak{h}_\mathbb{C} + \text{Ad}(x)(\mathfrak{h}_\mathbb{C})) \\ &= \mathfrak{c}'_\mathbb{C} \oplus (\mathfrak{h}_\mathbb{C} + \text{Ad}(x^{-1})(\mathfrak{h}_\mathbb{C})), \end{aligned}$$

where the intersection of the last two summands is $\mathfrak{m}'_\mathbb{C}$.

Proof. We are showing the claims for the first decomposition. The others follow by applying $\text{Ad}(x^{-1})$. Note that we assumed that $\text{Ad}(t)$ leave \mathfrak{c}' (and its complexification) invariant. Since $\text{Ad}(t)$ is orthogonal with respect to B , and because of Lemma 4.5, it therefore also leaves \mathfrak{m}' (and its complexification) invariant.

We evidently have the inclusion “ \supseteq ”, and the inclusion “ \subseteq ” follows from Proposition 4.9. So it remains to show that

$$\mathfrak{c}'_\mathbb{C} \cap (\mathfrak{h}_\mathbb{C} + \text{Ad}(x)(\mathfrak{h}_\mathbb{C})) = 0$$

and that

$$\mathfrak{h}_\mathbb{C} \cap \text{Ad}(x)(\mathfrak{h}_\mathbb{C}) = \mathfrak{m}'_\mathbb{C}.$$

For the first note that $\mathfrak{c}'_\mathbb{C}$ and $\mathfrak{h}_\mathbb{C}$ are orthogonal with respect to B because they lie in different eigenspaces of σ , which is orthogonal with respect to B . Since B restricted to $\mathfrak{c}'_\mathbb{C}$ is also non-degenerate, we conclude that $\mathfrak{c}'_\mathbb{C} \cap \mathfrak{h}_\mathbb{C} = 0$. Applying

$\text{Ad}(x)$ and the fact that $\mathfrak{c}'_{\mathbb{C}}$ is invariant under $\text{Ad}(x)$ yields the desired directness of the sum.

Let $Y \in \mathfrak{h}_{\mathbb{C}} \cap \text{Ad}(x)(\mathfrak{h}_{\mathbb{C}})$. We expand

$$Y = Y_{0,+} + Y_{0,-} + \sum_{\alpha \in \Sigma} Y_{\alpha}.$$

Then due to σ -invariance we have $Y_{0,-} = 0$ and $Y_{-\alpha} = \sigma(Y_{\alpha})$. Applying $\text{Ad}(x^{-1})$ yields

$$\text{Ad}(x^{-1})(Y) = \epsilon_0^{-1} \phi(Y_{0,+}) + \sum_{\alpha \in \Sigma^+} (x^{-\alpha} \phi(Y_{\alpha}) + x^{\alpha} \sigma(\phi(Y_{\alpha}))).$$

The fact that this is also σ -invariant and $x^{\alpha} \neq x^{-\alpha}$ (by Lemma 4.8) yields that $Y_{\alpha} = 0$, hence also $Y_{-\alpha}$, so that $Y = Y_{0,+} \in \mathfrak{m}'_{\mathbb{C}}$. \square

Corollary 4.11. *For $x \in C' \cap G_{rs}$ we can decompose*

$$U(\mathfrak{g}) = \text{Ad}(x^{-1})(U(\mathfrak{h}))S(\mathfrak{c}')U(\mathfrak{h}) = U(\mathfrak{h})S(\mathfrak{c}')\text{Ad}(x)(U(\mathfrak{h})).$$

with the only ambiguity being $U(\mathfrak{m}')$ acting on $\text{Ad}(x^{-1})(U(\mathfrak{h}))$ (resp. $U(\mathfrak{h}))$) from the left and on $U(\mathfrak{h})$ (resp. $\text{Ad}(x)(U(\mathfrak{h}))$) from the right.

Proof. From Lemma 4.10 we obtain that if \mathfrak{q} is a B -orthogonal complement of \mathfrak{m}' in \mathfrak{h} , we have $\mathfrak{g}_{\mathbb{C}} = \mathfrak{c}'_{\mathbb{C}} \oplus \mathfrak{m}_{\mathbb{C}} \oplus \mathfrak{q}_{\mathbb{C}} \oplus \text{Ad}(x)(\mathfrak{q}_{\mathbb{C}})$ (and the same expression with x inverted). By the Poincaré–Birkhoff–Witt theorem the multiplication map generates isomorphisms

$$U(\mathfrak{g}) \cong \text{Ad}(x^{-1})(U(\mathfrak{q})) \otimes U(\mathfrak{m}') \otimes S(\mathfrak{c}') \otimes U(\mathfrak{q}) \cong U(\mathfrak{q}) \otimes U(\mathfrak{m}') \otimes U(\mathfrak{c}') \otimes \text{Ad}(x)(U(\mathfrak{q})).$$

Applying the same to $U(\mathfrak{h})$ we see that $U(\mathfrak{h}) = U(\mathfrak{q}) \otimes U(\mathfrak{m}') = U(\mathfrak{m}') \otimes U(\mathfrak{q})$ and $\text{Ad}(x^{-1})(U(\mathfrak{h})) = \text{Ad}(x^{-1})(U(\mathfrak{q})) \otimes U(\mathfrak{m}')$, $\text{Ad}(x)(U(\mathfrak{h})) = U(\mathfrak{m}') \otimes \text{Ad}(x)(U(\mathfrak{q}))$,

so that the above decompositions become

$$\begin{aligned} U(\mathfrak{g}) &= \text{Ad}(x^{-1})(U(\mathfrak{h})) \otimes_{U(\mathfrak{m}')} (S(\mathfrak{c}') \otimes U(\mathfrak{h})) \\ U(\mathfrak{g}) &= (U(\mathfrak{h}) \otimes S(\mathfrak{c}')) \otimes_{U(\mathfrak{m}')} \text{Ad}(x)(U(\mathfrak{h})), \end{aligned}$$

where $U(\mathfrak{m}')$ acts only on the $U(\mathfrak{h})$ components. \square

Corollary 4.12. *There are maps*

$$\Pi, \tilde{\Pi}: U(\mathfrak{g}) \rightarrow C^{\infty}(C' \cap G_{rs}) \otimes S(\mathfrak{c}') \otimes (U(\mathfrak{h}) \otimes_{U(\mathfrak{m}')} U(\mathfrak{h}))$$

such that for

$$\Pi(p) = \sum_i f_i \otimes p_i \otimes u_i \otimes v_i \quad \tilde{\Pi}(u) = \sum_j \tilde{f}_j \otimes \tilde{p}_j \otimes \tilde{u}_j \otimes \tilde{v}_j$$

we have

$$p = \sum_i f_i(x) \text{Ad}(x^{-1})(u_i p_i) v_i = \sum_j \tilde{f}_j(x) \tilde{u}_j \tilde{p}_j \text{Ad}(x)(\tilde{v}_j)$$

for all $x \in C' \cap G_{rs}$. Here, $U(\mathfrak{m}')$ acts on the two $U(\mathfrak{h})$ -tensor factors as follows: the right action (on the 1st factor) is twisted by $\text{Ad}(t)$, and the left action is just the usual multiplication.

Proof. Follows from Corollary 4.11 and Proposition 4.9. For the $U(\mathfrak{m}')$ -actions note that we can write

$$p = \sum_i f_i(x) \text{Ad}(x^{-1})(u_i m_i p_i) v_i$$

where u_i, v_i are written entirely in terms of \mathfrak{q} , and $m_i \in U(\mathfrak{m}')$, then this equals

$$= \sum_i f_i(x) \text{Ad}(x^{-1})(u_i p_i) \text{Ad}(t^{-1})(m_i) v_i,$$

meaning that Π maps p to both

$$\sum_i f_i \otimes p_i \otimes u_i m_i \otimes v_i$$

and

$$\sum_i f_i \otimes p_i \otimes u_i \otimes \text{Ad}(t^{-1})v_i,$$

which should be equal. Similarly for $\tilde{\Pi}$. \square

Theorem 4.13. *Let W be a finite-dimensional H -bimodule and write π_L, π_R for the left and right actions, respectively (π_R is then a representation of H^{op}). There are maps*

$$R^W, \tilde{R}^W: U(\mathfrak{g}) \rightarrow C^\infty(C' \cap G_{rs}) \otimes S(\mathfrak{c}') \otimes \text{Hom}(W^{\mathfrak{m}'}, W)$$

such that for every $f \in E^W(G, H)$, $x \in C' \cap G_{rs}$, and $p \in U(\mathfrak{g})$ we have

$$\begin{aligned} (p \cdot f)(x) &= R^W(p)(x)(f|_{C'})(x) \\ (f \cdot p)(x) &= \tilde{R}^W(p)(x)(f|_{C'})(x). \end{aligned}$$

In particular, for $p \in U(\mathfrak{g})^{\mathfrak{m}'}$, the matrix parts of R^W, \tilde{R}^W lie in $\text{End}(W^{\mathfrak{m}'})$. Here, $Y \in \mathfrak{m}'$ acts as follows on $v \in W$:

$$Y \cdot_{\mathfrak{m}'} v = \text{Ad}(t)(Y) \cdot v - v \cdot Y.$$

Proof. Define R^W, \tilde{R}^W by post-composing $\Pi, \tilde{\Pi}$ with the representations π_L, π_R as follows: the last two tensor legs, $U(\mathfrak{h}) \otimes_{U(\mathfrak{m}')} U(\mathfrak{h})$ act as follows on $W^{\mathfrak{m}'}$:

$$(a \otimes b) \cdot \psi = a \cdot \psi \cdot b.$$

This is well-defined precisely because f is an \mathfrak{m}' -homomorphism twisted in the correct way.

We now show that R^W, \tilde{R}^W indeed yield the radial parts of the left and right actions on matrix-spherical functions. Let $f \in E^W(G, H)$, $x \in C' \cap G_{rs}$, $p \in U(\mathfrak{g})$, say

$$\Pi(p) = \sum_i f_i \otimes p_i \otimes u_i \otimes v_i,$$

so that

$$\forall x \in C' \cap G_{rs}: \sum_i f_i(x) \text{Ad}(x^{-1})(u_i p_i) v_i.$$

For any $x \in C' \cap G_{rs}$ we then have

$$(p \cdot f)(x) = \sum_i f_i(x) (\text{Ad}(x^{-1})(u_i p_i) v_i \cdot f)(x) = \sum_i f_i(x) (v_i \cdot f \cdot u_i p_i)(x)$$

by Lemma 2.2. Corollary 2.4 then allows us to write this as

$$\begin{aligned} (p \cdot f)(x) &= \sum_i f_i(x) u_i \cdot (f \cdot p_i)(x) \cdot v_i \\ &= R(p)(x) f(x), \end{aligned}$$

where we interpret $S(\mathfrak{c}')$ as differential operators acting on $C^\infty(G, W)$ via right-invariant vector fields.

To see that $R^W(p)$ for $p \in U(\mathfrak{g})^{\mathfrak{m}'}$ preserves pointwise \mathfrak{m}' -invariance, let $Y \in \mathfrak{m}'$, and note that

$$\begin{aligned} \text{Ad}(t)(Y) \circ (p \cdot f)(x) &= \sum_i f_i(x) \text{Ad}(t)(Y) u_i \cdot (p_i \cdot f)(x) \cdot v_i \\ &= \left(\sum_i f_i(x) Y \text{Ad}(x^{-1})(u_i) p_i v_i \cdot f \right)(x) \\ &= (Y p \cdot f)(x). \end{aligned}$$

Now, since p commutes with \mathfrak{m}' , this equals

$$\begin{aligned} (p Y \cdot f)(x) &= \left(\sum_i f_i(x) \text{Ad}(x^{-1})(u_i) p_i v_i Y \cdot f \right)(x) = \sum_i f_i(x) u_i \cdot (f \cdot p_i)(x) \cdot v_i Y \\ &= (p \cdot f)(x) \cdot Y. \end{aligned}$$

The statements about \tilde{R}^W follow analogously. \square

Recall from Lemma 2.7 that the restriction of a MSF maps not only to $W^{\mathfrak{m}'}$, but even to $W^{Z_{C'}}$. If we consider that invariance, we get more out of the previous result.

Definition 4.14. *Define*

$$M' := Z_H(\mathfrak{c}') \cap t^{-1} H t.$$

This is a closed subgroup of H with Lie algebra \mathfrak{m}' , since $\text{Ad}(t^{-1})$ leaves \mathfrak{m}' invariant.

Proposition 4.15. *The map $h \mapsto (tht^{-1}, h)$ is a (well-defined) isomorphism $M' \cong Z_{C'}$ that intertwines between the partially twisted action of M' on W and the action of $Z_{C'}$ on W . In particular, $W^{Z_{C'}}$ is a subspace of $W^{\mathfrak{m}'}$.*

Proof. Note that by definition of M' , both tht^{-1} and h are elements of H , so the map is well-defined. Furthermore, by Proposition 4.4, it is an isomorphism.

The partially twisted action of M' on W is defined so as to differentiate to the one of \mathfrak{m}' :

$$h \cdot_{M'} \psi = tht^{-1} \cdot \psi \cdot h^{-1},$$

which is exactly how $(tht^{-1}, h) \in Z_{C'}$ acts on $\psi \in W$.

This shows that $W^{Z_{C'}} = W^{M'}$ is a subspace of $W^{\mathfrak{m}'}$, which is equal if M' is the analytic subgroup generated by \mathfrak{m}' (i.e. is connected). \square

Corollary 4.16. *R^W and \tilde{R}^W restrict to maps*

$$U(\mathfrak{g})^{M'} \rightarrow C^\infty(C' \cap G_{rs}) \otimes S(\mathfrak{c}') \otimes \text{End}(W^{M'}).$$

Proof. Let $f \in E^W(G, H)$ and $p \in U(\mathfrak{g})^{M'}$. Let $m \in M'$ and $x \in C' \cap G_{rs}$, then

$$\begin{aligned} t m t^{-1} \cdot (p \cdot f)(x) \cdot m^{-1} &= \sum_i f_i(x) t m t^{-1} \cdot u_i \cdot (f \cdot p_i)(x) \cdot v_i \cdot m^{-1} \\ &= \sum_i f_i(x) (v_i \cdot m^{-1} \cdot f \cdot t m t^{-1} \cdot u_i p_i)(x) \\ &= \sum_i f_i(x) (m^{-1} \cdot \text{Ad}(m)(v_i) \cdot f \cdot \text{Ad}(t m t^{-1})(u_i p_i) \cdot t m t^{-1})(x) \\ &= \sum_i f_i(x) (\text{Ad}(m)(v_i) \cdot f \cdot \text{Ad}(t m t^{-1})(u_i p_i))(t m t^{-1} x m^{-1}). \end{aligned}$$

By Lemma 2.2, this equals

$$= \sum_i f_i(x) (\text{Ad}(mx^{-1})(u_i p_i) \text{Ad}(m)(v_i) \cdot f)(tmt^{-1}xm^{-1}).$$

By Propositions 4.15 and 4.4, this equals

$$\begin{aligned} &= \sum_i f_i(x) (\text{Ad}(m)(\text{Ad}(x^{-1})(u_i p_i) v_i) \cdot f)(x) \\ &= (\text{Ad}(m)(p) \cdot f)(x) = (p \cdot f)(x) \end{aligned}$$

since p is invariant under M' . \square

Corollary 4.17. *Let V, W be finite-dimensional H -modules and assume that V, \overline{W} (\overline{W} is the same as W , but with the \mathfrak{m}' -action twisted by $\text{Ad}(t)$) are semisimple \mathfrak{m}' -modules. Then*

$$(\text{Hom}(V, W))^{\mathfrak{m}'} = \text{Hom}_{\mathfrak{m}'}(V, \overline{W}) = \bigoplus_{\rho \in \widehat{\mathfrak{m}'}} \mathbb{C}^{[V:\rho][\overline{W}:\rho]} = : \mathcal{V}.$$

In that case, the maps R^W, \widetilde{R}^W map to

$$U(\mathfrak{g})^{\mathfrak{m}'} \rightarrow C^\infty(C' \cap G_{rs}) \otimes S(\mathfrak{c}') \otimes \text{End}(\mathcal{V}).$$

An analogous statement holds when we define \mathcal{V} in terms of the group M' instead of \mathfrak{m}' .

4.3. Radial Part of the Quadratic Casimir Element $\Omega_{\mathfrak{g}}$. In this subsection we will now compute Π of the quadratic Casimir element ($\widetilde{\Pi}$ will turn out to be the same). The application to MSF for a concrete H -bimodule W is then straightforward.

Definition 4.18. *Let $\alpha \in (\mathfrak{c}'_{\mathbb{C}})^*$. Write $C_\alpha \in \mathfrak{c}'_{\mathbb{C}}$ for the unique element X such that*

$$\forall Y \in \mathfrak{c}'_{\mathbb{C}}: \quad \alpha(Y) = B(X, Y).$$

This exists since due to Proposition 4.1(iv) and Lemma 4.5 B is non-degenerate when restricted to $\mathfrak{c}'_{\mathbb{C}}$.

Lemma 4.19. *Let $E \in \mathfrak{g}_\alpha$ and $H := E + \sigma(E)$.*

- (i) *Then $[E, \sigma(E)] = -B_\sigma(E, E)C_\alpha$.*
- (ii) *For $x \in C'$ we have $[H, \text{Ad}(x)(\phi(H))] = (x^\alpha - x^{-\alpha})B_\sigma(E, E)C_\alpha$.*
- (iii) *For $x \in C'$ we have $[\text{Ad}(x^{-1})(H), \phi(H)] = (x^\alpha - x^{-\alpha})B_\sigma(E, E)C_{t\alpha}$.*

Proof. (i) From Proposition 4.1(iii,ii) we see that $[E, \sigma(E)] \in \mathfrak{g}_0$. Furthermore, since σ is an involutive Lie algebra homomorphism, we have $\sigma([E, \sigma(E)]) = -[E, \sigma(E)]$, so that by Lemma 4.5, we have $[E, \sigma(E)] \in \mathfrak{c}'_{\mathbb{C}}$. Let $Y \in \mathfrak{c}'_{\mathbb{C}}$, then

$$\begin{aligned} B(Y, [E, \sigma(E)]) &= B([Y, E], \sigma(E)) = \alpha(Y)B(E, \sigma(E)) = -\alpha(Y)B_\sigma(E, E), \\ &\text{hence } [E, \sigma(E)] = -B_\sigma(E, E)C_\alpha. \end{aligned}$$

(ii) From the proof of Proposition 4.9 we know that

$$[H, \text{Ad}(x)(\phi(H))] = [E + \sigma(E), x^\alpha E + x^{-\alpha} \sigma(E)] = -(x^\alpha - x^{-\alpha})[E, \sigma(E)],$$

which together with (i) implies the claim.

(iii) From the proof of Proposition 4.9 we know that

$$\begin{aligned} [\text{Ad}(x^{-1})(H), \phi(H)] &= [x^{-\alpha} \phi(E) + x^\alpha \sigma(\phi(E)), \phi(E) + \sigma(\phi(E))] \\ &= -(x^\alpha - x^{-\alpha})[\phi(E), \sigma(\phi(E))]. \end{aligned}$$

Together with (i) applied to $\phi(E)$, the fact that $B_\sigma(\phi(E), \phi(E)) = B_\sigma(E, E)$, and the fact that $\phi(E) \in \mathfrak{g}_{t\alpha}$, the claim follows.

□

Corollary 4.20. *Let $E \in \mathfrak{g}_\alpha$ and $x \in C' \cap G_{rs}$. Write $H := E + \sigma(E)$. Then*

$$\begin{aligned} \{E, \sigma(E)\} &= -2 \frac{H^2 + \text{Ad}(x)(\phi(H)^2) - (x^\alpha + x^{-\alpha})H \text{Ad}(x)(\phi(H))}{(x^\alpha - x^{-\alpha})^2} \\ &\quad - B_\sigma(E, E) \frac{x^\alpha + x^{-\alpha}}{x^\alpha - x^{-\alpha}} C_\alpha \\ \{\phi(E), \sigma(\phi(E))\} &= -2 \frac{\phi(H)^2 + \text{Ad}(x^{-1})(H^2) - (x^\alpha + x^{-\alpha}) \text{Ad}(x^{-1})(H)\phi(H)}{(x^\alpha - x^{-\alpha})^2} \\ &\quad - B_\sigma(E, E) \frac{x^\alpha + x^{-\alpha}}{x^\alpha - x^{-\alpha}} C_{t\alpha} \end{aligned}$$

Proof. By Proposition 4.9 we have

$$\begin{aligned} (x^\alpha - x^{-\alpha})^2 \{E, \sigma(E)\} &= \{\text{Ad}(x)(\phi(H)) - x^{-\alpha}H, x^\alpha H - \text{Ad}(x)(\phi(H))\} \\ &= -2H^2 - 2 \text{Ad}(x)(\phi(H)^2) + (x^\alpha + x^{-\alpha})\{H, \text{Ad}(x)(\phi(H))\}. \end{aligned}$$

We can write the anticommutator as $2H \text{Ad}(x)(\phi(H)) - [H, \text{Ad}(x)(\phi(H))]$, so that by Lemma 4.19 we have

$$= -2H^2 - 2 \text{Ad}(x)(\phi(H)^2) + 2(x^\alpha + x^{-\alpha})H \text{Ad}(x)(\phi(H)) - B_\sigma(E, E) \left((x^\alpha)^2 - (x^{-\alpha})^2 \right) C_\alpha.$$

Similarly, we have

$$\begin{aligned} (x^\alpha - x^{-\alpha}) \{\phi(E), \sigma(\phi(E))\} &= \{x^\alpha \phi(H) - \text{Ad}(x^{-1})(H), \text{Ad}(x^{-1})(H) - x^{-\alpha} \phi(H)\} \\ &= -2\phi(H)^2 - 2 \text{Ad}(x^{-1})(H^2) + (x^\alpha + x^{-\alpha})\{\text{Ad}(x^{-1})(H), H\}. \end{aligned}$$

The anticommutator equals $2 \text{Ad}(x^{-1})(H)H - [\text{Ad}(x^{-1})(H), H]$, so that by Lemma 4.19, we have

$$\begin{aligned} &= -2\phi(H)^2 - 2 \text{Ad}(x^{-1})(H^2) + 2(x^\alpha + x^{-\alpha}) \text{Ad}(x^{-1})(H)H \\ &\quad - B_\sigma(E, E) \left((x^\alpha)^2 - (x^{-\alpha})^2 \right) C_{t\alpha}. \end{aligned} \quad \square$$

Proposition 4.21. *Let $\alpha \in \Sigma$, let X_1, \dots, X_r be an orthonormal basis of \mathfrak{g}_α with respect to B_σ . Define*

$$A_\alpha := \sum_{i=1}^r (X_i + \sigma(X_i)) \otimes (X_i + \sigma(X_i)) \in U(\mathfrak{h}) \otimes U(\mathfrak{h}).$$

Then A_α does not depend on the choice of basis. For $(h, h') \in N_{C'}$ we have

$$\begin{aligned} (\text{Ad}(h) \otimes \text{Ad}(h))(A_\alpha) &= A_{\text{Ad}^*(h)(\alpha)} \\ (\text{Ad}(h') \otimes \text{Ad}(h'))(A_\alpha) &= A_{\text{Ad}^*(h')(\alpha)} \end{aligned}$$

and

$$(\text{Ad}(h) \otimes \text{Ad}(h')\phi)(A_\alpha) = \pm(1 \otimes \phi)A_{\text{Ad}^*(h)(\alpha)}$$

depending on whether

$$\epsilon_\alpha(hth'^{-1})^{-\text{Ad}^*(h)(\alpha)} = \pm 1.$$

Proof. Let X_1, \dots, X_r and Y_1, \dots, Y_r be orthonormal bases, say

$$X_i = \sum_{j=1}^r a_{ij} Y_j,$$

then $a_{ij} = B_\sigma(X_i, Y_j)$ and in particular also $Y_i = \sum_{j=1}^r a_{ji} X_j$, so that

$$\sum_{i=1}^r a_{ij} a_{ik} = \sum_{i=1}^r B_\sigma(a_{ik} X_i, Y_j) = B_\sigma(Y_k, Y_j) = \delta_{jk}.$$

Consequently,

$$\begin{aligned} \sum_{i=1}^r (X_i + \sigma(X_i)) \otimes (X_i + \sigma(X_i)) &= \sum_{i=1}^r \sum_{j,k=1}^r a_{ij} a_{ik} (Y_j + \sigma(Y_j)) \otimes (Y_k + \sigma(Y_k)) \\ &= \sum_{j,k=1}^r \sum_{i=1}^r a_{ij} a_{ik} (Y_j + \sigma(Y_j)) \otimes (Y_k + \sigma(Y_k)) \\ &= \sum_{i=1}^r (Y_i + \sigma(Y_i)) \otimes (Y_i + \sigma(Y_i)). \end{aligned}$$

For any Lie algebra automorphism ψ that leaves \mathfrak{c}' and B invariant and commutes with σ (e.g. $\text{Ad}(h)$ or $\text{Ad}(h')$ for $(h, h') \in N_{C'}$), the family $\phi(X_1), \dots, \phi(X_r)$ is a B_σ -orthonormal basis of $\mathfrak{g}_{\phi^*\alpha}$, so

$$(\phi \otimes \phi)(A_\alpha) = A_{\phi^*\alpha}.$$

Lastly, for $(\text{Ad}(h) \otimes \text{Ad}(h'))\phi(A_\alpha)$ note that by Proposition 4.7(iv) we have

$$\epsilon_\alpha(\text{Ad}^*(h)(\alpha)) = \pm 1,$$

and in particular

$$\begin{aligned} \text{Ad}(h)(X_i) &= \pm \phi(\text{Ad}(h')(\phi(X_i))) \\ \text{Ad}(h)(\sigma(X_i)) &= \pm \sigma(\phi(\text{Ad}(h')(\phi(X_i)))) \end{aligned}$$

so that we have

$$\begin{aligned} (\text{Ad}(h) \otimes \text{Ad}(h'))\phi A_\alpha &= \pm \sum_{i=1}^r (\phi \text{Ad}(h')\phi \otimes \text{Ad}(h')\phi)(X_i + \sigma(X_i)) \otimes (X_i + \sigma(X_i)) \\ &= \pm (1 \otimes \phi) A_{\text{Ad}^*(h)(\alpha)} \end{aligned}$$

seeing as $\phi \text{Ad}(h')\phi$ is orthogonal with respect to B_σ and commutes with σ , so that the expression in the sum becomes the expression we used to define A_α , except that

$$\phi \text{Ad}(h')\phi(X_1), \dots, \phi \text{Ad}(h')\phi(X_r)$$

is now a B_σ -orthonormal basis of $\mathfrak{g}_{\text{Ad}^*(h)(\alpha)}$. \square

Theorem 4.22. *The decompositions of the quadratic Casimir element $\Omega_{\mathfrak{g}}$ with respect to the standard Cartan subset C' are as follows:*

$$\begin{aligned} \Pi(\Omega_{\mathfrak{g}}) = \tilde{\Pi}(\Omega_{\mathfrak{g}}) &= \Omega_{\mathfrak{c}'} + \sum_{\alpha \in \Sigma} \frac{n_\alpha}{2} \coth_\alpha C_\alpha + \Omega_{\mathfrak{m}'} \\ &+ \sum_{\alpha \in \Sigma} \frac{\text{csch}_\alpha^2}{4} (m(A_\alpha) \otimes 1 + 1 \otimes m(A_{t\alpha}) + 2(1 \otimes \phi)A_\alpha) \\ &- \sum_{\alpha \in \Sigma} \frac{\text{csch}_{\alpha/2}^2}{4} (1 \otimes \phi)A_\alpha. \end{aligned}$$

where $m: U(\mathfrak{h}) \otimes U(\mathfrak{h}) \rightarrow U(\mathfrak{h})$ is the multiplication map and $n_\alpha := \dim(\mathfrak{g}_\alpha)$.

Proof. Let $C_1, \dots, C_r \in \mathfrak{c}'_C$ and $M_1, \dots, M_s \in \mathfrak{m}'_C$ be orthonormal bases (with respect to B). For $\alpha \in \Sigma$ let $E_{\alpha,1}, \dots, E_{\alpha,n_\alpha} \in \mathfrak{g}_\alpha$ be an orthonormal basis with respect to B_σ . Without loss of generality assume that $E_{-\alpha,i} = \sigma(E_{\alpha,i})$. Then the following bases are dual to each other with respect to B :

$$\begin{aligned} C_1, \dots, C_r, M_1, \dots, M_s, (E_{\alpha,1}, \dots, E_{\alpha,n_\alpha})_{\alpha \in \Sigma} \\ C_1, \dots, C_r, M_1, \dots, M_s, (-\sigma(E_{\alpha,1}), \dots, -\sigma(E_{\alpha,n_\alpha}))_{\alpha \in \Sigma}. \end{aligned}$$

Thus,

$$\Omega_{\mathfrak{g}} = \sum_{i=1}^r C_i^2 + \sum_{i=1}^s M_i^2 - \sum_{\alpha \in \Sigma} \sum_{i=1}^{n_\alpha} E_{\alpha,i} \sigma(E_{\alpha,i}).$$

The first two sums are just $\Omega_{\mathfrak{c}'}$ and $\Omega_{\mathfrak{m}'}$, respectively. Symmetrising the summands over α somewhat we obtain

$$\Omega_{\mathfrak{g}} = \Omega_{\mathfrak{c}'} + \Omega_{\mathfrak{m}'} - \frac{1}{2} \sum_{\alpha \in \Sigma} \sum_{i=1}^{n_\alpha} \{E_{\alpha,i}, \sigma(E_{\alpha,i})\}.$$

Let $x \in C' \cap G_{rs}$. Substituting in the expression from Corollary 4.20 and setting $H_{\alpha,i} := E_{\alpha,i} + \sigma(E_{\alpha,i}) = E_{\alpha,i} + E_{-\alpha,i}$ we get

$$\begin{aligned} \Omega_{\mathfrak{g}} &= \Omega_{\mathfrak{c}'} + \Omega_{\mathfrak{m}'} + \frac{1}{2} \sum_{\alpha \in \Sigma} \sum_{i=1}^{n_\alpha} B_\sigma(E_{\alpha,i}, E_{\alpha,i}) \frac{x^\alpha + x^{-\alpha}}{x^\alpha - x^{-\alpha}} C_\alpha \\ &\quad + \sum_{\alpha \in \Sigma} \sum_{i=1}^{n_\alpha} \frac{H_{\alpha,i}^2 + \text{Ad}(x)(\phi(H_{\alpha,i})^2) - (x^\alpha + x^{-\alpha})H_{\alpha,i} \text{Ad}(x)(\phi(H_{\alpha,i}))}{(x^\alpha - x^{-\alpha})^2} \\ &= \Omega_{\mathfrak{c}'} + \Omega_{\mathfrak{m}'} + \sum_{\alpha \in \Sigma} \frac{n_\alpha}{2} \frac{x^\alpha + x^{-\alpha}}{x^\alpha - x^{-\alpha}} C_\alpha \\ &\quad + \sum_{\alpha \in \Sigma} \frac{m(A_\alpha) + \text{Ad}(x)(m(A_{t\alpha})) - (x^\alpha + x^{-\alpha})m(1 \otimes \text{Ad}(x)\phi)(A_\alpha)}{(x^\alpha - x^{-\alpha})^2}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \Omega_{\mathfrak{g}} &= \Omega_{\mathfrak{c}'} + \Omega_{\mathfrak{m}'} + \frac{1}{2} \sum_{\alpha \in \Sigma} \sum_{i=1}^{n_\alpha} B_\sigma(\phi(E_{\alpha,i}), \phi(E_{\alpha,i})) \frac{x^{t\alpha} + x^{t\alpha}}{x^{t\alpha} - x^{-t\alpha}} \text{Ad}(x^{-1})(C_{t\alpha}) \\ &\quad + \sum_{\alpha \in \Sigma} \sum_{i=1}^{n_\alpha} \frac{H_{\alpha,i}^2 + \text{Ad}(x^{-1})(\phi(H_{\alpha,i})^2) - (x^{t\alpha} + x^{-t\alpha}) \text{Ad}(x^{-1})(\phi(H_{\alpha,i}))H_{\alpha,i}}{(x^{t\alpha} - x^{-t\alpha})^2} \\ &= \Omega_{\mathfrak{c}'} + \Omega_{\mathfrak{m}'} + \sum_{\alpha \in \Sigma} \frac{n_\alpha}{2} \frac{x^\alpha + x^{-\alpha}}{x^\alpha - x^{-\alpha}} \text{Ad}(x^{-1})(C_\alpha) \\ &\quad + \sum_{\alpha \in \Sigma} \frac{m(A_{t\alpha}) + \text{Ad}(x^{-1})(m(A_\alpha)) - (x^\alpha + x^{-\alpha})m(\text{Ad}(x^{-1})\phi \otimes 1)A_{t\alpha}}{(x^\alpha - x^{-\alpha})^2}. \end{aligned}$$

Note that these expressions are both in the shape that we can read off $\Pi(\Omega_{\mathfrak{g}}), \widetilde{\Omega}_{\mathfrak{g}}$. Writing $\text{coth}_\alpha(x) := \frac{x^\alpha + x^{-\alpha}}{x^\alpha - x^{-\alpha}}$ and $\text{csch}_\alpha(x) := 2(x^\alpha - x^{-\alpha})^{-1}$, we thus obtain

$$\begin{aligned} \widetilde{\Pi}(\Omega_{\mathfrak{g}}) &= \Omega_{\mathfrak{c}'} + \Omega_{\mathfrak{m}'} + \sum_{\alpha \in \Sigma} \frac{n_\alpha}{2} \text{coth}_\alpha C_\alpha \\ &\quad + \sum_{\alpha \in \Sigma} \frac{\text{csch}_\alpha^2}{4} (m(A_\alpha) \otimes 1 + 1 \otimes m(A_{t\alpha})) \\ &\quad - \sum_{\alpha \in \Sigma} \frac{\text{csch}_\alpha \text{coth}_\alpha}{2} (1 \otimes \phi) A_\alpha \\ \Pi(\Omega_{\mathfrak{g}}) &= \Omega_{\mathfrak{c}'} + \Omega_{\mathfrak{m}'} + \sum_{\alpha \in \Sigma} \frac{n_\alpha}{2} \text{coth}_\alpha C_\alpha \\ &\quad + \sum_{\alpha \in \Sigma} \frac{\text{csch}_\alpha^2}{4} (1 \otimes m(A_{t\alpha}) + m(A_\alpha) \otimes 1) \\ &\quad - \sum_{\alpha \in \Sigma} \frac{\text{csch}_\alpha \text{coth}_\alpha}{2} (\phi \otimes 1) A_{t\alpha}, \end{aligned}$$

which equals $\tilde{\Pi}(\Omega_{\mathfrak{g}})$ because $(\phi \otimes \phi)A_\alpha = A_{t\alpha}$ (by Proposition 4.21). Lastly, noting that

$$\begin{aligned} \frac{\operatorname{csch}_\alpha(x) \operatorname{coth}_\alpha(x)}{2} &= \frac{x^\alpha + x^{-\alpha} + 2}{(x^\alpha - x^{-\alpha})} - \frac{\operatorname{csch}_\alpha^2(x)}{2} \\ &= \frac{(1+x^\alpha)(1+x^{-\alpha})}{(1+x^\alpha)(1-x^{-\alpha})(1+x^{-\alpha})(x^\alpha-1)} - \frac{\operatorname{csch}_\alpha^2(x)}{2} \\ &= \frac{\operatorname{csch}_{\alpha/2}^2(x)}{4} - \frac{\operatorname{csch}_\alpha^2(x)}{2}, \end{aligned}$$

where the square of $\operatorname{csch}_{\alpha/2}$ is a well-defined quantity obtained by multiplying out the product. In light of this we can also rewrite $\Pi(\Omega_{\mathfrak{g}}) = \tilde{\Pi}(\Omega_{\mathfrak{g}})$ as

$$\begin{aligned} \Pi(\Omega_{\mathfrak{g}}) &= \tilde{\Pi}(\Omega_{\mathfrak{g}}) = \Omega_{\mathfrak{c}'} + \sum_{\alpha \in \Sigma} \frac{n_\alpha}{2} \operatorname{coth}_\alpha C_\alpha + \Omega_{\mathfrak{m}'} \\ &\quad + \sum_{\alpha \in \Sigma} \frac{\operatorname{csch}_\alpha^2}{4} (m(A_\alpha) \otimes 1 + 1 \otimes m(A_{t\alpha}) + 2(1 \otimes \phi)A_\alpha) \\ &\quad - \sum_{\alpha \in \Sigma} \frac{\operatorname{csch}_{\alpha/2}^2}{4} (1 \otimes \phi)A_\alpha. \end{aligned}$$

□

Corollary 4.23. *In the case where $\operatorname{Ad}(t) = 1$, we have $(\exp(X)t)^\alpha = \exp(\alpha(X))$, and the radial part of $\Omega_{\mathfrak{g}}$ simplifies to*

$$\begin{aligned} \Pi(\Omega_{\mathfrak{g}}) &= \Omega_{\mathfrak{c}'} + \sum_{\alpha \in \Sigma} \frac{n_\alpha}{2} \operatorname{coth}_\alpha C_\alpha + \Omega_{\mathfrak{m}'} \\ &\quad + \sum_{\alpha \in \Sigma} \frac{\operatorname{csch}_\alpha^2}{4} (m(A_\alpha) \otimes 1 + 1 \otimes m(A_\alpha) + 2A_\alpha) \\ &\quad - \sum_{\alpha \in \Sigma} \frac{\operatorname{csch}_{\alpha/2}^2}{4} A_\alpha \end{aligned}$$

where $\operatorname{coth}_\alpha, \operatorname{csch}_\alpha$ are related to the usual hyperbolic functions: if $x = \exp(X)$, then

$$\operatorname{coth}_\alpha(x) = \operatorname{coth}(\alpha(X)), \quad \operatorname{csch}_\alpha(x) = \operatorname{csch}(\alpha(X)).$$

5. MATHEMATICAL SETUP FOR CONFORMAL BLOCKS

Let $p + q = d > 2$ be natural numbers with $p \geq q$. Most commonly, we will encounter $q = 0$ (*Euclidean*) or $q = 1$ (*Lorentzian*). Let furthermore η denote the standard bilinear form of signature (p, q) on $\mathbb{R}^{p,q} := \mathbb{R}^d$ and by abuse of notation also the one of signature $(p+1, q+1)$ on $\mathbb{R}^{p+1, q+1} = \mathbb{R}^{d+2}$.

We will use lower case Greek letters to denote indices pertaining to $\mathbb{R}^{p+1, q+1}$ and lower case Latin letters for $\mathbb{R}^{p,q}$. For both we shall use the usual index raising and lowering conventions of physics (with the pseudo-inner product η), such that $A_\mu = A^\mu$ if $\mu \leq p$ and $A_\mu = -A^\mu$ if $\mu > p$ (analogously for A_i).

5.1. Conformal Compactification.

Definition 5.1. *Write $q: \mathbb{R}^{p+1, q+1} \setminus \{0\} \rightarrow \mathbb{RP}^{d+1}$ for the projectivisation map. The real variety*

$$\widehat{\mathbb{R}^{p,q}} := \{q(v) \mid v \in \mathbb{R}^{p+1, q+1} \setminus \{0\}, \eta(v, v) = 0\}$$

is called the conformal compactification of $\mathbb{R}^{p,q}$.

Note that since $\widehat{\mathbb{R}^{p,q}}$ is Zariski closed, it is also closed in the Euclidean topology. Since $\mathbb{R}\mathbb{P}^{d+1}$ is compact, that entails that $\widehat{\mathbb{R}^{p,q}}$ is also compact, as the name would imply.

We first establish that $\widehat{\mathbb{R}^{p,q}}$ is a smooth manifold in a useful sense and that it is indeed a compactification of $\mathbb{R}^{p,q}$.

Lemma 5.2. *$\widehat{\mathbb{R}^{p,q}}$ is an embedded submanifold of $\mathbb{R}\mathbb{P}^{d+1}$. Furthermore, the map*

$$\iota: \mathbb{R}^{p,q} \rightarrow \widehat{\mathbb{R}^{p,q}}, \quad v \mapsto (1 - \eta(v, v) : 2v : 1 + \eta(v, v))$$

is a diffeomorphism with $\iota(\mathbb{R}^{p,q})$, which is dense in $\widehat{\mathbb{R}^{p,q}}$.

Proof. Note that $\widehat{\mathbb{R}^{p,q}}$ is regular since η is nondegenerate. Next, we define $f: \widehat{\mathbb{R}^{p,q}} \cap \{(v_0 : \dots : v_{d+1}) \mid v_0 + v_{d+1} \neq 0\} \rightarrow \mathbb{R}^{p,q}$ by

$$(v_0 : v : v_{d+1}) \mapsto \frac{v}{v_0 + v_{d+1}}.$$

This map is evidently well-defined and smooth. Note also that

$$f(\iota(v)) = v$$

for all $v \in \mathbb{R}^{p,q}$. If $(v_0 : v : v_{d+1}) \in \widehat{\mathbb{R}^{p,q}}$, we have $\eta(v, v) = -(v_0 + v_{d+1})(v_0 - v_{d+1})$, so that

$$\begin{aligned} \iota(f(v_0 : v : v_{d+1})) &= \iota\left(\frac{v}{v_0 + v_{d+1}}\right) \\ &= \left(1 - \frac{\eta(v, v)}{(v_0 + v_{d+1})^2} : \frac{2v}{v_0 + v_{d+1}} : 1 + \frac{v, v}{(v_0 + v_{d+1})^2}\right) \\ &= (v_0 + v_{d+1} + v_0 - v_{d+1} : 2v : v_0 + v_{d+1} - v_0 + v_{d+1}) \\ &= (v_0 : v : v_{d+1}). \end{aligned}$$

Consequently, ι is a diffeomorphism and f is its inverse. To see that $\iota(\mathbb{R}^{p,q})$ is dense, note that any element of the form

$$(v_0 : v : -v_0)$$

can be reached as follows: if $v = 0$, then the point $(1:0:-1) = \infty$ can be reached as

$$\begin{aligned} \lim_{t \rightarrow \infty} \iota(tw) &= \lim_{t \rightarrow \infty} (1 - t^2\eta(w, w) : 2tw : 1 + t^2\eta(w, w)) \\ &= \lim_{t \rightarrow \infty} (\eta(w, w) - t^{-2} : -2t^{-2}w : -\eta(w, w) - t^{-2}) \\ &= (1:0:-1) \end{aligned}$$

for any $w \in \mathbb{R}^{p,q}$ with $\eta(w, w) \neq 0$. If $v \neq 0$, there is a vector w such that $\eta(v, w) \neq 0$. By rescaling, we can choose that inner product to be $-v_0$. Note that $\eta(v, v) = 0$. Then

$$\begin{aligned} \lim_{t \rightarrow \infty} \iota(w + vt) &= \lim_{t \rightarrow \infty} (1 - \eta(w, w) + 2tv_0 : 2w + 2vt : 1 + \eta(w, w) - 2tv_0) \\ &= \lim_{t \rightarrow \infty} ((1 - \eta(w, w))t^{-1} + 2v_0 : 2t^{-1}w + 2v : (1 + \eta(w, w))t^{-1} - 2v_0) \\ &= (v_0 : v : -v_0). \end{aligned} \quad \square$$

5.2. Conformal Group G and its Structure.

Lemma 5.3. *$G := SO(p+1, q+1)_0$ is the biggest classical connected Lie group acting on $\widehat{\mathbb{R}^{p,q}}$ by projective transformations.*

Proof. The group of projective transformations of \mathbb{RP}^{d+1} is $PGL(d+2)$, i.e. $GL(d+2)/\mathbb{R}^\times$. Evidently, those that leave $\widehat{\mathbb{R}^{p,q}}$ invariant are

$$\begin{aligned} & \{g \in GL(d+2) \mid \forall v \in \mathbb{R}^{p+1,q+1}: \eta(v,v) = 0 \Rightarrow \eta(gv,gv) = 0\}/\mathbb{R}^\times \\ & = \{g \in GL(d+2) \mid \forall v \in \mathbb{C}^{p+1,q+1}: \eta(v,v) = 0 \Rightarrow \eta(gv,gv) = 0\}/\mathbb{R}^\times \\ & = \{g \in GL(d+2) \mid \mathbb{V}(\eta) \subseteq \mathbb{V}(\eta \circ g)\}/\mathbb{R}^\times, \end{aligned}$$

where we used that we can also complexify η and $\mathbb{R}^{p+1,q+1}$ and the fact that the real restriction of η 's zero set is Zariski dense. By Hilbert's Nullstellensatz, $\mathbb{V}(\eta) \subseteq \mathbb{V}(\eta \circ g)$ implies that the polynomial $\eta \circ g$ is a multiple of η (as the ideals generated by both are radical). Since they are both homogeneous of degree 2, this multiple must be a nonzero scalar, and since η, g have real coefficients, it must be a nonzero real, i.e.

$$\{g \in GL(d+2) \mid \exists C \in \mathbb{R}^\times: \eta \circ g = C\eta\}/\mathbb{R}^\times$$

is the group of projective transformations leaving $\widehat{\mathbb{R}^{p,q}}$ invariant. Since positive numbers have arbitrary roots in \mathbb{R} , we can get rid of some of the actions

$$\{g \in GL(d+2) \mid \eta \circ g = \pm\eta\}/\{\pm 1\},$$

thus this group is doubly covered by an extension of $O(p+1, q+1)$ by $\{\pm 1\}$. If we take the unit component, we get a group that is (potentially) doubly covered by $SO(p+1, q+1)_0 = G$. \square

We therefore see that it is reasonable to study G .

Lemma 5.4 ([4, §VII.2, example 2]). *Together with $K = SO(p+1) \times SO(q+1)$ and the maps*

$$\begin{aligned} \theta: \mathfrak{g} &\rightarrow \mathfrak{g}, & X &\mapsto -X^T = \eta X \eta \\ B: \mathfrak{g} \otimes \mathfrak{g} &\rightarrow \mathbb{R}, & X \otimes Y &\mapsto \text{tr}(XY), \end{aligned}$$

G is a reductive Lie group.

Definition 5.5. For $\mu, \nu, \sigma, \rho = 0, \dots, d+1$ define

$$(E_{\mu\nu})^\sigma_\rho = \delta_\nu^\sigma \eta_{\nu\rho}$$

(so that $(E_\mu^\nu) = \delta_\mu^\sigma \delta_\rho^\nu$) and

$$F_{\mu\nu} := E_{\mu\nu} - E_{\nu\mu}.$$

The set $(F_{\mu\nu})_{0 \leq \mu < \nu \leq d+1}$ is a basis of \mathfrak{g} and a short calculation will convince the reader that

$$[F_{\mu\nu}, F_{\rho\sigma}] = \eta_{\nu\rho} F_{\mu\sigma} + \eta_{\mu\sigma} F_{\nu\rho} - \eta_{\mu\rho} F_{\nu\sigma} - \eta_{\nu\sigma} F_{\mu\rho}.$$

Note in addition that $F_{\mu\nu} = -F_{\nu\mu}$ and that $\theta(F_{\mu\nu}) = F^{\mu\nu}$, which equals $F_{\mu\nu}$ if $\mu, \nu \leq p$ or $p < \mu, \nu$, and which equals $-F_{\mu\nu}$ otherwise.

Proposition 5.6. \mathfrak{k} is spanned by $F_{\mu\nu}$ where μ, ν are either both $\leq p$ or both $> p$ and \mathfrak{p} is spanned by $F_{\mu\nu}$ met $0 \leq \mu \leq p < \nu \leq d+1$. For $a: = 0, \dots, q$ define $D_a := F^{a,d+1-a} = -F_{a,d+1-a}$. Then

$$\mathfrak{a}_\mathfrak{p} := \text{span}\{D_0, \dots, D_q\}$$

is a maximal commutative subspace of \mathfrak{p} .

Proof. By previous observations, the $F_{\mu\nu}$ are contained in $\mathfrak{k}, \mathfrak{p}$, respectively. Since together they form a basis, and since $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, the appropriate $F_{\mu\nu}$ span $\mathfrak{k}, \mathfrak{p}$, respectively.

Note that two $F_{\mu\nu}$ commute automatically if their indices don't collide, so that \mathfrak{a}_p as defined is indeed commutative. Let now

$$X = \sum_{\mu \leq p < \nu} A^{\mu\nu} F_{\mu\nu} \in \mathfrak{p}$$

commute with \mathfrak{a}_p . In particular,

$$\begin{aligned} [D_a, X] &= \sum_{\mu \leq p < \nu} A^{\mu\nu} [F^{a, d+1-a}, F_{\mu\nu}] \\ &= \sum_{\mu \leq p < \nu} A^{\mu\nu} (-\delta_\mu^a F_\nu^{d+1-a} - \delta_\nu^{d+1-a} F_\mu^a) \\ &= - \sum_{\substack{\nu=p+1 \\ \nu \neq d+1-a}}^{d+1} A^{a, \nu} F_\nu^{d+1-a} - \sum_{\substack{\mu=0 \\ \mu \neq a}}^p A^{\mu, d+1-a} F_\mu^a. \end{aligned}$$

The first sum contains elements of $0 \oplus \mathfrak{so}(q+1)$ and the second sum contains elements of $\mathfrak{so}(p+1) \oplus 0$ and are therefore linearly independent. Since this is zero, we obtain $A^{\mu\nu} = 0$ for all $0 \leq \mu \leq p < \nu \leq d+1$ satisfying $\mu + \nu \neq d+1$. Consequently, $X \in \mathfrak{a}_p$. \square

Proposition 5.7. *Define $\epsilon_0, \dots, \epsilon_q \in \mathfrak{a}_p^*$ by $\epsilon_a(D_b) := \delta_{a,b}$. Then \mathfrak{g} has the following restricted root space decomposition with respect to \mathfrak{a}_p :*

$$\mathfrak{g} = \mathfrak{m}_p \oplus \mathfrak{a}_p \oplus \bigoplus_{\alpha \in R} \mathfrak{g}_\alpha$$

where \mathfrak{m}_p is spanned by $F_{\mu\nu}$ for $q < \mu < \nu \leq p$ and

$$R = \{\pm\epsilon_a \mid 0 \leq a \leq q\} \cup \{\epsilon_a \pm \epsilon_b, -\epsilon_a \pm \epsilon_b \mid 0 \leq a < b \leq q\}$$

is its restricted root system which is of type B_{q+1} . The root multiplicities are $p-q-1$ for the short roots and 1 for the long roots.

Proof. From [4, §VI.4] it is known that such a decomposition exists (for R the set of restricted roots) and is direct. It therefore suffices to find the roots and root spaces and \mathfrak{m}_p .

Short roots: Let $0 \leq a, b \leq q$ and $q < \mu \leq p$, then

$$\begin{aligned} [D_a, F^{b, \mu} \pm F^{d+1-b, \mu}] &= -\eta^{a,b} F^{d+1-b, \mu} \pm \eta^{d+1-a, d+1-b} F^{i, \mu} \\ &= \mp \delta_{a,b} (F^{b, \mu} \pm F^{d+1-b, \mu}) \\ &= (\mp \epsilon_b)(D_a)(F^{b, \mu} \pm F^{d+1-b, \mu}), \end{aligned}$$

so that $F^{b, \mu} \pm F^{d+1-b, \mu} \in \mathfrak{g}_{\mp \epsilon_b}$. The root multiplicity of $\pm \epsilon_b$ is therefore at least $p-q$.

Long roots: Let $a, b, c \in [0, q]$ with $i \neq j$, then

$$\begin{aligned} [D_c, F^{a,b} \pm F^{a, d+1-b} + F^{d+1-a, b} \pm F^{d+1-a, d+1-b}] \\ = (-\epsilon_a \mp \epsilon_b)(D_c)(F^{a,b} \pm F^{a, d+1-b} + F^{d+1-a, b} \pm F^{d+1-a, d+1-b}), \end{aligned}$$

so that we found an element of $\mathfrak{g}_{-\epsilon_a \mp \epsilon_b}$. By applying θ , we also find an element of $\mathfrak{g}_{\epsilon_a \pm \epsilon_b}$. The root multiplicity of $\epsilon_a \pm \epsilon_b, -\epsilon_a \mp \epsilon_b$ is therefore at least 1.

Zero spaces: For $q < \mu < \nu \leq p$ the indices of $F_{\mu\nu}$ are disjoint from $(a, d+1-a)$ for $a = 0, \dots, q$, therefore $[D_a, F_{\mu\nu}] = 0$. The dimension of \mathfrak{m}_p is therefore at least $\frac{(p-q)(p-q-1)}{2}$.

We therefore see that the roots from the claim are indeed roots of \mathfrak{g} . To see that we have indeed found all roots and fully described the root spaces, note that

$$\begin{aligned} \frac{(d+2)(d+1)}{2} &= \dim(\mathfrak{g}) = \dim(\mathfrak{m}_{\mathfrak{p}}) + \dim(\mathfrak{a}_{\mathfrak{p}}) + \sum_{\alpha \in R} \dim(\mathfrak{g}_{\alpha}) \\ &\geq \frac{(p-q)(p-q-1)}{2} + (q+1) + (q+1) \cdot 2 \cdot (p-q) + \frac{q(q+1)}{2} \cdot 4 \\ &= \frac{(d-2q)(d-2q-1) + 2(q+1) + 4(q+1)(d-2q) + 4q(q+1)}{2} \\ &= \frac{(d+2)(d+1)}{2}, \end{aligned}$$

so that all inequalities are sharp, and we are done. \square

Lemma 5.8. *G acts transitively on $\widehat{\mathbb{R}^{p,q}}$. Let Q be the stabiliser of $(1:0:1) = \iota(0)$. It is a closed subgroup. In particular $\widehat{\mathbb{R}^{p,q}} \cong G/Q$.*

Proof. Let $q(v) \in \widehat{\mathbb{R}^{p,q}}$ and decompose $v \in \mathbb{R}^{p+1,q+1}$ under $\mathbb{R}^{p+1,q+1} = \mathbb{R}^{p+1} \oplus \mathbb{R}^{q+1}$ as $v = v_1 \oplus v_2$. Since v is a null vector, we have $\|v_1\| = \|v_2\|$ (Euclidean norms). Since $SO(p+1)$ and $SO(q+1)$ work transitively on the respective unit spheres, there are $g \in SO(p+1), g' \in SO(q+1)$ such that

$$ge_0 = \frac{v_1}{\|v_1\|}, \quad g'e_q = \frac{v_2}{\|v_2\|}$$

(where e_0, \dots, e_p are the respective standard unit vectors). Let $h = g \oplus g'$, then $h \in SO(p+1, q+1)_0$ and

$$h(e_0 + e_{d+1}) = \begin{pmatrix} g & 0 \\ 0 & g' \end{pmatrix} \begin{pmatrix} e_0 \\ e_q \end{pmatrix} = \frac{1}{\|v_1\|} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix},$$

so that $h \cdot (1:0:1) = q(v)$ (where $(1:0:1)$ is split up as $1 + d + 1$).

The stabiliser Q is the preimage of the closed set $\{\iota(0)\}$ under the continuous function $G \rightarrow \widehat{\mathbb{R}^{p,q}}, g \mapsto g \cdot \iota(0)$, and hence closed. \square

We are now interested in what the corresponding Lie subalgebra \mathfrak{q} looks like.

Lemma 5.9. *Let*

$$\Gamma = R \setminus \{-\epsilon_0, -\epsilon_0 \pm \epsilon_a \mid 1 \leq a \leq q\},$$

then

$$\mathfrak{q} = \mathfrak{m}_{\mathfrak{p}} \oplus \mathfrak{a}_{\mathfrak{p}} \oplus \bigoplus_{\alpha \in \Gamma} \mathfrak{g}_{\alpha}.$$

Proof. We have $g \in Q$ iff $g(e_0 + e_{d+1}) \in \mathbb{R}(e_0 + e_{d+1})$. In particular, $X \in \mathfrak{q}$ iff the same condition holds. We shall first show $\mathfrak{a}_{\mathfrak{p}} \leq \mathfrak{q}$. Let $1 \leq a \leq q$, then D_a has zeroes in the 0th and $d+1$ -st columns, so that

$$D_a(e_0 + e_{d+1}) = 0,$$

whence $D_a \in \mathfrak{q}$. Lastly,

$$D_0(e_0 + e_{d+1}) = F^{0,d+1}(e_0 + e_{d+1}) = e_0 + e_{d+1},$$

so also $D_0 \in \mathfrak{q}$. Consequently $\mathfrak{a}_{\mathfrak{p}} \leq \mathfrak{q}$.

Now, since \mathfrak{q} is a Lie algebra, the algebra $\text{ad}(\mathfrak{a}_{\mathfrak{p}})$ leaves \mathfrak{q} invariant, so that

$$\mathfrak{q} = (\mathfrak{a}_{\mathfrak{p}} \oplus \mathfrak{m}_{\mathfrak{p}}) \cap \mathfrak{q} \oplus \bigoplus_{\alpha \in R} (\mathfrak{g}_{\alpha} \cap \mathfrak{q}).$$

We just need to see what those intersections are.

Zero: Let $q < \mu < \nu \leq p$, then $F_{\mu\nu}$ has zeroes in the 0th and $d+1$ -st column, hence

$$F_{\mu\nu}(e_0 + e_{d+1}) = 0,$$

so that $F_{\mu\nu} \in \mathfrak{q}$. Consequently, $\mathfrak{m}_p \oplus \mathfrak{a}_p \leq \mathfrak{q}$.
 $\pm\epsilon_0$: For $q < \mu \leq p$ we have

$$(F^{0,\mu} \pm F^{d+1,\mu})(e_0 + e_{d+1}) = (-1 \mp 1)e_\mu,$$

which is a multiple of $e_0 + e_{d+1}$ only in the --case. From Proposition 5.7 we conclude that $\mathfrak{g}_{\epsilon_0} \subseteq \mathfrak{q}$ and $\mathfrak{g}_{-\epsilon_0} \cap \mathfrak{q} = 0$.

Other short roots: For $a = 1, \dots, q$ and $q < \mu \leq p$ the matrix $F^{a,\mu} \pm F^{d+1-a,\mu}$ has zeroes in the 0th and $d+1$ -st columns, and is hence contained in \mathfrak{q} . Consequently, $\mathfrak{g}_{\mp\epsilon_a} \subseteq \mathfrak{q}$.

$\pm\epsilon_0 \pm \epsilon_a$: Let $1 \leq a \leq q$, then

$$\begin{aligned} (F^{0,a} \pm F^{0,d+1-a} + F^{d+1,a} \pm F^{d+1,d+1-a})(e_0 + e_{d+1}) &= -2e_a \pm 2e_{d+1-a} \neq 0 \\ (F^{0,a} \mp F^{0,d+1-a} - F^{d+1,a} \pm F^{d+1,d+1-a})(e_0 + e_{d+1}) &= 0, \end{aligned}$$

so that $\mathfrak{g}_{-\epsilon_0 \mp \epsilon_a} \cap \mathfrak{q} = 0$ and $\mathfrak{g}_{\epsilon_0 \pm \epsilon_a} \subseteq \mathfrak{q}$.

Other long roots: Let $1 \leq a < b \leq q$. The matrices contained in $\mathfrak{g}_{\epsilon_a \pm \epsilon_b}$ and $\mathfrak{g}_{-\epsilon_a \pm \epsilon_b}$ only have nonzero entries in the a -th, b -th, $d+1-a$ -th, $d+1-b$ -th columns, so in particular only zeroes in the 0th and $d+1$ -st column. Consequently, all these weight spaces are contained in \mathfrak{q} .

We found that $\mathfrak{g}_\alpha \cap \mathfrak{q} = \mathfrak{g}_\alpha$ unless $\alpha = -\epsilon_0$ or $\alpha = -\epsilon_0 \pm \epsilon_a$ for $1 \leq a \leq q$, as claimed. \square

We pick the positive subsystem

$$R^+ := \{\epsilon_a \mid 0 \leq a \leq q\} \cup \{\epsilon_a \pm \epsilon_b \mid 0 \leq a < b \leq q\}$$

and the corresponding set S of simple roots:

$$S := \{\epsilon_0 - \epsilon_1, \dots, \epsilon_{q-1} - \epsilon_q, \epsilon_q\}.$$

Let $S' := S \setminus \{\epsilon_0 - \epsilon_1\}$. Note that

$$\Gamma = R^+ \cup (R \cap \text{span}_{\mathbb{R}}(S')).$$

Corollary 5.10. $\mathfrak{q} \leq \mathfrak{g}$ is maximal parabolic with respect to our choices of positive subsystem and \mathfrak{a} .

Proof. \mathfrak{q} is parabolic since the minimal parabolic subalgebra

$$\mathfrak{m}_p \oplus \mathfrak{a}_p \oplus \bigoplus_{\alpha \in R^+} \mathfrak{g}_\alpha$$

is contained therein (recall that $R^+ \subseteq \Gamma$). By [4, Proposition VII.7.76], there is an inclusion-preserving 1-1 correspondence between subsets $S' \subseteq S$ and the parabolic subalgebras containing the minimal parabolic subalgebra that corresponds to our choices of \mathfrak{a}_p, R^+ . A correspondence which associates our S' to \mathfrak{q} . Any Lie algebra lying between \mathfrak{q} and \mathfrak{g} must therefore correspond to a subset lying between S' and S . Since these sets differ by only one element, there is no subset properly between S' and S . Consequently, there are no Lie algebras between \mathfrak{q} and \mathfrak{g} . \square

With this result in hand we shall now make some definitions in the usual parlance of parabolic subalgebras.

Definition 5.11. *We define the subalgebras*

$$\begin{aligned}\mathfrak{a} &:= \mathbb{R}D_0 \\ \mathfrak{a}_M &:= \text{span}_{\mathbb{R}}\{D_1, \dots, D_q\} \\ \mathfrak{m} &:= \text{span}_{\mathbb{R}}\{F_{\mu\nu} \mid 1 \leq \mu < \nu \leq d\} \\ \mathfrak{n} &:= \text{span}_{\mathbb{R}}\{K^i \mid 1 \leq i \leq d\} \\ \mathfrak{n}_M &:= \text{span}_{\mathbb{R}}\{F^{a,\nu} - F^{d+1-a,\nu} \mid 1 \leq a \leq q, 1 \leq \nu \leq d\} \\ \bar{\mathfrak{n}} &:= \text{span}_{\mathbb{R}}\{P^i \mid 1 \leq i \leq d\}\end{aligned}$$

where

$$K^i := F^{0,i} - F^{d+1,\mu}, \quad P^i := -F^{0,i} - F^{d+1,i}$$

($i = 1, \dots, d$). Let furthermore A, A_M, N, N_M be the analytic subgroups of $\mathfrak{a}, \mathfrak{a}_M, \mathfrak{n}, \mathfrak{n}_M$ and let $M := {}^0Z_G(\mathfrak{a})$ (notation from e.g. [4, §VII.7.2]).

Lemma 5.12. $Q = MAN$, which is the parabolic subgroup associated to \mathfrak{q} .

Proof. “ \subseteq ”: By [4, Proposition VII.7.83(b)], $MAN = N_G(\mathfrak{q})$, so it suffices to show that every $g \in Q$ normalises \mathfrak{q} . Let $X \in \mathfrak{q}$, then

$$\exp(t \text{Ad}(g)(X))\iota(0) = g \exp(tX)g^{-1}\iota(0) = \iota(0),$$

for all $t \in \mathbb{R}$. Taking the derivative at $t = 0$, we see that $\text{Ad}(g)(X) \in \mathfrak{q}$ as well.

“ \supseteq ”: Since $\mathfrak{a}, \mathfrak{n}$ fix $\iota(0)$, so do their analytic subgroups, i.e. $A, N \leq Q$. By [4, Proposition VII.7.82(c)] we have $M = (M \cap K)A_M N_M$. Now, both $\mathfrak{a}_M, \mathfrak{n}_M$ fix $\iota(0)$, hence so do their analytic subgroups. It therefore remains to check if $K \cap M \leq Q$. Assume

$$g = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \in K \cap M$$

(written as a $p+1$ -block and a $q+1$ -block). Then $\text{Ad}(g)(D_0) = 0$. If we block decompose D_0 this means,

$$\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \begin{pmatrix} 0 & e_1 e_{q+1}^T \\ e_{q+1} e_1^T & 0 \end{pmatrix} \begin{pmatrix} A^T & 0 \\ 0 & B^T \end{pmatrix} = \begin{pmatrix} 0 & Ae_1 (Be_{q+1})^T \\ Be_{q+1} (Ae_1)^T & 0 \end{pmatrix},$$

i.e. $Ae_1 (Be_{q+1}) = e_1 e_{q+1}^T$. I.e. there exists $\lambda \in \mathbb{R}$ such that $Ae_1 = \lambda e_1$ and $Be_{q+1} = \lambda^{-1} e_{q+1}$. Since ± 1 are the only two possible real eigenvalues an orthogonal matrix could have, we have $\lambda = \lambda^{-1}$. Consequently,

$$g\iota(0) = q \left(\begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix} \begin{pmatrix} e_1 \\ e_{q+1} \end{pmatrix} \right) = q(e_1 e_{q+1}) = \iota(0).$$

Thus $g \in Q$. Consequently, $K \cap M \leq Q$. \square

Lastly, for future reference, we shall also establish what exactly the group M looks like

Lemma 5.13. *For every $A \in SO(p, q)_0$, there exists $g \in M$ such that*

$$\forall x \in \mathbb{R}^{p,q}: \quad g \cdot \iota(x) = \iota(Ax).$$

Proof. Since $M \leq G$ is a closed Lie subgroup with Lie algebra \mathfrak{m} (which is isomorphic to $\mathfrak{so}(p, q)$), it contains the corresponding analytic subgroup, which is isomorphic to $SO(p, q)_0$. Let $g \in SO(p, q)_0$, the corresponding element of G is

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & g & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

which transforms $\iota(x)$ into $\iota(gx)$. \square

5.3. Point Configurations. When considering the action of G not on a single point of $\widehat{\mathbb{R}^{p,q}}$ but on tuples, we quickly run into singular strata, i.e. “thin” sets of orbits that prevent $\widehat{\mathbb{R}^{p,q}}/G$ from becoming a manifold. Part of these can be eliminated by considering $\text{Conf}(\widehat{\mathbb{R}^{p,q}}, n)$, the *configuration space of n points* in $\widehat{\mathbb{R}^{p,q}}$, which consists of tuples of *distinct* points. However, since our action is one by means of projective transformations and not all homeomorphisms, this is not sufficient and we have to resort to terminology from projective geometry.

Definition 5.14. *In projective geometry, a tuple $(q(v_1), \dots, q(v_m)) \in (\mathbb{R}\mathbb{P}^n)^m$ is said to be in general position if the vectors $v_1, \dots, v_m \in \mathbb{R}^{n+1}$ are linearly independent. (Here, $q: \mathbb{R}^{n+1} \setminus \{0\} \rightarrow \mathbb{R}\mathbb{P}^n$ is the projectivisation map.)*

Since our transformation group is not all of $PGL(n)$ either, but involves the inner product, we have to add even more conditions to obtain our smooth stratum.

Definition 5.15. *For the purposes of this paper, a tuple $(q(v_1), \dots, q(v_m)) \in (\widehat{\mathbb{R}^{p,q}})^m$ is said to be in general position if the vectors $v_1, \dots, v_m \in \mathbb{R}^{d+2}$ are linearly independent and no pair of them is orthogonal. Write $\text{GP}(\widehat{\mathbb{R}^{p,q}}, m) = \text{GP}(G/Q, m)$ for the set of all m -tuples of points in general position. It is an open dense subset as the map $(v_1, \dots, v_m) \mapsto (q(v_1), \dots, q(v_m))$ is a quotient map and since the conditions of being linearly independent and being pairwise non-orthogonal with respect to η are Zariski-open (and hence define a dense open subset).*

Another interpretation of the non-orthogonality condition can be found if we consider two embedded points from $\widehat{\mathbb{R}^{p,q}}$ and check if they satisfy the condition to be in general position.

Lemma 5.16. *Let $x, y \in \widehat{\mathbb{R}^{p,q}}$. Then $(\iota(x), \iota(y))$ is in general position iff x and y are not light-like separated or coincide, i.e. iff $\eta(x - y, x - y) \neq 0$.*

Proof. We have $\iota(x) = q(v)$ and $\iota(y) = q(w)$ for

$$v = \begin{pmatrix} 1 - \eta(x, x) \\ 2x \\ 1 + \eta(x, x) \end{pmatrix}, \quad w = \begin{pmatrix} 1 - \eta(y, y) \\ 2y \\ 1 + \eta(y, y) \end{pmatrix}.$$

“ \Rightarrow ”: Let $(\iota(x), \iota(y))$ be in general position, Note that

$$\begin{aligned} \eta(v, w) &= (1 - \eta(x, x))(1 - \eta(y, y)) + 4\eta(x, y) - (1 - \eta(x, x))(1 - \eta(y, y)) \\ &= 4\eta(x, y) - 2\eta(x, x) - 2\eta(y, y) \\ &= -2\eta(x - y, x - y). \end{aligned}$$

Since we know that $\eta(v, w) \neq 0$, we conclude that $\eta(x - y, x - y) \neq 0$, so that x, y are not light-like separated.

“ \Leftarrow ”: Since x, y are not light-like separated, the above calculation implies that $\eta(v, w) \neq 0$. Furthermore, we know $x \neq y$, so that $\iota(x) \neq \iota(y)$ and in particular, v, w are not multiples of each other. Since they are also both nonzero, they are linearly independent and consequently, $(q(v), q(w))$ are in general position. \square

Lemma 5.17. *G 's action on $\widehat{\mathbb{R}^{p,q}}$ extends naturally to $\text{GP}(\widehat{\mathbb{R}^{p,q}}, n)$.*

Proof. Let $g \in G$ and $(q(v_1), \dots, q(v_n))$ be in general position. Since $g \in GL(d+2)$, the vectors gv_1, \dots, gv_n are also linearly independent. Furthermore, for any two $i, j \in \{1, \dots, n\}$ we have

$$\eta(gv_i, gv_j) = \eta(v_i, v_j) \neq 0.$$

So $(q(gv_1), \dots, q(gv_n)) = (g \cdot q(v_1), \dots, g \cdot q(v_n))$ is again in general position. \square

Lemma 5.18. *The action of G on $\mathrm{GP}(G/Q, 2)$ is transitive with MA being a typical stabiliser.*

Proof. Let $(x, y) \in \mathrm{GP}(G/Q, 2)$ and pick $g \in G$ such that $g \cdot y = \infty = (1:0:-1)$. Write $g \cdot x = :q(v)$ with say $v = (v_0, \underline{v}, v_{d+1})$ (split as $1 + d + 1$). Since $(g \cdot x, g \cdot y)$ are in general position, we have

$$0 \neq \eta(v, (1, 0, -1)^T) = v_0 + v_{d+1}.$$

Recalling the proof of Lemma 5.2, we can thus conclude that $g \cdot x = \iota(u)$ where $u := \frac{\underline{v}}{v_0 + v_{d+1}}$. If we let the matrix u^\bullet equal the column vector u and correspondingly let u_\bullet be the diagonal matrix η multiplied with u (standard index raising/lowering notation), we obtain

$$\exp(-u \cdot P) = \exp\left(-\sum_{i=1}^d u_i P^i\right) = \begin{pmatrix} 1 - \frac{u^2}{2} & u_\bullet^T & -\frac{u^2}{2} \\ -u_\bullet & 1 & -u_\bullet \\ \frac{u^2}{2} & -u_\bullet^T & 1 + \frac{u^2}{2} \end{pmatrix}$$

($u^2 = \eta(u, u)$). Then we have

$$\begin{aligned} \exp(-u \cdot P) \cdot q(v) &= q\begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = \iota(0) \\ \exp(-u \cdot P) \cdot \infty &= q\begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} = \infty. \end{aligned}$$

Thus we have found an element $\exp(-u \cdot P)g \in G$ that maps (x, y) to $(\iota(0), \infty)$. This shows transitivity.

For a typical stabiliser we consider the point $(\iota(0), \infty)$ and let $H \leq G$ be its stabiliser. Let $w \in G$ be an element satisfying $g \cdot \iota(0) = \infty$. Then

$$H = \{g \in G \mid g \in Q, w^{-1}gw \in G\} = Q \cap wQw^{-1}.$$

In particular, because of what $\iota(0)$ and ∞ look like, we can choose H diagonal with its diagonal entries being ± 1 , say $w = \mathrm{diag}(w_0, \dots, w_{d+1})$ with $w_0 = -w_{d+1}$. Then $w^2 = 1$ and $w \in K$.

As a consequence, we have $\mathrm{Ad}(w)(D_0) = -D_0$ and $\mathrm{Ad}(w)(D_a) = \pm D_a$ for $a = 1, \dots, q$. This shows that $\mathrm{Ad}^*(w)(\Gamma) = -\Gamma$ and thus $wNw^{-1} = \overline{N}$. Furthermore, by [4, Proposition VII.7.82(a)], MA is the centraliser of \mathfrak{a} in G , thus if $g \in MA$ we have

$$\mathrm{Ad}(wgw^{-1})(D_0) = -\mathrm{Ad}(wg)(D_0) = -\mathrm{Ad}(w)(D_0) = D_0,$$

so that $wgw^{-1} \in MA$ as well, hence $wMAw^{-1} = MA$. Consequently, we have

$$Q \cap wQw^{-1} = MAN \cap wMAw^{-1}wNw^{-1} = MAN \cap M\overline{N}.$$

Let $g \in H$, say $g = man = m'a'\overline{n}'$, then $\overline{n}' = a'^{-1}m'^{-1}man \in \overline{N} \cap Q$, which is trivial by [4, Proposition VII.7.83(e)]. Consequently, $g \in MA$. \square

We can thus conclude that $\mathrm{GP}(G/Q, 2) \cong G/MA$ as smooth G -sets.

Lemma 5.19. *Let $n \geq 2$ and $(x_1, \dots, x_n) \in \mathrm{GP}(G/Q, n)$. Then there exists $g \in G$ such that*

$$g \cdot x_1 = \iota(0), \quad g \cdot x_2 = \infty, \quad g \cdot x_i = \iota(p_i) \quad (i = 3, \dots, n).$$

where $p_3, \dots, p_n \in \mathbb{R}^{p \cdot q}$ are linearly independent and none of p_i ($i = 3, \dots, n$) and $p_i - p_j$ ($i \neq j \in \{3, \dots, n\}$) is isotropic (i.e. light-like).

Proof. Write $x_i = q(v_i)$ for $v_1, \dots, v_n \in \mathbb{R}^{d+2} \setminus \{0\}$. We have $(x_1, x_2) \in \text{GP}(G/Q, 2)$, hence by Lemma 5.18, there exists $g \in G$ mapping (x_1, x_2) to $(\iota(0), \infty)$. Then since none of gv_3, \dots, gv_n is orthogonal to $gv_2 = (1, 0, -1)^T$, we can again employ the same argument as in the transitivity proof of Lemma 5.18, so that we find p_3, \dots, p_n such that $g \cdot x_i = \iota(p_i)$ ($i = 3, \dots, n$). By definition, the vectors

$$\begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, \begin{pmatrix} 1 - p_3^2 \\ 2p_3^\bullet \\ 1 + p_3^2 \end{pmatrix}, \dots, \begin{pmatrix} 1 - p_n^2 \\ 2p_n^\bullet \\ 1 + p_n^2 \end{pmatrix}$$

are linearly independent, which implies that p_3, \dots, p_n are linearly independent. Furthermore, by applying Lemma 5.16 to $(g \cdot x_i, g \cdot x_j)$ for $i, j \in \{1, 3, \dots, n\}$ in turn, we obtain that none of the vectors $p_i - p_j$ ($i \neq j \in \{3, \dots, n\}$) and $p_i - 0$ ($i = 3, \dots, n$) is isotropic. \square

For the rest of this paper we shall be concerned with four-point configurations, i.e. with $n = 4$.

Definition 5.20. Define $u, v: \text{GP}(G/Q, 4) \rightarrow \mathbb{R}$ by

$$(5.1) \quad u(q(v_1), \dots, q(v_4)) := \frac{\eta(v_1, v_2)\eta(v_3, v_4)}{\eta(v_1, v_3)\eta(v_2, v_4)}$$

$$(5.2) \quad v(q(v_1), \dots, q(v_4)) := \frac{\eta(v_1, v_4)\eta(v_2, v_3)}{\eta(v_1, v_3)\eta(v_2, v_4)}.$$

They are both well-defined, smooth, G -invariant and for the parameters $\iota(x_1), \dots, \iota(x_4)$ they reduce to the well-known expressions for the cross-ratios from e.g. [18, §III.C.3].

Proof. Firstly, note that both right-hand sides are homogeneous in v_1, \dots, v_4 of degree $(0, 0, 0, 0)$, and by definition neither $\eta(v_1, v_3)$ nor $\eta(v_2, v_4)$ are zero. Thus, the functions are well-defined and continuous. Since q is a smooth quotient map and our definition of u, v in terms of v_1, \dots, v_4 is also smooth, u, v are smooth maps on $\text{GP}(G/Q, 4)$. Since G leaves η invariant, we have $u(q(gv_1), \dots, q(gv_4)) = u(q(v_1), \dots, q(v_4))$ (same for v), so that u, v are G -invariant.

To see that our definition coincides with the usual one, recall that as shown in the proof of Lemma 5.16, for $q(v_i) = \iota(p_i)$ (for the usual choice of v_i) we have $\eta(v_i, v_j) = -2(p_i - p_j)^2$, so that

$$u(\iota(p_1), \dots, \iota(p_4)) = \frac{(p_1 - p_2)^2(p_3 - p_4)^2}{(p_1 - p_3)^2(p_2 - p_4)^2}$$

and similarly for v . \square

Corollary 5.21. Let (x_1, \dots, x_4) be in the same G -orbit as $(\iota(0), \infty, \iota(x), \iota(y))$ as in Lemma 5.19, then

$$u(x_1, \dots, x_4) = \frac{(x - y)^2}{x^2} \quad v(x_1, \dots, x_4) = \frac{y^2}{x^2}.$$

Lemma 5.22. The map $\psi: G \rightarrow \text{GP}(G/Q, 2)^{\times 2}$ mapping

$$g \mapsto (\iota(0), \infty, g\iota(0), g\infty) \cong (Q, wQ, gQ, gwQ)$$

($w \in K$ as in the proof for the stabiliser in Lemma 5.18) descends, when composed with the quotient map to $\text{GP}(G/Q, 2)^{\times 2}/G$, to a homeomorphism $MA \backslash G/MA \cong \text{GP}(G/Q, 2)^{\times 2}/G$.

Proof. Define

$$f: \text{GP}(G/Q, 2)^{\times 2}/G \rightarrow MA \backslash G/MA,$$

$$G(\iota(0), \infty, \iota(x), \iota(y)) \mapsto MA \exp(x \cdot P) \exp\left(\frac{y - x}{(y - x)^2} \cdot K\right) MA.$$

This is well-defined since every element of $\mathrm{GP}(G/Q, 2)^{\times 2}/G$ can be written in the form $G(\iota(0), \infty, \iota(x), \iota(y))$ and since any other choice of x, y would be related by MA , say there is $m \in MA$ and $\alpha \in \mathbb{R}$ such that $m \exp(\alpha D_0) \cdot \iota(x) = \iota(x')$ and the same for y, y' . Then

$$\mathrm{Ad}(m \exp(\alpha D_0))(x \cdot P) = x' \cdot P$$

and

$$\mathrm{Ad}(m \exp(\alpha D_0)) \frac{y-x}{(y-x)^2} \cdot K = \frac{y'-x'}{(y'-x')^2} \cdot K$$

due to $\exp(\alpha D_0)$ acting on $\iota(x)$ and the P^i as scaling by $\exp(-\alpha)$, and on the K^i as scaling by $\exp(\alpha)$. Consequently,

$$\begin{aligned} & MA \exp(x' \cdot P) \exp\left(\frac{y'-x'}{(y'-x')^2} \cdot K\right) MA \\ &= MA m \exp(\alpha D_0) \exp(x \cdot P) \exp\left(\frac{y-x}{(y-x)^2} \cdot K\right) m^{-1} \exp(-\alpha D_0) MA \\ &= MA \exp(x \cdot P) \exp\left(\frac{y-x}{(y-x)^2} \cdot K\right) MA. \end{aligned}$$

The map f is continuous by definition.

Write $\tilde{\Psi}: G \rightarrow \mathrm{GP}(G/Q, 2)^{\times 2}/G$ for the composition of ψ with the quotient map. Then $\tilde{\Psi}$ is MA -biinvariant:

$$\begin{aligned} \tilde{\Psi}(magm'a') &= G(\iota(0), \infty, magm'a' \cdot \iota(0), magm'a' \cdot \infty) \\ &= G(a^{-1}m^{-1}\iota(0), a^{-1}m^{-1} \cdot \infty, gm'a' \cdot \iota(0), gm'a' \cdot \infty) \end{aligned}$$

Since MA fixes $\iota(0), \infty$, this equals $\tilde{\Psi}(g)$. It therefore descends to a continuous map $\Psi: MA \backslash G / MA \rightarrow \mathrm{GP}(G/Q, 2)^{\times 2}/G$.

We shall now show that Ψ and f are inverses of each other. For this we consider

$$\begin{aligned} \Psi(f(G(\iota(0), \infty, \iota(x), \iota(y)))) &= \Psi\left(MA \exp(x \cdot P) \exp\left(\frac{y-x}{(y-x)^2} \cdot K\right) MA\right) \\ &= G(\iota(0), \infty, \iota(x), \iota(y)). \end{aligned}$$

Here we used that for $b := \frac{y-x}{(y-x)^2}$ we can compute the action of $g = \exp(x \cdot P) \exp(b \cdot K)$ on $\iota(0), \infty$ as follows:

$$\begin{aligned} g \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} &= \begin{pmatrix} 1 - \frac{x^2}{2} & -x^T & -\frac{x^2}{2} \\ x^\bullet & 1 & x^\bullet \\ \frac{x^2}{2} & x^T & 1 + \frac{x^2}{2} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 - x^2 \\ 2x \\ 1 + x^2 \end{pmatrix} \\ g \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} &= \begin{pmatrix} 1 - \frac{x^2}{2} & -x^T & -\frac{x^2}{2} \\ x^\bullet & 1 & x^\bullet \\ \frac{x^2}{2} & x^T & 1 + \frac{x^2}{2} \end{pmatrix} \begin{pmatrix} 1 - \frac{b^2}{2} & b^\bullet & \frac{b^2}{2} \\ -b^\bullet & 1 & b^\bullet \\ -\frac{b^2}{2} & b^T & 1 + \frac{b^2}{2} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} \\ &= \begin{pmatrix} 1 - b^2 + b^2 x^2 \\ -2(b + b^2 x) \\ -1 - b^2 - b^2 x^2 \end{pmatrix} \end{aligned}$$

the former of which projectivises to $\iota(x)$ and the latter to $\iota(y)$.

Let now $g \in G$ and let $MAhMA = f(\Psi(MAgMA))$, then $\Psi(MAgMA) = \Psi(f(\Psi(MAgMA)))$, i.e.

$$G(\iota(0), \infty, g \cdot \iota(0), g \cdot \infty) = G(\iota(0), \infty, h \cdot \iota(0), h \cdot \infty),$$

then there is $k \in G$ such that

$$k \cdot \iota(0) = \iota(0), \quad k \cdot \infty = \infty, \quad kg \cdot \iota(0) = h \cdot \iota(0), \quad kg \cdot \infty = h \cdot \infty.$$

This shows that both k and $h^{-1}kg$ fix $\iota(0), \infty$. By Lemma 5.18, this shows that $k, h^{-1}kg \in MA$. In other words:

$$g = k^{-1}h(h^{-1}kg) \in MAhMA.$$

Consequently, $f \circ \Psi$ is also the identity. \square

Using this homeomorphism, we can identify $\mathrm{GP}(G/Q, 4)/G$ with a dense open subset of $MA \backslash G/MA$, say $MA \backslash \tilde{G}/MA$ where $\tilde{G} \subseteq G$ is open, dense, and satisfies $MA\tilde{G}MA \subseteq \tilde{G}$ (more on this in Corollary 6.19).

5.4. 4-Point Functions as MSF.

Definition 5.23. Let (V_i, π_i) ($i = 1, \dots, 4$) be finite-dimensional Q -modules. A smooth function $f: G^{\times 4} \rightarrow V_1 \otimes \dots \otimes V_4$ is said to satisfy the Ward identities, if for all $g, g_1, \dots, g_4 \in G$ and $q_1, \dots, q_4 \in Q$ we have

$$f(gg_1q_1, \dots, gg_4q_4) = \pi_1(q_1)^{-1} \otimes \dots \otimes \pi_4(q_4)^{-1} f(g_1, \dots, g_4),$$

i.e. if f is a $G \times Q^{\times 4}$ -intertwiner for the following $G \times Q^{\times 4}$ -actions:

$$\text{on } G^{\times 4}: (g, q_1, \dots, q_4) \cdot (g_1, \dots, g_4) := (gg_1q_1^{-1}, \dots, gg_4q_4^{-1})$$

$$\text{on } V_1 \otimes \dots \otimes V_4: (g, q_1, \dots, q_4) \cdot (v_1 \otimes \dots \otimes v_4) := (q_1 \cdot v_1 \otimes \dots \otimes q_4 \cdot v_4).$$

Let $U \subseteq G^{\times 4}$ (also open and dense) be the preimage of $\mathrm{GP}(G/Q, 4)$ under the quotient map $G^{\times 4} \rightarrow (G/Q)^{\times 4}$. It is invariant under the action of $G \times Q^{\times 4}$, thus we can define an analogous notion of satisfying the Ward identities for functions defined on U .³

Fix an element $w \in G$ that maps $\iota(0)$ to ∞ and vice-versa. For simplicity, we can choose w to be a diagonal matrix that squares to 1. Then conjugation with w is an automorphism of MA (which inverts elements of A).

Theorem 5.24. Let $f: U \rightarrow V_1 \otimes \dots \otimes V_4$ solve the Ward identities. Write $W := V_1 \otimes \dots \otimes V_4$ and define $F: \tilde{G} \rightarrow W$ as $g \mapsto f(1, w, g, gw)$. Then F is a matrix-spherical function for the symmetric subgroup MA and the following left and right actions:

$$(ma) \cdot (v_1 \otimes \dots \otimes v_4) := (ma \cdot v_1) \otimes (wma \cdot v_2) \otimes v_3 \otimes v_4$$

$$(v_1 \otimes \dots \otimes v_4) \cdot (ma) := v_1 \otimes v_2 \otimes (m^{-1}a^{-1} \cdot v_3) \otimes (wm^{-1}a^{-1}w \cdot v_4).$$

The map $f \mapsto F$ is injective.

Proof. We begin by checking the biequivariance. Let $ma, m'a' \in MA$ and $g \in \tilde{G}$, then

$$\begin{aligned} F(magm'a') &= f(1, w, magm'a', magm'a'w) = f(a^{-1}m^{-1}, a^{-1}m^{-1}w, gm'a', gm'a'w) \\ &= \pi_1(ma) \otimes \pi_2(wma) \otimes \pi_3(m'a')^{-1} \otimes \pi_4(wm'a'w)^{-1} f(1, w, g, gw) \\ &= ma \cdot F(g) \cdot m'a'. \end{aligned}$$

For injectivity let F and F' arise from $f, f': U \rightarrow V_1 \otimes \dots \otimes V_4$ and $F = F'$. Let $(g_1, \dots, g_4) \in U$, then $(g_1Q, \dots, g_4Q) \in \mathrm{GP}(G/Q, 4)$. Then by Lemma 5.22 and definition of \tilde{G} , there is $h \in \tilde{G}$ such that

$$G\psi(h) = G(Q, wQ, hQ, hwQ) = G(g_1Q, g_2Q, g_3Q, g_4Q).$$

This implies that there are $g \in G$ and $q_1, \dots, q_4 \in Q$ such that

$$(1, w, h, hw) = (gg_1q_1, \dots, gg_4q_4).$$

³This definition can also be naturally put in the framework of generalised spherical functions associated to moduli spaces of principal connections on a corresponding star-shaped graph, see [19]. We thank Jasper Stokman for explaining this point of view to us.

Then

$$\begin{aligned}
f(g_1, \dots, g_4) &= f(g^{-1}q_1^{-1}, g^{-1}wq_2^{-1}, g^{-1}hq_3^{-1}, g^{-1}hwq_4^{-1}) \\
&= \pi_1(q_1) \cdots \pi_4(q_4) f(1, w, h, hw) \\
&= \pi_1(q_1) \cdots \pi_4(q_4) F(h) = \pi_1(q_1) \cdots \pi_4(q_4) F'(h) \\
&= f'(g_1, \dots, g_4).
\end{aligned}$$

Lastly, we need to show that $MA \leq G$ is symmetrising subgroup. Define $\sigma: G \rightarrow G$ be conjugation with the matrix $\text{diag}(-1, 1, \dots, 1, -1)$, then G^σ consists of the elements of σ that can be written as a block matrix of the following shape:

$$\begin{pmatrix} a & 0 & b \\ 0 & c & 0 \\ d & 0 & e \end{pmatrix}$$

for $a, b, d, e \in \mathbb{R}, c \in \mathbb{R}^{d \times d}$. Every element of MA can be written this way, consequently, $MA \leq G^\sigma$. Furthermore, the unit component of G^σ is the analytic subgroup for the subalgebra $\mathfrak{m} \oplus \mathfrak{a}$, which is contained in MA since MA has Lie algebra $\mathfrak{m} \oplus \mathfrak{a}$. Consequently, $(G^\sigma)_0 \leq MA \leq G^\sigma$. \square

Using the injection from Theorem 5.24, we can view solutions to the Ward identities as matrix-spherical functions for (G, MA) . Since we can no longer assume that these functions are defined on all G , we shall write $E^W(\tilde{G}, MA)$.

Lemma 5.25. *The set of functions $f: U \rightarrow V_1 \otimes \cdots \otimes V_4$ satisfying the Ward equations carries an action of $Z(U(\mathfrak{g}))$ that the injection from Theorem 5.24 intertwines with the right action described in Lemma 2.2.*

Proof. Let X be the set of functions $f: U \rightarrow V_1 \otimes \cdots \otimes V_4$ satisfying

$$\forall q_1, \dots, q_4 \in Q: f(g_1 q_1, \dots, g_4 q_4) = \pi_1(q_1)^{-1} \cdots \pi_4(q_4)^{-1} f(g_1, \dots, g_4).$$

X is acted upon from the right by G and by four copies of \mathfrak{g} :

$$(f \cdot h)(g_1, \dots, g_4) = f(hg_1, \dots, hg_4)$$

and the corresponding infinitesimal actions for each of the four inputs. This gives rise to an action of $(U(\mathfrak{g})^{\otimes 4})^G$ (diagonal action of G) on G -invariant of X , i.e. on solutions to the Ward identities. An algebra contained therein is $1 \otimes 1 \otimes \Delta(Z(U(\mathfrak{g})))$ (where Δ is the comultiplication).

Write Ψ for the injection from Theorem 5.24 (extended to all of X using the same formula). To see that Ψ intertwines these actions of $Z(U(\mathfrak{g}))$, note that for $X \in \mathfrak{g}$ and f satisfying the Ward identities we have

$$\begin{aligned}
\Psi(f \cdot X)(g) &= (f \cdot X)(1, w, g, gw) \\
&= (f \cdot (1 \otimes 1 \otimes X \otimes 1 + 1 \otimes 1 \otimes 1 \otimes X))(1, w, g, gw) \\
&= \frac{d}{dt} (f(1, w, \exp(tX)g, gw) + f(1, w, g, \exp(tX)gw)) \Big|_{t=0} \\
&= \frac{d}{dt} f(1, w, \exp(tX)g, \exp(tX)gw) \Big|_{t=0} \\
&= \frac{d}{dt} \Psi(f)(\exp(tX)g) \Big|_{t=0} \\
&= \Psi(f) \cdot X.
\end{aligned}$$

\square

Definition 5.26. *Let $\chi: Z(U(\mathfrak{g})) \rightarrow \mathbb{C}$ be a central character of \mathfrak{g} . A smooth function $f: U \rightarrow V_1 \otimes \cdots \otimes V_4$ (where V_1, \dots, V_4 are finite-dimensional Q -modules)*

or more generally $f \in E^W(\tilde{G}, MA)$ (for any finite-dimensional MA -bimodule W) is said to be a conformal block for χ if

$$\forall z \in Z(U(\mathfrak{g})): f \cdot z = \chi(z)f.^4$$

Conformal blocks are the fundamental building blocks used to decompose and put constraints on the correlation functions in conformal field theory. In order to find them, it is necessary to know what the above eigenvalue equation looks like for functions f satisfying the Ward identities, or more generally for (G, MA) -matrix-spherical functions. Since (G, MA) is a symmetric pair and G is a reductive Lie group, we can now employ the theory established in the first half of this paper.

6. OBTAINING THE CASIMIR EQUATION

In this section we shall focus on the eigenvalue equation for the quadratic Casimir element $\Omega_{\mathfrak{g}}$. In order to apply the results from Section 4, we first need to re-examine the structure of G . This time we will focus more on Cartan subsets and more generally, on the role that the involution σ plays for G and \mathfrak{g} .

6.1. Cartan Subsets $(C_i)_{i \in I}$ of G . Our first goal here will be choosing a fundamental Cartan subset C . Let us first describe the decomposition of \mathfrak{g} with respect to θ, σ . For that we will need to introduce a bit more notation.

Lemma 6.1. *We have $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p} = \mathfrak{h} \oplus \mathfrak{g}^{-\sigma} = \mathfrak{k}^{\sigma} \oplus \mathfrak{k}^{-\sigma} \oplus \mathfrak{p}^{\sigma} \oplus \mathfrak{p}^{-\sigma}$ with*

$$\begin{aligned} \mathfrak{k}^{\sigma} &= \text{span}\{F_{ij} \mid 1 \leq i, j \leq p \text{ or } p < i, j \leq d\} \\ \mathfrak{k}^{-\sigma} &= \text{span}\{F_{0,i}, F_{j,d+1} \mid 1 \leq i \leq p < j \leq d\} \\ \mathfrak{p}^{\sigma} &= \mathfrak{a} \oplus \text{span}\{F_{i,j} \mid 1 \leq i \leq p < j \leq d\} \\ \mathfrak{p}^{-\sigma} &= \text{span}\{F_{0,j}, F_{i,d+1} \mid 1 \leq i \leq p < j \leq d\}. \end{aligned}$$

Here $\mathfrak{h} = \mathfrak{g}^{\sigma} = \mathfrak{k}^{\sigma} \oplus \mathfrak{p}^{\sigma} = \mathfrak{m} \oplus \mathfrak{a}$ in previous notation.

Proof. The vector $F_{\mu,\nu}$ is σ -invariant if none or both of μ, ν are $0, d+1$, and σ -antiinvariant if exactly one index is $0, d+1$. Similarly, $F_{\mu,\nu}$ is θ -invariant if none or both of μ, ν are $> p$ and θ -antiinvariant if exactly one index is $> p$. This shows “ \supseteq ” for all four claimed equations. For “ \subseteq ” we recall that the sum

$$\mathfrak{g} = \mathfrak{k}^{\sigma} \oplus \mathfrak{k}^{-\sigma} \oplus \mathfrak{p}^{\sigma} \oplus \mathfrak{p}^{-\sigma}$$

is direct and that the spaces that make up the right-hand sides add up to \mathfrak{g} . \square

Definition 6.2. *If $q = 0$, define*

$$\mathfrak{t} := \mathbb{R}F_{01}, \quad \mathfrak{a} := \mathbb{R}F_{d,d+1};$$

if $q > 0$, define

$$\mathfrak{t} := \text{span}\{F_{0,1}, F_{d,d+1}\}, \quad \mathfrak{a} := 0.$$

Let $\mathfrak{c} := \mathfrak{t} \oplus \mathfrak{a}$.

Lemma 6.3. *$\mathfrak{t} \subseteq \mathfrak{k}^{-\sigma}$ is maximally commutative as is $\mathfrak{c} \subseteq \mathfrak{g}^{-\sigma}$.*

⁴In physics literature, one usually distinguishes between conformal blocks and conformal partial waves, depending on the boundary conditions and monodromy properties that one imposes on the (linear combinations of) eigenfunctions of the considered invariant differential operators. Since we do not discuss the solution theory in this paper, we also don’t distinguish between the two in our terminology.

Proof. For the first claim, we distinguish between $q = 0, q > 0$. If $q = 0$, suppose $X \in \mathfrak{k}^{-\sigma}$ commutes with $F_{0,1}$, say

$$X = \sum_{i=1}^d a_i F_{0,i}$$

(note that there is no $p < j \leq d$). Then

$$0 = [F_{0,1}, X] = - \sum_{i=1}^d a_i F_{1,i} = - \sum_{i=2}^d a_i F_{1,i}.$$

Since all the $F_{1,i}$ appearing in the last sum are linearly independent, we have $a_i = 0$ for $i = 2, \dots, d$, showing that $X = a_1 F_{0,1}$. This shows that $\mathfrak{t} \subseteq \mathfrak{k}^{-\sigma}$ is maximally commutative.

The $q > 0$ case is covered by the second claim, so we will show that one instead. Let $X \in \mathfrak{g}^{-\sigma}$ commute with $F_{0,1}, F_{d,d+1}$, say

$$X = \sum_{i=1}^d (a_i F_{0,i} + b_i F_{i,d+1}).$$

Then we have

$$\begin{aligned} 0 &= [F_{0,1}, X] = \sum_{i=1}^d (-a_i F_{1,i} + \delta_{i,1} b_i F_{0,d+1}) \\ &= b_1 F_{0,d+1} - \sum_{i=2}^d a_i F_{1,i} \\ &= [F_{d,d+1}, X] = \sum_{i=1}^d (a_i \delta_{i,d} F_{0,d+1} - b_i F_{i,d}) \\ &= a_d F_{0,d+1} - \sum_{i=1}^{d-1} b_i F_{i,d}. \end{aligned}$$

Since all $F_{\mu,\nu}$ appearing in the 2nd and 4th line are linearly independent, we conclude that all coefficients except for a_1 and b_d are zero, whence $X \in \mathfrak{t}$. \square

Definition 6.4. Write $C := \exp(\mathfrak{c})$ for the corresponding Cartan subset and $T := \exp(\mathfrak{t})$ for its compact torus. The elements of T are denoted by

$$t_\phi := \exp(\phi F_{0,1}) = \begin{pmatrix} \cos(\phi) & \sin(\phi) & 0 \\ -\sin(\phi) & \cos(\phi) & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

(written as $1 + 1 + d$ -block matrix) for $q = 0$ and

$$t_{\phi,\psi} := \exp(\phi F_{0,1} + \psi F_{d,d+1}) = \begin{pmatrix} \cos(\phi) & \sin(\phi) & 0 & 0 & 0 \\ -\sin(\phi) & \cos(\phi) & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & \cos(\psi) & -\sin(\psi) \\ 0 & 0 & 0 & \sin(\psi) & \cos(\psi) \end{pmatrix}$$

(written as $1 + 1 + (d - 2) + 1 + 1$ -block matrix).

Since Cartan subsets (with respect to C) are of the shape $\exp(\mathfrak{c}')t$ for $\mathfrak{c}' \subseteq \mathfrak{g}^{-\sigma} \cap \text{Ad}(t)(\mathfrak{g}^{-\sigma})$ of dimension 2, it is useful to know what this intersection looks like.

Proposition 6.5. (i) For $q = 0$ and $\phi \in \mathbb{R}$ the space $\mathfrak{g}^{-\sigma} \cap \text{Ad}(t_\phi)(\mathfrak{g}^{-\sigma})$ is spanned by \mathfrak{c} and

$$\begin{cases} F_{2,d+1}, \dots, F_{d-1,d+1} & \phi \notin \pi\mathbb{Z} \\ F_{0,2}, \dots, F_{0,d}, F_{1,d+1}, \dots, F_{d-1,d+1} & \phi \in \pi\mathbb{Z}. \end{cases}$$

(ii) For $q > 0$ and $\phi, \psi \in \mathbb{R}$, the space $\mathfrak{g}^{-\sigma} \cap \text{Ad}(t_{\phi,\psi})(\mathfrak{g}^{-\sigma})$ is spanned by \mathfrak{c} and

$$\begin{cases} 0 & \phi, \psi, \psi + \phi, \psi - \phi \notin \pi\mathbb{Z} \\ F_{0,d} \mp F_{1,d+1} & \phi, \psi, \psi \pm \phi \notin \pi\mathbb{Z}, \psi \mp \phi \in \pi\mathbb{Z} \\ F_{0,d}, F_{1,d+1} & \phi, \psi \in \frac{\pi}{2} + \pi\mathbb{Z}, \\ F_{2,d+1}, \dots, F_{d-1,d+1} & \phi \notin \pi\mathbb{Z}, \psi \in \pi\mathbb{Z}, \\ F_{0,2}, \dots, F_{0,d-1} & \psi \notin \pi\mathbb{Z}, \phi \in \pi\mathbb{Z}, \\ F_{0,2}, \dots, F_{0,d}, F_{1,d+1}, \dots, F_{d-1,d+1} & \phi, \psi \in \pi\mathbb{Z}. \end{cases}$$

Proof. (i) Let $X \in \mathfrak{g}^{-\sigma}$, say

$$X = \sum_{i=1}^d (a_i F_{0,i} + b_i F_{i,d+1}),$$

then

$$\begin{aligned} \text{Ad}(t_\phi)(X) &= a_1 F_{0,1} + b_1 (\cos(\phi) F_{1,d+1} + \sin(\phi) F_{0,d+1}) \\ &\quad + \sum_{i=2}^d (a_i (\cos(\phi) F_{0,i} - \sin(\phi) F_{1,i}) + b_i F_{i,d+1}), \end{aligned}$$

which is an element of $\mathfrak{g}^{-\sigma}$ iff $b_1 \sin(\phi) = a_i \sin(\phi) = 0$ ($i = 2, \dots, d$). If $\phi \notin \pi\mathbb{Z}$, we have $\sin(\phi) \neq 0$, so that $b_1 = a_2 = \dots = a_d = 0$, and thus X lies in the span of $F_{0,1}$ and $F_{i,d+1}$ ($i = 2, \dots, d$).

(ii) Let $X \in \mathfrak{g}^{-\sigma}$, say

$$X = \sum_{i=1}^d (a_i F_{0,i} + b_i F_{i,d+1}),$$

then

$$\begin{aligned} \text{Ad}(t_{\phi,\psi})(X) &= \sum_{i=2}^{d-1} (a_i (\cos(\phi) F_{0,i} - \sin(\phi) F_{1,i}) + b_i (\cos(\psi) F_{i,d+1} - \sin(\psi) F_{i,d})) \\ &\quad + a_1 F_{0,1} + b_d F_{d,d+1} \\ &\quad + a_d (\cos(\psi) \cos(\phi) F_{0,d} - \cos(\psi) \sin(\phi) F_{1,d} \\ &\quad \quad + \sin(\psi) \cos(\phi) F_{0,d+1} - \sin(\psi) \sin(\phi) F_{1,d+1}) \\ &\quad + b_1 (\cos(\psi) \cos(\phi) F_{1,d+1} + \cos(\psi) \sin(\phi) F_{0,d+1} \\ &\quad \quad - \sin(\psi) \cos(\phi) F_{1,d} - \sin(\psi) \sin(\phi) F_{0,d}) \\ &= \sum_{i=2}^{d-1} (a_i (\cos(\phi) F_{0,i} - \sin(\phi) F_{1,i}) + b_i (\cos(\psi) F_{i,d+1} - \sin(\psi) F_{i,d})) \\ &\quad + (a_d \cos(\psi) \cos(\phi) - b_1 \sin(\psi) \sin(\phi)) F_{0,d} \\ &\quad + (a_d \sin(\psi) \cos(\phi) + b_1 \cos(\psi) \sin(\phi)) F_{0,d+1} \\ &\quad - (a_d \cos(\psi) \sin(\phi) + b_1 \sin(\psi) \cos(\phi)) F_{1,d} \\ &\quad + (-a_d \sin(\psi) \sin(\phi) + b_1 \cos(\psi) \cos(\phi)) F_{1,d+1}, \end{aligned}$$

which is contained in $\mathfrak{g}^{-\sigma}$ iff

$$\begin{aligned} 0 &= a_i \sin(\phi) & (i = 2, \dots, d-1) \\ 0 &= b_i \sin(\psi) & (i = 2, \dots, d-1) \\ 0 &= \begin{pmatrix} \sin(\psi) \cos(\phi) & \cos(\psi) \sin(\phi) \\ \cos(\psi) \sin(\phi) & \sin(\psi) \cos(\phi) \end{pmatrix} \begin{pmatrix} a_d \\ b_1 \end{pmatrix}. \end{aligned}$$

If none of $\phi, \psi, \psi + \phi, \psi - \phi$ are contained in $\pi\mathbb{Z}$, we have $\sin(\phi), \sin(\psi) \neq 0$ and the matrix is regular. This shows that $a_i = b_i = 0$ ($i = 2, \dots, d-1$), as well as $a_d = b_1 = 0$, so that $X \in \mathfrak{c}$.

If $\phi, \psi, \psi \pm \phi \notin \pi\mathbb{Z}$ but $\psi \mp \phi \in \pi\mathbb{Z}$, the third condition reads

$$0 = \sin(\psi) \cos(\phi) \begin{pmatrix} 1 & \pm 1 \\ \pm 1 & 1 \end{pmatrix} \begin{pmatrix} a_d \\ b_1 \end{pmatrix},$$

where $\sin(\psi) \cos(\phi) \neq 0$ (by assumption $\sin(\psi) \neq 0$; if $\cos(\phi) = 0$, we had $2\phi \in \pi\mathbb{Z}$, but then $\psi \pm \phi = (\psi \mp \phi) \pm 2\phi \in \pi\mathbb{Z}$ as well). This shows that $a_i = b_i = 0$ ($i = 2, \dots, d-1$) and $a_d \pm b_1 = 0$. Consequently, X is spanned by \mathfrak{c} and $F_{0,d} \mp F_{1,d+1}$.

If $\phi, \psi \notin \pi\mathbb{Z}$ but $\psi + \phi, \psi - \phi \in \pi\mathbb{Z}$, we have $\phi, \psi \in \frac{\pi}{2} + \pi\mathbb{Z}$ and thus the conditions read

$$a_i = b_i = 0 \quad (i = 2, \dots, d-1),$$

so that X lies in the span of \mathfrak{c} and $F_{0,d}, F_{1,d+1}$.

If $\phi \notin \pi\mathbb{Z}$ but $\psi \in \pi\mathbb{Z}$, the conditions read

$$0 = a_i \quad (i = 2, \dots, d-1) \quad 0 = \cos(\psi) \sin(\phi) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} a_d \\ b_1 \end{pmatrix},$$

where neither $\cos(\psi)$ nor $\sin(\phi)$ are zero. Thus we have $a_i = 0$ ($i = 2, \dots, d-1$) and $a_d = b_1 = 0$. Thus, X lies in the span of \mathfrak{c} and $F_{2,d+1}, \dots, F_{d-1,d+1}$.

If conversely $\psi \notin \pi\mathbb{Z}$ and $\phi \in \pi\mathbb{Z}$, we get $b_i = 0$ ($i = 2, \dots, d-1$) and $a_d = b_1 = 0$, thus X lies in the span of \mathfrak{c} and $F_{0,2}, \dots, F_{0,d-1}$.

Lastly, if $\psi, \phi \in \pi\mathbb{Z}$, we also have $\psi + \phi, \psi - \phi \in \pi\mathbb{Z}$. Thus, all conditions are satisfied and thus X lies in the span of \mathfrak{c} and $F_{0,2}, \dots, F_{0,d}, F_{1,d+1}, \dots, F_{d-1,d+1}$. \square

We now first classify the Cartan subsets of G for the case $q = 0$.

Proposition 6.6. *If $q = 0$, C is the only Cartan subset of G (relative to C).*

Proof. Let $C' = \exp(\mathfrak{c}')t_\phi$ be a Cartan subset, with $\mathfrak{c}' = \mathfrak{t}' \oplus \mathfrak{a}'$.

If $\mathfrak{t}' = \mathfrak{t}$, we have $\mathfrak{c}' = \mathfrak{c}$, then t can be absorbed into $\exp(\mathfrak{c}')$. Consequently, $C' = C$.

Otherwise, $\mathfrak{t}' = 0$ and \mathfrak{a}' is a two-dimensional commutative subspace of $\mathfrak{p}^{-\sigma} \cap \text{Ad}(t_\phi)(\mathfrak{g}^{-\sigma})$ spanned say by $F_{d,d+1}$ and X . Expand $X \in \mathfrak{p}^{-\sigma}$ as

$$X = \sum_{i=1}^d a_i F_{i,d+1}.$$

Note that

$$0 = [F_{d,d+1}, X] = - \sum_{i=1}^d a_i F_{i,d} = - \sum_{i=1}^{d-1} a_i F_{i,d},$$

which implies that $a_1 = \dots = a_{d-1} = 0$, whence X is a multiple of $F_{d,d+1}$ and \mathfrak{a}' isn't two dimensional. Thus, there is no Cartan subset with $\mathfrak{t}' = 0$. \square

Interestingly, we will now demonstrate that there are more options for Cartan subsets in cases $q > 0$:

Proposition 6.7. *For $q > 0$, the Cartan subsets of G are $C' = \exp(\mathfrak{c}')t$ for*

- (i) $\mathfrak{c}' = \mathfrak{c}$ and $t = 1$;
- (ii) \mathfrak{c}' is spanned by $F_{0,1} \pm F_{d,d+1}$ and $F_{0,d} \mp F_{1,d+1}$ and t is 1 or $t_{0,\pi}$;
- (iii) $\mathfrak{t}' = \mathbb{R}F_{0,1}$ and \mathfrak{a}' is a 1-dimensional subspace of the span of $F_{2,d+1}, \dots, F_{p,d+1}$, and t is either 1 or $t_{0,\pi}$;
- (iv) $\mathfrak{t}' = \mathbb{R}F_{d,d+1}$ and \mathfrak{a}' is a 1-dimensional subspace of the span of $F_{0,p+1}, \dots, F_{0,d-1}$ and t is either 1 or $t_{\pi,0}$;
- (v) $\mathfrak{c}' = \mathbb{R}F_{0,d} \oplus \mathbb{R}F_{1,d+1}$ and $t = t_{\phi,\psi}$ for $\phi, \psi \in \frac{\pi}{2} + \pi\mathbb{Z}$;
- (vi) $\mathfrak{t}' = 0$ and \mathfrak{a}' is a two-dimensional commutative subalgebra of the span of $F_{0,p+1}, \dots, F_{0,d}, F_{1,d+1}, \dots, F_{p,d+1}$ with $t = t_{\phi,\psi}$ for $\phi, \psi \in \pi\mathbb{Z}$.

Proof. \mathfrak{t}' is a subspace of \mathfrak{t} , so it can have dimension 2, 1, or 0.

If $\dim(\mathfrak{t}') = 2$, then $\mathfrak{c}' = \mathfrak{t}' = \mathfrak{t} = \mathfrak{c}$. Furthermore, t can be absorbed into $\exp(\mathfrak{c}')$, whence $C' = C$, which is the 1st case

If $\dim(\mathfrak{t}') = 1$, say spanned by $X = aF_{0,1} + bF_{d,d+1}$. Then \mathfrak{a}' is spanned by Y , which lies in $\mathfrak{p}^{-\sigma}$. We can expand Y as

$$Y = \sum_{i=p+1}^d c_i F_{0,i} + \sum_{i=1}^p d_i F_{i,d+1},$$

then

$$\begin{aligned} [X, Y] &= - \sum_{i=p+1}^d ac_i F_{1,i} + ad_1 F_{0,d+1} - \sum_{i=1}^p bd_i F_{i,d} + bc_d F_{0,d+1} \\ &= -(ac_d + bd_1)F_{1,d} + (ad_1 + bc_d) - \sum_{i=p+1}^{d-1} ac_i F_{1,i} - \sum_{i=2}^p bd_i F_{i,d}. \end{aligned}$$

All basis vectors in the last equation are linearly independent, so we require

$$\begin{aligned} 0 &= ac_i & (i = p+1, \dots, d-1) \\ 0 &= bd_i & (i = 2, \dots, p) \\ 0 &\begin{pmatrix} a & b \\ b & a \end{pmatrix} \begin{pmatrix} c_d \\ d_1 \end{pmatrix}. \end{aligned}$$

If the matrix is regular, i.e. $a^2 - b^2 \neq 0$, and $a, b \neq 0$ as well, then $Y = 0$, which contradicts $\dim(\mathfrak{a}') = 1$. Consequently, at least one of $a, b, a+b, a-b$ must be zero. If more than one is zero, all others must be zero as well, which implies $X = 0$, which contradicts $\dim(\mathfrak{t}') = 1$.

In case $a \pm b = 0$, say X being a multiple of $F_{0,1} \mp F_{d,d+1}$, then Y must be a multiple of $F_{0,d} \pm F_{1,d+1}$. By Proposition 6.5(ii), this is an element of $\mathfrak{g}^{-\sigma} \cap \text{Ad}(t_{\phi,\psi})(\mathfrak{g}^{-\sigma})$ iff $\psi \pm \phi \in \pi\mathbb{Z}$. Then $t_{\phi,\psi} = t_{\phi,\mp\phi} t_{0,\psi\pm\phi}$, where $t_{\phi,\mp\phi} \in \exp(\mathfrak{c}')$ and $t_{0,\psi\pm\phi} = 1$ or $t_{0,\pi}$. This is the 2nd case with signs reversed.

In case $b = 0$, we can take $X = F_{0,1}$ without loss of generality. Then Y lies in the span of $F_{2,d+1}, \dots, F_{p,d+1}$. By Proposition 6.5(ii), such an element is contained in the proper intersection iff $\psi \in \pi\mathbb{Z}$. Furthermore, $t = t_{\phi,\psi} = t_{\phi,0} t_{0,\psi}$ with $t_{\phi,0} \in \exp(\mathfrak{c}')$ and $t_{0,\psi} = 1$ or $t_{0,\pi}$. This is the 3rd case.

In case $a = 0$, we can take $X = F_{d,d+1}$ without loss of generality. Then we have Y in the span of $F_{0,p+1}, \dots, F_{0,d-1}$. By Proposition 6.5(ii), such an element is contained in the proper intersection iff $\phi \in \pi\mathbb{Z}$. Furthermore, $t = t_{\phi,\psi} = t_{0,\psi} t_{\phi,0}$ with $t_{0,\psi} \in \exp(\mathfrak{c}')$ and $t_{\phi,0} = 1$ or $t_{\pi,0}$. This is the 4th case.

This covers all the cases for $\dim(\mathfrak{t}') = 1$, let's now turn our attention to $\mathfrak{t}' = 0$. In this case, \mathfrak{a}' has to be a two-dimensional subalgebra of $\mathfrak{p}^{-\sigma} \cap \text{Ad}(t_{\phi,\psi})(\mathfrak{g}^{-\sigma})$, which is spanned by

$$\begin{cases} 0 & \phi, \psi, \psi + \phi, \psi - \phi \notin \pi\mathbb{Z} \\ F_{0,d} \mp F_{1,d+1} & \phi, \psi, \psi \pm \phi \notin \pi\mathbb{Z}, \psi \mp \phi \in \pi\mathbb{Z} \\ F_{0,d}, F_{1,d+1} & \phi, \psi \in \frac{\pi}{2} + \pi\mathbb{Z}, \\ F_{2,d+1}, \dots, F_{p,d+1} & \phi \notin \pi\mathbb{Z}, \psi \in \pi\mathbb{Z}, \\ F_{0,p+1}, \dots, F_{0,d-1} & \psi \notin \pi\mathbb{Z}, \phi \in \pi\mathbb{Z}, \\ F_{0,p+1}, \dots, F_{0,d}, F_{1,d+1}, \dots, F_{p,d+1} & \phi, \psi \in \pi\mathbb{Z}. \end{cases}$$

This already excludes the first two lines. In case $\phi, \psi \in \frac{\pi}{2} + \pi\mathbb{Z}$, we can take $\mathfrak{a}' = \mathbb{R}F_{0,d} \oplus \mathbb{R}F_{1,d+1}$, which is a commutative subalgebra. This is the 5th case.

In case one of ϕ, ψ is contained in $\pi\mathbb{Z}$ but the other is not, there exists no 2-dimensional commutative subalgebra, so the only remaining possible case is $\phi, \psi \in \pi\mathbb{Z}$, which is the 6th case. \square

Next, we are interested which of these Cartan subsets are conjugate to each other. For this we need to investigate the action of

$$N_K^T := \{(h, h') \in (H \cap K)^2 \mid hTh'^{-1} = T\}$$

on T .

Lemma 6.8. *If $q > 1$, N_K^T acts on T as an affine reflection group generated by*

$$\begin{aligned} s_0: t_{\phi,\psi} &\mapsto t_{\pi-\phi, \pi-\psi} \\ s_1: t_{\phi,\psi} &\mapsto t_{\phi, -\psi} \\ s_2: t_{\phi,\psi} &\mapsto t_{-\phi, \phi}. \end{aligned}$$

Proof. Let $h \in K \cap H$, then h can be written as a $1 + p + q + 1$ -block matrix as

$$\begin{pmatrix} A & 0 & 0 & \\ 0 & B & 0 & 0 \\ 0 & 0 & C & 0 \\ 0 & 0 & 0 & D \end{pmatrix}$$

for $A, D \in \mathbb{R}$ and $B \in \mathbb{R}^{p \times p}, C \in \mathbb{R}^{q \times q}$, with $A^2 = 1, D^2 = 1$ and $B \in O(p), C \in O(q)$ with $\det(B) = A$ and $\det(C) = D$. Since H isn't all of G^σ , but only those elements that stabilise D_0 , we also have $A = D$. Consequently,

$$h = \begin{pmatrix} \pm 1 & 0 & 0 & 0 \\ 0 & A & 0 & 0 \\ 0 & 0 & B & 0 \\ 0 & 0 & 0 & \pm 1 \end{pmatrix}$$

with $A \in O(p), B \in O(q)$ with $\det(A) = \det(B) = \pm 1$. Furthermore, we can write $t_{\phi,\psi}$ in the same $1 + p + q + 1$ -block matrix form as

$$\begin{pmatrix} \cos(\phi) & \sin(\phi)e_1^T & 0 & 0 \\ -\sin(\phi)e_1 & 1 + (\cos(\phi) - 1)e_1e_1^T & 0 & 0 \\ 0 & 0 & 1 + (\cos(\psi) - 1)e_qe_q^T & -\sin(\psi)e_q \\ 0 & 0 & \sin(\psi)e_q^T & \cos(\psi) \end{pmatrix}.$$

If $h, h' \in K \cap H$ are block diagonal matrices with ϵ, A, B, ϵ and ζ, C, D, ζ , we have

$$ht_{\phi,\psi}h'^{-1} = \begin{pmatrix} X & 0 \\ 0 & Y \end{pmatrix}$$

with

$$X = \begin{pmatrix} \epsilon\zeta \cos(\phi) & \epsilon \sin(\phi)(Ce_1)^T \\ -\zeta \sin(\phi)Ae_1 & AC^T + (\cos(\phi) - 1)Ae_1(Ce_1)^T \end{pmatrix}$$

$$Y = \begin{pmatrix} BD^T + (\cos(\psi) - 1)Be_q(De_q)^T & -\zeta \sin(\psi)Be_q \\ \epsilon \sin(\psi)(De_q)^T & \epsilon\zeta \cos(\psi) \end{pmatrix}$$

Since this lies in T again, we have that Ae_1, Ce_1, e_1 are proportional, as are Be_q, De_q, e_q , say

$$Ae_1 = \alpha e_1, \quad Be_q = \beta e_q, \quad Ce_1 = \gamma e_1, \quad De_q = \delta e_q.$$

Since A, C, B, D are orthogonal, we have $\alpha, \beta, \gamma, \delta \in \{\pm 1\}$. Then

$$\epsilon\gamma = \alpha\zeta, \quad \epsilon\zeta = \alpha\gamma, \quad \epsilon\delta = \beta\zeta, \quad \epsilon\zeta = \beta\delta.$$

If $\epsilon\zeta = 1$, then $\alpha = \gamma$ and $\beta = \delta$. Then $ht_{\phi,\psi}h'^{-1} = t_{\epsilon\gamma\phi,\epsilon\delta\psi}$. All four possible combinations of signs of $\epsilon\gamma, \beta\zeta$ yield an action on T that is generated by s_1, s_2 .

If $\epsilon\zeta = -1$, we have $\alpha = -\gamma$ and $\beta = -\delta$, then $ht_{\phi,\psi}h'^{-1} = t_{\pi-\gamma\phi,\pi-\delta\psi}$. This transformation is definitely contained in the group generated by s_0, s_1, s_2 .

Conversely, s_0 is effected by

$$h = \text{diag}(1, -1, -1, 1, \dots, 1, -1, -1, 1), \quad h' = \text{diag}(-1, 1, -1, 1, \dots, 1, -1, 1, -1).$$

Since we assumed $q > 1$ (and hence also $p > 1$), all matrices are elements of K . Furthermore, they commute with D_0 , so that they are elements of $K \cap H$. The transformation s_1 is effected by

$$h = h' = \text{diag}(1, \dots, 1, -1, -1, 1)$$

and s_2 by

$$h = h' = \text{diag}(1, -1, -1, \dots, 1).$$

□

Lemma 6.9. *For $q = 1$, the action of N_K^T on T is generated by the reflection $t_{\phi,\psi} \mapsto t_{-\phi,\psi}$ and the translation $t_{\phi,\psi} \mapsto t_{\phi+\pi,\psi+\pi}$.*

Proof. An element $h \in H \cap K$ has the following $1 + p + 1 + 1$ -block matrix shape:

$$h = \begin{pmatrix} \epsilon & 0 & 0 & 0 \\ 0 & A & 0 & 0 \\ 0 & 0 & \epsilon & 0 \\ 0 & 0 & 0 & \epsilon \end{pmatrix}$$

for $A \in SO(p)$ and $\epsilon = \pm 1$. Two such elements (with variables ϵ, A and δ, B) give rise to an element of N_K^T if $Ae_1 = \alpha e_1, Be_1 = \beta e_1$ and $\epsilon\delta = \alpha\beta$ and $\epsilon\beta = \alpha\delta$. If $\epsilon\delta = 1$, we have $\alpha = \beta$, which corresponds to $t_{\phi,\psi} \mapsto t_{-\phi,\psi}$; and if $\epsilon\delta = -1$, we have $\alpha = -\beta$, which corresponds to $t_{\phi,\psi} \mapsto t_{\pi-\epsilon\beta\phi,\psi+\pi}$.

Conversely, the reflection is effected by

$$h = h' = \text{diag}(1, -1, -1, 1, \dots, 1),$$

which exists because $p = d - 1 > 1$ (by the original assumption that $d > 2$), and the translation by

$$h = 1, h' = \text{diag}(-1, -1, 1, \dots, 1, -1, -1).$$

□

Corollary 6.10. *For $q > 0$, the equivalence classes of Cartan subsets can be represented by:*

- (i) C ;
- (ii) $\exp(\mathbb{R}(F_{0,1} + F_{d,d+1}) \oplus \mathbb{R}(F_{0,d} - F_{1,d+1}))$;
- (iii) $\exp(\mathbb{R}(F_{0,1} + F_{d,d+1}) \oplus \mathbb{R}(F_{0,d} - F_{1,d+1}))t_{0,\pi}$;

- (iv) $\exp(\mathbb{R}F_{0,1} \oplus \mathbb{R}F_{2,d+1})$;
- (v) $\exp(\mathbb{R}F_{d,d+1} \oplus \mathbb{R}F_{0,d-1})$ (*this doesn't exist for $q = 1$*);
- (vi) $\exp(\mathbb{R}F_{0,d} \oplus \mathbb{R}F_{1,d+1})$;
- (vii) $\exp(\mathbb{R}F_{0,d} \oplus \mathbb{R}F_{1,d+1})t_{\pi/2, \pi/2}$;
- (viii) $\exp(\mathbb{R}F_{0,d} \oplus \mathbb{R}F_{1,d+1})t_{0, \pi}$.

Proof. The compact parts of the Cartan subsets described in Proposition 6.7 consists of $t_{\phi, \psi}$ with

- (i) no conditions, this is the first equivalence class.
- (ii) $\phi \equiv \pm\psi \pmod{2\pi\mathbb{Z}}$ or $\phi \equiv \pi \pm \psi \pmod{2\pi\mathbb{Z}}$. The first two cases (\pm) and cases three and four are related by $\phi \mapsto \phi$, which is a transformation that can be enacted using N_K^T (cf. Lemmas 6.8, 6.9). This yields the second and third equivalence class, respectively.
- (iii) $\psi \in \pi\mathbb{Z}$, i.e. $\psi \equiv 0 \pmod{2\pi\mathbb{Z}}$ or $\psi \equiv \pi \pmod{2\pi\mathbb{Z}}$. Both cases are related via a shift $(\psi, \phi) \mapsto (\psi + \pi, \phi + \pi)$, which can be effected using N_K^T . This yields the fourth equivalence class.
- (iv) The same for ϕ , which yields the fifth equivalence class.
- (v) (ϕ, ψ) being congruent (modulo $2\pi\mathbb{Z} \oplus 2\pi\mathbb{Z}$) to one of

$$\left(\frac{\pi}{2}, \frac{\pi}{2}\right), \left(\frac{3\pi}{2}, \frac{3\pi}{2}\right), \left(\frac{\pi}{2}, \frac{3\pi}{2}\right), \left(\frac{3\pi}{2}, \frac{\pi}{2}\right).$$

The first and second (and third and fourth) element are related via the shift $(\phi, \psi) \mapsto (\phi + \pi, \psi + \pi)$, and the first and third element are related using the reflection $(\phi, \psi) \mapsto (-\phi, \psi)$, thus they all are equivalent, which is the seventh equivalence class.

- (vi) (ϕ, ψ) being congruent (modulo $2\pi\mathbb{Z} \oplus 2\pi\mathbb{Z}$) to one of
- $$(0, 0), (\pi, \pi), (0, \pi), (\pi, 0).$$

Of these, the first and second (and third and fourth) are related via the translation, and no others are related via N_K^T , which is the sixth and eighth equivalence class, respectively.

□

6.2. Coordinates. We shall now introduce unified coordinates for all Cartan subsets and answer the question to what extent the functions u, v define a smooth manifold structure on $\text{GP}(G/Q, 4)/G$ (or $MA\backslash\hat{G}/MA$ via ψ).

Definition 6.11. *Let*

$$D := \left\{ (\chi_1, \chi_2) \in \mathbb{C}^2 \mid \chi_1, \chi_2, \frac{\chi_1 + \chi_2}{2}, \frac{\chi_1 - \chi_2}{2} \notin i\pi\mathbb{Z} \right\}.$$

Define $f, g: D \rightarrow \mathbb{C}^2$ by

$$\begin{aligned} f: (\chi_1, \chi_2) &\mapsto \begin{pmatrix} \text{sech}^2(\chi_1/2) \text{sech}^2(\chi_2/2) \\ \tanh^2(\chi_1/2) \tanh^2(\chi_2/2) \end{pmatrix} \\ g: (\chi_1, \chi_2) &\mapsto \begin{pmatrix} \sinh^2(\chi_1/2) \sinh^2(\chi_2/2) \\ \cosh^2(\chi_1/2) \cosh^2(\chi_2/2) \end{pmatrix}. \end{aligned}$$

Note that $\chi_i \notin i\pi\mathbb{Z}$ implies that $\frac{\chi_i}{2} \notin \left(\frac{i\pi}{2} + i\pi\mathbb{Z}\right)$, so that $\cosh\left(\frac{\chi_i}{2}\right) \neq 0$ and f is well-defined.

Lemma 6.12. *f is a local diffeomorphism.*

Proof. Note that

$$f(\chi_1, \chi_2) = \begin{pmatrix} \frac{1}{g_2(\chi_1, \chi_2)} \\ \frac{g_1(\chi_1, \chi_2)}{g_2(\chi_1, \chi_2)} \end{pmatrix},$$

so that f is a local diffeomorphism iff g is. We now compute the Jacobian of g :

$$g'(\chi_1, \chi_2) = \begin{pmatrix} \sinh\left(\frac{\chi_1}{2}\right) \cosh\left(\frac{\chi_1}{2}\right) \sinh^2\left(\frac{\chi_2}{2}\right) & \sinh^2\left(\frac{\chi_1}{2}\right) \sinh\left(\frac{\chi_2}{2}\right) \cosh\left(\frac{\chi_2}{2}\right) \\ \sinh\left(\frac{\chi_1}{2}\right) \cosh\left(\frac{\chi_1}{2}\right) \cosh^2\left(\frac{\chi_2}{2}\right) & \cosh^2\left(\frac{\chi_1}{2}\right) \sinh\left(\frac{\chi_2}{2}\right) \cosh\left(\frac{\chi_2}{2}\right) \end{pmatrix},$$

whose determinant is

$$\begin{aligned} \det(g'(\chi_1, \chi_2)) &= \sinh\left(\frac{\chi_1}{2}\right) \cosh\left(\frac{\chi_1}{2}\right) \sinh\left(\frac{\chi_2}{2}\right) \cosh\left(\frac{\chi_2}{2}\right) \\ &\quad \cdot \left(\sinh^2\left(\frac{\chi_2}{2}\right) \cosh^2\left(\frac{\chi_1}{2}\right) - \cosh^2\left(\frac{\chi_2}{2}\right) \sinh^2\left(\frac{\chi_1}{2}\right) \right) \\ &= 16 \sinh(\chi_1) \sinh(\chi_2) \sinh\left(\frac{\chi_2 - \chi_1}{2}\right) \sinh\left(\frac{\chi_2 + \chi_1}{2}\right). \end{aligned}$$

Since none of $\chi_1, \chi_2, \frac{\chi_1 + \chi_2}{2}, \frac{\chi_1 - \chi_2}{2}$ lies in the zero locus of \sinh , we have $\det(g'(\chi_1, \chi_2)) \neq 0$. By the inverse function theorem, g is a local diffeomorphism, hence so is f . \square

Since the complement of D is a union of (locally finitely many) codimension 2 real subspaces, D is still connected and therefore a covering space of $f(D)$. In particular, D carries an action of the fundamental groupoid of $f(D)$. A way of approaching this is to first consider some obvious symmetries of f and g , and then look what remains.

Lemma 6.13. *Define $s_0, s_1, s_2: D \rightarrow D$ by*

$$\begin{aligned} s_0(\chi_1, \chi_2) &\mapsto (\chi_1, 2\pi i - \chi_2) \\ s_1(\chi_1, \chi_2) &\mapsto (\chi_2, \chi_1) \\ s_2(\chi_1, \chi_2) &\mapsto (-\chi_1, \chi_2). \end{aligned}$$

These three transformations generate a Coxeter group \tilde{W} of type \tilde{C}_2 (and its (scaled) affine action on \mathbb{C}^2) of symmetries of f .

Proof. As before, $f \circ s_i = f$ iff $g \circ s_i = g$, so we check for g . Since \sinh^2, \cosh^2 are even and periodic with period πi , we see that s_0, s_2 are symmetries. Furthermore, g is symmetric with respect to exchanging χ_1, χ_2 , so that s_1 is also a symmetry.

To see that s_0, s_1, s_2 generate a Coxeter group of type \tilde{C}_2 , note that they are all involutions, that

$$\begin{aligned} s_0 s_1(\chi_1, \chi_2) &\mapsto (\chi_2, 2\pi i - \chi_1), \\ s_0 s_2(\chi_1, \chi_2) &\mapsto (-\chi_1, 2\pi i - \chi_2), \\ s_1 s_2(\chi_1, \chi_2) &\mapsto (\chi_2, -\chi_1) \end{aligned}$$

have orders 4, 2, 4, respectively, whence we see that they generate an affine Coxeter group of type \tilde{C}_2 . \square

Proposition 6.14. *Consequently, it suffices to study D/\tilde{W} , of which a fundamental domain is given by*

$$X = \bigsqcup_{I \subseteq \{0,1,2\}} X_I$$

with

$$\begin{aligned} X_\emptyset &= \mathbb{R}^2 + \{(a, b) \mid 0 < a < b < \pi\}i & X_{\{0\}} &= \mathbb{R} \times \mathbb{R}_{>0} + \{(a, \pi) \mid 0 < a < \pi\}i \\ X_{\{1\}} &= \{(a, b) \mid a < b\} + (0, \pi)i(1, 1) & X_{\{2\}} &= \mathbb{R}_{>0} \times \mathbb{R} + \{(0, a) \mid 0 < a < \pi\}i \\ X_{\{0,1\}} &= \{(a, b) \mid 0 < a < b\} + (\pi, \pi)i & X_{\{0,2\}} &= \mathbb{R}_{>0}^2 + (0, \pi)i \\ X_{\{1,2\}} &= \{(a, b) \mid 0 < a < b\}, \end{aligned}$$

where every X_I is fixed by \tilde{W}_I , the parabolic subgroup generated by s_i ($i \in I$).

Proof. We first focus on the imaginary parts. Since the action is just a rescaled version of the affine Weyl group of a B_2 root system, a fundamental domain is given by the fundamental alcove of said root system. This shows that the set

$$\tilde{X} := \{(\chi_1, \chi_2) \in D \mid 0 \leq \text{Im}(\chi_1) \leq \text{Im}(\chi_2) \leq \pi\}$$

(the preimage of a rescaled fundamental alcove under the projection onto the imaginary parts) touches every orbit, but not necessarily that it touches every orbit once: let $(\chi_1, \chi_2) \in \tilde{X}$ and $w \in \tilde{W}$, then we have either $w \cdot (\chi_1, \chi_2) \notin \tilde{X}$ or $\text{Im}(w \cdot (\chi_1, \chi_2)) = \text{Im}(\chi_1, \chi_2)$. However, since \tilde{W} also acts on the real parts, this is not enough to conclude that $w \cdot (\chi_1, \chi_2) = (\chi_1, \chi_2)$. In particular, if $\text{Im}(\chi_1, \chi_2)$ lies in the face stabilised by the parabolic subgroup \tilde{W}_I , we need to consider \tilde{W}_I 's action on the real parts and restrict to a fundamental domain for that, too.

We now proceed by the preimages of the faces of the fundamental alcove, indexed by its stabiliser subgroup \tilde{W}_I .

$I = \emptyset$: We have $0 < \text{Im}(\chi_1) < \text{Im}(\chi_2) < \pi$, i.e. $(\chi_1, \chi_2) \in X_\emptyset$. Since the stabiliser subgroup is the trivial group, a fundamental domain is given by allowing all real values. We consequently end up with X_\emptyset .

$I = \{0\}$: We have $0 < \text{Im}(\chi_1) < \text{Im}(\chi_2) = \pi$. The stabiliser subgroup is generated by s_0 , which acts like the reflection $s_1 s_2 s_1$ on the real parts. Consequently, a fundamental domain is given by requiring

$$0 < \text{Im}(\chi_1) < \text{Im}(\chi_2) = \pi, \quad \text{Re}(\chi_2) \geq 0.$$

Since $\chi_2 \neq \pi i$, we also know that $\text{Re}(\chi_2) > 0$ is in fact a strict inequality. This defines $X_{\{0\}}$.

$I = \{1\}$: We have $0 < \text{Im}(\chi_1) = \text{Im}(\chi_2) < \pi$. The stabiliser subgroup is generated by s_1 , which acts on the real parts by swapping them. A fundamental domain is therefore given by requiring

$$0 < \text{Im}(\chi_1) = \text{Im}(\chi_2) < \pi, \quad \text{Re}(\chi_1) \leq \text{Re}(\chi_2).$$

Since $\frac{\chi_1 - \chi_2}{2} \neq 0$, we additionally know that $\text{Re}(\chi_1) < \text{Re}(\chi_2)$, which defines $X_{\{1\}} (\subseteq D)$.

$I = \{2\}$: We have $0 = \text{Im}(\chi_1) < \text{Im}(\chi_2) < \pi$. The stabiliser subgroup is generated by s_2 , which acts on the real parts by negating the first. Consequently, a fundamental domain is given by

$$0 = \text{Im}(\chi_1) < \text{Im}(\chi_2) < \pi, \quad \text{Re}(\chi_1) \geq 0.$$

Since $\chi_1 \neq 0$ (as required by D), we also know that $\text{Re}(\chi_1) > 0$, which defines $X_{\{2\}}$.

$I = \{0, 1\}$: We have $\text{Im}(\chi_1) = \text{Im}(\chi_2) = \pi$. The stabiliser subgroup is generated by s_0, s_1 , which act on the real parts like the Weyl group of B_2 . Consequently, a fundamental domain is given by requiring

$$\text{Im}(\chi_1) = \text{Im}(\chi_2) = \pi, \quad 0 \leq \text{Re}(\chi_1) \leq \text{Re}(\chi_2).$$

Since $\chi_1 \neq \pi i$, we can in particular choose $0 < \text{Re}(\chi_1)$, and since $\frac{\chi_1 - \chi_2}{2} \neq 0$, we can also choose $\text{Re}(\chi_1) < \text{Re}(\chi_2)$, which defines $X_{\{0,1\}}$.

$I = \{0, 2\}$: We have $0 = \text{Im}(\chi_1) < \text{Im}(\chi_2) = \pi$. The stabiliser subgroup is generated by s_0, s_2 , which act on the real parts by negating one or the other. Consequently, a fundamental domain is given by

$$0 = \text{Im}(\chi_1) < \text{Im}(\chi_2) = \pi, \quad 0 \leq \text{Re}(\chi_1), \text{Re}(\chi_2).$$

Since $\chi_1 \neq 0, \chi_2 \neq \pi i$, we in particular have $0 < \text{Re}(\chi_1), \text{Re}(\chi_2)$, which describes $X_{\{0,2\}}$.

$I = \{1, 2\}$: We have $0 = \text{Im}(\chi_1) = \text{Im}(\chi_2)$. The stabiliser subgroup is generated by s_1, s_2 , which act on the real parts like the Weyl group of B_2 , so that a fundamental domain is given by

$$0 = \text{Im}(\chi_1) = \text{Im}(\chi_2), \quad 0 \leq \text{Re}(\chi_1) \leq \text{Re}(\chi_2).$$

Since $\chi_1, \frac{\chi_1 - \chi_2}{2} \neq 0$, we additionally require $0 < \text{Re}(\chi_1) < \text{Re}(\chi_2)$, which describes $X_{\{1,2\}}$.

$I = \{0, 1, 2\}$: There are no elements stabilised by s_0, s_1, s_2 .

Consequently, the union of all X_I touches every orbit exactly once, and every element of \tilde{X} (and hence of D) is related by \tilde{W} to an element of one of the X_I . \square

Lemma 6.15. *f is injective on X . In particular, \tilde{W} is the group of all symmetries of f .*

Proof. We show the result for g . Assume $(\chi_1, \chi_2), (\chi'_1, \chi'_2) \in D$ with $g(\chi_1, \chi_2) = g(\chi'_1, \chi'_2)$. We are going to show that $(\chi_1, \chi_2), (\chi'_1, \chi'_2)$ lie in the same \tilde{W} -orbit. We have

$$\begin{aligned} \frac{1}{4}(\cosh(\chi_1) - 1)(\cosh(\chi_2) - 1) &= \sinh^2\left(\frac{\chi_1}{2}\right) \sinh^2\left(\frac{\chi_2}{2}\right) \\ &= \sinh^2\left(\frac{\chi'_1}{2}\right) \sinh^2\left(\frac{\chi'_2}{2}\right) \\ &= \frac{1}{4}(\cosh(\chi'_1) - 1)(\cosh(\chi'_2) - 1), \\ \frac{1}{4}(\cosh(\chi_1) + 1)(\cosh(\chi_2) + 1) &= \cosh^2\left(\frac{\chi_1}{2}\right) \cosh^2\left(\frac{\chi_2}{2}\right) \\ &= \sinh^2\left(\frac{\chi'_1}{2}\right) \sinh^2\left(\frac{\chi'_2}{2}\right) \\ &= \frac{1}{4}(\cosh(\chi'_1) + 1)(\cosh(\chi'_2) + 1), \end{aligned}$$

which shows that $\cosh(\chi_1) + \cosh(\chi_2) = \cosh(\chi'_1) + \cosh(\chi'_2)$ and $\cosh(\chi_1) \cosh(\chi_2) = \cosh(\chi'_1) \cosh(\chi'_2)$. By standard algebra this shows that either $(\cosh(\chi_1), \cosh(\chi_2)) = (\cosh(\chi'_1), \cosh(\chi'_2))$ or $= (\cosh(\chi'_2), \cosh(\chi'_1))$. In both cases, (χ'_1, χ'_2) is at most an application of s_1 removed from satisfying the first equation i.e. $\cosh(\chi_1) = \cosh(\chi'_1), \cosh(\chi_2) = \cosh(\chi'_2)$. So without loss of generality, we can assume that the equation is satisfied.

The equation can be rewritten as $\exp(\chi_i) + \exp(-\chi_i) = \exp(\chi'_i) + \exp(-\chi'_i)$ for $i = 1, 2$. By the same algebra argument we conclude $\exp(\chi_i) = \exp(\chi'_i)$ or $= \exp(-\chi'_i)$ for $i = 1, 2$. This yields four possibilities, all of which can be related to $\exp(\chi_1) = \exp(\chi'_1), \exp(\chi_2) = \exp(\chi'_2)$ using the sign flips s_2 and $s_1 s_2 s_1$. Without loss of generality assume therefore that this equation holds.

Since $\exp: (\mathbb{C}, +) \rightarrow (\mathbb{C}^\times, \cdot)$ is a group homomorphism with kernel $2\pi i\mathbb{Z}$, this implies that $(\chi'_1, \chi'_2) - (\chi_1, \chi_2) \in (2\pi i\mathbb{Z})^2$. Such a translation can be effected using the elementary translations $s_0 s_1 s_2 s_1$ and $s_1 s_0 s_1 s_2$.

This shows that $(\chi_1, \chi_2), (\chi'_1, \chi'_2)$ lie in the same \tilde{W} -orbit. \square

Next, we investigate the preimage of \mathbb{R}^2 under f (equivalently, g).

Lemma 6.16. *The preimage of \mathbb{R}^2 under f consists of all the \tilde{W} -orbits of*

$$Y = \bigsqcup_{I \subseteq \{0,1,2\}} Y_I$$

where

$$\begin{aligned} Y_\emptyset &= i\{(a, b) \mid 0 < a < b < \pi\} & Y_{\{0\}} &= \{0\} \times \mathbb{R}_{>0} + i\{(a, \pi) \mid 0 < a < \pi\} \\ Y_{\{1\}} &= \mathbb{R}_{>0}(-1, 1) + (0, \pi)i(1, 1) & Y_{\{2\}} &= \mathbb{R}_{>0} \times \{0\} + i\{(0, a) \mid 0 < a < \pi\} \\ Y_{\{0,1\}} &= \{(a, b) \mid 0 < a < b\} + i(\pi, \pi) & Y_{\{0,2\}} &= \mathbb{R}_{>0}^2 + i(0, \pi) \\ Y_{\{1,2\}} &= \{(a, b) \mid 0 < a < b\}. \end{aligned}$$

Proof. Let $(\chi_1, \chi_2) \in D$, without loss of generality $(\chi_1, \chi_2) \in X_I$ for an $I \subseteq \{0, 1, 2\}$, and write

$$\begin{aligned} c_1 &:= \sinh\left(\frac{\chi_1}{2}\right) \sinh\left(\frac{\chi_2}{2}\right) \\ c_2 &:= \cosh\left(\frac{\chi_1}{2}\right) \cosh\left(\frac{\chi_2}{2}\right). \end{aligned}$$

We now need to investigate when c_1^2, c_2^2 are real numbers. For this to happen, c_i needs to be purely real or purely imaginary, making for four cases:

Both real: In this case, $c_1 \pm c_2$ are both also real numbers. We have

$$c_2 \pm c_1 = \cosh\left(\frac{\chi_1 \pm \chi_2}{2}\right).$$

If $\chi_1 = 2a + 2\phi$ and $\chi_2 = 2b + 2\psi$, this reads

$$c_2 \pm c_1 = \cosh(a \pm b) \cos(\phi \pm \psi) + i \sinh(a \pm b) \sin(\phi \pm \psi).$$

For this to be purely imaginary, we need $a \pm b$ or $\phi \pm \psi \in \pi\mathbb{Z}$.

If both $\phi + \psi, \phi - \psi \notin \pi\mathbb{Z}$, we need $a + b, a - b = 0$, which implies $a = b = 0$. Furthermore, the inequalities imply that $0 < \phi < \psi < \frac{\pi}{2}$, which in turn implies that $(\chi_1, \chi_2) \in X_\emptyset$, with zero real part, hence $(\chi_1, \chi_2) \in Y_\emptyset$.

If $\phi - \psi \in \pi\mathbb{Z}$ and $\phi + \psi \notin \pi\mathbb{Z}$, we need $a + b = 0$. Furthermore, we have $0 < \phi = \psi < \frac{\pi}{2}$, which implies that $(\chi_1, \chi_2) \in X_{\{1\}}$. Adding in the fact that $a = -b$, we obtain $(\chi_1, \chi_2) \in Y_{\{1\}}$.

If $\phi + \psi \in \pi\mathbb{Z}$, we can have either $\phi = \psi = 0$ or $\phi = \psi = \frac{\pi}{2}$. In both cases we also have $\phi - \psi \in \pi\mathbb{Z}$. In both cases there are no further restrictions on a, b , which implies $(\chi_1, \chi_2) \in X_{\{1,2\}} = Y_{\{1,2\}}$ or $X_{\{0,1\}} = Y_{\{0,1\}}$, respectively.

Both imaginary: In this case, $c_1 \pm c_2$ are purely imaginary numbers, which equal

$$c_2 \pm c_1 = \cosh(a \pm b) \cos(\phi \pm \psi) + i \sinh(a \pm b) \sin(\phi \pm \psi).$$

Since the cosh of any real number is nonzero, we obtain $\cos(\phi + \psi) = \cos(\phi - \psi) = 0$, meaning that $\phi + \psi, \phi - \psi \in \frac{\pi}{2} + \pi\mathbb{Z}$. Since we chose $0 \leq \phi \leq \psi \leq \frac{\pi}{2}$, the only possible combinations are $\phi = 0, \psi = \frac{\pi}{2}$, which implies that $(\chi_1, \chi_2) \in X_{\{0,2\}} = Y_{\{0,2\}}$.

c_1 imaginary, c_2 real: In this case $c_2 \pm c_1$ are complex conjugates of each other, meaning that

$$\cosh\left(\frac{\chi_1 + \chi_2}{2}\right) = \overline{\cosh\left(\frac{\chi_1 - \chi_2}{2}\right)} = \cosh\left(\frac{\chi_1 - \chi_2}{2}\right),$$

which shows that one of

$$\begin{aligned} 2\pi i\mathbb{Z} &\ni \frac{\chi_1 + \chi_2}{2} - \frac{\overline{\chi_1} - \overline{\chi_2}}{2} \\ &= a + b + i\phi + i\psi - (a - b - i\phi + i\psi) = 2b + 2i\phi \\ 2\pi i\mathbb{Z} &\ni \frac{\chi_1 + \chi_2}{2} + \frac{\overline{\chi_1} - \overline{\chi_2}}{2} \\ &= a + b + i\phi + i\psi + (a - b - i\phi + i\psi) = 2a + 2i\psi. \end{aligned}$$

If the first is true, we need $b = 0$ and $\phi \in \pi\mathbb{Z}$. $b = 0$ is only allowed for $X_\emptyset, X_{\{1\}}, X_{\{2\}}$, of which only $X_{\{2\}}$ allows $\phi \in \pi\mathbb{Z}$ (namely $\phi = 0$). Since $b = 0$, we in particular also have $(\chi_1, \chi_2) \in Y_{\{2\}}$.

If the second is true, we need $a = 0$ and $\psi \in \pi\mathbb{Z}$, which means that (χ_1, χ_2) is not contained in any X_I , which is a contradiction.

c_1 **real**, c_2 **imaginary**: In this case, $c_1 \pm c_2$ are complex conjugates of each other, meaning that

$$\cosh\left(\frac{\chi_1 + \chi_2}{2}\right) = -\overline{\cosh\left(\frac{\chi_1 - \chi_2}{2}\right)} = -\cosh\left(\frac{\overline{\chi_1 - \chi_2}}{2}\right),$$

which shows that one of

$$\begin{aligned} \pi i + 2\pi i\mathbb{Z} &\ni 2b + 2i\phi \\ \pi i + 2\pi i\mathbb{Z} &\ni 2a + 2i\psi. \end{aligned}$$

In the first case, we have $b = 0$ and $\phi \in \frac{\pi}{2} + \pi\mathbb{Z}$, i.e. $\phi = \frac{\pi}{2}$, which implies that (χ_1, χ_2) is not contained in any X_I , which is a contradiction.

In the second case, we have $a = 0$ and $\psi = \frac{\pi}{2}$, which implies that $(\chi_1, \chi_2) \in Y_{\{0\}} \subseteq X_{\{0\}}$.

We thus obtain the following:

c_1	c_2	Real Faces
re	re	$Y_\emptyset, Y_{\{1\}}, Y_{\{0,1\}}, Y_{\{1,2\}}$
re	im	$Y_{\{0\}}$
im	re	$Y_{\{2\}}$
im	im	$Y_{\{0,2\}}$.

□

Corollary 6.17. *Restricted to Y , f is a diffeomorphism onto $f(Y)$, which is given by*

$$\{(u, v) \in \mathbb{R}^2 \mid u, v, 1 + u^2 + v^2 - 2u - 2v - 2uv \neq 0\}.$$

Proof. Note that the submanifolds Y_I are real slices of D , meaning that infinitesimally, there is always one degree of freedom in χ_1 -direction and one in χ_2 -direction remaining. More concretely, this means that at any point $p \in Y$ we have $T_p Y \otimes \mathbb{C} = T_p D$. In particular, up to scalar factors, the Jacobian is the same as when we consider all of D . Consequently, by the definition of D , both f and g are local diffeomorphisms on Y_I . Y is now the disjoint union of (real) submanifolds of D of dimension 2, consequently it is itself a submanifold on which f is a local diffeomorphism. Furthermore, by Lemma 6.15, f is injective on X , hence in particular on Y , hence it is a diffeomorphism.

To see that $f(Y) \subset$ the set indicated, note that $\chi_1, \chi_2 \notin i\pi\mathbb{Z}$ implies that $\operatorname{sech}\left(\frac{\chi_i}{2}\right), \tanh\left(\frac{\chi_i}{2}\right)$ ($i = 1, 2$) are finite and nonzero. □

We now have a closer look at what the Cartan subsets look like when viewed through the lens of $\psi: \tilde{G} \rightarrow \operatorname{GP}(G/Q, 4)$ and the coordinate functions u, v . But first we need a shorthand to determine $(u, v)(\psi(g))$ from the entries of x for $x \in \tilde{G}$.

Lemma 6.18. *Let $x \in \tilde{G}$ be written as a $1 + d + 1$ -block matrix as follows:*

$$x = \begin{pmatrix} A & B & C \\ D & E & F \\ G & H & I \end{pmatrix}.$$

Then

$$\begin{aligned} u(\psi(x)) &= \frac{4}{(A-I)^2 - (C-G)^2} \\ v(\psi(x)) &= \frac{(A+I)^2 - (C+G)^2}{(A-I)^2 - (C-G)^2}. \end{aligned}$$

Proof. We have

$$\psi(x) = \left(\iota(0), \infty, q \begin{pmatrix} A+C \\ D+F \\ G+I \end{pmatrix}, q \begin{pmatrix} A-C \\ D-F \\ G-I \end{pmatrix} \right),$$

So by definition of u, v we have

$$\begin{aligned} u(\psi(x)) &= 2 \frac{A^2 - C^2 - G^2 + I^2 - \eta(D, D) - \eta(F, F)}{(A-I)^2 - (C-G)^2} \\ v(\psi(x)) &= \frac{(A+I)^2 - (C+G)^2}{(A-I)^2 - (C-G)^2}. \end{aligned}$$

Note that since $g \in O(p+1, q+1)$, the first and last column of g are vectors with square length 1 and -1 , respectively, whence

$$A^2 + \eta(D, D) - G^2 = 1 \quad C^2 + \eta(F, F) - I^2 = -1,$$

so that

$$u(\psi(x)) = \frac{4}{(A-I)^2 - (C-G)^2}.$$

We can thus infer $u(\psi(x)), v(\psi(x))$ from the four corner entries A, C, G, I . \square

Corollary 6.19. \tilde{G} consists of those matrices in G with corners A, C, G, I such that

$$(A-I)^2 - (C-G)^2 \neq 0 \neq (A+I)^2 - (C+G)^2,$$

so it is indeed a dense open subset.

Proof. Let $x \in G$, then $\psi(x) = (\iota(0), \infty, g \cdot \iota(0), g \cdot \infty) = (\iota(0), \infty, \iota(a), \iota(b))$ is in general position iff none of $\eta(x, x), \eta(y, y), \eta(x-y, x-y)$ is zero. By Corollary 5.21 this is the case if $u(\psi(g)), v(\psi(g))$ are both finite and nonzero. By Lemma 6.18, this is the case iff $(A-I)^2 - (C-G)^2, (A+I)^2 - (C+G)^2 \neq 0$. \square

Lemma 6.20. Let $q > 0$. The (representative) Cartan subsets C_I from Corollary 6.10 can be labelled as follows

- (i) $C_\emptyset = C$;
- (ii) $C_{\{0\}} = \exp(\mathbb{R}(F_{0,1} + F_{d,d+1}) \oplus \mathbb{R}(F_{0,d} - F_{1,d+1}))$;
- (iii) $C_{\{2\}} = \exp(\mathbb{R}(F_{0,1} + F_{d,d+1}) \oplus \mathbb{R}(F_{0,d} - F_{1,d+1}))t_{0,\pi}$;
- (iv) $C_{\{1\}} = \exp(\mathbb{R}F_{0,1} \oplus \mathbb{R}F_{2,d+1})$;
- (v) $C_{\{1\}'} = \exp(\mathbb{R}F_{d,d+1} \oplus \mathbb{R}F_{0,d-1})$;
- (vi) $C_{\{0,1\}} = \exp(\mathbb{R}F_{0,d} \oplus \mathbb{R}F_{1,d+1})$;
- (vii) $C_{\{0,2\}} = \exp(\mathbb{R}F_{0,d} \oplus \mathbb{R}F_{1,d+1})t_{\pi/2, \pi/2}$;
- (viii) $C_{\{1,2\}} = \exp(\mathbb{R}F_{0,d} \oplus \mathbb{R}F_{1,d+1})t_{0,\pi}$.

Then, for every I there exists a homeomorphism (“parametrisation”) from $\overline{Y_I}$ to a subset of C_I that maps Y_I to (a subset of) $C_I \cap \tilde{G}$, in a way that $(\chi_1, \chi_2) \in Y_I$ is mapped to x with $(u, v)(\psi(x)) = f(\chi_1, \chi_2)$.

Proof. In the following, we use the same numbering as in Corollary 6.10. For each Cartan subset C_I we compute $(u, v) \circ \psi$ for a typical element and match this with f^{-1} of an element of $\tilde{W}\overline{Y_I}$.

(i) For $x \in C$, say $x = t_{\phi, \psi}$ we have

$$\begin{pmatrix} A & C \\ G & I \end{pmatrix} = \begin{pmatrix} \cos(\phi) & 0 \\ 0 & \cos(\psi) \end{pmatrix}.$$

By Corollary 6.19, we have $x \in \tilde{G}$ iff

$$\cos^2(\phi) - \cos^2(\psi) = -\sin(\phi + \psi) \sin(\phi - \psi) \neq 0,$$

i.e. iff $\phi \pm \psi \notin \pi\mathbb{Z}$. Assume that is the case, then

$$\begin{aligned} u(\psi(x)) &= \frac{4}{(\cos(\phi) - \cos(\psi))^2} \\ &= \csc^2\left(\frac{\phi + \psi}{2}\right) \csc^2\left(\frac{\phi - \psi}{2}\right) \\ v(\psi(x)) &= \frac{(\cos(\phi) + \cos(\psi))^2}{(\cos(\phi) - \cos(\psi))^2} \\ &= \cot^2\left(\frac{\phi + \psi}{2}\right) \cot^2\left(\frac{\phi - \psi}{2}\right), \end{aligned}$$

such that $(u(\psi(x)), v(\psi(x))) = f(i(\phi + \psi + \pi), i(\phi - \psi + \pi))$. It is therefore $\overline{Y_\emptyset}$, which we can use to parametrise C_\emptyset as follows:

$$\overline{Y_\emptyset} \ni (\chi_1, \chi_2) \mapsto \exp\left(\frac{\chi_1 + \chi_2 - 2\pi i}{2i} F_{0,1} + \frac{\chi_1 - \chi_2}{2i} F_{d,d+1}\right).$$

In particular, if $(\chi_1, \chi_2) \in Y_\emptyset$, we have $\chi_1, \chi_2 \notin \pi i\mathbb{Z}$, which corresponds to $\phi \pm \psi$ not being elements of $\pi\mathbb{Z}$, so that we obtain an element of \tilde{G} .

(ii) For $x = \exp(a(F_{0,d} - F_{1,d+1}))t_{\phi, \phi}$ we have

$$\begin{pmatrix} A & C \\ G & I \end{pmatrix} = \begin{pmatrix} \cosh(a) \cos(\phi) & \sinh(a) \sin(\phi) \\ -\sinh(a) \sin(\phi) & \cosh(a) \cos(\phi) \end{pmatrix}.$$

We thus have $x \in \tilde{G}$ iff $\sinh(a) \sin(\phi) \neq 0$ and $\cosh(a) \cos(\phi) \neq 0$, i.e. iff $a \neq 0$ and $\phi \notin \frac{1}{2}\pi\mathbb{Z}$. Assume that is the case, then

$$\begin{aligned} u(\psi(x)) &= \frac{1}{-\sinh^2(a) \sin^2(\phi)} = -\operatorname{csch}^2(a) \csc^2(\phi) \\ v(\psi(x)) &= -\operatorname{coth}^2(a) \cot^2(\phi), \end{aligned}$$

such that $(u(\psi(x)), v(\psi(x))) = f(2i\phi + i\pi, 2a + i\pi)$. It is therefore $\overline{Y_{\{0\}}}$, which we can use to parametrise $C_{\{0\}}$ as follows:

$$\overline{Y_{\{0\}}} \ni (\chi_1, \chi_2) \mapsto \exp\left(\frac{\operatorname{Re}(\chi_2)}{2}(F_{0,d} - F_{1,d+1}) + \frac{\chi_1 - i\pi}{2i}(F_{0,1} + F_{d,d+1})\right).$$

In particular, for $(\chi_1, \chi_2) \in Y_{\{0\}}$, we obtain an element of \tilde{G} .

(iii) For $x = \exp(a(F_{0,d} - F_{1,d+1}))t_{\phi, \phi + \pi}$, the corners are

$$\begin{pmatrix} A & C \\ G & I \end{pmatrix} = \begin{pmatrix} \cosh(a) \cos(\phi) & -\sinh(a) \sin(\phi) \\ -\sinh(a) \sin(\phi) & -\cosh(a) \cos(\phi) \end{pmatrix}.$$

We thus have $x \in \tilde{G}$ iff $\cosh(a) \cos(\phi), \sinh(a) \sin(\phi) \neq 0$, i.e. iff $a \neq 0$ and $\phi \notin \frac{\pi}{2}\mathbb{Z}$. Assume that is the case, then

$$\begin{aligned} u(\psi(x)) &= \frac{1}{\cosh^2(a) \cos^2(\phi)} \\ v(\psi(x)) &= -\tanh^2(a) \tan^2(\phi), \end{aligned}$$

such that $(u(\psi(x)), v(\psi(x))) = f(2a, 2i\phi)$. It is therefore $\overline{Y_{\{2\}}}$ that we can use to parametrise $C_{\{2\}}$ as follows:

$$\overline{Y_{\{2\}}} \ni (\chi_1, \chi_2) \mapsto \exp\left(\frac{\operatorname{Re}(\chi_1)}{2}(F_{0,d} - F_{1,d+1}) + \frac{\chi_2}{2i}(F_{0,1} + F_{d,d+1})\right)t_{0,\pi}.$$

In particular, for $(\chi_1, \chi_2) \in Y_{\{2\}}$, we obtain an element of \tilde{G} .

(iv) For $x = \exp(\phi F_{0,1} + aF_{2,d+1})$ we have

$$\begin{pmatrix} A & C \\ G & I \end{pmatrix} = \begin{pmatrix} \cos(\phi) & 0 \\ 0 & \cosh(a) \end{pmatrix}.$$

We thus have $x \in \tilde{G}$ iff

$$\cos^2(\phi) - \cosh^2(a) = -\sin(\phi + ia)\sin(\phi - ia) \neq 0,$$

i.e. iff $\phi \pm ia \notin \pi\mathbb{Z}$. Assume that is the case, then

$$\begin{aligned} u(\psi(x)) &= \frac{4}{(\cos(\phi) - \cosh(a))^2} = \frac{4}{(\cosh(i\phi) - \cosh(a))^2} \\ &= \frac{1}{\sinh^2\left(\frac{a+i\phi}{2}\right)\sinh^2\left(\frac{a-i\phi}{2}\right)} \\ v(\psi(x)) &= \frac{(\cos(\phi) + \cosh(a))^2}{(\cos(\phi) - \cosh(a))^2} = \coth^2\left(\frac{a+i\phi}{2}\right)\coth^2\left(\frac{a-i\phi}{2}\right) \end{aligned}$$

such that

$$(u(\psi(x)), v(\psi(x))) = f(a + i\phi + i\pi, -a + i\phi + i\pi).$$

It is therefore $\overline{Y_{\{1\}}}$ that we can use to parametrise $C_{\{1\}}$ as follows:

$$\overline{Y_{\{1\}}} \ni (\chi_1, \chi_2) \mapsto \exp\left(\frac{\chi_1 + \chi_2 - 2i\pi}{2i}F_{0,1} + \frac{\operatorname{Re}(\chi_1 - \chi_2)}{2}F_{2,d+1}\right).$$

In particular, for $(\chi_1, \chi_2) \in Y_{\{1\}}$, we obtain an element of \tilde{G} .

(v) For $x = \exp(\phi F_{d,d+1} + aF_{0,d-1})$, the four corners are

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} \cosh(a) & 0 \\ 0 & \cos(\phi) \end{pmatrix},$$

which leads to the same conditions for $x \in \tilde{G}$ and the same cross-ratios u, v as the previous case. Consequently, we call this Cartan subset $C_{\{1\}}$, and parametrise it as

$$\overline{Y_{\{1\}}} \ni (\chi_1, \chi_2) \mapsto \exp\left(\frac{\chi_1 + \chi_2 - 2i\pi}{2i}F_{d,d+1} + \frac{\operatorname{Re}(\chi_1 - \chi_2)}{2}F_{0,d-1}\right),$$

where we get an element of \tilde{G} if $(\chi_1, \chi_2) \in Y_{\{1\}}$.

(vi) For $x = \exp(aF_{0,d} + bF_{1,d+1})$ the corners are

$$\begin{pmatrix} \cosh(a) & 0 \\ 0 & \cosh(b) \end{pmatrix}.$$

We thus have $x \in \tilde{G}$ iff

$$\cosh^2(a) - \cosh^2(b) = \sinh(a+b)\sinh(a-b) \neq 0,$$

i.e. iff $a \pm b \neq 0$. Assume that is the case, then

$$\begin{aligned} u(\psi(x)) &= \frac{4}{(\cosh(a) - \cosh(b))^2} = \frac{1}{\sinh^2\left(\frac{a+b}{2}\right)\sinh^2\left(\frac{a-b}{2}\right)} \\ v(\psi(x)) &= \frac{(\cosh(a) + \cosh(b))^2}{(\cosh(a) - \cosh(b))^2} = \coth^2\left(\frac{a+b}{2}\right)\coth^2\left(\frac{a-b}{2}\right), \end{aligned}$$

such that

$$(u(\psi(x)), v(\psi(x))) = f(a + b + i\pi, a - b + i\pi).$$

It is thus $\overline{Y_{\{0,1\}}}$ that we can use to parametrise $C_{\{0,1\}}$ as follows:

$$\overline{Y_{\{0,1\}}} \ni (\chi_1, \chi_2) \mapsto \exp\left(\frac{\operatorname{Re}(\chi_1 + \chi_2)}{2} F_{0,d} + \frac{\operatorname{Re}(\chi_1 - \chi_2)}{2} F_{1,d+1}\right),$$

where we obtain an element of \tilde{G} if $(\chi_1, \chi_2) \in Y_{\{0,1\}}$.

(vii) For $x = \exp(aF_{0,d} + bF_{1,d+1})t_{\pi/2, \pi/2}$ the corners are

$$\begin{pmatrix} A & C \\ G & I \end{pmatrix} = \begin{pmatrix} 0 & \sinh(a) \\ \sinh(b) & 0 \end{pmatrix}.$$

We thus have $x \in \tilde{G}$ iff

$$\sinh^2(a) - \sinh^2(b) = \sinh(a+b)\sinh(a-b) \neq 0,$$

i.e. iff $a \pm b \neq 0$. Assume that is the case, then

$$\begin{aligned} u(\psi(x)) &= \frac{4}{-(\sinh(a) - \sinh(b))^2} = -\operatorname{csch}^2\left(\frac{a-b}{2}\right) \operatorname{sech}^2\left(\frac{a+b}{2}\right) \\ v(\psi(x)) &= \frac{-(\sinh(a) + \sinh(b))^2}{-(\sinh(a) - \sinh(b))^2} = \operatorname{coth}^2\left(\frac{a-b}{2}\right) \tanh^2\left(\frac{a+b}{2}\right), \end{aligned}$$

which shows that

$$(u(\psi(x)), v(\psi(x))) = f(a+b, a-b+i\pi).$$

It is therefore $\overline{Y_{\{0,2\}}}$ that we can use to parametrise $C_{\{0,2\}}$ as follows:

$$\overline{Y_{\{0,2\}}} \ni (\chi_1, \chi_2) \mapsto \exp\left(\frac{\operatorname{Re}(\chi_1 + \chi_2)}{2} F_{0,d} + \frac{\operatorname{Re}(\chi_1 - \chi_2)}{2} F_{1,d+1}\right)t_{\pi/2, \pi/2}.$$

In particular, we obtain an element of \tilde{G} for $(\chi_1, \chi_2) \in Y_{\{0,2\}}$.

(viii) For $x = \exp(aF_{0,d} + bF_{1,d+1})t_{\pi/2, \pi/2}$ the corners are

$$\begin{pmatrix} A & C \\ G & I \end{pmatrix} = \begin{pmatrix} \cosh(a) & 0 \\ 0 & -\cosh(b) \end{pmatrix}.$$

We thus have $x \in \tilde{G}$ iff

$$\cosh^2(a) - \cosh^2(b) = \sinh(a+b)\sinh(a-b) \neq 0,$$

i.e. iff $a \pm b \neq 0$. Assume that is the case, then

$$\begin{aligned} u(\psi(x)) &= \frac{4}{(\cosh(a) + \cosh(b))^2} = \operatorname{sech}^2\left(\frac{a+b}{2}\right) \operatorname{sech}^2\left(\frac{a-b}{2}\right) \\ v(\psi(x)) &= \frac{(\cosh(a) - \cosh(b))^2}{(\cosh(a) + \cosh(b))^2} = \tanh^2\left(\frac{a+b}{2}\right) \tanh^2\left(\frac{a-b}{2}\right). \end{aligned}$$

This shows that

$$(u(\psi(x)), v(\psi(x))) = f(a+b, a-b).$$

It is therefore $\overline{Y_{\{1,2\}}}$ that we can use to parametrise $C_{\{1,2\}}$ as follows:

$$\overline{Y_{\{1,2\}}} \ni (\chi_1, \chi_2) \mapsto \exp\left(\frac{\operatorname{Re}(\chi_1 + \chi_2)}{2} F_{0,d} + \frac{\operatorname{Re}(\chi_1 - \chi_2)}{2} F_{1,d+1}\right)t_{0, \pi}.$$

In particular, we obtain an element of \tilde{G} for $(\chi_1, \chi_2) \in Y_{\{1,2\}}$.

□

Lemma 6.21. *Let $q = 0$. There is a homeomorphism (“parametrisation”) from $\overline{Y_{\{1\}}}$ to a subset of C that maps $Y_{\{1\}}$ to (a subset of) $C \cap \tilde{G}$ in such a way that $(\chi_1, \chi_2) \in Y_{\{1\}}$ is mapped to $x \in C$ with $(u, v)(\psi(x)) = f(\chi_1, \chi_2)$.*

Proof. We use the same techniques as in the proof of Lemma 6.20. Let $x \in C$, say

$$x = \exp(aF_{d,d+1})t_\phi,$$

then the four corners of the matrix x are

$$\begin{pmatrix} \cos(\phi) & 0 \\ 0 & \cosh(a) \end{pmatrix},$$

respectively. This leads to the same cross-ratios as in 6.20(iv,v), which shows the claim and suggests the following parametrisation:

$$\overline{Y_{\{1\}}} \ni (\chi_1, \chi_2) \mapsto \exp\left(\frac{\chi_1 + \chi_2 - 2i\pi}{2i}F_{0,1} + \frac{\chi_1 - \chi_2}{2}F_{d,d+1}\right). \quad \square$$

If we compare these Cartan subsets and the associated parameter regions Y_I with the causal configurations mentioned in [20, §5.1] (see also [8, §6.6.7]), we note that our parameters χ_1, χ_2 are related to the standard bootstrap variables z, \bar{z} used there as follows:

$$z = \operatorname{sech}^2\left(\frac{\chi_1}{2}\right), \quad \bar{z} = \operatorname{sech}^2\left(\frac{\chi_2}{2}\right)$$

up to potentially exchanging z, \bar{z} . This can be seen most easily from the fact that (see e.g. [18], (31)) u, v can be written as $u = z\bar{z}, v = (1-z)(1-\bar{z})$ combined with the Lemma 6.20, where we related u and v of a certain four-point configuration to the following functions of $\chi_{1/2}$:

$$u = \operatorname{sech}^2\left(\frac{\chi_1}{2}\right) \operatorname{sech}^2\left(\frac{\chi_2}{2}\right), \quad v = \left(1 - \operatorname{sech}^2\left(\frac{\chi_1}{2}\right)\right) \left(1 - \operatorname{sech}^2\left(\frac{\chi_2}{2}\right)\right).$$

We can thus read off the claimed relations as one possible way to realise these equations. After some arithmetic we arrive at the following correspondence between Y_I and causal regions from Qiao:

I	Causal region
\emptyset	E_{tu}
$\{0\}$	U
$\{1\}$	E_{stu}
$\{2\}$	T
$\{0, 1\}$	E_{su}
$\{0, 2\}$	S
$\{1, 2\}$	E_{st} .

In particular, we see that the Euclidean case $q = 0$ can be

completely described using the assumption that z, \bar{z} are non-real complex numbers that are conjugate to each other, which is how z, \bar{z} are usually introduced in the Euclidean setting, see e.g. [20, §3.2.1].

Remark 6.22. *Note that the “parametrisations” from Lemmas 6.20, 6.21 are not surjective. This corresponds to the fact that $(u, v) \circ \psi$ is not injective on $C_I \cap \tilde{G}$. For example, for $I = \emptyset$ we have $(u, v)(\psi(t_{\phi, \psi})) = (u, v)(\psi(t_{\phi', \psi'}))$ iff*

$$(\cos(\phi) + \cos(\psi))^2, (\cos(\phi) - \cos(\psi))^2$$

equal the corresponding quantities for ϕ', ψ' , which is the case if

$$(a, b) \in \{(a', b'), (b', a'), (-a', -b'), (-b', -a')\}$$

($a = \cos(\phi), b = \cos(\psi)$, and similarly for primed quantities). Since the cosine is even, each of these four possibilities contributes four more possibilities (two each for the relative signs that the corresponding angles could have), making for (generically) sixteen elements in the same fibre.

Nevertheless, we can extend all of these parametrisations to larger domains (consisting of several \tilde{W} -translates of $\overline{Y_1}$) such that they do become surjective.

6.3. Root Spaces in the Euclidean Setting. We now apply the method of Section 4 to compute the radial part of $\Omega_{\mathfrak{g}}$ in the Euclidean setting, i.e. for $q = 0$. In that case, we have just one Cartan subset C , meaning that we essentially obtain a KAK -decomposition $G_s = HCH$ and $G_{rs} = H(C \cap G_{rs})H$. We first establish a reduced root space decomposition of $\mathfrak{g}_{\mathbb{C}}$ with respect to \mathfrak{c} .

Proposition 6.23. *The root system $\Sigma(\mathfrak{g}; \mathfrak{c})$ is of type C_2 , where the short roots have multiplicity $d - 2$, the long roots 1, and 0 has multiplicity $\frac{(d-2)(d-3)}{2} + 2$. The root system is given by*

$$\left\{ \pm\epsilon_1, \pm\epsilon_2, \pm\frac{\epsilon_1 + \epsilon_2}{2}, \pm\frac{\epsilon_1 - \epsilon_2}{2} \right\},$$

where $\epsilon_{1/2}(aF_{0,1} + bF_{d,d+1}) = a \pm ib$. Moreover we have

$$x^{\frac{\epsilon_1 \pm \epsilon_2}{2}} = \mp \exp\left(\frac{\chi_1 \pm \chi_2}{2}\right)$$

when $x \in C$ is parametrised as in Lemma 6.21. Furthermore, the elements parametrised by $Y_{\{1\}}$ (i.e. not the boundary) lie in G_{rs} .

Proof. Let $X := aF_{0,1} + bF_{d,d+1} \in \mathfrak{c}_{\mathbb{C}}$. Let $i, j = 2, \dots, d - 1$, then

$$\begin{aligned} \operatorname{ad}(X)(F_{ij}) &= 0 \\ \operatorname{ad}(X)(F_{0,i} \pm iF_{1,i}) &= \pm ai(F_{0,i} \pm iF_{1,i}) \\ \operatorname{ad}(X)(F_{i,d} \pm F_{i,d+1}) &= \mp b(F_{i,d} \pm F_{i,d+1}) \\ \operatorname{ad}(X)(F_{0,d} \pm F_{0,d+1} + iF_{1,d} \pm iF_{1,d+1}) &= (\mp b + ai)(\dots) \\ \operatorname{ad}(X)(F_{0,d} \pm F_{0,d+1} - iF_{1,d} \mp iF_{1,d+1}) &= (\mp b - ai)(\dots). \end{aligned}$$

If we define $\epsilon_1(X) := ai + b$ and $\epsilon_2(X) := ai - b$, we obtain that $\pm\frac{\epsilon_1 \pm \epsilon_2}{2}, \pm\frac{\epsilon_1 - \epsilon_2}{2}, \pm\epsilon_1, \pm\epsilon_2$ are roots with multiplicity at least $d - 2, d - 2, 1, 1$, respectively. In addition the multiplicity of 0 is at least $\frac{(d-2)(d-3)}{2} + 2$. All of these vector spaces we have already found together have dimension $\frac{(d+2)(d+1)}{2} = \dim(\mathfrak{g})$. Therefore, these are all roots, and we have identified the full root spaces. Moreover,

$$\begin{aligned} x^{\frac{\epsilon_1 \pm \epsilon_2}{2}} &= \exp\left(i\frac{\chi_1 + \chi_2 - 2\pi i}{2i}\right) = -\exp\left(\frac{\chi_1 + \chi_2}{2}\right) \\ x^{\frac{\epsilon_1 - \epsilon_2}{2}} &= \exp\left(\frac{\chi_1 - \chi_2}{2}\right). \end{aligned}$$

From these formulae we can see that $x^{2\alpha} \neq 1$ for all $\alpha \in \Sigma$ because that condition is exactly equivalent to none of $\chi_1, \chi_2, \frac{\chi_1 \pm \chi_2}{2}$ being contained in $i\pi\mathbb{Z}$, which is how we defined $D \supset Y_{\{1\}}$. By Lemma 4.8, this suffices to show $x \in G_{rs}$. \square

Lemma 6.24. *Let W be a finite-dimensional H -bimodule and let $\Psi: E^W(\tilde{G}, H) \rightarrow C^\infty(Y_{\{1\}}, W^{Z_C})$ be the map obtained by restricting to C and then parametrising as described in Lemma 6.21. Let $C_{\epsilon_i} \in \mathfrak{c}_{\mathbb{C}}$ be the element dual (with respect to B) to ϵ_i ($i = 1, 2$). Then*

$$\Psi(C_{\epsilon_i} \cdot f) = \Psi(f \cdot C_{\epsilon_i}) = \partial_{\chi_i} \Psi(f). \quad (i = 1, 2)$$

Proof. We have

$$C_{\epsilon_{1/2}} = \frac{1}{2i}F_{0,1} \pm \frac{1}{2}F_{d,d+1},$$

so that

$$\begin{aligned}
\Psi(f \cdot C_{\epsilon_1})(\chi_1, \chi_2) &= \frac{d}{dt} f \left(\exp(tC_{\epsilon_1}) \exp \left(\frac{\chi_1 + \chi_2 - 2\pi i}{2i} F_{0,1} + \frac{\chi_1 - \chi_2}{2} F_{d,d+1} \right) \right) \Big|_{t=0} \\
&= \frac{d}{dt} \Psi(f)(\chi_1 + t, \chi_2) \Big|_{t=0} \\
&= \partial_{\chi_1} \Psi(f)(\chi_1, \chi_2) \\
\Psi(f \cdot C_{\epsilon_2})(\chi_1, \chi_2) &= \frac{d}{dt} \Psi(f)(\chi_1, \chi_2 + t) \Big|_{t=0} \\
&= \partial_{\chi_2} \Psi(f)(\chi_1, \chi_2).
\end{aligned}$$

□

Corollary 6.25. *The A_α operators from Proposition 4.21 in the Euclidean case are as follows:*

$$\begin{aligned}
A_{\frac{\epsilon_1 + \epsilon_2}{2}} &= \sum_{i=2}^{d-1} F_{1,i} \otimes F^{1,i} \\
A_{\frac{\epsilon_1 - \epsilon_2}{2}} &= \sum_{i=2}^{d-1} F_{i,d} \otimes F^{i,d} \\
A_{\epsilon_{1/2}} &= \frac{1}{2} (F_{0,d+1} \mp iF_{1,d}) \otimes (F^{0,d+1} \pm iF^{1,d})
\end{aligned}$$

(where $A_{-\alpha} = A_\alpha$).

Corollary 6.26. *Let $W = \mathbb{C}$ be an H -bimodule as follows: the group M from Definition 5.11 acts trivially, and the Lie algebra \mathfrak{a} acts as:*

$$\pi_L(D_0) = \alpha, \pi_R(D_0) = \beta.$$

Then

$$\pi_L(m(A_\gamma)) = -\frac{\alpha^2}{2}, \quad \pi(A_\gamma) = -\frac{\alpha\beta}{2}, \quad \pi_R(m(A_\gamma)) = -\frac{\beta^2}{2}$$

for $\gamma \in \{\pm\epsilon_1, \pm\epsilon_2\}$, and 0 otherwise.

6.4. Root Spaces in the Lorentzian Setting. We now want to use the method of Section 4 to compute the radial part of $\Omega_{\mathfrak{g}}$, at least acting on HC_iH for some Cartan subsets. For that we need to know which Cartan subsets satisfy the condition that $\text{Ad}(t)$ act by B -orthogonal involution and weight-space dependent constants. This is trivially true for cases (i), (ii), (iv), (v), (vi) from Corollary 6.10 (i.e. the Cartan sets we called $C_\emptyset, C_{\{0\}}, C_{\{1\}}, C_{\{1\}'}, C_{\{0,1\}}$). For the remaining ones, we refer to Lemma 6.30.

We first establish the (reduced) root space decompositions with respect to our five subalgebras. We shall do this by treating C_\emptyset explicitly and showing that there is an automorphism of $\mathfrak{g}_{\mathbb{C}}$ that maps \mathfrak{c}_\emptyset to all others. For that we use the following lemma.

Lemma 6.27. *Let $\mathfrak{c}_1, \mathfrak{c}_2$ be commutative Lie subalgebras of $\mathfrak{g}_{\mathbb{C}}$. Any Lie group isomorphism $\phi: \mathfrak{c}_1 \rightarrow \mathfrak{c}_2$ that pulls back the character of the defining representation of \mathfrak{c}_2 to that of \mathfrak{c}_1 can be effected by $\text{Ad}(O(\eta, \mathbb{C}))$.*

Proof. Write $V = \mathbb{C}^{d+2}$ for the defining representations of $\mathfrak{c}_1, \mathfrak{c}_2$. Let $\lambda, \mu \in \mathfrak{c}_1^*$ and let $v \in V_\lambda, w \in V_\mu$ (weight spaces). For $Z \in \mathfrak{c}_1$ we then have

$$(\lambda + \mu)(Z)\eta(v, w) = \eta(\lambda(Z)v, w) + \eta(v, \mu(Z)w) = \eta(Zv, w) + \eta(v, Zw) = 0,$$

as $\mathfrak{g}_{\mathbb{C}}$ consists of matrices that are antisymmetric with respect to η . Since this holds for all $Z \in \mathfrak{c}_1$, we conclude that $\eta(v, w)$ can only be nonzero if $\lambda + \mu = 0$. This shows that η is a nondegenerate pairing between the λ and the $-\lambda$ weight spaces for every λ . We now chose bases

$$(v_{\mu}^1, \dots, v_{\mu}^{n_{\mu}}), \quad (u_{\lambda}^1, \dots, u_{\lambda}^{n_{\lambda}})$$

of V_{μ} (and V_{λ}) for every nonzero weight $\mu \in \mathfrak{c}_1^*$ (and $\lambda \in \mathfrak{c}_2^*$) such that $(v_{\mu}^1, \dots, v_{\mu}^{n_{\mu}})$ and $(v_{-\mu}^1, \dots, v_{-\mu}^{n_{-\mu}})$ are dual with respect to η (note that η 's nondegeneracy as well as the fact that the orthogonal complement of $V_{\mu} \oplus V_{-\mu}$ implies that $n_{\mu} = n_{-\mu}$), similarly for the u . Choose furthermore an orthonormal basis $h_1, \dots, h_n \in V$ of the 0-weight space with respect to \mathfrak{c}_1 , and $k_1, \dots, k_m \in V$ of the 0-weight space with respect to \mathfrak{c}_2 . Since the defining representation is semisimple, these define two bases of V .

Since ϕ pulls back the character of the defining representation, we find that $n_{\lambda} = n_{\phi^*(\lambda)}$ for all $\lambda \in \mathfrak{c}_2^*$, and hence in particular also that $n = m$. Define now $g \in \text{End}(V)$ by mapping $v_{\phi^*(\lambda)}^i \mapsto u_{\lambda}^i$ for every nonzero weight λ of the \mathfrak{c}_2 -module V , and by mapping $h_i \mapsto k_i$. Since the linear map g maps two bases of V to each other, it is regular. Furthermore, we have

$$\begin{aligned} \eta(g(v_{\lambda}^i), g(v_{\mu}^j)) &= \eta(u_{\phi^*(\lambda)}^i, v_{\phi^*(\mu)}^j) \\ &= \delta_{\phi^*(\lambda+\mu), 0} \delta_{i,j} = \delta_{\lambda+\phi, 0} \delta_{i,j} \\ &= \eta(v_{\lambda}^i, v_{\mu}^j) \\ \eta(g(v_{\lambda}^i), g(h_j)) &= \eta(u_{\phi^*(\lambda)}^i, k_j) \\ &= 0 = \eta(v_{\lambda}^i, h_j) \\ \eta(g(h_i), g(h_j)) &= \eta(k_i, k_j) = \delta_{i,j} \\ &= \eta(h_i, h_j), \end{aligned}$$

so that g is orthogonal, i.e. $g \in O(V)$. Let $\lambda \in \mathfrak{c}_2^*$ be a weight of the defining representation and $v \in V_{\lambda}$, then $g^{-1}v \in V_{\phi^*(\lambda)}$ and for every $X \in \mathfrak{c}_1$ we have

$$gXg^{-1}v = \lambda(\phi(X))v = \phi(X)v.$$

By linearity this is true for all $v \in V$. Since the defining representation is faithful, we thus obtain $\phi(X) = \text{Ad}(g)(X)$. \square

Proposition 6.28. *For all I , the root system $\Sigma(\mathfrak{g}:\mathfrak{c}_I)$ is of type C_2 where the short roots have multiplicity $d-2$, the long roots 1, and 0 has multiplicity $\frac{(d-2)(d-3)}{2} + 2$. In every case, we can choose the root system to be*

$$\left\{ \pm\epsilon_1, \pm\epsilon_2, \frac{\epsilon_1 \pm \epsilon_2}{2}, -\frac{\epsilon_1 \pm \epsilon_2}{2} \right\}$$

such that

$$x^{\frac{\epsilon_1 \pm \epsilon_2}{2}} = \mp \exp\left(\frac{\chi_1 \pm \chi_2}{2}\right)$$

when $x \in C_I$ is parametrised as in Lemma 6.20 and when $I \subset \{0, 1\}$. For the remaining three choices of I , when C_I is not a subgroup, we can pick

$$\begin{aligned} \{2\}, \{1, 2\}: \quad x^{\frac{\epsilon_1 \pm \epsilon_2}{2}} &= \epsilon_{\epsilon_1 \pm \epsilon_2}^I \exp\left(\frac{\chi_1 \pm \chi_2}{2}\right) \\ \{0, 2\}: \quad x^{\frac{\epsilon_1 \pm \epsilon_2}{2}} &= \mp i \epsilon_{\epsilon_1 \pm \epsilon_2}^{\{0, 2\}} \exp\left(\frac{\chi_1 \pm \chi_2}{2}\right) \end{aligned}$$

(which determines all x^α if we can choose ϵ_I to be additive, i.e. so that it extends to a character of the root lattice). Furthermore, those elements parametrised by Y_I (i.e. not the rest of the closure) lie in G_{rs} .

Proof. We are going to start with finding a root space decomposition of C_\emptyset . Note that if we manage to show the existence of root spaces as claimed with dimension at least what is claimed, we have already described a subspace of \mathfrak{g} of dimension

$$\begin{aligned} \frac{(d-2)(d-3)}{2} + 2 + 4(d-2) + 4 &= \frac{d^2 - 5d + 6 + 4 + 8d - 8}{2} \\ &= \frac{d^2 + 3d + 2}{2} = \frac{(d+2)(d+1)}{2} \\ &= \dim(\mathfrak{g}), \end{aligned}$$

so we're done.

Let $a, b \in \mathbb{C}$ and $i, j = 2, \dots, d-1$. Then we have

$$\begin{aligned} \text{ad}(aF_{0,1} + bF_{d,d+1})(F_{i,j}) &= 0 \\ \text{ad}(aF_{0,1} + bF_{d,d+1})(F_{0,i} \pm iF_{1,i}) &= \pm ai(F_{0,i} \pm iF_{1,i}) \\ \text{ad}(aF_{0,1} + bF_{d,d+1})(F_{i,d} \pm iF_{i,d+1}) &= \mp bi(F_{i,d} \pm iF_{i,d+1}) \\ \text{ad}(aF_{0,1} + bF_{d,d+1})(F_{0,d} \pm iF_{0,d+1} \mp iF_{1,d} + F_{1,d+1}) &= \mp i(a+b)(\dots) \\ \text{ad}(aF_{0,1} + bF_{d,d+1})(F_{0,d} \pm iF_{0,d+1} \pm iF_{1,d} - F_{1,d+1}) &= \pm i(a-b)(\dots), \end{aligned}$$

showing that if we define $\epsilon_1(aF_{0,1} + bF_{d,d+1}) := ai + bi$ and $\epsilon_2(aF_{0,1} + bF_{d,d+1}) := ai - bi$, we obtain the claim. Note further that $x^{\frac{\epsilon_1 \pm \epsilon_2}{2}}$ for $x \in C_\emptyset$ have the claimed form.

Now, we want to employ Lemmas 6.27, 4.2. For that note that

$$aF_{0,1} + bF_{d,d+1} = \begin{pmatrix} 0 & a & 0 & 0 & 0 \\ -a & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -b \\ 0 & 0 & 0 & b & 0 \end{pmatrix}$$

(viewed as $1 + 1 + (d-2) + 1 + 1$ -blocks) which (generically) has eigenvalues $\pm ia$ and $\pm ib$ and 0 with multiplicities 1, 1, 1, 1, $d-2$, respectively. This means that the weights are the linear maps $\pm \frac{\epsilon_1 + \epsilon_2}{2}, \pm \frac{\epsilon_1 - \epsilon_2}{2}, 0$, where ϵ_1, ϵ_2 are linearly independent. Since we are talking about finite vector spaces, all other two-dimensional commutative subalgebras of $\mathfrak{g}_{\mathbb{C}}$ whose weights of the defining representation are of this form for linearly independent ϵ_1, ϵ_2 , and have weight space dimensions 1, 1, 1, 1, $d-2$, are conjugate via $O(\eta, \mathbb{C})$ by Lemma 6.27 and therefore have the same root system and root multiplicities by Lemma 4.2.

We now check the other subalgebras. In particular, we construct $\epsilon_1, \epsilon_2 \in \mathfrak{t}_I^*$ such that the weights of the defining representation are $\pm \frac{\epsilon_1 + \epsilon_2}{2}, \pm \frac{\epsilon_1 - \epsilon_2}{2}, 0$ (with multiplicities 1, 1, 1, 1, $d-2$), and such that $x^{\frac{\epsilon_1 \pm \epsilon_2}{2}}$ have the claimed forms, for $x \in C_I$ parametrised as in Lemma 6.20.

$I = \{0\}$: A generic element X of $\mathfrak{t}_{\{0\}}$ looks like

$$X = a(F_{0,1} + F_{d,d+1}) + b(F_{0,d} - F_{1,d+1}) = \begin{pmatrix} 0 & a & 0 & -b & 0 \\ -a & 0 & 0 & 0 & b \\ 0 & 0 & 0 & 0 & 0 \\ -b & 0 & 0 & 0 & -a \\ 0 & b & 0 & a & 0 \end{pmatrix},$$

which has eigenvalues $b \pm ia, -b \pm ia$ and 0 with multiplicities 1, 1, 1, 1, $d-2$, respectively. Define $\epsilon_1(X) := 2ia$ and $\epsilon_2(X) := 2b$, then the weights of

the defining representation are as claimed and $x^{\frac{\epsilon_1 \pm \epsilon_2}{2}}$ have the desired expressions. For future reference, we note that \mathfrak{c}_\emptyset is mapped to \mathfrak{c}_I by $\text{Ad}(g)$ with

$$g = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 & 0 & -i \\ 0 & 1 & 0 & -i & 0 \\ 0 & 0 & \sqrt{2} & 0 & 0 \\ 0 & i & 0 & -1 & 0 \\ i & 0 & 0 & 0 & -1 \end{pmatrix}$$

written as a $1 + 1 + (d - 2) + 1 + 1$ block matrix.

$I = \{2\}$: The algebra \mathfrak{c}_I is the same as for $I = \{0\}$, so all these notions carry over. In order to obtain the claimed expressions for $x^{\frac{\epsilon_1 \pm \epsilon_2}{2}}$, however, we need to exchange ϵ_1, ϵ_2 .

$I = \{1\}$: A generic element X of $\mathfrak{c}_{\{1\}}$ looks like

$$X = aF_{0,1} + bF_{2,d+1} = \begin{pmatrix} 0 & a & 0 & 0 & 0 \\ -a & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -b \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -b & 0 & 0 \end{pmatrix}$$

(written as a $1 + 1 + 1 + (d - 2) + 1$ -block matrix) with eigenvalues $\pm ia, \pm b, 0$ with multiplicities $1, 1, 1, 1, d - 2$, respectively. Define $\epsilon_1(X) := ia + b$ and $\epsilon_2(X) := ia - b$, then the weights of the defining representation are as claimed and $x^{\frac{\epsilon_1 \pm \epsilon_2}{2}}$ have the desired expression. For future reference, we note that \mathfrak{c}_\emptyset is mapped to \mathfrak{c}_I by $\text{Ad}(g)$ with

$$g = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -i \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}$$

written as a $(1 + 1 + 1 + (d - 2) + 1) \times (1 + 1 + (d - 2) + 1 + 1)$ block matrix.

$\{1\}'$: A generic element X of $\mathfrak{c}_{\{1\}'}$ looks like

$$X = aF_{d,d+1} + bF_{0,d-1} = \begin{pmatrix} 0 & 0 & -b & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -b & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -a \\ 0 & 0 & 0 & a & 0 \end{pmatrix}$$

(written as a $1 + (d - 2) + 1 + 1 + 1$ -block matrix), which has eigenvalues $\pm ia, \pm b, 0$ with multiplicities $1, 1, 1, 1, d - 2$, respectively. Let $\epsilon_1(X) := ia + b, \epsilon_2(X) := ia - b$, then the weights of the defining representation are as claimed and $x^{\frac{\epsilon_1 \pm \epsilon_2}{2}}$ have the desired expressions. For future reference, we note that \mathfrak{c}_\emptyset is mapped to \mathfrak{c}_I by $\text{Ad}(g)$ with

$$g = \begin{pmatrix} 0 & 0 & 0 & 0 & -i \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & i & 0 & 0 & 0 \\ i & 0 & 0 & 0 & 0 \end{pmatrix}$$

written as a $(1 + (d - 2) + 1 + 1 + 1) \times (1 + 1 + (d - 2) + 1 + 1)$ block matrix.

$I = \{0, 1\}, \{0, 2\}, \{1, 2\}$: A generic element X of $\mathfrak{c}_{\{0,1\}}$ looks like

$$X = aF_{0,d} + bF_{1,d+1} = \begin{pmatrix} 0 & 0 & 0 & -a & 0 \\ 0 & 0 & 0 & 0 & -b \\ 0 & 0 & 0 & 0 & 0 \\ -a & 0 & 0 & 0 & 0 \\ 0 & -b & 0 & 0 & 0 \end{pmatrix}$$

(written as a $1 + 1 + (d - 2) + 1 + 1$ -block matrix), which has eigenvalues $\pm a, \pm b, 0$ with multiplicities $1, 1, 1, 1, d - 2$, respectively. Defining $\epsilon_1(X) := a + b, \epsilon_2(X) := a - b$, the weights of the defining representation are as claimed and $x^{\frac{\epsilon_1 \pm \epsilon_2}{2}}$ have the desired expressions. For future reference, we note that \mathfrak{c}_\emptyset is mapped to \mathfrak{c}_I by $\text{Ad}(g)$ with

$$g = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -i \\ 0 & 0 & 1 & 0 & 0 \\ 0 & i & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}$$

written as a $1 + 1 + (d - 2) + 1 + 1$ block matrix.

Since the adjoint representation on $\mathfrak{g}_{\mathbb{C}}$ is the second exterior power of the defining representation, all roots in $\Sigma(\mathfrak{g}; \mathfrak{c}_I)$ can be expressed using the ϵ_1, ϵ_2 we defined (in the way claimed). Like in the proof of Proposition 6.23, we note that due to the expressions of x^α we arrived at, $x^\alpha \neq x^{-\alpha}$ ($\alpha \in \Sigma(\mathfrak{g}; \mathfrak{c}_I)$) is exactly equivalent to none of $\chi_1, \chi_2, \frac{\chi_1 \pm \chi_2}{2}$ being elements of $\pi i\mathbb{Z}$, which is equivalent with $(\chi_1, \chi_2) \in D$ (and hence in Y_I instead of its closure). \square

Having now established the root systems for the different Cartan subsets, we can now work out the corresponding root spaces.

Proposition 6.29. *The weight spaces of \mathfrak{g} with respect to \mathfrak{c}_I are spanned by*

$I = \emptyset$:

$$\begin{aligned} & 0: F_{i,j} \quad (i, j = 1, \dots, d-1) \\ & \pm \frac{\epsilon_1 + \epsilon_2}{2}: F_{0,i} \pm iF_{1,i} \quad (i = 1, \dots, d-1) \\ & \pm \frac{\epsilon_1 - \epsilon_2}{2}: F_{i,d} \mp iF_{i,d+1} \quad (i = 1, \dots, d-1) \\ & \pm \epsilon_1: F_{0,d} + F_{1,d+1} \pm i(-F_{0,d+1} + F_{1,d}) \\ & \pm \epsilon_2: F_{0,d} - F_{1,d+1} \pm i(F_{0,d+1} + F_{1,d}). \end{aligned}$$

$I = \{0\}, \{2\}$: ($\{2\}$ with ϵ_1, ϵ_2 exchanged)

$$\begin{aligned} & 0: F_{i,j} \quad (i, j = 1, \dots, d-1) \\ & \pm \frac{\epsilon_1 + \epsilon_2}{2}: F_{0,i} - iF_{i,d+1} \pm i(F_{1,i} - iF_{i,d}) \quad (i = 2, \dots, d-1) \\ & \pm \frac{\epsilon_1 - \epsilon_2}{2}: F_{1,i} + iF_{i,d} \mp i(F_{0,i} + iF_{i,d+1}) \quad (i = 2, \dots, d-1) \\ & \pm \epsilon_1: F_{0,d} + F_{1,d+1} \pm i(F_{0,d+1} + F_{1,d}) \\ & \pm \epsilon_2: F_{0,d+1} + F_{1,d} \pm (F_{0,1} - F_{d,d+1}). \end{aligned}$$

$I = \{1\}$:

$$\begin{aligned}
& 0:F_{i,j} \quad (i, j = 3, \dots, d) \\
& \pm \frac{\epsilon_1 + \epsilon_2}{2}:F_{0,i} \pm iF_{1,i} \quad (i = 3, \dots, d) \\
& \pm \frac{\epsilon_1 - \epsilon_2}{2}:F_{i,d+1} \pm F_{2,i} \quad (i = 3, \dots, d) \\
& \quad \pm \epsilon_1:F_{0,d+1} - iF_{1,2} \pm (-F_{0,2} + iF_{1,d+1}) \\
& \quad \pm \epsilon_2:F_{0,d+1} + iF_{1,2} \pm (F_{0,2} + iF_{1,d+1}).
\end{aligned}$$

$I = \{1\}'$:

$$\begin{aligned}
& 0:F_{i,j} \quad (i, j = 1, \dots, d-2) \\
& \pm \frac{\epsilon_1 + \epsilon_2}{2}:F_{i,d} \mp iF_{i,d+1} \quad (i = 1, \dots, d-2) \\
& \pm \frac{\epsilon_1 - \epsilon_2}{2}:F_{0,i} \pm F_{i,d-1} \quad (i = 1, \dots, d-2) \\
& \quad \pm \epsilon_1:F_{0,d} \mp F_{d-1,d} + iF_{d-1,d+1} \mp iF_{0,d+1} \\
& \quad \pm \epsilon_2:F_{0,d} \pm F_{d-1,d} - iF_{d-1,d+1} \mp iF_{0,d+1}.
\end{aligned}$$

$I = \{0, 1\}, \{0, 2\}, \{1, 2\}$:

$$\begin{aligned}
& 0:F_{i,j} \quad (i, j = 2, \dots, d-1) \\
& \pm \frac{\epsilon_1 + \epsilon_2}{2}:F_{0,i} \pm F_{i,d} \quad (i = 2, \dots, d-1) \\
& \pm \frac{\epsilon_1 - \epsilon_2}{2}:F_{1,i} \pm F_{i,d+1} \quad (i = 2, \dots, d-1) \\
& \quad \pm \epsilon_1:F_{0,1} + F_{d,d+1} \mp (F_{0,d+1} - F_{1,d}) \\
& \quad \pm \epsilon_2:F_{0,1} - F_{d,d+1} \pm (F_{0,d+1} + F_{1,d}).
\end{aligned}$$

Proof. We shall use the weight spaces of \mathfrak{c}_θ from the proof of Proposition 6.28 and apply $\text{Ad}(g_I)$ for the $g_I \in O(p+1, q+1; \mathbb{C})$ that were found in that same proof, which satisfy $\text{Ad}(g_I)(\mathfrak{c}_{\theta, \mathbb{C}}) = \mathfrak{c}_{I, \mathbb{C}}$. A similar proof as in the proof of in Proposition 4.21 can be employed to show that $\text{Ad}(g_I)(\mathfrak{g}_\alpha) = \mathfrak{g}_{\text{Ad}^*(g_I)(\alpha)}$ for $\alpha \in \Sigma(\mathfrak{g}:\mathfrak{c}_\theta)$. \square

With these weight space decompositions in hand, we can now ascertain that the Cartan subsets $C_{\{2\}}, C_{\{0,2\}}, C_{\{1,2\}}$ indeed satisfy the technical condition of Section 4.2 and that we can even choose ϵ^I to extend to group homomorphisms of the root lattice (so that $\alpha \mapsto x^\alpha$ also extends to a group homomorphism).

Lemma 6.30. *For $I = \{2\}, \{0, 2\}, \{1, 2\}$, we can write*

$$\text{Ad}(t_I)|_{\mathfrak{g}_\alpha} = \epsilon_\alpha^I \phi_I$$

for $\alpha \in \Sigma(\mathfrak{g}:\mathfrak{c}_I)$ such that C_I satisfies the condition at the beginning of Section 4.2. Furthermore, we can choose ϵ^I such that for $\alpha, \beta, \alpha + \beta \in \Sigma(\mathfrak{g}:\mathfrak{c}_I)$ we have $\epsilon_\alpha^I \epsilon_\beta^I = \epsilon_{\alpha+\beta}^I$. In particular, we can choose

$$\begin{aligned}
\epsilon_{\frac{\epsilon_1 \pm \epsilon_2}{2}}^{\{2\}} &= \mp 1 \\
\epsilon_{\frac{\epsilon_1 \pm \epsilon_2}{2}}^{\{0,2\}} &= \pm i \\
\epsilon_{\frac{\epsilon_1 \pm \epsilon_2}{2}}^{\{1,2\}} &= 1.
\end{aligned}$$

Proof. We begin with the case of $I = \{0, 2\}$ or $\{1, 2\}$. We construct B_σ -orthonormal bases of the weight spaces as is done in the proof of Theorem 4.22:

$$\begin{aligned} E_{\pm \frac{\epsilon_1 + \epsilon_2}{2}, i} &= \frac{1}{2\sqrt{\eta_{i,i}}} (\pm F_{0,i} + F_{i,d}) \\ E_{\pm \frac{\epsilon_1 - \epsilon_2}{2}, i} &= \frac{1}{2i\sqrt{\eta_{i,i}}} (F_{1,i} \pm F_{i,d+1}) \\ E_{\pm \epsilon_1} &= \frac{1}{2\sqrt{2}} (\mp (F_{0,1} + F_{d,d+1}) + F_{0,d+1} - F_{1,d}) \\ E_{\pm \epsilon_2} &= \frac{1}{2\sqrt{2}} (\pm (F_{0,1} - F_{d,d+1}) + F_{0,d+1} + F_{1,d}). \end{aligned}$$

These bases satisfy $\sigma(E_{\alpha,i}) = E_{-\alpha,i}$. For $I = \{1, 2\}$, the action of $\text{Ad}(t_I)$ on these basis elements is as follows:

$$\begin{aligned} E_{\pm \frac{\epsilon_1 + \epsilon_2}{2}, i} &\mapsto -E_{\mp \frac{\epsilon_1 + \epsilon_2}{2}, i} \\ E_{\pm \frac{\epsilon_1 - \epsilon_2}{2}, i} &\mapsto E_{\mp \frac{\epsilon_1 - \epsilon_2}{2}, i} \\ E_{\pm \epsilon_1} &\mapsto -E_{\mp \epsilon_1} \\ E_{\pm \epsilon_2} &\mapsto -E_{\mp \epsilon_2}. \end{aligned}$$

Having $\phi_{\{1,2\}}$ permute the basis (and be the identity map on $\mathfrak{m}_{I,\mathbb{C}}$, we can write this action in the desired way such that $\epsilon^{\{1,2\}}$ has the claimed form. For $I = \{0, 2\}$, the action of $\text{Ad}(t_I)$ on these basis elements is as follows:

$$\begin{aligned} \text{Ad}(t)(E_{\pm \frac{\epsilon_1 + \epsilon_2}{2}, i}) &= \mp i E_{\mp \frac{\epsilon_1 - \epsilon_2}{2}, i} \\ \text{Ad}(t)(E_{\pm \frac{\epsilon_1 - \epsilon_2}{2}, i}) &= \pm i E_{\mp \frac{\epsilon_1 + \epsilon_2}{2}, i} \\ \text{Ad}(t)(E_{\pm \epsilon_1}) &= -E_{\mp \epsilon_1} \\ \text{Ad}(t)(E_{\pm \epsilon_2}) &= E_{\pm \epsilon_2}. \end{aligned}$$

Having $\phi_{\{0,2\}}$ map

$$E_{\pm \frac{\epsilon_1 + \epsilon_2}{2}, i} \mapsto -E_{\mp \frac{\epsilon_1 - \epsilon_2}{2}, i}, \quad E_{\pm \frac{\epsilon_1 - \epsilon_2}{2}, i} \mapsto -E_{\mp \frac{\epsilon_1 + \epsilon_2}{2}, i}, \quad E_{\pm \epsilon_1} \mapsto -E_{\mp \epsilon_1}, \quad E_{\pm \epsilon_2} \mapsto -E_{\pm \epsilon_2},$$

and be the identity on $\mathfrak{m}_{I,\mathbb{C}}$, we obtain a decomposition of $\text{Ad}(t_I)$ with ϵ^I having the desired form.

For $I = \{2\}$ we note that $t_I^2 = 1$, so that $\text{Ad}(t_I)$ is a B -orthogonal involution. Furthermore, we have

$$\sigma \circ \text{Ad}(t_I) \circ \sigma = \text{Ad}(\sigma(t_I)) = \text{Ad}(t_I^{-1}) = \text{Ad}(t_I),$$

so that $\text{Ad}(t)$ also commutes with σ . Consequently, we can choose $\epsilon_\alpha^I = 1$ for all roots α and $\phi_I = \text{Ad}(t_I)$. \square

Corollary 6.31. *For $I = \{2\}, \{0, 2\}$ and $x \in C_I$ parametrised as in Lemma 6.20 we have*

$$x^{\frac{\epsilon_1 \pm \epsilon_2}{2}} = \exp\left(\frac{\chi_1 \pm \chi_2}{2}\right).$$

In all other cases we have

$$x^{\frac{\epsilon_1 \pm \epsilon_2}{2}} = \mp \exp\left(\frac{\chi_1 \pm \chi_2}{2}\right).$$

In every case, the function

$$\Sigma(\mathfrak{g}; \mathfrak{c}_I) \ni \alpha \mapsto x^\alpha$$

can be extended to a group homomorphism $\mathbb{Z}\Sigma \rightarrow \mathbb{C}^\times$. Furthermore, we always have

$$\begin{aligned} \coth_{\frac{\epsilon_1 \pm \epsilon_2}{2}}(x) &= \coth\left(\frac{\chi_1 \pm \chi_2}{2}\right) \\ \coth_{\epsilon_{1/2}}(x) &= \coth(\chi_{1/2}). \end{aligned}$$

Proof. Follows from Proposition 6.28 and Lemma 6.30. \square

Proposition 6.32. *The operators A_α look as follows:*

$I = \emptyset$:

$$\begin{aligned} A_{\frac{\epsilon_1 + \epsilon_2}{2}} &= \sum_{i=2}^{d-1} F_{1,i} \otimes F^{1,i}, & A_{\frac{\epsilon_1 - \epsilon_2}{2}} &= \sum_{i=2}^{d-1} F_{i,d} \otimes F^{i,d}, \\ A_{\epsilon_{1/2}} &= \frac{1}{2}(F_{0,d+1} \mp F_{1,d}) \otimes (F^{0,d+1} \mp F^{1,d}) \end{aligned}$$

$I = \{0\}, \{2\}$: (for $i = \{2\}$ we exchange ϵ_1, ϵ_2)

$$\begin{aligned} A_{\frac{\epsilon_1 \pm \epsilon_2}{2}} &= \frac{1}{2} \sum_{i=2}^{d-1} (F_{1,i} \mp iF_{i,d}) \otimes (F^{1,i} \pm iF^{i,d}) \\ A_{\epsilon_{1/2}} &= \frac{1}{2}(F_{0,d+1} \mp F_{1,d}) \otimes (F^{0,d+1} \mp F^{1,d}) \end{aligned}$$

$I = \{1\}$:

$$\begin{aligned} A_{\frac{\epsilon_1 + \epsilon_2}{2}} &= \sum_{i=3}^d F_{1,i} \otimes F^{1,i} & A_{\frac{\epsilon_1 - \epsilon_2}{2}} &= \sum_{i=3}^d F_{2,i} \otimes F^{2,i} \\ A_{\epsilon_{1/2}} &= \frac{1}{2}(F_{0,d+1} \mp iF_{1,2}) \otimes (F^{0,d+1} \pm iF^{1,2}) \end{aligned}$$

$I = \{1\}'$:

$$\begin{aligned} A_{\frac{\epsilon_1 + \epsilon_2}{2}} &= \sum_{i=1}^{d-2} F_{i,d} \otimes F^{i,d} & A_{\frac{\epsilon_1 - \epsilon_2}{2}} &= \sum_{i=1}^{d-2} F_{i,d-1} \otimes F^{i,d-1} \\ A_{\epsilon_{1/2}} &= \frac{1}{2}(F_{0,d+1} - iF_{d-1,d}) \otimes (F^{0,d+1} + iF^{d-1,d}) \end{aligned}$$

$I = \{0, 1\}, \{0, 2\}, \{1, 2\}$:

$$\begin{aligned} A_{\frac{\epsilon_1 + \epsilon_2}{2}} &= \sum_{i=2}^{d-1} F_{i,d} \otimes F^{i,d} & A_{\frac{\epsilon_1 - \epsilon_2}{2}} &= \sum_{i=2}^{d-1} F_{1,i} \otimes F^{1,i} \\ A_{\epsilon_{1/2}} &= \frac{1}{2}(F_{0,d+1} \mp F_{1,d}) \otimes (F^{0,d+1} \mp F^{1,d}). \end{aligned}$$

Corollary 6.33. *Let W be scalar as in Corollary 6.26, then*

$$\pi_{\mathbb{L}}(m(A_\gamma)) = -\frac{\alpha^2}{2}, \quad \pi(A_\gamma) = -\frac{\alpha\beta}{2}, \quad \pi_{\mathbb{R}}(m(A_\gamma)) = -\frac{\beta^2}{2}$$

if $\gamma \in \{\pm\epsilon_1, \pm\epsilon_2\}$, and zero otherwise.

Lemma 6.34. *Let W be a finite-dimensional H -bimodule, let I be the index of a Cartan subset, and let $\Psi_I: E^W(\tilde{G}, H) \rightarrow C^\infty(Y_I, W^{Z_{C_I}})$ be the map obtained by restricting to C_I and then parametrising as described in Lemma 6.20. Let $C_{\epsilon_i} \in \mathfrak{c}_{I,\mathbb{C}}$ be the dual element (with respect to B) to $\epsilon_i \in \mathfrak{c}_{I,\mathbb{C}}^*$. Then*

$$\Psi_I(\text{Ad}(t_I^{-1})(C_{\epsilon_i}) \cdot f) = \Psi_I(f \cdot C_{\epsilon_i}) = \partial_{\chi_i} \Psi_I(f)$$

for $i = 1, 2$.

Proof. The elements C_{ϵ_i} ($i = 1, 2$) are

$$\begin{aligned}
I = \emptyset: & \quad C_{\epsilon_1} = \frac{F_{0,1} + F_{d,d+1}}{2i}, & C_{\epsilon_2} &= \frac{F_{0,1} - F_{d,d+1}}{2i} \\
I = \{0\}: & \quad C_{\epsilon_1} = \frac{F_{0,1} + F_{d,d+1}}{2i}, & C_{\epsilon_2} &= \frac{F_{0,d} - F_{1,d+1}}{2} \\
I = \{2\}: & \quad C_{\epsilon_1} = \frac{F_{0,d} - F_{1,d+1}}{2}, & C_{\epsilon_2} &= \frac{F_{0,1} + F_{d,d+1}}{2i} \\
I = \{1\}: & \quad C_{\epsilon_1} = \frac{F_{0,1} + iF_{2,d+1}}{2i}, & C_{\epsilon_2} &= \frac{F_{0,1} - iF_{2,d+1}}{2i} \\
I = \{1\}': & \quad C_{\epsilon_1} = \frac{F_{d,d+1} + iF_{0,d-1}}{2i}, & C_{\epsilon_2} &= \frac{F_{d,d+1} - iF_{0,d-1}}{2i} \\
I = \{0, 1\}, \{0, 2\}, \{1, 2\}: & \quad C_{\epsilon_1} = \frac{F_{0,d} + F_{1,d+1}}{2}, & C_{\epsilon_2} &= \frac{F_{0,d} - F_{1,d+1}}{2}.
\end{aligned}$$

Consulting the parametrisations from Lemma 6.20 and ignoring issues of complexification, we obtain

$$\begin{aligned}
\Psi_I(t_I^{-1} \exp(sC_{\epsilon_1}) t_I \cdot f)(\chi_1, \chi_2) &= \Psi_I(f \cdot \exp(sC_{\epsilon_1}))(\chi_1, \chi_2) = \Psi_I(f)(\chi_1 + s, \chi_2) \\
\Psi_I(t_I^{-1} \exp(sC_{\epsilon_2}) t_I \cdot f)(\chi_1, \chi_2) &= \Psi_I(f \cdot \exp(sC_{\epsilon_2}))(\chi_1, \chi_2) = \Psi_I(f)(\chi_1, \chi_2 + s).
\end{aligned}$$

Taking the derivative with respect to s at $s = 0$ yields the claimed equality. \square

6.5. Matrix Case. If we combine Lemmas 6.24, 6.34, 6.21, 6.20 and Theorem 4.22, we obtain that

$$\begin{aligned}
\Psi_I(f \cdot \Omega_{\mathfrak{g}}) &= \Psi_I(\Omega_{\mathfrak{g}} \cdot f) = L(k)\Psi_I(f) + \pi_L(\Omega_{\mathfrak{m}_I})\Psi_I(f) \\
&+ \sum_{\gamma \in \Sigma(\mathfrak{g}; \mathfrak{c}_I)} \frac{\pi_L(m(A_\gamma)) + \pi_R(m(A_{\text{Ad}^*(t_I)(\gamma)})) + 2\pi(1 \otimes \phi_I)(A_\gamma)}{4 \sinh_\gamma^2} \Psi_I(f) \\
(6.1) \quad &- \sum_{\gamma \in \Sigma(\mathfrak{g}; \mathfrak{c}_I)} \frac{\pi(1 \otimes \phi_I)(A_\gamma)}{4 \sinh_{\gamma/2}^2} \Psi_I(f),
\end{aligned}$$

where

$$L(k) = \sum_{i=1}^2 \partial_{\xi_i} + \sum_{\alpha \in R^+} k_\alpha \frac{1 + e^{-\alpha}}{1 - e^{-\alpha}} \partial_\alpha$$

is the Laplacian from [11, Proposition 1.2.3] for the root system $R = 2\Sigma(\mathfrak{g}; \mathfrak{c}_I)$ of type C_2 and the multiplicities $k_{2\gamma} := \frac{n_\gamma}{2}$, with ξ_1, ξ_2 being an orthonormal basis of the underlying Euclidean space. In our case that Euclidean space is the real span of ϵ_1, ϵ_2 .

This equality holds both on a formal level where we associate our x^γ with their e^γ , but also on the level of functions, since by Proposition 6.23 and Corollary 6.31 the parametrisations from Lemmas 6.21 and 6.20 interact in such a way with our definitions of ϵ_1, ϵ_2 that

$$x^{\frac{\epsilon_1 \pm \epsilon_2}{2}} = \exp\left(\frac{\chi_1 \pm \chi_2}{2}\right)$$

up to signs, and these signs turn out not to play a role for \coth_γ ($\gamma \in \Sigma(\mathfrak{g}; \mathfrak{c}_I)$).

More concretely, the multiplicity vector is

$$k_{\pm 2\epsilon_{1/2}} = \frac{1}{2}, \quad k_{\pm \epsilon_1 \pm \epsilon_2} = \frac{d-2}{2}.$$

Later it will be useful to embed $2\Sigma(\mathfrak{g}; \mathfrak{c}_I)$ into a larger root system R of type BC_2 , by adding $\pm\epsilon_{1/2}$ and setting their k to zero. If we set

$$(6.2) \quad \begin{aligned} K_{2\gamma} &= \frac{\pi_L(m(A_\gamma)) + \pi_R(m(A_{\text{Ad}^*(t_I)(\gamma)})) + 2\pi(1 \otimes \phi_I)(A_\gamma)}{4}, \\ L_\gamma &= -\frac{\pi(1 \otimes \phi_I)(A_\gamma)}{4} \in \text{End}(W^{Z_{C_I}}) \end{aligned}$$

for $\gamma \in \Sigma(\mathfrak{g}; \mathfrak{c}_I)$ we can write the above as

$$\begin{aligned} \Psi_I(f \cdot \Omega_{\mathfrak{g}}) &= \Psi_I(\Omega_{\mathfrak{g}} \cdot f) = L(k)\Psi_I(f) + \pi_L(\Omega_{\mathfrak{m}_I})\Psi_I(f) \\ &+ \sum_{\gamma \in \Sigma(\mathfrak{g}; \mathfrak{c}_I)} \left(\frac{K_{2\gamma}}{\sinh^2 \gamma} + \frac{L_\gamma}{\sinh^2 \gamma/2} \right) \Psi_I(f). \end{aligned}$$

We now show that this matches [16, Equation 1.1]. For this, we consider the case $I = \{1\}$ and $q = 1$ and apply the permutation $(d+1, d, \dots, 1)$ to our indices to match conventions. In the notation of [16, Section 1.1] we then compute the matrices $K_{2\gamma}, L_\gamma$:

$$\begin{aligned} K_{2\epsilon_{1/2}} &= -\frac{1}{8}(D_{\mp}^{\prime 2} + D_{\pm}^2 - 2D'_{\pm}D_{\pm}) \\ K_{\epsilon_1 + \epsilon_2} &= \frac{1}{4}(M'_{2a}M'_{2a} + M_{2a}M_{2a} + 2M'_{2a}M_{2a}) \\ K_{\epsilon_1 - \epsilon_2} &= \frac{1}{4}(M'_{3a}M'_{3a} + M_{3a}M_{3a} + 2M'_{3a}M_{3a}) \\ L_{\epsilon_{1/2}} &= \frac{D'_{\mp}D_{\mp}}{8} \\ L_{\frac{\epsilon_1 + \epsilon_2}{2}} &= -\frac{M'_{2a}M_{2a}}{4} \\ L_{\frac{\epsilon_1 - \epsilon_2}{2}} &= -\frac{M'_{3a}M_{3a}}{4}. \end{aligned}$$

In particular, if we choose a parametrisation by t_1, t_2 in such a way that

$$x^{\frac{\epsilon_1 \pm \epsilon_2}{2}} = \exp(\pm t_{1/2}),$$

the sum

$$\sum_{\gamma \in \Sigma(\mathfrak{g}; \mathfrak{c}_I)} \left(\frac{K_{2\gamma}}{\sinh^2 \gamma} + \frac{L_\gamma}{\sinh^2 \gamma/2} \right)$$

corresponds to the matrix-valued function

$$\begin{aligned} &- \frac{D_-^{\prime 2} + D_-^2 - 2 \cosh(t_1 - t_2) D'_- D_-}{4 \sinh^2(t_1 - t_2)} - (- \leftrightarrow +) \\ &+ \frac{M'_{2a}M'_{2a} + M_{2a}M_{2a} - 2 \cosh(t_1) M'_{2a}M_{2a}}{2 \sinh^2(t_1)} \\ &+ \frac{M'_{3a}M'_{3a} + M_{3a}M_{3a} - 2 \cosh(t_2) M'_{3a}M_{3a}}{2 \sinh^2(t_2)}. \end{aligned}$$

Consequently, we have

$$\begin{aligned}
H^{\rho_l, \rho_r} &= 2 \sum_{\gamma \in \Sigma(\mathfrak{g}; \mathfrak{c}_I)} \left(\frac{K_{2\gamma}}{\sinh_\gamma^2} + \frac{L_\gamma}{\sinh_{\gamma/2}^2} \right) \\
&\quad + \partial_{t_1}^2 + \partial_{t_2}^2 - \frac{(d-2)(d-4)}{4} (\operatorname{csch}^2(t_1) + \operatorname{csch}^2(t_2)) + \frac{1}{2} (\operatorname{csch}^2(t_1 + t_2) + \operatorname{csch}^2(t_1 - t_2)) \\
(6.3) \quad &\quad - \frac{d^2 - 2d + 2}{2} - \frac{1}{2} L^{ab} L_{ab}.
\end{aligned}$$

Note that $(L_{ab})_{a < b}$ and $(\frac{1}{2} L^{ab})_{a < b}$ are dual bases of \mathfrak{m}_I , so that $\Omega_{\mathfrak{m}_I}$ is given by $-\frac{1}{4} L^{ab} L_{ab}$. Note furthermore that

$$\rho(k) = \frac{1}{2} ((d-1)\epsilon_1 + \epsilon_2),$$

whence

$$\|\rho(k)\|^2 = \frac{d^2 - 2d + 2}{4}.$$

Note lastly that the middle line of (6.3) is recognisable as twice the Hamiltonian form (see e.g. [11, Equation 2.1.9]) of the modified Laplacian $L(k) + \|\rho(k)\|^2$. This shows that

$$\begin{aligned}
H^{\rho_l, \rho_k} &= 2 \sum_{\gamma \in \Sigma(\mathfrak{g}; \mathfrak{c}_I)} \left(\frac{K_{2\gamma}}{\sinh_\gamma^2} + \frac{L_\gamma}{\sinh_{\gamma/2}^2} \right) \\
&\quad + 2\delta(k)^{1/2} L(k) \delta(k)^{-1/2} + 2\pi_L(\Omega_{\mathfrak{m}_I}) \\
&= 2\delta(k)^{1/2} \left(L(k) + \pi_L(\Omega_{\mathfrak{m}_I}) + \sum_{\gamma \in \Sigma(\mathfrak{g}; \mathfrak{c}_I)} \left(\frac{K_{2\gamma}}{\sinh_\gamma^2} + \frac{L_\gamma}{\sinh_{\gamma/2}^2} \right) \right) \delta(k)^{-1/2},
\end{aligned}$$

which is indeed twice the Hamiltonian form of the operator from (6.2).

6.6. Scalar Case. We now assume that W is a scalar bimodule as in the Corollaries 6.33, 6.26:

$$\pi_L(F_{0,d+1}) = \alpha, \quad \pi_R(F_{0,d+1}) = \beta$$

and all elements of M being mapped to 1. From Corollaries 6.26, 6.33, we find that

$$\begin{aligned}
\pi_L(m(A_\gamma)) &= -\frac{\alpha^2}{2} \\
\pi_R(m(A_{\operatorname{Ad}^*(t_I)(\gamma)})) &= -\frac{\beta^2}{2} \\
\pi(1 \otimes \phi_I)(A_\gamma) &= -\frac{\alpha\beta}{2}
\end{aligned}$$

for γ a long root, i.e. $\pm\epsilon_1, \pm\epsilon_2$, and zero in all other cases, for the Euclidean and all C_I (for $I \subseteq \{0, 1\}$ with and without prime, and $I = \{1, 2\}$) of the Lorentzian case. For the remaining two Lorentzian cases $I = \{2\}, \{0, 2\}$, the first two scalars are the same, but the third changes sign. Similarly, $\pi_L(\Omega_{\mathfrak{m}'}) = 0$. We thus obtain

$$K_{2\gamma} = -\frac{(\alpha \pm \beta)^2}{8}, \quad L_\gamma = \pm \frac{\alpha\beta}{8}$$

for $\gamma \in \Sigma(\mathfrak{g}; \mathfrak{c}_I)$ a long root and 0 otherwise, with “−” if $I = \{2\}$ or $\{0, 2\}$ and “+” otherwise.

Lemma 6.35. *If we write $e^{\frac{\epsilon_i}{2}}$ for the function mapping x to $\exp(\frac{\chi_i}{2})$, and we set*

$$\begin{aligned} l_{\pm\epsilon_1}^2 &= l_{\pm\epsilon_2}^2 = -\alpha\beta \\ l_{\pm(\epsilon_1+\epsilon_2)}^2 &= l_{\pm(\epsilon_1-\epsilon_2)}^2 = 0 \\ l_{\pm 2\epsilon_1}^2 &= l_{\pm 2\epsilon_2}^2 = -\left(\frac{\alpha-\beta}{2}\right)^2, \end{aligned}$$

we obtain

$$\Psi_I(\Omega_{\mathfrak{g}} \cdot f) = \Psi_I(f \cdot \Omega_{\mathfrak{g}}) = \left(L(k) + \sum_{\gamma \in R^+} \frac{l_{\gamma} B^*(\gamma, \gamma)}{(e^{\gamma/2} - e^{-\gamma/2})^2} \right) \Psi(f)$$

for all $f \in E^W(G, H)$, where R is now a root system of type BC_2 containing $2\Sigma(\mathfrak{g}; \mathfrak{c}_I)$ as its immultipliable roots.

Proof. By Corollary 6.31 and previous observations as to what $\pi(1 \otimes \phi_I)(A_{\gamma})$ is for the scalar representation, this is already the case for $I = \{2\}$ and $\{0, 2\}$. For the others we have

$$\sinh^2_{\frac{\epsilon_i}{2}}(x) = x^{\epsilon_i} + x^{-\epsilon_i} - 2 = -\exp(\chi_i) - \exp(-\chi_i) - 2 = -\cosh\left(\frac{\chi_i}{2}\right)^2,$$

so that the constant term reads

$$-\frac{(\alpha+\beta)^2}{4 \sinh^2(\chi_1)} - \frac{\alpha\beta}{4 \cosh^2\left(\frac{\chi_1}{2}\right)} + 1 \leftrightarrow 2.$$

Note that $\operatorname{csch}^2(2x) = \frac{\operatorname{csch}^2(x) - \operatorname{sech}^2(x)}{4}$, so that we have

$$\begin{aligned} -\frac{(\alpha+\beta)^2}{4 \sinh^2(\chi_i)} - \frac{\alpha\beta}{4 \cosh^2\left(\frac{\chi_i}{2}\right)} &= -\frac{(\alpha-\beta)^2}{4 \sinh^2(\chi_i)} - \alpha\beta \operatorname{csch}^2(\chi_i) - \frac{\alpha\beta}{4 \cosh^2\left(\frac{\chi_i}{2}\right)} \\ &= -\frac{(\alpha-\beta)^2}{4 \sinh^2(\chi_i)} - \frac{\alpha\beta}{4 \sinh^2\left(\frac{\chi_i}{2}\right)}. \end{aligned}$$

Consequently, the claim also holds for the other choices of I . \square

By [11, Corollary 2.1.2], we can absorb these last terms into $L(k)$ by conjugating as follows:

$$L(k) + \sum_{\alpha \in R^+} \frac{l_{\alpha}^2 B^*(\alpha, \alpha)}{(e^{\alpha/2} - e^{-\alpha/2})^2} = \delta(L(m) + \|\rho(m)\|^2 - \|\rho(k)\|^2) \delta^{-1}$$

where

$$\begin{aligned} m_{\pm\epsilon_1} &= m_{\pm\epsilon_2} = \alpha \\ m_{\pm(\epsilon_1+\epsilon_2)} &= m_{\pm(\epsilon_1-\epsilon_2)} = \frac{d-2}{2} \\ m_{\pm 2\epsilon_1} &= m_{\pm 2\epsilon_2} = \frac{1-\alpha+\beta}{2}, \end{aligned}$$

where

$$\|\rho(k)\|^2 = \frac{d^2 - 2d + 2}{4}, \quad \|\rho(m)\|^2 = \frac{(d+\beta-1)^2 + (d+1)^2}{4},$$

and where

$$\delta = \prod_{\gamma \in R^+} \left(e^{\gamma/2} - e^{-\gamma/2} \right)^{(m-k)\gamma},$$

which up to constants and in the parametrisation of Lemma 6.20 equals

$$\cosh^{-\frac{\alpha+\beta}{2}}\left(\frac{\chi_1}{2}\right) \cosh^{-\frac{\alpha+\beta}{2}}\left(\frac{\chi_2}{2}\right) \sinh^{\frac{\alpha+\beta}{2}}\left(\frac{\chi_1}{2}\right) \sinh^{\frac{\alpha+\beta}{2}}\left(\frac{\chi_2}{2}\right).$$

In conclusion, we obtain

$$(6.4) \quad \Psi_I(\Omega_{\mathfrak{g}} \cdot f) = \Psi_I(f \cdot \Omega_{\mathfrak{g}}) = \delta \left(L(m) + \frac{\beta(\beta + d)}{2} \right) \delta^{-1} \Psi_I(f),$$

which is exactly the observation in [10], thus explaining the appearance of a BC_2 root system and multiplicity vectors that contain both information on multiplicities and on the H -bimodule W . Notice that the classical theory of spherical functions for a non-compact Riemannian symmetric space from [11, Chapter 5] would have led one to expect a $C_2 = B_2$ root system with only multiplicity information.

6.7. Euclidean Spinor Case. For the simplest vector case we consider the case $p = 3, q = 0$, where V_2, V_3 are scalar representations and V_1, V_4 are the $\text{spin-}\frac{1}{2}$ representation of $\mathfrak{m} = \mathfrak{so}(3)$ (note that strictly speaking, this does not lift to a representation of M), given by

$$F_{1,2} \cong \frac{i}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad F_{1,3} \cong \frac{1}{2} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad F_{2,3} \cong \frac{i}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

This allows us to eventually describe spherical functions on $\text{Spin}(4, 1)$ without having to work on that group.

By picking our Weyl group generator w as the diagonal matrix $\text{diag}(-1, 1, -1, 1, 1)$, we can ensure that the \mathfrak{h} -bimodule $W = V_1 \otimes \cdots \otimes V_4$ from Theorem 5.24 becomes isomorphic to the \mathfrak{m} -bimodule $\text{End}(V)$ (with V being the $\text{spin-}\frac{1}{2}$ representation), where the generator $F_{0,4}$ of the scaling group acts as $\Delta_1 - \Delta_2 = : -2\alpha$ from the left and $\Delta_4 - \Delta_3 = : -2\beta$ from the right.

Let us for simplicity choose \mathfrak{c} to be generated by $F_{0,2}$ and $F_{3,4}$, instead of $F_{0,1}, F_{3,4}$ (in other words, we are exchanging 1 and 2). From Corollary 6.25 we then have

$$\begin{aligned} A_{\epsilon_1 + \epsilon_2} &= F_{1,2} \otimes F_{1,2} \\ A_{\epsilon_1 - \epsilon_2} &= F_{1,3} \otimes F_{1,3} \\ A_{\epsilon_{1/2}} &= -\frac{1}{2} (F_{0,4} \mp F_{2,3}) \otimes (F_{0,4} \mp iF_{2,3}). \end{aligned}$$

We see that $A_{\epsilon_{1/2}}$ acts by means of diagonal matrices, which simplifies computations.

Proposition 6.36. *In the above setting we have*

$$\begin{aligned} K_{\epsilon_1 + \epsilon_2} &= -\frac{1}{8} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix} & K_{\epsilon_1 - \epsilon_2} &= -\frac{1}{8} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix} \\ K_{2\epsilon_{1/2}} &= -\frac{1}{8} \text{diag} \begin{pmatrix} (2\alpha + 2\beta \pm 1)^2 \\ (2\alpha + 2\beta)^2 \\ (2\alpha + 2\beta)^2 \\ (2\alpha + 2\beta \mp 1)^2 \end{pmatrix} \\ L_{\frac{\epsilon_1 + \epsilon_2}{2}} &= \frac{1}{16} \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} & L_{\frac{\epsilon_1 - \epsilon_2}{2}} &= \frac{1}{16} \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \\ L_{\epsilon_{1/2}} &= \frac{1}{8} \text{diag} \begin{pmatrix} (2\alpha \pm \frac{1}{2})(2\beta \pm \frac{1}{2}) \\ (2\alpha \pm \frac{1}{2})(2\beta \mp \frac{1}{2}) \\ (2\alpha \mp \frac{1}{2})(2\beta \pm \frac{1}{2}) \\ (2\alpha \mp \frac{1}{2})(2\beta \mp \frac{1}{2}) \end{pmatrix}. \end{aligned}$$

This expression can indeed be matched with the differential operator obtained in [15, §4.2]. In order to do this, we need to apply the gauge transformation sketched from [15, Equation 4.18], which corresponds to conjugation with the matrix

$$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & -1 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix}.$$

Proposition 6.37. *The gauge transformation produces the following matrices:*

$$\begin{aligned} \tilde{K}_{\epsilon_1+\epsilon_2} &= -\frac{1}{4} \operatorname{diag} \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, & \tilde{K}_{\epsilon_1-\epsilon_2} &= -\frac{1}{4} \operatorname{diag} \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \\ \tilde{K}_{2\epsilon_{1/2}} &= -\frac{1}{2} \begin{pmatrix} (\alpha+\beta)^2 + \frac{1}{4} & 0 & 0 & \mp(\alpha+\beta) \\ 0 & (\alpha+\beta)^2 & 0 & 0 \\ 0 & 0 & (\alpha+\beta)^2 & 0 \\ \mp(\alpha+\beta) & 0 & 0 & (\alpha+\beta)^2 + \frac{1}{4} \end{pmatrix}, \\ \tilde{L}_{\frac{\epsilon_1+\epsilon_2}{2}} &= \frac{1}{16} \operatorname{diag} \begin{pmatrix} 1 \\ 1 \\ -1 \\ -1 \end{pmatrix}, & \tilde{L}_{\frac{\epsilon_1-\epsilon_2}{2}} &= \frac{1}{16} \operatorname{diag} \begin{pmatrix} 1 \\ -1 \\ 1 \\ -1 \end{pmatrix}, \\ \tilde{L}_{\epsilon_{1/2}} &= \frac{1}{8} \begin{pmatrix} 4\alpha\beta + \frac{1}{4} & 0 & 0 & \mp(\alpha+\beta) \\ 0 & 4\alpha\beta - \frac{1}{4} & \pm(\alpha-\beta) & 0 \\ 0 & \pm(\alpha-\beta) & 4\alpha\beta - \frac{1}{4} & 0 \\ \mp(\alpha+\beta) & 0 & 0 & 4\alpha\beta + \frac{1}{4} \end{pmatrix}. \end{aligned}$$

Defining

$$V_{(\alpha,\beta)}^{PT}(\gamma) := \frac{(\alpha+\beta)^2 + \frac{1}{4}}{\sinh^2 \gamma} - \frac{\alpha\beta}{\sinh^2 \gamma/2},$$

we discover that $R(\Omega_{\mathfrak{g}})$'s 0th term has a diagonal term that amounts to

$$-V_{(\alpha,\beta)}^{PT}(\epsilon_1) - V_{(\alpha,\beta)}^{PT}(\epsilon_2).$$

The remaining terms on the diagonal read

$$\frac{1}{16} \begin{pmatrix} \operatorname{csch}^2_{\frac{\epsilon_1}{2}} + \operatorname{csch}^2_{\frac{\epsilon_2}{2}} + 2 \operatorname{sech}^2_{\frac{\epsilon_1+\epsilon_2}{4}} + 2 \operatorname{sech}^2_{\frac{\epsilon_1-\epsilon_2}{4}} \\ -\operatorname{sech}^2_{\frac{\epsilon_1}{2}} - \operatorname{sech}^2_{\frac{\epsilon_2}{2}} + 2 \operatorname{sech}^2_{\frac{\epsilon_1+\epsilon_2}{4}} - 2 \operatorname{csch}^2_{\frac{\epsilon_1-\epsilon_2}{4}} \\ -\operatorname{sech}^2_{\frac{\epsilon_1}{2}} - \operatorname{sech}^2_{\frac{\epsilon_2}{2}} - 2 \operatorname{csch}^2_{\frac{\epsilon_1+\epsilon_2}{4}} + 2 \operatorname{sech}^2_{\frac{\epsilon_1-\epsilon_2}{4}} \\ \operatorname{csch}^2_{\frac{\epsilon_1}{2}} + \operatorname{csch}^2_{\frac{\epsilon_2}{2}} - 2 \operatorname{csch}^2_{\frac{\epsilon_1+\epsilon_2}{4}} - 2 \operatorname{csch}^2_{\frac{\epsilon_1-\epsilon_2}{4}} \end{pmatrix},$$

and the off-diagonal ones (top to bottom):

$$\begin{pmatrix} \frac{\alpha+\beta}{4} \left(-\operatorname{sech}^2_{\frac{\epsilon_1}{2}} + \operatorname{sech}^2_{\frac{\epsilon_2}{2}} \right) \\ \frac{\alpha-\beta}{4} \left(\operatorname{csch}^2_{\frac{\epsilon_1}{2}} - \operatorname{csch}^2_{\frac{\epsilon_2}{2}} \right) \\ \frac{\alpha-\beta}{4} \left(\operatorname{csch}^2_{\frac{\epsilon_1}{2}} - \operatorname{csch}^2_{\frac{\epsilon_2}{2}} \right) \\ \frac{\alpha+\beta}{4} \left(-\operatorname{sech}^2_{\frac{\epsilon_1}{2}} + \operatorname{sech}^2_{\frac{\epsilon_2}{2}} \right) \end{pmatrix}.$$

Up to the introduction of coordinates u_1, u_2 that associates concrete hypergeometric functions with our $\sinh_\gamma, \cosh_\gamma$ as follows:

$$\begin{aligned} f_{\frac{\epsilon_i}{2}} &\mapsto f\left(\frac{u_i}{2}\right) \quad (i = 1, 2) \\ f_{\frac{u_1 \pm u_2}{4}} &\mapsto f\left(\frac{u_1 \mp u_2}{2}\right), \end{aligned}$$

we thus obtain as 0th-order term almost the (negative of the) potential described in [15, Equations 4.16–17]. The difference is a scalar function (i.e. identity matrix)

$$\frac{\operatorname{csch}_{\epsilon_1}^2 + \operatorname{csch}_{\epsilon_2}^2}{4} + \frac{\operatorname{csch}_{\frac{\epsilon_1 + \epsilon_2}{2}}^2 + \operatorname{csch}_{\frac{\epsilon_1 - \epsilon_2}{2}}^2}{8},$$

which we, in line with the original reference [15], get from turning $L(k)$ into Hamiltonian form. In conclusion, we arrive exactly at the operator that was obtained in [15, Section 4.2].

7. BLOCKS OF CONFORMAL DEFECTS

As is detailed in [12, Chapter 3], (scalar) conformal blocks for two defects of dimension p in d Euclidean spacetime dimensions can be described as zonal spherical functions for the pair $(G, H) := (SO(d+1, 1)_0, SO(d-p) \times SO(p+1, 1)_0)$. This pair is symmetric: if we pick the involution σ to be conjugation with the diagonal matrix

$$\begin{pmatrix} 1_{d-p} & 0 \\ 0 & -1_{p+2} \end{pmatrix},$$

then $H = G^\sigma$. Furthermore, (G, K, θ, B) is a reductive Lie group, where $K = SO(d+1)$, the involution θ consists in conjugation with the diagonal matrix

$$\begin{pmatrix} 1_{d+1} & 0 \\ 0 & -1 \end{pmatrix},$$

and B is the trace form from the defining representation. Evidently, θ and σ commute and B is σ -invariant. Consequently, we can apply Matsuki's theory.

Proposition 7.1. *A fundamental Cartan subset C is given by $C = \exp(\mathfrak{c})$ where*

$$\mathfrak{c} = \begin{cases} \operatorname{span}\{F_{i+1, d-i} \mid i = 0, \dots, p\} \oplus \mathbb{R}F_{0, d+1} & 2p < d-1 \\ \operatorname{span}\{F_{i, d-i} \mid i = 0, \dots, d-p-1\} & 2p \geq d-1 \end{cases}.$$

Here, N , the rank of \mathfrak{c} , is given by $\min(p+2, d-p)$.

Proof. For $p+1 \geq d-p$, note that the claimed maximal commutative subalgebra $\mathfrak{c} = \mathfrak{t}$ indeed is a commutative subalgebra of $\mathfrak{k}^{-\sigma}$. The vector space $\mathfrak{k}^{-\sigma}$ is spanned by $F_{\mu, \nu}$ for $0 \leq \mu < d-p \leq \nu \leq d$. Let X be an element of it, say

$$X = \sum_{\mu=0}^{d-p-1} \sum_{\nu=d-p}^d a_{\mu, \nu} F_{\mu, \nu},$$

we then have

$$\begin{aligned} [F_{i, d-i}, X] &= \sum_{\mu, \nu} a_{\mu, \nu} (-\delta_{i, \mu} F_{d-\mu, \nu} + \delta_{d-i, \nu} F_{\mu, d-\nu}) \\ &= - \sum_{\nu \neq d-i} a_{i, \nu} F_{d-i, \nu} + \sum_{\mu \neq i} a_{\mu, d-i} F_{\mu, i}. \end{aligned}$$

This is zero iff (linear independence) $a_{i, \nu} = 0$ ($\nu \neq d-i$) and $a_{\mu, d-i} = 0$ ($\mu \neq i$). If we want this to hold true for all $i = 0, \dots, d-p-1$, we obtain for each possible first index $\mu = 0, \dots, d-p-1$ that $a_{\mu, \nu}$ is only allowed to be nonzero for $\mu + \nu =$

d . In other words, $X \in \mathfrak{t}$. This shows that \mathfrak{t} is indeed maximally commutative. Furthermore, by a similar argument there is no element in

$$\mathfrak{p}^{-\sigma} = \text{span}\{F_{i,d+1} \mid i = 0, \dots, d-p-1\}$$

that commutes with \mathfrak{t} , so that the algebra \mathfrak{a} is trivial. We thus see that \mathfrak{c} has rank $N = d-p$.

For $p+1 < d-p$, consider \mathfrak{t} , the compact part of the claimed algebra \mathfrak{c} . Let $X \in \mathfrak{k}^{-\sigma}$, say

$$X = \sum_{\mu=0}^{d-p-1} \sum_{\nu=d-p}^d a_{\mu,\nu} F_{\mu,\nu}.$$

Then

$$\begin{aligned} [F_{i+1,d-i}, X] &= \sum_{\mu,\nu} a_{\mu,\nu} (-\delta_{i+1,\mu} F_{d+1-\mu,\nu} + \delta_{d-i,\nu} F_{\mu,d+1-\nu}) \\ &= - \sum_{\nu \neq d-i} a_{i+1,\nu} F_{d-i,\nu} + \sum_{\mu \neq i+1} a_{\mu,d-i} F_{\mu,i+1}. \end{aligned}$$

Due to linear independence, this is zero iff $a_{i+1,\nu} = 0$ and $a_{\mu,d-i} = 0$ (for $\nu \neq d-i, \mu \neq i+1$). Consequently, X commutes with all of \mathfrak{t} if these conditions hold for all $i = 0, \dots, p$. This ensures that for all possible choices of the second index ν , we can only have $a_{\mu,\nu} \neq 0$ for $\mu + \nu = d+1$. Consequently, $X \in \mathfrak{t}$. The centraliser of \mathfrak{t} in $\mathfrak{p}^{-\sigma}$ is spanned by $F_{0,d+1}$ and $F_{p+2,d+1}, \dots, F_{d-p-1,d+1}$. Since no two linearly independent elements of this span commute, any maximal commutative subalgebra is therefore one-dimensional, and so we can choose $\mathbb{R}F_{0,d+1}$ as indicated. Furthermore, the rank of \mathfrak{c} is $p+2$, which is $\leq d-p$, hence $N = p+2$. \square

Starting from this choice of fundamental Cartan subset, we need to figure out what are the other standard Cartan subsets.

Proposition 7.2. *For $2p < d-1$, there are no other standard Cartan subsets. For $2p \geq d-1$, the remaining standard Cartan subsets are of the shape*

$$C_i := \exp(\mathfrak{c}_i), \quad \mathfrak{c}_i := \text{span}\{F_{j,d-j} \mid 0 \leq j \leq d-p-1, j \neq i\} \oplus \mathbb{R}F_{i,d+1}$$

$(i = 0, \dots, d-p-1)$ or $C'_i := C_i \exp(\pi F_{i,d-i})$.

Proof. Since no two linearly independent elements of $\mathfrak{p}^{-\sigma}$ commute, any commutative subalgebra \mathfrak{a} is at most one-dimensional. Consequently, there are no other choices of Cartan subset for $2p < d-1$.

For $2p \geq d-1$, let $C' = \exp(\mathfrak{a}')T'$ be a standard Cartan subset different from the fundamental Cartan subset C . Then \mathfrak{a} is one-dimensional, say spanned by

$$X = \sum_{i=0}^{d-p-1} a_i F_{i,d+1} \in \mathfrak{p}^{-\sigma}.$$

For any $t \in T'$ we then have $\text{Ad}(t^{-1})(X) \in \mathfrak{g}^{-\sigma}$ as well. If we write

$$t = \exp\left(\sum_{i=0}^{d-p-1} \phi_i F_{i,d-i}\right),$$

we have

$$\text{Ad}(t^{-1})(X) = \sum_{i=0}^{d-p-1} a_i (\cos(\phi_i) F_{i,d+1} + \sin(\phi_i) F_{d-i,d+1}).$$

This lies in $\mathfrak{g}^{-\sigma}$ iff $a_i \sin(\phi_i) = 0$ for all $i = 0, \dots, d-p-1$. Since $\dim(\mathfrak{t}') = d-p-1$, all but one of these equations have to be satisfied independently of ϕ_i . In other words: there is an $i = 0, \dots, d-p-1$ such that $a_i \neq 0$ (and $a_j = 0$ for $j \neq i$).

This then also implies $\sin(\phi_i) = 0$. Thus, $C' = C_i$ or C'_i , depending on whether $\phi_i \in 2\pi\mathbb{Z}$ or in $\pi + 2\pi\mathbb{Z}$. \square

Note that any orthogonal block matrix permuting the indices $i \leftrightarrow j$ and $d-i \leftrightarrow d-j$, we can immediately see that C_i and C_j are conjugate, as are C'_i, C'_j .

Proposition 7.3. *Both with respect to \mathfrak{c} and \mathfrak{c}_0 , \mathfrak{g} has a reduced root system of type B_N (or D_N in case $2p = d - 2$) with root multiplicities:*

$$n_{\text{short}} = |d - 2 - 2p|, \quad n_{\text{long}} = 1.$$

Proof. For $2p \geq d - 1$, we have for $0 \leq i, j, k \leq d - p - 1$:

$$\begin{aligned} [F_{i,d-i}, F_{j,k} \pm iF_{d-j,k} + iF_{j,d-k} \mp F_{d-j,d-k}] &= (\pm i\delta_{i,j} + i\delta_{i,k})(\cdots) \\ [F_{i,d-i}, F_{j,k} \pm iF_{d-j,k} - iF_{j,d-k} \pm F_{d-j,d-k}] &= (\pm i\delta_{i,j} - i\delta_{i,k})(\cdots) \\ [F_{i,d-i}, F_{j,m} \pm iF_{d-j,m}] &= \pm i\delta_{i,j}(\cdots) \\ [F_{i,d-i}, F_{m,n}] &= 0 \end{aligned}$$

for $m, n = d - p, \dots, p, d + 1$, which establishes that the reduced root system with respect to \mathfrak{c} is of type B_{d-p} and has root multiplicities $2p - d + 2$ and 1 for short and long roots, respectively. Note that the dimensions of the root spaces we found add up to $\dim(\mathfrak{g})$, so that we have indeed found all of them.

Similarly, we have

$$\begin{aligned} [F_{0,d+1}, F_{0,m} \pm F_{m,d+1}] &= \pm(F_{0,m} \pm F_{m,d+1}) \\ [F_{0,d+1}, F_{m,n}] &= 0 \end{aligned}$$

for $m, n = 1, \dots, d$, which yields a root system of type B_{d-p} with multiplicities $2p - d + 2$ and 1.

Lastly, for $2p < d - 1$ we have for $i = 0, \dots, p$:

$$\begin{aligned} [F_{i+1,d-i}, F_{j+1,k+1} \pm iF_{d-j,k+1} + iF_{j+1,d-k} \mp F_{d-j,d-k}] &= (\pm i\delta_{i,j} + i\delta_{i,k})(\cdots) \\ [F_{0,d+1}, F_{j+1,k+1} \pm iF_{d-j,k+1} + iF_{j+1,d-k} \mp F_{d-j,d-k}] &= 0 \\ [F_{i+1,d-i}, F_{j+1,k+1} \pm iF_{d-j,k+1} - iF_{j+1,d-k} \pm F_{d-j,d-k}] &= (\pm i\delta_{i,j} - i\delta_{i,k})(\cdots) \\ [F_{0,d+1}, F_{j+1,k+1} \pm iF_{d-j,k+1} - iF_{j+1,d-k} \pm F_{d-j,d-k}] &= 0 \\ [F_{i+1,d-i}, F_{0,j+1} \pm iF_{0,d-j} + F_{j+1,d+1} \pm iF_{d-j,d+1}] &= \pm i\delta_{i,j}(\cdots) \\ [F_{0,d+1}, F_{0,j+1} \pm iF_{0,d-j} + F_{j+1,d+1} \pm iF_{d-j,d+1}] &= \cdots \\ [F_{i+1,d-i}, F_{0,j+1} \pm iF_{0,d-j} - F_{j+1,d+1} \mp iF_{d-j,d+1}] &= \pm i\delta_{i,j}(\cdots) \\ [F_{0,d+1}, F_{0,j+1} \pm iF_{0,d-j} - F_{j+1,d+1} \mp iF_{d-j,d+1}] &= -\cdots \\ [F_{i+1,d-i}, F_{j+1,m} \pm iF_{d-j,m}] &= -i\delta_{i,j}(\cdots) \\ [F_{0,d+1}, F_{j+1,m} \pm iF_{d-j,m}] &= 0 \\ [F_{i+1,d-i}, F_{0,m} \pm F_{m,d+1}] &= 0 \\ [F_{0,d+1}, F_{0,m} \pm F_{m,d+1}] &= \pm(\cdots) \\ [F_{i+1,d-i}, F_{m,n}] &= 0 \\ [F_{0,d+1}, F_{m,n}] &= 0 \end{aligned}$$

where $m, n = p + 2, \dots, d - 1 - p$. This indeed establishes that we are dealing here with the root system of type B_{p+2} or D_{p+2} (in case $d - 2 = 2p$), with root multiplicities $d - 2p - 2$ and 1. \square

Lemma 7.4. *All standard Cartan subsets satisfy the technical condition of Section 4.2.*

Proof. For $2p < d - 1$ and the cases of C and C_i for $2p \geq d - 1$ there is nothing to show as the Cartan subsets in questions are subgroups and are therefore covered. It remains then to show that C'_i for $2p \geq d - 1$ satisfies the technical condition. This is a coset with “inhomogeneity” $t' = \exp(\pi F_{i,d-i})$. The map $\text{Ad}(t')$ leaves \mathfrak{c}_i invariant and squares to the identity, since $(t')^2 = 1$, so that it commutes with σ . Therefore it satisfies the conditions we impose on ϕ in the decomposition $\text{Ad}(t')|_{\mathfrak{g}_\alpha} = \epsilon_\alpha \phi$. \square

Theorem 7.5. *The quadratic Casimir element acts on conformal blocks for p -dimensional defects in d -dimensional Euclidean spacetime as the operator $L(k)$ from [11, Proposition 1.2.3] for a root system of type B_N (or D_N) with multiplicities*

$$k_{\text{short}} = \frac{|d - 2 - 2p|}{2}, \quad k_{\text{long}} = \frac{1}{2},$$

where $N = \min(p + 2, d - p)$. This exactly matches what was obtained in [12, Section 3] in the case $p = q$.

Proof. By Proposition 7.2 we need to consider C (for $2p < d - 1$) or C, C_i, C'_i (for $2p \geq d - 1$). By Lemma 7.4 we may apply Theorem 4.22. Since our H -bimodule is trivial, we ignore all elements of $U(\mathfrak{h}) \otimes_{U(\mathfrak{m}')} U(\mathfrak{h})$, so that the differential operator $R^{\mathbb{C}}(\Omega_{\mathfrak{g}})$ reduces to the Laplacian $L(k)$ for the root system $2\Sigma(\mathfrak{g}; \mathfrak{c}_I)$ where I is either the empty word or a number in $0, \dots, d - 1 - p$. By Proposition 7.3, this root system is of type B_N with multiplicities $|d - 2 - 2p|$ and 1, respectively (in case the short multiplicity is 0, it is of type D_N), which explains the choice of parameter k . \square

8. CONCLUSION AND OUTLOOK

In this paper we made a step towards developing a systematic theory of matrix-spherical functions for symmetric pairs (G, H) , where neither G nor H need to be compact. Namely, we used results from [5] to construct a group decomposition that generalises the usual KAK -decomposition, known from the standard theory of non-compact reductive Lie groups. We then used this new decomposition to explore the corresponding radial part reductions akin to the theory presented in [2], and in particular to establish a matrix-valued, general symmetric pair analogue of [11, Theorem 5.1.7].

Afterwards we applied this theory to some of the most basic examples arising in the study of conformal blocks in CFT. Firstly, we elaborated in detail on conformal blocks for 4-point functions, in particular on how the connection between the (quadratic) Casimir equation and the Calogero–Sutherland model [10] can be appropriately derived and on what the Casimir equation looks like for general non-scalar cases (both Euclidean and Lorentzian), paying special attention to mathematical subtleties on the way. And secondly, we also checked our results against the two simple benchmarks: the matrix Calogero–Sutherland Hamiltonians corresponding to the spinorial Casimir equation for three-dimensional spinning blocks [15] and to the conformal blocks for two p -dimensional scalar defects in Euclidean conformal field theory [12].

Highlighting the added value of this paper on the physics side, we would like to emphasize once more the need for a thorough understanding of the group decompositions and the corresponding radial part computations that were considered here. By now, the general form of the correspondence between conformal blocks, harmonic analysis and integrable models [6, 15] is rather clear, has been extended to various physically interesting settings, most importantly to the multipoint case in [21], and has started to bring practical results, providing the actual input for the machinery of numerical conformal bootstrap, especially in interesting not-yet-explored setups. However, if one ultimately strives for an exhaustive analysis of

the mathematics behind (the kinematics of) higher-dimensional conformal theories, it is fair to say that various pieces of this correspondence are not yet up to the standards of mathematical rigour. As a matter of fact, being perhaps at a risk of slightly overstating, one can regard this state of things as one of the crucial stumbling blocks on the way of properly using representation-theoretical tools in order to (analytically) address the questions of dynamics of higher-dimensional CFTs. It is precisely one of such shallower points that we have elaborated upon in the present paper: we clarified the specifics of the *KAK* decompositions used in computations of radial parts of conformal Casimirs and made the latter fully explicit in the present context. Elucidation of this specific issue, along with the proper analysis of the Casimir reduction in the Lorentzian setting, should be considered as the main added value of the present paper on the physics side.

Moving on to the outlook of possible further directions, the most interesting immediate development of our findings here would be a clarification of the connection of the present setup to the works of J. Stokman and N. Reshetikhin [19, 22, 23], who studied generalised spherical functions associated to the moduli spaces of principal connections on a finite graph. As we reviewed earlier, the CFT four-point functions are closely related to matrix-spherical functions for the symmetric pair (G, MA) , which from the above point of view can be thus regarded as a correspondence between such generalised spherical functions related to two different types of graphs. It would be interesting to understand correspondences of this type better and, in particular, to explore their possible analogues for multipoint conformal blocks. We also expect that the theory we developed here can be used in the Reshetikhin-Stokman setting to lift the restriction to compact subgroups, playing a role in [19].

On a slightly smaller scale, let us conclude by listing several concrete further directions that could additionally be addressed in the context of conformal block computations via Matsuki decompositions:

- It would be interesting to get explicit formulas also for the radial parts of other generators of the algebra of invariant differential operators in various setups. Conceptually this does not seem to be much less straightforward than for the quadratic Casimir, however, the amount of necessary calculations grows quite fast with the order of differential operator. For example, in the simplest case of the scalar 4-point blocks there is a well-known Casimir element of degree four, whose action has been calculated in [24, Equation 4.14].
- Another simple matrix-valued case that is feasible to treating in complete generality is $V_1 = V, V_2 = \mathbb{C}, V_3 = V^*, V_4 = \mathbb{C}$ for the defining representation V of $SO(p, q)$. V restricts to a sum of two trivial and one defining representation of M' , which means that $\text{End}_{M'}$ is five-dimensional.
- Our discussion of the blocks for conformal defects from [12] in the last section, of course, has also just scratched the surface. Most interestingly, Matsuki's decomposition naturally allows to consider different (symmetric) involutions of the two sides of a double coset, which thus fits perfectly into the framework of the conformal blocks for defects of different codimension $p \neq q$. In particular, we expect that it should be possible to rigorously reproduce and extend the results of [12, Section 3.3] using Matsuki's theory. It is needless to say that all these constructions can and, perhaps, should also be extended to other signatures (Lorentzian) and to the spinning case. Some of these setups have already been analysed in the physics literature, with the omissions similar to what we have addressed in this paper.

- Getting under control the radial part decompositions for the other types of non-compact symmetric spaces by using Matsuki’s decompositions is, of course, also an interesting, more distant goal. Some of these further setups also bear a clear relevance for physics, e.g. the group case is known to be related to the finite-temperature conformal blocks.

9. ACKNOWLEDGEMENTS

PS would like to thank his PhD supervisors Erik Koelink and Maarten van Puijssen for allowing him to pursue this side project. MI thanks Edward Berengoltz and Jasper Stokman for numerous discussions on the topics of this paper. The research of PS was funded by grant OCENW.M20.108 of the Dutch Research Council NWO.

REFERENCES

- [1] Garth Warner. *Harmonic analysis on semi-simple Lie groups. II*. Vol. Band 189. Die Grundlehren der mathematischen Wissenschaften. Springer-Verlag, New York-Heidelberg, 1972.
- [2] William Casselman and Dragan Milićić. “Asymptotic behavior of matrix coefficients of admissible representations”. In: *Duke Math. J.* 49.4 (1982), pp. 869–930. DOI: [10.1215/S0012-7094-82-04943-2](https://doi.org/10.1215/S0012-7094-82-04943-2).
- [3] Erik Koelink, Maarten van Puijssen, and Pablo Román. “Matrix elements of irreducible representations of $SU(n+1) \times SU(n+1)$ and multivariable matrix-valued orthogonal polynomials”. In: *J. Funct. Anal.* 278.7 (2020), pp. 108411, 48. DOI: [10.1016/j.jfa.2019.108411](https://doi.org/10.1016/j.jfa.2019.108411).
- [4] Anthony W. Knap. *Lie groups beyond an introduction*. Second. Vol. 140. Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, 2002.
- [5] Toshihiko Matsuki. “Double coset decompositions of reductive Lie groups arising from two involutions”. In: *J. Algebra* 197.1 (1997), pp. 49–91. DOI: [10.1006/jabr.1997.7123](https://doi.org/10.1006/jabr.1997.7123).
- [6] Volker Schomerus, Evgeny Sobko, and Mikhail Isachenkov. “Harmony of spinning conformal blocks”. In: *J. High Energy Phys.* 3 (2017), 085, front matter+22. DOI: [10.1007/JHEP03\(2017\)085](https://doi.org/10.1007/JHEP03(2017)085).
- [7] Petr Kravchuk, Jiaxin Qiao, and Slava Rychkov. “Distributions in CFT. Part I. Cross-ratio space”. In: *JHEP* 05 (2020), p. 137. DOI: [10.1007/JHEP05\(2020\)137](https://doi.org/10.1007/JHEP05(2020)137). arXiv: [2001.08778 \[hep-th\]](https://arxiv.org/abs/2001.08778).
- [8] Petr Kravchuk, Jiaxin Qiao, and Slava Rychkov. “Distributions in CFT. Part II. Minkowski space”. In: *J. High Energy Phys.* 8 (2021), Paper No. 094, 129. DOI: [10.1007/jhep08\(2021\)094](https://doi.org/10.1007/jhep08(2021)094).
- [9] F. A. Dolan and H. Osborn. “Conformal partial waves and the operator product expansion”. In: *Nuclear Phys. B* 678.1-2 (2004), pp. 491–507. DOI: [10.1016/j.nuclphysb.2003.11.016](https://doi.org/10.1016/j.nuclphysb.2003.11.016).
- [10] Mikhail Isachenkov and Volker Schomerus. “Superintegrability of d -dimensional conformal blocks”. In: *Phys. Rev. Lett.* 117.7 (2016), pp. 071602, 5. DOI: [10.1103/PhysRevLett.117.071602](https://doi.org/10.1103/PhysRevLett.117.071602).
- [11] Gerrit Heckman and Henrik Schlichtkrull. *Harmonic analysis and special functions on symmetric spaces*. Vol. 16. Perspectives in Mathematics. Academic Press, Inc., San Diego, CA, 1994.
- [12] M. Isachenkov et al. “Calogero-Sutherland approach to defect blocks”. In: *J. High Energy Phys.* 10 (2018), 204, front matter+43. DOI: [10.1007/jhep10\(2018\)204](https://doi.org/10.1007/jhep10(2018)204).

- [13] V. K. Dobrev et al. *Harmonic Analysis on the n -Dimensional Lorentz Group and Its Application to Conformal Quantum Field Theory*. Vol. 63. 1977. DOI: [10.1007/BFb0009678](https://doi.org/10.1007/BFb0009678).
- [14] M. Isachenkov and V. Schomerus. “Integrability of conformal blocks. Part I. Calogero-Sutherland scattering theory”. In: *JHEP* 07 (2018), p. 180. DOI: [10.1007/JHEP07\(2018\)180](https://doi.org/10.1007/JHEP07(2018)180). arXiv: [1711.06609](https://arxiv.org/abs/1711.06609) [hep-th].
- [15] Ilija Burić, Volker Schomerus, and Mikhail Isachenkov. “Conformal group theory of tensor structures”. In: *J. High Energy Phys.* 10 (2020), pp. 004, 38. DOI: [10.1007/jhep10\(2020\)004](https://doi.org/10.1007/jhep10(2020)004).
- [16] Ilija Buric and Volker Schomerus. “Universal spinning Casimir equations and their solutions”. In: *JHEP* 03 (2023), p. 133. DOI: [10.1007/JHEP03\(2023\)133](https://doi.org/10.1007/JHEP03(2023)133). arXiv: [2211.14340](https://arxiv.org/abs/2211.14340) [hep-th].
- [17] Pulkit Agarwal et al. “Embedding space approach to Lorentzian CFT amplitudes and causal spherical functions”. In: *Phys. Rev. D* 110.8 (2024), p. 086019. DOI: [10.1103/PhysRevD.110.086019](https://doi.org/10.1103/PhysRevD.110.086019). arXiv: [2302.06469](https://arxiv.org/abs/2302.06469) [hep-th].
- [18] David Poland, Slava Rychkov, and Alessandro Vichi. “The conformal bootstrap: Theory, numerical techniques, and applications”. In: *Rev. Mod. Phys.* 91 (1 2019).
- [19] N. Reshetikhin and J. V. Stokman. “Quantum superintegrable spin systems on graph connections”. In: (). arXiv: [2305.02767](https://arxiv.org/abs/2305.02767).
- [20] Jiaxin Qiao. “Classification of convergent OPE channels for Lorentzian CFT four-point functions”. In: *SciPost Phys.* 13.4 (2022), Paper No. 093, 74.
- [21] Ilija Buric et al. “From Gaudin Integrable Models to d -dimensional Multipoint Conformal Blocks”. In: *Phys. Rev. Lett.* 126.2 (2021), p. 021602. DOI: [10.1103/PhysRevLett.126.021602](https://doi.org/10.1103/PhysRevLett.126.021602). arXiv: [2009.11882](https://arxiv.org/abs/2009.11882) [hep-th].
- [22] N. Reshetikhin and J. V. Stokman. “ N -point spherical functions and asymptotic boundary KZB equations”. In: *Invent. Math.* 229 (2022), pp. 1–86. DOI: [10.1007/s00222-022-01102-3](https://doi.org/10.1007/s00222-022-01102-3).
- [23] N. Reshetikhin and J. V. Stokman. “Asymptotic boundary KZB operators and quantum Calogero-Moser spin chains”. In: *Hypergeometry, integrability and Lie theory*. Vol. 780. Contemp. Math. Amer. Math. Soc., [Providence], RI, 2022, pp. 205–241. DOI: [10.1090/conm/780/15691](https://doi.org/10.1090/conm/780/15691).
- [24] F. A. Dolan and H. Osborn. *Conformal Partial Waves: Further Mathematical Results*. 2012. arXiv: [1108.6194](https://arxiv.org/abs/1108.6194) [hep-th]. URL: <https://arxiv.org/abs/1108.6194>.

Email address: philip.schloesser@ru.nl

RADBOD UNIVERSITY, IMAPP-MATHEMATICS, HEYENDAALSEWEG 135, 6525 AJ NIJMEGEN, THE NETHERLANDS

Email address: m.isachenkov@uva.nl

UNIVERSITEIT VAN AMSTERDAM: KORTEWEG-DE VRIES INSTITUTE FOR MATHEMATICS, SCIENCE PARK 107, 1090 GE AMSTERDAM, AND INSTITUTE OF PHYSICS, SCIENCE PARK 904, 1098 XH AMSTERDAM, THE NETHERLANDS