High precision spectroscopy of trilobite Rydberg molecules

Markus Exner[®],^{1,*} Rohan Srikumar,^{2,*} Richard Blättner[®],¹

Matthew T. Eiles⁰,³ Peter Schmelcher,^{2,4} and Herwig Ott⁰^{1,†}

¹Department of Physics and Research Center OPTIMAS,

Rheinland-Pfälzische Technische Universität Kaiserslautern-Landau, 67663 Kaiserslautern, Germany

²Zentrum für Optische Quantentechnologien, Universität Hamburg,

Luruper Chaussee 149, 22761 Hamburg, Germany

³Max Planck Institute for the Physics of Complex Systems, Nöthnitzer Str. 38, 01187 Dresden, Germany

⁴ The Hamburg Centre for Ultrafast Imaging, Universität Hamburg,

Luruper Chaussee 149, 22761 Hamburg, Germany

(Dated: December 30, 2024)

We perform three-photon photoassociation to obtain high resolution spectra of ⁸⁷Rb trilobite dimers for the principal quantum numbers n = 22, 24, 25, 26, and 27. The large binding energy of the molecules in combination with a relative spectroscopic resolution of 10^{-4} provides a rigorous benchmark for existing theoretical models. A recently developed Green's function framework, which circumvents the convergence issues that afflicted previous studies, is employed to theoretically reproduce the vibrational spectrum of the molecule with high accuracy. The relatively large molecular binding energy are primarily determined by the low energy S-wave electron-atom scattering length, thereby allowing us to extract the ${}^{3}S_{1}$ scattering phase shift with unprecedented accuracy, at low energy regimes inaccessible to free electrons.

Introduction: The ability to introduce Rydberg excitations in ultracold atomic gases has set in motion a new branch of molecular physics, wherein atoms are bound over large distances via unconventional binding mechanisms [1–7]. Trilobite molecules are a peculiar category of such ultralong-range Rydberg molecules (ULRM) [1, 8-11], bound by a ground state atom scattering off a Rydberg electron with high angular momentum ℓ . These homonuclear dimers, which exhibit permanent electric dipole moments in the kilo-Debye range [12–15] and have bond-lengths of the order of micrometers, promise dynamical effects and field control not typical of conventional molecules [16–18]. Although the first experimental evidence for the trilobite dimer was obtained by mixing the trilobite wavefunction to low- ℓ Rydberg states [14, 19], it was only very recently that direct photoassociation of the molecule, accessing the trilobite potential energy curves (PECs) was made possible [15]. For more than a decade, numerous spectroscopic studies have been performed on ULRM [20–28] providing detailed insights into the spin-structure, potential energy landscape, and rovibronic spectra of these dimers. Since the low-energy scattering of the Rydberg electron forms the binding mechanism of the molecule, the depth of the potential wells and the vibrational binding energies are directly dependent on the size of the scattering partial wave L. In turn, high resolution spectroscopic data can provide precise values of the spin-dependent scattering lengths, ultimately testing the validity of the underlying interaction mechanism. Consequently, the ULRM has been used as a laboratory to extract electron-atom scattering lengths and resonance positions at low energy regimes inaccessible with free electrons [21].

In this work, we combine the newly formulated Green's

function method [29] with the advantages of pure trilobite photoassociation spectroscopy. Due to the large molecular binding energies and the high precision of spectroscopic measurement, we determine the potential energy depth with unprecedented accuracy, and extract the energy-dependent e^- Rb, ${}^{3}S_{1}$ scattering length. At the same time, we find indications that the predictive power of the underlying molecular Hamiltonian is being challenged, and further insight in the direction of nonadiabatic physics [30–32] and/or higher order scattering terms is necessary before we can unravel agreement between experiment and theory data within the resolution achieved.

methodology: The Interactions and Born-Oppenheimer PECs $U_i(R),$ and the associated electronic states $\phi_i(\vec{r}; R)$ are obtained by solving $H\phi_i(\vec{r};R) = U_i(R)\phi_i(\vec{r};R)$, where the electronic Hamiltonian H incorporates the fine-structure of the Rydberg electron, hyperfine-structure of the ground-state atom (with a total spin F=1,2 for ⁸⁷Rb), and the six different electron-atom scattering channels $({}^{1}S_{0}, {}^{3}S_{1}, {}^{1}P_{1}, {}^{3}P_{0,1,2}).$ Until recently, the standard method for determining the PECs was the diagonalization of H in a truncated basis [33]. Here, the scattering interaction between the Rydberg electron and perturber is characterized by a Fermi-type pseudopotential in S-wave [34] and P-wave channels [35]. While providing a valid semi-quantitative description of the system, the Dirac-delta form of the pseudopotential introduces unphysical basis size dependence of the PECs obtained utilizing diagonalization schemes [36]. Furthermore, the approach suffers from diverging scattering volumes in the ${}^{3}P_{I}$ scattering channels introduced by negative ion resonances. For this reason, the scattering phase shift extracted by



FIG. 1. Rydberg n=27, M=1/2 molecular Born-Oppenheimer potential curves obtained using the Green's function treatment (grey). The black curves highlight the trilobite potentials we experimentally probe, and for comparison the blue curves are potentials calculated by numerical diagonalization (using $n \in [25, 29]$ basis). On the right are two density plots of the Coulomb Green's function which represents the Rydberg electron distribution. Due to the *S*wave scattering and the admixture of high angular momentum states, the probability of the electron staying at the position of the ground state atom is maximised.

comparing experimental and theoretical binding energies obtained utilizing this method needs to be handled with care. Alternative approaches have been developed, but as these do not include all spin interactions, they are not sufficiently accurate to meet high-precision spectroscopy [37, 38]. However, the recently developed non-perturbative Green's function formulation (see [29] for detailed explanation) accounts for all spin interactions, and circumvents the aforementioned convergence issues.

Figure 1 shows the Born-Oppenheimer potential energy landscape of the ULRM obtained using the Green's function calculation for principal quantum number n=27, with the projection of total angular momentum along the internuclear axis taken to be M=1/2. The zero frequency is the asymptotic pair state $\text{Rb}(27f_{7/2}) + \text{Rb}(5\text{s},F = 1)$, from which the two potential curves emerge (black), corresponding to the pure triplet scattering channel (lower-PEC), and a singlet-triplet mixed channel (upper-PEC). For illustration the same two potentials were calculated with the diagonalisation method (blue dotted), which leads to 10% deviation in the binding energy. At 1050 a_0 the trilobite levels interact with the plummeting "butterfly" potential curves resulting from the shape resonance in the p-wave scattering, of which there are five of predominantly ${}^{3}P_{2}$ configuration, three of ${}^{3}P_{1}$, and a single ${}^{3}P_{0}$. In this work we experimentally access the vibrational bound states in the highlighted trilobite curves (hereby referred to as triplet trilobite and mixed trilobite PEC) for n = 22, 24, 25, 26, 27, using a direct three-photon excitation scheme to the $nf_{7/2}$ atomic resonance.

Experimental setup: We prepare the atomic sample starting from a magneto-optical trap (MOT) of ⁸⁷Rb atoms, which are subsequently transferred into a crossed dipole trap operating at $\lambda = 1064 \,\mathrm{nm}$. The spherical atomic ensemble has a temperature of $40\,\mu\text{K}$, a diameter of $40 \,\mu m$ and a peak density of $4 \times 10^{13} \,\mathrm{cm}^{-3}$. The atoms are prepared in the F = 1 hyperfine Employing a three-photon excitation ground state. scheme $(5s_{1/2} \rightarrow 5p_{3/2} \rightarrow 5d_{5/2} \rightarrow nf_{7/2})$ at wavelengths of 780 nm, 776 nm, and 1276 nm to 1288 nm, we can photoassociate pure trilobite Rydberg molecules utilizing its f-state admixture. For precise frequency control, all three excitation lasers are frequency stabilized to an ultra-low expansion cavity using a Pound-Drevel-Hall locking technique. Immediately following the excitation, the Rydberg atoms are ionized using a CO_2 laser. The resulting ions are guided by an electric field in a Wiley-McLaren configuration onto a time- and positionresolved multichannel plate detector, enabling the direct measurement of the momentum distribution prior to ionization. The generated ions are counted for each detuning step (2 MHz) and averaged over typically 10 experimental runs. To estimate the maximum error for the frequency, we use the accuracy of the free spectral range of 100 kHz, resulting in an error for the lowest trilobite state of $\frac{21\,{\rm GHz}}{1.4978\,{\rm GHz}} \cdot 100\,{\rm kHz} \approx 1.4\,{\rm MHz}$. On the other hand, we use the frequency step size of 2 MHz as the readout error. This results in a frequency uncertainty of 3.4 MHz, which corresponds to a deviation smaller than 0.01% at a binding energy of 21 GHz and thus illustrates the excellent accuracy of the experiment.

Spectral analysis: Figure 2 shows the observed spectra for all measured n, which are several GHz red-detuned from the $nf_{7/2}$ atomic resonance. Two different vibrational series can be identified, which still overlap at the principal quantum number of 22 but are clearly separated at n = 25. These originate from the mixed and the triplet F = 1 PECs, respectively. The vertical lines show the calculated binding energies for the triplet F = 1 (green) and the mixed (blue) trilobite states. The number of vibrational states in each potential is limited by the position of the crossing with the butterfly PECs, as tunneling towards shorter internuclear distances occurs, leading to



FIG. 2. Ion signal (red) as a function of the frequency detuning of the third excitation laser for all measured principal quantum numbers. The atomic resonance $nf_{7/2}$ defines the frequency zero. The calculated eigenenergies are shown as vertical lines for the triplet F = 1 (green) and the mixed (blue) potential

state-changing collisions or associative ionization [15, 39]. For n < 25 there is a single outer well beyond the *P*-wave crossing, whereas for n=25 and above, there is a second potential well. This explains the presence of double peaks in the spectrum for n = 25-27 that are absent in the n = 22or 24 spectra. The presence of the second potential well renders the assignment of the vibrational states challenging. Here, knowledge of the permanent dipole moment helps in the identification of states, as the dipole moment of the vibrational state increases with bond-length. To determine the dipole moment, a weak electric field is applied and the broadening of the state is measured [15]. The measured dipole moments for different vibrational states of the n=27 mixed trilobite potential is shown in Fig. 3. Their magnitudes accumulate around two different values, which clearly allows the assignment of the vibrational states to the two calculated potential wells with minima at $1100 a_0$ and $1275 a_0$. Note that the relative motional state of two ground-state atoms in our



FIG. 3. The trilobite spectrum (red) for the energy range of the mixed potential for a principal quantum number of n=27 with the potential energy curve (black) above including vibrational probability densities (blue, not to scale). The observed dipole moments of each measured peak is shown and assigned to the corresponding vibrational state.

setup varies slowly over the extent of the molecular vibrational states, implying minimal Franck-Condon overlap for odd molecular vibrational states. However, the molecular spectra shows that both odd and even vibrational states have comparable signal strength. This phenomenon is explained by considering the oscillations of the $nf_{7/2}$ admixture in the trilobite state (responsible for the dipole transition from $5d_{5/2}$), that violates the Franck-Condon approximation that electronic transitions are independent of the nuclear coordinates and facilitates the photoassociation of odd vibrational states.

Phase shift extraction: We now discuss the extraction of the low-energy S-wave scattering phase shift from the high resolution data experimentally obtained. By solving the unphysical basis-dependencies that afflicted other calculations, we are able modify and correct previous phase shifts in such a way that it reproduces the observed resonances more precisely, without dependence on any external model parameters. For this, we added a cubic polynomial to the phase shifts fitted in [21] and treated the coefficients as fit parameters. These were varied until the total difference between the theoretical and experimental bound state spectra from all n were simultaneously minimized. The coefficients of the aforementioned polynomial were modified to reduce the error between the experimental and theoretical estimates of binding energies of the vibrational ground-states for the experimentally probed range of n. Figure 4 (a) illustrates the trilobite potential energy curves as a function of the semiclassical electron momentum k. Note that $k(R) = \sqrt{2U_n(R) + 1/R}$ is calculated separately for each particular PEC $U_n(R)$,



FIG. 4. Potential energy structure and extracted S-wave phase shifts. (a) Illustrative sketch of the probed trilobite PECs as a function of k, limited between the energy zeros (k=0) and the P-wave crossings $(k \sim 0.025)$ for each PEC. (b) The PECs (solid blue/black lines) and the vibrational states (dashed) of the molecule, for n=26, in comparison with the ion-count spectra (red line), and the (c) S-wave phase shifts used to obtain them. The solid black lines show the phase shifts fitted anew, and the green lines are the phase shifts taken from a previous calculation and the potential energy obtained from them [21]. The shaded region portrays the uncertainty in the fitting of phase shifts (see text), leading to corresponding deviations in binding energy (dashed lines).

and the curves are vertically offset for an easier visual The overlapping k-ranges for multiple comparison. *n*-values provide several data points (vibrational states) in each k-window whose energetic positions are sensitive to the changes in phase shift. By minimizing the error between the computed and observed binding energies for the vibrational ground states localized in the outer-well, for all values of n simultaneously, we prevent possible overfits in our phase shift calculations. Figure 4 (b) shows the trilobite potential energy curves for n=26, calculated utilizing the phase shifts shown in Fig. 4 (c). The newly fit phase shift (solid black line, Fig. 4 (c)) deviates considerably from the previously fitted scattering phase shift [21]. The shaded blue region in Fig. 4 (b) and (c) is used to illustrate the sensitivity of the molecular binding energy on the ${}^{3}S_{1}$ scattering length. A 0.7% change in the S-wave phase shift in the relevant k-range (Fig. 4 (c)) induces an error of ~ 100 MHz in the binding energy of the n = 26 vibrational ground state (dashed lines Fig. 4 (b), nearly two orders of magnitude

larger than the experimental resolution. Apart from illustrating the sensitivity of our calculations, the shaded region also shows the range of S-wave phase shifts within which the theoretical and experimental data can be selectively matched perfectly for each n-value (case of overfitting). The inability to match observed and calculated binding energies simultaneously for all values of n, within experimental error bars, implies that there might be limitations in the description of the molecular interactions in the Hamiltonian. This necessitates a quantitative analysis of the error between the observed spectra and the numerical spectra calculated with the optimized phase shifts.

Figure 5 (a) portrays the relative error of the binding energies of all the measured vibrational states with respect to their theoretical counterparts. The vibrational states localized in the outer-wells of the n > 25 trilobite PECs are uniquely S-wave dominant and show an excellent maximum relative error < 0.32%. Although the inner-well states (marked \blacklozenge) are also S-wave dominant, the exact positions of the ${}^{3}P_{0,1,2}$ crossings finely affects the number of bound states observed. However, we see that their binding energies are also largely dependent on the ${}^{3}S_{1}$ phase shift, and were reproduced with a relative error $\leq 0.8\%$. The average of the absolute values of all error values shown in Fig. 5 (a), was calculated to be only 0.25%, validating and quantifying the high-accuracy of the calculated potential energy depth and the extracted ${}^{3}S_{1}$ phase shift. Furthermore, in Fig. 5 (b) we show the relative error between the experimental and theoretical estimates of vibrational energy splitting for both the inner-well states (marked \blacklozenge) and the outer-well states. This serves as a better estimate for the accuracy with which we probe the shape of the potential well. The difference between adjacent energy levels in both the triplet and the mixed PECs are obtained separately and the relative error is shown for each n-value. States localised in the outer-well exhibit maximal relative error 3.3%, with the inner-well counterparts featuring error rates less than 7%. Further improvements in the molecular Hamiltonian would be necessary to produce numerical results that reproduce the experimental data within the resolution achieved.

Conclusion: To summarize, we have measured pure trilobite Rydberg series for five different principal quantum numbers at a resolution of 10^{-4} . Permanent electric dipole moments of almost 3000 debye were measured and it was shown that this allows an unambiguous assignment of vibrational states in multi-well potentials. Furthermore, the recently developed Green's function method was used to analyze the most extensive experimental study of the trilobite ULRM to date. This allowed the binding energies and vibrational splittings to be theoretically reproduced with a relative error



FIG. 5. The percentage relative difference of the experimental results with respect to the numerical calculations given by $\Delta_R = 100\% \cdot (\epsilon_e - \epsilon_t)/\epsilon_t$ where ϵ_e and ϵ_t represent the experimental and theoretical data respectively for (a), the calculated vibrational binding energies and (b), the splitting between vibrational states localised in separate wells. The states localised in the inner-well are marked ' \blacklozenge '.

less than 0.8% and 7% respectively, providing us with an accurate estimate of the depth and shape of the potential energy curves. The vibrational ground states measured have large binding energies in comparison to the resolution, and are exceedingly S-wave dominant. This allowed for the extraction of low energy ${}^{3}S_{1}$ -wave scattering phase shift for 87 Rb with unprecedented accuracy in the relevant k-range. We argue that the accuracy achieved is largely limited by the predictive power of the Hamiltonian and an enhanced understanding of the molecular interactions is necessary to proceed further. The high resolution data provided is a strong benchmark to test future theoretical models of the ULRM, and can potentially lead to an even deeper insight into low-energy electron-atom scattering.

Acknowledgements: We would like to thank Max Althön for helpful discussions. This project is funded by the German science foundation DFG, project numbers 460443971 and 316211972.

Data availability: The data that support the findings of this study are available from the corresponding author upon reasonable request.

molecules, Phys. Rev. Lett. 85, 2458 (2000).

- [2] J. A. Fernández, H. R Sadeghpour, P. Schmelcher, und R. G Férez, Ultralong-Range Rb-KRb Rydberg Molecules: Selected Aspects of Electronic Structure, Orientation and Alignment, Journal of Physics: Conference Series 635, 012023 (2015).
- [3] M. Mayle, S. T. Rittenhouse, P. Schmelcher, und H. R. Sadeghpour, Electric field control in ultralong-range triatomic polar Rydberg molecules, Phys. Rev. A 85, 052511 (2012).
- [4] J. Zeiher, und S. Hollerith, Rydberg Macrodimers: Diatomic Molecules on the Micrometer Scale, J. Phys. Chem. A 127, 3925 (2023).
- [5] A. Duspayev, X. Han, M. A. Viray, L. Ma, L. Zhao, und G. Raithel, Long-range Rydberg-atom-ion molecules of Rb and Cs, Phys. Rev. Res. 3, 023114 (2021).
- [6] N. Zuber, V. S. V. Anasuri, M. Berngruber, Y. Q Zou, R. Löw, und T. Pfau, Observation of a molecular bond between ions and Rydberg atoms, Nature 52, 453 (2022).
- [7] D. Bosworth, F. Hummel, und P. Schmelcher, Charged ultralong-range Rydberg trimers, Physics. Rev. A 107, 022807 (2023).
- [8] M. T. Eiles, Trilobites, butterflies, and other exotic specimens of long-range Rydberg molecules, Journal of Physics B: Atomic, Molecular and Optical Physics 52, 113001 (2019).
- [9] C. Fey, F. Hummel, und P. Schmelcher, Ultralong-range Rydberg molecules, Mol. Phys. 118, e1679401 (2020).
- [10] F. B. Dunning, S. K. Kanungo, S. und Yoshida, Ultralong-range Rydberg molecules, J. Phys. B: At. Mol. Opt. Phys. 57, 212002 (2024).
- [11] E. L. Hamilton, C. H. Greene, und H. R. Sadeghpour, Shape-resonance-induced long-range molecular Rydberg states, Journal of Physics B: Atomic, Molecular and Optical Physics 35, L199 (2002).
- [12] W. Li, T. Pohl, J. M. Rost, Seth T. Rittenhouse, H. R. Sadeghpour, J. Nipper, B. Butscher, J. B. Balewski, V. Bendkowsky, R. Löw, und T. Pfau, A Homonuclear Molecule with a Permanent Electric Dipole Moment, Science **334**, 1110–1114 (2011).
- [13] M. I. Chibisov, A. A. Khuskivadze, und I. I. Fabrikant, Energies and dipole moments of long-range molecular Rydberg states, J. Phys. B: At. Mol. Opt. Phys. 35, L193 (2002).
- [14] D. Booth, S. T. Rittenhouse, J. Yang, H. R. Sadeghpour, und J. P. Shaffer, Production of trilobite Rydberg molecule dimers with kilo-Debye permanent electric dipole moments, Science **348**, 99–102 (2015).
- [15] M. Althön, M. Exner, R. Blättner, und H. Ott, Exploring the vibrational series of pure trilobite Rydberg molecules, Nature Communications 14, 8108 (2023).
- [16] F. Engel, S. K. Tiwari, T. Pfau, S. Wüster, und F. Meinert, In situ observation of chemistry in Rydberg molecules within a coherent solvent, arXiv preprint arXiv:2308.13762 (2023).
- [17] F. Hummel, K. Keiler, und P. Schmelcher, Electricfield-induced wave-packet dynamics and geometrical rearrangement of trilobite Rydberg molecules, Phys. Rev. A 103, 022827 (2021).
- [18] R. Srikumar, S. T. Rittenhouse, und P. Schmelcher, Internal diffraction dynamics of trilobite molecules, arXiv preprint arXiv:2408.02134 (2024).
- [19] K. S. Kleinbach, F. Meinert, F. Engel, W. J. Kwon, R. Löw, T. Pfau, und G. Raithel, Photoassociation of

^{*} These authors contributed equally to this work.

[†] Corresponding author: ott@physik.uni-kl.de

C. H. Greene, A. S. Dickinson, und H. R. Sadeghpour, Creation of polar and nonpolar ultra-long-range Rydberg

Trilobite Rydberg Molecules via Resonant Spin-Orbit Coupling, Phys. Rev. Lett. **118**, 223001 (2017).

- [20] V. Bendkowsky, B. Butscher, J. Nipper, J. P. Shaffer, R. Löw, und T. Pfau, Observation of ultralong-range Rydberg molecules, Nature 458, 1005–1008 (2009).
- [21] F. Engel, T. Dieterle, F. Hummel, C. Fey, P. Schmelcher, R. Löw, T. Pfau, und F. Meinert, Precision Spectroscopy of Negative-Ion Resonances in Ultralong-Range Rydberg Molecules, Phys. Rev. Lett. **123**, 073003 (2019).
- [22] T. Niederprüm, O. Thomas, T. Eichert, und H. Ott, Rydberg Molecule-Induced Remote Spin Flips, Phys. Rev. Lett. 117, 123002 (2016).
- [23] T. Niederprüm, O. Thomas, T. Eichert, C. Lippe, J. Pérez-Ríos, C. H. Greene, und H. Ott, Observation of pendular butterfly Rydberg molecules, Nature Communications 7, 12820 (2016).
- [24] M. Deiß, S. Haze, J. Wolf, L. Wang, F. Meinert, C. Fey, F. Hummel, P. Schmelcher, und J. H. Denschlag, Observation of spin-orbit-dependent electron scattering using long-range Rydberg molecules, Phys. Rev. Res. 2, 013047 (2020).
- [25] H. Saßmannshausen, F. Merkt, und J. Deiglmayr, Experimental Characterization of Singlet Scattering Channels in Long-Range Rydberg Molecules, Phys. Rev. Lett. 114, 133201 (2015).
- [26] M. Peper und J. Deiglmayr, Photodissociation of longrange Rydberg molecules, Phys. Rev. A 102, 062819 (2020).
- [27] M. Peper und J. Deiglmayr, Heteronuclear Long-Range Rydberg Molecules, Phys. Rev. Lett. **126**, 013001 (2021).
- [28] J. L. MacLennan, Y.-J. Chen, und G. Raithel, Deeply bound $(24D_J + 5S_{1/2})^{87}$ Rb and ⁸⁵Rb molecules for eight spin couplings, Phys. Rev. A **99**, 033407 (2019).
- [29] C. H. Greene, und M. T. Eiles, Green's-function treat-

ment of Rydberg molecules with spins, Phys. Rev. A **108**, 042805 (2023).

- [30] F. Hummel, P. Schmelcher, und M. T. Eiles, Vibronic interactions in trilobite and butterfly Rydberg molecules, Phys. Rev. Res. 5, 013114 (2023).
- [31] R. Srikumar, F. Hummel, und P. Schmelcher, Nonadiabatic interaction effects in the spectra of ultralong-range Rydberg molecules, Phys. Rev. A 108, 012809 (2023).
- [32] M. T. Eiles, und F. Hummel, Kato's theorem and ultralong-range Rydberg molecules, Phys. Rev. A 109, 022811 (2024).
- [33] M. T. Eiles, und C. H. Greene, Hamiltonian for the inclusion of spin effects in long-range Rydberg molecules, Phys. Rev. A 95, 042515 (2017).
- [34] E. Fermi, Sopra lo Spostamento per Pressione delle Righe Elevate delle Serie Spettrali, Nuovo Cim. 11, 157 (1934).
- [35] A. Omont, On the theory of collisions of atoms in rydberg states with neutral particles, J. Phys. France 38, 1343 (1977).
- [36] C. Fey, M. Kurz, P. Schmelcher, S. T. Rittenhouse, und H. R. Sadeghpour, A comparative analysis of binding in ultralong-range Rydberg molecules, New Journal of Physics 17, 055010 (2015).
- [37] M. Tarana, Long-range Rydberg molecule Rb₂: Twoelectron *R*-matrix calculations at intermediate internuclear distances, Phys. Rev. A **102**, 062802 (2020).
- [38] P. Giannakeas, M. T. Eiles, F. Robicheaux, und J. M. Rost, Generalized local frame-transformation theory for ultralong-range Rydberg molecules, Phys. Rev. A 102, 033315 (2020).
- [39] P. Geppert, M. Althön, D. Fichtner, and H. Ott, Diffusive-like redistribution in state-changing collisions between Rydberg atoms and ground state atoms, Nature Communications 12, 3900 (2021)