

P-wave bottom baryons of the $SU(3)$ flavor $\bar{\mathbf{3}}_F$

Yi-Jie Wang¹, Xuan Luo^{1,*}, Hua-Xing Chen^{1,†}, Er-Liang Cui^{2,‡}, Wei-Han Tan^{1,§} and Zhi-Yong Zhou^{1,¶}

¹*School of Physics, Southeast University, Nanjing 210094, China*

²*College of Science, Northwest A&F University, Yangling 712100, China*

We study the *P*-wave bottom baryons of the $SU(3)$ flavor $\bar{\mathbf{3}}_F$ and systematically calculate their strong decay properties, including their *D*-wave decays into ground-state bottom baryons with light pseudoscalar mesons and *S*-wave decays into ground-state bottom baryons with light vector mesons. Together with Refs. [54–57], a rather complete investigation has been performed to study their mass spectra and strong/radiative decay properties, through the methods of QCD sum rules and light-cone sum rules within the framework of heavy quark effective theory. Among various possibilities, we identify four Λ_b and four Ξ_b baryons, with limited decay widths and so capable of being observed in experiments. Their masses, mass splittings within the same multiplets, and strong/radiative decay widths are summarized in Table II for future experimental searching.

Keywords: excited bottom baryon, QCD sum rules, light-cone sum rules, heavy quark effective theory

I. INTRODUCTION

The singly bottom baryon is a three-quark system consisting of one heavy *bottom* quark and two light *up/down/strange* quarks [1–5]. Its research can provide crucial insights into the understanding of the non-perturbative QCD in the low energy region. Nowadays, all the ground-state bottom baryons, except the Ω_b^* baryon of $J^P = 3/2^+$, have been observed and well identified in experiments [6]. Besides, a significant number of excited bottom baryons have been reported, for examples:

- In 2012 the excited bottom baryons $\Lambda_b(5912)^0$ and $\Lambda_b(5920)^0$ were observed by LHCb [7] and CDF [8] in the $\Lambda_b^0\pi^+\pi^-$ invariant mass spectrum. Their masses and widths were measured to be [9]:

$$\Lambda_b(5912)^0 : M = 5912.19 \pm 0.17 \text{ MeV}, \\ \Gamma < 0.25 \text{ MeV at } 90\% \text{ CL}; \quad (1)$$

$$\Lambda_b(5920)^0 : M = 5920.09 \pm 0.17 \text{ MeV}, \\ \Gamma < 0.19 \text{ MeV at } 90\% \text{ CL}. \quad (2)$$

- In 2021 the CMS collaboration observed the $\Xi_b(6100)^-$ in the $\Xi_b^-\pi^+\pi^-$ invariant mass spectrum [10]. Later in 2023 the LHCb collaboration observed the $\Xi_b(6087)^0$ and $\Xi_b(6095)^0$ in the $\Xi_b^0\pi^+\pi^-$ invariant mass spectrum [11]. The $\Xi_b(6100)^-$ and $\Xi_b(6095)^0$ are probably isospin partners. Their masses and widths were measured to

be [9]:

$$\Xi_b(6100)^- : M = 6100.3 \pm 0.2 \pm 0.1 \pm 0.6 \text{ MeV}, \\ \Gamma = 0.94 \pm 0.30 \pm 0.08 \text{ MeV}; \quad (3)$$

$$\Xi_b(6095)^0 : M = 6095.4 \pm 0.15 \pm 0.03 \pm 0.5 \text{ MeV}, \\ \Gamma = 0.50 \pm 0.33 \pm 0.11 \text{ MeV}; \quad (4)$$

$$\Xi_b(6087)^0 : M = 6087.24 \pm 0.20 \pm 0.06 \pm 0.5 \text{ MeV}, \\ \Gamma = 2.43 \pm 0.51 \pm 0.10 \text{ MeV}. \quad (5)$$

These states are good candidates for the *P*-wave bottom baryons of the $SU(3)$ flavor $\bar{\mathbf{3}}_F$. Their detailed discussions can be found in the reviews [12–16]. Based on these observations, extensive theoretical investigations have been performed through a variety of phenomenological models, including various quark models [17–28], various hadronic molecular models [29–40], the quark pair creation model [41–43], the chiral perturbation theory [44–46], QCD sum rules [47–60], and Lattice QCD [61–64], etc.

We have studied the *P*-wave bottom baryons within the framework of heavy quark effective theory [65–67] in a series of works:

- In Ref. [53] we applied the QCD sum rule method [68–74] to systematically calculate their mass spectrum.
- In Ref. [54] we applied the light-cone sum rule method [75–84] to partly calculate their strong decay properties, including their *S*-wave decays into ground-state bottom baryons with light pseudoscalar mesons.
- In Ref. [55] we applied the light-cone sum rule method to systematically calculate their radiative decay properties.

Our results suggest that the $\Lambda_b(5912)^0$, $\Lambda_b(5920)^0$, $\Xi_b(6087)^0$, and $\Xi_b(6095)^0/\Xi_b(6100)^-$ can be well explained as the *P*-wave bottom baryons of the ρ -mode, where the orbital excitation is between the two light

*Electronic address: cnxluo@seu.edu.cn

†Electronic address: hxchen@seu.edu.cn

‡Electronic address: erliang.cui@nwafu.edu.cn

§Electronic address: tanwh@seu.edu.cn

¶Electronic address: zhouzhy@seu.edu.cn

quarks, as depicted in Fig. 1. This interpretation differs from the λ -mode interpretation supported by the quark model calculations [21–25], where the orbital angular momentum is between the bottom quark and the light diquark. Hence, further experimental and theoretical studies are necessary to improve our understanding on their internal structure.




FIG. 1: Jacobi coordinates $\vec{\lambda}$ and \vec{p} for the singly bottom baryon system.

In this paper we shall investigate the P -wave bottom baryons of the $SU(3)$ flavor $\bar{\mathbf{3}}_F$ and systematically calculate their strong decay properties, including their D -wave decays into ground-state bottom baryons with light pseudoscalar mesons and S -wave decays into ground-state bottom baryons with light vector mesons. Together with Refs. [53–55], we shall arrive at a rather complete investigation on the P -wave bottom baryons of the $SU(3)$ flavor $\bar{\mathbf{3}}_F$, allowing us to understand them as a whole. Note that similar investigations have been performed in Refs. [56, 57] to study the P -wave charmed and bottom baryons of the $SU(3)$ flavor $\mathbf{6}_F$.

This paper is organized as follows. In Sec. II we briefly introduce our notations to describe the P -wave bottom baryons of the $SU(3)$ flavor $\bar{\mathbf{3}}_F$. In Sec. III we study their strong decay properties through the light-cone sum rule method. In Sec. IV we discuss the obtained results and conclude this paper.

II. MASS SPECTRUM

In this section we briefly introduce our notations. A singly bottom baryon is composed of one heavy *bottom* quark and two light *up/down/strange* quarks. To well understand this system, we need to meticulously analyze the symmetries between the two light quarks, including the color, flavor, spin, and orbital degrees of freedom:

- The two light quarks always have the antisymmetric color structure $\bar{\mathbf{3}}_C$.
- Their $SU(3)$ flavor structure is either symmetric ($\mathbf{6}_F$) or antisymmetric ($\bar{\mathbf{3}}_F$).
- Their spin structure is either symmetric ($s_l = 1$) or antisymmetric ($s_l = 0$), with s_l the total spin of the two light quarks.

- Their orbital structure is either symmetric ($l_\rho = 0/2/4/\cdots$) or antisymmetric ($l_\rho = 1/3/5/\cdots$), with l_ρ the orbital angular momentum between the two light quarks, as depicted in Fig. 1. Besides, we use l_λ to denote the orbital angular momentum between the bottom quark and the light diquark. There may exist both the ρ -mode P -wave bottom baryons ($l_\rho = 1$ and $l_\lambda = 0$) and λ -mode ones ($l_\rho = 0$ and $l_\lambda = 1$).

In the context of the Pauli principle, the total symmetry of the two light quarks is antisymmetric. Consequently, we can categorize the P -wave bottom baryons into eight multiplets, four of which belong to the $SU(3)$ flavor $\bar{\mathbf{3}}_F$ representation, as illustrated in Fig. 2. We denote them as $[flavor, j_l, s_l, \rho/\lambda]$, with $j_l = l_\rho \otimes l_\lambda \otimes s_l$ the total angular momentum of the light component. Each multiplet contains one or two bottom baryons, having the total angular momenta $J = j_l \otimes s_Q = |j_l \pm 1/2|$.

The mass spectrum of the P -wave bottom baryons belonging to the $SU(3)$ flavor $\bar{\mathbf{3}}_F$ representation has been systematically studied in Ref. [54] through the QCD sum rule method within the framework of heavy quark effective theory, as summarized in Table I. In the next section we shall use these parameters to study their strong decay properties through the light-cone sum rule method still within the framework of heavy quark effective theory. Besides, we need the following parameters for the ground-state bottom baryons and light pseudoscalar/vector mesons:

- For the ground-state bottom baryons, we use

$$\begin{aligned} \Lambda_b(1/2^+) &: M = 5619.60 \text{ MeV}, \\ \Xi_b(1/2^+) &: M = 5793.20 \text{ MeV}, \\ \Sigma_b(1/2^+) &: M = 5813.4 \text{ MeV}, \\ \Sigma_b^*(3/2^+) &: M = 5833.6 \text{ MeV}, \\ \Xi'_b(1/2^+) &: M = 5935.02 \text{ MeV}, \\ \Xi_b^*(3/2^+) &: M = 5952.6 \text{ MeV}, \\ \Omega_b(1/2^+) &: M = 6046.1 \text{ MeV}, \\ \Omega_b^*(1/2^+) &: M = 6063 \text{ MeV}. \end{aligned}$$

- For the light pseudoscalar and vector mesons, we use

$$\begin{aligned} \pi(0^-) &: m = 138.04 \text{ MeV}, \\ K(0^-) &: m = 495.65 \text{ MeV}, \\ \rho(1^-) &: m = 775.21 \text{ MeV}, \\ &\Gamma = 148.2 \text{ MeV}, \\ g_{\rho\pi\pi} &= 5.94 \text{ GeV}^{-2}, \\ K^*(1^-) &: m = 893.57 \text{ MeV}, \\ &\Gamma = 49.1 \text{ MeV}, \\ g_{K^*K\pi} &= 3.20 \text{ GeV}^{-2}. \end{aligned}$$




FIG. 2: Categorization of the P -wave bottom baryons belonging to the $SU(3)$ flavor $\bar{\mathbf{3}}_F$ representation.

TABLE I: Mass spectrum of the P -wave bottom baryons belonging to the $SU(3)$ flavor $\bar{\mathbf{3}}_F$ representation, calculated through the QCD sum rule method within the framework of heavy quark effective theory. Decay constants in the last column satisfy $f_{\Xi_b^0} = f_{\Xi_b^-}$. Taken from Ref. [54].

Multiplets	B	ω_c (GeV)	Working region (GeV)	$\bar{\Lambda}$ (GeV)	Baryon (j^P)	Mass (GeV)	Difference (MeV)	Decay constant (GeV 4)
$[\bar{\mathbf{3}}_F, 1, 0, \lambda]$	Λ_b	2.19 ± 0.10	$0.26 < T < 0.40$	$1.53^{+0.14}_{-0.09}$	$\Lambda_b(1/2^-)$	$5.91^{+0.17}_{-0.13}$	4 ± 2	$0.081^{+0.020}_{-0.018} (\Lambda_b^0(1/2^-))$
	Λ_b	2.19 ± 0.10			$\Lambda_b(3/2^-)$	$5.91^{+0.17}_{-0.13}$		$0.038^{+0.009}_{-0.008} (\Lambda_b^0(3/2^-))$
	Ξ_b	2.36 ± 0.10	$0.25 < T < 0.43$	$1.68^{+0.17}_{-0.07}$	$\Xi_b(1/2^-)$	$6.10^{+0.20}_{-0.10}$	4 ± 2	$0.110^{+0.030}_{-0.016} (\Xi_b^-(1/2^-))$
	Ξ_b	2.36 ± 0.10			$\Xi_b(3/2^-)$	$6.10^{+0.20}_{-0.10}$		$0.052^{+0.014}_{-0.008} (\Xi_b^-(3/2^-))$
$[\bar{\mathbf{3}}_F, 0, 1, \rho]$	Λ_b	2.33 ± 0.10	$0.35 < T < 0.44$	$1.56^{+0.14}_{-0.16}$	$\Lambda_b(1/2^-)$	$5.92^{+0.17}_{-0.19}$	-	$0.059^{+0.013}_{-0.013} (\Lambda_b^0(1/2^-))$
	Ξ_b	2.29 ± 0.10	$T = 0.45, PC = 34\%$	$1.72^{+0.07}_{-0.07}$	$\Xi_b(1/2^-)$	$6.10^{+0.08}_{-0.08}$	-	$0.078^{+0.011}_{-0.009} (\Xi_b^-(1/2^-))$
$[\bar{\mathbf{3}}_F, 1, 1, \rho]$	Λ_b	2.17 ± 0.10	$0.26 < T < 0.39$	$1.60^{+0.11}_{-0.08}$	$\Lambda_b(1/2^-)$	$5.92^{+0.13}_{-0.10}$	8 ± 3	$0.169^{+0.036}_{-0.027} (\Lambda_b^0(1/2^-))$
	Λ_b	2.17 ± 0.10			$\Lambda_b(3/2^-)$	$5.92^{+0.13}_{-0.10}$		$0.080^{+0.017}_{-0.013} (\Lambda_b^0(3/2^-))$
	Ξ_b	2.32 ± 0.10	$0.27 < T < 0.41$	$1.75^{+0.11}_{-0.09}$	$\Xi_b(1/2^-)$	$6.09^{+0.13}_{-0.12}$	7 ± 3	$0.222^{+0.069}_{-0.039} (\Xi_b^-(1/2^-))$
	Ξ_b	2.32 ± 0.10			$\Xi_b(3/2^-)$	$6.09^{+0.13}_{-0.12}$		$0.105^{+0.033}_{-0.018} (\Xi_b^-(3/2^-))$
$[\bar{\mathbf{3}}_F, 2, 1, \rho]$	Λ_b	2.17 ± 0.10	$0.30 < T < 0.37$	$1.64^{+0.11}_{-0.11}$	$\Lambda_b(3/2^-)$	$5.93^{+0.13}_{-0.13}$	17 ± 8	$0.136^{+0.031}_{-0.029} (\Lambda_b^0(1/2^-))$
	Λ_b	2.17 ± 0.10	$\Lambda_b(5/2^-)$	$5.94^{+0.13}_{-0.13}$	$0.058^{+0.013}_{-0.012} (\Lambda_b^0(5/2^-))$			
	Ξ_b	2.31 ± 0.10	$0.31 < T < 0.40$	$1.80^{+0.09}_{-0.10}$	$\Xi_b(3/2^-)$	$6.10^{+0.15}_{-0.10}$	14 ± 7	$0.184^{+0.038}_{-0.036} (\Xi_b^-(3/2^-))$
	Ξ_b	2.31 ± 0.10	$\Xi_b(5/2^-)$	$6.11^{+0.15}_{-0.10}$	$0.078^{+0.160}_{-0.150} (\Xi_b^-(5/2^-))$			

III. STRONG DECAY PROPERTIES

In this section we study strong decay properties of the P -wave bottom baryons belonging to the $SU(3)$ flavor $\bar{\mathbf{3}}_F$

representation through the light-cone sum rule method within the framework of heavy quark effective theory. We shall investigate their S/D -wave decays into ground-state bottom baryons with light pseudoscalar/vector mesons:

$$(a1) \quad \Gamma[\Lambda_b[1/2^-, 1P] \rightarrow \Lambda_b + \pi] = \Gamma[\Lambda_b^0[1/2^-, 1P] \rightarrow \Lambda_b^0 + \pi^0], \quad (6)$$

$$(a2) \quad \Gamma[\Lambda_b[1/2^-, 1P] \rightarrow \Sigma_b + \pi] = 3 \times \Gamma[\Lambda_b^0[1/2^-, 1P] \rightarrow \Sigma_b^+ + \pi^- \rightarrow \Lambda_b^0 \pi^+ + \pi^-], \quad (7)$$

- (a3) $\Gamma[\Lambda_b[1/2^-, 1P] \rightarrow \Sigma_b^* + \pi \rightarrow \Lambda_b + \pi + \pi] = 3 \times \Gamma[\Lambda_b^0[1/2^-, 1P] \rightarrow \Sigma_b^{*+} + \pi^- \rightarrow \Lambda_b^0 + \pi^+ + \pi^-]$, (8)
- (a4) $\Gamma[\Lambda_b[1/2^-, 1P] \rightarrow \Lambda_b + \rho \rightarrow \Lambda_b + \pi + \pi] = \Gamma[\Lambda_b^0[1/2^-, 1P] \rightarrow \Lambda_b^0 + \pi^+ + \pi^-]$, (9)
- (a5) $\Gamma[\Lambda_b[1/2^-, 1P] \rightarrow \Sigma_b + \rho \rightarrow \Sigma_b + \pi + \pi] = 3 \times \Gamma[\Lambda_b^0[1/2^-, 1P] \rightarrow \Sigma_b^0 + \pi^+ + \pi^-]$, (10)
- (a6) $\Gamma[\Lambda_b[1/2^-, 1P] \rightarrow \Sigma_b^* + \rho \rightarrow \Sigma_b^* + \pi + \pi] = 3 \times \Gamma[\Lambda_b^0[1/2^-, 1P] \rightarrow \Sigma_b^{*0} + \pi^+ + \pi^-]$, (11)
- (b1) $\Gamma[\Lambda_b[3/2^-, 1P] \rightarrow \Lambda_b + \pi] = \Gamma[\Lambda_b^0[3/2^-] \rightarrow \Lambda_b^0 + \pi^0]$, (12)
- (b2) $\Gamma[\Lambda_b[3/2^-, 1P] \rightarrow \Sigma_b + \pi \rightarrow \Lambda_b + \pi + \pi] = 3 \times \Gamma[\Lambda_b^0[3/2^-, 1P] \rightarrow \Sigma_b^+ + \pi^- \rightarrow \Lambda_b^0 \pi^+ + \pi^-]$, (13)
- (b3) $\Gamma[\Lambda_b[3/2^-, 1P] \rightarrow \Sigma_b^* + \pi \rightarrow \Lambda_b + \pi + \pi] = 3 \times \Gamma[\Lambda_b^0[3/2^-, 1P] \rightarrow \Sigma_b^{*+} + \pi^- \rightarrow \Lambda_b^0 \pi^+ + \pi^-]$, (14)
- (b4) $\Gamma[\Lambda_b[3/2^-, 1P] \rightarrow \Lambda_b + \rho \rightarrow \Lambda_b + \pi + \pi] = \Gamma[\Lambda_b^0[3/2^-, 1P] \rightarrow \Lambda_b^0 + \pi^+ + \pi^-]$, (15)
- (b5) $\Gamma[\Lambda_b[3/2^-, 1P] \rightarrow \Sigma_b + \rho \rightarrow \Sigma_b + \pi + \pi] = 3 \times \Gamma[\Lambda_b^0[3/2^-, 1P] \rightarrow \Sigma_b^0 \pi^+ + \pi^-]$, (16)
- (b6) $\Gamma[\Lambda_b[3/2^-, 1P] \rightarrow \Sigma_b^* + \rho \rightarrow \Sigma_c^* + \pi + \pi] = 3 \times \Gamma[\Lambda_b^0[3/2^-, 1P] \rightarrow \Sigma_b^{*+} + \pi^+ + \pi^-]$, (17)
- (c1) $\Gamma[\Xi_b[1/2^-, 1P] \rightarrow \Lambda_b + \bar{K}] = \Gamma[\Xi_b^-[1/2^-, 1P] \rightarrow \Lambda_b^0 + K^-]$, (18)
- (c2) $\Gamma[\Xi_b[1/2^-, 1P] \rightarrow \Xi_b + \pi] = \frac{3}{2} \times \Gamma[\Xi_b^-[1/2^-, 1P] \rightarrow \Xi_b^0 + \pi^-]$, (19)
- (c3) $\Gamma[\Xi_b[1/2^-, 1P] \rightarrow \Sigma_b + \bar{K}] = 3 \times \Gamma[\Xi_b^-[1/2^-, 1P] \rightarrow \Sigma_b^0 + K^-]$, (20)
- (c4) $\Gamma[\Xi_b[1/2^-, 1P] \rightarrow \Xi'_b + \pi] = \frac{3}{2} \times \Gamma[\Xi_b^-[1/2^-, 1P] \rightarrow \Xi'^0_b + \pi^-]$, (21)
- (c5) $\Gamma[\Xi_b[1/2^-, 1P] \rightarrow \Sigma_b^* + K] = 3 \times \Gamma[\Xi_b^-[1/2^-, 1P] \rightarrow \Sigma_b^{*0} + K^-]$, (22)
- (c6) $\Gamma[\Xi_b[1/2^-, 1P] \rightarrow \Xi_b^* + \pi \rightarrow \Xi_b + \pi + \pi] = \frac{9}{2} \times \Gamma[\Xi_b^-[1/2^-, 1P] \rightarrow \Xi_b^{*0} + \pi^- \rightarrow \Xi_b^0 + \pi^0 + \pi^-]$, (23)
- (c7) $\Gamma[\Xi_b[1/2^-, 1P] \rightarrow \Lambda_b + \bar{K}^* \rightarrow \Lambda_b + \bar{K} + \pi] = 3 \times \Gamma[\Xi_b^-[1/2^-, 1P] \rightarrow \Lambda_b^0 + K^- + \pi^0]$, (24)
- (c8) $\Gamma[\Xi_b[1/2^-, 1P] \rightarrow \Xi_b + \rho \rightarrow \Xi_c + \pi + \pi] = \frac{3}{2} \times \Gamma[\Xi_b^-[1/2^-, 1P] \rightarrow \Xi_b^0 + \pi^0 + \pi^-]$, (25)
- (c9) $\Gamma[\Xi_b[1/2^-, 1P] \rightarrow \Sigma_b^* + \bar{K}^* \rightarrow \Sigma_b^* + \bar{K} + \pi] = 9 \times \Gamma[\Xi_b^-[1/2^-, 1P] \rightarrow \Sigma_b^{*0} + K^- + \pi^0]$, (26)
- (c10) $\Gamma[\Xi_b[1/2^-, 1P] \rightarrow \Xi_b^* + \rho \rightarrow \Xi_b^* + \pi + \pi] = \frac{3}{2} \times \Gamma[\Xi_b^-[1/2^-, 1P] \rightarrow \Xi_b^{*0} + \pi^0 + \pi^-]$, (27)
- (d1) $\Gamma[\Xi_b[3/2^-, 1P] \rightarrow \Lambda_b + \bar{K}] = \Gamma[\Xi_b^-[3/2^-, 1P] \rightarrow \Lambda_b^0 + K^-]$, (28)
- (d2) $\Gamma[\Xi_b[3/2^-, 1P] \rightarrow \Xi_b + \pi] = \frac{3}{2} \times \Gamma[\Xi_b^-[3/2^-, 1P] \rightarrow \Xi_b^0 + \pi^-]$, (29)
- (d3) $\Gamma[\Xi_b[3/2^-, 1P] \rightarrow \Sigma_b + \bar{K}] = 3 \times \Gamma[\Xi_b^-[3/2^-, 1P] \rightarrow \Sigma_b^0 + K^-]$, (30)
- (d4) $\Gamma[\Xi_b[3/2^-, 1P] \rightarrow \Xi'_b + \pi] = \frac{3}{2} \times \Gamma[\Xi_b^-[3/2^-, 1P] \rightarrow \Xi'^0_b + \pi^-]$, (31)
- (d5) $\Gamma[\Xi_b[3/2^-, 1P] \rightarrow \Sigma_b^* + \bar{K}] = 3 \times \Gamma[\Xi_b^-[3/2^-, 1P] \rightarrow \Sigma_b^{*0} + K^-]$, (32)
- (d6) $\Gamma[\Xi_b[3/2^-, 1P] \rightarrow \Xi_b^* + \pi \rightarrow \Xi_b + \pi + \pi] = \frac{9}{2} \times \Gamma[\Xi_b^{*-}[3/2^-, 1P] \rightarrow \Xi_b^{*0} + \pi^- \rightarrow \Xi_b^0 + \pi^0 + \pi^-]$, (33)
- (d7) $\Gamma[\Xi_b[3/2^-, 1P] \rightarrow \Lambda_b + \bar{K}^* \rightarrow \Lambda_b + \bar{K} + \pi] = 3 \times \Gamma[\Xi_b^-[3/2^-, 1P] \rightarrow \Lambda_b^0 + K^- + \pi^0]$, (34)
- (d8) $\Gamma[\Xi_b[3/2^-, 1P] \rightarrow \Xi_b + \rho \rightarrow \Xi_b + \pi + \pi] = \frac{3}{2} \times \Gamma[\Xi_b^-[3/2^-, 1P] \rightarrow \Xi_b^0 + \pi^0 + \pi^-]$, (35)
- (d9) $\Gamma[\Xi_b[3/2^-, 1P] \rightarrow \Sigma_b^* + \bar{K}^* \rightarrow \Sigma_b^{*0} + \bar{K} + \pi] = 9 \times \Gamma[\Xi_b^-[3/2^-, 1P] \rightarrow \Sigma_b^{*0} + K^- + \pi^0]$, (36)

$$(d10) \quad \mathbf{\Gamma} \left[\Xi_b[3/2^-, 1P] \rightarrow \Xi_b^* + \rho \rightarrow \Xi_b^* + \pi + \pi \right] = \frac{3}{2} \times \mathbf{\Gamma} \left[\Xi_b^- [3/2^-, 1P] \rightarrow \Xi_b^{*0} + \pi^0 + \pi^- \right]. \quad (37)$$

Their partial decay widths can be evaluated through the following Lagrangians:

$$\mathcal{L}_{X_b(1/2^-) \rightarrow Y_b(1/2^+)P}^S = g^S \bar{X}_b(1/2^-) Y_b(1/2^+) P, \quad (38)$$

$$\mathcal{L}_{X_b(3/2^-) \rightarrow Y_b(3/2^+)P}^S = g^S \bar{X}_b(3/2^-) Y_b^\mu(3/2^+) P, \quad (39)$$

$$\mathcal{L}_{X_b(1/2^-) \rightarrow Y_b(1/2^+)V}^S = g^S \bar{X}_b(1/2^-) \gamma_\mu \gamma_5 Y_b(1/2^+) V^\mu, \quad (40)$$

$$\mathcal{L}_{X_b(1/2^-) \rightarrow Y_b(3/2^+)V}^S = g^S \bar{X}_b(1/2^-) Y_b^\mu(3/2^+) V_\mu, \quad (41)$$

$$\mathcal{L}_{X_b(3/2^-) \rightarrow Y_b(1/2^+)V}^S = g^S \bar{X}_b^\mu(3/2^-) Y_b(1/2^+) V_\mu, \quad (42)$$

$$\mathcal{L}_{X_b(3/2^-) \rightarrow Y_b(3/2^+)V}^S = g^S \bar{X}_b^\nu(3/2^-) \gamma_\mu \gamma_5 Y_{b\nu}(3/2^+) V^\mu, \quad (43)$$

$$\mathcal{L}_{X_b(5/2^-) \rightarrow Y_b(3/2^+)V}^S = g^S \bar{X}_b^{\mu\nu}(5/2^-) Y_{b\mu}(3/2^+) V_\nu + g^S \bar{X}_b^{\nu\mu}(5/2^-) Y_{b\mu}(3/2^+) V_\nu, \quad (44)$$

$$\mathcal{L}_{X_b(1/2^-) \rightarrow Y_b(3/2^+)P}^D = g^D \bar{X}_b(1/2^-) \gamma_\mu \gamma_5 Y_{b\nu}(3/2^+) \partial^\mu \partial^\nu P, \quad (45)$$

$$\mathcal{L}_{X_b(3/2^-) \rightarrow Y_b(1/2^+)P}^D = g^D \bar{X}_{b\mu}(3/2^-) \gamma_\nu \gamma_5 Y_b(1/2^+) \partial^\mu \partial^\nu P, \quad (46)$$

$$\mathcal{L}_{X_b(3/2^-) \rightarrow Y_b(3/2^+)P}^D = g^D \bar{X}_{b\mu}(3/2^-) Y_{b\nu}(3/2^+) \partial^\mu \partial^\nu P, \quad (47)$$

$$\mathcal{L}_{X_b(5/2^-) \rightarrow Y_b(1/2^+)P}^D = g^D \bar{X}_{b\mu\nu}(5/2^-) Y_b(1/2^+) \partial^\mu \partial^\nu P, \quad (48)$$

$$\begin{aligned} & \mathcal{L}_{X_b(5/2^-) \rightarrow Y_b(3/2^+)P}^D \\ &= g^D \bar{X}_{b\mu\nu}(5/2^-) \gamma_\rho \gamma_5 Y_b^\mu(3/2^+) \partial^\nu \partial^\rho P \\ &+ g^D \bar{X}_{b\mu\nu}(5/2^-) \gamma_\rho \gamma_5 Y_b^\nu(3/2^+) \partial^\mu \partial^\rho P. \end{aligned} \quad (49)$$

In the above expressions, S and D denote the S - and D -wave decays, respectively; the fields $X_b^{(\mu\nu)}$ and $Y_b^{(\mu)}$ denote the P -wave and ground-state bottom baryons, respectively.

We select the P -wave bottom baryon $\Lambda_b^0(3/2^-)$ belonging to the $[\mathbf{3}_F, 1, 1, \rho]$ doublet as an example, and calculate its D -wave decay into $\Sigma_b^+ \pi^-$. To achieve this, we consider the two-point correlation function:

$$\begin{aligned} & \Pi^\alpha(\omega, \omega') \\ &= \int d^4x e^{-ik \cdot x} \langle 0 | J_{3/2-, -\Lambda_b^0, 1, 1, \rho}^\alpha(0) \bar{J}_{\Sigma_b^+}(x) | \pi^-(q) \rangle \\ &= \frac{1+\not{v}}{2} G_{\Lambda_b^0[\frac{3}{2}^-] \rightarrow \Sigma_b^+ \pi^-}^\alpha(\omega, \omega'), \end{aligned} \quad (50)$$

where $k' = k + q$, $\omega = v \cdot k$, and $\omega' = v \cdot k'$.

At the hadron level, we write $G_{\Lambda_b^0[\frac{3}{2}^-] \rightarrow \Sigma_b^+ \pi^-}^\alpha$ as

$$\begin{aligned} & G_{\Lambda_b^0[\frac{3}{2}^-] \rightarrow \Sigma_b^+ \pi^-}^\alpha(\omega, \omega') \\ &= g_{\Lambda_b^0[\frac{3}{2}^-] \rightarrow \Sigma_b^+ \pi^-}^D \frac{f_{\Lambda_b^0[\frac{3}{2}^-]} f_{\Sigma_b^+}}{(\bar{\Lambda}_{\Lambda_b^0[\frac{3}{2}^-]} - \omega')(\bar{\Lambda}_{\Sigma_b^+} - \omega)} \gamma_\alpha \gamma_5 + \dots, \end{aligned} \quad (51)$$

where \dots contains other possible amplitudes.

At the quark-gluon level, we calculate $G_{\Lambda_b^0[\frac{3}{2}^-] \rightarrow \Sigma_b^+ \pi^-}^\alpha$ using the method of operator product expansion(OPE):

$$\begin{aligned} & G_{\Lambda_b^0[\frac{3}{2}^-] \rightarrow \Sigma_b^+ \pi^-}^\alpha(\omega, \omega') \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega't} e^{iu\omega t} \times 4 \times \left(-\frac{f_\pi u_0}{4\pi^2 t^3} \phi_{2;\pi} + \frac{f_\pi u_0}{64\pi^2 t} \phi_{4;\pi}(u_0) + \frac{f_\pi m_\pi^2 u_0 t}{144(m_u)} \langle \bar{q}q \rangle \phi_3^\sigma(u_0) \right. \\ & - \frac{f_\pi m_\pi^2 u_0 t^3}{2304(m_u)} \langle g_s \sigma G s \rangle \phi_3^\sigma(u_0)) \times \gamma_\alpha \gamma_5 + \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega't(\alpha_2 + u\alpha_3)} e^{i\omega t(1-\alpha_2 - u\alpha_3)} \times \frac{1}{2} \times \left(\left(\frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) \alpha_3 u_0}{24\pi^2 t} \right. \right. \\ & + \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) \alpha_2 u_0}{24\pi^2 t} + \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) \alpha_3 u_0}{48\pi^2 t} + \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha}) \alpha_3 u_0}{48\pi^2 t} - \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) u_0}{24\pi^2 t} + \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) \alpha_2}{48\pi^2 t} + \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha}) \alpha_2}{48\pi^2 t} \\ & - \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha})}{48\pi^2 t} - \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha})}{48\pi^2 t}) - \left(\frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) u_0}{12\pi^2 t^2(v \cdot q)} + \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha}) u_0}{4\pi^2 t^2(v \cdot q)} + \frac{f_\pi \Psi_{4;\pi}(\underline{\alpha}) u_0}{24\pi^2 t^2(v \cdot q)} + \frac{f_\pi \tilde{\Psi}_{4;\pi}(\underline{\alpha}) u_0}{8\pi^2 t^2(v \cdot q)} + \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha})}{8\pi^2 t^2(v \cdot q)} \right. \\ & \left. \left. + \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha})}{24\pi^2 t^2(v \cdot q)} + \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha})}{24\pi^2 t^2(v \cdot q)} - \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha})}{24\pi^2 t^2(v \cdot q)} \right) \right) \times \gamma_\alpha \gamma_5. \end{aligned} \quad (52)$$

Then we perform the Borel transformation to both Eq. (51) at the hadron level and Eq. (53) at the quark-gluon level:

$$\begin{aligned}
& g_{\Lambda_b^0[3/2^-] \rightarrow \Sigma_b^+ \pi^-}^D f_{\Lambda_b^0[3/2^-]} f_{\Sigma_b^+} e^{-\frac{\bar{\Lambda}_{\Lambda_b^0[3/2^-]}}{T_1}} e^{-\frac{\bar{\Lambda}_{\Sigma_b^+}}{T_2}} \\
&= 4 \times \left(-\frac{f_\pi u_0}{4\pi^2} T^4 f_3(\frac{\omega_c}{T}) \phi_{2;\pi}(u_0)|_{u_0=\frac{1}{2}} + \frac{f_\pi u_0}{64\pi^2} T^2 f_1(\frac{\omega_c}{T}) \phi_{4;\pi}(u_0) u_0|_{u_0=\frac{1}{2}} + \frac{f_\pi m_\pi^2 u_0}{144(m_u)} \langle \bar{q}q \rangle \phi_3^\sigma(u_0)|_{u_0=\frac{1}{2}} \right. \\
&- \frac{f_\pi m_\pi^2 u_0}{2304(m_u)} T^{-2} \langle g_s \sigma G s \rangle \phi_3^\sigma(u_0)|_{u_0=\frac{1}{2}} + \frac{1}{2} \times (f_1(\frac{\omega_c}{T}) \int_0^{\frac{1}{2}} d\alpha_2 \int_{\frac{1}{2}-\alpha_2}^{1-\alpha_2} d\alpha_3 \frac{T^2}{\alpha_3} (\frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) \alpha_3 u_0}{24\pi^2} + \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) \alpha_2 u_0}{24\pi^2} \\
&+ \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) \alpha_3 u_0}{48\pi^2} + \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha}) \alpha_3 u_0}{48\pi^2} - \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) u_0}{24\pi^2} + \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) \alpha_2}{48\pi^2} + \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha}) \alpha_2}{48\pi^2} - \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha})}{48\pi^2} - \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha})}{48\pi^2})) \\
&- \frac{1}{2} \times (f_1(\frac{\omega_c}{T}) \int_0^{\frac{1}{2}} d\alpha_2 \int_{\frac{1}{2}-\alpha_2}^{1-\alpha_2} d\alpha_3 \frac{u_0 T^2}{\alpha_4} (\frac{f_\pi \Phi_{4;\pi}(\underline{\alpha}) u_0}{12\pi^2} + \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha}) u_0}{4\pi^2} + \frac{f_\pi \Psi_{4;\pi}(\underline{\alpha}) u_0}{24\pi^2} + \frac{f_\pi \tilde{\Psi}_{4;\pi}(\underline{\alpha}) u_0}{8\pi^2} \\
&+ \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha})}{8\pi^2} + \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha})}{24\pi^2} + \frac{f_\pi \Phi_{4;\pi}(\underline{\alpha})}{24\pi^2} - \frac{f_\pi \tilde{\Phi}_{4;\pi}(\underline{\alpha})}{24\pi^2})).
\end{aligned} \tag{53}$$

In the above expressions, $f_n(x) \equiv 1 - e^{-x} \sum_{k=0}^n \frac{x^k}{k!}$; the parameters ω and ω' are transformed to be T_1 and T_2 , respectively; we select the symmetric point $T_1 = T_2 = 2T$ so that $u_0 = \frac{T_1}{T_1+T_2} = \frac{1}{2}$; we choose $\omega_c = 2.17$ GeV to be the averaged threshold value of the $\Lambda_b^0(3/2^-)$ and $\Sigma_b^+(3/2^+)$ mass sum rules; we select 0.26 GeV $< T < 0.39$ GeV to be the Borel window of the $\Lambda_b^0(3/2^-)$ mass sum rule. The light-cone distribution amplitudes contained in the above sum rule expressions can be found in Refs. [77–84].

In the present work we use the following values for various quark and gluon parameters at the renormalization scale 1 GeV [68, 69, 85–90]:

$$\begin{aligned}
\langle \bar{q}q \rangle &= -(0.24 \pm 0.01)^3 \text{ GeV}^3, \\
\langle \bar{s}s \rangle &= (0.8 \pm 0.1) \times \langle \bar{q}q \rangle, \\
\langle g_s \bar{q} \sigma G q \rangle &= M_0^2 \times \langle \bar{q}q \rangle, \\
\langle g_s \bar{s} \sigma G s \rangle &= M_0^2 \times \langle \bar{s}s \rangle, \\
M_0^2 &= 0.8 \text{ GeV}^2, \\
\langle g_s^2 GG \rangle &= (0.48 \pm 0.14) \text{ GeV}^4.
\end{aligned} \tag{54}$$

As depicted in Fig. 3, we extract the coupling constant from Eq. (53) to be

$$\begin{aligned}
g_{\Lambda_b^0[3/2^-] \rightarrow \Sigma_b^+ \pi^-}^D &= 1.36^{+0.11}_{-0.10} {}^{+0.25}_{-0.15} {}^{+0.28}_{-0.23} {}^{+0.48}_{-0.49} \text{ GeV}^{-2} \\
&= 1.36^{+0.94}_{-0.70} \text{ GeV}^{-2},
\end{aligned}$$

where the uncertainties come from the Borel mass, parameters of $\Sigma_b^+(1/2^+)$, parameters of $\Lambda_b^0(3/2^-)$, and various QCD parameters given in Eqs. (54), respectively.

The D -wave decay widths of P -wave bottom baryons can be calculated through the Lagrangian given in Eq. (47):

$$\begin{aligned}
& \Gamma^D(\Lambda_b(3/2^-) \rightarrow \Sigma_b + \pi) \\
&= 3 \times \Gamma^D(\Lambda_b^0(3/2^-) \rightarrow \Sigma_b^+ + \pi^-)
\end{aligned} \tag{55}$$




FIG. 3: Coupling constant $g_{\Lambda_b^0[3/2^-] \rightarrow \Sigma_b^+ \pi^-}^D$ with respect to the Borel mass T . Here the bottom baryon doublet $[\bar{\mathbf{3}}_F, 1, 1, \rho]$ is investigated.

$$\begin{aligned}
&= \frac{3|\vec{p}_2|g_0^2}{32\pi m_0^2} \times \text{Tr}[(p/\not+m_0)\gamma_5 \gamma_\beta p_1(p/\not+m_1)\gamma_\beta \gamma_5] \\
&\times (g_{\alpha_1 \alpha_2} - \frac{1}{3}\gamma_{\alpha_2} \gamma_{\alpha_1} - \frac{p_0^{\alpha_2} \gamma_{\alpha_1} - p_0^{\alpha_1} \gamma_{\alpha_2}}{3m_0} - \frac{2p_0^{\alpha_2} p_0^{\alpha_1}}{3m_0^2}) \\
&\times p_2^{\alpha_1} p_2^{\alpha_2} p_2^{\beta_1} p_2^{\beta_2},
\end{aligned}$$

where 0, 1, and 2 denote $\Lambda_b^0(3/2^-)$, $\Sigma_b^0(1/2^+)$, and π^- , respectively. Numerically, we obtain

$$\Gamma_{\Lambda_b^0[3/2^-] \rightarrow \Sigma_c \pi}^D = 1.67^{+3.08}_{-1.27} \times 10^{-3} \text{ MeV}. \tag{56}$$

Similarly, we study the four $SU(3)$ flavor $\bar{\mathbf{3}}_F$ bottom baryon multiplets, $[\bar{\mathbf{3}}_F, 1, 0, \lambda]$, $[\bar{\mathbf{3}}_F, 0, 1, \rho]$, $[\bar{\mathbf{3}}_F, 1, 1, \rho]$, and $[\bar{\mathbf{3}}_F, 2, 1, \rho]$. We systematically study their D -wave decays into ground-state bottom baryons with light pseudoscalar mesons and S -wave decays into ground-state bottom baryons with light vector mesons. The relevant sum rule equations are listed in Appendix A, and the obtained results are summarized in Table II.

TABLE II: Decay properties of the P -wave bottom baryons belonging to the $SU(3)$ flavor $\bar{\mathbf{3}}_F$ representation. Radiative decay widths are given in the eighth column for the excited Λ_b^0 and Ξ_b^0 baryons [55].

Multiplets	B (j^P)	Mass (GeV)	Difference (MeV)	Decay channels	$g^{S/D}$ (GeV $^{-2}$)	Decay width (MeV)	Total width (MeV)	Candidate
$[\bar{\mathbf{3}}_F, 1, 0, \lambda]$	$\Lambda_b(\frac{1}{2}^-)$ $5.91^{+0.17}_{-0.13}$		4 ± 2	$\Lambda_b(\frac{1}{2}^-) \rightarrow \Sigma_b\pi \rightarrow \Lambda_b\pi\pi$	$g^S = 17.98^{+10.06}_{-11.67}$	$0 \sim 32000$	$\Gamma_{EM} = 0.45^{+5.90}_{-0.41}$ $\Gamma_{Total} = 0 \sim 32000$	—
				$\Lambda_b(\frac{1}{2}^-) \rightarrow \Sigma_b^*\pi \rightarrow \Lambda_b\pi\pi$	$g^D = 3.28^{+1.75}_{-1.05}$	$0.0^{+2.6}_{-0.0} \times 10^{-5}$		
				$\Lambda_b(\frac{1}{2}^-) \rightarrow \Lambda_b\rho \rightarrow \Lambda_b\pi\pi$	$g^S = 1.83^{+0.74}_{-0.52}$	$0.0^{+7.0}_{-0.0}$		
	$\Lambda_b(\frac{3}{2}^-)$ $5.91^{+0.17}_{-0.13}$		4 ± 2	$\Lambda_b(\frac{3}{2}^-) \rightarrow \Sigma_b^*\pi \rightarrow \Lambda_b\pi\pi$	$g^S = 8.62^{+4.34}_{-4.81}$	$0 \sim 6800$	$\Gamma_{EM} = 0.22^{+3.70}_{-0.22}$ $\Gamma_{Total} = 0 \sim 6800$	—
				$\Lambda_b(\frac{3}{2}^-) \rightarrow \Sigma_b\pi$	$g^D = 2.32^{+1.22}_{-0.73}$	$0 \sim 20$		
				$\Lambda_b(\frac{3}{2}^-) \rightarrow \Lambda_b\rho \rightarrow \Lambda\pi\pi$	$g^S = 2.58^{+1.02}_{-0.71}$	$0.0^{+0.5}_{-0.0}$		
	$\Xi_b(\frac{1}{2}^-)$ $6.10^{+0.20}_{-0.10}$		4 ± 2	$\Xi_b(\frac{1}{2}^-) \rightarrow \Xi'_b\pi$	$g^S = 12.05^{+6.72}_{-5.71}$	2800^{+8200}_{-2800}	$\Gamma_{EM} = 1.8^{+17.0}_{-1.8}$ $\Gamma_{Total} = 2800^{+8200}_{-2800}$	—
				$\Xi_b(\frac{1}{2}^-) \rightarrow \Xi_b^*\pi \rightarrow \Xi_b\pi\pi$	$g^D = 1.98^{+1.12}_{-0.67}$	$0.0^{+2.0}_{-0.0} \times 10^{-6}$		
				$\Xi_b(\frac{1}{2}^-) \rightarrow \Xi_b\rho \rightarrow \Xi_b\pi\pi$	$g^S = 1.32^{+0.47}_{-0.37}$	$0.0^{+1.4}_{-0.0}$		
	$\Xi_b(\frac{3}{2}^-)$ $6.10^{+0.20}_{-0.10}$		4 ± 2	$\Xi_b(\frac{3}{2}^-) \rightarrow \Xi'_b\pi$	$g^D = 2.42^{+1.28}_{-0.83}$	$0.0^{+2.0}_{-0.0}$	$\Gamma_{EM} = 0.70^{+6.20}_{-0.70}$ $\Gamma_{Total} = 330^{+2000}_{-330}$	—
				$\Xi_b(\frac{3}{2}^-) \rightarrow \Xi_b^*\pi$	$g^S = 5.73^{+2.94}_{-3.21}$	330^{+2000}_{-330}		
				$\Xi_b(\frac{3}{2}^-) \rightarrow \Xi_b\rho \rightarrow \Xi_b\pi\pi$	$g^D = 1.40^{+0.73}_{-0.49}$	$0.0^{+2.0}_{-0.0}$		
$[\bar{\mathbf{3}}_F, 0, 1, \rho]$	$\Lambda_b(\frac{1}{2}^-)$ $5.92^{+0.17}_{-0.19}$	—	—	—	—	—	—	—
	$\Xi_b(\frac{1}{2}^-)$ $6.10^{+0.08}_{-0.08}$	—	$\Xi_b(\frac{1}{2}^-) \rightarrow \Xi_b\pi$	$g^S = 13.90^{+5.00}_{-6.57}$	12000^{+10000}_{-9300}	$\Gamma_{EM} = 12.0^{+10.0}_{-9.0}$ $\Gamma_{Total} = 12000^{+10000}_{-9300}$	—	—
$[\bar{\mathbf{3}}_F, 1, 1, \rho]$	$\Lambda_b(\frac{1}{2}^-)$ $5.92^{+0.13}_{-0.10}$		8 ± 3	$\Lambda_b(\frac{1}{2}^-) \rightarrow \Sigma_b\pi \rightarrow \Lambda_b\pi\pi$	$g^S = 0.84^{+1.49}_{-0.51}$	$2.0^{+13.0}_{-2.0} \times 10^{-3}$	$\Gamma_{EM} = 0.03^{+0.08}_{-0.03}$ $\Gamma_{Total} = 0.03^{+0.09}_{-0.03}$	$\Lambda_b(5912)^0$
				$\Lambda_b(\frac{1}{2}^-) \rightarrow \Sigma_b^*\pi \rightarrow \Lambda_b\pi\pi$	$g^D = 0.49^{+1.14}_{-0.24}$	$2.3^{+22.0}_{-2.3} \times 10^{-8}$		
				$\Lambda_b(\frac{1}{2}^-) \rightarrow \Lambda_b\rho \rightarrow \Lambda_b\pi\pi$	$g^S = 1.97^{+3.20}_{-1.02}$	$1.4^{+8.4}_{-1.1} \times 10^{-4}$		
	$\Lambda_b(\frac{3}{2}^-)$ $5.92^{+0.13}_{-0.10}$		8 ± 3	$\Lambda_b(\frac{3}{2}^-) \rightarrow \Sigma_b^*\pi \rightarrow \Lambda_b\pi\pi$	$g^S = 0.80^{+1.54}_{-0.45}$	$4.7^{+29.0}_{-3.8} \times 10^{-3}$	$\Gamma_{EM} = 0.06^{+0.14}_{-0.05}$ $\Gamma_{Total} = 0.06^{+0.17}_{-0.05}$	$\Lambda_b(5920)^0$
				$\Lambda_b(\frac{3}{2}^-) \rightarrow \Sigma_b\pi$	$g^D = 0.79^{+0.33}_{-0.38}$	$9.7^{+26.0}_{-5.9} \times 10^{-6}$		
				$\Lambda_b(\frac{3}{2}^-) \rightarrow \Lambda_b\rho \rightarrow \Lambda\pi\pi$	$g^S = 1.36^{+0.61}_{-0.66}$	$1.7^{+2.1}_{-1.2} \times 10^{-3}$		
	$\Xi_b(\frac{1}{2}^-)$ $6.09^{+0.13}_{-0.12}$		7 ± 3	$\Xi_b(\frac{1}{2}^-) \rightarrow \Xi'_b\pi$	$g^S = 0.52^{+1.02}_{-0.34}$	$4.0^{+26.0}_{-4.0}$	$\Gamma_{EM} = 0.05^{+0.10}_{-0.05}$ $\Gamma_{Total} = 4.0^{+26.0}_{-4.0}$	$\Xi_b(6087)^0$
				$\Xi_b(\frac{1}{2}^-) \rightarrow \Xi_b^*\pi \rightarrow \Xi_b\pi\pi$	$g^D = 0.60^{+0.66}_{-0.31}$	$3.0^{+10.0}_{-2.1} \times 10^{-9}$		
				$\Xi_b(\frac{1}{2}^-) \rightarrow \Xi_b\rho \rightarrow \Xi_b\pi\pi$	$g^S = 1.40^{+2.41}_{-0.66}$	$1.9^{+12.0}_{-1.4} \times 10^{-4}$		
	$\Xi_b(\frac{3}{2}^-)$ $6.09^{+0.12}_{-0.12}$		7 ± 3	$\Xi_b(\frac{3}{2}^-) \rightarrow \Xi'_b\pi$	$g^D = 1.02^{+0.57}_{-0.46}$	$2.4^{+3.4}_{-1.7} \times 10^{-4}$	$\Gamma_{EM} = 0.09^{+0.17}_{-0.09}$ $\Gamma_{Total} = 0.70^{+5.60}_{-0.70}$	$\Xi_b(6095)^0$ $\Xi_b(6100)^-$
				$\Xi_b(\frac{3}{2}^-) \rightarrow \Xi_b^*\pi$	$g^S = 0.50^{+1.03}_{-0.30}$	$0.6^{+5.4}_{-0.6}$		
				$\Xi_b(\frac{3}{2}^-) \rightarrow \Xi_b\rho \rightarrow \Xi_b\pi\pi$	$g^D = 0.59^{+0.32}_{-0.27}$	$6.0^{+8.0}_{-4.2} \times 10^{-5}$		
$[\bar{\mathbf{3}}_F, 2, 1, \rho]$	$\Lambda_b(\frac{3}{2}^-)$ $5.93^{+0.13}_{-0.13}$		17 ± 8	$\Lambda_b(\frac{3}{2}^-) \rightarrow \Sigma_b\pi$	$g^D = 2.75^{+1.31}_{-1.26}$	$3.6^{+4.0}_{-3.6} \times 10^{-3}$	$\Gamma_{EM} = 0.08^{+0.56}_{-0.08}$ $\Gamma_{Total} = 0.0^{+11.0}_{-0.0}$	—
				$\Lambda_b(\frac{3}{2}^-) \rightarrow \Sigma_b^*\pi \rightarrow \Lambda_b\pi\pi$	$g^S = 0.38^{+0.56}_{-0.35}$	$0.0^{+11.0}_{-0.0}$		
	$\Lambda_b(\frac{5}{2}^-)$ $5.94^{+0.13}_{-0.13}$		17 ± 8	$\Lambda_b(\frac{5}{2}^-) \rightarrow \Sigma_b\pi$	$g^D = 2.59^{+1.25}_{-2.59}$	$9.0^{+4.3}_{-9.0} \times 10^{-5}$	$\Gamma_{EM} = 0.01^{+0.08}_{-0.01}$ $\Gamma_{Total} = 0.01^{+0.08}_{-0.01}$	—
				$\Lambda_b(\frac{5}{2}^-) \rightarrow \Sigma_b^*\pi$	$g^D = 0.25^{+0.12}_{-0.25}$	$1.5^{+1.8}_{-1.5} \times 10^{-6}$		
	$\Xi_b(\frac{3}{2}^-)$ $6.10^{+0.15}_{-0.10}$		14 ± 7	$\Xi_b(\frac{3}{2}^-) \rightarrow \Xi_b^*\pi$	$g^S = 0.23^{+0.20}_{-0.20}$	$0.40^{+3.00}_{-0.40}$	$\Gamma_{EM} = 0.15^{+0.55}_{-0.15}$ $\Gamma_{Total} = 1.4^{+12.0}_{-1.4}$	—
				$\Xi_b(\frac{3}{2}^-) \rightarrow \Xi_b\pi$	$g^D = 3.07^{+1.03}_{-0.89}$	$1.0^{+9.0}_{-1.0}$		
				$\Xi_b(\frac{3}{2}^-) \rightarrow \Xi'_b\pi$	$g^D = 1.89^{+0.84}_{-0.77}$	$0.00^{+0.40}_{-0.00}$		
	$\Xi_b(\frac{5}{2}^-)$ $6.11^{+0.15}_{-0.10}$		14 ± 7	$\Xi_b(\frac{5}{2}^-) \rightarrow \Xi_b\pi$	$g^D = 4.35^{+1.44}_{-1.27}$	$1.0^{+7.0}_{-1.0}$	$\Gamma_{EM} = 0.02^{+0.07}_{-0.02}$ $\Gamma_{Total} = 1.0^{+7.0}_{-1.0}$	—
				$\Xi_b(\frac{5}{2}^-) \rightarrow \Xi'_b\pi$	$g^D = 1.79^{+0.77}_{-0.70}$	$0.00^{+0.20}_{-0.00}$		
				$\Xi_b(\frac{5}{2}^-) \rightarrow \Xi_b^*\pi$	$g^D = 0.17^{+0.08}_{-0.08}$	$0.00^{+0.01}_{-0.00}$		

IV. DISCUSSIONS AND SUMMARY

In this paper we have studied strong decay properties of the P -wave bottom baryons belonging to the $SU(3)$ flavor $\bar{\mathbf{3}}_F$ representation through the light-cone sum rule method within the framework of heavy quark effective theory. We have systematically calculated their D -wave decays into ground-state bottom baryons with light pseudoscalar mesons and S -wave decays into ground-state bottom baryons with light vector mesons. The extracted decay widths are presented in Table II for the four $SU(3)$ flavor $\bar{\mathbf{3}}_F$ bottom baryon multiplets, $[\bar{\mathbf{3}}_F, 1, 0, \lambda]$, $[\bar{\mathbf{3}}_F, 0, 1, \rho]$, $[\bar{\mathbf{3}}_F, 1, 1, \rho]$, and $[\bar{\mathbf{3}}_F, 2, 1, \rho]$.

In addition, their mass spectrum has been calculated in Ref. [53], their S -wave decays into ground-state bottom baryons with light pseudoscalar mesons have been calculated in Ref. [54], and their radiative decays have been calculated in Ref. [55]. Altogether, a rather complete investigation has been performed on the P -wave bottom baryons of the $SU(3)$ flavor $\bar{\mathbf{3}}_F$, from which we can understand them as a whole:

- The $[\bar{\mathbf{3}}_F, 1, 0, \lambda]$ doublet contains four excited bottom baryons, $\Lambda_b(1/2^-)$, $\Lambda_b(3/2^-)$, $\Xi_b(1/2^-)$, and $\Xi_b(3/2^-)$. The former two Λ_b baryons can be used to explain the $\Lambda_b(5912)^0$ and $\Lambda_b(5920)^0$, but the latter two Ξ_b baryons are not easy to explain the $\Xi_b(6087)^0$ and $\Xi_b(6095)^0/\Xi_b(6100)^-$. Note that the large uncertainties of our theoretical results are due to the fact that the relevant phase spaces are highly uncertain.
- The $[\bar{\mathbf{3}}_F, 0, 1, \rho]$ singlet contains two excited bottom baryons, $\Lambda_b(1/2^-)$ and $\Xi_b(1/2^-)$. These two

baryons are not easy to be observed in experiments.

- The $[\bar{\mathbf{3}}_F, 1, 1, \rho]$ doublet contains four excited bottom baryons, $\Lambda_b(1/2^-)$, $\Lambda_b(3/2^-)$, $\Xi_b(1/2^-)$, and $\Xi_b(3/2^-)$. We can use this doublet to well explain the four excited bottom baryons $\Lambda_b(5912)^0$, $\Lambda_b(5920)^0$, $\Xi_b(6087)^0$, and $\Xi_b(6095)^0/\Xi_b(6100)^-$ as a whole.
- The $[\bar{\mathbf{3}}_F, 2, 1, \rho]$ doublet contains four excited bottom baryons, $\Lambda_b(3/2^-)$, $\Lambda_b(5/2^-)$, $\Xi_b(3/2^-)$, and $\Xi_b(5/2^-)$. We can use this doublet to further predict two Λ_b baryons and two Ξ_b baryons. Since this is a ρ -mode doublet, its experimental search can verify not only our approach, but also the existence of the ρ -mode.

Summarizing this paper, we systematically investigate the P -wave bottom baryons of the $SU(3)$ flavor $\bar{\mathbf{3}}_F$ to arrive at four Λ_b and four Ξ_b baryons, which have limited widths (less than 100 MeV) and so capable of being observed in experiments. Their masses, mass splittings within the same multiplets, and strong/radiative decay properties are summarized in Table II for future experimental searching.

Acknowledgments

This project is supported by the National Natural Science Foundation of China under Grants No. 12005172, No. 12075019, and No. 12375132, the Jiangsu Provincial Double-Innovation Program under Grant No. JSS-CRC2021488, and the Fundamental Research Funds for the Central Universities.

Appendix A: Sum rule equations

In this appendix we list the sum rule equations derived in the present study. For completeness, the sum rule equations derived in Ref. [54] are also listed here.

The sum rule equation for the $\Lambda_b^+(1/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 1, 0, \lambda]$ is

$$\begin{aligned} G_{\Lambda_b^+[\frac{1}{2}^-] \rightarrow \Sigma_b^+ \pi^-}(\omega, \omega') &= \frac{g_{\Lambda_b^+[\frac{1}{2}^-] \rightarrow \Sigma_b^+ \pi^-} f_{\Lambda_b^0[\frac{1}{2}^-]} f_{\Sigma_b^+}}{(\bar{\Lambda}_{\Lambda_b^0[\frac{1}{2}^-]} - \omega')(\bar{\Lambda}_{\Sigma_b^+} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 8 \times \left(\frac{3f_\pi m_\pi^2}{4\pi^2 t^4 (m_u + m_d)} \phi_{3;\pi}^p(u) - \frac{if_\pi m_\pi^2 v \cdot q}{8\pi^2 t^3 (m_u + m_d)} \phi_{3;\pi}^\sigma(u) + \frac{if_\pi}{16tv \cdot q} \langle \bar{q}q \rangle \psi_{4;\pi}(u) + \frac{if_\pi t}{256v \cdot q} \langle g_s \bar{q}\sigma G q \rangle \psi_{4;\pi}(u) \right). \end{aligned}$$

The sum rule equation for the $\Lambda_b^0(3/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 1, 0, \lambda]$ is

$$\begin{aligned} G_{\Lambda_b^0[\frac{3}{2}^-] \rightarrow \Sigma_b^{*+} \pi^-}(\omega, \omega') &= \frac{g_{\Lambda_b^0[\frac{3}{2}^-] \rightarrow \Sigma_b^{*+} \pi^-} f_{\Lambda_b^0[\frac{3}{2}^-]} f_{\Sigma_b^{*+}}}{(\bar{\Lambda}_{\Lambda_b^0[\frac{3}{2}^-]} - \omega')(\bar{\Lambda}_{\Sigma_b^{*+}} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 8 \times \left(\dots \right) \end{aligned}$$

$$-\frac{f_\pi m_\pi^2}{6\pi^2 t^4(m_u + m_d)} \phi_{3;\pi}^p(u) + \frac{if_\pi m_\pi^2 v \cdot q}{36\pi^2 t^3(m_u + m_d)} \phi_{3;\pi}^\sigma(u) - \frac{if_\pi}{72tv \cdot q} \langle \bar{q}q \rangle \psi_{4;\pi}(u) - \frac{if_\pi t}{1152v \cdot q} \langle g_s \bar{s}\sigma G s \rangle \psi_{4;\pi}(u).$$

The sum rule equations for the $\Xi_b^- (1/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 1, 0, \lambda]$ are

$$\begin{aligned} & G_{\Xi_b^- [\frac{1}{2}^-] \rightarrow \Xi_b^{*0} \pi^-}(\omega, \omega') = \frac{g_{\Xi_b^- [\frac{1}{2}^-] \rightarrow \Xi_b^{*0} \pi^-} f_{\Xi_b^- [\frac{1}{2}^-]} f_{\Xi_b^{*0}}}{(\Lambda_{\Xi_b^- [\frac{1}{2}^-]} - \omega')(\Lambda_{\Xi_b^{*0}} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 4 \times \left(\begin{array}{l} \frac{3f_\pi m_\pi^2}{4\pi^2 t^4(m_u + m_d)} \phi_{3;\pi}^p(u) - \frac{if_\pi m_\pi^2 v \cdot q}{8\pi^2 t^3(m_u + m_d)} \phi_{3;\pi}^\sigma(u) + \frac{if_\pi}{16tv \cdot q} \langle \bar{s}s \rangle \psi_{4;\pi}(u) + \frac{if_\pi t}{256v \cdot q} \langle g_s \bar{s}\sigma G s \rangle \psi_{4;\pi}(u) \\ + \frac{3if_\pi}{16\pi^2 t^3 v \cdot q} m_s \psi_{4;\pi}(u) + \frac{f_\pi m_\pi^2}{32(m_u + m_d)} m_s \langle \bar{s}s \rangle \phi_{3;\pi}^p(u) - \frac{if_\pi m_\pi^2 tv \cdot q}{192(m_u + m_d)} m_s \langle \bar{s}s \rangle \phi_{3;\pi}^\sigma(u) \end{array} \right), \\ & G_{\Xi_b^- [\frac{1}{2}^-] \rightarrow \Xi_b^0 \rho^-}(\omega, \omega') = \frac{g_{\Xi_b^- [\frac{1}{2}^-] \rightarrow \Xi_b^0 \rho^-} f_{\Xi_b^- [\frac{1}{2}^-]} f_{\Xi_b^0}}{(\Lambda_{\Xi_b^- [\frac{1}{2}^-]} - \omega')(\Lambda_{\Xi_b^0} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 4 \times \left(\begin{array}{l} \frac{if_\rho^\parallel m_\rho}{4\pi^2 t^4} \phi_{2;\rho}^\parallel(u) - \frac{if_\rho^\parallel m_\rho^3}{8\pi^2 t^4(v \cdot q)^2} \phi_{2;\rho}^\parallel(u) + \frac{if_\rho^\parallel m_\rho^3}{4\pi^2 t^4(v \cdot q)^2} \phi_{3;\rho}^\perp(u) - \frac{if_\rho^\parallel m_\rho^3}{8\pi^2 t^4(v \cdot q)^2} \psi_{4;\rho}^\parallel(u) + \frac{if_\rho^\parallel m_\rho^3}{64\pi^2 t^2} \phi_{4;\rho}^\parallel(u) \\ - \frac{if_\rho^\perp m_\rho^2}{48} \langle \bar{s}s \rangle \psi_{3;\rho}^\parallel(u) - \frac{if_\rho^\perp m_\rho^2 t^2}{768} \langle g_s \bar{s}\sigma G s \rangle \psi_{3;\rho}^\parallel(u) - \frac{if_\rho^\perp m_\rho^2}{16\pi^2 t^2} m_s \psi_{3;\rho}^\parallel(u) + \frac{if_\rho^\parallel m_\rho}{96} m_s \langle \bar{s}s \rangle \phi_{2;\rho}^\parallel(u) \\ - \frac{if_\rho^\parallel m_\rho^3}{192(v \cdot q)^2} m_s \langle \bar{s}s \rangle \phi_{2;\rho}^\parallel(u) + \frac{if_\rho^\parallel m_\rho^3}{96(v \cdot q)^2} m_s \langle \bar{s}s \rangle \phi_{3;\rho}^\perp(u) + \frac{if_\rho^\parallel m_\rho^3 t^2}{1536} m_s \langle \bar{s}s \rangle \phi_{4;\rho}^\parallel(u) - \frac{if_\rho^\parallel m_\rho^3}{192(v \cdot q)^2} m_s \langle \bar{s}s \rangle \psi_{4;\rho}^\parallel(u) \end{array} \right) \\ &+ \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega' t(\alpha_2 + u\alpha_3)} e^{iu\omega t(1 - \alpha_2 - u\alpha_3)} \times 4 \times \left(\begin{array}{l} - \frac{if_\rho^\parallel m_\rho^3}{8\pi^2 t^2} \Phi_{4;\rho}^\parallel(\underline{\alpha}) - \frac{if_\rho^\parallel m_\rho^3}{16\pi^2 t^2} \Psi_{4;\rho}^\parallel(\underline{\alpha}) - \frac{if_\rho^\parallel m_\rho^3}{8\pi^2 t^2} \tilde{\Phi}_{4;\rho}^\parallel(\underline{\alpha}) - \frac{if_\rho^\parallel m_\rho^3}{16\pi^2 t^2} \tilde{\Psi}_{4;\rho}^\parallel(\underline{\alpha}) - \frac{if_\rho^\parallel m_\rho^3 u}{4\pi^2 t^2} \Phi_{4;\rho}^\parallel(\underline{\alpha}) - \frac{if_\rho^\parallel m_\rho^3 u}{8\pi^2 t^2} \Psi_{4;\rho}^\parallel(\underline{\alpha}) \end{array} \right). \end{aligned}$$

The sum rule equations for the $\Xi_b^- (3/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 1, 0, \lambda]$ are

$$\begin{aligned} & G_{\Xi_b^- [\frac{3}{2}^-] \rightarrow \Xi_b^{*0} \pi^-}(\omega, \omega') = \frac{g_{\Xi_b^- [\frac{3}{2}^-] \rightarrow \Xi_b^{*0} \pi^-} f_{\Xi_b^- [\frac{3}{2}^-]} f_{\Xi_b^{*0}}}{(\Lambda_{\Xi_b^- [\frac{3}{2}^-]} - \omega')(\Lambda_{\Xi_b^{*0}} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 4 \times \left(\begin{array}{l} - \frac{f_\pi m_\pi^2}{6\pi^2 t^4(m_u + m_d)} \phi_{3;\pi}^p(u) + \frac{if_\pi m_\pi^2 v \cdot q}{36\pi^2 t^3(m_u + m_d)} \phi_{3;\pi}^\sigma(u) - \frac{if_\pi}{72tv \cdot q} \langle \bar{s}s \rangle \psi_{4;\pi}(u) - \frac{if_\pi t}{1152v \cdot q} \langle g_s \bar{s}\sigma G s \rangle \psi_{4;\pi}(u) \\ - \frac{if_\pi}{24\pi^2 t^3 v \cdot q} m_s \psi_{4;\pi}(u) - \frac{f_\pi m_\pi^2}{144(m_u + m_d)} m_s \langle \bar{s}s \rangle \phi_{3;\pi}^p(u) + \frac{if_\pi m_\pi^2 tv \cdot q}{864(m_u + m_d)} m_s \langle \bar{s}s \rangle \phi_{3;\pi}^\sigma(u) \end{array} \right), \\ & G_{\Xi_b^- [\frac{3}{2}^-] \rightarrow \Xi_b^0 \rho^-}(\omega, \omega') = \frac{g_{\Xi_b^- [\frac{3}{2}^-] \rightarrow \Xi_b^0 \rho^-} f_{\Xi_b^- [\frac{3}{2}^-]} f_{\Xi_b^0}}{(\Lambda_{\Xi_b^- [\frac{3}{2}^-]} - \omega')(\Lambda_{\Xi_b^0} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 4 \times \left(\begin{array}{l} \frac{if_\rho^\parallel m_\rho}{6\pi^2 t^4} \phi_{2;\rho}^\parallel(u) - \frac{if_\rho^\parallel m_\rho^3}{12\pi^2 t^4(v \cdot q)^2} \phi_{2;\rho}^\parallel(u) + \frac{if_\rho^\parallel m_\rho^3}{6\pi^2 t^4(v \cdot q)^2} \phi_{3;\rho}^\perp(u) - \frac{if_\rho^\parallel m_\rho^3}{12\pi^2 t^4(v \cdot q)^2} \psi_{4;\rho}^\parallel(u) + \frac{if_\rho^\parallel m_\rho^3}{96\pi^2 t^2} \phi_{4;\rho}^\parallel(u) \\ - \frac{if_\rho^\perp m_\rho^2}{72} \langle \bar{s}s \rangle \psi_{3;\rho}^\parallel(u) - \frac{if_\rho^\perp m_\rho^2 t^2}{1152} \langle g_s \bar{s}\sigma G s \rangle \psi_{3;\rho}^\parallel(u) - \frac{if_\rho^\perp m_\rho^2}{24\pi^2 t^2} m_s \psi_{3;\rho}^\parallel(u) + \frac{if_\rho^\parallel m_\rho}{144} m_s \langle \bar{s}s \rangle \phi_{2;\rho}^\parallel(u) \\ - \frac{if_\rho^\parallel m_\rho^3}{288(v \cdot q)^2} m_s \langle \bar{s}s \rangle \phi_{2;\rho}^\parallel(u) + \frac{if_\rho^\parallel m_\rho^3}{144(v \cdot q)^2} m_s \langle \bar{s}s \rangle \phi_{3;\rho}^\perp(u) + \frac{if_\rho^\parallel m_\rho^3 t^2}{2304} m_s \langle \bar{s}s \rangle \phi_{4;\rho}^\parallel(u) - \frac{if_\rho^\parallel m_\rho^3}{288(v \cdot q)^2} m_s \langle \bar{s}s \rangle \psi_{4;\rho}^\parallel(u) \end{array} \right) \end{aligned}$$

$$+ \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega't(\alpha_2+u\alpha_3)} e^{i\omega t(1-\alpha_2-u\alpha_3)} \times 4 \times \left(-\frac{if_\rho^\parallel m_\rho^3}{12\pi^2 t^2} \Phi_{4;\rho}^\parallel(\underline{\alpha}) - \frac{if_\rho^\parallel m_\rho^3}{24\pi^2 t^2} \Psi_{4;\rho}^\parallel(\underline{\alpha}) - \frac{if_\rho^\parallel m_\rho^3}{12\pi^2 t^2} \tilde{\Phi}_{4;\rho}^\parallel(\underline{\alpha}) - \frac{if_\rho^\parallel m_\rho^3}{24\pi^2 t^2} \tilde{\Psi}_{4;\rho}^\parallel(\underline{\alpha}) - \frac{if_\rho^\parallel m_\rho^3 u}{6\pi^2 t^2} \Phi_{4;\rho}^\parallel(\underline{\alpha}) - \frac{if_\rho^\parallel m_\rho^3 u}{12\pi^2 t^2} \Psi_{4;\rho}^\parallel(\underline{\alpha}) \right).$$

The sum rule equation for the $\Xi_b^- (1/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 0, 1, \rho]$ is

$$\begin{aligned} G_{\Xi_b^- [\frac{1}{2}^-] \rightarrow \Xi_b^0 \pi^-}(\omega, \omega') &= \frac{g_{\Xi_b^- [\frac{1}{2}^-] \rightarrow \Xi_b^0 \pi^-} f_{\Xi_b^- [\frac{1}{2}^-]} f_{\Xi_b^0}}{(\bar{\Lambda}_{\Xi_b^- [\frac{1}{2}^-]} - \omega')(\bar{\Lambda}_{\Xi_b^0} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 4 \times \left(-\frac{3f_\pi m_\pi^2}{4\pi^2 t^4 (m_u + m_d)} \phi_{3;\pi}^p(u) + \frac{if_\pi m_\pi^2 v \cdot q}{8\pi^2 t^3 (m_u + m_d)} \phi_{3;\pi}^\sigma(u) - \frac{if_\pi t}{16tv \cdot q} \langle \bar{s}s \rangle \psi_{4;\pi}(u) - \frac{if_\pi t}{256v \cdot q} \langle g_s \bar{s}\sigma G s \rangle \psi_{4;\pi}(u) \right. \\ &\quad \left. - \frac{3if_\pi}{16\pi^2 t^3 v \cdot q} m_s \psi_{4;\pi}(u) - \frac{f_\pi m_\pi^2}{32(m_u + m_d)} m_s \langle \bar{s}s \rangle \phi_{3;\pi}^p(u) + \frac{if_\pi m_\pi^2 tv \cdot q}{192(m_u + m_d)} m_s \langle \bar{s}s \rangle \phi_{3;\pi}^\sigma(u) \right). \end{aligned}$$

The sum rule equation for the $\Lambda_b^0 (1/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 1, 1, \rho]$ is

$$\begin{aligned} G_{\Lambda_b^0 [\frac{1}{2}^-] \rightarrow \Sigma_b^+ \pi^-}(\omega, \omega') &= \frac{g_{\Lambda_b^0 [\frac{1}{2}^-] \rightarrow \Sigma_b^+ \pi^-} f_{\Lambda_b^0 [\frac{1}{2}^-]} f_{\Sigma_b^+}}{(\bar{\Lambda}_{\Lambda_b^0 [\frac{1}{2}^-]} - \omega')(\bar{\Lambda}_{\Sigma_b^+} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 8 \times \left(\frac{3if_\pi v \cdot q}{2\pi^2 t^4} \phi_{2;\pi}(u) + \frac{3if_\pi v \cdot q}{32\pi^2 t^2} \phi_{4;\pi}(u) + \frac{3if_\pi}{2\pi^2 t^4 v \cdot q} \psi_{4;\pi}(u) + \frac{if_\pi m_\pi^2 v \cdot q}{24(m_u + m_d)} \langle \bar{q}q \rangle \phi_{3;\pi}^\sigma(u) \right. \\ &\quad \left. + \frac{if_\pi m_\pi^2 t^2 v \cdot q}{384(m_u + m_d)} \langle g_s \bar{q}\sigma G q \rangle \phi_{3;\pi}^\sigma(u) \right) \\ &+ \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega' t(\alpha_2+u\alpha_3)} e^{i\omega t(1-\alpha_2-u\alpha_3)} \times 8 \times \left(\frac{3if_\pi v \cdot q}{8\pi^2 t^2} \Phi_{4;\pi}(\underline{\alpha}) - \frac{if_\pi v \cdot q}{4\pi^2 t^2} \Psi_{4;\pi}(\underline{\alpha}) + \frac{if_\pi v \cdot q}{8\pi^2 t^2} \tilde{\Phi}_{4;\pi}(\underline{\alpha}) + \frac{if_\pi v \cdot q}{4\pi^2 t^2} \tilde{\Psi}_{4;\pi}(\underline{\alpha}) + \frac{if_\pi uv \cdot q}{4\pi^2 t^2} \Phi_{4;\pi}(\underline{\alpha}) + \frac{if_\pi uv \cdot q}{2\pi^2 t^2} \Psi_{4;\pi}(\underline{\alpha}) \right). \end{aligned}$$

One of the sum rule equations for the $\Lambda_b^0 (3/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 1, 1, \rho]$ has been given in Eq. (53), and the other is

$$\begin{aligned} G_{\Lambda_b^0 [\frac{3}{2}^-] \rightarrow \Sigma_b^{*+} \pi^-}(\omega, \omega') &= \frac{g_{\Lambda_b^0 [\frac{3}{2}^-] \rightarrow \Sigma_b^{*+} \pi^-} f_{\Lambda_b^0 [\frac{3}{2}^-]} f_{\Sigma_b^{*+}}}{(\bar{\Lambda}_{\Lambda_b^0 [\frac{3}{2}^-]} - \omega')(\bar{\Lambda}_{\Sigma_b^{*+}} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 8 \times \left(-\frac{f_\pi v \cdot q}{3\pi^2 t^4} \phi_{2;\pi}(u) - \frac{f_\pi v \cdot q}{48\pi^2 t^2} \phi_{4;\pi}(u) - \frac{f_\pi}{3\pi^2 t^4 v \cdot q} \psi_{4;\pi}(u) - \frac{f_\pi m_\pi^2 v \cdot q}{108(m_u + m_d)} \langle \bar{q}q \rangle \phi_{3;\pi}^\sigma(u) \right. \\ &\quad \left. - \frac{f_\pi m_\pi^2 t^2 v \cdot q}{1728(m_u + m_d)} \langle g_s \bar{q}\sigma G q \rangle \phi_{3;\pi}^\sigma(u) \right) \\ &+ \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega' t(\alpha_2+u\alpha_3)} e^{i\omega t(1-\alpha_2-u\alpha_3)} \times 8 \times \left(-\frac{f_\pi v \cdot q}{24\pi^2 t^2} \Phi_{4;\pi}(\underline{\alpha}) + \frac{5f_\pi v \cdot q}{72\pi^2 t^2} \Psi_{4;\pi}(\underline{\alpha}) - \frac{f_\pi v \cdot q}{72\pi^2 t^2} \tilde{\Phi}_{4;\pi}(\underline{\alpha}) - \frac{5f_\pi v \cdot q}{72\pi^2 t^2} \tilde{\Psi}_{4;\pi}(\underline{\alpha}) \right. \\ &\quad \left. - \frac{f_\pi uv \cdot q}{36\pi^2 t^2} \Phi_{4;\pi}(\underline{\alpha}) - \frac{7f_\pi uv \cdot q}{72\pi^2 t^2} \Psi_{4;\pi}(\underline{\alpha}) + \frac{f_\pi uv \cdot q}{12\pi^2 t^2} \tilde{\Phi}_{4;\pi}(\underline{\alpha}) + \frac{f_\pi uv \cdot q}{24\pi^2 t^2} \tilde{\Psi}_{4;\pi}(\underline{\alpha}) \right). \end{aligned}$$

The sum rule equation for the $\Xi_b^- (1/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 1, 1, \rho]$ is

$$\begin{aligned}
G_{\Xi_b^- [\frac{1}{2}^-] \rightarrow \Xi_b^0 \rho^-}(\omega, \omega') &= \frac{g_{\Xi_b^- [\frac{1}{2}^-] \rightarrow \Xi_b^0 \rho^-} f_{\Xi_b^- [\frac{1}{2}^-]} f_{\Xi_b^0}}{(\bar{\Lambda}_{\Xi_b^- [\frac{1}{2}^-]} - \omega')(\bar{\Lambda}_{\Xi_b^0} - \omega)} \\
&= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 4 \times \left(\right. \\
&\quad - \frac{f_\rho^\perp v \cdot q}{2\pi^2 t^4} \phi_{2;\rho}^\perp(u) + \frac{f_\rho^\perp m_\rho^2}{2\pi^2 t^4 v \cdot q} \phi_{2;\rho}^\perp(u) - \frac{f_\rho^\perp m_\rho^2}{2\pi^2 t^4 v \cdot q} \psi_{4;\rho}^\perp(u) - \frac{f_\rho^\perp m_\rho^2 v \cdot q}{32\pi^2 t^2} \phi_{4;\rho}^\perp(u) + \frac{f_\rho^\parallel m_\rho v \cdot q}{48} \langle \bar{s}s \rangle \psi_{3;\rho}^\perp(u) \\
&\quad + \frac{f_\rho^\parallel m_\rho t^2 v \cdot q}{768} \langle g_s \bar{s}\sigma G s \rangle \psi_{3;\rho}^\perp(u) + \frac{f_\rho^\parallel m_\rho v \cdot q}{16\pi^2 t^2} m_s \psi_{3;\rho}^\perp(u) - \frac{f_\rho^\perp v \cdot q}{48} m_s \langle \bar{s}s \rangle \phi_{2;\rho}^\perp(u) + \frac{f_\rho^\perp m_\rho^2}{48 v \cdot q} m_s \langle \bar{s}s \rangle \phi_{2;\rho}^\perp(u) \\
&\quad \left. - \frac{f_\rho^\perp m_\rho^2}{48 v \cdot q} m_s \langle \bar{s}s \rangle \psi_{4;\rho}^\perp(u) - \frac{f_\rho^\perp m_\rho^2 t^2 v \cdot q}{768} m_s \langle \bar{s}s \rangle \phi_{4;\rho}^\perp(u) \right) \\
&+ \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega' t(\alpha_2 + u\alpha_3)} e^{i\omega t(1-\alpha_2-u\alpha_3)} \times 4 \times \left(\right. \\
&\quad - \frac{f_\rho^\perp m_\rho^2 v \cdot q}{8\pi^2 t^2} \Psi_{4;\rho}^\perp(\underline{\alpha}) + \frac{f_\rho^\perp m_\rho^2 v \cdot q}{8\pi^2 t^2} \tilde{\Psi}_{4;\rho}^\perp(\underline{\alpha}) - \frac{f_\rho^\perp m_\rho^2 uv \cdot q}{8\pi^2 t^2} \Phi_{4;\rho}^{\perp 1}(\underline{\alpha}) + \frac{f_\rho^\perp m_\rho^2 uv \cdot q}{8\pi^2 t^2} \Phi_{4;\rho}^{\perp 2}(\underline{\alpha}) - \frac{f_\rho^\perp m_\rho^2 uv \cdot q}{4\pi^2 t^2} \tilde{\Psi}_{4;\rho}^\perp(\underline{\alpha}) \left. \right).
\end{aligned}$$

The sum rule equations for the $\Xi_b^- (3/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 1, 1, \rho]$ are

$$\begin{aligned}
G_{\Xi_b^- [\frac{3}{2}^-] \rightarrow \Xi_b^{*0} \pi^-}(\omega, \omega') &= \frac{g_{\Xi_b^- [\frac{3}{2}^-] \rightarrow \Xi_b^{*0} \pi^-} f_{\Xi_b^- [\frac{3}{2}^-]} f_{\Xi_b^{*0}}}{(\bar{\Lambda}_{\Xi_b^- [\frac{3}{2}^-]} - \omega')(\bar{\Lambda}_{\Xi_b^{*0}} - \omega)} \\
&= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 4 \times \left(\right. \\
&\quad - \frac{f_\pi v \cdot q}{3\pi^2 t^4} \phi_{2;\pi}(u) - \frac{f_\pi v \cdot q}{48\pi^2 t^2} \phi_{4;\pi}(u) - \frac{f_\pi}{3\pi^2 t^4 v \cdot q} \psi_{4;\pi}(u) - \frac{f_\pi m_\pi^2 v \cdot q}{108(m_u + m_d)} \langle \bar{s}s \rangle \phi_{3;\pi}^\sigma(u) \\
&\quad - \frac{f_\pi m_\pi^2 t^2 v \cdot q}{1728(m_u + m_d)} \langle g_s \bar{s}\sigma G s \rangle \phi_{3;\pi}^\sigma(u) - \frac{f_\pi m_\pi^2 v \cdot q}{36\pi^2 t^2(m_u + m_d)} m_s \phi_{3;\pi}^\sigma(u) - \frac{f_\pi v \cdot q}{72} m_s \langle \bar{s}s \rangle \phi_{2;\pi}(u) \\
&\quad \left. - \frac{f_\pi t^2 v \cdot q}{1152} m_s \langle \bar{s}s \rangle \phi_{4;\pi}(u) - \frac{f_\pi}{72 v \cdot q} m_s \langle \bar{s}s \rangle \psi_{4;\pi}(u) \right) \\
&+ \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega' t(\alpha_2 + u\alpha_3)} e^{i\omega t(1-\alpha_2-u\alpha_3)} \times 4 \times \left(\right. \\
&\quad - \frac{f_\pi v \cdot q}{24\pi^2 t^2} \Phi_{4;\pi}(\underline{\alpha}) + \frac{5f_\pi v \cdot q}{72\pi^2 t^2} \Psi_{4;\pi}(\underline{\alpha}) - \frac{f_\pi v \cdot q}{72\pi^2 t^2} \tilde{\Phi}_{4;\pi}(\underline{\alpha}) - \frac{5f_\pi v \cdot q}{72\pi^2 t^2} \tilde{\Psi}_{4;\pi}(\underline{\alpha}) \\
&\quad \left. - \frac{f_\pi uv \cdot q}{36\pi^2 t^2} \Phi_{4;\pi}(\underline{\alpha}) - \frac{7f_\pi uv \cdot q}{72\pi^2 t^2} \Psi_{4;\pi}(\underline{\alpha}) + \frac{f_\pi uv \cdot q}{12\pi^2 t^2} \tilde{\Phi}_{4;\pi}(\underline{\alpha}) + \frac{f_\pi uv \cdot q}{24\pi^2 t^2} \tilde{\Psi}_{4;\pi}(\underline{\alpha}) \right), \\
G_{\Xi_b^- [\frac{3}{2}^-] \rightarrow \Xi_b^0 \rho^-}(\omega, \omega') &= \frac{g_{\Xi_b^- [\frac{3}{2}^-] \rightarrow \Xi_b^0 \rho^-} f_{\Xi_b^- [\frac{3}{2}^-]} f_{\Xi_b^0}}{(\bar{\Lambda}_{\Xi_b^- [\frac{3}{2}^-]} - \omega')(\bar{\Lambda}_{\Xi_b^0} - \omega)} \\
&= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 4 \times \left(\right. \\
&\quad - \frac{f_\rho^\perp v \cdot q}{3\pi^2 t^4} \phi_{2;\rho}^\perp(u) - \frac{f_\rho^\perp m_\rho^2}{3\pi^2 t^4 v \cdot q} \phi_{2;\rho}^\perp(u) + \frac{f_\rho^\perp m_\rho^2}{3\pi^2 t^4 v \cdot q} \psi_{4;\rho}^\perp(u) + \frac{f_\rho^\perp m_\rho^2 v \cdot q}{48\pi^2 t^2} \phi_{4;\rho}^\perp(u) - \frac{f_\rho^\parallel m_\rho v \cdot q}{72} \langle \bar{s}s \rangle \psi_{3;\rho}^\perp(u) \\
&\quad - \frac{f_\rho^\parallel m_\rho t^2 v \cdot q}{1152} \langle g_s \bar{s}\sigma G s \rangle \psi_{3;\rho}^\perp(u) - \frac{f_\rho^\parallel m_\rho v \cdot q}{24\pi^2 t^2} m_s \psi_{3;\rho}^\perp(u) + \frac{f_\rho^\perp v \cdot q}{72} m_s \langle \bar{s}s \rangle \phi_{2;\rho}^\perp(u) - \frac{f_\rho^\perp m_\rho^2}{72 v \cdot q} m_s \langle \bar{s}s \rangle \phi_{2;\rho}^\perp(u) \\
&\quad \left. + \frac{f_\rho^\perp m_\rho^2}{72 v \cdot q} m_s \langle \bar{s}s \rangle \psi_{4;\rho}^\perp(u) + \frac{f_\rho^\perp m_\rho^2 t^2 v \cdot q}{1152} m_s \langle \bar{s}s \rangle \phi_{4;\rho}^\perp(u) \right) \\
&+ \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega' t(\alpha_2 + u\alpha_3)} e^{i\omega t(1-\alpha_2-u\alpha_3)} \times 4 \times \left(\right. \\
&\quad - \frac{f_\rho^\perp m_\rho^2 v \cdot q}{12\pi^2 t^2} \Psi_{4;\rho}^\perp(\underline{\alpha}) - \frac{f_\rho^\perp m_\rho^2 v \cdot q}{12\pi^2 t^2} \tilde{\Psi}_{4;\rho}^\perp(\underline{\alpha}) + \frac{f_\rho^\perp m_\rho^2 uv \cdot q}{12\pi^2 t^2} \Phi_{4;\rho}^{\perp 1}(\underline{\alpha}) - \frac{f_\rho^\perp m_\rho^2 uv \cdot q}{12\pi^2 t^2} \Phi_{4;\rho}^{\perp 2}(\underline{\alpha}) + \frac{f_\rho^\perp m_\rho^2 uv \cdot q}{6\pi^2 t^2} \tilde{\Psi}_{4;\rho}^\perp(\underline{\alpha}) \left. \right).
\end{aligned}$$

The sum rule equation for the $\Lambda_b^0(3/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 2, 1, \rho]$ is

$$\begin{aligned} G_{\Lambda_b^0[\frac{3}{2}^-] \rightarrow \Sigma_b^{*+} \pi^-}(\omega, \omega') &= \frac{g_{\Lambda_b^0[\frac{3}{2}^-] \rightarrow \Sigma_b^{*+} \pi^-} f_{\Lambda_b^0[\frac{3}{2}^-]} f_{\Sigma_b^{*+}}}{(\bar{\Lambda}_{\Lambda_b^0[\frac{3}{2}^-]} - \omega')(\bar{\Lambda}_{\Sigma_b^{*+}} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega' t(\alpha_2 + u\alpha_3)} e^{i\omega t(1 - \alpha_2 - u\alpha_3)} \times 8 \times \left(\right. \\ &\quad \frac{f_\pi v \cdot q}{24\pi^2 t^2} \Phi_{4;\pi}(\underline{\alpha}) + \frac{f_\pi v \cdot q}{24\pi^2 t^2} \Psi_{4;\pi}(\underline{\alpha}) - \frac{f_\pi v \cdot q}{24\pi^2 t^2} \tilde{\Phi}_{4;\pi}(\underline{\alpha}) - \frac{f_\pi v \cdot q}{24\pi^2 t^2} \tilde{\Psi}_{4;\pi}(\underline{\alpha}) \\ &\quad \left. - \frac{f_\pi uv \cdot q}{12\pi^2 t^2} \Phi_{4;\pi}(\underline{\alpha}) - \frac{f_\pi uv \cdot q}{24\pi^2 t^2} \Psi_{4;\pi}(\underline{\alpha}) + \frac{f_\pi uv \cdot q}{12\pi^2 t^2} \tilde{\Phi}_{4;\pi}(\underline{\alpha}) + \frac{f_\pi uv \cdot q}{24\pi^2 t^2} \tilde{\Psi}_{4;\pi}(\underline{\alpha}) \right). \end{aligned}$$

The sum rule equation for the $\Lambda_b^-(5/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 2, 1, \rho]$ is

$$\begin{aligned} G_{\Lambda_b^0[\frac{5}{2}^-] \rightarrow \Sigma_b^{*+} \rho^-}(\omega, \omega') &= \frac{g_{\Lambda_b^0[\frac{5}{2}^-] \rightarrow \Sigma_b^{*+} \rho^-} f_{\Lambda_b^+[\frac{5}{2}^-]} f_{\Sigma_b^{*+}}}{(\bar{\Lambda}_{\Lambda_b^0[\frac{5}{2}^-]} - \omega')(\bar{\Lambda}_{\Sigma_b^{*+}} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 8 \times \left(\right. \\ &\quad - \frac{3if_\rho^\parallel m_\rho^\parallel}{10\pi^2 t^4} \phi_{2;\rho}^\parallel(u) + \frac{3if_\rho^\parallel m_\rho^3}{20\pi^2 t^4 (v \cdot q)^2} \phi_{2;\rho}^\parallel(u) + \frac{3if_\rho^\parallel m_\rho}{10\pi^2 t^4} \phi_{3;\rho}^\perp(u) - \frac{3if_\rho^\parallel m_\rho^3}{10\pi^2 t^4 (v \cdot q)^2} \phi_{3;\rho}^\perp(u) - \frac{3f_\rho^\parallel m_\rho v \cdot q}{40\pi^2 t^3} \psi_{3;\rho}^\perp(u) \\ &\quad - \frac{3if_\rho^\parallel m_\rho^3}{160\pi^2 t^2} \phi_{4;\rho}^\parallel(u) + \frac{3if_\rho^\parallel m_\rho^3}{20\pi^2 t^4 (v \cdot q)^2} \psi_{4;\rho}^\parallel(u) + \frac{f_\rho^\perp m_\rho^2}{40tv \cdot q} \langle \bar{q}q \rangle \phi_{2;\rho}^\perp(u) + \frac{if_\rho^\perp m_\rho^2}{40} \langle \bar{q}q \rangle \psi_{3;\rho}^\parallel(u) - \frac{f_\rho^\perp m_\rho^2}{40tv \cdot q} \langle \bar{q}q \rangle \psi_{4;\rho}^\perp(u) \\ &\quad \left. + \frac{f_\rho^\perp m_\rho^2 t}{640v \cdot q} \langle g_s \bar{q} \sigma G q \rangle \phi_{2;\rho}^\perp(u) + \frac{if_\rho^\perp m_\rho^2 t^2}{640} \langle g_s \bar{q} \sigma G q \rangle \psi_{3;\rho}^\parallel(u) - \frac{f_\rho^\perp m_\rho^2 t}{640v \cdot q} \langle g_s \bar{q} \sigma G q \rangle \psi_{4;\rho}^\perp(u) \right) \\ &+ \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega' t(\alpha_2 + u\alpha_3)} e^{i\omega t(1 - \alpha_2 - u\alpha_3)} \times 8 \times \left(\right. \\ &\quad \left. \frac{3if_\rho^\parallel m_\rho^3}{40\pi^2 t^2} \Psi_{4;\rho}^\parallel(\underline{\alpha}) - \frac{3if_\rho^\parallel m_\rho^3}{40\pi^2 t^2} \tilde{\Psi}_{4;\rho}^\parallel(\underline{\alpha}) + \frac{3if_\rho^\parallel m_\rho^3 u}{20\pi^2 t^2} \Psi_{4;\rho}^\parallel(\underline{\alpha}) \right). \end{aligned}$$

The sum rule equation for the $\Xi_b^-(3/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 2, 1, \rho]$ is

$$\begin{aligned} G_{\Xi_b^-[\frac{3}{2}^-] \rightarrow \Xi_b^{*0} \pi^-}(\omega, \omega') &= \frac{g_{\Xi_b^-[\frac{3}{2}^-] \rightarrow \Xi_b^{*0} \pi^-} f_{\Xi_b^-[\frac{3}{2}^-]} f_{\Xi_b^{*0}}}{(\bar{\Lambda}_{\Xi_b^-[\frac{3}{2}^-]} - \omega')(\bar{\Lambda}_{\Xi_b^{*0}} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega' t(\alpha_2 + u\alpha_3)} e^{i\omega t(1 - \alpha_2 - u\alpha_3)} \times 4 \times \left(\right. \\ &\quad \frac{f_\pi v \cdot q}{24\pi^2 t^2} \Phi_{4;\pi}(\underline{\alpha}) + \frac{f_\pi v \cdot q}{24\pi^2 t^2} \Psi_{4;\pi}(\underline{\alpha}) - \frac{f_\pi v \cdot q}{24\pi^2 t^2} \tilde{\Phi}_{4;\pi}(\underline{\alpha}) - \frac{f_\pi v \cdot q}{24\pi^2 t^2} \tilde{\Psi}_{4;\pi}(\underline{\alpha}) \\ &\quad \left. - \frac{f_\pi uv \cdot q}{12\pi^2 t^2} \Phi_{4;\pi}(\underline{\alpha}) - \frac{f_\pi uv \cdot q}{24\pi^2 t^2} \Psi_{4;\pi}(\underline{\alpha}) + \frac{f_\pi uv \cdot q}{12\pi^2 t^2} \tilde{\Phi}_{4;\pi}(\underline{\alpha}) + \frac{f_\pi uv \cdot q}{24\pi^2 t^2} \tilde{\Psi}_{4;\pi}(\underline{\alpha}) \right). \end{aligned}$$

The sum rule equation for the $\Xi_b^-(5/2^-)$ belonging to $[\bar{\mathbf{3}}_F, 2, 1, \rho]$ is

$$\begin{aligned} G_{\Xi_b^-[\frac{5}{2}^-] \rightarrow \Xi_c^{*+} \rho^-}(\omega, \omega') &= \frac{g_{\Xi_b^-[\frac{5}{2}^-] \rightarrow \Xi_c^{*+} \rho^-} f_{\Xi_c^0[\frac{5}{2}^-]} f_{\Xi_b^{*0}}}{(\bar{\Lambda}_{\Xi_b^-[\frac{5}{2}^-]} - \omega')(\bar{\Lambda}_{\Xi_c^{*0}} - \omega)} \\ &= \int_0^\infty dt \int_0^1 du e^{i(1-u)\omega' t} e^{iu\omega t} \times 8 \times \left(\right. \\ &\quad - \frac{3if_\rho^\parallel m_\rho^\parallel}{10\pi^2 t^4} \phi_{2;\rho}^\parallel(u) + \frac{3if_\rho^\parallel m_\rho^3}{20\pi^2 t^4 (v \cdot q)^2} \phi_{2;\rho}^\parallel(u) + \frac{3if_\rho^\parallel m_\rho}{10\pi^2 t^4} \phi_{3;\rho}^\perp(u) - \frac{3if_\rho^\parallel m_\rho^3}{10\pi^2 t^4 (v \cdot q)^2} \phi_{3;\rho}^\perp(u) - \frac{3f_\rho^\parallel m_\rho v \cdot q}{40\pi^2 t^3} \psi_{3;\rho}^\perp(u) \\ &\quad - \frac{3if_\rho^\parallel m_\rho^3}{160\pi^2 t^2} \phi_{4;\rho}^\parallel(u) + \frac{3if_\rho^\parallel m_\rho^3}{20\pi^2 t^4 (v \cdot q)^2} \psi_{4;\rho}^\parallel(u) + \frac{f_\rho^\perp m_\rho^2}{40tv \cdot q} \langle \bar{s}s \rangle \phi_{2;\rho}^\perp(u) + \frac{if_\rho^\perp m_\rho^2}{40} \langle \bar{s}s \rangle \psi_{3;\rho}^\parallel(u) - \frac{f_\rho^\perp m_\rho^2}{40tv \cdot q} \langle \bar{s}s \rangle \psi_{4;\rho}^\perp(u) \\ &\quad \left. \right). \end{aligned}$$

$$\begin{aligned}
& + \frac{f_\rho^\perp m_\rho^2 t}{640 v \cdot q} \langle g_s \bar{s} \sigma G s \rangle \phi_{2;\rho}^\perp(u) + \frac{i f_\rho^\perp m_\rho^2 t^2}{640} \langle g_s \bar{s} \sigma G s \rangle \psi_{3;\rho}^\parallel(u) - \frac{f_\rho^\perp m_\rho^2 t}{640 v \cdot q} \langle g_s \bar{s} \sigma G s \rangle \psi_{4;\rho}^\perp(u) + \frac{3 f_\rho^\perp m_\rho^2}{40 \pi^2 t^3 v \cdot q} m_s \phi_{2;\rho}^\perp(u) \\
& + \frac{3 i f_\rho^\perp m_\rho^2}{40 \pi^2 t^2} m_s \psi_{3;\rho}^\parallel(u) - \frac{3 f_\rho^\perp m_\rho^2}{40 \pi^2 t^3 v \cdot q} m_s \psi_{4;\rho}^\perp(u) - \frac{i f_\rho^\parallel m_\rho}{80} m_s \langle \bar{s} s \rangle \phi_{2;\rho}^\parallel(u) + \frac{i f_\rho^\parallel m_\rho^3}{160 (v \cdot q)^2} m_s \langle \bar{s} s \rangle \phi_{2;\rho}^\parallel(u) \\
& + \frac{i f_\rho^\parallel m_\rho}{80} m_s \langle \bar{s} s \rangle \phi_{3;\rho}^\perp(u) - \frac{i f_\rho^\parallel m_\rho^3}{80 (v \cdot q)^2} m_s \langle \bar{s} s \rangle \phi_{3;\rho}^\perp(u) - \frac{f_\rho^\parallel m_\rho t v \cdot q}{320} m_s \langle \bar{s} s \rangle \psi_{3;\rho}^\perp(u) - \frac{i f_\rho^\parallel m_\rho^3 t^2}{1280} m_s \langle \bar{s} s \rangle \phi_{4;\rho}^\parallel(u) \\
& + \frac{i f_\rho^\parallel m_\rho^3}{160 (v \cdot q)^2} m_s \langle \bar{s} s \rangle \psi_{4;\rho}^\parallel(u) \\
& + \int_0^\infty dt \int_0^1 du \int \mathcal{D}\underline{\alpha} e^{i\omega' t(\alpha_2+u\alpha_3)} e^{i\omega t(1-\alpha_2-u\alpha_3)} \times 8 \times \left(\right. \\
& \left. \frac{3 i f_\rho^\parallel m_\rho^3}{40 \pi^2 t^2} \Psi_{4;\rho}^\parallel(\underline{\alpha}) - \frac{3 i f_\rho^\parallel m_\rho^3}{40 \pi^2 t^2} \tilde{\Psi}_{4;\rho}^\parallel(\underline{\alpha}) + \frac{3 i f_\rho^\parallel m_\rho^3 u}{20 \pi^2 t^2} \Psi_{4;\rho}^\parallel(\underline{\alpha}) \right).
\end{aligned}$$

- [1] M. Gell-Mann, A Schematic Model of Baryons and Mesons, Phys. Lett. **8**, 214-215 (1964). [doi:10.1016/S0031-9163\(64\)92001-3](https://doi.org/10.1016/S0031-9163(64)92001-3)
- [2] G. Zweig, An SU(3) model for strong interaction symmetry and its breaking. Version 1, CERN-TH-401.
- [3] D. Ebert, R. N. Faustov, V. O. Galkin, A. P. Martynenko and V. A. Saleev, Heavy baryons in the relativistic quark model [doi:10.1007/s002880050534](https://doi.org/10.1007/s002880050534), arXiv: [hep-ph/9607314](https://arxiv.org/abs/hep-ph/9607314).
- [4] S. M. Gerasyuta and D. V. Ivanov, Charmed baryons in bootstrap quark model, Nuovo Cim. A **112**, 261-276 (1999), [doi:10.1007/BF03035848](https://doi.org/10.1007/BF03035848), arXiv: [hep-ph/0101310](https://arxiv.org/abs/hep-ph/0101310).
- [5] M. Karliner, B. Keren-Zur, H. J. Lipkin and J. L. Rosner, The Quark Model and b Baryons, Annals Phys. **324**, 2 (2009). [doi:10.1016/j.aop.2008.05.003](https://doi.org/10.1016/j.aop.2008.05.003). arXiv: [0504.1575](https://arxiv.org/abs/0504.1575)
- [6] P. A. Zyla *et al.* [Particle Data Group], Review of Particle Physics, PTEP **2020** (2020) no.8, 083C01. [doi:10.1093/ptep/ptaa104](https://doi.org/10.1093/ptep/ptaa104).
- [7] R. Aaij *et al.* [LHCb], Observation of excited Λ_b^0 baryons, Phys. Rev. Lett. **109** (2012), 172003. [doi:10.1103/PhysRevLett.109.172003](https://doi.org/10.1103/PhysRevLett.109.172003), arXiv: [1205.3452](https://arxiv.org/abs/1205.3452).
- [8] T. A. Aaltonen *et al.* [CDF], Evidence for a Bottom Baryon Resonance Λ_b^{*0} in CDF Data, Phys. Rev. D **88**, no.7, 071101 (2013). [doi:10.1103/PhysRevD.88.071101](https://doi.org/10.1103/PhysRevD.88.071101), arXiv: [1308.1760](https://arxiv.org/abs/1308.1760).
- [9] S. Navas *et al.* [Particle Data Group], Review of particle physics, Phys. Rev. D **110** (2024) no.3, 030001. [doi:10.1103/PhysRevD.110.030001](https://doi.org/10.1103/PhysRevD.110.030001),
- [10] A. M. Sirunyan *et al.* [CMS], Observation of a New Excited Beauty Strange Baryon Decaying to $\Xi_b^- \pi^+ \pi^-$, Phys. Rev. Lett. **126** (2021) no.25, 252003. [doi:10.1103/PhysRevLett.126.252003](https://doi.org/10.1103/PhysRevLett.126.252003), arXiv: [2102.04524](https://arxiv.org/abs/2102.04524).
- [11] R. Aaij *et al.* [LHCb], Observation of New Baryons in the $\Xi_b^- \pi^+ \pi^-$ and $\Xi_b^0 \pi^+ \pi^-$ Systems, Phys. Rev. Lett. **131** (2023) no.17, 171901. [doi:10.1103/PhysRevLett.131.171901](https://doi.org/10.1103/PhysRevLett.131.171901), arXiv: [2307.13399](https://arxiv.org/abs/2307.13399).
- [12] V. Crede and W. Roberts, Progress towards understanding baryon resonances, Rept. Prog. Phys. **76** (2013), 076301. [doi:10.1088/0034-4885/76/7/076301](https://doi.org/10.1088/0034-4885/76/7/076301), arXiv: [1302.7299 \[nucl-ex\]](https://arxiv.org/abs/1302.7299).
- [13] H. X. Chen, W. Chen, X. Liu, Y. R. Liu and S. L. Zhu, A review of the open charm and open bottom systems, Rept. Prog. Phys. **80** (2017) no.7, 076201. [doi:10.1088/1361-6633/aa6420](https://doi.org/10.1088/1361-6633/aa6420), arXiv: [1609.08928 \[hep-ph\]](https://arxiv.org/abs/1609.08928).
- [14] H. Y. Cheng, Charmed baryon physics circa 2021, Chin. J. Phys. **78** (2022), 324-362. [doi:10.1016/j.cjph.2022.06.021](https://doi.org/10.1016/j.cjph.2022.06.021), arXiv: [2109.01216 \[hep-ph\]](https://arxiv.org/abs/2109.01216).
- [15] H. X. Chen, W. Chen, X. Liu, Y. R. Liu and S. L. Zhu, An updated review of the new hadron states, Rept. Prog. Phys. **86** (2023) no.2, 026201. [doi:10.1088/1361-6633/aca3b6](https://doi.org/10.1088/1361-6633/aca3b6), arXiv: [2204.02649 \[hep-ph\]](https://arxiv.org/abs/2204.02649).
- [16] M. Z. Liu, Y. W. Pan, Z. W. Liu, T. W. Wu, J. X. Lu and L. S. Geng, Three ways to decipher the nature of exotic hadrons: Multiplets, three-body hadronic molecules, and correlation functions, Phys. Rept. **1108**, 1-108 (2025). [doi:10.1016/j.physrep.2024.12.001](https://doi.org/10.1016/j.physrep.2024.12.001), arXiv: [2404.06399 \[hep-ph\]](https://arxiv.org/abs/2404.06399).
- [17] H. Garcilazo, J. Vijande and A. Valcarce, Faddeev study of heavy baryon spectroscopy, J. Phys. G **34** (2007), 961-976. [doi:10.1088/0954-3899/34/5/014](https://doi.org/10.1088/0954-3899/34/5/014), arXiv: [hep-ph/0703257](https://arxiv.org/abs/hep-ph/0703257).
- [18] D. Ebert, R. N. Faustov and V. O. Galkin, Masses of excited heavy baryons in the relativistic quark model, Phys. Lett. B **659** (2008), 612-620. [doi:10.1016/j.physletb.2007.11.037](https://doi.org/10.1016/j.physletb.2007.11.037), arXiv: [0705.2957](https://arxiv.org/abs/0705.2957).
- [19] W. Roberts and M. Pervin, Heavy baryons in a quark model, Int. J. Mod. Phys. A **23** (2008), 2817-2860. [doi:10.1142/S0217751X08041219](https://doi.org/10.1142/S0217751X08041219), arXiv: [0711.2492](https://arxiv.org/abs/0711.2492).
- [20] P. G. Ortega, D. R. Entem and F. Fernandez, Quark model description of the $\Lambda_c(2940)^+$ as a molecular D^*N state and the possible existence of the $\Lambda_b(6248)$, Phys. Lett. B **718** (2013), 1381-1384. [doi:10.1016/j.physletb.2012.12.025](https://doi.org/10.1016/j.physletb.2012.12.025), arXiv: [1210.2633](https://arxiv.org/abs/1210.2633).
- [21] T. Yoshida, E. Hiyama, A. Hosaka, M. Oka and K. Sadato, Spectrum of heavy baryons in the quark model, Phys. Rev. D **92** (2015) no.11, 114029. [doi:10.1103/PhysRevD.92.114029](https://doi.org/10.1103/PhysRevD.92.114029), arXiv: [10.1103/PhysRevD.92.114029](https://arxiv.org/abs/10.1103/PhysRevD.92.114029).

- [22] H. Nagahiro, S. Yasui, A. Hosaka, M. Oka and H. Noumi, Structure of charmed baryons studied by pionic decays, Phys. Rev. D **95** (2017) no.1, 014023. [doi:10.1103/PhysRevD.95.014023](#), [arXiv:1609.01085](#).
- [23] Y. Kawakami and M. Harada, Singly heavy baryons with chiral partner structure in a three-flavor chiral model, Phys. Rev. D **99** (2019) no.9, 094016. [doi:10.1103/PhysRevD.99.094016](#), [arXiv:1902.06774](#).
- [24] K. L. Wang, Y. X. Yao, X. H. Zhong and Q. Zhao, Strong and radiative decays of the low-lying S - and P -wave singly heavy baryons, Phys. Rev. D **96**, no.11, 116016 (2017). [doi:10.1103/PhysRevD.96.116016](#), [arXiv:1709.04268](#).
- [25] H. Z. He, W. Liang, Q. F. Lü and Y. B. Dong, Strong decays of the low-lying bottom strange baryons, Sci. China Phys. Mech. Astron. **64**, no.6, 261012 (2021). [doi:10.1007/s11433-021-1704-x](#), [arXiv:2102.07391](#).
- [26] K. L. Wang, Q. F. Lü and X. H. Zhong, Interpretation of the newly observed $\Sigma_b(6097)^{\pm}$ and $\Xi_b(6227)^-$ states as the P -wave bottom baryons, Phys. Rev. D **99** (2019) no.1, 014011. [doi:10.1103/PhysRevD.99.014011](#), [arXiv:1810.02205](#).
- [27] L. X. Gutiérrez-Guerrero, A. Bashir, M. A. Bedolla and E. Santopinto, Masses of Light and Heavy Mesons and Baryons: A Unified Picture, Phys. Rev. D **100** (2019) no.11, 114032. [doi:10.1103/PhysRevD.100.114032](#), [arXiv:1911.09213](#).
- [28] L. Y. Xiao, K. L. Wang, M. S. Liu and X. H. Zhong, Possible interpretation of the newly observed Ω_b states, Eur. Phys. J. C **80** (2020) no.3, 279. [doi:10.1140/epjc/s10052-020-7823-z](#), [arXiv:2001.05110](#).
- [29] C. Garcia-Recio, J. Nieves, O. Romanets, L. L. Salcedo and L. Tolos, Odd parity bottom-flavored baryon resonances, Phys. Rev. D **87**, 034032 (2013). [doi:10.1103/PhysRevD.87.034032](#), [arXiv:1210.4755](#).
- [30] W. H. Liang, C. W. Xiao and E. Oset, Baryon states with open beauty in the extended local hidden gauge approach, Phys. Rev. D **89** (2014) no.5, 054023. [doi:10.1103/PhysRevD.89.054023](#), [arXiv:1401.1441](#).
- [31] C. S. An and H. Chen, Observed Ω_c^0 resonances as pentaquark states, Phys. Rev. D **96** (2017) no.3, 034012. [doi:10.1103/PhysRevD.96.034012](#), [arXiv:1705.08571](#).
- [32] G. Montaña, A. Feijoo and À. Ramos, A meson-baryon molecular interpretation for some Ω_c excited states, Eur. Phys. J. A **54** (2018) no.4, 64. [doi:10.1140/epja/12018-12498-1](#), [arXiv:1709.08737](#).
- [33] V. R. Debastiani, J. M. Dias, W. H. Liang and E. Oset, Molecular Ω_c states generated from coupled meson-baryon channels, Phys. Rev. D **97** (2018) no.9, 094035. [doi:10.1103/PhysRevD.97.094035](#), [arXiv:1710.04231](#).
- [34] R. Chen, A. Hosaka and X. Liu, Searching for possible Ω_c -like molecular states from meson-baryon interaction, Phys. Rev. D **97** (2018) no.3, 036016. [doi:10.1103/PhysRevD.97.036016](#), [arXiv:1711.07650](#).
- [35] J. Nieves, R. Pavao and L. Tolos, Ω_c excited states within a $SU(6)_{\text{lf}} \times \text{HQSS}$ model, Eur. Phys. J. C **78** (2018) no.2, 114. [doi:10.1140/epjc/s10052-018-5597-3](#), [arXiv:1712.00327](#).
- [36] Y. Huang, C. j. Xiao, Q. F. Lü, R. Wang, J. He and L. Geng, Strong and radiative decays of $D\Xi$ molecular state and newly observed Ω_c states, Phys. Rev. D **97** (2018) no.9, 094013. [doi:10.1103/PhysRevD.97.094013](#), [arXiv:1801.03598](#).
- [37] J. Nieves, R. Pavao and L. Tolos, Ξ_c and Ξ_b excited states within a $SU(6)_{\text{lf}} \times \text{HQSS}$ model, Eur. Phys. J. C **80** (2020) no.1, 22. [doi:10.1140/epjc/s10052-019-7568-8](#), [arXiv:1911.06089](#).
- [38] W. H. Liang and E. Oset, Observed Ω_b spectrum and meson-baryon molecular states, Phys. Rev. D **101** (2020) no.5, 054033. [doi:10.1103/PhysRevD.101.054033](#), [arXiv:2001.02929](#).
- [39] Q. X. Yu, R. Pavao, V. R. Debastiani and E. Oset, Description of the Ξ_c and Ξ_b states as molecular states, Eur. Phys. J. C **79** (2019) no.2, 167. [doi:10.1140/epjc/s10052-019-6665-z](#), [arXiv:1811.11738](#).
- [40] Y. Huang, C. j. Xiao, L. S. Geng and J. He, Strong decays of the $\Xi_b(6227)$ as a $\Sigma_b\bar{K}$ molecule, Phys. Rev. D **99** (2019) no.1, 014008. [doi:10.1103/PhysRevD.99.014008](#), [arXiv:1811.10769](#).
- [41] B. Chen and X. Liu, Assigning the newly reported $\Sigma_b(6097)$ as a P -wave excited state and predicting its partners, Phys. Rev. D **98** (2018) no.7, 074032. [doi:10.1103/PhysRevD.98.074032](#), [arXiv:1810.00389](#).
- [42] P. Yang, J. J. Guo and A. Zhang, Identification of the newly observed $\Sigma_b(6097)^{\pm}$ baryons from their strong decays, Phys. Rev. D **99** (2019) no.3, 034018. [doi:10.1103/PhysRevD.99.034018](#), [arXiv:1810.06947](#).
- [43] W. Liang and Q. F. Lü, Strong decays of the newly observed narrow Ω_b structures, Eur. Phys. J. C **80** (2020) no.3, 198. [doi:10.1140/epjc/s10052-020-7759-3](#), [arXiv:2001.02221](#).
- [44] J. X. Lu, Y. Zhou, H. X. Chen, J. J. Xie and L. S. Geng, Dynamically generated $J^P = 1/2^- (3/2^-)$ singly charmed and bottom heavy baryons, Phys. Rev. D **92** (2015) no.1, 014036. [doi:10.1103/PhysRevD.92.014036](#), [arXiv:1409.3133](#).
- [45] H. Y. Cheng and C. K. Chua, Strong Decays of Charmed Baryons in Heavy Hadron Chiral Perturbation Theory, Phys. Rev. D **75** (2007), 014006. [doi:10.1103/PhysRevD.75.014006](#), [arXiv:hep-ph/0610283](#).
- [46] H. Y. Cheng and C. K. Chua, Strong Decays of Charmed Baryons in Heavy Hadron Chiral Perturbation Theory: An Update, Phys. Rev. D **92** (2015) no.7, 074014. [doi:10.1103/PhysRevD.92.074014](#), [arXiv:1508.05653](#).
- [47] T. M. Aliev, K. Azizi, Y. Sarac and H. Sundu, Determination of the quantum numbers of $\Sigma_b(6097)^{\pm}$ via their strong decays, Phys. Rev. D **99** (2019) no.9, 094003. [doi:10.1103/PhysRevD.99.094003](#), [arXiv:1811.05686](#).
- [48] T. M. Aliev, K. Azizi, Y. Sarac and H. Sundu, Structure of the $\Xi_b(6227)^-$ resonance, Phys. Rev. D **98** (2018) no.9, 094014. [doi:10.1103/PhysRevD.98.094014](#), [arXiv:1808.08032](#).
- [49] Z. G. Wang, Analysis of the $\Omega_b(6316)$, $\Omega_b(6330)$, $\Omega_b(6340)$ and $\Omega_b(6350)$ with QCD sum rules, Int. J. Mod. Phys. A **35** (2020) no.07, 2050043. [doi:10.1142/S0217751X20500438](#), [arXiv:2001.02961](#).
- [50] H. X. Chen, W. Chen, Q. Mao, A. Hosaka, X. Liu and S. L. Zhu, P -wave charmed baryons from QCD sum rules, Phys. Rev. D **91** (2015) no.5, 054034. [doi:10.1103/PhysRevD.91.054034](#), [arXiv:1502.01103](#).
- [51] H. X. Chen, Q. Mao, W. Chen, A. Hosaka, X. Liu and S. L. Zhu, Decay properties of P -wave charmed baryons from light-cone QCD sum rules, Phys. Rev. D **95** (2017) no.9, 094008. [doi:10.1103/PhysRevD.95.094008](#), [arXiv:1703.07703](#).
- [52] X. Liu, H. X. Chen, Y. R. Liu, A. Hosaka and S. L. Zhu, Bottom baryons, Phys. Rev. D **77** (2008), 014031. [doi:](#)

- [10.1103/PhysRevD.77.014031](https://doi.org/10.1103/PhysRevD.77.014031), [arXiv:0710.0123](https://arxiv.org/abs/0710.0123).
- [53] Q. Mao, H. X. Chen, W. Chen, A. Hosaka, X. Liu and S. L. Zhu, QCD sum rule calculation for P-wave bottom baryons, Phys. Rev. D **92** (2015) no.11, 114007. [doi:10.1103/PhysRevD.92.114007](https://doi.org/10.1103/PhysRevD.92.114007), [arXiv:1510.05267](https://arxiv.org/abs/1510.05267).
- [54] W. H. Tan, H. M. Yang and H. X. Chen, Predicted $\Xi_b(6087)^0$ and further predictions, Eur. Phys. J. C **84** (2024) no.4, 382. [doi:10.1140/epjc/s10052-024-12717-6](https://doi.org/10.1140/epjc/s10052-024-12717-6), [arXiv:2311.18380](https://arxiv.org/abs/2311.18380)
- [55] X. Luo, H. M. Yang and H. X. Chen, Radiative decays of P -wave bottom baryons from light-cone sum rules, [arXiv:2407.04433 \[hep-ph\]](https://arxiv.org/abs/2407.04433).
- [56] H. M. Yang, H. X. Chen, E. L. Cui, A. Hosaka and Q. Mao, Decay properties of P -wave bottom baryons within light-cone sum rules, Eur. Phys. J. C **80** (2020) no.2, 80. [doi:10.1140/epjc/s10052-020-7637-z](https://doi.org/10.1140/epjc/s10052-020-7637-z), [arXiv:1909.13575](https://arxiv.org/abs/1909.13575).
- [57] H. M. Yang and H. X. Chen, P -wave bottom baryons of the $SU(3)$ flavor **6**_F Phys. Rev. D **101** (2020) no.11, 114013. [doi:10.1103/PhysRevD.101.114013](https://doi.org/10.1103/PhysRevD.101.114013), [arXiv:2003.07488](https://arxiv.org/abs/2003.07488).
- [58] H. M. Yang, H. X. Chen, E. L. Cui and Q. Mao, Identifying the $\Xi_b(6100)^-$ as the P -wave bottom baryon of $J^P = \frac{3}{2}^-$, Phys. Rev. D **106**, no.3, 036018 (2022). [doi:10.1103/PhysRevD.106.036018](https://doi.org/10.1103/PhysRevD.106.036018), [arXiv:2205.07224](https://arxiv.org/abs/2205.07224).
- [59] E. L. Cui, H. M. Yang, H. X. Chen and A. Hosaka, Identifying the $\Xi_b(6227)$ and $\Sigma_b(6097)$ as P -wave bottom baryons of $J^P = 3/2^-$, Phys. Rev. D **99** (2019) no.9, 094021. [doi:10.1103/PhysRevD.99.094021](https://doi.org/10.1103/PhysRevD.99.094021), [arXiv:1903.10369](https://arxiv.org/abs/1903.10369).
- [60] H. X. Chen, E. L. Cui, A. Hosaka, Q. Mao and H. M. Yang, Excited Ω_b baryons and fine structure of strong interaction, Eur. Phys. J. C **80** (2020) no.3, 256. [doi:10.1140/epjc/s10052-020-7824-y](https://doi.org/10.1140/epjc/s10052-020-7824-y), [arXiv:2001.02147](https://arxiv.org/abs/2001.02147).
- [61] M. Padmanath, R. G. Edwards, N. Mathur and M. Pardon, Excited-state spectroscopy of singly, doubly and triply-charmed baryons from lattice QCD. [arXiv:1311.4806](https://arxiv.org/abs/1311.4806).
- [62] K. U. Can, H. Bahtiyar, G. Erkol, P. Gubler, M. Oka and T. T. Takahashi, Spectrum of the Charmed Baryons in 2+1-flavor Lattice QCD, JPS Conf. Proc. **26** (2019), 022028. [doi:10.7566/JSPCP.26.022028](https://doi.org/10.7566/JSPCP.26.022028).
- [63] T. Burch, Heavy hadrons on $N_f = 2$ and $2 + 1$ improved clover-Wilson lattices. [arXiv:1502.00675](https://arxiv.org/abs/1502.00675).
- [64] M. Padmanath and N. Mathur, Quantum Numbers of Recently Discovered Ω_c^0 Baryons from Lattice QCD, Phys. Rev. Lett. **119** (2017) no.4, 042001. [doi:10.1103/PhysRevLett.119.042001](https://doi.org/10.1103/PhysRevLett.119.042001), [arXiv:1704.00259](https://arxiv.org/abs/1704.00259).
- [65] B. Grinstein, The Static Quark Effective Theory, Nucl. Phys. B **339** (1990), 253-268. [doi:10.1016/0550-3213\(90\)90349-I](https://doi.org/10.1016/0550-3213(90)90349-I).
- [66] E. Eichten and B. R. Hill, An Effective Field Theory for the Calculation of Matrix Elements Involving Heavy Quarks, Phys. Lett. B **234** (1990), 511-516. [doi:10.1016/0370-2693\(90\)92049-0](https://doi.org/10.1016/0370-2693(90)92049-0).
- [67] A. F. Falk, H. Georgi, B. Grinstein and M. B. Wise, Heavy Meson Form-factors From QCD, Nucl. Phys. B **343** (1990), 1-13. [doi:10.1016/0550-3213\(90\)90591-Z](https://doi.org/10.1016/0550-3213(90)90591-Z).
- [68] S. Narison, QCD as a Theory of Hadrons : From Partons to Confinement, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **17** (2007), 1-812 Oxford University Press, 2005,. [doi:10.1017/9781009290296](https://doi.org/10.1017/9781009290296), [arXiv:hep-ph/0205006](https://arxiv.org/abs/hep-ph/0205006).
- [69] V. Gimenez, V. Lubicz, F. Mescia, V. Porretti and J. Reyes, Operator product expansion and quark condensate from lattice QCD in coordinate space, Eur. Phys. J. C **41** (2005), 535-544. [doi:10.1140/epjc/s2005-02250-9](https://doi.org/10.1140/epjc/s2005-02250-9), [arXiv:hep-lat/0503001](https://arxiv.org/abs/hep-lat/0503001).
- [70] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, QCD and Resonance Physics. Theoretical Foundations, Nucl. Phys. B **147**, 385-447 (1979). [doi:10.1016/0550-3213\(79\)90022-1](https://doi.org/10.1016/0550-3213(79)90022-1),
- [71] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, QCD and Resonance Physics: Applications, Nucl. Phys. B **147**, 448-518 (1979) [doi:10.1016/0550-3213\(79\)90023-3](https://doi.org/10.1016/0550-3213(79)90023-3),
- [72] L. J. Reinders, H. Rubinstein and S. Yazaki, Hadron Properties from QCD Sum Rules, Phys. Rept. **127**, 1 (1985). [doi:10.1016/0370-1573\(85\)90065-1](https://doi.org/10.1016/0370-1573(85)90065-1),
- [73] M. Nielsen, F. S. Navarra and S. H. Lee, New Charmonium States in QCD Sum Rules: A Concise Review, Phys. Rept. **497**, 41-83 (2010). [doi:10.1016/j.physrep.2010.07.005](https://doi.org/10.1016/j.physrep.2010.07.005), [arXiv:0911.1958](https://arxiv.org/abs/0911.1958).
- [74] P. Gubler and D. Satow, Recent Progress in QCD Condensate Evaluations and Sum Rules, Prog. Part. Nucl. Phys. **106**, 1-67 (2019). [doi:10.1016/j.ppnp.2019.02.005](https://doi.org/10.1016/j.ppnp.2019.02.005), [arXiv:1812.00385](https://arxiv.org/abs/1812.00385).
- [75] V. M. Braun and I. E. Filyanov, QCD Sum Rules in Exclusive Kinematics and Pion Wave Function, Z. Phys. C **44** (1989), 157. [doi:10.1007/BF01548594](https://doi.org/10.1007/BF01548594).
- [76] V. L. Chernyak and I. R. Zhitnitsky, B meson exclusive decays into baryons, Nucl. Phys. B **345** (1990), 137-172. [doi:10.1016/0550-3213\(90\)90612-H](https://doi.org/10.1016/0550-3213(90)90612-H),
- [77] P. Ball, Theoretical update of pseudoscalar meson distribution amplitudes of higher twist: The Nonsinglet case, JHEP **01** (1999), 010. [doi:10.1088/1126-6708/1999/01/010](https://doi.org/10.1088/1126-6708/1999/01/010), [arXiv:hep-ph/9812375](https://arxiv.org/abs/hep-ph/9812375).
- [78] P. Ball, V. M. Braun and A. Lenz, Higher-twist distribution amplitudes of the K meson in QCD, JHEP **05** (2006), 004. [doi:10.1088/1126-6708/2006/05/004](https://doi.org/10.1088/1126-6708/2006/05/004), [arXiv:hep-ph/0603063](https://arxiv.org/abs/hep-ph/0603063).
- [79] P. Ball and R. Zwicky, $B_{d,s} \rightarrow \rho, \omega, K^*, \phi$ decay form-factors from light-cone sum rules revisited, Phys. Rev. D **71** (2005), 014029. [doi:10.1103/PhysRevD.71.014029](https://doi.org/10.1103/PhysRevD.71.014029), [arXiv:hep-ph/0412079](https://arxiv.org/abs/hep-ph/0412079).
- [80] P. Ball and V. M. Braun, Exclusive semileptonic and rare B meson decays in QCD, Phys. Rev. D **58** (1998), 094016. [doi:10.1103/PhysRevD.58.094016](https://doi.org/10.1103/PhysRevD.58.094016), [arXiv:hep-ph/9805422](https://arxiv.org/abs/hep-ph/9805422).
- [81] P. Ball, V. M. Braun, Y. Koike and K. Tanaka, Higher twist distribution amplitudes of vector mesons in QCD: Formalism and twist - three distributions, Nucl. Phys. B **529** (1998), 323-382. [doi:10.1016/S0550-3213\(98\)00356-3](https://doi.org/10.1016/S0550-3213(98)00356-3), [arXiv:hep-ph/9802299](https://arxiv.org/abs/hep-ph/9802299).
- [82] P. Ball and V. M. Braun, Higher twist distribution amplitudes of vector mesons in QCD: Twist - 4 distributions and meson mass corrections, Nucl. Phys. B **543** (1999), 201-238. [doi:10.1016/S0550-3213\(99\)00014-0](https://doi.org/10.1016/S0550-3213(99)00014-0), [arXiv:hep-ph/9810475](https://arxiv.org/abs/hep-ph/9810475).
- [83] P. Ball and G. W. Jones, Twist-3 distribution amplitudes of K^* and phi mesons, JHEP **03** (2007), 069. [doi:10.1088/1126-6708/2007/03/069](https://doi.org/10.1088/1126-6708/2007/03/069), [arXiv:hep-ph/0702100](https://arxiv.org/abs/hep-ph/0702100).
- [84] P. Ball, V. M. Braun and A. Lenz, Twist-4 distribution amplitudes of the K^* and phi mesons in QCD, JHEP **08** (2007), 090. [doi:10.1088/1126-6708/2007/08/090](https://doi.org/10.1088/1126-6708/2007/08/090), [arXiv:hep-ph/0707.1201](https://arxiv.org/abs/hep-ph/0707.1201).
- [85] M. Jamin, Flavor symmetry breaking of the quark

- condensate and chiral corrections to the Gell-Mann-Oakes-Renner relation, Phys. Lett. B **538** (2002), 71-76. [doi:10.1016/S0370-2693\(02\)01951-2](https://doi.org/10.1016/S0370-2693(02)01951-2), [arXiv:hep-ph/0201174](https://arxiv.org/abs/hep-ph/0201174).
- [86] B. L. Ioffe and K. N. Zyablyuk, Gluon condensate in charmonium sum rules with three loop corrections, Eur. Phys. J. C **27** (2003), 229-241. [doi:10.1140/epjc/s2002-01099-8](https://doi.org/10.1140/epjc/s2002-01099-8), [arXiv:hep-ph/0207183](https://arxiv.org/abs/hep-ph/0207183).
- [87] A. A. Ovchinnikov and A. A. Pivovarov, QCD sum rule calculation of the quark gluon condensate, Sov. J. Nucl. Phys. **48** (1988), 721-723.
- [88] K. C. Yang, W. Y. P. Hwang, E. M. Henley and L. S. Kisslinger, QCD sum rules and neutron proton mass difference, Phys. Rev. D **47** (1993), 3001-3012. [doi:10.1103/PhysRevD.47.3001](https://doi.org/10.1103/PhysRevD.47.3001).
- [89] M. Shifman and B. Ioffe, At the frontier of particle physics. Handbook of QCD. Vol. 1-3, World Scientific, 2001. [doi:10.1142/4544](https://doi.org/10.1142/4544).
- [90] W. Y. P. Hwang and K. C. Yang, QCD sum rules: Delta-N and Sigma0-Lambda mass splittings, Phys. Rev. D **49** (1994), 460-465. [doi:10.1103/PhysRevD.49.460](https://doi.org/10.1103/PhysRevD.49.460).