

ASYMPTOTIC EXPANSIONS FOR THE RECIPROCAL HARDY-LITTLEWOOD LOGARITHMIC INTEGRALS

GLENN BRUDA

ABSTRACT. Defining a family of recurrences, we generalize Comtet's formula for the generating function of the enumeration of indecomposable permutations. Consequently, we generalize Panaitopol's asymptotic expansion for the prime counting function, obtaining asymptotic expansions salient to the first Hardy-Littlewood conjecture.

1. INTRODUCTION

Preliminarily, a *permutation* σ of length $n \in \mathbb{Z}^+$ is a bijection $\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$. Given a permutation σ of length n , the one-line notation of σ is $m_1 m_2 \cdots m_n$, where each $m_i = \sigma(i)$. For example, $\sigma_1 = 12345$ and $\sigma_2 = 23451$ are permutations of length 5, where σ_1 is the identity mapping and $\sigma_2(k) = k + 1 \pmod{5}$ for all $k \in \{1, 2, 3, 4, 5\}$. A *proper prefix* of a permutation σ of length n is a restriction $\sigma|_{\{1, 2, \dots, k\}}$ for some $1 \leq k < n$. For example, the proper prefixes of 23451 are 2345, 234, 23, 2.

As seen in [6], an *indecomposable permutation*¹ is a permutation such that no proper prefix of itself is a permutation. For example, 321, 312, 231 are all indecomposable since 2, 3, 31, 32, 23 are not permutations. On the other hand, 123, 132, 213 are not indecomposable since 1, 12, 21 are all permutations. A thorough introduction to indecomposable permutations also appears in [6] and the OEIS sequence of their enumeration is seen at [10].

All power series in this paper should be considered as formal power series over the integers. A *formal power series* over a ring R is an infinite series of monomials

$$c_0 + c_1 x + c_2 x^2 + c_3 x^3 + \cdots$$

such that each coefficient $c_i \in R$. An introduction to formal power series may be found in [9].

Comtet introduced indecomposable permutations in [2], motivated by determining the coefficients of the reciprocal of $\sum_{n \geq 0} n! x^n$. In particular, he found that

$$1 - \left(\sum_{n \geq 0} n! x^n \right)^{-1} = \sum_{n \geq 1} \mathcal{I}_n x^n,$$

where \mathcal{I}_n is the number of indecomposable permutations of length n . We generalize this result in Proposition 2.2, determining the coefficients of the reciprocal of $\sum_{n \geq 0} (n + k - 1)! x^n$ for any $k \in \mathbb{Z}^+$.

Date: December 31, 2024.

Email: glenn.bruda@ufl.edu.

¹Also known as a *connected permutation* or an *irreducible permutation* according to [10].

We use Proposition 2.2 to determine the asymptotic behavior of the reciprocals of a class of integrals appearing in Conjecture 1.3. In particular, we obtain asymptotic expansions for said reciprocal functions.

Definition 1.1 (Asymptotic expansion, [8, Section 2.2]). *Let $\{\varphi_n\}_{n \in \mathbb{N}}$ be a sequence of real-valued functions such that $\varphi_{n+1}(x) = o(\varphi_n(x))$ as $x \rightarrow \infty$ for every $n \in \mathbb{N} := \{0, 1, 2, \dots\}$. Let f be a real-valued function. Suppose for every $N \in \mathbb{N}$ that*

$$f(x) - \sum_{n=0}^N a_n \varphi_n(x) = o(\varphi_N(x))$$

as $x \rightarrow \infty$, where $\{a_n\}$ is a sequence of real numbers. Then we say that $\sum_{n \geq 0} a_n \varphi_n(x)$ is the asymptotic expansion of $f(x)$ with respect to the asymptotic sequence $\varphi_n(x)$, writing

$$f(x) \sim \sum_{n \geq 0} a_n \varphi_n(x).$$

In [11], Panaitopol showed that the enumeration of indecomposable permutations appeared in an asymptotic expansion of the prime counting function. An extensive discussion on asymptotic expansions of the prime counting function may be found in [5].

Proposition 1.2 (Panaitopol, [11]). *Letting $\pi(x)$ denote the prime counting function, we have the asymptotic expansion*

$$\frac{1}{\pi(x)} \sim \frac{\log x}{x} \left(1 - \frac{1}{\log x} - \sum_{n \geq 2} \frac{\mathcal{I}_n}{(\log x)^n} \right),$$

where \mathcal{I}_n denotes the number of indecomposable permutations of length n .

We remark that Proposition 1.2 is equivalent to the $k = 1$ case of Theorem 2.7 since $\text{li}(x)^{-1}$ and $\pi(x)^{-1}$ have the same asymptotic expansion with respect to the asymptotic sequence $x^{-1}(\log x)^{-n+1}$. Indeed, letting $n \in \mathbb{N}$ be arbitrary and appealing to the classical estimate

$$\pi(x) = \text{li}(x) + O\left(xe^{-a\sqrt{\log x}}\right) \text{ for some } a > 0$$

of de la Vallée Poussin [3], we have

$$\begin{aligned} & \frac{x}{\pi(x) \log^{-n+1}(x)} - \frac{x}{\text{li}(x) \log^{-n+1}(x)} = x \log^{n-1}(x) \left(\frac{1}{\pi(x)} - \frac{1}{\text{li}(x)} \right) \\ &= x \log^{n-1}(x) \left(\frac{1}{\text{li}(x) + O(xe^{-a\sqrt{\log x}})} - \frac{1}{\text{li}(x)} \right) = x \log^{n-1}(x) \left(\frac{O(xe^{-a\sqrt{\log x}})}{\text{li}(x)(\text{li}(x) + O(xe^{-a\sqrt{\log x}}))} \right) \\ &= x \log^{n-1}(x) O\left(\frac{xe^{-a\sqrt{\log x}}}{\text{li}(x)^2}\right) = x \log^{n-1}(x) O\left(\frac{xe^{-a\sqrt{\log x}}}{x^2/\log^2(x)}\right) = O\left(e^{-a\sqrt{\log x}} \log^{n+1}(x)\right), \end{aligned}$$

which converges to zero for any $a > 0$ since the last function is globally nonnegative and decreasing for $x > \exp((2n+2)^2/a^2)$. With this remark noted, Proposition 1.2 may be quickly obtained from Comtet's generating function formula by following the $k = 1$ case of the proof of Theorem 2.7.

The famous prime number theorem, equivalently asserting that the prime counting function is asymptotic to the logarithmic integral, is generalized by the first Hardy-Littlewood conjecture.

Conjecture 1.3 (First Hardy-Littlewood Conjecture, [1, 7, 12]). *Let $0 < m_1 < m_2 < \dots < m_k$ be integers. Unless $P = (p, p + 2m_1, p + 2m_2, \dots, p + 2m_k)$ forms a complete residue class with respect to some prime, the number of primes $p \leq x$ such that $p, p + 2m_1, p + 2m_2, \dots, p + 2m_k$ are prime, denoted $\pi_P(x)$, is asymptotic² to*

$$2^k \prod_{\substack{q \text{ prime} \\ q \geq 3}} \left(1 - \frac{1}{q}\right)^{-k-1} \left(1 - \frac{w(q; 2m_1, 2m_2, \dots, 2m_k)}{q}\right) \int_2^x \frac{1}{(\log t)^{k+1}} dt,$$

where $w(q; 2m_1, 2m_2, \dots, 2m_k)$ is the number of distinct residues of $0, 2m_1, 2m_2, \dots, 2m_k$ modulo q .

Like the prime number theorem, Conjecture 1.3 does not concern an asymptotic expansion; rather, only the lowest order asymptotic term of $\pi_P(x)$ is regarded. Theorem 2.7, the main result of this paper, gives the asymptotic expansion

$$\left(\int_0^x \frac{1}{(\log t)^k} dt\right)^{-1} \sim \frac{(\log x)^k}{x} \left(1 - \frac{k}{\log x} - \sum_{n \geq 1} \frac{a_n^{(k)}}{(\log x)^{n+1}}\right)$$

for any $k \in \mathbb{Z}^+$, where the dashed integral denotes the Cauchy principal value and the sequence $a_n^{(k)}$ is as defined in Definition 2.1. In particular, Theorem 2.7 provides auxiliary information for a potential strengthening of Conjecture 1.3 such that we may eventually obtain (or at least conjecture) a complete asymptotic expansion for $\pi_P(x)$.

2. RESULTS

To prove Theorem 2.7, we first generalize Comtet's generating function formula in Proposition 2.2. To this end, we define the following family of recurrences.

Definition 2.1. *For $k \in \mathbb{Z}^+$, let $a_n^{(k)}$ be the sequence defined by the recurrence*

$$(k-1)!a_n^{(k)} = (n+k)! - k(n+k-1)! - \sum_{m=0}^{n-2} (n-m+k-2)!a_{m+1}^{(k)}$$

with $a_0^{(k)} = 1$.

As noted in [5, pg.357], $a_n^{(1)}$ is the number of indecomposable permutations of length $n+1$. The sequences for $k=2, 3, 4$ and their respective OEIS A-numbers are listed in Table 1.

Proposition 2.2. *Let $k \in \mathbb{Z}^+$. Then*

$$1 - (k-1)! \left(\sum_{n \geq 0} (n+k-1)!x^n\right)^{-1} = kx + \sum_{n \geq 1} a_n^{(k)} x^{n+1},$$

taken as formal power series.

²We say that functions f, g are *asymptotic* if $\lim_{x \rightarrow \infty} f(x)/g(x) = 1$.

TABLE 1. The first few terms of $a_n^{(2)}$, $a_n^{(3)}$, $a_n^{(4)}$ and their OEIS A-numbers.

n	0	1	2	3	4	5	6	7	8	9
$a_n^{(2)}$ (A111537)	1	2	8	44	296	2312	20384	199376	2138336	24936416
$a_n^{(3)}$ (A111546)	1	3	15	99	783	7083	71415	789939	9485343	122721723
$a_n^{(4)}$ (A111556)	1	4	24	184	1664	17024	192384	2366144	31362304	444907264

Proof. Let $k \in \mathbb{Z}^+$. Let $I_k(x) := kx + \sum_{n \geq 1} a_n^{(k)} x^{n+1}$ and for $m \in \mathbb{N}$, let $A_m(x) := \sum_{n \geq 0} (n+m)! x^n$. Then by the definition of $a_n^{(k)}$,

$$\begin{aligned}
(k-1)!I_k(x) &= k!x - \sum_{n \geq 1} (k-1)!a_n^{(k)} x^{n+1} \\
&= k!x + x(A_k(x) - k!) - kx(A_{k-1}(x) - (k-1)!) - x \sum_{n \geq 1} \sum_{m=0}^{n-2} (n-m+k-2)! a_{m+1}^{(k)} x^n \\
&= k!x + xA_k(x) - kxA_{k-1}(x) - x^3 \sum_{n \geq 0} \sum_{m=0}^n (n-m+k)! a_{m+1}^{(k)} x^n.
\end{aligned}$$

Noting that

$$\sum_{n \geq 0} a_{n+1}^{(k)} x^n = x^{-2} \sum_{n \geq 1} a_n^{(k)} x^{n+1} = x^{-2}(I_k(x) - kx),$$

we obtain

$$\begin{aligned}
(k-1)!I_k(x) &= k!x + xA_k(x) - kxA_{k-1}(x) - x(I_k(x) - kx)A_k(x) \\
&= k!x - kxA_{k-1}(x) - x(I_k(x) - kx - 1)A_k(x).
\end{aligned}$$

It then follows that

$$I_k(x) = \frac{k!x + (1+kx)xA_k(x) - kxA_{k-1}(x)}{(k-1)! + xA_k(x)}.$$

We may now simplify the above since

$$(k-1)! + xA_k(x) = A_{k-1}(x).$$

Indeed, we thusly have

$$\begin{aligned}
I_k(x) &= \frac{k!x + (1+kx)(A_{k-1}(x) - (k-1)!) - kxA_{k-1}(x)}{A_{k-1}(x)} \\
&= 1 - \frac{(k-1)!}{A_{k-1}(x)} \\
&= 1 - (k-1)! \left(\sum_{n \geq 0} (n+k-1)! x^n \right)^{-1},
\end{aligned}$$

as desired. \square

Now with Proposition 2.2 proven, we define the *exponential integral* and the *Hardy-Littlewood logarithmic integrals*. Then, after invoking the standard expansion stated in Proposition 2.4 and inductively proving Lemma 2.6, we finally prove Theorem 2.7.

Definition 2.3 (Exponential integral). *We define $\text{Ei} : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ by*

$$\text{Ei}(x) := \int_{-\infty}^x \frac{e^t}{t} dt,$$

where the dashed integral denotes the Cauchy principal value.

Proposition 2.4 ([4, Equation 6.12.2]). *As $x \rightarrow \infty$, we have the asymptotic expansion*

$$\text{Ei}(x) \sim \frac{e^x}{x} \sum_{n \geq 0} \frac{n!}{x^n}.$$

Definition 2.5 (Hardy-Littlewood logarithmic integrals). *For $k \in \mathbb{Z}^+$, we define $\text{li}_k : \mathbb{R} \setminus \{1\} \rightarrow \mathbb{R}$ by*

$$\text{li}_k(x) := \int_0^x \frac{1}{(\log t)^k} dt,$$

where the dashed integral denotes the Cauchy principal value. Note that $\text{li}_1(x)$ is the usual logarithmic integral, denoted $\text{li}(x)$.

Lemma 2.6. *Let $k \in \mathbb{Z}^+$. Then*

$$\text{li}_k(x) = \frac{1}{(k-1)!} \left(\text{li}(x) - x \sum_{i=0}^{k-2} \frac{i!}{(\log x)^{i+1}} \right).$$

Proof. This result lends a simple inductive proof via integration by parts. For $k = 1$, the claim follows from the definition of the logarithmic integral. For the inductive step, note that

$$\text{li}_k(x) = \frac{x}{(\log x)^k} + \int_0^x \frac{k}{(\log t)^{k+1}} dt.$$

So by inductive hypothesis,

$$\begin{aligned} \text{li}_{k+1}(x) &= \int_0^x \frac{1}{(\log t)^{k+1}} dt = \frac{1}{k!} \left(\text{li}(x) - x \sum_{i=0}^{k-2} \frac{i!}{(\log x)^{i+1}} \right) - \frac{x}{k(\log x)^k} \\ &= \frac{1}{k!} \left(\text{li}(x) - x \sum_{i=0}^{k-1} \frac{i!}{(\log x)^{i+1}} \right). \end{aligned}$$

□

Theorem 2.7. *Let $k \in \mathbb{Z}^+$. Then we have the asymptotic expansion*

$$\frac{1}{\text{li}_k(x)} \sim \frac{(\log x)^k}{x} \left(1 - \frac{k}{\log x} - \sum_{n \geq 1} \frac{a_n^{(k)}}{(\log x)^{n+1}} \right).$$

Proof. Let $k \in \mathbb{Z}^+$ and $A(x) := \sum_{n \geq 0} n!x^n$. By Proposition 2.2,

$$\begin{aligned} kx + \sum_{n \geq 1} a_n^{(k)} x^{n+1} &= 1 - (k-1)! \left(\sum_{n \geq 0} (n+k-1)! x^n \right)^{-1} \\ &= 1 - (k-1)! \left(\sum_{n \geq k-1} n! x^{n-k+1} \right)^{-1} \\ &= 1 - \frac{(k-1)! x^{k-1}}{\sum_{n \geq k-1} n! x^n} \\ &= 1 - \frac{(k-1)! x^{k-1}}{A(x) - \sum_{i=0}^{k-2} i! x^i}. \end{aligned}$$

After a change of variable, we obtain

$$\frac{(k-1)! x^{1-k}}{A(1/x) - \sum_{i=0}^{k-2} i! x^{-i}} = 1 - \frac{k}{x} - \sum_{n \geq 1} \frac{a_n^{(k)}}{x^{n+1}}.$$

Proposition 2.4 gives the relation $A(1/x) \sim x e^{-x} \text{Ei}(x)$; thus,

$$\begin{aligned} 1 - \frac{k}{x} - \sum_{n \geq 1} \frac{a_n^{(k)}}{x^{n+1}} &\sim \frac{(k-1)! x^{1-k}}{x e^{-x} \text{Ei}(x) - \sum_{i=0}^{k-2} i! x^{-i}} \\ &= \frac{(k-1)!}{x^k e^{-x} \left(\text{Ei}(x) - e^x \sum_{i=0}^{k-2} i! x^{-i-1} \right)}. \end{aligned}$$

Noting that $\text{Ei}(x) = \text{li}(e^x)$, by Lemma 2.6 we have

$$x^k e^{-x} \left(1 - \frac{k}{x} - \sum_{n \geq 1} \frac{a_n^{(k)}}{x^{n+1}} \right) \sim \frac{(k-1)!}{\text{Ei}(x) - e^x \sum_{i=0}^{k-2} i! x^{-i-1}} = \frac{1}{\text{li}_k(e^x)}.$$

Thus,

$$\frac{1}{\text{li}_k(x)} \sim \frac{(\log x)^k}{x} \left(1 - \frac{k}{\log x} - \sum_{n \geq 1} \frac{a_n^{(k)}}{(\log x)^{n+1}} \right),$$

as desired. □

ACKNOWLEDGEMENTS

We thank Steven J. Miller for his guidance and for recommending trying to generalize the $k = 1$ case of Theorem 2.7.

REFERENCES

- [1] Soren Laing Aletheia-Zomlefer, Lenny Fukshansky, and Stephan Ramon Garcia, *The Bateman-Horn conjecture: Heuristic, history, and applications*, *Expositiones Mathematicae* **38** (2020), no. 4, 430–479.
- [2] L. Comtet, *Sur les coefficients de l'inverse de la série formelle $\sum n!x^n$* , *C. R. Acad. Sci. Paris Sér. A-B* **275** (1972), A569–A572.
- [3] Charles Jean de la Vallée Poussin, *Sur la fonction zeta de riemann et le nombre des nombres premiers inferieur a une limite donnée*, *Mémoires couronnés de L'Académie de Belgique* **59** (1899).
- [4] *NIST Digital Library of Mathematical Functions*. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [5] Jesse Elliott, *Asymptotic expansions of the prime counting function*, *Journal of Number Theory* **238** (2022), 353–395.
- [6] Alice L.L. Gao, Sergey Kitaev, and Philip B. Zhang, *On pattern avoiding indecomposable permutations*, *Integers* **18** (2018).
- [7] G. H. Hardy and J. E. Littlewood, *Some problems of 'Partitio numerorum'; III: On the expression of a number as a sum of primes*, *Acta Mathematica* **44** (1923), 1–70.
- [8] John K. Hunter, *Asymptotic Analysis and Singular Perturbation Theory*, 2004. University of California at Davis.
- [9] Ivan Niven, *Formal Power Series*, *The American Mathematical Monthly* **76** (1969), no. 8, 871–889, available at <https://doi.org/10.1080/00029890.1969.12000359>.
- [10] *Number of connected permutations of length n*, 2024. Entry A003319 in The On-Line Encyclopedia of Integer Sequences, <https://oeis.org/A003319>.
- [11] L. Panaitopol, *A formula for $\pi(x)$ applied to a result of Koninck-Ivić*, *Nieuw Arch. Wiskd.* **1** (2000), no. 1, 55–56.
- [12] László Tóth, *On the Asymptotic Density of Prime k -tuples and a Conjecture of Hardy and Littlewood*, *Computational Methods in Science and Technology* **25** (2019), no. 3, 143–148.