

# K-MODULI OF PURE STATES OF FOUR QUBITS

IVAN CHELTSOV, MAKSYM FEDORCHUK, KENTO FUJITA, ANNE-SOPHIE KALOGHIROS

ABSTRACT. We find all K-polystable limits of divisors in  $(\mathbb{P}^1)^4$  of degree  $(1, 1, 1, 1)$  and explicitly describe the associated irreducible component of the K-moduli space.

## 1. INTRODUCTION

Smooth Fano 3-folds have been classified by Iskovskikh, Mori, Mukai into 105 families. Fano 3-folds in each deformation family can be parametrised by an irreducible rational variety. For instance, the family  $\mathbb{N}^{\circ} 4.1$  contains smooth Fano 3-folds of Picard rank 4 and anticanonical degree 24. These 3-folds are smooth divisors of degree  $(1, 1, 1, 1)$  in  $(\mathbb{P}^1)^4$ . Moreover, it follows from [11, 17, 29] that each smooth member of this deformation family can be given in  $\mathbb{P}_{x_1, y_1}^1 \times \mathbb{P}_{x_2, y_2}^1 \times \mathbb{P}_{x_3, y_3}^1 \times \mathbb{P}_{x_4, y_4}^1$  by the equation

$$(\heartsuit) \quad \frac{a+d}{2}(x_1x_2x_3x_4 + y_1y_2y_3y_4) + \frac{a-d}{2}(x_1x_2y_3y_4 + y_1y_2x_3x_4) + \\ + \frac{b+c}{2}(x_1y_2x_3y_4 + y_1x_2y_3x_4) + \frac{b-c}{2}(x_1y_2y_3x_4 + y_1x_2x_3y_4) = 0$$

for some  $a, b, c, d \in \mathbb{C}$  such that  $(a^2 - b^2)(a^2 - c^2)(a^2 - d^2)(b^2 - c^2)(b^2 - d^2)(c^2 - d^2) \neq 0$ . So, smooth Fano 3-folds in the family  $\mathbb{N}^{\circ} 4.1$  can be parametrised by an open subset in  $\mathbb{P}^3$ . These 3-folds together with their singular limits appeared in many problems in algebraic geometry [4, 5, 7, 9, 10, 19] and also in mathematical physics [11, 13, 16, 17, 18, 25, 29]. For example, all smooth 3-folds in this family are known to be K-stable [4].

On the other hand, we know that there exists a projective moduli space  $M_{3,24}^{\text{Kps}}$  whose closed points parametrise K-polystable Fano 3-folds of (anticanonical) degree 24 [24]. The goal of this paper is to describe the irreducible component of this space whose points parametrise smooth members of the family  $\mathbb{N}^{\circ} 4.1$  together with their K-polystable limits.

**Main Theorem.** *Let  $\mathbf{M}$  be the irreducible component of the moduli space  $M_{3,24}^{\text{Kps}}$  whose points parametrise K-polystable limits of smooth Fano 3-folds in the family  $\mathbb{N}^{\circ} 4.1$ . Then*

- $\mathbf{M}$  is the blow-up at a smooth point of  $\mathbb{P}(1, 3, 4, 6)$  with weights  $(1, 2, 3)$ ;
- if  $X$  is a Fano 3-fold parameterised by a closed point in  $\mathbf{M}$ , then
  - either  $X$  is a divisor in  $(\mathbb{P}^1)^4$  given by  $(\heartsuit)$  such that

$$(a : b : c : d) \notin \{(1 : 0 : 0 : 0), (0 : 1 : 0 : 0), (0 : 0 : 1 : 0), (0 : 0 : 0 : 1), (1 : \pm 1 : \pm 1 : \pm 1)\},$$

- or  $X$  is a divisor in  $\mathbb{P}(1_{s_1}, 1_{t_1}, 2_{w_1}) \times \mathbb{P}(1_{s_2}, 1_{t_2}, 2_{w_2})$  given by

$$(\diamond) \quad w_1w_2 = as_1t_1s_2t_2 + \frac{b+c}{4}(s_1^2s_2^2 + t_1^2t_2^2) + \frac{b-c}{4}(s_1^2t_2^2 + t_1^2s_2^2)$$

for any  $(a : b : c) \in \mathbb{P}^2$ .

For more details about the 3-fold  $\mathbf{M}$  in this theorem, see Proposition 5.3.

*Remark 1.1.* Let  $X$  be one of the K-polystable Fano 3-folds described in Main Theorem, and let  $\tilde{X} \rightarrow X$  be its standard resolution of singularities which is given by the ordinary blow up of the singular locus. Then the following are equivalent

- (1) the Intermediate Jacobian of the 3-fold  $\tilde{X}$  is not trivial,
- (2) the Intermediate Jacobian of the 3-fold  $\tilde{X}$  is a smooth elliptic curve,
- (3) either  $X$  is smooth or  $X$  is a divisor in  $(\mathbb{P}(1, 1, 2))^2$  given by  $(\diamond)$  such that

$$\text{Sing}(X) = \{s_1 = t_1 = w_2 = 0\} \cup \{s_2 = t_2 = w_1 = 0\}.$$

This reminds a similar phenomenon discovered in [12] for degenerations of cubic 3-folds, which is also discussed in [20, Remark 5.6].

The structure of the paper is the following. In Section 2, we describe the GIT moduli space of divisors of degree  $(1, 1, 1, 1)$  in  $(\mathbb{P}^1)^4$ , and we describe singularities of divisors  $(\heartsuit)$ . In Section 3, we study basic properties of divisors in  $(\mathbb{P}(1, 1, 2))^2$  given by the equation  $(\diamond)$ . In Section 4, we prove K-polystability of the following Fano 3-folds:

- all irreducible divisors in  $(\mathbb{P}^1)^4$  given by  $(\heartsuit)$ ;
- all divisors in  $(\mathbb{P}(1, 1, 2))^2$  given by  $(\diamond)$ .

In Section 5, we complete the proof of Main Theorem. In Section 6, we study K-semistable degenerations of smooth Fano 3-folds in the family  $\mathbb{N}^{\circ}4.1$ . Finally, in Appendix A, we show that singular K-polystable divisors of degree  $(1, 1, 1, 1)$  in  $(\mathbb{P}^1)^4$  and their K-polystable limits can be constructed as anticanonical models of very simple blow ups of  $\mathbb{P}^3$ .

**Acknowledgements.** Ivan Cheltsov has been supported by EPSRC grant EP/V054597/1 and by Simons Collaboration grant *Moduli of varieties*. Kento Fujita has been supported by JSPS KAKENHI Grant Number 22K03269, Royal Society International Collaboration Award ICA\1\23109 and Asian Young Scientist Fellowship. Anne-Sophie Kaloghiros has been supported by EPSRC grant EP/V056689/1.

## 2. DIVISORS OF DEGREE $(1, 1, 1, 1)$ IN $(\mathbb{P}^1)^4$

In this section, we study divisors of degree  $(1, 1, 1, 1)$  in  $\mathbb{P}_{x_1, y_1}^1 \times \mathbb{P}_{x_2, y_2}^1 \times \mathbb{P}_{x_3, y_3}^1 \times \mathbb{P}_{x_4, y_4}^1$ . Let  $V = H^0((\mathbb{P}^1)^4, \mathcal{O}(1, 1, 1, 1))$ . Set  $\Gamma_0 = \text{SL}_2(\mathbb{C})^4$ , set

$$\Gamma = \Gamma_0 \rtimes \mathfrak{S}_4 = \text{SL}_2(\mathbb{C})^4 \rtimes \mathfrak{S}_4,$$

and consider the natural actions of these two groups on the space  $V$ . In this section, we will study the projective GIT quotient  $\mathbb{P}(V)^{\text{ss}} // \Gamma$ . Here, we use classical notation for the projectivization  $\mathbb{P}(V)$  — the variety of one-dimensional subspaces in  $V$ .

Note that, physicists often use the symbol SLOCC in place of  $\Gamma_0$ , because this reductive group is called as *Stochastic Local Operations with Classical Communication*.

**2.1. Normal forms.** Forms in  $V$  were studied extensively in the context of quantum computing [11, 16, 17, 18, 25, 29], as they represent entanglement of systems of four qubits. In particular, it was shown that such forms can be put in normal forms, which gives

**Corollary 2.1** ([29, 11]). *Let  $f$  be a non-zero form in  $V$  such that  $\{f = 0\}$  is smooth. Then the orbit  $\Gamma_0.f$  contains the form*

$$(G_{a,b,c,d}) \quad \frac{a+d}{2}(x_1x_2x_3x_4 + y_1y_2y_3y_4) + \frac{a-d}{2}(x_1x_2y_3y_4 + y_1y_2x_3x_4) + \\ + \frac{b+c}{2}(x_1y_2x_3y_4 + y_1x_2y_3x_4) + \frac{b-c}{2}(x_1y_2y_3x_4 + y_1x_2x_3y_4)$$

for some  $(a, b, c, d) \in \mathbb{C}^4$ .

Now, we consider the following vector subspace:

$$\mathcal{G} = \{f \in V \mid f \text{ is the form } (G_{a,b,c,d}) \text{ for some } (a, b, c, d) \in \mathbb{C}^4\} \subset V.$$

Let  $G$  be the subgroup in  $\text{Aut}((\mathbb{P}^1)^4)$  generated by the following involutions:

$$\begin{aligned} \tau_1 &: ((x_1 : y_1), (x_2 : y_2), (x_3 : y_3), (x_4 : y_4)) \mapsto ((x_2 : y_2), (x_1 : y_1), (x_4 : y_4), (x_3 : y_3)), \\ \tau_2 &: ((x_1 : y_1), (x_2 : y_2), (x_3 : y_3), (x_4 : y_4)) \mapsto ((x_4 : y_4), (x_3 : y_3), (x_2 : y_2), (x_1 : y_1)), \\ \tau_3 &: ((x_1 : y_1), (x_2 : y_2), (x_3 : y_3), (x_4 : y_4)) \mapsto ((y_1 : x_1), (y_2 : x_2), (y_3 : x_3), (y_4 : x_4)), \\ \tau_4 &: ((x_1 : y_1), (x_2 : y_2), (x_3 : y_3), (x_4 : y_4)) \mapsto ((x_1 : -y_1), (x_2 : -y_2), (x_3 : -y_3), (x_4 : -y_4)). \end{aligned}$$

Then  $G \simeq (\mathbb{Z}/2\mathbb{Z})^4$ , every form in  $\mathcal{G}$  is  $G$ -invariant,  $\mathcal{G}$  contains all  $G$ -invariant forms in  $V$ , and  $\mathcal{G}$  contains defining polynomials of all smooth divisors of degree  $(1, 1, 1, 1)$  in  $(\mathbb{P}^1)^4$ .

For sufficiently general form  $f \in \mathcal{G}$ , the  $G$ -invariant divisor  $\{f = 0\} \subset (\mathbb{P}^1)^4$  is smooth. The following lemma describes reducible divisors  $\{f = 0\}$  for  $f \in \mathcal{G}$ .

**Lemma 2.2.** *Let  $f$  be a non-zero form in  $\mathcal{G}$ . Then*

$$f \text{ is reducible} \iff \begin{cases} a^2 = b^2 = c^2 = d^2 \text{ or} \\ a = b = c = 0 \text{ or} \\ a = b = d = 0 \text{ or} \\ a = c = d = 0 \text{ or} \\ b = c = d = 0 \end{cases} \iff f \in \Gamma.(x_1x_2 - y_1y_2)(x_3x_4 - y_3y_4),$$

where  $(x_1x_2 - y_1y_2)(x_3x_4 - y_3y_4)$  is the form  $(G_{a,b,c,d})$  with  $(a, b, c, d) = (0, 0, 0, 1)$ .

*Proof.* Direct computation. □

To study the GIT quotient  $\mathbb{P}(V)^{\text{ss}} // \Gamma$ , we need the following lemma.

**Lemma 2.3.** *Let  $f$  and  $f'$  be non-zero forms in  $\mathcal{G}$ , let  $X = \{f = 0\}$  and  $X' = \{f' = 0\}$ . Then  $X$  and  $X'$  are isomorphic if and only if  $f$  and  $f'$  lie in one  $\Gamma$ -orbit.*

*Proof.* By Lemma 2.2, we may assume that both  $X$  and  $X'$  are irreducible. In fact, they are normal by Theorem 2.8, but we do not need this. Then the restriction homomorphism

$$\text{Pic}((\mathbb{P}^1)^4) \rightarrow \text{Pic}(X)$$

is an isomorphism by the Lefschetz hyperplane section theorem.

For  $i \in \{1, 2, 3, 4\}$ , let  $\text{pr}^i: X \rightarrow (\mathbb{P}^1)^3$  be the projection to all but the  $i^{\text{th}}$  factor of  $(\mathbb{P}^1)^4$ . Then  $\text{pr}^i$  is birational, so the Mori cone  $\overline{\text{NE}}(X)$  is spanned by all  $\text{pr}^i$ -exceptional curves. Since  $\text{Nef}(X)$  is the dual cone of  $\overline{\text{NE}}(X)$ , it is spanned by the pullbacks

$$\text{pr}_1^*(\mathcal{O}_{\mathbb{P}^1}(1)), \quad \text{pr}_2^*(\mathcal{O}_{\mathbb{P}^1}(1)), \quad \text{pr}_3^*(\mathcal{O}_{\mathbb{P}^1}(1)), \quad \text{pr}_4^*(\mathcal{O}_{\mathbb{P}^1}(1)),$$

where  $\text{pr}_i: X \rightarrow \mathbb{P}^1$  is the projection to the  $i^{\text{th}}$  factor of  $(\mathbb{P}^1)^4$ . Using this, we see that any isomorphism  $X \simeq X'$  is induced by an automorphism of  $(\mathbb{P}^1)^4$ .  $\square$

We now consider the action of  $\Gamma_0$  and  $\Gamma$  on forms in  $\mathcal{G}$ . For convenience, we set

$$\begin{cases} u_1 = x_1x_2x_3x_4 + x_1x_2y_3y_4 + y_1y_2x_3x_4 + y_1y_2y_3y_4, \\ u_2 = x_1x_2x_3x_4 - x_1x_2y_3y_4 - y_1y_2x_3x_4 + y_1y_2y_3y_4, \\ u_3 = x_1y_2x_3y_4 + x_1y_2y_3x_4 + y_1x_2x_3y_4 + y_1x_2y_3x_4, \\ u_4 = x_1y_2x_3y_4 - x_1y_2y_3x_4 - y_1x_2x_3y_4 + y_1x_2y_3x_4. \end{cases}$$

Then, we can rewrite the form  $(G_{a,b,c,d})$  in  $\mathcal{G}$  as

$$\frac{a}{2}u_1 + \frac{d}{2}u_2 + \frac{b}{2}u_3 + \frac{c}{2}u_4.$$

Let  $\Gamma_0^{\mathcal{G}}$  and  $\Gamma^{\mathcal{G}}$  be the images in  $\text{GL}(\mathcal{G})$  of the subgroups of the groups  $\Gamma_0$  and  $\Gamma$  that leave the subspace  $\mathcal{G}$  invariant, respectively. By [16, Section II], the group  $\Gamma_0^{\mathcal{G}}$  is generated by

$$\begin{aligned} (u_1, u_2, u_3, u_4) &\mapsto (-u_1, -u_2, u_3, u_4), \\ (u_1, u_2, u_3, u_4) &\mapsto (u_2, u_1, u_3, u_4), \\ (u_1, u_2, u_3, u_4) &\mapsto (u_2, u_3, u_4, u_1), \end{aligned}$$

which implies that  $\Gamma_0^{\mathcal{G}} \simeq (\mathbb{Z}/2\mathbb{Z})^3 \rtimes \mathfrak{S}_4$ , and  $\Gamma_0^{\mathcal{G}}$  is isomorphic to the Weyl group  $W(D_4)$ . Similarly, the group  $\Gamma^{\mathcal{G}}$  is generated by  $\Gamma_0^{\mathcal{G}}$ , the transformation

$$(u_1, u_2, u_3, u_4) \mapsto (u_1, u_2, u_3, -u_4)$$

and the transformation

$$(u_1, u_2, u_3, u_4) \mapsto \left( \frac{u_1 + u_2 + u_3 + u_4}{2}, \frac{u_1 + u_2 - u_3 - u_4}{2}, \frac{u_1 - u_2 + u_3 - u_4}{2}, \frac{u_1 - u_2 - u_3 + u_4}{2} \right).$$

The group  $\Gamma^{\mathcal{G}}$  is isomorphic to the Weyl group  $W(F_4)$ , and its GAP ID is [1152,157478]. Since  $-1 \in \Gamma^{\mathcal{G}}$  acts trivially on  $\mathbb{P}(\mathcal{G})$ , we obtain an effective action of the group

$$\Gamma^{\mathcal{G}}/\langle -1 \rangle \simeq W(F_4)/\langle -1 \rangle \simeq (\mathfrak{A}_4 \times \mathfrak{A}_4) \times (\mathbb{Z}/2\mathbb{Z})^2$$

on  $\mathbb{P}(\mathcal{G}) \simeq \mathbb{P}_{a,b,c,d}^3$ . The GAP ID of the group  $W(F_4)/\langle -1 \rangle$  is [576,8654].

In the following, we identify  $\Gamma^{\mathcal{G}} = W(F_4)$ ,  $\Gamma^{\mathcal{G}}/\langle -1 \rangle = W(F_4)/\langle -1 \rangle$  and  $\mathbb{P}(\mathcal{G}) = \mathbb{P}_{a,b,c,d}^3$ .

**2.2. GIT moduli space.** Now, we recall some results on the GIT quotient  $\mathbb{P}(V)^{\text{ss}} // \Gamma$ . Recall that  $V = H^0((\mathbb{P}^1)^4, \mathcal{O}(1, 1, 1, 1))$ , and fix its basis: for  $i, j, k, l \in \{0, 1\}$ , we let

$$a_m = x_1^i y_1^{1-i} x_2^j y_2^{1-j} x_3^k y_3^{1-k} x_4^l y_4^{1-l}$$

for  $m = 8i + 4j + 2k + l$ . Therefore, for example, we have  $a_0 = x_1x_2x_3x_4$  and  $a_3 = x_1x_2y_3y_4$ . Let  $W = V^\vee$ , and let  $\mathcal{S}$  be the symmetric algebra of  $W$ . Then it follows from [25] that the  $\Gamma_0$ -invariant subring of  $\mathcal{S}$  is  $\mathcal{S}^{\Gamma_0} = \mathbb{C}[H, L, M, D]$ , where

$$H = a_0a_{15} - a_1a_{14} - a_2a_{13} + a_3a_{12} - a_4a_{11} + a_5a_{10} + a_6a_9 - a_7a_8,$$

$$L = \begin{vmatrix} a_0 & a_4 & a_8 & a_{12} \\ a_1 & a_5 & a_9 & a_{13} \\ a_2 & a_6 & a_{10} & a_{14} \\ a_3 & a_7 & a_{11} & a_{15} \end{vmatrix}, \quad M = \begin{vmatrix} a_0 & a_8 & a_2 & a_{10} \\ a_1 & a_9 & a_3 & a_{11} \\ a_4 & a_{12} & a_6 & a_{14} \\ a_5 & a_{13} & a_7 & a_{15} \end{vmatrix}$$

and

$$D = \begin{vmatrix} a_0a_9 - a_1a_8 & a_0a_{11} + a_2a_9 - a_1a_{10} - a_3a_8 & a_2a_{11} - a_3a_{10} \\ a_0a_{13} + a_4a_9 - a_1a_{12} - a_5a_8 & (a_0a_{15} + a_2a_{13} + a_4a_{11} + a_6a_9) & a_2a_{15} + a_6a_{11} - a_3a_{14} - a_7a_{10} \\ a_4a_{13} - a_5a_{12} & -a_1a_{14} - a_3a_{12} - a_5a_{10} - a_7a_8 & a_4a_{15} + a_6a_{13} - a_5a_{14} - a_7a_{12} & a_6a_{15} - a_7a_{14} \end{vmatrix}.$$

To describe  $\mathcal{S}^\Gamma = (\mathcal{S}^{\Gamma_0})^{\mathfrak{S}_4}$ , let  $R, S$  and  $T$  be the forms of degrees 6, 8 and 12 given by

$$(2.4) \quad \begin{aligned} R &= H(L - M) + 3D, \\ S &= \frac{1}{12}H^4 - \frac{2}{3}H^2L + \frac{2}{3}H^2M - 2HD + \frac{4}{3}(L^2 + LM + M^2), \\ T &= \frac{1}{216}H^6 - \frac{1}{18}H^4(L - M) - \frac{1}{6}H^3D + \frac{1}{9}H^2(2L^2 - LM + 2M^2) \\ &\quad + \frac{2}{3}H(L - M)D - \frac{8}{27}(L^3 - M^3) - \frac{4}{9}LM(L - M) + D^2. \end{aligned}$$

Observe that  $H, R, S, T$  are algebraically independent.

**Proposition 2.5** ([27, 11, 16]). *One has  $\mathcal{S}^\Gamma = \mathbb{C}[H, R, S, T]$ , so that*

$$\mathbb{P}(V)^{\text{ss}} // \Gamma \simeq \mathbb{P}(1, 3, 4, 6).$$

*Proof.* Let us give a proof for readers' convenience. Under the natural embedding

$$\mathfrak{S}_3 = \{\text{id}, (1, 2, 3), (1, 3, 2), (1, 2), (1, 3), (2, 3)\} \subset \mathfrak{S}_4,$$

we have  $\mathfrak{S}_4 = (\mathbb{Z}/2\mathbb{Z})^2 \rtimes \mathfrak{S}_3$ , where  $(\mathbb{Z}/2\mathbb{Z})^2 \triangleleft \mathfrak{S}_4$  is the Klein four-group

$$\{\text{id}, (1, 2)(3, 4), (1, 3)(2, 4), (1, 4)(2, 3)\}.$$

We can directly check the following:

- We have  $H^\sigma = H$  and  $R^\sigma = R$  for every  $\sigma \in \mathfrak{S}_4$ .
- We have  $L^\sigma = L$ ,  $M^\sigma = M$  and  $D^\sigma = D$  for every  $\sigma \in (\mathbb{Z}/2\mathbb{Z})^2$ .

Thus, we have  $(\mathcal{S}^{\Gamma_0})^{(\mathbb{Z}/2\mathbb{Z})^2} = \mathbb{C}[H, L, M, D]$  and  $\mathcal{S}^\Gamma = (\mathcal{S}^{\Gamma_0})^{\mathfrak{S}_3}$ .

Observe that the subring  $\mathbb{C}[L, M] \subset \mathbb{C}[H, L, M, D]$  is  $\mathfrak{S}_3$ -invariant, since

$$\begin{aligned} L^{(1,2)} &= -L, & L^{(1,3)} &= M - L, & L^{(2,3)} &= -M, & L^{(1,2,3)} &= L - M, & L^{(1,3,2)} &= M, \\ M^{(1,2)} &= M - L, & M^{(1,3)} &= -M, & M^{(2,3)} &= -L, & M^{(1,2,3)} &= L, & M^{(1,3,2)} &= L - M. \end{aligned}$$

Note that the polynomials  $L^2 + LM + M^2$  and  $2L^3 + 3L^2M - 3LM^2 - 2M^3$  are  $\mathfrak{S}_3$ -invariant. Moreover, the Hilbert polynomial of the invariant ring  $\mathbb{C}[L, M]^{\mathfrak{S}_3}$  is equal to

$$\frac{1}{6} \left( \frac{1}{(1-t)^2} + \frac{3}{1-t^2} + \frac{2}{1+t+t^2} \right) = \frac{1}{(1-t^2)(1-t^3)}$$

by Molien's formula, where we take the degrees of  $L$  and  $M$  to be 1. This gives

$$\mathbb{C}[L, M]^{\mathfrak{S}_3} = \mathbb{C} [L^2 + LM + M^2, 2L^3 + 3L^2M - 3LM^2 - 2M^3],$$

because  $L^2 + LM + M^2$  and  $2L^3 + 3L^2M - 3LM^2 - 2M^3$  are algebraically independent. This shows that

$$\mathbb{C}[H, L, M, D]^{\mathfrak{S}_3} = \mathbb{C} [H, R, L^2 + LM + M^2, 2L^3 + 3L^2M - 3LM^2 - 2M^3].$$

On the other hand, the equality

$$\mathbb{C} [H, R, L^2 + LM + M^2, 2L^3 + 3L^2M - 3LM^2 - 2M^3] = \mathbb{C}[H, R, S, T]$$

is trivial from the definitions of  $S$  and  $T$ .  $\square$

Let  $\phi: \mathbb{P}(V)^{\text{ss}} \rightarrow \mathbb{P}(V)^{\text{ss}} // \Gamma$  be the GIT quotient morphism. Then  $\mathbb{P}(\mathcal{G}) \subset \mathbb{P}(V)^{\text{ss}}$  by

**Proposition 2.6** ([11]). *The subset  $\Gamma \cdot \mathbb{P}(\mathcal{G}) \subset \mathbb{P}(V)$  consists of all GIT polystable points with respects to the action  $\Gamma \curvearrowright V$ .*

Thus, we can consider the restriction of  $\phi$  to  $\mathbb{P}(\mathcal{G})$ :

$$\phi|_{\mathbb{P}(\mathcal{G})}: \mathbb{P}(\mathcal{G}) = \mathbb{P}_{a,b,c,d}^3 \rightarrow \mathbb{P}(V)^{\text{ss}} // \Gamma \simeq \mathbb{P}(1, 3, 4, 6).$$

In the following, we identify  $\mathbb{P}(V)^{\text{ss}} // \Gamma = \mathbb{P}(1, 3, 4, 6)$ .

**Proposition 2.7.** *The restriction  $\phi|_{\mathbb{P}(\mathcal{G})}$  is the quotient of  $\mathbb{P}(\mathcal{G})$  by the group  $W(\mathbb{F}_4)/\langle -1 \rangle$ , which is the composition  $\phi_3 \circ \phi_2 \circ \phi_1$ , where  $\phi_1: \mathbb{P}^3 \rightarrow \mathbb{P}^3$  is a degree 8 morphism given by*

$$(a : b : c : d) \mapsto (a^2 : b^2 : c^2 : d^2),$$

the morphism  $\phi_2: \mathbb{P}^3 \rightarrow \mathbb{P}(1, 2, 3, 4)$  is a morphism of degree 24 given by

$$(p_1 : p_2 : p_3 : p_4) \mapsto (p_1 + p_2 + p_3 + p_4 : p_1p_2 + p_1p_3 + p_1p_4 + p_2p_3 + p_2p_4 + p_3p_4 : \\ : p_2p_3p_4 + p_1p_3p_4 + p_1p_2p_4 + p_1p_2p_3 : p_1p_2p_3p_4),$$

and  $\phi_3: \mathbb{P}(1, 2, 3, 4) \rightarrow \mathbb{P}(1, 3, 4, 6)$  is a morphism of degree 3 given by

$$(s_1 : s_2 : s_3 : s_4) \mapsto \left( \frac{1}{2}s_1 : \frac{1}{32}(s_1^3 - 4s_1s_2 + 24s_3) : \frac{1}{12}(12s_4 + s_2^2 - 3s_1s_3) : \right. \\ \left. : \frac{1}{432}(27s_1^2s_4 - 72s_2s_4 + 2s_2^3 - 9s_1s_2s_3 + 27s_3^2) \right)$$

where  $s_1, s_2, s_3, s_4$  are coordinates on  $\mathbb{P}(1, 2, 3, 4)$  of weights 1, 2, 3, 4, respectively.

*Proof.* Direct computations.  $\square$

For any non-zero  $f \in \mathcal{G}$  that is the form  $(G_{a,b,c,d})$  for  $(a, b, c, d) \in \mathbb{C}^4$ , set

$$X_{(a:b:c:d)} = \{f = 0\} \subset (\mathbb{P}^1)^4.$$

For convenience, we set  $\mathbb{P} = \mathbb{P}_{a,b,c,d}^3$  and

$$\begin{aligned} \mathbb{P}^{\text{Sing}} &= \{(a, b, c, d) \in \mathbb{P} \mid X_{(a:b:c:d)} \text{ is singular}\}, \\ \mathbb{P}^{\text{Red}} &= \{(a, b, c, d) \in \mathbb{P} \mid X_{(a:b:c:d)} \text{ is reducible}\}, \\ \mathbb{P}^{\text{Curv}} &= \{(a, b, c, d) \in \mathbb{P} \mid \dim(\text{Sing}(X_{(a:b:c:d)})) = 1\}, \\ \mathbb{P}^{6\mathbb{A}_1} &= \{(a, b, c, d) \in \mathbb{P} \mid X_{(a:b:c:d)} \text{ has exactly 6 isolated } \mathbb{A}_1 \text{ singularities}\}, \\ \mathbb{P}^{4\mathbb{A}_1} &= \{(a, b, c, d) \in \mathbb{P} \mid X_{(a:b:c:d)} \text{ has exactly 4 isolated } \mathbb{A}_1 \text{ singularities}\}, \\ \mathbb{P}^{2\mathbb{A}_1} &= \{(a, b, c, d) \in \mathbb{P} \mid X_{(a:b:c:d)} \text{ has exactly 2 isolated } \mathbb{A}_1 \text{ singularities}\}. \end{aligned}$$

**Theorem 2.8.** *The boundary of the moduli space  $\mathbb{P}^{\text{Sing}}$  admits a stratification*

$$\mathbb{P}^{\text{Sing}} = \mathbb{P}^{\text{Red}} \sqcup \mathbb{P}^{\text{Curv}} \sqcup \mathbb{P}^{6\mathbb{A}_1} \sqcup \mathbb{P}^{4\mathbb{A}_1} \sqcup \mathbb{P}^{2\mathbb{A}_1},$$

and the closure of the strata satisfy:

$$\overline{\mathbb{P}^{2\mathbb{A}_1}} = \mathbb{P}^{\text{Sing}}, \quad \overline{\mathbb{P}^{\text{Curv}}} \setminus \mathbb{P}^{\text{Curv}} = \mathbb{P}^{\text{Red}}, \quad \overline{\mathbb{P}^{4\mathbb{A}_1}} \setminus \mathbb{P}^{4\mathbb{A}_1} = \mathbb{P}^{\text{Red}} \sqcup \mathbb{P}^{6\mathbb{A}_1}.$$

Moreover, in the notation of (2.4), the following assertions hold:

(1)  $\phi(\mathbb{P}^{\text{Sing}})$  is the hypersurface

$$\{S^3 - 27T^2 = 0\} \subset \mathbb{P}(1, 3, 4, 6),$$

and the divisor  $X_{(a:b:c:d)}$  is singular if and only if

$$(a^2 - b^2)(a^2 - c^2)(a^2 - d^2)(b^2 - c^2)(b^2 - d^2)(c^2 - d^2) = 0,$$

(2)  $\phi(\mathbb{P}^{\text{Red}}) = (2 : 2 : 0 : 0) \in \mathbb{P}(1, 3, 4, 6)$ , and  $X_{(a:b:c:d)}$  is reducible if and only if

$$a^2 = b^2 = c^2 = d^2 \text{ or } \sigma(b) = \sigma(c) = \sigma(d) = 0 \text{ for some } \sigma \in \mathfrak{S}_4,$$

(3)  $\phi(\mathbb{P}^{6\mathbb{A}_1}) = \left(2 : 0 : \frac{4}{3} : \frac{8}{27}\right)$ , and  $X_{(a:b:c:d)}$  has 6 ordinary double points if and only if

$$\sigma(a) = \sigma(b) = 0 \text{ and } \sigma(c)^2 = \sigma(d)^2 \text{ for some } \sigma \in \mathfrak{S}_4.$$

(4)  $\phi(\overline{\mathbb{P}^{\text{Curv}}}) = \{S = T = 0\}$ , and  $X_{(a:b:c:d)}$  is singular along a curve if and only if

$$\sigma(b)^2 = \sigma(c)^2 = \sigma(d)^2 \neq 0 \text{ for some } \sigma \in \mathfrak{S}_4,$$

(5)  $\phi(\mathbb{P}^{4\mathbb{A}_1})$  is the set of points

$$\left\{ \left( \frac{1}{2}(t+1) : \frac{1}{32}(t+1)(t-1)^2 : \frac{1}{12}t^2 : \frac{1}{216}t^3 \right) \mid t \in \mathbb{C} \setminus \{0, 1\} \right\} \subset \mathbb{P}(1, 3, 4, 6),$$

and  $(a : b : c : d) \in \overline{\mathbb{P}^{4\mathbb{A}_1}}$  if and only if there is  $\sigma \in \mathfrak{S}_4$  such that either  $\sigma(a)^2 = \sigma(b)^2$  and  $\sigma(c)^2 = \sigma(d)^2$ , or  $\sigma(a) = \sigma(b) = 0$ .

*Proof.* Follows from direct computation, see [17]. □

Let us translate Theorem 2.8 into a simpler form. To do this, consider the plane

$$\{a + b = 0\} \subset \mathbb{P}(\mathcal{G}) = \mathbb{P}_{a,b,c,d}^3.$$

If  $(a : b : c : d) \in \{a + b = 0\}$ , then  $X_{(a:b:c:d)}$  is singular. Vice versa, if  $X_{(a:b:c:d)}$  is singular, then the plane  $\{a + b = 0\}$  contains a point in the  $W(\mathbb{F}_4)/\langle -1 \rangle$ -orbit of  $(a : b : c : d)$ . Therefore, in Theorem 2.8, we can assume that  $a + b = 0$  up to the action of  $W(\mathbb{F}_4)/\langle -1 \rangle$ , and we can restate the result in terms of points in the plane  $\{a + b = 0\}$ . Namely, set

$$\mathbb{P}_{a+b=0}^{\text{Red}} = \mathbb{P}^{\text{Red}} \cap \{a + b = 0\},$$

$$\mathbb{P}_{a+b=0}^{\text{Curv}} = \mathbb{P}^{\text{Curv}} \cap \{a + b = 0\},$$

$$\mathbb{P}_{a+b=0}^{6\mathbb{A}_1} = \mathbb{P}^{6\mathbb{A}_1} \cap \{a + b = 0\},$$

$$\mathbb{P}_{a+b=0}^{4\mathbb{A}_1} = \mathbb{P}^{4\mathbb{A}_1} \cap \{a + b = 0\},$$

$$\mathbb{P}_{a+b=0}^{2\mathbb{A}_1} = \mathbb{P}^{2\mathbb{A}_1} \cap \{a + b = 0\}.$$

**Corollary 2.9.** *The plane  $\{a + b = 0\} \subset \mathbb{P}(\mathcal{G}) = \mathbb{P}_{a,b,c,d}^3$  admits the stratification*

$$\{a + b = 0\} = \mathbb{P}_{a+b=0}^{\text{Red}} \sqcup \mathbb{P}_{a+b=0}^{\text{Curv}} \sqcup \mathbb{P}_{a+b=0}^{6\mathbb{A}_1} \sqcup \mathbb{P}_{a+b=0}^{4\mathbb{A}_1} \sqcup \mathbb{P}_{a+b=0}^{2\mathbb{A}_1},$$

which satisfies

$$\overline{\mathbb{P}_{a+b=0}^{2\mathbb{A}_1}} = \{a + b = 0\}, \quad \overline{\mathbb{P}_{a+b=0}^{\text{Curv}}} \setminus \mathbb{P}_{a+b=0}^{\text{Curv}} = \mathbb{P}_{a+b=0}^{\text{Red}}, \quad \overline{\mathbb{P}_{a+b=0}^{4\mathbb{A}_1}} \setminus \mathbb{P}_{a+b=0}^{4\mathbb{A}_1} = \mathbb{P}_{a+b=0}^{\text{Red}} \sqcup \mathbb{P}_{a+b=0}^{6\mathbb{A}_1}.$$

Moreover, under the canonical isomorphism  $\{a + b = 0\} \simeq \mathbb{P}_{a,c,d}^2$ , the following holds:

$$\begin{aligned} \overline{\mathbb{P}_{a+b=0}^{\text{Curv}}} &= \{(a+c)(a-c)(a+d)(a-d) = 0\}, \\ \overline{\mathbb{P}_{a+b=0}^{4\mathbb{A}_1}} &= \{a(c+d)(c-d) = 0\}, \\ \mathbb{P}_{a+b=0}^{\text{Red}} &= \{(1:1:1), (-1:1:1), (1:-1:1), (1:1:-1), (0:1:0), (0:0:1)\}, \\ \mathbb{P}_{a+b=0}^{6\mathbb{A}_1} &= \{(1:0:0), (0:1:1), (0:1:-1)\}. \end{aligned}$$

Now, up to the action of the group  $W(F_4)/\langle -1 \rangle$  on  $\mathbb{P}(\mathcal{G}) = \mathbb{P}_{a,b,c,d}^3$ , we can explicitly describe all singular divisors  $X_{(a:b:c:d)} \subset (\mathbb{P}^1)^4$  as follows:

- the divisor  $X_{(1:-1:1:1)}$  is reducible and is given by

$$(x_1x_4 - y_1y_4)(x_2x_3 - y_2y_3) = 0,$$

so  $X_{(1:-1:1:1)}$  is a union of two divisors of degree  $(1, 0, 0, 1)$  and  $(0, 1, 1, 0)$ ;

- the divisor  $X_{(0:0:1:1)}$  has 6 isolated  $\mathbb{A}_1$  singularities at

$$\begin{aligned} &((1:-1), (1:-1), (1:1), (1:1)), \quad ((1:1), (1:1), (1:1), (1:1)), \\ &((1:1), (1:-1), (1:-1), (1:1)), \quad ((1:-1), (1:-1), (1:1), (1:-1)), \\ &((1:-1), (1:-1), (1:-1), (1:-1)), \quad ((1:1), (1:1), (1:-1), (1:-1)), \end{aligned}$$

and  $X_{(0:0:1:1)}$  is toric — the toric K-polystable Fano 3-fold with reflexible ID 1530 in the Graded Ring Data Base of toric canonical Fano 3-folds [14];

- the divisor  $X_{(1:-1:1:d)}$  with  $d \neq \pm 1$  is singular along the degree  $(1, 1, 1, 1)$  curve

$$\{((u:v), (v:u), (u:v), (v:u) \mid (u:v) \in \mathbb{P}^1\} \subset (\mathbb{P}^1)^4,$$

and  $X_{(1:-1:1:d)}$  has  $\mathbb{A}_1$  singularity along this curve;

- the divisor  $X_{(0:0:1:d)}$  with  $d \neq \pm 1$  and  $d \neq 0$  has 4 isolated  $\mathbb{A}_1$  singularities

$$\begin{aligned} &((1:-1), (1:-1), (1:1), (1:1)), \quad ((1:1), (1:1), (1:1), (1:1)), \\ &((1:-1), (1:-1), (1:-1), (1:-1)), \quad ((1:1), (1:1), (1:-1), (1:-1)), \end{aligned}$$

- general divisor  $X_{(a:-a:c:d)}$  has 2 isolated  $\mathbb{A}_1$  singularities at

$$((1:-1), (1:-1), (1:1), (1:1)), \quad ((1:1), (1:1), (1:1), (1:1)).$$

### 3. DIVISORS OF DEGREE $(2, 2)$ IN $\mathbb{P}(1, 1, 2) \times \mathbb{P}(1, 1, 2)$

In Section 2, we described all GIT polystable degenerations of smooth Fano 3-folds in the deformation family  $\mathcal{N}^{\mathbb{A}_1}$ . We will prove later in Section 4, that all these degenerations are K-polystable provided that they are irreducible. However, one point of the GIT moduli space corresponds to the orbit of the divisor

$$\{(x_1x_2 - y_1y_2)(x_3x_4 - y_3y_4) = 0\} \subset \mathbb{P}_{x_1,y_1}^1 \times \mathbb{P}_{x_2,y_2}^1 \times \mathbb{P}_{x_3,y_3}^1 \times \mathbb{P}_{x_4,y_4}^1,$$



which is a union of divisors of degree  $(1, 1, 0, 0)$  and  $(0, 0, 1, 1)$ . It is not K-polystable, and we have to resolve this issue. This can be done as follows.

In the notations of Section 2, the GIT moduli space is the global quotient

$$\mathbb{P}_{a,b,c,d}^3 / (W(F_4) / \langle -1 \rangle),$$

and the reducible divisor corresponds to the orbit of the point  $(0 : 0 : 0 : 1) \in \mathbb{P}_{a,b,c,d}^3$ . When we approach the point  $(0 : 0 : 0 : 1)$  along a line in  $\mathbb{P}_{a,b,c,d}^3$ , we can reparametrise the corresponding one-dimensional subfamily of divisors  $X_{(a:b:c:d)} \subset (\mathbb{P}^1)^4$  and find their unique K-polystable limit that replaces the reducible one. To find such reparametrisations, we consider the embedding  $X_{(a:b:c:d)} \hookrightarrow \mathbb{P}^3 \times \mathbb{P}^3$  via the map

$$\begin{aligned} & ((x_1 : y_1), (x_2 : y_2), (x_3 : y_3), (x_4 : y_4)) \mapsto \\ \mapsto & ((x_1x_2 - y_1y_2 : 2x_1y_2 : 2x_2y_1 : x_1x_2 + y_1y_2), (x_3x_4 - y_3y_4 : 2x_3y_4 : 2x_4y_3 : x_3x_4 + y_3y_4)). \end{aligned}$$

Then the image of  $X_{(a:b:c:d)} \subset \mathbb{P}_{u_1, u_2, u_3, u_4}^3 \times \mathbb{P}_{v_1, v_2, v_3, v_4}^3$  is given by

$$\begin{cases} u_1^2 + u_2u_3 - u_4^2 = 0, \\ v_1^2 + v_2v_3 - v_4^2 = 0, \\ du_1v_1 - \frac{b}{4}(u_2 + u_3)(v_2 + v_3) - \frac{c}{4}(u_2 - u_3)(v_2 - v_3) - au_4v_4 = 0. \end{cases}$$

Now, if we approach the point  $(0 : 0 : 0 : 1) \in \mathbb{P}_{a,b,c,d}^3$  along the line

$$L_{(a:b:c)} = \{(sa : sb : sc : t) \mid (s : t) \in \mathbb{P}^1\} \subset \mathbb{P}_{a,b,c,d}^3,$$

we obtain the one parameter family

$$\begin{cases} u_1^2 + u_2u_3 - u_4^2 = 0, \\ v_1^2 + v_2v_3 - v_4^2 = 0, \\ u_1v_1 - s \left( \frac{b}{4}(u_2 + u_3)(v_2 + v_3) - \frac{c}{4}(u_2 - u_3)(v_2 - v_3) - au_4v_4 \right) = 0, \end{cases}$$

which degenerates to a reducible 3-fold when  $s$  approaches 0. We can reparametrise it by changing the parameter  $s \mapsto s^2$ , and then changing coordinates as  $u_1 \mapsto su_1$ ,  $v_1 \mapsto sv_1$ , which gives the one-parameter family

$$\begin{cases} s^2u_1^2 + u_2u_3 - u_4^2 = 0, \\ s^2v_1^2 + v_2v_3 - v_4^2 = 0, \\ u_1v_1 - \frac{b}{4}(u_2 + u_3)(v_2 + v_3) - \frac{c}{4}(u_2 - u_3)(v_2 - v_3) - au_4v_4 = 0. \end{cases}$$

When  $s$  approaches 0, the 3-folds in this family degenerate to the singular 3-fold

$$(3.1) \quad \begin{cases} u_2u_3 - u_4^2 = 0, \\ v_2v_3 - v_4^2 = 0, \\ u_1v_1 - \frac{b}{4}(u_2 + u_3)(v_2 + v_3) - \frac{c}{4}(u_2 - u_3)(v_2 - v_3) - au_4v_4 = 0. \end{cases}$$

Note that  $\mathbb{P}(1_{s_1}, 1_{t_1}, 2_{w_1}) \times \mathbb{P}(1_{s_2}, 1_{t_2}, 2_{w_2}) \simeq \{u_2u_3 - u_4^2 = 0, v_2v_3 - v_4^2 = 0\}$  via the map

$$((s_1, t_1, w_1), (s_2, t_2, w_1)) \mapsto ((w_1, s_1^2, t_1^2, s_1t_1), (w_2, s_2^2, t_2^2, s_2t_2)),$$

so (3.1) is isomorphic to the 3-fold in  $\mathbb{P}(1_{s_1}, 1_{t_1}, 2_{w_1}) \times \mathbb{P}(1_{s_2}, 1_{t_2}, 2_{w_2})$  given by  $(\diamond)$ .

For every  $(a : b : c) \in \mathbb{P}^2$ , the Fano 3-fold  $(\diamond)$  has canonical Gorenstein singularities. Moreover, we will see in Section 5 that this 3-fold is K-polystable for every  $(a : b : c) \in \mathbb{P}^2$ .

*Remark 3.2.* A more conceptual and fancy explanation of what we just did is the following. Consider the family of complete intersections in  $\mathbb{P}_{u_1, u_2, u_3, u_4}^3 \times \mathbb{P}_{v_1, v_2, v_3, v_4}^3$  given by

$$\begin{cases} d^2u_1^2 + u_2u_3 - u_4^2 = 0, \\ d^2v_1^2 + v_2v_3 - v_4^2 = 0, \\ d^2u_1v_1 - \frac{b}{4}(u_2 + u_3)(v_2 + v_3) - \frac{c}{4}(u_2 - u_3)(v_2 - v_3) - au_4v_4 = 0 \end{cases}$$

over the base  $T = \text{Spec}(\mathbb{C}[a, b, c, d])$ . This family is invariant under the  $\mathbb{C}^*$ -action with weights  $(-1, 0, 0, 0)$  on  $(u_1, u_2, u_3, u_4)$ , with weights  $(-1, 0, 0, 0)$  on  $(v_1, v_2, v_3, v_4)$ , and with weights  $(0, 0, 0, 1)$  on the coordinates of the base  $(a, b, c, d)$ . The invariant base of the family under this  $\mathbb{C}^*$ -action is  $\text{Spec}(T^{\mathbb{C}^*}) \simeq \mathbb{C}^3$ . Let

$$\begin{aligned} Z^+ &= \mathbb{C}_{a,b,c,d}^4 \setminus \{d = 0\}, \\ Z^- &= \mathbb{C}_{a,b,c,d}^4 \setminus \{a = b = c = 0\}. \end{aligned}$$

Then there are morphisms  $\alpha: Z^+/\mathbb{C}^* \rightarrow \text{Spec}(T^{\mathbb{C}^*})$  and  $\beta: Z^-/\mathbb{C}^* \rightarrow \text{Spec}(T^{\mathbb{C}^*})$  such that

- $\alpha$  is an isomorphism and gives an affine neighborhood of  $(0 : 0 : 0 : 1) \in \mathbb{P}_{a,b,c,d}^3$ ,
- $\beta$  is a blow-up of the point  $(0 : 0 : 0 : 1)$ .

Therefore, blowing up  $\mathbb{B} \rightarrow \mathbb{P}_{a,b,c,d}^3$  at the  $W(F_4)/\langle -1 \rangle$ -orbit of the point  $(0 : 0 : 0 : 1)$ , we obtain a new deformation family

$$\mathcal{X} \rightarrow \mathbb{B}$$

that generically parametrises smooth Fano 3-folds in the family  $\mathbb{N}^4.1$ .

Let  $\mathbb{E}$  be the exceptional divisor of this blow up over  $(0 : 0 : 0 : 1)$ . Then  $\mathbb{E}$  is canonically isomorphic to  $\mathbb{P}_{a,b,c}^2$ , and closed points of  $\mathbb{E}$  parametrise singular divisors of degree  $(2, 2)$  in the product  $\mathbb{P}(1_{s_1}, 1_{t_1}, 2_{w_1}) \times \mathbb{P}(1_{s_2}, 1_{t_2}, 2_{w_2})$  that are given by  $(\diamond)$ .

*Remark 3.3.* If  $\Theta$  is the stabilizer in  $W(F_4)/\langle -1 \rangle$  of  $(0 : 0 : 0 : 1)$ , then

$$\Theta \simeq \mathfrak{S}_4 \times (\mathbb{Z}/2\mathbb{Z})$$

the group  $\Theta$  is isomorphic to the Weyl group  $W(B_3)$ , and it acts on  $\mathbb{E}$  with kernel  $(\mathbb{Z}/2\mathbb{Z})$ , so that its image in  $\text{Aut}(\mathbb{E})$  is isomorphic to  $\mathfrak{S}_4$ . This gives  $\mathbb{E}/\Theta \simeq \mathbb{P}(1, 2, 3)$ .

Now, let  $X_{(a:b:c)}$  be the hypersurface in  $\mathbb{P}(1_{s_1}, 1_{t_1}, 2_{w_1}) \times \mathbb{P}(1_{s_2}, 1_{t_2}, 2_{w_2})$  given by  $(\diamond)$  for some  $(a : b : c) \in \mathbb{P}^2$ , and let  $G$  be the subgroup of  $\text{Aut}(X)$  generated by the involutions

$$\begin{aligned} \tau_1 &: ((s_1 : t_1 : w_1), (s_2 : t_2 : w_2)) \mapsto ((s_2 : t_2 : w_1), (s_1 : t_1 : w_2)), \\ \tau_2 &: ((s_1 : t_1 : w_1), (s_2 : t_2 : w_2)) \mapsto ((t_1 : s_1 : w_1), (t_2 : s_2 : w_2)), \\ \tau_3 &: ((s_1 : t_1 : w_1), (s_2 : t_2 : w_2)) \mapsto ((s_1 : -t_1 : w_1), (s_2 : -t_2 : w_2)), \end{aligned}$$

and by the automorphisms

$$\sigma_\lambda : ((s_1 : t_1 : w_1), (s_2 : t_2 : w_2)) \mapsto \left( (s_1 : t_1 : \lambda w_1), (s_2 : t_2 : \frac{w_2}{\lambda}) \right),$$

for  $\lambda \in \mathbb{C}^*$ . Then  $G \simeq \mathbb{C}^* \rtimes (\mathbb{Z}/2\mathbb{Z})^3$ . Observe that  $X$  is singular along the curves

$$\begin{aligned} C_1 &= \{s_1 = t_1 = w_2 = 0\}, \\ C_2 &= \{s_2 = t_2 = w_1 = 0\}. \end{aligned}$$

To describe all possible singularities of  $X_{(a:b:c)}$ , we set

$$\begin{aligned} \mathbb{E}^{\text{Sing}+} &= \{(a : b : c) \in \mathbb{E} \mid \text{Sing}(X_{(a:b:c)}) \neq C_1 \cup C_2\}, \\ \mathbb{E}^{\text{Curv}+} &= \{(a : b : c) \in \mathbb{E} \mid \text{Sing}(X_{(a:b:c)}) \setminus (C_1 \cup C_2) \text{ is a curve}\}, \\ \mathbb{E}^{4\mathbb{A}_1+} &= \{(a : b : c) \in \mathbb{E} \mid X_{(a:b:c)} \text{ has 4 isolated } \mathbb{A}_1 \text{ singularities outside } C_1 \cup C_2\}, \\ \mathbb{E}^{2\mathbb{A}_1+} &= \{(a : b : c) \in \mathbb{E} \mid X_{(a:b:c)} \text{ has 2 isolated } \mathbb{A}_1 \text{ singularities outside } C_1 \cup C_2\}. \end{aligned}$$

Here and in the following, we identify  $\mathbb{E} = \mathbb{P}_{a,b,c}^2$ .

**Proposition 3.4.** *We have the stratification*

$$\mathbb{E}^{\text{Sing}+} = \mathbb{E}^{\text{Curv}+} \sqcup \mathbb{E}^{4\mathbb{A}_1+} \sqcup \mathbb{E}^{2\mathbb{A}_1+},$$

satisfying that

$$\overline{\mathbb{E}^{2\mathbb{A}_1+}} = \mathbb{E}^{\text{Sing}+}, \quad \mathbb{E}^{\text{Sing}+} \setminus \mathbb{E}^{2\mathbb{A}_1+} = \mathbb{E}^{\text{Curv}+} \sqcup \mathbb{E}^{4\mathbb{A}_1+} \sqcup \mathbb{E}^{2\mathbb{A}_1+}.$$

Moreover, one has

$$\begin{aligned} \mathbb{E}^{\text{Sing}+} &= \{(a^2 - b^2)(a^2 - c^2)(b^2 - c^2) = 0\}, \\ \mathbb{E}^{\text{Curv}+} &= \{(1 : 1 : 1), (1 : 1 : -1), (1 : -1 : -1), (1 : -1 : 1)\}, \\ \mathbb{E}^{4\mathbb{A}_1+} &= \{(1 : 0 : 0), (0 : 1 : 0), (0 : 0 : 1)\}. \end{aligned}$$

Furthermore, the subsets  $\mathbb{E}^{\text{Curv}+}$ ,  $\mathbb{E}^{4\mathbb{A}_1+}$ ,  $\overline{\mathbb{E}^{2\mathbb{A}_1+}}$  are cut out on the divisor  $\mathbb{E}$  by the strict transforms of the subsets  $\mathbb{P}^{\text{Curv}}$ ,  $\overline{\mathbb{P}^{4\mathbb{A}_1}}$ ,  $\overline{\mathbb{P}^{2\mathbb{A}_1}}$  defined in Section 2, respectively.

*Proof.* Direct computations. □

Let us describe  $\text{Sing}(X) \setminus (C_1 \cup C_2)$  explicitly. If  $(a : b : c) \in \mathbb{E}^{\text{Curv}+}$ , then

$$\text{Sing}(X) \setminus (C_1 \cup C_2) = \begin{cases} \{w_1 = w_2 = s_1 s_2 + t_1 t_2 = 0\} & \text{if } (a : b : c) = (1 : 1 : 1), \\ \{w_1 = w_2 = s_1 t_2 + t_1 s_2 = 0\} & \text{if } (a : b : c) = (1 : 1 : -1), \\ \{w_1 = w_2 = s_1 s_2 - t_1 t_2 = 0\} & \text{if } (a : b : c) = (1 : -1 : -1), \\ \{w_1 = w_2 = s_1 t_2 - t_1 s_2 = 0\} & \text{if } (a : b : c) = (1 : -1 : 1), \end{cases}$$

and  $X_{(a:b:c)}$  has  $\mathbb{A}_1$  singularity along this curve. Moreover, if  $(a : b : c) = (1 : 0 : 0)$ , then

$$\begin{aligned} \text{Sing}(X) \setminus (C_1 \cup C_2) &= \left\{ ((1 : 0 : 0), (1 : 0 : 0)), (1 : 0 : 0), (0 : 1 : 0), \right. \\ &\quad \left. ((0 : 1 : 0), (1 : 0 : 0)), ((0 : 1 : 0), (0 : 1 : 0)) \right\}. \end{aligned}$$

If  $(a : b : c) = (0 : 1 : 0)$ , then

$$\text{Sing}(X) \setminus (C_1 \cup C_2) = \left\{ ((i : 1 : 0), (i : 1 : 0)), ((i : 1 : 0), (-i : 1 : 0)), \right. \\ \left. ((-i : 1 : 0), (i : 1 : 0)), ((-i : 1 : 0), (-i : 1 : 0)) \right\}.$$

If  $(a : b : c) = (0 : 0 : 1)$ , then

$$\text{Sing}(X) \setminus (C_1 \cup C_2) = \left\{ ((1 : 1 : 0), (1 : 1 : 0)), ((1 : 1 : 0), (1 : -1 : 0)), \right. \\ \left. ((1 : -1 : 0), (1 : 1 : 0)), ((1 : -1 : 0), (1 : -1 : 0)) \right\}.$$

Note that  $X_{(1:0:0)}$ ,  $X_{(0:1:0)}$ ,  $X_{(0:0:1)}$  are isomorphic to the  $\mathbb{K}$ -polystable toric Fano 3-fold with reflexive ID 610 in the Graded Rings Database [14]. If  $(a : b : c) \in \mathbb{E}^{2\mathbb{A}_1^+}$ , then

$$\text{Sing}(X) \setminus (C_1 \cup C_2) = \begin{cases} ((1 : 0 : 0), (1 : 0 : 0)) \cup ((0 : 1 : 0), (0 : 1 : 0)) & \text{if } b + c = 0, \\ ((1 : 0 : 0), (0 : 1 : 0)) \cup ((0 : 1 : 0), (1 : 0 : 0)) & \text{if } b - c = 0, \\ ((1 : 1 : 0), (1 : -1 : 0)) \cup ((1 : -1 : 0), (1 : 1 : 0)) & \text{if } a - b = 0, \\ ((1 : i : 0), (1 : i : 0)) \cup ((1 : -i : 0), (1 : -i : 0)) & \text{if } a - c = 0, \\ ((1 : i : 0), (1 : -i : 0)) \cup ((1 : -i : 0), (1 : i : 0)) & \text{if } a + c = 0, \\ ((1 : 1 : 0), (1 : 1 : 0)) \cup ((1 : -1 : 0), (1 : -1 : 0)) & \text{if } a + b = 0. \end{cases}$$

In the next section we will show that  $X_{(a:b:c)}$  is  $\mathbb{K}$ -polystable for every  $(a : b : c) \in \mathbb{P}^2$ .

#### 4. $\mathbb{K}$ -POLYSTABLE DEGENERATIONS

Since all smooth elements of family  $\mathbb{N}^{\circ}4.1$  are known to be  $\mathbb{K}$ -polystable [4], we now investigate singular  $\mathbb{K}$ -polystable limits of elements in the family.

**4.1. Case  $(\heartsuit)$ .** Let  $X$  be an divisor in  $\mathbb{P}_{x_1, y_1}^1 \times \mathbb{P}_{x_2, y_2}^1 \times \mathbb{P}_{x_3, y_3}^1 \times \mathbb{P}_{x_4, y_4}^1$  that is given by zeroes of the form  $(G_{a,b,c,d})$  for some  $(a, b, c, d) \neq (0, 0, 0, 0)$ . Suppose that  $X$  is irreducible.

**Theorem 4.1.** *The Fano 3-fold  $X$  is  $\mathbb{K}$ -polystable.*

For each  $i \in \{1, 2, 3, 4\}$ , let  $\text{pr}^i : X \rightarrow (\mathbb{P}^1)^3$  be the projection to all but the  $i^{\text{th}}$  factor. Then each  $\text{pr}^i$  is a birational morphism (if  $X$  is smooth, then each  $\text{pr}^i$  blows up a smooth elliptic curve). Let  $E_i$  be  $\text{pr}^i$ -exceptional surface.

**Lemma 4.2.** *One has  $\text{Sing}(X) = E_1 \cap E_2 \cap E_3 \cap E_4$*

*Proof.* Write the equation of  $X$  as  $x_i f_i + y_i g_i = 0$  for forms  $f_i$  and  $g_i$  in  $H^0((\mathbb{P}^1)^3, \mathcal{O}(1, 1, 1))$ . Then  $E_i = \{f_i = g_i = 0\}$ . Now, apply the Jacobian criterion.  $\square$

Let us use notations of Section 2. Recall that  $\text{Aut}(X)$  contains the subgroup

$$G = \langle \tau_1, \tau_2, \tau_3, \tau_4 \rangle \simeq (\mathbb{Z}/2\mathbb{Z})^4.$$

**Lemma 4.3.** *Suppose that  $X$  has isolated singularities. Then  $X$  is  $\mathbb{K}$ -polystable.*

*Proof.* We suppose that  $X$  is not  $K$ -polystable. Then it follows from [15, 22, 31] that there exists a  $G$ -invariant prime divisor  $\mathbf{F}$  over  $X$  such that  $\beta(\mathbf{F}) \leq 0$ . Let  $Z$  be its center on  $X$ . Then  $Z$  is not a point, because  $G$  does not fix points in  $X$ . Thus, either  $Z$  is an irreducible curve or an irreducible surface. Hence, since  $X$  has isolated singularities by assumption, it follows from Lemma 4.2 that  $Z$  is not contained in  $E_1 \cup E_2 \cup E_3 \cup E_4$ .

Note also that  $Z$  is mapped surjectively to  $\mathbb{P}^1$  via the projection  $X \rightarrow \mathbb{P}^1$  to any factor, because the subgroup  $\langle \tau_3, \tau_4 \rangle \subset G$  does not fix points in  $\mathbb{P}^1$ .

Let  $P$  be a general point in  $Z$ , and let  $S$  be the fiber of the projection  $X \rightarrow \mathbb{P}^1$  to the first factor of  $(\mathbb{P}^1)^4$  that contains  $P$ . Then  $P \notin E_1 \cup E_2 \cup E_3 \cup E_4$ , and  $S$  is a smooth sextic del Pezzo surface. Fix  $u \in \mathbb{R}_{\geq 0}$ . Then  $-K_X - uS$  is pseudo-effective  $\iff u \leq 2$ . For  $u \in [0, 2]$ , let  $P(u)$  be the positive part of the Zariski decomposition of  $-K_X - uS$ , and let  $N(u)$  be its negative part. Note that this decomposition exists on  $X$ : we have

$$P(u) \sim_{\mathbb{R}} \begin{cases} -K_X - uS & \text{for } 0 \leq u \leq 1, \\ -K_X - uS - (u-1)E_1 & \text{for } 1 \leq u \leq 2, \end{cases}$$

and

$$N(u) = \begin{cases} 0 & \text{for } 0 \leq u \leq 1, \\ (u-1)E_1 & \text{for } 1 \leq u \leq 2. \end{cases}$$

Integrating, we get  $S_X(S) = \frac{1}{24} \int_0^2 (P(u)^3) du = \frac{11}{16}$ , so that  $\beta(S) = 1 - S_X(S) = \frac{5}{16} > 0$ .

Thus, since  $\delta(S) = 1$  [2], applying [8, Lemma 26], known as the Nemuro Lemma, we get

$$\delta_P(X) \geq \frac{(-K_X)^3}{3} \frac{\delta(S)}{2(-K_S)^2} = \frac{2}{3}.$$

But the proof of the Nemuro Lemma gives  $\delta_P(X) \geq \frac{16}{15}$ . Indeed, recall from [1, 2] that

$$\delta_P(X) \geq \min \left\{ \frac{1}{S_X(S)}, \inf_{\substack{F/S \\ P \in C_S(F)}} \frac{A_S(F)}{S(W_{\bullet, \bullet}^S; F)} \right\},$$

where the infimum is taken by all prime divisors over  $S$  whose center on  $S$  contains  $P$ , and

$$S(W_{\bullet, \bullet}^S; F) = \frac{3}{(-K_X)^3} \int_0^2 \left( P(u)|_S \right)^2 \text{ord}_F(N(u)) du + \frac{3}{(-K_X)^3} \int_0^2 \int_0^\infty \text{vol}(P(u)|_S - vF) dv du.$$

We claim that  $S(W_{\bullet,\bullet}^S; F) \leq \frac{15}{16}A_S(F)$  for every prime divisor over the surface  $S$  whose center on  $S$  contains  $P$ . Indeed, since  $P \notin E_1$ , we have

$$\begin{aligned} S(W_{\bullet,\bullet}^S; F) &= \frac{3}{(-K_X)^3} \int_0^2 \int_0^\infty \text{vol}(P(u)|_S - vF) dvdu = \\ &= \frac{1}{8} \int_0^1 \int_0^\infty \text{vol}(-K_S - vF) dvdu + \frac{1}{8} \int_1^2 \int_0^\infty \text{vol}((2-u)(-K_S) - vF) dvdu = \\ &= \frac{1}{8} \int_0^1 \int_0^\infty \text{vol}(-K_S - vF) dvdu + \frac{1}{8} \int_1^2 (2-u)^3 \int_0^\infty \text{vol}(-K_S - vF) dvdu. \end{aligned}$$

But it follows from the definition of  $\delta(S)$  that

$$\frac{1}{(-K_S)^2} \int_0^\infty \text{vol}(-K_S - vF) \leq \frac{A_S(F)}{\delta(S)}.$$

Thus, we conclude that

$$\begin{aligned} S(W_{\bullet,\bullet}^S; F) &\leq \frac{1}{8} \int_0^1 \frac{(-K_S)^2}{\delta(S)} A_S(F) du + \frac{1}{8} \int_1^2 (2-u)^3 \frac{(-K_S)^2}{\delta(S)} A_S(F) du = \\ &= \frac{3}{4} A_S(F) + \frac{3}{4} A_S(F) \int_1^2 (2-u)^3 du = \frac{15}{16} A_S(F). \end{aligned}$$

This gives  $\delta_P(X) \geq \frac{16}{15}$  as claimed. On the other hand, we have  $\delta_P(X) \leq 1$ , since  $\beta(\mathbf{F}) \leq 0$ . The obtained contradiction completes the proof of the lemma.  $\square$

Hence, to complete the proof of Theorem 4.1, we may assume that  $\dim(\text{Sing}(X)) = 1$ .

Recall from Section 2 that  $\text{Sing}(X)$  consists of a single curve, and  $X$  has ordinary double singularity along this curve. Let  $\eta: \tilde{X} \rightarrow X$  be the blow-up of the curve  $\text{Sing}(X)$ , let  $F$  the exceptional divisor of the blow-up  $\eta$ , and let  $\tilde{E}_1, \tilde{E}_2, \tilde{E}_3, \tilde{E}_4$  be the proper transforms on the 3-fold  $\tilde{X}$  of the surfaces  $E_1, E_2, E_3, E_4$ , respectively. Then it follows from Appendix A that we have the following  $G$ -equivariant commutative diagram:

$$\begin{array}{ccc} & \tilde{X} & \\ \pi \swarrow & & \searrow \eta \\ \mathbb{P}^3 & \text{---} \text{---} \text{---} \text{---} \chi \text{---} \text{---} \text{---} \text{---} \text{---} & X \end{array}$$

where  $\pi$  is a birational morphism contracting  $\tilde{E}_1, \tilde{E}_2, \tilde{E}_3, \tilde{E}_4$  to skew lines  $L_1, L_2, L_3, L_4$ , respectively, and  $\pi(F)$  is a smooth quadric surface containing those lines. The rational map  $\chi$  is the product of the linear projections  $\mathbb{P}^3 \dashrightarrow \mathbb{P}^1$  from  $L_1, L_2, L_3, L_4$ . Note that

$$\eta^*(-K_X) \sim -K_{\tilde{X}} \sim 2F + \tilde{E}_1 + \tilde{E}_2 + \tilde{E}_3 + \tilde{E}_4.$$

From this diagram, we immediately see that  $\text{Aut}^0(X) \simeq \text{PGL}_2(\mathbb{C})$ , and  $\text{Aut}(X)$  contains a subgroup isomorphic to  $\text{PGL}_2(\mathbb{C}) \times (\mathbb{Z}/2\mathbb{Z})^2$  such that the factor  $(\mathbb{Z}/2\mathbb{Z})^2$  permutes transitively the surfaces  $\tilde{E}_1, \tilde{E}_2, \tilde{E}_3, \tilde{E}_4$ . Then  $F$  is the only  $\text{Aut}(X)$ -invariant prime divisor over  $X$ , so it is enough to show that  $\beta(F) > 0$  to prove that  $X$  is K-polystable [31].

Fix  $u \in \mathbb{R}_{\geq 0}$ ,  $-K_X - uF$  is pseudo-effective if and only if  $u \leq 2$ , since

$$\eta^*(-K_X) - uF \sim_{\mathbb{R}} (2-u)F + \tilde{E}_1 + \tilde{E}_2 + \tilde{E}_3 + \tilde{E}_4.$$

For  $u \in [0, 2]$ , let  $P(u)$  be the positive part of the Zariski decomposition of  $-K_X - uF$ , and let  $N(u)$  be its negative part. Note that this decomposition exists on  $X$ : we have

$$P(u) \sim_{\mathbb{R}} \begin{cases} (2-u)F + \tilde{E}_1 + \tilde{E}_2 + \tilde{E}_3 + \tilde{E}_4 & \text{for } 0 \leq u \leq 1, \\ (2-u)(F + \tilde{E}_1 + \tilde{E}_2 + \tilde{E}_3 + \tilde{E}_4) & \text{for } 1 \leq u \leq 2, \end{cases}$$

and

$$N(u) = \begin{cases} 0 & \text{for } 0 \leq u \leq 1, \\ (u-1)(\tilde{E}_1 + \tilde{E}_2 + \tilde{E}_3 + \tilde{E}_4) & \text{for } 1 \leq u \leq 2. \end{cases}$$

This gives  $S_X(F) = \frac{1}{24} \int_0^2 P(u)^3 du = \frac{1}{24} \int_0^1 8u^3 - 24u^2 + 24u du + \frac{1}{24} \int_1^2 8(2-u)^3 u du = \frac{5}{6}$ , which implies that  $\beta(F) = 1 - S_X(F) = \frac{1}{6} > 0$ . This shows that  $X$  is K-polystable.

4.2. **Case** ( $\diamond$ ). Let  $X$  be a divisor in  $\mathbb{P}(1_{s_1}, 1_{t_1}, 2_{w_1}) \times \mathbb{P}(1_{s_2}, 1_{t_2}, 2_{w_2})$  given by ( $\diamond$ ).

**Theorem 4.4.** *The Fano 3-fold  $X$  is K-polystable.*

Let us use notations of Section 3. Recall that  $\text{Aut}(X)$  has subgroup  $G \simeq \mathbb{C}^* \rtimes (\mathbb{Z}/2\mathbb{Z})^3$ , which is generated by the involutions  $\tau_1, \tau_2, \tau_3$ , and automorphisms  $\sigma_\lambda$ , where  $\lambda \in \mathbb{C}^*$ . Let  $\rho: X \dashrightarrow \mathbb{P}^1 \times \mathbb{P}^1$  be the rational map

$$((s_1 : t_1 : w_1), (s_2 : t_2 : w_2)) \mapsto ((s_1 : t_1), (s_2 : t_2)),$$

let  $\eta: \tilde{X} \rightarrow X$  be the blow-up of the curves  $C_1$  and  $C_2$ , and let  $E_1$  and  $E_2$  be the exceptional surfaces of  $\eta$  that are mapped  $C_1$  and  $C_2$ , respectively. Then  $\eta$  and  $\rho$  are  $G$ -equivariant, and we have the following  $G$ -equivariant commutative diagram:

$$(4.5) \quad \begin{array}{ccc} & \tilde{X} & \\ \eta \swarrow & & \searrow \pi \\ X & \dashrightarrow_{\rho} & \mathbb{P}^1 \times \mathbb{P}^1 \end{array}$$

where  $\pi$  is a (non-standard) conic bundle, of which  $E_1$  and  $E_2$  are sections.

**Lemma 4.6.** *If  $\tilde{X}$  has non-isolated singularities,  $X$  is K-polystable.*

*Proof.* We may assume that  $(a : b : c) = (1 : -1 : 1)$ . Then  $\text{Sing}(X) = C_1 \cup C_2 \cup C'$ , where

$$C' = \{s_1 t_2 - s_2 t_1 = w_1 = w_2 = 0\}.$$

Observe that  $C' = F_1 \cap F_2$ , where

$$F_1 = \{s_1 t_2 - s_2 t_1 = w_1 = 0\},$$

$$F_2 = \{s_1 t_2 - s_2 t_1 = w_2 = 0\}.$$

Let  $\tilde{F}_1$  and  $\tilde{F}_2$  be the strict transforms of  $F_1$  and  $F_2$  and  $\tilde{C}' = \tilde{F}_1 \cap \tilde{F}_2$  their intersection. Under  $\pi$ , the image of  $\tilde{C}'$  is a smooth  $(1, 1)$ -curve and  $\pi^*(\pi(\tilde{C}')) = \tilde{F}_1 + \tilde{F}_2$ .

Let  $\phi: \hat{X} \rightarrow \tilde{X}$  be the blow-up of the curve  $\tilde{C}'$ , let  $E'$  be the  $\phi$ -exceptional surface, and let  $\hat{E}_1, \hat{E}_2, \hat{F}_1, \hat{F}_2$  be the strict transforms on  $\hat{X}$  of the surfaces  $E_1, E_2, F_1, F_2$ , respectively. Then it follows from Appendix A that there is  $G$ -equivariant commutative diagram

$$\begin{array}{ccccc}
 & & \hat{X} & & \\
 & \swarrow \psi & & \searrow \phi & \\
 \bar{X} & & & & \tilde{X} \\
 \swarrow \theta & & \searrow \varpi & \swarrow \pi & \searrow \eta \\
 \mathbb{P}^3 & \overset{\chi}{\dashrightarrow} & \mathbb{P}^1 \times \mathbb{P}^1 & \overset{\rho}{\dashleftarrow} & X
 \end{array}$$

where:  $\psi$  is a birational morphism that contracts  $\hat{F}_1$  and  $\hat{F}_2$ ,  $\theta$  is a birational morphism that contracts  $\psi(\hat{E}_1)$  and  $\psi(\hat{E}_2)$  to two skew lines,  $\chi$  is the product of the linear projections from  $\psi(\hat{E}_1)$  and  $\psi(\hat{E}_2)$ , and  $\varpi$  is a  $\mathbb{P}^1$ -bundle. Then the surface  $\theta \circ \psi(E')$  is a smooth quadric that contains the lines  $\psi(\hat{E}_1)$  and  $\psi(\hat{E}_2)$ . Thus, we have

$$(\eta \circ \phi)^*(-K_X) \sim -K_{\hat{X}} \sim 2E' + \hat{E}_1 + \hat{E}_2 + 2\hat{F}_1 + 2\hat{F}_2.$$

The map  $\psi$  is the blow-up of the curves  $\psi(E') \cap \psi(\hat{E}_1)$  and  $\psi(E') \cap \psi(\hat{E}_2)$ . Then

$$\text{Aut}(X) \cong \text{PGL}_2(\mathbb{C}) \times (\mathbb{C}^* \rtimes (\mathbb{Z}/2\mathbb{Z})),$$

and the surface  $E'$  is the only  $\text{Aut}(X)$ -invariant prime divisor over  $X$ . Hence, it is enough to check that  $\beta(E') > 0$  to show that  $X$  is K-polystable [31].

Fix  $u \in \mathbb{R}_{\geq 0}$ . Then  $-K_X - uE'$  is pseudo-effective if and only if  $u \leq 2$ , since

$$(\eta \circ \phi)^*(-K_X) - uE' \sim_{\mathbb{R}} (2 - u)E' + \hat{E}_1 + \hat{E}_2 + 2\hat{F}_1 + 2\hat{F}_2.$$

For  $u \in [0, 2]$ , let  $P(u)$  be the positive part of the Zariski decomposition of  $-K_X - uE'$ , and let  $N(u)$  be its negative part. Note that this decomposition exists on  $X$ : we have

$$P(u) \sim_{\mathbb{R}} \begin{cases} (2 - u)E' + \hat{E}_1 + \hat{E}_2 + 2\hat{F}_1 + 2\hat{F}_2 & \text{for } 0 \leq u \leq 1, \\ (2 - u)(E' + \hat{E}_1 + \hat{E}_2 + 2\hat{F}_1 + 2\hat{F}_2) & \text{for } 1 \leq u \leq 2, \end{cases}$$

and

$$N(u) = \begin{cases} 0 & \text{for } 0 \leq u \leq 1, \\ (u - 1)(\hat{E}_1 + \hat{E}_2 + 2\hat{F}_1 + 2\hat{F}_2) & \text{for } 1 \leq u \leq 2. \end{cases}$$

This gives  $S_X(E') = \frac{1}{24} \int_0^2 (P(u))^3 du = \frac{1}{24} \int_0^1 8u^3 - 24u^2 + 24du + \frac{1}{24} \int_1^2 8(2 - u)^3 u du = \frac{5}{6}$ , which implies that  $\beta(E') = 1 - S_X(E') = \frac{1}{6} > 0$ . This shows that  $X$  is K-polystable.  $\square$

Hence, to prove Theorem 4.4, we may assume that  $\tilde{X}$  has isolated singularities:

$$(a : b : c) \neq (1 : \pm 1 : \pm 1).$$

Then one of the following cases holds:

- $(a : b : c) \notin \mathbb{E}^{\text{Sing}^+}$ , and  $\tilde{X}$  is smooth;



- $(a : b : c) \in \mathbb{E}^{2A_1+}$ , and  $\tilde{X}$  has 2 ordinary double points;
- $(a : b : c) \in \mathbb{E}^{4A_1+}$ , and  $X$  is toric.

If  $X$  is toric, it is K-polystable (see Section 3). Hence, we may assume that  $\tilde{X}$  is smooth, or  $\tilde{X}$  has 2 ordinary double points. Set

$$F_1 = \left\{ as_1t_1s_2t_2 + \frac{b+c}{4}(s_1^2s_2^2 + t_1^2t_2^2) + \frac{b-c}{4}(s_1^2t_2^2 + t_1^2s_2^2) = w_1 = 0 \right\}$$

$$F_2 = \left\{ as_1t_1s_2t_2 + \frac{b+c}{4}(s_1^2s_2^2 + t_1^2t_2^2) + \frac{b-c}{4}(s_1^2t_2^2 + t_1^2s_2^2) = w_2 = 0 \right\}.$$

Denote by  $\tilde{F}_1$  and  $\tilde{F}_2$  the strict transforms of  $F_1$  and  $F_2$  on  $\tilde{X}$ , respectively. Then

$$\eta^*(-K_X) \sim -K_{\tilde{X}} \sim \tilde{F}_1 + \tilde{F}_2 + E_1 + E_2.$$

If  $\tilde{X}$  is smooth, then  $F_1$  and  $F_2$  are irreducible. If  $\tilde{X}$  has 2 ordinary double points, then these surfaces have two components.

**Lemma 4.7.** *Let  $S$  be a  $G$ -invariant irreducible surface in  $X$ . Then*

$$S \sim_{\mathbb{Q}} \frac{n}{2}(-K_X)$$

for some  $n \in \mathbb{N}$ .

*Proof.* Left to the reader.  $\square$

**Lemma 4.8.** *If  $\tilde{X}$  is smooth, then  $F_1 \cap F_2$  is the only  $G$ -invariant irreducible curve in  $X$ . Similarly, if  $\tilde{X}$  is singular, then  $X$  does not contain  $G$ -invariant irreducible curves.*

*Proof.* If  $Z$  is a  $G$ -invariant irreducible curve in  $X$ , it is pointwise fixed by the  $\mathbb{C}^*$ -action, because  $G$  does not fix points in  $\mathbb{P}^1 \times \mathbb{P}^1$ . Then  $Z = F_1 \cap F_2$ , so  $\tilde{X}$  is smooth, since otherwise the intersection  $F_1 \cap F_2$  would be reducible.  $\square$

Now, we are ready to prove that  $X$  is K-polystable. Suppose that  $X$  is not K-polystable. Then, by [15, 22, 31], there exists a  $G$ -invariant prime divisor  $\mathbf{F}$  over  $X$  such that  $\beta(\mathbf{F}) \leq 0$ . Let  $Z$  be the center of this divisor on  $X$ . Then  $Z$  is not a point, since  $G$  does not fix points. Similarly, we see that  $Z$  is not a surface by Lemma 4.7. Thus, we see that  $Z$  is a curve. Then  $\tilde{X}$  is smooth, and  $Z = F_1 \cap F_2$  by Lemma 4.8.

Let  $\tilde{Z}$  be the strict transform on  $\tilde{X}$  of the curve  $Z$ , let  $\xi: \hat{X} \rightarrow \tilde{X}$  be the ordinary blow-up along  $\tilde{Z}$ , and let  $E$  be the exceptional divisor of  $\xi$ . Then the  $G$ -action lifts to  $\hat{X}$ , and  $E$  does not contain  $G$ -invariant irreducible curves. Hence,  $E$  is the only  $G$ -invariant prime divisor over  $X$ . So, it follows from [31] that  $\beta(E) \leq 0$ . Let us compute  $\beta(E)$ .

Fix  $u \in \mathbb{R}_{\geq 0}$ . If  $u \in [0, 1]$ , then  $\xi^*(-K_X) - uE$  is nef. If  $u \in [1, 2]$ , then the positive part of the Zariski decomposition of  $\xi^*(-K_X) - uE$  is

$$(u-1) \left( \hat{E}_1 + \hat{E}_2 + \hat{F}_1 + \hat{F}_2 \right),$$

where  $\hat{E}_1, \hat{E}_2, \hat{F}_1, \hat{F}_2$  are the strict transforms of the surfaces  $E_1, E_2, F_1, F_2$ , respectively. This gives

$$\text{vol}(\xi^*(-K_X) - uE) = \begin{cases} 8(3 - 3u^2 + u^3) & \text{if } u \in [0, 1], \\ 8(2 - u)^3 & \text{if } u \in [1, 2]. \end{cases}$$

Thus, integrating, we get  $S_X(E) = \frac{5}{6} < A_X(E) = 2$ , so  $\beta(E) > 0$  — a contradiction.

This finishes the proof of Theorem 4.4.

## 5. K-MODULI SPACE

To describe the scheme-structure of the K-moduli space, we need the following result:

**Proposition 5.1.** *Let  $X$  be a complete intersection in  $\mathbb{P}^3 \times \mathbb{P}^3$  of three divisors that have degrees  $(2, 0)$ ,  $(0, 2)$  and  $(1, 1)$ . Suppose that  $X$  has canonical Gorenstein singularities. Then the deformations of  $X$  are unobstructed.*

*Proof.* The proof is almost same as the proof of [6, Proposition 2.9]. Namely, it follows from [28, Theorem 2.3.2 and 2.4.1], [28, Corollary 2.4.2], [26, Propositions 2.4 and 2.6] that it is enough to show that  $\text{Ext}_{\mathcal{O}_X}^2(\Omega_X^1, \mathcal{O}_X) = 0$ . As in [26, §1.2], we know that

$$\text{Ext}_{\mathcal{O}_X}^2(\Omega_X^1, \mathcal{O}_X) \simeq H^1(X, \Omega_X^1 \otimes \omega_X)^\vee.$$

From the exact sequence

$$0 \rightarrow \mathcal{N}_{X/\mathbb{P}^3 \times \mathbb{P}^3}^\vee \rightarrow \Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1|_X \rightarrow \Omega_X^1 \rightarrow 0,$$

we have

$$H^1(X, \Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1|_X \otimes \omega_X) \simeq H^1(X, \Omega_X^1 \otimes \omega_X).$$

Indeed, for  $i \in \{1, 2\}$ , we have

$$\begin{aligned} H^i\left(X, \mathcal{N}_{X/\mathbb{P}^3 \times \mathbb{P}^3}^\vee \otimes \omega_X\right) &\simeq \\ &\simeq H^i(X, \mathcal{O}_X(-2, -2)) \oplus H^i(X, \mathcal{O}_X(-3, -1)) \oplus H^i(X, \mathcal{O}_X(-1, -3)) = 0 \end{aligned}$$

by the Kodaira-type vanishing theorem, where  $\mathcal{O}_X(a, b)$  is the restriction to  $X$  of the line bundle on  $\mathbb{P}^3 \times \mathbb{P}^3$  of degree  $(a, b)$ .

Let  $Q_1$  and  $Q_2$  be the divisors in  $\mathbb{P}^3 \times \mathbb{P}^3$  containing  $X$  that have degree  $(2, 0)$  and  $(0, 2)$ , respectively. Set  $V = Q_1 \cap Q_2$ . Then it follows from the exact sequence of sheaves

$$0 \rightarrow \mathcal{O}_V(-1, -1) \rightarrow \mathcal{O}_V \rightarrow \mathcal{O}_X \rightarrow 0$$

that we have an exact sequence of cohomology groups

$$\begin{aligned} H^1(V, (\Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-1, -1))|_V) &\rightarrow \\ &\rightarrow H^1(X, \Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1|_X \otimes \omega_X) \rightarrow \\ &\rightarrow H^2(V, (\Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-2, -2))|_V). \end{aligned}$$

Now, using the exact sequences

$$0 \rightarrow \mathcal{O}_{Q_1}(0, -2) \rightarrow \mathcal{O}_{Q_1} \rightarrow \mathcal{O}_V \rightarrow 0,$$

and

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-2, 0) \rightarrow \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3} \rightarrow \mathcal{O}_{Q_1} \rightarrow 0,$$

we get the exact sequences

$$\begin{aligned} H^i(Q_1, (\Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-i, -i))|_{Q_1}) &\rightarrow \\ &\rightarrow H^i(V, (\Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-i, -i))|_V) \rightarrow \\ &\rightarrow H^{i+1}(Q_1, (\Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-i, -i-2))|_{Q_1}), \end{aligned}$$

$$\begin{aligned}
H^i(\mathbb{P}^3 \times \mathbb{P}^3, \Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-i, -i)) &\rightarrow \\
&\rightarrow H^i(Q \times \mathbb{P}^3, (\Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-i, -i))|_{Q_1}) \rightarrow \\
&\rightarrow H^{i+1}(\mathbb{P}^3 \times \mathbb{P}^3, \Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-i-2, -i)),
\end{aligned}$$

and

$$\begin{aligned}
H^{i+1}(\mathbb{P}^3 \times \mathbb{P}^3, \Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-i, -i-2)) &\rightarrow \\
&\rightarrow H^{i+1}(Q_1, (\Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-i, -i-2))|_{Q_1}) \rightarrow \\
&\rightarrow H^{i+2}(\mathbb{P}^3 \times \mathbb{P}^3, \Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-i-2, -i-2))
\end{aligned}$$

for  $i \in \{1, 2\}$ . Thus, it is enough to show that

$$\begin{aligned}
H^i(\mathbb{P}^3 \times \mathbb{P}^3, \Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-i, -i)) &= 0, \\
H^{j+1}(\mathbb{P}^3 \times \mathbb{P}^3, \Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \otimes \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-j, -j-2)) &= 0
\end{aligned}$$

for  $i = 1, 2, 3, 4$  and  $j = 1, 2$ , which follows from the Euler exact sequence

$$0 \rightarrow \Omega_{\mathbb{P}^3 \times \mathbb{P}^3}^1 \rightarrow \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(-1, 0)^{\oplus 4} \oplus \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}(0, -1)^{\oplus 4} \rightarrow \mathcal{O}_{\mathbb{P}^3 \times \mathbb{P}^3}^{\oplus 2} \rightarrow 0.$$

This completes the proof of the proposition.  $\square$

Now, let  $\mathbf{M}$  be the irreducible component of the K-moduli space  $M_{3,24}^{\text{Kps}}$  whose points parametrise smooth Fano 3-folds in the family  $\mathfrak{N}^{\circ}4.1$  and their K-polystable degenerations. Recall from Section 2 that the GIT moduli space of divisors of degree  $(1, 1, 1, 1)$  in  $(\mathbb{P}^1)^4$  for the natural action of the group  $\Gamma = \text{SL}_4^4(\mathbb{C}) \rtimes \mathfrak{S}_4$  is isomorphic to

$$\mathbb{P}_{a,b,c,d}^3 / (W(\text{F}_4) / \langle -1 \rangle) \simeq \mathbb{P}(1, 3, 4, 6),$$

where  $\mathbb{P}_{a,b,c,d}^3$  is identified with the projectivisation of the vector space of forms  $(G_{a,b,c,d})$ , and the action of the finite group  $W(\text{F}_4) / \langle -1 \rangle \simeq (\mathfrak{A}_4 \times \mathfrak{A}_4) \rtimes (\mathbb{Z}/2\mathbb{Z})^2$  together with the corresponding quotient map  $\mathbb{P}_{a,b,c,d}^3 \rightarrow \mathbb{P}(1, 3, 4, 6)$  are explicitly described in Section 2. Recall that the orbit of the point  $(0 : 0 : 0 : 1) \in \mathbb{P}_{a,b,c,d}$  parametrise reducible divisors.

**Theorem 5.2.** *The following assertions hold:*

- there exists  $W(\text{F}_4) / \langle -1 \rangle$ -equivariant commutative diagram

$$\begin{array}{ccc}
\mathbb{B} & \longrightarrow & \mathbf{M} \\
\downarrow & & \downarrow \\
\mathbb{P}_{a,b,c,d}^3 & \longrightarrow & \mathbb{P}(1, 3, 4, 6)
\end{array}$$

where  $\mathbb{B} \rightarrow \mathbb{P}_{a,b,c,d}^3$  is the blow up of the  $W(\text{F}_4) / \langle -1 \rangle$ -orbit of the point  $(0 : 0 : 0 : 1)$ , both horizontal arrows are the quotients by the action of the group  $W(\text{F}_4) / \langle -1 \rangle$ , and  $\mathbf{M} \rightarrow \mathbb{P}(1, 3, 4, 6)$  is a weighted blow up of a smooth point with weights  $(1, 2, 3)$ ,

- $\mathbf{M}$  is a connected component of  $M_{3,24}^{\text{Kps}}$ .

*Proof.* In Section 3, we have constructed a flat deformation family  $\mathcal{X} \rightarrow \mathbb{B}$  of K-polystable 3-folds of the forms  $(\heartsuit)$  or  $(\diamondsuit)$ . Thus, we can canonically get a finite morphism

$$\mathbb{B} \rightarrow M_{3,24}^{\text{Kps}}.$$

Since the rational map  $\mathbb{P}_{a,b,c,d}^3 \dashrightarrow M_{3,24}^{\text{Kps}}$  and the blow-up  $\mathbb{B} \rightarrow \mathbb{P}_{a,b,c,d}^3$  are both equivariant for the  $W(F_4)/\langle -1 \rangle$ -action, the composition  $\mathbb{B} \rightarrow M_{3,24}^{\text{Kps}}$  is also  $W(F_4)/\langle -1 \rangle$ -equivariant. By the universal property for the quotient of finite groups, we get the morphism

$$\mathbb{B}/(W(F_4)/\langle -1 \rangle) \rightarrow M_{3,24}^{\text{Kps}}.$$

This morphism is finite, and it is birational onto its image. Moreover, by Proposition 5.1, every point in the image is normal, so the morphism is an isomorphism onto a connected component  $\mathbf{M}$  of  $M_{3,24}^{\text{Kps}}$  by Zariski's main theorem.  $\square$

Let us conclude this section with a more detailed structure of the moduli space  $\mathbf{M}$ .

**Proposition 5.3.** *The 3-fold  $\mathbf{M}$  is isomorphic to the weighted blow-up of  $\mathbb{P}(1_H, 3_R, 4_S, 6_T)$  at the point  $(2 : 2 : 0 : 0)$  with weights  $(1, 2, 3)$  at the regular sequence of parameters*

$$\mathbf{r} = \frac{4R - H^3}{H^3}, \quad \mathbf{s} = \frac{S}{H^4}, \quad \mathbf{t} = \frac{T}{H^6},$$

and  $\mathbf{M}$  is isomorphic to the toric 3-fold  $X_\Sigma$ , where  $\Sigma$  is the 3-dimensional fan whose one-dimensional cones have primitive generators

$$\mathbf{v}_0 = (1, 2, 3), \quad \mathbf{v}_1 = (1, 0, 0), \quad \mathbf{v}_2 = (0, 1, 0), \quad \mathbf{v}_3 = (0, 0, 1), \quad \mathbf{v}_4 = (-3, -4, -6),$$

and whose three-dimensional cones are

$$\begin{aligned} &\text{Cone}\{\mathbf{v}_0, \mathbf{v}_1, \mathbf{v}_2\}, && \text{Cone}\{\mathbf{v}_0, \mathbf{v}_1, \mathbf{v}_3\}, \\ &\text{Cone}\{\mathbf{v}_0, \mathbf{v}_2, \mathbf{v}_3\}, && \text{Cone}\{\mathbf{v}_4, \mathbf{v}_1, \mathbf{v}_2\}, \\ &\text{Cone}\{\mathbf{v}_4, \mathbf{v}_1, \mathbf{v}_3\}, && \text{Cone}\{\mathbf{v}_4, \mathbf{v}_2, \mathbf{v}_3\}. \end{aligned}$$

*Proof.* Let  $\mathbb{E}$  be the exceptional divisor of the blow-up  $\mathbb{B} \rightarrow \mathbb{P}_{a,b,c,d}^3$ , and let  $\mathcal{I}$  be the ideal sheaf on  $\mathbb{P}(1_H, 3_R, 4_S, 6_T)$  with co-support  $\{(2 : 2 : 0 : 0)\}$  defined locally by

$$\mathcal{I} = (\mathbf{r}^6, \mathbf{r}^4\mathbf{s}, \mathbf{r}^3\mathbf{t}, \mathbf{r}^2\mathbf{s}^2, \mathbf{rst}, \mathbf{s}^3, \mathbf{t}^2).$$

Then  $X_\Sigma$  is the blow-up of  $\mathbb{P}(1_H, 3_R, 4_S, 6_T)$  along  $\mathcal{I}$ . It follows from Proposition 2.7 that

$$\eta^*(\mathcal{I}) \cdot \mathcal{O}_{\mathbb{B}} = \mathcal{O}_{\mathbb{B}}(-12\mathbb{E}),$$

where  $\eta$  is the composition  $\mathbb{B} \rightarrow \mathbb{P}_{a,b,c,d}^3 \rightarrow \mathbb{P}(1_H, 3_R, 4_S, 6_T)$ . This gives  $\mathbf{M} \simeq X_\Sigma$ , because the morphism  $\eta$  naturally factors through  $X_\Sigma$ .  $\square$

## 6. K-SEMISTABLE LIMITS

In Section 5, we have described all K-polystable Fano 3-folds that are limits of smooth divisors in  $(\mathbb{P}^1)^4$  of degree  $(1, 1, 1, 1)$ . Now, we prove the following result:

**Theorem 6.1.** *Let  $X$  be a K-polystable Fano 3-fold admitting a  $\mathbb{Q}$ -Gorenstein smoothing to a divisor in  $(\mathbb{P}^1)^4$  of degree  $(1, 1, 1, 1)$ . Then  $X$  is isomorphic to an irreducible complete intersection of three divisors in  $\mathbb{P}^3 \times \mathbb{P}^3$  of degree  $(2, 0)$ ,  $(0, 2)$ ,  $(1, 1)$ .*

*Proof.* Let  $\pi: \mathcal{X} \rightarrow \mathbb{A}_t^1$  be the K-polystable degeneration of  $X$ , and let  $X_t$  be the fiber of the morphism  $\pi$  over the point  $t \in \mathbb{A}_t^1$ . Then

$$X_t \simeq X$$

for any  $t \neq 0$ , and  $X_0$  is one of K-polystable Fano 3-folds described in Main Theorem. Since  $X_0$  has canonical Gorenstein singularities, so is  $\mathcal{X}$ . Therefore, the 3-fold  $X_t$  also has canonical Gorenstein singularities for every  $t \in \mathbb{A}^1$ .

On the other hand, we know from Main Theorem that  $\text{Pic}(X_0)$  contains two Cartier divisors  $H_1$  and  $H_2$  such that  $-K_X \sim H_1 + H_2$ ,  $H_1^3 = H_2^3 = 0$ ,  $H_1^2 \cdot H_2 = H_1 \cdot H_2^2 = 4$ , both linear system  $|H_1|$  and  $|H_2|$  are base point free, and give two morphisms  $X_0 \rightarrow \mathbb{P}^3$ . Moreover, arguing as in the proof of [30, Lemma 2.17], we conclude that the restriction homomorphism  $\text{Pic}(\mathcal{X}) \rightarrow \text{Pic}(X_0)$  is an isomorphism. Let  $H_1^\mathcal{X}$  and  $H_2^\mathcal{X}$  be Cartier divisors on  $\mathcal{X}$  such that  $H_1^\mathcal{X}|_{X_0} = H_1$  and  $H_2^\mathcal{X}|_{X_0} = H_2$ , let  $H_{1,t} = H_1^\mathcal{X}|_{X_t}$  and  $H_{2,t} = H_2^\mathcal{X}|_{X_t}$ . Then

$$h^i(X_t, H_{1,t}) = h^i(X_t, H_{2,t}) = 0$$

for every  $i \geq 1$  by the Kawamata–Viehweg vanishing theorem. Then

$$h^0(X_t, H_{1,t}) = h^0(X_t, H_{2,t}) = 4$$

by the Riemann–Roch theorem. This implies that  $\pi_*(H_i^\mathcal{X})$  is locally free sheaf of rank 4. Again by the Kawamata–Viehweg vanishing theorem, the restriction

$$\pi_*(H_i^\mathcal{X}) \rightarrow (\pi_*(H_i^\mathcal{X}))|_{X_t}$$

is surjective for any  $t \in \mathbb{A}^1$ , so the evaluation homomorphism  $\pi^*\pi_*(H_i^\mathcal{X}) \rightarrow H_i^\mathcal{X}$  is surjective. Thus, for  $i \in \{1, 2\}$ , the divisor  $H_i^\mathcal{X}$  gives a morphism

$$\phi_i: \mathcal{X} \rightarrow \mathbb{P}_{\mathbb{A}^1}(\pi_*(H_i^\mathcal{X})) \cong \mathbb{P}^3 \times \mathbb{A}^1,$$

and its restriction  $\phi_{i,t}: X_t \rightarrow \mathbb{P}^3$  is given by the complete linear series  $|H_{i,t}|$ .

For each  $i \in \{1, 2\}$  and every  $t \in \mathbb{A}^1$ , let  $Q_{i,t} \subset \mathbb{P}^3$  be the image of the morphism  $\phi_{i,t}$ . Since  $(H_{1,t}^3) = (H_{2,t}^3) = 0$  and  $-K_{X_t} \cdot H_{1,t}^2 = -K_{X_t} \cdot H_{2,t}^2 = 4$ , we see that  $Q_{i,t}$  is a surface, and the degree of  $Q_{i,t}$  is either 2 or 4. If the degree is 4, then

$$-K_{X_t} \cdot \ell = 1$$

for a general fiber  $\ell$  of  $\phi_{i,t}: X_t \rightarrow Q_{i,t}$ . But the Stein factorization of  $\phi_{i,t}$  gives a conic bundle, so  $-K_{X_t} \cdot \ell$  is divisible by 2. Thus, we see that  $Q_{i,t}$  is a normal quadric surface.

Let  $Q_i^\mathcal{X} \subset \mathbb{P}^3 \times \mathbb{A}^1$  be the scheme-theoretic image of  $\phi_i$ . Since  $Q_i^\mathcal{X}$  is an irreducible and reduced Cartier divisor on  $\mathbb{P}^3 \times \mathbb{A}^1$ , it is Cohen–Macaulay. Therefore, the scheme-theoretic fiber of  $Q_i^\mathcal{X} \rightarrow \mathbb{A}^1$  over  $t \in \mathbb{A}^1$  is the surface  $Q_{i,t}$ , because this is true for a general  $t \in \mathbb{A}^1$ .

Let us consider the product morphism

$$\phi = \phi_1 \times \phi_2: \mathcal{X} \rightarrow \mathbb{P}^3 \times \mathbb{P}^3 \times \mathbb{A}^1$$

over  $\mathbb{A}^1$ . Then  $\phi$  factors through  $\phi: \mathcal{X} \rightarrow Q_1^\mathcal{X} \times_{\mathbb{A}^1} Q_2^\mathcal{X}$ , and  $\phi$  is finite, since

$$H_1^\mathcal{X} + H_2^\mathcal{X} \sim_{\mathbb{A}^1} -K_{\mathcal{X}/\mathbb{A}^1}.$$

Moreover, the restriction of  $\phi$  at  $0 \in \mathbb{A}^1$  is a closed embedding, so  $\phi$  is a closed embedding near  $X_0$  (cf. [23, §5.1, Exercise 1.25]). Then  $\phi: \mathcal{X} \hookrightarrow Q_1^\mathcal{X} \times_{\mathbb{A}^1} Q_2^\mathcal{X}$  is a closed embedding.

Let us consider the reflexive sheaf  $\mathcal{O}_{Q_1^\mathcal{X} \times_{\mathbb{A}^1} Q_2^\mathcal{X}}(\phi(\mathcal{X}))$  on  $Q_1^\mathcal{X} \times_{\mathbb{A}^1} Q_2^\mathcal{X}$ . Note that

$$\mathcal{O}_{Q_1^\mathcal{X} \times_{\mathbb{A}^1} Q_2^\mathcal{X}}(\phi(\mathcal{X}))|_{Q_{1,0} \times Q_{2,0}} \simeq \mathcal{O}_{Q_{1,0} \times Q_{2,0}}(X_0)$$

by [21, Lemma 12.1.8]. Since  $\mathcal{O}_{Q_{1,0} \times Q_{2,0}}(X_0)$  is locally free, so is  $\mathcal{O}_{Q_1^\mathcal{X} \times_{\mathbb{A}^1} Q_2^\mathcal{X}}(\phi(\mathcal{X}))$  around the fiber over  $0 \in \mathbb{A}^1$ . Therefore, for any  $t \in \mathbb{A}^1$ , the image of the embedding

$$X_t \hookrightarrow Q_{1,t} \times Q_{2,t}$$

is a Cartier divisor. Thus, since  $-K_{X_t} \sim H_{1,t} + H_{2,t}$ , we have  $X_t \in |\mathcal{O}_{Q_{1,t} \times Q_{2,t}}(1, 1)|$ .  $\square$

Now, we set

$$\mathbb{H} = \mathbb{P}\left(H^0(\mathbb{P}^3 \times \mathbb{P}^3, \mathcal{O}(2, 0))\right) \times \mathbb{P}\left(H^0(\mathbb{P}^3 \times \mathbb{P}^3, \mathcal{O}(0, 2))\right) \times \mathbb{P}\left(H^0(\mathbb{P}^3 \times \mathbb{P}^3, \mathcal{O}(1, 1))\right),$$

Then  $\mathbb{H} \simeq \mathbb{P}^9 \times \mathbb{P}^9 \times \mathbb{P}^{15}$  is the space parametrising complete intersections in  $\mathbb{P}^3 \times \mathbb{P}^3$  of three divisors of degree  $(2, 0)$ ,  $(0, 2)$ ,  $(1, 1)$ . Let  $\mathbb{H}^{\text{Kss}} \subset \mathbb{H}$  be the locus whose closed points parametrise K-semistable complete intersections. Then

$$\text{codim}(\mathbb{H} \setminus \mathbb{H}^{\text{Kss}}) \geq 2.$$

Let  $\lambda_{\text{CM}}$  be the CM line bundle on  $\mathbb{H}^{\text{Kss}}$  of the corresponding universal family of complete intersections, and let  $\mathcal{X}^{\mathbb{H}} \rightarrow \mathbb{H}$  be the universal family of complete intersections. Then

$$\begin{aligned} ((-K_{\mathcal{X}^{\mathbb{H}}/\mathbb{H}})^4 \cdot \mathcal{O}_{\mathbb{H}}(1, 0, 0)|_{\mathcal{X}^{\mathbb{H}}}^8 \cdot \mathcal{O}_{\mathbb{H}}(0, 1, 0)|_{\mathcal{X}^{\mathbb{H}}}^9 \cdot \mathcal{O}_{\mathbb{H}}(0, 0, 1)|_{\mathcal{X}^{\mathbb{H}}}^{15}) &= -76, \\ ((-K_{\mathcal{X}^{\mathbb{H}}/\mathbb{H}})^4 \cdot \mathcal{O}_{\mathbb{H}}(1, 0, 0)|_{\mathcal{X}^{\mathbb{H}}}^9 \cdot \mathcal{O}_{\mathbb{H}}(0, 1, 0)|_{\mathcal{X}^{\mathbb{H}}}^8 \cdot \mathcal{O}_{\mathbb{H}}(0, 0, 1)|_{\mathcal{X}^{\mathbb{H}}}^{15}) &= -76, \\ ((-K_{\mathcal{X}^{\mathbb{H}}/\mathbb{H}})^4 \cdot \mathcal{O}_{\mathbb{H}}(1, 0, 0)|_{\mathcal{X}^{\mathbb{H}}}^9 \cdot \mathcal{O}_{\mathbb{H}}(0, 1, 0)|_{\mathcal{X}^{\mathbb{H}}}^9 \cdot \mathcal{O}_{\mathbb{H}}(0, 0, 1)|_{\mathcal{X}^{\mathbb{H}}}^{14}) &= -72. \end{aligned}$$

This implies that  $\lambda_{\text{CM}}$  is the restriction of the line bundle  $\mathcal{O}_{\mathbb{H}}(76, 76, 72)$ .

#### APPENDIX A. FOUR LINES IN $\mathbb{P}^3$

Recall from Section 2 that  $\mathbb{P}^{\text{Sing}}$  is the divisor in  $\mathbb{P}_{a,b,c,d}^3$  whose closed points parametrise singular divisors in  $\mathbb{P}_{x_1, y_1}^1 \times \mathbb{P}_{x_2, y_2}^1 \times \mathbb{P}_{x_3, y_3}^1 \times \mathbb{P}_{x_4, y_4}^1$  that are given by

$$\begin{aligned} \text{(A.1)} \quad \frac{a+d}{2}(x_1 x_2 x_3 x_4 + y_1 y_2 y_3 y_4) + \frac{a-d}{2}(x_1 x_2 y_3 y_4 + y_1 y_2 x_3 x_4) + \\ + \frac{b+c}{2}(x_1 y_2 x_3 y_4 + y_1 x_2 y_3 x_4) + \frac{b-c}{2}(x_1 y_2 y_3 x_4 + y_1 x_2 x_3 y_4) = 0. \end{aligned}$$

Recall from Section 3 that  $\mathbb{B}$  is the blow up of  $\mathbb{P}_{a,b,c,d}^3$  along 12 points parametrising reducible divisors, and  $\mathbb{E} = \mathbb{P}_{a,b,c}^2$  is one of the exceptional divisors of this blow up whose closed points parametrise divisors in  $\mathbb{P}(1_{s_1}, 1_{t_1}, 2_{w_1}) \times \mathbb{P}(1_{s_2}, 1_{t_2}, 2_{w_2})$  that are given by

$$\text{(A.2)} \quad w_1 w_2 = a s_1 t_1 s_2 t_2 + \frac{b+c}{4}(s_1^2 s_2^2 + t_1^2 t_2^2) + \frac{b-c}{4}(s_1^2 t_2^2 + t_1^2 s_2^2).$$

Let  $\widetilde{\mathbb{P}^{\text{Sing}}}$  be the strict transform on  $\mathbb{B}$  of the divisor  $\mathbb{P}^{\text{Sing}}$ . If  $X$  is a singular Fano 3-fold which is parametrised by a point of  $\widetilde{\mathbb{P}^{\text{Sing}}}$ , it can be described in a simple geometric way. For instance, if  $X$  is a divisor (A.1) with two singular points, then we have the following commutative diagram:

$$\text{(A.3)} \quad \begin{array}{ccc} & Y & \\ \pi \swarrow & & \searrow \eta \\ \mathbb{P}^3 & \overset{\chi}{\dashrightarrow} & X \end{array}$$

where  $\pi$  is a blow up of four disjoint lines  $L_1, L_2, L_3, L_4$  that are not contained in a quadric surface,  $\chi$  is induced by the composition of linear projections  $\mathbb{P}^3 \dashrightarrow \mathbb{P}^1$  from these lines, and  $\eta$  is a small contraction of the strict transforms of the two lines in  $\mathbb{P}^3$  that intersect the lines  $L_1, L_2, L_3, L_4$ . If  $X$  is a divisor (A.1) singular along a curve, we still has (A.3),

but in this case the lines  $L_1, L_2, L_3, L_4$  are contained in a quadric surface, and  $\eta$  contracts the strict transform of this surface.

**A.1. Disjoint lines.** In this section, we will show the existence of the diagram (A.3) in the case when  $X$  is a divisor (A.1) whose singular locus is either two points or a curve.

Let  $G \simeq (\mathbb{Z}/2\mathbb{Z})^3$  be the subgroup in  $\text{Aut}(\mathbb{P}^3_{z_0, z_1, z_2, z_3})$  generated by the involutions:

$$\begin{aligned}\tau_1 &: (z_0 : z_1 : z_2 : z_3) \mapsto (z_1 : z_0 : z_3 : z_2); \\ \tau_2 &: (z_0 : z_1 : z_2 : z_3) \mapsto (z_0 : -z_1 : z_2 : -z_3); \\ \tau_3 &: (z_0 : z_1 : z_2 : z_3) \mapsto (z_2 : z_3 : z_0 : z_1).\end{aligned}$$

Fix  $(c_0 : c_1 : c_2 : c_3) \in \mathbb{P}^3$  such that  $c_0^2 \neq c_2^2$ , and let  $L_1$  be the line in  $\mathbb{P}^3$  that passes through the points  $(c_0 : c_1 : c_2 : c_3)$  and  $(c_2 : c_3 : c_0 : c_1) = \tau_3(c_0 : c_1 : c_2 : c_3)$ . Note that

$$(c_2 : c_3 : c_0 : c_1) \neq (c_0 : c_1 : c_2 : c_3).$$

Similarly, we define the lines  $L_2, L_3, L_4$  as follows:  $L_2 = \tau_1(L_1)$ ,  $L_3 = \tau_2(L_2)$ ,  $L_4 = \tau_2(L_1)$ . Then  $L_1 = \{f_1 = g_1 = 0\}$ ,  $L_2 = \{f_2 = g_2 = 0\}$ ,  $L_3 = \{f_3 = g_3 = 0\}$ ,  $L_4 = \{f_4 = g_4 = 0\}$ , where

$$\begin{aligned}f_1 &= (c_2c_3 - c_0c_1)z_0 + (c_0^2 - c_2^2)z_1 + (c_1c_2 - c_0c_3)z_2, \\ g_1 &= (c_1c_2 - c_0c_3)z_0 + (c_2c_3 - c_0c_1)z_2 + (c_0^2 - c_2^2)z_3, \\ f_2 &= (c_0^2 - c_2^2)z_0 + (c_2c_3 - c_0c_1)z_1 + (c_1c_2 - c_0c_3)z_3, \\ g_2 &= (c_1c_2 - c_0c_3)z_1 + (c_0^2 - c_2^2)z_2 + (c_2c_3 - c_0c_1)z_3, \\ f_3 &= (c_0^2 - c_2^2)z_0 + (c_0c_1 - c_2c_3)z_1 + (c_0c_3 - c_1c_2)z_3, \\ g_3 &= (c_0c_3 - c_1c_2)z_1 + (c_0^2 - c_2^2)z_2 + (c_0c_1 - c_2c_3)z_3, \\ f_4 &= (c_2c_3 - c_0c_1)z_0 - (c_0^2 - c_2^2)z_1 + (c_1c_2 - c_0c_3)z_2, \\ g_4 &= (c_1c_2 - c_0c_3)z_0 + (c_2c_3 - c_0c_1)z_2 - (c_0^2 - c_2^2)z_3.\end{aligned}$$

Set

$$\begin{aligned}D_{12} &= ((c_0 + c_1)^2 - (c_2 + c_3)^2)((c_0 - c_1)^2 - (c_2 - c_3)^2), \\ D_{13} &= ((c_0 - c_2)^2 + (c_1 - c_3)^2)((c_0 + c_2)^2 + (c_1 + c_3)^2), \\ D_{14} &= (c_0^2 - c_2^2)(c_1^2 - c_3^2).\end{aligned}$$

Then the lines  $L_1, L_2, L_3, L_4$  are disjoint if and only if

$$(A.4) \quad D_{12}D_{13}D_{14} \neq 0.$$

Similarly, the lines  $L_1, L_2, L_3, L_4$  are distinct but each of them intersects with exactly two of the others if and only if

$$(A.5) \quad \begin{aligned}D_{12} = 0, D_{13} = 0, D_{14} \neq 0, \text{ or} \\ D_{12} = 0, D_{14} = 0, D_{13} \neq 0, \text{ or} \\ D_{13} = 0, D_{14} = 0, D_{12} \neq 0.\end{aligned}$$

From now on, we assume that the lines  $L_1, L_2, L_3, L_4$  are disjoint.

Let  $\chi: \mathbb{P}^3 \dashrightarrow (\mathbb{P}^1)^4$  be the map given by

$$(A.6) \quad (z_0 : z_1 : z_2 : z_3) \mapsto ((f_1 : g_1), (f_2 : g_2), (f_3 : g_3), (f_4 : g_4)),$$

and let  $X$  be the closure of its image in  $(\mathbb{P}^1)^4$ . Then  $\chi$  is  $G$ -equivariant, and  $X$  is a divisor in  $(\mathbb{P}^1)^4$  of degree  $(1, 1, 1, 1)$  which is given by the following equation:

$$\begin{aligned} & 4(c_0c_3 - c_1c_2)^2(x_1x_2x_3x_4 + y_1y_2y_3y_4) - \\ & - (c_0^2 - c_1^2 - c_2^2 + c_3^2)^2(x_1x_2y_3y_4 + x_3x_4y_1y_2) + \\ & + (c_0^2 + c_1^2 - c_2^2 - c_3^2)^2(x_1x_3y_2y_4 + x_2x_4y_1y_3) - \\ & - 4(c_0c_1 - c_2c_3)^2(x_1x_4y_2y_3 + x_2x_3y_1y_4) = 0. \end{aligned}$$

This is equation (A.1) with

$$(A.7) \quad \begin{cases} a = ((c_0 + c_1)^2 - (c_2 + c_3)^2)((c_2 - c_3)^2 - (c_0 - c_1)^2), \\ b = ((c_0 + c_1)^2 - (c_2 + c_3)^2)((c_0 - c_1)^2 - (c_2 - c_3)^2), \\ c = (c_0^2 + c_1^2 - c_2^2 - c_3^2)^2 + 4(c_0c_1 - c_2c_3)^2, \\ d = 4(c_0c_3 - c_1c_2)^2 + (c_0^2 - c_1^2 - c_2^2 + c_3^2), \end{cases}$$

so that we have  $a + b = 0$ . Moreover, we have  $G$ -equivariant commutative diagram (A.3), where  $\eta$  is the birational morphism contracting strict transforms of all lines in  $\mathbb{P}^3$  that intersect the lines  $L_1, L_2, L_3, L_4$ . In particular, if  $L_1, L_2, L_3, L_4$  are contained in a quadric, then  $\eta$  contracts a divisor, and  $X$  is singular along a curve. This does not happen if

$$(A.8) \quad (c_0^2 + c_1^2 - c_2^2 - c_3^2)(c_0^2 - c_1^2 - c_2^2 + c_3^2)(c_0c_1 - c_2c_3)(c_0c_3 - c_1c_2) \neq 0.$$

Moreover, if  $L_1, L_2, L_3, L_4$  are not contained in a quadric, then there exist exactly two distinct lines in  $\mathbb{P}^3$  that intersect all lines  $L_1, L_2, L_3, L_4$ .

We have constructed (A.3) such that  $X$  is a divisor (A.1) that either has two nodes or a curve of singularities. However, it is not clear that we can obtain all divisors (A.1) that have these properties. This follows from the following lemmas.

**Lemma A.9.** *Let  $\rho: \mathbb{P}_{z_0, z_1, z_2, z_3}^3 \dashrightarrow \mathbb{P}_{x, y, z, t}^3$  be the rational map given by*

$$(z_0 : z_1 : z_2 : z_3) \mapsto (2(z_0z_3 - z_1z_2) : (z_0^2 - z_1^2 - z_2^2 + z_3^2) : 2(z_0z_1 - z_2z_3) : (z_0^2 + z_1^2 - z_2^2 - z_3^2)),$$

let  $\sigma: \mathbb{P}_{x, y, z, t}^3 \rightarrow \mathbb{P}_{a, b, c, d}^3$  be the morphism given by

$$(x : y : z : t) \mapsto (x^2 - y^2 : -z^2 + t^2 : z^2 + t^2 : x^2 + y^2).$$

Take any point  $(a : b : c : d) \in \mathbb{P}^3 \setminus (1 : -1 : 1 : 1)$  such that  $a + b = 0$ . Then there exists a point  $(c_0 : c_1 : c_2 : c_3) \in \mathbb{P}^3$  outside of the indeterminacy locus of  $\sigma \circ \rho$  such that

$$\sigma \circ \rho(c_0 : c_1 : c_2 : c_3) = (a : b : c : d)$$

and  $c_0^2 \neq c_2^2$ .

*Proof.* Observe that the image of  $\rho$  is the quadric surface  $Q = \{x^2 - y^2 - z^2 + t^2 = 0\}$ , the image  $\sigma(Q)$  is the plane  $\{a + b = 0\}$ , and  $\rho$  fits the following commutative diagram:

$$\begin{array}{ccc} & W & \\ \varpi \swarrow & & \searrow \varphi \\ \mathbb{P}_{z_0, z_1, z_2, z_3}^3 & \dashrightarrow & \mathbb{P}_{s_1, t_1}^1 \times \mathbb{P}_{s_2, t_2}^1 \simeq Q \end{array}$$



where  $\varpi$  is the blow up of the lines  $\{z_0 - z_2 = z_3 - z_1 = 0\}$  and  $\{z_0 + z_2 = z_1 + z_3 = 0\}$ , and  $\varphi$  is a  $\mathbb{P}^1$ -bundle. Moreover, the map  $\mathbb{P}^3_{z_0, z_1, z_2, z_3} \dashrightarrow \mathbb{P}^1_{s_i, t_i}$  is given by

$$(z_0 : z_1 : z_2 : z_3) \mapsto ((z_0 - z_2 : z_3 - z_1), (z_1 + z_3 : z_0 + z_2)),$$

and the isomorphism  $\mathbb{P}^1_{s_1, t_1} \times \mathbb{P}^1_{s_2, t_2} \simeq Q$  is given by

$$((s_1 : t_1), (s_2 : t_2)) \mapsto (s_1 s_2 + t_1 t_2 : s_1 t_2 + s_2 t_1 : s_1 s_2 - t_1 t_2 : s_1 t_2 - s_2 t_1).$$

Since  $(a : b : c : d) \neq (1 : -1 : 1 : 1)$ , there exists a point  $((s_1 : t_1), (s_2 : t_2))$  with  $s_1 t_2 \neq 0$  which is mapped by  $\sigma$  to  $(a : b : c : d)$ . Since  $s_1 t_2 \neq 0$ , the image of the fiber of  $\varphi$  over the point  $((s_1 : t_1), (s_2 : t_2))$  is not contained by any plane of the form  $\{z_0 - \lambda z_2 = 0\}$ , which implies the required assertion.  $\square$

**Corollary A.10.** *Let  $(a : b : c : d)$  be a point in  $\mathbb{P}^3$  such that  $a + b = 0$ , and (A.1) defines a divisor  $X \subset (\mathbb{P}^1)^4$  such that  $X$  has either two singular points or a curve of singularities. Then there is  $(c_0 : c_1 : c_2 : c_3) \in \mathbb{P}^3$  satisfying (A.4) and (A.7).*

*Proof.* By Corollary 2.9,  $a(c^2 - d^2) \neq 0$ , so the assertion follows from Lemma A.9.  $\square$

**A.2. Intersecting lines.** What we did in Appendix A.1 works also in some cases when the lines  $L_1, L_2, L_3, L_4$  are not disjoint. Namely, let  $X \subset (\mathbb{P}^1)^4$  be the divisor (A.1) for

$$(a : b : c : d) = (-u : u : 1 : 1)$$

with some  $u \in \mathbb{C} \setminus \{\pm 1\}$ . If  $u \neq 0$ , then  $X$  has 4 ordinary double points by Theorem 2.8. Similarly, if  $u = 0$ , then  $X$  has 6 ordinary double points. Moreover, by Lemma A.9, there exists  $(c_0 : c_1 : c_2 : c_3) \in \mathbb{P}^3$  such that  $c_0^2 \neq c_2^2$  and

$$\sigma \circ \rho(c_0 : c_1 : c_2 : c_3) = (-u : u : 1 : 1).$$

Replacing  $(c_2, c_3)$  with  $(-c_2, -c_3)$  if necessary, we may assume that  $c_1 = c_3$ . Then

$$(u : 1) = (4c_1^2 - (c_0 + c_2)^2 : 4c_1^2 + (c_0 + c_2)^2).$$

Note that  $L_1 \cap L_4 = (-1 : 0 : 1 : 0)$  and  $L_2 \cap L_3 = (0 : 1 : 0 : -1)$ . If  $u \neq 0$ , then

$$(L_1 \cup L_4) \cap (L_2 \cup L_3) = \emptyset.$$

Similarly, if  $u = 0$ , then  $L_1 \cap L_2 = (1 : \pm 1 : 1 : \pm 1)$  and  $L_3 \cap L_4 = (-1 : \pm 1 : -1 : \pm 1)$ .

The image of the rational map  $\chi : \mathbb{P}^3 \dashrightarrow (\mathbb{P}^1)^4$  given by (A.6) defines our divisor  $X$ . Moreover, if  $u \neq 0$ , we have the following  $G$ -equivariant commutative diagram:

$$(A.11) \quad \begin{array}{ccc} U & \xleftarrow{\pi_2} & Y \\ \pi_1 \downarrow & & \downarrow \eta \\ \mathbb{P}^3 & \xrightarrow{\chi} & X \end{array}$$

where  $\pi_1$  blows up the points  $L_1 \cap L_4$  and  $L_2 \cap L_3$ ,  $\pi_2$  blows up the strict transforms of the lines  $L_1, L_2, L_3, L_4$ , and  $\eta$  is the anticanonical morphism. Note that  $\eta$  contracts the strict transforms of the two lines in  $\mathbb{P}^3$  that intersect  $L_1, L_2, L_3, L_4$  — these are the line that passes through  $L_1 \cap L_4$  and  $L_2 \cap L_3$ , and the intersection line of the planes containing the curves  $L_1 + L_4$  and  $L_2 + L_3$ , respectively. The morphism  $\eta$  also contracts the strict transforms of the two lines in the  $\pi_1$ -exceptional divisors that pass through the intersection points with strict transforms of  $L_1, L_2, L_3, L_4$ .

The diagram (A.11) exists for  $u = 0$ . In this case,  $\pi_1$  is the blow up of the intersection points  $L_1 \cap L_4$ ,  $L_2 \cap L_3$ ,  $L_1 \cap L_2$ ,  $L_3 \cap L_4$ , and  $\eta$  contracts the strict transforms of the two lines in  $\mathbb{P}^3$  that intersect  $L_1, L_2, L_3, L_4$  — the line that passes through  $L_1 \cap L_4$  and  $L_2 \cap L_3$ , and the line that passes through  $L_1 \cap L_2$  and  $L_3 \cap L_4$ . In addition to these two curves, the morphism  $\eta$  also contracts the strict transforms of the four lines in the  $\pi_1$ -exceptional divisors that pass through the intersection points with strict transforms of  $L_1, L_2, L_3, L_4$ .

**A.3. Merging lines.** The construction described in Appendix A.1 degenerate in the case when two pair of lines among  $L_1, L_2, L_3, L_4$  merge together. To be precise, let us fix two disjoint lines  $L_1$  and  $L_2$  in  $\mathbb{P}^3$ , and let  $\pi_1: U \rightarrow \mathbb{P}^3$  be the blow up of the lines  $L_1$  and  $L_2$ . Then we have the following commutative diagram

$$\begin{array}{ccc} & U & \\ \pi_1 \swarrow & & \searrow \nu \\ \mathbb{P}^3 & \dashrightarrow_{\rho} & \mathbb{P}^1 \times \mathbb{P}^1 \end{array}$$

where  $\rho$  is given by the product of projections from  $L_1$  and  $L_2$ , and  $\nu$  is a  $\mathbb{P}^1$ -bundle. Let  $E_1$  and  $E_2$  be the  $\pi_1$ -exceptional surfaces such that  $\pi_1(E_1) = L_1$  and  $\pi_1(E_2) = L_2$ , let  $L_3 \subset E_1$  and  $L_4 \subset E_2$  be (possibly reducible) curves such that  $\nu(L_3)$  and  $\nu(L_4)$  are curves of degree  $(1, 1)$ , and one of the following cases holds:

- (1)  $L_3$  and  $L_4$  are smooth and the intersection  $\nu(L_3) \cap \nu(L_4)$  consists of two points;
- (2)  $L_3$  and  $L_4$  are singular and the intersection  $\nu(L_3) \cap \nu(L_4)$  consists of two points;
- (3)  $L_3$  and  $L_4$  are smooth and  $\nu(L_3) = \nu(L_4)$ .

Let  $\pi_2: Y \rightarrow U$  be the blow up of the curves  $L_3$  and  $L_4$ , and let  $\tilde{E}_1$  and  $\tilde{E}_2$  be the strict transforms on  $Y$  of the surfaces  $E_1$  and  $E_2$ , respectively. Then  $Y$  is a weak Fano 3-fold, and the linear system  $| -K_Y |$  gives a birational morphism  $\eta: Y \rightarrow X$  such that  $X$  is a divisor in  $\mathbb{P}(1, 1, 2) \times \mathbb{P}(1, 1, 2)$  of degree  $(2, 2)$  that can be given by (A.2), and we have the following commutative diagram:

$$\begin{array}{ccccc} & & U & \xleftarrow{\pi_2} & Y \\ \pi_1 \swarrow & & \searrow \nu & & \searrow \eta \\ \mathbb{P}^3 & \dashrightarrow_{\chi} & \mathbb{P}^1 \times \mathbb{P}^1 & \dashleftarrow_{\rho} & X \end{array}$$

where the map  $\rho$  is given by the natural projection  $\mathbb{P}(1, 1, 2) \times \mathbb{P}(1, 1, 2) \dashrightarrow \mathbb{P}^1 \times \mathbb{P}^1$ .

The morphism  $\nu$  contracts  $\tilde{E}_1$  and  $\tilde{E}_2$  such that  $\eta(\tilde{E}_1)$  and  $\eta(\tilde{E}_2)$  are curves in  $\text{Sing}(X)$ , and  $\eta$  also contracts strict transforms of the fibers of the  $\mathbb{P}^1$ -bundle  $\nu$  that intersect both curves  $L_3$  and  $L_4$ . In particular, if the intersection  $\nu(L_3) \cap \nu(L_4)$  consists of two points, there are exactly two such fibers — the fibers over the points  $\nu(L_3) \cap \nu(L_4)$ . Similarly, if

$$\nu(L_3) = \nu(L_4),$$

then such fibers span a surface in  $Y$ , whose image in  $\mathbb{P}^3$  is a smooth quadric surface, so the morphism  $\eta$  contracts this surface to another curve of singularities of the 3-fold  $X$ . In both cases, the 3-fold  $X$  is one of the divisors (A.2) parametrised by points in  $\widetilde{\mathbb{P}^{\text{Sing}}(X)} \cap \mathbb{E}$ . Moreover, we can obtain all such divisors using this construction.

## REFERENCES

- [1] H. Abban, Z. Zhuang, *K-stability of Fano varieties via admissible flags*, Forum of Mathematics Pi **10** (2022), 1–43.
- [2] C. Araujo, A.-M. Castravet, I. Cheltsov, K. Fujita, A.-S. Kaloghiros, J. Martinez-Garcia, C. Shramov, H. Süß, N. Viswanathan, *The Calabi problem for Fano threefolds*, Lecture Notes in Mathematics, Cambridge University Press, 2023.
- [3] K. Ascher, K. DeVleming, Yu. Liu, *Wall crossing for K-moduli spaces of plane curves*, to appear in Proc. of LMS.
- [4] G. Belousov, K. Loginov, *K-stability of Fano threefolds of rank 4 and degree 24*, Eur. J. Math. **90** (2023), 80.
- [5] D. Boozer, *The moduli space of stable rank 2 parabolic bundles over an elliptic curve with 3 marked points*, Proc. Am. Math. Soc. **150** (2022), 4645–4653.
- [6] I. Cheltsov, T. Duarte Guerreiro, K. Fujita, I. Krylov, J. Martinez-Garcia, *K-stability of Casagrande-Druel varieties*, to appear in Journal für die Reine und Angewandte Mathematik.
- [7] I. Cheltsov, A. Dubouloz, T. Kishimoto, *Toric G-solid Fano threefolds*, Selecta Mathematica **29** (2023), paper 22, 45 pages.
- [8] I. Cheltsov, K. Fujita, T. Kishimoto, T. Okada, *K-stable divisors in  $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^2$  of degree (1, 1, 2)*, Nagoya Math. J. **251** (2023), 686–714.
- [9] I. Cheltsov, L. Marquand, Y. Tschinkel, Z. Zhang, *Equivariant geometry of singular cubic threefolds, II*, arXiv:2405.02744, 2024.
- [10] I. Cheltsov, Y. Tschinkel, Z. Zhang, *Equivariant geometry of singular cubic threefolds*, to appear in Forum of Mathematics, Sigma.
- [11] O. Chterental, D. Djokovic, *Normal forms and tensor ranks of pure states of four qubits*, Linear Algebra Research Advancee, G. D. Ling (Ed.), Chapter 4, pp. 133–167, Nova Science Publishers, New York, 2007.
- [12] A. Collino, *The fundamental group of the Fano surface, I*, Lecture Notes in Mathematics **947** (1982), 209–218.
- [13] H. Dietrich, W. de Graaf, A. Marrani, M. Origlia, *Classification of four qubit states and their stabilisers under SLOCC operations*, J. of Physics A: Math. Theor. **55** (2022), 095302.
- [14] A. Kasprzyk, *Canonical toric Fano threefolds*, Canadian Journal of Mathematics **62** (2010), 1293–1309.
- [15] K. Fujita, *A valuative criterion for uniform K-stability of  $\mathbb{Q}$ -Fano varieties*, J. Reine Angew. Math. **751** (2019), 309–338.
- [16] G. Gour, N. Wallach, *On symmetric SL-invariant polynomials in four qubits*, Symmetry: representation theory and its applications, 259–267, Progr. Math., **257**, Birkhäuser/Springer, New York, 2014.
- [17] F. Holweck, J.-G. Luque, M. Planat, *Singularity of type  $D_4$  arising from four-qubit systems*, J. Physics A **47** (2014), 135301.
- [18] F. Holweck, J.-G. Luque, J.-Y. Thibon, *Entanglement of four-qubit systems: a geometric atlas with polynomial compass II (the tame world)*, J. Math. Phys. **58** (2017), 022201.
- [19] A. Kuznetsov, Yu. Prokhorov, *Rationality over non-closed fields of Fano threefolds with higher geometric Picard rank*, Journal of the Institute of Mathematics of Jussieu **23** (2024), 207–247.
- [20] J. Kollár, R. Laza, G. Sacca, C. Voisin, *Remarks on degenerations of hyper-Kähler manifolds*, Ann. Inst. Fourier **68** (2018), 2837–2882.
- [21] J. Kollár, S. Mori, *Classification of three-dimensional flips*, J. Amer. Math. Soc. **5** (1992), no.3, 533–703.
- [22] C. Li, *K-semistability is equivariant volume minimization*, Duke Math. Journal **166** (2017), 3147–3218.
- [23] Q. Liu, *Algebraic geometry and arithmetic curves*, Oxf. Grad. Texts Math. **6**, Oxford University Press, Oxford, 2002.
- [24] Y. Liu, C. Xu and Z. Zhuang, *Finite generation for valuations computing stability thresholds and applications to K-stability*, Ann. of Math. **196** (2022), 507–566.
- [25] J.-G. Luque, J. Thibon, *Polynomial invariants of four qubits*, Phys. Rev. A **67**, 043203.

- [26] T. Sano, *On deformations of  $\mathbb{Q}$ -Fano 3-folds*, Algebraic Geom. **25** (2016), 141–176.
- [27] L. Schläfli, *Über die Resultante eines Systemes mehrerer algebraischer Gleichungen. Ein Beitrag zur Theorie der Elimination*, Denkschriften der Kaiserlichen Akademie der Wissenschaften/Mathematisch–Naturwissenschaftliche Classe **4** (1852), 1–74.
- [28] E. Sernesi, *Deformations of algebraic schemes*, Grundlehren Math. Wiss., **334**, Springer-Verlag, Berlin, 2006.
- [29] F. Verstraete, J. Dehaene, B. De Moor, H. Verschelde, *Four qubits can be entangled in nine different ways*, Phys. Rev. A **65** (2002), 052112.
- [30] Z. Zhuang, *Product theorem for  $K$ -stability*, Adv. Math. **371** (2020), 107250, 18 pp.
- [31] Z. Zhuang, *Optimal destabilizing centers and equivariant  $K$ -stability*, Invent. Math. **226** (2021), 195–223.
- [32] C. Xu,  *$K$ -stability of Fano varieties: an algebro-geometric approach*, EMS Surv. Math. Sci. **8** (2021), 265–354.

*Ivan Cheltsov*

University of Edinburgh, Edinburgh, Scotland  
i.cheltsov@ed.ac.uk

*Maksym Fedorchuk*

Boston College, Chestnut Hill, Massachusetts  
maksym.fedorchuk@bc.edu

*Kento Fujita*

Osaka University, Osaka, Japan  
fujita@math.sci.osaka-u.ac.jp

*Anne-Sophie Kaloghiros*

Brunel University London, Uxbridge, England  
anne-sophie.kaloghiros@brunel.ac.uk