

Schouten like metrics on five dimensional nilpotents Lie groups.

M. L. Foka¹, M. B. N. Djiadeu², T.B. Bouetou³,

e-mail: 1. lanndrymaris@gmail.com, 2. michel.djiadeu@facsciences-uy1.cm,

3. tbouetou@gmail.com,

University of Yaounde 1, Faculty of Science, Department of Mathematics, P.O. Box 812, Yaounde,
Republic of Cameroon.

Abstract

The prescribed Ricci curvature problem consists in finding a Riemannian metric g to satisfy the equation $Ric(g) = T$, for some fixed symmetric $(0, 2)$ -tensor field T on a differential manifold M . In this paper, we define Schouten like metric as a particular solution of a prescribed Ricci curvature problem and we classify them on five dimensional nilpotent Lie groups by establishing a link with algebraic Schouten solitons. This also permit us to make a classification of five dimensional nilsoliton.

keywords: Lie groups, Lie algebra, Ricci operator, nilsoliton, algebraic Schouten solitons, Schouten like metric.

MSC: 53C20, 53C35, 53C30

1 Introduction

In 2023, SIYAO Liu introduced in [6] the notion of Algebraic Schouten solitons based on the general definition of the Schouten tensor in [4]. He was strongly motivated by [4, 13]. However the Ricci curvature problem is a problem that arouses great interest in Riemannian geometry. Indeed, it is about the search for Riemannian metrics on a differentiable manifold whose Ricci curvature is a tensor field of type $(0,2)$ prescribed on the manifold. For a review of the literature on this problem, see the references [2, 3, 9, 11]. One of the most recent works on this problem is that of FOKA .M .L and al in [5]. Using the definition of Schouten solitons introduced in [4], we introduce the notion of Schouten likes metrics as the solutions of a prescribed Ricci curvature problem in which we impose a particular form on the prescribed tensor field. In this paper we devote ourselves to the classification of Schouten likes metrics on five-dimensional nilpotent Lie groups using the recent classification of five-dimensional nilpotent Lie groups given in [5]. We give as a direct correlative of this classification, a classification of the dimension five Nilsolitons whose first classification is given by MOGHADDAM and Hamid Reza Salimi in [10].

To carry out our work we organize it as follows. In Section 2 we unfold preliminary tools necessary for the establishment of our results. Section 3 will be dedicated to the study proper of Schouten likes metrics on nilpotent Lie groups of dimension five.

2 Preliminaries

In this section, we fix our notation and recall some useful facts that will be used to establish our results in the next section. Let (G, g) be a finite dimensional Lie group equipped with a left-invariant riemannian metric, and \mathfrak{g} its Lie algebra.

For any $u \in \mathfrak{g}$, we denote by $ad_u, J_u : \mathfrak{g} \rightarrow \mathfrak{g}$ the endomorphisms given by $ad_u v = [u, v]$ and $J_u v = ad_v^* u$ where ad_v^* is the transpose of ad_v . We denote also by $B : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ the Killing form given by

$$B(u, v) = tr(ad_u \circ ad_v) \quad \forall u, v \in \mathfrak{g}.$$

The mean curvature vector on \mathfrak{g} is defined by

$$\langle H, u \rangle = tr(ad_u), \quad \forall u \in \mathfrak{g} \quad (\langle \cdot, \cdot \rangle = g(e)).$$

The Ricci curvature tensor ric_g and the Ricci operator Ric_g of (G, g) are given by (see [1]),

$$ric_g(u, v) = -\frac{1}{2}B(u, v) - \frac{1}{2}tr(ad_u \circ ad_v^*) - \frac{1}{4}tr(J_u \circ J_v) - \frac{1}{2}(\langle ad_H u, v \rangle + \langle ad_H v, u \rangle)$$

and

$$ric_g(u, v) = g(Ric_g(u), v) \quad \text{for } u, v \in \mathfrak{g} \quad (1)$$

Remark 1 *If*

i- *If (G, g) is a Riemannian nilpotent Lie group. Then its Ricci curvature tensor is given by*

$$ric_g(u, v) = -\frac{1}{2}tr(ad_u \circ ad_v^*) - \frac{1}{4}tr(J_u \circ J_v). \quad (2)$$

ii- *Since ric_g is symmetric, Ric_g is self-adjoint.*

Definition 1 *A derivation $D \in Der(\mathfrak{g})$ is said to be symmetric with respect to the metric g if it satisfies the following condition:*

$$g(D(u), v) = g(u, D(v)) \quad \forall u, v \in \mathfrak{g}. \quad (3)$$

A generalized definition of the Schouten tensor is given in [6] by

$$S(X, Y) = ric_g(X, Y) - s\lambda_0 g(X, Y) \quad (4)$$

where s denotes the scalar curvature and λ_0 is a real number.

Definition 2 *A left-invariant Riemannian metric g on a Lie group G of Lie algebra \mathfrak{g} is called a nilsoliton if it satisfies*

$$Ric_g \in \mathbb{R}Id_{\mathfrak{g}} \oplus Der(\mathfrak{g}). \quad (5)$$

Where Ric_g is the Ricci operator of (G, g) and $Der(\mathfrak{g})$ the set of derivations of \mathfrak{g} .

Definition 3 *A left-invariant Riemannian metric g on a Lie group G of Lie algebra \mathfrak{g} is called an algebraic Schouten soliton associated with the connection ∇ if it satisfies*

$$Ric_g - (s\lambda_0 + c)Id \in Der(\mathfrak{g}). \quad (6)$$

Where ∇ is the Levi-Civita connection of (G, g) , Ric_g the Ricci operator of (G, g) , s the scalar curvature of (G, g) , c, λ_0 and c are real numbers and $Der(\mathfrak{g})$ the set of derivations of \mathfrak{g} .

Proposition 1 *If $D = Ric_g - (s\lambda_0 + c)Id$ is a derivation of \mathfrak{g} then it's symmetric with respect to g .*

Proof: Let's suppose that $D = Ric_g - (s\lambda_0 + c)Id$ is a derivation of \mathfrak{g} . Let $u, v \in \mathfrak{g}$.

$$\begin{aligned}
g(D(u), v) &= g(Ric_g(u) - (s\lambda_0 + c)u, v) \\
&= g(Ric_g(u), v) - (s\lambda_0 + c)g(u, v) \\
&= ric_g(u, v) - (s\lambda_0 + c)g(u, v) \\
&= ric_g(v, u) - (s\lambda_0 + c)g(v, u) \\
&= g(Ric_g(v), u) - (s\lambda_0 + c)g(v, u) \\
&= g(Ric_g(v) - (s\lambda_0 + c)v, u) \quad \text{because } Ric_g \text{ is self-adjoint} \\
&= g(D(v), u) \\
&= g(u, D(v))
\end{aligned}$$

■

Proposition 2 *Nilsolitons and algebraic Schouten solitons are invariant under isometry and scaling.*

Proof: Indeed, Let g and h be two left-invariant riemannian metrics on G such that there exist $c > 0$, $\varphi \in Aut(\mathfrak{g})$ satisfying $h = c\varphi.g$. Then $Ric_h = c^2\varphi Ric_g \varphi^{-1}$.

As $\varphi Der(\mathfrak{g}) \varphi^{-1} = Der(\mathfrak{g})$, it follows that $Ric_g \in \mathbb{R}Id_{\mathfrak{g}} \oplus Der(\mathfrak{g})$ if and only if $Ric_h \in \mathbb{R}Id_{\mathfrak{g}} \oplus Der(\mathfrak{g})$.

The proof is similar for algebraic Schouten solitons. ■

Remark 2 *Nilsolitons constitute a particular class of algebraic Schouten solitons. This because by taking $\lambda_0 = 0$ in the definition of algebraic Schouten solitons, we obtain Nilsolitons.*

Definition 4 *A left-invariant Riemannian metric g on a Lie group G of Lie algebra \mathfrak{g} is called a Schouten like metric if it satisfies*

$$ric_g(\cdot, \cdot) = (s\lambda_0 + c)g(\cdot, \cdot) + g(D(\cdot), \cdot). \quad (7)$$

Where s is the scalar curvature, λ_0 and c are real numbers, and D is symmetric derivation of \mathfrak{g} with respect to g .

Lemma 1 *A left-invariant Riemannian metric g on a Lie group G of Lie algebra \mathfrak{g} is a Schouten like metric if and only if it's an algebraic Schouten soliton*

Proof: Let g be a left-invariant Riemannian metric on a Lie group G of Lie algebra \mathfrak{g} .

If g is a Schouten like metric on G then for all $X, Y \in \mathfrak{g}$,

$$ric_g(X, Y) = (s\lambda_0 + c)g(X, Y) + g(D(X), Y) \quad \lambda_0, c \in \mathbb{R}.$$

From the definition of the Ricci operator, we have

$$g(Ric_g(X), Y) = (s\lambda_0 + c)g(X, Y) + g(D(X), Y).$$

Therefore the fact that g is bilinear gives,

$$g(Ric_g(X), Y) = g((s\lambda_0 + c)X + D(X), Y).$$

Since g is non-degenerated, $Ric_g(X) - (s\lambda_0 + c)X = D(X)$ for all $X \in \mathfrak{g}$. So g is an algebraic Schouten soliton on G .

Conversely, let us assume that g is an algebraic Schouten soliton on G . Then there exist $c \in \mathbb{R}$ and $D \in \text{Der}(\mathfrak{g})$ such that

$$\text{Ric}_g - (s\lambda_0 + c)\text{Id} = D, \quad \lambda_0 \in \mathbb{R}. \quad (8)$$

Therefore, for all $X, Y \in \mathfrak{g}$,

$$g(\text{Ric}_g(X), Y) = g((s\lambda_0 + c)X + D(X), Y).$$

Then, for all $X, Y \in \mathfrak{g}$,

$$\text{ric}_g(X, Y) = g(\text{Ric}_g(X), Y) = (s\lambda_0 + c)g(X, Y) + g(D(X), Y).$$

Moreover, Proposition 1 ensures the symmetry of D . Consequently, g qualifies as a Schouten-like metric on G . ■

Marius Landry Foka et al., [5] characterised the set of all inner products on each five-dimensional nilpotent Lie algebra using the Milnor-type theorem technique. For a detailed exploration of this technique, refer to [7, 8, 12]. The characterization is encapsulated in the following result.

Proposition 3 [5]

Let \mathfrak{g} be a five-dimensional nilpotent Lie algebra. For every inner product $\langle \cdot, \cdot \rangle$ on \mathfrak{g} , there exist $\eta > 0$, and an orthonormal basis $\mathcal{B} = \{v_1, v_2, v_3, v_4, v_5\}$ with respect to $\eta\langle \cdot, \cdot \rangle$, such that all non zero commutators are given by Table 1:

3 Schouten like metrics on Five dimensional nilpotent Lie groups

Throughout this section, by \mathfrak{g} we shall denote any of the five-dimensional nilpotent Lie algebra cited in Table 1 and G the five-dimensional simply connected nilpotent Lie group having Lie algebra \mathfrak{g} .

From Lemma 1, one can observe that the study of Schouten like metrics can be done through Schouten solitons. For this purpose let introduce the following endomorphism D of \mathfrak{g} with g a left-invariant Riemannian metric on G

$$\begin{aligned} D : \mathfrak{g} &\longrightarrow \mathfrak{g} \\ X &\longmapsto (\text{Ric}_g - (\lambda_0 s + c)\text{Id}_{\mathfrak{g}})X. \end{aligned}$$

Where λ_0 and c are real constants and Ric_g the Ricci operator of (G, g) . The metric g will be a Schouten like metric if and only if D satisfy the following equation:

$$D[X, Y] = [DX, Y] + [X, DY] \quad \text{for } X, Y \in \mathfrak{g}. \quad (9)$$

The corollaries of the following proposition come directly from the Remark 2.

Proposition 4 An inner product $\langle \cdot, \cdot \rangle$ on $\mathfrak{g} = A_{5,4}$ is a Schouten like metric if and only if $\alpha = 0$ and $\beta = \gamma$.

Proof: Let $\langle \cdot, \cdot \rangle$ be an inner product on $A_{5,4}$. From [5],

$$\text{Ric}_{\langle \cdot, \cdot \rangle} = -\frac{1}{2} \begin{pmatrix} \alpha^2 + \beta^2 & \alpha\gamma & 0 & 0 & 0 \\ \alpha\gamma & \gamma^2 & 0 & 0 & 0 \\ 0 & 0 & \alpha^2 + \gamma^2 & \alpha\beta & 0 \\ 0 & 0 & \alpha\beta & \beta^2 & 0 \\ 0 & 0 & 0 & 0 & -\alpha^2 - \beta^2 - \gamma^2 \end{pmatrix}. \quad (10)$$

Table 1:

Lie algebra	non zero commutation relation	conditions
$5A_1$	none	none
$A_{5,4}$	$[v_1, v_3] = \alpha v_5, [v_1, v_4] = \beta v_5$ $[v_2, v_3] = \gamma v_5$	$\beta, \gamma > 0, \alpha \in \mathbb{R}$
$A_{3,1} \oplus 2A_1$	$[v_1, v_2] = \alpha v_5$	$\alpha > 0$
$A_{4,1} \oplus A_1$ first case	$[v_1, v_2] = \alpha v_3 + \gamma v_5, [v_1, v_3] = \beta v_5$	$\alpha, \beta > 0, \gamma \in \mathbb{R}$
$A_{4,1} \oplus A_1$ second case	$[v_1, v_2] = \alpha v_3 + \gamma v_4, [v_1, v_3] = \beta v_5$	$\alpha, \beta > 0, \gamma \in \mathbb{R}$
$A_{5,6}$	$[v_1, v_2] = \alpha v_3 + \beta v_4, [v_1, v_3] = \gamma v_4 + \delta v_5$ $[v_1, v_4] = \varepsilon v_5, [v_2, v_3] = \sigma v_5$	$\alpha < 0, \gamma, \varepsilon, \sigma > 0, \beta, \delta \in \mathbb{R}$
$A_{5,5}$	$[v_1, v_2] = \alpha v_4 + \beta v_5, [v_1, v_3] = \gamma v_5$ $[v_2, v_3] = \delta v_5, [v_2, v_4] = \varepsilon v_5$	$\alpha, \gamma, \varepsilon > 0, \beta, \delta \in \mathbb{R}$
$A_{5,3}$	$[v_1, v_2] = \alpha v_3 + \beta v_4, [v_1, v_3] = \gamma v_4 + \delta v_5$ $[v_2, v_3] = \varepsilon v_5$	$\alpha, \gamma, \varepsilon > 0, \beta, \delta \in \mathbb{R}$
$A_{5,1}$	$[v_1, v_2] = \alpha v_4 + \beta v_5, [v_1, v_3] = \gamma v_5$	$\alpha, \gamma > 0, \beta \in \mathbb{R}$
$A_{5,2}$	$[v_1, v_2] = \alpha v_3 + \beta v_4, [v_1, v_3] = \gamma v_4$ $[v_1, v_4] = \delta v_5$	$\alpha, \gamma, \delta > 0, \beta \in \mathbb{R}$

Therefore, we can write D as

$$\begin{cases} Dv_1 = -\frac{1}{2}[(1 - \lambda_0)\alpha^2 + (1 - \lambda_0)\beta^2 - \lambda_0\gamma^2 + 2c]v_1 - \frac{1}{2}\alpha\gamma v_2 \\ Dv_2 = -\frac{1}{2}\alpha\gamma v_1 - \frac{1}{2}[-\lambda_0\alpha^2 - \lambda_0\beta^2 + (1 - \lambda_0)\gamma^2 + 2c]v_2 \\ Dv_3 = -\frac{1}{2}[(1 - \lambda_0)\alpha^2 - \lambda_0\beta^2 + (1 - \lambda_0)\gamma^2 + 2c]v_3 - \frac{1}{2}\alpha\beta v_4 \\ Dv_4 = -\frac{1}{2}\alpha\beta v_3 - \frac{1}{2}[-\lambda_0\alpha^2 + (1 - \lambda_0)\beta^2 - \lambda_0\gamma^2 + 2c]v_4 \\ Dv_5 = \frac{1}{2}[(1 + \lambda_0)\alpha^2 + (1 + \lambda_0)\beta^2 + (1 + \lambda_0)\gamma^2 - 2c]v_5 \end{cases} \quad (11)$$

Hence, by (9), \langle, \rangle is an algebraic Schouten soliton if and only if the following polynomial system of equations is satisfied

$$\begin{aligned}\alpha[(3 - \lambda_0)\alpha^2 + (3 - \lambda_0)\beta^2 + (3 - \lambda_0)\gamma^2 + 2c] &= 0 \\ \beta[(3 - \lambda_0)\alpha^2 + (3 - \lambda_0)\beta^2 + (1 - \lambda_0)\gamma^2 + 2c] &= 0 \\ \gamma[(3 - \lambda_0)\alpha^2 + (1 - \lambda_0)\beta^2 + (3 - \lambda_0)\gamma^2 + 2c] &= 0 \\ \alpha\beta\gamma &= 0.\end{aligned}\tag{12}$$

Since $\beta, \gamma > 0$, we obtain from the fourth equation of system (12), that $\alpha = 0$. Therefore, the second and third equations of system (12) become

$$\begin{aligned}(3 - \lambda_0)\beta^2 + (1 - \lambda_0)\gamma^2 + 2c &= 0, \\ (1 - \lambda_0)\beta^2 + (3 - \lambda_0)\gamma^2 + 2c &= 0.\end{aligned}$$

Hence $2\beta^2 - 2\gamma^2 = 0$. i.e. $\beta = \gamma$. ■

Corollary 1 *An inner product \langle, \rangle on $\mathfrak{g} = A_{5,4}$ is a nilsoliton if and only if $\alpha = 0$ and $\beta = \gamma$.*

Proposition 5 *Any inner product on $A_{3,1} \oplus 2A_1$ is a Schouten like metric.*

Proof: Let \langle, \rangle be an inner product on $A_{3,1} \oplus 2A_1$. In [5], the Ricci operator is given by

$$Ric_{\langle, \rangle} = -\frac{1}{2}diag(\alpha^2, \alpha^2, 0, 0, -\alpha^2).\tag{13}$$

We can write D as:

$$\begin{cases} Dv_1 = -\frac{1}{2}[(1 - \lambda_0)\alpha^2 + 2c]v_1 \\ Dv_2 = -\frac{1}{2}[(1 - \lambda_0)\alpha^2 + 2c]v_2 \\ Dv_3 = -\frac{1}{2}(\lambda_0 + 2c)v_3 \\ Dv_4 = -\frac{1}{2}(\lambda_0 + 2c)v_4 \\ Dv_5 = \frac{1}{2}[(1 + \lambda_0)\alpha^2 - 2c]v_5. \end{cases}\tag{14}$$

By using (9), and making tedious calculations, we have:

$$\alpha[(3 - \lambda_0)\alpha^2 + 2c] = 0.\tag{15}$$

Then taking $c = -\frac{\lambda_0 - 3}{2}\alpha^2$, we have $Ric_{\langle, \rangle} - (\lambda_0 s + c)Id \in \mathbb{R} \oplus Der(\mathfrak{g})$. ■

Corollary 2 *Any inner product on $A_{3,1} \oplus 2A_1$ is a nilsoliton.*

Proposition 6 *An inner product \langle, \rangle on $\mathfrak{g} = A_{4,1} \oplus A_1$ is a Schouten like metri if and only if $\gamma = 0$ and $\alpha = \beta$.*

Proof: Let \langle, \rangle be an inner product on $A_{4,1} \oplus A_1$. From reference [5],

$$Ric_{\langle, \rangle} = -\frac{1}{2} \begin{pmatrix} \alpha^2 + \beta^2 + \gamma^2 & 0 & 0 & 0 \\ 0 & \alpha^2 + \gamma^2 & 0 & 0 \\ 0 & \beta\gamma & \beta^2 - \alpha^2 & -\alpha\gamma \\ 0 & 0 & 0 & -\beta^2 - \gamma^2 \end{pmatrix} \text{ or } Ric_{\langle, \rangle} = -\frac{1}{2} \begin{pmatrix} \alpha^2 + \beta^2 + \gamma^2 & 0 & 0 & 0 \\ 0 & \alpha^2 + \gamma^2 & 0 & 0 \\ 0 & 0 & \alpha^2 - \beta^2 & -\alpha\gamma \\ 0 & 0 & -\alpha\gamma & -\gamma^2 \\ 0 & 0 & 0 & -\beta^2 \end{pmatrix}.$$

i) If

$$Ric_{\langle, \rangle} = -\frac{1}{2} \begin{pmatrix} \alpha^2 + \beta^2 + \gamma^2 & 0 & 0 & 0 & 0 \\ 0 & \alpha^2 + \gamma^2 & \beta\gamma & 0 & 0 \\ 0 & \beta\gamma & \beta^2 - \alpha^2 & 0 & -\alpha\gamma \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\alpha\gamma & 0 & -\beta^2 - \gamma^2 \end{pmatrix}.$$

A simple calculation shows that:

$$\begin{cases} Dv_1 = -\frac{1}{2}[(1 - \lambda_0)\alpha^2 + (1 - \lambda_0)\beta^2 + (1 - \lambda_0)\gamma^2 + 2c]v_1 \\ Dv_2 = -\frac{1}{2}[(1 - \lambda_0)\alpha^2 - \lambda_0\beta^2 + (1 - \lambda_0)\gamma^2 + 2c]v_2 - \frac{1}{2}\beta\gamma v_3 \\ Dv_3 = -\frac{1}{2}\beta\gamma v_2 - \frac{1}{2}[-(1 + \lambda_0)\alpha^2 + (1 - \lambda_0)\beta^2 - \lambda_0\gamma^2 + 2c]v_3 + \frac{1}{2}\alpha\gamma v_5 \\ Dv_4 = -\frac{1}{2}[\lambda_0\alpha^2 + \lambda_0\beta^2 + \lambda_0\gamma^2 - 2c]v_4 \\ Dv_5 = \frac{1}{2}\alpha\gamma v_3 + \frac{1}{2}[\lambda_0\alpha^2 + (1 + \lambda_0)\beta^2 + (1 + \lambda_0)\gamma^2 - 2c]v_5. \end{cases}$$

Hence, (9) yields, \langle, \rangle is a Schouten like metric if and only if the following system of equations is satisfied

$$\begin{aligned} \alpha[(3 - \lambda_0)\alpha^2 - \lambda_0\beta^2 + (3 - \lambda_0)\gamma^2 + 2c] &= 0 \\ \beta[-\lambda_0\alpha^2 + (3 - \lambda_0)\beta^2 + (3 - \lambda_0)\gamma^2 + 2] &= 0 \\ \gamma[(2 - \lambda_0)\alpha^2 + (3 - \lambda_0)\beta^2 + (3 - \lambda_0)\gamma^2 + 2c] &= 0 \\ \alpha\beta\gamma &= 0. \end{aligned} \quad (16)$$

Since $\alpha, \beta > 0$, then from the fourth equation of system (16), $\gamma = 0$. Therefore, the system become

$$\begin{aligned} (3 - \lambda_0)\alpha^2 - \lambda_0\beta^2 + 2c &= 0, \\ -\lambda_0\alpha^2 + (3 - \lambda_0)\beta^2 + 2c &= 0. \end{aligned}$$

Then, $\alpha = \beta$.

ii) If

$$Ric_{\langle, \rangle} = -\frac{1}{2} \begin{pmatrix} \alpha^2 + \beta^2 + \gamma^2 & 0 & 0 & 0 & 0 \\ 0 & \alpha^2 + \gamma^2 & 0 & 0 & 0 \\ 0 & 0 & \alpha^2 - \beta^2 & -\alpha\gamma & 0 \\ 0 & 0 & -\alpha\gamma & -\gamma^2 & 0 \\ 0 & 0 & 0 & 0 & -\beta^2 \end{pmatrix},$$

we can then write D as

$$\begin{cases} Dv_1 = -\frac{1}{2}[(1 - 3\lambda_0)\alpha^2 + (1 + \lambda_0)\beta^2 + (1 - \lambda_0)\gamma^2 + 2c]v_1 \\ Dv_2 = -\frac{1}{2}[(1 - 3\lambda_0)\alpha^2 + \lambda_0\beta^2 + (1 - \lambda_0)\gamma^2 + 2c]v_2 \\ Dv_3 = -\frac{1}{2}[(1 - 3\lambda_0)\alpha^2 + (\lambda_0 - 1)\beta^2 - \lambda_0\gamma^2 + 2c]v_3 + \frac{1}{2}\alpha\gamma v_4 \\ Dv_4 = \frac{1}{2}\alpha\gamma v_3 + \frac{1}{2}[3\lambda_0\alpha^2 - \lambda_0\beta^2 + (1 + \lambda_0)\gamma^2 - 2c]v_4 \\ Dv_5 = +\frac{1}{2}[3\lambda_0\alpha^2 + (1 - \lambda_0)\beta^2 + \lambda_0\gamma^2 - 2c]v_5. \end{cases}$$

Thus, (9) now acquiesces, \langle, \rangle is a Schouten like metric if and only if the subsequent system is satisfied

$$\begin{aligned} \alpha[(1 - 3\lambda_0)\alpha^2 + (2 + \lambda_0)\beta^2 + (3 - \lambda_0)\gamma^2 + 2c] &= 0 \\ \beta[(2 - 3\lambda_0)\alpha^2 + (1 + \lambda_0)\beta^2 + (1 + 3\lambda_0)\gamma^2 + 2] &= 0 \\ \gamma[3(1 - \lambda_0)\alpha^2 + (1 + \lambda_0)\beta^2 + (3 - \lambda_0)\gamma^2 + 2c] &= 0 \\ \alpha\beta\gamma &= 0. \end{aligned} \quad (17)$$

From the fourth equation of system (17), one have $\gamma = 0$. Consequently, the system become

$$\begin{aligned}(1 - 3\lambda_0)\alpha^2 + (2 + \lambda_0)\beta^2 + 2c &= 0, \\ (2 - 3\lambda_0)\alpha^2 + (1 + \lambda_0)\beta^2 + 2c &= 0.\end{aligned}$$

Therefore, $\alpha = \beta$.

■

Corollary 3 An inner product \langle, \rangle on $\mathfrak{g} = A_{4,1} \oplus A_1$ is a nilsoliton if and only if $\gamma = 0$ and $\alpha = \beta$.

Proposition 7 The Lie algebra $A_{5,6}$ does not admit Schouten like metrics.

Proof: Let \langle, \rangle be an inner product on $A_{5,6}$. In [5], the Ricci operator is given by

$$Ric_{\langle, \rangle} = -\frac{1}{2} \begin{pmatrix} \alpha^2 + \beta^2 + \gamma^2 + \delta^2 + \varepsilon^2 & \delta\sigma & 0 & 0 & 0 \\ \delta\sigma & \alpha^2 + \beta^2 & \beta\gamma & 0 & 0 \\ 0 & \beta\gamma & \gamma^2 + \delta^2 + \sigma^2 - \alpha^2 & \delta\varepsilon - \alpha\beta & 0 \\ 0 & 0 & \delta\varepsilon - \alpha\beta & \varepsilon^2 - \beta^2 - \gamma^2 & -\delta\gamma \\ 0 & 0 & 0 & -\delta\gamma & -\delta^2 - \varepsilon^2 - \sigma^2 \end{pmatrix}. \quad (18)$$

Assuming that \langle, \rangle is a Schouten like metric, there exist a real number c and a derivation D on $A_{5,6}$ such that $Ric_{\langle, \rangle} = (\lambda_0 s + c)Id + D$. A simple calculation shows that

$$\begin{cases} Dv_1 = -\frac{1}{2}[(1 - \lambda_0)\alpha^2 + (1 - \lambda_0)\beta^2 + (1 - \lambda_0)\gamma^2 + (1 - \lambda_0)\delta^2 + (1 - \lambda_0)\varepsilon^2 + 2c]v_1 - \frac{1}{2}\varepsilon\sigma v_2 \\ Dv_2 = -\frac{1}{2}\delta\sigma v_1 - \frac{1}{2}[(1 - \lambda_0)\alpha^2 + (1 - \lambda_0)\beta^2 - \lambda_0\gamma^2 - \lambda_0\delta^2 - \lambda_0\varepsilon^2 + 2c]v_2 - \frac{1}{2}\beta\gamma v_3 \\ Dv_3 = -\frac{1}{2}\beta\gamma v_2 - \frac{1}{2}[-(1 + \lambda_0)\alpha^2 - \lambda_0\beta^2 + (1 - \lambda_0)\gamma^2 + (1 - \lambda_0)\delta^2 - \lambda_0\varepsilon^2 + \sigma^2 + 2c]v_3 - \frac{1}{2}(\delta\varepsilon - \alpha\beta)v_4 \\ Dv_4 = -\frac{1}{2}(\delta\varepsilon - \alpha\beta)v_3 + \frac{1}{2}[\lambda_0\alpha^2 + (1 + \lambda_0)\beta^2 + (1 + \lambda_0)\gamma^2 + \lambda_0\delta^2 + (\lambda_0 - 1)\varepsilon^2 - 2c]v_4 + \frac{1}{2}\delta\gamma v_5 \\ Dv_5 = \frac{1}{2}\delta\gamma v_4 + \frac{1}{2}[\lambda_0\alpha^2 + \lambda_0\beta^2 + \lambda_0\gamma^2 + (1 + \lambda_0)\delta^2 + (1 + \lambda_0)\varepsilon^2 + \sigma^2 - 2c]v_5. \end{cases}$$

Let apply (9), to the following system:

$$\begin{aligned}[v_1, v_5] &= 0 \\ [v_2, v_4] &= 0.\end{aligned}$$

We obtaint:

$$\begin{aligned}\delta\gamma\varepsilon &= 0 \\ \sigma(2\delta\varepsilon - \alpha\beta) &= 0\end{aligned}$$

Therefore $\beta = \delta = 0$ inasmuch as $\alpha < 0$, $\gamma, \sigma, \varepsilon > 0$. Then, the relation

$[v_1, v_3] = \gamma v_4 + \delta v_5$ becomes:

$$-\frac{1}{2}\gamma[-\lambda_0\alpha^2 + (3 - \lambda_0)\gamma^2 - \lambda_0\varepsilon^2 + \sigma^2 + 2c]v_4 + \frac{1}{2}\varepsilon\sigma^2 v_5 = 0. \quad (19)$$

Hence, $\varepsilon\sigma^2 = 0$. This contradict the fact that $\varepsilon, \sigma > 0$. ■

Corollary 4 The Lie algebra $A_{5,6}$ does not admit nilsoliton.

Proposition 8 An inner product \langle, \rangle on $\mathfrak{g} = A_{5,5}$ is a Schouten like metric if and only if $\beta = \delta = 0$ and $\alpha = \varepsilon = \sqrt{2}\gamma$.

Proof: Let \langle , \rangle be an inner product on $A_{5,5}$. In [5], the Ricci operator is given by

$$Ric_{\langle , \rangle} = -\frac{1}{2} \begin{pmatrix} \alpha^2 + \beta^2 + \gamma^2 & \gamma\delta & -\beta\delta & -\beta\varepsilon & 0 \\ \gamma\delta & \alpha^2 + \beta^2 + \delta^2 + \varepsilon^2 & \beta\gamma & 0 & 0 \\ -\beta\delta & \beta\gamma & \gamma^2 + \delta^2 & \delta\varepsilon & 0 \\ -\beta\varepsilon & 0 & \delta\varepsilon & \varepsilon^2 - \alpha^2 & -\alpha\beta \\ 0 & 0 & 0 & -\alpha\beta & -\beta^2 - \gamma^2 - \delta^2 - \varepsilon^2 \end{pmatrix}. \quad (20)$$

A simple calculation shows that

$$\begin{cases} Dv_1 = -\frac{1}{2}[(1-\lambda_0)\alpha^2 + (1-\lambda_0)\beta^2 + (1-\lambda_0)\gamma^2 - \lambda_0\delta^2 - \lambda_0\varepsilon^2 + 2c]v_1 - \frac{1}{2}\gamma\delta v_2 + \frac{1}{2}\beta\delta v_3 + \frac{1}{2}\beta\varepsilon v_4 \\ Dv_2 = -\frac{1}{2}\gamma\delta v_1 - \frac{1}{2}[(1-\lambda_0)\alpha^2 + (1-\lambda_0)\beta^2 - \lambda_0\gamma^2 + (1-\lambda_0)\delta^2 + (1-\lambda_0)\varepsilon^2 + 2c]v_2 - \frac{1}{2}\beta\gamma v_3 \\ Dv_3 = \frac{1}{2}\beta\delta v_1 - \frac{1}{2}\beta\gamma v_2 - \frac{1}{2}[-\lambda_0\alpha^2 - \lambda_0\beta^2 + (1-\lambda_0)\gamma^2 + (1-\lambda_0)\delta^2 - \lambda_0\varepsilon^2 + 2c]v_3 + \frac{1}{2}\delta\varepsilon v_4 \\ Dv_4 = \frac{1}{2}\beta\varepsilon v_1 - \frac{1}{2}\delta\varepsilon v_3 - \frac{1}{2}[-(1+\lambda_0)\alpha^2 - \lambda_0\beta^2 - \lambda_0\gamma^2 - \lambda_0\delta^2 + (1-\lambda_0)\varepsilon^2 + 2c]v_4 + \frac{1}{2}\alpha\beta v_5 \\ Dv_5 = \frac{1}{2}\alpha\beta v_4 + \frac{1}{2}[\lambda_0\alpha^2 + (1+\lambda_0)\beta^2 + (1+\lambda_0)\gamma^2 + (1+\lambda_0)\delta^2 + (1+\lambda_0)\varepsilon^2 - 2c]v_5. \end{cases}$$

In this way, (9) is satisfied if and only if

$$\begin{aligned} \alpha[(3-\lambda_0)\alpha^2 + (3-\lambda_0)\beta^2 + (1-\lambda_0)\gamma^2 + (1-\lambda_0)\delta^2 - \lambda_0\varepsilon^2 + 2c] &= 0 \\ \beta[(3-\lambda_0)\alpha^2 + (3-\lambda_0)\beta^2 + (3-\lambda_0)\gamma^2 + (3-\lambda_0)\delta^2 + (3-\lambda_0)\varepsilon^2 + 2c] &= 0 \\ \gamma[(1-\lambda_0)\alpha^2 + (3-\lambda_0)\beta^2 + (3-\lambda_0)\gamma^2 + (3-\lambda_0)\delta^2 + (1-\lambda_0)\varepsilon^2 + 2c] &= 0 \\ \delta[(1-\lambda_0)\alpha^2 + (3-\lambda_0)\beta^2 + (3-\lambda_0)\gamma^2 + (3-\lambda_0)\delta^2 + (3-\lambda_0)\varepsilon^2 + 2c] &= 0 \\ \varepsilon[\lambda_0\alpha^2 + (3-\lambda_0)\beta^2 + (1-\lambda_0)\gamma^2 + (3-\lambda_0)\delta^2 + (3-\lambda_0)\varepsilon^2 + 2c] &= 0 \\ \alpha\beta\gamma &= 0 \\ \alpha\beta\delta &= 0 \\ \alpha\beta\varepsilon &= 0 \\ \alpha\delta\varepsilon &= 0 \\ \gamma\delta\varepsilon &= 0 \\ \beta\gamma\varepsilon &= 0. \end{aligned} \quad (21)$$

Since $\gamma, \varepsilon > 0$, we have $\beta = \delta = 0$. The previous system is transform into Since $\gamma, \varepsilon > 0$, we have $\beta = \delta = 0$. System (24) is equivalent to

$$\begin{aligned} (3-\lambda_0)\alpha^2 + (1-\lambda_0)\gamma^2 - \lambda_0\varepsilon^2 + 2c &= 0 \\ (1-\lambda_0)\alpha^2 + (3-\lambda_0)\gamma^2 + (1-\lambda_0)\varepsilon^2 + 2c &= 0 \\ \lambda_0\alpha^2 + (1-\lambda_0)\gamma^2 + (3-\lambda_0)\varepsilon^2 + 2c &= 0. \end{aligned} \quad (22)$$

Therefore, $\alpha = \varepsilon = \sqrt{2}\gamma$. ■

Corollary 5 An inner product \langle , \rangle on $\mathfrak{g} = A_{5,5}$ is a nilsoliton if and only if $\beta = \delta = 0$ and $\alpha = \varepsilon = \sqrt{2}\gamma$.

Proposition 9 An inner product \langle , \rangle on $\mathfrak{g} = A_{5,3}$ is a Schouten like metric if and only if $\beta = \delta = 0$ and $\varepsilon = \gamma = \frac{\sqrt{3}}{2}\alpha$.

Proof: Let \langle , \rangle be an inner product on $A_{5,3}$. In [5], the Ricci operator is given by

$$Ric_{\langle , \rangle} = -\frac{1}{2} \begin{pmatrix} \alpha^2 + \beta^2 + \gamma^2 + \delta^2 & \delta\varepsilon & 0 & 0 & 0 \\ \delta\varepsilon & \alpha^2 + \beta^2 + \varepsilon^2 & \beta\gamma & 0 & 0 \\ 0 & \beta\gamma & \gamma^2 + \delta^2 + \varepsilon^2 - \alpha^2 & -\alpha\beta & 0 \\ 0 & 0 & -\alpha\beta & -\beta^2 - \gamma^2 & -\delta\gamma \\ 0 & 0 & 0 & -\delta\gamma & -\delta^2 - \varepsilon^2 \end{pmatrix}. \quad (23)$$

A simple calculation shows that

$$\begin{cases} Dv_1 = -\frac{1}{2}[(1-\lambda_0)\alpha^2 + (1-\lambda_0)\beta^2 + (1-\lambda_0)\gamma^2 + (1-\lambda_0)\delta^2 - \lambda_0\varepsilon^2 + 2c]v_1 - \frac{1}{2}\delta\varepsilon v_2 \\ Dv_2 = -\frac{1}{2}\delta\varepsilon v_1 - \frac{1}{2}[(1-\lambda_0)\alpha^2 + (1-\lambda_0)\beta^2 - \lambda_0\gamma^2 - \lambda_0\delta^2 + (1-\lambda_0)\varepsilon^2 + 2c]v_2 - \frac{1}{2}\beta\gamma v_3 \\ Dv_3 = -\frac{1}{2}\beta\gamma v_2 - \frac{1}{2}[-(1+\lambda_0)\alpha^2 - \lambda_0\beta^2 + (1-\lambda_0)\gamma^2 + (1-\lambda_0)\delta^2 + (1-\lambda_0)\varepsilon^2 + 2c]v_3 + \frac{1}{2}\alpha\beta v_4 \\ Dv_4 = \frac{1}{2}\alpha\beta v_3 + \frac{1}{2}[\lambda_0\alpha^2 + (1+\lambda_0)\beta^2 + (1+\lambda_0)\gamma^2 + \lambda_0\delta^2 + \lambda_0\varepsilon^2 - 2c]v_4 + \frac{1}{2}\delta\gamma v_5 \\ Dv_5 = \frac{1}{2}\delta\gamma v_4 + \frac{1}{2}[\lambda_0\alpha^2 + \lambda_0\beta^2 + \lambda_0\gamma^2 + (1+\lambda_0)\delta^2 + (1+\lambda_0)\varepsilon^2 - 2c]v_5. \end{cases}$$

In this way, (9) is satisfied if and only if

$$\begin{aligned} \alpha[(3-\lambda_0)\alpha^2 + (3-\lambda_0)\beta^2 - \lambda_0\gamma^2 - \lambda_0\delta^2 - \lambda_0\varepsilon^2 + 2c] &= 0 \\ \beta[(3-\lambda_0)\alpha^2 + (3-\lambda_0)\beta^2 + (3-\lambda_0)\gamma^2 + (1-\lambda_0)\delta^2 + (1-\lambda_0)\varepsilon^2 + 2c] &= 0 \\ \gamma[-\lambda_0\alpha^2 + (3-\lambda_0)\beta^2 + (3-\lambda_0)\gamma^2 + (3-\lambda_0)\delta^2 + (1-\lambda_0)\varepsilon^2 + 2c] &= 0 \\ \delta[-\lambda_0\alpha^2 + (1-\lambda_0)\beta^2 + (3-\lambda_0)\gamma^2 + (3-\lambda_0)\delta^2 + (3-\lambda_0)\varepsilon^2 + 2c] &= 0 \\ \varepsilon[-\lambda_0\alpha^2 + (1-\lambda_0)\beta^2 + (1-\lambda_0)\gamma^2 + (3-\lambda_0)\delta^2 + (3-\lambda_0)\varepsilon^2 + 2c] &= 0 \\ \alpha\beta\gamma &= 0 \\ \alpha\beta\delta &= 0 \\ \alpha\beta\varepsilon &= 0 \\ \beta\gamma\delta &= 0 \\ \gamma\delta\varepsilon &= 0. \end{aligned} \tag{24}$$

Since $\gamma, \varepsilon > 0$, we have $\beta = \delta = 0$. System (24) is equivalent to

$$\begin{aligned} (3-\lambda_0)\alpha^2 - \lambda_0\gamma^2 - \lambda_0\varepsilon^2 + 2c &= 0 \\ -\lambda_0\alpha^2 + (3-\lambda_0)\gamma^2 + (1-\lambda_0)\varepsilon^2 + 2c &= 0 \\ -\lambda_0\alpha^2 + (1-\lambda_0)\gamma^2 + (3-\lambda_0)\varepsilon^2 + 2c &= 0. \end{aligned} \tag{25}$$

consequently, $\gamma = \varepsilon = \frac{\sqrt{3}}{2}\alpha$. ■

Corollary 6 An inner product \langle, \rangle on $\mathfrak{g} = A_{5,3}$ is a nilsoliton if and only if $\beta = \delta = 0$ and $\varepsilon = \gamma = \frac{\sqrt{3}}{2}\alpha$.

Proposition 10 An inner product \langle, \rangle on $\mathfrak{g} = A_{5,1}$ is an algebraic Schouten soliton if and only if $\beta = 0$ and $\alpha = \gamma$.

Proof: Let \langle, \rangle be an inner product on $A_{5,1}$. By [5],

$$Ric_{\langle, \rangle} = -\frac{1}{2} \begin{pmatrix} \alpha^2 + \beta^2 + \gamma^2 & 0 & 0 & 0 & 0 \\ 0 & \alpha^2 + \beta^2 & \beta\gamma & 0 & 0 \\ 0 & \beta\gamma & \gamma^2 & 0 & 0 \\ 0 & 0 & 0 & -\alpha^2 & -\alpha\beta \\ 0 & 0 & 0 & -\alpha\beta & -\beta^2 - \gamma^2 \end{pmatrix}. \tag{26}$$

Then,

$$\begin{cases} Dv_1 = -\frac{1}{2}[(1-\lambda_0)\alpha^2 + (1-\lambda_0)\beta^2 + (1-\lambda_0)\gamma^2 + 2c]v_1 \\ Dv_2 = -\frac{1}{2}[(1-\lambda_0)\alpha^2 + (1-\lambda_0)\beta^2 + \lambda_0\gamma^2 + 2c]v_2 - \frac{1}{2}\beta\gamma v_3 \\ Dv_3 = -\frac{1}{2}\beta\gamma v_2 - \frac{1}{2}[-\lambda_0\alpha^2 - \lambda_0\beta^2 + (1-\lambda_0)\gamma^2 + 2c]v_3 \\ Dv_4 = \frac{1}{2}[(1+\lambda_0)\alpha^2 + \lambda_0\beta^2 + \lambda_0\gamma^2 - 2c]v_4 + \frac{1}{2}\alpha\beta v_5 \\ Dv_5 = \frac{1}{2}\alpha\beta v_4 + \frac{1}{2}[\lambda_0\alpha^2 + (1+\lambda_0)\beta^2 + (1+\lambda_0)\gamma^2 - 2c]v_5. \end{cases}$$

Ergo, Equation (9) is satisfied if and only if

$$\begin{aligned}
\alpha[(3 - \lambda_0)\alpha^2 + (3 - \lambda_0)\beta^2 + (1 - \lambda_0)\gamma^2 + 2c] &= 0 \\
\beta[(3 - \lambda_0)\alpha^2 + (3 - \lambda_0)\beta^2 + (3 - \lambda_0)\gamma^2 + 2] &= 0 \\
\gamma[(1 - \lambda_0)\alpha^2 + (3 - \lambda_0)\beta^2 + (3 - \lambda_0)\gamma^2 + 2c] &= 0 \\
\alpha\beta\gamma &= 0.
\end{aligned} \tag{27}$$

By the fourth equation in the previous system, we get $\beta = 0$. Hence, system (27) becomes

$$\begin{aligned}
(3 - \lambda_0)\alpha^2 + (1 - \lambda_0)\gamma^2 + 2c &= 0, \\
(1 - \lambda_0)\alpha^2 + (3 - \lambda_0)\gamma^2 + 2c &= 0.
\end{aligned}$$

And then, one obtains $\alpha = \gamma$. ■

Corollary 7 An inner product \langle, \rangle on $\mathfrak{g} = A_{5,1}$ is a nilsoliton if and only if $\beta = 0$ and $\alpha = \gamma$.

Proposition 11 An inner product \langle, \rangle on $\mathfrak{g} = A_{5,2}$ is a Schouten like metric if and only if $\beta = 0$ and $\alpha = \delta = \frac{\sqrt{3}}{2}\gamma$.

Proof: Let \langle, \rangle be an inner product on $A_{5,2}$. According to [5],

$$\text{Ric}_{\langle, \rangle} = -\frac{1}{2} \begin{pmatrix} \alpha^2 + \beta^2 + \gamma^2 + \delta^2 & 0 & 0 & 0 & 0 \\ 0 & \alpha^2 + \beta^2 & \beta\gamma & 0 & 0 \\ 0 & \beta\gamma & \gamma^2 - \alpha^2 & -\alpha\beta & 0 \\ 0 & 0 & -\alpha\beta & \delta^2 - \beta^2 - \gamma^2 & 0 \\ 0 & 0 & 0 & 0 & -\delta^2 \end{pmatrix}. \tag{28}$$

We have

$$\begin{cases} Dv_1 = -\frac{1}{2}[(1 - \lambda_0)\alpha^2 + (1 - \lambda_0)\beta^2 + (1 - \lambda_0)\gamma^2 + (1 - \lambda_0)\delta^2 + 2c]v_1 \\ Dv_2 = -\frac{1}{2}[(1 - \lambda_0)\alpha^2 + (1 - \lambda_0)\beta^2 - \lambda_0\gamma^2 - \lambda_0\delta^2 + 2c]v_2 - \frac{1}{2}\beta\gamma v_3 \\ Dv_3 = -\frac{1}{2}\beta\gamma v_2 - \frac{1}{2}[-(1 + \lambda_0)\alpha^2 - \lambda_0\beta^2 + (1 - \lambda_0)\gamma^2 - \lambda_0\delta^2 + 2c]v_3 + \frac{1}{2}\alpha\beta v_4 \\ Dv_4 = \frac{1}{2}\alpha\beta v_3 - \frac{1}{2}[-\lambda_0\alpha^2 - (1 + \lambda_0)\beta^2 - (1 + \lambda_0)\gamma^2 + (1 - \lambda_0)\delta^2 + 2c]v_4 \\ Dv_5 = \frac{1}{2}[\lambda_0\alpha^2 + \lambda_0\beta^2 + \lambda_0\gamma^2 + (1 + \lambda_0)\delta^2 - 2c]v_5. \end{cases}$$

Thus, Equation (9) is satisfied if and only if

$$\begin{aligned}
\delta[(1 - \lambda_0)\alpha^2 - \lambda_0\beta^2 - \lambda_0\gamma^2 + (3 - \lambda_0)\delta^2 + 2c] &= 0 \\
\beta[(3 - \lambda_0)\alpha^2 + (3 - \lambda_0)\beta^2 + (3 - \lambda_0)\gamma^2 - \lambda_0\delta^2 + 2c] &= 0 \\
\alpha[(3 - \lambda_0)\alpha^2 + (3 - \lambda_0)\beta^2 + (1 - \lambda_0)\gamma^2 - \lambda_0\delta^2 + 2c] &= 0 \\
\gamma[-\lambda_0\alpha^2 + (3 - \lambda_0)\beta^2 + (3 - \lambda_0)\gamma^2 - \lambda_0\delta^2 + 2c] &= 0 \\
\alpha\beta\gamma &= 0 \\
\alpha\beta\delta &= 0.
\end{aligned} \tag{29}$$

Due to the fact that $\alpha, \gamma > 0$, the fifth equation of system (29) is equivalent to $\beta = 0$. Accordingly

$$\begin{aligned}
(3 - \lambda_0)\alpha^2 - \lambda_0\gamma^2 + (1 - \lambda_0)\delta^2 + 2c &= 0, \\
-\lambda_0\alpha^2 + (3 - \lambda_0)\gamma^2 - \lambda_0\delta^2 + 2c &= 0, \\
(1 - \lambda_0)\alpha^2 - \lambda_0\gamma^2 + (3 - \lambda_0)\delta^2 + 2c &= 0.
\end{aligned}$$

Then

$$\begin{aligned}3\alpha^2 - 3\gamma^2 + \delta^2 &= 0, \\ \alpha^2 - 3\gamma^2 + 3\delta^2 &= 0, \\ 2\alpha^2 - 2\delta^2 + 2c &= 0.\end{aligned}$$

Therefore, the first and second equations among the previous three equations give $\alpha = \delta$. Then the last one gives $\alpha = \frac{\sqrt{3}}{2}\gamma$. ■

Corollary 8 *An inner product \langle , \rangle on $\mathfrak{g} = A_{5,2}$ is a nilsoliton if and only if $\beta = 0$ and $\alpha = \delta = \frac{\sqrt{3}}{2}\gamma$.*

References

- [1] M.Boucetta. Curvature of left invariant riemannian metrics on lie groups. *advance in mathematics*, 1976. *Manuscripta Math.* **135**(1-2) (2011) 229-243.
- [2] T. Buttsworth. The prescribe Ricci curvature problem on three-dimensional Lie groups. *Mathematische Nachrichten.* 2018;1-13.
- [3] T. Buttsworth, A. Pulemotov. The Prescribed Ricci Curvature Problem for Homogeneous Metrics. In: Dearthcott O, Tuschmann W, Nikolayevsky Y, Leistner T, Crowley D, eds. *Differential Geometry in the Large*. London Mathematical Society Lecture Note Series. Cambridge University Press; 2020:169-192.
- [4] Calvino-Louzao, E.; Hervella, L.M.; Seoane-Bascoy, J.; Vazquez-Lorenzo, R. Homogeneous Cotton solitons. *J. Phys. A Math. Theor.* 2013, **46**, 285204
- [5] Marius Landry FOKA et al., The prescribed Ricci curvature problem on 5-dimensional nilpotent Lie groups, *International Journal of Geometric Methods in Modern Physics*, <https://doi.org/10.1142/S0219887825500550>.
- [6] Liu, Siyao. "Algebraic Schouten Solitons of Three-Dimensional Lorentzian Lie Groups." *Symmetry* **15.4** (2023): 866.
- [7] T. Hashinaga and H. Tamaru, Three-dimensional solvsolitons and the minimality of the corresponding submanifolds, preprint, arXiv:1501.05513.
- [8] T. Hashinaga, H. Tamaru and K. Terada, Milnor-type theorems for left-invariant Riemannian metrics on Lie groups, *J. Math. Soc. Japan* **68**(2) (2016) 669-684.
- [9] J. Milnor. Curvature of left invariant metrics on lie groups. *advance in mathematics*, 1976.
- [10] MOGHADDAM, Hamid Reza Salimi. On the classification of five-dimensional nilsolitons. arXiv preprint arXiv:1912.13322, 2019.
- [11] A. Pulemotov. Metrics with prescribed Ricci curvature near the boundary of a manifold. *Math.*
- [12] A. Takahara and H. Tamaru, The space of left-invariant metrics on a Lie group up to isometry and scaling, *Manuscripta Math.* **135**(1-2) (2011) 229-243.
- [13] WEARS, Thomas H. On algebraic solitons for geometric evolution equations on three-dimensional Lie groups. 2016.