

# Kendall and Spearman Rank Correlations for Skew-Elliptical Copulas

Ye Lu\*

December 2024

## Abstract

In this paper, we derive explicit formulas of Kendall's tau and Spearman's rho rank correlations for two general classes of skew-elliptical copulas: normal location-scale mixture copulas and skew-normal scale mixture copulas, which encompass some widely used skew- $t$  and skew-normal copulas. These formulas establish mappings from copula parameters to rank correlation coefficients, potentially facilitating robust rank-based estimation of skew-elliptical copula models. Additionally, we investigate and compare the impacts of asymmetry parameters on the properties of both rank correlations within these two classes of skew-elliptical models. Notably, the introduction of asymmetry in normal location-scale mixture copulas restricts the attainable range of the rank correlations from  $[-1, 1]$  – as observed under elliptical symmetry – to a strict subset of  $[-1, 1]$ . In contrast, the entire interval  $[-1, 1]$  remains attainable for skew-normal scale mixture copulas.

---

\*School of Economics, University of Sydney. Email: ye.lu1@sydney.edu.au.

# 1 Introduction

Rank correlations are measures of association between two variables that play a crucial role in many empirical methods. Compared to Pearson’s linear correlation, rank correlations are invariant under potentially nonlinear, rank-preserving transformations of the measurement scale (such as log transformations), and they are robust to outliers and violations of normality; see [Kruskal \(1958\)](#) for a survey. These attributes make rank correlations highly desirable measures of dependence in practice. Among various rank correlation measures, Kendall’s tau, denoted by  $\tau$ , and Spearman’s rho, denoted by  $\rho_S$ , are the most commonly employed.<sup>1</sup>

Since rank correlations depend solely on the copula of a bivariate distribution, another practical benefit is that they can be used to fit copula-based models, which are widely used in a variety of applications, to empirical data. When the mapping from copula parameters to a rank correlation coefficient is explicitly known, the classical method of moments (MM) or generalized method of moments (GMM) can be employed to estimate the copula parameter using empirical rank correlation estimates.<sup>2</sup> A simple yet widely used application of this approach involves estimating the (pseudo-)correlation coefficient<sup>3</sup>  $\rho$  by inverting the well-known relation  $\tau = \frac{2}{\pi} \arcsin \rho$  for the class of *elliptical* copulas, using an estimate of  $\tau$ . Compared to the standard sample correlation, the rank-based estimator is more robust to various marginal distribution characteristics, such as infinite variance and heteroskedasticity.<sup>4</sup>

In the presence of asymmetry – a paramount feature in empirical data across many applications – several commonly used Archimedean copulas, such as Clayton and Gumbel copulas, along with their non-exchangeable versions, can effectively model the required

---

<sup>1</sup>The so-called *quadrant measure* is another rank correlation measure discussed by [Kruskal \(1958\)](#). In the bivariate Gaussian case, it coincides with Kendall’s tau. Since the quadrant measure is less commonly used than Kendall’s and Spearman’s rank correlations, we focus on the latter two in this paper.

<sup>2</sup>If the explicit form of such a mapping is unavailable, the simulated method of moments (SMM) can be used instead; see, for example, the approach considered by [Oh & Patton \(2013\)](#).

<sup>3</sup>Although the usual correlation coefficient of a bivariate elliptical distribution may not exist due to infinite variances, the pseudo-correlation coefficient is always well-defined.

<sup>4</sup>For discussions and examples, see, e.g., [McNeil et al. \(2005, Section 3.3.4\)](#) in simpler settings, and [Hansen & Luo \(2023\)](#) for an application in financial econometrics to construct an estimator of realized correlation robust to time-varying volatility.

asymmetric dependence. The rank correlation formulas for these copulas have been derived in the literature. However, the application of Archimedean copulas is significantly limited to two to three dimensions, as extending them to higher dimensions is considerably more complex. For further discussion, see [McNeil et al. \(2005, Section 5.4.3\)](#) and [McNeil & Nešlehová \(2009\)](#).

In contrast, elliptical copulas, including Gaussian and Student's  $t$  copulas, are implicit copulas derived from elliptical distributions via Sklar's theorem ([Sklar, 1959](#)), and thus have straightforward high-dimensional extensions. The *asymmetric* generalizations of elliptical distributions are often referred to as *skew-elliptical* distributions and their corresponding copulas are known as skew-elliptical copulas. Skew-elliptical copulas can accommodate highly flexible forms of asymmetry while maintaining tractability in higher dimensions. Consequently, they have become increasingly popular tools in high-dimensional distributional modeling.

Among all developments, two classes of skew-elliptical distributions – arising from different methods of modulating elliptical symmetry – have garnered the most attention. The first class is obtained by mixing multivariate normal distributions with different locations as well as different scales, referred to as *normal location-scale mixtures*. In these models, the location-mixture coefficients serve as asymmetry parameters. A notable subclass within this category is the generalized hyperbolic (GH) distributions; see, for example, [Blaesild & Jensen \(1981\)](#). A further special case, known as the GH skew- $t$  distribution, was highlighted by [Demarta & McNeil \(2005\)](#), and has since found extensive applications. Some important applications in financial econometrics include [Aas & Haff \(2006\)](#), [Christoffersen et al. \(2012\)](#), [Creal & Tsay \(2015\)](#), [Lucas et al. \(2017\)](#), and [Oh & Patton \(2023\)](#), among others.

The second class of skew-elliptical distributions arises from mixing multivariate skew-normal distributions with different scales, referred to as *skew-normal scale mixtures*.<sup>5</sup> Similar to the first class of models, asymmetry is introduced through a vector of skewness parameters whose dimension matches that of the model. Within this category, the

---

<sup>5</sup>As explained by [Azzalini \(2014, Section 6.1.3\)](#), the skew-normal scale mixture is closely related to the *modulated elliptical distribution* introduced by [Branco & Dey \(2001\)](#). Specifically, if the parent elliptical distribution of the latter is a normal scale mixture with a particular mixing distribution, then the modulated elliptical distribution is a skew-normal scale mixture using the same mixing distribution.

skew-normal distribution itself and the so-called AC skew- $t$  distribution – named after [Azzalini & Capitanio \(2003\)](#) – are the most significant special cases. Their applications have been advanced by [Kollo & Pettere \(2010\)](#), [Azzalini \(2014\)](#), [Yoshida \(2018\)](#), and [Deng et al. \(2024\)](#), among others.

Despite the widespread use of skew-elliptical distributions and their copulas, the explicit expressions and properties of their rank correlations remain largely unexplored.<sup>6</sup> Asymmetry in these distributions arises from an additional skewness (or asymmetry) vector, prompting the natural question of how this skewness vector affect rank correlations and their properties across different classes of skew-elliptical models. Moreover, the mapping from copula parameters – including both pseudo-correlations and skewness parameters – to rank correlations could enable rank-based estimation for skew-elliptical copula models.

In this paper, we derive explicit formulas for Kendall’s tau and Spearman’s rho in terms of copula parameters for the aforementioned two general classes of skew-elliptical copulas: the normal location-scale mixture copulas and skew-normal scale mixture copulas. We demonstrate that, for both models, the rank correlations can be expressed as expectations of mixtures of zero-mean normal cumulative distribution functions (cdf’s) and can be efficiently evaluated using quasi-Monte Carlo (QMC) integration methods. This enables a clear mapping from copula parameters to each type of rank correlation. Additionally, we specialize our results to derive Kendall’s tau and Spearman’s rho for notable special cases, specifically the GH skew- $t$ , skew-normal and AC skew- $t$  copulas.

Building on these formulas, we carefully examine the effects of the asymmetry/skewness parameters on both rank correlations, and how these effects cause deviations from the elliptical case when asymmetry is present. For both classes of bivariate skew-elliptical copulas, we demonstrate that Kendall’s tau and Spearman’s rho are symmetric in the components of the skewness vector and invariant under a sign change of the skewness vector. Furthermore, in the single-skew case – where only one of the skewness parameter is nonzero – both Kendall’s tau and Spearman’s rho are odd functions of the pseudo-correlation coefficient, a property also observed for elliptical copulas. Additionally, increasing the level of asymmetry decreases the magnitude of both rank correlations in the single-skew case.

---

<sup>6</sup>To the best of my knowledge, the only research in this area is by [Heinen & Valdesogo \(2022\)](#), who studied Kendall and Spearman rank correlations for the bivariate skew-normal distribution.

However, we also found some significant differences in the properties of Kendall’s tau and Spearman’s rho between the two classes of skew-elliptical copulas. For example, the effects of asymmetry on the rank correlations in the equi-skew case, where the skewness parameters take the same value, differ in direction for the two classes of models: increasing the equi-skewness parameter *increases* the rank correlations for normal location-scale mixture copulas, but *decreases* the rank correlations for skew-normal scale mixture copulas.

The most notable difference pertains to the attainable range of Kendall’s tau and Spearman’s rho in the presence of asymmetry. Specifically, when using normal location-scale copulas – the class of skew-elliptical copulas to which GH skew- $t$  copulas belong – the attainable ranges for both rank correlations become *strict* subsets of the interval  $[-1, 1]$ . This sharply contrasts with the elliptical case, where the full interval  $[-1, 1]$  is attainable for rank correlations. Additionally, unlike for elliptical copulas, Kendall’s tau and Spearman’s rho are no longer always  $-1$  or  $1$  when the pseudo-correlation coefficient equals  $-1$  or  $1$ , respectively. Conversely, these properties of rank correlations for elliptical copulas are preserved when asymmetry is introduced in the other class of skew-elliptical copulas – the skew-normal scale mixture copulas.

The rest of the paper is organized as follows. After providing notation, Section 2 gives the definitions of Kendall’s tau and Spearman’s rho, and reviews the known results of these rank correlations in the context of bivariate elliptical copulas. Section 3 formally defines the two general classes of skew-elliptical distributions and their associated copulas that are the focus of this study. Section 4 and Section 5 present the main results of this paper, namely Kendall’s and Spearman’s rho and their properties for the two previously defined classes of skew-elliptical copulas. A thorough comparison of the results between these two classes is also provided. Section 6 concludes the paper, and the mathematical proofs are contained in the Appendix.

Before concluding this section, we clarify the notation used throughout the paper. In general, we use  $\rho$  to denote a correlation coefficient,  $P$  a correlation matrix, and  $\varrho$  the *correlation operator*. In particular, when the correlation operator operates on a  $d \times d$  covariance matrix  $\Sigma = [\sigma_{ij}]$ , written  $\varrho(\Sigma)$ , it returns  $\text{diag}(\Sigma)^{-1/2} \Sigma \text{diag}(\Sigma)^{-1/2}$ , the corresponding correlation matrix; when  $d = 2$ ,  $\varrho(\Sigma)$  gives the pseudo-correlation coefficient  $\sigma_{12}/\sqrt{\sigma_{11}\sigma_{22}}$ . Moreover, when  $\varrho$  operates on a correlation coefficient  $\rho \in (-1, 1)$ , written  $\varrho(\rho)$ , it returns

a  $2 \times 2$  correlation matrix with off-diagonal entries  $\rho$ . For the common probability distributions, we use  $N$  and  $IG$  to denote the normal distribution and inverse-gamma distribution, respectively;  $U(0, 1)^d$  the uniform distribution on the  $d$ -dimensional hypercube;  $\text{Ga}(\alpha, \beta)$  the gamma distribution whose density at  $x > 0$  is given by  $\beta^\alpha \Gamma^{-1}(\alpha) x^{\alpha-1} \exp(-\beta x)$  for  $\alpha > 0, \beta > 0$ ; and  $\text{IG}(\alpha, \beta)$  the inverse-gamma distribution whose density at  $x > 0$  is given by  $\beta^\alpha \Gamma^{-1}(\alpha) x^{-\alpha-1} \exp(-\beta/x)$  for  $\alpha > 0, \beta > 0$ . We denote by  $\phi(x)$  and  $\Phi(x)$ , for  $x \in \mathbb{R}$ , the pdf and cdf of the univariate standard normal distribution, respectively. The pdf of the univariate standard skew-normal distribution with skewness parameter  $\alpha \in \mathbb{R}$  is  $\phi^s(x; \alpha) := 2\phi(x)\Phi(\alpha x)$ . For a  $d \times d$  covariance matrix  $\Sigma$ , we denote by  $\phi_d(x; \Sigma)$  and  $\Phi_d(x; \Sigma)$ , for  $x \in \mathbb{R}^d$ , the pdf and cdf of the  $d$ -dimensional normal distribution  $N(0, \Sigma)$ . When  $d = 2$  and  $\rho$  is a correlation coefficient, we use the shorthand notation  $\phi_2(x; \rho)$  and  $\Phi_d(x; \rho)$  for  $\phi_2(x; \varrho(\rho))$  and  $\Phi_2(x; \varrho(\rho))$ , respectively. Finally, unless specified otherwise, all vectors stated in this paper are assumed to be column vectors.

## 2 Kendall and Spearman rank correlations

As the primary measures of rank correlation, Kendall's tau and Spearman's rho serve as indicators of concordance for pairs of random variables. Their formal definitions are provided below.

**Definition 1.** *Let  $X_1$  and  $X_2$  be two random variables with marginal distributions  $F_1$  and  $F_2$ , respectively. Kendall's tau for  $X_1$  and  $X_2$  is defined as*

$$\tau = \mathbb{E} \text{sign}(X_1 - X_1^*)(X_2 - X_2^*),$$

where  $(X_1^*, X_2^*)$  is an independent copy<sup>7</sup> of  $(X_1, X_2)$ . Spearman's rho is defined by


$$\rho_S = \text{Corr}(F_1(X_1), F_2(X_2)).$$

It follows directly from the definitions that Kendall's tau and Spearman's rho are symmetric measures taking values in the interval  $[-1, 1]$ . Furthermore, if  $X_1$  and  $X_2$  have

---

<sup>7</sup>Two random vectors are independent copies of each other if they are independent and share the same joint distribution.

Figure 1: Kendall’s tau for elliptical copulas and Spearman’s rho for Student’s  $t$  and Gaussian copulas.



Notes: The left panel presents the mapping from the pseudo-correlation coefficient  $\rho$  to Kendall’s tau, as shown in equation (2.1), for all bivariate elliptical copulas. The dark solid line in the right panel depicts the mapping from  $\rho$  to Spearman’s rho, as given in equation (2.2), specifically for bivariate Gaussian copulas. Additionally, the red and yellow dashed lines in the right panel illustrate the relationships between Spearman’s rho and  $\rho$  for bivariate Student’s  $t$  copulas with degrees of freedom 1 and 4, respectively. The corresponding formulas can be found in Heinen & Valdesogo (2020, Proposition 4).

continuous margins and a unique copula  $C$ , then both rank correlations depend solely on their copula:

$$\tau = 4 \int_0^1 \int_0^1 C(u_1, u_2) dC(u_1, u_2) - 1 \quad \text{and} \quad \rho_S = 12 \int_0^1 \int_0^1 C(u_1, u_2) du_1 du_2 - 3.$$

See, for example, McNeil et al. (2005, Proposition 5.29). For this reason, we focus on copulas in our discussion.

Kendall’s tau for the bivariate Gaussian copula with correlation coefficient  $\rho$  is well established and is given by

$$\tau = \frac{2}{\pi} \arcsin \rho. \tag{2.1}$$

This theoretical relationship extends more generally to all bivariate elliptical copulas with a pseudo-correlation coefficient  $\rho$  (see, for example, McNeil et al., 2005, Proposition 5.37). The left panel of Figure 1 illustrates the graph of this one-to-one mapping from  $\rho$  to  $\tau$ .

There is also a well-known relationship between Spearman’s rho and the linear correlation parameter  $\rho$  for the bivariate Gaussian copula:

$$\rho_S = \frac{6}{\pi} \arcsin \frac{\rho}{2}. \quad (2.2)$$

However, this relationship does not extend to other non-Gaussian elliptical copulas. In a recent paper, [Heinen & Valdesogo \(2020\)](#) demonstrate that Spearman’s rho for bivariate scale mixtures of normals – a significant class of elliptical distributions that includes Student’s  $t$  distribution – can be expressed as a correlation mixture of Spearman’s rho for the Gaussian case. They also show that within this class, for any value of the correlation parameter, Spearman’s rho for the Gaussian case has the greatest absolute value.

The right panel of [Figure 1](#) presents the mappings from the (pseudo-)correlation coefficient  $\rho$  to Spearman’s rank correlation  $\rho_S$  for the Gaussian copula (depicted as a dark solid line) and for two  $t$  copulas with degrees of freedom 1 and 4 (depicted as colored dashed lines). It is evident that the graph of Spearman’s rho as a function of  $\rho$  for the Gaussian copula closely aligns with the 45-degree line,<sup>8</sup> and the graph for the  $t$  copula with 4 degrees of freedom is already very close to that for the Gaussian case.

As mentioned in the Introduction, the primary practical application of the theoretical relationship between rank correlations and copula parameter(s) is to provide rank-based estimation procedures using the method of moments (MM) or generalized method of moments (GMM). By substituting empirical rank correlation values into these relationships, moment equations can be derived to estimate some or all copula parameters. This approach offers at least two advantages. First, since rank correlations are robust to outliers and model deviations, the corresponding moment estimators inherit this robustness. Second, this approach eliminates the need to estimate marginal distributions when used to fit (or partially fit) copulas to empirical data, thereby simplifying the modeling process and reducing potential sources of estimation error.

In contrast to the extensive applications of Kendall’s and Spearman’s rank correlations for elliptically symmetric copulas, much less is known about their counterparts for asym-

---

<sup>8</sup>For the Gaussian copula, Spearman’s rank correlation  $\rho_S$  is sometimes used to approximate the linear correlation  $\rho$ ; see e.g. [Iman & Conover \(1982\)](#). The approximation error satisfies  $|\rho_S - \rho| = |6 \arcsin(\rho/2)/\pi - \rho| \leq \pi^{-1}(\pi - 3) |\rho| \leq 0.0181$ .



metric generalizations of elliptical copulas. The only notable development in this area is by Heinen & Valdesogo (2022), who derived Kendall’s tau and Spearman’s rho in terms of the correlation and skewness parameters for the bivariate skew normal distribution (Azzalini & Dalla Valle, 1996; Azzalini & Capitanio, 1999).

### 3 Two classes of skew-elliptical copulas

Skew-elliptical copulas are implicit copulas derived from skew-elliptical distributions, which are asymmetric generalizations of elliptical distributions. This paper focuses on two main classes of skew-elliptical distributions and their associated copulas. The first class, introduced in Section 3.1, is based on location and scale mixtures of normal distributions. The second class, presented in Section 3.2, originates from scale mixtures of skew-normal distributions.

#### 3.1 Normal location-scale mixtures

Scale mixtures of normal distributions constitute a significant subclass of elliptical distributions. A common approach for introducing asymmetry is through an extension to location-scale mixtures, formally defined as follows.

**Definition 2.** *Let  $d \geq 1$  be a positive integer. Given  $\mu, \beta \in \mathbb{R}^d$ , a  $d \times d$  positive definite matrix  $\Sigma$ , and a univariate distribution  $F$  on  $(0, \infty)$ , a  $d$ -dimensional random vector  $X$  is said to be a location-scale mixture of normals, denoted  $X \sim \text{MN}(\mu, \Sigma, \beta, F)$ , if  $X$  can be expressed as*

$$X = \mu + W\beta + \sqrt{W}Z,$$

*where  $W$  and  $Z$  are independent random variables, with  $W \sim F$  and  $Z \sim \text{N}(0, \Sigma)$ .*

The distribution of  $X$  in the above definition reduces to the scale mixture of normals – hence becomes elliptically symmetric – if and only if  $\beta = 0$ . Consequently, the vector  $\beta$  in the location mixture component acts as the *skewness/asymmetry* parameter.

A notable subclass of this class of skew-elliptical distributions is the *generalized hyperbolic* (GH) distribution introduced by [Barndorff-Nielsen \(1977\)](#) for the univariate case and extended by [Blaesild & Jensen \(1981\)](#) for the multivariate case. This subclass is derived by selecting the mixing distribution  $F$  to be a *generalized inverse Gaussian* (GIG) distribution ([Barndorff-Nielsen & Halgreen, 1977](#)), denoted by  $N^-(\lambda, \chi, \psi)$  with parameters  $\lambda \in \mathbb{R}, \chi, \psi \geq 0$ . The pdf of the GIG distribution is given by

$$f(x; \lambda, \chi, \psi) = \frac{(\psi/\chi)^{\lambda/2}}{2K_\lambda(\sqrt{\psi\chi})} x^{\lambda-1} \exp\left(-\frac{\psi x + \chi/x}{2}\right) \quad (x > 0), \quad (3.1)$$

where  $K_\lambda$  is a modified Bessel function of the third kind (also referred to as the modified Bessel function of the second kind, or the modified Hankel function) with index  $\lambda$ . The parameters must satisfy (i)  $\chi > 0, \psi \geq 0$  if  $\lambda < 0$ ; (ii)  $\chi > 0, \psi > 0$  if  $\lambda = 0$ ; and (iii)  $\chi \geq 0, \psi > 0$  if  $\lambda > 0$ . In general, if  $W \sim N^-(\lambda, \chi, \psi)$ , then  $W^{-1} \sim N^-(-\lambda, \psi, \chi)$ . The GIG density (3.1) contains the gamma and inverse-gamma densities as special *limit* cases, corresponding to  $\chi = 0$  and  $\psi = 0$ , respectively.<sup>9</sup>

The GH family is highly flexible, encompassing many known distributions as special cases. For example, when  $\lambda = (d+1)/2$ , it corresponds to a  $d$ -dimensional hyperbolic distribution. When  $\lambda > 0, \psi > 0$  and  $\chi = 0$ , the distribution is known as the variance-gamma, generalized Laplace, or Bessel function distribution. Setting  $\lambda = -1/2$  yields the normal inverse gamma (NIG) distribution.

Particular attention is given to the case where  $\lambda = -\nu/2, \chi = \nu$  for some  $\nu > 0$  and  $\psi = 0$ . Under this parameter setting, the mixing distribution is  $N^-(-\nu/2, 0, \nu) \stackrel{d}{=} \text{IG}(\nu/2, \nu/2)$ , and the GH family reduces to a class of asymmetric or skew  $t$  distributions, known as the *GH skew- $t$*  distribution. This class of skew- $t$  distributions and their associated copulas were first introduced by [Demarta & McNeil \(2005\)](#) and has since been widely applied in risk management and financial econometrics, as highlighted in the Introduction.

Definition 3 below defines the normal location-scale mixture copula. Since standardizing the margins involves strictly increasing transformations that do not alter the copula, the copula of  $\text{MN}(\mu, \Sigma, \beta, F)$  is identical to that of  $\text{MN}(0, \varrho(\Sigma), \text{diag}(\Sigma)^{-1/2}\beta, F)$ . As a result,

---

<sup>9</sup>These limiting cases can be evaluated using the asymptotic relations  $K_\lambda(x) \sim \Gamma(|\lambda|)2^{|\lambda|-1}x^{-|\lambda|}$  as  $x \rightarrow 0^+$  for  $\lambda \neq 0$ . The fact that  $K_\lambda(x) = K_{-\lambda}(x)$  is also useful. In this way, it can be verified that  $N^-(\lambda, \chi, 0) \stackrel{d}{=} \text{IG}(-\lambda, \chi/2)$  for  $\lambda < 0, \chi > 0$ , and  $N^-(\lambda, 0, \psi) \stackrel{d}{=} \text{Ga}(\lambda, \psi/2)$  for  $\lambda > 0, \psi > 0$ .

the normal location-scale mixture copula is fully characterized by a correlation matrix, a skewness vector<sup>10</sup>, and a mixing distribution.

**Definition 3.** Let  $d \geq 2$  be a positive integer. Given  $\beta \in \mathbb{R}^d$ , a  $d \times d$  correlation matrix  $P$  and a univariate distribution  $F$  on  $(0, \infty)$ , define the normal location-scale mixture copula  $C_{\text{mn}}(P, \beta, F)$  as the copula of the normal location-scale mixture  $\text{MN}(0, P, \beta, F)$ . In the case  $d = 2$ , we denote it by  $C_{\text{mn}}(\rho, \beta, F)$ , where  $\rho$  is the pseudo-correlation coefficient.

If the mixing distribution  $F$  in Definition 3 is specified as  $\text{IG}(\nu/2, \nu/2)$  for some  $\nu > 0$ , the resulting copula is the so-called GH skew- $t$  copula with degrees of freedom  $\nu$ . The bivariate GH skew- $t$  copula is written as  $C_{\text{GHt}}(\rho, \beta, \nu)$ .

## 3.2 Skew-normal scale mixtures

An alternative popular approach to introducing asymmetry in elliptical distributions is to consider the *skew-normal* distribution and its scale mixtures; see, for example, the discussion in [Azzalini \(2014, Sections 5 and 6\)](#). Following [Azzalini \(2014, Section 5.1.1\)](#), we define the multivariate skew-normal distribution as follows.

**Definition 4.** A  $d$ -dimensional normalized skew-normal distribution  $\text{SN}(0, P, \alpha)$  with a positive definite correlation matrix  $P$  and skewness vector  $\alpha \in \mathbb{R}^d$  is defined by its pdf:

$$2 \phi_d(x; P) \Phi(\alpha^\top x) \quad (x \in \mathbb{R}^d).$$

Given a location vector  $\mu \in \mathbb{R}^d$  and a positive definite covariance matrix  $\Sigma$ , we say a variable  $Y$  follows a  $\text{SN}(\mu, \Sigma, \alpha)$  distribution, if it can be written as  $Y = \mu + D^{1/2}X$ , where  $D := \text{diag}(\Sigma)$  and  $X \sim \text{SN}(0, \varrho(\Sigma), \alpha)$ .

When the skewness vector  $\alpha$  is zero, the skew-normal distribution  $\text{SN}(0, P, 0)$  simplifies to the normal distribution  $\text{N}(0, P)$ . Moreover, similar to the skewness vector  $\beta$  in the normal location-scale mixture model, the skewness vector  $\alpha$  given in Definition 4 is *unrestricted*,

---

<sup>10</sup>The skewness vector in the normal location-scale mixture copula is not necessarily identical to the skewness vector in the original normal location-scale mixture distribution where the copula is derived from. It is instead a scaled version.

allowing its value to vary freely in  $\mathbb{R}^d$ . This flexibility is a beneficial feature of the  $(P, \alpha)$ -parameterization of the normalized skew-normal distribution. However, it is worth noting that although the skew-normal distribution is closed under marginalization, the marginal distribution of the  $i$ -th component of  $\text{SN}(0, P, \alpha)$  is *not* simply  $\text{SN}(0, 1, \alpha_i)$ , where  $\alpha_i$  denotes the  $i$ -th component of  $\alpha$ . An alternative parameterization with a simpler transformation of parameters under marginalization is presented in the following remark.

**Remark 1.** *The normalized skew-normal distribution in Definition 4 can alternatively be parameterized by  $(P, \delta)$ , where the  $d$ -dimensional vector  $\delta$  should satisfy  $\delta^\top P^{-1} \delta < 1$  and each entry of  $\delta$  lies in  $(-1, 1)$ . There is a one-to-one correspondence between  $(P, \alpha)$  and  $(P, \delta)$ , as the skewness vectors are related by*

$$\delta = \frac{P\alpha}{\sqrt{1 + \alpha^\top P\alpha}} \quad \text{and} \quad \alpha = \frac{P^{-1}\delta}{\sqrt{1 - \delta^\top P^{-1}\delta}}. \quad (3.2)$$

Clearly,  $\alpha = 0$  if and only if  $\delta = 0$ , and changing the sign of one skewness vector induces the same sign change in the other. Under the  $(P, \delta)$  parameterization, the  $i$ -th marginal distribution of  $\text{SN}(0, P, \delta)$  is simply  $\text{SN}(0, 1, \delta_i)$ , where  $\delta_i$  is the  $i$ -th component of  $\delta$ . Applying the second mapping given in (3.2) for the univariate case, we can deduce that the corresponding marginal skewness parameter is  $\alpha_i^\dagger = \delta_i / \sqrt{1 - \delta_i^2}$ , which is not necessarily equal to the  $i$ -th component of  $\alpha$ .

Next, we define the normal scale mixtures, which arise from mixing multivariate skew-normal distributions with different scales.

**Definition 5.** *Given  $\mu, \alpha \in \mathbb{R}^d$ , a  $d \times d$  positive definite matrix  $\Sigma$ , and a univariate distribution  $F$  on  $(0, \infty)$ , a  $d$ -dimensional random vector  $X$  is said to be a scale mixture of skew-normals, denoted  $X \sim \text{MSN}(\mu, \Sigma, \alpha, F)$ , if  $X$  can be expressed as*

$$X = \mu + \sqrt{W}Z,$$

where  $W$  and  $Z$  are independent random variables, with  $W \sim F$  and  $Z \sim \text{SN}(0, \Sigma, \alpha)$ .

As highlighted in the Introduction, within this class of skew-elliptical distributions, particular emphasis is often placed on a notable subclass known as the *AC skew- $t$*  distribution<sup>11</sup>,

---

<sup>11</sup>The AC skew- $t$  distribution, named after [Azzalini & Capitanio \(2003, Section 4\)](#), was independently

which arises when the mixing distribution  $F$  is chosen as the inverse-gamma distribution  $\text{IG}(\nu/2, \nu/2)$  with  $\nu > 0$ .

Again, since standardizing the margins does not alter the copula, the copula of  $X$ , where  $X \sim \text{MSN}(\mu, \Sigma, \alpha, F)$ , is the same as the copula of  $D^{-1/2}(X - \mu)$ , with  $D = \text{diag}(\Sigma)$ . By the representation in Definition 5 and the normalization of skew-normal distributions, we have  $D^{-1/2}(X - \mu) \stackrel{d}{=} \sqrt{W}D^{-1/2}\text{SN}(0, \Sigma, \alpha) \stackrel{d}{=} \sqrt{W}\text{SN}(0, \varrho(\Sigma), \alpha) \stackrel{d}{=} \text{MSN}(0, \varrho(\Sigma), \alpha, F)$ . Therefore, the skew-normal scale mixture copula is fully characterized by a correlation matrix, a skewness vector, and a mixing distribution. Its formal definition is as follows.

**Definition 6.** *Let  $d \geq 2$  be a positive integer. Given  $\alpha \in \mathbb{R}^d$ , a  $d \times d$  correlation matrix  $P$  and a univariate distribution  $F$  on  $(0, \infty)$ , define the skew-normal scale mixture copula  $C_{\text{msn}}(P, \alpha, F)$  to be the copula of the scale mixture of skew-normals  $\text{MSN}(0, P, \alpha, F)$ . In the case  $d = 2$ , we denote it by  $C_{\text{msn}}(\rho, \alpha, F)$  where  $\rho$  is the pseudo-correlation coefficient.*

Because the one-to-one correspondence between  $\alpha$  and  $\delta$  in (3.2) (see Remark 1), the copula  $C_{\text{msn}}(P, \alpha, F)$  can be equivalently represented by  $C_{\text{msn}}(P, \delta, F)$ . The alternative parameterization of skewness by  $\delta$  is instrumental for analyzing the rank correlations in bivariate skew-normal scale mixture copulas, as discussed in Section 5. Accordingly, we now focus on the bivariate setting of Remark 1 and summarize the key points in the following remark.

**Remark 2.** *In the bivariate case, by substituting  $\varrho(\rho)$  for  $P$  in (3.2), we obtain the following relations between the skewness vectors  $\alpha = (\alpha_1, \alpha_2)^\top$  and  $\delta = (\delta_1, \delta_2)^\top$ :*

$$\delta_i = \frac{\alpha_i + \rho\alpha_{-i}}{\sqrt{1 + \alpha_1^2 + \alpha_2^2 + 2\rho\alpha_1\alpha_2}} \quad \text{and} \quad \alpha_i = \frac{(\delta_i - \rho\delta_{-i})/\sqrt{1 - \rho^2}}{\sqrt{1 - \rho^2 - \delta_1^2 - \delta_2^2 + 2\rho\delta_1\delta_2}}, \quad (3.3)$$

for  $i = 1, 2$  and  $\rho \in (-1, 1)$ . By convention,  $-i$  denotes the index other than  $i$ . From (3.3), it follows that  $\alpha_1 = \alpha_2$  if and only if  $\delta_1 = \delta_2$ . However, if one of  $\alpha_1$  and  $\alpha_2$  is zero, it need not be the case that one of  $\delta_1$  and  $\delta_2$  is zero – and vice versa. As noted in Remark 1, the skewness vector  $\delta$  must satisfy  $\delta^\top \varrho(\rho)^{-1} \delta < 1$ , which boils down to  $\delta_1^2 + \delta_2^2 - 2\rho\delta_1\delta_2 < 1 - \rho^2$ , for all  $\rho \in (-1, 1)$ .

---

proposed by Gupta (2003). For a detailed discussion of the properties of this class of multivariate skew- $t$  distributions and a historical perspective, see Azzalini (2014, Section 6.2).

As a degenerated special case, the skew-normal copula is nested in this class of skew-elliptical copulas. The bivariate skew-normal copula is denoted by  $C_{\text{sn}}(\rho, \alpha)$ . On the other hand, if the mixing distribution  $F$  is specified as  $\text{IG}(\nu/2, \nu/2)$  for some  $\nu > 0$ , the resulting copula is the AC skew- $t$  copula with degrees of freedom  $\nu$ . Its bivariate form is denoted by  $C_{\text{ACt}}(\rho, \alpha, \nu)$ .

## 4 Rank correlations of normal location-scale mixtures

In this section, we start by presenting the formulas and properties of Kendall's and Spearman's rank correlations for bivariate normal location-scale mixture copulas, defined in Definition 3. Next, we specialize these results to the GH skew- $t$  copula, an important and widely applicable special case. In the last subsection, we focus on the equi-skew and single-skew settings where we highlight their particular relevance.

### 4.1 Formulas of Kendall's tau and Spearman's rho

The following theorem gives the formulas of Kendall's tau and Spearman's rho for the bivariate normal location-scale mixture copula defined in Definition 3. In the theorem, we use  $\Phi_2(x_1, x_2; \rho)$  to denote the cdf of the bivariate normal distribution  $\text{N}(0, \varrho(\rho))$ .

**Theorem 1.** *Suppose  $X_1$  and  $X_2$  are two random variables with continuous margins and normal location-scale mixture copula  $C_{\text{mn}}(\rho, \beta, F)$  where  $\rho \in [-1, 1]$ ,  $\beta = (\beta_1, \beta_2) \in \mathbb{R}^2$ , and  $F$  is a univariate continuous distribution on  $(0, \infty)$ . Then,*

(i) *Kendall's tau of  $X_1$  and  $X_2$  is given by*

$$\tau(\rho, \beta, F) = 4 \mathbb{E} \Phi_2(\beta_1 V, \beta_2 V; \rho) - 1, \quad (4.1)$$

where  $V = (W^* - W)/\sqrt{W^* + W}$  with  $W, W^* \sim i.i.d. F$ ;

(ii) *Spearman's rho of  $X_1$  and  $X_2$  is given by*

$$\rho_S(\rho, \beta, F) = 12 \mathbb{E} \Phi_2(\beta_1 V_1, \beta_2 V_2; \rho V_3) - 3, \quad (4.2)$$

where  $V_i = (W_i - W_3)/\sqrt{W_i + W_3}$ , for  $i = 1, 2$ , and  $V_3 = W_3/\sqrt{(W_1 + W_3)(W_2 + W_3)}$ , with  $W_1, W_2, W_3 \sim i.i.d. F$ .

Both Kendall's tau and Spearman's rho, as described in equations (4.1) and (4.2), are expressed in terms of expectations of mixtures of the bivariate normal cdf, where the distributions of the mixing variables  $V$  and  $(V_1, V_2, V_3)$  are derived from the mixing distribution  $F$  specified in the copula. The following remark summarizes some properties of the distributions of  $V$  and  $(V_1, V_2, V_3)$ , which will become useful later when we examine the properties of these rank correlations.

**Remark 3.** *By the constructions of  $V$  and  $(V_1, V_2, V_3)$  in Theorem 1, we can deduce the following properties of their distributions: (i)  $V \stackrel{d}{=} -V$ , (ii)  $(V_1, V_2, V_3) \stackrel{d}{=} (V_2, V_1, V_3)$ , and (iii)  $V_i \stackrel{d}{=} -V_i$  for  $i = 1, 2$ .*

For Kendall's tau given in (4.1), the mixture involves a single random variable  $V$  whose distribution is symmetric about zero, as noted in Remark 3 (i). The pdf of  $V$  can be derived from the pdf of  $F$  using the transformation technique,<sup>12</sup> and the expectation on the right-hand side of (4.1) can then be expressed as a one-dimensional integral. However, since the closed form of the pdf of  $V$  is generally unavailable, it is more practical to interpret the expectation as a two-dimensional integral with respect to the joint probability measures of the independent pair  $(W, W^*)$ . This approach enables efficient evaluation using *quasi-Monte-Carlo* (QMC) integration techniques.

To demonstrate this, we rewrite the formula of Kendall's tau in (4.1) as

$$\begin{aligned} \tau(\rho, \beta, F) &= 4 \int_0^\infty \int_0^\infty \Phi_2 \left( \frac{\beta_1(w^* - w)}{\sqrt{w^* + w}}, \frac{\beta_2(w^* - w)}{\sqrt{w^* + w}}; \rho \right) f(w)f(w^*) dw dw^* - 1 \\ &= 4 \mathbb{E} \Phi_2(\beta_1 g(U_1, U_2), \beta_2 g(U_1, U_2); \rho) - 1, \end{aligned} \quad (4.3)$$

where  $(U_1, U_2) \sim U(0, 1)^2$  and  $g(u_1, u_2) := [q(u_2) - q(u_1)]/\sqrt{q(u_1) + q(u_2)}$ , with  $q := F^{-1}$  being the quantile function of  $F$ . The expectation on the right-hand side of (4.3) can then

---

<sup>12</sup>To see this, let  $f$  and  $f_V$  denote the pdf's of  $F$  and  $V$ , respectively. Define the transformation  $(U, V) = (\sqrt{W + W^*}, (W^* - W)/\sqrt{W^* + W})$ , where  $U$  serves as an auxiliary variable and the joint density of  $(W, W^*)$  is  $f(w)f(w^*)1\{w > 0, w^* > 0\}$ . From the inverse transformation  $(W, W^*) = ((U^2 - UV)/2, (U^2 + UV)/2)$ , we can show that the support of  $(U, V)$  is  $\{u > 0, |v| < u\}$  and the Jacobian determinant of the inverse transformation is  $U^2$ . Then, we have  $f_V(v) = \int_0^\infty f((u^2 - uv)/2) f((u^2 + uv)/2) u^2 du$ , for  $v \in \mathbb{R}$ .

be evaluated efficiently via QMC methods by sampling  $(U_1, U_2)$  from a low-discrepancy sequence over  $(0, 1)^2$ , such as Sobol or Halton sequences, and then approximating the expectation as the average of the computed values from the sample.

Similarly, we may rewrite the formula of Spearman's rho in (4.2) as

$$\rho_S(\rho, \beta, F) = 12 \mathbb{E} \Phi_2(\beta_1 h_1(U), \beta_2 h_2(U); \rho h_3(U)) - 3,$$

where  $U := (U_1, U_2, U_3) \sim U(0, 1)^3$ , and  $h_1, h_2$  and  $h_3$  are defined as, respectively,

$$h_i(u) = \frac{q(u_i) - q(u_3)}{\sqrt{q(u_i) + q(u_3)}}, \quad i = 1, 2, \quad \text{and} \quad h_3(u) = \frac{q(u_3)}{\sqrt{[q(u_1) + q(u_3)][q(u_2) + q(u_3)]}},$$

where  $u = (u_1, u_2, u_3)$ , and  $q$  is again the quantile function of  $F$ . Then, we can evaluate Spearman's rho through a sampling of  $(U_1, U_2, U_3)$  from a low-discrepancy sequence over  $(0, 1)^3$  and then approximating the expectation as the average of all the computed values. It is known that QMC integration is preferred over Monte-Carlo (MC) integration, especially when the dimension of the integration is relatively high, as in this case, where we need to evaluate a three-dimension integral.

Lastly, to appreciate the role played by the skewness vector  $\beta$  in the rank correlations for this class of skew-elliptical copula, we connect the formulas in (4.1) and (4.2) to their counterparts in elliptical case. Specifically, when  $\beta = 0$ , the copula  $C_{\text{mn}}(\rho, 0, F)$  is the elliptically symmetric normal scale mixture copula with pseudo-correlation  $\rho$  and mixing distribution  $F$ . In this case, formula (4.1) in Theorem 1 (i) simplifies to

$$\tau(\rho, 0, F) = 4 \Phi_2(0, 0; \rho) - 1 = \frac{2}{\pi} \arcsin \rho, \quad (4.4)$$

where the second equality follows from the identity  $\Phi_2(0, 0; \rho) = 1/4 + (\arcsin \rho)/(2\pi)$ , as stated in McNeil et al. (2005, Proposition 5.35, p. 215). This mapping from  $\rho$  to  $\tau$  is well expected and was illustrated in the left panel of Figure 1.

On the other hand, formula (4.2) in Theorem 1 (ii) implies that Spearman's rho of the normal scale mixture copula is

$$\rho_S(\rho, 0, F) = 12 \mathbb{E} \Phi_2(0, 0; \rho V_3) - 3 = \frac{6}{\pi} \mathbb{E} \arcsin \rho V_3, \quad (4.5)$$



which, unlike Kendall's tau above, depends on the mixing distribution  $F$  through  $V_3$ . This formula was previously derived by [Heinen & Valdesogo \(2020, Proposition 1\)](#). The bivariate Gaussian copula can be considered as a scale normal mixture copula in which the mixing distribution  $F$  is concentrated at 1. In that case,  $V_3 = 1/2$  almost surely, and then  $\rho_S = (6/\pi) \arcsin(\rho/2)$ , as presented earlier in equation (2.2).

## 4.2 Properties of Kendall's tau and Spearman's rho

Since Kendall's tau and Spearman's rho presented in Theorem 1 are both expressed in terms of  $\Phi_2(\cdot; \rho)$ , the cdf of the bivariate normal  $N(0, \varrho(\rho))$ , their properties can be analyzed using the characteristics of this bivariate normal cdf. In particular, for  $(x_1, x_2) \in \mathbb{R}^2$  and  $\rho \in (-1, 1)$ , two useful representations of  $\Phi(x_1, x_2; \rho)$  are

$$\Phi_2(x_1, x_2; \rho) = \Phi(x_1)\Phi(x_2) + \int_0^\rho \phi_2(x_1, x_2; r) dr, \quad (4.6)$$

and

$$\begin{aligned} \Phi_2(x_1, x_2; \rho) &= \frac{1}{2} [\Phi(x_1) + \Phi(x_2)] - a(x_1, x_2) \\ &\quad - T\left(x_1, \frac{x_2 - \rho x_1}{x_1 \sqrt{1 - \rho^2}}\right) - T\left(x_2, \frac{x_1 - \rho x_2}{x_2 \sqrt{1 - \rho^2}}\right), \end{aligned} \quad (4.7)$$

where  $T$  is Owen's T function, and  $a(x, y) := \frac{1}{2} \mathbf{1}\{xy < 0 \text{ or if } xy = 0 \text{ and } x \text{ or } y < 0\}$ .<sup>13</sup> The decomposition of  $\Phi_2(x_1, x_2; \rho)$  given in (4.7) is particularly useful for analyzing the properties of Spearman's rho, whose expression in (4.2) involves more intricate mixing structure. We also note the following partial derivatives:

$$\frac{\partial}{\partial x_1} \Phi_2(x_1, x_2; \rho) = \phi(x_1) \Phi\left(\frac{x_2 - \rho x_1}{\sqrt{1 - \rho^2}}\right), \quad (4.8)$$

$$\frac{\partial}{\partial \rho} \Phi_2(x_1, x_2; \rho) = \phi_2(x_1, x_2; \rho). \quad (4.9)$$

---

<sup>13</sup>See, for example [Owen \(1956, Equation 3.6, p. 1078\)](#) for the representation in (4.6), and [Azzalini \(2014, \(B.18\), p. 234\)](#) for the representation in (4.7).

Drawing on the formulas for Kendall's tau and Spearman's rho given in Theorem 1, the properties of the mixing distributions highlighted in Remark 3, and the properties of the bivariate normal cdf  $\Phi_2(\cdot; \rho)$ , we can summarize several basic properties of the rank correlations of the normal location-scale mixture copula in the following proposition.

**Proposition 1.** *Kendall's tau and Spearman's rho for the bivariate normal location-scale mixture copula  $C_{\text{mn}}(\rho, \beta, F)$  possess the following properties:*

(i) *Symmetry in components:*

$$\tau(\rho, (\beta_1, \beta_2), F) = \tau(\rho, (\beta_2, \beta_1), F) \quad \text{and} \quad \rho_S(\rho, (\beta_1, \beta_2), F) = \rho_S(\rho, (\beta_2, \beta_1), F).$$

(ii) *Invariance to a sign change of the skewness vector:*

$$\tau(\rho, \beta, F) = \tau(\rho, -\beta, F) \quad \text{and} \quad \rho_S(\rho, \beta, F) = \rho_S(\rho, -\beta, F).$$

(iii) *Strict monotonicity in  $\rho$ :*

$$\frac{\partial}{\partial \rho} \tau(\rho, \beta, F) > 0 \quad \text{and} \quad \frac{\partial}{\partial \rho} \rho_S(\rho, \beta, F) > 0.$$

All the properties stated in Proposition 1 can be understood intuitively. For properties (i) and (ii), note that, given any two variables  $X_1$  and  $X_2$ , the rank correlation between  $X_1$  and  $X_2$  is always equal to that between  $X_2$  and  $X_1$ , and also equal to that between  $-X_1$  and  $-X_2$ . In the current context, we can easily demonstrate that if  $(X_1, X_2) \sim \text{MN}(0, \varrho(\rho), (\beta_1, \beta_2), F)$ , so that the copula of  $(X_1, X_2)$  is  $C_{\text{mn}}(\rho, (\beta_1, \beta_2), F)$ , then  $(X_2, X_2) \sim \text{MN}(0, \varrho(\rho), (\beta_2, \beta_1), F)$  and  $(-X_1, -X_2) \sim \text{MN}(0, \varrho(\rho), -(\beta_1, \beta_2), F)$ .

Moreover, the strict monotonicity in  $\rho$  described in property (iii) aligns with the behavior observed for elliptical copulas, as illustrated in Figure 1. This monotonic relationship highlights that both Kendall's tau and Spearman's rho effectively capture increasing dependence as the pseudo-correlation  $\rho$  grows, even in the presence of asymmetry. However, as we will demonstrate later, certain properties for elliptical copulas – such as Kendall's tau and Spearman's rho taking value 1, 0 or  $-1$  whenever the pseudo-correlation  $\rho$  takes value 1, 0 or  $-1$ , respectively – do not extend to this class of skew-elliptical copulas.

### 4.3 Rank correlations for GH skew- $t$ copula

As noted earlier, the widely used GH skew- $t$  copula emerges as a special case of the normal location-scale mixture copula when the mixing distribution  $F$  is chosen as  $\text{IG}(\nu/2, \nu/2)$  with  $\nu > 0$ . This leads to the following corollary to Theorem 1, which summarizes the Kendall's and Spearman's rank correlations for the GH skew- $t$  copula.


**Corollary 1.** *For the bivariate GH skew- $t$  copula  $C_{\text{GH}t}(\rho, \beta, \nu)$ , Kendall's tau and Spearman's rho are given by (4.1) and (4.2) in Theorem 1, respectively, with the mixing distribution  $F$  specified as  $\text{IG}(\nu/2, \nu/2)$ .*

Using the QMC method described at the end of Section 3.1, we evaluate Kendall's and Spearman's rank correlations for the GH skew- $t$  copulas under various parameter settings and present the functional relationships between the pseudo-correlation  $\rho$  and each rank correlation in Figure 2. For comparison, the benchmark  $t$  copula case, corresponding to  $\beta_1 = \beta_2 = 0$ , is depicted with a dark solid line.

As clearly shown in Figure 2, the asymmetry introduced by nonzero skewness parameters  $\beta_1$  and  $\beta_2$  can significantly alter the behavior of both  $\tau$  and  $\rho_S$  as functions of  $\rho$ . Notably, while both rank correlations remain monotonically increasing in  $\rho$ , the full interval  $[-1, 1]$  are no longer always attainable, as in the elliptical case. For example, as shown by the red dotted lines in the left two panels, both Kendall's tau and Spearman's rho are strictly positive for the GH skew- $t$  copula  $C_{\text{GH}t}(\rho, (1, 2), 4)$  for any  $\rho \in [-1, 1]$ . Furthermore, the property that rank correlations take the values  $-1$  or  $1$  whenever  $\rho$  equals  $-1$  or  $1$ , respectively, generally ceases to hold in the presence of asymmetry.

Figure 2 also shows that when one skewness parameter is fixed at a *positive* value ( $\beta_1 = 1$ ), both rank correlations *increase* as the other skewness parameter ( $\beta_2$ ) increases. By property (ii) in Proposition 1, we can infer that when one skewness parameter is fixed at a *negative* value, both rank correlations *decreases* as the other skewness parameter increases. Moreover, when one of the skewness parameters is zero ( $\beta_2 = 0$ ), both Kendall's tau and Spearman's rho take the value zero when  $\rho = 0$ , as in the case for elliptical copulas. We will show in Section 4.4.2 below that this is true in general for the entire class of skew-elliptical copulas under the single-skew setting .

Figure 2: Kendall's tau and Spearman's rho for GH skew  $t$  copula.



Notes: Kendall's tau and Spearman's rho are shown as functions of the pseudo-correlation  $\rho$  for the bivariate GH skew- $t$  copula  $C_{\text{GH}t}(\rho, (\beta_1, \beta_2), \nu)$ . The left panels show the case  $\nu = 4$ , and the right panels  $\nu = 10$ . The formulas are given by (4.1) and (4.2) in Theorem 1, with the mixing distribution  $F$  specified as  $\text{IG}(\nu/2, \nu/2)$ . In each panel, various settings of skewness parameters  $(\beta_1, \beta_2)$  are considered, and the dark solid line represents the benchmark  $t$  copula case ( $\beta_1 = \beta_2 = 0$ ).

#### 4.4 Equi-skew and single-skew cases

Building on the general result of Theorem 1, we now examine two special cases: the *equi-skew* setting ( $\beta_1 = \beta_2$ ) and the *single-skew* setting (one of  $\beta_1$  and  $\beta_2$  is zero). We present the corresponding results in Propositions 2 and 3, respectively, and the graphical illustrations in Figures 3 and 4, using the GH skew- $t$  copula as an example.

#### 4.4.1 Equi-skew case

To appreciate the importance of studying the equi-skew case, it is useful to recall that elliptical copulas possess two forms of symmetry: *exchangeability* and *radial symmetry*. In the equi-skew setting where  $\beta_1 = \beta_2 = b \in \mathbb{R}$ , the copula  $C_{\text{nm}}(\rho, (b, b), F)$  retains the exchangeability of elliptical copulas, yet it breaks their radial symmetry. Therefore, the equi-skew setting is useful when allowing radial asymmetry is sufficient or when one wishes to maintain exchangeability within the model. Moreover, imposing equi-skewness naturally reduces the number of parameters within this class of skew-elliptical copulas, which is particularly advantageous in high-dimensional applications where parsimony is crucial for practical implementation and interpretation.<sup>14</sup>

The following proposition presents the expressions and properties of Kendall's tau and Spearman's rho for the bivariate normal location-scale mixture copula under equal skewness. Since both rank correlations are invariant under a sign change of the skewness vector, as noted in Proposition 1 (ii), we shall, without loss of generality, focus on the  $b > 0$  case in the discussion.

**Proposition 2.** (*Equi-skewness*) Suppose  $X_1$  and  $X_2$  are two random variables with continuous margins and copula  $C_{\text{nm}}(\rho, (b, b), F)$  where  $\rho \in (-1, 1]$ ,  $b > 0$ , and  $F$  is a univariate distribution on  $(0, \infty)$ . Then, we have the following.

(i) Kendall's tau of  $X_1$  and  $X_2$  is given by  $\tau(\rho, b, F) = 4 \mathbb{E} \Phi_2(bV, bV; \rho) - 1$ , where  $V$  is defined in Theorem 1 (i). If additionally  $\mathbb{E} V^2 < \infty$ , then

$$\frac{\partial}{\partial b} \tau(\rho, b, F) > 0 \quad \text{and} \quad \frac{\partial^2}{\partial b \partial \rho} \tau(\rho, b, F) < 0,$$

(ii) Spearman's rho of  $X_1$  and  $X_2$  is given by  $\rho_S(\rho, b, F) = 12 \mathbb{E} \Phi_2(bV_1, bV_2; \rho V_3) - 3$ , where  $V_1, V_2, V_3$  are defined in Theorem 1 (ii).


(iii) When  $\rho = 1$ , we have  $\tau(1, b, F) = \rho_S(1, b, F) = 1$  for any  $b$  and  $F$ .

In Figure 3, we present Kendall's tau and Spearman's rho for GH skew- $t$  copulas under the equi-skew setting. It is evident from the figure that increasing the level of asymmetry

---

<sup>14</sup>For example, both Christoffersen et al. (2012) and Oh & Patton (2023) use the GH skew- $t$  copula under equi-skewness or grouped equi-skewness assumptions in their high-dimensional applications.

Figure 3: Kendall's tau and Spearman's rho for GH skew  $t$ -copula under equi-skewness.




Notes: Kendall's tau and Spearman's rho are presented as functions of the pseudo-correlation  $\rho$  for the bivariate GH skew- $t$  copula under equi-skewness ( $\beta_1 = \beta_2$ ). The left panels show the case  $\nu = 4$ , and the right panels  $\nu = 10$ . The formulas are given in Proposition 2 (i) and (ii), with the mixing distribution  $F$  specified as  $IG(\nu/2, \nu/2)$ . In each panel, various levels of the equi-skewness are considered, and the dark solid line represents the benchmark  $t$  copula case ( $\beta_1 = \beta_2 = 0$ ).

risers both rank correlations  $\tau$  and  $\rho_S$  regardless of the value of  $\rho$ . However, as  $\rho$  grows, the magnitude of this asymmetry-driven increase in the rank correlations diminishes. We demonstrate Proposition 2 (i), by identifying the signs of the partial derivatives, that this pattern holds in general for Kendall's tau within this class of skew-elliptical copulas under mild assumptions on the mixing distribution.

Moreover, as stated in Proposition 2 (iii), both Kendall's tau and Spearman's rho take the value 1 when  $\rho = 1$  under the equi-skew setting. This is confirmed for the GH skew- $t$

Figure 4: Kendall's tau and Spearman's rho for GH skew- $t$  copula under single-skewness.



Notes: Kendall's tau and Spearman's rho are presented as functions of the pseudo-correlation  $\rho$  for the bivariate GH skew- $t$  copula under single-skewness ( $\beta_2 = 0$ ). The left panels show the case  $\nu = 4$ , and the right panels  $\nu = 10$ . The formulas are given in Proposition 3 (i), with the mixing distribution  $F$  specified as  $\text{IG}(\nu/2, \nu/2)$ . In each panel, various levels of the single-skewness are considered, and the dark solid line represents the benchmark  $t$  copula case ( $\beta_1 = \beta_2 = 0$ ).

copulas shown in Figure 3. However, Figure 3 also shows that neither rank correlation approaches  $-1$  as  $\rho$  tends to  $-1$ .

#### 4.4.2 Single-skew case

Next, we consider the single-skew setting, where one of the skewness parameters is zero. The following proposition summarizes the expressions and properties of Kendall's tau and

Spearman's rho under this condition. Due to properties (i) and (ii) in Proposition 1, we focus on  $\beta_1 = b > 0$  and  $\beta_2 = 0$  in our discussion, without loss of generality.

**Proposition 3.** *(Single-skewness) Suppose  $X_1$  and  $X_2$  are two random variables with continuous margins and copula  $C_{\text{nm}}(\rho, (b, 0), F)$  where  $\rho \in [-1, 1]$ ,  $b > 0$ , and  $F$  is a univariate distribution on  $(0, \infty)$ . Then, we have the following.*

(i) *Kendall's tau and Spearman's rho of  $X_1$  and  $X_2$  are, respectively,  $4 \mathbb{E} \Phi_2(bV, 0; \rho) - 1$  and  $12 \mathbb{E} \Phi_2(bV_1, 0; \rho V_3) - 3$ , where  $V, V_1$  and  $V_3$  are defined in Theorem 1.*

(ii) *Kendall's tau and Spearman's rho of  $X_1$  and  $X_2$  are both zero when  $\rho = 0$ , and they are odd functions of  $\rho$ .*

(iii) *Kendall's tau and Spearman's rho of  $X_1$  and  $X_2$  are both strictly decreasing (or strictly increasing) in  $b$  when  $\rho > 0$  (or when  $\rho < 0$ ).*

As stated in Proposition 3 (iii), in the single-skew case,  $\tau = \rho_S = 0$  whenever  $\rho = 0$ , and both  $\tau$  and  $\rho_S$  are odd functions of  $\rho$ . This property aligns with the behavior of  $\tau$  and  $\rho_S$  for elliptical copulas. However, as clearly illustrated in Figure 4 for the GH skew- $t$  copula under single-skewness, the presence of the skewness causes the values of  $\tau$  and  $\rho_S$  to deviate from  $-1$  or  $1$  as  $\rho$  approaches boundary values  $-1$  or  $1$ . In particular, as shown in Proposition 3 (iii), both rank correlations are strictly decreasing (increasing) in the level of the single skewness when  $\rho$  is positive (negative).

## 5 Rank correlations of skew-normal scale mixtures

In this section, we turn our attention to another class of skew-elliptical copulas – the skew-normal scale mixture copula defined in Definition 6 – and establish the formulas and properties of Kendall's tau and Spearman's rho for this class. We then specialize the results to the skew-normal and AC skew- $t$  copulas. We also discuss the equi-skew and single-skew settings for this class of skew-elliptical copulas in the final subsection.



## 5.1 Formulas of Kendall's tau and Spearman's rho

Before presenting the results in Theorem 2, we first introduce the following  $4 \times 4$  and  $5 \times 5$  correlation-matrix-valued functions, which will play a key role in the theorem. In particular, for  $\rho \in [-1, 1]$  and  $u = (u_1, u_2)$  with  $|u_i| < 1$  for  $i = 1, 2$ , we define

$$P_\tau(\rho, u, v) = \left[ \begin{array}{cc|cc} 1 & & & \\ \rho & 1 & & \\ \hline u_1 v_1 & u_2 v_1 & 1 & \\ u_1 v_2 & u_2 v_2 & 0 & 1 \end{array} \right], \quad \text{where } v = (v_1, v_2), \quad (5.1)$$

and

$$P_S(\rho, u, v) = \left[ \begin{array}{cc|ccc} 1 & & & & \\ \rho v_3 & 1 & & & \\ \hline u_1 v_1 & 0 & 1 & & \\ 0 & u_2 v_2 & 0 & 1 & \\ u_1 v_1^- & u_2 v_2^- & 0 & 0 & 1 \end{array} \right], \quad \text{where } v = (v_1, v_2, v_1^-, v_2^-, v_3), \quad (5.2)$$

where the upper-triangular entries are omitted in the correlation matrix. The values of  $u$  and  $v$  in (5.1) and (5.2) must satisfy certain conditions to ensure that  $P_\tau(\rho, u, v)$  and  $P_S(\rho, u, v)$  form valid correlation matrices.

The following theorem presents formulas of Kendall's tau and Spearman's rho for bivariate skew-normal scale mixture copulas. In the theorem, we use  $\Phi_d(0; \Sigma)$  to denote the cdf of the  $d$ -dimensional normal distribution  $N(0; \Sigma)$  evaluated at the zero vector.

**Theorem 2.** *Suppose  $X_1$  and  $X_2$  are two random variables with continuous margins and skew-normal scale mixture copula  $C_{\text{msn}}(\rho, \alpha, F)$  where  $\rho \in [-1, 1]$ ,  $\alpha = (\alpha_1, \alpha_2) \in \mathbb{R}^2$ , and  $F$  is a univariate distribution on  $(0, \infty)$ . Let  $P_\tau$  and  $P_S$  be the two correlation-matrix-valued functions defined in (5.1) and (5.2), respectively. Furthermore, let  $\delta = (\delta_1, \delta_2)$  be the alternative skewness vector that corresponds to  $\alpha$  via (3.3). Then,*

(i) *Kendall's tau of  $X_1$  and  $X_2$  is given by*

$$\tau(\rho, \alpha, F) = 16 \mathbb{E} \Phi_4(0; P_\tau(\rho, \delta, V)) - 1, \quad (5.3)$$

where  $V = (V_1, V_2)$  in which  $V_1 = \sqrt{W_2/(W_1 + W_2)}$  and  $V_2 = -\sqrt{W_1/(W_1 + W_2)}$ , with  $W_1, W_2 \sim i.i.d. F$ ;

(ii) Spearman's rho of  $X_1$  and  $X_2$  is given by

$$\rho_S(\rho, \alpha, F) = 96 \mathbb{E} \Phi_5(0; P_S(\rho, \delta, V)) - 3, \quad (5.4)$$

where  $V = (V_1, V_2, V_1^-, V_2^-, V_3)$  in which  $V_i = \sqrt{W_i/(W_i + W_3)}$ ,  $V_i^- = -\sqrt{W_3/(W_i + W_3)}$ , for  $i = 1, 2$ , and  $V_3 = W_3/\sqrt{(W_1 + W_3)(W_2 + W_3)}$ , with  $W_1, W_2, W_3 \sim i.i.d. F$ .

Compared to the formulas for the normal location-scale mixture copula presented in Theorem 1, Kendall's tau and Spearman's rho for the skew-normal scale mixture copula, as given in Theorem 2, are also expressed in terms of expectations of mixtures of the normal cdf and can be computed via QMC methods. The following remark highlights some properties of the distributions of the mixing variables  $(V_1, V_2)$  and  $(V_1, V_2, V_1^-, V_2^-, V_3)$  in Theorem 2.

**Remark 4.** The distribution of  $(V_1, V_2)$  in Theorem 2 (i) satisfies  $(V_1, V_2) \stackrel{d}{=} -(V_2, V_1)$ . Moreover, the distribution of  $(V_1, V_2, V_1^-, V_2^-, V_3)$  in Theorem 2 (ii) satisfies  $(V_i, V_i^-) \stackrel{d}{=} -(V_i^-, V_i)$ , for  $i = 1, 2$ .

Unlike Theorem 1, which expresses  $\tau$  and  $\rho_S$  for normal location-scale mixture copulas using only the bivariate normal cdf, Theorem 2 for skew-normal scale mixture copulas involves four- and five-dimensional normal cdf's, whose evaluation often relies on QMC integration algorithms in most software packages. Corollary 2 to Theorem 2, presented below, offers alternative formulas for  $\tau$  and  $\rho_S$  that rely on the bivariate normal cdf.

In Corollary 2, the coefficients  $\alpha_1^\dagger$ ,  $\alpha_2^\dagger$  and  $\rho^\dagger$  are defined as:

$$\alpha_i^\dagger := \frac{\delta_i}{\sqrt{1 - \delta_i^2}} = \frac{\alpha_i + \rho\alpha_{-i}}{\sqrt{1 + (1 - \rho^2)\alpha_{-i}^2}} \quad (i = 1, 2) \quad (5.5)$$

and

$$\rho^\dagger := \frac{\rho - \delta_1\delta_2}{\sqrt{(1 - \delta_1^2)(1 - \delta_2^2)}}, \quad (5.6)$$

where  $(\delta_1, \delta_2)$  are related to  $\rho$  and  $(\alpha_1, \alpha_2)$  via (3.3).<sup>15</sup> Again, by convention,  $-i$  in (5.5) denotes the index other than  $i$ , for  $i = 1, 2$ .

**Corollary 2.** *Suppose  $X_1$  and  $X_2$  are two random variables with continuous margins and skew-normal scale mixture copula  $C_{\text{msn}}(\rho, \alpha, F)$  where  $\rho \in [-1, 1]$ ,  $\alpha = (\alpha_1, \alpha_2) \in \mathbb{R}^2$ , and  $F$  is a univariate distribution on  $(0, \infty)$ . Let  $\alpha_1^\dagger, \alpha_2^\dagger$  and  $\rho^\dagger$  be defined by (5.5) and (5.6), respectively. Then, we have the following.*

(i) *Kendall's tau of  $X_1$  and  $X_2$  given in (5.3) can be alternatively expressed as*

$$\tau(\rho, \alpha, F) = 4 \mathbb{E} \Phi_2(\alpha_1^\dagger Z, \alpha_2^\dagger Z; \rho^\dagger) - 1, \quad (5.7)$$

where  $Z := V_1 Y_1 + V_2 Y_2$ ,  $V = (V_1, V_2)$  is as defined in Theorem 2 (i), and  $(Y_1, Y_2)$  is a pair of independent half-normal random variables conditional on  $V$ .

(ii) *Spearman's rho of  $X_1$  and  $X_2$  given in (5.4) can be alternatively expressed as*

$$\rho_S(\rho, \alpha, F) = 12 \mathbb{E} \Phi_2(\alpha_1^\dagger Z_1, \alpha_2^\dagger Z_2; \rho^\dagger V_3) - 3, \quad (5.8)$$

where  $Z_1 := V_1 Y_1 + V_1^- Y_3$ ,  $Z_2 := V_2 Y_2 + V_2^- Y_3$ ,  $V = (V_1, V_2, V_1^-, V_2^-, V_3)$  is as defined in Theorem 2 (ii), and  $(Y_1, Y_2, Y_3)$  are mutually independent half-normal random variables conditional on  $V$ .

In Corollary 2, the dimension of the normal cdf in the rank correlation formulas is reduced from four or five (as in Theorem 2) to only two, at a cost of a more intricate construction of mixing variables. Nevertheless, evaluating the expectations in (5.7) and (5.8) is not difficult via QMC integration.

Importantly, the alternative expressions of  $\tau$  and  $\rho_S$  in Corollary 2 closely parallel those in Theorem 1 for the other skew-elliptical copula class. In Remark 5 below, we clarify the similarities and differences between the rank correlation formulas in Theorem 1 and Corollary 2, and relate them to the common and unique properties of the rank correlations across different skew-elliptical models, as further discussed in Section 5.2.

---

<sup>15</sup>As noted in Remark 1,  $\alpha_1^\dagger$  and  $\alpha_2^\dagger$  defined in (5.5) are the marginal  $\alpha$ -skewness parameters of the bivariate skew-normal distribution  $\text{SN}(0, \varrho(\rho), \alpha)$ . However, since the rank correlations do not depend on marginal distributions, we do not emphasize this interpretation of  $\alpha_1^\dagger$  and  $\alpha_2^\dagger$  in the following discussion.

**Remark 5.** (i) Although the distributions of the mixing variables  $Z$  and  $(Z_1, Z_2)$  in Corollary 2 are clearly different from those of  $V$  and  $(V_1, V_2)$  in Theorem 1, their marginal distributions are all symmetric about zero. As a result, Kendall's tau and Spearman's rho for both classes of skew-elliptical copulas are invariant under the sign change of the skewness vector, as shown in Proposition 1 (ii) and will be shown in Proposition 4 (ii).

(ii) A subtle but important difference in the rank correlation formulas in Corollary 2, compared to those in Theorem 1, lies in the nature of the coefficients/parameters. In (5.7) and (5.8), the coefficients  $\alpha_1^\dagger, \alpha_2^\dagger$  and  $\rho^\dagger$  that multiply the mixing variables are functional transformations of the skewness parameters  $\alpha$  and correlation parameter  $\rho$ , rather than these parameters themselves. This distinction is crucial to understanding why, unlike the case for normal location-scale mixture copulas, the entire interval  $[-1, 1]$  is attainable for  $\tau$  and  $\rho_S$  in the class of skew-normal scale mixture copulas (Proposition 4 (iii)).

Before concluding this subsection, we note that just as the normal location-scale mixture copula  $C_{mn}(\rho, \beta, F)$  reduces to the scale normal mixture copula when  $\beta = 0$ , the skew-normal scale mixture copula  $C_{msn}(\rho, \alpha, F)$  reduces to the *same* class of elliptical copulas when  $\alpha = 0$ . Substituting  $\alpha = 0$  into equations (5.7) and (5.8) yields, respectively,

$$\tau(\rho, 0, F) = 4\Phi_2(0, 0; \rho) - 1 = \frac{2}{\pi} \arcsin \rho$$

and

$$\rho_S(\rho, 0, F) = 12 \mathbb{E} \Phi_2(0, 0; \rho V_3) - 1 = \frac{6}{\pi} \mathbb{E} \arcsin \rho V_3,$$

which match the previously derived formulas in (4.4) and (4.5).

## 5.2 Properties of Kendall's tau and Spearman's rho

Using the representations of  $\tau$  and  $\rho_S$  in terms of the bivariate normal cdf, as given in Corollary 2, we carry out some analyses similar to those in Section 4.2 and establish the following basic properties for bivariate skew-normal scale mixture copulas.

**Proposition 4.** *Kendall's tau and Spearman's rho for the bivariate scale skew-normal normal mixture copula  $C_{msn}(\rho, \alpha, F)$ , where  $\rho \in [-1, 1]$  and  $\alpha = (\alpha_1, \alpha_2) \in \mathbb{R}^2$ , possess the following properties:*

(i) *Symmetry in components:*

$$\tau(\rho, (\alpha_1, \alpha_2), F) = \tau(\rho, (\alpha_2, \alpha_1), F) \quad \text{and} \quad \rho_S(\rho, (\alpha_1, \alpha_2), F) = \tau(\rho_S, (\alpha_2, \alpha_1), F).$$

(ii) *Invariance to a sign change of the skewness vector:*

$$\tau(\rho, \alpha, F) = \tau(\rho, -\alpha, F) \quad \text{and} \quad \rho_S(\rho, \alpha, F) = \rho_S(\rho, -\alpha, F).$$

(iii)  $\tau(-1, \alpha, F) = \rho_S(-1, \alpha, F) = -1$  and  $\tau(1, \alpha, F) = \rho_S(1, \alpha, F) = 1$ .

Properties (i) and (ii) in Proposition 4 parallel their counterparts for normal location-scale mixture copulas, stated in Parts (i) and (ii) of Proposition 1. We note here that both two properties can also be formulated in terms of skewness vector  $\delta$ .

Unlike for normal location-scale mixture copulas, property (iii) in Proposition 4 shows that, despite the asymmetry introduced in skew-normal scale mixture copulas, both Kendall's tau and Spearman's rho still attain  $-1$  or  $1$  when the pseudo-correlation coefficient  $\rho$  equals  $-1$  or  $1$ , mirroring the elliptical case. Moreover, since  $\tau$  and  $\rho_S$  are continuous functions of  $\rho$ , the entire interval  $[-1, 1]$  is attainable for these rank correlations, again mirroring the elliptical case.

Unlike (i) and (ii), property (iii) is more naturally expressed in terms of the skewness vector  $\alpha$ , rather than  $\delta$ , since  $\delta$  cannot vary freely from  $\rho$  (see Remark 1). Indeed, from (3.3) it follows that  $\delta_1 = \delta_2$  when  $\rho = 1$  and  $\delta_1 = -\delta_2$  when  $\rho = -1$ . These relationships are key to proving property (iii).

Although Part (iii) of Proposition 1 shows that both rank correlations increase strictly with  $\rho$ , holding the skewness vector  $\beta$  and mixing distribution fixed, for normal location-scale mixture copulas, establishing a similar property theoretically for skew-normal scale mixture copulas appears challenging in general.<sup>16</sup> Nevertheless, graphical illustrations of Kendall's tau and Spearman's rho against  $\rho$  for AC skew- $t$  copulas and skew-normal

---

<sup>16</sup>Although Heinen & Valdesogo (2022) state in their Lemma 1 and Lemma 3 that the partial derivatives of Kendall's tau and Spearman's rho with respect to  $\rho$  are strictly positive for bivariate skew-normal copulas, those derivative only make sense if  $\delta_1$  and  $\delta_2$  remain fixed while  $\rho$  varies. However, because the condition that  $\delta_1$  and  $\delta_2$  must satisfy changes as  $\rho$  changes (see Remark 2), one cannot hold  $\delta_1$  and  $\delta_2$  fixed while varying  $\rho$  in general.

copulas (Figures 5 and 6, respectively) suggest that both rank correlations do indeed increase strictly with  $\rho$  given each skewness vector  $\alpha$ . Moreover, under equi-skewness and single-skewness conditions, we do prove that Kendall's tau is a strictly increasing function of  $\rho$  for skew-normal scale mixture copulas (Proposition 5 (i) and Proposition 6 (ii)).

### 5.3 Rank correlations for skew-normal and AC skew- $t$ copulas

As mentioned at the end of Section 3.2, two important special cases of the skew-normal scale mixture copula arise when the mixing distribution is either  $\text{IG}(\nu/2, \nu/2)$  for some  $\nu > 0$  or degenerated at 1. In the former case, we obtain the AC skew- $t$  copula; in the latter, the skew-normal copula. In this section, we specialize the results of Theorem 2 and Corollary 2 to these two cases.

First, the following corollary gives Kendall's tau and Spearman's rho for the bivariate AC skew- $t$  copula.


**Corollary 3.** *For the bivariate AC skew- $t$  copula  $C_{\text{ACt}}(\rho, \alpha, \nu)$  where  $\rho \in [-1, 1]$ ,  $\alpha = (\alpha_1, \alpha_2) \in \mathbb{R}^2$  and  $\nu > 0$ , Kendall's tau and Spearman's rho are given by, respectively, (5.3) and (5.4) in Theorem 2, or (5.7) and (5.8) in Corollary 2, with the mixing distribution  $F$  specified as  $\text{IG}(\nu/2, \nu/2)$ .*

Next, by substituting the degenerated value 1 for all the mixing variables  $W$ 's in Theorem 2 and Corollary 2, we obtain the following corollary, which provides alternative expressions of Kendall's tau and Spearman's rho for the bivariate skew-normal copula. As a side note, the skew-normal copula can be viewed as the limiting case of the AC skew- $t$  copula as  $\nu \rightarrow \infty$ .

**Corollary 4.** *For the bivariate skew-normal copula  $C_{\text{sn}}(\rho, \alpha)$  where  $\rho \in [-1, 1]$  and  $\alpha = (\alpha_1, \alpha_2) \in \mathbb{R}^2$ , Kendall's tau and Spearman's rho are given by, respectively,*

$$\begin{aligned}\tau(\rho, \alpha) &= 16 \Phi_4(0; P_\tau(\rho, \delta, (c, -c))) - 1, \\ \rho_S(\rho, \alpha) &= 96 \Phi_5(0; P_S(\rho, \delta, (c, c, -c, -c, c^2))) - 3,\end{aligned}$$

Figure 5: Kendall's tau and Spearman's rho of AC skew  $t$  copula.




Notes: Kendall's tau and Spearman's rho are shown as functions of the pseudo-correlation  $\rho$  for the bivariate AC skew- $t$  copula  $C_{\text{AC}t}(\rho, (\alpha_1, \alpha_2), \nu)$ . The left panels show the case  $\nu = 1$ , and the right panels  $\nu = 10$ . Equivalent formulas are given in Theorem 2 and Corollary 2, with the mixing distribution  $F$  specified as  $\text{IG}(\nu/2, \nu/2)$ . In each panel, various settings of skewness parameters  $(\alpha_1, \alpha_2)$  are considered, and the dark solid line represents the benchmark  $t$  copula case ( $\alpha_1 = \alpha_2 = 0$ ). Here, we show the  $\nu = 1$  case instead of  $\nu = 4$  (as in Figure 2), because the difference between  $\nu = 4$  and  $\nu = 10$  is visually negligible for AC skew- $t$  copulas.

where  $c = 1/\sqrt{2}$ , and  $\delta$  is the alternative skewness vector that corresponds to  $\alpha$  via (3.3). Alternatively, given  $\alpha_1^\dagger, \alpha_2^\dagger$  and  $\rho^\dagger$  defined in (5.5) and (5.6), we have

$$\begin{aligned}\tau(\rho, \alpha) &= 4 \mathbb{E} \Phi_2(\alpha_1^\dagger Z, \alpha_2^\dagger Z; \rho^\dagger) - 1, \\ \rho_S(\rho, \alpha) &= 12 \mathbb{E} \Phi_2(\alpha_1^\dagger Z_1, \alpha_2^\dagger Z_2; \rho^\dagger/2) - 3,\end{aligned}$$

Figure 6: Kendall's tau and Spearman's rho of skew-normal copula.



Notes: Kendall's tau and Spearman's rho are presented as functions of the pseudo-correlation  $\rho$  for the bivariate skew-normal copula  $C_{\text{sn}}(\rho, (\alpha_1, \alpha_2))$ . The formulas are given in Corollary 4. In each panel, various settings of skewness parameters  $(\alpha_1, \alpha_2)$  are considered, and the dark solid line represents the benchmark  $t$  copula case ( $\alpha_1 = \alpha_2 = 0$ ).

where  $Z = (Y_1 - Y_2)/\sqrt{2}$ ,  $Z_1 = (Y_1 - Y_3)/\sqrt{2}$ , and  $Z_2 = (Y_2 - Y_3)/\sqrt{2}$ , with  $(Y_1, Y_2, Y_3)$  being mutually independent half-normal random variables.

In Corollary 4, the first set of formulas for  $\tau$  and  $\rho_S$ , expressed via four- and five-dimensional normal orthant probabilities, aligns with Propositions 1 and 3 of Heinen & Valdesogo (2022), who investigate the rank correlations for bivariate skew-normal distributions. The alternative formulas, involving only bivariate normal cdf's, are not provided in their paper.

We present  $\tau$  and  $\rho_S$  as functions of the pseudo-correlation  $\rho$  for bivariate AC skew- $t$  and skew-normal copulas in Figures 5 and 6, respectively. In contrast to the GH skew- $t$  copula case in Figure 2, the most notable difference in Figures 5 and 6 is that both ends of the curves are tied at  $(-1, -1)$  and  $(1, 1)$ , as indicated by Proposition 4 (iii). Moreover, both rank correlations increase strictly with  $\rho$ , and, unlike for GH skew- $t$  copulas, always span the full interval  $[-1, 1]$ . Additionally, when one skewness parameter ( $\alpha_1$  in the figures) is held fixed, increasing the other ( $\alpha_2$ ) *decreases* both rank correlations – opposite to the behavior observed in GH skew- $t$  copulas.



## 5.4 Equi-skew and single-skew cases

Just as what in Section 4.4 for normal location-scale mixture copulas, we also consider two special cases – the equi-skew setting ( $\alpha_1 = \alpha_2$ ) and the single-skew setting (one of  $\alpha_1$  and  $\alpha_2$  is zero) – for skew-normal scale mixture copulas. The corresponding results appear in Propositions 5 and 6, respectively, and Figures 7 and 8 illustrate these cases via the AC skew- $t$  copula.

### 5.4.1 Equi-skew case

In the equi-skew case, where  $\alpha_1 = \alpha_2 = a$  for some  $a \in \mathbb{R}$ , it follows from (3.3) that

$$\delta_1 = \delta_2 = \bar{\delta} := \frac{a(1 + \rho)}{\sqrt{1 + 2a^2(1 + \rho)}}. \quad (5.9)$$

Moreover, it follows from (5.5) that

$$\alpha_1^\dagger = \alpha_2^\dagger = a^\dagger := \frac{a(1 + \rho)}{\sqrt{1 + a^2(1 - \rho^2)}}. \quad (5.10)$$


It is straightforward to verify that (i)  $\text{sgn}(\bar{\delta}) = \text{sgn}(a^\dagger) = \text{sgn}(a)$ , (ii)  $\partial\bar{\delta}/\partial a > 0$  and  $\partial a^\dagger/\partial a > 0$ , and (iii)  $\text{sgn}(\partial\bar{\delta}/\partial\rho) = \text{sgn}(\partial a^\dagger/\partial\rho) = \text{sgn}(a)$ , for all  $a \in \mathbb{R}$  and  $\rho \in (-1, 1)$ . Furthermore, when  $\alpha_1 = \alpha_2 = a$ , it follows from (5.6) and (5.9) that  $\rho^\dagger$  takes the value

$$\rho^\dagger := \frac{1 + \rho}{1 + a^2(1 - \rho^2)} - 1. \quad (5.11)$$

Since both rank correlations are invariant under a sign change of  $\alpha$  – or equivalently, the scalar  $a$  in this equi-skew scenario – we shall, without loss of generality, focus on the case  $a > 0$  in the following proposition.

**Proposition 5.** *(Equi-skewness) Suppose  $X_1$  and  $X_2$  are two random variables with continuous margins and copula  $C_{\text{msn}}(\rho, (a, a), F)$  with  $\rho \in (-1, 1)$ , equi-skewness parameter  $a > 0$  and  $F$  being a univariate distribution on  $(0, \infty)$ . Let  $a^\dagger$  and  $\rho^\dagger$  be defined by (5.10) and (5.11), respectively. Then, we have the following.*

Figure 7: Kendall's tau and Spearman's rho of AC skew  $t$ -copula under equi-skewness.



Notes: Kendall's tau and Spearman's rho are presented as functions of the pseudo-correlation  $\rho$  for the bivariate AC skew- $t$  copula under equi-skewness ( $\alpha_1 = \alpha_2$ ). The left panels show the case  $\nu = 1$ , and the right panels  $\nu = 10$ . In each panel, various settings of skewness parameters ( $\alpha_1, \alpha_2$ ) are considered, and the dark solid line represents the benchmark  $t$  copula case ( $\alpha_1 = \alpha_2 = 0$ ). Here, we show the  $\nu = 1$  case instead of  $\nu = 4$  (as in Figure 3), because the difference between  $\nu = 4$  and  $\nu = 10$  is visually negligible for AC skew- $t$  copulas.

(i) Kendall's tau of  $X_1$  and  $X_2$  can be expressed as  $4 \mathbb{E} \Phi_2(a^\dagger Z, a^\dagger Z; \rho^\dagger) - 1$ , where  $Z$  is defined in Corollary 2 (i). It is strictly increasing in  $\rho$  and strictly decreasing in  $a$ .

(ii) Spearman's rho of  $X_1$  and  $X_2$  can be expressed as  $12 \mathbb{E} \Phi_2(a^\dagger Z_1, a^\dagger Z_2; \rho^\dagger V_3) - 3$ , where  $(Z_1, Z_2)$  and  $V_3$  are defined, respectively, in Corollary 2 (ii) and Theorem 2 (ii).

Figure 7 displays Kendall's tau and Spearman's rho as functions of the pseudo-correlation  $\rho$  for AC skew- $t$  copulas under the equi-skew setting. All the functions are strictly in-

creasing in  $\rho$ , and their ends meet at  $(-1, -1)$  and  $(1, 1)$ . Unlike the GH skew- $t$  copula case shown in Figure 3, increasing the level of asymmetry (i.e.  $a$  or  $\bar{\delta}$  in the equi-skew case) in AC skew- $t$  copulas reduces both rank correlations for any fixed  $\rho$ .

#### 5.4.2 Single-skew case

Unlike the equi-skew case discussed in Section 5.4.1, defining what we mean by “single-skew” for skew-normal scale mixture copulas is not straightforward. As noted in Remark 2, setting one of the skewness parameters in  $(\alpha_1, \alpha_2)$  to zero does not necessarily imply that one of  $(\delta_1, \delta_2)$  is zero – and vice versa. Here, we define the single-skew scenario by requiring one of  $\alpha_1$  and  $\alpha_2$  to be zero, since  $\alpha_1$  and  $\alpha_2$  are unconstrained. Specifically, without loss of generality, let  $\alpha_1 = a$ , for some  $a \in \mathbb{R}$ , and  $\alpha_2 = 0$ . In this case, from (3.3) we obtain

$$\delta_1 = \delta_\circ \quad \text{and} \quad \delta_2 = \delta_\circ \rho,$$

where  $\delta_\circ := a/\sqrt{1+a^2}$ . Note that  $\text{sgn}(\delta_\circ) = \text{sgn}(a)$  and  $\partial\delta_\circ/\partial a > 0$ . Moreover, it follows from (5.6) and (5.5) that in this case,

$$\rho^\dagger = \frac{\rho}{\sqrt{1+a^2(1-\rho^2)}} \tag{5.12}$$

and


$$\alpha_1^\dagger = a, \quad \alpha_2^\dagger = \frac{a\rho}{\sqrt{1+a^2(1-\rho^2)}} = a\rho^\dagger. \tag{5.13}$$

Again, since both rank correlations are invariant under a sign change of the skewness vector, as stated in Proposition 4 (ii), we focus on the  $a > 0$  case in the following proposition. The corresponding results for  $a < 0$  can be deduced accordingly.

**Proposition 6.** *(Single-skewness) Suppose  $X_1$  and  $X_2$  are two random variables with continuous margins and copula  $C_{\text{msn}}(\rho, (a, 0), F)$  with  $\rho \in (-1, 1)$ , the single-skewness parameter  $a > 0$  and  $F$  being a univariate distribution on  $(0, \infty)$ . Let  $\rho^\dagger$  be defined by (5.12). Then, we have the following.*

(i) *Kendall’s tau and Spearman’s rho of  $X_1$  and  $X_2$  are both zero when  $\rho = 0$ , and they are odd functions of  $\rho$ .*

Figure 8: Kendall's tau and Spearman's rho of AC skew  $t$ -copula under single-skewness.



Notes: Kendall's tau and Spearman's rho are presented as functions of the pseudo-correlation  $\rho$  for the bivariate AC skew- $t$  copula under single-skewness ( $\alpha_2 = 0$ ). The left panels show the case  $\nu = 1$ , and the right panels  $\nu = 10$ . In each panel, various settings of skewness parameters ( $\alpha_1, \alpha_2$ ) are considered, and the dark solid line represents the benchmark  $t$  copula case ( $\alpha_1 = \alpha_2 = 0$ ).

(ii) Kendall's tau of  $X_1$  and  $X_2$  can be expressed as  $4 \mathbb{E} \Phi_2(aZ, a\rho^\dagger Z; \rho^\dagger) - 1$ , where  $Z$  is defined in Corollary 2 (i). It is strictly increasing in  $\rho$ , and strictly decreasing (or increasing) in  $a$  when  $\rho > 0$  (or when  $\rho < 0$ ).

(iii) Spearman's rho of  $X_1$  and  $X_2$  can be expressed as  $12 \mathbb{E} \Phi_2(aZ_1, a\rho^\dagger Z_2; \rho^\dagger V_3) - 3$ , where  $(Z_1, Z_2)$  and  $V_3$  are defined, respectively, in Corollary 2 (ii) and Theorem 2 (ii).

Just as in the case for normal location-scale mixture copulas shown in Proposition 3, Kendall's tau and Spearman's rho for skew-normal scale mixture copulas are also odd

functions of  $\rho$  under single-skewness. This property is demonstrated clearly in Figure 8, which depicts  $\tau$  and  $\rho_S$  as functions of  $\rho$  for AC skew- $t$  copulas under various parameter settings. The figure also shows that, as the level of single-skewness increases, the magnitudes of the rank correlations decrease – similar to what is observed for GH skew- $t$  copulas.

## 6 Concluding remarks

We have derived explicit formulas for Kendall’s tau and Spearman’s rho for two general classes of skew-elliptical copulas, the normal location-scale mixture copulas and skew-normal scale mixture copulas. The results for the widely used GH skew- $t$ , AC skew- $t$  and skew-normal copulas can be derived as special cases of our general results. In particular, our formulas provide mappings from copula parameters to rank correlations through efficient numerical integration algorithms, thereby paving the way for future research on estimating copula parameters using (generalized) method of moments for these classes of skew-elliptical copula models, especially in high-dimensional applications.

The theoretical exploration of the effects of asymmetry vectors on the rank correlations for different classes of skew-elliptical copulas in this paper has yielded interesting results. Most notably, we show that the presence of asymmetry parameters in the normal location-scale mixture copulas alters the attainable range of Kendall’s tau and Spearman’s rho from  $[-1, 1]$  – as observed under elliptical symmetry – to a *strict* subset of  $[-1, 1]$ . In contrast, the entire interval of  $[-1, 1]$  remains attainable for skew-normal scale mixture copulas. This fact may impose constraints on the application of the former class of skew-elliptical copulas – which includes the GH skew- $t$  copula as a special case – since it is not always possible to construct a joint distribution with arbitrary empirical value of Kendall’s tau or Spearman’s rho rank correlation with  $[-1, 1]$ .

## References

- Aas, K., & Haff, I. H. (2006). The generalized hyperbolic skew student's  $t$ -distribution. *Journal of Financial Econometrics*, 4(2), 275–309.
- Azzalini, A. (2014). *The skew-normal and related families* (Vol. 3). Cambridge University Press.
- Azzalini, A., & Capitanio, A. (1999). Statistical applications of the multivariate skew normal distribution. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 61(3), 579–602.
- Azzalini, A., & Capitanio, A. (2003). Distributions generated by perturbation of symmetry with emphasis on a multivariate skew  $t$ -distribution. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 65(2), 367–389.
- Azzalini, A., & Dalla Valle, A. (1996). The multivariate skew-normal distribution. *Biometrika*, 83(4), 715–726.
- Barndorff-Nielsen, O. (1977). Exponentially decreasing distributions for the logarithm of particle size. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 353(1674), 401–419.
- Barndorff-Nielsen, O., & Halgreen, C. (1977). Infinite divisibility of the hyperbolic and generalized inverse gaussian distributions. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 38(4), 309–311.
- Blaesild, P., & Jensen, J. L. (1981). Multivariate distributions of hyperbolic type. In *Statistical Distributions in Scientific Work*. Springer Netherlands.
- Branco, M. D., & Dey, D. K. (2001). A general class of multivariate skew-elliptical distributions. *Journal of Multivariate Analysis*, 79(1), 99–113.
- Christoffersen, P., Errunza, V., Jacobs, K., & Langlois, H. (2012). Is the potential for international diversification disappearing? a dynamic copula approach. *The Review of Financial Studies*, 25(12), 3711–3751.
- Creal, D. D., & Tsay, R. S. (2015). High dimensional dynamic stochastic copula models. *Journal of Econometrics*, 189(2), 335–345.
- Demarta, S., & McNeil, A. J. (2005). The  $t$  copula and related copulas. *International Statistical Review*, 73(1), 111–129.

- Deng, L., Smith, M. S., & Maneesoonthorn, W. (2024). Large skew- $t$  copula models and asymmetric dependence in intraday equity returns. *Journal of Business & Economic Statistics*, 1–17.
- Gupta, A. K. (2003). Multivariate skew  $t$ -distribution. *Statistics*, 37(4), 359–363.
- Hansen, P. R., & Luo, Y. (2023). Robust estimation of realized correlation: New insight about intraday fluctuations in market betas. *arXiv preprint arXiv:2310.19992*.
- Heinen, A., & Valdesogo, A. (2020). Spearman rank correlation of the bivariate Student  $t$  and scale mixtures of normal distributions. *Journal of Multivariate Analysis*, 179, 104650.
- Heinen, A., & Valdesogo, A. (2022). The kendall and spearman rank correlations of the bivariate skew normal distribution. *Scandinavian Journal of Statistics*, 49(4), 1669–1698.
- Iman, R. L., & Conover, W.-J. (1982). A distribution-free approach to inducing rank correlation among input variables. *Communications in Statistics-Simulation and computation*, 11(3), 311–334.
- Kollo, T., & Pettere, G. (2010). Parameter estimation and application of the multivariate skew  $t$ -copula. In *Copula Theory and Its Applications* (pp. 289–298).
- Kruskal, W. H. (1958). Ordinal measures of association. *Journal of the American Statistical Association*, 53(284), 814–861.
- Lucas, A., Schwaab, B., & Zhang, X. (2017). Modeling financial sector joint tail risk in the euro area. *Journal of Applied Econometrics*, 32, 171–191.
- McNeil, A. J., Frey, R., & Embrechts, P. (2005). *Quantitative risk management: concepts, techniques and tools*. Princeton university press.
- McNeil, A. J., & Nešlehová, J. (2009). Multivariate archimedean copulas,  $d$ -monotone functions and  $\ell_1$ -norm symmetric distributions. *The Annals of Statistics*, 37(5B), 3059–3097.
- Oh, D. H., & Patton, A. J. (2013). Simulated method of moments estimation for copula-based multivariate models. *Journal of the American Statistical Association*, 108(502), 689–700.
- Oh, D. H., & Patton, A. J. (2023). Dynamic factor copula models with estimated cluster assignments. *Journal of Econometrics*, 237(2), 105374.

- Owen, D. B. (1956). Tables for computing bivariate normal probabilities. *The Annals of Mathematical Statistics*, 27(4), 1075–1090.
- Plackett, R. L. (1954). A reduction formula for normal multivariate integrals. *Biometrika*, 41(3/4), 351–360.
- Sklar, M. (1959). Fonctions de répartition à n dimensions et leurs marges. In *Annales de l'isup* (Vol. 8, pp. 229–231).
- Yoshihara, T. (2018). Maximum likelihood estimation of skew-t copulas with its applications to stock returns. *Journal of Statistical Computation and Simulation*, 88(13), 2489–2506.



# Appendices

## A Useful lemmas

**Lemma 1.** *Let  $X_1$  and  $X_2$  be two random variables with continuous marginal distributions  $F_1$  and  $F_2$ . Then,*

(i) *Kendall's tau of  $X_1$  and  $X_2$  is given by  $\tau = 4 \mathbb{P}\{X_1 < X_1^*, X_2 < X_2^*\} - 1$ , where  $(X_1^*, X_2^*)$  is an independent copy of  $(X_1, X_2)$ ;*

(ii) *Spearman's rho of  $X_1$  and  $X_2$  is given by  $\rho_S = 12 \mathbb{P}\{X_1 < X_1^\circ, X_2 < X_2^\circ\} - 3$ , where  $(X_1^\circ, X_2^\circ)$  is a pair of independent random variables independent of  $(X_1, X_2)$  and having the same marginal distributions  $F_1$  and  $F_2$ .*

*Proof.* By Definition 1, Kendall's tau of  $X_1$  and  $X_2$  can be written as the probability of concordance minus the probability of discordance of the pairs  $(X_1, X_2)$  and  $(X_1^*, X_2^*)$ :

$$\tau = \mathbb{P}\{(X_1 - X_1^*)(X_2 - X_2^*) > 0\} - \mathbb{P}\{(X_1 - X_1^*)(X_2 - X_2^*) < 0\}.$$

Since  $\mathbb{P}\{(X_1 - X_1^*)(X_2 - X_2^*) < 0\} = 1 - \mathbb{P}\{(X_1 - X_1^*)(X_2 - X_2^*) > 0\}$ , we then have

$$\tau = 2 \mathbb{P}\{(X_1 - X_1^*)(X_2 - X_2^*) > 0\} - 1 = 4 \mathbb{P}\{X_1 < X_1^*, X_2 < X_2^*\} - 1,$$

which proves Part (i). To prove Part (ii), note that  $\rho_S = 12 \int_0^1 \int_0^1 C(u_1, u_2) du_1 du_2 - 3$ , where  $C$  is the copula of  $X_1$  and  $X_2$ , by Proposition 5.29 in McNeil et al. (2005). As a result, for  $(U_1, U_2) := (F_1(X_1), F_2(X_2))$ , and  $(U_1^\circ, U_2^\circ)$  being a pair of independent  $U(0, 1)$  random variables that are independent of  $(U_1, U_2)$ , we have

$$\rho_S = 12 \mathbb{P}\{U_1 < U_1^\circ, U_2 < U_2^\circ\} - 3 = 12 \mathbb{P}\{X_1 < X_1^\circ, X_2 < X_2^\circ\} - 3,$$

where  $X_1^\circ := F_1^{-1}(U_1^\circ)$  and  $X_2^\circ := F_2^{-1}(U_2^\circ)$  are independent random variables with marginal distributions  $F_1$  and  $F_2$ , respectively. This completes the proof of Part (ii) of the lemma.  $\square$

**Lemma 2.** *For any  $x \in \mathbb{R}$  and  $\rho \in (-1, 1]$ , we have  $\Phi_2(x, x; \rho) = \Phi^s(x; \alpha_\rho)$ , where  $\Phi^s(\cdot; \alpha_\rho)$  is the univariate standard skew-normal cdf with skewness parameter  $\alpha_\rho = \sqrt{(1 - \rho)/(1 + \rho)} > 0$ .*

*Proof.* Suppose  $(X_1, X_2)$  has cdf  $\Phi_2(\cdot; \rho)$  with  $\rho \in (-1, 1)$ . Then, for  $x \in \mathbb{R}$ , we have

$$\Phi_2(x, x; \rho) = 2 \mathbb{P}\{X_1 \leq X_2, X_2 \leq x\} = 2 \int_{-\infty}^x \mathbb{P}\{X_1 \leq y \mid X_2 = y\} \phi(y) dy. \quad (\text{A.1})$$

Moreover, since  $(X_1 \mid X_2 = y) \sim N(\rho y, 1 - \rho^2)$ , we have

$$\mathbb{P}(X_1 \leq y \mid X_2 = y) = \mathbb{P}\left\{ \frac{X_1 - \rho y}{\sqrt{1 - \rho^2}} \leq \frac{(1 - \rho)y}{\sqrt{1 - \rho^2}} \mid X_2 = y \right\} = \Phi(\alpha_\rho y). \quad (\text{A.2})$$

It then follows from (A.1) and (A.2) that

$$\Phi_2(x, x; \rho) = \int_{-\infty}^x 2\phi(y)\Phi(\alpha_\rho y) dy = \int_{-\infty}^x \phi^s(y; \alpha_\rho) dy = \Phi^s(x; \alpha_\rho)$$

for  $x \in \mathbb{R}$  and  $\rho \in (-1, 1)$ . When  $\rho = 1$ , we have  $\alpha_\rho = 0$ . Moreover, it can be easily shown that  $\Phi_2(x, y; 1) = \Phi(\min\{x, y\})$ , and, therefore,  $\Phi_2(x, y; 1) = \Phi(x) = \Phi^s(x; 0)$  for any  $x \in \mathbb{R}$ . The proof of the lemma is then complete.  $\square$

**Lemma 3.** *Suppose  $X$  is a real valued symmetrically (about zero) distributed random variable with a well-defined pdf and finite first moment, and  $\phi^s(\cdot; \alpha)$  is the pdf of the univariate standard skew-normal distribution with  $\alpha > 0$ . Then, for any  $b > 0$ ,*

(i)  $\mathbb{E}X\phi^s(bX; \alpha) > 0$ ;

(ii)  $\mathbb{E}X\phi^s(bX; \alpha)$  is strictly increasing in  $\alpha$ , if, additionally,  $0 < \mathbb{E}X^2 < \infty$ .

*Proof.* Let  $f$  denote the pdf of  $X$ , which is a positive even function on  $\mathbb{R}$  by the symmetry of  $X$ . Then, by the change of variables,

$$\begin{aligned} \mathbb{E}X\phi^s(bX; \alpha) &= \int_{-\infty}^{\infty} x\phi^s(bx; \alpha)f(x) dx \\ &= b^{-2} \int_{-\infty}^{\infty} yf(y/b)\phi^s(y; \alpha) dy = b^{-2} \mathbb{E}g(Y), \end{aligned} \quad (\text{A.3})$$

where  $Y$  is a univariate standard skew-normal random variable with skewness  $\alpha$ , and  $g(x) := xf(x/b)$  for  $x \in \mathbb{R}$ . Since  $\alpha > 0$ , we have  $\mathbb{E}g(Y) > 0$ , as  $Y$  is strictly positively skewed and  $g$  is an odd function on  $\mathbb{R}$ . This proves Part (i) of the lemma.

To show Part (ii) of the lemma, note that if  $0 < \mathbb{E}X^2 < \infty$ , then taking partial derivative with

respect to  $\alpha$  on both sides of (A.3) yields (noticing that  $\phi^s(bx; \alpha) = 2\phi(bx)\Phi(\alpha bx)$ )

$$\frac{\partial}{\partial \alpha} \mathbb{E}X\phi^s(bX; \alpha) = 2b \int_{-\infty}^{\infty} x^2 \phi(bx)\phi(\alpha bx)f(x) dx = 2b \mathbb{E}X^2\phi(bX)\phi(\alpha bX) > 0,$$

as desired to be shown.  $\square$

**Lemma 4.** *Let  $(X_1, X_2) \sim \text{SN}(0, \varrho(\rho), \alpha)$  where  $\rho \in (-1, 1)$  and  $\alpha \in \mathbb{R}^2$ , and  $(X_1^*, X_2^*)$  be an independent copy of  $(X_1, X_2)$ . Let  $a, b$  be two strictly positive constants. We have*

$$\mathbb{P}\{aX_1^* > bX_1, aX_2^* > bX_2\} = 4\Phi_4(0; P_\tau(\rho, \delta, v)),$$

where  $P_\tau$  is defined by (5.1),  $\delta = (\delta_1, \delta_2)^\top = \varrho(\rho)\alpha/\sqrt{1 + \alpha^\top \varrho(\rho)\alpha}$ , and  $v = (v_1, v_2)$  with  $v_1 = a/\sqrt{a^2 + b^2}$  and  $v_2 = -b/\sqrt{a^2 + b^2}$ .

*Proof.* Let  $Z = (Z_1, \dots, Z_6)^\top$  be a 6-dimensional normal random vector with mean zero and covariance matrix (upper-triangular entries omitted)

$$P_Z = \left[ \begin{array}{ccc|ccc} 1 & & & & & \\ \rho & 1 & & & & \\ \delta_1 & \delta_2 & 1 & & & \\ \hline 0 & 0 & 0 & 1 & & \\ 0 & 0 & 0 & \rho & 1 & \\ 0 & 0 & 0 & \delta_1 & \delta_2 & 1 \end{array} \right] = \begin{bmatrix} P_3 & 0 \\ 0 & P_3 \end{bmatrix}, \quad \text{where } P_3 = \begin{bmatrix} \varrho(\rho) & \delta \\ \delta^\top & 1 \end{bmatrix}.$$

Note that, by the properties of the multivariate normal distribution and the stochastic representation of the skew-normal distribution via conditioning (see e.g. [Azzalini, 2014](#), Section 5.1.3), we have the representations  $(X_1, X_2) \stackrel{d}{=} (Z_4, Z_5) | \{Z_6 > 0\}$  and  $(X_1^*, X_2^*) \stackrel{d}{=} (Z_1, Z_2) | \{Z_3 > 0\}$ . Next, for  $a, b > 0$ , we define

$$\begin{bmatrix} Y_1 \\ Y_2 \end{bmatrix} \stackrel{d}{=} \left( a \begin{bmatrix} Z_1 \\ Z_2 \end{bmatrix} - b \begin{bmatrix} Z_4 \\ Z_5 \end{bmatrix} \right) \Big| \{Z_3 > 0, Z_6 > 0\}. \quad (\text{A.4})$$

Then, to prove the lemma it suffices to show that  $\mathbb{P}\{Y_1 > 0, Y_2 > 0\} = 4\Phi_4(0; P_\tau(\rho, \delta, v))$ . To see this, we define a 4-dimensional normal random vector  $W = (W_1, \dots, W_4)^\top$  by

$$W = \begin{bmatrix} A & B \\ C & D \end{bmatrix} Z$$

where

$$A = \begin{bmatrix} a & 0 & 0 \\ 0 & a & 0 \end{bmatrix}, \quad B = \begin{bmatrix} -b & 0 & 0 \\ 0 & -b & 0 \end{bmatrix}, \quad C = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \quad D = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

so that

$$W_1 = aZ_1 - bZ_4, \quad W_2 = aZ_2 - bZ_5, \quad W_3 = Z_3, \quad W_4 = Z_6. \quad (\text{A.5})$$

By construction,  $W \sim N(0, \Sigma_W)$ , where

$$\Sigma_W = \begin{bmatrix} AP_1A^\top + BP_1B^\top & AP_1C^\top + BP_1D^\top \\ CP_1A^\top + DP_1B^\top & CP_1C^\top + DP_1D^\top \end{bmatrix} = \left[ \begin{array}{cc|cc} a^2 + b^2 & (a^2 + b^2)\rho & a\delta_1 & -b\delta_1 \\ (a^2 + b^2)\rho & a^2 + b^2 & a\delta_2 & -b\delta_2 \\ \hline a\delta_1 & a\delta_2 & 1 & 0 \\ -b\delta_1 & -b\delta_2 & 0 & 1 \end{array} \right].$$

By (A.4) and (A.5), we have  $(Y_1, Y_2) \stackrel{d}{=} (W_1, W_2) | \{W_3 > 0, W_4 > 0\}$ , and consequently,

$$\begin{aligned} \mathbb{P}\{Y_1 > 0, Y_2 > 0\} &= \mathbb{P}\{W_1 > 0, W_2 > 0 | W_3 > 0, W_4 > 0\} \\ &= \frac{\mathbb{P}\{W_1 > 0, W_2 > 0, W_3 > 0, W_4 > 0\}}{\mathbb{P}\{W_3 > 0, W_4 > 0\}} = 4\Phi_4(0; \varrho(\Sigma_W)). \end{aligned}$$

The proof is then complete upon noticing that  $\varrho(\Sigma_W) = P_\tau(\rho, \delta, v)$  stated in the lemma.  $\square$

**Lemma 5.** *Let  $(X_1, X_2) \sim \text{SN}(0, \varrho(\rho), \alpha)$  where  $\rho \in (-1, 1)$  and  $\alpha \in \mathbb{R}^2$ . Let  $(X_1^\circ, X_2^\circ)$  be a bivariate skew-normal vector satisfying  $(X_1^\circ, X_2^\circ) \perp (X_1, X_2)$ ,  $X_1^\circ \perp X_2^\circ$  and  $X_i^\circ \stackrel{d}{=} X_i$  for  $i=1,2$ . Moreover, let  $a_1, a_2, b$  be three strictly positive constants. We have*

$$\mathbb{P}\{a_1X_1^\circ > bX_1, a_2X_2^\circ > bX_2\} = 8\Phi_5(0; P_S(\rho, \delta, v)),$$

where  $P_S$  is defined by (5.2),  $\delta = (\delta_1, \delta_2)^\top = \varrho(\rho)\alpha/\sqrt{1 + \alpha^\top \varrho(\rho)\alpha}$ ,  $v = (v_1, v_2, v_1^-, v_2^-, v_3)$  with  $v_1 = a_1/\sigma_1$ ,  $v_2 = a_2/\sigma_2$ ,  $v_1^- = -b/\sigma_1$ ,  $v_2^- = -b/\sigma_2$  and  $v_3 = b^2/(\sigma_1\sigma_2)$ , where  $\sigma_1 = \sqrt{a_1^2 + b^2}$  and  $\sigma_2 = \sqrt{a_2^2 + b^2}$ .

*Proof.* Let  $Z = (Z_1, \dots, Z_7)^\top$  be a 7-dimensional normal random vector with mean zero and

covariance matrix (upper-triangular entries omitted)

$$P_Z = \left[ \begin{array}{cccc|ccc} 1 & & & & & & & \\ 0 & 1 & & & & & & \\ 0 & 0 & 1 & & & & & \\ 0 & 0 & \rho & 1 & & & & \\ \hline \delta_1 & 0 & 0 & 0 & 1 & & & \\ 0 & \delta_2 & 0 & 0 & 0 & 1 & & \\ 0 & 0 & \delta_1 & \delta_2 & 0 & 0 & 1 & \end{array} \right] = \begin{bmatrix} \mathbf{B} & \mathbf{C} \\ \mathbf{C}^\top & \mathbf{I}_3 \end{bmatrix}, \quad \mathbf{B} = \begin{bmatrix} \mathbf{I}_2 & 0 \\ 0 & \varrho(\rho) \end{bmatrix}, \quad \mathbf{C}^\top = \begin{bmatrix} \text{diag}(\delta) & 0 \\ 0 & \delta^\top \end{bmatrix}.$$

By properties of the multivariate normal distribution and the stochastic representation of the skew-normal distribution via conditioning (Azzalini, 2014, Section 5.1.3), we have the representations  $(X_1, X_2) \stackrel{d}{=} (Z_3, Z_4) | \{Z_7 > 0\}$ ,  $X_1^\circ \stackrel{d}{=} (Z_1 | Z_5 > 0)$  and  $X_2^\circ \stackrel{d}{=} (Z_2 | Z_6 > 0)$ . Next, for  $a_1, a_2, b > 0$ , define

$$\begin{bmatrix} Y_1 \\ Y_2 \end{bmatrix} =_d \left( \begin{bmatrix} a_1 Z_1 \\ a_2 Z_2 \end{bmatrix} - b \begin{bmatrix} Z_3 \\ Z_4 \end{bmatrix} \right) \Big| \{Z_5 > 0, Z_6 > 0, Z_7 > 0\}. \quad (\text{A.6})$$

Then, to prove the lemma it suffices to show that  $\mathbb{P}\{Y_1 > 0, Y_2 > 0\} = 8\Phi_5(0; P_S(\rho, \delta, v))$ . To see this, define a 5-dimensional normal random vector  $W = (W_1, \dots, W_5)^\top$  by

$$W = \begin{bmatrix} \mathbf{A} & 0 \\ 0 & \mathbf{I}_3 \end{bmatrix} Z, \quad \text{where } \mathbf{A} = \begin{bmatrix} a_1 & 0 & -b & 0 \\ 0 & a_2 & 0 & -b \end{bmatrix} = \begin{bmatrix} \text{diag}(a_1, a_2) & -b\mathbf{I}_2 \end{bmatrix},$$

so that

$$W_1 = a_1 Z_1 - b Z_3, \quad W_2 = a_2 Z_2 - b Z_4, \quad W_3 = Z_5, \quad W_4 = Z_6, \quad W_5 = Z_7. \quad (\text{A.7})$$

By construction,  $W \sim \mathbf{N}(0, \Sigma_W)$ , where

$$\Sigma_W = \begin{bmatrix} \mathbf{A}\mathbf{B}\mathbf{A}^\top & \mathbf{A}\mathbf{C} \\ \mathbf{C}^\top\mathbf{A}^\top & \mathbf{I}_3 \end{bmatrix} = \left[ \begin{array}{cc|ccc} a_1^2 + b^2 & b^2\rho & a_1\delta_1 & 0 & -b\delta_1 \\ b^2\rho & a_2^2 + b^2 & 0 & a_2\delta_2 & -b\delta_2 \\ \hline a_1\delta_1 & 0 & 1 & 0 & 0 \\ 0 & a_2\delta_2 & 0 & 1 & 0 \\ -b\delta_1 & -b\delta_2 & 0 & 0 & 1 \end{array} \right].$$

By (A.6) and (A.7), we have  $(Y_1, Y_2) \stackrel{d}{=} (W_1, W_2) | \{W_3 > 0, W_4 > 0, W_5 > 0\}$ , and then

$$\begin{aligned} \mathbb{P}\{Y_1 > 0, Y_2 > 0\} &= \mathbb{P}\{W_1 > 0, W_2 > 0 | W_3 > 0, W_4 > 0, W_5 > 0\} \\ &= \frac{\mathbb{P}\{W_1 > 0, W_2 > 0, W_3 > 0, W_4 > 0, W_5 > 0\}}{\mathbb{P}\{W_3 > 0, W_4 > 0, W_5 > 0\}} = 8 \Phi_5(0; \varrho(\Sigma_W)). \end{aligned}$$

The proof is then complete upon noticing that  $\varrho(\Sigma_W) = P_S(\rho, \delta, v)$  stated in the lemma.  $\square$

**Lemma 6.** *Let  $P$  be a  $d \times d$  correlation matrix whose upper-left  $2 \times 2$  sub-matrix is given by  $\varrho(\rho)$  with  $\rho \in (-1, 1)$ . Then, we have*

$$\frac{\partial \Phi_d(0; P)}{\partial \rho} = \begin{cases} (2\pi\sqrt{1-\rho^2})^{-1} > 0, & d = 2, \\ (2\pi\sqrt{1-\rho^2})^{-1} \Phi_{d-2}(0; P/\varrho(\rho)) > 0, & d > 2, \end{cases}$$

where  $P/\varrho(\rho)$  is the Schur complement of  $\varrho(\rho)$  in  $P$ .

*Proof.* In the case of  $d = 2$ , we have  $P = \varrho(\rho)$  and hence  $\Phi_2(0; P) = \Phi_2(0; \rho)$  and  $\phi_2(x; P) = \phi_2(x; \rho)$ . Then,

$$\frac{\partial \Phi_2(0; \rho)}{\partial \rho} = \int_{(-\infty, 0]^2} \frac{\partial \phi_2(x; \rho)}{\partial \rho} dx = \int_{(-\infty, 0]^2} \frac{\partial^2 \phi_2(x; \rho)}{\partial x_1 \partial x_2} dx = \phi_2(0, 0; \rho) = \frac{1}{2\pi\sqrt{1-\rho^2}} > 0,$$

where the second equality above holds due to  $\partial \phi(x; \rho)/\partial \rho = \partial^2 \phi(x; \rho)/\partial x_1 \partial x_2$  (see e.g. Equation (3) in Plackett, 1954). If  $d > 2$ , note that

$$\frac{\partial \Phi_d(0; P)}{\partial \rho} = \int_{(-\infty, 0]^d} \frac{\partial^2 \phi_d(x; P)}{\partial x_1 \partial x_2} dx = \int_{(-\infty, 0]^{d-2}} \phi_d(0, 0, x_3, \dots, x_d; P) dx_3 \cdots dx_d,$$

and then, by the conditioning property of multivariate normal distributions, we deduce that

$$\phi_d(0, 0, x_3, \dots, x_d; P) = \phi_2(0, 0; \rho) \phi_{d-2}(x_3, \dots, x_d; P/\varrho(\rho)) = \frac{\phi_{d-2}(x_3, \dots, x_d; P/\varrho(\rho))}{2\pi\sqrt{1-\rho^2}}.$$

This leads to the claim in the lemma for  $d > 2$  case, and completes the proof of the lemma.  $\square$

**Lemma 7.** *Let  $P_\tau(\rho, \delta, v)$  be the correlation matrix-valued function defined in (5.1), where  $\rho \in (-1, 1)$ ,  $\delta = (\delta_1, \delta_2)$  and  $v = (v_1, v_2)$ , and let  $V = (V_1, V_2)$  be the random vector defined in Theorem 2 (i). We have*

$$\frac{\partial \Phi_4(0; P_\tau(\rho, \delta, V))}{\partial \delta_1} \stackrel{d}{=} \frac{V_1 \arcsin \tilde{V}(\delta_1, \delta_2)}{2\pi^2 \sqrt{1 - V_1^2 \delta_1^2}}$$

and

$$\frac{\partial \Phi_4(0; P_\tau(\rho, \delta, V))}{\partial \delta_2} \stackrel{d}{=} \frac{V_1 \arcsin \tilde{V}(\delta_2, \delta_1)}{2\pi^2 \sqrt{1 - V_1^2 \delta_2^2}},$$

where

$$\tilde{V}(\delta_1, \delta_2) := \frac{V_2(\delta_2 - \rho\delta_1)}{\sqrt{1 - \delta_1^2} \sqrt{1 - \rho^2 - (\delta_1^2 - 2\rho\delta_1\delta_2 + \delta_2^2)V_1^2}}.$$

*Proof.* Consider the following four matrices:

$$P_a = \left[ \begin{array}{cc|cc} 1 & & & \\ a & 1 & & \\ \rho & b & 1 & \\ c & 0 & d & 1 \end{array} \right], \quad P_b = \left[ \begin{array}{cc|cc} 1 & & & \\ b & 1 & & \\ \rho & a & 1 & \\ d & 0 & c & 1 \end{array} \right],$$

$$P_c = \left[ \begin{array}{cc|cc} 1 & & & \\ c & 1 & & \\ a & 0 & 1 & \\ \rho & d & b & 1 \end{array} \right], \quad P_d = \left[ \begin{array}{cc|cc} 1 & & & \\ d & 1 & & \\ b & 0 & 1 & \\ \rho & c & a & 1 \end{array} \right],$$

where  $a = v_1\delta_1$ ,  $b = v_1\delta_2$ ,  $c = v_2\delta_1$  and  $d = v_2\delta_2$ . It is clear that

$$\Phi_4(0; P_\tau(\rho, \delta, v)) = \Phi_4(0; P_a) = \Phi_4(0; P_b) = \Phi_4(0; P_c) = \Phi_4(0; P_d),$$

and, by total differentiation and Lemma 6, we can deduce that

$$\frac{\partial}{\partial \delta_1} \Phi_4(0; P_\tau) = \frac{v_1 \Phi_2(0; P_a/\varrho(a))}{2\pi\sqrt{1-a^2}} + \frac{v_2 \Phi_2(0; P_c/\varrho(c))}{2\pi\sqrt{1-c^2}}, \quad (\text{A.8})$$

$$\frac{\partial}{\partial \delta_2} \Phi_4(0; P_\tau) = \frac{v_1 \Phi_2(0; P_b/\varrho(b))}{2\pi\sqrt{1-b^2}} + \frac{v_2 \Phi_2(0; P_d/\varrho(d))}{2\pi\sqrt{1-d^2}}. \quad (\text{A.9})$$

Some matrix algebra yields

$$\Phi_2(0; P_x/\varrho(x)) = \frac{1}{4} + \frac{\arcsin u_x(a, b, c, d)}{2\pi} \quad (x = a, b, c, d), \quad (\text{A.10})$$

where

$$\begin{aligned}
u_a(a, b, c, d) &= \frac{d(1 - a^2) - c(\rho - ab)}{\sqrt{1 - a^2 - c^2} \sqrt{1 - \rho^2 - (a^2 - 2\rho ab + b^2)}}, \\
u_b(a, b, c, d) &= \frac{c(1 - b^2) - d(\rho - ab)}{\sqrt{1 - b^2 - d^2} \sqrt{1 - \rho^2 - (a^2 - 2\rho ab + b^2)}}, \\
u_c(a, b, c, d) &= \frac{b(1 - c^2) - a(\rho - cd)}{\sqrt{1 - a^2 - c^2} \sqrt{1 - \rho^2 - (c^2 - 2\rho cd + d^2)}}, \\
u_d(a, b, c, d) &= \frac{a(1 - d^2) - b(\rho - cd)}{\sqrt{1 - b^2 - d^2} \sqrt{1 - \rho^2 - (c^2 - 2\rho cd + d^2)}}.
\end{aligned}$$

Using the fact that  $(V_1, V_2) \stackrel{d}{=} -(V_2, V_1)$ , we can verify:

$$\tilde{V}(\delta_1, \delta_2) = u_a(V_1\delta_1, V_1\delta_2, V_2\delta_1, V_2\delta_2) \stackrel{d}{=} -u_c(V_1\delta_1, V_1\delta_2, V_2\delta_1, V_2\delta_2), \quad (\text{A.11})$$

$$\tilde{V}(\delta_2, \delta_1) = u_b(V_1\delta_1, V_1\delta_2, V_2\delta_1, V_2\delta_2) \stackrel{d}{=} -u_d(V_1\delta_1, V_1\delta_2, V_2\delta_1, V_2\delta_2). \quad (\text{A.12})$$

Noticing  $V_1 \stackrel{d}{=} -V_2$ , the first statement in the lemma holds due to (A.8), (A.10), (A.11), and the second statement holds due to (A.9), (A.10), (A.12).  $\square$

**Lemma 8.** *Let  $P(u)$  be a  $4 \times 4$  correlation matrix-valued function defined as follows:*

$$P(u) = \begin{bmatrix} \varrho(u) & L^\top \\ L & R \end{bmatrix}, \quad u \in (-1, 1),$$

where  $L$  and  $R$  are  $2 \times 2$  matrices that ensure  $P(u)$  is a proper correlation matrix. Given  $\rho \in (-1, 1)$  such that  $P(\rho)$  is a proper correlation matrix, we have

$$\Phi_4(0; P(\rho)) - \Phi_0(0; P(0)) = \frac{\arcsin \rho}{4\pi} + \frac{1}{4\pi^2} \int_{\pi/2}^{\arccos \rho} \arccos \varrho(S_\theta) d\theta,$$

where, for any  $\theta \in [0, \pi]$ ,  $S_\theta := R - L\varrho(\cos \theta)^{-1}L^\top$  is a  $2 \times 2$  covariance matrix.

*Proof.* To prove the lemma, we first denote by  $P_{22,1}(u)$  the Schur complement of  $\varrho(u)$  in  $P(u)$  for any  $u \in (-1, 1)$ . Next, we define a real-valued function  $f(t) = \Phi_4(0; P(u(t)))$  where  $u(t) = \rho t$  for  $t \in [0, 1]$ . It follows that

$$\Phi_4(0; P(\rho)) - \Phi_0(0; P(0)) = f(1) - f(0) = \int_0^1 f'(t) dt,$$



and

$$f'(t) = \frac{\partial \Phi_4(0; P(u(t)))}{\partial u} \cdot u'(t) = \frac{\rho \cdot \Phi_2(0; P_{22.1}(\rho t))}{2\pi\sqrt{1-\rho^2 t^2}},$$

where the second equality holds from Lemma 6 and  $u(t) = \rho t$ . Consequently, we have

$$\begin{aligned} \Phi_4(0; P(\rho)) - \Phi_0(0; P(0)) &= \int_0^1 \frac{\rho \cdot \Phi_2(0; P_{22.1}(\rho t))}{2\pi\sqrt{1-\rho^2 t^2}} dt = \frac{1}{2\pi} \int_0^\rho \frac{\Phi_2(0; P_{22.1}(s))}{\sqrt{1-s^2}} ds \\ &= -\frac{1}{2\pi} \int_{\pi/2}^{\arccos \rho} \Phi_2(0; P_{22.1}(\cos \theta)) d\theta, \end{aligned}$$

which further simplifies to

$$-\frac{1}{4\pi^2} \int_{\pi/2}^{\arccos \rho} \arccos -\varrho(S_\theta) d\theta,$$

by the fact that  $P_{22.1}(\cos \theta) = S_\theta$  and  $\Phi_2(0; \Sigma) = (2\pi)^{-1} \arccos -\varrho(\Sigma)$  for any  $2 \times 2$  covariance matrix  $\Sigma$ . The claim in the lemma then follows immediately after applying the identities  $\arccos(-x) = \pi - \arccos(x)$  and  $\arcsin(x) + \arccos(x) = \pi/2$  to the above expression.  $\square$

## B Proofs of Theorems and Propositions in Section 4

**Proof of Theorem 1.** Since Kendall's and Spearman's rank correlations depend solely on the copula of  $X_1$  and  $X_2$ , it suffices to consider  $X = (X_1, X_2)^\top \sim \text{MN}(0, \varrho(\rho), \beta, F)$  which has the representation  $X = W\beta + \sqrt{W}Z$  for some  $Z \sim \text{N}(0, \varrho(\rho))$ ,  $W \sim F$ , and  $W$  is independent of  $Z$ .

To derive Kendall's tau of  $X_1$  and  $X_2$  stated in Part (i) of the theorem, we start by defining  $X^* := W^*\beta + \sqrt{W^*}Z^*$  where  $W^*$  and  $Z^*$  are independent copies of  $W$  and  $Z$ , respectively, and  $W^*$  is independent of  $Z^*$ . Then, by construction,  $X^*$  is an independent copy of  $X$ , and

$$\begin{bmatrix} X^* \\ X \end{bmatrix} \Big| \{W, W^*\} \sim \text{N} \left( \begin{bmatrix} W^*\beta \\ W\beta \end{bmatrix}, \begin{bmatrix} W^*\varrho(\rho) & 0 \\ 0 & W\varrho(\rho) \end{bmatrix} \right). \quad (\text{B.1})$$

Now, we define  $Y = (Y_1, Y_2)^\top := X - X^*$ . Note that, by Lemma 1 (i), Kendall's tau of  $X_1$  and  $X_2$  is given by  $\tau = 4\mathbb{P}\{Y_1 < 0, Y_2 < 0\} - 1$ . To obtain the joint probability of  $Y_1 < 0$  and  $Y_2 < 0$ , we first deduce from (B.1) that

$$Y \mid \{W, W^*\} \sim \text{N}((W - W^*)\beta, (W + W^*)\varrho(\rho)), \quad (\text{B.2})$$

and then note that  $\mathbb{P}\{Y_1 < 0, Y_2 < 0 | W, W^*\} = \mathbb{P}\{Z_1 < V\beta_1, Z_2 < V\beta_2 | W, W^*\}$ , with  $Z_i := [Y_i - (W - W^*)\beta_i] / \sqrt{W + W^*}$  for  $i = 1, 2$ , and  $V$  is defined in Part (i) of the theorem. Since (B.2) implies that  $(Z_1, Z_2)^\top | \{W, W^*\} \sim N(0, \varrho(\rho))$ , it then follows that

$$\mathbb{P}\{Y_1 < 0, Y_2 < 0 | W, W^*\} = \mathbb{P}\{Z_1 < V\beta_1, Z_2 < V\beta_2 | W, W^*\} = \Phi_2(V\beta_1, V\beta_2; \rho).$$

By the law of iterated expectations, we have  $\mathbb{P}\{Y_1 < 0, Y_2 < 0\} = \mathbb{E} \Phi_2(V\beta_1, V\beta_2; \rho)$ . Part (i) of the theorem then holds immediately due to Lemma 1 (i).

To derive Spearman's rho of  $X_1$  and  $X_2$  stated in Part (ii) of the theorem, we define  $X^\circ := (X_1^\circ, X_2^\circ)^\top$  with  $X_i^\circ = \beta_i W_i + \sqrt{W_i} Z_i^\circ$  for  $i = 1, 2$ , where  $W_1, W_2 \sim \text{i.i.d. } F$  and are independent of  $W$ , and  $Z_1^\circ, Z_2^\circ \sim \text{i.i.d. } N(0, 1)$  and are independent of  $W_1, W_2, W$  and  $Z$ . Then, by construction,  $X^\circ$  is independent of  $X$ , has independent margins, and  $X_i^\circ \stackrel{d}{=} X_i$  for  $i = 1, 2$ . Moreover,

$$\begin{bmatrix} X^\circ \\ X \end{bmatrix} \Big| \{W, W_1, W_2\} \sim N \left( \begin{bmatrix} W_1 \beta_1 \\ W_2 \beta_2 \\ W \beta \end{bmatrix}, \begin{bmatrix} \text{diag}(W_1, W_2) & 0 \\ 0 & W \varrho(\rho) \end{bmatrix} \right).$$

Next, we define  $Y^\circ = (Y_1^\circ, Y_2^\circ)^\top := X - X^\circ$ . By Lemma 1 (ii), Spearman's tau of  $X_1$  and  $X_2$  is given by  $\rho_S = 12 \mathbb{P}\{Y_1^\circ < 0, Y_2^\circ < 0\} - 3$ . To obtain the joint probability of  $Y_1^\circ < 0$  and  $Y_2^\circ < 0$ , we first note that  $Y^\circ | \{W, W_1, W_2\} \sim N(\mu_Y, \Sigma_Y)$  where

$$\mu_Y = \begin{bmatrix} (W - W_1)\beta_1 \\ (W - W_2)\beta_2 \end{bmatrix}, \quad \Sigma_Y = \begin{bmatrix} W_1 & 0 \\ 0 & W_2 \end{bmatrix} + W \begin{bmatrix} 1 & \rho \\ \rho & 1 \end{bmatrix} = \begin{bmatrix} W_1 + W & W\rho \\ W\rho & W_2 + W \end{bmatrix}.$$

Define  $Z_i^\circ = [Y_i^\circ - (W - W_i)\beta_i] / \sqrt{W + W_i}$  for  $i = 1, 2$ . It then follows that

$$(Z_1^\circ, Z_2^\circ)^\top | \{W, W_1, W_2\} \sim N(0, \varrho(V_3\rho)) \quad \text{where} \quad V_3 = W / \sqrt{(W_1 + W)(W_2 + W)},$$

and, for  $V_i := (W_i - W) / \sqrt{W_i + W}$  for  $i = 1, 2$ ,

$$\begin{aligned} \mathbb{P}\{Y_1^\circ < 0, Y_2^\circ < 0 | W, W_1, W_2\} &= \mathbb{P}\{Z_1^\circ < V_1\beta_1, Z_2^\circ < V_2\beta_2 | W, W_1, W_2\} \\ &= \Phi_2(V_1\beta_1, V_2\beta_2; V_3\rho). \end{aligned}$$

By the law of iterated expectations, we have  $\mathbb{P}\{Y_1^\circ < 0, Y_2^\circ < 0\} = \mathbb{E} \Phi_2(V_1\beta_1, V_2\beta_2; V_3\rho)$ . Part (ii) of the theorem then holds immediately due to Lemma 1 (ii).  $\square$

**Proof of Proposition 1.** Part (i) of the proposition follows directly from the expressions of  $\tau$  and  $\rho_S$  given in Theorem 1 and the exchangeability of the bivariate normal cdf:  $\Phi_2(x_1, x_2; \rho) = \Phi_2(x_2, x_1; \rho)$ .

Next, we prove Part (ii) of the proposition. First,  $\tau(\rho, \beta, F) = \tau(\rho, -\beta, F)$  follows directly from the expression of  $\tau$  in (4.1) and the fact that  $V \stackrel{d}{=} -V$ . To show that  $\rho_S(\rho, \beta, F) = \rho_S(\rho, -\beta, F)$ , it suffices to prove that

$$\mathbb{E} \Phi_2(\beta_1 V_1, \beta_2 V_2; \rho V_3) = \mathbb{E} \Phi_2(-\beta_1 V_1, -\beta_2 V_2; \rho V_3), \quad (\text{B.3})$$

where the mixing variables  $(V_1, V_2, V_3)$  are defined in Theorem 1 (ii). Evaluating the decomposition of  $\Phi_2(\cdot; \rho)$  in (4.7) at  $(-x_1, -x_2)$  and then using the identity  $T(h, a) = T(-h, a)$ , see e.g. (B.19) in Azzalini (2014, p. 235), we have

$$\begin{aligned} \Phi_2(-x_1, -x_2; \rho) &= \frac{1}{2} [\Phi(-x_1) + \Phi(-x_2)] - a(-x_1, -x_2) \\ &\quad - T\left(x_1, \frac{x_2 - \rho x_1}{x_1 \sqrt{1 - \rho^2}}\right) - T\left(x_2, \frac{x_1 - \rho x_2}{x_2 \sqrt{1 - \rho^2}}\right). \end{aligned} \quad (\text{B.4})$$

Since the last two terms in (B.4) and in (4.7) are identical, equation (B.3) holds if  $\mathbb{E} \Phi(\beta_i V_i) = \mathbb{E} \Phi(-\beta_i V_i)$  for  $i = 1, 2$ , and  $\mathbb{E} a(\beta_1 V_1, \beta_2 V_2) = \mathbb{E} a(-\beta_1 V_1, -\beta_2 V_2)$ . Since these conditions are guaranteed by the fact that  $V_i \stackrel{d}{=} -V_i$  for  $i = 1, 2$  and by the definition of function  $a$ , we can show that (B.3) holds, and hence  $\rho_S(\rho, \beta, F) = \rho_S(\rho, -\beta, F)$ .

Lastly, we prove Part (iii) of the proposition. It follows from  $\partial \Phi_2(x_1, x_2; \rho) / \partial \rho = \phi_2(x_1, x_2; \rho)$ , given in (4.9), that  $\partial \tau / \partial \rho = 4 \mathbb{E} \phi_2(\beta_1 V, \beta_2 V; \rho)$  and  $\partial \rho_S / \partial \rho = 12 \mathbb{E} V_3 \phi_2(\beta_1 V_1, \beta_2 V_2; \rho V_3)$ . Both partial derivatives are strictly positive because the normal pdf  $\phi_2$  is strictly positive and  $V_3$  is a strictly positive random variable. Therefore, the statements in Part (iii) are proved.  $\square$

**Proof of Corollary 1.** The results follow immediately from Theorem 1 and the definition of GH skew- $t$  copulas.  $\square$

**Proof of Proposition 2.** We first prove Part (i) and the first half of Part (iii). The expression of  $\tau$  in Part (i) follows directly from Theorem 1 (i), Lemma 2. For the partial derivatives, we have

$$\frac{\partial}{\partial b} \tau(\rho, b, F) = 4 \mathbb{E} \frac{\partial}{\partial b} \Phi^s(bV; \alpha_\rho) = 4 \mathbb{E} V \phi^s(bV; \alpha_\rho) > 0,$$

where the inequality holds due to Lemma 3 (i). Furthermore, Lemma 3 (ii) implies that  $\partial^2\tau/\partial b\partial\alpha_\rho > 0$ . Since  $\partial\alpha_\rho/\partial\rho = -(1-\rho)^{-1/2}(1+\rho)^{-3/2} < 0$ , we then have  $\partial^2\tau/\partial b\partial\rho < 0$ . This completes the proof of Part (i). To show that  $\tau(1, b, F) = 1$  stated in Part (iii), note that  $\alpha_\rho = 0$  when  $\rho = 1$ . Thus, we have

$$\tau(1, b, F) = 4 \mathbb{E} \Phi^s(bV; 0) - 1 = 4 \mathbb{E} \Phi(bV) - 1 = 4 \times \frac{1}{2} - 1 = 1.$$

In the above derivation,  $\mathbb{E} \Phi(bV) = 1/2$  because  $\Phi(x) + \Phi(-x) = 1$  for all  $x \in \mathbb{R}$  and  $V \stackrel{d}{=} -V$ .

The expression for  $\rho_S$  in Part (ii) of the proposition follows directly from Theorem 1 (ii). To show that  $\rho_S(1, b, F) = 1$  stated in the second half of Part (iii) of the proposition, it suffices to demonstrate that  $\mathbb{E} \Phi_2(bV_1, bV_2, V_3) = 1/3$  for all  $b$  and  $F$ . To this end, we apply the decomposition of the bivariate normal cdf given in (4.7) to get

$$\begin{aligned} \Phi_2(bV_1, bV_2; V_3) &= \frac{1}{2} [\Phi(bV_1) + \Phi(bV_2)] - a(bV_1, bV_2) \\ &\quad - T\left(bV_1, \frac{V_2/V_1 - V_3}{\sqrt{1 - V_3^2}}\right) - T\left(bV_2, \frac{V_1/V_2 - V_3}{\sqrt{1 - V_3^2}}\right). \end{aligned}$$

Taking expectations on both sides of the above equation and noting that  $(V_1, V_2, V_3) \stackrel{d}{=} (V_2, V_1, V_3)$ , we obtain

$$\mathbb{E} \Phi_2(bV_1, bV_2; V_3) = \mathbb{E} \Phi(bV_1) - \mathbb{E} T(S_1, S_2) - \frac{1}{2} \mathbb{P}\{V_1 V_2 < 0\}, \quad (\text{B.5})$$

where  $S_1 := bV_1$  and  $S_2 := (V_2/V_1 - V_3)/\sqrt{1 - V_3^2}$ .

We analyze the terms on the right-hand side of (B.5) subsequently. First,  $\mathbb{E} \Phi(bV_1) = 1/2$  because  $bV_1 \stackrel{d}{=} -bV_1$  and  $\Phi(x) + \Phi(-x) = 1$  for all  $x \in \mathbb{R}$ . For the second term, note that, by the relationship between  $(V_1, V_2, V_3)$  and  $(W_1, W_2, W_3) \sim \text{i.i.d. } F$ , and the definitions of  $S_1$  and  $S_2$ , we deduce that  $(S_1, S_2) \stackrel{d}{=} (-S_1, -S_2)$ . Using the properties of Owen's T function, specifically  $T(h, -a) = -T(h, a)$  and  $T(-h, a) = T(h, a)$ , we have  $T(S_1, S_2) \stackrel{d}{=} T(-S_1, -S_2) = -T(S_1, S_2)$ . Consequently,  $\mathbb{E} T(S_1, S_2) = -\mathbb{E} T(S_1, S_2)$ , which implies  $\mathbb{E} T(S_1, S_2) = 0$ . Lastly, consider

$$\begin{aligned} \frac{1}{2} \mathbb{P}\{V_1 V_2 < 0\} &= \frac{1}{2} \mathbb{P}\{(W_1 - W_3)(W_2 - W_3) < 0\} = \mathbb{P}\{W_1 < W_3, W_2 > W_3\} \\ &= \mathbb{E} \mathbb{P}\{W_1 < W_3, W_2 > W_3 \mid W_3\} \\ &= \mathbb{E} F(W_3)(1 - F(W_3)) = \frac{1}{6}, \end{aligned}$$

where we used the fact that  $W_1, W_2, W_3 \sim \text{i.i.d. } F$  and  $F(W) \sim U(0, 1)$ . Combining these

results, we obtain  $\mathbb{E} \Phi_2(bV_1, bV_2; V_3) = 1/2 - 0 - 1/6 = 1/3$ , as desired. This completes the proof of Part (iii) of the proposition, and the proof of the proposition is complete.  $\square$

**Proof of Proposition 3.** Part (i) of the proposition follows immediately from substituting  $\beta_1 = b$  and  $\beta_2 = 0$  in the formulas of Kendall's tau and Spearman's rho in Theorem 1.

Next, we prove Part (ii). By equation (4.6) and a change of variables,

$$\Phi_2(x_1, x_2; \rho) = \Phi(x_1)\Phi(x_2) + \frac{1}{2\pi} \int_0^{\arcsin \rho} \exp\left(-\frac{x_1^2 + x_2^2 - 2x_1x_2 \sin \theta}{2 \cos^2 \theta}\right) d\theta,$$

from which we can easily deduce that

$$\Phi_2(0, x; \rho) + \Phi_2(0, x; -\rho) = \Phi(x), \quad (\text{B.6})$$

for any  $x \in \mathbb{R}$  and  $\rho \in (-1, 1)$ . Then, by Theorem 1 (i) and (B.6),

$$\begin{aligned} \tau(-\rho, (0, b), F) &= 4 \mathbb{E} \Phi_2(0, bV; -\rho) - 1 \\ &= 4 \mathbb{E} \Phi(bV) - 4 \mathbb{E} \Phi_2(0, bV; \rho) - 1 \\ &= 4 \mathbb{E} \Phi(bV) - 2 - \tau(\rho, (0, b), F). \end{aligned}$$

Since  $V \stackrel{d}{=} -V$  and  $\Phi(x) + \Phi(-x) = 1$ , we have  $\mathbb{E} \Phi(bV) = 1/2$ . It then follows immediately that  $\tau(-\rho, (0, b), F) = -\tau(\rho, (0, b), F)$ . Similarly, we can show  $\rho_S(-\rho, (0, b), F) = -\rho_S(\rho, (0, b), F)$  by using Theorem 1 (ii) and (B.6) and noticing that  $V_2 \stackrel{d}{=} -V_2$ . This completes the proof of Part (ii) of the proposition.

To prove Part (iii), it suffices to show that  $\tau(\rho, (0, b), F)$  and  $\rho_S(\rho, (0, b), F)$  are both strictly increasing in  $b$  for  $b > 0$  and  $\rho < 0$ , provided Part (ii) of the proposition and the fact that  $\tau$  and  $\rho_S$  are invariant under the sign change of the skewness vector. Taking derivatives with respect to  $\beta_1$  on both sides of (4.1) and (4.2) and using the partial derivative given by (4.8) yields

$$\begin{aligned} \frac{\partial}{\partial \beta_1} \tau(\rho, \beta, F) &= 4 \mathbb{E} V \phi(\beta_1 V) \Phi\left(\frac{(\beta_2 - \rho \beta_1) V}{\sqrt{1 - \rho^2}}\right), \\ \frac{\partial}{\partial \beta_1} \rho_S(\rho, \beta, F) &= 12 \mathbb{E} V_1 \phi(\beta_1 V_1) \Phi\left(\frac{\beta_2 V_2 - \rho \beta_1 V_1 V_3}{\sqrt{1 - \rho^2 V_3^2}}\right). \end{aligned}$$

Evaluated at  $\beta_1 = b$  and  $\beta_2 = 0$ , together with the definition of  $\phi^s$ , we obtain:

$$\frac{\partial}{\partial b} \tau(\rho, (0, b), F) = 2 \mathbb{E} V \phi^s(bV; \alpha(\rho)) \quad \text{and} \quad \frac{\partial}{\partial b} \rho_S(\rho, (0, b), F) = 6 \mathbb{E} V_1 \phi^s(bV_1; \alpha(\rho V_3)),$$

where  $\alpha(r) := -r/\sqrt{1-r^2}$  for any  $r \in (-1, 1)$ . Since  $\rho < 0$  and  $V_3 > 0$  almost surely,  $\alpha(\rho) > 0$  and  $\alpha(\rho V_3) > 0$  almost surely. Moreover, as both  $V$  and  $V_1$  are symmetrically distributed, Lemma 3 (i) applies so that both partial derivatives above are strictly positive. This proves Part (iii) of the proposition.  $\square$

## C Proofs of Theorems and Propositions in Section 5

**Proof of Theorem 2.** Since Kendall's and Spearman's rank correlations depend solely on the copula of  $X_1$  and  $X_2$ , it suffices to consider  $X = (X_1, X_2) \sim \text{MSN}(0, \varrho(\rho), \alpha, F)$  which has the representation  $X = \sqrt{W}Z$  for some  $Z = (Z_1, Z_2) \sim \text{SN}(0, \varrho(\rho), \alpha)$ ,  $W \sim F$ , and  $W$  is independent of  $Z$ .

To derive Kendall's tau of  $X_1$  and  $X_2$  stated in Part (i) of the theorem, we start by defining  $X^* = (X_1^*, X_2^*)^\top := \sqrt{W^*}Z^*$  where  $W^*$  and  $Z^*$  are independent copies of  $W$  and  $Z$ , respectively, and  $W^*$  is independent of  $Z^*$ . Then, by construction,  $X^*$  is an independent copy of  $X$ . Then, by Lemma 1 (i), Kendall's tau of  $X_1$  and  $X_2$  is given by

$$\begin{aligned} \tau &= 4 \mathbb{P}\{X_1 < X_1^*, X_2 < X_2^*\} - 1 \\ &= 4 \mathbb{E} \mathbb{P}\{\sqrt{W}Z_1 < \sqrt{W^*}Z_1^*, \sqrt{W}Z_2 < \sqrt{W^*}Z_2^* \mid W, W^*\} - 1. \end{aligned} \quad (\text{C.1})$$

Applying Lemma 4 to  $(Z_1, Z_2)$  and  $(Z_1^*, Z_2^*)$ , we have

$$\mathbb{P}\{\sqrt{W}Z_1 < \sqrt{W^*}Z_1^*, \sqrt{W}Z_2 < \sqrt{W^*}Z_2^* \mid W, W^*\} = 4 \Phi_4(0; P_\tau(\rho, \delta, S)), \quad (\text{C.2})$$

where  $S = (\sqrt{W^*/(W^* + W)}, -\sqrt{W/(W^* + W)})$ . Part (i) of the theorem then holds due to (C.1), (C.2), and the fact that  $S \stackrel{d}{=} V$  for  $V$  being defined in Part (i) of the theorem.

To derive Spearman's rho of  $X_1$  and  $X_2$  given in Part (ii) of the theorem, we define  $(Z_1^\circ, Z_2^\circ)$  to be a bivariate skew-normal vector satisfying  $(Z_1^\circ, Z_2^\circ) \perp (Z_1, Z_2)$ ,  $Z_1^\circ \perp Z_2^\circ$  and  $Z_i^\circ \stackrel{d}{=} Z_i$  for  $i = 1, 2$ . Next, define  $X_1^\circ := \sqrt{W_1}Z_1^\circ$  and  $X_2^\circ := \sqrt{W_2}Z_2^\circ$ , where  $W_1, W_2 \sim \text{i.i.d. } F$  and are independent of  $W$ ,  $Z_1^\circ$  and  $Z_2^\circ$ . By construction,  $(X_1^\circ, X_2^\circ) \perp (X_1, X_2)$ ,  $X_1^\circ \perp X_2^\circ$ , and  $X_i^\circ \stackrel{d}{=} X_i$

for  $i = 1, 2$ . By Lemma 1 (ii), Spearman's rho of  $X_1$  and  $X_2$  is

$$\begin{aligned}\rho_S &= 12 \mathbb{P}\{X_1 < X_1^\circ, X_2 < X_2^\circ\} - 3 \\ &= 12 \mathbb{E} \mathbb{P}\{\sqrt{W}Z_1 < \sqrt{W_1}Z_1^\circ, \sqrt{W}Z_2 < \sqrt{W_2}Z_2^\circ \mid W, W_1, W_2\} - 3.\end{aligned}\quad (\text{C.3})$$

Applying Lemma 5 to  $(Z_1, Z_2)$  and  $(Z_1^\circ, Z_2^\circ)$ , we have

$$\mathbb{P}\{\sqrt{W}Z_1 < \sqrt{W_1}Z_1^\circ, \sqrt{W}Z_2 < \sqrt{W_2}Z_2^\circ \mid W, W_1, W_2\} = 8\Phi_5(0; P_S(\rho, \delta, T)), \quad (\text{C.4})$$

where  $T \stackrel{d}{=} V$  for  $V$  being defined in Part (ii) of the theorem. Then, Part (ii) of the theorem follows from (C.3) and (C.4), and the proof of the whole theorem is complete.  $\square$

**Proof of Corollary 2.** We prove the alternative expression (5.7) for Kendall's tau first. Define a four-dimensional random vector  $X = (X_1, X_2, X_3, X_4)$  satisfying  $X \mid V \sim N(0, P_\tau(\rho, \delta, V))$  where  $P_\tau$  is defined by (5.1). Then, we have

$$\begin{aligned}\Phi_4(0; P_\tau(\rho, \delta(\rho, \alpha), V)) &= \mathbb{P}\{X_1 < 0, X_2 < 0, X_3 < 0, X_4 < 0 \mid V\} \\ &= \frac{1}{4} \mathbb{P}\{X_1 < 0, X_2 < 0 \mid X_3 < 0, X_4 < 0, V\},\end{aligned}\quad (\text{C.5})$$

where in the second equality we used  $\mathbb{P}\{X_3 < 0, X_4 < 0 \mid V\} = (1/2)^2 = 1/4$ , since  $(X_3, X_4) \mid V \sim N(0, I_2)$ . Consequently, by the original expression (5.3) for Kendall's tau and (C.5), we have

$$\tau(\rho, \alpha, F) = 4 \mathbb{E} \mathbb{P}\{X_1 < 0, X_2 < 0 \mid X_3 < 0, X_4 < 0, V\} - 1. \quad (\text{C.6})$$

Using the conditioning property of the multivariate normal distribution and the identity  $V_1^2 + V_2^2 = 1$ , we can deduce that  $(X_1, X_2) \mid \{X_3, X_4, V\} \sim N(\mu, \Sigma)$ , where

$$\mu = \begin{bmatrix} (V_1 X_3 + V_2 X_4) \delta_1 \\ (V_1 X_3 + V_2 X_4) \delta_2 \end{bmatrix} \quad \text{and} \quad \Sigma = \begin{bmatrix} 1 - \delta_1^2 & \rho - \delta_1 \delta_2 \\ \rho - \delta_1 \delta_2 & 1 - \delta_2^2 \end{bmatrix}.$$

Therefore, we have

$$\mathbb{P}\{X_1 < 0, X_2 < 0 \mid X_3, X_4, V\} = \Phi_2(-\alpha_1^\dagger(V_1 X_3 + V_2 X_4), -\alpha_2^\dagger(V_1 X_3 + V_2 X_4); \rho^\dagger), \quad (\text{C.7})$$

where  $\alpha_1^\dagger, \alpha_2^\dagger$  and  $\rho^\dagger$  are defined by (5.5) and (5.6). Since  $(X_3, X_4) \mid V$  is a pair of independent standard normal random variates, equation (5.7) stated in the corollary follows from the definition of  $(Y_1, Y_2)$ , along with (C.6) and (C.7).

Next, we prove the alternative expression (5.8) for Spearman's rho. Define a five-dimensional random vector  $X = (X_1, X_2, X_3, X_4, X_5)$  satisfying  $X | V \sim N(0, P_S(\rho, \delta, V))$  where  $P_S$  is defined by (5.2). Then, we have

$$\begin{aligned} \Phi_5(0; P_S(\rho, \delta(\rho, \alpha), V)) &= \mathbb{P}\{X_1 < 0, X_2 < 0, X_3 < 0, X_4 < 0, X_5 < 0 | V\} \\ &= \frac{1}{8} \mathbb{P}\{X_1 < 0, X_2 < 0 | X_3 < 0, X_4 < 0, X_5 < 0, V\}, \end{aligned} \quad (\text{C.8})$$

where in the second equality we used  $\mathbb{P}\{X_3 < 0, X_4 < 0, X_5 < 0 | V\} = (1/2)^3 = 1/8$ , since  $(X_3, X_4, X_5) | V \sim N(0, I_3)$ . Consequently, it follows from (5.4) and (C.8) that

$$\rho_S(\rho, \alpha, F) = 12 \mathbb{E} \mathbb{P}\{X_1 < 0, X_2 < 0 | X_3 < 0, X_4 < 0, X_5 < 0, V\} - 3. \quad (\text{C.9})$$

By the conditioning property of the multivariate normal distribution and the identities  $V_i^2 + (V_i^-)^2 = 1$ , for  $i = 1, 2$ , and  $V_1^- V_2^- = V_3$ , we obtain  $(X_1, X_2) | \{X_3, X_4, X_5, V\} \sim N(\mu, \Sigma)$ , where

$$\mu = \begin{bmatrix} (V_1 X_3 + V_1^- X_5) \delta_1 \\ (V_2 X_4 + V_2^- X_5) \delta_2 \end{bmatrix} \quad \text{and} \quad \Sigma = \begin{bmatrix} 1 - \delta_1^2 & V_3(\rho - \delta_1 \delta_2) \\ V_3(\rho - \delta_1 \delta_2) & 1 - \delta_2^2 \end{bmatrix}.$$

Therefore, we have

$$\mathbb{P}\{X_1 < 0, X_2 < 0 | X_3, X_4, X_5, V\} = \Phi_2(-\alpha_1^\dagger(V_1 X_3 + V_1^- X_5), -\alpha_2^\dagger(V_2 X_4 + V_2^- X_5); \rho^\dagger V_3). \quad (\text{C.10})$$

Since  $(X_3, X_4, X_5) | V$  are mutually independent standard normal random variates, equation (5.8) in the corollary follows from the definition of  $(Y_1, Y_2, Y_3)$ , along with (C.9) and (C.10).  $\square$

**Proof of Proposition 4.** To prove the proposition, we begin with the expressions for Kendall's tau and Spearman's rho given in (5.7) and (5.8), respectively, in Corollary 2. Part (i) of the proposition follows directly from the exchangeability of bivariate normal cdf and the fact that  $\alpha_1^\dagger(\rho, (\alpha_1, \alpha_2)) = \alpha_2^\dagger(\rho, (\alpha_2, \alpha_1))$  and  $\alpha_2^\dagger(\rho, (\alpha_1, \alpha_2)) = \alpha_1^\dagger(\rho, (\alpha_2, \alpha_1))$ , which can be inferred from (5.5).

To prove Part (ii), we first note that  $\tau(\rho, \alpha, F) = \tau(\rho, -\alpha, F)$  follows directly from the expression of  $\tau$  in (5.7) and the fact that  $Z \stackrel{d}{=} -Z$ . Moreover, from (5.8), it is clear that to show  $\rho_S(\rho, \alpha, F) = \rho_S(\rho, -\alpha, F)$ , it suffices to prove

$$\mathbb{E} \Phi_2(\alpha_1^\dagger Z_1, \alpha_2^\dagger Z_2; \rho^\dagger V_3) = \mathbb{E} \Phi_2(-\alpha_1^\dagger Z_1, -\alpha_2^\dagger Z_2; \rho^\dagger V_3),$$



where the mixing variables  $(Z_1, Z_2, V_3)$  are defined in Corollary 2 (ii). This equality can be established using the same proof strategy employed in the proof of Proposition 1 (ii), along with the fact that  $Z_i \stackrel{d}{=} -Z_i$ , for  $i = 1, 2$ .

To prove Part (iii), we first notice that, for any  $\alpha \in \mathbb{R}^2$ , we have  $\delta_1(-1, \alpha) = -\delta_2(-1, \alpha)$  and  $\delta_1(1, \alpha) = \delta_2(1, \alpha)$ . Therefore,  $\alpha_1^\dagger = -\alpha_2^\dagger := \alpha^\circ$  and  $\rho^\dagger = -1$ , when  $\rho = -1$ ; and  $\alpha_1^\dagger = \alpha_2^\dagger := \alpha_\circ$  and  $\rho^\dagger = 1$ , when  $\rho = 1$ . To show the result for Kendall's tau, we substitute  $\rho = -1$  into equation (5.7) to obtain

$$\tau(-1, \alpha, F) = 4 \mathbb{E} \Phi_2(\alpha^\circ Z, -\alpha^\circ Z; -1) - 1.$$

Then, using the identity  $\Phi_2(x, -x; -1) = \Phi(x) - \Phi(x) = 0$  for all  $x \in \mathbb{R}$ , we deduce that  $\tau(-1, \alpha, F) = 4 \times 0 - 1 = -1$ . On the other hand, substituting  $\rho = 1$  into equation (5.7) and then applying Lemma 2 for  $\rho = 1$  case yields

$$\tau(1, \alpha, F) = 4 \mathbb{E} \Phi_2(\alpha_\circ Z, \alpha_\circ Z; 1) - 1 = 4 \mathbb{E} \Phi^s(\alpha_\circ Z; 0) - 1 = 4 \mathbb{E} \Phi(\alpha_\circ Z) - 1,$$

where  $Z = V_1 Y_1 + V_2 Y_2$ , as defined in Corollary 2 (i). By  $\alpha_\circ Z \stackrel{d}{=} -\alpha_\circ Z$  and the identity  $\Phi(\alpha_\circ Z) + \Phi(-\alpha_\circ Z) = 1$ , we obtain  $\mathbb{E} \Phi(\alpha_\circ Z) = 1/2$ , and hence  $\tau(1, \alpha, F) = 4 \times 1/2 - 1 = 1$ .

To prove the corresponding result for Spearman's rho, we substitute  $\rho = \pm 1$  into equation (5.8) to get

$$\begin{aligned} \rho_S(1, \alpha, F) &= 12 \mathbb{E} \Phi_2(\alpha_\circ Z_1, \alpha_\circ Z_2; V_3) - 3, \\ \rho_S(-1, \alpha, F) &= 12 \mathbb{E} \Phi_2(\alpha^\circ Z_1, -\alpha^\circ Z_2; -V_3) - 3, \end{aligned} \tag{C.11}$$

where  $Z_1 = V_1 Y_1 + V_1^- Y_3$  and  $Z_2 = V_2 Y_2 + V_2^- Y_3$  are defined in Corollary 2 (ii). Using the identity  $\Phi_2(x, y; \rho) + \Phi_2(x, -y; -\rho) = \Phi(x)$ , we can rewrite the above expression for  $\rho_S(-1, \alpha, F)$  as

$$\rho_S(-1, \alpha, F) = 12 [\mathbb{E} \Phi(\alpha^\circ Z_1) - \mathbb{E} \Phi_2(\alpha^\circ Z_1, \alpha^\circ Z_2; V_3)] - 3. \tag{C.12}$$

By  $\alpha^\circ Z_1 \stackrel{d}{=} -\alpha^\circ Z_1$  and the identity  $\Phi(\alpha^\circ Z) + \Phi(-\alpha^\circ Z) = 1$ , we have

$$\mathbb{E} \Phi(\alpha^\circ Z_1) = \frac{1}{2}. \tag{C.13}$$

Furthermore, following the same proof strategy used to show  $\rho_S(1, b, F) = 1$  in Proposition 2

under the equi-skewness condition, we can show that

$$\mathbb{E} \Phi_2(\alpha_\circ Z_1, \alpha_\circ Z_2; V_3) = \frac{1}{3}. \quad (\text{C.14})$$

Substituting (C.13) and (C.14) into equations (C.11) and (C.12), we obtain  $\rho_S(1, \alpha, F) = 1$  and  $\rho_S(-1, \alpha, F) = -1$  for all  $\alpha \in \mathbb{R}^2$  and  $F$ . This complete the proof for Part (iii) of the proposition, and also the proof of the whole proposition.  $\square$

**Proof of Proposition 5.** The expressions of  $\tau$  of  $\rho_S$  stated in Part (i) and Part (ii) of the proposition follow directly from Corollary 2 and the equi-skew setting. To analyze  $\partial\tau/\partial\rho$ , we deduce from (4.8) and (4.9) that

$$\frac{\partial}{\partial x} \Phi_2(x, x; \rho) = 2\phi(x)\Phi(\alpha_\rho x) = \phi^s(x; \alpha_{\rho^\dagger}) \quad \text{and} \quad \frac{\partial}{\partial \rho} \Phi_2(x, x; \rho) = \phi_2(x, x; \rho),$$

where  $\alpha_{\rho^\dagger} := \sqrt{(1 - \rho^\dagger)/(1 + \rho^\dagger)} > 0$ . Then, we have

$$\frac{\partial}{\partial \rho} \mathbb{E} \Phi_2(a^\dagger Z, a^\dagger Z; \rho^\dagger) = \frac{\partial a^\dagger}{\partial \rho} \mathbb{E} \phi^s(a^\dagger Z; \alpha_{\rho^\dagger}) Z + \frac{\partial \rho^\dagger}{\partial \rho} \mathbb{E} \phi_2(a^\dagger Z, a^\dagger Z; \rho^\dagger).$$

In the above expression, we note that (i)  $\partial a^\dagger/\partial \rho > 0$  when  $a > 0$ , (ii)  $\mathbb{E} \phi^s(a^\dagger Z; \alpha_{\rho^\dagger}) Z > 0$  when  $a > 0$ , by Lemma 3 (i) and the fact that  $Z \stackrel{d}{=} -Z$ , (iii)  $\partial \rho^\dagger/\partial \rho > 0$ , and (iv)  $\phi_2$  is a strictly positive function. Consequently, we can deduce that  $\partial \mathbb{E} \Phi_2(a^\dagger Z, a^\dagger Z; \rho^\dagger)/\partial \rho > 0$ , and hence  $\partial\tau(\rho, a, F)/\partial \rho > 0$ .

Next, we prove  $\partial\tau/\partial a < 0$ . Since  $\partial\bar{\delta}/\partial a > 0$  and  $\bar{\delta}$  shares the same sign as  $a$ , to show  $\partial\tau/\partial a < 0$  it suffices to show

$$\frac{\partial}{\partial \bar{\delta}} \mathbb{E} \Phi_4(0; P_\tau(\rho, (\bar{\delta}, \bar{\delta}), V)) < 0 \quad \text{for} \quad \bar{\delta} > 0. \quad (\text{C.15})$$

By Lemma 7, we have

$$\frac{\partial}{\partial \bar{\delta}} \Phi_4(0; P_\tau(\rho, (\bar{\delta}, \bar{\delta}), V)) \stackrel{d}{=} \frac{V_1 \arcsin \tilde{V}(\bar{\delta}, \bar{\delta})}{\pi^2 \sqrt{1 - V_1^2 \bar{\delta}^2}}.$$

Since  $V_1 > 0$  and  $\tilde{V}(\bar{\delta}, \bar{\delta}) = V_2(1 - \rho)\bar{\delta}[(1 - \bar{\delta}^2)(1 - \rho^2 - 2V_1^2\bar{\delta}^2(1 - \rho))]^{-1/2} < 0$  when  $\bar{\delta} > 0$ , we can easily deduce (C.15) upon verifying the conditions of interchanging expectation and differentiation. This completes the proof of the proposition.  $\square$

**Proof of Proposition 6.** We prove Part (i) of the proposition first. Note that, in the equi-skew case with  $\rho = 0$ , it follows from (5.12) and (5.13) that  $\rho^\dagger = 0$ ,  $a_1^\dagger = a$  and  $a_2^\dagger = 0$ . Then, applying Corollary 2, we have

$$\begin{aligned}\tau(0, (a, 0), F) &= 4 \mathbb{E} \Phi_2(aZ, 0; 0) - 1 = 2 \mathbb{E} \Phi(aZ) - 1, \\ \rho_S(0, (a, 0), F) &= 12 \mathbb{E} \Phi_2(aZ_1, 0; 0) - 3 = 6 \mathbb{E} \Phi(aZ_1) - 3.\end{aligned}$$

Since  $Z \stackrel{d}{=} -Z$  and  $Z_1 \stackrel{d}{=} -Z_1$ , we can deduce that  $\mathbb{E} \Phi(aZ) = \mathbb{E} \Phi(aZ_1) = 1/2$ , using the identity that  $\Phi(x) + \Phi(-x) = 1$  for all  $x \in \mathbb{R}$ . Consequently,  $\tau(0, (a, 0), F) = \rho_S(0, (a, 0), F) = 0$ .

Next, we show that  $\tau(\rho, (0, a), F)$  is an odd functions of  $\rho$ , or  $\tau(\rho, (0, a), F) + \tau(-\rho, (0, a), F) = 0$ , for any  $a \in \mathbb{R}$  and  $F$ . Using Corollary 2 (i), it suffices to show that

$$\mathbb{E} [\Phi_2(aZ, a\rho^\dagger Z; \rho^\dagger) + \Phi_2(aZ, -a\rho^\dagger Z; -\rho^\dagger)] = \frac{1}{2}, \quad (\text{C.16})$$

where  $\rho^\dagger$  is defined by (5.12). Applying the representation of  $\Phi_2(x_1, x_2; \rho)$  in (4.6) and then using the identities  $\phi_2(x_1, x_2; -\rho) = \phi_2(x_1, -x_2; \rho)$  and  $\phi(x) + \phi(-x) = 1$ , we can deduce that the left-hand side of (C.16) is equal to  $\mathbb{E} \Phi(aZ)$ . Since  $\mathbb{E} \Phi(aZ) = 1/2$ , as shown above, equation (C.16) holds. Analogously, we can show that

$$\mathbb{E} [\Phi_2(aZ_1, a\rho^\dagger Z_2; \rho^\dagger V_3) + \Phi_2(aZ_1, -a\rho^\dagger Z_2; -\rho^\dagger, V_3)] = \frac{1}{2},$$

where  $Z_1, Z_2$  and  $V_3$  are defined in Corollary 2 (ii), and, consequently,  $\rho_S(\rho, (0, a), F)$  is an odd functions of  $\rho$  as well. This completes the proof of Part (i) of the proposition.

Provided that Kendall's tau is an odd function of  $\rho$ , to prove Part (ii) of the proposition, we may focus on the case of  $\rho > 0$ , and show that  $\partial\tau/\partial\rho > 0$  and  $\partial\tau/\partial a < 0$ , when  $\rho > 0$  and  $a > 0$ . To analyze  $\partial\tau/\partial\rho$ , we first deduce from (4.8) and (4.9) that

$$\begin{aligned}\frac{\partial}{\partial x_2} \Phi_2(x_1, x_2; \rho) &= \phi(x_2) \Phi\left(\frac{x_1 - \rho x_2}{\sqrt{1 - \rho^2}}\right) = \frac{1}{2} \phi^s\left(x_2; \frac{x_1/x_2 - \rho}{\sqrt{1 - \rho^2}}\right), \\ \frac{\partial}{\partial \rho} \Phi_2(x_1, x_2; \rho) &= \phi_2(x_1, x_2; \rho).\end{aligned}$$

Then, we have

$$\frac{\partial}{\partial \rho} \mathbb{E} \Phi_2(aZ, a\rho^\dagger Z; \rho^\dagger) = \frac{1}{2} \frac{\partial \rho^\dagger}{\partial \rho} \mathbb{E} [a \phi^s(a\rho^\dagger Z; \alpha_{\rho^\dagger}) Z + 2\phi_2(aZ, a\rho^\dagger Z; \rho^\dagger)]$$

where  $\alpha_{\rho^\dagger} := (1 - \rho^{\dagger 2})/\rho^\dagger \sqrt{1 - \rho^{\dagger 2}}$ . Since  $\alpha_{\rho^\dagger} > 0$  and  $a\rho^\dagger > 0$ , when both  $\rho$  and  $a$  are strictly positive, it follows from Lemma 3 (i) that  $\mathbb{E} \phi^s(a\rho^\dagger Z; \alpha_{\rho^\dagger}) Z > 0$ . Along with  $\partial\rho^\dagger/\partial\rho > 0$  and  $\phi_2(\cdot; \rho^\dagger) > 0$ , we can deduce that  $\partial\mathbb{E} \Phi_2(aZ, a\rho^\dagger Z; \rho^\dagger)/\partial\rho > 0$  and hence  $\partial\tau/\partial\rho > 0$ , as desired to be shown.

To analyze  $\partial\tau/\partial a$ , we first note that  $\partial\delta_\circ/\partial a > 0$  and  $\text{sgn}(\delta_\circ) = \text{sgn}(a)$ . Therefore, to show  $\partial\tau/\partial a < 0$  when  $\rho > 0$  and  $a > 0$ , it suffices to prove that

$$\frac{\partial}{\partial\delta_\circ} \mathbb{E} \Phi_4(0; P_\tau(\rho, (\delta_\circ, \delta_\circ\rho), V)) < 0 \quad \text{when } \rho > 0 \text{ and } \delta_\circ > 0, \quad (\text{C.17})$$

using the expression of Kendall's tau given by Theorem 2 (i). By Lemma 7, we have

$$\frac{\partial}{\partial\delta_\circ} \Phi_4(0; P_\tau(\rho, (\delta_\circ, \delta_\circ\rho), V)) =_d \frac{\rho V_1 \arcsin \tilde{V}(\delta_\circ\rho, \delta_\circ)}{2\pi^2 \sqrt{1 - (V_1\delta_\circ\rho)^2}}.$$

Since  $V_1 > 0$  and  $\tilde{V}(\delta_\circ\rho, \delta_\circ) = V_2\delta_\circ(1 - \rho^2)\{(1 - \rho^2\delta_\circ^2)[(1 - \rho^2)(1 - V_1^2\delta_\circ^2)]\}^{-1/2} < 0$  when  $\delta_\circ > 0$ , we can deduce (C.17) after verifying the conditions of interchanging expectation and differentiation. This completes Part (ii) of the proposition.  $\square$