A BIALGEBRA THEORY OF ANTI-LEIBNIZ ALGEBRAS

BO HOU AND ZHANPENG CUI

Abstract. We introduce the notion of an anti-Leibniz bialgebra which is equivalent to a Manin triple of anti-Leibniz algebras, is equivalent to a matched pair of anti-Leibniz algebras. The study of some special anti-Leibniz bialgebras leads to the introduction of the anti-Leibniz Yang-Baxter equation in an anti-Leibniz algebra. A symmetric (or an invariant) solution of the anti-Leibniz Yang-Baxter equation gives an anti-Leibniz bialgebra. The notion of a relative Rota-Baxter operator of an anti-Leibniz algebra is introduced to construct symmetric solutions of the anti-Leibniz Yang-Baxter equation. Moreover, we introduce the notions of factorizable anti-Leibniz bialgebras and skew-symmetric Rota-Baxter anti-Leibniz algebras, and show that a factorizable anti-Leibniz bialgebra leads to a factorization of the underlying anti-Leibniz algebra. There is a one-to-one correspondence between factorizable anti-Leibniz bialgebras and skew-quadratic Rota-Baxter anti-Leibniz algebras. Finally, we constrict anti-Leibniz bialgebras form Leibniz bialgebras by the tensor product and constrict infinite-dimensional anti-Leibniz bialgebras form finite-dimensional anti-Leibniz bialgebras by the completed tensor product.

CONTENTS

1. Introduction

A bialgebra structure on a given algebraic structure is obtained as a coalgebra structure together which gives the same algebraic structure on the dual space with a set of compatibility conditions between the multiplications and comultiplications. One of the most famous examples of bialgebras is the Lie bialgebra [\[14\]](#page-28-1) and more importantly there have been a lot of bialgebra theories for other algebra structures that essentially follow the approach of Lie bialgebras such as antisymmetric infinitesimal bialgebras [\[1,](#page-28-2) [4,](#page-28-3) [5\]](#page-28-4), left-symmetric bialgebras [\[3\]](#page-28-5), Jordan bialgebras [\[29\]](#page-29-0), Poisson bialgebras [\[24\]](#page-28-6), transposed Poisson bialgebras [\[22\]](#page-28-7), Leibniz bialgebra [\[28\]](#page-29-1) and mock-Lie bialgebras [\[11\]](#page-28-8). In this paper, we give a bialgebra theory for anti-Leibniz algebras.

Leibniz algebra is a non-commutative analogue of Lie algebra, which was first considered in [\[23\]](#page-28-9) by the study of the periodicity in algebraic K-theory. A mock-Lie algebra (also called Jacobi-Jordan algebra, Lie-Jordan algebra) is a commutative algebra satisfying the Jacobi identity, which is a special Jordan algebra. In recent years, mock-Lie algebra has attracted the attention of many scholars $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$ $[7, 8, 10, 11, 12, 16, 18, 19, 30]$. Recently, the notion of anti-Leibniz algebras as a noncommutative

²⁰¹⁰ *Mathematics Subject Classification.* 17A30, 16T10, 16T25, 17B40.

Key words and phrases. anti-Leibniz algebra, anti-Leibniz bialgebra, Yang-Baxter equation, relative Rota-Baxter operator, factorizable anti-Leibniz bialgebras, infinite-dimensional anti-Leibniz bialgebras.

version of mock-Lie algebras was introduced in [\[9\]](#page-28-17). In this paper, we still take a similar approach as the study on Lie bialgebras. The notion of anti-Leibniz bialgebra is thus introduced as an equivalent structure of a Manin triple of anti-Leibniz algebras, which is interpreted in terms of matched pairs of anti-Leibniz algebras. One reason for the usefulness of the Lie bialgebra is that it has a coboundary theory, which leads to the construction of Lie bialgebras from solutions of the classical Yang-Baxter equations. An important class of quasi-triangular Lie bialgebras, factorizable Lie bialgebras are used to connect classical *r*-matrices with certain factorization problems, and have various applications in integrable systems $[25, 27]$ $[25, 27]$. Recently, the factorizable Lie bialgebras, factorizable antisymmetric infinitesimal bialgebras and factorizable Leibniz bialgebras have been studied in [\[17\]](#page-28-18), [\[26\]](#page-29-5) and [\[6\]](#page-28-19). Here we also introduce the notion of quasi-triangular anti-Leibniz bialgebra and the anti-Leibniz Yang-Baxter equation in an anti-Leibniz algebra. We give the relative Rota-Baxter operators of anti-Leibniz algebras and construct a solution of the Yang-Baxter equation in an anti-Leibniz algebra. Moreover, the notion of factorizable anti-Leibniz bialgebra is introduced, and a one-to-one correspondence between factorizable anti-Leibniz bialgebras and skew-quadratic Rota-Baxter anti-Leibniz algebras of nonzero weight is given. We know very little about specific examples of anti-Leibniz algebras and anti-Leibniz bilgebras. Note that the tensor product of a Leibniz algebra and an anti-commutative anti-associative algebra is an anti-Leibniz algebra, we take this result to the level of the bialgebras, and provide some specific examples of anti-Leibniz bialgebras. The bialgebra theory of anti-Leibniz algebras is based on finitedimensional anti-Leibniz algebras. For the infinite-dimensional anti-Leibniz bialgebras, we provide a construction method using the completed tensor products.

The paper is organized as follows. In Section [2,](#page-1-0) we recall some basic definitions of anti-Leibniz algebras and bimodules over anti-Leibniz algebras. We introduce the notions of matched pairs of anti-Leibniz algebras, Manin triples of anti-Leibniz algebras and anti-Leibniz bialgebras. The equivalence between them is interpreted. In Section [3,](#page-6-0) we consider some special classes of anti-Leibniz bialgebras, such as triangular anti-Leibniz bialgebras and factorizable anti-Leibniz bialgebras, and introduce the notion of the anti-Leibniz Yang-Baxter equation in an anti-Leibniz algebra. The notion of relative Rota-Baxter operators of anti-Leibniz algebras is introduced to provide symmetric solutions of the anti-Leibniz Yang-Baxter equation in semi-direct product anti-Leibniz algebras and hence gives rise to anti-Leibniz bialgebras. Moreover, we establish the factorizable theories for anti-Leibniz bialgebras, and give a one-to-one correspondence between factorizable anti-Leibniz bialgebras and skew-quadratic Rota-Baxter anti-Leibniz algebras. At end of this section, we give some examples of anti-Leibniz bialgebras by Leibniz bialgebras. In Section [4,](#page-23-0) we introduce the notions of completed tensor product of Z-graded vector spaces, completed anti-Leibniz coalgebras, and propose a method for constructing infinite-dimensional anti-Leibniz bialgebras using the completed tensor product of finite-dimensional anti-Leibniz bialgebras and quadratic Z-graded commutative associative algebra.

Throughout this paper, we fix \Bbbk a field and characteristic zero. All the vector spaces and algebras are of finite dimension over \Bbbk , and all tensor products are also taking over \Bbbk .

2. Anti-Leibniz algebras and anti-Leibniz bialgebras

In this section, we recall the background on anti-Leibniz algebras and bimodules over anti-Leibniz algebras. We introduce the notions of matched pairs of anti-Leibniz algebras, Manin triples of anti-Leibniz algebras and anti-Leibniz bialgebras. The equivalence between them is interpreted.

Definition 2.1 ([\[9\]](#page-28-17)). An anti-Leibniz algebra *over* \bf{k} *is a* \bf{k} -vector space A with a multiplication \cdot : $A \otimes A \rightarrow A$, such that for any $a_1, a_2, a_3 \in A$,

$$
a_1(a_2a_3) + (a_1a_2)a_3 + a_2(a_1a_3) = 0,
$$

where $a_1a_2 := a_1 \cdot a_2$ *for simply.*

By the definition, it is easy to see that $(a_1a_2)a_3 = (a_2a_1)a_3$ for any elements a_1, a_2, a_3 in an anti-Leibniz algebra (*A*, ·). Anti-Leibniz algebra is a noncommutative version of the mock-Lie algebras. Recall that

a mock-Lie algebra is a vector space *V* together with a bilinear map $[-, -] : V \times V \rightarrow V$, such that for any $v_1, v_2, v_3 \in V$,

$$
[v_1, v_2] = [v_2, v_1], \qquad [v_1, [v_2, v_3]] + [v_2, [v_3, v_1]] + [v_3, [v_1, v_2]] = 0.
$$

Since $[v_1, v_2] = [v_2, v_1]$, the second equation can be rewrite as $[v_1, [v_2, v_3]] + [[v_1, v_2], v_3] + [v_2, [v_1, v_3]] = 0$ or $[v_1, [v_2, v_3]] + [[v_1, v_2], v_3] + [[v_1, v_3], v_2] = 0$. The anti-Leibniz algebra given in Definition [2.1](#page-1-1) is also called a *left anti-Leibniz algebra*. A *right anti-Leibniz algebra* (A, \cdot) is given by $a_1(a_2a_3) + (a_1a_2)a_3 + (a_2a_4)^2$ $(a_1a_3)a_2 = 0$, for any $a_1, a_2, a_3 \in A$. It is easy to see that (A, \cdot) is a left anti-Leibniz algebra if and only if its opposite algebra is right anti-Leibniz. For classification problems that anti-Leibniz algebra, we still know very little. Here we give some simple examples.

Example 2.2. Let vector space $A = \mathbb{k}\{e_1, e_2\}$. Then the non trivial anti-Leibniz algebraic structure on A *is only isomorphic to the algebra given by*

$$
\Lambda_1^2: e_1e_1=e_2;
$$

 Λ^2 2 : $e_1e_1 = -ae_1 - \frac{a^2}{b}$ $\frac{a^2}{b}e_2$, $e_1e_2 = e_2e_1 = be_1 + ae_2$, $e_2e_2 = -\frac{b^2}{a}$ $\frac{b^2}{a}e_1 - be_2$, with 0 ≠ *a*, *b* ∈ k.

That is to say, all 2*-dimensional anti-Leibniz algebras are commutative.*

Let vector space $A = \mathbb{k}\{e_1, e_2, e_3\}$. Define $e_1e_2 = e_3 = e_2e_2$. Then A is a 3-dimensional (noncommuta*tive) anti-Leibniz algebra under this product.*

Let (A, \cdot) and (B, \cdot') be two anti-Leibniz algebras. We call (B, \cdot') is a subalgebra of (A, \cdot) if *B* is a subspace of *A* and \cdot is the restriction of \cdot on *B*. A linear map $f : A \rightarrow B$ is called a *homomorphism of anti-Leibniz algebras* if for any $a_1, a_2 \in A$, $f(a_1 \cdot a_2) = f(a_1) \cdot f(a_2)$. A homomorphism f is said to be an isomorphism if *f* is a bijection. Now we introduce the definition of module over an anti-Leibniz algebra.

Definition 2.3. *Let* (*A*, ·) *be an anti-Leibniz algebra, M be a* k*-vector space. If there exists a pair of bilinear maps* $I, r : A \rightarrow gl(M)$ *, such that for any* $a_1, a_2 \in A$ *and* $m \in M$ *,*

$$
I(a_1a_2)(m) + I(a_1)(I(a_2)(m)) + I(a_2)(I(a_1)(m)) = 0,
$$

\n
$$
I(a_1)(r(a_2)(m)) + r(a_2)(I(a_1)(m)) + r(a_1a_2)(m) = 0,
$$

\n
$$
r(a_1a_2)(m) + r(a_2)(r(a_1)(m)) + I(a_1)(r(a_2)(m)) = 0,
$$

we call (M, I, r) *(or simply M) is a bimodule over* (A, \cdot) *.*

Let (A, \cdot) be an anti-Leibniz algebra, (M, I, r) be a bimodule over (A, \cdot) . Then we get

$$
r(a_1)(r(a_2)(m)) = r(a_1)(I(a_2)(m)),
$$

for any $a_1, a_2 \in A$ and $m \in M$. Now, let (M, I, r) and (N, I', r') be two bimodules over (A, \cdot) and f : $M \rightarrow N$ be a linear map. Then *f* is called a *morphism of bimodule* if $f(1(a)(m)) = 1'(a)(f(m))$ and $f(r(a)(m)) = r'(a)(f(m))$ for any $a \in A$ and $m \in M$. (M, I, r) and (N, I', r') are called *isomorphic as bimodule* if the morphism *f* is an isomorphism. Moreover, if we denote I_A , $r_A : A \rightarrow \mathfrak{gl}(A)$ by $l_A(a_1)(a_2) = a_1a_2$ and $\mathbf{r}_A(a_1)(a_2) = a_2a_1$ for any $a_1, a_2 \in A$, then it is easy to see that (A, I_A, \mathbf{r}_A) is a bimodule over (A, \cdot) , which is called the *regular bimodule* over (A, \cdot) . By direct calculations, we can give an equivalent condition as follows.

Proposition 2.4. *Let* (A, \cdot) *be an anti-Leibniz algebra, M be a* k-vector space, $I, r : A \rightarrow \mathfrak{gl}(M)$ *be two bilinear maps. Then* (M, I, r) *is a bimodule over* (A, \cdot) *if and only if* $A \oplus M$ *is an anti-Leibniz algebra under the following operation:*

$$
(a_1, m_1)(a_2, m_2) = (a_1a_2, \, I(a_1)(m_2) + \mathfrak{r}(a_2)m_1),
$$

for all $a_1, a_2 \in A$ *and* $m_1, m_2 \in M$. This anti-Leibniz algebra is called the semi-direct product of (A, \cdot) *by bimodule* (M, I, r) *, denoted by* $A \times M$ *.*

Let *V* be a vector space. Denote the standard pairing between the dual space V^* and *V* by

$$
\langle -, - \rangle: \quad V^* \otimes V \to \mathbb{k}, \quad (\xi, v) := \xi(v),
$$

for any $\xi \in V^*$ and $v \in V$. Let V, W be two vector spaces. For a linear map $\varphi : V \to W$, the transpose map $\varphi^* : W^* \to V^*$ is defined by

$$
\langle \varphi^*(\xi), v \rangle := \langle \xi, \varphi(v) \rangle,
$$

for any $v \in V$ and $\xi \in W^*$. Let (A, \cdot) be an anti-Leibniz algebra and V be a vector space. For a linear map $\psi : A \to \mathfrak{gl}(V)$, the linear map $\psi^* : A \to \mathfrak{gl}(V^*)$ is defined by

$$
\langle \psi^*(a)(\xi), v \rangle := \langle \xi, \psi(a)(v) \rangle,
$$

for any $a \in A$, $v \in V$, $\xi \in V^*$. That is, $\psi^*(a) = \psi(a)^*$ for all $a \in A$. Then, for any bimodule (M, I, r) over anti-Leibniz algebra (A, \cdot) , it is easy to see that, $(M^*, I^*, I^* - r^*)$ is again a bimodule over (A, \cdot) . In particular, we get (A^*, \mathbf{l}_A^*) A^*_{A} , **l**_A^{*} – **r**_A^{*} A_A^*) is a bimodule over (A, \cdot) , which is called the *coregular bimodule*.

We give the definition of matched pair of anti-Leibniz algebras.

Definition 2.5. Let (A, \cdot) and (B, \cdot') be two anti-Leibniz algebras, (B, I_A, r_A) and (A, I_B, r_B) be bimodules *over* (A, \cdot) *and* (B, \cdot') *respectively. If for any* $a, a_1, a_2 \in A$ *and* $b, b_1, b_2 \in B$,

(2.1) r*A*(*a*)(*b*1*b*2) + *b*1r*A*(*a*)(*b*2) + *b*2r*A*(*a*)(*b*1) + r*A*(l*B*(*b*2)(*a*))(*b*1) + r*A*(l*B*(*b*1)(*a*))(*b*2) = 0,

$$
(2.2) \qquad I_A(a)(b_1b_2) + I_A(a)(b_1)b_2 + b_1I_A(a)(b_2) + I_A(\mathfrak{r}_B(b_1)(a))(b_2) + \mathfrak{r}_A(\mathfrak{r}_B(b_2)(a))(b_1) = 0,
$$

(2.3)
$$
I_A(a)(b_1)b_2 + I_A(\mathfrak{r}_B(b_1)(a))(b_2) - \mathfrak{r}_A(a)(b_1)b_2 - I_A(I_B(b_1)(a))(b_2) = 0,
$$

(2.4) r*B*(*b*)(*a*1*a*2) + *a*1r*B*(*b*)(*a*2) + *a*2r*B*(*b*)(*a*1) + r*B*(l*A*(*a*2)(*b*))(*a*1) + r*B*(l*A*(*a*1)(*b*))(*a*2) = 0,

$$
(2.5) \tI_B(b)(a_1a_2) + I_B(b)(a_1)a_2 + a_1I_B(b)(a_2) + I_B(\mathfrak{r}_A(a_1)(b))(a_2) + \mathfrak{r}_B(\mathfrak{r}_A(a_2)(b))(a_1) = 0,
$$

$$
(2.6) \tI_B(b)(a_1)a_2 + I_B(\mathfrak{r}_A(a_1)(b))(a_2) - \mathfrak{r}_B(b)(a_1)a_2 - I_B(I_A(a_1)(b))(a_2) = 0,
$$

where $a_1a_2 := a_1 \cdot a_2$ *and* $b_1b_2 := b_1 \cdot b_2$, we called $(A, B, I_A, r_A, I_B, r_B)$ *is a* matched pair of anti-Leibniz algebras of (A, \cdot) and (B, \cdot) .

For the matched pair of anti-Leibniz algebras, we have the following equivalent description.

Proposition 2.6. Let (A, \cdot) and (B, \cdot') be two anti-Leibniz algebras, I_A , $r_A : A \to \mathfrak{gl}(B)$ and I_B , $r_B : B \to B$ gl(A) *be four linear maps. Then* $(A, B, I_A, r_A, I_B, r_B)$ *is a matched pair of anti-Leibniz algebras of* (A, \cdot) *and* (*B*, · ′) *if and only if A* ⊕ *B under the product*

$$
(2.7) \qquad (a_1, b_1) * (a_2, b_2) = (a_1a_2 + I_B(b_1)(a_2) + r_B(b_2)(a_1), \ b_1b_2 + I_A(a_1)(b_2) + r_A(a_2)(b_1)),
$$

for all $a_1, a_2 \in A$ and $b_1, b_2 \in B$, is also an anti-Leibniz algebra. This anti-Leibniz algebra is called a crossed product *of* (A, \cdot) *and* (B, \cdot') *, denoted by* $A \Join B$ *.*

Proof. It is straightforward. □

In particular, consider the matched pair of anti-Leibniz algebra structures on *A* and *A* ∗ , we have the following corollary.

Corollary 2.7. Let (A, \cdot) and (A^*, \cdot') be two anti-Leibniz algebras. Then (A, A^*, I^*_A) A^* , **l**_A^{*} – **r**_A^{*} $A^*_{A^*}, I^*_{A^*} - \mathbf{r}^*_{A^*}$) *is a* matched pair of anti-Leibniz algebras of (A, \cdot) and (A^*, Δ^*) if and only if for any $a_1, a_2 \in A$ and $\xi \in A^*$,

(2.8)
$$
(\mathbf{l}_{A^*}^* - \mathbf{r}_{A^*}^*)(\xi)(a_1a_2) + a_1(\mathbf{l}_{A^*}^* - \mathbf{r}_{A^*}^*)(\xi)(a_2) + a_2(\mathbf{l}_{A^*}^* - \mathbf{r}_{A^*}^*)(\xi)(a_1) + (\mathbf{l}_{A^*}^* - \mathbf{r}_{A^*}^*)(\mathbf{l}_A^*(a_2)(\xi))(a_1) + (\mathbf{l}_{A^*}^* - \mathbf{r}_{A^*}^*)(\mathbf{l}_A^*(a_1)(\xi))(a_2) = 0,
$$

(2.9) $I^*_{A^*}(\xi)(a_1a_2) + I^*_{A^*}(\xi)(a_1)a_2 + a_1I^*_{A^*}(\xi)(a_2)$

$$
+ I_{A^*}^*((I_A^* - \mathbf{r}_A^*)(a_1)(\xi))(a_2) + (I_{A^*}^* - \mathbf{r}_A^*)(I_A^* - \mathbf{r}_A^*)(a_2)(\xi))(a_1) = 0,
$$

(2.10)
$$
\mathbf{r}_A^*(\xi)(a_1)a_2 - I_{A^*}^*(\mathbf{r}_A^*(a_1)(\xi))(a_2) = 0.
$$

$$
f_{\rm{max}}
$$

Proof. Taking $I_A = I_A^*$ A^* , $r_A = I^*_A - r^*_A$ A_A^* , $I_B = I_{A^*}^*$ and $r_B = I_{A^*}^* - r_{A^*}^*$ in Eqs. [\(2.4\)](#page-3-0)-[\(2.6\)](#page-3-1), we get Eqs. [\(2.8\)](#page-3-2)-[\(2.10\)](#page-3-3) respectively. Note that for any $a_1, a_2 \in A$ and $\xi_1, \xi_2 \in A^*$,

$$
\langle (\mathbf{r}_{A^*}^*(\xi_1)(a_1)a_2, \xi_2 \rangle - \langle \mathbf{l}_{A^*}^*(\mathbf{r}_A^*(a_1)(\xi_1))(a_2), \xi_2 \rangle = \langle \mathbf{r}_{A^*}^*(\xi_1)(a_1)a_2, \xi_2 \rangle - \langle a_2, \mathbf{r}_A^*(a_1)(\xi_1)\xi_2 \rangle
$$

= $\langle a_2, \mathbf{l}_{A^*}^*(\mathbf{r}_{A^*}^*(\xi_1)(a_1))(\xi_2) \rangle - \langle a_2, \mathbf{r}_A^*(a_1)(\xi_1)\xi_2 \rangle$,

we get Eq. $(2.8) \Leftrightarrow$ Eq. (2.3) . Similarly, if Eq. (2.3) holds, we can deduce Eq. $(2.9) \Leftrightarrow$ $(2.9) \Leftrightarrow$ Eq. (2.2) and Eq. $(2.8) \Leftrightarrow$ $(2.8) \Leftrightarrow$ Eq. (2.1) . Thus we obtain this corollary.

Let B(−, −) be a bilinear form on an anti-Leibniz algebra (*A*, ·).

- $\mathfrak{B}(-, -)$ is called *nondegenerate* if $\mathfrak{B}(a_1, a_2) = 0$ for any $a_2 \in A$, then $a_1 = 0$;
- B(−, −) is called *skew-symmetric* if B(*a*1, *a*2) = −B(*a*2, *a*1), for any *a*1, *a*² ∈ *A*;
- $-$ B(−, −) is called *invariant* if B(a_1a_2 , a_3) = B(a_1 , $a_2a_3 a_3a_2$), for any $a_1, a_2, a_3 \in A$.

Let B(−, −) be a skew-symmetric invariant bilinear form on an anti-Leibniz algebra (*A*, ·). Then, we also have $\mathfrak{B}(a_1a_2, a_3) = -\mathfrak{B}(a_1a_3, a_2) = \mathfrak{B}(a_2, a_1a_3)$, for any $a_1, a_2, a_3 \in A$.

Lemma 2.8. Let (A, \cdot) be an anti-Leibniz algebra and (A, I_A, r_A) be the regular bimodule over (A, \cdot) . *Then* (A, I_A, r_A) *and* (A^*, I_A^*) A^*_{A} , **l**_A^{*} – **r**_A^{*} *A*) *are isomorphic as bimodules over* (*A*, ·) *if and only if there exists a nondegenerate skew-symmetric invariant bilinear form* B(−, −) *on* (*A*, ·)*.*

Proof. Suppose that there exists a nondegenerate skew-symmetric invariant bilinear form $\mathfrak{B}(-, -)$ on (*A*, ·). Then there exists a linear isomorphism $\varphi : A \to A^*$ defined by $\langle \varphi(a_1), a_2 \rangle = \mathfrak{B}(a_1, a_2)$. Moreover, note that, for any $a_1, a_2, a_3 \in A$,

$$
\langle \varphi(\mathbf{l}_A(a_1)(a_2)), a_3 \rangle = \mathfrak{B}(a_1 a_2, a_3) = \mathfrak{B}(a_2, a_1 a_3) = \langle \mathbf{l}_A^*(a_1)(\varphi(a_2)), a_3 \rangle,
$$

$$
\langle \varphi(\mathbf{r}_A(a_1)(a_2)), a_3 \rangle = \mathfrak{B}(a_2 a_1, a_3) = \mathfrak{B}(a_2, a_1 a_3 - a_3 a_1) = \langle (\mathbf{l}_A^* - \mathbf{r}_A^*)(a_1)(\varphi(a_2)), a_3 \rangle,
$$

we get φ is an isomorphism. Conversely, in a similar way, we can get the conclusion.

Now we give the definition of Manin triple of anti-Leibniz algebras.

Definition 2.9. *Let* (*A*, ·) *be an anti-Leibniz algebra. Suppose that there is an anti-Leibniz algebra structure* (*A* ∗ , · ′) *on the dual space A*[∗] *of A and there is an anti-Leibniz algebra structure* ∗ *on the direct* $sum A \oplus A^*$ *such that* (A, \cdot) *and* (A^*, \cdot') *are subalgebras of* $(A \oplus A^*, *)$ *. If the natural nondegenerate symmetric bilinear form on A* ⊕ *A* [∗] *given by*

(2.11)
$$
\mathfrak{B}_d((a_1,\xi_1), (a_2,\xi_2)) := \langle \xi_1, a_2 \rangle - \langle \xi_2, a_1 \rangle,
$$

for any $a_1, a_2 \in A$ and $\xi_1, \xi_2 \in A^*$, *is invariant, then* $(A \oplus A^*, A, A^*)$ *is called a* (standard) Manin triple of anti-Leibniz algebras *associated to the standard bilinear form* \mathcal{B}_d *, and it is denoted by* (($A \oplus$ A^* , \mathfrak{B}_d , A , A^*) *for simply.*

For the matched pair of anti-Leibniz algebras and (standard) Manin triple of anti-Leibniz algebras, we have the following proposition.

Proposition 2.10. *Let* (*A*, ·) *be an anti-Leibniz algebra. Suppose that there is an anti-Leibniz algebra structure* (A^*, \cdot') *on* A^* . Then there exists an anti-Leibniz algebra structure on the vector space $A \oplus A^*$ such that $(A ⊕ A^*, A, A^*)$ is a (standard) Manin triple of anti-Leibniz algebras with respect to $\mathfrak{B}_d(−, −)$ *defined by* (2.11) *if and only if* (A, A^*, I_A^*) A^*_{A} , **l**_A^{*} – **r**_A^{*} *A* , l ∗ *^A*[∗] , l ∗ *^A*[∗] −r ∗ *^A*[∗]) *is a matched pair of anti-Leibniz algebras.*

Proof. If (A, A^*, \mathbf{I}_A^*) A^*_{A} , I^*_{A} – r^*_{A} $A^*_{A^*}, I^*_{A^*}, I^*_{A^*} - \mathbf{r}^*_{A^*}$) is a matched pair of anti-Leibniz algebras, we have an anti-Leibniz algebra $(A \oplus A^*, *)$, where the product $*$ is given by [\(2.7\)](#page-3-8). It is easy to see that (A, \cdot) and (A^*, \cdot') are subalgebras of $(A \oplus A^*, *)$. We only need to show that $\mathfrak{B}_d(-, -)$ is invariant. Indeed, for any

 $a_1, a_2, a_3 \in A$ and $\xi_1, \xi_2, \xi_3 \in A^*$,

$$
\mathcal{B}_{d}((a_{1},\xi_{1}) * (a_{2},\xi_{2}), (a_{3},\xi_{3}))
$$
\n
$$
= \langle \xi_{1}\xi_{2}, a_{3}\rangle + \langle I_{A}^{*}(a_{1})(\xi_{2}), a_{3}\rangle + \langle I_{A}^{*}(a_{2})(\xi_{1})
$$
\n
$$
- \mathbf{r}_{A}^{*}(a_{2})(\xi_{1}), a_{3}\rangle - \langle \xi_{3}, a_{1}a_{2}\rangle - \langle \xi_{3}, I_{A}^{*}(\xi_{1})(a_{2})\rangle - \langle \xi_{3}, I_{A}^{*}(\xi_{2})(a_{1}) + \mathbf{r}_{A}^{*}(\xi_{2})(a_{1})\rangle
$$
\n
$$
= \langle \xi_{1}, a_{2}a_{3}\rangle - \langle \xi_{1}\xi_{3}, a_{2}\rangle - \langle \xi_{2}\xi_{3}, a_{1}\rangle + \langle \xi_{2}, a_{1}a_{3}\rangle
$$
\n
$$
- \langle \xi_{1}, a_{3}a_{2}\rangle + \langle \xi_{1}\xi_{2}, a_{3}\rangle + \langle \xi_{3}\xi_{2}, a_{1}\rangle - \langle \xi_{3}, a_{1}a_{2}\rangle
$$
\n
$$
= \mathfrak{B}_{d}((a_{1},\xi_{1}), (a_{2},\xi_{2}) * (a_{3},\xi_{3}) - (a_{3},\xi_{3}) * (a_{2},\xi_{2})).
$$

Thus, $((A \oplus A^*, \mathfrak{B}_d), A, A^*)$ is a (standard) Manin triple of anti-Leibniz algebras.

Conversely, if $((A \oplus A^*, \mathfrak{B}_d), A, A^*)$ is a (standard) Manin triple of anti-Leibniz algebras, then $\mathfrak{B}_d(-, -)$ induces an isomorphism from $A \oplus A^*$ to $A \oplus A^*$. That is, the anti-Leibniz algebra structure on $A \oplus A^*$ just the crossed product $A \bowtie A^*$. Thus (A, A^*, I^*_A) A^*_{A} , **l**_A^{*} – **r**_A^{*} $A^*_{A^*}, I^*_{A^*} - \mathbf{r}^*_{A^*}$ is a matched pair of anti-Leibniz \triangle algebras.

Finally, we consider anti-Leibniz bialgebras. Let *A* be a vector space and $\Delta : A \to A \otimes A$ be a linear map. Then (*A*, ∆) is called an *anti-Leibniz coalgebra* if

$$
(\Delta \otimes id)\Delta + (id \otimes \Delta)\Delta + (\tau \otimes id)(id \otimes \Delta)\Delta = 0,
$$

where the twisting map $\tau : A \otimes A \to A \otimes A$ is given by $\tau(a_1 \otimes a_2) = a_2 \otimes a_1$ for all $a_1, a_2 \in A$. Clearly, the notion of anti-Leibniz coalgebras is the dualization of the notion of anti-Leibniz algebras, that is, (A, Δ) is an anti-Leibniz coalgebra if and only if (A^*, Δ^*) is an anti-Leibniz algebra.

Definition 2.11. An anti-Leibniz bialgebra *is a triple* (A, \cdot, Δ) *containing an anti-Leibniz algebra* (A, \cdot) *and an anti-Leibniz coalgebra* (A, Δ) *such that for any a*₁, *a*₂ ∈ *A*,

(2.12)
$$
\Delta(a_1a_2) + ((id \otimes \mathbf{r}_A(a_2) - \mathbf{r}_A(a_2) \otimes id + \mathbf{l}_A(a_2) \otimes id)(id \otimes id - \tau))\Delta(a_1) + (id \otimes \mathbf{l}_A(a_1) + \mathbf{l}_A(a_1) \otimes id)\Delta(a_2) = 0,
$$

(2.13)
$$
(\mathbf{r}_A(a_1) \otimes \mathrm{id})\Delta(a_2) - \tau((\mathbf{r}_A(a_2) \otimes \mathrm{id})\Delta(a_1)) = 0.
$$

Example 2.12. Consider the anti-Leibniz algebra $\Lambda_1^2 = (\mathbb{k}\{e_1, e_2\}, \cdot)$ given in Example [2.2,](#page-2-0) define linear $map \ \overline{\Lambda} : \Lambda_1^2 \to \Lambda_1^2 \otimes \Lambda_1^2$ 1 *by* ∆(*e*1) = *ke*² ⊗ *e*2*, k* ∈ k*, and* ∆(*e*2) = 0*, then we get a* 2*-dimensional anti-Leibniz bialgebra.*

Let (A, \cdot, Δ) and (B, \cdot', Δ') be two anti-Leibniz bialgebras. A linear map $\psi : A \to B$ is a *homomorphism of anti-Leibniz bialgebras* if ψ satisfies, for any $a_1, a_2 \in A$, $\psi(a_1 \cdot a_2) = \psi(a_1) \cdot' \psi(a_2)$ and $(\psi \otimes \psi) \Delta = \Delta' \psi$.

Proposition 2.13. *Let* (A, \cdot) *be an anti-Leibniz algebra. Suppose that there is a linear map* $\Delta : A \to A \otimes A$ *such that* (*A*, ∆) *is an anti-coassociative coalgebra. Then the triple* (*A*, ·, ∆) *is an anti-Leibniz bialgebra if and only if* (A, A^*, I^*_A) A^*_{A} , **l**_A^{*} – **r**_A^{*} ^{*}_A, **l**^{*}_A_{*}, **l**^{*}_A_{*} \bullet *is a matched pair of anti-Leibniz algebras, where* **(***A***^{*}, ∆^{*})** *is the dual algebra of anti-coassociative coalgebra* (A, Δ) *.*

Proof. For any $a_1, a_2 \in A$ and $\xi_1, \xi_2 \in A^*$, since

$$
\langle \mathbf{r}_{A^*}^*(\xi_2)(a_1)a_2, \xi_1 \rangle - \langle \mathbf{l}_{A^*}^*(\mathbf{r}_A^*(a_1)(\xi_2))(a_2), \xi_1 \rangle
$$

= $\langle a_1, \mathbf{r}_A^*(a_2)(\xi_1)\xi_2 \rangle - \langle a_2, \mathbf{r}_A^*(a_1)(\xi_2)\xi_1 \rangle$
= $\langle (\mathbf{r}_A(a_2) \otimes \text{id})\Delta(a_1), \xi_1 \otimes \xi_2 \rangle - \langle ((\mathbf{r}_A(a_1) \otimes \text{id})\Delta(a_2), \xi_2 \otimes \xi_1 \rangle,$

we get Eq. $(2.10) \Leftrightarrow$ Eq. (2.13) . Moreover, note that

$$
\langle I_{A^*}^*(\xi_2)(a_1a_2), \xi_1 \rangle + \langle I_{A^*}^*(\xi_2)(a_1)a_2, \xi_1 \rangle + \langle a_1 I_{A^*}^*(\xi_2)(a_2), \xi_1 \rangle + \langle I_{A^*}^*(I_A^*(a_1)(\xi_2))(a_2), \xi_1 \rangle - \langle I_{A^*}^*(\mathbf{r}_A^*(a_1)(\xi_2))(a_2), \xi_1 \rangle + \langle I_{A^*}^*(I_A^*(a_2)(\xi_2))(a_1), \xi_1 \rangle - \langle \mathbf{r}_{A^*}^*(I_A^*(a_2)(\xi_2))(a_1), \xi_1 \rangle - \langle I_{A^*}^*(\mathbf{r}_A^*(a_2)(\xi_2))(a_1), \xi_1 \rangle + \langle \mathbf{r}_{A^*}^*(\mathbf{r}_A^*(a_2)(\xi_2))(a_1), \xi_1 \rangle
$$

 $= \langle \Delta(a_1a_2), \xi_2 \otimes \xi_1 \rangle + \langle (id \otimes r_A(a_2))\Delta(a_1), \xi_2 \otimes \xi_1 \rangle + \langle (id \otimes l_A(a_1))\Delta(a_2), \xi_2 \otimes \xi_1 \rangle$ + $\langle (l_A(a_1) \otimes id) \Delta(a_2), \xi_2 \otimes \xi_1 \rangle - \langle (r_A(a_1) \otimes id) \Delta(a_2), \xi_2 \otimes \xi_1 \rangle + \langle (l_A(a_2) \otimes id) \Delta(a_1), \xi_2 \otimes \xi_1 \rangle$ $-\langle (id \otimes I_A(a_2))\Delta(a_1), \xi_1 \otimes \xi_2 \rangle - \langle (r_A(a_2) \otimes id)\Delta(a_1), \xi_2 \otimes \xi_1 \rangle + \langle (id \otimes r_A(a_2))\Delta(a_1), \xi_1 \otimes \xi_2 \rangle,$

we get Eq. (2.9) holds if and only if

$$
0 = \Delta(a_1a_2) + (\mathrm{id} \otimes \mathbf{r}_A(a_2))\Delta(a_1) + (\mathrm{id} \otimes \mathbf{l}_A(a_1))\Delta(a_2) + (\mathbf{l}_A(a_1) \otimes \mathrm{id})\Delta(a_2) - (\mathbf{r}_A(a_1) \otimes \mathrm{id})\Delta(a_2)
$$

+ $(\mathbf{l}_A(a_2) \otimes \mathrm{id})\Delta(a_1) - \tau(\mathrm{id} \otimes \mathbf{l}_A(a_2))\Delta(a_1) - (\mathbf{r}_A(a_2) \otimes \mathrm{id})\Delta(a_1) + \tau(\mathrm{id} \otimes \mathbf{r}_A(a_2))\Delta(a_1)$
= $\Delta(a_1a_2) + (\mathrm{id} \otimes \mathbf{r}_A(a_2))\Delta(a_1) + (\mathrm{id} \otimes \mathbf{l}_A(a_1))\Delta(a_2) + (\mathbf{l}_A(a_1) \otimes \mathrm{id})\Delta(a_2) - \tau(\mathbf{r}_A(a_2) \otimes \mathrm{id})\Delta(a_1)$
+ $(\mathbf{l}_A(a_2) \otimes \mathrm{id})\Delta(a_1) - \tau(\mathrm{id} \otimes \mathbf{l}_A(a_2))\Delta(a_1) - (\mathbf{r}_A(a_2) \otimes \mathrm{id})\Delta(a_1) + \tau(\mathrm{id} \otimes \mathbf{r}_A(a_2))\Delta(a_1)$
= $\Delta(a_1a_2) + ((\mathrm{id} \otimes \mathbf{r}_A(a_2) - \mathbf{r}_A(a_2) \otimes \mathrm{id} + \mathbf{l}_A(a_2) \otimes \mathrm{id})((\mathrm{id} \otimes \mathrm{id} - \tau))\Delta(a_1)$
+ $(\mathrm{id} \otimes \mathbf{l}_A(a_1) + \mathbf{l}_A(a_1) \otimes \mathrm{id})\Delta(a_2).$

That is, Eq. $(2.9) \Leftrightarrow$ Eq. (2.12) if Eq. (2.9) holds. Similarly, we also have Eq. $(2.8) \Leftrightarrow$ $(2.8) \Leftrightarrow$ Eq. (2.12) if Eq. [\(2.9\)](#page-3-5) holds. Thus, (A, \cdot, Δ) is an anti-Leibniz bialgebra if and only if (A, A^*, I^*_{A}) A^*_{A} , **l**_A^{*} – **r**_A^{*} $A^*_{A^*}, I^*_{A^*} - \mathbf{r}^*_{A^*}$) is a matched pair of anti-Leibniz algebras.

Combining Propositions [2.10](#page-4-1) and [2.13,](#page-5-2) we have the following conclusion.

Theorem 2.14. *Let* (A, \cdot) *be an anti-Leibniz algebra. Suppose that there is a linear map* $\Delta : A \to A \otimes A$ *such that* (*A*, ∆) *is an anti-Leibniz coalgebra. Then the following conditions are equivalent:*

- (i) (A, A^*, \mathbf{r}_A^*) $_A^*, \mathbf{l}_A^*$ *A* , r ∗ *^A*[∗] , l ∗ *^A*[∗]) *is a matched pair of anti-Leibniz algebras;*
- (*ii*) $((A \oplus A^*, \mathfrak{B}_d^*, A, A^*)$ *is a (standard) Manin triple of anti-Leibniz algebras* (A, \cdot) *and* (A^*, Δ^*) ;
- (*iii*) (*A*, ·, ∆) *is an anti-Leibniz bialgebra.*

As a direct conclusion, we have

Corollary 2.15. *Let* (A, \cdot, Δ) *be an anti-Leibniz bialgebra. Then* $(A^*, \Delta^*, \overline{\Delta})$ *is also an anti-Leibniz bialgebra, where* $\bar{\Delta}: A^* \to A^* \otimes A^*$ *is given by* $\langle \bar{\Delta}(\xi), a_1 \otimes a_2 \rangle = \langle \xi, a_1 a_2 \rangle$, for any $\xi \in A^*$ *and* $a_1, a_2 \in A$.

3. Some special anti-Leibniz bialgebras and anti-Leibniz Yang-Baxter equation

In this section, we consider some special classes of anti-Leibniz bialgebras called quasi-triangular anti-Leibniz bialgebras, triangular anti-Leibniz bialgebras, factorizable anti-Leibniz bialgebras, and introduce the notion of the Yang-Baxter equation in an anti-Leibniz algebra. The notion of relative Rota-Baxter operators of anti-Leibniz algebras is introduced to provide antisymmetric solutions of the Yang-Baxter equation in semi-direct product anti-Leibniz algebras and hence give rise to anti-Leibniz bialgebras.

3.1. Triangular anti-Leibniz bialgebras. Here, we consider a special class of anti-Leibniz bialgebras (A, \cdot, Δ_r) , where the comultiplication Δ_r is given by

(3.1)
$$
\Delta_r(a) := ((\mathbf{r}_A - \mathbf{l}_A)(a) \otimes \mathrm{id} + \mathrm{id} \otimes \mathbf{r}_A(a))(r),
$$

for any $a \in A$ and some $r \in A \otimes A$. In this case, we also call (A, \cdot, Δ_r) is an anti-Leibniz bialgebra induced by *r*. Such anti-Leibniz bialgebras are quite similar as the coboundary Lie bialgebras for Lie algebras and the coboundary infinitesimal bialgebras for associative algebras. Now, we consider the anti-Leibniz coalgebra (A, Δ_r) .

Proposition 3.1. Let (A, \cdot) be an anti-Leibniz algebra and $r = \sum_i x_i \otimes y_i \in A \otimes A$. Define a linear map Δ_r : *A* → *A* ⊗ *A by Eq.* [\(3.1\)](#page-6-2)*.* Then (*A*, Δ_r) *is an anti-Leibniz coalgebra if and only if*

$$
(3.2) \quad ((\mathbf{l}_A - \mathbf{r}_A)(a) \otimes \mathrm{id} \otimes \mathrm{id})([\![r, r]\!]) + (\mathrm{id} \otimes \mathrm{id} \otimes \mathbf{r}_A(a) + \mathrm{id} \otimes (\mathbf{l}_A - \mathbf{r}_A)(a) \otimes \mathrm{id})((\![r, r]\!])
$$

$$
- (\mathrm{id} \otimes \mathrm{id} \otimes \mathbf{r}_A(a) - \mathrm{id} \otimes (\mathbf{l}_A - \mathbf{r}_A)(a) \otimes \mathrm{id}) (\sum_i ((\mathbf{r}_A - \mathbf{l}_A)(x_i) \otimes \mathrm{id} + \mathrm{id} \otimes \mathbf{r}_A(x_i))(r - \tau(r)) \otimes y_i)
$$

+
$$
\sum_j ((\mathbf{l}_A - \mathbf{r}_A)(x_j) \otimes id \otimes id)(\tau(((\mathbf{r}_A - \mathbf{l}_A)(a) \otimes id + id \otimes \mathbf{r}_A(a))(r - \tau(r))) \otimes y_j),
$$

for any $a \in A$ *, where we denote by*

$$
\llbracket r, r \rrbracket := r_{12}r_{13} + r_{12}r_{23} - r_{23}r_{12} - r_{23}r_{13}
$$
\n
$$
= \sum_{i,j} x_i x_j \otimes y_i \otimes y_j + x_i \otimes y_i x_j \otimes y_j - x_j \otimes x_i y_j \otimes y_i - x_j \otimes x_i \otimes y_i y_j.
$$

Proof. First, for any $a \in A$, we have

$$
(\Delta_r \otimes id)\Delta_r(a) + (id \otimes \Delta_r)\Delta_r(a) + (\tau \otimes id)(id \otimes \Delta_r)\Delta_r(a)
$$

=
$$
\sum_{i,j} x_j \otimes y_j x_i \otimes y_i a - x_i x_j \otimes y_j \otimes y_i a + x_j x_i \otimes y_j \otimes y_i a - x_j \otimes y_j (ax_i) \otimes y_i
$$

+
$$
(ax_i)x_j \otimes y_j \otimes y_i - x_j (ax_i) \otimes y_j \otimes y_i + x_j \otimes y_j (x_i a) \otimes y_i - (x_i a)x_j \otimes y_j \otimes y_i
$$

+
$$
x_j(x_i a) \otimes y_j \otimes y_i + x_i \otimes x_j \otimes y_j (y_i a) - x_i \otimes (y_i a)x_j \otimes y_j + x_i \otimes x_j (y_i a) \otimes y_j
$$

-
$$
ax_i \otimes x_j \otimes y_j y_i + ax_i \otimes y_i x_j \otimes y_j - ax_i \otimes x_j y_i \otimes y_j + x_i a \otimes x_j \otimes y_j y_i
$$

-
$$
x_i a \otimes y_i x_j \otimes y_j + x_i a \otimes x_j y_i \otimes y_j + x_j \otimes x_i \otimes y_j (y_i a) - (y_i a)x_j \otimes x_i \otimes y_j
$$

+
$$
x_j (y_i a) \otimes x_i \otimes y_j - x_j \otimes ax_i \otimes y_j y_i + y_j x_j \otimes ax_i \otimes y_j - x_j y_i \otimes ax_i \otimes y_j
$$

+
$$
x_j \otimes x_i a \otimes y_j y_i - y_i x_j \otimes x_i a \otimes y_j + x_j y_i \otimes x_i a \otimes y_j.
$$

By the anti-Leibniz identity, we get

A := X *i*, *j x^j* ⊗ *yjxⁱ* ⊗ *yia* − *xix^j* ⊗ *y^j* ⊗ *yia* + *xjxⁱ* ⊗ *y^j* ⊗ *yia* + *xⁱ* ⊗ *x^j* ⊗ *yj*(*yia*) + *x^j* ⊗ *xⁱ* ⊗ *yj*(*yia*) =(id ⊗ id ⊗ r*A*(*a*)) (τ ⊗ id) [[*r*,*r*]] + X *j* ((r*^A* − l*A*)(*xj*) ⊗ id + id ⊗ r*A*(*xj*))(*r* − τ(*r*)) ⊗ *yja*, B := X *i*, *j* −*xj*(*axi*) ⊗ *y^j* ⊗ *yⁱ* − *axⁱ* ⊗ *x^j* ⊗ *yjyⁱ* + *axⁱ* ⊗ *yix^j* ⊗ *y^j* − *axⁱ* ⊗ *xjyⁱ* ⊗ *yj*) =(l*A*(*a*) ⊗ id ⊗ id) [[*r*,*r*]] + X *i*, *j* (*axi*)*x^j* ⊗ *yⁱ* ⊗ *y^j* , C := X *i*, *j* −*x^j* ⊗ *axⁱ* ⊗ *yjyⁱ* + *yix^j* ⊗ *axⁱ* ⊗ *y^j* − *xjyⁱ* ⊗ *axⁱ* ⊗ *yj*) =(id ⊗ l*A*(*a*) ⊗ id)(τ ⊗ id) [[*r*,*r*]] − X *i*, *j yⁱ* ⊗ *a*(*xixj*) ⊗ *y^j* , D := X *i*, *j xj*(*xia*) ⊗ *y^j* ⊗ *yⁱ* + *xia* ⊗ *x^j* ⊗ *yjyⁱ* − *xia* ⊗ *yix^j* ⊗ *y^j* + *xia* ⊗ *xjyⁱ* ⊗ *yj*) = − (r*A*(*a*) ⊗ id ⊗ id) [[*r*,*r*]] − X *i*, *j xj*(*xia*) ⊗ *yⁱ* ⊗ *y^j* , E := X *i*, *j x^j* ⊗ *xia* ⊗ *yjyⁱ* − *yix^j* ⊗ *xia* ⊗ *y^j* + *xjyⁱ* ⊗ *xia* ⊗ *yj*) = − (id ⊗r*A*(*a*) ⊗ id) (τ ⊗ id) [[*r*,*r*]] + X *i*, *j yⁱ* ⊗ (*xixj*)*a* ⊗ *y^j* ,

Second, since $(ax_i)x_j \otimes y_j \otimes y_i - (x_ia)x_j \otimes y_j \otimes y_i = 0$, we obtain

$$
(\Delta_r \otimes id)\Delta_r(a) + (id \otimes \Delta_r)\Delta_r(a) + (\tau \otimes id)(id \otimes \Delta_r)\Delta_r(a)
$$

= $\mathcal{A} + \mathcal{B} + C + \mathcal{D} + \mathcal{E} + \left(\sum_{i,j} x_j(y_i a) \otimes x_i \otimes y_j - (y_i a)x_j \otimes x_i \otimes y_j\right)$
 $- x_j \otimes y_j(ax_i) \otimes y_i + x_j \otimes y_j(x_i a) \otimes y_i - x_i \otimes (y_i a)x_j \otimes y_j + x_i \otimes x_j(y_i a) \otimes y_j)$
= $((\mathbf{I}_A - \mathbf{r}_A)(a) \otimes id \otimes id)([[r, r]]) + (id \otimes id \otimes \mathbf{r}_A(a) + id \otimes (\mathbf{I}_A - \mathbf{r}_A)(a) \otimes id)(([[r, r]]))$

+
$$
\left(\sum_{i,j} x_j(y_ia) \otimes x_i \otimes y_j - (y_ia)x_j \otimes x_i \otimes y_j - x_j \otimes y_j(ax_i) \otimes y_i + x_j \otimes y_j(x_ia) \otimes y_i\right) - x_i \otimes (y_ia)x_j \otimes y_j + x_i \otimes x_j(y_ia) \otimes y_j + (ax_i)x_j \otimes y_i \otimes y_j - y_i \otimes a(x_ix_j) \otimes y_j - x_j(x_ia) \otimes y_i \otimes y_j + y_i \otimes (x_ix_j)a \otimes y_j\right) + \sum_j ((\mathbf{r}_A - \mathbf{l}_A)(x_j) \otimes id + id \otimes \mathbf{r}_A(x_j))(r - \tau(r)) \otimes y_j a.
$$

Finally, note that

$$
- \sum_{j} ((\mathbf{I}_{A} - \mathbf{r}_{A})(x_{j}) \otimes \mathrm{id} \otimes \mathrm{id}) \tau (((\mathbf{r}_{A} - \mathbf{I}_{A})(a) \otimes \mathrm{id} + \mathrm{id} \otimes \mathbf{r}_{A}(a))(r - \tau(r))) \otimes y_{j}
$$

=
$$
\sum_{i,j} x_{j}(x_{i}a) \otimes y_{i} \otimes y_{j} - x_{j}(y_{i}a) \otimes x_{i} \otimes y_{j} + x_{j}y_{i} \otimes ax_{i} \otimes y_{j} - x_{j}y_{i} \otimes x_{i}a \otimes y_{j}
$$

$$
- x_{j}x_{i} \otimes ay_{i} \otimes y_{j} + x_{j}x_{i} \otimes y_{i}a \otimes y_{j} - (x_{i}a)x_{j} \otimes y_{i} \otimes y_{j} + (y_{i}a)x_{j} \otimes x_{i} \otimes y_{j}
$$

$$
- y_{i}x_{j} \otimes ax_{i} \otimes y_{j} + y_{i}x_{j} \otimes x_{i}a \otimes y_{j} + x_{i}x_{j} \otimes ay_{i} \otimes y_{j} - x_{i}x_{j} \otimes y_{i}a \otimes y_{j},
$$

and

$$
- (\mathrm{id} \otimes (\mathbf{l}_A + \mathbf{r}_A)(a) \otimes \mathrm{id}) \Big(\sum_j ((\mathbf{r}_A - \mathbf{l}_A)(x_j) \otimes \mathrm{id} + \mathrm{id} \otimes \mathbf{r}_A(x_j))(r - \tau(r)) \otimes y_j \Big)
$$

=
$$
\sum_{i,j} y_i x_j \otimes a x_i \otimes y_j - y_i x_j \otimes x_i a \otimes y_j - x_i x_j \otimes a y_i \otimes y_j + x_i x_j \otimes y_i a \otimes y_j
$$

$$
- x_j y_i \otimes a x_i \otimes y_j + x_j y_i \otimes x_i a \otimes y_j + x_j x_i \otimes a y_i \otimes y_j - x_j x_i \otimes y_i a \otimes y_j
$$

$$
- x_i \otimes a(y_i x_j) \otimes y_j + x_i \otimes (y_i x_j) a \otimes y_j + y_i \otimes a(x_i x_j) \otimes y_j - y_i \otimes (x_i x_j) a \otimes y_j,
$$

we get

$$
(\Delta_r \otimes id)\Delta_r(a) + (id \otimes \Delta_r)\Delta_r(a) + (\tau \otimes id)(id \otimes \Delta_r)\Delta_r(a)
$$

= $((\mathbf{l}_A - \mathbf{r}_A)(a) \otimes id \otimes id)([\![r, r]\!]) + (id \otimes id \otimes \mathbf{r}_A(a) + id \otimes (\mathbf{l}_A - \mathbf{r}_A)(a) \otimes id)([\![r, r]\!])$
- $(id \otimes id \otimes \mathbf{r}_A(a) - id \otimes (\mathbf{l}_A - \mathbf{r}_A)(a) \otimes id)(\sum_i ((\mathbf{r}_A - \mathbf{l}_A)(x_i) \otimes id + id \otimes \mathbf{r}_A(x_i))(r - \tau(r)) \otimes y_i)$
+ $\sum_j ((\mathbf{l}_A - \mathbf{r}_A)(x_j) \otimes id \otimes id)(\tau(((\mathbf{r}_A - \mathbf{l}_A)(a) \otimes id + id \otimes \mathbf{r}_A(a))(r - \tau(r))) \otimes y_j).$

Thus, (A, Δ_r) is an anti-Leibniz coalgebra if and only if Eq. [\(4.1\)](#page-24-0) holds.

Next, we consider the anti-Leibniz bialgebra (A, \cdot, Δ_r) .

Proposition 3.2. Let (A, \cdot) be an anti-Leibniz algebra and $r = \sum_i x_i \otimes y_i \in A \otimes A$. Define a linear map Δ_r : $A \rightarrow A \otimes A$ by Eq. [\(3.1\)](#page-6-2). Then Eq. [\(2.12\)](#page-5-1) if and only if

$$
(3.3) \qquad ((\mathbf{r}_A - \mathbf{l}_A)(a_2) \otimes id)\tau + (id \otimes id - \tau)(\mathbf{r}_A(a_2) \otimes id))((\mathbf{r}_A - \mathbf{l}_A)(a_1) \otimes id + id \otimes \mathbf{r}_A(a_1))(r - \tau(r)))
$$

+ $\tau(\mathbf{r}_A(a_1) \otimes id)((\mathbf{r}_A - \mathbf{l}_A)(a_2) \otimes id + id \otimes \mathbf{r}_A(a_2))(r - \tau(r))) = 0,$

and Eq. [\(2.13\)](#page-5-0) *if and only if*

(3.4)
$$
(\mathbf{r}_A(a_1) \otimes id)((\mathbf{r}_A - \mathbf{l}_A)(a_2) \otimes id + id \otimes \mathbf{r}_A(a_2))(r - \tau(r))) = 0,
$$

for any $a_1, a_2 \in A$.

Proof. First, by the definition of Δ_r , for any $a_1, a_2 \in A$, we have

$$
\Delta_r(a_1a_2) + ((id \otimes \mathbf{r}_A(a_2) - \mathbf{r}_A(a_2) \otimes id + \mathbf{l}_A(a_2) \otimes id)(id - \tau))\Delta_r(a_1)
$$

+
$$
(id \otimes \mathbf{l}_A(a_1) + \mathbf{l}_A(a_1) \otimes id)\Delta_r(a_2)
$$

=
$$
\sum_i x_i(a_1a_2) \otimes y_i - (a_1a_2)x_i \otimes y_i + x_i \otimes y_i(a_1a_2) - a_1x_i \otimes y_i a_2 + x_i a_1 \otimes y_i a_2
$$

+
$$
x_i \otimes (y_i a_1)a_2 + (a_1x_i)a_2 \otimes y_i - (x_i a_1)a_2 \otimes y_i - x_i a_2 \otimes y_i a_1 - a_2(a_1x_i) \otimes y_i
$$

$$
\Box
$$

+
$$
a_2(x_ia_1) \otimes y_i + a_2x_i \otimes y_ia_1 + y_i \otimes (a_1x_i)a_2 - y_i \otimes (x_ia_1)a_2 - y_ia_1 \otimes x_ia_2
$$

\n- $y_ia_2 \otimes a_1x_i + y_ia_2 \otimes x_ia_1 + (y_ia_1)a_2 \otimes x_i + a_2y_i \otimes a_1x_i - a_2y_i \otimes x_ia_1$
\n- $a_2(y_ia_1) \otimes x_i - a_2x_i \otimes a_1y_i + x_ia_2 \otimes a_1y_i + x_i \otimes a_1(y_ia_2) - a_1(a_2x_i) \otimes y_i$
\n+ $a_1(x_ia_2) \otimes y_i + a_1x_i \otimes y_ia_2$.

Note that

$$
\sum_{i} x_{i} \otimes y_{i}(a_{1}a_{2}) + x_{i} \otimes a_{1}(y_{i}a_{2}) + x_{i} \otimes (y_{i}a_{1})a_{2} = 0 = \sum_{i} y_{i} \otimes (a_{1}x_{i})a_{2} - y_{i} \otimes (x_{i}a_{1})a_{2},
$$
\n
$$
\sum_{i} -(a_{1}a_{2})x_{i} \otimes y_{i} - a_{1}(a_{2}x_{i}) \otimes y_{i} - a_{2}(a_{1}x_{i}) \otimes y_{i} = 0 = \sum_{i} (a_{1}x_{i})a_{2} \otimes y_{i} - (x_{i}a_{1})a_{2} \otimes y_{i},
$$
\n
$$
\sum_{i} x_{i}a_{1} \otimes y_{i}a_{2} - y_{i}a_{1} \otimes x_{i}a_{2} = -\tau ((\mathbf{r}_{A}(a_{2}) \otimes \text{id})(((\mathbf{r}_{A} - \mathbf{I}_{A})(a_{1}) \otimes \text{id} + \text{id} \otimes \mathbf{r}_{A}(a_{1}))(r - \tau(r))))),
$$
\n
$$
\sum_{i} y_{i}a_{2} \otimes x_{i}a_{1} - x_{i}a_{2} \otimes y_{i}a_{1} = \tau ((\mathbf{r}_{A}(a_{1}) \otimes \text{id})(((\mathbf{r}_{A} - \mathbf{I}_{A})(a_{2}) \otimes \text{id} + \text{id} \otimes \mathbf{r}_{A}(a_{2}))(r - \tau(r))))),
$$
\n
$$
\sum_{i} a_{2}y_{i} \otimes a_{1}x_{i} - a_{2}x_{i} \otimes a_{1}y_{i} + a_{2}x_{i} \otimes y_{i}a_{1}
$$
\n
$$
= -(1_{A}(a_{2}) \otimes \text{id}) (\tau (((\mathbf{r}_{A} - \mathbf{I}_{A})(a_{1}) \otimes \text{id} + \text{id} \otimes \mathbf{r}_{A}(a_{1}))(r - \tau(r))))),
$$
\n
$$
\sum_{i} x_{i}a_{2} \otimes a_{1}y_{i} - y_{i}a_{2} \otimes a_{1}x_{i} + x_{i}(a_{1}a_{2}) \otimes y_{i} + a_{1}(x_{i}a_{2}) \otimes y_{i} + (y
$$

we get Eq. [\(2.12\)](#page-5-1) holds if and only if Eq. [\(3.3\)](#page-8-0) holds. Finally, since for any $a_1, a_2 \in A$,

$$
(\mathbf{r}_A(a_1) \otimes \mathrm{id})\Delta_r(a_2) - \tau((\mathbf{r}_A(a_2) \otimes \mathrm{id})\Delta_r(a_1))
$$

=
$$
\sum_i x_i a_1 \otimes y_i a_2 - y_i a_1 \otimes x_i a_2
$$

=
$$
(\mathbf{r}_A(a_1) \otimes \mathrm{id})(((\mathbf{r}_A - \mathbf{l}_A)(a_2) \otimes \mathrm{id} + \mathrm{id} \otimes \mathbf{r}_A(a_2))(r - \tau(r)))
$$

 Δ

we obtain this proposition. \Box

Definition 3.3. *Let* (A, \cdot) *be an anti-Leibniz algebra and* $r ∈ A ⊗ A$ *. Then the equation*

(3.5)
$$
[[r, r]] := r_{12}r_{13} + r_{12}r_{23} - r_{23}r_{12} - r_{23}r_{13} = 0
$$

is called the anti-Leibniz Yang-Baxter equation *(or* aLYBE*) in the anti-Leibniz algebra* (*A*, ·)*.*

An element r ∈ *A* ⊗ *A is called an* anti-Leibniz *r*-matrix *if r is a solution of the* aLYBE*, is called* invariant, *if for any* $a \in A$,

(3.6)
$$
((\mathbf{r}_A - \mathbf{l}_A)(a) \otimes id + id \otimes \mathbf{r}_A(a))(r) = 0.
$$

By Propositions [3.1](#page-6-3) and [3.2,](#page-8-1) we can provide sufficient and necessary conditions for (A, \cdot, Δ_r) to be an anti-Leibniz bialgebra.

Corollary 3.4. *Let* (A, \cdot) *be an anti-Leibniz algebra,* $r = \sum_i x_i \otimes y_i \in A \otimes A$ *, and* $\Delta_r : A \to A \otimes A$ *be linear map given by Eq.* [\(3.1\)](#page-6-2). Then (A, \cdot, Δ_r) *is an anti-Leibniz bialgebra if and only if Eqs.* [\(4.1\)](#page-24-0)*,* [\(3.3\)](#page-8-0) *and* [\(3.4\)](#page-8-2) *hold. In particular,*

(*i*) *if* $r - \tau(r)$ *is invariant and* r *is a solution of the aLYBE in* (A, \cdot) *, or*

(*ii*) *if* r *is a symmetric solution of the* aLYBE *in* (A, \cdot) *,*

then (*A*, ·, ∆*r*) *is an anti-Leibniz bialgebra, which is called the anti-Leibniz bialgebra induced by the solution r.*

Definition 3.5. *Let* (A, \cdot) *be an anti-Leibniz algebra and* $r \in A \otimes A$ *. If* r *is a solution of the* aLYBE *in* (A, \cdot) *and r* − τ (*r*) *is invariant, the induced anti-Leibniz bialgebra* (A, \cdot, Δ_r) *is called* quasi-triangular; *if*

r is a symmetric solution of the aLYBE *in* (*A*, ·)*, the induced anti-Leibniz bialgebra* (*A*, ·, ∆*r*) *is called* triangular*.*

Example 3.6. *Let* $(A = \mathbb{k}\{e_1, e_2\},\cdot)$ *be the 2-dimensional anti-Leibniz algebra* Λ_1^2 1 *given in Example [2.2,](#page-2-0) i.e.,* $e_1e_1 = e_2$. Then $r = e_1 \otimes e_2 + e_2 \otimes e_1$ *is a symmetric solution of the aLYBE in* (A, \cdot) *. Thus we* get a triangular anti-Leibniz bialgebra (A, \cdot, Δ_r) , where the linear map $\Delta_r : A \to A \otimes A$ is given by $\Delta_r(e_1) = e_2 \otimes e_2$, $\Delta(e_2) = 0$. This triangular anti-Leibniz bialgebra just the 2-dimensional anti-Leibniz *bialgebra in Example [2.12.](#page-5-3)*

For any vector space *V*, we define $\tau_{13} \in \mathfrak{gl}(V \otimes V \otimes V)$ by $\tau_{13}(v_1 \otimes v_2 \otimes v_3) = v_3 \otimes v_2 \otimes v_1$ for any $v_1, v_2, v_3 \in V$. Then by direct calculations, we have

Proposition 3.7. *Let* (A, \cdot) *be an anti-Leibniz algebra and* $r = \sum_i x_i \otimes y_i \in A \otimes A$. *Then,*

$$
[\![\tau(r), \ \tau(r)]\!] = -\tau_{13}[\![r, \ r]\!].
$$

That is to say, r is a solution of the aLYBE in (A, \cdot) *if and only if* $\tau(r)$ *is a solution of the aLYBE in* (A, \cdot) *.*

Thus, by corollary [3.4](#page-9-0) and Proposition [3.7,](#page-10-0) we have

Corollary 3.8. *If* (A, \cdot, Δ_r) *is a quasi-triangular (resp. triangular) anti-Leibniz bialgebra, then* $(A, \cdot, \Delta_{\tau(r)})$ *is also a quasi-triangular (resp. triangular) anti-Leibniz bialgebra.*

For the construction of anti-Leibniz bialgebras, we have

Theorem 3.9. *Let* (*A*, ·, ∆) *be an anti-Leibniz bialgebra. Then there is a canonical anti-Leibniz bialgebra structure on* $A \oplus A^*$ *such that both* $\iota_A : A \to A \oplus A^*$ *, a* $\mapsto (a, 0)$ *, and* $\iota_{A^*} : A^* \to A \oplus A^*$ *,* $\xi \mapsto (0, \xi)$ *, are homomorphisms of anti-Leibniz bialgebras, where the anti-Leibniz bialgebra structure on A* ⊕ *A* ∗ *is* $(A \oplus A^*, *, \Delta_r)$, $\Delta_{\tilde{r}}$ *is given by Eq.* [\(3.1\)](#page-6-2) *for some* $\tilde{r} \in (A \oplus A^*) \otimes (A \oplus A^*)$.

Proof. Let $r \in A \otimes A^* \subset (A \oplus A^*) \otimes (A \oplus A^*)$ correspond to the identity map id: $A \to A$. That is, $\tilde{r} = \sum_{i} e_i \otimes f_i$, where $\{e_1, e_2, \dots, e_n\}$ is a basis of A and $\{f_1, f_2, \dots, f_n\}$ be its dual basis. Since (A, \cdot, Δ) be an anti-Leibniz bialgebra, there is an anti-Leibniz algebra structure on *A*^{*}, i.e., $\xi_1 \xi_2 = \Delta^* (\xi_1 \otimes \xi_2)$, and an anti-Leibniz algebra structure ∗ on *A* ⊕ *A* [∗] which is given by

$$
(a_1, \xi_1) * (a_2, \xi_2) = (a_1 a_2 + \mathbf{I}_{A^*}^*(\xi_1)(a_2) + (\mathbf{I}_{A^*}^* - \mathbf{r}_{A^*}^*)(\xi_2)(a_1), \xi_1 \xi_2 + \mathbf{I}_{A}^*(a_1)(\xi_2) + (\mathbf{I}_{A}^* - \mathbf{r}_{A}^*)(a_2)(\xi_1)),
$$

for any $a_1, a_2 \in A$ and $\xi_1, \xi_2 \in A^*$. We denote $A \bowtie A^* := (A \oplus A^*, *)$ and show that there is a quasitriangular anti-Leibniz bialgebra structure on $A \Join A^*$, i.e., Eqs. [\(3.5\)](#page-9-1) and [\(3.6\)](#page-9-2) hold. Note that for any $(e_s, f_t), (e_k, f_l), (e_p, f_q) \in A \oplus A^*$

$$
\langle \tilde{r}_{12}\tilde{r}_{13} + \tilde{r}_{12}\tilde{r}_{23} - \tilde{r}_{23}\tilde{r}_{12} - \tilde{r}_{23}\tilde{r}_{13}, (e_s, f_t) \otimes (e_k, f_l) \otimes (e_p, f_q) \rangle
$$

\n
$$
= \sum_{i,j} \langle e_i * e_j \otimes f_i \otimes f_j + e_i \otimes f_i * e_j \otimes f_j - e_j \otimes e_i * f_j \otimes f_i - e_j \otimes e_i \otimes f_i * f_j,
$$

\n
$$
(e_s, f_t) \otimes (e_k, f_l) \otimes (e_p, f_q) \rangle
$$

\n
$$
= \sum_{i,j} \langle f_t, e_i e_j \rangle \langle f_i, e_k \rangle \langle f_j, e_p \rangle + \langle f_t, e_i \rangle \langle f_i f_l, e_j \rangle \langle f_i, e_p \rangle + \langle f_t, e_i \rangle \langle f_l, e_j e_k \rangle \langle f_j, e_p \rangle
$$

\n
$$
= \langle f_t, e_i \rangle \langle f_l, e_k e_j \rangle \langle f_j, e_p \rangle - \langle f_t, e_j \rangle \langle f_j f_l, e_i \rangle \langle f_i, e_p \rangle + \langle f_t, e_j \rangle \langle f_l f_j, e_i \rangle \langle f_i, e_p \rangle
$$

\n
$$
= \langle f_t, e_j \rangle \langle f_j, e_i e_k \rangle \langle f_i, e_p \rangle - \langle f_t, e_j \rangle \langle f_l, e_i \rangle \langle f_i f_j, e_p \rangle
$$

\n
$$
= 0,
$$

we get $[\![\tilde{r}, \tilde{r}]\!] = 0$. Similarly, one can check that $((\mathbf{r}_{A \oplus A^*} - \mathbf{l}_{A \oplus A^*})(x) \otimes id + id \otimes \mathbf{r}_{A \oplus A^*}(x))(\tilde{r} - \tau(\tilde{r})) = 0$ for any $x \in A \bowtie A^*$. Hence, there is a quasi-triangular anti-Leibniz bialgebra structure on $A \bowtie A^*$ by Corollary [3.4.](#page-9-0)

For any $e_i \in A$, note that

$$
\Delta_{\tilde{r}}(e_j) = ((\mathbf{r}_{A \oplus A^*} - \mathbf{I}_{A \oplus A^*})(e_j) \otimes id + id \otimes \mathbf{r}_{A \oplus A^*}(e_j)) \Big(\sum_i e_i \otimes f_i \Big)
$$

\n
$$
= \sum_{i,k} e_i e_j \otimes f_i - e_j e_i \otimes f_i + e_i \otimes \langle f_i f_k, e_j \rangle e_k + e_i \otimes \langle f_i, e_j e_k \rangle f_k - e_i \otimes \langle f_i, e_k e_j \rangle f_k
$$

\n
$$
= \sum_{i,k} \langle f_i f_k, e_j \rangle e_i \otimes e_k
$$

\n
$$
= \Delta(e_j).
$$

Therefore, $\iota_A : A \to A \oplus A^*$ is a homomorphism of anti-Leibniz bialgebras. Similarly, $\iota_{A^*} : A^* \to A \oplus A^*$ is also a homomorphism of anti-Leibniz bialgebras. The proof is finished.

With the anti-Leibniz bialgebra structure given in the theorem above, $(A \oplus A^*, *, \Delta_{\tilde{r}})$ is called the *double of anti-Leibniz bialgebra* (*A*, ·, ∆).

Example 3.10. *Let* $(A = \mathbb{k}\{e_1, e_2\}, \cdot, \Delta)$ *be the* 2-dimensional anti-Leibniz bialgebra in Example [2.12,](#page-5-3) i.e., $e_1e_1 = e_2$, $\Delta(e_1) = e_2 \otimes e_2$. Denote $\{f_1, f_2\}$ the dual basis of $\{e_1, e_2\}$ and $\tilde{r} = e_1 \otimes f_1 + e_2 \otimes f_2$. Then *we get a* 4*-dimensional anti-Leibniz bialgebra* ($D(A) = \mathbb{k}{e_1, e_2, f_1, f_2}, \dots, \Delta_i}$ *, where*

$$
e_1 * e_1 = e_2,
$$
 $f_2 * f_2 = f_1,$ $e_1 * f_2 = f_1,$ $f_2 * e_1 = e_2,$
 $\Delta_{\tilde{r}}(e_1) = e_2 \otimes e_2,$ $\Delta_{\tilde{r}}(f_2) = f_1 \otimes f_1.$

3.2. Relative Rota-Baxter operator and anti-Leibniz Yang-Baxter equation. In this subsection, we are discussing the relationship between (relative) Rota-Baxter operators on an anti-Leibniz algebra and some special resolutions of the aLYBE, and show that each relative Rota-Baxter operator induces an anti-Leibniz bialgebra. First, we recall the definition of relative Rota-Baxter operator.

Definition 3.11. *Let* (*A*, ·) *be an anti-Leibniz algebra and* (*M*, l,r) *be a bimodule over* (*A*, ·)*. A linear map* $R : M \to A$ *is called a* relative Rota-Baxter operator of (A, \cdot) associated to (M, I, r) *if R satisfies*

$$
R(m_1)R(m_2) = R\Big(\mathfrak{l}(R(m_1))(m_2) + \mathfrak{r}(R(m_2))m_1\Big),
$$

for any $m_1, m_2 \in M$. In particular, if the bimodule (M, I, r) *is the regular bimodule* (A, I_A, r_A) *, the operator R is called a* Rota-Baxter operator *on* (*A*, ·)*.*

Let *V* be a vector space. For all $r \in V \otimes V$, there is a linear map $r^{\sharp}: V^* \to V$ in the following way:

$$
\langle \xi_1 \otimes \xi_2, r \rangle = \langle r^{\sharp}(\xi_1), \xi_2 \rangle,
$$

for any $\xi_1, \xi_2 \in V^*$. If $r = \sum_i x_i \otimes y_i \in V \otimes V$, it is easy to see that $r^{\sharp}(\xi) = \sum_i \langle \xi, x_i \rangle y_i$, for any $\xi \in V^*$. We say that $r \in V \otimes V$ is *nondegenerate* if the map r^{\sharp} is a linear isomorphism.

Proposition 3.12. Let (A, \cdot) be an anti-Leibniz algebra and $r = \sum_i x_i \otimes y_i \in A \otimes A$. Then r is a solution *of the* aLYBE *in* (A, \cdot) *if and only if for any* $\xi_1, \xi_2 \in A^*$,

$$
\tau(r)^{\sharp}(\xi_1)\tau(r)^{\sharp}(\xi_2)=\tau(r)^{\sharp}\big(\mathbf{I}_{A}^{*}(r^{\sharp}(\xi_1))(\xi_2)+(\mathbf{I}_{A}^{*}-\mathbf{r}_{A}^{*})(\tau(r)^{\sharp}(\xi_2))(\xi_1)\big).
$$

Proof. For any $\xi_1, \xi_2, \xi_3 \in A^*$, we have

$$
\langle \tau(r)^{\sharp}(\xi_1)\tau(r)^{\sharp}(\xi_2), \xi_3 \rangle = \langle r, \ \mathbf{r}_A^*(\tau(r)^{\sharp}(\xi_2))(\xi_3) \otimes \xi_1 \rangle = \sum_i \langle x_i, \ \mathbf{r}_A^*(\tau(r)^{\sharp}(\xi_2))(\xi_3) \rangle \langle y_i, \xi_1 \rangle
$$

\n
$$
= \sum_i \langle r, \ \mathbf{l}_A^*(x_i)(\xi_3) \otimes \xi_2 \rangle \langle y_i, \xi_1 \rangle = \sum_{i,j} \langle x_j, \ \mathbf{l}_A^*(x_i)(\xi_3) \rangle \langle y_j, \xi_2 \rangle \langle y_i, \xi_1 \rangle
$$

\n
$$
= \sum_{i,j} \langle y_i \otimes y_j \otimes x_i x_j, \ \xi_1 \otimes \xi_2 \otimes \xi_3 \rangle.
$$

Similarly, we have

$$
\langle \tau(r)^{\sharp}(\mathbf{I}_{A}^{*}(r^{\sharp}(\xi_{1}))(\xi_{2})), \xi_{3} \rangle = \sum_{i,j} \langle x_{j} \otimes y_{j}y_{i} \otimes x_{i}, \xi_{1} \otimes \xi_{2} \otimes \xi_{3} \rangle,
$$

$$
\langle \tau(r)^{\sharp}(\mathbf{I}_{A}^{*}(\tau(r)^{\sharp}(\xi_{2}))(\xi_{1})), \xi_{3} \rangle = \sum_{i,j} \langle x_{j}y_{i} \otimes y_{j} \otimes x_{i}, \xi_{1} \otimes \xi_{2} \otimes \xi_{3} \rangle,
$$

$$
\langle \tau(r)^{\sharp}(\mathbf{r}_{A}^{*}(\tau(r)^{\sharp}(\xi_{2}))(\xi_{1})), \xi_{3} \rangle = \sum_{i,j} \langle y_{i}x_{j} \otimes y_{j} \otimes x_{i}, \xi_{1} \otimes \xi_{2} \otimes \xi_{3} \rangle.
$$

Hence, we have

$$
\langle \tau(r)^{\sharp}(\xi_{1})\tau(r)^{\sharp}(\xi_{2}) - \tau(r)^{\sharp}(\mathbf{I}_{A}^{*}(r^{\sharp}(\xi_{1}))(\xi_{2}) + (\mathbf{I}_{A}^{*} - \mathbf{r}_{A}^{*})(\tau(r)^{\sharp}(\xi_{2}))(\xi_{1})), \xi_{3} \rangle
$$

=
$$
\sum_{i,j} \langle y_{i} \otimes y_{j} \otimes x_{i}x_{j} - x_{j} \otimes y_{j}y_{i} \otimes x_{i} - x_{j}y_{i} \otimes y_{j} \otimes x_{i} + y_{i}x_{j} \otimes y_{j} \otimes x_{i}, \xi_{1} \otimes \xi_{2} \otimes \xi_{3} \rangle,
$$

=
$$
\sum_{i,j} \langle (\mathrm{id} \otimes \tau)((\tau \otimes \mathrm{id})([\![r,r]\!])) , \xi_{1} \otimes \xi_{2} \otimes \xi_{3} \rangle.
$$

That is to say, this proposition is hold. \square

If $r \in A \otimes A$ is symmetric, i.e., $r = \tau(r)$, then we have

Corollary 3.13. *Let* (*A*, ·) *be an anti-Leibniz algebra and r* ∈ *A*⊗*A be symmetric. Then r is a solution of the* aLYBE *in* (*A*, ·) *if and only if r*[♯] *is a relative Rota-Baxter operator of* (*A*, ·) *associated to the coregular bimodule* (A[∗], **l**[∗]₄ A^*_{A} , **l**_A^{*} – **r**_A^{*} *A*)*.*

Let (A, \cdot, \mathcal{B}) be an anti-Leibniz algebra with a nondegenerate skew-symmetric invariant bilinear form. Then under the natural bijection $Hom(A \otimes A, \mathbb{k}) \cong Hom(A, A^*)$, the bilinear form $\mathfrak{B}(-, -)$ corresponds to a linear map $\varphi : A \to A^*$, which is given by $\langle \varphi(a_1), a_2 \rangle = \mathfrak{B}(a_1, a_2)$, for any $a_1, a_2 \in A$. Define a 2-tensor $r_{\mathfrak{B}}$ to be the tensor form of φ^{-1} , i.e., $\langle \xi_1 \otimes \xi_2, r_{\mathfrak{B}} \rangle = \langle \varphi^{-1}(\xi_1), \xi_2 \rangle$. Then, by direct calculations, we have

Lemma 3.14. *Let* (A, \cdot, \mathfrak{B}) *be an anti-Leibniz algebra with a nondegenerate bilinear form. Then* $\mathfrak{B}(-, -)$ *is skew-symmetric invariant if and only if r_B <i>is skew-symmetric invariant.*

For any $r \in A \otimes A$, define a linear map $R_r : A \to A$, $a \mapsto R_r(a) := r^{\sharp}(\varphi(a))$, then we have the following proposition and corollary.

Proposition 3.15. *Let* (*A*, ·, B) *be an anti-Leibniz algebra with a nondegenerate skew-symmetric invariant bilinear form and* $R : A^* \to A$ *be a linear map. Then* R is a relative Rota-Baxter operator of (A, \cdot) associated to (A[∗], l[∗]_A</sub> A^*_{A} , **l**_A^{*} – **r**_A^{*} A_A^*) *if and only if* $R_r: A \to A$ *is a Rota-Baxter operator on* (A, \cdot) *.*

Proof. For any $a_1, a_2 \in A$, by Lemma [2.8,](#page-4-2) we have $R_r(a_1)R_r(a_2) = r^{\sharp}(\varphi(a_1))r^{\sharp}(\varphi(a_2))$ and

$$
R_r(a_1R_r(a_2) + R_r(a_1)a_2) = r^{\sharp}(\varphi(a_1r^{\sharp}(\varphi(a_2)) + r^{\sharp}(\varphi(a_1))a_2))
$$

= $r^{\sharp}(\mathbf{r}_A^*(r^{\sharp}(\varphi(a_1)))(\varphi(a_2)) + (\mathbf{l}_A^* - \mathbf{r}_A^*)(r^{\sharp}(\varphi(a_2)))(\varphi(a_1))).$

Thus, *R* is a relative Rota-Baxter operator if and only if R_r is a Rota-Baxter operator.

As a direct conclusion of Corollary [3.13](#page-12-0) and Proposition [3.15,](#page-12-1) we have

Corollary 3.16. *Let* (*A*, ·, B) *be an anti-Leibniz algebra with a nondegenerate skew-symmetric invariant bilinear form and r* \in *A* \otimes *A be symmetric. Then, r is a solution of the aLYBE in* (A, \cdot) *if and only if* R_r *is a Rota-Baxter operator on* (A, \cdot) .

More generally, if $r \in A \otimes A$ such that $r - \tau(r)$ is invariant, we consider when r is a solution of the aLYBE in (A, \cdot) .

Proposition 3.17. *Let* (A, \cdot) *be an anti-Leibniz algebra. Suppose* $r \in A \otimes A$ *such that* $r - \tau(r)$ *is invariant. Then, r is a solution of the aLYBE in* (A, \cdot) *if and only if for any* $\xi_1, \xi_2 \in A^*$,

$$
r^{\sharp}(\xi_1)r^{\sharp}(\xi_2) = r^{\sharp}\big(\mathbf{l}_A^*(r^{\sharp}(\xi_1))(\xi_2) + (\mathbf{l}_A^* - \mathbf{r}_A^*)(r^{\sharp}(\xi_2))(\xi_1) - \mathbf{l}_A^*((r^{\sharp} - \tau(r)^{\sharp})(\xi_1))(\xi_2)\big),
$$

Proof. First, since $r - \tau(r)$ is skew-symmetric, it is easy to see that $r - \tau(r)$ is invariant if and only if l[∗] $A^*_{A}((r^{\sharp} - \tau(r)^{\sharp})(\xi_1))(\xi_2) = (\mathbf{I}_A^* - \mathbf{r}_A^*)$ χ_A^*)($(r^{\sharp} - \tau(r)^{\sharp})(\xi_2)$)(ξ_1), for any $\xi_1, \xi_2 \in A^*$. Thus, by direct calculations, we have

$$
r^{\sharp}\left(\mathbf{l}_{A}^{*}(r^{\sharp}(\xi_{1}))(\xi_{2})+(\mathbf{l}_{A}^{*}-\mathbf{r}_{A}^{*})(r^{\sharp}(\xi_{2}))(\xi_{1})-\mathbf{l}_{A}^{*}((r^{\sharp}-\tau(r)^{\sharp})(\xi_{1}))(\xi_{2})\right) =r^{\sharp}\left(\mathbf{l}_{A}^{*}(r^{\sharp}(\xi_{1}))(\xi_{2})+(\mathbf{l}_{A}^{*}-\mathbf{r}_{A}^{*})(r^{\sharp}(\xi_{2}))(\xi_{1})-(\mathbf{l}_{A}^{*}-\mathbf{r}_{A}^{*})((r^{\sharp}-\tau(r)^{\sharp})(\xi_{2}))(\xi_{1})\right) =r^{\sharp}\left(\mathbf{l}_{A}^{*}(r^{\sharp}(\xi_{1}))(\xi_{2})+(\mathbf{l}_{A}^{*}-\mathbf{r}_{A}^{*})(\tau(r)^{\sharp}(\xi_{2}))(\xi_{1})\right).
$$

Second, since $r - \tau(r)$ is invariant, by Proposition [3.12,](#page-11-1) *r* is a solution of the aLYBE in (A, \cdot) if and only if $r^{\sharp}(\xi_1)r^{\sharp}(\xi_2) = r^{\sharp}(\mathbf{l}_{A}^*)$ $A^*(r^{\sharp}(\xi_1))(\xi_2) + (\mathbf{l}_A^* - \mathbf{r}_A^*)$ \int_A^*)($\tau(r)^\sharp(\xi_2)$)(ξ_1)). Thus, we get this proposition.

Let (A, \cdot) be an anti-Leibniz algebra. By the Definition [3.3,](#page-9-3) it is easy to see that $r \in A \otimes A$ is invariant if and only if $r^{\sharp}(\xi)a = r^{\sharp}((\mathbf{l}_{A}^{*} - \mathbf{r}_{A}^{*}))$ $A^*(A)(a)(\xi)$ for any $a \in A$ and $\xi \in A^*$.

Corollary 3.18. *Let* (*A*, ·, B) *be an anti-Leibniz algebra with a nondegenerate skew-symmetric invariant bilinear form and r* ∈ *A* ⊗ *A be invariant. Then, r is a solution of the* aLYBE *in* (*A*, ·) *if and only if for any* $a_1, a_2 \in A$,

$$
R_r(a_1)R_r(a_1) = R_r(R_r(a_1)a_2 + a_1R_r(a_2) - a_1R_r(a_2) + a_1\tau(r)^{\sharp}(\varphi(a_2))
$$

where R_r : $A \to A$ *is defined by* $R_r(a) = r^{\sharp}(\varphi(a))$ *.*

Proof. Since $r_{\mathcal{B}}$ is skew-symmetric invariant in this case, we get $\varphi(r^{\sharp}(\xi_1)\varphi^{-1}(\xi_2)) = \mathbf{I}_{A}^{*}$ $A^*(r^{\sharp}(\xi_1))(\xi_2)$ and $\varphi(\varphi^{-1}(\xi_1)r^{\sharp}(\xi_2)) = (\mathbf{l}_A^* - \mathbf{r}_A^*)$ A_{A}^{*})($r^{\sharp}(\xi_2)$)(ξ_1). For any $a_1, a_2 \in A$, suppose $a_1 = \varphi^{-1}(\xi_1)$ and $a_2 = \varphi^{-1}(\xi_2)$, $\xi_1, \xi_2 \in A^*$. Then, we have

$$
R_r(a_1)R_r(a_1) = r^{\sharp}(\xi_1)r^{\sharp}(\xi_2),
$$

\n
$$
R_r(a_1R_r(a_2)) = r^{\sharp}(\mathbf{1}_A^*(r^{\sharp}(\xi_1))(\xi_2)),
$$

\n
$$
R_r(R_r(a_1)a_2) = r^{\sharp}((\mathbf{1}_A^* - \mathbf{r}_A^*)(r^{\sharp}(\xi_2))(\xi_1)),
$$

\n
$$
R_r(a_1\tau(r)^{\sharp}(\varphi(a_2)) - a_1R_r(a_2)) = r^{\sharp}(\mathbf{1}_A^*((\tau(r)^{\sharp} - r^{\sharp})(\xi_1))(\xi_2)).
$$

By Proposition [3.17,](#page-13-0) we get the conclusion. \square

In the Lie algebra context, it is well-known that the dual description of a classical *r*-matrix is a symplectic structure on a Lie algebra. For the anti-Leibniz algebras, we have

Proposition 3.19. *Let* (A, \cdot) *be an anti-Leibniz algebra and* $r \in A \otimes A$ *. Suppose that r is symmetric and nondegenerate. The linear isomorphism r*[♯] : *A* [∗] → *A define a bilinear form* ω(−, −) *on* (*A*, ·) *by* $\omega(a_1, a_2) = \langle (r^{\sharp})^{-1}(a_1), a_2 \rangle$ for any $a_1, a_2 \in A$. Then r is a solution of the aLYBE in (A, \cdot) if and only if *the bilinear form* $\omega(-, -)$ *satisfies*

(3.7)
$$
\omega(a_2a_3, a_1) + \omega(a_1a_3, a_2) - \omega(a_3a_1, a_2) - \omega(a_2a_1, a_3) = 0,
$$

for any $a_1, a_2, a_3 \in A$.

Proof. Let $r = \sum_i x_i \otimes y_i \in A \otimes A$ be symmetric. Then $r^{\sharp}(\xi) = \sum_i \langle \xi, x_i \rangle y_i = \sum_i \langle \xi, y_i \rangle x_i$ for any $\xi \in A^*$ and the bilinear form $\omega(-, -)$ is symmetric, i.e., $\omega(a_1, a_2) = \omega(a_2, a_1)$ for any $a_1, a_2 \in A$. Since *r* is nondegenerate, for any $a_1, a_2, a_3 \in A$, there are $\xi_1, \xi_2, \xi_3 \in A^*$ such that $r^{\sharp}(\xi_i) = a_i$ for $i = 1, 2, 3$. Thus, we have

$$
\omega(a_2a_3, a_1) + \omega(a_1a_3, a_2) - \omega(a_3a_1, a_2) - \omega(a_2a_1, a_3)
$$

= $\langle r^{\sharp}(\xi_2)r^{\sharp}(\xi_3), \xi_1 \rangle + \langle r^{\sharp}(\xi_1)r^{\sharp}(\xi_3), \xi_2 \rangle - \langle r^{\sharp}(\xi_3)r^{\sharp}(\xi_1), \xi_2 \rangle - \langle r^{\sharp}(\xi_2)r^{\sharp}(\xi_1), \xi_3 \rangle$
= $\sum_{i,j} (\langle \xi_1, x_i x_j \rangle \langle \xi_2, y_i \rangle \langle \xi_3, y_j \rangle + \langle \xi_1, x_i \rangle \langle \xi_2, y_i x_j \rangle \langle \xi_3, y_j \rangle - \langle \xi_1, x_j \rangle \langle \xi_2, x_i \rangle \langle \xi_3, y_i y_j \rangle)$
= $\langle \xi_1 \otimes \xi_2 \otimes \xi_3, r_{12}r_{13} + r_{12}r_{23} - r_{23}r_{12} - r_{23}r_{13} \rangle.$

That is, $\omega(a_2a_3, a_1) + \omega(a_1a_3, a_2) - \omega(a_3a_1, a_2) - \omega(a_2a_1, a_3) = 0$ for any $a_1, a_2, a_3 \in A$ if and only if $\langle \xi_1 \otimes \xi_2 \otimes \xi_3, \quad [r, r] \rangle = 0$ for any $\xi_1, \xi_2, \xi_3 \in A^*$, if and only if $[[r, r]] = 0$. The proof is finished. \square

Next, we show that each relative Rota-Baxter operator induce an anti-Leibniz bialgebra.

Proposition 3.20. *Let* (*A*, ·) *be an anti-Leibniz algebra and* (*M*, l,r) *be a bimodule over* (*A*, ·)*. Let P* : \tilde{M} → *A* be a linear map which is identified as an element in $(A \ltimes M^*) \otimes (A \ltimes M^*)$ through Hom $(M, A) \cong$ $A \otimes M^* \subset (A \ltimes M^*) \otimes (A \ltimes M^*)$. Then $r := P + \tau(P)$ is a symmetric solution of the aLYBE in semi*direct product anti-Leibniz algebra* $A \times M^*$ *if and only if P is a relative Rota-Baxter operator of* (A, \cdot) *associated to* (*M*, l,r)*.*

Proof. Let $\{e_1, e_2, \dots, e_n\}$ be a basis of *M* and $\{f_1, f_2, \dots, f_n\}$ be its dual basis in *M*^{*}. Then *P* = $\sum_{i=1}^{n} P(e_i) \otimes f_i \in (A \ltimes M^*) \otimes (A \ltimes M^*)$, $r = P - \tau(P) = \sum_{i=1}^{n} (P(e_i) \otimes f_i + f_i \otimes P(e_i))$, and

$$
r_{12}r_{13} = \sum_{i,j=1}^{n} P(e_i)P(e_j) \otimes f_i \otimes f_j - I^*(P(e_i))(f_j) \otimes f_i \otimes P(e_j) - (I^* - r^*)(P(e_j))(f_i) \otimes P(e_i) \otimes f_j,
$$

\n
$$
r_{12}r_{23} = \sum_{i,j=1}^{n} P(e_i) \otimes (I^* - r^*)(P(e_j))(f_i) \otimes f_j - f_i \otimes P(e_i)P(e_j) \otimes f_j + f_i \otimes I^*(P(e_i))(f_j) \otimes P(e_j),
$$

\n
$$
r_{23}r_{12} = \sum_{i,j=1}^{n} P(e_j) \otimes I^*(P(e_i))(f_j) \otimes f_i - f_j \otimes P(e_i)P(e_j) \otimes f_i + f_j \otimes (I^* - r^*)(P(e_j))(f_i) \otimes P(e_i),
$$

\n
$$
r_{23}r_{13} = \sum_{i,j=1}^{n} -f_j \otimes P(e_i) \otimes (I^* - r^*)(P(e_j))(f_i) - P(e_j) \otimes f_i \otimes I^*(P(e_i))(f_j) + f_i \otimes f_j \otimes P(e_i)P(e_j).
$$

Note that $I^*(P(e_i))(f_j) = \sum_{k=1}^{r} f_j(I(P(e_i))(e_k))f_k$, we get

$$
\sum_{i,j=1}^n P(e_j) \otimes I^*(P(e_i))(f_j) \otimes f_i = \sum_{i,j,k=1}^n f_j(I(P(e_i))(e_k))P(e_j) \otimes f_k \otimes f_i
$$

=
$$
\sum_{i,j=1}^n P\Big(\sum_{k=1}^n f_k(I(P(e_i))(e_j))e_k\Big) \otimes f_j \otimes f_i = \sum_{i,j=1}^n P(I(P(e_i))(e_j)) \otimes f_j \otimes f_i.
$$

Similarly, we also have $\sum_{i,j=1}^n f_j \otimes I^*(P(e_j))(f_i) \otimes P(e_i) = f_j \otimes f_i \otimes P(I(P(e_i))(e_j)), \sum_{i,j=1}^n P(e_i) \otimes I^*(P(e_j))(f_i) \otimes$ $f_j = \sum_{i,j=1}^n P(\mathfrak{r}(P(e_j))(e_i)) \otimes f_i \otimes f_j$ and $\sum_{i,j=1}^n f_i \otimes \mathfrak{r}^*(P(e_i))(f_j) \otimes P(e_i) = f_i \otimes f_j \otimes P(\mathfrak{r}(P(e_j))(e_i)).$ Thus

$$
r_{12}r_{13} + r_{12}r_{23} - r_{23}r_{12} - r_{23}r_{13} = \sum_{i,j=1}^{n} \Big(\big(P(e_i)P(e_j) - P(\mathfrak{r}(P(e_j))(e_i) - P(\mathfrak{t}(P(e_i))(e_j)) \big) \otimes f_i \otimes f_j + f_i \otimes f_j \otimes \big(P(\mathfrak{t}(P(e_i))(e_j)) + P(\mathfrak{r}(P(e_j))(e_i)) - P(e_i)P(e_j) \big) \Big).
$$

Thus, *r* is a solution of the aLYBE in $A \ltimes M^*$ if and only if *P* is a relative Rota-Baxter operator of (A, \cdot) associated to (M, I, r) .

As a direct conclusion, we have

Corollary 3.21. Let (A, \cdot) be an anti-Leibniz algebra and (M, I, r) be a bimodule over (A, \cdot) . If $P : M \to$ *A is a relative Rota-Baxter operator of* (*A*, ·) *associated to* (*M*, l,r)*, then there is an anti-Leibniz algebra* $(A \ltimes M^*, \Delta_r)$, where Δ_r is given by Eq. [\(3.1\)](#page-6-2) for $r = P + \tau(P)$.

3.3. Factorizable anti-Leibniz bialgebras and skew-quadratic Rota-Baxter anti-Leibniz algebras. In this subsection, we establish the factorizable theories for anti-Leibniz bialgebras. Let (A, \cdot) be an anti-Leibniz algebra and $r \in A \otimes A$. We have defined a linear map $r^{\sharp}: A^* \to A$ by

$$
\langle r^{\sharp}(\xi_1), \xi_2 \rangle = \langle \xi_1 \otimes \xi_2, r \rangle.
$$

for any $\xi_1, \xi_2 \in A^*$. If (A, \cdot, Δ_r) is an anti-Leibniz bialgebra, then the anti-Leibniz algebra structure \cdot_r on *A*^{*} dual to the comultiplication Δ_r defined by Eq. [\(3.1\)](#page-6-2) is given by

$$
\xi_1 \cdot_r \xi_2 = \mathbf{I}_A^* (r^{\sharp}(\xi_1))(\xi_2) + \mathbf{I}_A^* (\tau(r)^{\sharp}(\xi_2))(\xi_1) - \mathbf{r}_A^* (\tau(r)^{\sharp}(\xi_2))(\xi_1),
$$

for any $\xi_1, \xi_2 \in A^*$. Let (A, \cdot) be an anti-Leibniz algebra. In Subsection [3.2,](#page-11-0) we have get that an element $r \in A \otimes A$ is invariant if and only if $r^{\sharp}(\xi)a = r^{\sharp}((\mathbf{l}_{A}^{*} - \mathbf{r}_{A}^{*}))$ $A^*(A)(a)(\xi)$ for any $a \in A$ and $\xi \in A^*$.

Lemma 3.22. *Let* (A, \cdot) *be an anti-Leibniz algebra and* $r \in A \otimes A$ *. Denote by* $I = r^{\sharp} - \tau(r)^{\sharp} : A^* \to A$ *. Then* $r - \tau(r)$ *is invariant if and only if* $I(\mathbf{l}_A^* - \mathbf{r}_A^*)$ \mathbf{F}_{A}^{*} (a) = $\mathbf{r}_{A}(a)$ *I* or $I\mathbf{r}_{A}^{*}$ $A^*_{A}(a) = (\dot{I}^*_{A} - \dot{r}^*_{A})$ $_A^*(A)(a)$ *I*, for any $a \in A$.

Proof. It is easy to see that $\mathcal{I}^* = -\mathcal{I}$ if we identify $(A^*)^*$ with *A*, and $\langle \mathcal{I}(\xi), \eta \rangle = (r - \tau(r))(\xi_1 \otimes \xi_2)$ for any $\xi_1, \xi_2 \in A^*$. Since $r - \tau(r)$ is invariant if and only if $\mathbf{r}_A(a)((r^{\sharp} - \tau(r)^{\sharp})(\xi)) = (r^{\sharp} - \tau(r)^{\sharp})((\mathbf{l}_A^* - \mathbf{r}_A^*)^*$ $_{A}^{*})$ $(a)(\xi)$ and $\overline{I}^* = -\overline{I}$ if we identify $(A^*)^*$ with *A*, by the dual, we get $\overline{I}(\mathbf{l}_A^* - \mathbf{r}_A^*)$ $\mathbf{A}^*(A)$ = $\mathbf{r}_A(a)$ is equivalent to $\bm{\mathcal{I}} \, \mathbf{r}^*_A$ $A_A^*(a) = (\mathbf{l}_A - \mathbf{r}_A)(a)$ for any $a \in A$. The proof is finished.

Proposition 3.23. *Let* (A, \cdot) *be an anti-Leibniz algebra and* $r \in A \otimes A$ *such that* $r - \tau(r)$ *is invariant. The following are equivalent.*

- (*i*) r *is a solution of the aLYBE in* (A, \cdot) *;*
- (*ii*) (A^*, \cdot_r) *is an anti-Leibniz algebra and* r^{\sharp} : $(A^*, \cdot_r) \to (A, \cdot)$ *is a homomorphism of anti-Leibniz algebras;*
- (*iii*) (A^*, \cdot_r) *is an anti-Leibniz algebra and* $\tau(r)^{\sharp} : (A^*, \cdot_r) \to (A, \cdot)$ *is a homomorphism of anti-Leibniz algebras.*

Proof. (*i*) \Leftrightarrow (*ii*). If *r* is a solution of the aLYBE in (A, \cdot) , then (A, \cdot, Δ_r) is a quasi-triangular anti-Leibniz bialgebra, and so that (A^*, \cdot_r) is an anti-Leibniz algebra. Since the product \cdot_r is given by $\xi_1 \cdot_r \xi_2 =$ l[∗] $A^*_{A}(r^{\bar{\sharp}}(\xi_1))(\xi_2) + \mathbf{l}_A^*$ $A^*_{A}(\tau(r)^{\sharp}(\xi_2))(\xi_1) - \mathbf{r}^*_{A}$ $A^*(\tau(r)^\sharp(\xi_2))(\xi_1)$, this equivalence is following Proposition [3.12.](#page-11-1)

(*i*) ⇔ (*iii*). By a direct calculation, for any $\xi_1, \xi_2, \xi_3 \in A^*$, we have

$$
\langle r^{\sharp}(\xi_1)r^{\sharp}(\xi_2), \xi_3 \rangle = \sum_{i,j} \langle x_j \otimes x_i \otimes y_j y_i, \xi_1 \otimes \xi_2 \otimes \xi_3 \rangle,
$$

$$
\langle r^{\sharp}(\mathbf{r}_A^*(\tau(r)^{\sharp}(\xi_2))(\xi_1)), \xi_3 \rangle = \sum_{i,j} \langle x_i x_j \otimes y_j \otimes y_i, \xi_1 \otimes \xi_2 \otimes \xi_3 \rangle,
$$

$$
\langle r^{\sharp}(\mathbf{l}_A^*(\tau(r)^{\sharp}(\xi_2))(\xi_1)), \xi_3 \rangle = \sum_{i,j} \langle x_j x_i \otimes y_j \otimes y_i, \xi_1 \otimes \xi_2 \otimes \xi_3 \rangle,
$$

$$
\langle r^{\sharp}(\mathbf{l}_A^*(r^{\sharp}(\xi_1))(\xi_2)), \xi_3 \rangle = \sum_{i,j} \langle x_j \otimes y_j x_i \otimes y_i, \xi_1 \otimes \xi_2 \otimes \xi_3 \rangle.
$$

Hence,

$$
\langle r^{\sharp}(\xi_{1} \cdot_{r} \xi_{2}) - r^{\sharp}(\xi_{1})r^{\sharp}(\xi_{2}), \xi_{3} \rangle
$$

= $\langle r^{\sharp}(\mathbf{1}_{A}^{*}(\tau(r)^{\sharp}(\xi_{2}))(\xi_{1})) + r^{\sharp}(\mathbf{1}_{A}^{*}(r^{\sharp}(\xi_{1}))(\xi_{2})) - r^{\sharp}(\mathbf{1}_{A}^{*}(\tau(r)^{\sharp}(\xi_{2}))(\xi_{1})) - r^{\sharp}(\xi_{1})r^{\sharp}(\xi_{2}), \xi_{3} \rangle$
=
$$
\sum_{i,j} \langle x_{j}x_{i} \otimes y_{j} \otimes y_{i} + x_{j} \otimes y_{j}x_{i} \otimes y_{i} - x_{i}x_{j} \otimes y_{j} \otimes y_{i} - x_{j} \otimes x_{i} \otimes y_{j}y_{i}, \xi_{1} \otimes \xi_{2} \otimes \xi_{3} \rangle
$$

= $\langle ((\mathbf{r}_{A} - \mathbf{I}_{A})(x_{j}) \otimes id + id \otimes \mathbf{r}_{A}(x_{j}))(r - \tau(r)) \otimes y_{j} + (\tau \otimes id)([\![r, r]\!]), \xi_{1} \otimes \xi_{2} \otimes \xi_{3} \rangle.$

That is to say, if $r - \tau(r)$ is invariant and *r* is a solution of the aLYBE in (A, \cdot) , then r^{\sharp} is a homomorphism of anti-Leibniz algebras. Conversely, it is directly available from the above calculation.

Triangular anti-Leibniz bialgebras is an important class of anti-Leibniz bialgebra. Another important class of anti-Leibniz bialgebra is the factorizable anti-Leibniz bialgebras.

Definition 3.24. *A quasi-triangular anti-Leibniz bialgebra* (*A*, ·, ∆) *is called* factorizable *if the linear map* $I = r^{\sharp} - \tau(r)^{\sharp} : A^* \to A$ *is a linear isomorphism of vector spaces.*

Clearly, quasi-triangular anti-Leibniz bialgebras contain triangular anti-Leibniz bialgebras and factorizable anti-Leibniz bialgebras as two subclasses. Moreover, it is easy to see that a quasi-triangular anti-Leibniz bialgebra (A, \cdot, Δ) is a triangular anti-Leibniz bialgebra if and only if $\mathcal{I} = r^{\sharp} - \tau(r)^{\sharp}: A^* \to A$ is the zero map. Factorizable anti-Leibniz bialgebras are however concerned with the opposite case. Following, for convenience, we consider the linear map $\mathcal{I} = r^{\sharp} - \tau(r)^{\sharp} : A^* \to A$ as a composition of maps as follows:

 $A^* \xrightarrow{r^{\sharp} \oplus \tau(r)^{\sharp}} A \oplus A \xrightarrow{(a_1, a_2) \mapsto a_1 - a_2} A.$

The following result justifies the terminology of a factorizable anti-Leibniz bialgebra.

Proposition 3.25. *Let* (A, \cdot) *be an anti-Leibniz algebra and* $r \in A \otimes A$ *. If there is an anti-Leibniz bialgebra* (A, ⋅, Δ_{*r*})</sub> *induced by r and it is factorizable, then* Im(r [#] ⊕ τ(r)[#]) *is an anti-Leibniz subalgebra of the direct sum anti-Leibniz algebra* $A \oplus A$, which is isomorphism to the anti-Leibniz algebra (A^*, \cdot_r) . *Moreover, any a* \in *A has a unique decomposition a* = $a_+ + a_-,$ where $a_+ \in \text{Im}(r^{\sharp})$ *and* $a_- \in \text{Im}(r(r)^{\sharp}).$

Proof. Since (A, \cdot, Δ_r) is quasi-triangular anti-Leibniz bialgebra, both r^{\sharp} and $\tau(r)^{\sharp}$ are homomorphisms of anti-Leibniz algebras. Therefore, $\text{Im}(r^{\sharp} \oplus \tau(r)^{\sharp})$ is an anti-Leibniz subalgebra of the direct sum anti-Leibniz algebra $A \oplus A$. Since $I : A^* \to A$ is a linear isomorphism, it follows that $r^{\sharp} \oplus \tau(r)^{\sharp}$ is injective, then the anti-Leibniz algebra $\text{Im}(r^{\sharp} \oplus \tau(r)^{\sharp})$ is isomorphic to the anti-Leibniz algebra (A^*, \cdot_r) . Moreover, since \overline{I} is an isomorphism again, for any $a \in A$, we have

$$
a = (r^{\sharp} - \tau(r)^{\sharp})(\mathcal{I}^{-1}(a)) = r^{\sharp}(\mathcal{I}^{-1}(a)) - \tau(r)^{\sharp}(\mathcal{I}^{-1}(a)),
$$

which implies that $a = a_+ + a_-$, where $a_+ = r^{\sharp}(\mathcal{I}^{-1}(a))$ and $a_- = -\tau(r)^{\sharp}(\mathcal{I}^{-1}(a))$. The uniqueness also follows from the fact that I is an isomorphism.

In Theorem [3.9,](#page-10-1) for any anti-Leibniz bialgebra (A, \cdot, Δ) , we construct a new anti-Leibniz bialgebra $(A \oplus A^*, *, \Delta_{\tilde{r}})$, which is called the double of anti-Leibniz bialgebra (A, \cdot, Δ) . The comultiplication $\Delta_{\tilde{r}}$ is given by Eq. [\(3.1\)](#page-6-2) for $r = \sum_i e_i \otimes f_i$, where $\{e_1, e_2, \dots, e_n\}$ is a basis of A and $\{f_1, f_2, \dots, f_n\}$ be its dual basis. The importance of factorizable anti-Leibniz bialgebras in the study of anti-Leibniz bialgebras can also be observed from the following proposition that the double space of an arbitrary anti-Leibniz bialgebra admits a factorizable anti-Leibniz bialgebra structure.

Proposition 3.26. With the above notations, the double anti-Leibniz bialgebra $(A \Join A^*, *, \Delta_{\tilde{r}})$ of any *anti-Leibniz bialgebra* (*A*, ·, ∆) *is factorizable.*

Proof. Let $\tilde{r} = \sum_i e_i \otimes f_i$, where $\{e_1, e_2, \dots, e_n\}$ is a basis of A and $\{f_1, f_2, \dots, f_n\}$ be its dual basis in A^* . Since $\tilde{r} - \tau(\tilde{r}) = \sum_{i=1}^{n} (e_i \otimes f_i - f_i \otimes e_i)$, for any $(\xi, a) \in A^* \oplus A$, we get $\tilde{I}(\xi, a) = (-a, \xi) \in A \oplus A^*$, where $\tilde{\mathcal{I}} = (\tilde{r}^{\sharp} - \tau(\tilde{r})^{\sharp})$. Thus, for any $(a_1, \xi_1), (a_2, \xi_2) \in A \oplus A^*$,

$$
\mathbf{r}_{A \bowtie A^*}^*(a_1, \xi_1)(\tilde{I}(\xi_2, a_2))
$$
\n= $(-a_2a_1 + \mathbf{I}_{A^*}^*(\xi_2)(a_1) + (\mathbf{I}_{A^*}^* - \mathbf{r}_{A^*}^*)(\xi_1)(-a_2), \xi_2 \cdot_{A^*} \xi_1 + \mathbf{I}_{A}^*(-a_2)(\xi_1) + (\mathbf{I}_{A}^* - \mathbf{r}_{A}^*)(a_1)(\xi_2))$
\n= $\tilde{I}(\xi_2 \cdot_{A^*} \xi_1 + \mathbf{I}_{A}^*(-a_2)(\xi_1) + (\mathbf{I}_{A}^* - \mathbf{r}_{A}^*)(a_1)(\xi_2), a_2a_1 - \mathbf{I}_{A^*}^*(\xi_2)(a_1) + (\mathbf{r}_{A^*}^* - \mathbf{I}_{A^*}^*)(\xi_1)(-a_2))$
\n= $\tilde{I}(((\mathbf{I}_{A \bowtie A^*}^* - \mathbf{r}_{A \bowtie A^*}^*)(a_1, \xi_1))(\xi_2, a_2)).$

By Lemma [3.22,](#page-15-1) we get $\tilde{r} - \tau(\tilde{r})$ is invariant. Thus, $(A \Join A^*, \Lambda_{\tilde{r}})$ is a quasi-triangular anti-Leibniz bialgebra. Finally, since $I(\xi, a) = (-a, \xi)$ is a linear isomorphism, we get that $(A \Join A^*, \times, \Delta_{\tilde{r}})$ is a factorizable anti-Leibniz bialgebra.

Proposition [3.26](#page-16-0) provides a method for constructing factorizable anti-Leibniz bialgebras.

Example 3.27. *Consider the* 2*-dimensional anti-Leibniz bialgebra* (*A*, ·, ∆) *given by Example [3.6,](#page-10-2) where A* = $\mathbb{k}\{e_1, e_2\}$, the nonzero multiplication and comultiplication: $e_1e_1 = e_2$ and $\Delta(e_1) = e_2 \otimes e_2$. Denote by $\{f_1, f_2\}$ the dual basis of $\{e_1, e_2\}$ and $\tilde{r} = e_1 \otimes f_1 + e_2 \otimes f_2 \in A \otimes A^* \subset (A \oplus A^*) \otimes (A \oplus A^*)$. Then we *get a* 4*-dimensional anti-Leibniz bialgebra* (*A* ⊕ *A* ∗ , ∗, ∆*^r*˜)*, where*

$$
e_1 * e_1 = e_2,
$$
 $f_2 * f_2 = f_1,$ $e_1 * f_2 = f_1,$ $f_2 * e_1 = e_2,$
 $\Delta_{\tilde{r}}(e_1) = e_2 \otimes e_2,$ $\Delta_{\tilde{r}}(f_2) = f_1 \otimes f_1.$

One can check $(A \oplus A^*, *, \Delta_{\tilde{r}})$ *be a quasi-triangular anti-Leibniz bialgebra. Note that* $r - \tau(r) = e_1 \otimes f_1 +$ $e_2 \otimes f_2 - f_1 \otimes e_1 - f_2 \otimes e_2$, we get that the linear map \tilde{I} : $(A \oplus A^*)^* \to A \oplus A^*$ is given by

$$
\tilde{I}(e_1^{\star}) = f_1, \qquad \tilde{I}(e_2^{\star}) = f_2, \qquad \tilde{I}(f_1^{\star}) = -e_1, \qquad \tilde{I}(f_2^{\star}) = -e_2,
$$

where $\{e_1^{\star}\}$ $\boldsymbol{\dot{e}}_1^{\star}, e_2^{\star}$ $\frac{\star}{2}$, f_1^{\star} f_1^{\star}, f_2^{\star} $\{x^*\}\$ *in* $(A\oplus A^*)^*$ *is the dual basis of* $\{e_1, e_2, f_1, f_2\}$ *in* $A\oplus A^*$ *. Hence, we get* \tilde{I} *is a linear isomorphism, and so that* $(A \oplus A^*, *, \Delta_{\tilde{r}})$ *is factorizable anti-Leibniz bialgebra.*

Next, we will give a characterization of factorizable anti-Leibniz bialgebras by the Rota-Baxter operator on anti-Leibniz algebras with a nondegenerate skew-symmetric invariant bilinear form. Let (*A*, ·) be an anti-Leibniz algebra and $\lambda \in \mathbb{k}$. A linear map $R : A \rightarrow A$ is called a *Rota-Baxter operator of weight* λ on (A, \cdot) if

$$
R(a_1)R(a_2) = R(R(a_1)a_2 + a_1R(a_2) + \lambda a_1a_2),
$$

for any $a_1, a_2 \in A$. Let (A, \cdot, R) be an anti-Leibniz algebra with a Rota-Baxter operator R of weight λ . Then there is a new multiplication \cdot_R on *A* defined by

$$
a_1 \cdot_R a_2 = R(a_1)a_2 + a_1R(a_2) + \lambda a_1 a_2,
$$

for any $a_1, a_2 \in A$. This anti-Leibniz algebra is denoted by A_R . It is easy to see that *R* is a homomorphism of anti-Leibniz algebras from (A, \cdot_R) to (A, \cdot) .

Definition 3.28. A linear map $R : A \rightarrow A$ is called a Rota-Baxter operator of weight λ on an anti-Leibniz algebra (A, \cdot) with a nondegenerate skew-symmetric invariant bilinear form $\mathfrak{B}(-, -)$ *if R is a Rota-Baxter operator of weight* λ *on* (A, \cdot) *and for any* $a_1, a_2 \in A$,

$$
\mathfrak{B}(R(a_1), a_2) + \mathfrak{B}(a_1, R(a_2)) + \lambda \mathfrak{B}(a_1, a_2) = 0.
$$

The quadruple (*A*, ·, **B**, *R*) *is called a* skew-quadratic Rota-Baxter anti-Leibniz algebra of weight λ.

By direct calculations, we have

Proposition 3.29. *Let* (*A*, ·) *be an anti-Leibniz algebra,* B(−, −) *be a nondegenerate skew-symmetric invariant bilinear form on* (A, \cdot) *and* $R : A \rightarrow A$ *is a linear map. Then,* $(A, \cdot, \mathcal{B}, R)$ *is a skew-quadratic Rota-Baxter anti-Leibniz algebra of weight* λ *if and only if* (*A*, ·, −B, −(λ id+*R*)) *is a skew-quadratic Rota-Baxter anti-Leibniz algebra of weight* λ*.*

The following theorem shows that there is a one-to-one correspondence between factorizable anti-Leibniz bialgebras and skew-quadratic Rota-Baxter anti-Leibniz algebras of nonzero weight.

Theorem 3.30. Let (A, \cdot) be an anti-Leibniz algebra and $r \in A \otimes A$. Denote $I = r^{\sharp} - \tau(r)^{\sharp}$ and define *a bilinear form* \mathfrak{B}_I *by* $\mathfrak{B}_I(a_1, a_2) = \langle I^{-1}(a_1), a_2 \rangle$, *for any* $a_1, a_2 \in A$. *If there is an anti-Leibniz bialgebra* (A, \cdot, Δ) *induced by r and it is factorizable, then* $(A, \cdot, \mathfrak{B}_I, R = \lambda \tau(r)^{\sharp} \mathcal{I}^{-1}$ *) is a skew-quadratic Rota-Baxter anti-Leibniz algebras of weight* λ*.*

Conversely, for any skew-quadratic Rota-Baxter anti-Leibniz algebra (*A*, ·, B, *R*) *of weight* λ*, we have* a linear isomorphism $I_{\mathfrak{B}}: A^* \to A$ by $\langle T_{\mathfrak{B}}^{-1}(a_1), a_2 \rangle = \mathfrak{B}(a_1, a_2)$, for any $a_1, a_2 \in A$. If $\lambda \neq 0$, we define

$$
r^{\sharp} := \frac{1}{\lambda}(R + \lambda \operatorname{id}) \mathcal{I}_{\mathfrak{B}} : A^* \longrightarrow A,
$$

and define $r \in A \otimes A$ by $\langle r^{\sharp}(\xi_1), \xi_2 \rangle = \langle r, \xi_1 \otimes \xi_2 \rangle$, for any $\xi_1, \xi_2 \in A^*$. Then r is a solution of the aLYBE i *n*(A , \cdot) *and gives rise to a factorizable anti-Leibniz bialgebra* (A , \cdot , Δ _r), where Δ _r *is given by Eq.* [\(3.1\)](#page-6-2).

Proof. Since, r^{\sharp} , $\tau(r)^{\sharp}$: $(A^*, \cdot_r) \to (A, \cdot)$ are homomorphisms of anti-Leibniz algebras, for all $a_1, a_2, a_3 \in$ *A*, we have

$$
(3.8) \quad I(I^{-1}(a_1) \cdot_r I^{-1}(a_2)) = (r^{\sharp} - \tau(r)^{\sharp})(I^{-1}(a_1) \cdot_r I^{-1}(a_2))
$$

$$
= (I + \tau(r)^{\sharp})(I^{-1}(a_1))(I + \tau(r)^{\sharp})(I^{-1}(a_2)) - \tau(r)^{\sharp}(I^{-1}(a_1))\tau(r)^{\sharp}(I^{-1}(a_2))
$$

$$
= \tau(r)^{\sharp}(I^{-1}(a_1))a_2 + a_1\tau(r)^{\sharp}(I^{-1}(a_2)) + a_1a_2.
$$

Therefore, for any $a_1, a_2 \in A$,

$$
R(R(a_1)a_2 + a_1R(a_2) + \lambda a_1a_2) = \lambda^2 \tau(r)^{\sharp} \mathcal{I}^{-1}(\tau(r)^{\sharp} (\mathcal{I}^{-1}(a_1))a_2 + a_1 \tau(r)^{\sharp} (\mathcal{I}^{-1}(a_2)) + a_1a_2)
$$

= $\lambda^2 \tau(r)^{\sharp} (\mathcal{I}^{-1}(a_1) \cdot_r \mathcal{I}^{-1}(a_2))$
= $R(a_1)R(a_2)$,

which implies that *R* is a Rota-Baxter operator of weight λ on (A, \cdot) .

Next, we prove that $\mathfrak{B}_I(-, -)$ is nondegenerate skew-symmetric invariant. $\mathfrak{B}_I(-, -)$ is nondegenerate since \overline{I} is an isomorphism. Moreover, since $(I^{-1})^* = -I^{-1}$, we get

$$
\mathfrak{B}_{\mathcal{I}}(a_1,a_2)=\langle \mathcal{I}^{-1}(a_1), a_2\rangle=-\langle a_1, \mathcal{I}^{-1}(a_2)\rangle=-\mathfrak{B}_{\mathcal{I}}(a_2,a_1),
$$

for any $a_1, a_2 \in A$. Hence, \mathfrak{B}_I is skew-symmetric. Moreover, since the $r - \tau(r)$ is invariant, by Lemma [3.22,](#page-15-1) we have $I^{-1}r_A(a) = (l_A^* - r_A^*)$ $_A^{\ast}(a)$ I^{-1} . Thus,

$$
\mathfrak{B}_I(a_1, a_2a_3) - \mathfrak{B}_I(a_1, a_2a_3) = \langle \mathcal{I}^{-1}(\mathbf{r}_A(a_2)(a_1)) - (\mathbf{l}_A^* - \mathbf{r}_A^*)(a_2)(\mathcal{I}^{-1}(a_1))), a_3 \rangle = 0,
$$

for any $a_1, a_2, a_3 \in A$, which implies that \mathfrak{B}_I is invariant. Moreover, since $(\tau(r)^\sharp)^* = r^\sharp$, we get

$$
\mathfrak{B}_{I}(a_{1}, R(a_{2})) + \mathfrak{B}_{I}(R(a_{1}), a_{2}) + \lambda \mathfrak{B}_{I}(a_{1}, a_{2})
$$
\n
$$
= \lambda \langle I^{-1}(a_{1}), \tau(r)^{\sharp}(I^{-1}(a_{2})) \rangle + \lambda \langle I^{-1}(\tau(r)^{\sharp}(I^{-1}(a_{1}))), a_{2} \rangle + \lambda \langle I^{-1}(a_{1}), a_{2} \rangle
$$
\n
$$
= -\lambda \langle I^{-1}(a_{1}), a_{2} \rangle + \lambda \langle I^{-1}(a_{1}), a_{2} \rangle
$$
\n
$$
= 0,
$$

for $a_1, a_2 \in A$. Thus, $(A, \cdot, \mathfrak{B}_I, R)$ is a skew-quadratic Rota-Baxter anti-Leibniz algebras of weight λ .

Conversely, suppose that $(A, \cdot, \mathfrak{B}, R)$ is a skew-quadratic Rota-Baxter anti-Leibniz algebra of weight λ. First, since $\mathfrak{B}(-, -)$ is skew-symmetric and $\mathfrak{B}(a_1, R(a_2)) + \mathfrak{B}(R(a_1), a_2) + \lambda \mathfrak{B}(a_1, a_2) = 0$, we get $\overline{I}^*_{\scriptscriptstyle{9}}$ $\frac{1}{\mathfrak{B}} = -\mathcal{I}_{\mathfrak{B}}$ and

$$
\langle \mathcal{I}_{\mathfrak{B}}^{-1}(a_1), R(a_2) \rangle + \langle \mathcal{I}_{\mathfrak{B}}^{-1}(R(a_1)), a_2 \rangle + \lambda \langle \mathcal{I}_{\mathfrak{B}}^{-1}(a_1), a_2 \rangle = 0,
$$

for any $a_1, a_2 \in A$, which implies that $R^* \mathcal{I}_{\mathcal{B}}^{-1} + \mathcal{I}_{\mathcal{B}}^{-1}R + \lambda \mathcal{I}_{\mathcal{B}}^{-1} = 0$, and so that, $\mathcal{I}_{\mathcal{B}} R^* + R \mathcal{I}_{\mathcal{B}} + \lambda \mathcal{I}_{\mathcal{B}} = 0$. Therefore, we get $\tau(r)^{\sharp} = (r^{\sharp})^* = -\frac{1}{4}$ $\frac{1}{\lambda}(I_{\mathfrak{B}}R^* + \lambda I_{\mathfrak{B}}) = \frac{1}{\lambda}$ $\frac{1}{4}RT_{\mathfrak{B}}$, and $\mathcal{I}_{\mathfrak{B}} = r^{\sharp} - \tau(r)^{\sharp}$. Define a multiplication ·*^r* on *A* [∗] by

$$
\xi_1 \cdot_r \xi_2 = \mathbf{I}_A^* (r^{\sharp}(\xi_1))(\xi_2) + \mathbf{I}_A^* (\tau(r)^{\sharp}(\xi_2))(\xi_1) - \mathbf{r}_A^* (\tau(r)^{\sharp}(\xi_2))(\xi_1),
$$

for any $\xi_1, \xi_2 \in A^*$. Second, we will show that $\frac{1}{\lambda} \mathcal{I}_{\mathfrak{B}} : (A, \cdot_r) \to (A, \cdot_R)$ is an isomorphism of anti-Leibniz algebras. Indeed, for any $a_1, a_2, a_3 \in A$, since $\mathfrak{B}(a_1a_2, a_3) = \mathfrak{B}(a_1, a_2a_3 - a_3a_2)$, we get $\langle \mathcal{I}_{\mathcal{B}}^{-1}(\mathbf{r}_A(a_2)(a_1)), a_3 \rangle = \langle (\mathbf{l}_A^* - \mathbf{r}_A^*)$ $A(A)(a_2)(\mathcal{I}_{\mathcal{B}}^{-1}(a_1)), a_3$, and so that, $\mathbf{r}_A(a)\mathcal{I}_{\mathcal{B}} = \mathcal{I}_{\mathcal{B}}((\mathbf{l}_A^* - \mathbf{r}_A^*)$ $_A^*(a_1)$). That is, $r - \tau(r)$ is invariant. Thus, for any $\xi_1, \xi_2 \in A^*$ we have

$$
I_{\mathfrak{B}}(\xi_1 \cdot_r \xi_2) = I_{\mathfrak{B}}\Big(\mathbf{I}_A^*(r^{\sharp}(\xi_1))(\xi_2) + \mathbf{I}_A^*(\tau(r)^{\sharp}(\xi_2))(\xi_1) - \mathbf{r}_A^*(\tau(r)^{\sharp}(\xi_2))(\xi_1)\Big) = \mathbf{r}_A(r^{\sharp}(\xi_1))(I_{\mathfrak{B}}(\xi_2)) + \mathbf{r}_A(\tau(r)^{\sharp}(\xi_2))(I_{\mathfrak{B}}(\xi_1)) = r^{\sharp}(\xi_1)r^{\sharp}(\xi_2) - r^{\sharp}(\xi)\tau(r)^{\sharp}(\xi_2) + r^{\sharp}(\xi_1)\tau(r)^{\sharp}(\xi_2) - \tau(r)^{\sharp}(\xi)\tau(r)^{\sharp}(\xi_2) = r^{\sharp}(\xi_1)r^{\sharp}(\xi_2) - \tau(r)^{\sharp}(\xi_1)\tau(r)^{\sharp}(\xi_2),
$$

and

$$
I_{\mathfrak{B}}(\xi_1) \cdot_P I_{\mathfrak{B}}(\xi_2)
$$

= $R(I_{\mathfrak{B}}(\xi_1))I_{\mathfrak{B}}(\xi_2) + I_{\mathfrak{B}}(\xi_1)P(I_{\mathfrak{B}}(\xi_2)) + \lambda I_{\mathfrak{B}}(\xi_1)I_{\mathfrak{B}}(\xi_2)$
= $\lambda \tau(r)^{\sharp}(\xi_1)(r^{\sharp}(\xi_2) - \tau(r)^{\sharp}(\xi_2)) + \lambda(r^{\sharp}(\xi_1) - \tau(r)^{\sharp}(\xi_1))\tau(r)^{\sharp}(\xi_2) + \lambda(r^{\sharp}(\xi_1) - \tau(r)^{\sharp}(\xi_1))(r^{\sharp}(\xi_2) - \tau(r)^{\sharp}(\xi_2))$
= $\lambda r^{\sharp}(\xi_1)r^{\sharp}(\xi_2) - \lambda \tau(r)^{\sharp}(\xi_1)\tau(r)^{\sharp}(\xi_2),$

Thus, $\frac{1}{\lambda} \mathcal{I}_{\mathfrak{B}}$ is an anti-Leibniz algebra isomorphism from (A^*, \cdot_r) to (A, \cdot_P) .

Finally, since $R + \lambda$ id, $R : (A, \cdot_P) \to (A, \cdot)$ are homomorphisms of anti-Leibniz algebras, we deduce that $r^{\sharp} = \frac{1}{4}$ $\frac{1}{\lambda}(R + \lambda \text{ id})\mathcal{I}_{\mathfrak{B}}$ and $\tau(r)^{\sharp} = \frac{1}{\lambda}$ $\frac{1}{4}R\mathcal{I}_{\mathfrak{B}}$ are homomorphisms of anti-Leibniz algebras from (A^*, \cdot_r) to (*A*, ·). Hence, by Proposition [3.23,](#page-15-2) we get that *r* is a solution of the aLYBE in (*A*, ·), and so that (*A*, ·, Δ) is a is a factorizable anti-Leibniz bialgebra, since $\mathcal{I}_{\mathfrak{B}} = r^{\sharp} - \tau(r)^{\sharp}$ is an isomorphism.

Thus, if $\lambda \neq 0$, we have the following commutative diagram for the factorizable anti-Leibniz bialgebras and skew-quadratic Rota-Baxter anti-Leibniz algebra:

Moreover, following Theorem [3.30,](#page-17-0) we have two corollaries.

Corollary 3.31. *Let* (*A*, ·) *be an anti-Leibniz algebra and r* ∈ *A*⊗*A. If there is an anti-Leibniz bialgebra* (*A*, ·, ∆) *induced by r and it is factorizable, then* −λ id −*P is also a Rota-Baxter operator of weight* λ *on quadratic anti-Leibniz algebra* $(A, \cdot, \mathfrak{B}_I)$, where $I = r^{\sharp} - \tau(r)^{\sharp}$, $P = \lambda \tau(r)^{\sharp} I^{-1} : A \to A$ and the bilinear *form* \mathfrak{B}_I *is defined by* $\mathfrak{B}_I(a_1, a_2) = \langle I^{-1}(a_1), a_2 \rangle$, for any $a_1, a_2 \in A$.

Proof. It can be obtained by direct calculation.

Corollary 3.32. Let (A, \cdot) be an anti-Leibniz algebra and $r \in A \otimes A$. Define $I = r^{\sharp} - \tau(r)^{\sharp}$ and $P = \lambda \tau(r)^{\sharp} \mathcal{I}^{-1} : A \to A$ where $0 \neq \lambda \in \mathbb{k}$. If there is an anti-Leibniz bialgebra (A, \cdot, Δ) induced by r and *it is factorizable, then* (A, \cdot_P, Δ_I) *is an anti-Leibniz bialgebra, where*

$$
\Delta_I^*(\xi_1 \otimes \xi_2) = -\frac{1}{\lambda} \mathcal{I}^{-1}(\mathcal{I}(\xi_1) \mathcal{I}(\xi_2)),
$$

for any $\xi_1, \xi_2 \in A^*$. Moreover, $\frac{1}{\lambda}I : A^* \to A$ gives an isomorphism of anti-Leibniz bialgebras from $(A^*, \cdot, \Delta_{A^*})$ to (A, \cdot_P, Δ_I) , where Δ_{A^*} is the dual of product in (A, \cdot) , i.e., $\Delta_{A^*}^*(a_1 \otimes a_2) = a_1 a_2$, for any a_1, a_2 ∈ *A*.

Proof. First, we show that $\frac{1}{\lambda}I$ is an anti-Leibniz algebra isomorphism from (A^*, \cdot_r) to (A, \cdot_P) . In fact, for any $\xi_1, \xi_2 \in A^*$, taking $a_1 = I(\xi_1) \in A$ and $a_2 = I(\xi_2) \in A$, by Eq. [\(3.8\)](#page-18-0), we have

$$
\frac{1}{\lambda}I(\xi_1 \cdot_r \xi_2) = \frac{1}{\lambda^2} (P(I(\xi_1))I(\xi_2) + I(\xi_1)P(I(\xi_2)) + \lambda I(\xi_1)I(\xi_2)) = \frac{1}{\lambda}I(\xi_1) \cdot_p \frac{1}{\lambda}I(\xi_2).
$$

So, $\frac{1}{\lambda}I$ is an anti-Leibniz algebra isomorphism.

Second, denote the dual of Δ_I by \cdot_I . Since $(\frac{1}{\lambda}I)^* = -\frac{1}{\lambda}$ $\frac{1}{\lambda}$ *I*, we get

$$
(\frac{1}{\lambda}I)^{*}(\xi_1 \cdot_I \xi_2) = \frac{1}{\lambda}I(\xi_1)\frac{1}{\lambda}I(\xi_2) = (\frac{1}{\lambda}I)^{*}(\xi_1)(\frac{1}{\lambda}I)^{*}(\xi_2),
$$

which means $(\frac{1}{\lambda}I)^*$: $(A^*, \cdot_I) \to (A, \cdot)$ is an isomorphism of anti-Leibniz algebras. Hence, $\frac{1}{\lambda}I$: $(A^*, \Delta_{A^*}) \to (A, \Delta_I)$ is an isomorphism of anti-Leibniz coalgebras. Therefore, $\frac{1}{\lambda}I$ is an isomorphism of anti-Leibniz bialgebras and (A, \cdot_P, Δ_I) is an anti-Leibniz bialgebra. $□$

$$
\Box
$$

3.4. Anti-Leibniz bialgebras via Leibniz bialgebras. Here, we constrict the anti-Leibniz bialgebras form Leibniz bialgebras and anti-commutative anti-associative algebras. By using this method, we can construct a large number of examples for anti-Leibniz algebra and anti-Leibniz bialgebra. Recall that a *(left)* Leibniz algebra is a pair $(L, [-,-])$, where L is a vector space and $[-,-] : L \otimes L \to L$ is a multiplication such that

$$
[x_1, [x_2, x_3]] = [[x_1, x_2], x_3] + [x_2, [x_1, x_3]],
$$

for any $x_1, x_2, x_3 \in L$. An *anti-commutative anti-associative algebra* (B, \cdot) is a vector space *B* with a bilinear operations $\cdot : B \otimes B \to B$, $(b_1, b_2) \mapsto b_1b_2 := b_1 \cdot b_2$, such that

$$
b_1b_2 = -b_2b_1
$$
, and $b_1(b_2b_3) = -(b_1b_2)b_3$,

for any $b_1, b_2, b_3 \in B$. The anti-commutative anti-associative algebra is also called the dual mock-Lie algebra, since the operad of anti-commutative anti-associative algebra is the Koszul dual to the mock-Lie operad.

Lemma 3.33. *Let* (*L*, [−, −]) *be a Leibniz algebra,* (*B*, ·) *be an anti-commutative anti-associative algebra. Define a binary operation on A* := $L \otimes B$ *by*

$$
(x_1 \otimes b_1) \circ (x_2 \otimes b_2) = [x_1, x_2] \otimes (b_1 b_2),
$$

for any $x_1, x_2 \in L$ and $b_1, b_2 \in B$. Then (A, \circ) is an anti-Leibniz algebra, which is called the anti-Leibniz *algebra induced by* $(L, [-,-])$ *and* (B, \cdot) *.*

Proof. For any $x_1, x_2, x_3 \in L$ and $b_1, b_2, b_3 \in B$, note that

$$
(x_1 \otimes b_1) \circ ((x_2 \otimes b_2) \circ (x_3 \otimes b_3)) + ((x_1 \otimes b_1) \circ (x_2 \otimes b_2)) \circ (x_3 \otimes b_3)
$$

+ $(x_2 \otimes b_2) \circ ((x_1 \otimes b_1) \circ (x_3 \otimes b_3))$
= $[x_1, [x_2, x_3]] \otimes (b_1(b_2b_3)) + [[x_1, x_2], x_3] \otimes ((b_1b_2)b_3) + [x_2, [x_1, x_3]] \otimes (b_2(b_1b_3))$
= $((x_1, [x_2, x_3]] - [[x_1, x_2], x_3] - [x_2, [x_1, x_3]]) \otimes (b_1(b_2b_3))$
= 0,

we get (A, \circ) is an anti-Leibniz algebra.

For classification problems that anti-Leibniz algebra, we still know very little. But for the classification of Leibniz algebras, some results have been obtained [\[2\]](#page-28-20). Recall that the non trivial anticommutative anti-associative algebra of dimensional two is only isomorphic to the algebra $(B = \mathbb{k}\{e_1, e_2\}, \cdot)$, where the nonzero multiplication is given by $e_1e_1 = e_2$. Two dimensional Leibniz algebras have been classified by Cuvier [\[13\]](#page-28-21). More precisely, the non trivial Leibniz algebraic structure on $L = \mathbb{k}{x_1, x_2}$ is only isomorphic to the algebra with nonzero multiplication as follows:

$$
L_1: [x_1, x_1] = x_2; \qquad L_2: [x_1, x_2] = -[x_2, x_1] = x_2; \qquad L_1: [x_1, x_2] = x_1 = [x_2, x_2].
$$

Thus, by Lemma [3.33,](#page-20-1) we get some 4-dimensional anti-Leibniz algebra as follows: $A = \mathbb{k}{a_1, a_2, a_3, a_4}$ with nonzero multiplication

$$
A_1: a_1 \circ a_1 = a_2;
$$
 $A_2: a_1 \circ a_2 = -a_2 \circ a_1 = a_3;$ $A_1: a_1 \circ a_2 = a_3 = a_2 \circ a_2.$

Let *B* be a vector space and $\Delta : B \to B \otimes B$ be a linear map. Then (B, Δ) is called an *anticocommutative anti-coassociative coalgebra* if

$$
\tau \Delta = -\Delta
$$
, and $(id \otimes \Delta)\Delta = -(\Delta \otimes id)\Delta$.

Clearly, the notion of anti-cocommutative anti-coassociative coalgebras is the dualization of the notion of anti-commutative anti-associative algebras, that is, (B, Δ) is an anti-cocommutative anti-coassociative coalgebra if and only if (*B*^{*}, ∆^{*}) is an anti-commutative anti-associative algebra. Recall that a *Leibniz coalgebra* is a pair (L, δ) , where *L* is a vector space and $\delta : L \to L \otimes L$ is a comultiplication such that

$$
(\delta \otimes id)\delta(x) + (\tau \otimes id)(id \otimes \delta)\delta(x) - (id \otimes \delta)\delta(x) = 0,
$$

$$
\qquad \qquad \Box
$$

for any $x \in L$. Now, we give the dual version of Lemma [3.33.](#page-20-1)

Lemma 3.34. *Let* (*L*, δ) *be a Leibniz coalgebra,* (*B*, ∆) *be an anti-cocommutative anti-coassociative coalgebra. Define* $A := L \otimes B$ *and a linear map* $\tilde{\Delta} : A \rightarrow A \otimes A$ *by*

$$
\tilde{\Delta}(x \otimes b) = \delta(x) \bullet \Delta(b) = \sum_{(x)} \sum_{(b)} (x_{(1)} \otimes b_{(1)}) \otimes (x_{(2)} \otimes b_{(2)}),
$$

for any $x \in L$ *and* $b \in B$, *where* $\Delta(b) = \sum_{(b)} b_{(1)} \otimes b_{(2)}$ *and* $\delta(x) = \sum_{(x)} x_{(1)} \otimes x_{(2)}$ *in the Sweedler notation. Then* $(A, \tilde{\Delta})$ *is an anti-Leibniz algebra.*

Proof. For any *x i* i_1, x_2^i i_{2}^{i}, x_{3}^{i} j_3 \in *L* and b_1^j j_1^j, b_2^j $^{j}_{2}, b^{j}_{3}$ $\frac{J}{3} \in B$, we denote

$$
\Big(\sum_i x_1^i \otimes x_2^i \otimes x_3^i\Big) \bullet \Big(\sum_j b_1^j \otimes b_2^j \otimes b_3^j\Big) = \sum_{i,j} (x_1^i \otimes b_1^j) \otimes (x_2^i \otimes b_2^j) \otimes (x_3^i \otimes b_3^j).
$$

Then, for any $x \in L$ and $b \in B$, we have

$$
(\tilde{\Delta} \otimes id)\tilde{\Delta}(x \otimes b) + (id \otimes \tilde{\Delta})\tilde{\Delta}(x \otimes b) + (\tau \otimes id)(id \otimes \tilde{\Delta})\tilde{\Delta}(x \otimes b)
$$

= $(\delta \otimes id)\delta(x) \cdot (\Delta \otimes id)\Delta(b) + (id \otimes \delta)\delta(x) \cdot (id \otimes \Delta)\Delta(b) + (\tau \otimes id)(id \otimes \delta)\delta(x) \cdot (\tau \otimes id)(id \otimes \Delta)\Delta(b)$
= $((\delta \otimes id)\delta(x) + (\tau \otimes id)(id \otimes \delta)\delta(x) - (id \otimes \delta)\delta(x)) \cdot (\Delta \otimes id)\Delta(b)$
= 0.

Thus, (A, Δ) is an anti-Leibniz coalgebra.

Let (B, \cdot) be an anti-commutative anti-associative algebra. A bilinear form $\overline{\omega}(-, -)$ on (B, \cdot) is called *nondegenerate* if $\varpi(b_1, b_2) = 0$ for any $b_2 \in B$, then $b_1 = 0$; is called *symmetric* if $\varpi(b_1, b_2) = \varpi(b_2, b_1)$; is called *invariant* if $\varpi(b_1b_2, b_3) = \varpi(b_1, b_2b_3)$, for any $b_1, b_2, b_3 \in B$. A *quadratic anti-commutative anti-associative algebra*, denoted by (B, ϖ) , is an anti-commutative anti-associative algebra (B, \cdot) together with a symmetric invariant nondegenerate bilinear form.

Lemma 3.35. Let (B, ϖ) be a quadratic anti-commutative anti-associative algebra. Suppose $\Delta_{\varpi}: B \to$ *B* ⊗ *B be the dual of the multiplication of* (B, ϖ) *under the bilinear form* $\varpi(-, -)$ *, i,e.,*

$$
\varpi(\Delta_{\varpi}(b_1), b_2 \otimes b_3) = \varpi(b_1, b_2b_3),
$$

 \int *for any* $b_1, b_2, b_3 \in B$, where $\varpi(\Delta_{\varpi}(b_1), b_2 \otimes b_3) = \sum_{(b_1)} \varpi((b_1)_{(1)}, b_2) \varpi((b_1)_{(2)}, b_3)$ if $\Delta_{\varpi}(b_1) = \sum_{(b_1)} (b_1)_{(1)}$ ⊗ $(b_1)_{(2)}$. Then $(B, \Delta_{\overline{\omega}})$ *is an anti-cocommutative anti-coassociative coalgebra.*

Proof. For any $b \in B$ and $b_1 \otimes b_2 \otimes b_3 \in B \otimes B \otimes B$, by the definition of Δ_{ϖ} , we get $\varpi(\tau \Delta_{\varpi}(b), b_1 \otimes b_2) =$ $\overline{\omega}(b, b_2b_1) = -\overline{\omega}(b, b_1b_2) = \overline{\omega}(-\Delta_{\overline{\omega}}(b), b_1 \otimes b_2)$ and

$$
\begin{aligned} \varpi((\Delta_{\varpi} \otimes \mathrm{id})\Delta_{\varpi}(b), \ \ b_1 \otimes b_2 \otimes b_3) &= \varpi(b, \ (b_1b_2)b_3) \\ &= -\varpi(b, \ b_1(b_2b_3)) \\ &= \varpi(-(\mathrm{id} \otimes \Delta_{\varpi})\Delta_{\varpi}(b), \ \ b_1 \otimes b_2 \otimes b_3). \end{aligned}
$$

That is, $(\Delta_{\overline{\omega}} \otimes id)\Delta_{\overline{\omega}} = -(id \otimes \Delta_{\overline{\omega}})\Delta_{\overline{\omega}}$ and $\Delta_{\overline{\omega}} = -\tau\Delta_{\overline{\omega}}$. Thus $(B, \Delta_{\overline{\omega}})$ is an anti-cocommutative anticoassociative coalgebra.

Example 3.36. Consider the 2-dimensional anti-commutative anti-associative algebra ($B = \mathbb{k}\{e_1, e_2\}$, ·), *where the nonzero multiplication is given by* $e_1e_1 = e_2$ *. Define a bilinear form* $\varpi(-,-)$ *on* (B,\cdot) *by* $\overline{\omega}(e_1, e_2) = \overline{\omega}(e_2, e_1) = 1$ and $\overline{\omega}(e_1, e_1) = \overline{\omega}(e_2, e_2) = 0$. Then $(B, \overline{\omega})$ is a quadratic anti-commutative *anti-associative algebra. Thus, we get an anti-cocommutative anti-coassociative coalgebra* (B, Δ_{ϖ}) *, where* $\Delta_{\varpi}(e_1) = e_2 \otimes e_2$ *and* $\Delta_{\varpi}(e_2) = 0$ *.*

$$
\qquad \qquad \Box
$$

Recall that a *Leibniz bialgebra* is a triple $(L, [-, -], \delta)$, where $(L, [-, -])$ is a Leibniz algebra, (L, δ) is a Leibniz coalgebra and the following equations hold:

$$
\tau(\mathbf{r}_L(y) \otimes id)\delta(x) = (\mathbf{r}_L(x) \otimes id)\delta(y),
$$

$$
\delta([x, y]) = (id \otimes \mathbf{r}_L(y) - (\mathbf{l}_L + \mathbf{r}_L)(y) \otimes id)(id \otimes id + \tau)\delta(x) + (id \otimes \mathbf{l}_L(x) + \mathbf{l}_L(x) \otimes id)\delta(y),
$$

for all $x, y \in L$.

Theorem 3.37. Let $(L, [-, -], \delta)$ be a Leibniz bialgebra and (B, ϖ) be a quadratic anti-commutative *anti-associative algebra. Suppose* $(A = L \otimes B, \circ)$ *is the anti-Leibniz algebra induced by* $(L, [-, -])$ *by* (B, \cdot) , Δ_{ϖ} : $B \to B \otimes B$ *is the dual of the multiplication of* (B, ϖ) *under the bilinear form* $\varpi(-, -)$ *, and* $\hat{\Delta}: A \to A \otimes A$ *is the linear map defined by* $\hat{\Delta}(x \otimes b) = \delta(x) \cdot \Delta_{\varpi}(b)$ *for* $x \in L$, $b \in B$. Then $(A, \circ, \hat{\Delta})$ *is an anti-Leibniz bialgebra, which is called the anti-Leibniz bialgebra induced by* $(L, [-, -], \delta)$ *and* (B, ϖ) *.*

Proof. Since $\varpi(-,-)$ is an invariant bilinear form, for any *b*, *b*', *c*, *d* \in *B*, we have

$$
\varpi\Big(\sum_{(b)} b_{(1)}b' \otimes b_{(2)}, c \otimes d\Big) = \varpi\Big(\sum_{(b)} b_{(1)} \otimes b_{(1)}, b'c \otimes d\Big) \n= \varpi(b, (b'c)d) \n= -\varpi(\Delta_{\varpi}(bb'), c \otimes d).
$$

That is to say, $\Delta_{\varpi}(bb') = -\sum_{(b)} b_{(1)}b' \otimes b_{(2)}$. Moreover, since (B, \cdot) is anti-commutative and Δ_{ϖ} is anticocommutative, we get $-\Delta_{\overline{\omega}}(b'b) = \Delta_{\overline{\omega}}(bb') = \sum_{(b')} b'_{(1)} b \otimes b'_{(2)} = \sum_{(b)} b_{(2)} \otimes b_{(1)} b' = \sum_{(b)} b'b_{(1)} \otimes b_{(2)} =$ $-\sum_{(b)} b_{(1)}b' \otimes b_{(2)} = \sum_{(b)} b_{(2)} \otimes b_{(1)}b' = -\sum_{(b)} b' b_{(2)} \otimes b_{(1)} = \sum_{(b)} b' b_{(2)}b' \otimes b_{(1)} = -\sum_{(b)} b'_{(1)} \otimes b b'_{(2)} =$ $-\sum_{(b')} bb'_{(1)} \otimes b'_{(2)}$. Thus, for any *x* ⊗ *b*, *x*' ⊗ *b*' ∈ *A*,

$$
(\mathbf{r}_A(x \otimes b) \otimes \mathrm{id}) \tilde{\Delta}(x' \otimes b') - \hat{\tau}((\mathbf{r}_A(x' \otimes b') \otimes \mathrm{id}) \tilde{\Delta}(x \otimes b))
$$

=
$$
\sum_{(x')} \sum_{(b')} ([x'_{(1)}, x] \otimes x'_{(2)}) \bullet (b'_{(1)}b \otimes b'_{(2)}) - \sum_{(x)} \sum_{(b)} (x_{(2)} \otimes [x_{(1)}, x']) \bullet (b_{(2)} \otimes b_{(1)}b')
$$

=
$$
((\mathbf{r}_L(x) \otimes \mathrm{id}) \delta(x') - \tau((\mathbf{r}_L(x') \otimes \mathrm{id}) \delta(x))) \bullet \Delta_{\varpi}(bb')
$$

= 0,

and

$$
\tilde{\Delta}((x \otimes b) \circ (a' \otimes c')) + ((id \otimes r_{\mathcal{A}}(x' \otimes b') + (l_A - r_A)(x' \otimes b') \otimes id)(id \otimes id - \tau))\tilde{\Delta}(x \otimes b)
$$
\n
$$
+ (id \otimes l_A(x \otimes b) + l_A(x \otimes b) \otimes id)\tilde{\Delta}(x' \otimes b')
$$
\n
$$
= \delta([x, x']) \bullet \Delta_{\varpi}(bb') + \sum_{(x)} \sum_{(b)} (x_{(1)} \otimes [x_{(2)}, x']) \bullet (b_{(1)} \otimes b_{(2)}b') + \sum_{(x)} \sum_{(b)} ([x', x_{(1)}] \otimes x_{(2)}) \bullet (b' b_{(1)} \otimes b_{(2)})
$$
\n
$$
- \sum_{(x)} \sum_{(b)} ([x_{(1)}, x'] \otimes x_{(2)}) \bullet (b_{(1)}b' \otimes b_{(2)}) - \sum_{(x)} \sum_{(b)} (x_{(2)} \otimes [x_{(1)}, x']) \bullet (b_{(2)} \otimes b_{(1)}b')
$$
\n
$$
- \sum_{(x)} \sum_{(b)} ((x', x_{(2)}] \otimes x_{(1)}) \bullet (b' b_{(2)} \otimes b_{(1)}) + \sum_{(x)} \sum_{(b)} ((x_{(2)}, x'] \otimes x_{(1)}) \bullet (b_{(2)}b' \otimes b_{(1)})
$$
\n
$$
+ \sum_{(x)} \sum_{(b')} (x'_{(1)} \otimes [x, x'_{(2)}]) \bullet (b'_{(1)} \otimes b'_{(2)}) + \sum_{(x)} \sum_{(b')} ((x, x'_{(1)}] \otimes x'_{(2)}) \bullet (b b'_{(1)} \otimes b'_{(2)})
$$
\n
$$
= (\Delta([x, x']) - ((id \otimes r_L(x') + r_L(x') \otimes id + l_L(x') \otimes id)(id \otimes id + \tau))\delta(x)) \bullet \Delta_{\varpi}(bb')
$$
\n
$$
- ((id \otimes l_L(x) + l_L(x) \otimes id) \delta(x')) \bullet \Delta_{\varpi}(bb')
$$
\n
$$
= 0,
$$

since $(L, [-,-], \delta)$ is a Leibniz bialgebra. By the definition, we get that $(A, \circ, \tilde{\Delta})$ is a completed anti-Leibniz bialgebra.

Example 3.38. *In* [\[20\]](#page-28-22)*, the authors have given some examples of triangular Leibniz bialgebras. Here we using Theorem* [3.37](#page-22-0) *to give some examples of anti-Leibniz bialgebras. We fix* (*B*, ·) *as the* 2*-dimensional* *anti-commutative anti-associative algebra given in Example* [3.36](#page-21-0), *i.e.*, $B = \mathbb{k}{e_1, e_2}$ *and* $e_1e_1 = e_2$ *By the bilinear form* $\varpi(-,-)$ *on* (B, \cdot) *defined in Example* [3.36](#page-21-0)*, we get an anti-cocommutative anticoassociative coalgebra* $(B, \Delta_{\overline{\omega}})$ *, where* $\Delta_{\overline{\omega}}(e_1) = e_2 \otimes e_2$ *.*

(*I*) Let $(L, [-,-], \delta)$ be a 2-dimensional Leibniz bialgebra with basis $\{x_1, x_2\}$, and $(A = L \otimes B, \circ, \tilde{\Delta})$ be *the induced* 4-dimensional anti-Leibniz bialgebra with basis $\{a_1, a_2, a_3, a_4\}$.

- (i) If $[x_1, x_1] = x_2$, $\delta(x_1) = k(x_2 \otimes x_2)$, then $a_1 \circ a_1 = a_2$ and $\Delta(a_1) = k(a_2 \otimes a_2)$, where $k \in \mathbb{R}$. That *is, the Leibniz bialgebra* (*L*, [−, −], δ) *is also an anti-Leibniz bialgebra.*
- (ii) If $[x_1, x_2] = x_1 = [x_2, x_1]$, $\delta(x_1) = \delta(x_2) = k(x_1 \otimes x_2 x_2 \otimes x_1)$, then $a_1 \circ a_2 = a_3 = a_2 \circ a_2$ and $\Delta(a_1) = \Delta(a_1) = k(a_2 \otimes a_4 - a_4 \otimes a_2)$, where $k \in \mathbb{R}$.

(*II*) Let $(L, [-,-], \delta)$ be a 3-dimensional Leibniz bialgebra with basis $\{x_1, x_2, x_3\}$, and $(A = L \otimes B, \circ, \Delta)$ *be the induced* 6-dimensional anti-Leibniz bialgebra with basis $\{a_1, a_2, a_3, a_4, a_5, a_6\}$.

- (i) If $[x_1, x_3] = x_1 + x_2$, $[x_3, x_3] = x_1$, $\delta(x_3) = k(x_1 \otimes x_1 + x_2 \otimes x_1) + l(x_1 \otimes x_2 + x_2 \otimes x_2)$, then $a_1 \circ a_5 = a_2 + a_4$, $a_5 \circ a_5 = a_2$ and $\tilde{\Delta}(a_5) = k(a_2 \otimes a_2 + a_4 \otimes a_2) + l(a_1 \otimes a_4 + a_4 \otimes a_4)$, where $k, l \in \mathbb{k}$.
- (ii) If $[x_2, x_3] = x_2 = -[x_3, x_2]$, $[x_3, x_3] = x_1$, $\delta(x_2) = k(x_1 \otimes x_2)$, $\delta(x_3) = k(x_1 \otimes x_1) l(x_1 \otimes x_2)$, then $a_3 \circ a_5 = a_4 = -a_5 \circ a_3$, $a_5 \circ a_5 = a_3$, $\tilde{\Delta}(a_3) = k(a_2 \otimes a_4)$ and $\tilde{\Delta}(a_5) = k(a_2 \otimes a_2) - l(a_2 \otimes a_4)$, *where* $k, l \in \mathbb{k}$.
- (iii) If $[x_2, x_2] = x_1 = [x_3, x_3]$, $\delta(x_2) = k(x_1 \otimes x_1)$, then $a_3 \circ a_3 = a_2 = a_5 \circ a_5$ and $\tilde{\Delta}(a_3) = k(a_2 \otimes a_2)$, *where* $k \in \mathbb{K}$.

4. Infinite-dimensional anti-Leibniz bialgebras

In this section, we constrict infinite-dimensional anti-Leibniz bialgebras form finite-dimensional anti-Leibniz bialgebras by the completed tensor product. In this section, a commutative algebra means a commutative associative algebra, and a cocommutative coalgebra means a cocommutative coassociative coalgebra for simply.

Definition 4.1. A Z-graded commutative algebra *(resp. a* Z-graded anti-Leibniz algebra) is a commuta*tive algebra* (C, \cdot) *(resp. an anti-Leibniz algebra* (A, \cdot) *) with a linear decomposition* $C = \bigoplus_{i \in \mathbb{Z}} C_i$ *(resp.* $A=\oplus_{i\in\mathbb{Z}}A_i$) such that each C_i (resp. A_i) is finite-dimensional and $C_i\cdot C_j\subseteq C_{i+j}$ (resp. $A_i\cdot A_j\subseteq A_{i+j}$) for *all* $i, j \in \mathbb{Z}$.

Clearly, the Laurent polynomial algebra $\mathbb{K}[\mathfrak{t}, \mathfrak{t}^{-1}]$ is a Z-graded commutative algebra. We now consider a tensor product of anti-Leibniz algebras and Z-graded commutative algebras.

Lemma 4.2. Let (A, \cdot) be a finite-dimensional anti-Leibniz algebra, $(C = \bigoplus_{i \in \mathbb{Z}} C_i, \cdot)$ be a Z-graded *commutative algebra. Define a binary operation on* $\mathcal{A} := A \otimes C$ by

$$
(a_1 \otimes c_1) \diamond (a_2 \otimes c_2) = (a_1 a_2) \otimes (c_1 c_2),
$$

for any $a_1, a_2 \in A$ *and* $c_1, c_2 \in C$. Then (A, \diamond) *is a* Z-graded *anti-Leibniz algebra, which is called* affine anti-Leibniz algebra *from* (A, \cdot) *by* $(C = \bigoplus_{i \in \mathbb{Z}} C_i, \cdot)$. Moreover, if $(C = \bigoplus_{i \in \mathbb{Z}} C_i, \cdot) = \mathbb{k}[t, t^{-1}]$ *is the Laurent polynomial algebra, then* (A, ⋄) *is a* Z*-graded anti-Leibniz algebra if and only if* (*A*, ·) *is an anti-Leibniz algebra.*

Proof. It is straightforward.

To carry out the affine anti-Leibniz coalgebra, we need to extend the codomain of the coproduct ∆ to allow infinite sums. Let $U = \bigoplus_{i \in \mathbb{Z}} U_i$ and $V = \bigoplus_{i \in \mathbb{Z}} V_i$ be Z-graded vector spaces. We call the *completed tensor product* of *U* and *V* to be the vector space

$$
U\,\hat{\otimes}\,V:=\prod_{i,j\in\mathbb{Z}}U_i\otimes V_j.
$$

$$
\qquad \qquad \Box
$$

If *U* and *V* are finite-dimensional, then $U \hat{\otimes} V$ is just the usual tensor product $U \otimes V$. In general, an element in $U \hat{\otimes} V$ is an infinite formal sum $\sum_{i,j \in \mathbb{Z}} \overline{X}_{ij}$ with $X_{ij} \in U_i \otimes V_j$. So $X_{ij} = \sum_{\alpha} u_{i,\alpha} \otimes v_{j,\alpha}$ for pure tensors $u_{i,\alpha} \otimes v_{j,\alpha} \in U_i \otimes V_j$ with α in a finite index set. Thus a general term of $U \otimes V$ is a possibly infinite sum $\sum_{i,j,\alpha} u_{i\alpha} \otimes v_{j\alpha}$, where *i*, $j \in \mathbb{Z}$ and α is in a finite index set (which might depend on *i*, *j*). With these notations, for linear maps $f: U \to U'$ and $g: V \to V'$, define

$$
f \hat{\otimes} g : U \hat{\otimes} V \to U' \hat{\otimes} V', \qquad \sum_{i,j,\alpha} u_{i,\alpha} \otimes v_{j,\alpha} \mapsto \sum_{i,j,\alpha} f(u_{i,\alpha}) \otimes g(v_{j,\alpha}).
$$

Also the twist map τ has its completion

$$
\hat{\tau}: V \hat{\otimes} V \to V \hat{\otimes} V, \qquad \sum_{i,j,\alpha} u_{i,\alpha} \otimes v_{j,\alpha} \mapsto \sum_{i,j,\alpha} v_{j,\alpha} \otimes u_{i,\alpha}.
$$

Finally, we define a (completed) coproduct to be a linear map

$$
\hat{\Delta}: V \to V \hat{\otimes} V, \qquad \hat{\Delta}(v) := \sum_{i,j,\alpha} v_{1,i,\alpha} \otimes v_{2,j,\alpha}.
$$

Then we have the well-defined map

$$
(\hat{\Delta}\hat{\otimes} \text{id})\hat{\Delta}(v) = (\hat{\Delta}\hat{\otimes} \text{id})\Big(\sum_{i,j,\alpha} v_{1,i,\alpha} \otimes v_{2,j,\alpha}\Big) := \sum_{i,j,\alpha} \hat{\Delta}(v_{1,i,\alpha}) \otimes v_{2,j,\alpha} \in V \hat{\otimes} V \hat{\otimes} V.
$$

Definition 4.3. (*i*) *A* completed cocommutative coalgebra *is a pair* $(C, \bar{\Delta})$ *where* $C = \bigoplus_{i \in \mathbb{Z}} C_i$ *is a* Z*graded vector space and* $\overline{\Delta}: C \to C \hat{\otimes} C$ *is a linear map satisfying*

$$
\hat{\tau}\bar{\Delta} = \bar{\Delta}, \qquad (\bar{\Delta}\,\hat{\otimes}\,\mathrm{id})\bar{\Delta} = (\mathrm{id}\,\hat{\otimes}\,\bar{\Delta})\bar{\Delta}.
$$

 (iii) *A* completed anti-Leibniz coalgebra *is a pair* $(A, \hat{\Delta})$ *where* $A = \bigoplus_{i \in \mathbb{Z}} A_i$ *is a* Z-graded vector space *and* $\hat{\Delta}$: *A* → *A* $\hat{\otimes}$ *A is a linear map satisfying*

$$
(\hat{\Delta} \otimes id)\hat{\Delta} + (id \otimes \hat{\Delta})\hat{\Delta} + (\hat{\tau} \otimes id)(id \otimes \hat{\Delta})\hat{\Delta} = 0.
$$

Example 4.4. Consider the Laurent polynomial algebra k[t, t⁻¹], there is a naturally completed cocom*mutative coalgebra structure on* $\mathbb{K}[\tilde{t}, t^{-1}]$, which is given by $\bar{\Delta}(t^k) = \sum_{i \in \mathbb{Z}} t^i \otimes t^{k-i}$ for all $k \in \mathbb{Z}$. This *completed cocommutative coalgebra is called* Laurent polynomial coalgebra*.*

For completed anti-Leibniz coalgebra, we give the dual version of Lemma [4.2.](#page-23-1)

Lemma 4.5. *Let* (A, Δ) *be a finite-dimensional anti-Leibniz coalgebra,* $(C, \overline{\Delta})$ *be a completed cocommutative coalgebra and* $\mathcal{A} := A \otimes C$. Define a linear map $\hat{\Delta} : \mathcal{A} \to \mathcal{A} \hat{\otimes} \mathcal{A}$ by

(4.1)
$$
\hat{\Delta}(a \otimes c) = \Delta(a) \bullet \bar{\Delta}(c) = \sum_{(a)} \sum_{i,j,\alpha} (a_{(1)} \otimes c_{1,i,\alpha}) \otimes (a_{(2)} \otimes c_{2,j,\alpha}),
$$

for any $a \in A$ *and* $c \in C$, where $\Delta(a) = \sum_{(a)} a_{(1)} \otimes a_{(2)}$ *in the Sweedler notation and* $\bar{\Delta}(c) = \sum_{i,j,a} c_{1,i,a} \otimes$ $c_{2,i,\alpha}$ *. Then* $(\mathcal{A}, \hat{\Delta})$ *is a completed anti-Leibniz coalgebra.*

 \bar{F} urthermore, if A is finite-dimensional, (C, $\bar{\Delta}$) is the Laurent polynomial coalgebra (k[t, t⁻¹], $\bar{\Delta}$), then (A, ∆ˆ) *is a completed anti-Leibniz coalgebra if and only if* (*A*, ∆) *is an anti-Leibniz coalgebra.*

Proof. First, for any $\sum_l a'_l \otimes a''_l \otimes a''_l'$ $\ell_l'' \in A \otimes A \otimes A$ and $\sum_{i,j,k,\alpha} c'_{i,\alpha} \otimes c''_{j,\alpha} \otimes c''_{k,\alpha}$ $\gamma_{k,\alpha}^{\prime\prime\prime}\in C\,\hat{\otimes}\,C\,\hat{\otimes}\,C,$ we denote

$$
\Big(\sum_{l} a'_{l} \otimes a''_{l} \otimes a''_{l}\Big) \bullet \Big(\sum_{i,j,k,\alpha} c'_{i,\alpha} \otimes c''_{j,\alpha} \otimes c''_{k,\alpha}\Big) = \sum_{l} \sum_{i,j,k,\alpha} (a'_{l} \otimes c'_{i,\alpha}) \otimes (a''_{l} \otimes c''_{j,\alpha}) \otimes (a''_{l} \otimes c''_{k,\alpha}).
$$

Using the above notations, for any $a \otimes c \in \mathcal{A}$, we have

$$
(\hat{\Delta}\hat{\otimes} \text{id})\hat{\Delta}(a\otimes c) + (\text{id}\hat{\otimes}\hat{\Delta})\hat{\Delta}(a\otimes c) + (\hat{\tau}\hat{\otimes} \text{id})(\text{id}\hat{\otimes}\hat{\Delta})\hat{\Delta}(a\otimes c)
$$

= $(\Delta\otimes \text{id})\delta(a) \bullet (\bar{\Delta}\hat{\otimes} \text{id})\bar{\Delta}(c) + (\text{id}\otimes\Delta)\Delta(a) \bullet (\text{id}\hat{\otimes}\bar{\Delta})\bar{\Delta}(c) + (\tau\otimes \text{id})(\text{id}\otimes\Delta)\Delta(a) \bullet (\hat{\tau}\hat{\otimes} \text{id})(\text{id}\hat{\otimes}\bar{\Delta})\bar{\Delta}(c)$
= $((\Delta\otimes \text{id})\Delta(a) + (\text{id}\otimes\Delta)\Delta(a) + (\tau\otimes \text{id})(\text{id}\otimes\Delta)\Delta(a)) \bullet (\bar{\Delta}\hat{\otimes} \text{id})\bar{\Delta}(c)$
= 0.

That is, $(\mathcal{A}, \hat{\Delta})$ is a completed anti-Leibniz coalgebra.

Second, if $(C, \bar{\Delta}) = (\Bbbk[t, t^{-1}], \bar{\Delta})$ and $(\mathcal{A}, \hat{\Delta})$ is a completed anti-Leibniz coalgebra constructed as above, we abbreviated $a \otimes t^k$ as at^k for $a \in A$, then

$$
(\hat{\Delta}\hat{\otimes}\text{id})\hat{\Delta}(at^{k}) = \sum_{(a)}\sum_{i,j}(a_{11}t^{j})\otimes(a_{12}t^{i-j})\otimes(a_{2}t^{k-i}),
$$

\n
$$
(\text{id}\hat{\otimes}\hat{\Delta})\hat{\Delta}(at^{k}) = \sum_{(a)}\sum_{i,j}(a_{1}t^{i})\otimes(a_{21}t^{j})\otimes(a_{22}t^{k-i-j}),
$$

\n
$$
(\hat{\tau}\hat{\otimes}\text{id})(\text{id}\hat{\otimes}\hat{\Delta})\hat{\Delta}(at^{k}) = \sum_{(a)}\sum_{i,j}(a_{21}t^{j})\otimes(a_{1}t^{i})\otimes(a_{22}t^{k-i-j}).
$$

Let $i = j = 0$. Comparing the coefficients of $1 \otimes 1 \otimes t^k$, we obtain $\sum_{(a)} a_{11} \otimes a_{12} \otimes a_2 + a_1 \otimes a_{21} \otimes a_{22} + a_1 \otimes a_2$ $a_{21} \otimes a_1 \otimes a_{22}$, i.e., $(\Delta \otimes id)\Delta + (id \otimes \Delta)\Delta + (\tau \otimes id)(id \otimes \Delta)\Delta$. Thus (A, Δ) is an anti-Leibniz coalgebra. The proof is finished. \Box

Finally, we extend the affine anti-Leibniz algebras and the affine anti-Leibniz coalgebras to the affine anti-Leibniz bialgebra.

Definition 4.6. Let $\varpi(-,-)$ be a bilinear form on a Z-graded commutative algebra $(C = \bigoplus_{i \in \mathbb{Z}} C_i, \cdot)$. *Then*

(*i*) $\varpi(-,-)$ *is called* invariant, *if* $\varpi(c_1c_2, c_3) = \varpi(c_1, c_2c_3)$, for any $c_1, c_2, c_3 \in C$;

(*ii*) $\varpi(-,-)$ *is called* graded, *if there exists* $m \in \mathbb{Z}$ *such that* $\varpi(C_i, C_j) = 0$ *when* $i + j + m \neq 0$ *.*

A quadratic \mathbb{Z} -graded commutative algebra, *denoted by* ($C = \bigoplus_{i \in \mathbb{Z}} C_i$, ϖ), *is a* \mathbb{Z} -graded commutative *algebra together with a symmetric invariant nondegenerate graded bilinear form.*

Clearly, in Definition [4.6,](#page-25-0) if $C = C_0$, then a quadratic Z-graded commutative algebra is just a quadratic commutative algebra.

 $\bf{Example 4.7.}$ Consider the Laurent polynomial algebra $\mathbb{k}[\mathfrak{t},\mathfrak{t}^{-1}]$, there is a natural bilinear form $\varpi(-,-)$ ω ω $\mathbb{E}[t, t^{-1}]$ *which is given by* $\varpi(t^i, t^j) = \delta_{i,j}$, for $i, j \in \mathbb{Z}$, where $\delta_{i,j}$ is the Kronecker delta. It is easy to see that this bilinear form is a symmetric invariant nondegenerate graded bilinear form. Thus, $(\Bbbk [\mathfrak{t},\mathfrak{t}^{-1}],\varpi)$ *is a quadratic* Z*-graded commutative algebra, and it is called a* quadratic Z-graded Laurent polynomial algebra*.*

For a quadratic Z-graded commutative algebra ($C = \bigoplus_{i \in \mathbb{Z}} C_i$, ϖ), the nondegenerate symmetric bilinear form $\overline{\omega}(-, -)$ induces bilinear forms

$$
(\underbrace{C\,\hat{\otimes}\,C\,\hat{\otimes}\,\cdots\,\hat{\otimes}\,C}_{n\text{-fold}})\otimes(\underbrace{C\otimes C\,\otimes\cdots\otimes C}_{n\text{-fold}})\longrightarrow \Bbbk,
$$

for all $n \ge 2$, which is denoted by $\bar{\varpi}(-,-)$ and defined by

$$
\overline{\varpi}\Big(\sum_{i_1,\dots,i_n,\alpha}c'_{1,i_1,\alpha}\otimes\cdots\otimes c'_{n,i_n,\alpha},\quad c_1\otimes\cdots\otimes c_n\Big)=\sum_{i_1,\dots,i_n,\alpha}\prod_{j=1}^n\varpi(c'_{j,i_j,\alpha},\ c_j).
$$

One can check that $\bar{\varpi}(-, -)$ is *left nondegenerate*, i.e., if

$$
\bar{\varpi}\Big(\sum_{i_1,\dots,i_n,\alpha}c'_{1,i_1,\alpha}\otimes\cdots\otimes c'_{n,i_n,\alpha},\quad c_1\otimes\cdots\otimes c_n\Big)=\bar{\varpi}\Big(\sum_{i_1,\dots,i_n,\alpha}c''_{1,i_1,\alpha}\otimes\cdots\otimes c''_{n,i_n,\alpha},\quad c_1\otimes\cdots\otimes c_n\Big)
$$

for any homogeneous elements $c_1, c_2, \dots, c_n \in C$, then

$$
\sum_{i_1,\cdots,i_n,\alpha}c'_{1,i_1,\alpha}\otimes\cdots\otimes c'_{n,i_n,\alpha}=\sum_{i_1,\cdots,i_n,\alpha}c''_{1,i_1,\alpha}\otimes\cdots\otimes c''_{n,i_n,\alpha}.
$$

Lemma 4.8. *Let* (*C* = ⊕*i*∈^Z*Cⁱ* , ̟) *be a quadratic* Z*-graded commutative associative algebra. Suppose* $\bar{\Delta}: C \to C \hat{\otimes} C$ be the dual of the multiplication of C under the left nondegenerate bilinear form $\bar{\varpi}(-,-)$, *i,e.,*

(4.2)
$$
\bar{\varpi}(\bar{\Delta}(c_1), c_2 \otimes c_3) = \varpi(c_1, c_2c_3),
$$

for any $c_1, c_2, c_3 \in C$. Then $(C, \overline{\Delta})$ *is a completed cocommutative coalgebra.*

Proof. For any $c \in C$ and $c_1 \otimes c_2 \otimes c_3 \in C_i \otimes C_j \otimes C_k$, by the definition of $\overline{\Delta}$, we get

$$
\overline{\omega}((\overline{\Delta}\hat{\otimes} id)(\overline{\Delta}(c)), c_1 \otimes c_2 \otimes c_3) = \omega(c, (c_1c_2)c_3)) = \omega(c, c_1(c_2c_3))
$$

= $\overline{\omega}((id \hat{\otimes} \overline{\Delta})(\overline{\Delta}(c)), c_1 \otimes c_2 \otimes c_3).$

That is, $(\bar{\Delta}\hat{\otimes} \mathrm{id})\bar{\Delta} = (\mathrm{id}\hat{\otimes}\bar{\Delta})\bar{\Delta}$. Similarly, one can check that $\bar{\Delta} = \hat{\tau}\bar{\Delta}$. Thus $(C, \bar{\Delta})$ is a completed cocommutative coalgebra.

Example 4.9. *Consider the quadratic* Z-graded commutative Laurent polynomial algebra ($\mathbb{K}[\mathfrak{t},\mathfrak{t}^{-1}], \varpi$) π *which is given in Example* [4.7](#page-25-1), by Lemma [4.8](#page-26-0), we get a completed cocommutative coalgebra (\k[t, t⁻¹], $\bar{\Delta}$), *where* $\overline{\Delta}$ *is given by:* $\overline{\Delta}(t^k) = \sum_{i \in \mathbb{Z}} t^i \otimes t^{k-i}$ *for all* $k \in \mathbb{Z}$ *. This completed cocommutative coalgebra* $(k[t, t^{-1}], \bar{\Delta})$ is just the Laurent polynomial coalgebra given in Example [4.4](#page-24-1).

We now give the notion and results on completed anti-Leibniz bialgebras.

Definition 4.10. *A* completed anti-Leibniz bialgebra *is a triple* $(A = \bigoplus_{i \in \mathbb{Z}} A_i, \cdot, \hat{\Delta})$ *such that* $(A = \bigoplus_{i \in \mathbb{Z}} A_i, \cdot)$ *is a* Z*-graded anti-Leibniz algebra,* (*A* = ⊕*i*∈^Z*Aⁱ* , ∆ˆ) *is a completed anti-Leibniz coalgebra, and the following compatibility condition holds:*

$$
(\mathbf{r}_A(a_1) \hat{\otimes} \text{ id})\hat{\Delta}(a_2) - \hat{\tau}((\mathbf{r}_A(a_2) \hat{\otimes} \text{ id})\hat{\Delta}(a_1)) = 0.
$$

 $\hat{\Delta}(a_1a_2) + ((id \hat{\otimes} \mathbf{r}_A(a_2) + (\mathbf{l}_A - \mathbf{r}_A)(a_2) \hat{\otimes} id)(id \hat{\otimes} id - \hat{\tau}))\hat{\Delta}(a_1) + (id \hat{\otimes} \mathbf{l}_A(a_1) + \mathbf{l}_A(a_1) \hat{\otimes} id)\hat{\Delta}(a_2) = 0,$

for any $a_1, a_2 \in A$.

In the definition above, if $A = A_0$, the completed anti-Leibniz bialgebra is just the anti-Leibniz bialgebra given in Definition [2.11.](#page-5-4)

Theorem 4.11. *Let* (A, \cdot, Δ) *be a finite-dimensional anti-Leibniz bialgebra and* $(C = \bigoplus_{i \in \mathbb{Z}} C_i, \omega)$ *be a quadratic* ℤ-graded commutative associative algebra. Suppose that $(A, ∘)$ *is the* ℤ-graded anti-Leibniz a *lgebra from* (A, \cdot) *by* $(C = \bigoplus_{i \in \mathbb{Z}} C_i, \cdot)$ *in Lemma* $\overline{4.2}$ $\overline{4.2}$ $\overline{4.2}$, $\overline{\Delta}$: $C \to C \hat{\otimes} C$ *is the linear map defined by Eq.* [\(4.2\)](#page-26-1) *and* ∆ˆ : A → A ⊗ Aˆ *is the linear map defined by Eq.* [\(4.1\)](#page-24-0)*. Then* (A, ⋄, ∆ˆ) *is a completed anti-Leibniz bialgebra, which is called an* affine anti-Leibniz bialgebra *from* (A, \cdot, Δ) *by* $(C = \bigoplus_{i \in \mathbb{Z}} C_i, \omega)$ *.*

Furthermore, if A is finite-dimensional and $(C = \bigoplus_{i \in \mathbb{Z}} C_i, \varpi)$ *is the quadratic* Z-graded Laurent *polynomial algebra* (k[t, t⁻¹], *ω*), then (A, \circ , $\hat{\Delta}$) is a completed anti-Leibniz bialgebra constructed as *above if and only if* (*A*, ·, ∆) *is an anti-Leibniz bialgebra.*

Proof. First, since the bilinear form $\varpi(-,-)$ on *C* is invariant and (C, \cdot) is an associative algebra, for any $c, c', x, y \in C$, we have

$$
\overline{\varpi}\Big(\sum_{i,j,\alpha}c'_{1,i,\alpha}c\otimes c'_{2,j,\alpha},\ x\otimes y\Big)=\overline{\varpi}\Big(\sum_{i,j,\alpha}c'_{1,i,\alpha}\otimes c'_{2,j,\alpha},\ cx\otimes y\Big)
$$

$$
=\varpi(c',\ cxy)=\varpi(c'c,\ xy)=\overline{\varpi}(\overline{\Delta}(c'c),\ x\otimes y).
$$

That is, $\bar{\Delta}(c'c) = \sum_{i,j,\alpha} c'_1$ $C'_{1,i,\alpha}c \otimes c'_2$ $Z_{2,i,\alpha}$. Note that (C, \cdot) is commutative and $\overline{\Delta}$ is cocommutative, we have $\bar{\Delta}(c'c) = \bar{\Delta}(cc') = \sum_{i,j,\alpha} c'_{1,i,\alpha} \otimes cc'_{2,j,\alpha} = \sum_{i,j,\alpha} c_{1,i,\alpha}c' \otimes c_{2,j,\alpha} = \sum_{i,j,\alpha} cc'_{1,i,\alpha} \otimes c'_{2,j,\alpha} = \sum_{i,j,\alpha} c'_{1,i,\alpha} \otimes c'_{2,j,\alpha}$ $\int_{2,j,\alpha}^{\prime} c =$ $\sum_{i,j,\alpha} c_{2,j,\alpha} c' \otimes c_{1,i,\alpha} = \sum_{i,j,\alpha} c_{2,j,\alpha} \otimes c' c_{1,i,\alpha}$. Thus, for any $a \otimes c, a' \otimes c' \in \mathcal{A}$,

$$
(\mathbf{r}_{\mathcal{A}}(a \otimes c) \hat{\otimes} \text{id})\hat{\Delta}(a' \otimes c') - \hat{\tau}((\mathbf{r}_{\mathcal{A}}(a' \otimes c') \hat{\otimes} \text{id})\hat{\Delta}(a \otimes c))
$$

=
$$
\sum_{(a')} \sum_{i,j,a} (a'_{(1)}a \otimes a'_{(2)}) \bullet (c'_{1,i,a}c \otimes c'_{2,j,a}) - \sum_{(a)} \sum_{i,j,a} (a_{(2)} \otimes a_{(1)}a') \bullet (c_{2,j,a} \otimes c_{1,i,a}c')
$$

=
$$
((\mathbf{r}_{A}(a) \otimes \text{id})\Delta(a') - \tau((\mathbf{r}_{A}(a') \otimes \text{id})\Delta(a))) \bullet \bar{\Delta}(cc')
$$

= 0,

and

$$
\hat{\Delta}((a \otimes c) \diamond (a' \otimes c')) + ((id \hat{\otimes} r_{\mathcal{A}}(a' \otimes c') + (l_{\mathcal{A}} - r_{\mathcal{A}})(a' \otimes c') \hat{\otimes} id)(id \hat{\otimes} id - \hat{\tau}))\hat{\Delta}(a \otimes c)
$$
\n
$$
+ (id \hat{\otimes} l_{\mathcal{A}}(a \otimes c) + l_{\mathcal{A}}(a \otimes c) \hat{\otimes} id)\hat{\Delta}(a' \otimes c')
$$
\n
$$
= \Delta(aa') \bullet \bar{\Delta}(cc') + \sum_{(a)} \sum_{i,j,a} (a_{(1)} \otimes a_{(2)}a') \bullet (c_{1,i,a} \otimes c_{2,j,a}c') + \sum_{(a)} \sum_{i,j,a} (a'a_{(1)} \otimes a_{(2)}) \bullet (c'c_{1,i,a} \otimes c_{2,j,a})
$$
\n
$$
- \sum_{(a)} \sum_{i,j,a} (a_{(1)}a' \otimes a_{(2)}) \bullet (c_{1,i,a}c' \otimes c_{2,j,a}) - \sum_{(a)} \sum_{i,j,a} (a_{(2)} \otimes a_{(1)}a') \bullet (c_{2,i,a} \otimes c_{1,j,a}c')
$$
\n
$$
- \sum_{(a)} \sum_{i,j,a} (a'a_{(2)} \otimes a_{(1)}) \bullet (c'c_{2,i,a} \otimes c_{1,j,a}) + \sum_{(a)} \sum_{i,j,a} (a_{(2)}a' \otimes a_{(1)}) \bullet (c_{2,i,a}c' \otimes c_{1,j,a})
$$
\n
$$
+ \sum_{(a')} \sum_{i,j,a} (a'_{(1)} \otimes a a'_{(2)}) \bullet (c'_{1,i,a} \otimes c'_{2,j,a}) + \sum_{(a')} \sum_{i,j,a} (a a'_{(1)} \otimes a'_{(2)}) \bullet (c c'_{1,i,a} \otimes c'_{2,j,a})
$$
\n
$$
= (\Delta(aa') + ((id \otimes r_{A}(a') - r_{A}(a') \otimes id + l_{A}(a') \otimes id)(id \otimes id - \tau))\Delta(a)) \bullet \bar{\Delta}(cc')
$$
\n
$$
+ ((id \otimes l_{A}(a) + l_{A}(a) \otimes id)\Delta(a')) \bullet \bar{\Delta}(cc')
$$

since (A, \cdot, Δ) is an anti-Leibniz bialgebra. By the definition, we get that $(\mathcal{A}, \diamond, \hat{\Delta})$ is a completed anti-Leibniz bialgebra.

Second, if $(\mathcal{A}, \diamond, \hat{\Delta})$ is a completed anti-Leibniz bialgebra, where $(C = \bigoplus_{i \in \mathbb{Z}} C_i, \omega)$ is the quadratic Z-graded Laurent polynomial algebra (k[t, t⁻¹], ϖ), denote $a \otimes t^k$ by at^k for $a \in A$, then for any $a, a' \in A$ and $k, l \in \mathbb{Z}$, we have

$$
0 = (\mathbf{r}_{\mathcal{A}}(at^k) \hat{\otimes} \text{id}) \hat{\Delta}(a't^l) - \hat{\tau}((\mathbf{r}_{\mathcal{A}}(a't^l) \hat{\otimes} \text{id}) \hat{\Delta}(at^k))
$$

= ((\mathbf{r}_A(a) \otimes \text{id})\Delta(a') - \tau((\mathbf{r}_A(a') \otimes \text{id})\Delta(a))) \bullet \bar{\Delta}(t^{k+l}).

Thus, we get Eq. (2.13) holds. Similarly, we can get Eq. (2.12) also holds. That is, (A, \cdot, Δ) is an anti-Leibniz bialgebra.

Example 4.12. (*I*) *Let* (Λ_1^2) 1 , ∆) *be the* 2*-dimension anti-Leibniz bialgebra given in Example* [2.12](#page-5-3)*. Con*sider the Z-graded vector space $\Lambda_1^2 \otimes \Bbbk[t, t^{-1}]$, we denote the homogeneous element in $\Lambda_1^2 \otimes \Bbbk[t, t^{-1}]$ by a^{t} , where $a \in \Lambda^2$. Then the multiplication \circ on $\Lambda^2_1 \otimes \mathbb{k}[t, t^{-1}]$ is given by $e_1 t^i \circ e_1 t^j = e_2 t^{i+j}$, and the completed comultiplication $\hat{\Delta}$ on $\Lambda_1^2 \otimes \Bbbk[t, t^{-1}]$ is given by $\hat{\Delta}(e_1 t^k) = \sum_{i \in \Bbb Z} e_2 t^i \otimes e_2 t^{k-i}$, for any i, j, $k \in \Bbb Z$. *This anti-Leibniz bialgebra is also a Leibniz bialgebra, is also an anti-associative bialgebra (see* [\[15\]](#page-28-23)*).*

(*II*) Let $(A = \mathbb{k}\{e_1, e_2, e_3\}, \cdot, \Delta$ be an anti-Leibniz bialgebra with nonzero multiplication and comul*tiplication:* $e_2e_2 = e_1 = e_3e_3$ *and* $\Delta(e_2) = e_1 \otimes e_1$ *. Then we get an infinite-dimensional anti-Leibniz bialgebraic structure on* $A \otimes k[t, t^{-1}]$ *under the multiplication* $e_2t^i \circ e_2t^j = e_1t^{i+j} = e_3t^i * e_3t^j$ *and com* p *leted comultiplication* $\hat{\Delta}(e_2 t^k) = \sum_{i \in \mathbb{Z}} e_1 t^i \otimes e_1 t^{k-i}$, for any $i, j, k \in \mathbb{Z}$.

Remark 4.13. *In this section, we using the* Z*-graded commutative associative algebra to construct infinite-dimensional anti-Leibniz bialgebra. In fact, we can using the* Z*-graded perm algebra to construct infinite-dimensional anti-Leibniz bialgebra. Recently, Lin, Zhou and Bai construct a* Z*-graded*

perm algebra in [\[21\]](#page-28-24) *as follows. Let* $P = \{f_1 \partial_1 + f_2 \partial_2 \mid f_1, f_2 \in \mathbb{R}[\mathbf{x}_1^{\pm}]$ x_1^{\pm} , x_2^{\pm} 2]} *and define a binary operation* · : *P* ⊗ *P* → *P by*

$$
(x_1^{i_1}x_2^{i_2}\partial_s)\cdot (x_1^{j_1}x_2^{j_2}\partial_t)=\delta_{s,1}x_1^{i_1+j_1+1}x_2^{i_2+j_2}\partial_t+\delta_{s,2}x_1^{i_1+j_1}x_2^{i_2+j_2+1}\partial_t,
$$

for any $i_1, i_2, j_1, j_2 \in \mathbb{Z}$ and $s, t \in \{1, 2\}$. Then (P, \cdot) *is a* Z-graded perm algebra with the linear decom*position* $P = \bigoplus_{i \in \mathbb{Z}} P_i$, where

$$
P_i = \Big\{ \sum_{k=1}^2 f_k \partial_k \mid f_k \text{ is a homogeneous polynomial with } \deg(f_k) = i - 1, \ k = 1, 2 \Big\},
$$

for all i ∈ Z*. If we replace the Laurent polynomial algebra with this* Z*-graded perm algebra, the conclusions of this chapter can also be obtained.*

Acknowledgements. This work was financially supported by National Natural Science Foundation of China (No. 11801141).

REFERENCES

- [1] M. Aguiar, *On the associative analog of Lie bialgebras*, J. Algebra 244 (2001), 492–532.
- [2] S. Ayupov, B. Omirov, I. Rakhimov, *Leibniz algebras: structure and classification*, CRC Press, Taylor & Francis Group, 2020.
- [3] C. Bai, *Left-symmetric bialgebras and an analogue of the classical Yang-Baxter equation*, Commun. Contemp. Math. 10 (2008), 221–260.
- [4] C. Bai, *Double constructions of Frobenius algebras, Connes cocycles and their duality*, J. Noncommut. Geom. 4 (2010), 475–530.
- [5] C. Bai, L. Guo, X. Ni, O*-operators on associative algebras and associative Yang-Baxter equations,* Pacific J. Math. 256 (2012), 257–289.
- [6] C. Bai, G. Liu, Y. Sheng, R. Tang, *Quasi-triangular, factorizable Leibniz bialgebras and relative Rota-Baxter operators,* arXiv: 2410.03089.
- [7] A. Baklouti, S. Benayadi, *Symplectic Jacobi-Jordan algebras*, Linear and Multilinear Algebra 69 (2019), 1–22.
- [8] A. Baklouti, S. Benayadi, A. Makhlouf and S. Mansour, *Cohomology and deformations of Jacobi-Jordan algebras*, [arXiv:2109.12364.](http://arxiv.org/abs/2109.12364)
- [9] S. Braiek, T. Chtioui, S. Mabrouk, *Anti-Leibniz algebras: A non-commutative version of mock-Lie algebras*, J. Geom. Phys. 209 (2025), 105385, 15PP.
- [10] K. Benali, *Bialgebras via Manin triples of compatible mock-Lie algebras*, Comm. Algebra *52* (2024), 1–14.
- [11] K. Benali, T. Chtioui, A. Hajjaji, S. Mabrouk, *Bialgebras, the Yang-Baxter equation and Manin triples for mock-Lie algebras*, Acta et Commentationes Universitatis Tartuensis de Mathematica 27 (2023), 211–233.
- [12] D. Burde, A. Fialowski, *Jacobi-Jordan algebras*, Linear Algebra Appl. 459 (2014), 586–594.
- [13] C. Cuvier, *Algèbres de Leibnitz: définitions, propriétés*, Ann. Sci. Ecole Norm. Sup. 27 (4) (1994), 1–45.
- [14] V.G. Drinfeld, *Hamiltonian structures of Lie groups, Lie bialgebras and the geometric meaning of the classical Yang-Baxter equations*, Soviet Math. Dokl. 27 (1983), 68–71.
- [15] B. Hou, Z. Cui, *A bialgebra theory of anti-associative algebras*, in preparation.
- [16] B. Hou, Z. Cui, *Factorizable mock-Lie bialgebras and Factorizable Rota-Baxter mock-Lie bialgebras*, in preparation.
- [17] H. Lang, Y. Sheng, *Factorizable Lie bialgebras, quadratic Rota-Baxter Lie algebras and Rota-Baxter Lie bialgebras*, Comm. Math. Phy. 397(2) (2023), 763–791.
- [18] I. Laraiedh, *Rota-Baxter mock-Lie bialgebras and related structures*, Int. J. Geom. Methods Mod. Phys. 21 (2024), 79–96.
- [19] I. Laraiedh, *Di*ff*erential mock-Lie bialgebras and admissible mock-Lie Yang-Baxter equations*, J. Geom. Phys. 198 (2024), 105135, 16PP.
- [20] H. Li, T. Ma, S. Wang, *When Leibniz algebras are Nijenhuis?*, arXiv: 2310.14267.
- [21] Y. Lin, P. Zhou, C. Bai, *Infinite-dimensional Lie bialgebras via a*ffi*nization of perm bialgebras and pre-Lie bialgebras*, J. Algebra 663 (2025), 210–258.
- [22] G. Liu, C. Bai, *A bialgebra theory for transposed Poisson algebras via anti-pre-Lie bialgebras and anti-pre-Lie-Poisson bialgebras*, Commun. Contemp. Math. 26 (2024), 2350050, 49pp.
- [23] J.L. Loday, *Une version non commutative des alg`ebres de Lie: les alg`ebres de Leibniz*, Enseign. Math. 39 (1993), 269–293.
- [24] X. Ni, C. Bai, *Poisson bialgebras*, J. Math. Phys. 54 (2013), 023515, 14pp.
- [25] N. Reshetikhin, M.A. Semenov-Tian-Shansky, *Quantum R-matrices and factorization problems*, J. Geom. Phys. 5 (1988), 533–550.
- [26] Y. Sheng, Y. Wang, *Quasi-triangular and factorizable antisymmetric infinitesimal bialgebras*, J. Algebra 628 (2023), 415–433.
- [27] M.A. Semenov-Tian-Shansky, *Integrable systems and factorization problems*, Oper. Theory, Adv. Appl. 141 (2003), 155–218.
- [28] R. Tang, Y. Sheng, *Leibniz bialgebras, relative Rota-Baxter operators, and the classical Leibniz Yang-Baxter equation*, J. Noncommut. Geom. 16 (2022), 1179–1211.
- [29] V.N. Zhelyabin, *Jordan bialgebras and their relation to Lie bialgebras*, Algebra Logic, 36 (1997), 1–15.
- [30] P. Zusmanovich, *Special and exceptional mock-Lie algebras*, Linear Algebra Appl. 518 (2017), 79–96.

SCHOOL OF MATHEMATICS AND STATISTICS, HENAN UNIVERSITY, KAIFENG 475004, CHINA *Email address*: bohou1981@163.com

SCHOOL OF MATHEMATICS AND STATISTICS, HENAN UNIVERSITY, KAIFENG 475004, CHINA *Email address*: czp15824833068@163.com