

MAIN CONJECTURES FOR NON-CM ELLIPTIC CURVES AT GOOD ORDINARY PRIMES

XIAOJUN YAN AND XIUWU ZHU

ABSTRACT. Let E/\mathbb{Q} be an elliptic curve and $p > 2$ be a prime of good ordinary reduction for E . Assume that the residue representation associated with (E, p) is irreducible. In this paper, we prove more cases on several Iwasawa main conjectures for E . As applications, we prove more general cases of p -converse theorem and p -part BSD formula when the rank is less than or equal to 1.

CONTENTS

1. Introduction	1
1.1. Main result	2
1.2. Strategy	3
1.3. Acknowledgment	3
2. Selmer groups	3
2.1. Selmer groups for Iwasawa algebras	3
2.2. Selmer groups for Hida families	5
3. p -adic L -functions	6
3.1. A three variable p -adic L -function	6
3.2. Two variable p -adic L -functions: type I	7
3.3. Cyclotomic p -adic L -function	7
3.4. Two variable p -adic L -functions: type II	8
3.5. BDP p -adic L -function	8
4. Main conjectures	8
4.1. Two variable Iwasawa main conjectures	8
4.2. A three variable Iwasawa main conjecture	9
4.3. Proof of Theorem 4.2	10
4.4. Mazur's main conjecture	10
4.5. Anticyclotomic main conjectures	10
4.6. Applications	11
Conflict of interest	12
References	12

1. INTRODUCTION

Let E be an elliptic curve over \mathbb{Q} . Fix an odd prime p and embeddings $\iota_p : \bar{\mathbb{Q}} \hookrightarrow \bar{\mathbb{Q}}_p$, $\iota_\infty : \bar{\mathbb{Q}} \hookrightarrow \mathbb{C}$. Let $T_p E$ be the p -adic Tate module and $\rho_E : G_{\mathbb{Q}} \rightarrow \text{Aut}_{\mathbb{Z}_p}(T_p E)$ the associated p -adic Galois representation. Suppose that E has good ordinary reduction at p .

Let K be an imaginary quadratic field. Suppose that $p = \mathfrak{p}\bar{\mathfrak{p}}$ splits in K , where \mathfrak{p} is the prime induced by ι_p . Let K_∞^+ , K_∞^- be the cyclotomic, respectively, anticyclotomic \mathbb{Z}_p -extension of K , and $K_\infty = K_\infty^+ K_\infty^-$. Then we have the Iwasawa algebras

$$\Lambda_K^\pm = \mathbb{Z}_p[[\text{Gal}(K_\infty^\pm/K)]], \quad \Lambda_K = \mathbb{Z}_p[[\text{Gal}(K_\infty/K)]].$$

As in Castella-Grossi-Skinner [CGS23], we have

- (1) the two-variable p -adic L -function $\mathcal{L}_p^{\text{PR}}(E/K) \in \Lambda_K$, which interpolates $L(E/K, \chi^{-1}, 1)$ for finite order characters χ of $\text{Gal}(K_\infty/K)$.
- (2) the Λ_K -module Selmer group $H_{\mathcal{F}_{\text{ord}}}^1(K, T_p E \otimes \Lambda_K^\vee)$ with ordinary Selmer conditions at $v|p$, where $(\cdot)^\vee$ is the Pontryagin dual.

Also, we have

- (1) another type two-variable p -adic L -function $\mathcal{L}_p^{\text{Gr}}(E/K) \in \Lambda_K^{\text{ur}} := \Lambda_K \widehat{\otimes}_{\mathbb{Z}_p} \mathbb{Z}_p^{\text{ur}}$, which interpolates roughly $L(E/K, \chi, 1)$ for χ Hecke characters over K with infinity type (b, a) , $a \leq -1$, $b \geq 1$, where \mathbb{Z}_p^{ur} is the completion of the ring of integers of the maximal unramified extension of \mathbb{Q}_p .
- (2) the Λ_K -module Selmer group $H_{\mathcal{F}_{\text{Gr}}}^1(K, T_p E \otimes \Lambda_K^\vee)$ with relaxed Selmer conditions at $v = \mathfrak{p}$ and strict Selmer conditions at $v = \bar{\mathfrak{p}}$.

Conjecture 1.1. *Suppose that the residue representation $\bar{\rho}_E|_{G_K}$ is irreducible.*

- (1) $H_{\mathcal{F}_{\text{ord}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee$ is Λ_K -torsion and

$$\text{Char}_{\Lambda_K}(H_{\mathcal{F}_{\text{ord}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee) = (\mathcal{L}_p^{\text{PR}}(E/K)).$$
- (2) $H_{\mathcal{F}_{\text{Gr}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee$ is Λ_K -torsion and

$$\text{Char}_{\Lambda_K}(H_{\mathcal{F}_{\text{Gr}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee) \Lambda_K^{\text{ur}} = (\mathcal{L}_p^{\text{Gr}}(E/K)).$$

1.1. Main result.

Theorem 1.2. *If the Heegner hypothesis holds (in particular, $\text{sign}(E/K) = -1$) and $\bar{\rho}_E|_{G_K}$ is absolutely irreducible, then*

- (1) $H_{\mathcal{F}_{\text{ord}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee$ is Λ_K -torsion and

$$\text{Char}_{\Lambda_K}(H_{\mathcal{F}_{\text{ord}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee) \subset (\mathcal{L}_p^{\text{PR}}(E/K)).$$
- (2) $H_{\mathcal{F}_{\text{Gr}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee$ is Λ_K -torsion and

$$\text{Char}_{\Lambda_K}(H_{\mathcal{F}_{\text{Gr}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee) \Lambda_K^{\text{ur}} \subset (\mathcal{L}_p^{\text{Gr}}(E/K)).$$

Moreover, if

(Im) *there exists $\tau \in \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}(\mu_{p^\infty}))$ such that $T_p E/(\rho_E(\tau) - 1)T_p E$ is free of \mathbb{Z}_p -rank one holds, the Conjecture 1.1 is true.*

Skinner-Urban [SU14] first proved Conjecture 1.1 (1) used Eisenstein congruences on $GU(2, 2)$ under certain conditions; particularly under the assumption that the sign is $+1$. Wan [Wan20, Wan21] proved Conjecture 1.1 (2) using Eisenstein congruences on $GU(3, 1)$ under different conditions, specifically under the assumption that the sign is -1 and E is semistable. In [BSTW24], using the Beilinson-Flach elements and explicit reciprocity law, Burungale-Skinner-Tian-Wan proved the equivalence between Conjecture 1.1 (1) and 1.1 (2).

In this paper, our proof of Theorem 1.2 is based on Skinner-Urban's work and a simple observation. The other ingredients are the equivalence of Conjecture 1.1 (1) and 1.1 (2), and Hsieh's result on non-vanishing of the μ -invariants of BDP p -adic L -function [Hsi14].

We observe that, Skinner-Urban actually proved that the left-hand side of Conjecture 1.1 (1) is contained in right-hand side, after tensoring the fractional field of Λ_K^+ . By [BSTW24], this shows that the left hand side of Conjecture 1.1 (2) is contained in the right hand side, after tensoring the fractional field of Λ_K^+ . Therefore, by [Hsi14], if the Heegner hypothesis holds, the inclusion relation in Conjecture 1.1 (2) holds in general. Then as [SU14], by using Kato's result on Mazur's main conjecture [Kat04], we could complete the proof under the condition (Im).

Remark 1.3. Recently, using Wan's results on main conjecture for Hilbert modular forms [Wan15] and the base change method, Burungale-Castella-Skinner [BCS24] proved the rational version of Conjecture 1.1 assuming $p > 3$ and $\bar{\rho}_E$ is irreducible, and the integral version if (Im) also holds. Their results don't cover ours, nor do ours cover theirs.

As an application, we prove more cases of several one variable main conjectures, the p -part of the BSD formula and the p -converse theorem.

Corollary 1.4. *Let E/\mathbb{Q} be an elliptic curve of conductor N , $p \nmid 2N$ a prime. Assume that E has ordinary reduction at p , if $r \leq 1$, the following are equivalent*

- (1) $\text{corank}_{\mathbb{Z}_p} \text{Sel}_{p^\infty}(E/\mathbb{Q}) = r$,
- (2) $\text{ord}_{s=1} L(E/\mathbb{Q}, s) = r$.

Under any of the above conditions, if (Im) also holds, then the p -part of the BSD formula for E holds.

Remark 1.5. There have been several works on the p -part of the BSD formula and the p -converse theorem for elliptic curves. For example, in our case, see [SU14], [Zha14], [JSW17], [BSTW24], [BCS24] for some previous works on p -part BSD formula, and [Ski20], [Zha14], [BSTW24], [BCGS23] for some previous works on p -converse theorem.

1.2. Strategy. By [CGS23, Proposition 3.2.1] or [BSTW24, Proposition 9.18], the use of Beilinson-Flach elements, combined with the reciprocity law, establishes a connection between different Main Conjectures. Consequently, the inclusion relations (and their opposites) in Theorem 1.2 (1) and (2) are shown to be equivalent.

By Skinner-Urban [SU14], especially Theorem 7.7 and Proposition 13.6 (1), we have

Theorem 1.6. *Suppose that the residue representation $\bar{\rho}_E : G_{\mathbb{Q}} \rightarrow \text{Aut}(E[p])$ is irreducible. Then for any height one prime P ,*

$$\text{ord}_P(\text{Char}_{\Lambda_K} H_{\mathcal{F}_{\text{ord}}}^1(K, T_p E \otimes \Lambda_K^{\vee})^{\vee}) \geq \text{ord}_P(\mathcal{L}_p^{\text{PR}}(E/K))$$

unless $P = P^+ \Lambda_K$ for some $P^+ \subset \Lambda_K^+$.

Similarly, as [CGS23, Proposition 3.2.1], we have

Theorem 1.7. *Under the assumption of above theorem, we have that for any height one prime P ,*

$$\text{ord}_P(\text{Char}_{\Lambda_K^{\text{ur}}} H_{\mathcal{F}_{\text{Gr}}}^1(K, T_p E \otimes \Lambda_K^{\vee})^{\vee}) \geq \text{ord}_P(\mathcal{L}_p^{\text{Gr}}(E/K))$$

unless $P = P^+ \Lambda_K^{\text{ur}}$ for some $P^+ \subset \Lambda_K^{\text{ur},+}$.

However, by the result on the μ -invariant of the BDP p -adic L -function [Hsi14], we have that if $P \subset \Lambda_K^{\text{ur}}$ is a height one primes of the form $P = P^+ \Lambda_K^{\text{ur}}$ for some $P^+ \subset \Lambda_K^{\text{ur},+}$, then

$$\text{ord}_P(\mathcal{L}_p^{\text{Gr}}(E/K)) = 0.$$

Hence, Theorem 1.2 (2) holds, implying that Theorem 1.2 (1) also holds. Moreover, if (Im) is satisfied, then, similarly to [SU14, Theorem 3.30], we conclude that Conjecture 1.1 (2) is true, and consequently, so is Conjecture 1.1 (1).

1.3. Acknowledgment. The authors are grateful to Ye Tian and Wei He for their continuous encouragement and many helpful communications and to Ashay Burungale for his valuable comments.

2. SELMER GROUPS

Let K be an imaginary quadratic field of discriminant D_K . Let $p > 2$ be a prime such that $p = \mathfrak{p}\bar{\mathfrak{p}}$ splits in K . Let \mathbb{Q}_{∞} be the \mathbb{Z}_p -extension of \mathbb{Q} . Let K_{∞} be the \mathbb{Z}_p^2 -extension of K , and K_{∞}^+ (respectively, K_{∞}^-) be the cyclotomic (respectively, anticyclotomic) \mathbb{Z}_p -extension of K . Let

$$\Gamma_{\mathbb{Q}} := \text{Gal}(\mathbb{Q}_{\infty}/\mathbb{Q}), \quad \Gamma_K := \text{Gal}(K_{\infty}/K), \quad \Gamma_K^{\pm} := \text{Gal}(K_{\infty}^{\pm}/K).$$

Then $\text{Gal}(K_{\infty}^+/K)$ is identified with $\text{Gal}(\mathbb{Q}_{\infty}/\mathbb{Q})$. Let γ^{\pm} be a topological generator of Γ_K^{\pm} . Let

$$\Lambda_{\mathbb{Q}} := \mathbb{Z}_p[[\Gamma_{\mathbb{Q}}]], \quad \Lambda_K := \mathbb{Z}_p[[\Gamma_K]], \quad \Lambda_K^{\pm} := \mathbb{Z}_p[[\Gamma_K^{\pm}]]$$

be the corresponding Iwasawa algebras, and

$$\varepsilon_{\mathbb{Q}} : G_{\mathbb{Q}} \twoheadrightarrow \Gamma_{\mathbb{Q}} \hookrightarrow \Lambda_{\mathbb{Q}}^{\times}, \quad \varepsilon_K : G_K \twoheadrightarrow \Gamma_K \hookrightarrow \Lambda_K^{\times}, \quad \varepsilon_{K,\pm} : G_K \twoheadrightarrow \Gamma_K^{\pm} \hookrightarrow \Lambda_K^{\pm,\times}$$

the natural characters. For a discrete \mathbb{Z}_p -module M , let $M^{\vee} := \text{Hom}_{\text{cts}}(M, \mathbb{Q}_p/\mathbb{Z}_p)$ be its Pontryagin dual. The module $\Lambda_{\mathbb{Q}}^{\vee}$ is equipped with a $G_{\mathbb{Q}}$ -action via $\varepsilon_{\mathbb{Q}}^{-1}$. Similarly, the modules Λ_K^{\vee} and $\Lambda_K^{\pm,\vee}$ are equipped with G_K -actions.

We normalize the reciprocity map in class field theory using the arithmetic Frobenius. Specifically, we require that uniformizers are mapped to the arithmetic Frobenius via the reciprocity map. In this way, we identify Hecke characters with Galois characters in this paper.

2.1. Selmer groups for Iwasawa algebras. Let E/\mathbb{Q} be an elliptic curve of conductor N prime to D_K . Assume that E has good ordinary reduction at p .

Discrete Selmer groups. Let F be \mathbb{Q} or K , w a prime of F above p , and \mathbb{F}_w the residue field at w . Let $\tilde{E}_{/\mathbb{F}_w}$ be the reduction of E at w , and denote $\mathcal{F}_w^+ T_p E := \ker(T_p E \rightarrow T_p \tilde{E}_{/\mathbb{F}_w})$ as the kernel of the reduction map.

Definition 2.1. *For Λ any one of $\Lambda_{\mathbb{Q}}, \Lambda_K$ or Λ_K^{\pm} , we define*

- (1) $H_{\text{ord}}^1(F_w, T_p E \otimes \Lambda^{\vee}) = \text{Im}(H^1(F_w, \mathcal{F}_w^+ T_p E \otimes \Lambda^{\vee}) \rightarrow H^1(F_w, T_p E \otimes \Lambda^{\vee}))$,
- (2) $H_{\text{rel}}^1(F_w, T_p E \otimes \Lambda^{\vee}) = H^1(F_w, T_p E \otimes \Lambda^{\vee})$,
- (3) $H_{\text{str}}^1(F_w, T_p E \otimes \Lambda^{\vee}) = 0$.

Let Σ be a set of places of F such that Σ contains all places of F dividing $pN\infty$. In the anticyclotomic case, assume moreover that every finite place in Σ splits in K . Let F_Σ be the maximal extension of F unramified outside Σ , and let $G_{F,\Sigma} := \text{Gal}(F_\Sigma/F)$.

In the case $F = \mathbb{Q}$, for $a \in \{\text{ord}, \text{str}, \text{rel}\}$ and $M = T_p E \otimes \Lambda_{\mathbb{Q}}^\vee$, let

$$H_{\mathcal{F}_a}^1(\mathbb{Q}, M) = \text{Ker} \left(H^1(G_{\mathbb{Q},\Sigma}, M) \rightarrow \prod_{q \in \Sigma, q \nmid p} H^1(\mathbb{Q}_q, M) \times \frac{H^1(\mathbb{Q}_p, M)}{H_a^1(\mathbb{Q}_p, M)} \right),$$

and define

$$\mathcal{X}_{\mathcal{F}_a}(E/\mathbb{Q}_\infty) = H_{\mathcal{F}_a}^1(\mathbb{Q}, T_p E \otimes \Lambda_{\mathbb{Q}}^\vee)^\vee.$$

In the case $F = K$, for $a, b \in \{\text{ord}, \text{str}, \text{rel}\}$, $\Lambda \in \{\Lambda_K, \Lambda_K^-, \Lambda_K^+\}$, and $M = T_p E \otimes \Lambda^\vee$, let

$$H_{\mathcal{F}_{a,b}}^1(K, M) = \text{Ker} \left(H^1(G_{K,\Sigma}, M) \rightarrow \prod_{q \in \Sigma, q \nmid p} H^1(K_q, M) \times \frac{H^1(K_{\mathfrak{p}}, M)}{H_a^1(K_{\mathfrak{p}}, M)} \times \frac{H^1(K_{\bar{\mathfrak{p}}}, M)}{H_b^1(K_{\bar{\mathfrak{p}}}, M)} \right).$$

For simplicity, we write

- (1) $H_{\mathcal{F}_a}^1(K, M) = H_{\mathcal{F}_{a,a}}^1(K, M)$ for $a \in \{\text{ord}, \text{str}, \text{rel}\}$,
- (2) $H_{\text{Gr}}^1(K, M) = H_{\mathcal{F}_{\text{rel}, \text{str}}}^1(K, M)$,

And for $a \in \{\text{ord}, \text{str}, \text{rel}, \text{Gr}\}$, we define

$$\mathcal{X}_{\mathcal{F}_a}(E/K_\infty) = H_{\mathcal{F}_a}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee, \quad \mathcal{X}_{\mathcal{F}_a}(E/K_\infty^\pm) = H_{\mathcal{F}_a}^1(K, T_p E \otimes \Lambda_K^{\pm, \vee})^\vee.$$

For any prime \mathfrak{q} of K , we denote the inertia subgroup of $G_{K_{\mathfrak{q}}}$ by $I_{\mathfrak{q}}$.

Lemma 2.2. *For any $\mathfrak{q} \nmid p$, we have*

- (1) $H^1(G_{K_{\mathfrak{q}}}/I_{\mathfrak{q}}, (T_p E \otimes \Lambda_K^\vee)^{I_{\mathfrak{q}}}) = 0$.
- (2) $\text{Char}_{\Lambda_K}((H^1(I_{\mathfrak{q}}, T_p E \otimes \Lambda_K^\vee)^{G_{K_{\mathfrak{q}}}})^\vee) = (P_{\mathfrak{q}}(\varepsilon_K(\text{Frob}_{\mathfrak{q}}^{-1})))$, where

$$P_{\mathfrak{q}}(X) = \det(1 - \text{Nm}(\mathfrak{q})^{-1} X \cdot \text{Frob}_{\mathfrak{q}} | V_p E^{I_{\mathfrak{q}}}).$$

Proof. For simplicity, we let $M := T_p E \otimes \Lambda_K^\vee$, $N := T_p E \otimes \Lambda_K$. Then we have the natural G_K -perfect pairing

$$M \times N \rightarrow \mathbb{Q}_p/\mathbb{Z}_p(1).$$

It induces the perfect local Tate pairing

$$H^1(K_{\mathfrak{q}}, M) \times H^1(K_{\mathfrak{q}}, N) \rightarrow \mathbb{Q}_p/\mathbb{Z}_p, \quad \text{for every place } \mathfrak{q}$$

via cup product. Let $\mathfrak{q} \nmid p$ be a prime of K . Since $G_{K_{\mathfrak{q}}}/I_{\mathfrak{q}} = \langle \text{Frob}_{\mathfrak{q}} \rangle \simeq \widehat{\mathbb{Z}}$ has cohomological dimension 1, we have that under local Tate pairing, $H^1(G_{K_{\mathfrak{q}}}/I_{\mathfrak{q}}, N^{I_{\mathfrak{q}}})$ and $H^1(G_{K_{\mathfrak{q}}}/I_{\mathfrak{q}}, M^{I_{\mathfrak{q}}})$ annihilate each other. Hence, by the inflation-restriction exact sequence

$$0 \rightarrow H^1(G_{K_{\mathfrak{q}}}/I_{\mathfrak{q}}, N^{I_{\mathfrak{q}}}) \rightarrow H^1(G_{K_{\mathfrak{q}}}, N) \rightarrow H^1(I_{\mathfrak{q}}, N)^{G_{K_{\mathfrak{q}}}} \rightarrow 0,$$

to prove (1), we only need to prove that $H^1(I_{\mathfrak{q}}, N)^{G_{K_{\mathfrak{q}}}} = 0$. Since $\mathfrak{q} \nmid p$, we know that there is a unique closed subgroup $J_{\mathfrak{q}}$ of $I_{\mathfrak{q}}$ such that $I_{\mathfrak{q}}/J_{\mathfrak{q}} \simeq \mathbb{Z}_p(1)$ as $\text{Gal}(K_{\mathfrak{q}}^{\text{ur}}/K_{\mathfrak{q}})$ -module, and $J_{\mathfrak{q}}$ has profinite degree prime to p . By the inflation-restriction exact sequence again, we have

$$H^1(I_{\mathfrak{q}}, N) = H^1(I_{\mathfrak{q}}/J_{\mathfrak{q}}, N^{J_{\mathfrak{q}}}).$$

Let $N_0 := T_p E^{J_{\mathfrak{q}}}$. Since $\varepsilon_K(I_{\mathfrak{q}}) = 1$, we can find that $N^{J_{\mathfrak{q}}} = N_0 \otimes \Lambda_K$. Let $\sigma_{\mathfrak{q}}$ be a topological generator of $I_{\mathfrak{q}}/J_{\mathfrak{q}} \simeq \mathbb{Z}_p(1)$. For any $\phi \in H^1(I_{\mathfrak{q}}/J_{\mathfrak{q}}, N_0 \otimes \Lambda_K)^{G_{K_{\mathfrak{q}}}}$, $g \in G_{K_{\mathfrak{q}}}$, we have $(g\phi)(\sigma_{\mathfrak{q}}) = \phi(\sigma_{\mathfrak{q}})$. Hence, there exists an integer $r = r(g)$ (we can assume $r > 0$) such that

$$g^{-1}\phi(\sigma_{\mathfrak{q}}) = \phi(g^{-1}\sigma_{\mathfrak{q}}g) = \phi(\sigma_{\mathfrak{q}}^r) = (1 + \sigma_{\mathfrak{q}} + \cdots + \sigma_{\mathfrak{q}}^{r-1})\phi(\sigma_{\mathfrak{q}}).$$

Since \mathfrak{q} is not completely split in K_∞/K , and $\varepsilon_K(G_{K_{\mathfrak{q}}})$ is not trivial, we can choose g such that $\varepsilon_K(g) \neq 1$. However, since $\varepsilon_K(\sigma_{\mathfrak{q}}) = 1$, it follows that $\phi(\sigma_{\mathfrak{q}}) = 0$ which implies that $H^1(I_{\mathfrak{q}}, N)^{G_{K_{\mathfrak{q}}}} = 0$.

Now we prove (2). As before, we have

$$(H^1(I_{\mathfrak{q}}, M)^{G_{K_{\mathfrak{q}}}})^\vee \simeq H^1(G_{K_{\mathfrak{q}}}/I_{\mathfrak{q}}, N^{I_{\mathfrak{q}}}).$$

Since

$$H^1(G_{K_{\mathfrak{q}}}/I_{\mathfrak{q}}, N^{I_{\mathfrak{q}}}) \simeq (T_p E^{I_{\mathfrak{q}}} \otimes \Lambda_K)/(1 - \text{Frob}_{\mathfrak{q}}^{-1}),$$

we have that

$$\text{Fitt}_{\Lambda_K}((H^1(I_{\mathfrak{q}}, M)^{G_{K_{\mathfrak{q}}}})^\vee) = (P_{\mathfrak{q}}(\varepsilon_K^{-1}(\text{Frob}_{\mathfrak{q}})))$$

which is principal and therefore divisorial. However, $\text{Char}_{\Lambda_K}((H^1(I_{\mathfrak{q}}, M)^{G_{K_{\mathfrak{q}}}})^{\vee})$ is the minimal divisorial ideal containing $\text{Fitt}_{\Lambda_K}((H^1(I_{\mathfrak{q}}, M)^{G_{K_{\mathfrak{q}}}})^{\vee})$, hence we have

$$\text{Char}_{\Lambda_K}((H^1(I_{\mathfrak{q}}, M)^{G_{K_{\mathfrak{q}}}})^{\vee}) = \text{Fitt}_{\Lambda_K}((H^1(I_{\mathfrak{q}}, M)^{G_{K_{\mathfrak{q}}}})^{\vee}) = (P_{\mathfrak{q}}(\varepsilon_K^{-1}(\text{Frob}_{\mathfrak{q}}))).$$

□

Lemma 2.3. *The natural Λ_K^- -module homomorphism*

$$\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty})/(\gamma^+ - 1)\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}) \rightarrow \mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}^-)$$

is pseudo-isomorphism.

Proof. By [Wan21, Proposition 3.1] and Lemma 2.2 (1) we obtain the desired result. Note that in our case, every finite place in Σ splits in K . □

We also consider imprimitive Selmer groups. For $M = T_p E \otimes \Lambda_K^{\vee}$, we define

$$H_{\mathcal{F}_{\text{ord}}^{\Sigma}}^1(K, M) = \text{Ker} \left(H^1(G_{K, \Sigma}, M) \rightarrow \prod_{\mathfrak{q}|p} \frac{H^1(K_{\mathfrak{q}}, M)}{H_{\text{ord}}^1(K_{\mathfrak{q}}, M)} \right)$$

and $\mathcal{X}_{\text{ord}}^{\Sigma}(E/K_{\infty}) = H_{\mathcal{F}_{\text{ord}}^{\Sigma}}^1(K, M)^{\vee}$.

In the proof of [SU14, Lemma 3.16], the following conclusion is essentially demonstrated.

Lemma 2.4. $\mathcal{X}_{\text{ord}}^{\Sigma}(E/K_{\infty})$ is Λ_K -torsion.

Therefore, we have the following corollary.

Corollary 2.5. $\mathcal{X}_{\text{ord}}(E/K_{\infty})$ is Λ_K -torsion, and

$$\text{Char}_{\Lambda_K}(\mathcal{X}_{\text{ord}}^{\Sigma}(E/K_{\infty})) \supset \text{Char}_{\Lambda_K}(\mathcal{X}_{\text{ord}}(E/K_{\infty})) \prod_{\mathfrak{q} \in \Sigma - \{\mathfrak{p}, \bar{\mathfrak{p}}\}} (P_{\mathfrak{q}}(\varepsilon_K(\text{Frob}_{\mathfrak{q}}^{-1}))).$$

Compact Selmer groups. We also consider compact Selmer groups. For Λ any one of Λ_K or Λ_K^{\pm} , let $N = T_p E \otimes_{\mathbb{Z}_p} \Lambda$, where Λ is equipped with the natural G_K -action. For $a \in \{\text{ord}, \text{rel}, \text{str}\}$, and $\mathfrak{q}|p$ a prime above p , we define the local conditions $H_a^1(K_{\mathfrak{q}}, N) \subset H^1(K_{\mathfrak{q}}, N)$ similarly to Definition 2.1, and for $a, b \in \{\text{ord}, \text{rel}, \text{str}\}$, let

$$H_{\mathcal{F}_{a,b}}^1(K, N) = \text{Ker} \left(H^1(G_{K, \Sigma}, N) \rightarrow \frac{H^1(K_{\mathfrak{p}}, N)}{H_a^1(K_{\mathfrak{p}}, N)} \times \frac{H^1(K_{\bar{\mathfrak{p}}}, N)}{H_b^1(K_{\bar{\mathfrak{p}}}, N)} \right).$$

For simplicity, we write

$$S_{a,b}(E/K_{\infty}) = H_{\mathcal{F}_{a,b}}^1(K, T_p E \otimes_{\mathbb{Z}_p} \Lambda_K), \quad S_a(E/K_{\infty}) = H_{\mathcal{F}_{a,a}}^1(K, T_p E \otimes_{\mathbb{Z}_p} \Lambda_K)$$

for $a, b \in \{\text{ord}, \text{str}, \text{rel}\}$, and similarly for E/K_{∞}^{\pm} with Λ_K replaced by Λ_K^{\pm} .

2.2. Selmer groups for Hida families. Let W be an indeterminate and $\Lambda_W = \mathbb{Z}_p[[W]]$. We identify Λ_W with Λ_K^+ by identifying the topological generator γ^+ with $1 + W$, which induces $\varepsilon_W : G_K \rightarrow \Lambda_W^{\times}$.

Let $L \subset \mathbb{Q}_p$ be a finite extension of \mathbb{Q}_p with integer ring \mathcal{O} . Let \mathbb{I} be a local reduced finite integral extension of $\Lambda_{W, \mathcal{O}} := \Lambda_W \otimes_{\mathbb{Z}_p} \mathcal{O}$. Recall that $\phi \in \text{Hom}_{\text{cont } \mathcal{O}\text{-alg}}(\mathbb{I}, \bar{\mathbb{Q}}_p)$ is called arithmetic if $\phi(1 + W) = \zeta_{\phi}(1 + p)^{k_{\phi}-2}$ for some p -power root of unity ζ_{ϕ} and integer k_{ϕ} . Let $t_{\phi} > 0$ be the integer such that ζ_{ϕ} is primitive $p^{t_{\phi}-1}$ -th root of unity, and $\chi_{\phi} : \mathbb{A}_{\mathbb{Q}}^{\times}/\mathbb{Q}^{\times} \rightarrow \mu_{p^{\infty}}$ be the unique character such that $\chi_{\phi, p}(1 + p) = \zeta_{\phi}^{-1}$ and has p -power conductor. Define

$$\mathfrak{X}_{\mathbb{I}, \mathcal{O}}^a = \{ \phi \in \text{Hom}_{\text{cont } \mathcal{O}\text{-alg}}(\mathbb{I}, \bar{\mathbb{Q}}_p) : \phi \text{ is arithmetic, } k_{\phi} \geq 2 \}.$$

Let \mathbf{f} be an \mathbb{I} -adic ordinary eigenform of tame level N and trivial character, i.e., a q -expansion $\mathbf{f} = \sum_{n=0}^{\infty} \mathbf{a}_n(\mathbf{f})q^n \in \mathbb{I}[[q]]$ such that for all $\phi \in \mathfrak{X}_{\mathbb{I}, \mathcal{O}}^a$,

$$\mathbf{f}_{\phi} = \sum_{n=0}^{\infty} \phi(\mathbf{a}_n(\mathbf{f}))q^n \in M_{k_{\phi}}(Np^{t_{\phi}}, \omega^{k_{\phi}-2}\chi_{\phi}; \phi(\mathbb{I}))$$

is ordinary. Here, ω is the Teichmüller character.

Assume \mathbf{f} satisfies the following condition

(irred \mathbf{f}) the residue representation of $\rho_{\mathbf{f}_{\phi}}$ is irreducible for some (hence all) $\phi \in \mathfrak{X}_{\mathbb{I}, \mathcal{O}}^a$.

Then there exists a continuous \mathbb{I} -linear Galois representation $(\rho_{\mathbf{f}}, T_{\mathbf{f}})$ with $T_{\mathbf{f}}$ a free \mathbb{I} -module of rank two and $\rho_{\mathbf{f}} : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_{\mathbb{I}}(T_{\mathbf{f}})$ a continuous representation characterized by the property that $\rho_{\mathbf{f}}$ is unramified at all primes $\ell \nmid Np$ and satisfies

$$\mathrm{tr}\rho_{\mathbf{f}}(\mathrm{Frob}_{\ell}) = a_{\ell}(\mathbf{f}), \quad \ell \nmid Np,$$

and

$$\det(\rho_{\mathbf{f}}) = \epsilon \varepsilon_W,$$

where ϵ is the p -adic cyclotomic character.

Let $F_{\mathbb{I}}$ be the ring of fractions of \mathbb{I} and $V_{\mathbf{f}} := T_{\mathbf{f}} \otimes_{\mathbb{I}} F_{\mathbb{I}}$. Since \mathbf{f} is ordinary, there exists an $F_{\mathbb{I}}$ -line $V_{\mathbf{f}}^+ \subset V_{\mathbf{f}}$ which is stable under the action of $G_{\mathbb{Q}_p}$. Furthermore, $G_{\mathbb{Q}_p}$ acts on $V_{\mathbf{f}}^- := V_{\mathbf{f}}/V_{\mathbf{f}}^+$ via the unramified character $\delta_{\mathbf{f}}$ characterized by $\delta_{\mathbf{f}}(\mathrm{Frob}_p) = a_p(\mathbf{f})$. Then $T_{\mathbf{f}}^+ := T_{\mathbf{f}} \cap V_{\mathbf{f}}^+$ is a free \mathbb{I} -summand of $T_{\mathbf{f}}$ of rank one.

Put $\mathbb{I}_K := \mathbb{I}[[\Gamma_K]]$. Let $\mathbf{M} = T_{\mathbf{f}} \otimes_{\mathbb{I}} \mathbb{I}_K^{\vee}$, $\mathbf{M}^- = T_{\mathbf{f}}^- \otimes_{\mathbb{I}} \mathbb{I}_K^{\vee}$ where \mathbb{I}_K^{\vee} is equipped with a G_K -action via ε_K^{-1} . Let Σ be a set of places of K containing all places of K dividing $pN\infty$. We define

$$H_{\mathcal{F}_{\mathrm{ord}}^{\Sigma}}^1(K, \mathbf{M}) := \ker \left(H^1(G_{K, \Sigma}, \mathbf{M}) \rightarrow \prod_{\mathfrak{q}|p} H^1(I_{\mathfrak{q}}, \mathbf{M}^-) \right).$$

Write $\mathcal{X}_{\mathrm{ord}}^{\Sigma}(\mathbf{f}/K_{\infty}) := H_{\mathcal{F}_{\mathrm{ord}}^{\Sigma}}^1(K, \mathbf{M})^{\vee}$. It is \mathbb{I}_K -torsion by [SU14, Lemma 3.16].

Let E/\mathbb{Q} be an elliptic curve such that E has good ordinary reduction at p . Let \mathbf{f} be an \mathbb{I} -adic ordinary eigenform and $\phi \in \mathfrak{X}_{\mathbb{I}, A}^a$ such that $\phi(1+W) = 1$ and $\phi(\mathbf{f})$ is equal to p -stabilization of the newform associated to E .

Lemma 2.6. *For every $\mathfrak{q}|p$, we have $H^1(G_{K_{\mathfrak{q}}}/I_{\mathfrak{q}}, (T_p \tilde{E}/\mathbb{F}_{\mathfrak{q}} \otimes \Lambda_K^{\vee})^{I_{\mathfrak{q}}}) = 0$.*

Proof. Let $\alpha_{\mathfrak{q}} : G_{K_{\mathfrak{q}}} \rightarrow \mathbb{Q}_p^{\times}$ be the unramified character associated to $T_p \tilde{E}/\mathbb{F}_{\mathfrak{q}}$, then $G_{K_{\mathfrak{q}}}$ acts on $T_p \tilde{E}/\mathbb{F}_{\mathfrak{q}} \otimes \Lambda_K^{\vee}$ via $\alpha_{\mathfrak{q}} \varepsilon_K^{-1}$, and $\alpha_{\mathfrak{q}}(\mathrm{Frob}_{\mathfrak{q}})$ is not a unit. Let $C_{\mathfrak{q}} \subset \Lambda_K$ be the ideal generated by $\{\varepsilon_K(g) - 1 : g \in I_{\mathfrak{q}}\}$, and $\sigma_{\mathfrak{q}} \in G_{K_{\mathfrak{q}}}$ be a lifting of $\mathrm{Frob}_{\mathfrak{q}}$. Then we can find that

$$H^1(G_{K_{\mathfrak{q}}}/I_{\mathfrak{q}}, (T_p \tilde{E}/\mathbb{F}_{\mathfrak{q}} \otimes \Lambda_K^{\vee})^{I_{\mathfrak{q}}}) \simeq \mathrm{Hom}_{\mathrm{cts}} \left((\Lambda_K/C_{\mathfrak{q}})^{\varepsilon_K(\sigma_{\mathfrak{q}}) = \alpha_{\mathfrak{q}}^{-1}(\mathrm{Frob}_{\mathfrak{q}})}, \mathbb{Q}_p/\mathbb{Z}_p \right) = 0,$$

where $(\Lambda_K/C_{\mathfrak{q}})^{\varepsilon_K(\sigma_{\mathfrak{q}}) = \alpha_{\mathfrak{q}}^{-1}(\mathrm{Frob}_{\mathfrak{q}})}$ is the submodule of $\Lambda_K/C_{\mathfrak{q}}$ killed by $\varepsilon_K(\sigma_{\mathfrak{q}}) - \alpha_{\mathfrak{q}}^{-1}(\mathrm{Frob}_{\mathfrak{q}})$. \square

Corollary 2.7. *Let $\mathfrak{p}_{\phi} := \ker \phi$, then $\mathcal{X}_{\mathrm{ord}}^{\Sigma}(\mathbf{f}/K_{\infty})/\mathfrak{p}_{\phi} \mathcal{X}_{\mathrm{ord}}^{\Sigma}(\mathbf{f}/K_{\infty}) \otimes_{\mathbb{I}, \phi} \mathcal{O} \simeq \mathcal{X}_{\mathrm{ord}}^{\Sigma}(E/K_{\infty})$.*

Proof. By [SU14, Proposition 3.7] and above lemma. \square

3. p -ADIC L -FUNCTIONS

3.1. A three variable p -adic L -function. Let \mathbf{f} be an \mathbb{I} -adic ordinary eigenform of tame level N and trivial character as in Section 2.2. Suppose that L contains $\mathbb{Q}[\mu_{Np}, i, D_K^{1/2}]$. Let

$$\mathfrak{X}_{\mathbb{I}_K, \mathcal{O}}^a := \{\phi \in \mathrm{Hom}_{\mathrm{cont}} \mathcal{O}\text{-alg}(\mathbb{I}_K, \bar{\mathbb{Q}}_p) : \phi|_{\mathbb{I}} \in \mathfrak{X}_{\mathbb{I}, \mathcal{O}}^a, \phi(\gamma^+) = \zeta_+(1+p)^{k_{\phi}|_{\mathbb{I}}-2}, \phi(\gamma^-) = \zeta_-\}$$

where ζ_{\pm} are p -power roots of unity. For each $\phi \in \mathfrak{X}_{\mathbb{I}_K, \mathcal{O}}^a$, let k_{ϕ} , t_{ϕ} and χ_{ϕ} denote the corresponding objects for $\phi|_{\mathbb{I}}$. Define

$$\xi_{\phi} := \phi \circ (\varepsilon_K/\varepsilon_W), \quad \theta_{\phi} := \omega^{2-k_{\phi}} \chi_{\phi}^{-1} \xi_{\phi}.$$

These are finite order idele class characters of \mathbb{A}_K^{\times} . For an idele class character ψ , denote its conductor by \mathfrak{f}_{ψ} . Let

$$\mathfrak{X}'_{\mathbb{I}_K, \mathcal{O}} := \{\phi \in \mathfrak{X}_{\mathbb{I}_K, \mathcal{O}}^a : p|\mathfrak{f}_{\xi_{\phi}}, p^{t_{\phi}}|\mathrm{Nm}(\mathfrak{f}_{\xi_{\phi}}), p|\mathfrak{f}_{\theta_{\phi}}\}.$$

Theorem 3.1. [SU14, Section 3.4.5] *Let Σ be a finite set of primes containing all those dividing pND_K . Let \mathbf{f} be an \mathbb{I} -adic ordinary eigenform of tame level N and trivial character. Assume that \mathbf{f} satisfies (irred $_{\mathbf{f}}$). Then there exists $\mathcal{L}_{\mathbf{f}, K}^{\Sigma} \in \mathbb{I}_K$ such that for any $\phi \in \mathfrak{X}'_{\mathbb{I}_K, \mathcal{O}}$, we have*

$$\begin{aligned} \mathcal{L}_{\mathbf{f}, K}^{\Sigma}(\phi) &= u_{\mathbf{f}_{\phi}} a_p(\mathbf{f}_{\phi})^{-\mathrm{ord}_p(\mathrm{Nm}(\mathfrak{f}_{\theta_{\phi}}))} \\ &\quad \times \frac{((k_{\phi} - 2)!)^2 \mathfrak{g}(\theta_{\phi}^{-1}) \mathrm{Nm}(\mathfrak{f}_{\theta_{\phi}} \delta_K)^{k_{\phi}-2} L^{\Sigma}(\mathbf{f}_{\phi}/K, \theta_{\phi}^{-1}, k_{\theta_{\phi}} - 1)}{(-2\pi i)^{2k_{\phi}-2} \Omega_{\mathbf{f}_{\phi}}^+ \Omega_{\mathbf{f}_{\phi}}^-}, \end{aligned}$$

where $u_{\mathbf{f}_{\phi}}$ is a p -adic unit depending only on \mathbf{f}_{ϕ} , $\mathfrak{g}(\theta_{\phi}^{-1})$ is the (global) Gauss sum, δ_K is the differential ideal, and $\Omega_{\mathbf{f}_{\phi}}^{\pm}$ are the canonical periods of \mathbf{f}_{ϕ} .

Remark 3.2. Note that we use arithmetic Frobenius to normalize the reciprocity map of class field theory, while [SU14] used geometric Frobenius.

3.2. Two variable p -adic L -functions: type I. Let $f = \sum_n a_n q^n \in S_2(\Gamma_0(N))$ be a newform with $p \nmid a_p$, and $c_f \in \mathbb{Z}_p$ be the congruence number of f as in [Hid81, section 7] or [Rib83].

Theorem 3.3. [CGS23, Theorem 1.2.1] *There exists an element $\mathcal{L}_p^I(f/K) \in c_f^{-1} \Lambda_K$ such that for any finite order nontrivial character ξ of Γ_K ,*

$$\mathcal{L}_p^I(f/K)(\xi) = W(\xi) p^{\text{ord}_p(\text{Nm}(f_\xi))/2} \alpha_p^{-\text{ord}_p(\text{Nm}(f_\xi))} \left(1 - \frac{p}{\alpha_p^2}\right)^{-1} \left(1 - \frac{1}{\alpha_p^2}\right)^{-1} \frac{L(f/K, \xi^{-1}, 1)}{8\pi^2 \langle f, f \rangle},$$

where α_p is the p -adic unit root of $x^2 - a_p x + p$, $\langle f, g \rangle = \int_{\Gamma_0(N) \backslash \mathcal{H}} \overline{f(\tau)} g(\tau) d\tau$ is the Petersson inner product on $S_2(\Gamma_1(N))$, $W(\xi)$ is the Artin root number.

Let E/\mathbb{Q} be an elliptic curve such that E has good ordinary reduction at p . Let $f_E \in S_2(\Gamma_0(N))$ be the newform associated to E , and $\pi_E : X_0(N) \rightarrow E$ a modular parametrization.

Definition 3.4 (Perrin-Riou's p -adic L -function). *We define Perrin-Riou's p -adic L -function to be*

$$\mathcal{L}_p^{\text{PR}}(E/K) := \left(1 - \frac{p}{\alpha_p^2}\right) \left(1 - \frac{1}{\alpha_p^2}\right) \cdot \frac{\deg(\pi_E)}{c_E^2} \cdot \mathcal{L}_p^I(f_E/K) \in \Lambda_K,$$

where c_E is the Manin constant.

Let \mathbf{f} be an \mathbb{I} -adic ordinary eigenform and $\phi \in \mathfrak{X}_{\mathbb{I}, \mathcal{O}}^a$ as in Section 2.2, such that $\phi(1+W) = 1$ and \mathbf{f}_ϕ equal to p -stabilization of f_E .

Proposition 3.5. *Assume that the residue representation $\bar{\rho}_E : G_{\mathbb{Q}} \rightarrow \text{Aut}(E[p])$ is irreducible, then*

$$\mathcal{L}_{\mathbf{f}, K}^{\Sigma}(\phi \otimes \text{id}) = \alpha \cdot \prod_{\mathfrak{q} \in \Sigma, \mathfrak{q} \nmid p} P_{\mathfrak{q}}(\varepsilon_K(\text{Frob}_{\mathfrak{q}}^{-1})) \cdot \mathcal{L}_p^{\text{PR}}(E/K),$$

where

$$P_{\mathfrak{q}}(X) = \det(1 - \text{Nm}(\mathfrak{q})^{-1} X \cdot \text{Frob}_{\mathfrak{q}} | V_p E^{I_{\mathfrak{q}}})$$

is the local Euler factor, α is a p -adic units which is independent of ξ , and $\phi \otimes \text{id} : \mathbb{I}_K = \mathbb{I} \hat{\otimes} \Lambda_K \rightarrow \phi(\mathbb{I}) \otimes \Lambda_K$ is the natural map.

Proof. We have that $W(\xi) = g(\bar{\xi})/\sqrt{\text{Nm}(\bar{f}_\xi)}$. By [Maz78, Corollary 4.1], c_E is a p -adic unit. By [CGS23, Lemma 3.1.2], $\deg(\pi_E) = c_{f_E}$ up to a p -adic unit. By [SZ14, Lemma 9.5],

$$\frac{\langle f_E, f_E \rangle}{c_{f_E}} = i(2\pi i)^2 \Omega_{f_E}^+ \Omega_{f_E}^-.$$

These give the equality of the proposition. \square

3.3. Cyclotomic p -adic L -function. Let E/\mathbb{Q} be an elliptic curve such that E has good ordinary reduction at p . Pick generators δ^\pm of $H_1(E, \mathbb{Z})^\pm$, and define the Néron periods Ω_E^\pm by

$$\Omega_E^\pm = \int_{\delta^\pm} \omega_E,$$

where ω_E is a minimal differential on E . We normalize the δ^\pm so that $\Omega_E^+ \in \mathbb{R}_{>0}$ and $\Omega_E^- \in i\mathbb{R}_{>0}$.

Let a_p be the Fourier coefficient of the newform f_E associated to E , and α_p be the p -adic unit root of $x^2 - a_p x + p$ as before.

Theorem 3.6. [CGS23, Theorem 1.1.1] *There exists an element $\mathcal{L}_p^{\text{MSD}}(E/\mathbb{Q}) \in \Lambda_{\mathbb{Q}}$ such that for any finite order character χ of $\Gamma_{\mathbb{Q}}$ of conductor p^r with $r > 0$, we have*

$$\mathcal{L}_p^{\text{MSD}}(E/\mathbb{Q})(\chi) = \frac{p^r}{g(\bar{\chi}) \alpha_p^r} \cdot \frac{L(E, \bar{\chi}, 1)}{\Omega_E^+},$$

where $g(\bar{\chi}) = \sum_{a \bmod p^r} \bar{\chi}(a) e^{2\pi i a/p^r}$ is the Gauss sum, and

$$\mathcal{L}_p^{\text{MSD}}(E/\mathbb{Q})(1) = (1 - \alpha_p^{-1})^2 \cdot \frac{L(E, 1)}{\Omega_E^+}.$$

Let $\mathcal{L}_p^{\text{PR}}(E/K)^+ \in \Lambda_{\mathbb{Q}}$ be the image of $\mathcal{L}_p^{\text{PR}}(E/K)$ under the map induced by the projection $\Gamma_K \rightarrow \Gamma_K^+ \simeq \Gamma_{\mathbb{Q}}$.

Proposition 3.7. [CGS23, Proposition 1.2.4] *We have*

$$\mathcal{L}_p^{\text{PR}}(E/K)^+ = \mathcal{L}_p^{\text{MSD}}(E/\mathbb{Q}) \cdot \mathcal{L}_p^{\text{MSD}}(E^K/\mathbb{Q})$$

up to a unit in $\Lambda_{\mathbb{Q}}^{\times}$, where E^K is the twist of E by the quadratic character corresponding to K/\mathbb{Q} .

3.4. Two variable p -adic L -functions: type II.

Theorem 3.8. [CGS23, Theorem 1.4.1] *There exists an element $\mathcal{L}_p^{\text{II}}(f/K) \in \text{Frac}\Lambda_K$ such that for every character ξ of Γ_K which is crystalline at both \mathfrak{p} and $\bar{\mathfrak{p}}$, and of infinity type (b, a) with $a \leq -1, b \geq 1$, we have*

$$\mathcal{L}_p^{\text{II}}(f/K)(\xi) = \frac{2^{a-b} i^{b-a-1} \Gamma(b+1) \Gamma(b) N^{a+b+1}}{(2\pi)^{2b+1} \langle \theta_{\xi_b}, \theta_{\xi_b} \rangle} \cdot \frac{\mathcal{E}(\xi, f, 1)}{(1 - \xi^{1-\tau}(\bar{\mathfrak{p}}))(1 - p^{-1} \xi^{1-\tau}(\bar{\mathfrak{p}}))} \cdot L(f/K, \xi, 1),$$

where θ_{ξ_b} is the theta series associated to the Hecke character $\xi_b = \xi| \cdot |^{-b}$, τ is the complex conjugation, and

$$\mathcal{E}(\xi, f, 1) = (1 - p^{-1} \xi(\bar{\mathfrak{p}}) \alpha_p) (1 - \xi(\bar{\mathfrak{p}}) \alpha_p^{-1}) (1 - p^{-1} \xi^{-1}(\mathfrak{p}) \alpha_p) (1 - \xi^{-1}(\mathfrak{p}) \alpha_p^{-1}).$$

Let $\Lambda_K^{\text{ur}} := \Lambda_K \widehat{\otimes} \mathbb{Z}_p^{\text{ur}}$, where \mathbb{Z}_p^{ur} is the completion of the ring of integers of the maximal unramified extension of \mathbb{Q}_p .

Theorem 3.9. [dS87, Chapter 2, Theorem 4.14] *There exists an element $\mathcal{L}_p(K) \in \Lambda_K^{\text{ur}}$ such that for every character ξ of Γ_K of infinity type (j, k) with $0 \leq -j \leq k$, we have*

$$\mathcal{L}_p(K)(\xi) = \frac{\Omega_p^{k-j}}{\Omega_K^{k-j}} \cdot \Gamma(k) \cdot \left(\frac{\sqrt{D_K}}{2\pi} \right)^j \cdot (1 - \xi^{-1}(\mathfrak{p}) p^{-1}) (1 - \xi(\bar{\mathfrak{p}})) \cdot L(\xi, 0),$$

where Ω_p and Ω_K are CM periods attached to K .

Definition 3.10 (Greenberg's p -adic L -function). *Let*

$$\mathcal{L}_p^{\text{Gr}}(f/K) := h_K \cdot \mathcal{L}_p(K)' \cdot \mathcal{L}_p^{\text{II}}(f/K),$$

where h_K is the class number of K , and $\mathcal{L}_p(K)'$ is the image of $\mathcal{L}_p(K)$ under the map $\Lambda_K^{\text{ur}} \rightarrow \Lambda_K^{\text{ur}}$ given by $\gamma \mapsto \gamma^{1-\tau}$ for $\gamma \in \Gamma_K$.

Lemma 3.11. [CGS23, Lemma 1.4.4] *The Greenberg's p -adic L -function $\mathcal{L}_p^{\text{Gr}}(f/K)$ is integral, i.e., belongs to Λ_K^{ur} .*

3.5. BDP p -adic L -function. Assume that D_K is odd and not equal to -3 , and the Heegner hypothesis holds. Fix an integral ideal $\mathfrak{n} \subset \mathcal{O}_K$ with $\mathcal{O}_K/\mathfrak{n} \simeq \mathbb{Z}/N\mathbb{Z}$. Let $\Lambda_K^{\text{ur}, -} := \Lambda_K^- \widehat{\otimes} \mathbb{Z}_p^{\text{ur}}$.

Theorem 3.12. [CGLS22, Theorem 2.1.1] *There exists an element $\mathcal{L}_p^{\text{BDP}}(f/K) \in \Lambda_K^{\text{ur}, -}$ characterized by the following interpolation property: for every character ξ of Γ_K^- crystalline at both \mathfrak{p} and $\bar{\mathfrak{p}}$ and corresponding to a Hecke character of K of infinity type $(n, -n)$ with $n \in \mathbb{Z}_{>0}$ and $n \equiv 0 \pmod{p-1}$, we have*

$$\mathcal{L}_p^{\text{BDP}}(f/K)(\xi) = \frac{\Omega_p^{4n}}{\Omega_K^{4n}} \cdot \frac{\Gamma(n) \Gamma(n+1) \xi(\mathfrak{n}^{-1})}{4(2\pi)^{2n+1} \sqrt{D_K}^{-2n-1}} \cdot (1 - a_p \xi(\bar{\mathfrak{p}}) p^{-1} + \xi(\bar{\mathfrak{p}})^2 p^{-1})^2 \cdot L(f/K, \xi, 1).$$

Denote by $\mathcal{L}_p^{\text{Gr}}(f/K)^-$ the image of $\mathcal{L}_p^{\text{Gr}}(f/K)$ under the natural projection $\Lambda_K^{\text{ur}} \rightarrow \Lambda_K^{\text{ur}, -}$. We have the following proposition.

Proposition 3.13. [CGS23, Proposition 1.4.5] $\mathcal{L}_p^{\text{Gr}}(f/K)^- \cdot \Lambda_K^{\text{ur}, -} = \mathcal{L}_p^{\text{BDP}}(f/K) \cdot \Lambda_K^{\text{ur}, -}$.

4. MAIN CONJECTURES

4.1. Two variable Iwasawa main conjectures. Let E/\mathbb{Q} be an elliptic curve of conductor N , $p > 2$ a prime such that E has good ordinary reduction at p , K an imaginary quadratic field such that $p = \mathfrak{p}\bar{\mathfrak{p}}$ split in K . Assume that residue representation $\bar{\rho}_E|_{G_K} : G_K \rightarrow \text{Aut}(E[p])$ is irreducible.

The following two variable Iwasawa main conjectures are considered.

Conjecture 4.1. (1) $H_{\mathcal{F}_{\text{ord}}}^1(K, T_p E \otimes \Lambda_K^{\vee})^{\vee}$ is Λ_K -torsion and

$$\text{Char}_{\Lambda_K}(H_{\mathcal{F}_{\text{ord}}}^1(K, T_p E \otimes \Lambda_K^{\vee})^{\vee}) = (\mathcal{L}_p^{\text{PR}}(E/K)).$$

(2) $H_{\mathcal{F}_{\text{Gr}}}^1(K, T_p E \otimes \Lambda_K^{\vee})^{\vee}$ is Λ_K -torsion and

$$\text{Char}_{\Lambda_K}(H_{\mathcal{F}_{\text{Gr}}}^1(K, T_p E \otimes \Lambda_K^{\vee})^{\vee}) \Lambda_K^{\text{ur}} = (\mathcal{L}_p^{\text{Gr}}(E/K)).$$

Recall the main theorem of this paper as stated below.

Theorem 4.2. *Suppose that the residue representation $\bar{\rho}_E|_{G_K} : G_K \rightarrow \text{Aut}(E[p])$ is absolutely irreducible. If the Heegner hypothesis holds (in particular, $\text{sign}(E/K) = -1$), then*

(1) $H_{\mathcal{F}_{\text{ord}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee$ is Λ_K -torsion and

$$\text{Char}_{\Lambda_K}(H_{\mathcal{F}_{\text{ord}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee) \subset (\mathcal{L}_p^{\text{PR}}(E/K)).$$

(2) $H_{\mathcal{F}_{\text{Gr}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee$ is Λ_K -torsion and

$$\text{Char}_{\Lambda_K}(H_{\mathcal{F}_{\text{Gr}}}^1(K, T_p E \otimes \Lambda_K^\vee)^\vee) \Lambda_K^{\text{ur}} \subset (\mathcal{L}_p^{\text{Gr}}(E/K)).$$

Moreover, if (Im) holds, the Conjecture 4.1 is true.

We will prove this theorem in the next two subsections.

4.2. A three variable Iwasawa main conjecture. Let \mathbf{f} be an \mathbb{I} -adic ordinary eigenform of tame level N and trivial character as in Section 2.2. Suppose that $L \supset \mathbb{Q}[\mu_{Np}, i, D_K^{1/2}]$, \mathbb{I} is a normal domain and \mathbf{f} satisfies (irred $_{\mathbf{f}}$).

Conjecture 4.3. [SU14] *Let Σ be a finite set of primes containing all those dividing pND_K . Then*

$$\text{Char}_{\mathbb{I}_K}(\mathcal{X}_{\text{ord}}^\Sigma(\mathbf{f}/K_\infty)) = (\mathcal{L}_{\mathbf{f}, K}^\Sigma).$$

Theorem 4.4 (Skinner-Urban). *Under the condition of Conjecture 4.3, we have that for any height one prime P of $\mathbb{I}_K = \mathbb{I}[[\Gamma_K]]$,*

$$\text{ord}_P(\text{Char}_{\mathbb{I}_K}(\mathcal{X}_{\text{ord}}^\Sigma(\mathbf{f}/K_\infty))) \geq \text{ord}_P(\mathcal{L}_{\mathbf{f}, K}^\Sigma),$$

unless $P = P^+ \mathbb{I}[[\Gamma_K]]$ for some height one prime P^+ of $\mathbb{I}[[\Gamma_K^+]]$.

Proof. Let $\mathbf{D} = (A, \mathbf{f}, 1, 1, \Sigma)$ be a p -adic Eisenstein datum defined in [SU14], with $A \supset \mathbb{Z}_p[i, D_K]$ a finite \mathbb{Z}_p -algebra. Let $\Lambda_{\mathbf{D}} := \mathbb{I}[[\Gamma_K]][[\Gamma_K^-]]$ as in [SU14]. The proof is essentially in section 7.4 of [SU14] except that we do not consider the prime $P = P^+ \Lambda_{\mathbf{D}}$ for some height one prime P^+ of $\mathbb{I}[[\Gamma_K^+]]$. By [SU14, Proposition 13.6(1)], assume Σ is large enough, then there is a p -adic Eisenstein series $\mathbf{E}_{\mathbf{D}}$ with coefficients (associated to formally q -expansion) belong to $\Lambda_{\mathbf{D}}$, and that there exists a set $\mathcal{C}_{\mathbf{D}}$ of some coefficients of $\mathbf{E}_{\mathbf{D}}$ satisfying that for any height one prime P of $\Lambda_{\mathbf{D}}$, if $P \supset \mathcal{C}_{\mathbf{D}}$, then $P = P^+ \Lambda_{\mathbf{D}}$ for some height one prime P^+ of $\mathbb{I}[[\Gamma_K^+]]$. Now we apply [SU14, Theorem 7.7]. Consider two p -adic L -functions \mathcal{L}_1^Σ and $\mathcal{L}_{\mathbf{f}, K, 1}^\Sigma := \mathcal{L}_{\mathbf{f}, K}^\Sigma$ in [SU14, Theorem 7.7]. Since $\mathcal{L}_1^\Sigma \in \mathbb{I}[[\Gamma_K^+]]$ by definition, we have that

$$\text{ord}_P(\text{Char}_{\mathbb{I}_K}(\mathcal{X}_{\text{ord}}^\Sigma(\mathbf{f}/K_\infty))) \geq \text{ord}_P(\mathcal{L}_{\mathbf{f}, K}^\Sigma)$$

for any height one prime P of $\Lambda_{\mathbf{D}}$ such that P is not of the form $P^+ \Lambda_{\mathbf{D}}$ for some height one prime P^+ of $\mathbb{I}[[\Gamma_K^+]]$. Now it is easy to see that our conclusion comes from above. \square

Remark 4.5. Since we do not consider the prime $P = P^+ \Lambda_{\mathbf{D}}$ for some height one prime P^+ of $\mathbb{I}[[\Gamma_K^+]]$, we don't need Proposition 13.6 (2) in [SU14]. Therefore, we do not need the hypotheses on N and $\bar{\rho}_{\mathbf{f}}|_{I_\ell}$ for $\ell|N$. Here, $\bar{\rho}_{\mathbf{f}} := \rho_{\mathbf{f}} \bmod \mathfrak{m}_{\mathbb{I}}$, and $\mathfrak{m}_{\mathbb{I}}$ is the maximal ideal of \mathbb{I} . See also the remark after Theorem 3.26 in [SU14].

Corollary 4.6. *There exists a nontrivial multiplicative set $S \subset \Lambda_K^+ \subset \Lambda_K$ such that*

$$S^{-1} \text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{ord}}}(E/K_\infty)) \subset (\mathcal{L}_p^{\text{PR}}(E/K))$$

holds in $S^{-1} \Lambda_K$.

Proof. By [Roh84], the p -adic L -function $\mathcal{L}_{\mathbf{f}, K}^\Sigma$ doesn't belong to $f p_\phi \mathbb{I}_K$. Let P_1, \dots, P_n be height one primes of \mathbb{I}_K such that $\text{ord}_{P_i} \mathcal{L}_{\mathbf{f}, K}^\Sigma > 0$, then $P_i = P_i^+ \mathbb{I}_K$ for some height one prime P_i^+ of $\mathbb{I}[[\Gamma_K^+]]$. We have $P_i \not\subset \mathfrak{p}_\phi \mathbb{I}_K$. Choose $h_i \in P_i \setminus \mathfrak{p}_\phi \mathbb{I}_K$, and let T be the multiplicative set generated by $\{h_i : i = 1, \dots, n\}$, and $S = \phi(T)$. Then by Theorem 4.4, [SU14, Corollary 3.8], Corollary 2.7, Corollary 2.5 and Proposition 3.5, we complete the proof. \square

4.3. **Proof of Theorem 4.2.** Similarly, as [BSTW24, Proposition 9.18] (see also [CGS23, Proposition 3.2.1]), the following theorem holds.

Theorem 4.7. *For every nontrivial multiplicative set $S \subset \Lambda_K$, the following are equivalent*

- (1) $S^{-1}\text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{ord}}}(E/K_{\infty})) \subset (\mathcal{L}_p^{\text{PR}}(E/K))$.
- (2) $S^{-1}\text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}))\Lambda_K^{\text{ur}} \subset (\mathcal{L}_p^{\text{Gr}}(E/K))$.

The same result holds for the opposite divisibilities.

By Theorem 4.7 and Corollary 4.6, there exists a nontrivial multiplicative set $S \subset \Lambda_K^+ \subset \Lambda_K$ such that

$$S^{-1}\text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}))\Lambda_K^{\text{ur}} \subset (\mathcal{L}_p^{\text{Gr}}(E/K)).$$

We may assume that S is generated by prime elements in the unique factorization domain Λ_K . For height one prime $P \subset \Lambda_K^{\text{ur}}$, if $P \cap S \neq \emptyset$, we have $P = P^+ \Lambda_K^{\text{ur}}$ for some $P^+ \subset \Lambda_K^{\text{ur},+}$. However, by [Hsi14, Theorem B] and Proposition 3.13,

$$\mu(\mathcal{L}_p^{\text{Gr}}(E/K)^-) = \mu(\mathcal{L}_p^{\text{BDP}}(E/K)) = 0,$$

where $\mu(\cdot)$ denotes the μ -invariant. Hence,

$$\text{ord}_P(\mathcal{L}_p^{\text{Gr}}(E/K)) = 0$$

if $P \subset \Lambda_K^{\text{ur}}$ is a height one prime of the form $P = P^+ \Lambda_K^{\text{ur}}$ for some $P^+ \subset \Lambda_K^{\text{ur},+}$. It implies that

$$\text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}))\Lambda_K^{\text{ur}} \subset (\mathcal{L}_p^{\text{Gr}}(E/K)).$$

Moreover, if (Im) holds, similarly to [SU14, Theorem 3.30], by using [Kat04, Theorem 17.4] and a commutative algebra lemma ([SU14, Lemma 3.2]), we have that Conjecture 1.1 (2) is true. Therefore Conjecture 1.1 (1) holds. Note that in [SU14], the condition $\text{Im}(\rho_E) \supset \text{SL}_2(\mathbb{Z}_p)$ is used, but it can be replaced by (Im) as discussed in the last paragraph of [Ski16, page 187].

4.4. **Mazur's main conjecture.** Let E/\mathbb{Q} be an elliptic curve of conductor N , $p > 2$ a prime such that E has good ordinary reduction at p . Suppose that the residue representation $\bar{\rho}_E : G_{\mathbb{Q}} \rightarrow \text{Aut}(E[p])$ is irreducible.

Conjecture 4.8. *(Mazur's main conjecture) $\mathcal{X}_{\mathcal{F}_{\text{ord}}}(E/\mathbb{Q}_{\infty})$ is $\Lambda_{\mathbb{Q}}$ -torsion and*

$$\text{Char}_{\Lambda_{\mathbb{Q}}}(\mathcal{X}_{\mathcal{F}_{\text{ord}}}(E/\mathbb{Q}_{\infty})) = (\mathcal{L}_p^{\text{MSD}}(E/\mathbb{Q})).$$

Choose an imaginary quadratic field K such that p is split in K and (E, K) satisfies the Heegner hypothesis. Similarly to [SU14, Theorem 3.33], by Theorem 4.2 and descent arguments, we have the following theorem. Note that by Serre ([Ser87, Section 3.3]), the condition $\bar{\rho}_E$ is irreducible implies that $\bar{\rho}_E$ is absolutely irreducible in the case $p > 2$.

Theorem 4.9. *$\mathcal{X}_{\mathcal{F}_{\text{ord}}}(E/\mathbb{Q}_{\infty})$ is $\Lambda_{\mathbb{Q}}$ -torsion and*

$$\text{Char}_{\Lambda_{\mathbb{Q}}}(\mathcal{X}_{\mathcal{F}_{\text{ord}}}(E/\mathbb{Q}_{\infty})) \otimes \mathbb{Q}_p = (\mathcal{L}_p^{\text{MSD}}(E/\mathbb{Q}))$$

in $\Lambda_{\mathbb{Q}} \otimes \mathbb{Q}_p$. Moreover, if (Im) holds, we have

$$\text{Char}_{\Lambda_{\mathbb{Q}}}(\mathcal{X}_{\mathcal{F}_{\text{ord}}}(E/\mathbb{Q}_{\infty})) = (\mathcal{L}_p^{\text{MSD}}(E/\mathbb{Q}))$$

in $\Lambda_{\mathbb{Q}}$.

4.5. **Anticyclotomic main conjectures.** Let E/\mathbb{Q} be an elliptic curve of conductor N , $p > 2$ a prime such that E has good ordinary reduction at p , K an imaginary quadratic field such that $p = \mathfrak{p}\bar{\mathfrak{p}}$ split in K and (E, K) satisfies the Heegner hypothesis. Assume that residue representation $\bar{\rho}_E|_{G_K} : G_K \rightarrow \text{Aut}(E[p])$ is irreducible.

Conjecture 4.10. *(BDP main conjecture) $\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}^-)$ is Λ_K -torsion and*

$$\text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{ord}}}(E/K_{\infty}^-))\Lambda_K^{\text{ur}} = (\mathcal{L}_p^{\text{BDP}}(E/K)).$$

Fix a modular parametrization $\pi : X_0(N) \rightarrow E$. In [PR87], Perrin-Riou constructed an element $\kappa \in S_{\text{ord}}(E/K_{\infty}^-)$ via the Kummer images of Heegner points on $X_0(N)$, which is Λ_K^- -non-torsion by Cornut-Vatsal [CV07]. Then she formulated an anticyclotomic main conjecture as follows.

Conjecture 4.11. *(Heegner point main conjecture) $S_{\text{ord}}(E/K_{\infty}^-)$ and $\mathcal{X}_{\text{ord}}(E/K_{\infty}^-)$ are both Λ_K^- -rank one, and*

$$\text{Char}_{\Lambda_K^-}(\mathcal{X}_{\text{ord}}(E/K_{\infty}^-)_{\text{tor}}) = \text{Char}_{\Lambda_K^-}(S_{\text{ord}}(E/K_{\infty}^-)/\Lambda_K^- \cdot \kappa)^2.$$

Theorem 4.12. (1) $\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}^-)$ is Λ_K -torsion and

$$\text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{ord}}}(E/K_{\infty}))\Lambda_K^{\text{ur}} \otimes \mathbb{Q}_p = (\mathcal{L}_p^{\text{BDP}}(E/K))$$

holds in $\Lambda_K \otimes \mathbb{Q}_p$. Moreover, if (Im) holds, then

$$\text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{ord}}}(E/K_{\infty}))\Lambda_K^{\text{ur}} = (\mathcal{L}_p^{\text{BDP}}(E/K)).$$

(2) $S_{\text{ord}}(E/K_{\infty}^-)$ and $\mathcal{X}_{\text{ord}}(E/K_{\infty}^-)$ are both Λ_K^- -rank one, and

$$\text{Char}_{\Lambda_K^-}(\mathcal{X}_{\text{ord}}(E/K_{\infty}^-)_{\text{tor}}) \otimes \mathbb{Q}_p = \text{Char}_{\Lambda_K^-}(S_{\text{ord}}(E/K_{\infty}^-)/\Lambda_K^- \cdot \kappa)^2 \otimes \mathbb{Q}_p$$

holds in $\Lambda_K \otimes \mathbb{Q}_p$. Moreover, if (Im) holds, then

$$\text{Char}_{\Lambda_K^-}(\mathcal{X}_{\text{ord}}(E/K_{\infty}^-)_{\text{tor}}) = \text{Char}_{\Lambda_K^-}(S_{\text{ord}}(E/K_{\infty}^-)/\Lambda_K^- \cdot \kappa)^2.$$

We prove this theorem in the remaining part of this subsection. First recall a theorem in [CGS23].

Theorem 4.13. [CGS23, Theorem 5.5.2] $S_{\text{ord}}(E/K_{\infty}^-)$ and $\mathcal{X}_{\text{ord}}(E/K_{\infty}^-)$ are both Λ_K^- -rank one, and

$$\text{Char}_{\Lambda_K^-}(\mathcal{X}_{\text{ord}}(E/K_{\infty}^-)_{\text{tor}}) \otimes \mathbb{Q}_p \supset \text{Char}_{\Lambda_K^-}(S_{\text{ord}}(E/K_{\infty}^-)/\Lambda_K^- \cdot \kappa)^2 \otimes \mathbb{Q}_p$$

holds in $\Lambda_K^- \otimes \mathbb{Q}_p$.

Similarly to [BCK21, Theorem 5.2], we have the following theorem.

Theorem 4.14. $\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}^-)$ is Λ_K -torsion and for every nontrivial multiplicative set $S \subset \Lambda_K^-$, the following are equivalent

- (1) $S^{-1}\text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}^-))\Lambda_K^{\text{ur},-} \supset (\mathcal{L}_p^{\text{BDP}}(E/K))$.
- (2) $S^{-1}\text{Char}_{\Lambda_K^-}(\mathcal{X}_{\text{ord}}(E/K_{\infty}^-)_{\text{tor}}) \supset S^{-1}\text{Char}_{\Lambda_K^-}(S_{\text{ord}}(E/K_{\infty}^-)/\Lambda_K^- \cdot \kappa)^2$.

The same result holds for the opposite divisibilities.

Since $\mathcal{L}_p^{\text{BDP}}(E/K)$ is nonzero ([BCK21, Corollary 4.5]), by the argument in Section 4.3, there exists a nontrivial multiplicative set $S \subset \Lambda_K^+ \subset \Lambda_K$ such that for any $s \in S$, $\gamma^+ - 1 \nmid s$, and

$$S^{-1}\text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}))\Lambda_K^{\text{ur}} \subset (\mathcal{L}_p^{\text{Gr}}(E/K)).$$

By Lemma 2.3, we have

$$\text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}^-))\Lambda_K^{\text{ur}} \otimes \mathbb{Q}_p \subset (\mathcal{L}_p^{\text{BDP}}(E/K)).$$

The other direction divisibility is given by combining Theorem 4.13 and Theorem 4.14.

Moreover, if (Im) holds, then

$$\text{Char}_{\Lambda_K}(\mathcal{X}_{\mathcal{F}_{\text{Gr}}}(E/K_{\infty}))\Lambda_K^{\text{ur}} \subset (\mathcal{L}_p^{\text{Gr}}(E/K)).$$

The remaining part can be deduced from above divisibility and Mazur's main conjecture (Theorem 4.9). See the proof in [BSTW24, Section 12.2.1] for details.

4.6. Applications. Let E/\mathbb{Q} be an elliptic curve of conductor N , $p > 2$ a prime such that E has good ordinary reduction at p .

Theorem 4.15. For an integer $r \leq 1$, the following are equivalent

- (1) $\text{rank}_{\mathbb{Z}} E(\mathbb{Q}) = r$ and $\#\text{III}(E/\mathbb{Q}) < \infty$,
- (2) $\text{corank}_{\mathbb{Z}_p} \text{Sel}_{p^\infty}(E/\mathbb{Q}) = r$,
- (3) $\text{ord}_{s=1} L(E/\mathbb{Q}, s) = r$.

Under any of the above, if (Im) also holds, then p -part BSD formula for E holds, i.e.,

$$\left| \frac{L^{(r)}(1, E)}{r! \cdot \Omega_E R_E} \right|_p = \left| \frac{\#\text{III}(E)[p^\infty] \cdot \prod_{\ell|N} c_\ell(E)}{(\#E(\mathbb{Q})_{\text{tor}})^2} \right|_p,$$

where R_E is the regulator of $E(\mathbb{Q})$, Ω_E the Néron period, $c_\ell(E)$ the Tamagawa number at a prime ℓ , and $|\cdot|_p$ the p -adic absolute value.

Proof. (1) \Rightarrow (2) is trivial. (3) \Rightarrow (1) is given by Gross-Zagier ([GZ86]) and Kolyvagin ([Kol90]). For (2) \Rightarrow (3), we can choose an imaginary quadratic field K such that $\text{ord}_{s=1} L(E^K, s) \leq 1$ and (E, K) satisfies the Heegner hypothesis by [FH95]. Similarly to [Wan21, Theorem 1.9], by applying descent arguments to (the rational part of) Theorem 4.12 (2) and using Gross-Zagier formula, we have

$$\text{corank}_{\mathbb{Z}_p} \text{Sel}_{p^\infty}(E/K) = 1 \text{ implies } \text{ord}_{s=1} L(E/K, s) = 1,$$

therefore we have (2) \Rightarrow (3) holds.

The rank 0 p -part BSD formula comes from (the integral part of) Theorem 4.9 and descent arguments (see [SU14, Section 3.6.1] for details). Choosing an imaginary quadratic field K as above, the rank 1 p -part BSD formula comes from (the integral part of) Theorem 4.12 and descent arguments (see [JSW17] for details). \square

CONFLICT OF INTEREST

On behalf of all authors, the corresponding author states that there is no conflict of interest.

REFERENCES

- [BCGS23] Ashay Burungale, Francesc Castella, Giada Grossi, and Christopher Skinner, *Non-vanishing of kolyvagin systems and iwawata theory*, arXiv preprint arXiv:2312.09301 (2023).
- [BCK21] Ashay Burungale, Francesc Castella, and Chan-Ho Kim, *A proof of Perrin-Riou’s Heegner point main conjecture*, Algebra Number Theory **15** (2021), no. 7, 1627–1653 (English).
- [BCS24] Ashay Burungale, Francesc Castella, and Christopher Skinner, *Base change and Iwasawa Main Conjectures for GL_2* , arXiv preprint arXiv:2405.00270 (2024).
- [BSTW24] Ashay Burungale, Christopher Skinner, Ye Tian, and Xin Wan, *Zeta elements for elliptic curves and applications*, Preprint, arXiv:2409.01350 [math.NT] (2024), 2024.
- [CGLS22] Francesc Castella, Giada Grossi, Jaehoon Lee, and Christopher Skinner, *On the anticyclotomic Iwasawa theory of rational elliptic curves at Eisenstein primes*, Invent. Math. **227** (2022), no. 2, 517–580. MR 4372220
- [CGS23] Francesc Castella, Giada Grossi, and Christopher Skinner, *Mazur’s main conjecture at Eisenstein primes*, Preprint, arXiv:2303.04373 [math.NT] (2023), 2023.
- [CV07] Christophe Cornut and Vinayak Vatsal, *Nontriviality of Rankin-Selberg L -functions and CM points, L -functions and Galois representations*, London Math. Soc. Lecture Note Ser., vol. 320, Cambridge Univ. Press, Cambridge, 2007, pp. 121–186. MR 2392354
- [dS87] Ehud de Shalit, *Iwasawa theory of elliptic curves with complex multiplication*, Perspectives in Mathematics, vol. 3, Academic Press, Inc., Boston, MA, 1987, p -adic L functions. MR 917944
- [FH95] Solomon Friedberg and Jeffrey Hoffstein, *Nonvanishing theorems for automorphic L -functions on $\text{GL}(2)$* , Ann. of Math. (2) **142** (1995), no. 2, 385–423. MR 1343325
- [GZ86] Benedict H. Gross and Don B. Zagier, *Heegner points and derivatives of L -series*, Invent. Math. **84** (1986), no. 2, 225–320. MR 833192
- [Hid81] Haruzo Hida, *Congruence of cusp forms and special values of their zeta functions*, Invent. Math. **63** (1981), no. 2, 225–261. MR 610538
- [Hsi14] Ming-Lun Hsieh, *Special values of anticyclotomic rankin-selberg l -functions*, Documenta Mathematica **19** (2014), 709–767.
- [JSW17] Dimitar Jetchev, Christopher Skinner, and Xin Wan, *The Birch and Swinnerton-Dyer formula for elliptic curves of analytic rank one*, Camb. J. Math. **5** (2017), no. 3, 369–434. MR 3684675
- [Kat04] Kazuya Kato, *p -adic Hodge theory and values of zeta functions of modular forms.*, Cohomologies p -adiques et applications arithmétiques (III), Paris: Société Mathématique de France, 2004, pp. 117–290 (English).
- [Kol90] V. A. Kolyvagin, *Euler systems*, The Grothendieck Festschrift, Vol. II, Progr. Math., vol. 87, Birkhäuser Boston, Boston, MA, 1990, pp. 435–483. MR 1106906
- [Maz78] B. Mazur, *Rational isogenies of prime degree (with an appendix by D. Goldfeld)*, Invent. Math. **44** (1978), no. 2, 129–162. MR 482230
- [PR87] Bernadette Perrin-Riou, *Fonctions L p -adiques, théorie d’Iwasawa et points de Heegner*, Bull. Soc. Math. France **115** (1987), no. 4, 399–456. MR 928018
- [Rib83] Kenneth A. Ribet, *Mod p Hecke operators and congruences between modular forms*, Invent. Math. **71** (1983), no. 1, 193–205. MR 688264
- [Roh84] David E Rohrlich, *On L -functions of elliptic curves and cyclotomic towers*, Inventiones mathematicae **75** (1984), no. 3, 409–423.
- [Ser87] Jean-Pierre Serre, *Sur les représentations modulaires de degré 2 de $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$* , Duke Math. J. **54** (1987), no. 1, 179–230. MR 885783
- [Ski16] Christopher Skinner, *Multiplicative reduction and the cyclotomic main conjecture for GL_2* , Pacific J. Math. **283** (2016), no. 1, 171–200. MR 3513846
- [Ski20] ———, *A converse to a theorem of gross, zagier, and kolyvagin*, Annals of Mathematics **191** (2020), no. 2, 329–354.
- [SU14] Christopher Skinner and Eric Urban, *The Iwasawa main conjectures for GL_2* , Inventiones mathematicae **195** (2014), 1–277.
- [SZ14] Christopher Skinner and Wei Zhang, *Indivisibility of Heegner points in the multiplicative case*, arXiv preprint arXiv:1407.1099 (2014).

- [Wan15] Xin Wan, *The Iwasawa main conjecture for Hilbert modular forms*, Forum Math. Sigma **3** (2015), Paper No. e18, 95. MR 3482263
- [Wan20] ———, *Iwasawa main conjecture for Rankin-Selberg p -adic L -functions*, Algebra Number Theory **14** (2020), no. 2, 383–483. MR 4195651
- [Wan21] ———, *Heegner point Kolyvagin system and Iwasawa main conjecture*, Acta Math. Sin., Engl. Ser. **37** (2021), no. 1, 104–120 (English).
- [Zha14] Wei Zhang, *Selmer groups and the indivisibility of heegner points*, Cambridge Journal of Mathematics **2** (2014), no. 2, 191–253.

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA.
Email address: xjyan95@amss.ac.cn

BEIJING INSTITUTE OF MATHEMATICAL SCIENCES AND APPLICATIONS, BEIJING 101408, CHINA.

DEPARTMENT OF MATHEMATICS AND YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY;
Email address: xwzhu@bimsa.cn