

Dephasing-induced jumps in non-Hermitian disordered lattices

Emmanouil T. Kokkinakis^{1,2}, Konstantinos G. Makris^{1,2}, and Eleftherios N. Economou^{1,2}

¹ *Department of Physics, University of Crete, 70013 Heraklion, Greece and*

² *Institute of Electronic Structure and Laser (IESL), FORTH, 71110 Heraklion, Greece*

(Dated: December 31, 2024)

Changes in the wavefunction's phase during propagation in a random Hermitian lattice, a process known as dephasing, results in diffusion rather than Anderson localization. However, when non-Hermiticity is introduced, the wave behavior changes drastically. In particular, we demonstrate that in weakly disordered non-Hermitian lattices, dephasing induces unexpected eigenmode localization which results in abrupt jumps between spatially distant regions. These jumps, which are absent under purely coherent conditions, emerge from the interplay between complex disorder and dephasing.

I. INTRODUCTION

In recent years, photonics has been established as the most successful platform for exploring non-Hermitian systems [1], due to advanced experimental techniques that enable precise manipulation of non-Hermiticity [2–5]. This is in contrast to most other areas of physics, where controlling openness poses substantial practical challenges. The introduction of concepts of parity-time (\mathcal{PT}) symmetry [6–15] and exceptional-points [16–21] in optics has led to an intense research activity in the emerging field of non-Hermitian photonics over the last decade [22–46]. The main approach for realizing non-Hermitian Hamiltonians involves designing structures with spatially distributed loss and/or gain materials. In semiclassical optics, such dissipative and active materials, are easily accessible, unlike in condensed matter or quantum systems. In addition to systems with on-site gain/loss potentials, extensive theoretical and experimental research has focused on non-Hermitian lattices with asymmetric couplings [47–70].

In contrast to these relatively recent advances, Anderson localization is a phenomenon that has been extensively studied for over six decades [71, 72]. As first introduced by P. W. Anderson in his seminal work [73], it predicts the absence of diffusion of waves propagating through lattices with random on-site disorder. Although initially rooted in the context of solid-state physics, Anderson localization has since had significant impact over a wide range of fields, including disordered photonics, Bose-Einstein condensates, imaging, and acoustics [74–77]. In these areas, direct experimental investigations are not hindered by many-body interactions or temperature-dependent effects, allowing for direct observations of localization phenomena. With notable exceptions such as random lasers [78–80], most studies have focused on conservative systems governed by Hermitian Hamiltonians.

Recently, attention has been drawn to non-Hermitian disordered problems [81–105], driven by the synergy of Anderson localization and non-Hermiticity. This novel approach to non-Hermitian Anderson localization [95–98, 101] allows for precisely controlled experiments, opening up a largely unexplored area of localization in com-

plex media. Notable examples of intricate wave dynamics include constant-intensity waves despite strong localization [81, 82] and sudden Anderson jumps [99–103] in disordered or quasiperiodic lattices. These concepts have been experimentally demonstrated in acoustics [106] and optics [107], offering a different perspective in Anderson localization in open systems.

However, wave propagation in conservative lattices is profoundly affected by their interaction with the environment. One possible way to effectively model such an interaction is by randomize the wavefunction's phase, an effect known as dephasing. In periodic systems, such as lattices described by the Haken-Strobl-Reineker (HSR) model [108], dephasing leads to dynamics similar to that of a Brownian particle [109–112]. In one- and two-dimensional disordered lattices, it is well-established that any non-zero level of disorder results in Anderson localization, thereby preventing transport. Nevertheless, studies examining dephasing, evolving disorder, or iterative measurements, suggest that the delicate interference essential to Anderson localization is disrupted, resulting in delocalization and incoherent, diffusive motion through classical inter-site hopping [112–127]. Very recently, the effect of dephasing on non-Hermitian systems [128, 129] and Hermitian quasiperiodic systems beyond the Anderson model [130, 131] has started to gain attention, yet many aspects remain largely unexplored.

This work aims to investigate the effects of dephasing on wave propagation in non-Hermitian lattices with complex on-site random disorder, where Anderson jumps can emerge under coherent dynamics [99–101]. However, in the weak-disorder regime, these jumps are not apparent due to the large spatial extent of different eigenstates that are consequently overlapping. Counter-intuitively, in weakly disordered non-Hermitian lattices, rapid dephasing induces eigenmode localization, resulting in jumpy wavefunction dynamics. We demonstrate that given the potential distributions and initial conditions, the position and duration of each jump can be analytically predicted a priori. Additionally, we discuss the evolution of the derivative of optical power, which governs the system's dynamics, and the degree of localization of eigenmodes, as well as their dependence on the dephasing rate.

II. MODEL UNDER COHERENT AND INCOHERENT DYNAMICS

Our study begins with a one-dimensional (1D) non-Hermitian Anderson lattice consisting of N evanescently coupled waveguides (indexed by $n \in \{1, 2, \dots, N\}$) that have complex on-site (gain and/or loss) strength. Thus, the Hamiltonian of the system is described by

$$\hat{H} = \sum_{n=1}^{N-1} |n\rangle \langle n+1| + |n+1\rangle \langle n| + \sum_{n=1}^N \epsilon_n |n\rangle \langle n|, \quad (1)$$

where $\epsilon_n = a_n + ib_n \in \mathbb{C}$ take random values from a rectangular distribution, with $a_n \in [-W_R/2, W_R/2]$ and $b_n \in [-W_I/2, W_I/2]$. We refer to $W_R, W_I \in \mathbb{R}^+$ as the real and imaginary disorder strength, respectively.

Under the paraxial approximation, the evolution of a wavefunction $|\psi\rangle \equiv \sum_{n=1}^N \psi_n |n\rangle$ in this lattice is described by the coupled-mode equations

$$\hat{H} |\psi\rangle = -i \frac{d|\psi\rangle}{dz}, \quad (2)$$

where we assume that the waveguide channels form a linear chain with open boundary conditions (OBCs), i.e. $\psi_0 = \psi_{N+1} = 0$.

The eigenstates of the right eigenvalue problem associated with this Hamiltonian are expressed as $|\psi_j(z)\rangle = |u_j^R\rangle e^{i\omega_j z}$, where $|u_j^R\rangle$ is one right eigenvector of the set $\{|u_m^R\rangle\}$, and ω_j is the corresponding eigenvalue, i.e., $\hat{H} |u_j^R\rangle = \omega_j |u_j^R\rangle$. The left eigenvalue problem associated with this Hamiltonian reads as $\hat{H}^\dagger |u_j^L\rangle = \omega_j^* |u_j^L\rangle$. The right and left eigenvectors of the Hamiltonian satisfy the biorthogonality condition $\langle u_k^L | u_j^R \rangle = \delta_{kj}$ while its symmetry ($\hat{H} = \hat{H}^T$) implies that $|u_j^L\rangle^* = |u_j^R\rangle$. Therefore, $|\psi\rangle$ can be expressed as

$$|\psi(z)\rangle = \sum_{j=1}^N c_{j,0} e^{-\text{Im}(\omega_j)z} e^{i\text{Re}(\omega_j)z} |u_j^R\rangle \quad (3)$$

where $c_{j,0} \equiv \langle u_j^L | \psi(0) \rangle$, and $|\psi(0)\rangle = \sum_{n=1}^N \psi_n(0) |n\rangle$ is the initial condition. The non-Hermiticity of the Hamiltonian results in the non-conservation of optical power, defined as $\mathcal{P}(z) \equiv \langle \psi | \psi \rangle$. To capture the wavepacket's dynamics, we introduce a normalized wavefunction at each propagation distance z , given by $|\phi\rangle \equiv |\psi\rangle / \sqrt{\mathcal{P}(z)}$. This normalization can be implemented experimentally [100] and ensures that the spatial profile of the wavefunction remains unchanged.

Regarding the dephasing, which is the focus of our study, we consider randomizing the phase of the wavefunction in a periodic fashion at each site n at propagation distances $z_a = a \cdot l$, where $a \in \mathbb{N}$ and l is the dephasing period. This is mathematically expressed as

$$\psi_n(z_a^+) = e^{i\theta_n^{(a)}} \psi_n(z_a^-). \quad (4)$$

where $\theta_n^{(a)} \in [0, 2\pi]$ are randomly selected phases. After statistical averaging over multiple realizations of the totally uncorrelated stochastic phases, the dynamical evolution of the average probability density $P_n(z) \equiv \overline{|\psi_n(z)|^2}$ is described by

$$P_n(z = al) = \mathcal{S}^a P_n(z = 0), \quad (5)$$

where $\mathcal{S}(l)$ is the propagator matrix of the fully incoherent system, with elements given by $\mathcal{S}_{kj}(l) = |(e^{iHl})_{kj}|^2$. Thus, the matrix \mathcal{S} , which we will refer to as the incoherent propagator, is non-negative since $\mathcal{S}_{kj} \geq 0$. Since the Hamiltonian \hat{H} is symmetric ($\hat{H} = \hat{H}^T$), the propagator matrix $U \equiv e^{iHl}$ is also symmetric, i.e. $(e^{iHl})_{kj} = (e^{iHl})_{jk}$. Consequently, \mathcal{S} is symmetric since $\mathcal{S}_{kj} = |(e^{iHl})_{kj}|^2 = |(e^{iHl})_{jk}|^2 = \mathcal{S}_{jk}$. So, the eigenvalue equation for the symmetric and real-valued (Hermitian) matrix \mathcal{S} is

$$\mathcal{S} |v_j\rangle = \beta_j |v_j\rangle, \quad (6)$$

From Eq. 5, it follows that

$$P(z = al) = \sum_{j=1}^N d_{j,0} \beta_j^a |v_j\rangle, \quad (7)$$

where $P(z) \equiv \sum_{n=1}^N P_n(z) |n\rangle$ and $d_{j,0} \equiv \langle v_j | P(z = 0) \rangle$ (the derivation of Eqs. 5 and 7 is provided in Appendix A).

In the case of $W_I = 0$, where \hat{H} is Hermitian ($\hat{H}^\dagger = \hat{H}$), the propagator matrix $U = e^{i\hat{H}l}$ is unitary ($UU^\dagger = \hat{1}$). Consequently, the relation $\sum_{k=1}^N \mathcal{S}_{kj} = \sum_{j=1}^N \mathcal{S}_{kj} = 1$ for all rows and columns indexed by k and j holds, and since \mathcal{S} is non-negative, it is by definition a doubly stochastic matrix. Thus, it can be directly shown that \mathcal{S} has an eigenvalue $\beta_1 = 1$ corresponding to the eigenstate $|v_1\rangle \sim (1, 1, \dots, 1)^T$. All other eigenvalues of \mathcal{S} have absolute values $|\beta_m| \leq 1$, thus β_1 is the so called Perron-Frobenius eigenvalue [132]. Therefore, it follows from Eq. 7 that the projections of the average probability function onto eigenstates $\{|v_m\rangle\}$ for $m \neq 1$ eventually vanish, and in the limit $z \rightarrow \infty$, any initial excitation diffracts and adopts a homogeneous distribution in the lattice, i.e., $|P(z \rightarrow \infty)\rangle = |v_1\rangle$.

For $W_I > 0$, the non-unitarity of U results in a non-stochastic matrix \mathcal{S} , thus lifting the constraint $|\beta_m| \leq 1$, and the dynamical characteristics of the system change significantly.

At this point, it is useful to introduce the inverse participation ratio (IPR), a widely used measure of wavefunction localization, that we will use in the following sections. For a one-dimensional wavefunction $|\xi\rangle$, the IPR is defined as

$$\text{IPR} \equiv \frac{\sum_{n=1}^N |\xi_n|^4}{\left(\sum_{n=1}^N |\xi_n|^2\right)^2} \quad (8)$$

where $\xi_n \equiv \langle n|\xi \rangle$. This metric ranges from $1/N$ for a fully extended state, where $\xi_n = 1/\sqrt{N}$ for all n , to 1 for a completely localized state, where $\xi_n = \delta_{nm}$.

III. ABSENCE OF JUMPS UNDER COHERENT PROPAGATION AT WEAK DISORDER

Before further discussing the effect of dephasing, it is beneficial to review what happens under fully coherent conditions in a weakly non-Hermitian disordered lattice. In fact, a relevant phenomenon in such lattices, is the appearance of abrupt jumps between spatially distant regions of the lattice [99–101], as depicted in Fig. 1a. These jumps occur between eigenstates that are localized in different regions and exhibit unequal rates of amplification or dissipation.

The dynamics in such systems are governed by the evolution of the projection coefficients $|c_j| = |c_{j,0}|e^{-\text{Im}(\omega_j)z}$ of Eq. 3. A jump occurs when the projection $|c_k|$ of an eigenstate $|u_k^R\rangle$ surpasses the projection $|c_r|$ of the previously dominant eigenstate $|u_r^R\rangle$ (Fig. 1b), where the condition $\text{Im}(\omega_k) < \text{Im}(\omega_r)$ holds. Ultimately, the eigenstate with the highest gain or the lowest loss ($\min\{\text{Im}(\omega_j)\}$) dominates the system's dynamics (Fig. 1c). However, these transitions are abrupt only in the high disorder regime, as lower disorder typically results in the presence of more extended eigenstates, which are spatially overlapping,

A comparison of the evolution of the normalized wavefunction $|\phi\rangle$ for a single realization of high and low complex disorder, $W_R = W_I = 6$ and $W_R = W_I = 1$, respectively, in a lattice of $N = 50$ waveguide channels under coherent dynamics (Eq. 2) is shown in Fig. 1a (for strong disorder) and Fig. 1d (for low disorder). For both cases, we have considered a single-channel excitation at the middle of the lattice as initial condition, i.e., $\psi_n(0) = \delta_{n,25}$. In the high disorder case, the jump between distant regions of the lattice, seen in Fig. 1a, corresponds to the crossing between dominant eigenstates, as indicated in Fig. 1b. These eigenstates are highly localized, as evidenced by their large IPR (Fig. 1c).

In the case of weak disorder, although there are two propagation distances at which transitions between dominant modes occur during the dynamics (Fig. 1e), no jumps are observed, as shown in Fig. 1d. The absence of jumps arises because none of the three involved modes (nor any other eigenmodes of \hat{H}) are strongly localized. This is evidenced by their very low (IPR) values, as illustrated in Fig. 1f.

IV. EMERGENCE OF JUMPS DUE TO DEPHASING

In the regime of fully incoherent dynamics, the evolution of the average probability density $P(z)$ is described by Eq. 7. Thus, the magnitude of the projections

$d_j \equiv \langle v_j|P(z)\rangle$ of $P(z)$ on the eigenstates $|v_j\rangle$ of the incoherent propagator \mathcal{S} evolve as $|d_j| = |d_{j,0}|\beta_j^{z/l}$ so that

$$\ln|d_j| = \ln|d_{j,0}| + \frac{\ln\beta_j}{l} \cdot z \quad (9)$$

From Eq. 9 it is evident that after a finite propagation distance z_{cr} the eigenmode $|v_{\max}\rangle$ with largest eigenvalue $\beta_{\max} \equiv \max\{\beta_j\}$ will dominate the dynamics, giving the most significant contribution to the expansion of Eq. 7. Under fully incoherent dynamics, the average optical power of the system can be defined as

$$\overline{\mathcal{P}}(z) \equiv \sum_{n=1}^N P_n(z) \stackrel{(7)}{=} \sum_{n=1}^N \sum_{j=1}^N d_{j,0} v_{n,j} \beta_j^{z/l} \quad (10)$$

where $v_{n,j} \equiv \langle n|v_j\rangle$, so when one mode $|v_m\rangle$ dominates the dynamics, the above Eq. 10 simplifies to

$$\overline{\mathcal{P}} \approx \beta_m^{z/l} \sum_{n=1}^N d_{m,0} v_{n,m} \quad (11)$$

and it directly follows that

$$\frac{d \ln \overline{\mathcal{P}}(z)}{dz} = \frac{\ln \beta_m}{l} \quad (12)$$

In Fig. 2a, it is shown that for the same initial condition and disorder realization as in the bottom row of Fig. 1, the wave dynamics changes dramatically due to dephasing with a period $l = 0.01$. This is evident from the abrupt transitions between highly localized regions, which, as shown in Fig. 2b, occur at propagation distances corresponding to crossings between dominant projection coefficients $|d_m|$, whose evolution is described by Eq. 9. In Fig. 2c, the real spectrum of \mathcal{S} is illustrated, where as indicated by the high values of IPR, the corresponding eigenmodes $|v_j\rangle$, are highly localized. Thus, in the fully incoherent regime, the localization of the eigenmodes of \mathcal{S} facilitates the occurrence of abrupt jumps. Additionally, as shown in Figure 3b, the derivative of the logarithm of optical power $\mathcal{P}(z)$ for a single realization of random phases matches the expected values from Eq. 12. The latter are calculated based on the eigenvalue β_m corresponding to the dominant eigenmode $|v_m\rangle$ for each propagation interval.

It is important to note that the evolution of both the wavefunction $|\phi\rangle$ and the derivative of the optical power, $d \ln \mathcal{P}/dz$, (as shown in Fig. 2a and Fig. 3b) are numerically calculated for a single realization of random phases. Despite this, they perfectly agree with the theoretical predictions given by Eq. 9 and Eq. 12, respectively. Although the aforementioned equations were derived under the assumption of statistical averaging, the high dephasing frequency $1/l = 100$ which is employed in this study, effectively minimizes any significant differences in the wavefunction's evolution across different realizations of random phases.




FIG. 1. Comparison of coherent dynamics (Eq. 2) for single realizations of strong disorder ($W_R = W_I = 6$, top row) and weak disorder ($W_R = W_I = 1$, bottom row) in lattices with $N = 50$ channels. (a)/(d) Evolution of the normalized wavefunction $|\phi\rangle$ under single-channel excitation at $n = 25$ for the strong/weak disorder case, respectively. (b)/(e) Logarithmic magnitudes of the projection coefficients $\ln|c_j|$ for selected eigenstates, sorted by descending $\text{Im}(\omega_j)$, in the strong/weak disorder case. (c)/(f) Eigenvalue spectrum of \hat{H} in the complex plane for the strong/weak disorder case, with each eigenvalue ω_j colored according to the inverse participation ratio (IPR) of the corresponding right eigenstate $|u_j^R\rangle$. Arrows indicate the crossings between dominant projection coefficients as seen in (b) and (e).

V. DEPENDENCE OF POWER DERIVATIVE AND DEGREE OF LOCALIZATION ON DEPHASING RATE

Since \hat{H} is non-Hermitian, the optical power \mathcal{P} is not conserved under incoherent dynamics as well. In Fig. 3, we calculate the evolution of the derivative of the logarithm of optical power for the same single realization of weak complex disorder ($W_R = W_I = 1$) as in Fig. 1(bottom row) and Fig. 2, considering both coherent (Fig. 3a) and incoherent dynamics with rapid dephasing ($l = 0.01$, Fig. 3b). It is clear that dephasing significantly changes the evolution of optical power. This observation naturally raises the question of whether and how this physical quantity depends on the dephasing period l in the fully incoherent regime. According to Eq. 12, it is expected that beyond a finite distance $z_{cr}(l)$, where the eigenmode $|u_{\max}\rangle$ becomes dominant, the average derivative of the optical power logarithm will remain constant. Consequently, this long-term power derivative depends on the dephasing period l and the eigenvalue β_{\max} , which is itself a function of l , according to Eq. 12.

As shown in Fig. 4a, the long-term rate of power derivative $d \ln \overline{\mathcal{P}}(z > z_{cr})/dz = \ln \beta_{\max}/l$ for the same single potential realization as Fig. 3, rises as the dephasing period l decreases. This rate eventually saturates for dephasing period less than $l \sim 10^{-3}$.

It is also important to examine how the degree of localization of the eigenstates $|v_j\rangle$ of the incoherent propagator \mathcal{S} , as quantified by the inverse participation ratio IPR, depends on the dephasing period l . As shown in Fig. 4b, using the same weak ($W_R = W_I = 1$) disorder realization as in the previous figures, decreasing the dephasing period l increases the mean inverse participation ratio (IPR) of the eigenstates $|v_j\rangle$, such that $\langle \text{IPR} \rangle \rightarrow 1$ for $l < 10^{-4}$. This indicates that below a certain value of the dephasing period l , all eigenstates of \mathcal{S} become perfectly localized at single sites. Consequently, the incoherent dynamics of a single-channel excitation in such a lattice will primarily exhibit a jumpy evolution between distant single sites.

This behavior is not unique to the specific potential realization used in the previous analysis. As illustrated in Fig. 5, complete localization of the eigenmodes of




FIG. 2. Single realization (same with bottom row of Fig.1) of weak disorder ($W_R = W_I = 1$) under fully incoherent dynamics (Eq. 4), for a lattice of $N = 50$ channels and dephasing period $l = 0.01$ (a) Evolution of normalized wavefunction $|\phi\rangle$ under single channel excitation at $n = 25$, for a single random-phases realization (b) Logarithmic magnitudes of projection coefficients $\ln|d_j|$ for specific eigenstates, sorted in ascending β_j (c) Eigenvalue spectrum of \mathcal{S} . Each point β_j has color associated with the IPR of the corresponding eigenstate $|v_j\rangle$. Arrows indicate the crossings between dominant projection coefficients as seen in (b).




FIG. 3. Evolution of the derivative of the power logarithm for a single realization of weak disorder ($W_R = W_I = 1$) corresponding to Fig. 1 (bottom row) and Fig. 2, under (a) coherent dynamics (no dephasing) and (b) incoherent dynamics with dephasing period ($l = 0.01$). At (b) the blue line represents the result from a single random-phase realization, while the red lines indicate the expected values from Eq. 12.




FIG. 4. (a) Long-term average power logarithm derivative as predicted by Eq. 12 for the potential realization in Fig. 1 (bottom row) and Fig. 2. As l decreases, it saturates at $d \ln \bar{\mathcal{P}}(z)/dz \approx 0.8852$. (b) Mean inverse participation ratio $\langle \text{IPR} \rangle$ of the eigenmodes $|v_j\rangle$ of the incoherent propagator $\mathcal{S}(l)$, for the same potential realization. As l decreases, this saturates at $\langle \text{IPR} \rangle = 1$, indicating localization to single sites.

VI. DISCUSSION AND CONCLUSIONS

the incoherent propagator occurs for all values of complex disorder strength $W \equiv W_R = W_I$, beyond a critical dephasing period $l_{cr}(W)$. For each value of disorder strength depicted in Fig. 5, 1000 different realizations of the random complex disorder were averaged to calculate the mean inverse participation ratio of the eigenmodes of \mathcal{S} , ensuring statistically reliable results.

In conclusion, we have investigated the impact of dephasing on wave propagation in non-Hermitian lattices with complex on-site weak disorder. Our analysis reveals that, while Anderson jumps are absent in the coherent case, dephasing, conversely, induces jumpy propagation. These phenomena should be experimentally observable in currently existing photonic discrete-time quantum walk




FIG. 5. Statistically averaged mean inverse participation ratio $\overline{\text{IPR}}$ of the eigenmodes $|v_j\rangle$ of the incoherent propagator $\mathcal{S}(l)$, for various values of dephasing period l and disorder strength $W_R = W_I \equiv W$. For each value of W , 1000 random realizations of potential are used for statistical averaging.

platforms that use light-pulse propagation in synthetic-mesh lattices [100]. Importantly, we demonstrate that the position and duration of each jump can be predicted analytically based on the given potential distributions and initial conditions. That comes in complete contradiction to the effect of dephasing on Hermitian disordered systems, where its presence destroys Anderson localization and leads to diffusion. Additionally, we examine the behavior of the optical power and show the saturation of its derivative with increasing dephasing frequency. The effect of dephasing rate on eigenmode localization is also analyzed. These findings contribute to a deeper understanding of wave dynamics in non-Hermitian systems and are relevant for future theoretical and experimental studies in this area.

ACKNOWLEDGMENTS

The authors acknowledge the financial support provided by the European Research Council (ERC) under the Consolidator Grant Agreement No. 101045135 (Beyond_Anderson). This research project was also co-funded by the Stavros Niarchos Foundation (SNF) and the Hellenic Foundation for Research and Innovation (H.F.R.I.) through the 5th Call of the ‘‘Science and Society’’ Action, titled ‘‘Always strive for excellence – Theodoros Papazoglou’’ (Project Number: 11496, ‘‘PSEUDOTOPPOS’’).

Appendix A: Incoherent propagation equation

As also discussed in [129], it follows from Eq. 2 and Eq. 4, that the wavefunction’s values at site n between two consecutive dephasing events, i.e. $z_{a-1} = (a-1)l$

and $z_a = al$, where $a \in \mathbb{N}$, are connected through

$$\psi_n(z_a) = e^{i\theta_n^{(a)}} \sum_{m=1}^N U_{n,m} \psi_m(z_{a-1}) \quad (\text{A1})$$

and $U \equiv e^{iHl}$. Recursively, the following equations hold

$$\begin{aligned} \psi_n(z_a) &= e^{i\theta_n^{(a)}} \sum_{m_a=1}^N U_{n,m_a} \psi_{m_a}(z_{a-1}) \\ \psi_{m_a}(z_{a-1}) &= e^{i\theta_{m_a}^{(a-1)}} \sum_{m_{a-1}=1}^N U_{m_a,m_{a-1}} \psi_{m_{a-1}}(z_{a-2}) \\ &\vdots \\ \psi_{m_2}(z_1) &= e^{i\theta_{m_2}^{(1)}} \sum_{m_1=1}^N U_{m_2,m_1} \psi_{m_1}(z_0) \end{aligned} \quad (\text{A2})$$

so by consecutive substitutions one gets the expressions

$$\begin{aligned} \psi_n(z_a) &= \sum_{m_1, m_2, \dots, m_a}^N e^{i\theta_n^{(a)} + i\theta_{m_a}^{(a-1)} + \dots + i\theta_{m_2}^{(1)}} \times \\ &U_{n,m_a} U_{m_a,m_{a-1}} \dots U_{m_2,m_1} \psi_{m_1}(z_0) \end{aligned} \quad (\text{A3})$$

and

$$\begin{aligned} |\psi_n(z_a)|^2 &= \sum_{m_1, \rho_1, \dots, m_a, \rho_a}^N e^{i\theta_{m_a}^{(a-1)} - i\theta_{\rho_a}^{(a-1)} + \dots + i\theta_{m_2}^{(1)} - i\theta_{\rho_2}^{(1)}} \times \\ &U_{n,m_a} U_{n,\rho_a}^* \dots U_{m_2,m_1} U_{\rho_2,\rho_1}^* \psi_{m_1}(z_0) \psi_{\rho_1}^*(z_0) \end{aligned} \quad (\text{A4})$$

After statistical averaging over different realizations of the totally uncorrelated stochastic phases $\theta_n^{(a)} \in [0, 2\pi]$, only terms with $m_i = \rho_i$ survive in the sum of Eq. A4, so

$$P_n(z) \equiv \overline{|\psi_n(z)|^2} = \sum_{m_1, \dots, m_a}^N |U_{n,m_a}|^2 \dots |U_{m_2,m_1}|^2 P_{m_1}(z_0) \quad (\text{A5})$$

From this equation, it readily holds that

$$P_n(z_a) = \sum_{m=1}^N |U_{n,m}|^2 P_m(z_{a-1}) \quad (\text{A6})$$

This difference equation describes the dynamics of any system in the totally incoherent regime and is mathematically equivalent to

$$\psi_n(z_a) = \sum_{m=1}^N U_{n,m} \psi_m(z_{a-1}) \quad (\text{A7})$$

which is the well known expression for a system under coherent dynamics described by the Schrödinger-type equation Eq. 2. Thus, in the fully incoherent regime, the proper propagator matrix \mathcal{S} for the dynamics of the system has elements given by $\mathcal{S}_{nm} = |U_{n,m}|^2$. Recursively from Eq.A5, we conclude to the expression

$$P_n(z = al) = \mathcal{S}^a P_n(z = 0), \quad (\text{A8})$$

which is Eq.5. Applying Cayley-Hamilton's theorem to the eigenvalue equation of the incoherent propagator \mathcal{S} we have

$$\mathcal{S}^a |v_j\rangle = \beta_j^a |v_j\rangle, \quad (\text{A9})$$

We can express $P(z=0) \equiv \sum_{n=1}^N P_n(z=0) |n\rangle$ in terms of the eigenstates of \mathcal{S} as

$$P(z=0) = \sum_{j=1}^N d_{j,0} |v_j\rangle, \quad (\text{A10})$$

where $d_{j,0} \equiv \langle v_j | P(z=0) \rangle$. From Eqs. A8, A9 and A10 it follows that

$$P(z=al) = \sum_{j=1}^N d_{j,0} \beta_j^a |v_j\rangle, \quad (\text{A11})$$

which is Eq. 7 of the main text.

-
- [1] D. N. Christodoulides, F. Lederer, and Y. Silberberg, Discretizing light behaviour in linear and nonlinear waveguide lattices, *Nature (London)* **424**, 817 (2003).
- [2] D. S. Wiersma, P. Bartolini, A. Lagendijk, and R. B. Rijnveld, Localization of light in a disordered medium, *Nature (London)* **390**, 671 (1997).
- [3] A. Lagendijk, B. v. Tiggelen, and D. S. Wiersma, Fifty years of Anderson localization, *Phys. Today* **62**, 24 (2009).
- [4] T. Schwartz, G. Bartal, S. Fishman, and M. Segev, Transport and Anderson localization in disordered two-dimensional photonic lattices, *Nature (London)* **446**, 52 (2007).
- [5] Y. Lahini, A. Avidan, F. Pozzi, M. Sorel, R. Morandotti, D. N. Christodoulides, and Y. Silberberg, Anderson Localization and Nonlinearity in One-Dimensional Disordered Photonic Lattices, *Phys. Rev. Lett.* **100**, 013906 (2008).
- [6] C. M. Bender and S. Boettcher, Real Spectra in Non-Hermitian Hamiltonians Having \mathcal{PT} Symmetry, *Phys. Rev. Lett.* **80**, 5243 (1998).
- [7] C. M. Bender, S. Boettcher, P. N. Meisinger, \mathcal{PT} symmetric quantum mechanics, *J. Math. Phys.* **40**, 2201 (1999).
- [8] C. M. Bender, D. C. Brody, and H. F. Jones, Complex Extension of Quantum Mechanics, *Phys. Rev. Lett.* **89**, 270401 (2002).
- [9] K. G. Makris, R. El-Ganainy, D. N. Christodoulides, and Z. H. Musslimani, Beam Dynamics in \mathcal{PT} Symmetric Optical Lattices, *Phys. Rev. Lett.* **100**, 103904 (2008).
- [10] R. El-Ganainy, K. G. Makris, D. N. Christodoulides, and Z. H. Musslimani, Theory of coupled optical \mathcal{PT} -symmetric structures, *Opt.Lett.* **32**, 2632 (2007).
- [11] Z. H. Musslimani, K. G. Makris, R. El-Ganainy, and D. N. Christodoulides, Optical Solitons in \mathcal{PT} Periodic Potentials, *Phys. Rev. Lett.* **100**, 030402 (2008).
- [12] A. Guo, G. J. Salamo, D. Duchesne, R. Morandotti, M. Volatier-Ravat, V. Aimez, G. A. Siviloglou, and D. N. Christodoulides, Observation of \mathcal{PT} -Symmetry Breaking in Complex Optical Potentials, *Phys. Rev. Lett.* **103**, 093902 (2009).
- [13] C. E. Rüter, K. G. Makris, R. El-Ganainy, D. N. Christodoulides, M. Segev, and D. Kip, Observation of parity-time symmetry in optics, *Nat. Physics* **6**, 192 (2010).
- [14] K. G. Makris, R. El-Ganainy, D. N. Christodoulides, and Z. H. Musslimani, \mathcal{PT} -symmetric optical lattices, *Phys. Rev. A* **81**, 063807 (2010).
- [15] C. M. Bender, Making sense of non-Hermitian Hamiltonians, *Rep. Prog. Phys.* **70**, 947 (2007).
- [16] N. Moiseyev, *Non-Hermitian Quantum Mechanics* (Cambridge, New York, 2011).
- [17] M. V. Berry, Physics of non-Hermitian degeneracies, *Czech. J. Phys.* **54**, 1039 (2004).
- [18] W. D. Heiss, Exceptional points of non-Hermitian operators, *J. Phys. A: Math. Gen.* **37**, 2455 (2004).
- [19] J. Wiersig, S.-W. Kim, and M. Hentschel, Asymmetric scattering and nonorthogonal mode patterns in optical microspirals, *Phys. Rev. A* **78**, 053809 (2008).
- [20] S. B. Lee, J. Yang, S. Moon, S. Y. Lee, J. B. Shim, S. W. Kim, J. H. Lee, and K. An, Observation of an Exceptional Point in a Chaotic Optical Microcavity, *Phys. Rev. Lett.* **103**, 134101 (2009).
- [21] S. M. A. Miri and A. Alu, Exceptional points in optics and photonics, *Science* **363**, eaar7709 (2019).
- [22] R. El-Ganainy, K. G. Makris, M. Khajavikhan, Z. H. Musslimani, S. Rotter, and D. N. Christodoulides, Non-Hermitian physics and \mathcal{PT} symmetry, *Nat. Physics* **14**, 11-19 (2018).
- [23] S. Longhi, Parity-time symmetry meets photonics: A new twist in non-Hermitian optics, *Europhys. Lett.* **120**, 64001 (2018).
- [24] A. Regensburger, C. Bersch, M. A. Miri, G. Onishchukov, D. N. Christodoulides, and U. Peschel, Parity-time synthetic photonic lattices, *Nature (London)* **488**, 167 (2012).
- [25] L. Feng, YL. Xu, W. S. Fegadolli, MH. Lu, J. E. Oliveira, V. R. Almeida, YF. Chen, and A. Scherer, Experimental demonstration of a unidirectional reflectionless parity-time metamaterial at optical frequencies, *Nat. Mater.* **12**, 108 (2013).
- [26] J. Zhang, B. Peng, Ş. K. Özdemir, K. Pichler, D. O. Krimer, G. Zhao, F. Nori, Yx. Liu, S. Rotter, and L. Yang, A phonon laser operating at an exceptional point, *Nat. Photonics* **12**, 479 (2018).
- [27] H. Hodaei, M. A. Miri, M. Heinrich, D. N. Christodoulides, and M. Khajavikhan, Parity-time-symmetric microring lasers, *Science* **346**, 975 (2014).
- [28] V. V. Konotop, J. Yang, and D. A. Zezyulin, Nonlinear waves in \mathcal{PT} -symmetric systems, *Rev. Mod. Phys.* **88**, 035002 (2016).
- [29] L. Feng, R. El-Ganainy, and L. Ge, Non-Hermitian photonics based on parity-time symmetry, *Nat. Photon.* **11**, 752 (2017).

- [30] D. F. Pile and D. N. Christodoulides, Gaining with loss, *Nat. Photon.* **11**, 742 (2017)
- [31] G. Gbur and K. G. Makris, Introduction to non-Hermitian photonics in complex media: \mathcal{PT} -symmetry and beyond, *Photon. Res.* **6**, PTS1 (2018)
- [32] S. K. Özdemir, S. Rotter, F. Nori, and L. Yang, Parity-time symmetry and exceptional points in photonics, *Nat. Mater.* **18**, 783 (2019).
- [33] Y. Chong, L. Ge, and A. D. Stone, \mathcal{PT} -Symmetry Breaking and Laser-Absorber Modes in Optical Scattering Systems, *Phys. Rev. Lett.* **106**, 093902 (2011).
- [34] L. Ge, Y. Chong, and A. D. Stone, Conservation relations and anisotropic transmission resonances in one-dimensional \mathcal{PT} -symmetric photonic heterostructures, *Phys. Rev. A* **85**, 023802 (2012).
- [35] M. Wimmer, A. Regensburger, M.-A. Miri, C. Bersch, D. N. Christodoulides, and U. Peschel, Observation of optical solitons in \mathcal{PT} -symmetric lattices, *Nat. Commun.* **6**, 7782 (2015).
- [36] A. Szameit, M. C. Rechtsman, O. Bahat-Treidel, and M. Segev, \mathcal{PT} -symmetry in honeycomb photonic lattices, *Phys. Rev. A* **84**, 021806 (2011).
- [37] J. M. Zeuner, M. C. Rechtsman, Y. Plotnik, Y. Lumer, S. Nolte, M. S. Rudner, M. Segev, and A. Szameit, Observation of a Topological Transition in the Bulk of a Non-Hermitian System, *Phys. Rev. Lett.* **115**, 040402 (2015).
- [38] B. Peng, S. K. Özdemir, F. Lei, F. Monifi, M. Gianfreda, G. L. Long, S. Fan, F. Nori, C. M. Bender, and L. Yang, Parity-time-symmetric whispering-gallery microcavities, *Nat. Phys.* **10**, 394 (2014).
- [39] L. Feng, Z. J. Wong, R.-M. Ma, Y. Wang, and X. Zhang, Single-mode laser by parity-time symmetry breaking, *Science* **346**, 972 (2014).
- [40] B. Peng, S. Özdemir, S. Rotter, H. Yilmaz, M. Liertzer, F. Monifi, C. Bender, F. Nori, and L. Yang, Loss-induced suppression and revival of lasing, *Science* **346**, 328 (2014).
- [41] S. Assaworarith, X. Yu, and S. Fan, Robust wireless power transfer using a nonlinear parity-time-symmetric circuit, *Nature (London)* **546**, 387 (2017).
- [42] K. G. Makris, Z. H. Musslimani, D. N. Christodoulides, and S. Rotter, Constant-intensity waves and their modulation instability in non-Hermitian potentials, *Nat. Commun.* **6**, 7257 (2015).
- [43] K. G. Makris, Z. H. Musslimani, D. N. Christodoulides, and S. Rotter, Constant Intensity Supermodes in Non-Hermitian Lattices, *IEEE J. Sel. Top. Quantum Electron.* **22**, 42 (2016).
- [44] A. Brandstötter, K. G. Makris, and S. Rotter, and S. Rotter, Scattering-free pulse propagation through invisible non-Hermitian media, *Phys. Rev. B* **99**, 115402 (2019).
- [45] S. Longhi, Topological Phase Transition in non-Hermitian Quasicrystals, *Phys. Rev. Lett.* **122**, 237601 (2019)
- [46] I. Komis and K. G. Makris, Angular excitation of exceptional points and pseudospectra of photonic lattices, *J. Phys. Photonics* **6**, 045011 (2024)
- [47] N. Hatano and D. R. Nelson, Localization Transitions in Non-Hermitian Quantum Mechanics, *Phys. Rev. Lett.* **77**, 570 (1996).
- [48] N. Hatano and D. R. Nelson, Vortex pinning and non-Hermitian quantum mechanics, *Phys. Rev. B* **56**, 8651 (1997).
- [49] H. Jiang, L.-J. Lang, C. Yang, S.-L. Zhu, and S. Chen, Interplay of non-Hermitian skin effects and Anderson localization in nonreciprocal quasiperiodic lattices, *Phys. Rev. B* **100**, 054301 (2019).
- [50] Y. G. Liu, P. S. Jung, M. Parto, D. N. Christodoulides, and M. Khajavikhan, Gain-induced topological response via tailored long-range interactions, *Nat. Physics* **17**, 704 (2021).
- [51] Y. G. Liu, Y. Wei, O. Hemmatyar, G. G. Pyrialakos, P. S. Jung, D. N. Christodoulides, and M. Khajavikhan, Complex skin modes in non-Hermitian coupled laser arrays. *Light: Sci.Appl.* **11**, 336 (2022).
- [52] Z. Gao, X. Qiao, M. Pan, S. Wu, J. Yim, K. Chen, B. Midya, L. Ge, and L. Feng, Two-Dimensional Reconfigurable Non-Hermitian Gauged Laser Array, *Phys. Rev. Lett.* **130**, 263801 (2023).
- [53] S. Yao and Z. Wang, Edge States and Topological Invariants of Non-Hermitian Systems, *Phys. Rev. Lett.* **121**, 086803 (2018).
- [54] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa, and M. Ueda, Topological Phases of Non-Hermitian Systems, *Phys. Rev. X* **8**, 031079 (2018).
- [55] T. E. Lee, Anomalous Edge State in a Non-Hermitian Lattice, *Phys. Rev. Lett.* **116**, 133903 (2016).
- [56] G. Harari, M. A. Bandres, Y. Lumer, M. C. Rechtsman, Y. D. Chong, M. Khajavikhan, D. N. Christodoulides, and M. Segev, Topological insulator laser: theory, *Science* **359**, 4003 (2018).
- [57] M. A. Bandres, S. Wittek, G. Harari, M. Parto, J. Ren, M. Segev, D. N. Christodoulides, and M. Khajavikhan, Topological insulator laser: Experiments, *Science* **359**, 4005 (2018).
- [58] N. Okuma, K. Kawabata, K. Shiozaki, and M. Sato, Topological Origin of Non-Hermitian Skin Effects, *Phys. Rev. Lett.* **124**, 086801 (2020).
- [59] L. Li, C. H. Lee, S. Mu, and J. Gong, Critical non-Hermitian skin effect, *Nat. Commun.*, **11**, 5491 (2020).
- [60] S. Longhi, Spectral deformations in non-Hermitian lattices with disorder and skin effect: A solvable model, *Phys. Rev. B* **103**, 144202 (2021).
- [61] M. Ezawa, Dynamical nonlinear higher-order non-Hermitian skin effects and topological trap-skin phase, *Phys. Rev. B* **105**, 125421 (2022).
- [62] W. N. Faugno and T. Ozawa, Interaction-Induced Non-Hermitian Topological Phases from a Dynamical Gauge Field, *Phys. Rev. Lett.* **129**, 180401 (2022).
- [63] B. Zhu, Q. Wang, D. Leykam, H. Xue, Q. J. Wang, and Y. D. Chong, Anomalous Single-Mode Lasing Induced by Nonlinearity and the Non-Hermitian Skin Effect *Phys. Rev. Lett.* **129**, 013903 (2022).
- [64] S. Weidemann, M. Kremer, T. Helbig, T. Hofmann, A. Stegmaier, M. Greiter, R. Thomale, and A. Szameit, Topological funneling of light, *Science* **368**, 311 (2020).
- [65] I. Komis, Z. H. Musslimani, and K. G. Makris, Skin solitons, *Opt. Lett.* **48**, 6525 (2023).
- [66] S. Longhi, Phase transitions in a non-Hermitian Aubry - André - Harper model, *Phys. Rev. B*, **103**, 054203 (2021).
- [67] H. Chen and Y. Zhang, Real spectra in one-dimensional single-band non-Hermitian Hamiltonians, *Phys. Rev. B* **108** 115114 (2023).

- [68] S. Gandhi and J. N. Bandyopadhyay, Topological triple phase transition in non-hermitian quasicrystals with complex asymmetric hopping, *Phys. Rev. B* **108**, 014204 (2023).
- [69] H. Xiao and Q.-B. Zeng, Coexistence of non-hermitian skin effect and extended states in one-dimensional non-reciprocal lattices, *Phys. Rev. B* **110**, 024205 (2024).
- [70] E. T. Kokkinakis, K. G. Makris, and E. N. Economou, Anderson localization versus hopping asymmetry in a disordered lattice, *Phys. Rev. A* **110**, 053517 (2024).
- [71] E. Akkermans and G. Montambaux, *Mesoscopic Physics of Electrons and Photons*. (Cambridge University Press, 2007).
- [72] P. W. Anderson, The question of classical localization A theory of white paint?, *Philos. Mag. B* **52**, 505 (1985).
- [73] P. W. Anderson, Absence of Diffusion in Certain Random Lattices, *Phys. Rev.* **109**, 1492 (1958).
- [74] M. Segev, Y. Silberberg, and D. N. Christodoulides, Anderson localization of light, *Nat. Photonics* **7**, 197 (2013).
- [75] D. S. Wiersma, Disordered photonics, *Nat. Photonics* **7**, 188 (2013).
- [76] B. Damski, J. Zakrzewski, L. Santos, P. Zoller, and M. Lewenstein, Atomic Bose and Anderson Glasses in Optical Lattices, *Phys. Rev. Lett.* **91**, 080403 (2003).
- [77] C. A. Condat and T. R. Kirkpatrick, Observability of acoustical and optical localization, *Phys. Rev. Lett.* **58**, 226 (1987).
- [78] D. S. Wiersma. The physics and applications of random lasers, *Nat. Physics* **4**, 359 (2008).
- [79] H. Cao, Y. G. Zhao, S. T. Ho, E. W. Seelig, Q. H. Wang, and R. P. H. Chang, Random Laser Action in Semiconductor Powder, *Phys. Rev. Lett.* **82**, 2278 (1999).
- [80] C. Vanneste and P. Sebbah, Selective Excitation of Localized Modes in Active Random Media, *Phys. Rev. Lett.* **87**, 183903 (2001).
- [81] K. G. Makris, A. Brandstötter, P. Ambichl, Z. H. Muslimani, and S. Rotter, Wave propagation through disordered media without backscattering and intensity variations, *Light: Sci. Appl.* **6** (2017).
- [82] K. G. Makris, I. Krešić, A. Brandstötter, and S. Rotter, Scattering-free channels of invisibility across non-Hermitian media, *Optica* **7**, 619 (2020).
- [83] A. F. Tzortzakakis, K. G. Makris, and E. N. Economou, Non-Hermitian disorder in two-dimensional optical lattices, *Phys. Rev. B* **101**, 014202 (2020).
- [84] Y. Huang and B. I. Shklovskii, Anderson transition in three-dimensional systems with non-Hermitian disorder, *Phys. Rev. B* **101**, 014204 (2020).
- [85] A. F. Tzortzakakis, K. G. Makris, S. Rotter, and E. N. Economou, Shape-preserving beam transmission through non-Hermitian disordered lattices, *Phys. Rev. A* **102**, 033504 (2020).
- [86] Y. Liu, Y. Wang, Z. Zheng, and S. Chen, Exact non-Hermitian mobility edges in one-dimensional quasicrystal lattice with exponentially decaying hopping and its dual lattice, *Phys. Rev. B* **103**, 134208 (2021).
- [87] Y. Liu, Q. Zhou, and S. Chen, Localization transition, spectrum structure, and winding numbers for one-dimensional non-Hermitian quasicrystals, *Phys. Rev. B* **104**, 024201 (2021).
- [88] X. Luo, T. Ohtsuki, and R. Shindou, Transfer matrix study of the Anderson transition in non-Hermitian systems, *Phys. Rev. B* **104**, 104203 (2021).
- [89] X. Luo, T. Ohtsuki, and R. Shindou, Universality Classes of the Anderson Transitions Driven by Non-Hermitian Disorder, *Phys. Rev. Lett.* **126**, 090402 (2021).
- [90] X. Luo, Z. Xiao, K. Kawabata, T. Ohtsuki, and R. Shindou, Unifying the Anderson transitions in Hermitian and non-Hermitian systems, *Phys. Rev. Res.* **4**, L022035 (2022).
- [91] Q. Tang and Y. He, Mobility edges in non-Hermitian models with slowly varying quasiperiodic disorder, *Phys. Rev. B* **109**, 224204 (2024).
- [92] Y. -M. Sun, X. -Yu Wang, and L. -Jun Zhai, Hybrid scaling properties of the localization transition in a non-Hermitian disordered Aubry-André model, *Phys. Rev. B* **110**, 054202 (2024).
- [93] J. -R. Li, S. -F. Zhang, L. -L. Zhang, and W. -J. Gong, Localization in the one-dimensional quantum chain with nonreciprocal disorder, *Phys. Rev. B* **110**, 085409 (2024).
- [94] L. Wang, J. Liu, Z. Wang, and S. Chen, Exact complex mobility edges and flagellate-like spectra for non-Hermitian quasicrystals with exponential hoppings, *Phys. Rev. B* **110**, 144205 (2024).
- [95] P. O. Sukhachov and A. V. Balatsky, Non-Hermitian impurities in Dirac systems, *Phys. Rev. Res.* **2**, 013325 (2020).
- [96] Y. Liu, X. P. Jiang, J. Cao, and S. Chen, Non-Hermitian mobility edges in one-dimensional quasicrystals with parity-time symmetry, *Phys. Rev. B* **101**, 174205 (2020).
- [97] K. Kawabata and S. Ryu, Nonunitary Scaling Theory of Non-Hermitian Localization, *Phys. Rev. Lett.* **126**, 166801 (2021).
- [98] H. Liu, Z. Su, ZQ. Zhang, and H. Jiang, Topological Anderson insulator in two-dimensional non-Hermitian systems, *Chin. Phys. B* **29**, 050502 (2020).
- [99] A. F. Tzortzakakis, K. G. Makris, A. Szameit, and E. N. Economou, Transport and spectral features in non-Hermitian open systems, *Phys. Rev. Res.* **3**, 013208 (2021).
- [100] S. Weidemann, M. Kremer, S. Longhi, and A. Szameit, Coexistence of dynamical delocalization and spectral localization through stochastic dissipation, *Nat. Photonics* **15**, 576 (2021).
- [101] A. Leventis, K. G. Makris, and E. N. Economou, Non-Hermitian jumps in disordered lattices, *Phys. Rev. B* **106**, 064205 (2022).
- [102] S. Longhi, Anderson Localization in Dissipative Lattices, *Ann. Phys. (Berlin)* **535**, 2200658 (2023).
- [103] A. Ghatak, D. H. Kaltsas, M. Kulkarni, and K. G. Makris, Diffraction and pseudospectra in non-Hermitian quasiperiodic lattices, arXiv:2410.09185 [cond-mat.stat-mech] (2024).
- [104] A. P. Acharya, A. Chakrabarty, and D. K. Sahu, S. Datta, Localization, \mathcal{PT} symmetry breaking, and topological transitions in non-Hermitian quasicrystals, *Phys. Rev. B* **105**, 014202 (2022).
- [105] S. Longhi, Lifshitz tail states in non-Hermitian disordered photonic lattices, arXiv:2412.09106 [physics.optics] (2024).
- [106] C. E. Rivet, A. Brandstötter, K. G. Makris, H. Lissek, S. Rotter, and R. Fleury, Constant-pressure sound waves in non-Hermitian disordered media, *Nat. Physics* **14**, 942 (2018).

- [107] A. Steinfurth, I. Krešić, S. Weidemann, M. Kremer, K. G. Makris, M. Heinrich, S. Rotter, and A. Szameit, Observation of photonic constant-intensity waves and induced transparency in tailored non-Hermitian lattices. *Sci. Advances* **8**, 21 (2022).
- [108] V. M. Kenkre and P. Reineker, *Exciton Dynamics in Molecular Crystals and Aggregates*, Springer(Berlin), (1982).
- [109] A. Madhukar and W. Post, Exact Solution for the Diffusion of a Particle in a Medium with Site Diagonal and Off-Diagonal Dynamic Disorder, *Phys. Rev. Lett.* **39**, 1424 (1977).
- [110] J. B. Witkoskie, S. Yang, and J. Cao, Brownian motion in dynamically disordered media, *Phys. Rev. E* **66**, 051111 (2002).
- [111] A. Amir, Y. Lahini, and H. B. Perets, Classical diffusion of a quantum particle in a noisy environment, *Phys. Rev. E* **79**, 050105(R) (2009).
- [112] J. M. Moix, M. Khasin, and J. Cao, Coherent quantum transport in disordered systems: I. The influence of dephasing on the transport properties and absorption spectra on one-dimensional systems, *New J. Phys.* **15**, 085010 (2013).
- [113] D. E. Logan and P. G. Wolynes, Dephasing and Anderson localization in topologically disordered systems, *Phys. Rev. B* **36**, 4135 (1987).
- [114] D. A. Evensky, R. T. Scalettar, and P. G. Wolynes, Localization and dephasing effects in a time-dependent Anderson Hamiltonian, *J. Phys. Chem.* **94**(3), 1149 (1990).
- [115] A. Schreiber, K. N. Cassemiro, V. Potoček, A. Gábris, I. Jex, and C. Silberhorn, Decoherence and Disorder in Quantum Walks: From Ballistic Spread to Localization, *Phys. Rev. Lett.* **106**, 180403 (2011).
- [116] S. Gopalakrishnan, K. R. Islam, and M. Knap, Noise-Induced Subdiffusion in Strongly Localized Quantum Systems, *Phys. Rev. Lett.* **119**, 046601 (2017).
- [117] Y. Rath and F. Mintert, Prominent interference peaks in the dephasing Anderson model, *Phys. Rev. Research.* **2**, 023161 (2020).
- [118] J. C. Flores, Diffusion in disordered systems under iterative measurement, *Phys. Rev. B* **60**, 30 (1999).
- [119] S. A. Gurvitz, Delocalization in the Anderson Model due to a Local Measurement, *Phys. Rev. Lett.* **85**, 812 (2000).
- [120] J. Košík, V. Bužek, and M. Hillery, Quantum walks with random phase shifts, *Phys. Rev. A* **74**, 022310 (2006).
- [121] N. Kamiya, Quantum-to-classical reduction of quantum master equations, *Prog. Theor. Exp. Phys.* **2015**, 043A02 (2015).
- [122] M. A. Broome, A. Fedrizzi, B. P. Lanyon, I. Kassal, A. Aspuru-Guzik, and A. G. White, Discrete single-photon quantum walks with tunable decoherence, *Phys. Rev. Lett.* **104**, 153602 (2010).
- [123] M. Znidaric and M. Horvat, Transport in a disordered tight-binding chain with dephasing, *Eur. Phys. J. B* **86**, 67 (2013).
- [124] H. Yamada and K. S. Ikeda, Dynamical delocalization in one-dimensional disordered systems with oscillatory perturbation, *Phys. Rev. E* **59**, 5214 (1999).
- [125] J. Cao, R. J. Silbey, Optimization of exciton trapping in energy transfer processes, *J. Phys. Chem. A* **113**(50), 13825 (2009).
- [126] S. Hoyer, M. Sarovar, and K. B. Whaley, Limits of quantum speedup in photosynthetic light harvesting, *New J. Phys.* **12**, 065041 (2010).
- [127] A. M. Jayannavar and N. Kumar, Mooij correlation in disordered metals, *Phys. Rev. B* **37**, 573 (1988).
- [128] S. Longhi, Robust Anderson transition in non-Hermitian photonic quasicrystals, *Opt. Lett.* **49**, 1373 (2024).
- [129] S. Longhi, Incoherent non-Hermitian skin effect in photonic quantum walks, *Light: Sci. Appl.* **13**, 95 (2024).
- [130] D. S. Bhakuni, T. L. M. Lezama, Y. B. Lev, Noise-induced transport in the Aubry-André-Harper model, *SciPost Phys. Core* **7**, 023 (2024).
- [131] S. Longhi, Dephasing-Induced Mobility Edges in Quasicrystals, *Phys. Rev. Lett.* **132**, 236301 (2024).
- [132] G.F. Lawler, *Introduction to Stochastic Processes, Second Edition*, Taylor & Francis (2006).