Orbital maneuvers for a space probe around Titania

Jadilene Xavier^{1*}, Antônio Bertachini A. Prado^{2†}, Silvia Giuliatti Winter^{1†} and Andre Amarante^{1†}

^{1*}Grupo de Dinâmica Orbital & Planetologia, São Paulo State University (Unesp), School of Engineering and Sciences, Guaratinguetá, CEP 12516-410, São Paulo, Brazil.
²National Institute for Space Research, Av. dos Astronautas, 1758, São José dos Campos, SP, Brazil. Professor, Academy of Engineering, RUDN University, Miklukho-Maklaya street 6, Moscow, Russia, 117198.

*Corresponding author(s). E-mail(s): jadilene.rodrigues@unesp.br; Contributing authors: antonio.prado@inpe.br; giuliatti.winter@unesp.br; andre.amarante@unesp.br; †These authors contributed equally to this work.

Abstract

For most space missions, it is interesting that the probe remains for a considerable time around the mission target. The longer the lifetime of a mission, the greater the chances of collecting information about the orbited body. In this work, we present orbital maneuvers that aim to show how to avoid a collision of a space probe with the surface of Titania. Through an expansion of the gravitational potential to the second order, the asymmetry of the gravitational field due to the coefficient C_{22} of Titania, the zonal coefficient J_2 , and the gravitational perturbation of Uranus are considered. Two models of coplanar bi-impulse maneuvers are presented. The first maneuver consists of transferring an initial elliptical orbit to a final circular orbit, and the second has the objective of transferring an initial elliptical orbit to a final orbit that is also elliptical. The lag in the inclination and semi-major axis of the orbits is investigated before performing the maneuvers. To point out the best scenarios for carrying out the maneuvers, a study is presented for different points of an orbit where transfers could be made. In addition, a maneuver strategy is presented to correct the variation of the periapsis argument. The results show that maneuvers performed a few days after integration are more economical than maneuvers performed later, a few days before the collision. The economy of the maneuvers is also demonstrated through an analysis of the ratio of the increase in speed to the lifetime.

Keywords: orbits, maneuvers, lifetime, astrodynamics, numerical simulations, planetary satellite

1 Introduction

In recent years there has been an increasing interest in sending space missions to the gaseous planets of the solar system. According to [1], NASA and ESA are planning possible missions to the ice giants from 2024 to 2037. These possible missions

can help to understand the origin and evolution of these systems. In addition, some missions are directed towards some specific natural satellites, such as the JUpiter ICy moons Explorer (JUICE) mission, launched in 2023, which will investigate Jupiter and its moons, Ganymede, Europa, and Callisto [2].

Several works look for better orbits around natural satellites. The purpose of these works is to provide essential data to help future space missions that target these satellites. However, sending a probe into space is a process that demands a lot of time and investment. It is desirable that, during a mission, the probe remains in orbit around the observed body for a long period of time. Thus, searching for long-duration orbits and avoiding collisions with the mission target is very important to achieve success during the observation of the desired object.

The work by [3] presents a study on orbits around Titania, one of the moons of Uranus, an ice-giant planet. The authors investigate orbits for a space probe around Titania with various eccentricities values. Perturbations due to the third body and the gravitational coefficients of Titania are considered, and through lifetime maps, orbits with long lifetimes are found. The results of numerical simulations showed that orbits with the longest lifetimes were those whose eccentricity is equal to 10^{-3} . The authors also show that values other than zero for the argument of periapsis and the longitude of the ascending node can prolong the probe's lifetime. Furthermore, a study on lifetime sensitivity caused by possible errors in Titania's gravitational coefficients concluded that orbits with eccentricity equal to 10^{-3} are the most affected by these potential errors.

Long-duration orbits are also studied in [4]. In this case, the natural satellite is Io, one of Jupiter's Galilean satellites. The authors were able to detect collisions and possible escapes of the probe. Orbits with significant lifetimes ranged from 6 months to 2.3 years. In addition, the study showed that orbits around Io are more susceptible to perturbations due to their short distance from the perturber. Finally, they emphasized the importance of assigning non-zero values for the argument of periapsis and the longitude of the ascending node.

The work developed by [5] presents a study of the lifetime of a probe in a low-altitude and highly inclined orbit around the natural satellite Europa. The results showed that the obliquity of the third body and the nodal phase affect the lifetime of the probe.

Studying long-duration orbits around natural satellites also requires finding how to avoid a probe collision with the proposed mission target. In this sense, in [6], corrective orbital maneuvers are proposed for a spacecraft around Europa. The results showed that the lifetime can be extended through appropriate strategies of corrective maneuvers. Orbital maneuvers are also investigated in [7]. After analyzing long-term orbits around Saturn's natural satellite, Titan, the authors present the best results for such maneuvers to be carried out.

Given the above mentioned, in this paper we present models for orbital maneuvers to avoid collision of a probe with the surface of Titania. The first maneuver proposes to transfer the spacecraft from an initial elliptical orbit to a final circular orbit. In the second model, the transfer is from an initial elliptical orbit to a final elliptical orbit, both of which are coplanar. For this second model, we performed maneuvers at various points of the orbit to analyze if it is more feasible for a maneuver to take place close to the collision or within a few days of the mission's lifetime. We also present a maneuver model for correcting small variations of the argument of periapsis. These maneuvers were performed considering points where the deviation of the semi-major axis and the eccentricity are small.

2 Mathematical Model

The system used here includes Titania as the central body, a space probe in orbit around Titania, and Uranus as the third body. Uranus is assumed to be in a Keplerian orbit around Titania with a radius of 25.362×10^3 km and a mass 8.68×10^{25} kg (https://ssd.jpl.nasa.gov/). The orbital elements used in the numerical simulations are presented in Table 1. We also consider the two main gravitational terms of Titania, J_2 and C_{22} , with values equal to 1.13×10^{-4} and 3.38×10^{-5} , respectively [8]. They are the most critical terms in the gravitational field after the Keplerian term. These terms cause significant variations in some orbital elements, such as the eccentricity and inclination of the orbit. In this way, they become the most effective terms to model the irregular shape of a body [3, 9, 10]. In the case of Titania, it is worth noting that these terms are the only ones available in the literature. In addition to these coefficients, the mass of Titania $(35.27 \times 10^{20} \text{ kg})$ and its radius (788.9 km) are considered in this work. The equation of motion of the probe can be written in the following form [11]:

$$\ddot{\vec{r}} = -\frac{G(M_T + m)\vec{r}}{\vec{r}^3} + GM_U \left(\frac{\vec{r}_U - \vec{r}}{|\vec{r}_U - \vec{r}|^3} - \frac{\vec{r}_U}{\vec{r}_U^3} \right) + \vec{P}_T$$
(1)

$$P_{Tx} = -\frac{G(M_T + m)J_2x}{2r^5} \left[3 - 15\left(\frac{z}{r}\right)^2 \right] + \frac{3G(M_T + m)C_{22}x}{r^5} \left[2 - \frac{5(x^2 - y^2)}{r^2} \right]$$
(2)

$$P_{Ty} = -\frac{G(M_T + m)J_2y}{2r^5} \left[3 - 15\left(\frac{z}{r}\right)^2 \right] - \frac{3G(M_T + m)C_{22}y}{r^5} \left[2 + \frac{5(x^2 - y^2)}{r^2} \right]$$
(3)

$$P_{Tz} = -\frac{G(M_T + m)J_2z}{2r^5} \left[9 - 15\left(\frac{z}{r}\right)^2 \right] - \frac{15G(M_T + m)C_{22}z}{r^7} (x^2 - y^2)$$
(4)

where \vec{P}_T is the expansion of the gravitational potential up to the second order for Titania; P_{Tx} , P_{Ty} and P_{Tz} are components of the P_T vector. m is the mass of the space probe, M_T the mass of Titania and M_U the mass of Uranus. The terms \vec{r} and \vec{r}_U are the radius vector of the probe and Uranus, respectively.

Table 1 Parameters of Titania with respect to Uranus.

Parameter	Value		
Semi-major axis (km)	435.8×10^{3}		
Eccentricity	1.18×10^{-3}		
Inclination (\circ)	1.0×10^{-1}		
Argument of periapsis (°)	1.64×10^{2}		
Longitude of ascending node (°)	1.67×10^2		
Mean anomaly (\circ)	2.05×10^{2}		
JPL.Website: https://ssd.jpl.nas	a.gov/.		

First, the orbits were **numerically** integrated considering the system described by equations 1-4. During these **numerical** integrations, the maneuvers were performed. To integrate the orbits and to carry out the maneuvers, modifications were made to the Mercury package [12].

2.1 Orbital Maneuvers

During the decay process of an orbit, its orbital elements change. Therefore, when trying to perform a maneuver, we must consider some of these variables, such as the semi-major axis (a). eccentricity (e), and inclination (I). For example, suppose a maneuver is performed close to the probe's collision with the mission target's surface. In that case, the orbit will have a high eccentricity and, possibly, a considerable variation in its semimajor axis. Thus, we present in this section two models of maneuvers. First, consider the transfer from an initial elliptical orbit to a final circular orbit and later, transferring the spacecraft from an initial elliptical orbit to a final elliptical orbit, thus correcting the original altitude of the orbit without considering the inclination. Both maneuvers are coplanar. The final orbit is the desired orbit for the mission, while the initial orbit is the orbit after the perturbations acted and changed the initial orbit.

2.1.1 Maneuvering from an elliptical to a circular orbit.

The first maneuver presented here involves transferring the space probe from an elliptical to a final circular orbit. The first impulse (ΔV_1) is applied at the apoapsis of the elliptical orbit, making the probe to enter a transfer ellipse. The second impulse (ΔV_2) is applied when the probe is the apoapsis of the transfer orbit to make the probe occupy the desired final circular orbit. The total increment in velocity is given by the sum of the modules (ΔV_1) and (ΔV_2) . The time for carrying out the orbital transfer is half the period of the transfer orbit. The equations for performing this maneuver are described as follows:

$$V_{ap} = \sqrt{\frac{2\mu}{a(1+e)} - \frac{\mu}{a}}$$
 (5)

$$V_{p1} = \sqrt{\frac{2\mu}{a(1+e)} - \frac{2\mu}{a(1+e) + r_{cir}}}$$
 (6)

$$\Delta V_1 = V_{p1} - V_{ap} \tag{7}$$

$$V_2 = \sqrt{\frac{2\mu}{r_{cir}} - \frac{2\mu}{a(1+e) + r_{cir}}}$$
 (8)

Springer Nature 2021 LATEX template

$$V_{cir} = \sqrt{\frac{\mu}{r_{cir}}} \tag{9}$$

$$\Delta V_2 = V_{cir} - V_2 \tag{10}$$

$$\Delta V = |\Delta V_1| + |\Delta V_2| \tag{11}$$

where V_{ap} is the velocity at the apoapsis of the initial orbit, V_{p1} the velocity at point 1 in the transfer orbit, ΔV_1 the first impulse, V_2 is the velocity at the apoapsis (point 2) of the transfer orbit, V_{cir} the velocity in the final circular orbit, ΔV_2 the second impulse and ΔV the total impulse.

2.1.2 Maneuvering between elliptical orbits

Another maneuver presented in this work will be from an initial elliptical orbit to another final orbit, also elliptical. These orbits are coplanar. It happens when the nominal orbit of the spacecraft is elliptic. For this maneuver, the first impulse (ΔV_1) is applied at the periapsis of the initial elliptical orbit. Then, the probe will occupy a transfer orbit. The second impulse (ΔV_2) is applied at the apoapsis of the transfer orbit, and the spacecraft will be placed into its final elliptical orbit. The sum of (ΔV_1) and (ΔV_2) results in the total increment of velocity for performing the maneuver. The equations for carrying out such a maneuver are given by:

$$V_{p1} = \sqrt{\frac{2\mu}{r_{p1}} - \frac{2\mu}{r_{p1} + r_{a1}}} \tag{12}$$

$$V_{pt} = \sqrt{\frac{2\mu}{r_{p1}} - \frac{2\mu}{r_{p1} + r_{ta}}} \tag{13}$$

$$\Delta V_1 = V_{pt} - V_{p1} \tag{14}$$

$$V_{at} = \sqrt{\frac{2\mu}{r_{ta}} - \frac{2\mu}{r_{p1} + r_{ta}}} \tag{15}$$

$$V_{a2} = \sqrt{\frac{2\mu}{r_{ta}} - \frac{2\mu}{r_{p2} + r_{ta}}} \tag{16}$$

$$\Delta V_2 = V_{a2} - V_{at} \tag{17}$$

$$\Delta V = |\Delta V_1| + |\Delta V_2| \tag{18}$$

where r_{p1} and r_{a1} are the initial orbit's periapsis and apoapsis. Furthermore, r_{ta} is the apoapsis of the transfer ellipse, r_{p2} is the periapsis of the final orbit, $r_{a2} = r_{ta}$ is the apoapsis of the desired final orbit. Regarding the increase in velocity, we have V_{p1} equal to the velocity at point 1 of the initial orbit, V_{pt} the velocity at point 1 of the transfer ellipse, ΔV_1 the first impulse, V_{at} the velocity at the apoapsis of the transfer ellipse, V_{a2} the velocity at point 2 concerning the final desired orbit, ΔV_2 the second impulse and ΔV the total impulse.

3 Analyzing the Results

The first results presented are about the first maneuver, from an initial elliptical orbit to a final circular orbit. This is the situation that occurs when the nominal orbit is circular and the orbit after the pertubations is elliptical. Then, the radii of the final desired orbits were chosen according to the regions with the most prolonged orbital duration, according to the analysis presented in [3]. This radius ranges from 1000-2000 km, in intervals of 200 km. The choice of the right moment to carry out the maneuver was made as follows: first, we chose a circular orbit with an initial semimajor axis, $(a_i = R_i)$, ranging from 1000-2000 km, the we analyzed the decay of this orbit and, when reaching a significant value of variation from its starting position, the maneuver is performed. We take the orbital elements of this point and consider them as elements of the initial orbit (black orbit, Figure 1). Small changes in an orbit's initial conditions can shorten its lifetime, and the orbit reaches high eccentricities and inclinations. Thus, some maneuvers are performed after a significant change in its initial position to return the orbit to its original altitude position, disregarding the inclination. Four initial inclinations were considered: 60°, 70°, 80° and 90°. Figure 1 shows a diagram of how the maneuvers will be carried out.




Fig. 1 Representation of the desired final circular orbits, where R_f is the radius of these orbits. Purple for $R_f=1000$ km, green for $R_f=1200$ km, blue for $R_f=1400$ km, red for $R_f=1600$ km, light green for $R_f=1800$ km and dark purple for $R_f=2000$ km. Initial orbit represented by the black line

To plan the maneuvers, we took into account the gravitational coefficients of Titania, including J_2 and C_{22} , as well as the gravitational coefficients of Uranus. After that, we executed the maneuvers necessary to achieve the desired orbits, utilizing equations 6-11.

The orbits with the largest deviations of 20-30% of the initial value (black orbit, Figure 1), are those with initial inclinations of 60°, 70° and 80° for all simulations. In this case, when the orbits reach the deviation, the lifetime comes close to 240-250 days, and the ΔV needed to reach the circular orbit of radius 1000 km again is around $1\times10^{-2}-6\times10^{-2}$ km/s. Considering that orbits with this inclination have a total lifetime of 252 days, it is concluded that the maximum inclination variation occurs a few days before the collision.

In the case of an orbit with an initial inclination of 90° , the maximum variation of I is 5% for all final orbits. However, the ΔV ranges from $4\times 10^{-2}-1\times 10^{-1}$ km/s and the decay times range from 1-17 days. The smallest values of ΔV are for maneuvers to reach final orbits with a radius of 1000-1200 km. These orbits also have the highest rate of inclination variations.

The variations in the semi-major axis are quite small, between 0.1-0.9 % of the initial value. For the four inclinations analyzed, 60° , 70° , 80° and 90° , there are cases of maximum variations of the semi-major axis of 0.6 - 0.9 %. In these cases, the ΔV ranges from $8 \times 10^{-2} - 1.1 \times 10^{-1}$ km/s. The

cases with the lowest value of ΔV are for final orbits with a 1000-1200 km radius.

Some missions choose to correct the space-craft's position with a short lifetime before a considerable variation in its position. For example, Figure 2 presents a study that seeks to find a minimum ΔV to perform an orbital transfer before a large variation of its initial position. Thus, now the maneuvers of interest are not the ones with a significant change in the inclination or in the semi-major axis. As in the previous cases, for the four inclinations analyzed, the minimum ΔV is calculated for cases where the final orbits have a 1000-2000 km radius. As shown in Figure




Fig. 2 Minimum ΔV as a function of the circular radius of the final orbit. Analyzing four initial inclinations 60° , 70° , 80° and 90° in orange, red, green, and black colors, respectively. Initial values: $a_i = R_i = 1000 - 2000$ km, $e_i = 0$ and $\omega = \Omega = 0^{\circ}$. The radius of intended final orbits is 1000 - 2000 km.

2, the smallest value of ΔV is approximately 5.4×10^{-5} km/s, for a final orbit with a radius of 1050 km, for all inclinations. For the first inclination of 60°, the maneuver was performed after 5 days, equivalent to approximately 45 orbital periods. At that point, the semi-major axis and eccentricity had values equal to 1.0498×10^3 km and 2.8×10^{-4} , respectively. The ratio of change of the semi-major axis is 0.016 % of its initial value and 0.45 % for the inclination.

In the case of the orbit with an inclination of 70°, this maneuver was carried out after 14 days, when a and e are 1.0492×10^3 km and 2.3×10^{-4} , respectively. The inclination variation for this ΔV is 0.4%, and for the semi-major axis 0.01 %.

For I equal to 80° and 90°, the minimum ΔV is found after 5 days. In the case of $I=80^{\circ}$, at the

Table 2 Values of ΔV for maneuvers with significant offset values in the inclination and in the semimajor axis. Initial conditions used in maneuvers such as initial inclination, orbital duration time, apoapsis radius (r_{apo}) , final circular orbit radius (r_{circ}) , semimajor axis offset (Def_a) , and the inclination (Def_I) in percentage, and the equivalent ΔV .

I(°)	Time (Orbital period) ¹	r _{Apo} (km)	r _{circ} (km)	Def_a (%)	Def_I (%)	$\Delta V (km/s)$
$I_i = 60$	1735	1010	1000	0.057	24.9	2.5332×10^{-3}
$I_i = 70$	2172	1042	1000	0.44	28.3	1.0192×10^{-2}
$I_i = 80$	2264	1107	1000	0.07	25.4	2.5823×10^{-2}
$I_i = 90$	2236	1145	1000	0.06	2.0	3.5056×10^{-2}
$\overline{I_i} = 60$	1735	1264	1200	0.067	14.7	1.1916×10^{-2}
$I_i = 70$	2136	1585	1200	0.06	20	7.0039×10^{-2}
$I_i = 80$	1872	1562	1200	0.13	13	6.6109×10^{-2}
$I_i = 90$	1800	1538	1200	0.04	1.8	6.1449×10^{-2}
$I_i = 60$	1934	1990	1400	0.084	14.5	8.5579×10^{-2}
$I_i = 70$	1581	1990	1400	0.11	13.3	8.5653×10^{-2}
$I_i = 80$	1436	1948	1400	0.11	7.6	7.9481×10^{-2}
$I_i = 90$	1400	1912	1400	0.09	1.9	7.4107×10^{-2}
$I_i = 60$	1462	2348	1600	0.072	14.6	8.8944×10^{-2}
$I_i = 70$	1227	2383	1600	0.19	11	9.3866×10^{-2}
$I_i = 80$	1118	2302	1600	0.18	6.0	8.3743×10^{-2}
$I_i = 90$	1127	2332	1600	0.3	2.2	8.7669×10^{-2}
$I_i = 60$	1144	2700	1800	0.22	14.6	9.0549×10^{-2}
$I_i = 70$	963	2653	1800	0.5	8.6	8.6478×10^{-2}
$I_i = 80$	918	2784	1800	0.3	6.8	9.9839×10^{-2}
$I_i = 90$	890	2626	1800	0.3	2.4	8.3109×10^{-2}
$I_i = 60$	953	3094	2000	0.76	17.4	9.6332×10^{-2}
$I_i = 70$	800	3125	2000	0.4	12	9.8090×10^{-2}
$I_i = 80$	727	2990	2000	0.4	6.0	8.5558×10^{-2}
$I_i = 90$	727	3006	2000	0.4	3.2	8.7159×10^{-2}

¹The orbital period is measured from the period of the orbit closest to the surface of the central body.

moment of the maneuver, the value of the semimajor axis is 1.0497×10^3 km and the eccentricity 2.0×10^{-4} . The inclination and semi-major axis variations are 0.36~% and 0.02~%, respectively. For an inclination of 90°, the maneuver is performed when $a = 1.0497 \times 10^3$ km and $e = 2.03 \times 10^{-4}$. The variation on the inclination is approximately 0.3 % and on the semi-major axis 0.023 %. Within this minimum range, it is possible to identify that; for a final orbit with a radius of 950 km, the ΔV required to perform the maneuver is greater than the other cases analyzed. For orbits with initial inclinations of 80° and 90°, these values are close to 5.7×10^{-4} km/s for $I = 80^{\circ}$ and $5.55 \times 10^{-4} \text{ km/s for } I = 90^{\circ}. \text{ For } I = 70^{\circ} \text{ this}$ orbit can be achieved with $\Delta V = 7.1 \times 10^{-4}$ km/s.

In order to analyze the best values of ΔV used to perform orbital maneuvers, Table 2 shows some values of ΔV used for orbits lagged in inclination and semimajor axis. For each inclination, $I=60^\circ$, $I=70^\circ$, $I=80^\circ$ and $I=90^\circ$, values of ΔV are presented, as well as the initial conditions for carrying out the maneuvers. Cases are presented for final orbits with radii equal to 1000, 1200, 1400, 1600, 1800, and 2000 km.

The next discussions are about the results obtained for maneuvers between two elliptical orbits. In [3], the orbital decay of objects with eccentricity equal to 10^{-4} , 10^{-3} , 10^{-2} and 10^{-1} are studied. The authors concluded that the lifetime was longer for orbits with an eccentricity 10^{-3} . Thus, the intended final orbits investigated

in this work have the same eccentricity as the orbits studied in that work (10^{-4} , 10^{-3} , 10^{-2} and 10^{-1}). We call these eccentricities e_f . Figure 3 shows these orbits.




Fig. 3 Representation of final elliptical orbits. Orange for eccentricity 10^{-1} , green for 10^{-2} , pink for 10^{-3} and blue for 10^{-4} .

A set of numerical simulations was made to perform the maneuvers for each eccentricity.

Six points in the orbit were chosen to carry out the maneuvers for each value of e. This choice aims to analyze whether it is more feasible to make one maneuver at different times, resulting in several maneuvers before the collision, or only one maneuver moments before the spacecraft collides with the surface of the moon. Such maneuvers are called m_P , m_1 , m_2 , m_3 , m_4 and $\mathbf{m_5}$. Being m_P , the main maneuver is because this maneuver is performed at the closest point to the surface of Titania, that is, closest to the collision. The approximate periapsis for this maneuver is 790 km for all values of e. The point for maneuver m_5 is farthest from the surface of Titania, with the maneuver performed with a few days of decay compared to m_P . The periapsis for m_5 is approximately 965 km. The m_1 , m_2 , m_3 , and m_4 maneuvers are carried out at intermediate points between m_P and m_5 . The periapsis of m_1 is around 825 km, $m_2 \approx 860$ km, $m_3 \approx 895$ km, $m_4 \approx 930 \text{ km}$ and $m_5 \approx 965 \text{ km}$. Points vary every 35 km.

All simulations were made considering a=1000 km, $I=90^{\circ}$, $\omega=\Omega=0^{\circ}$, the gravitational coefficients J_2 and C_{22} of Titania and the gravitational effects of Uranus.

The first set of maneuvers was done for $e = 10^{-4}$, giving a total lifetime of 268.536 days. All points of the orbit considered for the maneuvers are represented in Figure 4 (at the top). To




Fig. 4 Projection of orbital positions for the six maneuvers performed $(m_P, m_1, m_2, m_3, m_4 \text{ and } m_5)$ for $e=10^{-4}$ (at the top). Variation of orbital elements e, I, and periapsis radius as a function of lifetime and maneuvering time (at the bottom). Initial conditions: a=1000 km, $e=10^{-4}$, $I=90^{\circ}$, $\omega=\Omega=0^{\circ}$.

relate the variations of some parameters when performing the maneuvers, Figure 4 (at the bottom) shows the position of the periapsis (r_p) , e and I as a function of time. The day each maneuver was performed can also be visualized in this figure.

For the main maneuver m_P , the parameters are: $a = 9.993 \times 10^2$ km, $e = 2.094 \times 10^{-1}$, $I = 87.9^{\circ}$, periapsis equal to 790.0 km, apoapsis equal to 1208.59 km and maneuver time of 267.26 days. Considering these values, we decay approximately 21% in the periapsis, 2.3% in the inclination, and 0.1% in the semi-major axis. There is also an increase in the eccentricity of three orders. The m_P maneuver was performed about a day before the collision. The estimated ΔV for this transfer was $\approx 5.26 \times 10^{-2}$ km/s. The representation of the location of this maneuver can be seen in Figure 4, in a dotted red line.

The second maneuver (m_1) , for $e = 10^{-4}$, was performed within 259.87 days of numerical integration, approximately 8 days before the collision. At this chosen point, the periapsis of the orbit was 825.09 km, approximately 17.5\% of its initial value. The other initial conditions were: $a = 9.989 \times 10^2 \text{ km}, e = 1.740 \times 10^{-1}, I = 88.3^{\circ},$ and apoapsis equal to 1172.88 km. To perform this maneuver, the ΔV used was approximately 4.34×10^{-2} km/s. Analyzing the variation of the orbital elements for this maneuver, it is possible to observe that, during 259.87 days, the periapsis decreased by almost 180 km, the inclination by 1.4° and the semi-major axis by 2 km. The maneuver m_1 is represented by the dotted blue line in Figure 4.

The maneuver (m_2) was performed after 248.59 days, about 19 days before the collision. In this maneuver, the ΔV required to reach the orbit with an eccentricity of 10^{-4} was $3.45 \times$ 10^{-2} km/s. The following initial conditions were used to execute the maneuver: $a = 9.991 \times 10^2$ km, $e = 1.392 \times 10^{-1}, I = 88.5^{\circ}, \text{ periapsis equal to}$ 860.01 km, and apoapsis equal to 1138.35 km. The periapsis had the highest percentage of change of all the variations of the orbital elements during the decay of the orbit. It decayed almost 14%. The semi-major axis and inclination decreased by 0.1% and 1.66%, respectively. In the case of eccentricity, there was an increase of three orders of magnitude about its initial value. The line orange represents this maneuver in Figure 4.

In figure 4 the color representing the maneuver (m_3) for $e=10^{-4}$ is green. The lifetime for this maneuver was 234.86 days, almost 33 days before the collision. In this maneuver, the periapsis decay was approximately 10%, from an initial value of 999.9 km to 895 km. The eccentricity increased by two orders of magnitude, to $e=9.970\times10^{-2}$, and the apoapsis increased by almost 103 km. The semi-major axis and inclination had small variations, 0.1% and 1.22%, respectively. The maneuver m_3 has a smaller offset of the orbital elements concerning the previous maneuvers. However, the point where the maneuver is carried out may be of interest to a mission, and the ΔV necessary is 2.56×10^{-2} km/s.

In the case of the maneuver (m_4) , represented in Figure 4 by the light blue color, the time to perform the maneuver was 218.20 days. The maneuver preceded 50.34 days of the collision, and its periapsis was 930.16 km, almost a drop of 7% compared to its initial value. Regarding the inclination, the variation was small, around 1.33%. The eccentricity with a value of 6.95×10^{-2} and a ΔV of 1.70×10^{-2} km/s was used to reach the same origin position.

In pink, the last maneuver for $e=10^{-4}$ shown in Figure 4 was performed when the orbit reached 187.84 days. As already mentioned, this maneuver is performed within fewer days since the start of the mission than the others and, therefore, has a smaller periapsis decay. The initial conditions for this point are $a=9.991\times10^2$ km, $e=3.420\times10^{-2}$, $I=89^\circ$, periapsis equal to 965.01 km and apoapsis equal to 1033.37 km. The periapsis offset was $\approx 3.5\%$ and the inclination $\approx 1.11\%$, with a velocity increment $\Delta V=8.31\times10^{-3}$ km/s.

Comparing all maneuvers performed for $e=10^{-4}$, it is possible to observe that the best maneuver is (m_5) . The ΔV of 8.31×10^{-3} km/s is much smaller for the number of days of life. Compared to the other maneuvers, (m_P) for example, if two maneuvers (m_5) are performed, the orbit can have a lifetime of up to 374 days with $2\times\Delta V$ equal to 1.662×10^{-2} km/s. While two maneuvers (m_P) , we will have a lifetime of ≈ 534 days but a total ΔV of 1.052×10^{-1} km/s, almost 6 times higher than the total ΔV used in (m_5) . If we do six maneuvers at the point of the maneuver (m_5) , we would have a longer lifetime and lower fuel consumption. This comparison holds for the other maneuvers as well.

The following discussions are for calculating ΔV for $e=10^{-3}$. Since the results are closer to the case presented for $e=10^{-4}$, we will omit the figure with the projections of the maneuvering points. As in the previous case, six points in the orbit were chosen to carry out the maneuver. In this case, the orbital duration time was 313 days.

The m_1 maneuver took place with a duration of 304.44 days, 8.56 days before the collision. For this maneuver, a periapsis of 825.18 km was considered, 17.4% smaller than its initial value, and an apoapsis of 1157 km, almost 15% larger than the value before the decay. The other initial conditions were: $a = 9.999 \times 10^2$ km, $e = 1.747 \times 10^{-1}$, $I = 87.9^{\circ}$. The value of ΔV was approximately 4.33×10^{-2} km/s. Comparing the value of ΔV found for m_1 in the previous case, it is noticed that the values are very close, and the differences are irrelevant.

For maneuver m_2 , we used the following initial conditions: maneuver time of 295.15 days, almost 18 days before the collision. $a = 9.996 \times 10^2$ km, $e = 1.396 \times 10^{-1}$, $I = 88.0^{\circ}$, periapsis equal to 860.10 km and apoapsis equal to 1139.22 km. Regarding the variations of the parameters, we had a decay of almost 14% in the periapsis and 2.2% for the inclination. With these initial conditions, we found a ΔV of $\approx 3.43 \times 10^{-2}$ km/s. As in the previous maneuvers, the value of ΔV for m_2 is very close to that of ΔV found for the same maneuver analyzed in the case $e = 10^{-4}$.

In m_3 , the ΔV found was 2.54×10^{-2} km/s, in a time of 282.04 days, which preceded 30.96 days of the collision. At this point, the orbit has a semi-major axis of 9.990×10^2 km, eccentricity 1.041×10^{-1} , and inclination equal to 88.2°, with a periapsis variation of 10.4% and apoapsis with an increase of almost 10.2%. The maneuver m_3 also has a similar value to the one analyzed for $e = 10^{-4}$. For m_4 , the value of ΔV found was $\approx 1.68 \times 10^{-2}$ km/s for a lifetime of 264.93 days. In these maneuvers, the periapsis had a decay of almost 7% and an inclination of $\approx 1.5\%$. Periapsis is equal to 930.03 km, apoapsis is equal to 1069.74 km, and eccentricity is equal to $6.986 \times$ 10^{-2} . For the same maneuver analyzed in 10^{-4} , we can say that the values of ΔV are closer.

The initial conditions for the orbital transfer for m_5 were considered: $a=9.989\times 10^2$ km, $e=3.388\times 10^{-2}$, $I=88.9^\circ$, periapsis at 965.05 km and apoapsis at 1032.76 km. The results showed that $\Delta V=8.01\times 10^{-3}$ km/s, a decay of $\approx 3.4\%$ from the periapsis and $\approx 1.22\%$ for inclination and a lifetime of 234.19 days, almost 79 days before the collision.

Carrying out a similar analysis to the one made for $e=10^{-4}$, this time between m_5 and m_1 , we note that the maneuver m_5 is the most viable of all. While m_5 spent 8.01×10^{-3} km/s to perform a maneuver after 234 days of life, m_1 spent 4.33×10^{-2} km/s for 304.44 days of life. If we do two maneuvers m_5 , we will have the probe in orbit for 468.38 days spending a total ΔV of 1.602×10^{-2} km/s. In the case of two maneuvers m_1 , the total ΔV would be 8.66×10^{-2} km/s for a little more than 608 days of life, almost 5.4 times greater than the ΔV used for two maneuvers m_5 .

Graphically, the results shown for $e = 10^{-2}$ are similar to those for $e = 10^{-1}$, which we will show below, and, for this reason, we omitted it

here. For $e=10^{-2}$, the maneuver m_P was performed with a ΔV of 4.99×10^{-2} km/s. Compared to the values of ΔV for m_P in the cases for e equal to 10^{-4} and 10^{-3} , this value is slightly lower, around 5%. In this maneuver, the periapsis offset was 20.2%, and only 0.6% for the inclination. The other initial conditions used were: $a=9.988\times 10^2$ km, $e=2.089\times 10^{-1}$, $I=89.4^\circ$, apoapsis equal to 1207.00 km. The maneuver occurred at 152.68 days, 0.5 days before the collision with the central body.

In the case of m_1 , the maneuver time preceded almost seven days of the collision (145.90 days), with a decay of almost 17% at the periapsis (825.01 km) and 1.2% in the inclination. The ΔV needed to perform the transfer was $\approx 4.10 \times 10^{-2}$ km/s. The semi-major axis was 9.995×10^2 km, eccentricity equal to 1.745×10^{-1} and apoapsis equal to 1174.00 km. The maneuver m_1 also has a slightly lower value than the maneuvers analyzed in the previous cases, around 5%, for the same periapsis.

The turnaround time for m_2 was 135.00 days, 18 days before the end of the numerical integration. To perform this maneuver, a ΔV of $\approx 2.92 \times 10^{-2}$ km/s was found. Comparing with the ΔV for the cases $e=10^{-4}$ and $e=10^{-3}$, ΔV is now almost 15% smaller. The initial elliptical orbit has a $a=9.991\times 10^2$ km, $e=1.389\times 10^{-1}$, $I=89.6^\circ$, periapsis equal to 860.26 km ($\approx 13\%$ less than the initial value) and apoapsis equal to 1137.90 km ($\approx 13\%$ greater than the initial value).

For the maneuver m_3 performed approximately 30 days before the collision, the orbit had the following initial conditions: $a=9.989\times 10^2$ km, $e=1.039\times 10^{-1}$, $I=89.5^\circ$, periapsis equal to 895.07 km, with a decay of almost 9% and an apoapsis of 1102.80 km, almost 93 km greater than its initial value. The total impulse in velocity was $\approx 2.31\times 10^{-2}$ km/s. This value is almost 10% less than the values of ΔV found in cases $e=10^{-4}$ and $e=10^{-3}$.

The maneuver represented in light blue, m_4 , happened in 104.85 days. To find the increase in velocity ΔV at this point, a periapsis with a variation of 6% was considered. The inclination had a small variation, about 0.3%, until the moment of the maneuver. With these initial conditions, semi-major axis equal to 9.993 \times 10² km, $e=6.931\times10^{-2}$, $I=89.7^{\circ}$, periapsis equal to 930.07 km and apoapsis equal to 1068.6 km, a ΔV

of 1.45×10^{-2} km/s was found. This value is also considerably smaller than the ones found for m_4 in the previous cases.

The last maneuver, m_5 , for $e = 10^{-2}$, was performed after 74.11 days of life. The initial conditions were: $a = 9.999 \times 10^2$ km, $e = 3.482 \times 10^{-2}$, $I = 89.8^{\circ}$, periapsis equal to 965.14 km with decay of 2.5% and apoapsis equal to 1034.7 km. In this case, the ΔV was 6.04×10^{-3} km/s.

In all maneuvers performed for a final orbit with eccentricity 10^{-2} , the ΔV presented lower values than those found for the same maneuvers with $e=10^{-4}$ and $e=10^{-3}$. In terms of percentage, we can say that $m_P\approx 5\%$, $m_1\approx 5\%$, $m_2\approx 15\%$, $m_3\approx 9\%$, m_4 between 15% and 17% and m_5 between 25% and 27% smaller for $e=10^{-2}$ compared to the previous cases.

Maneuver m_5 appears, again, as the best maneuver found. Here, we will compare this maneuver with m_2 . Doing two maneuvers m_5 will give an orbit with a lifetime of approximately 148 days, with a total ΔV of 1.208×10^{-2} km/s. However, doing two maneuvers m_2 , the orbit will have a lifetime of 270 days using a total ΔV of 6.38×10^{-2} km/s. Although the lifetime is longer for m_2 , the ΔV for m_5 is almost 5.2 times smaller.

Finally, we present in Figure 5 the results for the maneuvers performed for $e=10^{-1}$. For this case, we consider only four maneuvers, m_P , m_1 , m_2 , and m_3 . This choice is justified because the initial periapsis of the orbits starts at 990 km, so the points for maneuvers m_4 (periapsis equal to 930 km) and m_5 (periapsis equal to 965 km) are not possible. The colors and points are the same considered in the previous cases for m_P to m_3 . The total integration time, in this case, was 52 days.

The m_P maneuver was performed considering a periapsis of 790 km and an apoapsis =1209.5 km. We adopted semi-major equal to 9.998×10^2 km, $e=2.098 \times 10^{-1}$ and $I=89.5^{\circ}$. The total impulse of 2.74×10^{-2} km/s was given almost a day before the collision. At the time of the maneuver, the periapsis offset was approximately 12.2% and 0.55% for the inclination. Compared to the values of ΔV analysed for m_P in the previous cases, the value found is 45% smaller than the value found for $e=10^{-2}$, almost 48% smaller for $e=10^{-3}$ and for $e=10^{-4}$.

In the case of m_1 , the increase in velocity was given almost eight days before the collision, in




Fig. 5 Projection of orbital positions for four maneuvers performed $(m_P, m_1, m_2 \text{ and } m_3)$ for $e = 10^{-1}$ (at the top). Variation of orbital elements e and I and periapsis radius as a function of lifetime and maneuvering time (at the bottom). Initial conditions: a = 1000 km, $e = 10^{-1}$, $I = 90^{\circ}$, $\omega = \Omega = 0^{\circ}$. Simulation time equal to 52 days.

42.12 days. The value of $\Delta V = 1.84 \times 10^{-2}$ km/s is approximately 55% smaller than the value found for the case $e = 10^{-2}$, 57% smaller for $e = 10^{-3}$ and 57.6% smaller for $e = 10^{-4}$. For this transfer, the value a of the orbit was 9.994×10^2 km, $e = 1.744 \times 10^{-1}$, $I = 89.5^{\circ}$, periapsis equal to 825.07 km, apoapsis equal to 1173.7 km and maneuver time 42.12 days.

The time to perform the maneuver m_2 was 30.46 days, a little more than 21 days before the collision. The value of ΔV used to perform the orbit transfer was $\approx 9.81 \times 10^{-3}$ km/s. At the time of the maneuver, the initial orbit had a periapsis of 860.0 km, 4.44% variation about its initial value. The value of a equal to 1.0×10^3 km, $e = 1.399 \times 10^{-1}$, $I = 89.9^{\circ}$ and apoapsis of 1140.0 km, almost 4% higher than its initial value. The ΔV used in this maneuver is significantly

smaller than those found for m_2 in the previous cases. Compared to the maneuvers m_2 performed for the previous cases, this maneuver had a ΔV 71.5% and 71.4% lower compared to the cases $e = 10^{-4}$ and $e = 10^{-3}$, respectively. For $e = 10^{-2}$ this difference was 66.4% smaller.

The last maneuver (m_3) for $e = 10^{-1}$ had the smallest value of ΔV found in relation to all analyzed cases, $\approx 1.19 \times 10^{-3}$ km/s. The high value of the eccentricity makes the collision with the central body happen within a few days of integration, so the maneuver is performed within a few days of life. In this case, the transfer occurred with only 11.53 days of orbital duration. To perform this maneuver, the following initial conditions were considered, $a = 9.998 \times 10^2$ km, $e = 1.048 \times 10^{-1}$, $I = 89.5^{\circ}$, apoapsis equal to 895.0 km and an apoapsis of 1104.0 km. As mentioned, this maneuver had the lowest cost to transfer from one elliptical orbit to another. Compared to the same maneuver performed for $e = 10^{-4}$, this maneuver has a ΔV 95% smaller for $e = 10^{-3}$ this percentage is 95.3% and, 94.8% for $e = 10^{-2}$.

Making two maneuvers m_P will give a ΔV of 5.48×10^{-2} km/s and a little over 100 days of life. Making two maneuvers m_3 would give a ΔV of 2.38×10^{-3} km/s for ≈ 23 days. The cost of the maneuver m_P is much higher than m_3 , since the ΔV used in the maneuver with a few days of life is 23 times smaller. Making another analysis of m_P with another maneuver, now m_2 , we confirm that, for this case, m_3 is also more economical. To perform two maneuvers m_2 , the ΔV found is $\approx 1.962 \times 10^{-2}$ km/s to achieve a lifetime slightly greater than 60 days. This value is eight times greater than the ΔV of two maneuvers m_3 . Thus, it is concluded that the best maneuver for $e = 10^{-1}$ is m_3 .

The results discussed in this section were essential to show the best conditions for performing maneuvers involving two coplanar elliptical orbits. Our simulations showed that the most economical maneuvers are those performed with little variations of the orbital elements. In the figures where the relationship between the variations of these elements and the maneuver time is presented, it is evident that these maneuvers are the most viable.

Another essential fact taken from the results presented in this section is that the lower-cost




Fig. 6 Fuel economy analysis of an orbital maneuver through the $(\Delta V/t)$ versus lifetime relationship. Minimum value of $(\Delta V/t)$ for all analyzed cases. a) Case studied for $e=10^{-4}$. b) Case studied for $e=10^{-2}$. Initial conditions: a=1000 km, $I=90^{\circ}$, $\omega=0^{\circ}$ and $\Omega=0^{\circ}$.

maneuvers are those for final orbits with eccentricity equal to 10^{-2} and 10^{-1} . The maneuvers performed to reach a final orbit with these values of e showed low values of ΔV compared to the other maneuvers performed for final orbits with e equal to 10^{-4} and 10^{-3} . The maneuvers whose final destination was an orbit with $e = 10^{-1}$ showed a value of ΔV up to 95% lower than the maneuvers performed for the other values of e analyzed. Furthermore, it is evident that the maneuvers for final orbits with $e = 10^{-3}$ and $e = 10^{-4}$ have very close ΔV . Thus, it is more advantageous to carry out the transfer to a final orbit with $e = 10^{-3}$ since the lifetime for an orbit with this eccentricity is greater, about 45 days more for an orbit with $e = 10^{-4}$. All parameters used in performing the maneuvers discussed in this section are shown in Table 3.

Another way of showing how economical a maneuver is concerning the others is to evaluate

Table 3 Values of ΔV and initial conditions used in carrying out the maneuvers m_P , m_1 , m_2 and m_3 for final values of eccentricity $e=10^{-1}$ and m_P , m_1 , m_2 , m_3 , m_4 and m_5 for $e=10^{-2}$, $e=10^{-3}$ and $e=10^{-4}$. $\mathbf{T_m}$ is the time the maneuver was performed and $\mathbf{T_c}$ is the collision time

er	ΔV (km/s)	a _i (km)	ei	$I_i(^{\circ})$	r _{pi} (km)	r _{api} (km)	T _m (Days)	T _c (Days)	
10-1	$2.74 \times 10^{-2} \text{ (m}_P)$	9.998×10^{2}	2.098×10^{-1}	89.5	790.0	1209.5	50.83	52	
	$1.84 \times 10^{-2} \text{ (m}_1\text{)}$	9.994×10^{2}	1.744×10^{-1}	89.5	825.07	1173.7	42.12		
	$9.81 \times 10^{-3} \text{ (m}_2\text{)}$	1.0×10^{3}	1.399×10^{-1}	89.9	860.0	1140.0	30.46		
	$1.19 \times 10^{-3} \text{ (m}_3\text{)}$	9.998×10^{2}	1.048×10^{-1}	89.5	895.0	1104.0	11.53		
	$4.99 \times 10^{-2} (m_P)$	9.988×10^{2}	2.089×10^{-1}	89.4	790.0	1207.0	152.68		
	$4.10 \times 10^{-2} \text{ (m}_1\text{)}$	9.995×10^{2}	1.745×10^{-1}	88.9	825.01	1174.0	145.9	153. 2	
10^{-2}	$2.92 \times 10^{-2} \text{ (m}_2\text{)}$	9.991×10^{2}	1.389×10^{-1}	89.6	860.26	1137.90	135.0		
10 -	$2.31 \times 10^{-2} \text{ (m}_3\text{)}$	9.989×10^{2}	1.039×10^{-1}	89.5	895.07	1102.80	122.35		
	$1.45 \times 10^{-2} \text{ (m}_4\text{)}$	9.993×10^{2}	6.931×10^{-2}	89.7	930.07	1068.60	104.85		
	$6.04 \times 10^{-3} \text{ (m}_5\text{)}$	9.999×10^{2}	3.482×10^{-2}	89.8	965.14	1034.70	74.110		
	$5.23 \times 10^{-2} \text{ (m}_P)$	9.988×10^{2}	2.089×10^{-1}	87.6	790.15	1207.56	312.57		
	$4.33 \times 10^{-2} \text{ (m}_1\text{)}$	9.999×10^{2}	1.747×10^{-1}	87.9	825.18	1174.67	304.44		
10^{-3}	$3.43 \times 10^{-2} \text{ (m}_2\text{)}$	9.996×10^{2}	1.396×10^{-1}	88.0	860.10	1139.22	295.15	313	
10	$2.54 \times 10^{-2} \text{ (m}_3\text{)}$	9.99×10^{2}	1.041×10^{-1}	88.2	895.0	1103.01	282.04	313	
	$1.68 \times 10^{-2} \text{ (m}_4\text{)}$	9.998×10^{2}	6.986×10^{-2}	88.6	930.03	1069.74	264.93		
	$8.01 \times 10^{-3} \text{ (m}_5\text{)}$	9.989×10^{2}	3.388×10^{-2}	88.9	965.05	1032.76	234.19		
10-4	$5.26 \times 10^{-2} \text{ (m}_P\text{)}$	9.993×10^{2}	2.094×10^{-1}	87.7	790.0	1208.59	267.26		
	$4.34 \times 10^{-2} \text{ (m}_1\text{)}$	9.989×10^{2}	1.74×10^{-1}	88.3	825.09	1172.88	259.87		
	$3.45 \times 10^{-2} \text{ (m}_2\text{)}$	9.991×10^{2}	1.392×10^{-1}	88.5	860.01	1138.35	248.59	268,536	
	$2.56 \times 10^{-2} \text{ (m}_3\text{)}$	9.988×10^{2}	9.97×10^{-2}	88.9	895.0	1103.21	234.86	200.330	
	$1.7 \times 10^{-2} \text{ (m}_4\text{)}$	9.996×10^{2}	6.951×10^{-2}	88.8	930.16	1069.15	218.20		
	$8.31 \times 10^{-3} \text{ (m}_5\text{)}$	9.991×10^{2}	3.42×10^{-2}	89.0	965.01	1033.37	187.40		

the ΔV spent per day to perform the maneuver $(\Delta V/t)$ as a function of the lifetime. This analysis is shown in previous works such as [7] to find better points for carrying out orbital maneuvers of vehicles around natural satellites. For example, figure 6 shows this relationship as a function of a lifetime for some values of the eccentricities.

Figure 6 shows the amount of fuel per day that a maneuver requires to correct the position of an orbit. The figure also presents a minimum value of $\Delta V/t$; however, it is necessary to analyze the feasibility of this value as a function of the number of days of lifetime.

In Figure 6a, where $e = 10^{-4}$, the best values are found in a range of 20-100 days. In 25 days, a minimum value of 2.47×10^{-6} km/(s.day) is found. At this point, the ΔV is 6.19×10^{-5} km/s. With 12 maneuvers like this, we already achieve the same lifetime as the maneuver m_P performed one day before the collision and with a much smaller minimum fuel expenditure.

In this case, for $e=10^{-3}$, the orbital duration is longer, causing the variation of orbital elements to be slower. Thus, the minimum $\Delta V/t$ value is found in a range of 25-150 days, more precisely in 95 days. This value of 1.37×10^{-7} km/(s.day) is equivalent to a ΔV of 1.31×10^{-5} km/s, as in the previous case, which is the best ΔV in terms of lifetime. Only a few maneuvers with this value are necessary to increase in lifetime significantly.

In the analysis presented in Figure 6b, for $e = 10^{-2}$, the smallest values of $\Delta V/t$ are seen for a few days of integration. Due to the high eccentricity value, the minimum values of $\Delta V/t$ are around 5 to 40 days. In 20 days, we have the smallest value, 5.76×10^{-6} km/(s.day). At this point,

 ΔV is 1.15×10^{-4} km/s. Comparing the ΔV found for the maneuver m_P made days before the collision and the ΔV related to the highlighted point in Figure 6b, it is much more feasible to make several maneuvers with the minimum value than to perform the maneuver days before the collision. Making eight maneuvers of 1.15×10^{-4} km/s, we achieve the orbital duration of the maneuver m_P , using much less fuel.

For $e=10^{-1}$ in this figure, it is possible to note that $\Delta V/t$ behaves similarly to those presented in the previous cases. The minimum value found is 1.82×10^{-5} km/(s.day), in five days of integration. The related ΔV is 9.11×10^{-5} km/s, and in this case, it is also the best ΔV value for this case.

4 Maneuvers to correct the periapsis argument ω

Recently, [3, 4, 7] presented a study showing that values different from zero for the argument of periapsis (ω) and the longitude of the ascending node (Ω) are essential to increase the lifetime of an orbit. These works also present values of (ω) and (Ω) capable of prolonging the orbital lifetime. In [6, 13], maneuver models for correcting changes in the argument of periapsis are proposed. These maneuvers are essential because they ensure that this angle remains close to the values found in the regions of the most extended orbital lifetime and thus prolong the duration of the orbit.

Depending on the initial parameters considered for launching a spacecraft, the argument of periapsis may not be located in the desired geographic position, and therefore corrections become necessary.

Figure 7 shows the rotation of an ellipse by an angle θ , which also represents the variation of ω .




Fig. 7 Rotation of an elliptical orbit by an angle θ . θ is also the ω value to be corrected in the maneuver.

If it is desired to change the periapsis argument by a value θ without changing the semi-major axis a and the eccentricity e, the maneuver must be performed in the plane of the orbit. The realization of a maneuver like this is given by [14]:

$$\Delta V = 2\sqrt{\frac{\mu}{a(1-e^2)}}e\sin\left(\frac{\theta}{2}\right) \tag{19}$$

where a is the semi-major axis, e is the eccentricity, θ is the desired change in the argument of periapsis, P is the common point of the two orbits, the initial orbit and the orbit to be occupied after the maneuver, and ΔV is the total velocity to be added to the orbit to correct ω . However, it is important to note that equation 19 is only valid for constant values of a and e, which becomes ineffective for high values of these parameters.

Given the above, values of ΔV will be calculated for some cases of variation in the argument of periapsis. To effectively use equation 19, ΔV will be calculated over a time interval where variations in the semi-major axis and eccentricity are tiny. For the cases analyzed in this work, the best time to perform the periapsis argument correction maneuver would be within 50 days, after which the periapsis argument will have undergone significant changes, as well as the eccentricity [6]. Therefore, ΔV will be calculated at a point before or within 50 days and the final desired orbits will have the same eccentricity values analyzed in the previous section $(e_f = 10^{-4}, 10^{-3}, 10^{-2}, 10^{-1})$.

For an orbit with a final eccentricity of 10^{-4} , a=1000 km, $I=90^{\circ}$, and $\omega=\Omega=0^{\circ}$, the semi-major axis suffers a small variation during the whole integration, around 1 km. However, the eccentricity increases around 80 days, and then the maneuver will be performed in 50 days. When the orbit reaches this lifetime, its parameters are the semi-major axis 9.997×10^2 km, $e=9.7\times 10^{-4}$, $\theta=2.32\times 10^2$ degrees (variation of periapsis argument). Considering these parameters, a ΔV of 1.92×10^{-3} km/s is obtained.

For $e_f = 10^{-3}$, the other initial conditions are the same as for $e = 10^{-4}$. In this case, the semimajor axis also has a slight variation, around 1 km. Although the eccentricity remains very low until around 100 days of integration, the periapsis argument has a high variation, so the maneuver will be performed at 50 days. At 50 days, the orbit

has a semi-major axis of 9.99×10^2 km, an eccentricity of 8.6×10^{-4} , and θ equals 8.0 degrees. Therefore, the ΔV for this point in the orbit was 5.85×10^{-5} km/s. In the previous section, the results indicated that the most viable maneuver would be for an orbit with e equal to 10^{-3} , since orbits with this eccentricity have a longer useful life and remain with their elements with low variation for a long period. This is also shown here, as $\Delta V = 5.85 \times 10^{-5}$ km/s is a considerable value for such a maneuver.

The best interval for performing the maneuver for $e_f = 10^{-2}$ is up to 20 days. In this time, the orbit has a semi-major axis of 9.99×10^2 km, an eccentricity of 1.23×10^{-2} , θ equal to 22 degrees, and thus, $\Delta V = 2.29 \times 10^{-3}$ km/s. The initial parameters are the same as those used in the previous cases.

In the case of $e_f = 10^{-1}$, since the initial eccentricity is very high, the best interval for maneuver execution is up to 8 days. The semi-major axis and the eccentricity within this interval show little variations from their initial values, remaining almost constant. Thus, any point within this interval can have its values used in equation 19 to calculate the desired ΔV . For example, with eight days of integration, the semi-major axis is 9.99×10^2 km, the eccentricity is 1.0×10^{-1} , $\theta = 1.8^{\circ}$, and thus the ΔV required is 6.68×10^{-3} km/s.

The results presented in this section demonstrate a maneuver's feasibility to correct the periapsis argument. The ΔV_s found for different eccentricities suggest that the most viable maneuver is for an orbit with e equal to 10^{-3} , since the value found for this case is of the order of 10^{-5} km/s. In this case, it would require approximately 0.058 m/s to correct only 1° of variation in the argument of periapsis.

In addition, we have shown that, as presented in the previous section, maneuvers are more effective when performed with a short duration of the orbit.

A summary of the entire discussion in this section is presented in Table 4.

In Figure 8 we provide a representation of a maneuver with one of the best values of ΔV , its initial orbit (black line), transfer ellipse (red line) and the desired final elliptical orbit (green). Values for performing this maneuver are provided in Table 3 and in the description of the figure.

Table 4 Values of ΔV and initial conditions used in carrying out the maneuvers to correct the variation of the periapsis argument $(\theta = \omega)$. Where T_m is the time to perform the maneuver and e_f is the exectricity of the intended final orbit.

$\mathbf{e_f}$	$\Delta V (km/s)$	a _i (km)	$\mathbf{e_{i}}$	$\theta = \omega(^{\circ})$	T _m (Days)
$e = 10^{-1}$	6.68×10^{-3}	9.99×10^{2}	1.0×10^{-1}	1.8	8
$e = 10^{-2}$	2.29×10^{-3}	9.99×10^{2}	1.23×10^{-2}	22	20
$e = 10^{-3}$	5.85×10^{-5}	9.99×10^{2}	8.6×10^{-4}	8	50
$e = 10^{-4}$	1.92×10^{-3}	9.997×10^{2}	9.7×10^{-4}	232	50




Fig. 8 Representation of a maneuver using a minimum ΔV equal to 8.01×10^{-3} . Initial conditions: $a = 9.989 \times 10^{2}$ km, $e_i = 3.388 \times 10^{-2}$, $I_i = 88.9^{\circ}$.

5 Final Comments

In this work, we present a study of maneuvers for high inclination. To investigate which orbit gives a more significant variation over its orbital duration, we consider four initial inclinations 60° , 70° , 80° and 90° , with final orbit radii between 1000-2000 km.

Our results showed that orbits with an initial inclination of 70° presented a greater deflection than the other inclinations. We also observed that the best maneuvers were found for smaller orbit radii, between 1000-1050 km. We found a minimum ΔV of 5.4×10^{-05} km/s for all inclinations.

We also presented a set of simulations for maneuvers between two coplanar elliptical orbits, with final eccentricities of 10^{-4} , 10^{-3} , 10^{-2} , and 10^{-1} . We consider a few points in the orbit to compare the feasibility of one maneuver another, with six points for the values of $e = 10^{-4}$, $e = 10^{-3}$, and $e = 10^{-2}$, and three points for $e = 10^{-1}$.

This set of simulations showed that maneuvers carried out within a few days of life were more economical than those carried out near collision. These results are confirmed by analyzing the $\Delta V/t$

ratio, which showed that the velocity per maneuver time was indeed lower for orbits with a few days of duration and, consequently, with a lower variation of the orbital elements.

Finally, we present an option for a maneuver capable of correcting the argument of periapsis variation. We calculate the ΔV at the points of interest. Our results corroborate the previous results showing that maneuvers carried out within a few days of integration are more feasible than those carried out near collision.

Acknowledgments. The authors thank Improvement Coordination Higher Education Personnel - Brazil (CAPES) - Financing Code 001. The Center for Mathematical Sciences Applied to Industry (Ce-MEAI), FAPESP (Proc 2013/07375-0 and Proc 2016/23542-1). SMGW also thanks CNPq (Proc 313043/2020-5) for the financial support. ABAP thanks CNPq (Proc 309089/2021-2). This publication has been supported by the RUDN University Scientific Projects Grant System, project N° 202235-2-000.

Availability of data. All data generated or analyzed during this study are included in this published article in the form of figures.

References

- Hofstadter, M., Simon, A., Atreya, S., Banfield, D., Fortney, J.J., Hayes, A., Hedman, M., Hospodarsky, G., Mandt, K., Masters, A., Showalter, M., Soderlund, K.M., Turrini, D., Turtle, E., Reh, K., Elliott, J., Arora, N., Petropoulos, A.: Uranus and neptune missions: A study in advance of the next planetary science decadal survey. Planetary and Space Science 177, 104680 (2019). https://doi.org/10.1016/j.pss.2019.06.004
- [2] Grasset, O., Dougherty, M.K., Coustenis, A., Bunce, E., Erd, C., Titov, D.V., Blanc, M., Coates, A., Drossart, P., Fletcher, L., Hussmann, H., Jaumann, R., Krupp, N., Lebreton, J.-P., Prieto-Ballesteros, O., Tortora, P., Tosi, F., Van Hoolst, T.: Jupiter icy moons explorer (juice): An esa mission to orbit ganymede and to characterise the jupiter system. Planetary and Space Science 78, 1–21 (2013). https://doi.org/10.1016/j.pss.2012.12.002

- [3] Xavier, J., Prado, A., Winter, S., Amarante, A.: Mapping long-term natural orbits about titania, a satellite of uranus (2022)
- [4] Ferreira, T.C.F.C., Prado, A.F.B.A., Giuliatti Winter, S.M., Ferreira, L.S.: Mapping Natural Orbits around Io. Symmetry 14(7), 1478 (2022). https://doi.org/10.3390/sym14071478
- [5] Cinelli, M.: Probe lifetime around natural satellites with obliquity. Astrodynamics **6** (2022). https://doi.org/10.1007/s42064-022-0145-1
- [6] Cinelli, M., Ortore, E., Circi, C.: Long lifetime orbits for the observation of europa. Journal of Guidance, Control, and Dynamics (2019)
- [7] Ferreira, L.S., Sfair, R., Prado, A.F.B.A.: Lifetime and Dynamics of Natural Orbits around Titan. Symmetry 14(6), 1243 (2022). https://doi.org/10.3390/sym14061243
- [8] Chen, E.M.A., Nimmo, F., Glatzmaier, G.A.: Tidal heating in icy satellite oceans. Icarus 229, 11–30 (2014). https://doi.org/10.1016/ j.icarus.2013.10.024
- [9] Tzirti, S., Tsiganis, K., Varvoglis, H.: Quasicritical orbits for artificial lunar satellites. Celestial Mechanics and Dynamical Astronomy 104(3), 227–239 (2009). https://doi. org/10.1007/s10569-009-9207-4
- [10] Tzirti, S., Tsiganis, K., Varvoglis, H.: Effect of 3rd-degree gravity harmonics and Earth perturbations on lunar artificial satellite orbits. Celestial Mechanics and Dynamical Astronomy 108(4), 389–404 (2010). https: //doi.org/10.1007/s10569-010-9313-3
- [11] Paskowitz, M.E., Scheeres, D.J.: Design of Science Orbits About Planetary Satellites: Application to Europa. Journal of Guidance Control Dynamics **29**(5), 1147–1158 (2006). https://doi.org/10.2514/1.19464
- [12] Chambers, J.E.: A hybrid symplectic integrator that permits close encounters between massive bodies. MNRAS **304**(4), 793–799

- [13] Zhang, G., Zhang, H., Cao, X.: New solutions to impulsive correction for argument of perigee using gauss's variational equations. Journal of Aerospace Engineering 32, 04019071 (2019). https://doi.org/10.1061/(ASCE)AS.1943-5525.0001073
- [14] Sidi, M.J.: Spacecraft Dynamics and Control: A Practical Engineering Approach. Cambridge University Press, Cambridge (1997)