

MONOGENIC S_3 -CUBIC POLYNOMIALS AND CLASS NUMBERS OF ASSOCIATED QUADRATIC FIELDS

LENNY JONES

ABSTRACT. We say that a monic polynomial $f(x) \in \mathbb{Z}[x]$ is *monogenic* if $f(x)$ is irreducible over \mathbb{Q} and $\{1, \theta, \theta^2, \dots, \theta^{\deg f - 1}\}$ is a basis for the ring of integers of $\mathbb{Q}(\theta)$, where $f(\theta) = 0$. In this article, we use a theorem due to Kishi and Miyake to give an explicit description of an infinite set \mathcal{F} of cubic polynomials $f(x) \in \mathbb{Z}[x]$, such that $f(x)$ satisfies the three conditions: $\text{Gal}_{\mathbb{Q}}(f) \simeq S_3$, the class number of the unique quadratic subfield of the splitting field of $f(x)$ is divisible by 3 and $f(x)$ is monogenic. Moreover, if $g(x)$ is any cubic polynomial satisfying these three conditions, then there exists a unique $f(x) \in \mathcal{F}$ such that $f(x)$ and $g(x)$ generate the same cubic extension over \mathbb{Q} .

1. INTRODUCTION

We say that a monic polynomial $f(x) \in \mathbb{Z}[x]$ is *monogenic* if $f(x)$ is irreducible over \mathbb{Q} and $\{1, \theta, \theta^2, \dots, \theta^{\deg f - 1}\}$ is a basis for the ring of integers of $\mathbb{Q}(\theta)$, where $f(\theta) = 0$. Hence, $f(x)$ is monogenic if and only if $\mathbb{Z}_K = \mathbb{Z}[\theta]$. For the minimal polynomial $f(x)$ of an algebraic integer θ over \mathbb{Q} , it is well known [1] that

$$(1.1) \quad \Delta(f) = [\mathbb{Z}_K : \mathbb{Z}[\theta]]^2 \Delta(K),$$

where $\Delta(f)$ and $\Delta(K)$ are the respective discriminants over \mathbb{Q} of $f(x)$ and the number field K . Thus, from (1.1), $f(x)$ is monogenic if and only if $\Delta(f) = \Delta(K)$.

In 2000, Kishi and Miyake [5] gave the following parameterization of the quadratic fields whose class numbers are divisible by 3:

Theorem 1.1. *Let $u, w \in \mathbb{Z}$ with $\gcd(u, w) = 1$, such that $d := 4uw^3 - 27u^2$ is not a square and one of the following sets of conditions holds:*

$$(1.2) \quad 3 \nmid w;$$

$$(1.3) \quad 3 \mid w, \quad uw \not\equiv 3 \pmod{9}, \quad u \equiv w \pm 1 \pmod{9};$$

$$(1.4) \quad 3 \mid w, \quad uw \equiv 3 \pmod{9}, \quad u \equiv w \pm 1 \pmod{27}.$$

Define

$$(1.5) \quad f(x) := x^3 - uwx - u^2.$$

If $f(x)$ is irreducible over \mathbb{Q} , then the roots of $f(x) = 0$ generate an unramified cyclic cubic extension of the quadratic field $\mathbb{Q}(\sqrt{d})$. Conversely, every quadratic field K whose class number is divisible by 3 and every unramified cyclic cubic extension of K are given in this way by a suitable pair of integers u and w .

In this article, we use Theorem 1.1 to give an explicit description of an infinite set \mathcal{F} of cubic polynomials $f(x) \in \mathbb{Z}[x]$, such that $f(x)$ satisfies the three conditions:

$$(1.6) \quad \begin{aligned} & \text{(i)} \quad \text{Gal}_{\mathbb{Q}}(f) \simeq S_3, \\ & \text{(ii)} \quad 3 \mid h_K, \\ & \text{(iii)} \quad f(x) \text{ is monogenic,} \end{aligned}$$

where h_K is the class number of the unique quadratic subfield K of the splitting field of $f(x)$. Moreover, if $g(x)$ is any cubic polynomial satisfying conditions (1.6), then there exists a unique $f(x) \in \mathcal{F}$ such that $f(x)$ and $g(x)$ generate the same cubic extension over \mathbb{Q} . Since Theorem 1.1 characterizes cubic polynomials $f(x)$ satisfying conditions (i) and (ii) of (1.6), we can limit our search for cubic polynomials satisfying all three of the condition in (1.6) to the trinomials $f(x)$ as defined in (1.5) of Theorem 1.1. Our main theorem is as follows:

Theorem 1.2. *Let u, w and d be integers that satisfy the conditions set forth in Theorem 1.1, and let $f(x)$ be as defined in (1.5). Suppose that $f(x)$ is irreducible over \mathbb{Q} . Then*

$$(1.7) \quad f(x) \text{ is monogenic if and only if } u = 1 \text{ and } d \text{ is squarefree.}$$

Furthermore, if $g(x)$ is any cubic polynomial satisfying conditions (1.6), then there exists a unique trinomial in the infinite set

$$(1.8) \quad \mathcal{F} = \{f(x) : u = 1 \text{ and } d \text{ is squarefree}\}$$

such that $f(x)$ and $g(x)$ generate the same cubic extension over \mathbb{Q} .

2. PRELIMINARIES

The theorem below is an adaptation of a theorem due to Jakhar, Khanduja and Sangwan [4, Theorem 1.1], which is used to determine the monogenicity of an irreducible trinomial, to the specific cubic trinomial $f(x) = x^3 - uwx - u^2$. We use the notation $q^e \parallel W$ to mean that q^e is the exact power of the prime q that divides the integer W .

Theorem 2.1. *Let $u, w \in \mathbb{Z}$ and let $f(x) = x^3 - uwx - u^2$ be irreducible over \mathbb{Q} . Let $K = \mathbb{Q}(\theta)$, where $f(\theta) = 0$, and let \mathbb{Z}_K denote the ring of integers of K . A prime factor q of $\Delta(f) = u^3(4w^3 - 27u)$ does not divide $[\mathbb{Z}_K : \mathbb{Z}[\theta]]$ if and only if q satisfies one of the following conditions:*

- (1) when $q \mid uw$ and $q \mid u^2$, then $q^2 \nmid u^2$;
- (2) when $q \mid uw$ and $q \nmid u^2$, then

$$\text{either } q \mid A_2 \text{ and } q \nmid B_1 \quad \text{or} \quad q \nmid A_2 (u^2 A_2^3 + B_1^3),$$

$$\text{where } A_2 = -uw/q \text{ and } B_1 = \frac{-u^2 + (u^2)^{q^j}}{q} \text{ with } q^j \parallel 3;$$

- (3) when $q \nmid u^3w$, then $q^2 \nmid \Delta(f)$.

3. PROOF OF THEOREM 1.2

Proof. As previously mentioned, since $f(x) = x^3 - uwx - u^2$ is irreducible over \mathbb{Q} , we have by Theorem 1.1 that $f(x)$ satisfies conditions (i) and (ii) of (1.6). A straightforward calculation shows that

$$\Delta(f) = 4u^3w^3 - 27u^4.$$

We also note that replacing u with $-u$ and w with $-w$ in (1.5) yields the same trinomial $f(x)$. Consequently, we can assume without loss of generality that $u > 0$.

We begin with the proof of (1.7), and we assume first that $f(x)$ is monogenic. Let q be a prime divisor of $\Delta(f)$, and suppose that $f(\theta) = 0$. If $u > 1$, then condition (1) of Theorem 2.1 is not satisfied for every prime divisor q of u ; in which case, $q \mid [\mathbb{Z}_K : \mathbb{Z}[\theta]]$ and $f(x)$ is not monogenic. Thus, $u = 1$ so that

$$(3.1) \quad f(x) = x^3 - wx - 1 \quad \text{and} \quad \Delta(f) = d = 4w^3 - 27.$$

Observe then from (3.1) that $f(x)$ is irreducible over \mathbb{Q} if and only if $w \notin \{0, 2\}$ by the Rational Root Theorem.

If $q = 3$, then $3 \mid w$ from (3.1). From condition (2) of Theorem 2.1, we see that $B_1 = 0$, which implies that $3 \nmid A_2$ since $f(x)$ is monogenic. Hence, we have that $w \equiv \pm 3 \pmod{9}$. However, with $u = 1$, examining the two possibilities for w reveals that neither condition (1.3) nor condition (1.4) of Theorem 1.1 can be satisfied. For example, if $w \equiv -3 \pmod{9}$, then $uw \equiv -3 \not\equiv 3 \pmod{9}$, but $u - w \equiv 4 \not\equiv \pm 1 \pmod{9}$. Thus, $3 \nmid \Delta(f)$ and $3 \nmid w$, so that condition (1.2) of Theorem 1.1 holds. As a consequence, we have shown that no cubic trinomial $f(x)$ of the form in (3.1) is monogenic when $3 \mid w$.

Suppose next that q is a prime divisor of d such that $q^2 \mid d$. By the previous discussion, we have that $q \neq 3$ and $q \nmid w$. But then condition (3) of Theorem 2.1 is not satisfied, contradicting the fact that $f(x)$ is monogenic. Hence, d is squarefree, which completes the proof in this direction.

Conversely, suppose that $u = 1$ and d is squarefree. Let q be a prime divisor of $\Delta(f)$. Then, from (3.1), conditions (1) and (3) of Theorem 2.1 are easily seen to be satisfied. Suppose then that $q \mid w$. Then $q = 3$ from (3.1). Since $q \nmid u$, we have from condition (2) of Theorem 2.1 that $A_2 = \pm w/3$, $B_1 = 0$ and

$$A_2(u^2 A_2^3 + B_1^3) = \frac{w^4}{3^4} \not\equiv 0 \pmod{3},$$

since d is squarefree, which completes the proof of (1.7).

Next, let \mathcal{F} be the set as defined in (1.8). Then $f(x)$ and $\Delta(f)$ are as given in (3.1). By (1.7), the set \mathcal{F} consists of all cubic trinomials satisfying conditions (1.6). We claim that this set is infinite. Let $G(t) := 4t^3 - 27 \in \mathbb{Z}[t]$. It is easy to verify that $G(t)$ is irreducible over \mathbb{Q} . Then, it follows from a result of Erdős [2] that there exist infinitely many integers z such that $G(z)$ is squarefree. (In fact, there exist infinitely many primes p such that $G(p)$ is squarefree [3].) Consequently, there exist infinitely many integers w such that d is squarefree, and hence, \mathcal{F} is infinite.

Let

$$f_1(x) = x^3 - w_1x - 1 \quad \text{and} \quad f_2(x) = x^3 - w_2x - 1$$

be elements of \mathcal{F} such that $f_1(x) \neq f_2(x)$ and $K_i := \mathbb{Q}(\theta_i)$, where $f_i(\theta_i) = 0$. To establish the next part of the theorem it is enough, by Theorem 1.1, to show that $K_1 \neq K_2$. Assume, by way of contradiction, that $K_1 = K_2$. Then, since $f_i(x)$ is monogenic, we have that $\Delta(f_i) = \Delta(K_i)$, and so it follows from (1.1) that

$$(3.2) \quad 4w_1^3 - 27 = 4w_2^3 - 27.$$

Since $w_1 \neq w_2$, it is easy to see that (3.2) has no integer solutions, which completes the proof of the theorem. \square

For $|w| \leq 10$, we give in Table 1 the values of w and d for trinomials $f(x)$ in \mathcal{F} (from Theorem 1.2), and the class numbers h_K of the unique quadratic subfield $K = \mathbb{Q}(\sqrt{d})$ of the splitting field of $f(x)$.

w	-10	-7	-5	-4	-1	1	4	5	7	8	10
d	-4027	-1399	-527	-283	-31	-23	229	473	1345	2021	3973
h_K	9	27	18	3	3	3	3	3	6	3	6

TABLE 1. Values of w and d for $f(x) \in \mathcal{F}$ and class numbers of $\mathbb{Q}(\sqrt{d})$

REFERENCES

- [1] H. Cohen, *A Course in Computational Algebraic Number Theory*, Springer-Verlag, 2000.
- [2] P. Erdős, *Arithmetical properties of polynomials*, J. London Math. Soc. **28** (1953), 416–425.
- [3] H. Helfgott, *Square-free values of $f(p)$, f cubic*, Acta Math. **213** (2014), no. 1, 107–135.
- [4] A. Jakhar, S. Khanduja and N. Sangwan, *Characterization of primes dividing the index of a trinomial*, Int. J. Number Theory **13** (2017), no. 10, 2505–2514.
- [5] Y. Kishi and K. Miyake, *Parametrization of the quadratic fields whose class numbers are divisible by three*. (English summary) J. Number Theory **80** (2000), no. 2, 209–217.

PROFESSOR EMERITUS, DEPARTMENT OF MATHEMATICS, SHIPPENSBURG UNIVERSITY, SHIPPENSBURG, PENNSYLVANIA 17257, USA

Email address, Lenny Jones: doctorlennyjones@gmail.com